

# A GENERAL CORRELATION INEQUALITY FOR LEVEL SETS OF SUMS OF INDEPENDENT RANDOM VARIABLES USING THE BERNOULLI PART WITH APPLICATIONS TO THE ALMOST SURE LOCAL LIMIT THEOREM

MICHEL J. G. WEBER

**ABSTRACT.** Let  $X = \{X_j, j \geq 1\}$  be a sequence of independent, square integrable variables taking values in a common lattice  $\mathcal{L}(v_0, D) = \{v_k = v_0 + Dk, k \in \mathbb{Z}\}$ . Let  $S_n = X_1 + \dots + X_n$ ,  $S_0 = 0$ ,  $a_n = \mathbb{E} S_n$ , and  $\sigma_n^2 = \text{Var}(S_n) \rightarrow \infty$  with  $n$ . Assume that for each  $j$ ,  $\vartheta_{X_j} = \sum_{k \in \mathbb{Z}} \mathbb{P}\{X_j = v_k\} \wedge \mathbb{P}\{X_j = v_{k+1}\} > 0$ .

Using the Bernoulli part, we prove a general sharp correlation inequality extending the one we obtained in the i.i.d. case in [55]: Let  $0 < \vartheta_j \leq \vartheta_{X_j}$  and assume that  $\nu_n = \sum_{j=1}^n \vartheta_j \uparrow \infty$ ,  $n \rightarrow \infty$ . Let  $\kappa_j \in \mathcal{L}(jv_0, D)$ ,  $j = 1, 2, \dots$  be a sequence of integers such that

$$(1) \quad \frac{\kappa_j - a_j}{\sigma_j} = \mathcal{O}(1) \quad (2) \quad \text{for all } j \geq i \geq 0, \quad (\sigma_j^2 - \sigma_i^2)^{1/2} \mathbb{P}\{S_j - S_i = \kappa_j - \kappa_i\} = \mathcal{O}(1).$$

Then there exists a constant  $C$  such that for all  $1 \leq m < n$ ,

$$\begin{aligned} & \left| \sigma_n \sigma_m \left[ \mathbb{P}\{S_n = \kappa_n, S_m = \kappa_m\} - \mathbb{P}\{S_n = \kappa_n\} \mathbb{P}\{S_m = \kappa_m\} \right] \right| \\ & \leq \frac{C}{D^2} \max \left( \frac{\sigma_n}{\sqrt{\nu_n}}, \frac{\sigma_m}{\sqrt{\nu_m}} \right)^3 \left\{ \nu_n^{1/2} \prod_{j=m+1}^n \vartheta_j + \frac{\nu_n^{1/2}}{(\nu_n - \nu_m)^{3/2}} + \frac{1}{\sqrt{\frac{\nu_n}{\nu_m} - 1}} \right\}. \end{aligned}$$

We derive a nearly optimal almost sure local limit theorem: Assume that  $M(t) = \sum_{1 \leq n < t} \frac{1}{\sigma_n \sqrt{\nu_n}} \uparrow \infty$  with  $t$ . Given any  $R > 1$ , let  $\mathcal{M} = \{M_j = M(R^j), j \geq 1\}$ ,  $I_l = [R^l, R^{l+1}[$ ,  $l \geq 1$ .

We prove under moderate and simple conditions, that the series

$$\sum_{\substack{k \geq 1 \\ I_k \cap \mathcal{M} \neq \emptyset}} \frac{1}{R^k} \sup_{\substack{j \geq 1 \\ M_j \in I_k}} \left| \sum_{1 \leq n < R^j} \frac{\vartheta_n (\mathbf{1}_{\{S_n = \kappa_n\}} - \mathbb{P}\{S_n = \kappa_n\})}{\sqrt{\nu_n}} \right|$$

converges almost surely. Further if  $\lim_{n \rightarrow \infty} \sigma_n \mathbb{P}\{S_n = \kappa_n\} = \frac{D}{\sqrt{2\pi}} e^{-\frac{\kappa^2}{2}}$ ,  $M(t)$  is slowly varying near infinity, then the ASLLT holds,

$$\lim_{N \rightarrow \infty} \frac{1}{\sum_{1 \leq n < N} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}}} \sum_{1 \leq n < N} \frac{\mathbf{1}_{\{S_n = \kappa_n\}}}{\sqrt{\nu_n}} \stackrel{\text{a.s.}}{=} \frac{D}{\sqrt{2\pi}} e^{-\frac{\kappa^2}{2}}.$$

Applications are given, notably to the Cramér model.

## 1. INTRODUCTION

Throughout let  $X = \{X_i, i \geq 1\}$  denotes a sequence of independent variables taking values in a common lattice  $\mathcal{L}(v_0, D)$ , namely defined by the sequence  $v_k = v_0 + Dk$ ,

2010 *Mathematics Subject Classification*: Primary: 60F15, 60G50 ; Secondary: 60F05.

*Key words and phrases*: Local limit theorem, almost sure version, independent random variables, lattice distributed random variables, Bernoulli-part of a random variable, square integrability, correlation inequality, uniform asymptotic distribution, quasi-orthogonal system, i.i.d. random variables, Cramér model, a.s. convergent series.

$k \in \mathbb{Z}$ , where  $v_0$  and  $D > 0$  are real numbers,  $D$  is called the *span* of the lattice. Let  $S_n = X_1 + \dots + X_n$ ,  $n \geq 1$ . Then  $S_n$  takes values in the lattice  $\mathcal{L}(v_0 n, D)$ . We assume that the random variables  $X_i$  are square integrable, and that  $\sigma_n^2 = \text{Var}(S_n) \rightarrow \infty$  with  $n$ . Let also  $a_n = \mathbb{E} S_n$ , for each  $n$ . In this work, we consider the problem of estimating the probability of the level sets

$$\mathbb{P}\{S_n = N\}$$

$N \in \mathcal{L}(v_0 n, D)$ . This is obviously a quite important problem, which attracted a lot of attention. The solution, when it exists, is described by a famous limit theorem, the local limit theorem (LLT). The sequence  $X$  satisfies a *local limit theorem* (in the usual sense) if

$$(1.1) \quad \sup_{N=v_0 n + Dk} \left| \sigma_n \mathbb{P}\{S_n = N\} - \frac{D}{\sqrt{2\pi}} e^{-\frac{(N-a_n)^2}{2\sigma_n^2}} \right| = o(1), \quad n \rightarrow \infty.$$

Remark that the series

$$(1.2) \quad \sum_{N=v_0 n + Dk} \left( \mathbb{P}\{S_n = N\} - \frac{D}{\sqrt{2\pi} B_n} e^{-\frac{(N-M_n)^2}{2B_n^2}} \right),$$

is obviously convergent, whereas nothing can be deduced concerning its order from the very definition of the local limit theorem. However one can draw from Poisson summation formula that

$$(1.3) \quad \sum_{N=v_0 n + Dk} \left( \mathbb{P}\{S_n = N\} - \frac{D}{\sqrt{2\pi} B_n} e^{-\frac{(N-M_n)^2}{2B_n^2}} \right) = \mathcal{O}(D/B_n),$$

see [51], Remark 1.1.

In the iid case the probability  $\mathbb{P}\{S_n = N\}$  can be efficiently estimated by using Gnedenko's local limit theorem [23], which asserts that

$$(1.4) \quad \sup_{N \in \mathcal{L}(v_0 n, D)} \left| \sigma \sqrt{n} \mathbb{P}\{S_n = N\} - \frac{D}{\sqrt{2\pi}} e^{-\frac{(N-n\mu)^2}{2n\sigma^2}} \right| = o(1),$$

if and only if the span  $D$  is maximal, (there are no other real numbers  $v'_0$  and  $D' > D$  for which  $\mathbb{P}\{X \in \mathcal{L}(v'_0, D')\} = 1$ ). See [43] (Theorem 1 and proof of Theorem 2, p. 193–195).

The study of this fine limit theorem is intrinsically more complicated than the one of the central limit theorem; conditions on arithmetical properties of the support of a random variable are for instance always present. It was investigated by numerous authors, we only list some recent, and refer to our monograph [48] jointly written with Szewczak ( $\approx 240$  references). We cite Gamkrelidze [13, 14, 15, 16] (convergence in variation, counterexamples), Mukhin [39, 40, 41] (necessary and sufficient condition, with a recent correction in Weber [49]), Doney [10, 11, 12] (local large deviations), Breuillard [2] (diophantine measures and Edgeworth expansions), MacDonald [36, 37], Dabrowski and McDonald [5] (Bernoulli part extraction), Giuliano and Weber [21, 22] (effective rate), Macht and Wolf [44] (using Hölder-Continuity), Röllin and Ross [46] (using Landau-Kolmogorov inequalities), Jacod, Kowalski and Nikeghbali [31] (using Mod- $\phi$  convergence), Dolgopyat [8], Dolgopyat and Hafouta [9], Hafouta and Kifer [28, 29] (under tightness assumptions), and Rousseau-Egele [47], Broise [3], Calderoni, Campanino and Capocaccia [4], Gouëzel [25] (LLT for ergodic sums of expanding maps).

The study of almost sure versions of the local limit theorem is more recent, and is one motivation of this paper. This notion was introduced in 2002 by Denker and Koch in [7]

(sections 1,2), in analogy with the usual almost sure central limit theorem: “A *stationary sequence of random variables*  $\{X_n, n \geq 1\}$  *taking values in*  $\mathbb{R}$  *or*  $\mathbb{Z}$  *with partial sums*  $S_n = X_1 + \dots + X_n$  *satisfies an almost sure local limit theorem, in short ASLLT, if there exist sequences*  $\{a_n, n \geq 1\}$  *in*  $\mathbb{R}$  *and*  $\{b_n, n \geq 1\}$  *in*  $\mathbb{R}^+$  *satisfying*  $b_n \rightarrow \infty$ , *such that*

$$(1.5) \quad \lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n=1}^N \frac{b_n}{n} \chi\{S_n \in k_n + I\} \stackrel{\text{a.s.}}{=} g(\kappa)|I| \quad \text{as} \quad \frac{k_n - a_n}{b_n} \rightarrow \kappa,$$

where  $g$  denotes some density and  $I \subset \mathbb{R}$  is some bounded interval. Further  $|I|$  denotes the length of the interval  $I$  in the case where  $X_1$  is real valued and the counting measure of  $I$  otherwise.”

The above definition is however incomplete, as remarked in Weber [55], Section 4. Assume that  $\mathbb{P}\{X_i \in \mathcal{L}(v_0, D)\} = 1$ , for each  $i$ . As  $g$  is a density, there are reals  $\kappa$  such that  $g(\kappa) \neq 0$ . Now if  $\{k_n, n \geq 1\}$  is such that  $\frac{k_n - a_n}{b_n} \rightarrow \kappa$ , then any sequence  $\{\kappa_n, n \geq 1\}$ ,  $\kappa_n = k_n + u_n$  where  $u_n$  are uniformly bounded also satisfies this. But we can arrange the  $u_n$  so that  $\kappa_n \notin \mathcal{L}(v_0, D)$  for all  $n$ . Picking  $I = [-\delta, \delta]$  with  $\delta < 1/2$ , we get

$$\lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n=1}^N \frac{b_n}{n} \chi\{S_n \in k_n + I\} \stackrel{\text{a.s.}}{=} 0 \neq g(\kappa)|I|,$$

hence a contradiction. It is thus necessary to assume

$$(1.6) \quad \kappa_n \in \mathcal{L}(nv_0, D), \quad n = 1, 2, \dots,$$

also to change  $|I|$  for  $\#\{I \cap \mathcal{L}(v_0, D)\}$ . Then (1.5) modified becomes,

$$(1.7) \quad \lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n=1}^N \frac{b_n}{n} \chi\{S_n \in k_n + I\} \stackrel{\text{a.s.}}{=} g(\kappa) \#\{I \cap \mathcal{L}(v_0, D)\}, \quad \text{as} \quad \frac{k_n - a_n}{b_n} \rightarrow \kappa,$$

where  $I$  is a bounded interval.

Translating it in the independent case, we consider ( $b_n = \sigma_n$ ,  $g(x) = \frac{D}{\sqrt{2\pi}} e^{-x^2/2}$ ), this means that the ASLLT holds, by definition, if

$$(1.8) \quad \lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n=1}^N \frac{\sigma_n}{n} \chi\{S_n = k_n\} \stackrel{\text{a.s.}}{=} \frac{D}{\sqrt{2\pi}} e^{-x^2/2}, \quad \text{whenever} \quad \frac{k_n - a_n}{\sigma_n} \rightarrow \kappa.$$

However, as we shall see in Section 5, (Theorem 5.11) that formulation is not appropriate, and the right one we prove turns up to be more complicated, involving notably the sequence of parameters  $\{\vartheta_n, n \geq 1\}$ . One must nevertheless admit that the right formulation is difficult to guess, the i.i.d. case being generally weakly informative of the independent non identically distributed case. As mentioned by the authors in [7], p.146, the existence of almost sure local limit theorems is of fundamental interest. It seems reasonable to expect applications, notably at the interface with Number Theory.

The inherent second order study, which has its own interest, is much more difficult than for establishing the almost sure central limit theorem. The almost sure local limit theorems are very recent and already cover the i.i.d. case, the stable case, Markov chains, the model of the Dickman function. When the random variables are identically distributed, the ASLLT states as follows.

**Theorem 1.1.** *Let  $X$  be a square integrable random variable taking values on the lattice  $\mathcal{L}(v_0, D) = \{v_0 + kD, k \in \mathbb{Z}\}$  with maximal span  $D$ . Let  $\mu = \mathbb{E} X$ ,  $\sigma^2 = \text{Var}(X) > 0$ . Let also  $\{X_k, k \geq 1\}$  be independent copies of  $X$ , and put  $S_n = X_1 + \dots + X_n$ ,  $n \geq 1$ . Then*

$$\lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n \leq N} \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}} \stackrel{\text{a.s.}}{=} \frac{D}{\sqrt{2\pi}\sigma} e^{-\kappa^2/(2\sigma^2)},$$

for any sequence of integers  $\kappa_n \in \mathcal{L}(nv_0, D)$ ,  $n = 1, 2, \dots$  such that

$$(1.9) \quad \lim_{n \rightarrow \infty} \frac{\kappa_n - n\mu}{\sqrt{n}} = \kappa > 0.$$

Note that by (1.4), if  $\kappa_n \in \mathcal{L}(nv_0, D)$  is a sequence which verifies condition (1.9), then

$$(1.10) \quad \lim_{n \rightarrow \infty} \sqrt{n} \mathbb{P}\{S_n = \kappa_n\} = \frac{D}{\sqrt{2\pi}\sigma} e^{-\frac{\kappa^2}{2\sigma^2}}.$$

Theorem 1.1 was announced in Denker and Koch [7] (Corollary 2). The succinct proof given however contains a gap, see Weber [55]. A complete proof was given in [55]. Later, in the well-written paper [18] addressing the same problem in the stable i.i.d. case, more precisely, for specific classes of stable i.i.d. random variables, Giuliano and Szewczak, recovered that result as a particular case. One interesting aspect of the approach used in [18] is that it is based on Fourier analysis, and is thus different from ours. See also Giuliano and Szewczak [19], for a result of this kind concerning Markov chains. See Section 6 for a detailed exposition and a new improvement.

An ASLLT with rate was primarily proved in Giuliano-Weber [22].

**Theorem 1.2** ([22], Theorem 1). *Assume that  $\mathbb{E} X^{2+\varepsilon} < \infty$  for some positive  $\varepsilon$ . Then,*

$$\lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n \leq N} \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}} \stackrel{\text{a.s.}}{=} \frac{D}{\sqrt{2\pi}\sigma} e^{-\frac{\kappa^2}{2\sigma^2}},$$

for any sequence of integers  $\{\kappa_n, n \geq 1\}$  such that (1.10) holds. Moreover, if (1.9) is sharpened as follows,

$$\frac{\kappa_n - n\mu}{\sqrt{n}} = \kappa + \mathcal{O}_\eta((\log n)^{-1/2+\eta}),$$

then

$$\frac{1}{\log N} \sum_{n \leq N} \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}} \stackrel{\text{a.s.}}{=} \frac{D}{\sqrt{2\pi}\sigma} e^{-\frac{\kappa^2}{2\sigma^2}} + \mathcal{O}_\eta((\log N)^{-1/2+\eta}).$$

The proof of Theorem 1.1 mainly relies upon on sharp correlation inequalities for the associated level sets,

$$(1.11) \quad |\mathbb{P}\{S_m = k_m, S_n = k_n\} - \mathbb{P}\{S_m = k_m\}\mathbb{P}\{S_n = k_n\}|,$$

which are also established in [55], and are much harder to get than the correlation inequalities

$$(1.12) \quad |\mathbb{P}\{S_m < k_m, S_n < k_n\} - \mathbb{P}\{S_m < k_m\}\mathbb{P}\{S_n < k_n\}|,$$

needed for the proof of global a.s. central limit theorems, see e.g. Lacey and Philipp [34].

**1.1. General problem investigated.** It is a well-known in probability theory that the iid case and the independent case are always quite a different matter; one for instance can take the example of the law of the iterated logarithm ([43]). As announced, we consider the independent, non necessarily identically distributed case, and our main objective will be to establish in this very large setting a general correlation inequality, next to apply it to prove the corresponding ASLLT. In fact we will prove new general stronger forms of it, valid in a more larger setting than of the LLT, this one being not assumed, and we also show that the validity conditions we found are nearly optimal. The search of that correlation inequality has revealed new facts, which result from the investigation of a more general, and in the same time, more complex case.

**1.2. Characteristic of a random variable.** Let  $Y$  be a random variable such that  $\mathbb{P}\{Y \in \mathcal{L}(v_0, D)\} = 1$ . We do not assume that the span  $D$  is maximal. Introduce the following characteristic,

$$(1.13) \quad \vartheta_Y = \sum_{k \in \mathbb{Z}} \mathbb{P}\{Y = v_k\} \wedge \mathbb{P}\{Y = v_{k+1}\},$$

where  $a \wedge b = \min(a, b)$ . Obviously  $\vartheta_Y$  depends on  $D$ . Note that we always have the relation

$$(1.14) \quad 0 \leq \vartheta_Y < 1.$$

When  $Y$  has finite variance  $\sigma^2$ , we have the following important liaison inequality,

$$(1.15) \quad \sigma^2 \geq \frac{D^2}{4} \vartheta_Y,$$

from which it follows that (since  $X_i$  are independent,  $\text{Var}(X_1 + \dots + X_n) = \text{Var}(X_1) + \dots + \text{Var}(X_n)$ ),

$$(1.16) \quad \sigma_n^2 - \sigma_m^2 \geq \frac{D^2}{4} \sum_{j=m+1}^n \vartheta_{X_j}, \quad n \geq m \geq 1.$$

For the proofs of (1.14), (1.15), we refer the reader to the article Giuliano-Weber [22], subsection 2.1, or to the recent monograph Szewczak-Weber [48], see (1.136) and after, also on the equivalence with the “smoothness” characteristic

$$(1.17) \quad \delta_X = \sum_{m \in \mathbb{Z}} |\mathbb{P}\{X = m\} - \mathbb{P}\{X = m - 1\}|,$$

introduced by Gamkrelidze in [17].

**1.3. General Assumption.** Throughout this work we assume that

$$(1.18) \quad \vartheta_{X_j} > 0, \quad \text{for each } j \geq 1.$$

We select  $\vartheta_j$  so that

$$(1.19) \quad 0 < \vartheta_j \leq \vartheta_{X_j} \quad \text{for each } j \geq 1,$$

and we assume that

$$(1.20) \quad \nu_n := \sum_{j=1}^n \vartheta_j \uparrow \infty, \quad \text{as } n \rightarrow \infty.$$

The sequences  $\{\text{Var}(X_j), j \geq 1\}$  and  $\{\vartheta_{X_j}, j \geq 1\}$  we consider, are otherwise *arbitrary*, and are the main parameters in this study. Condition (1.18) is natural in our setting. Assume the local limit theorem (1.1) to be applicable to the sequence  $X$ . Let  $\{\kappa_n, n \geq 1\}$  be a sequence such that

$$(1.21) \quad \lim_{n \rightarrow \infty} \frac{\kappa_n - a_n}{\sigma_n} = \kappa.$$

The local limit theorem implies

$$(1.22) \quad \lim_{n \rightarrow \infty} \sigma_n \mathbb{P}\{S_n = \kappa_n\} = \frac{D}{\sqrt{2\pi}} e^{-\frac{\kappa^2}{2}}.$$

And so

$$\lim_{n \rightarrow \infty} \sigma_n \mathbb{P}\{S_n = \ell_n\} = \frac{D}{\sqrt{2\pi}} e^{-\frac{\kappa^2}{2}}, \quad (\ell_n \equiv \kappa_n \text{ or } \ell_n \equiv \kappa_n + D).$$

Then for some  $n_\kappa < \infty$ ,  $\mathbb{P}\{S_n = \kappa_n\} \wedge \mathbb{P}\{S_n = \kappa_n + 1\} > 0$  if  $n \geq n_\kappa$ . Changing  $X$  for  $X' = S_{n_\kappa}$ , we see that the new sequence  $X'$  satisfies (1.18). This was used in the course of the proof of the ASLLT in [55]. It is true in general, and has some degree of importance, at the light of assumption (1.18). The second order theory of a probabilistic system is a key element of its study. Combined with criteria of almost everywhere convergence, it allows one to prove almost sure convergence results. This is the standard approach for treating these questions. In this case here, we study the correlations properties of the system of set's indicators

$$\mathcal{T}(\kappa) = \left\{ \left( \mathbf{1}_{\{S_n = \kappa_n\}} - \mathbb{P}\{S_n = \kappa_n\} \right), \quad n \geq 1 \right\},$$

where  $\kappa = \{\kappa_n, n \geq 1\}$ , and  $\kappa_n \in \mathcal{L}(nv_0, D)$  for each  $n$ .

The general correlation inequality as well as some corollaries are stated in the next Section. The proof is technically complicated and is given in Section 4. An obvious reason is that in addition to the sequence of variances  $\text{Var}(X_j)$ , the sequence of characteristics  $\vartheta_{X_j}$  is involved in *all* estimates. In section 3, preliminary results are collected. We prove in Section 5 that for independent square integrable random variables, the almost sure local limit theorem still holds, under fairly reasonable conditions, but the proof is more involving. We also prove in Section 6, an almost sure local limit theorem with speed of convergence in the i.i.d. square integrable case, and show the almost sure convergence of tightly related random series.

*Notation.* Throughout the paper, the letter  $C$  denotes a universal constant whose value may change at each occurrence, and  $C_{\alpha, \beta, \dots}$  denotes a constant depending only on the parameters  $\alpha, \beta, \dots$ .

## 2. A GENERAL CORRELATION INEQUALITY

We prove the following result in which a new fact is that we do not assume a local limit theorem to be applicable, neither an integral limit theorem to hold.

**Theorem 2.1 (Main Result).** *Let  $\kappa_j \in \mathcal{L}(jv_0, D)$ ,  $j = 1, 2, \dots$ , be such that*

$$(1) \quad \frac{\kappa_j - a_j}{\sigma_j} = \mathcal{O}(1),$$

$$(2.1) \quad (2) \quad \text{for all } j \geq i \geq 0, \quad (\sigma_j^2 - \sigma_i^2)^{1/2} \mathbb{P}\{S_j - S_i = \kappa_j - \kappa_i\} = \mathcal{O}(1).$$

Then there exists a constant  $C$  such that for all  $1 \leq m < n$ ,

$$(2.2) \quad \begin{aligned} & \sigma_n \sigma_m \left| \mathbb{P}\{S_n = \kappa_n, S_m = \kappa_m\} - \mathbb{P}\{S_n = \kappa_n\} \mathbb{P}\{S_m = \kappa_m\} \right| \\ & \leq \frac{C}{D^2} \max\left(\frac{\sigma_n}{\sqrt{\nu_n}}, \frac{\sigma_m}{\sqrt{\nu_m}}\right)^3 \left\{ \nu_n^{1/2} \left( \prod_{j=m+1}^n \vartheta_j + \frac{1}{(\nu_n - \nu_m)^{3/2}} \right) + \frac{1}{\sqrt{\frac{\nu_n}{\nu_m} - 1}} \right\}. \end{aligned}$$

The proof is delicate and long. Some comments are in order.

**Comments 2.2.** (1) It is necessary to observe that  $\vartheta_{X_j}$  cannot be too small. This follows from the compensation effect existing between assumptions (1.19) and (1.20).

(2) Condition (1.20) implies that  $\{S_n, n \geq 1\}$  is asymptotically uniformly distributed, a.u.d. in short. See Weber [51], Th.3.3. It is known that the local limit theorem is applicable to the sequence  $\{S_n, n \geq 1\}$  only if the a.u.d. property is satisfied.

(3) It is also known that if  $X$  satisfies a local limit theorem in the strong form, it is necessary that Rozanov's condition be fulfilled, namely that

$$(2.3) \quad \sum_{k=1}^{\infty} \min_{0 \leq m < q} \mathbb{P}\{X_k \not\equiv m \pmod{q}\} = \infty, \quad \text{for all integers } q \geq 2.$$

That condition is also sufficient in some important examples, in particular if  $X_j$  have stable limit distribution, see Mitalauskas [38].

(4) Choosing  $\vartheta_j$  smaller (see (1) however) may increase the size of the correlation bound, up to some extend, and on a case-by-case basis.

(5) Although  $0 < \vartheta_{X_j} < 1$  for each  $j$ , we may have  $\vartheta_{X_j}$  arbitrary close to 1, for infinitely many  $j$ , and so assumption (2.4) below is no longer true if we pick  $\vartheta_j$  near  $\vartheta_{X_j}$ , for each  $j$  (according to previous remark). It is interesting to estimate  $\prod_{j=m+1}^n \vartheta_{X_j}$  in this case on an example.

**Example 2.3.** Let  $X_j$  be independent random variables, each defined as follows,

$$\mathbb{P}\{X_j = v_m\} = \begin{cases} 0 & \text{if } m \leq n_j \text{ or if } m > n_j + b_j, \\ \frac{1}{b_j} & \text{if } m = n_j + 1, \dots, n_j + b_j, \end{cases}$$

where  $n_j, b_j$  are positive integers,  $n_j$  can be all equal,  $b_j \rightarrow \infty$  with  $j$ , and the series  $\sum_{j \geq 1} \frac{1}{b_j}$  diverges. We have  $\vartheta_{X_j} = \sum_{k \in \mathbb{Z}} \mathbb{P}\{X_j = v_k\} \wedge \mathbb{P}\{X_j = v_{k+1}\} = \sum_{m=n_j+1}^{n_j+b_j-1} \mathbb{P}\{X_j = v_m\} \wedge \mathbb{P}\{X_j = v_{m+1}\}$ , and so  $\vartheta_{X_j} = 1 - \frac{1}{b_j}$ , for each  $j$ . Thus for some  $b > 0$ ,

$$\prod_{j=m+1}^n \vartheta_{X_j} = \prod_{j=m+1}^n \left(1 - \frac{1}{b_j}\right).$$

If for instance  $b_j = \frac{1}{j}$ , this product contributes for  $\frac{m}{n}$ . Choosing  $\vartheta_j$  sufficiently close to  $\vartheta_{X_j}$ , for each  $j$ , we have  $\prod_{j=m+1}^n \vartheta_j \approx \frac{m}{n}$ , which, for  $m = o(n)$ , is bigger than  $\frac{1}{(\nu_n - \nu_m)^{3/2}} \sim \frac{1}{(n-m)^{3/2}}$  in the right-term of inequality (2.2).

The following Corollaries are immediate consequences of Theorem 2.1.

**Corollary 2.4.** *Under assumptions of Theorem 2.1, suppose that  $\vartheta_n$  are chosen so that*

$$(2.4) \quad \tau := \sup_{j \geq 1} \vartheta_j < 1.$$

*Then we have the simplified bound,*

$$(2.5) \quad \begin{aligned} & \sigma_n \sigma_m \left| \mathbb{P}\{S_n = \kappa_n, S_m = \kappa_m\} - \mathbb{P}\{S_n = \kappa_n\} \mathbb{P}\{S_m = \kappa_m\} \right| \\ & \leq \frac{C_\tau}{D^2} \max \left( \frac{\sigma_n}{\sqrt{\nu_n}}, \frac{\sigma_m}{\sqrt{\nu_m}} \right)^3 \left\{ \frac{1}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} + \frac{\sqrt{\nu_n}}{(\nu_n - \nu_m)^{3/2}} \right\}. \end{aligned}$$

Note that we do not require the below different condition

$$(2.6) \quad \tau_X := \sup_{j \geq 1} \vartheta_{X_j} < 1,$$

to be hold. See also at this regard (4) in Comments 2.2. Condition (2.4) trivially holds in the i.i.d. case, choosing  $\vartheta = \vartheta_{X_1} > 0$ . Thus the previous estimate contains the correlation estimate in the i.i.d. square integrable case, with weaker requirements; in particular we do not assume a LLT to be applicable unlike in [55].

We state:

**Corollary 2.5 (i.i.d. case).** *Let  $X$  be a square integrable random variable taking values on the lattice  $\mathcal{L}(v_0, D) = \{v_0 + kD, k \in \mathbb{Z}\}$  with maximal span  $D$ . Let  $\mu = \mathbb{E}X$ ,  $\sigma^2 = \text{Var}(X) > 0$  and assume that  $\vartheta_X > 0$ . Let also  $\{X_k, k \geq 1\}$  be independent copies of  $X$ , and put  $S_n = X_1 + \dots + X_n$ ,  $n \geq 1$ . Let  $\kappa_j \in \mathcal{L}(jv_0, D)$ ,  $j = 1, 2, \dots$  be such that*

$$(2.7) \quad (1) \quad \frac{\kappa_j - j\mu}{\sigma\sqrt{j}} = \mathcal{O}(1), \quad (2) \quad \sigma\sqrt{j} \mathbb{P}\{S_j = \kappa_j\} = \mathcal{O}(1).$$

*Then for all  $1 \leq m < n$ ,*

$$(2.8) \quad \begin{aligned} & \sigma^2 \sqrt{nm} \left| \mathbb{P}\{S_n = \kappa_n, S_m = \kappa_m\} - \mathbb{P}\{S_n = \kappa_n\} \mathbb{P}\{S_m = \kappa_m\} \right| \\ & \leq \frac{C_\vartheta}{D^2} \left\{ \frac{n^{1/2}}{(n-m)^{3/2}} + \frac{1}{\sqrt{\frac{n}{m}} - 1} \right\}. \end{aligned}$$

Now we pass to another Corollary, which the proof is given in Section 4.

**Corollary 2.6.** *Let  $0 < c < 1$ . Under assumptions of Theorem 2.1 and (2.4), the following assertions are fulfilled:*

(1) *There exists a constant  $C_{\tau,c}$  such that for all  $1 \leq \nu_m \leq c\nu_n$ ,*

$$\begin{aligned} & \sigma_n \sigma_m \left| \mathbb{P}\{S_n = \kappa_n, S_m = \kappa_m\} - \mathbb{P}\{S_n = \kappa_n\} \mathbb{P}\{S_m = \kappa_m\} \right| \\ & \leq \frac{C_{\tau,c}}{D^2} \max \left( \frac{\sigma_n}{\sqrt{\nu_n}}, \frac{\sigma_m}{\sqrt{\nu_m}} \right)^3 \sqrt{\frac{\nu_m}{\nu_n}}. \end{aligned}$$

(2) *Further if,*

$$(2.9) \quad \sigma_n = \mathcal{O}(\sqrt{\nu_n}),$$

*then for some other constant  $C'_{\tau,c}$ , we have for all  $1 \leq \nu_m \leq c\nu_n$ ,*

$$(2.10) \quad \sigma_n \sigma_m \left| \mathbb{P}\{S_n = \kappa_n, S_m = \kappa_m\} - \mathbb{P}\{S_n = \kappa_n\} \mathbb{P}\{S_m = \kappa_m\} \right| \leq \frac{C'_{\tau,c}}{D^2} \sqrt{\frac{\nu_m}{\nu_n}}.$$



**Remark 2.7.** Assumption (2.9) implies in view of (1.16) that  $\sigma_n \asymp \sqrt{\nu_n}$ .

Thus for the case

$$(2.11) \quad \sigma_n \asymp \sqrt{\nu_n}, \quad \sup_{j \geq 1} \vartheta_j < 1.$$

estimate (2.10) holds, and provides a very handable correlation bound.

**Remark 2.8.** We indicate here a simple correlation bound, involving only the sequence  $\{\sigma_n, n \geq 1\}$ . For  $n > m \geq 1$ ,

$$\sigma_m \sigma_n \left| \mathbb{P}\{S_m = \kappa_m, S_n = \kappa_n\} - \mathbb{P}\{S_m = \kappa_m\} \mathbb{P}\{S_n = \kappa_n\} \right| \leq C \left( \frac{\sigma_n}{\sigma_{n-m}} + 1 \right).$$

The proof is elementary, see (4.5).

### 3. PRELIMINARIES

**3.1. The Bernoulli part extraction of a random variable.** Let  $X$  be a random variable such that  $\mathbb{P}\{X \in \mathcal{L}(v_0, D)\} = 1$ . We assume that

$$(3.1) \quad \vartheta_X > 0.$$

Let  $f(k) = \mathbb{P}\{X = v_k\}$ ,  $k \in \mathbb{Z}$ . Let also  $0 < \vartheta \leq \vartheta_X$ . One can associate to  $\vartheta$  and  $X$  a sequence  $\{\tau_k, k \in \mathbb{Z}\}$  of non-negative reals such that

$$(3.2) \quad \tau_{k-1} + \tau_k \leq 2f(k), \quad \sum_{k \in \mathbb{Z}} \tau_k = \vartheta.$$

Take for instance  $\tau_k = \frac{\vartheta}{\nu_X} (f(k) \wedge f(k+1))$ . Now define a pair of random variables  $(V, \varepsilon)$  as follows:

$$(3.3) \quad \begin{cases} \mathbb{P}\{(V, \varepsilon) = (v_k, 1)\} = \tau_k, \\ \mathbb{P}\{(V, \varepsilon) = (v_k, 0)\} = f(k) - \frac{\tau_{k-1} + \tau_k}{2}. \end{cases} \quad (\forall k \in \mathbb{Z})$$

By assumption this is well-defined, and the margin laws verify

$$(3.4) \quad \begin{cases} \mathbb{P}\{V = v_k\} &= f(k) + \frac{\tau_k - \tau_{k-1}}{2}, \\ \mathbb{P}\{\varepsilon = 1\} &= \vartheta = 1 - \mathbb{P}\{\varepsilon = 0\}. \end{cases}$$

Indeed,  $\mathbb{P}\{V = v_k\} = \mathbb{P}\{(V, \varepsilon) = (v_k, 1)\} + \mathbb{P}\{(V, \varepsilon) = (v_k, 0)\} = f(k) + \frac{\tau_k - \tau_{k-1}}{2}$ . Further  $\mathbb{P}\{\varepsilon = 1\} = \sum_{k \in \mathbb{Z}} \mathbb{P}\{(V, \varepsilon) = (v_k, 1)\} = \sum_{k \in \mathbb{Z}} \tau_k = \vartheta$ .

**Lemma 3.1.** *Let  $L$  be a Bernoulli random variable which is independent of  $(V, \varepsilon)$ , and put  $Z = V + \varepsilon DL$ . We have  $Z \stackrel{\mathcal{D}}{=} X$ .*

*Proof.* Plainly,

$$\begin{aligned} \mathbb{P}\{Z = v_k\} &= \mathbb{P}\{V + \varepsilon DL = v_k, \varepsilon = 1\} + \mathbb{P}\{V + \varepsilon DL = v_k, \varepsilon = 0\} \\ &= \frac{\mathbb{P}\{V = v_{k-1}, \varepsilon = 1\} + \mathbb{P}\{V = v_k, \varepsilon = 1\}}{2} + \mathbb{P}\{V = v_k, \varepsilon = 0\} \\ &= \frac{\tau_{k-1} + \tau_k}{2} + f(k) - \frac{\tau_{k-1} + \tau_k}{2} = f(k). \end{aligned}$$

□

Consider now independent random variables  $X_j, j = 1, \dots, n$ , and assume that

$$(3.5) \quad \vartheta_{X_j} > 0, \quad j = 1, \dots, n.$$

Let  $0 < \vartheta_j \leq \vartheta_{X_j}, j = 1, \dots, n$ . One can associate to them a sequence of independent vectors  $(V_j, \varepsilon_j, L_j), j = 1, \dots, n$  such that

$$(3.6) \quad \{V_j + \varepsilon_j D L_j, j = 1, \dots, n\} \stackrel{\mathcal{D}}{=} \{X_j, j = 1, \dots, n\}.$$

Further the sequences  $\{(V_j, \varepsilon_j), j = 1, \dots, n\}$  and  $\{L_j, j = 1, \dots, n\}$  are independent. For each  $j = 1, \dots, n$ , the law of  $(V_j, \varepsilon_j)$  is defined according to (3.3) with  $\vartheta = \vartheta_j$ . And  $\{L_j, j = 1, \dots, n\}$  is a sequence of independent Bernoulli random variables. Set

$$(3.7) \quad S_n = \sum_{j=1}^n X_j, \quad W_n = \sum_{j=1}^n V_j, \quad M_n = \sum_{j=1}^n \varepsilon_j L_j, \quad B_n = \sum_{j=1}^n \varepsilon_j.$$

**Proposition 3.2.** *We have*

$$\{S_k, 1 \leq k \leq n\} \stackrel{\mathcal{D}}{=} \{W_k + D M_k, 1 \leq k \leq n\}.$$

And  $M_n \stackrel{\mathcal{D}}{=} \sum_{j=1}^{B_n} L_j$ .

We also will need the following local limit theorem for Bernoulli sums.

**Proposition 3.3.** *Let  $\mathcal{B}_n = \beta_1 + \dots + \beta_n, n = 1, 2, \dots$  where  $\beta_i$  are i.i.d. Bernoulli r.v.'s ( $\mathbb{P}\{\beta_i = 0\} = \mathbb{P}\{\beta_i = 1\} = 1/2$ ). There exists a numerical constant  $C_0$  such that for all positive  $n$*

$$\sup_k \left| \mathbb{P}\{\mathcal{B}_n = k\} - \sqrt{\frac{2}{\pi n}} e^{-\frac{(2k-n)^2}{2n}} \right| \leq \frac{C_0}{n^{3/2}}.$$

**3.2. A concentration inequality for sums of independent random variables.** We also need the following Lemma ([35], Theorem 2.3).

**Lemma 3.4.** *Let  $X_1, \dots, X_n$  be independent random variables, with  $0 \leq X_k \leq 1$  for each  $k$ . Let  $S_n = \sum_{k=1}^n X_k$  and  $\mu = \mathbb{E} S_n$ . Then for any  $\epsilon > 0$ ,*

$$(a) \quad \mathbb{P}\{S_n \geq (1 + \epsilon)\mu\} \leq e^{-\frac{\epsilon^2 \mu}{2(1+\epsilon/3)}},$$

$$(b) \quad \mathbb{P}\{S_n \leq (1 - \epsilon)\mu\} \leq e^{-\frac{\epsilon^2 \mu}{2}}.$$

**3.3. An estimate for quadratic forms.** We refer to Weber [52], inequality (2.19).

**Lemma 3.5.** *For any system of complex numbers  $\{x_i\}$  and  $\{\alpha_{i,j}\}$ ,*

$$(3.8) \quad \left| \sum_{1 \leq i < j \leq n} x_i x_j \alpha_{i,j} \right| \leq \frac{1}{2} \sum_{i=1}^n |x_i|^2 \left( \sum_{i < \ell \leq n} |\alpha_{i,\ell}| + \sum_{1 \leq \ell < i} |\alpha_{\ell,i}| \right).$$

Also,

$$(3.9) \quad \left| \sum_{\substack{1 \leq i, j \leq n \\ i \neq j}} x_i x_j \alpha_{i,j} \right| \leq \frac{1}{2} \sum_{i=1}^n |x_i|^2 \left( \sum_{\substack{\ell=1 \\ \ell \neq i}}^n (|\alpha_{i,\ell}| + |\alpha_{\ell,i}|) \right).$$

In particular, if  $\alpha_{i,j} = \alpha_{j,i}$ , then

$$(3.10) \quad \left| \sum_{\substack{1 \leq i, j \leq n \\ i \neq j}} x_i x_j \alpha_{i,j} \right| \leq \sum_{i=1}^n |x_i|^2 \left( \sum_{\substack{\ell=1 \\ \ell \neq i}}^n |\alpha_{i,\ell}| \right).$$

*Proof.* We have

$$\begin{aligned}
 \left| \sum_{1 \leq i < j \leq n} x_i x_j \alpha_{i,j} \right| &\leq \sum_{1 \leq i < j \leq n} \left( \frac{|x_i|^2 + |x_j|^2}{2} \right) |\alpha_{i,j}| \\
 (3.11) \qquad \qquad \qquad &\leq \frac{1}{2} \sum_{i=1}^n |x_i|^2 \left( \sum_{i < \ell \leq n} |\alpha_{i,\ell}| + \sum_{1 \leq \ell < i} |\alpha_{\ell,i}| \right).
 \end{aligned}$$

Operating similarly for the sum  $\sum_{1 \leq j < i \leq n} x_i x_j \alpha_{i,j}$  gives the result.  $\square$

**3.4. Background on quasi-orthogonal systems.** We recall some classical facts on the notion of quasi-orthogonality in Hilbert spaces, of much relevance in the present work, and which are taken from our book [57], p. 22. Let  $(H, \|\cdot\|)$  be a real or complex Hilbert space, and let  $\{f_n, n \geq 1\}$  be orthonormal vectors in the inner product space  $H$ . The fact that  $\frac{1}{n} \sum_{i=1}^n f_i \rightarrow 0$  in  $H$  follows for instance from Ky Fan's result. Recall Bessel's inequality: *If  $\{e_i, 1 \leq i \leq n\}$  are orthogonal vectors in the inner product space  $H$ , then*

$$\sum_{i=1}^n |\langle x, e_i \rangle|^2 \leq \|x\|^2 \quad \text{for any } x \in H.$$

There are many generalizations of this inequality. In relation with Bessel's inequality, Bellman introduced the notion of a quasi-orthogonal system (see Kac, Salem and Zygmund [33]). A sequence  $\underline{f} = \{f_n, n \geq 1\}$  in  $H$  is called a quasi-orthogonal system if the quadratic form on  $\ell_2$  defined by  $\{x_h, h \geq 1\} \mapsto \|\sum_h x_h f_h\|^2$  is bounded. A necessary and sufficient condition for  $\underline{f}$  to be quasi-orthogonal is that the series  $\sum c_n f_n$  converges in  $H$ , for any sequence  $\{c_n, n \geq 1\}$  such that  $\sum c_n^2 < \infty$ .

As observed in [33], “*every theorem on orthogonal systems whose proof depends only on Bessel's inequality, holds for quasi-orthogonal systems*”.

In particular for  $H = L^2(X, \mathcal{A}, \mu)$ ,  $(X, \mathcal{A}, \mu)$  a probability space, Rademacher–Menchov's Theorem 3.6 applies, and so the series  $\sum c_n f_n$  converges almost everywhere, provided that  $\sum c_n^2 \log^2 n < \infty$ . This is easily seen from the fact that  $\underline{f}$  is quasi-orthogonal if and only if there exists a constant  $L$  depending on  $\underline{f}$  only, such that

$$(3.12) \qquad \qquad \qquad \left\| \sum_{i \leq n} y_i f_i \right\| \leq L \left( \sum_{i \leq n} |y_i|^2 \right)^{1/2}.$$

There is a useful sufficient condition for quasi-orthogonality: *In order for  $\underline{f}$  to be quasi-orthogonal, it is sufficient that*

$$(3.13) \qquad \qquad \qquad \sup_{i \geq 1} \left( \sum_{\ell \geq 1} |\langle f_i, f_\ell \rangle| \right) < \infty.$$

This is Lemma 7.4.4 in Weber [58], among other sources. Indeed, (3.10) in Lemma 3.5 implies

$$(3.14) \qquad \qquad \qquad \left| \sum_{1 \leq i, j \leq n} x_i x_j \alpha_{i,j} \right| \leq \sum_{i=1}^n |x_i|^2 \left( \sum_{1 \leq \ell \leq n} |\alpha_{i,\ell}| \right).$$

Thereby,

$$(3.15) \qquad \left\| \sum_{i \leq n} y_i f_i \right\|^2 \leq \sum_{i=1}^n |y_i|^2 \left( \sum_{1 \leq \ell \leq n} |\langle f_i, f_\ell \rangle| \right) \leq \left( \sup_{i \geq 1} \sum_{\ell \geq 1} |\langle f_i, f_\ell \rangle| \right) \cdot \sum_{i=1}^n |y_i|^2.$$

Thus if (3.13) is fulfilled, (3.12) readily follows from the previous calculation.

**3.5. Rademacher–Menshov’s theorem.** This well-known result states as follows.

**Theorem 3.6.** a) *Let  $\{a_k, k \geq 1\}$  be a sequence of reals such that*

$$\sum_{k \geq 1} |a_k|_2^2 \log^2 k < \infty.$$

*Then for any orthonormal sequence  $\{\xi_k, k \geq 1\}$  in  $L^2(\mathbb{P})$ ,  $(\Omega, \mathcal{B}, \mathbb{P})$  a probability space, the series  $\sum_{k \geq 1} a_k \xi_k$  converges  $\mathbb{P}$ -almost surely.*

b) *If  $\{w(n), n \geq 1\}$  is an arbitrary positive monotone increasing sequence of numbers with  $w(n) = o(\log n)$ , then there exists an everywhere divergent orthonormal series  $\sum_{k \geq 1} a_k \psi_k$  whose coefficients satisfy the condition*

$$\sum_{k \geq 1} c_k^2 w(k)^2 = \infty.$$

We refer to Alexits [1], (see p.80), Kashin and Saakyan [32], Olevskii [42] concerning orthogonal systems in Fourier analysis, and Weber [57], [58]. Less known, and less applied, are the refinements of this result, obtained notably by Tandori, see [32], and those obtained using the majorizing measure method, which are nevertheless of a self-evident practical interest in the study of limit theorems in Probability Theory. There are many different proofs available. See also, more specifically Weber [56] and Ch. 7 in [58], and references therein.

**3.6. A useful almost sure convergence criterion.** For the proof of the ASLLT (Theorem 5.11), the following almost sure convergence criterion is of particular relevance. This one has the advantage, comparatively to other known criteria, to allow one to control convergence of series of random variables inherent to questions of this kind.

**Theorem 3.7.** *Let  $\xi = \{\xi_l, l \geq 1\}$  be a sequence of real random variables with partial sums  $S_n = \sum_{l=1}^n \xi_l$ ,  $n \geq 1$ . Assume that the following assumption is satisfied:*

*For some  $\gamma > 1$ ,*

$$(3.16) \quad \mathbb{E} \left| \sum_{i \leq l \leq j} \xi_l \right|^\gamma \leq \Phi \left( \sum_{l \leq j} u_l \right) \Psi \left( \sum_{i \leq l \leq j} u_l \right) \quad (\forall 1 \leq i \leq j < \infty),$$

*where  $\Phi, \Psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  are non-decreasing,  $\Psi(x)/x^{1+\eta}$  is non-decreasing for some  $\eta \geq 0$  and  $u = \{u_i, i \geq 1\}$  is a sequence of non negative reals such that  $U_j = \sum_{l=1}^j u_l \uparrow \infty$ . Further assume that for some real  $M > 1$ , the series*

$$(3.17) \quad s^\gamma = s_M^\gamma = \sum_{\substack{l \geq 1 \\ [M^l, M^{l+1}[ \cap \mathcal{U} \neq \emptyset}} \frac{\Psi(M^l) \Phi(M^l) L(l)^{\varepsilon(\eta)}}{M^{\gamma l}}$$

*converges, where  $\mathcal{U} = \{U_j, j \geq 1\}$ ,  $I(l) = I_M(l) = [M^l, M^{l+1}[$ ,  $L(l) = L_M(l) = 1 + \log \left( \# \{ [M^l, M^{l+1}[ \cap \mathcal{U} \} \right)$ ,  $l \geq 1$ , and  $\varepsilon(\eta) = 0$  or  $1$ , according to  $\eta > 0$  or  $\eta = 0$ .*

*Put,*

$$(3.18) \quad \mathcal{S}^\gamma = \mathcal{S}_M^\gamma = \sum_{\substack{k \geq 1 \\ [M^k, M^{k+1}[ \cap \mathcal{U} \neq \emptyset}} \sup_{M^k \leq U_j < M^{k+1}} \left( \frac{|S_j|}{U_j} \right)^\gamma.$$

Then,

$$(3.19) \quad \|\mathcal{S}_M\|_\gamma \leq MK_\gamma s, \quad \text{and in particular} \quad \mathbb{P}\left\{\lim_{j \rightarrow \infty} \frac{S_j}{U_j} = 0\right\} = 1,$$

where  $K_\gamma$  is a constant depending on  $\gamma$  only.

Let  $1 < M_1 \leq M$ . Then

$$(3.20) \quad \|\mathcal{S}_{M_1}\|_\gamma^\gamma \leq 2^\gamma \|\mathcal{S}_M\|_\gamma^\gamma.$$

This is a slight variant of the criterion given in Weber [54, Th. 8.2] generalizing Gál-Koksma's, in which the reals  $u_i$  are assumed to be integers.

Several remarks are in order.

**Remark 3.8.** The above Theorem provides with (3.19), fine estimates of  $|S_j|/U_j$ . Functions  $\Phi, \Psi$  which are present in assumption (3.16), appear in the course of the proof where increment's assumptions are combined with Lemma 3.12 to control intermediate local maxima, but not in (3.18). However assumption (3.17), which is used to achieve the proof, completely relies on these functions, and appear in the first part of (3.19).

**Remark 3.9.** It is quite interesting to observe that the bound obtained *tightly depend* on the way the sequence  $\mathcal{U}$  is asymptotically distributed, which is reflected by (3.17). Indeed, only those  $k$  such that  $[M^k, M^{k+1}[\cap \mathcal{U} \neq \emptyset$  have to be counted. This seems to have been overlooked, in particular by the author. This remark also applies to similar criteria.

**Remark 3.10.** An analog bound can be derived from the proof for partially observed sequences  $\{S_n, n \in \mathcal{N}\}$ ,  $\mathcal{N}$  a growing sequence of integers.

**Remark 3.11** (bounded case). When  $\mathcal{U} = \{U_i, i \geq 1\}$  is a bounded sequence, assumption (3.16) takes the simpler form: for some  $\gamma > 1$ ,

$$(3.21) \quad \mathbb{E} \left| \sum_{i \leq l \leq j} \xi_l \right|^\gamma \leq C \Psi \left( \sum_{i \leq l \leq j} u_l \right) \quad (\forall 1 \leq i \leq j < \infty).$$

This readily implies that  $S_n = \sum_{l=1}^n \xi_l$ ,  $n \geq 1$  is a Cauchy sequence in  $L^\gamma(\mathbb{P})$ , thus converging to some element  $S \in L^\gamma(\mathbb{P})$ . Further, assuming a little more than the convergence of the series  $\sum_{l=1}^\infty m_l$ , the series  $\sum_{l=1}^\infty \xi_l$  also converges almost surely. More precisely we have the implication,

$$(3.22) \quad \sum_{l=1}^\infty u_l (\log l)^\gamma < \infty \implies \text{the series } \sum_{l=1}^\infty \xi_l \text{ converges almost surely.}$$

The case  $\gamma = 2$  contains Rademacher-Menshov's Theorem. See Remark 8.3.5 and (8.3.27) in Weber [57], and Chapter 8 for detailed study of these questions. See also Remark ?? where this is applied.

For the sake of completeness, we give a detailed proof of Theorem 3.7.

*Proof.* Let  $\kappa = \kappa(M) = \{\kappa_p(M), p \geq 1\}$  be the sequence defined by  $\kappa_p = \kappa_p(M) = k$ , if  $I_k$  is the  $p$ -th interval such that  $I_k \cap \mathcal{U} \neq \emptyset$ . Let  $\mathcal{L}_p = \mathcal{L}_p(M)$  be the set of indices defined by  $L \in \mathcal{L}_p \Leftrightarrow U_L \in I_{\kappa_p}$ . Pick arbitrarily some index in  $\mathcal{L}_p$ , which we write  $L_p^* = L_p^*(M)$ .

On the one hand

$$(3.23) \quad \sum_p \frac{\|S_{L_p^*}\|_\gamma^\gamma}{M^{\gamma(\kappa_p+1)}} \leq \sum_p \frac{\Psi(M^{\kappa_p+1})\Phi(M^{\kappa_p+1})}{M^{\gamma(\kappa_p+1)}}.$$

And on the other, for  $i, j \in \mathcal{L}_p$ ,  $i \leq j$ ,

$$(3.24) \quad \mathbb{E} \frac{|S_j - S_i|^\gamma}{\Phi(M^{\kappa_p+1})} \leq \frac{\Psi(U_j - U_i)}{(U_j - U_i)^{1+\eta}} (U_j - U_i)^{1+\eta} \leq \frac{\Psi(M^{\kappa_p+1})}{(M^{\kappa_p+1})^{1+\eta}} (U_j - U_i)^{1+\eta}.$$

Thus

$$(3.25) \quad \mathbb{E} |S_j - S_i|^\gamma \leq \Phi(M^{\kappa_p+1}) \Psi(M^{\kappa_p+1}) \left( \frac{U_j}{M^{\kappa_p+1}} - \frac{U_i}{M^{\kappa_p+1}} \right)^{1+\eta}.$$

We use the following Lemma based on metric entropy chaining.

**Lemma 3.12** ([54], Lemma 3.4). *Let  $\gamma > 1$ ,  $0 < \beta \leq 1$  and consider a finite collection of random variables  $(X_1, \dots, X_N) \subset L^\gamma(\mathbb{P})$ , and reals  $0 \leq t_1 \leq t_2 \leq \dots \leq t_N \leq 1$  such that*

$$(3.26) \quad \|X_j - X_i\|_\gamma \leq (t_j - t_i)^\beta \quad (\forall 1 \leq i \leq j \leq N).$$

*Then, there exists a constant  $K_{\beta, \gamma}$  depending on  $\beta, \gamma$  only, such that*

$$(3.27) \quad \left\| \sup_{1 \leq i, j \leq N} |X_i - X_j| \right\|_\gamma \leq \begin{cases} K_{\beta, \gamma} & \text{if } \beta\gamma > 1, \\ K_{\beta, \gamma} \log N & \text{if } \beta\gamma = 1, \\ K_{\beta, \gamma} N^{\frac{1}{\gamma} - \beta} & \text{if } \beta\gamma < 1. \end{cases}$$

We deduce from (3.25) and Lemma 3.12,

$$(3.28) \quad \left\| \sup_{i, j \in \mathcal{L}_p} |S_i - S_j| \right\|_\gamma \leq \begin{cases} K_{\eta, \gamma} \Phi(M^{\kappa_p+1})^{1/\gamma} \Psi(M^{\kappa_p+1})^{1/\gamma} & \text{if } \eta > 0, \\ K_{\eta, \gamma} \Phi(M^{\kappa_p+1})^{1/\gamma} \Psi(M^{\kappa_p+1})^{1/\gamma} \log \#(\mathcal{L}_p) & \text{if } \eta = 0, \end{cases}$$

where  $K_{\eta, \gamma}$  depend on  $\eta, \gamma$  only.

Assume that  $\eta = 0$ . Then

$$(3.29) \quad \sum_p \frac{\left\| \sup_{i, j \in \mathcal{L}_p} |S_i - S_j| \right\|_\gamma^\gamma}{M^{\gamma(\kappa_p+1)}} \leq K_\gamma \sum_p \frac{\Phi(M^{\kappa_p+1}) \Psi(M^{\kappa_p+1}) (1 + \log \#(\mathcal{L}_p))}{M^{\gamma(\kappa_p+1)}}.$$

From (3.23) and (3.29) and the elementary inequality  $(a+b)^y \leq 2^{y-1}(a^y + b^y)$ ,  $a \geq 0, b \geq 0$  and  $y \geq 1$  (which is a plain application of Hölder's inequality), we deduce that,

$$(3.30) \quad \sum_p \frac{\left\| \sup_{j \in \mathcal{L}_p} |S_j| \right\|_\gamma^\gamma}{M^{\gamma(\kappa_p+1)}} \leq K_\gamma \sum_p \frac{\Phi(M^{\kappa_p+1}) \Psi(M^{\kappa_p+1}) (1 + \log \#(\mathcal{L}_p))}{M^{\gamma(\kappa_p+1)}}.$$

Observing for  $L \in \mathcal{L}_p$  that  $M^{\kappa_p+1} \leq MU_L$ , we obtain, in view of the definition of  $\mathcal{L}_p$ , the following bound,

$$\begin{aligned} \|S_M\|_\gamma^\gamma &= \sum_{\substack{k \geq 1 \\ [M^k, M^{k+1}] \cap \mathcal{U} \neq \emptyset}} \left\| \sup_{M^k \leq U_j < M^{k+1}} \left( \frac{|S_j|}{U_j} \right) \right\|_\gamma^\gamma \\ &= \sum_p \left\| \sup_{j \in \mathcal{L}_p} \left( \frac{|S_j|}{U_j} \right) \right\|_\gamma^\gamma \leq M \sum_p \frac{\left\| \sup_{j \in \mathcal{L}_p} |S_j| \right\|_\gamma^\gamma}{M^{\gamma(\kappa_p+1)}} \\ &\leq M K_\gamma \sum_p \Phi(M^{\kappa_p+1}) \Psi(M^{\kappa_p+1}) (1 + \log \#(\mathcal{L}_p)) / M^{\gamma(\kappa_p+1)} \\ &\leq M K_\gamma s^\gamma. \end{aligned}$$

Further  $\sup_{j \in \mathcal{L}_p} |S_j|/M^{\kappa_p}$  tends to 0 almost surely, as  $p$  tends to infinity. And so (3.19) follows. The case  $\eta > 0$  is identical.

Let  $1 < M_1 \leq M$ . For any integer  $k \geq 1$ , there exists an integer  $l \geq 1$  such that  $[M_1^k, M_1^{k+1}[ \subset [M^l, M^{l+2}[$ . Moreover if  $[M_1^k, M_1^{k+1}[ \cap \mathcal{U} \neq \emptyset$ , then  $[M^l, M^{l+1}[ \cap \mathcal{U} \neq \emptyset$  and/or  $[M^{l+1}, M^{l+2}[ \cap \mathcal{U} \neq \emptyset$ .

From the following inequality,

$$(3.31) \quad \left\| \sup_{M_1^k \leq U_j < M_1^{k+1}} \left( \frac{|S_j|}{U_j} \right) \right\|_\gamma^\gamma \leq \left\{ \left\| \sup_{M^l \leq U_j < M^{l+1}} \left( \frac{|S_j|}{U_j} \right) \right\|_\gamma + \left\| \sup_{M^{l+1} \leq U_j < M^{l+2}} \left( \frac{|S_j|}{U_j} \right) \right\|_\gamma \right\}^\gamma \\ \leq 2^{\gamma-1} \left\{ \left\| \sup_{M^l \leq U_j < M^{l+1}} \left( \frac{|S_j|}{U_j} \right) \right\|_\gamma^\gamma + \left\| \sup_{M^{l+1} \leq U_j < M^{l+2}} \left( \frac{|S_j|}{U_j} \right) \right\|_\gamma^\gamma \right\},$$

and definition of  $\mathcal{S}_M$ , it follows that

$$(3.32) \quad \|\mathcal{S}_{M_1}\|_\gamma^\gamma \leq 2^\gamma \|\mathcal{S}_M\|_\gamma^\gamma.$$

This implies by the same argument used before that  $\sup_{j \in \mathcal{L}_p} |S_j|/M_1^{\kappa'_p}$  tends to 0 almost surely, as  $p$  tends to infinity,  $\kappa'_p = \kappa_p(M_1)$  being defined with respect to the sequence  $\{I_{M_1}(l), l \geq 1\}$  and  $\mathcal{U}$ . This achieves the proof.  $\square$

#### 4. PROOFS OF THEOREM 2.1 AND OF COROLLARY 2.6

Assumption (1.18) implies that condition (3.5) is realized. Let  $0 < \vartheta_j \leq \vartheta_{X_j}$ . By Proposition 3.2, we may associate to  $\{X_k, k \geq 1\}$  a sequence  $\{(V_j, \varepsilon_j, L_j), j \geq 1\}$  of independent copies of  $(V, \varepsilon, L)$  such that

$$\{V_j + \varepsilon_j D L_j, j \geq 1\} \stackrel{\mathcal{D}}{=} \{X_j, j \geq 1\}.$$

Further  $\{(V_j, \varepsilon_j), j \geq 1\}$  and  $\{L_j, j \geq 1\}$  are independent sequences. And  $\{L_j, j \geq 1\}$  is a sequence of independent Bernoulli random variables. Set

$$(4.1) \quad S_n = \sum_{j=1}^n X_j, \quad W_n = \sum_{j=1}^n V_j, \quad M_n = \sum_{j=1}^n \varepsilon_j L_j, \quad B_n = \sum_{j=1}^n \varepsilon_j.$$

We notice that  $M_n$  is a sum of exactly  $B_n$  Bernoulli random variables. We have the representation

$$\{S_n, n \geq 1\} \stackrel{\mathcal{D}}{=} \{W_n + D M_n, n \geq 1\}.$$

And  $M_n \stackrel{\mathcal{D}}{=} \sum_{j=1}^{B_n} L_j$ .

We denote by  $\mathbb{E}_{(V, \varepsilon)}$ ,  $\mathbb{P}_{(V, \varepsilon)}$  (resp.  $\mathbb{E}_L$ ,  $\mathbb{P}_L$ ) the expectation and probability symbols relatively to the  $\sigma$ -algebra generated by the sequence  $\{(V_j, \varepsilon_j), j = 1, \dots, n\}$  (resp.  $\{L_j, j = 1, \dots, n\}$ ). These algebra are independent.

Put

$$(4.2) \quad Y_n = \sigma_n (\mathbf{1}_{\{S_n = \kappa_n\}} - \mathbb{P}\{S_n = \kappa_n\}).$$

Let  $n > m \geq 1$ . We have a simple expression for the correlation

$$(4.3) \quad \sigma_m \sigma_n \left( \mathbb{P}\{S_m = \kappa_m, S_n = \kappa_n\} - \mathbb{P}\{S_m = \kappa_m\} \mathbb{P}\{S_n = \kappa_n\} \right) = \mathbb{E} Y_n Y_m,$$

which we can more conveniently rewrite as follows,

$$(4.4) \quad \mathbb{E} Y_n Y_m = \sigma_m \mathbb{P}\{S_m = \kappa_m\} \sigma_n \left( \mathbb{P}\{S_n - S_m = \kappa_n - \kappa_m\} - \mathbb{P}\{S_n = \kappa_n\} \right).$$

Since by (2.1)-(2),

$$(\sigma_j^2 - \sigma_i^2)^{1/2} \mathbb{P}\{S_j - S_i = \kappa_j - \kappa_i\} = \mathcal{O}(1),$$

we have the bound

$$\begin{aligned} & \sigma_m \mathbb{P}\{S_m = \kappa_m\} \sigma_n \left| \mathbb{P}\{S_n - S_m = \kappa_n - \kappa_m\} - \mathbb{P}\{S_n = \kappa_n\} \right| \\ & \leq C \left( \frac{\sigma_n}{(\sigma_n^2 - \sigma_m^2)^{1/2}} \right) (\sigma_n^2 - \sigma_m^2)^{1/2} \mathbb{P}\{S_n - S_m = \kappa_n - \kappa_m\} + \sigma_n \mathbb{P}\{S_n = \kappa_n\} \leq C \left( \frac{\sigma_n}{(\sigma_n^2 - \sigma_m^2)^{1/2}} + 1 \right). \end{aligned}$$

Whence for  $n > m \geq 1$ ,

$$(4.5) \quad |\mathbb{E} Y_n Y_m| \leq C \left( \frac{\sigma_n}{(\sigma_n^2 - \sigma_m^2)^{1/2}} + 1 \right).$$

We therefore have to estimate  $|A|$  where

$$\begin{aligned} (4.6) \quad A &:= \sigma_n \left( \mathbb{P}\{S_n - S_m = \kappa_n - \kappa_m\} - \mathbb{P}\{S_n = \kappa_n\} \right) \\ &= \sigma_n \mathbb{E} \left( \mathbf{1}_{\{B_n \leq \theta \nu_n\}} + \mathbf{1}_{\{B_n > \theta \nu_n\}} \right) \left( \mathbf{1}_{\{S_n - S_m = \kappa_n - \kappa_m\}} - \mathbf{1}_{\{S_n = \kappa_n\}} \right). \end{aligned}$$

Further,

$$(4.7) \quad \mathbb{E} Y_n^2 = \sigma_n^2 \mathbb{P}\{S_n = \kappa_n\} (1 - \mathbb{P}\{S_n = \kappa_n\}) = \mathcal{O}(\sigma_n).$$

Let  $0 < \varepsilon < 1$  and set  $\rho = e^{-\frac{\varepsilon^2}{2}}$ ,  $\theta = 1 - \varepsilon$ . By Lemma 3.4,

$$(4.8) \quad \mathbb{P}\{B_n \leq \theta \nu_n\} \leq \rho^{\nu_n}, \quad \mathbb{P}\{B_n - B_m \leq \theta(\nu_n - \nu_m)\} \leq \rho^{\nu_n - \nu_m},$$

for all  $n \geq m \geq 1$ .

We can write in view of Proposition 3.2

$$\begin{aligned} (4.9) \quad & \sigma_n \mathbb{E} \mathbf{1}_{\{B_n > \theta \nu_n\}} \left( \mathbf{1}_{\{S_n - S_m = \kappa_n - \kappa_m\}} - \mathbf{1}_{\{S_n = \kappa_n\}} \right) \\ &= \sigma_n \mathbb{E}_{(V, \varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n\}} \left( \mathbb{P}_L \left\{ D \sum_{j=m+1}^n \varepsilon_j L_j = \kappa_n - \kappa_m - (W_n - W_m) \right\} \right. \\ & \quad \left. - \mathbb{P}_L \left\{ D \sum_{j=1}^n \varepsilon_j L_j = \kappa_n - W_n \right\} \right). \end{aligned}$$

To treat the independent case, it is necessary to introduce the sets

$$A(n, m) = \{\varepsilon_j = 0, \quad m < j \leq n\}.$$

On  $A(n, m)$  we have  $\sum_{j=m+1}^n \varepsilon_j L_j = 0$ . Thus

$$\left\{ D \sum_{j=m+1}^n \varepsilon_j L_j = \kappa_n - \kappa_m - (W_n - W_m) \right\} = \left\{ W_n - W_m = \kappa_n - \kappa_m \right\}.$$

So that (4.9) may be continued with

$$\begin{aligned} &= \sigma_n \left\{ \mathbb{E}_{(V, \varepsilon)} \mathbf{1}_{\{(B_n > \theta \nu_n) \cap A(n, m)\}} \left( \mathbf{1}_{\{W_n - W_m = \kappa_n - \kappa_m\}} \right. \right. \\ & \quad \left. \left. - \mathbb{P}_L \left\{ D \sum_{j=1}^n \varepsilon_j L_j = \kappa_n - W_n \right\} \right) \right\} \end{aligned}$$



$$\begin{aligned}
& + \sigma_n \left\{ \mathbb{E}_{(V, \varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n\} \cap A(n, m)^c} \left[ \mathbb{P}_L \left\{ D \sum_{j=m+1}^n \varepsilon_j L_j = \kappa_n - \kappa_m - \right. \right. \right. \\
& \quad \left. \left. \left. (W_n - W_m) \right\} - \mathbb{P}_L \left\{ D \sum_{j=1}^n \varepsilon_j L_j = \kappa_n - W_n \right\} \right] \right\} \\
(4.10) \quad & := A' + A''.
\end{aligned}$$

We have the easy bound

$$(4.11) \quad |A'| \leq \sigma_n \mathbb{P}\{A(n, m)\} = \sigma_n \prod_{j=m+1}^n \vartheta_j.$$

Concerning  $A''$ , we note that

$$\sum_{j=1}^n \varepsilon_j L_j \stackrel{\mathcal{D}}{=} \sum_{j=1}^{B_n} L_j, \quad \sum_{j=m+1}^n \varepsilon_j L_j \stackrel{\mathcal{D}}{=} \sum_{j=B_m+1}^{B_n} L_j.$$

By applying Proposition 3.3 we obtain,

$$(4.12) \quad \sup_z \left| \sqrt{N} \mathbb{P}\left\{ \sum_{j=1}^N L_j = z \right\} - \frac{2}{\sqrt{2\pi}} e^{-\frac{(z - (N/2))^2}{(N/2)}} \right| = o\left(\frac{1}{N}\right).$$

It follows that

$$\left| \mathbb{P}_L \left\{ D \sum_{j=1}^{B_n} L_j = \kappa_n - W_n \right\} - \frac{2e^{-\frac{(\kappa_n - W_n - (B_n/2))^2}{D^2(B_n/2)}}}{\sqrt{2\pi B_n}} \right| = o\left(\frac{1}{B_n^{3/2}}\right).$$

Further since on the set  $A(n, m)^c$ ,  $\varepsilon_j = 1$  for some  $m < j \leq n$ , we have  $B_n > B_m$ , then

$$\begin{aligned}
& \left| \mathbb{P}_L \left\{ D \sum_{j=1}^{B_n - B_m} L_j = \kappa_n - \kappa_m - (W_n - W_m) \right\} - \frac{2e^{-\frac{(\kappa_n - \kappa_m - (W_n - W_m) - (B_n - B_m)/2)^2}{D^2(B_n - B_m)/2}}}{\sqrt{2\pi(B_n - B_m)}} \right| \\
& = o\left(\frac{1}{(B_n - B_m)^{3/2}}\right).
\end{aligned}$$

Therefore,

$$\begin{aligned}
|A''| & \leq \sigma_n \left| \mathbb{E}_{(V, \varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n\} \cap A(n, m)^c} \left\{ \frac{2e^{-\frac{(\kappa_n - W_n - (B_n/2))^2}{D^2(B_n/2)}}}{\sqrt{2\pi B_n}} \right. \right. \\
& \quad \left. \left. - \frac{2e^{-\frac{(\kappa_n - \kappa_m - (W_n - W_m) - (B_n - B_m)/2)^2}{D^2(B_n - B_m)/2}}}{\sqrt{2\pi(B_n - B_m)}} \right\} \right| \\
& \quad + C_0 \sigma_n \mathbb{E}_{(V, \varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n, B_n > B_m\}} \left\{ \frac{1}{B_n^{3/2}} + \left( \frac{1}{(B_n - B_m)^{3/2}} \right) \right\} \\
(4.13) \quad & := A''_1 + C_0 A''_2,
\end{aligned}$$

where the constant  $C_0$  comes from (4.12). The second term is easily estimated. Indeed,

$$\begin{aligned}
A''_2 & = \sigma_n \mathbb{E}_{(V, \varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n, B_n > B_m\}} \left( \frac{1}{B_n^{3/2}} + \frac{1}{(B_n - B_m)^{3/2}} \right) \\
& \leq 2\sigma_n \mathbb{P}\{B_n - B_m \leq \theta(\nu_n - \nu_m)\}
\end{aligned}$$

$$\begin{aligned}
& + \sigma_n \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n, B_n - B_m > \theta(\nu_n - \nu_m)\}} \left( \frac{1}{B_n^{3/2}} + \frac{1}{(B_n - B_m)^{3/2}} \right) \\
(4.14) \quad & \leq C \sigma_n \left\{ \rho^{\nu_n - \nu_m} + \frac{1}{(\theta \nu_n)^{3/2}} + \frac{1}{(\theta(\nu_n - \nu_m))^{3/2}} \right\}.
\end{aligned}$$

We now estimate  $A_1''$ , which we bound as follows:

$$\begin{aligned}
A_1'' &= \sigma_n \left| \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{(B_n > \theta \nu_n) \cap A(n,m)^c\}} \left\{ \frac{2e^{-\frac{(\kappa_n - W_n - (B_n/2))^2}{D^2(B_n/2)}}}{\sqrt{2\pi B_n}} \right. \right. \\
&\quad \left. \left. - \frac{2e^{-\frac{(\kappa_n - \kappa_m - (W_n - W_m) - (B_n - B_m)/2)^2}{D^2(B_n - B_m)/2}}}{\sqrt{2\pi(B_n - B_m)}} \right\} \right| \\
&\leq C \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{(B_n > \theta \nu_n) \cap A(n,m)^c\}} \left\{ \left( \frac{\sigma_n}{\sqrt{B_n}} \right) \left[ \sqrt{\frac{B_n}{B_n - B_m}} - 1 \right] \right. \\
&\quad \left. \times e^{-\frac{(\kappa_n - \kappa_m - (W_n - W_m) - (B_n - B_m)/2)^2}{D^2(B_n - B_m)/2}} \right\} \\
&\quad + C \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{(B_n > \theta \nu_n) \cap A(n,m)^c\}} \left( \frac{\sigma_n}{\sqrt{B_n}} \right) \\
&\quad \times \left| e^{-\frac{(\kappa_n - W_n - (B_n/2))^2}{D^2(B_n/2)}} - e^{-\frac{(\kappa_n - \kappa_m - (W_n - W_m) - (B_n - B_m)/2)^2}{D^2(B_n - B_m)/2}} \right| \\
&\leq C_\theta \left( \frac{\sigma_n}{\sqrt{\nu_n}} \right) \left\{ \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n, B_n > B_m\}} \left[ \sqrt{\frac{B_n}{B_n - B_m}} - 1 \right] \right. \\
&\quad + \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n, B_n > B_m\}} \\
&\quad \left. \times \left| e^{-\frac{(\kappa_n - W_n - (B_n/2))^2}{D^2(B_n/2)}} - e^{-\frac{(\kappa_n - \kappa_m - (W_n - W_m) - (B_n - B_m)/2)^2}{D^2(B_n - B_m)/2}} \right| \right\} \\
(4.15) \quad &:= C_\theta \left( \frac{\sigma_n}{\sqrt{\nu_n}} \right) \{A_{11}'' + A_{12}''\}.
\end{aligned}$$

As  $\nu_n \uparrow \infty$  with  $n$ , it follows from Kolmogorov's law of the iterated logarithm that

$$(4.16) \quad \limsup_{n \rightarrow \infty} \frac{B_n - \nu_n}{\sqrt{\nu_n \log \log \nu_n}} \stackrel{\text{a.s.}}{=} 1.$$

Thus

$$B_* = \sup_{n \geq 1} \frac{B_n}{\nu_n} < \infty,$$

almost surely. Moreover  $\|B_*\|_p < \infty$  for  $1 < p < \infty$ .

Consider  $A_{11}''$ . Using the inequality  $\sqrt{x} - \sqrt{y} \leq \sqrt{x - y}$  if  $x \geq y \geq 0$ , we have on the set  $\{B_n > B_m, B_n - B_m > \theta(\nu_n - \nu_m)\}$ ,

$$\begin{aligned}
\sqrt{\frac{B_n}{B_n - B_m}} - 1 &= \frac{\sqrt{B_n} - \sqrt{B_n - B_m}}{\sqrt{B_n - B_m}} \leq \frac{\sqrt{B_m}}{\sqrt{B_n - B_m}} \leq B_*^{1/2} \frac{\sqrt{\nu_m}}{\sqrt{\theta(\nu_n - \nu_m)}} \\
&= B_*^{1/2} \frac{1}{\sqrt{\theta} \sqrt{\frac{\nu_n}{\nu_m} - 1}} \leq B_*^{1/2} \frac{1}{\sqrt{\theta} (\sqrt{\frac{\nu_n}{\nu_m}} - 1)}.
\end{aligned}$$

Thus

$$\begin{aligned}
A''_{11} &= \mathbb{E}_{(V,\varepsilon)} \left( \mathbf{1}_{\left\{ \frac{B_n - B_m}{\theta(\nu_n - \nu_m)} > 1 \right\}} + \mathbf{1}_{\left\{ \frac{B_n - B_m}{\theta(\nu_n - \nu_m)} \leq 1 \right\}} \right) \mathbf{1}_{\{B_n > \theta\nu_n, B_n > B_m\}} \left[ \sqrt{\frac{B_n}{B_n - B_m}} - 1 \right] \\
&\leq \frac{\mathbb{E}_{(V,\varepsilon)} B_*^{1/2}}{\sqrt{\theta}(\sqrt{\frac{\nu_n}{\nu_m}} - 1)} + \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\left\{ \frac{B_n - B_m}{\theta(\nu_n - \nu_m)} \leq 1 \right\}} \mathbf{1}_{\{B_n > B_m\}} \left[ \frac{\sqrt{B_m}}{\sqrt{B_n - B_m}} \right] \\
(4.17) \quad &\leq \frac{\mathbb{E}_{(V,\varepsilon)} B_*^{1/2}}{\sqrt{\theta}(\sqrt{\frac{\nu_n}{\nu_m}} - 1)} + \sqrt{\nu_m} \mathbb{E}_{(V,\varepsilon)} B_*^{1/2} \mathbf{1}_{\left\{ \frac{B_n - B_m}{\theta(\nu_n - \nu_m)} \leq 1 \right\}}.
\end{aligned}$$

By Hölder's inequality, for  $\alpha > 1$ ,

$$\begin{aligned}
\mathbb{E}_{(V,\varepsilon)} B_*^{1/2} \mathbf{1}_{\left\{ \frac{B_n - B_m}{\theta(\nu_n - \nu_m)} \leq 1 \right\}} &\leq (\mathbb{E}_{(V,\varepsilon)} (B_*)^{\alpha/2})^{1/\alpha} \mathbb{P}^{1-1/\alpha} \left\{ \frac{B_n - B_m}{\theta(\nu_n - \nu_m)} \leq 1 \right\} \\
&\leq (\mathbb{E}_{(V,\varepsilon)} (B_*)^{\alpha/2})^{1/\alpha} \rho^{(1-1/\alpha)(\nu_n - \nu_m)}.
\end{aligned}$$

Therefore

$$(4.18) \quad A''_{11} \leq \frac{\mathbb{E}_{(V,\varepsilon)} B_*^{1/2}}{\sqrt{\theta}(\sqrt{\frac{\nu_n}{\nu_m}} - 1)} + \|B_*^{1/2}\|_\alpha \sqrt{\nu_m} \rho^{(1-1/\alpha)(\nu_n - \nu_m)}.$$

We now turn to  $A''_{12}$ . Put

$$\kappa'_n = \kappa_n - W_n - (DB_n/2).$$

Then

$$\begin{aligned}
A''_{12} &= \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{B_n > \theta\nu_n, B_n > B_m\}} \left| e^{-\frac{(\kappa_n - W_n - (B_n/2))^2}{D^2(B_n/2)}} \right. \\
&\quad \left. - e^{-\frac{(\kappa_n - \kappa_m - (W_n - W_m) - (B_n - B_m)/2)^2}{D^2(B_n - B_m)/2}} \right| \\
&= \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{B_n > \theta\nu_n, B_n > B_m\}} \left| e^{-\frac{\kappa_n'^2}{D^2(B_n/2)}} - e^{-\frac{(\kappa_n' - \kappa_m')^2}{D^2(B_n - B_m)/2}} \right|.
\end{aligned}$$

At first,

$$(4.19) \quad \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{B_n > \theta\nu_n, 0 < B_n - B_m \leq \theta(\nu_n - \nu_m)\}} \left| e^{-\frac{\kappa_n'^2}{D^2(B_n/2)}} - e^{-\frac{(\kappa_n' - \kappa_m')^2}{D^2(B_n - B_m)/2}} \right| \leq 2\rho^{\nu_n - \nu_m}.$$

Concerning the integration over  $\{B_n > \theta\nu_n, B_n - B_m > \theta(\nu_n - \nu_m)\}$ , we get by letting

$$b_n = \frac{\kappa_n'}{\sqrt{B_n}},$$

and using the inequality  $|e^{-u} - e^{-v}| \leq |u - v|$  for reals  $u \geq 0, v \geq 0$ ,

$$\begin{aligned}
&\frac{D^2}{2} \left| e^{-\frac{\kappa_n'^2}{D^2(B_n/2)}} - e^{-\frac{(\kappa_n' - \kappa_m')^2}{D^2(B_n - B_m)/2}} \right| \\
&\leq \left| -\frac{(\kappa_n' - \kappa_m')^2}{(B_n - B_m)} + \frac{\kappa_n'^2}{B_n} \right| = \left| -\frac{(\sqrt{B_n}b_n - \sqrt{B_m}b_m)^2}{B_n - B_m} + b_n^2 \right| \\
&= \left| \frac{-B_nb_n^2 - B_mb_m^2 + 2\sqrt{B_nB_m}b_nb_m + B_nb_n^2 - B_mb_n^2}{B_n - B_m} \right| \\
&= \left| \frac{-(b_n - b_m)^2 + 2b_nb_m(\sqrt{\frac{B_n}{B_m}} - 1)}{\frac{B_n}{B_m} - 1} \right|
\end{aligned}$$

$$(4.20) \quad \leq \frac{2(b_n^2 + b_m^2) + 2|b_m||b_n|(\sqrt{\frac{B_n}{B_m}} - 1)}{\frac{B_n}{B_m} - 1}.$$

Hence

$$(4.21) \quad \begin{aligned} & \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n, B_n - B_m > \theta(\nu_n - \nu_m)\}} \left| e^{-\frac{\kappa_n'^2}{D^2(B_n/2)}} - e^{-\frac{(\kappa_n' - \kappa_m')^2}{D^2(B_n - B_m)/2}} \right| \\ & \leq C \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n, B_n - B_m > \theta(\nu_n - \nu_m)\}} \left\{ \frac{b_n^2 + b_m^2}{\frac{B_n}{B_m} - 1} + \frac{|b_m||b_n|}{\sqrt{\frac{B_n}{B_m}} - 1} \right\}. \end{aligned}$$

One next establishes the following three estimates:

$$(4.22) \quad \begin{aligned} \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n, B_n - B_m > \theta(\nu_n - \nu_m)\}} \frac{|b_n||b_m|}{\sqrt{\frac{B_n}{B_m}} - 1} & \leq \left( \frac{\sigma_n \sigma_m}{\sqrt{\nu_n \nu_m}} \right) \frac{C_\theta}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} \\ \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n, B_n - B_m > \theta(\nu_n - \nu_m)\}} \frac{|b_n|^2}{\frac{B_n}{B_m} - 1} & \leq \left( \frac{\sigma_n^2}{\nu_n} \right) \frac{C_\theta}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} \\ \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n, B_n - B_m > \theta(\nu_n - \nu_m)\}} \frac{|b_m|^2}{\frac{B_n}{B_m} - 1} & \leq \left( \frac{\sigma_m^2}{\nu_m} \right) \frac{C_\theta}{\sqrt{\frac{\nu_n}{\nu_m}} - 1}. \end{aligned}$$

Let  $S'_m = W_m + (DB_m/2)$ , we note that  $S'_m - a_m = S'_m - \mathbb{E}_{(V,\varepsilon)} S'_m$ , since  $\mathbb{E} S_m = \mathbb{E}_{(V,\varepsilon)} S'_m = a_m$ . Besides by (2.1)-(1),  $|\kappa_j - a_j| = \mathcal{O}(\sigma_j)$ ; thus

$$(4.23) \quad |b_j| = \frac{|\kappa_j - a_j - (W_j + (DB_j/2) - a_j)|}{\sqrt{B_j}} \leq \frac{C}{\sqrt{B_j}} \left[ \sigma_j + |S'_j - \mathbb{E}_{(V,\varepsilon)} S'_j| \right].$$

We note that  $\frac{1}{\sqrt{\frac{B_n}{B_m}} - 1} \leq \frac{\sqrt{B_m}(\sqrt{B_n} + \sqrt{B_m})}{\theta(\nu_n - \nu_m)}$ , on the set  $\{B_n > \theta \nu_n, B_n - B_m > \theta(\nu_n - \nu_m)\}$ .

Thus

$$\begin{aligned} & \frac{|b_n||b_m|}{\sqrt{\frac{B_n}{B_m}} - 1} \leq \frac{C}{\sqrt{B_n B_m}} \frac{\sqrt{B_m}(\sqrt{B_n} + \sqrt{B_m})}{\theta(\nu_n - \nu_m)} \\ & \quad \times \left[ \sigma_n + |S'_n - \mathbb{E}_{(V,\varepsilon)} S'_n| \right] \left[ \sigma_m + |S'_m - \mathbb{E}_{(V,\varepsilon)} S'_m| \right] \\ & \leq \frac{2C\sigma_n\sigma_m}{\theta(\nu_n - \nu_m)} \left[ 1 + \frac{|S'_n - \mathbb{E}_{(V,\varepsilon)} S'_n|}{\sigma_n} \right] \left[ 1 + \frac{|S'_m - \mathbb{E}_{(V,\varepsilon)} S'_m|}{\sigma_m} \right] \\ & \leq \left( \frac{\sigma_n\sigma_m}{\sqrt{\nu_n\nu_m}} \right) \frac{2C\sqrt{\nu_m}(\sqrt{\nu_n} + \sqrt{\nu_m})}{\theta(\nu_n - \nu_m)} \left[ 1 + \frac{|S'_n - \mathbb{E}_{(V,\varepsilon)} S'_n|}{\sigma_n} \right] \left[ 1 + \frac{|S'_m - \mathbb{E}_{(V,\varepsilon)} S'_m|}{\sigma_m} \right] \\ & = \left( \frac{\sigma_n\sigma_m}{\sqrt{\nu_n\nu_m}} \right) \frac{2C}{\theta(\sqrt{\frac{\nu_n}{\nu_m}} - 1)} \left[ 1 + \frac{|S'_n - \mathbb{E}_{(V,\varepsilon)} S'_n|}{\sigma_n} \right] \left[ 1 + \frac{|S'_m - \mathbb{E}_{(V,\varepsilon)} S'_m|}{\sigma_m} \right]. \end{aligned}$$

By the Cauchy-Schwarz inequality,

$$(4.24) \quad \begin{aligned} & \mathbb{E}_{(V,\varepsilon)} \frac{|S'_n - \mathbb{E}_{(V,\varepsilon)} S'_n|}{\sigma_n} \frac{|S'_m - \mathbb{E}_{(V,\varepsilon)} S'_m|}{\sigma_m} \\ & \leq \left[ \mathbb{E}_{(V,\varepsilon)} \frac{|S'_n - \mathbb{E}_{(V,\varepsilon)} S'_n|^2}{\sigma_n^2} \right]^{1/2} \left[ \mathbb{E}_{(V,\varepsilon)} \frac{|S'_m - \mathbb{E}_{(V,\varepsilon)} S'_m|^2}{\sigma_m^2} \right]^{1/2} \leq C. \end{aligned}$$

Since

$$\mathbb{E}_{(V,\varepsilon)} \frac{|b_n||b_m|}{\sqrt{\frac{B_n}{B_m}} - 1} \leq \left( \frac{\sigma_n \sigma_m}{\sqrt{\nu_n \nu_m}} \right) \frac{2C}{\theta(\sqrt{\frac{\nu_n}{\nu_m}} - 1)} \mathbb{E}_{(V,\varepsilon)} \left[ 1 + \frac{|S'_n - \mathbb{E}_{(V,\varepsilon)} S'_n|}{\sigma_n} \right] \left[ 1 + \frac{|S'_m - \mathbb{E}_{(V,\varepsilon)} S'_m|}{\sigma_m} \right]$$

we deduce

$$(4.25) \quad \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n, B_n - B_m > \theta(\nu_n - \nu_m)\}} \frac{|b_n||b_m|}{\sqrt{\frac{B_n}{B_m}} - 1} \leq \left( \frac{\sigma_n \sigma_m}{\sqrt{\nu_n \nu_m}} \right) \frac{2C}{\theta(\sqrt{\frac{\nu_n}{\nu_m}} - 1)}.$$

Now on the set  $\{B_n > \theta \nu_n, B_n - B_m > \theta(\nu_n - \nu_m)\}$ ,

$$\begin{aligned} \frac{|b_n|^2}{\frac{B_n}{B_m} - 1} &\leq \frac{C}{B_n} \left[ \sigma_n + |S'_n - \mathbb{E}_{(V,\varepsilon)} S'_n| \right]^2 \frac{B_m}{B_n - B_m} \\ &\leq C \frac{\sigma_n^2 B_m}{B_n(\theta(\nu_n - \nu_m))} \left[ 1 + \frac{|S'_n - \mathbb{E}_{(V,\varepsilon)} S'_n|}{\sigma_n} \right]^2 \\ &\leq C \left( \frac{\sigma_n^2}{\nu_n} \right) \frac{B_m}{\theta^2(\nu_n - \nu_m)} \left[ 1 + \frac{|S'_n - \mathbb{E}_{(V,\varepsilon)} S'_n|}{\sigma_n} \right]^2 \\ &\leq C B_* \left( \frac{\sigma_n^2}{\nu_n} \right) \frac{\nu_m}{\theta^2(\nu_n - \nu_m)} \left[ 1 + \frac{|S'_n - \mathbb{E}_{(V,\varepsilon)} S'_n|}{\sigma_n} \right]^2 \\ &= C B_* \left( \frac{\sigma_n^2}{\nu_n} \right) \frac{1}{\theta^2(\frac{\nu_n}{\nu_m} - 1)} \left[ 1 + \frac{|S'_n - \mathbb{E}_{(V,\varepsilon)} S'_n|}{\sigma_n} \right]^2 \\ &\leq C_{\theta,\rho} B_* \left( \frac{\sigma_n^2}{\nu_n} \right) \frac{1}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} \left[ 1 + \frac{|S'_n - \mathbb{E}_{(V,\varepsilon)} S'_n|}{\sigma_n} \right]^2. \end{aligned}$$

Therefore

$$\begin{aligned} &\mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n, B_n - B_m > \theta(\nu_n - \nu_m)\}} \frac{|b_n|^2}{\frac{B_n}{B_m} - 1} \\ &\leq C_{\theta,\rho} \left( \frac{\sigma_n^2}{\nu_n} \right) \frac{1}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} \mathbb{E}_{(V,\varepsilon)} B_* \left[ 1 + \frac{|S'_n - \mathbb{E}_{(V,\varepsilon)} S'_n|}{\sigma_n} \right]^2 \\ &\leq C_{\theta,\rho} \left( \frac{\sigma_n^2}{\nu_n} \right) \frac{1}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} \left[ 1 + \frac{1}{\sigma_n^2} \mathbb{E}_{(V,\varepsilon)} |S'_n - \mathbb{E}_{(V,\varepsilon)} S'_n|^2 \right] \\ (4.26) \quad &\leq \left( \frac{\sigma_n^2}{\nu_n} \right) \frac{C_{\theta,\rho}}{\sqrt{\frac{\nu_n}{\nu_m}} - 1}. \end{aligned}$$

Next  $|b_m| \leq \frac{C}{\sqrt{B_m}} \left[ \sigma_m + |S'_m - \mathbb{E}_{(V,\varepsilon)} S'_m| \right]$ , and so on the set  $\{B_n - B_m > \theta(\nu_n - \nu_m)\}$ ,

$$\begin{aligned} \frac{|b_m|^2}{\frac{B_n}{B_m} - 1} &\leq \frac{C}{B_m} \left[ \sigma_m + |S'_m - \mathbb{E}_{(V,\varepsilon)} S'_m| \right]^2 \frac{B_m}{B_n - B_m} \\ &\leq C \frac{\sigma_m^2}{\theta(\nu_n - \nu_m)} \left[ 1 + \frac{|S'_m - \mathbb{E}_{(V,\varepsilon)} S'_m|}{\sigma_m} \right]^2 \\ &\leq C \left( \frac{\sigma_m^2}{\nu_m} \right) \frac{\nu_m}{\theta(\nu_n - \nu_m)} \left[ 1 + \frac{|S'_m - \mathbb{E}_{(V,\varepsilon)} S'_m|}{\sigma_m} \right]^2 \\ &\leq C \left( \frac{\sigma_m^2}{\nu_m} \right) \frac{1}{\theta(\frac{\nu_n}{\nu_m} - 1)} \left[ 1 + \frac{|S'_m - \mathbb{E}_{(V,\varepsilon)} S'_m|}{\sigma_m} \right]^2 \end{aligned}$$

$$\leq \left( \frac{\sigma_m^2}{\nu_m} \right) \frac{C_{\theta,\rho}}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} \left[ 1 + \frac{|S'_m - \mathbb{E}_{(V,\varepsilon)} S'_m|}{\sigma_m} \right]^2.$$

Therefore

$$(4.27) \quad \begin{aligned} & \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n, B_n - B_m > \theta(\nu_n - \nu_m)\}} \frac{|b_m|^2}{\frac{B_n}{B_m} - 1} \\ & \leq \left( \frac{\sigma_m^2}{\nu_m} \right) \frac{C_{\theta,\rho}}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} \left[ 1 + \frac{|S'_m - \mathbb{E}_{(V,\varepsilon)} S'_m|}{\sigma_m} \right]^2 \leq \left( \frac{\sigma_m^2}{\nu_m} \right) \frac{C_{\theta,\rho}}{\sqrt{\frac{\nu_n}{\nu_m}} - 1}. \end{aligned}$$

We thus arrive at

$$(4.28) \quad \mathbb{E}_{(V,\varepsilon)} \mathbf{1}_{\{B_n > \theta \nu_n, B_n - B_m > \theta(\nu_n - \nu_m)\}} \left| e^{-\frac{\kappa'_n{}^2}{D^2(B_n/2)}} - e^{-\frac{(\kappa'_n - \kappa'_m)^2}{D^2(B_n - B_m)/2}} \right| \leq \left( \frac{\sigma_n}{\sqrt{\nu_n}} + \frac{\sigma_m}{\sqrt{\nu_m}} \right)^2 \frac{C_{\theta,\rho}}{\sqrt{\frac{\nu_n}{\nu_m}} - 1},$$

Consequently,

$$(4.29) \quad |A''_{12}| \leq 2\rho^{\nu_n - \nu_m} + \left( \frac{\sigma_n}{\sqrt{\nu_n}} + \frac{\sigma_m}{\sqrt{\nu_m}} \right)^2 \frac{C_{\theta,\rho}}{\sqrt{\frac{\nu_n}{\nu_m}} - 1}.$$

As by (4.18),

$$|A''_{11}| \leq \frac{\mathbb{E}_{(V,\varepsilon)} B_*}{\sqrt{\theta}(\sqrt{\frac{\nu_n}{\nu_m}} - 1)} + \|B_*\|_\alpha \sqrt{\nu_m} \rho^{(1-1/\alpha)(\nu_n - \nu_m)},$$

it follows that

$$(4.30) \quad |A''_1| \leq \frac{\mathbb{E}_{(V,\varepsilon)} B_*^{1/2}}{\sqrt{\theta}(\sqrt{\frac{\nu_n}{\nu_m}} - 1)} + \|B_*^{1/2}\|_\alpha \sqrt{\nu_m} \rho^{(1-1/\alpha)(\nu_n - \nu_m)} + 2\rho^{\nu_n - \nu_m} + \left( \frac{\sigma_n}{\sqrt{\nu_n}} + \frac{\sigma_m}{\sqrt{\nu_m}} \right)^2 \frac{C_{\theta,\rho}}{\sqrt{\frac{\nu_n}{\nu_m}} - 1}.$$

We have

$$|A| \leq A' + C_0 |A''_2| + C_\theta \left( \frac{\sigma_n}{\sqrt{\nu_n}} \right) A''_1.$$

Recalling that  $|A'| \leq \sigma_n \prod_{j=m+1}^n \vartheta_j$ , and  $|A''_2| \leq C \sigma_n \left\{ \rho^{\nu_n - \nu_m} + \frac{1}{(\theta \nu_n)^{3/2}} + \frac{1}{(\theta(\nu_n - \nu_m))^{3/2}} \right\}$ , by (4.11) and (4.14) respectively, we obtain

$$(4.31) \quad \begin{aligned} |A| & \leq \sigma_n \prod_{j=m+1}^n \vartheta_j + C_0 C \sigma_n \left\{ \rho^{\nu_n - \nu_m} + \frac{1}{(\theta \nu_n)^{3/2}} + \frac{1}{(\theta(\nu_n - \nu_m))^{3/2}} \right\} \\ & + C_\theta \left( \frac{\sigma_n}{\sqrt{\nu_n}} \right) \left\{ \frac{\mathbb{E}_{(V,\varepsilon)} B_*}{\sqrt{\theta}(\sqrt{\frac{\nu_n}{\nu_m}} - 1)} + \|B_*\|_\alpha \sqrt{\nu_m} \rho^{(1-1/\alpha)(\nu_n - \nu_m)} + 2\rho^{\nu_n - \nu_m} \right. \\ & \left. + \left( \frac{\sigma_n}{\sqrt{\nu_n}} + \frac{\sigma_m}{\sqrt{\nu_m}} \right)^2 \frac{C_{\theta,\rho}}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} \right\}. \end{aligned}$$

We note that

$$\rho^{\nu_n - \nu_m} + \frac{1}{(\theta \nu_n)^{3/2}} + \frac{1}{(\theta(\nu_n - \nu_m))^{3/2}}$$

is less than  $\frac{1}{(\nu_n - \nu_m)^{3/2}}$ , up to a constant  $C_{\theta,\rho,\alpha}$ . Let  $\sigma(n, m) = \max\left(\frac{\sigma_n}{\sqrt{\nu_n}}, \frac{\sigma_m}{\sqrt{\nu_m}}, 1\right)$ .

By considering separately the cases  $\nu_n - \nu_m \geq 1$ ,  $0 < \nu_n - \nu_m < 1$ , we thus get

$$|A| \leq \sigma_n \prod_{j=m+1}^n \vartheta_j + C_{\theta,\rho,\alpha} \frac{\sigma_n}{(\nu_n - \nu_m)^{3/2}}$$

$$(4.32) \quad +C_{\theta,\rho,\alpha} \left( \frac{\sigma_n}{\sqrt{\nu_n}} \right) \left\{ \frac{1}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} + \sqrt{\nu_m} \rho^{(1-1/\alpha)(\nu_n-\nu_m)} + \left( \frac{\sigma_n}{\sqrt{\nu_n}} + \frac{\sigma_m}{\sqrt{\nu_m}} \right)^2 \frac{1}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} \right\}.$$

Now by (1.15),  $1 \leq \frac{2}{D} \frac{\sigma}{\sqrt{\vartheta_Y}}$ , and so

$$\sigma(n, m) \leq \frac{C}{D} \max \left( \frac{\sigma_n}{\sqrt{\nu_n}}, \frac{\sigma_m}{\sqrt{\nu_m}} \right).$$

Thus

$$(4.33) \quad \begin{aligned} |A| &\leq \sigma_n \prod_{j=m+1}^n \vartheta_j + C_{\theta,\rho,\alpha} \left( \frac{\sigma_n}{\sqrt{\nu_n}} \right) \frac{\sqrt{\nu_n}}{(\nu_n - \nu_m)^{3/2}} \\ &\quad + C_{\theta,\rho,\alpha} \left( \frac{\sigma_n}{\sqrt{\nu_n}} \right) \left\{ \frac{1}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} + \frac{\sqrt{\nu_m}}{(\nu_n - \nu_m)^{3/2}} + \left( \frac{C}{D} \max \left( \frac{\sigma_n}{\sqrt{\nu_n}}, \frac{\sigma_m}{\sqrt{\nu_m}} \right) \right)^2 \frac{1}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} \right\} \\ &\leq \sigma_n \prod_{j=m+1}^n \vartheta_j + C_{\theta,\rho,\alpha} \left( \frac{\sigma_n}{\sqrt{\nu_n}} \right) \left\{ \frac{1}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} + \frac{\sqrt{\nu_n}}{(\nu_n - \nu_m)^{3/2}} \right\} \\ &\quad + C_{\theta,\rho,\alpha} D \left( \frac{C}{D} \max \left( \frac{\sigma_n}{\sqrt{\nu_n}}, \frac{\sigma_m}{\sqrt{\nu_m}} \right) \right)^3 \left\{ \frac{1}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} \right\} \\ &\leq C_{\theta,\rho,\alpha} \frac{C}{D^2} \max \left( \frac{\sigma_n}{\sqrt{\nu_n}}, \frac{\sigma_m}{\sqrt{\nu_m}} \right)^3 \\ &\quad \times \left\{ \nu_n^{1/2} \prod_{j=m+1}^n \vartheta_j + \frac{1}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} + \frac{\sqrt{\nu_n}}{(\nu_n - \nu_m)^{3/2}} \right\}. \end{aligned}$$

Using (4.4), (4.6) we obtain (the value of  $\alpha$  being irrelevant and  $\theta, \rho$  depending only on  $\varepsilon$ , the value of which being irrelevant too)

$$(4.34) \quad \begin{aligned} |\mathbb{E} Y_n Y_m| &\leq \frac{C}{D^2} \max \left( \frac{\sigma_n}{\sqrt{\nu_n}}, \frac{\sigma_m}{\sqrt{\nu_m}} \right)^3 \\ &\times \left\{ \nu_n^{1/2} \prod_{j=m+1}^n \vartheta_j + \frac{1}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} + \frac{\sqrt{\nu_n}}{(\nu_n - \nu_m)^{3/2}} \right\} \end{aligned}$$

if  $n > m \geq 1$ . This together with (4.2), proves (2.2).

Finally if  $\tau = \sup_{j \geq 1} \vartheta_{X_j} < 1$ , we pick  $\vartheta_j$  so that

$$0 < \vartheta_j \leq \vartheta_{X_j} \leq \tau < 1, \quad j = 1, 2, \dots,$$

and note that

$$\log \frac{1}{\vartheta_j} \geq \log \frac{1}{\tau} \geq \left( \frac{\log \frac{1}{\tau}}{\tau} \right) \vartheta_j,$$

Thus

$$\prod_{j=m+1}^n \vartheta_j = e^{-\sum_{j=m+1}^n \log \frac{1}{\vartheta_j}} \leq \frac{C(M)}{(\sum_{j=m+1}^n \log \frac{1}{\vartheta_j})^M} \leq \frac{C(\tau, M)}{(\sum_{j=m+1}^n \vartheta_j)^M} = \frac{C(\cdot, M)}{(\nu_n - \nu_m)^M}.$$

Taking  $M = 3/2$  provides the bound

$$\prod_{j=m+1}^n \vartheta_j \leq \frac{C(\tau)}{(\nu_n - \nu_m)^{3/2}}.$$

By carrying it back to estimate (4.34), we get

$$(4.35) \quad |\mathbb{E} Y_n Y_m| \leq \frac{C_{\tau,c}}{D^2} \max\left(\frac{\sigma_n}{\sqrt{\nu_n}}, \frac{\sigma_m}{\sqrt{\nu_m}}\right)^3 \left\{ \frac{1}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} + \frac{\sqrt{\nu_n}}{(\nu_n - \nu_m)^{3/2}} \right\},$$

which is estimate (2.5). This completes the proof of Theorem 2.1.

**4.1. Proof of Corollary 2.6.** Let  $0 < c < 1$ . Let  $\nu_m \leq c \nu_n$ . Then

$$\frac{1}{\sqrt{\nu_n/\nu_m} - 1} \leq \left(\frac{1}{1 - \sqrt{c}}\right) \sqrt{\frac{\nu_m}{\nu_n}} = C_c \sqrt{\frac{\nu_m}{\nu_n}}.$$

Further

$$\frac{\sqrt{\nu_n}}{(\nu_n - \nu_m)^{3/2}} = \sqrt{\frac{\nu_n}{\nu_n - \nu_m}} \frac{1}{(\nu_n - \nu_m)} = \frac{1}{(1 - \nu_m/\nu_n)^{3/2}} \frac{1}{\nu_n} \leq \frac{1}{(1 - c)^{3/2}} \frac{1}{\nu_n}.$$

By incorporating these estimates into (4.32) we get

$$(4.36) \quad |\mathbb{E} Y_n Y_m| \leq \frac{C_{\tau,c}}{D^2} \max\left(\frac{\sigma_n}{\sqrt{\nu_n}}, \frac{\sigma_m}{\sqrt{\nu_m}}\right)^3 \sqrt{\frac{\nu_m}{\nu_n}}.$$

This proves Corollary 2.6.

In the following Example, we study the correlation of the system of random variables (recalling (4.2))

$$(4.37) \quad \frac{\vartheta_n Y_n}{\sigma_n \sqrt{\nu_n}} = \frac{\vartheta_n (\mathbf{1}_{\{S_n = \kappa_n\}} - \mathbb{P}\{S_n = \kappa_n\})}{\sqrt{\nu_n}}, \quad n \geq 2.$$

We consider the case

$$\lim_{n \rightarrow \infty} \vartheta_{X_n} = 0.$$

**4.2. Example.** Let

$$(4.38) \quad \text{Var}(X_n) = \vartheta_{X_n} = \log n - \log(n-1), \quad n \geq 1.$$

We also choose

$$D = 2, \quad \vartheta_n = \vartheta_{X_n}, \quad n \geq 1.$$

As  $\nu_n = \log n$ , assumption 1.3 is satisfied. By (4.38),

$$(4.39) \quad \sigma_n = \sqrt{\log n} = \sqrt{\nu_n}.$$

This is compatible with (1.16), since we assumed  $D = 2$ . We have  $\frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}} = \frac{1}{n \log n}$ , thus the series  $\sum_{n \geq 1} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}}$  is divergent.

In the case considered,

$$(4.40) \quad \frac{Y_n}{n \log n} = \frac{\mathbf{1}_{\{S_n = \kappa_n\}} - \mathbb{P}\{S_n = \kappa_n\}}{n \sqrt{\log n}}, \quad n \geq 2.$$

We examine the magnitude of the  $L^2$ -increments.



**Lemma 4.1.** *There exists an absolute constant  $C$  such that for any integers  $1 < M_1 < M_2$ ,*

$$(4.41) \quad \mathbb{E} \left| \sum_{M_1 \leq m \leq M_2} \frac{Y_m}{m \log m} \right|^2 \leq C \sum_{M_1 \leq m \leq M_2} \frac{1}{m \log m}.$$

*Proof.* Let  $b > 1$  be some fixed real. We consider several cases.

**Case 1:** ( $m = n$ ) By using (4.7), we have  $\mathbb{E} Y_n^2 \leq C \sqrt{n}$ , so that

$$\frac{\mathbb{E} Y_m^2}{m \log m} \leq \frac{C}{\sqrt{m} \log m} \leq C.$$

**Case 2:** ( $m + 1 \leq n < m^b$ ) By using (4.5),

$$(4.42) \quad \begin{aligned} \sum_{m+1 \leq n < m^b} \frac{|\mathbb{E} Y_m Y_n|}{n \log n} &\leq C \sum_{m+1 \leq n < m^b} \left( \frac{1}{n \sqrt{\log n} \sqrt{\log(n-m)}} + \frac{1}{n \log n} \right) \\ &\leq C_b \left( \frac{1}{\sqrt{\log m}} \sum_{1 \leq h \leq m^b} \frac{1}{h \sqrt{\log h}} + (\log \log m^b - \log \log m) \right) \\ &\leq C_b \left( \frac{1}{\sqrt{\log m}} \sqrt{\log m} + \log \frac{b \log m}{\log m} \right) \leq C_b. \end{aligned}$$

**Case 3:** ( $n \geq m^b$ ) By Corollary 2.4,

$$(4.43) \quad \begin{aligned} |\mathbb{E} Y_n Y_m| &= \sigma_n \sigma_m \left| \mathbb{P}\{S_n = \kappa_n, S_m = \kappa_m\} - \mathbb{P}\{S_n = \kappa_n\} \mathbb{P}\{S_m = \kappa_m\} \right| \\ &\leq \frac{C_\tau}{D^2} \max \left( \frac{\sigma_n}{\sqrt{\nu_n}}, \frac{\sigma_m}{\sqrt{\nu_m}} \right)^3 \left\{ \frac{1}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} + \frac{\sqrt{\nu_n}}{(\nu_n - \nu_m)^{3/2}} \right\} \\ &\leq C \left\{ \frac{1}{\sqrt{\frac{\nu_n}{\nu_m}} - 1} + \frac{\sqrt{\nu_n}}{(\nu_n - \nu_m)^{3/2}} \right\} \leq C \frac{1}{\sqrt{\frac{\nu_n}{\nu_m}} - 1}. \end{aligned}$$

Thus

$$(4.44) \quad \sum_{n > m^b} \frac{|\mathbb{E} Y_m Y_n|}{n \log n} \leq C \sum_{n > m^b} \frac{1}{n(\log n) \sqrt{\frac{\log n}{\log m} - 1}}.$$

We have  $\frac{\log n}{\log m} \geq (1 - 1/b)$ . Thus

$$\frac{1}{\sqrt{\frac{\log n}{\log m} - 1}} \leq \left( \frac{1}{1 - \sqrt{c}} \right) \sqrt{\frac{\log m}{\log n}} = C_c \sqrt{\frac{\log m}{\log n}}.$$

So that

$$(4.45) \quad \sum_{n > m^b} \frac{|\mathbb{E} Y_m Y_n|}{n \log n} \leq C_b \sum_{n > m^b} \frac{1}{n(\log n)} \sqrt{\frac{\log m}{\log n}} = C_b \sqrt{\log m} \sum_{n > m^b} \frac{1}{n(\log n)^{3/2}}.$$

This sum is a special case of the sum

$$\sum_{n > N} \frac{\vartheta_n}{\nu_n^\delta \nu_n},$$

and we can apply Lemma 5.6 with  $\delta = \frac{1}{2}$ . Consequently (choosing  $N = m^b$ )

$$\sqrt{\log m^b} \sum_{n > m^b} \frac{1}{n(\log n)^{3/2}} \leq C_b.$$

We deduce

$$(4.46) \quad \sum_{n > m^b} \frac{|\mathbb{E} Y_m Y_n|}{n \log n} \leq C_b \sqrt{\log m} \sum_{n > m^b} \frac{1}{n(\log n)^{3/2}} \leq C_b.$$

By combining,

$$(4.47) \quad \sum_{n \geq m} \frac{|\mathbb{E} Y_m Y_n|}{n \log n} \leq C_b.$$

Now noticing that

$$(4.48) \quad \mathbb{E} \left| \sum_{M_1 \leq m \leq M_2} \frac{Y_m}{m \log m} \right|^2 \leq \sum_{M_1 \leq m \leq M_2} \frac{\mathbb{E} Y_m^2}{(m \log m)^2} + 2 \sum_{M_1 \leq m \leq M_2} \frac{1}{m \log m} \left( \sum_{m < n \leq M_2} \frac{|\mathbb{E} Y_m Y_n|}{n \log n} \right).$$

□

## 5. A GENERAL ASLLT WITH ALMOST SURE CONVERGENT SERIES

In this Section we derive from Theorem 2.1 and Corollary 2.6, a sharp almost sure local limit theorem for sums of independent random variables. This one is stated in Theorem 5.11, assertion (iv), and thereby extends the ASLLT proved in the i.i.d. case (Theorem 1.1), to the independent non-identically distributed case. This result is deduced from new general stronger forms in (i), (ii) and (iii), in which the almost sure convergence of the naturally associated series, namely

$$\sum_{k \geq 1} \sup \left\{ \frac{\left| \sum_{2^j \leq n < 2^{j+1}} \frac{\mathbf{1}_{\{S_n = \kappa_n\}} - \mathbb{P}\{S_n = \kappa_n\}}{\sqrt{\nu_n}} \right|}{\sum_{1 \leq n < 2^j} \frac{1}{\sigma_n \sqrt{\nu_n}}} \right\}, \quad 2^k \leq \sum_{1 \leq n < 2^j} \frac{1}{\sigma_n \sqrt{\nu_n}} < 2^{k+1} \Bigg\},$$

is established. For proving these results *no* LLT is assumed; and thus these are in that sense *disconnected* from LLT. However if the LLT is applicable, our assumptions can be weakened and the ASLLT becomes stronger: compare Claims (iii) and (iv) in Theorem 5.11.

**5.1. Basic Assumption.** We assume throughout this Section that

*The series  $\sum_{n \geq 1} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}}$  is divergent.*

**Remark 5.1.** It will be later proved that in the case this assumption is not satisfied, by assuming only a little more than the convergence of the series  $\sum_{n \geq 1} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}}$ , a strong convergence result holds, although radically different from the almost sure local limit property.

**5.2. Introducing function  $h$ .** Let function  $h$  be defined as follows:

$$(5.1) \quad h(x) = \max_{1 \leq m \leq x} \left( \frac{\sigma_m^2}{\nu_m} \right), \quad x \geq 1.$$

**Remark 5.2.** By inequality (1.16),  $\sigma_n^2 \geq \frac{D^2}{4} \nu_n$ , for  $n \geq 1$ , so that

$$h(x) \geq D^2/4, \quad x \geq 1.$$

Now if

$$(5.2) \quad \sigma_n^2 = \mathcal{O}(\nu_n),$$

one can take  $h(x) \equiv \text{const.}$ , and this shows that the basic assumption is fulfilled. This follows from assertion (i) of Th. 162 in [27]. Condition (5.2) is obviously fulfilled in the i.i.d. case.

**5.3. A system of weights.** Introduce a system of weights associated with sequences  $\{\vartheta_n, n \geq 1\}$ ,  $\{\sigma_n, n \geq 1\}$ , and of key importance for controlling  $\mathbb{E} Z_i^2$ .

Set

$$(5.3) \quad \omega(m) = \sum_{\nu_m < \nu_n < 2\nu_m} \frac{\vartheta_n}{(\sigma_n^2 - \sigma_m^2)^{1/2} \sqrt{\nu_n}} \quad m \geq 1.$$

From (1.16) follows that

$$(5.4) \quad \omega(m) \leq C_\vartheta \sum_{\nu_m < \nu_n < 2\nu_m} \frac{\vartheta_n}{\sqrt{\nu_n - \nu_m} \sqrt{\nu_n}}.$$

We begin with giving some examples for which the following condition

$$(5.5) \quad \omega_X^* := \sup_{m \geq 1} \omega(m) < \infty,$$

is satisfied.

**Example 5.3** (i.i.d. case). In this case ( $\vartheta_n = C$ ,  $\nu_n = n$ ,  $\sigma_n = \sqrt{n}$ ), we have

$$\omega(m) = C \sum_{m < n < 2m} \frac{1}{\sqrt{n} \sqrt{n - m}} \leq \frac{C}{\sqrt{m}} \sum_{h=1}^m \frac{1}{\sqrt{h}} \leq \frac{C}{\sqrt{m}} \sqrt{m} = C.$$

Thus  $\omega_X^*$  is finite.

**Example 5.4** (Cramér's probabilistic model). This well-known model basically consists with a sequence of independent random variables  $\xi = \{\xi_j, j \geq 3\}$ , and associated partial sums  $S_n = \sum_{j=3}^n \xi_j$ , each  $\xi_j$  being defined by

$$(5.6) \quad \mathbb{P}\{\xi_j = 1\} = \frac{1}{\log j}, \quad \mathbb{P}\{\xi_j = 0\} = 1 - \frac{1}{\log j}.$$

Thus for  $j \geq 3$ ,  $\mathbb{P}\{\xi_j = k\} \wedge \mathbb{P}\{\xi_j = k+1\} = \frac{1}{\log j}$ , if  $k = 0$ , equals 0 for  $k \in \mathbb{Z}_*$ ; so the span is 1, and  $\vartheta_{\xi_j} = \frac{1}{\log j}$ . Choose  $\vartheta_j = \vartheta_{\xi_j}$ , then  $\nu_n = \sum_{j=3}^n \frac{1}{\log j}$ . We have  $\mathbb{E} S_n = \sum_{j=3}^n \frac{1}{\log j}$ ,  $\sigma_n^2 = \sum_{j=3}^n (1 - \frac{1}{\log j})(\frac{1}{\log j})$ . Moreover  $h(n) = \mathcal{O}(1)$ .

We have for  $n > m$ ,  $m$  large, using  $(x/\log x)' = 1/\log x - 1/(\log x)^2$ ,

$$\sigma_n^2 - \sigma_m^2 = \sum_{j=m+1}^n \left( \frac{1}{\log j} - \frac{1}{(\log j)^2} \right) \sim \int_m^n \left( \frac{x}{\log x} \right)' dx = \frac{n}{\log n} - \frac{m}{\log m} \geq (1 - o(1)) \frac{n - m}{\log n}.$$

Using also that  $\nu_n \geq \frac{n}{\log n} - o(1)$ ,  $2\nu_m \sim \nu_{2m}$ , we deduce

$$\begin{aligned}
 \omega(n) &\leq \sum_{\nu_m < \nu_n < 2\nu_m} \frac{\vartheta_n}{(\sigma_n^2 - \sigma_m^2)^{1/2} \sqrt{\nu_n}} \leq C \sum_{\nu_m < \nu_n < 2\nu_m} \frac{1}{\log n} \sqrt{\frac{\log n}{n}} \sqrt{\frac{\log n}{n-m}} \\
 (5.7) \quad &\leq C \sum_{\nu_m < \nu_n < 2\nu_m} \frac{1}{\sqrt{n-m}} \leq C \int_m^n \frac{dx}{\sqrt{x(x-m)}} \leq C \int_0^1 \frac{du}{\sqrt{u(u+1)}} = C.
 \end{aligned}$$

So that  $\omega_\xi^* < \infty$ .

By Proposition 3.1 in [53], the local limit theorem holds,

$$\left| \mathbb{P}\{S_n = \kappa\} - \frac{1}{\sqrt{2\pi\sigma_n}} e^{-\frac{(\kappa - m_n)^2}{2\sigma_n^2}} \right| \leq C \frac{(\log n)^{3/2}}{n},$$

for all  $\kappa \in \mathbb{Z}$  such that  $|\kappa - m_n| \leq C \frac{n^{3/4}}{\log n}$ .

In the next Example however we can only show that  $\omega(m) = \mathcal{O}(\log \log m)$ .

**Example 5.5** (Example 4.2 continued). Here  $\vartheta_n = \frac{1}{n}$ ,  $\nu_n = \log n$ ,  $\sigma_n = \sqrt{\log n}$ . By using Cauchy-Schwarz's inequality,

$$\begin{aligned}
 \omega(m) &= \sum_{\nu_m < \nu_n < 2\nu_m} \frac{1}{n \sqrt{(\log n)(\log n - \log m)}} \\
 &= \sum_{m+1 \leq n < m^2} \frac{1}{n \sqrt{\log n} \sqrt{\log(n-m)}} \\
 &\leq \left( \int_m^{m^2} \frac{dx}{x \log x} \right)^{1/2} \left( \int_m^{m^2} \frac{dx}{x(\log x - \log m)} \right)^{1/2} \\
 &= \sqrt{\log 2} \left( \int_1^m \frac{dt}{t \log t} \right)^{1/2} \\
 &= \sqrt{(\log 2) \log \log m}.
 \end{aligned}$$

The next lemma is of relevance for estimating the rectangle sums appearing in the first Step of the proof of Theorem 5.11.

**Lemma 5.6.** *We have the following estimates,*

(i) *Let  $0 < \delta < 1$ . Then for all  $N \geq 1$ ,*

$$\sum_{n > N} \frac{\delta \vartheta_n}{\nu_{n-1}^\delta \nu_n} \leq \frac{1}{\nu_N^\delta}.$$

(ii) *Further, for  $M > N \geq 1$  such that  $\nu_N > e$ ,*

$$(5.8) \quad \left| \sum_{N < n \leq M} \frac{\vartheta_n}{\nu_n} \right| \leq |\log \nu_M - \log \nu_N|.$$

*Proof.* (i) We proceed as in the proof of Th. 162 in Hardy, Littlewood and Pólya [27]. By Th. 41 in [27],  $\delta \vartheta_n \nu_n^{\delta-1} = \delta \nu_n^{\delta-1} (\nu_n - \nu_{n-1}) \leq \nu_n^\delta - \nu_{n-1}^\delta$ . So that

$$\sum_{n > N} \frac{\delta \vartheta_n}{\nu_{n-1}^\delta \nu_n} = \sum_{n > N} \frac{\delta \vartheta_n \nu_n^{\delta-1}}{\nu_{n-1}^\delta \nu_n^\delta} = \sum_{n > N} \frac{\delta \nu_n^{\delta-1} (\nu_n - \nu_{n-1})}{\nu_{n-1}^\delta \nu_n^\delta} \leq \sum_{n > N} \frac{\nu_n^\delta - \nu_{n-1}^\delta}{\nu_{n-1}^\delta \nu_n^\delta}$$

$$(5.9) \quad = \sum_{n>N} \left( \frac{1}{\nu_{n-1}^\delta} - \frac{1}{\nu_n^\delta} \right) = \frac{1}{\nu_N^\delta}.$$

(ii) Now, at first for  $M > N \geq 1$  such that  $\nu_N > e$ , we have

$$(5.10) \quad \begin{aligned} \sum_{N < n \leq M} \frac{\delta \vartheta_n}{\nu_{n-1}^\delta \nu_n} &= \sum_{N < n \leq M} \frac{\delta \vartheta_n \nu_n^{\delta-1}}{\nu_{n-1}^\delta \nu_n^\delta} = \sum_{N < n \leq M} \frac{\delta \nu_n^{\delta-1} (\nu_n - \nu_{n-1})}{\nu_{n-1}^\delta \nu_n^\delta} \\ &\leq \sum_{N < n \leq M} \frac{\nu_n^\delta - \nu_{n-1}^\delta}{\nu_{n-1}^\delta \nu_n^\delta} = \sum_{N < n \leq M} \left( \frac{1}{\nu_{n-1}^\delta} - \frac{1}{\nu_n^\delta} \right) \\ &= \frac{1}{\nu_N^\delta} - \frac{1}{\nu_M^\delta} = \delta (\log \nu_M - \log \nu_N) e^{-\theta(N,M)}, \end{aligned}$$

for some *positive* real  $\theta(N, M)$  such that  $\delta \log \nu_N < \theta(N, M) < \delta \log \nu_M$ , where we used Lagrange's Theorem in the last line of inequalities. It follows that

$$(5.11) \quad \left| \sum_{N < n \leq M} \frac{\delta \vartheta_n}{\nu_{n-1}^\delta \nu_n} \right| \leq \delta (\log \nu_M - \log \nu_N).$$

By dividing both sides by  $\delta$ , letting next  $\delta$  tend to 0 in the left-term, we obtain

$$(5.12) \quad \left| \sum_{N < n \leq M} \frac{\vartheta_n}{\nu_n} \right| \leq |\log \nu_M - \log \nu_N|.$$

□

**5.4. Associated functions  $k$ ,  $M$ ,  $\Phi$  and sequence  $\mathcal{M}$ .** Let

$$(5.13) \quad M(t) = \sum_{1 \leq \nu_n < t} \frac{\vartheta_m}{\sigma_m \sqrt{\nu_m}}, \quad t \geq 1$$

and set for  $x \geq M(1)$ ,  $M^{-1}(x) = \sup\{t \geq 1 : M(t) \leq x\}$ .

Put for any positive integer  $i$ ,

$$(5.14) \quad m_i = \sum_{2^i \leq \nu_n < 2^{i+1}} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}}.$$

Let the sequence  $\mathcal{M}$  be defined as follows,

$$(5.15) \quad \mathcal{M} = \{M_J, J \geq 1\} \quad \text{where} \quad M_J = M(2^J) = \sum_{1 \leq i < J} m_i, \quad J \geq 1.$$

The almost sure local limit theorem in view, will tightly rely upon the asymptotic distribution of the sequence  $\mathcal{M}$ .

It will be deduced from the study of the asymptotic almost sure behavior of the sequence

$$(5.16) \quad \mathcal{Z} = \{Z_i, i \geq 1\} \quad \text{where for each } i, \quad Z_i = \sum_{2^i \leq \nu_n < 2^{i+1}} \frac{\vartheta_n Y_n}{\sigma_n \sqrt{\nu_n}},$$

recalling that  $Y_n = \sigma_n(\mathbf{1}_{\{S_n = \kappa_n\}} - \mathbb{P}\{S_n = \kappa_n\})$ . We will apply Theorem 3.7 with the choices  $\xi_l = Z_l$  and  $u_l = m_l$ ,  $l \geq 1$ . We thus need assumptions (3.16), (3.17) to be fulfilled.

Introduce function  $k(x)$

$$(5.17) \quad k(x) = h(x)\omega(x), \quad x \geq 1.$$

Now let the non-decreasing function  $\Phi$  be defined as follows:

$$(5.18) \quad \Phi(x) = k \circ M^{-1}(x).$$

**Example 5.7.** Assume that  $h(x) = \mathcal{O}(1)$ , and  $\omega_X^* = \mathcal{O}(1)$  (condition (5.5)), then

$$\Phi(x) = \mathcal{O}(1).$$

In Examples (5.3), (5.4), condition (5.5) is satisfied.

**5.5. The key metrical estimate.** We control the  $L^2$ -norm of increments  $\sum_{I \leq i < J} Z_i$  and prove the following estimate.

**Theorem 5.8.** *Assume that*

$$(5.19) \quad m_i = \mathcal{O}(1), \quad i \rightarrow \infty.$$

*There exists a constant  $C_X$  such that for any  $J \geq I \geq 1$ ,*

$$(5.20) \quad \mathbb{E} \left| \sum_{I \leq i < J} Z_i \right|^2 \leq C_X h(2^{J+1}) \sum_{I \leq i \leq J} \left( \sum_{2^i < \nu_m \leq 2^{i+1}} \frac{\vartheta_m \max(1, \omega(m))}{\sigma_m \sqrt{\nu_m}} \right).$$

*In particular,*

$$(5.21) \quad \mathbb{E} \left| \sum_{I \leq i < J} Z_i \right|^2 \leq C_X \Phi(M_J) \left( \sum_{I \leq i \leq J} m_i \right).$$

*Proof.* Let  $0 < c < 1$ . By Corollary 2.6, and assumption (1.3), there exists a constant  $C_c$  such that for all  $1 \leq \nu_m \leq c\nu_n$ ,

$$|\mathbb{E} Y_n Y_m| \leq C_{c,D} h(n) \sqrt{\frac{\nu_m}{\nu_n}}.$$

Recall that

$$\mathbb{E} Y_n^2 = \sigma_n^2 \mathbb{P}\{S_n = \kappa_n\} (1 - \mathbb{P}\{S_n = \kappa_n\}) = \mathcal{O}(\sigma_n)$$

by assumption (2.1)-(2).

*Step 1.* We first bound the rectangle sums

$$(5.22) \quad \sum_{I \leq i < j-1 < J} |\mathbb{E} Z_i Z_j|.$$

Let  $1 \leq i < j-1$ . Then

$$(5.23) \quad \begin{aligned} |\mathbb{E} Z_i Z_j| &\leq \sum_{\substack{2^i \leq \nu_m < 2^{i+1} \\ 2^j \leq \nu_n < 2^{j+1}}} \frac{\vartheta_m \vartheta_n}{\sigma_m \sqrt{\nu_m} \sigma_n \sqrt{\nu_n}} |\mathbb{E} Y_n Y_m| \\ &\leq C_D h(2^{j+1}) \sum_{\substack{2^i \leq \nu_m < 2^{i+1} \\ 2^j \leq \nu_n < 2^{j+1}}} \frac{\vartheta_m \vartheta_n}{\sigma_n \sigma_m \sqrt{\nu_m \nu_n}} \sqrt{\frac{\nu_m}{\nu_n}} \\ &= C_D h(2^{j+1}) \sum_{2^i \leq \nu_m < 2^{i+1}} \frac{\vartheta_m}{\sigma_m \sqrt{\nu_m}} \left( \sqrt{\nu_m} \sum_{2^j \leq \nu_n < 2^{j+1}} \frac{\vartheta_n}{\sigma_n \nu_n} \right). \end{aligned}$$

Noting that  $\nu_n \geq \nu_m$  implies  $n \geq m$  by assumption (1.19), so that  $\nu_n \geq 2^j$  and  $\nu_m < 2^{i+1}$ , imply  $n \geq m$ , we have

$$\begin{aligned}
& \sum_{I \leq i < j-1 < J} |\mathbb{E} Z_i Z_j| \\
& \leq C_D h(2^{J+1}) \sum_{I \leq i < j-1 < J} \sum_{2^i \leq \nu_m < 2^{i+1}} \frac{\vartheta_m}{\sigma_m \sqrt{\nu_m}} \left( \sqrt{\nu_m} \sum_{2^j \leq \nu_n < 2^{j+1}} \frac{\vartheta_n}{\sigma_n \nu_n} \right) \\
& \leq C_D h(2^{J+1}) \sum_{I \leq i \leq J} \sum_{2^i \leq \nu_m < 2^{i+1}} \frac{\vartheta_m}{\sigma_m \sqrt{\nu_m}} \left( \sqrt{\nu_m} \sum_{n \geq m} \frac{\vartheta_n}{\sigma_n \nu_n} \right) \\
(5.24) \quad & \leq C_D h(2^{J+1}) \left( \max_{m \leq 2^{J+1}} \sqrt{\nu_m} \sum_{n \geq m} \frac{\vartheta_n}{\sigma_n \nu_n} \right) \left( \sum_{I \leq i \leq J} m_i \right).
\end{aligned}$$

To control the mid-term, we appeal to Lemma 5.6. By applying it with  $\delta = \frac{1}{2}$ , we get

$$(5.25) \quad \nu_m^{1/2} \sum_{n \geq m} \frac{\vartheta_n}{\sigma_n \nu_n} \leq C \sqrt{\nu_m} \sum_{n \geq m} \frac{\vartheta_n}{\nu_n^{3/2}} \leq C \nu_m^{1/2} \frac{1}{\nu_m^{1/2}} = C.$$

Therefore

$$(5.26) \quad \sum_{I \leq i < j-1 < J} |\mathbb{E} Z_i Z_j| \leq C_D h(2^{J+1}) \left( \sum_{I \leq i \leq J} m_i \right).$$

*Step 2.* Now we bound the square sum

$$(5.27) \quad \sum_{I \leq i \leq J} \mathbb{E} Z_i^2.$$

We have

$$\begin{aligned}
\mathbb{E} |Z_i|^2 &= \mathbb{E} \left| \sum_{2^i \leq \nu_m < 2^{i+1}} \frac{\vartheta_m Y_m}{\sigma_m \sqrt{\nu_m}} \right|^2 \\
&= 2 \sum_{2^i \leq \nu_m < \nu_n < 2^{i+1}} \frac{\vartheta_m \vartheta_n \mathbb{E} Y_m Y_n}{\sigma_m \sigma_n \sqrt{\nu_m \nu_n}} + \sum_{2^i \leq \nu_m < 2^{i+1}} \frac{\vartheta_m \mathbb{E} Y_m^2}{\sigma_m^2 \nu_m} \\
(5.28) \quad &:= I_1 + I_2.
\end{aligned}$$

At first we deduce from (4.7),

$$I_2 = \sum_{2^i \leq \nu_m < 2^{i+1}} \frac{\vartheta_m \mathbb{E} Y_m^2}{\sigma_m^2 \nu_m} \leq C \sum_{2^i \leq \nu_m < 2^{i+1}} \frac{\vartheta_m}{\sigma_m \nu_m} \leq C \sum_{2^i \leq \nu_m < 2^{i+1}} \frac{\vartheta_m}{\sigma_m \sqrt{\nu_m}}.$$

Now

$$(5.29) \quad |I_1| \leq 2 \sum_{2^i \leq \nu_m < \nu_n < 2^{i+1}} \frac{\vartheta_m \vartheta_n |\mathbb{E} Y_m Y_n|}{\sigma_m \sigma_n \sqrt{\nu_m \nu_n}}$$

By (4.5)

$$|\mathbb{E} Y_m Y_n| \leq C_\vartheta \left( \frac{\sigma_n}{\sqrt{\sigma_n^2 - \sigma_m^2}} + 1 \right), \quad (n > m \geq 1)$$

so that

$$\sum_{2^i \leq \nu_m < \nu_n < 2^{i+1}} \frac{\vartheta_m \vartheta_n |\mathbb{E} Y_m Y_n|}{\sigma_m \sigma_n \sqrt{\nu_m \nu_n}}$$

$$\begin{aligned}
&\leq C_\vartheta \sum_{2^i \leq \nu_m < \nu_n < 2^{i+1}} \frac{\vartheta_m \vartheta_n}{\sigma_m \sigma_n \sqrt{\nu_m \nu_n}} \left( \frac{\sigma_n}{\sqrt{\sigma_n^2 - \sigma_m^2}} + 1 \right) \\
&\leq C_\vartheta \sum_{2^i < \nu_m \leq 2^{i+1}} \frac{\vartheta_m}{\sigma_m \sqrt{\nu_m}} \left\{ \sum_{\nu_m < \nu_n < 2^{i+1}} \frac{\vartheta_n}{\sqrt{\nu_n}} \left( \frac{1}{\sqrt{\sigma_n^2 - \sigma_m^2}} \right) \right\} \\
&\quad + C_\vartheta \left( \sum_{2^i \leq \nu_m < 2^{i+1}} \frac{\vartheta_m}{\sigma_m \sqrt{\nu_m}} \right)^2 \\
&= C_\vartheta \left( \sum_{2^i < \nu_m \leq 2^{i+1}} \frac{\vartheta_m \omega(m)}{\sigma_m \sqrt{\nu_m}} \right) + C_\vartheta \left( \sum_{2^i \leq \nu_m < 2^{i+1}} \frac{\vartheta_m}{\sigma_m \sqrt{\nu_m}} \right)^2 \\
(5.30) \quad &\leq C_\vartheta \left( \sum_{2^i < \nu_m \leq 2^{i+1}} \frac{\vartheta_m \max(1, \omega(m))}{\sigma_m \sqrt{\nu_m}} \right),
\end{aligned}$$

since  $\sum_{2^i \leq \nu_m < 2^{i+1}} \frac{\vartheta_m}{\sigma_m \sqrt{\nu_m}} = \mathcal{O}(1)$  by assumption.

We therefore get

$$(5.31) \quad \mathbb{E} Z_i^2 \leq C_\vartheta \left( \sum_{2^i < \nu_m \leq 2^{i+1}} \frac{\vartheta_m \max(1, \omega(m))}{\sigma_m \sqrt{\nu_m}} \right),$$

and

$$(5.32) \quad \sum_{I \leq i \leq J} \mathbb{E} Z_i^2 \leq C_\vartheta \sum_{I \leq i \leq J} \left( \sum_{2^i < \nu_m \leq 2^{i+1}} \frac{\vartheta_m \max(1, \omega(m))}{\sigma_m \sqrt{\nu_m}} \right).$$

Further

$$\begin{aligned}
\mathbb{E} |Z_i Z_{i+1}| &\leq \|Z_i\|_2 \|Z_{i+1}\|_2 \leq \frac{1}{2} \sum_{h \in \{i, i+1\}} \|Z_h\|_2^2 \\
(5.33) \quad &\leq C_\vartheta \left( \sum_{2^i < \nu_m \leq 2^{i+1}} \frac{\vartheta_m \max(1, \omega(m))}{\sigma_m \sqrt{\nu_m}} \right).
\end{aligned}$$

*Step 3.* By combining (5.32) and (5.33) with estimate (5.26) we get,

$$\begin{aligned}
\mathbb{E} \left| \sum_{I \leq i < J} Z_i \right|^2 &\leq C_D h(2^{J+1}) \sum_{I \leq i \leq J} \left( \sum_{2^i < \nu_m \leq 2^{i+1}} \frac{\vartheta_m \max(1, \omega(m))}{\sigma_m \sqrt{\nu_m}} \right) \\
&\leq C_D h(2^{J+1}) \left( \max_{m \leq 2^{J+1}} (1, \omega(m)) \right) \sum_{I \leq i \leq J} \left( \sum_{2^i < \nu_m \leq 2^{i+1}} \frac{\vartheta_m}{\sigma_m \sqrt{\nu_m}} \right) \\
(5.34) \quad &= C_D \Phi(M_J) \left( \sum_{I \leq i \leq J} m_i \right).
\end{aligned}$$

At this stage we make a useful observation.

**Remark 5.9** (*R*-geometric blocks). Let  $R > 1$  and consider the sequence  $\mathcal{Z}(R) = \{Z_{R,i}, i \geq 1\}$  of sums of *R*-geometric blocks defined as follows,

$$(5.35) \quad Z_{R,i} = \sum_{R^i \leq \nu_n < R^{i+1}} \frac{\vartheta_n Y_n}{\sigma_n \sqrt{\nu_n}}, \quad i \geq 1.$$



Correspondingly, we note

$$(5.36) \quad m_{R,i} = \sum_{R^i \leq \nu_n < R^{i+1}} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}}, \quad i \geq 1,$$

and  $\mathcal{M}_R$  the sequence

$$(5.37) \quad \mathcal{M}_R = \{M_{R,J}, J \geq 1\} \quad \text{where} \quad M_{R,J} = M(R^J) = \sum_{1 \leq i < J} m_{R,i}, \quad J \geq 1.$$

For controlling rectangle sums (5.22) of 2-geometric blocks,

$$\sum_{I \leq i < j-1 < J} |\mathbb{E} Z_i Z_j|,$$

we have used Corollary 2.6, and assumption (1.3). Next we controlled separately  $\mathbb{E} |Z_i Z_{i+1}|$  in (5.33), by using Cauchy-Schwarz's inequality and an elementary inequality. If in place we had rectangle sums of  $R$ -geometric blocks, with  $1 < R < 2$ , then this can be done quite similarly by first controlling the modified rectangle sums

$$(5.38) \quad \sum_{I \leq i < j - \delta(R) < J} |\mathbb{E} Z_i Z_j|,$$

for some suitable integer  $\delta(R) > 1$  depending on  $R$  only. Next we can control  $\mathbb{E} |Z_i Z_{i+h}|$ ,  $h = 1, \dots, \delta(R)$  correspondingly as before. Finally we control the square sums of  $R$ -geometric blocks the same way as in *Step 2* for the sum (5.27). Consequently we control by using the same proof the sums of  $R$ -geometric blocks. For simplicity of the notation we only write it for 2-geometric blocks.

By Remark 5.9, we also have

$$(5.39) \quad \mathbb{E} \left| \sum_{I \leq i < J} Z_{R,i} \right|^2 \leq C_{D,R} \Phi(M(R^J)) \left( \sum_{I \leq i \leq J} m_{R,i} \right).$$

□

**Subsequence case.** Let  $\mathcal{N} \subseteq \mathbb{N}$  be an increasing sequence of positive integers and consider the  $\mathcal{N}$ -restricted blocks

$$(5.40) \quad Z_i^{\mathcal{N}} = \sum_{\substack{2^i \leq \nu_n < 2^{i+1} \\ n \in \mathcal{N}}} \frac{\vartheta_n Y_n}{\sigma_n \sqrt{\nu_n}}, \quad i \geq 1.$$

Let also

$$(5.41) \quad m_i^{\mathcal{N}} = \sum_{\substack{2^i \leq \nu_n < 2^{i+1} \\ n \in \mathcal{N}}} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}}, \quad i \geq 1,$$

and define correspondingly  $\mathcal{M}^{\mathcal{N}}$ , and  $\mathcal{M}_j^{\mathcal{N}}$ ,  $j \geq 1$ , as before.

**Theorem 5.10** (Subsequences). *Assume that*

$$(5.42) \quad m_i^{\mathcal{N}} = \mathcal{O}(1), \quad i \rightarrow \infty.$$

There exists a constant  $C_X$  such that for any  $J \geq I \geq 1$ ,

$$(5.43) \quad \mathbb{E} \left| \sum_{I \leq i < J} Z_i^{\mathcal{N}} \right|^2 \leq C_X h(2^{J+1}) \sum_{I \leq i \leq J} \left( \sum_{\substack{2^i < \nu_m \leq 2^{i+1} \\ m \in \mathcal{N}}} \frac{\vartheta_m \max(1, \omega(m))}{\sigma_m \sqrt{\nu_m}} \right).$$

In particular,

$$(5.44) \quad \mathbb{E} \left| \sum_{I \leq i < J} Z_i^{\mathcal{N}} \right|^2 \leq C_X \Phi(M_J) \left( \sum_{I \leq i \leq J} m_i^{\mathcal{N}} \right).$$

It is of matter to indicate that applications to corresponding subsequence-ASLLT's, will require the following modification of assumption 5.46 to hold:

$$(5.45) \quad \sum_{n \in \mathcal{N}} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}} = \infty,$$

also that  $m_i^{\mathcal{N}}$  might tend to 0, even in the i.i.d. case, and (see (4.2)) that  $\kappa_n$  defines  $Y_n$ .

*Proof.* It is a simple adaptation of the previous proof. In (5.23), we bounded the correlation  $|\mathbb{E} Z_i Z_j|$  by the sum

$$\sum_{\substack{2^i \leq \nu_m < 2^{i+1} \\ 2^j \leq \nu_n < 2^{j+1}}} \frac{\vartheta_m \vartheta_n}{\sigma_m \sqrt{\nu_m} \sigma_n \sqrt{\nu_n}} |\mathbb{E} Y_n Y_m|,$$

next  $\sum_{I \leq i < j-1 < J} |\mathbb{E} Z_i Z_j|$  by the sum

$$\sum_{I \leq i < j-1 < J} \sum_{\substack{2^i \leq \nu_m < 2^{i+1} \\ 2^j \leq \nu_n < 2^{j+1}}} \frac{\vartheta_m \vartheta_n}{\sigma_m \sqrt{\nu_m} \sigma_n \sqrt{\nu_n}} |\mathbb{E} Y_n Y_m|.$$

which is shown in (5.24) to be bounded by  $\sum_{I \leq i \leq J} m_i$ . Similarly, in (5.29), we controlled  $|I_1|$  by the sum

$$\sum_{2^i \leq \nu_m < \nu_n < 2^{i+1}} \frac{\vartheta_m \vartheta_n |\mathbb{E} Y_m Y_n|}{\sigma_m \sigma_n \sqrt{\nu_m \nu_n}}$$

with terms  $|\mathbb{E} Y_m Y_n|$ , which is next bounded in (5.30), by

$$C_{\vartheta} \left( \sum_{2^i < \nu_m \leq 2^{i+1}} \frac{\vartheta_m \max(1, \omega(m))}{\sigma_m \sqrt{\nu_m}} \right).$$

The control of the sum  $I_2$  raises no peculiar problem. It is clear that all this is transferable *ipso facto* to the subsequence case by pointing at each place where  $\nu_n$  appears, that  $n \in \mathcal{N}$ , and next collecting the modified bounds.  $\square$

**5.6. The ASLLT.** We now are in position to state the main result of this Section.

**Theorem 5.11.** *Let  $\{\vartheta_n, n \geq 1\}$  be chosen so that (2.4) holds. Suppose that*

$$(5.46) \quad \text{The series } \sum_{n \geq 1} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}} \text{ is divergent.}$$

*We also assume that the following condition is fulfilled:*

$$(5.47) \quad \text{The series } s^2 := \sum_{\substack{l \geq 1 \\ [2^l, 2^{l+1}[ \cap \mathcal{M} \neq \emptyset}} \frac{\Phi(2^l) (1 + \log \# \{[2^l, 2^{l+1}[ \cap \mathcal{M}\})}{2^l} \text{ is convergent.}$$

Let  $\kappa_n \in \mathcal{L}(nv_0, D)$ ,  $n = 1, 2, \dots$  be a sequence of integers such that (2.1) holds.

(i) We have (recalling that  $M_j = \sum_{1 \leq n < 2^j} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}}$ ),

$$\left\| \sum_{\substack{k \geq 1 \\ [2^k, 2^{k+1}] \cap \mathcal{M} \neq \emptyset}} \sup_{2^k \leq M_j < 2^{k+1}} \left\{ \frac{\left| \sum_{1 \leq n < 2^j} \frac{\vartheta_n (\mathbf{1}_{\{S_n = \kappa_n\}} - \mathbb{P}\{S_n = \kappa_n\})}{\sqrt{\nu_n}} \right|}{M_j} \right\} \right\|_2 \leq Ks.$$

Further

$$(5.48) \quad \lim_{j \rightarrow \infty} \frac{1}{\sum_{1 \leq n < 2^j} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}}} \sum_{1 \leq n < 2^j} \frac{\vartheta_n (\mathbf{1}_{\{S_n = \kappa_n\}} - \mathbb{P}\{S_n = \kappa_n\})}{\sqrt{\nu_n}} \stackrel{\text{a.s.}}{=} 0.$$

(ii) Assume that

$$(5.49) \quad \lim_{n \rightarrow \infty} \sigma_n \mathbb{P}\{S_n = \kappa_n\} = \gamma, \quad (0 < \gamma < \infty).$$

Then we have

$$(5.50) \quad \lim_{j \rightarrow \infty} \frac{1}{\sum_{1 \leq n < 2^j} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}}} \sum_{1 \leq n < 2^j} \frac{\vartheta_n \mathbf{1}_{\{S_n = \kappa_n\}}}{\sqrt{\nu_n}} \stackrel{\text{a.s.}}{=} \gamma.$$

(iii) Assume that

$$(5.51) \quad \lim_{n \rightarrow \infty} \sigma_n \mathbb{P}\{S_n = \kappa_n\} = \frac{D}{\sqrt{2\pi}} e^{-\frac{\kappa^2}{2}},$$

and that  $M(t)$  is slowly varying near infinity. Then the ASLLT holds,

$$(5.52) \quad \lim_{N \rightarrow \infty} \frac{1}{\sum_{1 \leq n < N} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}}} \sum_{1 \leq n < N} \frac{\vartheta_n \mathbf{1}_{\{S_n = \kappa_n\}}}{\sqrt{\nu_n}} \stackrel{\text{a.s.}}{=} \frac{D}{\sqrt{2\pi}} e^{-\frac{\kappa^2}{2}}.$$

(iv) In particular, if the LLT is applicable to the sequence  $X$ , then for any sequence  $\{\kappa_n, n \geq 1\}$  such that

$$\lim_{n \rightarrow \infty} \frac{\kappa_n - a_n}{\sigma_n} = \kappa.$$

In addition if  $M(t)$  is slowly varying near infinity, then (5.52) holds.

(v) (Optimality of Assumption (5.46)) Assume that  $\Phi(x) = \mathcal{O}(1)$  and

$$(5.53) \quad \sum_{n=1}^{\infty} \frac{\vartheta_n (\log \log n)^2}{\sigma_n \sqrt{\nu_n}} < \infty.$$

Then a strong but different almost sure convergence result takes place, namely that the sequence

$$\sum_{n < 2^J} \frac{\mathbf{1}_{\{S_n = \kappa_n\}} - \mathbb{P}\{S_n = \kappa_n\}}{\sqrt{\nu_n}} \quad J = 1, 2, \dots,$$

converges almost surely.

*Proof.* By Theorem 5.8 the sequence  $\mathcal{Z}$  satisfies condition (3.16) with the choice  $\gamma = 2$ ,  $u_l = m_l$ ,  $\Psi(x) = Cx$  for some suitable finite, positive constant  $C$ , and  $\Phi(x)$  defined in (5.18). In view of assumption (5.47), Theorem 3.7 is in force.

Therefore

$$\mathcal{S}^2 = \sum_{k \geq 1} \sup \left\{ \frac{\left| \sum_{1 \leq \nu_n < 2^{j+1}} \frac{\vartheta_n Y_n}{\sigma_n \sqrt{\nu_n}} \right|}{\sum_{1 \leq \nu_n < 2^j} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}}} : 2^k \leq \sum_{1 \leq \nu_n < 2^j} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}} < 2^{k+1} \right\}^2,$$

satisfies

$$\|\mathcal{S}\|_2 \leq Ks, \quad \text{and} \quad \lim_{j \rightarrow \infty} \frac{1}{\sum_{1 \leq \nu_n < 2^j} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}}} \sum_{1 \leq i < j} \sum_{2^i \leq \nu_n < 2^{i+1}} \frac{\vartheta_n Y_n}{\sigma_n \sqrt{\nu_n}} \stackrel{\text{a.s.}}{=} 0.$$

Recalling (4.2),  $Y_n = \sigma_n(\mathbf{1}_{\{S_n = \kappa_n\}} - \mathbb{P}\{S_n = \kappa_n\})$ , the latter means

$$(5.54) \quad \lim_{j \rightarrow \infty} \frac{1}{\sum_{1 \leq \nu_n < 2^j} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}}} \sum_{1 \leq \nu_n < 2^j} \frac{\vartheta_n(\mathbf{1}_{\{S_n = \kappa_n\}} - \mathbb{P}\{S_n = \kappa_n\})}{\sqrt{\nu_n}} \stackrel{\text{a.s.}}{=} 0.$$

In particular, (5.50) is immediate. Now if

$$(5.55) \quad \lim_{n \rightarrow \infty} \sigma_n \mathbb{P}\{S_n = \kappa_n\} = \frac{D}{\sqrt{2\pi}} e^{-\frac{\kappa^2}{2}},$$

then

$$(5.56) \quad \lim_{j \rightarrow \infty} \frac{1}{\sum_{1 \leq \nu_n < 2^j} \frac{\vartheta_n}{\sigma_n \sqrt{\nu_n}}} \sum_{1 \leq \nu_n < 2^j} \frac{\vartheta_n \mathbf{1}_{\{S_n = \kappa_n\}}}{\sqrt{\nu_n}} \stackrel{\text{a.s.}}{=} \frac{D}{\sqrt{2\pi}} e^{-\frac{\kappa^2}{2}}.$$

This proves assertions (i) and (ii) of the Theorem.

As to assertions (iii)-(iv), we use the fact (see Remark 5.9 and estimate (5.39)) that the proof given remains true if instead of sieving by the geometric sequence  $2^i, i \geq 1$ , we sieve with the geometric sequence  $R^i, i \geq 1$ , where  $R > 1$  is arbitrary; that is (5.48), (5.50) are true for these sequences. Next using (1.10) and that  $M(t)$  is slowly varying near infinity allows to conclude to (5.52).

(v) The attentive reader will have certainly observed that assumption (5.46) is not required in the proof of Theorem 5.8, but only in the application of Proposition 3.7. Consider the case where  $\sum_{j=1}^{\infty} m_j < \infty$  and assume that  $\Phi(x) = \mathcal{O}(1)$ . See Example 5.7-(i). In that case the above inequality takes the simpler form

$$\mathbb{E} \left| \sum_{I \leq i \leq J} Z_i \right|^2 \leq C_{c,D} \left( \sum_{I \leq i \leq J} m_i \right).$$

Assume only a little more, namely that  $\sum_{i=1}^{\infty} m_i (\log i)^2 < \infty$ , which by definition of  $m_i$  in (5.14) means that

$$\sum_{n=1}^{\infty} \frac{\vartheta_n (\log \log n)^2}{\sigma_n \sqrt{\nu_n}} < \infty,$$

which is (5.53). We use Remark 3.11, and more precisely, implication (3.22), thank to which it follows that the series  $\sum_{l=1}^{\infty} Z_l$ , namely the sequence

$$\sum_{n < 2^J} \frac{Y_n}{\sigma_n \sqrt{\nu_n}} \quad J = 1, 2, \dots,$$

converges almost surely. Therefore as  $Y_n = \sigma_n(\mathbf{1}_{\{S_n=\kappa_n\}} - \mathbb{P}\{S_n = \kappa_n\})$ , the sequence

$$\sum_{n < 2^J} \frac{\mathbf{1}_{\{S_n=\kappa_n\}} - \mathbb{P}\{S_n = \kappa_n\}}{\sqrt{\nu_n}} \quad J = 1, 2, \dots,$$

converges almost surely. Consequently, if assumption (5.53) is fulfilled, a strong but different almost sure convergence result takes place.  $\square$

**5.7. The quasi-orthogonal case.** We prove the following

**Theorem 5.12.** *Assume that  $h(x) = \mathcal{O}(1)$ ,  $m_i \asymp (1)$  and further that condition (5.5) is satisfied. Then the sequence  $\{Z_i, i \geq 1\}$  is a quasi-orthogonal system.*

*Proof.* Let  $i, j$  be positive integers such that either  $j > i + 1$  or  $j < i - 1$ . Obviously

$$\begin{aligned} |\mathbb{E} Z_i Z_j| &\leq \sum_{\substack{2^i \leq \nu_m < 2^{i+1} \\ 2^j \leq \nu_n < 2^{j+1}}} \frac{\vartheta_m \vartheta_n}{\sigma_m \sqrt{\nu_m} \sigma_n \sqrt{\nu_n}} |\mathbb{E} Y_n Y_m| \\ &\leq C_D \sum_{\substack{2^i \leq \nu_m < 2^{i+1} \\ 2^j \leq \nu_n < 2^{j+1}}} \frac{\vartheta_m \vartheta_n}{\sigma_n \sigma_m \sqrt{\nu_m \nu_n}} \sqrt{\frac{\nu_m}{\nu_n}} \\ &\leq C_D 2^{-(j-i)/2} \left( \sum_{2^i \leq \nu_m < 2^{i+1}} \frac{\vartheta_m}{\sigma_m \sqrt{\nu_m}} \right) \left( \sum_{2^j \leq \nu_n < 2^{j+1}} \frac{\vartheta_n}{\sigma_n \nu_n} \right) \\ &= C_D 2^{-(j-i)/2} m_i m_j. \end{aligned}$$

Therefore,

$$\sum_{|j-i|>1} \frac{|\mathbb{E} Z_i Z_j|}{m_i m_j} \leq C_D H.$$

In view of (5.31) and the assumptions made,

$$\mathbb{E} Z_i^2 \leq C_\vartheta \left( \sum_{2^i < \nu_m \leq 2^{i+1}} \frac{\vartheta_m}{\sigma_m \sqrt{\nu_m}} \right) = C_\vartheta m_i.$$

Since  $m_i \asymp (1)$  we have,

$$\mathbb{E} \left( \frac{Z_i}{m_i} \right)^2 \leq \frac{C_\vartheta}{m_i} \leq C_\vartheta.$$

Consequently

$$(5.57) \quad \sup_{i \geq 1} \sum_{j \geq 1} |\mathbb{E} Z_i Z_j| \leq C_D H.$$

This shows by using criterion (3.13), that the sequence  $\{Z_i, i \geq 1\}$  is a quasi-orthogonal system.  $\square$

Note that (see Example 5.7-(i)) in the i.i.d. case,  $h(x) \equiv C$  (by Remark 5.2), and condition (5.5) holds. Further

$$m_i = \sum_{2^i \leq \nu_m < 2^{i+1}} \frac{\vartheta_m}{\sigma_m \sqrt{\nu_m}} = \sum_{2^i \leq \nu_m < 2^{i+1}} \frac{\nu}{m} = \text{const.}$$

Therefore we have

**Corollary 5.13** (i.i.d. case). *The corresponding sequence  $\{Z_i, i \geq 1\}$  is a quasi-orthogonal system.*

This is no longer true when passing to subsequences! See Theorem 5.10 and comments hereafter.

Another consequence is

**Corollary 5.14** (a.s. convergent series and asllt). *Let  $b > 3/2$ . Then the series*

$$(5.58) \quad \sum_{j>1} \frac{1}{j^{1/2}(\log j)^b} \left( \sum_{m=2^{j-1}}^{2^j-1} \frac{\mathbf{1}_{\{S_m=\kappa_m\}} - \mathbb{P}\{S_m = \kappa_m\}}{\sqrt{m}} \right),$$

*converges almost surely.*

*Also,*

$$(5.59) \quad \lim_{N \rightarrow \infty} \frac{1}{N^{1/2}(\log N)^b} \sum_{j=1}^{2^N} \frac{\mathbf{1}_{\{S_m=\kappa_m\}} - \mathbb{P}\{S_m = \kappa_m\}}{\sqrt{m}} = 0,$$

*almost surely.*

*Further,*

$$\lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n \leq N} \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n=\kappa_n\}} \stackrel{a.s.}{=} \frac{D}{\sqrt{2\pi}\sigma} e^{-\kappa^2/(2\sigma^2)},$$

*for any sequence of integers  $\kappa_n \in \mathcal{L}(nv_0, D)$ ,  $n = 1, 2, \dots$  such that (1.9) holds.*

The first claim ([55]) cannot be derived from Gál–Koksma’s criterion ([45], p.134), unlike to the second one. The third one is Theorem 1.1.

*Proof.* It is very short. As Rademacher–Menchov’s Theorem applies to quasi-orthogonal systems, the series

$$\sum_j \frac{Z_j}{j^{1/2}(\log j)^b}$$

converges almost surely if  $b > 3/2$ . By Kronecker’s Lemma,  $\frac{1}{N^{1/2}(\log N)^b} \sum_{j=1}^N Z_j \rightarrow 0$ , as  $N$  tends to infinity, almost surely, whence the second claim. By arguing as in Remark 5.9, this remains true along any  $R$ -geometric sequence,  $R > 1$ . Now by Gnedenko’s Theorem (1.4), if  $\kappa_n \in \mathcal{L}(nv_0, D)$  is a sequence which verifies condition (1.9), then (1.10) holds. The proof is achieved by applying the following well-known elementary Lemma.

**Lemma 5.15.** *Let  $\{f_n, n \geq 1\}$  be a sequence of nonnegative numbers. If for each  $r > 1$ , the averages  $r^{-k} \sum_{n \leq r^k} f_n$  converges, as  $k \rightarrow \infty$ . Then for each  $r > 1$  this limit is the same, call it  $L$ , and we have  $\lim_{N \rightarrow \infty} N^{-1} \sum_{n \leq N} f_n = L$ .*

□

We pass to another example.

**Corollary 5.16** (Cramér model). *Let  $\xi = \{\xi_j, j \geq 1\}$  be the sequence defined in Example 5.4. Then the corresponding sequence  $\{Z_i, i \geq 1\}$  is a quasi-orthogonal system.*

*Proof.* Recall that  $\vartheta_m = \frac{1}{\log m}$ , and  $\sigma_m^2 \sim \nu_m \sim \frac{m}{\log m}$ , as  $m \rightarrow \infty$ . Thus  $h(x) \equiv C$ , further

$$\begin{aligned} m_i &\sim \sum_{2^i \leq \nu_m < 2^{i+1}} \frac{\log m}{(\log m)m} \sim \sum_{2^i \leq \frac{m}{\log m} < 2^{i+1}} \frac{1}{m} \\ &\sim \sum_{i2^i \leq m < (i+1)2^{i+1}} \frac{1}{m} \sim \log(i+1) + (i+1)\log 2 - \log i - i\log 2 \sim \text{const.} \end{aligned}$$

□

## 6. REVISITING THE ASLLT IN THE I.I.D. CASE

The goal of this Section is to prove that in the i. i. d. case, an almost sure local limit theorem with explicit speed of convergence (Theorem 6.1) expressed with an a.s. convergent series. Let  $X$  be an  $\mathcal{L}(v_0, D)$ -valued square integrable random variable, with maximal span  $D$ ,  $\mu = \mathbb{E} X$ ,  $\sigma^2 = \text{Var}(X) > 0$ . Assume that  $\vartheta_X > 0$  and choose  $\vartheta = \vartheta_X$ . Let  $S_n = X_1 + \dots + X_n$ ,  $X_k$  being independent copies of  $X$ . Thus  $\nu_n = \vartheta n$ ,  $\sigma_n = \sigma\sqrt{n}$  and assumptions 1.3 and 5.46 are satisfied.

Let  $\kappa_j \in \mathcal{L}(jv_0, D)$ ,  $j = 1, 2, \dots$  be such that assumption (2.7) is satisfied. The system of random variables (4.37) writes:

$$(6.1) \quad \frac{\vartheta_n Y_n}{\sigma_n \sqrt{\nu_n}} = \frac{\vartheta Y_n}{n} = \vartheta \frac{\mathbf{1}_{\{S_n = \kappa_n\}} - \mathbb{P}\{S_n = \kappa_n\}}{\sqrt{n}}, \quad n \geq 1,$$

$$(6.2) \quad Z_i = \sum_{m=2^{i-1}}^{2^i-1} \frac{Y_m}{m} = \sigma \sum_{m=2^{i-1}}^{2^i-1} \frac{\mathbf{1}_{\{S_m = \kappa_m\}} - \mathbb{P}\{S_m = \kappa_m\}}{\sqrt{m}}, \quad i \geq 1.$$

A natural question is whether it is possible in (5.58) to substitute to  $\mathbb{P}\{S_m = \kappa_m\}$ , its limit  $\frac{D}{\sqrt{2\pi}\sigma} e^{-\kappa^2/2\sigma^2}$ . Let

$$(6.3) \quad \delta_n = \left| \sigma\sqrt{n} \mathbb{P}\{S_n = \kappa_n\} - \frac{D}{\sqrt{2\pi}} e^{-\frac{(\kappa_n - n\mu)^2}{2n\sigma^2}} \right|.$$

By Gnedenko's Theorem (1.4),  $\delta_n = o(1)$ . Recall that  $\vartheta_X$  is defined in (1.13).

**Theorem 6.1.** *Assume that  $\vartheta_X > 0$ . Let  $\kappa_n = n\mu + \kappa\sqrt{n}(1 + \varepsilon_n)$ ,  $\varepsilon_n \rightarrow 0$ .*

(i) *We have*

$$\frac{\sum_{1 \leq n < 2^{N+1}} \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}}}{\sum_{1 \leq n < 2^{N+1}} \frac{1}{n}} = \frac{D}{\sigma\sqrt{2\pi}} e^{-\frac{\kappa^2}{2\sigma^2}} + \mathcal{O}\left(\frac{1}{N} \sum_{1 \leq n < 2^{N+1}} \frac{\delta_n + \varepsilon_n}{n}\right) + o\left(\frac{(\log N)^b}{N^{1/2}}\right).$$

(ii) *Assume that*

$$\sum_j \frac{1}{j^{1/2}(\log j)^b} \sum_{2^j \leq n < 2^{j+1}} \frac{\delta_n + \varepsilon_n}{n} < \infty.$$

*This is fulfilled if  $\max(\delta_n, \varepsilon_n) = \mathcal{O}(\log^{-1/2} n)$ . Then the series*

$$\sum_j \frac{1}{j^{1/2}(\log j)^b} \left\{ \sum_{2^j \leq n < 2^{j+1}} \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}} - \left( \frac{D \log 2}{\sigma\sqrt{2\pi}} e^{-\frac{\kappa^2}{2\sigma^2}} \right) \right\}$$

*converges almost surely.*

*Proof.* We have

$$\left| e^{-\frac{(\kappa_n - n\mu)^2}{2n\sigma^2}} - e^{-\frac{\kappa^2}{2\sigma^2}} \right| \leq \frac{1}{2\sigma^2} \left| \frac{(\kappa_n - n\mu)^2}{n} - \kappa^2 \right| = \frac{\kappa^2}{2\sigma^2} |(1 + \varepsilon_n)^2 - 1| \leq \frac{3\kappa^2}{2\sigma^2} \varepsilon_n.$$

Thus

$$\begin{aligned} & \left| \sigma\sqrt{n}\mathbb{P}\{S_n = \kappa_n\} - \frac{D}{\sqrt{2\pi}}e^{-\frac{\kappa^2}{2\sigma^2}} \right| \\ & \leq \left| \sigma\sqrt{n}\mathbb{P}\{S_n = \kappa_n\} - \frac{D}{\sqrt{2\pi}}e^{-\frac{\kappa^2}{2\sigma^2}}e^{-\frac{(\kappa_n - n\mu)^2}{2n\sigma^2}} \right| + \frac{3D\kappa^2}{2\sqrt{2\pi}\sigma^2} \varepsilon_n \leq \delta_n + \frac{3\kappa^2}{2\sigma^2} \varepsilon_n. \end{aligned}$$

So that

$$\begin{aligned} \frac{Y_n}{n} &= \frac{1}{n} \left( \sqrt{n} \left\{ \mathbf{1}_{\{S_n = \kappa_n\}} - \mathbb{P}\{S_n = \kappa_n\} \right\} \right) = \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}} - \frac{\sqrt{n}\mathbb{P}\{S_n = \kappa_n\}}{n} \\ &= \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}} - \left( \frac{D}{\sigma\sqrt{2\pi}} e^{-\frac{\kappa^2}{2\sigma^2}} \right) \frac{1}{n} + \mathcal{O}\left(\frac{\delta_n + \varepsilon_n}{n}\right). \end{aligned}$$

(i) By Theorem 5.14, if  $b > 3/2$ ,

$$\frac{1}{N^{1/2}(\log N)^b} \sum_{j=1}^N Z_j \rightarrow 0,$$

as  $N$  tends to infinity, almost surely. But

$$\begin{aligned} & \frac{1}{N^{1/2}(\log N)^b} \sum_{j=1}^N Z_j = \frac{1}{N^{1/2}(\log N)^b} \sum_{1 \leq n < 2^{N+1}} \frac{Y_n}{n} \\ &= \frac{1}{N^{1/2}(\log N)^b} \sum_{1 \leq n < 2^{N+1}} \left( \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}} - \left( \frac{D}{\sigma\sqrt{2\pi}} e^{-\frac{\kappa^2}{2\sigma^2}} \right) \frac{1}{n} + \mathcal{O}\left(\frac{\delta_n + \varepsilon_n}{n}\right) \right) \\ &= \frac{1}{N^{1/2}(\log N)^b} \left( \sum_{1 \leq n < 2^{N+1}} \frac{1}{n} \right) \left\{ \frac{\sum_{1 \leq n < 2^{N+1}} \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}}}{\sum_{1 \leq n < 2^{N+1}} \frac{1}{n}} - \left( \frac{D}{\sigma\sqrt{2\pi}} e^{-\frac{\kappa^2}{2\sigma^2}} \right) \right\} \\ & \quad + \mathcal{O}\left( \frac{1}{N^{1/2}(\log N)^b} \sum_{1 \leq n < 2^{N+1}} \frac{\delta_n + \varepsilon_n}{n} \right) \rightarrow 0, \end{aligned}$$

as  $N$  tends to infinity, almost surely. Therefore  $(\sum_{1 \leq n < 2^{N+1}} \frac{1}{n} \sim CN)$ ,

$$\frac{\sum_{1 \leq n < 2^{N+1}} \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}}}{\sum_{1 \leq n < 2^{N+1}} \frac{1}{n}} = \frac{D}{\sigma\sqrt{2\pi}} e^{-\frac{\kappa^2}{2\sigma^2}} + \mathcal{O}\left(\frac{1}{N} \sum_{1 \leq n < 2^{N+1}} \frac{\delta_n + \varepsilon_n}{n}\right) + o\left(\frac{(\log N)^b}{N^{1/2}}\right).$$

(ii) Now assume that

$$\sum_j \frac{1}{j^{1/2}(\log j)^b} \sum_{2^j \leq n < 2^{j+1}} \frac{\delta_n + \varepsilon_n}{n} < \infty.$$

Then the series

$$\sum_j \frac{1}{j^{1/2}(\log j)^b} \left\{ \sum_{2^j \leq n < 2^{j+1}} \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}} - \left( \frac{D \log 2}{\sigma\sqrt{2\pi}} e^{-\frac{\kappa^2}{2\sigma^2}} \right) \right\}$$

converges almost surely.



Indeed, first the series

$$\sum_j \frac{1}{j^{1/2}(\log j)^b} \left\{ \sum_{2^j \leq n < 2^{j+1}} \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}} - \left( \sum_{2^j \leq n < 2^{j+1}} \frac{1}{n} \right) \left( \frac{D}{\sigma \sqrt{2\pi}} e^{-\frac{\kappa^2}{2\sigma^2}} \right) \right\}$$

converges almost surely, since

$$\begin{aligned} \sum_j \frac{Z_j}{j^{1/2}(\log j)^b} &= \sum_j \frac{1}{j^{1/2}(\log j)^b} \sum_{2^j \leq n < 2^{j+1}} \frac{Y_n}{n} \\ &= \sum_j \frac{1}{j^{1/2}(\log j)^b} \left\{ \sum_{2^j \leq n < 2^{j+1}} \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}} - \left( \sum_{2^j \leq n < 2^{j+1}} \frac{1}{n} \right) \left( \frac{D}{\sigma \sqrt{2\pi}} e^{-\frac{\kappa^2}{2\sigma^2}} \right) \right\} \\ &\quad + \mathcal{O} \left( \sum_j \frac{1}{j^{1/2}(\log j)^b} \sum_{2^j \leq n < 2^{j+1}} \frac{\delta_n + \varepsilon_n}{n} \right). \end{aligned}$$

Besides

$$\sum_{2^j \leq n < 2^{j+1}} \frac{1}{n} = \log 2 + \mathcal{O}(2^{-j}),$$

so that the simplified series

$$\sum_j \frac{1}{j^{1/2}(\log j)^b} \left\{ \sum_{2^j \leq n < 2^{j+1}} \frac{1}{\sqrt{n}} \mathbf{1}_{\{S_n = \kappa_n\}} - \left( \frac{D \log 2}{\sigma \sqrt{2\pi}} e^{-\frac{\kappa^2}{2\sigma^2}} \right) \right\}$$

converges almost surely. □

## 7. CONCLUDING REMARKS

In a previous attempt, for proving the ASLLT (Theorem 5.11), we considered slightly different block sums,

$$Z_i = \sum_{2^i \leq n < 2^{i+1}} \frac{Y_n}{\sigma_n \sqrt{\nu_n}},$$

and operated differently in order to control their covariances; in particular the diagonal case was not treated separately. We bounded  $|\mathbb{E} Z_i Z_j|$  from above, by proceeding as follows: for  $1 \leq i \leq j$ ,

$$|\mathbb{E} Z_i Z_j| \leq \sum_{\substack{2^i \leq m < 2^{i+1} \\ 2^j \leq n < 2^{j+1} \\ 1 \leq \nu_m \leq c\nu_n}} \frac{|\mathbb{E} Y_m Y_n|}{\sigma_m \sqrt{\nu_m} \sigma_n \sqrt{\nu_n}} + \sum_{\substack{2^i \leq m < 2^{i+1} \\ 2^j \leq n < 2^{j+1} \\ \nu_m > c\nu_n}} \frac{|\mathbb{E} Y_m Y_n|}{\sigma_m \sqrt{\nu_m} \sigma_n \sqrt{\nu_n}}.$$

the first sum can be estimated efficiently, the estimation of the second imposed restrictive assumptions. We have not tried any further, and switched to the one implemented in this paper. Maybe the preceding approach can still be improved, at the price of some new argument, similar to the ones used here. We don't believe it would bring more.

## REFERENCES

- [1] Alexits, G., *Convergence problems of orthogonal series*. Internat. Ser. Monogr. Pure Appl. Math. **20**, (1961), Pergamon Press, New York, Oxford, Paris.
- [2] Breuillard E., *Distribution diophantiennes et théorème limite local sur  $\mathbb{R}^d$* , Probab. Theory Related Fields **132** (2005), 39–73.
- [3] A. Broise, Transformations dilatantes de l'intervalle et théorèmes limites, *Astérisque* **238** (1996), 1–109.
- [4] Calderoni P., Campanino M. and Capocaccia D., *A local limit theorem for a sequence of interval transformations*, Ergodic Theory Dynam. Systems **5** (1985), 185–201.
- [5] Dabrowski A. R. and McDonald, D., *An application of the Bernoulli part to local limit theorems for moving averages on stationary sequences*, The Canadian Journal of Statistics **24** 3 (1996), 293–305.
- [6] Davis, B., and MacDonald, D., An elementary proof of the local central limit theorem, *J. Theoretical Prob.* **8** (1995), no. 3, 695–701.
- [7] Denker, M., and Koch, S., Almost sure local limit theorems, *Statistica Neerlandica* **56** (2002), no. 2, 143–151.
- [8] Dolgopyat D., *A local limit theorem for sums of independent random vectors*, Electron. J. Probab. **21** 39 (2016), 1–15.
- [9] Dolgopyat D. and Hafouta Y., *Edgeworth expansions for independent bounded integer valued random variables*, Stoch. Process. Appl. **152** (2022), 486–532.
- [10] Doney R. A., *A large deviation local limit theorem*, Math. Proc. Cambridge Philos. Soc. **105** (1989), 575–577.
- [11] Doney, R. A. *One-sided local large deviation and renewal theorems in the case of infinite mean*, Probab. Theory Related Fields **107** (1997), 461–465.
- [12] Doney, R. A. *A local limit theorem for moderate deviations*, Bull. London Math. Soc. **33** (2001), 100–108.
- [13] Gamkrelidze N. G., *On an inequality for multidimensional characteristic function*, Theor. Probab. Appl. **30** 1 (2001), 133–135.
- [14] Gamkrelidze N. G., *On One Inequality for Characteristic Functions*, in: Yu. V. Prokhorov, V. Statulevičius, (eds.), Limit Theorems of Probability Theory, Springer, 2000, 275–280.
- [15] Gamkrelidze N. G., *On a local limit theorem in strong sense*, Statist. Probab. Lett. **35** (1997), 79–83.
- [16] Gamkrelidze N. G., *On a probabilistic property of the Fibonacci sequence*, The Fibonacci Quartely **33** 2 (1995), 147–152.
- [17] N. G. Gamkrelidze, *A measure of “smoothness” of multidimensional distributions of integer-valued random vectors*, Theor. Probab. Appl. **30** 2 (1985), 427–431.
- [18] Giuliano, R., and Szewczak, Z. S., A general correlation inequality and the Almost Sure Local Limit Theorem for random sequences in the domain of attraction of a stable law, *Stoch. Proc. Appl.* **124** (2014), 1612–1626.
- [19] Giuliano-Antonini, R. and Szewczak, Z. S., An almost sure local limit theorem for Markov chains, *Statist. Probab. Lett.* **83** (2013) 2, 573–579.
- [20] Giuliano, R., and Weber, M., Local limit theorems in some random models from Number Theory, *Stoch. Anal. & Appl.* **34** (2016), no. 6, 941–960.
- [21] Giuliano, R., and Weber, M., Almost sure local limit theorems with rate, *Stoch. Anal. & Appl.* **29** (2011), no. 5, 779–798.
- [22] Giuliano, R., and Weber, M., Approximate local limit theorems with effective rate and application to random walks in random scenery, *Bernoulli* **23** (2017) (4B), 3268–3310.
- [23] Gnedenko, B. V., On a local limit theorem in the theory of probability, *Uspekhi Mat. Nauk.* (N.S.) **3** (1948), no. 3(25), 187–194.
- [24] Gnedenko B.V., *Course in the Theory of Probability*, 5th Ed. ”Nauka”, Moscow, English transl. of 4th ed., (1967) Chelsea, New-York.
- [25] Gouëzel S., *Berry-Esseen theorem and local limit theorem for non uniformly expanding maps*, Ann. I. H. Poincaré-PR **41** (2005), 997–1024.
- [26] Szewczak Z. S., *A de Moivre theorem revisited*, Statist. Probab. Lett. **181** (2022) 109–260.
- [27] Hardy G., Littlewood J. E., Pólya G., *Inequalities*, (1934). Cambridge Math. Library.

- [28] Hafouta Y. and Kifer Y., *A nonconventional local limit theorem*, J. Theoret. Probab. **29** (2016), 1524–1553.
- [29] Hafouta Y. and Kifer Y., *Nonconventional limit theorems and random dynamics*, World Scientific, 2018.
- [30] Ibragimov I.A., Linnik Y.V., *Independent and Stationary Sequences of Random Variables*, (1971) Wolters-Noordhoff Publishing Groningen, The Netherlands.
- [31] Jacod J., Kowalski E. and Nikeghbali A., *Mod-Gaussian convergence: new limit theorems in probability and number theory*, Forum Math. **23** 4 (2011), 835–873.
- [32] Kashin, B. S., and Saakyan, A. A. (1989) *Orthogonal series*, Transl. Math. Monogr. **75**, Amer. Math. Soc., Providence, R.I.
- [33] Kac, M., Salem, R., and Zygmund, A., A gap theorem, *Trans. Amer. Math. Soc.* **63** (1948), 235–243.
- [34] Lacey, M. and Philipp, W., A note on the almost everywhere central limit theorem. *Stat. Prob. Letters* **9** (1990), 201–205.
- [35] MacDiarmid, C., (1998). Concentration, *Prob. Methods for Algorithmic Discrete Math.*, 195–248, Algorithms Combin. **16**, Springer, Berlin.
- [36] MacDonald, D., A local limit theorem for large deviations of sums of independent, non-identically distributed random variables, *Annals of Prob.* **7** (1979) no. 3, 526–531.
- [37] MacDonald, D., On local limit theorems for integer valued random variables, *Theor. Prob. Appl.* **33** (1979), 352–355.
- [38] Mitalauskas, A. A., Local limit theorems for stable limit distributions, *Theor. Prob. Appl.* **VII** (1962) (2), 180–185.
- [39] A. B. Mukhin, *Some necessary and sufficient conditions for the validity of the local limit theorem*, Dokl. Akad. Nauk UzSSR **8** (1984), 7–8 (in Russian).
- [40] A. B. Mukhin, *Local limit theorems for distributions of sums of independent random vectors*, Theor. Prob. Appl. **29** (1984), 369–375.
- [41] A. B. Mukhin, *Local limit theorems for lattice random variables*, Theor. Prob. Appl. **36** 4 (1991), 698–713.
- [42] Olevskii, A. M. (1975) *Fourier series with respect to general orthogonal systems*, Ergeb. Math. Grenzgeb. **86**, Springer-Verlag, Berlin, Heidelberg, New York.
- [43] Petrov, V. V., *Sums of Independent Random Variables*, Ergebnisse der Math. und ihre Grenzgebiete **82**, Springer, 1975.
- [44] Macht W. and Wolf W., *The Local Limit Theorem and Hölder-Continuity*, Probab. Theory Related Fields **82** 2 (1989), 295–305.
- [45] Philipp, W., Stout W., *Almost sure invariance principles for partial sums of weakly dependent random variables*, Memoirs Amer. Math. Soc. **161** 1975.
- [46] Röllin A. and Ross N., *Local limit theorems via Landau–Kolmogorov inequalities*, Bernoulli, **21** 2 (2015), 851–881.
- [47] Rousseau-Egele J., *Un Theoreme de la Limite Locale Pour une Classe de Transformations Dilatantes et Monotones par Morceaux*, Ann. Probab. **11** 3 (1983), 772–788.
- [48] Szewczak, Z. and Weber M., *Classical and Almost Sure Local Limit Theorems*, 103 p., *Dissert. Math.*, No **589**, 1–97, (2024).
- [49] Weber, M., On Mukhin’s necessary and sufficient condition for the validity of the local limit theorem, *Forum Math.* **36**, No 1, 1–4, (2024).
- [50] Weber, M., A uniform semi-local limit theorem along sets of multiples for sums of i.i.d. square integrable random variables, *Funct. Approx. Comment. Math.* **70** (2), 177–199 (2024).
- [51] Weber, M., A strengthened asymptotic uniform distribution, *Probab. Math. Stat.* **43**, No. 2, 263–285 (2023).
- [52] Weber, M., An arithmetical approach to the convergence problem of series of dilated functions and its connection with the Riemann Zeta function, *J. Number Th.*, **162** (2016), 137–179.
- [53] Weber, M., Critical probabilistic characteristics of the Cramér model for primes and arithmetical properties, *Indian J. of Pure and Applied Math.*, (2024). <https://doi.org/10.1.1007/s12188-024-00278-0>.
- [54] Weber, M., Some examples of application of the metric entropy method, *Acta Math. Hungar.* **105** (2004), 39–83

- [55] Weber, M., A sharp correlation inequality with an application to almost sure local limit theorem, *Prob. and Math. Stat.* **31** (2011), Fasc. 1, 79–98.
- [56] Weber M., Some theorems related to almost sure convergence of orthogonal series, *Indag. Math.* (N.S.) **11** (2000).
- [57] Weber M., *Dynamical Systems and Processes*, European Mathematical Society Publishing House, IRMA Lectures in Mathematics and Theoretical Physics **14**, (2009), xiii+761p.
- [58] Weber M., *Entropie métrique et convergence presque partout*, Travaux en Cours **58**, (1998), Hermann, Paris.

IRMA, UMR 7501, UNIVERSITÉ LOUIS-PASTEUR ET C.N.R.S., 7 RUE RENÉ DESCARTES, 67084 STRASBOURG CEDEX, FRANCE. E-MAIL: [michel.weber@math.unistra.fr](mailto:michel.weber@math.unistra.fr)