

CHROMATIC SYMMETRIC FUNCTIONS OF CONJOINED GRAPHS

ETHAN Y.J. QI, DAVION Q.B. TANG, AND DAVID G.L. WANG*

ABSTRACT. We introduce path-conjoined graphs defined for two rooted graphs by joining their roots with a path, and investigate the chromatic symmetric functions of its two generalizations: spider-conjoined graphs and chain-conjoined graphs. By using the composition method developed by Zhou and the third author recently, we obtain neat positive e_I -expansions for the chromatic symmetric functions of clique-path-cycle graphs, path-clique-path graphs, and clique-clique-path graphs. We pose the e -positivity conjecture for hat-chains.

1. INTRODUCTION

The chromatic symmetric function of a graph was introduced by Stanley [22] as a generalization of Birkhoff's chromatic polynomial. It attracted a considerable number of studies from various branches of mathematics, including combinatorics, representation and knot theory, and from mathematical physics, see Chmutov, Kazarian, and Lando [4], Kato [13], Loeb [16], Noble and Welsh [18].

One of the most popular topics in this field is [Stanley and Stembridge](#)'s conjecture [24]. In virtue of [Guay-Paquet](#)'s reduction [10], it is equivalent to the e -positivity of unit interval graphs. More precisely, the chromatic symmetric function of any unit interval graph has only nonnegative e -coefficients. The conjecture charmed representation experts because of a close relation between chromatic quasisymmetric functions that were introduced by Shareshian and Wachs [21] and the representations of symmetric groups on cohomology of regular semisimple Hessenberg varieties, see Brosnan and Chow [3]. In October 2024, Hikita [12] resolved [Stanley and Stembridge](#)'s conjecture.

A natural generalization of the conjecture is to characterize all e -positive graphs, as mentioned by Stanley [22, Section 5]. Though seemingly unachievable is it, more and more families of graphs have been confirmed to be e -positive. The most noted e -positive graphs are complete graphs, paths, cycles, and graphs whose independent numbers are 2. A bit less well-known e -positive graphs are lollipops, tadpoles and generalized bulls, which can be derived from a direct application of [Gebhard and Sagan](#)'s (e) -positivity strengthening, see [9, Theorem 7.6], and also Dahlberg and van Willigenburg [6], Li, Li, Wang, and Yang [14]. Here strengthening means that the (e) -positivity is stronger than the e -positivity. More e -positive graphs can be found from Banaian, Celano, Chang-Lee, Colmenarejo, Goff, Kimble, Kimpel, Lentfer, Liang, and Sundaram [2], Cho and Huh [5], Foley, Hoàng, and Merkel [8], Hamel, Hoàng, and Tuero [11], Li and Yang [15], Tsujie [31], Wang [32], Wang and Wang [33].

Along the way of confirming the e -positivity of graphs arose various approaches. Notable routes include (i) [Stanley](#)'s generating function and recurrence relation method, (ii) the strategy of expressing a chromatic symmetric function as a sum of e -positive symmetric functions, often as a sum of known e -positive chromatic symmetric functions; a typical example is the proof for the 3-spiders of the form $S(n+1, n, 1)$, see Dahlberg, Foley, and van Willigenburg [7, Page 2681], (iii) [Gebhard and Sagan](#)'s (e) -positivity strengthening that takes advantage of the algebra NCSym of symmetric functions in

2020 *Mathematics Subject Classification.* 05E05, 05A15.

Key words and phrases. chromatic symmetric functions, the composition method, conjoined graphs, e -positivity, hat-chains.

*Wang is the corresponding author, and is supported by the NSFC (Grant No. 12171034).

noncommutative variables, (iv) understanding the e -coefficients of a chromatic symmetric function combinatorially and constructing sign-reversing involutions, see Tom [28] for instance, (v) the composition method which is inspired from Shareshian and Wachs [21]'s formula for the chromatic symmetric function of paths, initiated implicitly by Thibon and Wang [26] in proving the Schur positivity of the 3-spiders of the forms $S(a, 2, 1)$ and $S(a, 4, 1)$, and formulated by Wang and Zhou [34].

In this paper, we continue explore the power of the composition method. A key idea of the composition method is to extend the defining domain of the elementary symmetric functions e_λ from partitions λ to compositions I . An e_I -expansion of a symmetric function f is an expression

$$(1.1) \quad f = \sum_{I \vdash n} c_I e_I$$

such that

$$(1.2) \quad f = \sum_{\lambda \vdash n} \left(\sum_{\rho(I) = \lambda} c_I \right) e_\lambda,$$

where $\rho(I)$ is the partition obtained from the composition I by rearranging its parts. We call Eq. (1.1) a *positive e_I -expansion* if $c_I \geq 0$ for all compositions I of n . In order to show that f is e -positive, it suffices to find a positive e_I -expansion of f . Every e -positive symmetric function has definitely a positive e_I -expansion, while the non-uniqueness of e_I -expansions suggests a flexibility in invoking combinatorial ideas to expose or construct a positive e_I -expansion.

One family of the foregoing e -positive graphs that are unit interval graphs is *K -chains*, see Section 2. Its e -positivity was due to Gebhard and Sagan [9], and rediscovered by McDonough, Pylyavskyy, and Wang [17], Tom [27]. We would like to seek for a neat positive e_I -expansion for K -chains. Here the word neat means that essentially less compositions appear in the expansion, or aesthetically less technical conditions are required for the compositions that appear. This has been done for two particular kinds of K -chains: barbells and generalized bulls, see [34]. In this paper, we continue this course and light on another two K -chains: path-clique-paths (PKPs) and clique-clique-paths (KKPs), see Theorems 5.3 and 5.4 respectively. In fact, all generalized bulls are PKPs. In the same fashion, we find a positive e_I -expansions for clique-path-cycles (KPCs), see Theorem 3.3. This is the first positive e_I -expansion of KPC graphs to the best of our knowledge. It is transparent that the composition method is distinct from the involution one, and our positive e_I -expansions for PKPs and KKP are different from Tom's, see Theorem 2.9. We should mention that the e -positivity of all these graphs are derivable by using Gebhard and Sagan's (e)-positivity routine.

Ad hoc algebraic combinatorial techniques are called for in coping with PKPs, KKP and KPCs, nevertheless, these graphs have a common structure: a path part as a bridge that links some *node graphs* such like cliques or cycles. We are therefore led to the graph operation of *path-conjoining*, see Section 3. We then generalize it to spider- and chain-conjoining operations, and express the chromatic symmetric function of these amalgamated graphs in terms of those of the node graphs.

This paper is organized as follows. Section 2 consists of preliminary results that we will need in the sequel. In Section 3, we study the chromatic symmetric function of path-conjoined graphs, and specify our results to those with a clique node and those with a cycle node. Section 4 is devoted to spider-conjoined graphs. In particular, we investigate some pineapple graphs which appeared in Topcu, Sorgun, and Haemers [30], see Corollary 4.7. In Section 5, we consider 2-chain-conjoined graphs with the middle node being a clique or a cycle, see Theorems 5.1 and 5.5. We end this paper with the e -positivity conjecture for hat-chain graphs.

2. PRELIMINARIES

This section contains basic knowledge on chromatic symmetric functions. We adopt notion and notations of Stanley [23]. Let n be a positive integer. A *composition* of n is a sequence of positive integers with sum n , commonly denoted $I = i_1 \cdots i_l \models n$, with *size* $|I| = n$, *length* $\ell(I) = l$, and *parts* i_1, \dots, i_l . A *prefix* of I is a composition of the form $i_1 \cdots i_k$, where $0 \leq k \leq \ell(I)$. We write $I = v^s$ if all parts have the same value v . For notational convenience, we denote the k th last part i_{l+1-k} by i_{-k} , and denote the composition obtained by removing the k th part by $I \setminus i_k$, i.e.,

$$I \setminus i_k = i_1 \cdots i_{k-1} i_{k+1} \cdots i_{-1}.$$

When a capital letter like I and J stands for a composition, we use its small letter counterpart with integer subscripts to denote the parts. A *partition* of n is a multiset of positive integers λ_i with sum n , denoted $\lambda = \lambda_1 \lambda_2 \cdots \models n$, where $\lambda_1 \geq \lambda_2 \geq \cdots \geq 1$.

A *symmetric function* of homogeneous degree n over the field \mathbb{Q} of rational numbers is a formal power series

$$f(x_1, x_2, \dots) = \sum_{\lambda = \lambda_1 \lambda_2 \cdots \models n} c_\lambda \cdot x_1^{\lambda_1} x_2^{\lambda_2} \cdots,$$

where $c_\lambda \in \mathbb{Q}$, such that $f(x_1, x_2, \dots) = f(x_{\pi(1)}, x_{\pi(2)}, \dots)$ for any permutation π . One basis of the vector space of homogeneous symmetric functions of degree n over \mathbb{Q} is the set of elementary symmetric functions $e_\lambda = e_{\lambda_1} e_{\lambda_2} \cdots$, where

$$e_k = \sum_{1 \leq i_1 < \cdots < i_k} x_{i_1} \cdots x_{i_k}.$$

A symmetric function f is said to be *e-positive* if every e_λ -coefficient is nonnegative.

For any composition I , there is a unique partition $\rho(I)$ that consists of the parts of I . As a consequence, one may extend the defining domain of elementary symmetric functions from partitions to compositions as $e_I = e_{\rho(I)}$. An *e_I-expansion* of a symmetric function f is an expression Eq. (1.1) that satisfies Eq. (1.2). It is said to be *positive* if $c_I \geq 0$ for all compositions $I \models n$. In order to show that f is *e*-positive, it suffices to find a positive *e_I*-expansion of f .

Stanley [22] introduced the chromatic symmetric function for a simple graph G as

$$X_G = \sum_{\kappa} \prod_{v \in V(G)} x_{\kappa(v)},$$

where κ runs over proper colorings of G . It is a homogeneous symmetric function of degree $|G|$, the order of G . For instance, the chromatic symmetric function of the complete graph K_n is $X_{K_n} = n! e_n$.

Orellana and Scott [19, Theorem 3.1, Corollaries 3.2 and 3.3] established the *triple-deletion* properties for chromatic symmetric functions as follows.

Proposition 2.1 (Orellana and Scott, the triple-deletion properties). *Let G be a graph with a stable set T of order 3. Denote the edges linking the vertices in T by e_1, e_2 and e_3 . For any set $S \subseteq \{1, 2, 3\}$, denote by G_S the graph with vertex set $V(G)$ and edge set $E(G) \cup \{e_j : j \in S\}$. Then*

$$(2.1) \quad X_{G_{12}} = X_{G_1} + X_{G_{23}} - X_{G_3}, \quad \text{and}$$

$$(2.2) \quad X_{G_{123}} = X_{G_{13}} + X_{G_{23}} - X_{G_3}.$$

Using Proposition 2.1, Aliniaeifard, Wang, and van Willigenburg [1, Proposition 6] showed that certain sequence of chromatic symmetric functions forms an arithmetic progression. A *clique* of a graph G is a subset of vertices of G such that every two distinct vertices in the clique are adjacent. The *closed neighborhood* of a vertex v of G is the union of the set of vertices that are adjacent to v and v itself.

Proposition 2.2 (Aliniaeifard et al., the arithmetic progression property). *Suppose that G contains a clique on the vertices v_1, \dots, v_k , which have the same closed neighborhoods. Let w be a vertex of G that is not adjacent to any vertex v_j . Then the sequence X_{G_0}, \dots, X_{G_k} forms an arithmetic progression, where $G_0 = G$ and G_j for $1 \leq j \leq k$ is obtained from G_{j-1} by adding the edge wv_j .*

In applying Proposition 2.2, we define a *tribe-vertex pair* of a graph G to be a pair (K, x) in which

- (1) K is a clique of G ,
- (2) x is a vertex that may or may not be in G , and
- (3) all vertices in K have the same closed neighborhoods in the graph $G - x$.

Corollary 2.3. *Let (K, x) be a tribe-vertex pair of a graph G . Let G_i be the graph obtained from G by connecting exactly i vertices from K to x . Then*

$$(i - j)X_{G_k} + (j - k)X_{G_i} + (k - i)X_{G_j} = 0, \quad \text{for any } 0 \leq i \leq j \leq k \leq |K|.$$

Proof. By Proposition 2.2, the chromatic symmetric functions X_{G_j} form an arithmetic progression. In other words, if we let $m = |K|$, then for any $0 \leq i \leq m$,

$$(2.3) \quad X_{G_i} = \frac{m-i}{m}X_{G_0} + \frac{i}{m}X_{G_m}.$$

Let $F = (i - j)X_{G_k} + (j - k)X_{G_i} + (k - i)X_{G_j}$. Substituting Eq. (2.3) and its analogues obtained by replacing i with j and k respectively into F , we can express F as a linear combination of X_{G_0} and X_{G_m} . In that expression, the coefficient of X_{G_0} in F is

$$(i - j)\frac{m - k}{m} + (j - k)\frac{m - i}{m} + (k - i)\frac{m - j}{m} = 0,$$

and the coefficient of X_{G_m} in F is

$$(i - j)\frac{k}{m} + (j - k)\frac{i}{m} + (k - i)\frac{j}{m} = 0.$$

This proves $F = 0$ as desired. \square

Here is an analogue of Corollary 2.3.

Corollary 2.4. *Let (K, x) be a tribe-vertex pair of a graph H . Suppose that x connects to all vertices of K . Let H_i be the graph obtained from H by removing i edges from K to x . Then*

$$(i - j)X_{H_k} + (j - k)X_{H_i} + (k - i)X_{H_j} = 0, \quad \text{for any } 0 \leq i \leq j \leq k \leq |K|.$$

Proof. Suppose that $|K| = m$. In Corollary 2.3, if we take $G = H$, then H_i becomes G_{m-i} , and

$$((m - k) - (m - j))X_{G_{m-i}} + ((m - j) - (m - i))X_{G_{m-k}} + ((m - i) - (m - k))X_{G_{m-j}} = 0.$$


Simplifying it yields the desired identity. \square

Shareshian and Wachs [21, Table 1] discovered a captivating formula using Stanley's generating function for Smirnov words, see also Shareshian and Wachs [20, Theorem 7.2].

Proposition 2.5 (Shareshian and Wachs). $X_{P_n} = \sum_{I \models n} w_I e_I$ for any $n \geq 1$, where

$$(2.4) \quad w_I = i_1(i_2 - 1)(i_3 - 1) \cdots (i_{-1} - 1).$$

The *lollipop* K_m^l (resp., *tadpole* C_m^l) is the graph obtained by identifying a vertex of the complete graph K_m (resp., the cycle C_m) and an end of the path P_{l+1} , see the leftmost (resp., middle) illustration in Fig. 1. Tom [27, Thm. 2] computed the chromatic symmetric function of melting lollipops. We restate his result for lollipops as a specialization.

FIGURE 1. The lollipop K_m^l , the tadpole C_m^l , and the 3-spider $S(abc)$.

Proposition 2.6 (Lollipops, [Tom](#)). *For any $n \geq a \geq 1$,*

$$X_{K_a^{n-a}} = (a-1)! \sum_{I \models n, i_{-1} \geq a} w_I e_I.$$

Wang and Zhou [\[34\]](#) provided a formula for tadpoles. For any number $a \leq n$, define

$$(2.5) \quad \sigma_I(a) = \min\{i_1 + \dots + i_k : 0 \leq k \leq \ell(I), i_1 + \dots + i_k \geq a\}.$$

The a -surplus of I is the number

$$(2.6) \quad \Theta_I(a) = \sigma_I(a) - a.$$

Proposition 2.7 (Tadpoles, [Wang and Zhou](#)). *For any $0 \leq l \leq n-2$,*

$$X_{C_{n-l}^l} = \sum_{I \models n} \Theta_I(l+1) w_I e_I.$$

For any composition $I \models n-1$, the *spider* $S(I)$ is the tree of order n obtained by identifying an end of the paths $P_{i_1+1}, P_{i_2+1}, \dots$. The identified vertex is said to be the *center* of $S(I)$. We say that $S(I)$ is an $\ell(I)$ -spider, see the rightmost illustration in Fig. 1 for the 3-spider $S(abc)$. Zheng [\[35, Lemma 4.4\]](#) expressed the chromatic symmetric function of 3-spiders in terms of chromatic symmetric functions of paths.

Proposition 2.8 (3-Spiders, [Zheng](#)). *For any composition $abc \models n-1$,*

$$X_{S(abc)} = \sum_{i=0}^c X_{P_i} X_{P_{n-i}} - \sum_{i=b+1}^{b+c} X_{P_i} X_{P_{n-i}}.$$

Let K_{i_1}, \dots, K_{i_l} be completed graphs, where $i_k \geq 2$ for all k . Suppose that u_k and v_k are distinct vertices of K_{i_k} . The K -chain $K_{i_1} + \dots + K_{i_l}$ is the graph obtained by identifying v_k and u_{k+1} for all $1 \leq k \leq l-1$. Tom [\[27, Thm. 3\]](#) discovered an elegant positive e_I -expansion for K -chains. A *weak composition* of n is a sequence of nonnegative integers whose sum is n .

Theorem 2.9 (K -chains, [Tom](#)). *Let $\gamma = \gamma_1 \dots \gamma_l$ be a composition with each part at least 2. If K_γ is the K -chain with maximal clique sizes γ_i , then*

$$X_{K_\gamma} = (\gamma_1 - 2)! \dots (\gamma_{l-1} - 2)! (\gamma_l - 1)! \sum_{\alpha \in A_\gamma} \left(\alpha_1 \prod_{i=2}^l |\alpha_i - \gamma_{i-1} + 1| \right) e_\alpha,$$

where $l = \ell(\gamma)$ and A_γ is the set of weak compositions $\alpha = \alpha_1 \dots \alpha_l$ of $|\gamma| - l + 1$ such that for each $2 \leq i \leq l$, either

- $\alpha_i < \gamma_{i-1}$ and $\alpha_i + \dots + \alpha_l < \gamma_i + \dots + \gamma_l - (l-i)$, or
- $\alpha_i \geq \gamma_{i-1}$ and $\alpha_i + \dots + \alpha_l \geq \gamma_i + \dots + \gamma_l - (l-i)$.

3. PATH-CONJOINED GRAPHS

A *rooted graph* (G, u) is a pair consisting of a graph G and a vertex u of G . We call u the *root* of G . For any rooted graphs (G, u) and (H, v) , we define the *path-conjoined graph* $P^k(G, H)$ to be the graph obtained by adding a path of length k which links u and v . It has order $|G| + |H| + k - 1$, see Fig. 2. We call G and H the *node graphs* of $P^k(G, H)$.




FIGURE 2. The path-conjoined graph $P^k(G, H)$.

A graph is *vertex-transitive* if for any vertices x and y , there exists an automorphism f such that $f(x) = y$. We say that G is a rooted graph without mentioning its root, whenever G is vertex-transitive or the root is clear from context. For instance, path-conjoined graphs $P^k(K_g, K_h)$ are barbells, see [34, Theorem 3.7]. When one of the nodes, say, H , is a singleton K_1 , we write

$$G^k = P^k(G, K_1),$$

and call it a *tailed graph*. Then lollipops are tailed cliques, and tadpoles are tailed cycles.

The path-conjoining operation by a path of length k is equivalent to joining a graph with a new edge P_2 for k times, as considered by Gebhard and Sagan [9] and Tom [27]. As will be seen, one benefits from the viewpoint of recognizing a path as a whole; in this way the number of compositions that appear in our formula decreases, in comparison with that in Theorem 2.9.

The path-conjoined graphs $P^k(G, H)$ with one node, say, G , being a clique or a cycle is of particular interests. We express the chromatic symmetric function of such a graph in terms of those of tailed graphs with node H , see Propositions 3.1 and 3.2.

Proposition 3.1 (KPGs). *Let H be a rooted graph. Then for any $k \geq 0$ and $g \geq 1$,*

$$X_{P^k(K_g, H)} = (g-1)! \sum_{l=0}^{g-1} (1-l)e_l X_{H^{k+g-1-l}}.$$

Proof. When $g = 1$, the desired formula holds by the definition $G^k = P^k(G, K_1)$ for tailed graphs. Let $g \geq 2$. In Corollary 2.4, if we take $K = K_g$, and take x to be the root of K_g , then $G_0 = G$, $G_{g-2} = P^{k+1}(K_{g-1}, H)$, and G_{g-1} is the disjoint union of K_{g-1} and H^k . It follows that

$$X_{P^k(K_g, H)} = (g-1)X_{P^{k+1}(K_{g-1}, H)} - (g-2)X_{K_{g-1}}X_{H^k} \quad \text{for } g \geq 2,$$

in which $X_{K_{g-1}} = (g-1)!e_{g-1}$. The desired formula follows by using this relation iteratively. \square

Proposition 3.2 (CPGs). *Let H be a rooted graph. Then for any $k \geq 0$ and $g \geq 2$,*

$$X_{P^k(C_g, H)} = (g-1)X_{H^{k+g-1}} - \sum_{l=1}^{g-2} X_{C_{g-l}}X_{H^{k+l-1}}.$$

Proof. The desired formula holds for $g = 2$ since H^{k+1} is the underlying graph of $P^k(C_2, H)$. For $g \geq 3$, let u be the root of C_g , with neighbors x and y on C_g . Applying Proposition 2.1 to the triangle uxy , we obtain

$$X_{P^k(C_g, H)} = X_{P^{k+1}(C_{g-1}, H)} + X_{H^{k+g-1}} - X_{C_{g-1}}X_{H^k} \quad \text{for } g \geq 3.$$

We can then infer the desired formula by using this relation iteratively for g down to 3. \square

The particular case for when $H = K_1$ appeared in Li et al. [14, Formula (3.11)].

A *clique-path-cycle* graph is a graph of the form $P^b(K_a, C_c)$. We call them KPC graphs for short. Now we show their e -positivity by using Proposition 3.1 and the composition method.

Theorem 3.3 (KPCs). *For any $a \geq 1$, $b \geq 0$ and $c \geq 2$, the chromatic symmetric function of the KPC graph $P^b(K_a, C_c)$ has the positive e_I -expansion*

$$X_{P^b(K_a, C_c)} = (a-1)! \sum_{K \models n} c_K w_K e_K,$$

where $n = a + b + c - 1$, w_K is defined by Def. (2.4), and

$$c_K = \begin{cases} 0, & \text{if } \ell(K) \geq 2 \text{ and } k_2 < a, \\ k_2 - a - b + \frac{k_2 - k_1}{k_2 - 1}, & \text{if } k_1 \leq a - 1 \text{ and } k_2 \geq a + b, \\ \Theta_K(a+b), & \text{otherwise.} \end{cases}$$

Here the function $\Theta_K(\cdot)$ is defined by Def. (2.6). As a consequence, all KPC graphs are e -positive.

Proof. Let $G = P^b(K_a, C_c)$. Since $c \geq 2$, by using Propositions 2.7 and 3.1, we can deduce that

$$\frac{X_G}{(a-1)!} = \sum_{l=0}^{a-1} (1-l)e_l X_{C_c^{a+b-1-l}} = \sum_{K \models n} \Theta_K(a+b) w_K e_K - \sum_{K \models n, \ell(K) \geq 2, k_2 \leq a-1} q_K w_K e_K,$$

where $q_K = \Theta_{K \setminus k_2}(a+b-k_2)$. Now we deal with the coefficient in the last sum.

- (1) If $k_1 < a + b - k_2$, then $q_K = \Theta_K(a+b)$.
- (2) If $a + b - k_2 \leq k_1 < a + b$, then $q_K = k_1 + k_2 - (a+b) = \Theta_K(a+b)$.
- (3) If $k_1 \geq a + b$, then $q_K = k_1 + k_2 - (a+b) = \Theta_K(a+b) + k_2$.

Therefore, the last sum can be rewritten as

$$\sum_{\substack{K \models n, \ell(K) \geq 2 \\ k_2 \leq a-1}} q_K w_K e_K = \sum_{\substack{K \models n, \ell(K) \geq 2 \\ k_2 \leq a-1}} \Theta_K(a+b) w_K e_K + \sum_{\substack{K \models n, \ell(K) \geq 2 \\ k_1 \geq a+b, k_2 \leq a-1}} k_2 w_K e_K.$$

It follows that

$$(3.1) \quad \frac{X_G}{(a-1)!} = (c-1)ne_n + \sum_{K \in \mathcal{A}_n} \Theta_K(a+b) w_K e_K - \sum_{K \in \mathcal{B}_n} k_2 w_K e_K,$$

where

$$\begin{aligned} \mathcal{A}_n &= \{K \models n: \ell(K) \geq 2, k_2 \geq a\}, \quad \text{and} \\ \mathcal{B}_n &= \{K \models n: \ell(K) \geq 2, k_1 \geq a+b, 2 \leq k_2 \leq a-1\}. \end{aligned}$$

For any $K \in \mathcal{B}_n$, let $f(K)$ be the composition obtained by exchanging the first two parts of K . Then f is a bijection between \mathcal{B}_n and the set

$$f(\mathcal{B}_n) = \{K \models n: \ell(K) \geq 2, 2 \leq k_1 \leq a-1, k_2 \geq a+b\}.$$

Then we can rewrite the negative sum in Eq. (3.1) as

$$\sum_{K \in \mathcal{B}_n} k_2 w_K e_K = \sum_{K \in f(\mathcal{B}_n)} \frac{k_1 - 1}{k_2 - 1} \cdot k_2 w_K e_K.$$

It is clear that $f(\mathcal{B}_n) \subseteq \mathcal{A}_n$. For $K \in f(\mathcal{B}_n)$, we have

$$\Theta_K(a+b) - \frac{k_2(k_1-1)}{k_2-1} = (k_2 - a - b) + \frac{k_2 - k_1}{k_2 - 1} > 0.$$

It follows that

$$\frac{X_G}{(a-1)!} = (c-1)ne_n + \sum_{K \in \mathcal{A}_n \setminus f(\mathcal{B}_n)} \Theta_K(a+b)w_K e_K + \sum_{K \in f(\mathcal{B}_n)} \left(k_2 - a - b + \frac{k_2 - k_1}{k_2 - 1} \right) w_K e_K,$$

which can be recast as the desired formula. \square

Theorem 3.3 reduces to Proposition 2.6 for $c = 2$, and to Proposition 2.7 for both $a = 1$ and $a = 2$. For example, taking $(a, k, b) = (4, 2, 4)$, we can calculate

$$\begin{aligned} X_{P^2(K_4, C_4)} = & 18e_{1422} + 162e_{144} + 30e_{162} + 126e_{18} + 48e_{252} + 132e_{27} + 54e_{342} + 54e_{36} \\ & + 288e_{45} + 270e_{54} + 162e_9. \end{aligned}$$

A *kayak paddle* is a path-conjoined graph with cycle nodes. [Aliniaeifard et al.](#) extended the appendability technique of (e) -positivity and proved the e -positivity of some chain-like graphs including kayak paddles, see [1, Proposition 34].

Proposition 3.4 ([Aliniaeifard et al.](#)). *All kayak paddles are e -positive.*

Tang and Wang [25] discovered a positive e_I -expansion for the chromatic symmetric function of kayak paddles.

4. SPIDER-CONJOINED GRAPHS

Let $\tau = \tau_1 \cdots \tau_l$ be a weak composition. Let $\{s_1, \dots, s_l\}$ be the set of leaves of the spider $S(\tau)$ with center c , in which the leg containing s_i has length τ_i . For any rooted graphs $(G_1, u_1), \dots, (G_l, u_l)$, we define the l -spider-conjoined graph $G = S^\tau(G_1, \dots, G_l)$ to be the graph obtained by identifying u_i with s_i for all i . It has order

$$|G| = |G_1| + \cdots + |G_l| + |\tau| - l + 1.$$

In particular, 1-spider-conjoined graphs are tailed graphs, and 2-spider-conjoined graphs are path-conjoined graphs. We say that G has *center* c and *nodes* G_i , and omit the root of G_i whenever G_i is vertex-transitive or the root is clear from context.

The chromatic symmetric function of l -spider-conjoined graphs for $l \geq 4$ can be expressed in terms of those for 3-, 2-, and 1-spider-conjoined graphs, when τ has at least two positive parts.

Proposition 4.1. *Let $\tau = \tau_1 \cdots \tau_l$ be a weak composition with $l \geq 3$ and $\tau_1, \tau_2 \geq 1$. Then for any l -spider-conjoined graph $G = S^\tau(G_1, \dots, G_l)$,*

$$X_G = X_{S^{(\tau_1-1)\tau_21}}(G_1, G_2, H) + X_{G_1^{\tau_1-1}} X_{S^{\tau_2 \cdots \tau_l}(G_2, \dots, G_l)} - X_{P^{\tau_1+\tau_2-1}(G_1, G_2)} X_H,$$

where $H = S^{\tau_3 \cdots \tau_l}(G_3, \dots, G_l)$ is considered to be a rooted graph whose root is the center of G .

Proof. Applying Eq. (2.1) to the triangle formed by the center of G and its neighbors on the G_1 - and G_2 -legs, we obtain the desired formula. \square

With Proposition 4.1 in hand, one may concentrate on 3-spider-conjoined graphs.

Proposition 4.2. *Let G, H and J be rooted graphs. Then for any $g, j \geq 0$ and $h \geq 1$,*

$$X_{S^{ghj}(G, H, J)} = X_{S^{(g+j)h0}(G, H, J)} + \sum_{i=1}^j (X_{P^{g+h+i-1}(G, H)} X_{J^{j-i}} - X_{G^{g+i-1}} X_{P^{h+j-i}(H, J)}).$$

Proof. It is trivial for $j = 0$. For $j \geq 1$, taking $(G_1, G_2, G_3) = (J, H, G)$ in Proposition 4.1 yields

$$X_{S^{ghj}(G,H,J)} = X_{S^{(g+1)h(j-1)}(G,H,J)} + X_{P^{g+h}(G,H)}X_{J^{j-1}} - X_{G^g}X_{P^{h+j-1}(H,J)}.$$

Using it iteratively for j down to 1, we can derive the desired formula. \square

For any $1 \leq k \leq l$, we denote

$$S^\tau(G_1, \dots, G_k, K_1, K_1, \dots, K_l) = S^{\tau_1 \dots \tau_k}_{\tau_{k+1} \dots \tau_l}(G_1, \dots, G_k)$$

for compactness. See Fig. 3 for the graph $S_j^{gh}(G, H)$.




FIGURE 3. The spider-conjoined graph $S_j^{gh}(G, H)$.

For when one or two of the nodes of a 3-spider-conjoined graph are the singletons K_1 , we write the specifications out. It will be used in the proofs of Theorems 4.5 and 5.5.

Corollary 4.3. *Let G and H be rooted graphs. Then for any $g, j \geq 0$ and $h \geq 1$,*

$$\begin{aligned} X_{S_j^{gh}(G,H)} &= \sum_{i=0}^j X_{P_i} X_{P^{g+h+j-i}(G,H)} - \sum_{i=1}^j X_{G^{g+i-1}} X_{H^{h+j-i}}, \quad \text{and} \\ X_{S_{hj}^g(G)} &= \sum_{i=0}^j X_{P_i} X_{G^{g+j-i+h}} - \sum_{i=1}^j X_{P_{i+h}} X_{G^{g+j-i}}. \end{aligned}$$

Proof. They follow by taking $J = K_1$ and taking $J = H = K_1$ in Proposition 4.2, respectively. \square

The second formula in Corollary 4.3 for $G = K_1$ reduces to Proposition 2.8.

In the remaining of Section 4, we consider 3-spider-conjoined graphs with a clique node and those with a cycle node; see Theorems 4.5 and 4.8, respectively. The following Lemma 4.4 will be used in proving Theorems 4.5 and 5.2.

Lemma 4.4. *For $a \geq 0$,*

$$\sum_{l=0}^a (1-l)e_l X_{P_{n-l}} = \begin{cases} X_{K_{a+1}^{n-1-a}}/a!, & \text{if } a \leq n-1; \\ e_n, & \text{if } a = n. \end{cases}$$

Proof. Let $0 \leq a \leq n-1$. By Propositions 2.5 and 2.6, we can deduce that

$$\sum_{l=2}^a (l-1)e_l X_{P_{n-l}} = \sum_{K \models n, k_{-1} \leq a} w_K e_K = X_{P_n} - \sum_{K \models n, k_{-1} \geq a+1} w_K e_K = X_{P_n} - X_{K_{a+1}^{n-a-1}}/a!.$$

Rearranging the terms yields the desired formula for $a < n$. Now, taking $a = n-1$, we obtain

$$\sum_{l=0}^{n-1} (1-l)e_l X_{P_{n-l}} = \frac{X_{K_n}}{(n-1)!} = n e_n.$$

Adding both sides by $(1-n)e_n$ yields the desired formula for $a = n$. \square

Here is the reduction for the chromatic symmetric function of 3-spider-conjoined graphs $S^{ghk}(G, H, K)$ with a clique node $K = K_m$.

Theorem 4.5. *Let G and H be rooted graphs. For any $m, h \geq 1$ and $g, k \geq 0$,*

$$\begin{aligned} \frac{X_{S^{ghk}(G, H, K_m)}}{(m-1)!} &= \sum_{z=0}^{m-1} e_{m-1-z} X_{P^{g+h+k+z}(G, H)} - \sum_{\substack{a+b+c=k+m-2 \\ a, b, c \geq 0, a \leq m-1}} (1-a)e_a X_{G^{b+g}} X_{H^{c+h}} \\ &\quad + \frac{1}{(m-1)!} \sum_{z=0}^{k-1} X_{K_m^z} X_{P^{g+h+k-z-1}(G, H)}. \end{aligned}$$

Proof. Applying Proposition 3.1 to the spider $S_0^{gh}(G, H)$ and the clique K_m , we obtain

$$X_{S^{ghk}(G, H, K_m)} = (m-1)! \sum_{l=0}^{m-1} (1-l)e_l X_{S_{k+m-1-l}^{gh}(G, H)}.$$

We write $q = k + m - 1$ and $P^l(G, H) = P^l$ for short. Using Corollary 4.3, we can deduce that

$$(4.1) \quad \frac{X_{S^{ghk}(G, H, K_m)}}{(m-1)!} = \sum_{l=0}^{m-1} (1-l)e_l \left(\sum_{z=0}^{q-l} X_{P_{q-l-z}} X_{P^{g+h+z}} - \sum_{z=0}^{q-l-1} X_{G^{g+q-l-z-1}} X_{H^{h+z}} \right).$$

Let A and B be the first and second double sum on the right side of Eq. (4.1), respectively. By interchanging the order of summation in A , we obtain

$$\begin{aligned} A &= \sum_{l=0}^{m-1} (1-l)e_l \sum_{z=0}^{q-l} X_{P_{q-l-z}} X_{P^{g+h+z}} = \sum_{z=0}^q X_{P^{g+h+z}} \sum_{l=0}^{\min(m-1, q-z)} (1-l)e_l X_{P_{q-l-z}} \\ &= \sum_{z=0}^q X_{P^{g+h+q-z}} \sum_{l=0}^{\min(m-1, z)} (1-l)e_l X_{P_{z-l}} \\ &= \sum_{z=0}^{m-1} X_{P^{g+h+q-z}} \sum_{l=0}^z (1-l)e_l X_{P_{z-l}} + \sum_{z=m}^{k+m-1} X_{P^{g+h+q-z}} \sum_{l=0}^{m-1} (1-l)e_l X_{P_{z-l}}. \end{aligned}$$

Using Lemma 4.4 and Proposition 2.6, we can further transform it as

$$\begin{aligned} A &= \sum_{z=0}^{m-1} e_z X_{P^{g+h+q-z}} + \sum_{z=m}^{k+m-1} X_{P^{g+h+q-z}} \frac{X_{K_m^{z-m}}}{(m-1)!} \\ &= \sum_{z=0}^{m-1} e_z X_{P^{g+h+q-z}} + \frac{1}{(m-1)!} \sum_{z=0}^{k-1} X_{P^{g+h+q-z-m}} X_{K_m^z}. \end{aligned}$$

On the other hand, the double sum B can be recast as the compact form

$$B = \sum_{l=0}^{m-1} (1-l)e_l \sum_{z=0}^{q-l-1} X_{G^{g+q-l-z-1}} X_{H^{h+z}} = \sum_{\substack{a+b+c=k+m-2 \\ a, b, c \geq 0, a \leq m-1}} (1-a)e_a X_{G^{b+g}} X_{H^{c+h}}.$$

The sum $A + B$ is the desired formula. \square

When $H = K_m$, one may further simplify the formula in Theorem 4.5.

Corollary 4.6. *Let G be a rooted graph. For any $k, g \geq 0$ and $m, h \geq 1$,*

$$X_{S_h^{gk}(G, K_m)} = (m-1)! \left(\sum_{z=0}^{m-1} e_z X_{G^{g+h+k+m-z-1}} - \sum_{z=1}^{m-1} \frac{X_{K_z^h}}{(z-1)!} X_{G^{g+k+m-z-1}} \right)$$

$$+ \sum_{z=0}^{k-1} \left(X_{K_m^z} X_{G^{g+h+k-z-1}} - X_{K_m^{h+z}} X_{G^{g+k-z-1}} \right).$$

Proof. In Theorem 4.5, taking $H = K_1$, we obtain

$$\begin{aligned} \frac{X_{S_h^{gk}(G, K_m)}}{(m-1)!} &= \sum_{z=0}^{m-1} e_z X_{G^{g+h+k+m-1-z}} - \sum_{\substack{a+b+c=k+m-2 \\ a, b, c \geq 0, a \leq m-1}} (1-a)e_a X_{G^{b+g}} X_{P_{c+h+1}} \\ &\quad + \frac{1}{(m-1)!} \sum_{z=0}^{k-1} X_{K_m^z} X_{G^{g+h+k-z-1}}. \end{aligned}$$

Let A be the second sum, namely, the one with parameters a, b , and c . Then

$$\begin{aligned} A &= \sum_{i=0}^{m-1} (1-i)e_i \sum_{j=0}^{k+m-2-i} X_{G^{g+j}} X_{P_{h+k+m-i-j-1}} \\ &= \sum_{j=0}^{k+m-2} X_{G^{g+j}} \sum_{i=0}^{\min(m-1, k+m-j-2)} (1-i)e_i X_{P_{h+k+m-i-j-1}} \\ &= \sum_{j=0}^{k+m-2} X_{G^{g+k+m-j-2}} \sum_{i=0}^{\min(m-1, j)} (1-i)e_i X_{P_{h-i+j+1}}. \end{aligned}$$

By Lemma 4.4, we can further derive that

$$\begin{aligned} A &= \sum_{j=0}^{k+m-2} X_{G^{g+k+m-j-2}} \frac{X_{K_{\min(m, j+1)}^{h+j+1-\min(m, j+1)}}}{\min(m-1, j)!} \\ &= \sum_{j=0}^{m-2} X_{G^{g+k+m-j-2}} \frac{X_{K_{j+1}^h}}{j!} + \sum_{j=m-1}^{k+m-2} X_{G^{g+k+m-j-2}} \frac{X_{K_m^{h+j-m+1}}}{(m-1)!} \\ &= \sum_{j=1}^{m-1} X_{G^{g+k+m-j-1}} \frac{X_{K_j^h}}{(j-1)!} + \sum_{j=0}^{k-1} X_{G^{g+k-j+1}} \frac{X_{K_m^{h+j}}}{(m-1)!}. \end{aligned}$$

Substituting it back to the first formula in this proof, we obtain the desired one. \square

The pineapple graph Pi_m^λ studied in [30] is the graph $S_\lambda^0(K_m)$. Now we can give a formula for pineapples with $\ell(\lambda) = 2$. It is a graph obtained from a rooted clique (K_m, u) by adding two disjoint paths with an end u , whose lengths are λ_1 and λ_2 respectively.

Corollary 4.7. *For any $g \geq 0$ and $h, m \geq 1$,*

$$X_{S_{gh}^0(K_m)} = (m-1)! \left(\sum_{z=0}^{m-1} e_z X_{P_{g+h+m-z}} - \sum_{z=1}^{m-1} \frac{X_{K_z^h}}{(z-1)!} X_{P_{g+m-z}} \right).$$

Proof. Immediately by taking $G = K_1$ and $k = 0$ in Corollary 4.6. \square

We end this section with a reduction for the chromatic symmetric function of 3-spider-conjoined graphs $S^{ghk}(G, H, K)$ with a cycle node $K = C_m$.

Theorem 4.8. *Let G and H be rooted graphs. For any $g, h \geq 0$ and $m \geq 2$,*

$$X_{S^{ghk}(G, H, C_m)} = (m-1)X_{S_{k+m-1}^{gh}(G, H)} - \sum_{l=1}^{m-2} X_{S_{k+l-1}^{gh}(G, H)} X_{C_{m-l}}.$$

Proof. In the graph $W = S_0^{gh}(G, H)$, we set its root to be the spider center. Then

$$W^k = S_k^{gh}(G, H) \quad \text{and} \quad P^k(W, C_m) = S^{ghk}(G, H, C_m).$$

Taking $G = W$ in Proposition 3.2 yields the desired formula. \square

5. CHAIN-CONJOINED GRAPHS

This section is devoted to chain-conjoined graphs. A *double rooted graph* (G, u, v) is a triple consisting of a graph G and two vertices u and v of G . For our purpose, we restrict u and v to be distinct. We call u and v its *roots*. For any double rooted graphs $(G_0, u_0, v_0), \dots, (G_l, u_l, v_l)$, and any weak composition $\tau = \tau_1 \cdots \tau_l$, we define the *l-chain-conjoined graph* $G = C^\tau(G_0, \dots, G_l)$ to be the graph obtained by adding a path of length τ_i linking v_i and u_{i+1} for all i . It has order

$$|G| = |G_0| + \cdots + |G_l| + |\tau| - l,$$

see Fig. 4. Then 1-chain-conjoined graphs are path-conjoined graphs. We call G_i the *nodes* of G , and




FIGURE 4. The chain-conjoined graph $C(G_1, k_1, G_2, k_2, \dots, G_{l-1}, k_{l-1}, G_l)$.

omit the roots of G_i whenever G_i is vertex-transitive or the roots of G_i are clear from context. Note that the roots u_1 and v_l are useless in our configuration.

We consider 2-chain-conjoined graphs with the middle node $G_2 = K_m$ for $m \geq 2$, denoted

$$K_m^{gh}(G, H) = C^{gh}(G, K_m, H).$$

Theorem 5.1. *Let G and H be rooted graphs. Then for any $g, h \geq 0$ and $m \geq 2$,*

$$\frac{X_{K_m^{gh}(G, H)}}{(m-2)!} = \sum_{z=1}^{m-1} ze_{m-1-z} X_{P^{g+h+z}(G, H)} + (m-2) \sum_{\substack{a+b+c=m-2 \\ a, b, c \geq 0}} (a-1)e_a X_{G^{b+g}} X_{H^{c+h}}.$$

Proof. The desired formula holds for $m = 2$ since $K_2^{gh}(G, H) = P^{g+h+1}(G, H)$. Suppose that $m \geq 3$. We label the vertices of the clique K_m in G by $uv_1v_2 \cdots v_{m-1}$, where u (resp., v_{m-1}) is the root for conjoining K_m with G (resp., with H). Let $K_{m,k}^{gh}$ be the graph obtained from $K_m^{gh} = K_m^{gh}(G, H)$ by removing the edges uv_1, \dots, uv_k . By Corollary 2.4,

$$X_{K_m^{gh}} = (m-2)X_{K_{m,m-3}^{gh}} - (m-3)X_{K_{m,m-2}^{gh}}, \quad \text{for } m \geq 3.$$

Write $\mathfrak{K}_m^{gh} = S^{gh0}(G, H, K_m)$. Then the vertex in K_m that connects G with a path coincides the vertex that connects H with a path, and $K_{m,m-2}^{gh} = \mathfrak{K}_{m-1}^{(g+1)h}$. Applying Eq. (2.2) to the graph $K_{m,m-3}^{gh}$ with respect to the triangle uvv_{m-2} , we obtain

$$X_{K_{m,m-3}^{gh}} + X_{K_{m,m-1}^{gh}} = X_{K_{m-1}^{(g+1)h}} + X_{\mathfrak{K}_{m-1}^{(g+1)h}}, \quad \text{for } m \geq 3.$$

Since $K_{m,m-1}^{gh}$ is the disjoint union of the graphs G^g and $P^h(H, K_{m-1})$, we can deduce that

$$X_{K_m^{gh}} = (m-2)X_{K_{m-1}^{(g+1)h}} - (m-2)X_{G^g} X_{P^h(H, K_{m-1})} + X_{\mathfrak{K}_{m-1}^{(g+1)h}}$$

$$\begin{aligned}
&= (m-2) \left((m-3)X_{K_{m-2}^{(g+2)h}} - (m-3)X_{G^{g+1}}X_{P^h(H, K_{m-2})} + X_{\mathfrak{K}_{m-2}^{(g+2)h}} \right) \\
&\quad - (m-2)X_{G^g}X_{P^h(H, K_{m-1})} + X_{\mathfrak{K}_{m-1}^{(g+1)h}} = \dots \\
&= (m-2)!X_{K_2^{(g+m-2)h}} - \sum_{k=1}^{m-2} (m-2)_k X_{G^{g+k-1}}X_{P^h(H, K_{m-k})} + \sum_{k=1}^{m-2} (m-2)_{k-1} X_{\mathfrak{K}_{m-k}^{(g+k)h}} \\
&= \sum_{k=1}^{m-1} (m-2)_{k-1} X_{\mathfrak{K}_{m-k}^{(g+k)h}} - \sum_{k \geq 1} (m-2)_k X_{G^{g+k-1}}X_{P^h(H, K_{m-k})} \\
&= \sum_{k=0}^{m-2} (m-2)_k X_{S^{(g+k+1)h0}(G, H, K_{m-k-1})} - \sum_{k \geq 1} (m-2)_k X_{G^{g+k-1}}X_{P^h(H, K_{m-k})},
\end{aligned}$$

where $(m-2)_k = (m-2)(m-3)\cdots(m-k-1)$ stands for the falling factorial. By Theorem 4.5 and Proposition 3.1, we can infer that

$$\begin{aligned}
(5.1) \quad X_{K_m^{gh}} &= (m-2)! \sum_{k=0}^{m-2} \sum_{z=0}^{m-k-2} e_z X_{P^{g+h+m-1-z}(G, H)} \\
&\quad - (m-2)! \sum_{k=0}^{m-2} \sum_{\substack{a+b+c=g+h+m-2 \\ a \geq 0, b \geq g+k+1, c \geq h}} (1-a)e_a X_{G^b} X_{H^c} \\
&\quad - (m-2)! \sum_{k=1}^{m-2} X_{G^{g+k-1}} (m-k-1) \sum_{l=0}^{m-k-1} (1-l)e_l X_{H^{h+m-k-1-l}}.
\end{aligned}$$

Since the first double sum contains no parameter k , it can be simplified as

$$\sum_{z=0}^{m-2} (m-1-z)e_z X_{P^{g+h+m-1-z}(G, H)} = \sum_{z=1}^{m-1} z e_{m-1-z} X_{P^{g+h+z}(G, H)}.$$

We write $Q = (1-a)e_a X_{G^b} X_{H^c}$ for short. Since the upper bound of k for the second double sum is essentially $m-3$, we can process the last two double sums by bootstrapping as

$$\begin{aligned}
&\sum_{k=0}^{m-3} \sum_{\substack{a+b+c=g+h+m-2 \\ a \geq 0, b \geq g+k+1, c \geq h}} Q + \sum_{\substack{a+b+c=g+h+m-2 \\ a \geq 0, b \geq g, c \geq h}} (g+m-b-2)Q \\
&= \sum_{k=1}^{m-3} \sum_{\substack{a+b+c=g+h+m-2 \\ a \geq 0, b \geq g+k+1, c \geq h}} Q + \sum_{\substack{a+b+c=g+h+m-2 \\ a \geq 0, b \geq g+1, c \geq h}} (g+m-b-1)Q + (m-2) \sum_{\substack{a+c=h+m-2 \\ a \geq 0, c \geq h}} Q|_{b=g} \\
&= \sum_{k=2}^{m-3} \sum_{\substack{a+b+c=g+h+m-2 \\ a \geq 0, b \geq g+k+1, c \geq h}} Q + \sum_{\substack{a+b+c=g+h+m-2 \\ a \geq 0, b \geq g+2, c \geq h}} (g+m-b)Q + (m-2) \left(\sum_{\substack{a+c=h+m-3 \\ a \geq 0, c \geq h}} Q|_{b=g+1} + \sum_{\substack{a+c=h+m-2 \\ a \geq 0, c \geq h}} Q|_{b=g} \right) \\
&= \dots \\
&= \sum_{\substack{a+b+c=g+h+m-2 \\ a \geq 0, b \geq g+m-2, c \geq h}} (g+m-b+m-4)Q + (m-2) \sum_{i=0}^{m-3} \sum_{\substack{a+c=h+m-2-i \\ a \geq 0, c \geq h}} Q|_{b=g+i} \\
&= (m-2) \sum_{i=0}^{m-2} \sum_{\substack{a+c=h+m-2-i \\ a \geq 0, c \geq h}} Q|_{b=g+i} = (m-2) \sum_{k=0}^{m-2} \sum_{a=0}^{m-2-k} (1-a)e_a X_{G^{g+k}} X_{H^{h+m-2-k-a}}.
\end{aligned}$$

Substituting the previous two formulas into Eq. (5.1), we obtain the desired formula. \square

Theorem 5.1 reduces to a useful formula for PKPG graphs by taking $H = K_1$.

Theorem 5.2 (PKPGs). *For any $g, h \geq 0$ and $m \geq 2$,*

$$\frac{X_{K_m^{gh}(G, K_1)}}{(m-2)!} = \sum_{z=1}^{m-1} ze_{m-1-z} X_{G^{g+h+z}} - (m-2) \sum_{z=1}^{m-1} \frac{X_{K_z^h}}{(z-1)!} X_{G^{m-1-z+g}}.$$

Proof. Taking $H = K_1$ in Theorem 5.1, we obtain

$$\begin{aligned} \frac{X_{K_m^{gh}(G, K_1)}}{(m-2)!} - \sum_{z=1}^{m-1} ze_{m-1-z} X_{G^{g+h+z}} &= (m-2) \sum_{\substack{a+b+c=g+h+m-2 \\ a \geq 0, b \geq g, c \geq h}} (a-1)e_a X_{G^b} X_{P_{c+1}} \\ &= (2-m) \sum_{b=g}^{g+m-2} X_{G^b} \sum_{a=0}^{g+m-2-b} (1-a)e_a X_{P_{g+h+m-1-a-b}}. \end{aligned}$$

Using Lemma 4.4, we can simplify the rightmost side of the equation above as

$$(2-m) \sum_{b=g}^{g+m-2} X_{G^b} \frac{X_{K_{g+m-2-b+1}^h}}{(g+m-2-b)!} = (2-m) \sum_{b=0}^{m-2} \frac{X_{G^{m-2-b+g}} X_{K_{b+1}^h}}{b!}.$$

Rearranging the terms yields the desired formula. \square

By using the composition method with the aid of Theorem 5.2, we now derive positive e_I -expansions for the chromatic symmetric functions of two kinds of K -chains, see Theorems 5.3 and 5.4. Let

$$(5.2) \quad \begin{cases} f_1(I, a) = (a-1)w_I e_I, \\ f_2(I, a) = (a-2)i_{-1}w_{I \setminus i_{-1}} e_I, \quad \text{and} \\ f_3(I, a) = (i_{-1} - a + 1)w_{I \setminus i_{-1}} e_I, \end{cases}$$

where the symbol $I \setminus i_{-1}$ stands for the composition obtained from I by removing the last part. Then

$$(5.3) \quad f_1(I, a) - f_2(I, a) - f_3(I, a) = \begin{cases} (a-1)e_n, & \text{if } I = n, \\ 0, & \text{otherwise.} \end{cases}$$

A *path-clique-path* graph is a graph of the form $K_m^{gh}(K_1, K_1)$.

Theorem 5.3 (PKPs). *Let $n = g + h + m$, where $g, h \geq 0$ and $m \geq 2$. Then the path-clique-path graph $K_m^{gh}(K_1, K_1)$ has the positive e_I -expansion*

$$\frac{X_{K_m^{gh}(K_1, K_1)}}{(m-2)!} = (m-1)e_n + \sum_{I \models n, \Theta_I(h+1) \geq m-1} f_2(I, m) + \sum_{I \models n, i_{-1} \geq m-1} f_3(I, m),$$

where f_2 and f_3 are defined by Def. (5.2).

Proof. Let $G = K_m^{gh}(K_1, K_1)$. By Theorem 5.2 and Propositions 2.5 and 2.6,

$$\begin{aligned} \frac{X_G}{(m-2)!} &= \sum_{z=1}^{m-1} ze_{m-1-z} X_{P_{g+h+z+1}} - (m-2) \sum_{z=1}^{m-1} \frac{X_{P_{m-z+g}} X_{K_z^h}}{(z-1)!} \\ &= \sum_{z=0}^{m-1} (m-1-z)e_z X_{P_{n-z}} - (m-2) \sum_{z=1}^{m-1} \sum_{J \models m-z+g} w_J e_J \sum_{I \models h+z, i_{-1} \geq z} w_I e_I. \end{aligned}$$

It follows that

$$(5.4) \quad \frac{X_G}{(m-2)!} = \sum_{z=0}^{m-2} \sum_{I \models n} (m-1-z)w_I e_{Iz} - (m-2) \sum_{z=1}^{m-1} \sum_{IJ \models n, I \models h+z, i_{-1} \geq z} w_I w_J e_{IJ}.$$

For the first double sum, we separate its summand for $z = 0$ and rewrite it as

$$(5.5) \quad \sum_{z=0}^{m-2} \sum_{I \models n} (m-1-z) w_I e_{Iz} = \sum_{K \models n} (m-1) w_K e_K + \sum_{K \models n, k_{-1} \leq m-2} (m-1-k_{-1}) w_{K \setminus k_{-1}} e_K.$$

In the second double sum, the composition J can be equivalently replaced with its reversal \bar{J} ; namely,

$$\sum_{z=1}^{m-1} \sum_{IJ \models n, I \models h+z, i_{-1} \geq z} w_I w_J e_{IJ} = \sum_{z=1}^{m-1} \sum_{IJ \models n, I \models h+z, i_{-1} \geq z} w_I w_{\bar{J}} e_{IJ}.$$

We regard IJ as a whole competition K . On the one hand, since $|I| = h+z \leq h+i_{-1}$, we find $i_1 + \dots + i_{-2} \leq h$. Since $|I| = h+z \geq h+1$, the composition I must be the shortest prefix of K such that $|I| \geq h+1$. On the other hand, $w_I w_{\bar{J}} e_{IJ} = k_{-1} w_{K \setminus k_{-1}} e_K$. As a result, we can rewrite the second double sum as

$$(5.6) \quad \sum_{z=1}^{m-1} \sum_{IJ \models n, I \models h+z, i_{-1} \geq z} w_I w_J e_{IJ} = \sum_{K \models n, \Theta_K(h+1) \leq m-2} k_{-1} w_{K \setminus k_{-1}} e_K.$$

Substituting Eqs. (5.5) and (5.6) into Eq. (5.4), we obtain

$$\frac{X_G}{(m-2)!} = \sum_{K \models n} f_1(K, m) - \sum_{K \models n, \Theta_K(h+1) \leq m-2} f_2(K, m) - \sum_{K \models n, k_{-1} \leq m-2} f_3(K, m).$$

We proceed in 4 cases. Let $K \models n$, $x = \Theta_K(h+1)$, $y = k_{-1}$, and $I = K \setminus k_{-1}$. Denote by c_K the coefficient of e_K in $X_G/(m-2)!$. Here we distinguish c_K and c_H if K and H are distinct compositions. Then we have the following arguments.

- (1) If $x, y \geq m-1$, then $c_K = (m-1) w_K$.
- (2) If $x \geq m-1 > y$, then $c_K = (m-1) w_K + (m-1-y) w_I = (m-2) y w_I$.
- (3) If $x, y < m-1$, then $c_K = 0$ by the previous item.
- (4) If $x < m-1 \leq y$, then $c_K = (m-1) w_K - (m-2) y w_I = (y-m+1) w_I$.

Gathering them together, we obtain

$$\frac{X_{K_m^{gh}(K_1, K_1)}}{(m-2)!} = \sum_{\substack{K \models n, k_{-1} \leq m-2 \\ \Theta_K(h+1) \geq m-1}} f_2(K, m) + \sum_{\substack{K \models n, k_{-1} \geq m-1 \\ \Theta_K(h+1) \geq m-1}} f_1(K, m) + \sum_{\substack{K \models n, k_{-1} \geq m-1 \\ \Theta_K(h+1) \leq m-2}} f_3(K, m).$$

By Eq. (5.3), one may recast it as desired. \square

For example, by Theorem 5.3 we can calculate

$$X_{K_4^{21}(K_1, K_1)} = 24e_{142} + 16e_{151} + 24e_{16} + 2e_{421} + 6e_{43} + 48e_{52} + 30e_{61} + 42e_7.$$

A *clique-clique-path* graph is a graph of the form $K_{b+1}^{0a}(K_c, K_1)$.

Theorem 5.4 (KKPs). *Let $n = a + b + c$, where $a \geq 0$, $b \geq 1$ and $c \geq 1$. Then the chromatic symmetric function of the clique-clique-path graph $K_{b+1}^{0a}(K_c, K_1)$ has the positive e_I -expansion*

$$\frac{X_{K_{b+1}^{0a}(K_c, K_1)}}{(b-1)!(c-1)!} = \sum_{\substack{K \models n \\ k_{-1} \geq b+c}} b w_K e_K + \sum_{\substack{K \models n, \ell(K) \geq 2, k_{-1}+k_{-2} \geq b+c \\ k_{-1} \leq \min(b-1, c-1) \text{ or } k_{-1} \geq \max(b+1, c)}} |b-k_{-1}| w_{K \setminus k_{-1}} e_K,$$

where the symbol $I \setminus i_{-1}$ stands for the composition obtained from I by removing the last part, and the function w is defined by Def. (2.4).

Proof. Let $G = X_{K_{b+1}^{0a}(K_c, K_1)}$. By Theorem 5.2 and Proposition 2.6,

$$\begin{aligned} \frac{X_G}{(b-1)!} &= \sum_{z=1}^b z e_{b-z} X_{K_c^{a+z}} - (b-1) \sum_{z=1}^b X_{K_c^{b-z}} \sum_{J \models a+z, j_{-1} \geq z} w_J e_J \\ &= \sum_{z=1}^b z e_{b-z} \sum_{I \models a+c+z, i_{-1} \geq c} (c-1)! w_I e_I - (b-1)(c-1)! \sum_{z=1}^b \sum_{\substack{I \models b+c-z, i_{-1} \geq c \\ J \models a+z, j_{-1} \geq z}} w_I w_J e_{IJ}. \end{aligned}$$

Separating the term for $z = b$ from the first double sum, we can rewrite it as

$$\frac{X_G}{(b-1)!(c-1)!} = Y_1 + Y_2 - Y_3,$$

where $Y_1 = b \sum_{K \models n, k_{-1} \geq c} w_K e_K$,

$$\begin{aligned} Y_2 &= \sum_{z=1}^{b-1} \sum_{\substack{I \models a+c+z \\ i_{-1} \geq c}} z w_I e_{i_1 \cdots i_{-2}(b-z)i_{-1}} = \sum_{\substack{K \models n, k_{-1} \geq c \\ \ell(K) \geq 2, k_{-2} \leq b-1}} (b-k_{-2}) w_{K \setminus k_{-2}} e_K, \quad \text{and} \\ Y_3 &= (b-1) \sum_{z=1}^b \sum_{\substack{I \models a+b+1-z, i_{-1} \geq b-z+1 \\ J \models c+z-1, j_{-1} \geq c}} w_I w_J e_{IJ}. \end{aligned}$$

In the last expression, since $a+1 \leq |I| \leq a+i_{-1}$, the composition I must be the shortest prefix of the composition IJ such that $|I| \geq a+1$. It follows that

$$(5.7) \quad Y_3 = (b-1) \sum_{K=IJ \models n, j_{-1} \geq c, |I|=\sigma_K(a+1)} w_I w_J e_K,$$

where the function $\sigma_K(\cdot)$ is defined by Def. (2.5). We claim that

$$(5.8) \quad Y_3 = \sum_{\substack{K \models n, c \leq k_{-1} \leq b+c-1 \\ \ell(K) \geq 2, k_{-1}+k_{-2} \geq b+c}} (b-1)k_{-1} w_{K \setminus k_{-1}} e_K + \sum_{\substack{K \models n, k_{-1} \geq c \\ k_{-1}+k_{-2} \leq b+c-1}} (b-1)k_{-2} w_{K \setminus k_{-2}} e_K.$$

In fact, we separate the sum in Eq. (5.7) according to whether $\ell(J) = 1$.

(1) For $\ell(J) = 1$, we have

$$w_I w_J = k_{-1} w_{K \setminus k_{-1}}, \quad k_{-1} \leq b+c-1, \quad \ell(K) \geq 2, \quad \text{and} \quad k_{-1} + k_{-2} \geq b+c.$$

Conversely, if these three inequalities hold, then $|k_1 \cdots k_{-3}| \leq a$. Since $|I| = \sigma_K(a+1)$, we find $k_1 \cdots k_{-3}$ must be a proper prefix of I . Since J is not empty, we derive that $I = k_1 \cdots k_{-2}$ and $\ell(J) = 1$. Thus the terms in Eq. (5.7) with $\ell(J) = 1$ sum to the first sum in Eq. (5.8).

(2) For $\ell(J) \geq 2$, define J' to be the composition obtained by moving j_{-2} to the beginning of J , i.e., $J' = j_{-2} j_1 j_2 \cdots j_{-3} j_{-1}$. When J runs over the set $\{J \models n - |I| : j_{-1} \geq c\}$, so does J' , and vice versa. If we write $K = IJ'$, then

$$w_I w_{J'} = k_{-2} w_{K \setminus k_{-2}} \quad \text{and} \quad k_{-1} + k_{-2} \leq b+c-1.$$

Conversely, if $k_{-1} + k_{-2} \leq b+c-1$, then $|k_1 \cdots k_{-3}| \geq a+1$, which implies that I is a prefix of $k_1 \cdots k_{-3}$, and $\ell(J) \geq 2$. Thus the terms in Eq. (5.7) with $\ell(J) \geq 2$ sum to the second sum in Eq. (5.8).

This proves Eq. (5.8). Now we compute $Y_1 - Y_3$. Since for $i \in \{1, 2\}$,

$$bw_K - (b-1)k_{-i} w_{K \setminus k_{-i}} = w_{K \setminus k_{-i}} (b(k_{-i} - 1) - (b-1)k_{-i}) = (k_{-i} - b) w_{K \setminus k_{-i}},$$

we obtain $Y_1 - Y_3 = Q_1 + Q_2 + Q_3$, where

$$\begin{aligned} Q_1 &= \sum_{K \models n, k_{-1} \geq b+c} bw_K e_K, \\ Q_2 &= \sum_{\substack{K \models n, c \leq k_{-1} \leq b+c-1 \\ \ell(K) \geq 2, k_{-1}+k_{-2} \geq b+c}} (k_{-1} - b) w_{K \setminus k_{-1}} e_K, \quad \text{and} \\ Q_3 &= \sum_{\substack{K \models n, k_{-1} \geq c \\ \ell(K) \geq 2, k_{-1}+k_{-2} \leq b+c-1}} (k_{-2} - b) w_{K \setminus k_{-2}} e_K. \end{aligned}$$

Note that in Q_3 , we always have $k_{-2} \leq b-1$. Therefore,

$$\begin{aligned} Y_2 + Q_3 &= \sum_{\substack{K \models n, k_{-1} \geq c \\ \ell(K) \geq 2, k_{-2} \leq b-1}} (b - k_{-2}) w_{K \setminus k_{-2}} e_K + \sum_{\substack{K \models n, k_{-1} \geq c \\ \ell(K) \geq 2, k_{-1}+k_{-2} \leq b+c-1}} (k_{-2} - b) w_{K \setminus k_{-2}} e_K \\ &= \sum_{\substack{K \models n, k_{-1} \geq c \\ \ell(K) \geq 2, k_{-2} \leq b-1 \\ k_{-1}+k_{-2} \geq b+c}} (b - k_{-2}) w_{K \setminus k_{-2}} e_K = \sum_{\substack{K \models n, \ell(K) \geq 2, \\ k_{-2} \leq b-1 \\ k_{-1}+k_{-2} \geq b+c}} (b - k_{-2}) w_{K \setminus k_{-2}} e_K. \end{aligned}$$

We then deal with the summands in Q_2 with $k_{-1} \leq b-1$ by exchanging k_{-2} and k_{-1} . As a result, we can write $Q_2 = Q_{21} + Q_{22}$, where

$$\begin{aligned} Q_{21} &= \sum_{\substack{K \models n, \max(b, c) \leq k_{-1} \leq b+c-1 \\ \ell(K) \geq 2, k_{-1}+k_{-2} \geq b+c}} (k_{-1} - b) w_{K \setminus k_{-1}} e_K, \quad \text{and} \\ Q_{22} &= \sum_{\substack{K \models n, c \leq k_{-2} \leq b-1 \\ \ell(K) \geq 2, k_{-1}+k_{-2} \geq b+c}} (k_{-2} - b) w_{K \setminus k_{-2}} e_K. \end{aligned}$$

We observe that Q_{22} cancels out a part of $Y_2 + Q_3$ as

$$\begin{aligned} (Y_2 + Q_3) + Q_{22} &= \sum_{\substack{K \models n, \ell(K) \geq 2 \\ k_{-2} \leq b-1 \\ k_{-1}+k_{-2} \geq b+c}} (b - k_{-2}) w_{K \setminus k_{-2}} e_K + \sum_{\substack{K \models n, \ell(K) \geq 2 \\ c \leq k_{-2} \leq b-1 \\ k_{-1}+k_{-2} \geq b+c}} (k_{-2} - b) w_{K \setminus k_{-2}} e_K \\ &= \sum_{\substack{K \models n, \ell(K) \geq 2 \\ k_{-2} \leq \min(b, c) - 1 \\ k_{-1}+k_{-2} \geq b+c}} (b - k_{-2}) w_{K \setminus k_{-2}} e_K = \sum_{\substack{K \models n, \ell(K) \geq 2 \\ k_{-1} \leq \min(b, c) - 1 \\ k_{-1}+k_{-2} \geq b+c}} (b - k_{-1}) w_{K \setminus k_{-1}} e_K. \end{aligned}$$

The parameters k_{-1} and k_{-2} are exchanged again in the last step. Collecting the results in

$$\frac{X_G}{(b-1)!(c-1)!} = Q_1 + Q_{21} + (Y_2 + Q_3 + Q_{22}),$$

we obtain the desired formula. \square

For example, by Theorem 5.4, we can calculate

$$X_{K_7^{01}(K_3, K_1)} = 48e_{126} + 720e_{162} + 1152e_{171} + 1680e_{18} + 192e_{27} + 144e_{36} + 1008e_{72} + 1536e_{81} + 2160e_9.$$

At the end of this paper, we consider 2-chain-conjoined graphs with the middle node $G_2 = C_m$ for $m \geq 3$, with an additional condition that the two nodes of G_2 are adjacent, denoted

$$C_m^{gh}(G, H) = C^{gh}(G, C_m, H).$$

In particular, the 2-chain-conjoined graph $C_m^{ab}(K_1, K_1)$ is the hat graph $H_{a,m,b}$ studied in [34]. The adjacency condition for the roots of C_m is essential for the e -positivity. For instance, the graph $G = C^{11}(K_1, C_4, K_1)$ in which the roots of C_4 are non-adjacent is not e -positive as

$$X_G = 18e_6 + 22e_{51} - 2e_{42} + 6e_{411} + 9e_{33} + 4e_{321} - 2e_{222} + e_{2211}.$$

Theorem 5.5 (GCHs). *Let (C_m, x, y) be a double-rooted cycle with adjacent roots x and y , where $m \geq 2$. For any rooted graphs (G, v_G) and (H, v_H) and any $g, h \geq 0$, let $C_m^{gh}(G, H)$ be the graph obtained by adding a path of length g linking x and v_G and another path of length h linking y and v_H . Then*

$$X_{C_m^{gh}(G, H)} = \sum_{\substack{a+b=g+h+m-1 \\ a \geq 0, b \geq g+h+1}} (a+1) X_{P_a} X_{P^b(G, H)} + \sum_{\substack{a+b+c=g+h+m-2 \\ a \geq g, b \geq h, c \geq 2}} X_{G^a} X_{H^b} X_{C_c} - (m-2) \sum_{\substack{a+b=g+h+m-2 \\ a \geq g, b \geq h}} X_{G^a} X_{H^b}.$$

Proof. We omit the variables G and H for short and write

$$P^k(G, H) = P^k, \quad S_m^{gh}(G, H) = S_m^{gh} \quad \text{and} \quad C_m^{gh}(G, H) = C_m^{gh}.$$

For convenience, we label the vertices of the clique C_m in G by v_0, \dots, v_{m-1} , where $u = v_0$ and $v = v_{m-1}$. Applying the triple-deletion property to the triangle uvv_1 , we obtain

$$X_{C_m^{gh}} = X_{C_{m-1}^{(g+1)h}} + X_{S_{m-2}^{(g+1)h}} - X_{G^g} X_{P^h(H, C_{m-1})}, \quad \text{for } m \geq 3.$$

Iteratively using it, we can infer that

$$(5.9) \quad X_{C_m^{gh}} = \sum_{k=1}^{m-1} X_{S_{m-1-k}^{(g+k)h}} - \sum_{k=1}^{m-2} X_{G^{g+k-1}} X_{P^h(H, C_{m-k})}.$$

This leads us to deal with the graphs $S_{m-1-k}^{(g+k)h}$ and $P^h(H, C_{m-k})$, respectively. On one hand, in Corollary 4.3, replacing k with $m-1-k$ and replacing g with $g+k$, we obtain

$$X_{S_{m-1-k}^{(g+k)h}} = \sum_{l=0}^{m-1-k} X_{P_{m-1-k-l}} X_{P^{g+h+k+l}} - \sum_{l=0}^{m-2-k} X_{G^{g+m-2-l}} X_{H^{h+l}}.$$

It follows that

$$(5.10) \quad \begin{aligned} \sum_{k=1}^{m-1} X_{S_{m-1-k}^{(g+k)h}} &= \sum_{k=1}^{m-1} \sum_{l=k}^{m-1} X_{P_{m-1-l}} X_{P^{g+h+l}} - \sum_{k=1}^{m-1} \sum_{l=0}^{m-2-k} X_{G^{g+m-2-l}} X_{H^{h+l}} \\ &= \sum_{l=1}^{m-1} (m-l) X_{P_{m-1-l}} X_{P^{g+h+l}} - \sum_{l=0}^{m-3} (m-2-l) X_{G^{g+m-2-l}} X_{H^{h+l}}. \end{aligned}$$

On the other hand, in Proposition 3.2, taking $k = h$, $G = H$, and setting m to be $m-k$, we obtain

$$X_{P^h(H, C_{m-k})} = (m-k-1) X_{H^{h+m-k-1}} - \sum_{l=1}^{m-k-2} X_{H^{h+l-1}} X_{C_{m-k-l}}.$$

It follows that

$$(5.11) \quad \begin{aligned} \sum_{k=1}^{m-2} X_{G^{g+k-1}} X_{P^h(H, C_{m-k})} &= \sum_{k=1}^{m-2} (m-k-1) X_{G^{g+k-1}} X_{H^{h+m-k-1}} - \sum_{k=1}^{m-2} \sum_{l=1}^{m-k-2} X_{G^{g+k-1}} X_{H^{h+l-1}} X_{C_{m-k-l}} \\ &= \sum_{l=1}^{m-2} l \cdot X_{G^{g+m-l-2}} X_{H^{h+l}} - \sum_{l=1}^{m-3} \sum_{k=1}^{m-2-l} X_{G^{g+k-1}} X_{H^{h+l-1}} X_{C_{m-k-l}}. \end{aligned}$$

Substituting Eqs. (5.10) and (5.11) into Eq. (5.9), we obtain

$$X_{C_m^{gh}} = \sum_{l=1}^{m-1} (m-l) X_{P_{m-1-l}} X_{P^{g+h+l}} - \sum_{l=0}^{m-3} (m-2-l) X_{G^{g+m-2-l}} X_{H^{h+l}}$$

$$\begin{aligned}
& - \sum_{l=1}^{m-2} l X_{G^{g+m-l-2}} X_{H^{h+l}} + \sum_{l=1}^{m-3} \sum_{k=1}^{m-2-l} X_{G^{g+k-1}} X_{H^{h+l-1}} X_{C_{m-k-l}} \\
& = \sum_{l=1}^{m-1} (m-l) X_{P_{m-1-l}} X_{P^{g+h+l}} + \sum_{l=0}^{m-4} \sum_{k=0}^{m-4-l} X_{G^{g+k}} X_{H^{h+l}} X_{C_{m-k-l-2}} - (m-2) \sum_{l=0}^{m-2} X_{G^{g+m-2-l}} X_{H^{h+l}} \\
& = \sum_{k=1}^{m-1} k \cdot X_{P_{k-1}} X_{P^{g+h+m-k}} + \sum_{\substack{k+l \leq m-4, k, l \geq 0}} X_{G^{g+k}} X_{H^{h+l}} X_{C_{m-k-l-2}} - (m-2) \sum_{l=1}^{m-1} X_{G^{g+m-1-l}} X_{H^{h+l-1}},
\end{aligned}$$

which simplifies to the desired formula. \square

A *hat-chain* is a $(l+1)$ -chain-conjoined graph of the form $C^{\tau_0 \tau_1 \cdots \tau_l}(K_1, C_{m_1}, C_{m_2}, \dots, C_{m_l}, K_1)$, where C_j are double-rooted cycles with adjacent roots, see Fig. 5.




FIGURE 5. The hat-chain $C^{\tau_0 \tau_1 \cdots \tau_l}(K_1, C_{m_1}, C_{m_2}, \dots, C_{m_l}, K_1)$.

Conjecture 5.6. *All hat-chains are e -positive.*

Wang and Zhou [34] confirmed its truth for $l = 1$, i.e., when the hat-chain is a single hat. When $l = 2$ and $\tau_0 = \tau_2 = 0$, the hat-chains becomes kayak paddles, and Conjecture 5.6 is true by Proposition 3.4. In the first version of our manuscript (arXiv:2406.01418v1), the conjecture for hat-chains was stated with the restriction $\tau_0 = \tau_l = 0$. In fact, it can be considered equivalent to Conjecture 5.6 if one allows $m_1 = m_l = 2$. Recently, Tom and Vailaya [29] found an involution proof for confirming Conjecture 5.6.

ACKNOWLEDGMENT

The third author is appreciative of the hospitality of the LIGM at Université Gustave Eiffel, where he was visiting Professor Jean-Yves Thibon in 2024. This paper was written and submitted during that period. The authors thank Watson Wang for catching an error in an early version of Theorem 4.5.

REFERENCES

- [1] F. Aliniaeifard, V. Wang, and S. van Willigenburg. The chromatic symmetric function of a graph centred at a vertex. *Electron. J. Combin.*, 31(4):#P4.22, 2024. [3](#), [4](#), [8](#)
- [2] E. Banaian, K. Celano, M. Chang-Lee, L. Colmenarejo, O. Goff, J. Kimble, L. Kimpel, J. Lentfer, J. Liang, and S. Sundaram. The e -positivity of the chromatic symmetric function for twinned paths and cycles. *Discrete Math.*, 348(12):114687, 2025. [1](#)
- [3] P. Brosnan and T. Y. Chow. Unit interval orders and the dot action on the cohomology of regular semisimple Hessenberg varieties. *Adv. Math.*, 329:955–1001, 2018. [1](#)
- [4] S. Chmutov, M. Kazarian, and S. Lando. Polynomial graph invariants and the KP hierarchy. *Selecta Math. (N.S.)*, 26:Article no. 34, 2020. [1](#)
- [5] S. Cho and J. Huh. On e -positivity and e -unimodality of chromatic quasi-symmetric functions. *SIAM J. Discrete Math.*, 33(4):2286–2315, 2019. [1](#)
- [6] S. Dahlberg and S. van Willigenburg. Lollipop and lariat symmetric functions. *SIAM J. Discrete Math.*, 32(2):1029–1039, 2018. [1](#)
- [7] S. Dahlberg, A. Foley, and S. van Willigenburg. Resolving Stanley’s e -positivity of claw-contractible-free graphs. *J. European Math. Soc.*, 22(8):2673–2696, 2020. [1](#)

- [8] A. M. Foley, C. T. Hoàng, and O. D. Merkel. Classes of graphs with e -positive chromatic symmetric function. *Electron. J. Combin.*, 26(3):#P3.51, 2019. [1](#)
- [9] D. D. Gebhard and B. E. Sagan. A chromatic symmetric function in noncommuting variables. *J. Alg. Combin.*, 13(3):227–255, 2001. [1](#), [2](#), [6](#)
- [10] M. Guay-Paquet. A modular relation for the chromatic symmetric functions of (3+1)-free posets. arXiv:1306.2400, 2013. [1](#)
- [11] A. M. Hamel, C. T. Hoàng, and J. E. Tuero. Chromatic symmetric functions and H -free graphs. *Graphs Combin.*, 35(4):815–825, 2019. [1](#)
- [12] T. Hikita. On the Stanley–Stembridge conjecture. *Sém. Lothar. Combin.*, Article #31:12 pp., 2025. [1](#)
- [13] S. Kato. A geometric realization of the chromatic symmetric function of a unit interval graph. arXiv:2410.12231, 2024. [1](#)
- [14] E. Y. H. Li, G. M. X. Li, D. G. L. Wang, and A. L. B. Yang. The twinning operation on graphs does not always preserve e -positivity. *Taiwanese J. Math.*, 25(6):1089–1111, 2021. [1](#), [7](#)
- [15] G. M. X. Li and A. L. B. Yang. On the e -positivity of (claw, $2K_2$)-free graphs. *Electron. J. Combin.*, 28(2):#P2.40, 2021. [1](#)
- [16] M. Loeb. Chromatic polynomial, q -binomial counting and colored Jones function. *Adv. Math.*, 211(2):546–565, 2007. [1](#)
- [17] J. McDonough, P. Pylyavskyy, and S. Wang. The Stanley–Stembridge conjecture for $\mathbf{2} + \mathbf{1} + \mathbf{1}$ -avoiding unit interval orders: a diagrammatic proof. arXiv:2404.07280, 2024. [2](#)
- [18] S. D. Noble and D. J. A. Welsh. A weighted graph polynomial from chromatic invariants of knots. *Ann. Inst. Fourier (Grenoble)*, 49(3):1057–1087, 1999. [1](#)
- [19] R. Orellana and G. Scott. Graphs with equal chromatic symmetric functions. *Discrete Math.*, 320:1–14, 2014. [3](#)
- [20] J. Shareshian and M. L. Wachs. Eulerian quasisymmetric functions. *Adv. Math.*, 225(6):2921–2966, 2010. [4](#)
- [21] J. Shareshian and M. L. Wachs. Chromatic quasisymmetric functions. *Adv. Math.*, 295:497–551, 2016. [1](#), [2](#), [4](#)
- [22] R. P. Stanley. A symmetric function generalization of the chromatic polynomial of a graph. *Adv. Math.*, 111(1):166–194, 1995. [1](#), [3](#)
- [23] R. P. Stanley. *Enumerative Combinatorics, Vol. 1*, volume 49 of *Cambridge Stud. in Adv. Math.* Camb. Univ. Press, Cambridge, 2nd edition, 2011. It is a thoroughly revised version of the 1st edition published in 1986. [3](#)
- [24] R. P. Stanley and J. R. Stembridge. On immanants of Jacobi–Trudi matrices and permutations with restricted position. *J. Combin. Theory Ser. A*, 62(2):261–279, 1993. [1](#)
- [25] D. Q. B. Tang and D. G. L. Wang. Positive e -expansions of the chromatic symmetric functions of KPKPs, twinned lollipops, and kayak paddles. arXiv:2408.01385, 2024. [8](#)
- [26] J.-Y. Thibon and D. G. L. Wang. A noncommutative approach to the Schur positivity of chromatic symmetric functions. arXiv:2305.07858, 2023. [2](#)
- [27] F. Tom. A signed e -expansion of the chromatic symmetric function and some new e -positive graphs. In *Proceedings of the 36th Conference on Formal Power Series and Algebraic Combinatorics, Sém. Lothar. Combin.*, volume 91B, Article #48, Bochum, 2024. [2](#), [4](#), [5](#), [6](#)
- [28] F. Tom. A signed e -expansion of the chromatic quasisymmetric function. *Combin. Theory*, 5(2), 2025. [2](#)
- [29] F. Tom and A. Vailaya. Adjacent cycle-chains are e -positive. arXiv:2410.21762, to appear in *Electron. J. Combin.*, 2025. [19](#)
- [30] H. Topcu, S. Sorgun, and W. H. Haemers. On the spectral characterization of pineapple graphs. *Linear Algebra Appl.*, 507:267–273, 2016. [2](#), [11](#)
- [31] S. Tsujie. The chromatic symmetric functions of trivially perfect graphs and cographs. *Graphs Combin.*, 34(5):1037–1048, 2018. [1](#)
- [32] D. G. L. Wang. All cycle-chords are e -positive. *Ann. Comb.*, 2025. doi: 10.1007/s00026-025-00753-2. [1](#)
- [33] D. G. L. Wang and M. M. Y. Wang. The e -positivity of two classes of cycle-chord graphs. *J. Alg. Combin.*, 57(2):495–514, 2023. [1](#)
- [34] D. G. L. Wang and J. Z. Zhou. A composition method for neat formulas of chromatic symmetric functions. *Adv. Appl. Math.*, 167:102886, 2025. [2](#), [5](#), [6](#), [18](#), [19](#)
- [35] K. Zheng. On the e -positivity of trees and spiders. *J. Combin. Theory Ser. A*, 189:105608, 2022. [5](#)

(Ethan Y.J. Qi) SCHOOL OF MATHEMATICS AND STATISTICS, BEIJING INSTITUTE OF TECHNOLOGY, BEIJING 102400, P. R. CHINA.

Email address: ethane@bit.edu.cn

(Davion Q.B. Tang) SCHOOL OF MATHEMATICS AND STATISTICS, BEIJING INSTITUTE OF TECHNOLOGY, BEIJING 102400, P. R. CHINA

Email address: davion@bit.edu.cn

(David G.L. Wang) SCHOOL OF MATHEMATICS AND STATISTICS & MIIT KEY LABORATORY OF MATHEMATICAL THEORY AND COMPUTATION IN INFORMATION SECURITY, BEIJING INSTITUTE OF TECHNOLOGY, BEIJING 102400, P. R. CHINA.

Email address: glw@bit.edu.cn