

MODULAR OPERADS, ITERATED DISTRIBUTIVE LAWS AND A NERVE THEOREM FOR CIRCUIT ALGEBRAS

SOPHIE RAYNOR

ABSTRACT. Circuit algebras are a symmetric version of Jones’s planar algebras. They originated in quantum topology as a framework for encoding virtual crossings. Oriented circuit algebras are equivalent to wheeled props. This paper extends existing results for modular operads to construct a graphical calculus and monad for general circuit algebras and prove an abstract nerve theorem. Specialisations of these results to wheeled props follow as straightforward corollaries. The proof relies on a subtle interplay between distributive laws and abstract nerve theory, and provides extra insights into the underlying structures.

1. INTRODUCTION

Circuit algebras are a symmetric version of Jones’s planar algebras, introduced in [3] as a framework for studying finite-type invariants of virtual knotted objects (see also [12, 25, 49]). They are graded symmetric monoids equipped with a unary contraction operation satisfying suitable associativity and distributivity axioms. These axioms imply that circuit algebras may be equivalently described as algebras for an operad of wiring diagrams, as symmetric monoidal functors from a category of Brauer diagrams, or as modular operads in a monoid subcategory of symmetric species [42].

Besides quantum topology, these structures arise in many areas of mathematics, including invariant theory [15, 42] and string theory [45]. In particular, oriented circuit algebras are equivalent to wheeled props [13, 42] which govern algebraic structures with trace such as those that appear in contexts including Batalin–Vilkovisky quantisation and deformation theory (see e.g., [15, 36, 38]).

This paper provides a detailed technical account of circuit algebra combinatorics. Building on analogous results for modular operads [41], a monad and graphical calculus for circuit algebras is constructed and a nerve theorem in the style of Weber [6, 51] is proved.

Theorem 1.1 (Equation (7.9)). *The category \mathfrak{CA} of circuit algebras is equivalent to the Eilenberg–Moore (EM) category of algebras for a monad \mathbb{O} on Joyal and Kock’s graphical species category \mathbf{GS} [27].*

Let $\mathbf{psh}(\mathbf{C})$ denote the category of presheaves (contravariant \mathbf{Set} -valued functors) on a category \mathbf{C} .

Theorem 1.2 (Equation (8.4)). *There is a full subcategory $\Xi^\times \hookrightarrow \mathfrak{CA}$, whose objects are undirected graphs. The nerve functor $N: \mathfrak{CA} \rightarrow \mathbf{psh}(\Xi^\times)$ induced by the inclusion $\Xi^\times \hookrightarrow \mathfrak{CA}$ is fully faithful.*

A presheaf P on Ξ^\times is equivalent to the nerve of a circuit algebra if and only if it is “Segal”.

Put simply, the graphs encode ways of combining circuit algebra operations, and the “Segal condition” says that complex combinations of these operations are completely determined by their constituent parts (see Section 2.3 and e.g., [21]).

Date: December 5, 2025.

The author acknowledges the support of Australian Research Council grants DP160101519 and FT160100393.

These theorems apply simultaneously to a very general class of circuit algebras that includes (non)oriented and coloured versions. A monad and nerve theorem for wheeled props, in terms of a category of directed graphs, follow as immediate corollaries (see Equation (8.6)).

The proof of Equation (1.2) uses Weber’s abstract nerve theory [6, 51]. However, unlike the nerve theorems for categories and operads described in the examples of [51, Section 4], the Segal condition for circuit algebras in Equation (8.4) (as with the Segal condition for modular operads in [41, Theorem 8.2]) cannot be directly obtained with Weber’s machinery: while Theorems 1.1 and 1.2 imply that the circuit algebra monad \mathbb{O} is “nervous” in the sense of [7], it does not “have arities” in the sense of [6, 51].

In [41, Introduction & Section 6], I explained how the so-called “problem of loops”, that is related to contractions of modular operadic units (the unit trace), means that the monad for unital modular operads does not admit a straightforward description in terms of the “graph-substitution” notion that underlies analogous constructions for operads [39] or prop(erad)s [22, 31, 36]. This is also an issue for constructing the monad \mathbb{O} of Equation (1.1).

The central principle underlying this work (and [41]) is that, under certain conditions, it is possible to build up a description of complicated algebraic structures by suitably combining simpler ones. In particular, if the simpler structures are governed by monads, we may be able to reframe this as a problem of “composing” these monads. Distributive laws [5] tell us whether and how monads can be composed and how they interact in the composite. Iterated distributive laws [10] describe conditions under which multiple monads may be composed.

In [41, Section 7], the modular operad monad was obtained as a composition $\mathbb{D}\mathbb{T}$, via a distributive law, of monads \mathbb{T} (governing composition) and \mathbb{D} (governing units) on \mathbf{GS} . For circuit algebras (Section 7.2), it is even necessary to extend the construction in [41] to include a third monad \mathbb{L} (governing the monoidal structure) and iterated distributive laws.

Theorem 1.3 (Equation (7.8) & Equation (7.9)). *There is a triple of monads \mathbb{L}, \mathbb{D} and \mathbb{T} on \mathbf{GS} and iterated distributive laws between them such that the induced composite $\mathbb{L}\mathbb{D}\mathbb{T}$ is the circuit algebra monad \mathbb{O} from Equation (1.1).*

The decomposition in Equation (1.3) is crucial to the proof of Equation (1.2). Weber’s nerve machinery [6, 51] relies on the structure of algebras for a monad \mathbb{M} on a category \mathbf{C} being fixed by some proper subcategory $\mathbf{D} \subsetneq \mathbf{C}$, that is said to *provide arities* for \mathbb{M} . This is only satisfied under very special conditions: the circuit algebra monad \mathbb{O} , for example, does not have arities. However, if $\mathbb{M} = \mathbb{M}_n \dots \mathbb{M}_1$ is a composite monad, then each subset of $\{\mathbb{M}_1, \dots, \mathbb{M}_n\}$ describes a monad on \mathbf{C} . The proof of Equation (1.2) utilises the idea that, a suitable extension of this monad, if it exists, may have arities.

The iterated distributive laws used to prove Theorems 1.1–1.3 illuminate the graphical combinatorics of circuit algebras, and dovetail nicely with variations of circuit algebras that appear in the literature: Algebras for \mathbb{L} are monoids in \mathbf{GS} (Equation (3.20)) so Equation (1.3) implies that circuit algebras are equivalent to modular operads ($\mathbb{D}\mathbb{T}$ -algebras) in the monoid subcategory of \mathbf{GS} . Moreover, algebras for the monad $\mathbb{L}\mathbb{T}$ are nonunital circuit algebras, whose underlying modular operadic multiplication is not required to have a multiplicative unit (Equation (5.14) and Equation (7.4)). By Section 5, these are symmetric monoidal functors from subcategories of “downward” Brauer diagrams [44] (see also [42] and [32, 47]). In particular, these structures have applications to the representation theory of the infinite dimensional orthogonal, symplectic and general linear groups (see [44] and Sections 3.3 & 6.2 of [42]). These relationships are explored in the companion paper [42]. A partial summary is provided in [42, Table 1].

This paper aligns with a larger body of recent works aimed at describing the underlying combinatorics of operad-like structures, such as cyclic and modular operads [17, 20, 23, 24, 29, 41, 47, 50], properads and props [22, 30, 46]. A key motivation for much of this research is to find models that characterise generalised operads and their algebras *up-to-homotopy*. The present work is no exception.

Recently, Dancso, Halacheva and Robertson have used circuit algebras to relate the graded Kashiwara-Vergne and graded Grothendieck-Teichmüller groups, KRV and GRT [14]. However, associativity of the circuit algebra operations is not preserved under passage to pro-unipotent completion and, to relate the ungraded groups GT and KV directly, it is necessary to weaken the circuit algebra axioms (see [14, Remark 1.1]).

Equation (1.2) characterises circuit algebras via a *strong* Segal condition on graphical presheaves. So, it is natural to define homotopy circuit algebras as (simplicial set valued) presheaves that satisfy a weakened Segal condition. Indeed, [24] and [41] described models for homotopy modular operads – using different methods and slightly different graphical categories – as weak Segal presheaves. Unfortunately, neither of these results can be generalised straightforwardly to obtain a model structure on $\mathbf{psh}(\Xi^\times)$. It is for future work to explore whether such a model structure exists, or if the weak Segal condition is sufficient to describe a practical homotopy theory for circuit algebras.

The nonunital case, which avoids loops, is better understood: Stoeckl [46] has constructed a model for nonunital homotopy wheeled props, building on Ward’s proof that the operad for nonunital modular operads is Koszul [50], and Kaufmann and Ward [29] have shown that the operad governing (nonoriented) nonunital circuit algebras (there called “Schwarz modular operads”) is Koszul. It would be interesting to compare the model for nonunital homotopy circuit algebras so-obtained with the Segal characterisation suggested by the nonunital case of Equation (1.2). It is not expected that these results can be extended to unital circuit algebras. However, the iterated distributive law of Equation (1.3) together with Markl’s notion of distributive laws for operads in [37] – used in [29] to prove that the (monochrome) nonunital circuit algebra operad is Koszul – gives a possible approach for exploring Koszulity of the operad for unital circuit algebras. (See also Equation (7.12) and [42, Introduction].) Batanin and Berger’s theory of *tame polynomial monads* [4], together with the distributive law decomposition of the circuit algebra monad, could also shed light on the homotopy theory.

Finally, a word on potential applications of this work outside mathematics. My own interest in circuit algebras, modular operads, and other operadic structures governed by graphs was originally motivated by questions involving complex networks with structure at multiple scales: Given a circuit algebra A , the bar construction for \mathbb{LDT} and A may be viewed as a rulebook for *zooming in and out* of networks decorated by A . Indeed, there is an increasing body of work exploring this type of approach to applications in fields such as systems engineering, dynamical systems, software engineering and neuroscience (see e.g., [40]). Just as they provide the main difficulty for constructing the circuit algebra monad, cycles, or feedback loops, are the drivers of complexity in many real world networks (see e.g., [18, 19]). The approach of this work (and [41]) highlights the combinatorial structure of loops and may therefore shed light on the properties of feedback loops as they appear in diverse applications.

1.1. Details and overview. Section 2 provides a relatively informal introduction to iterated distributive law and the abstract nerve machinery of [6], on which the proofs of Theorems 1.1-1.3 are built, as well as some technical remarks on the relationship of this work to the extended algebraic patterns of [11] and extensions of strongly cartesian monads described in [6, Proposition 4.4]. Most of this section should be accessible to readers with just a basic knowledge of category theory.

In Section 3, circuit algebras and modular operads are introduced as V -valued presheaves over \mathbf{GS} that satisfy certain axioms. It is important to note that the companion paper [42] and e.g., [3, 14, 23–

25] are concerned with circuit algebras (and modular operads) “enriched” in a symmetric monoidal category (\mathbf{X}, \otimes, I) . From this perspective, different flavours of enriched circuit algebras are described as algebras over different operads. By contrast, this paper gives constructions “internal” to a category \mathbf{V} with sufficient (co)limits. The sets of colours (or generating objects) in the enriched version are replaced by “object objects” (palettes) in the internal version. This enables all (internal) circuit algebras to be obtained as algebras for a single monad. Of course, the two definitions are equivalent when $\mathbf{X} = \mathbf{V}$ is the category of sets.

The technical work of the paper starts at Section 4 which is largely a review of basic graph constructions, using the formalism of Joyal and Kock’s “(Feynman) graphs” [27]. (See also [23, 24, 41]).

A monad \mathbb{T}^\times for nonunital circuit algebras is constructed in Section 5. This is almost identical to the monad discussed by Kock in [31] and very similar to the nonunital modular operad monad \mathbb{T} constructed in [41]. Section 6 reviews the construction in [41] of the modular operad monad \mathbb{DT} by composing \mathbb{T} with a unit monad \mathbb{D} , and gives an explicit description of the monad \mathbb{T}_* , that extends \mathbb{T} to the EM category of algebras for \mathbb{D} , \mathbf{GS}_* .

Despite the similarities between \mathbb{T}^\times and \mathbb{T} , the monad for unital circuit algebras cannot be obtained directly from a distributive law involving \mathbb{T}^\times . Instead, it is necessary to supplement the construction of [41] by observing that \mathbb{T}^\times is, itself a composite monad. This happens in Section 7, in which an iterated distributive law is described and it is shown that algebras for the resulting composite \mathbb{LDT} of three monads on \mathbf{GS} are, indeed, equivalent to circuit algebras.

Finally, Section 8 contains a detailed description of the graphical category Ξ^\times for circuit algebras, and this leads easily to the proof of the nerve theorem (Equation (1.2)). There is a “lifted” monad $\tilde{\mathbb{L}}\mathbb{T}_*$ on \mathbf{GS}_* whose algebras are circuit algebras. Equation (1.2) is then implied by the observation that $\tilde{\mathbb{L}}\mathbb{T}_*$ has arities on \mathbf{GS}_* and hence satisfies the conditions of the abstract nerve machinery of [6]. Three different proofs of this theorem are discussed (c.f., Proof of Equation (8.4) and Remarks 8.17 and 8.18).

Acknowledgements. I thank Marcy Robertson for motivating me to write this paper and for her unfailing guidance and support. The ideas in this work were incubated and nourished during my time at the Centre of Australian Category Theory, Macquarie University, Dharug Country. I acknowledge the support, friendship, and mathematical input of my friends and colleagues there. Thanks, in particular, to Steve Lack and John Power. Thanks also to Kurt Stoeckl for great suggestions and to Bek McCann and my other students and colleagues at James Cook University, Bindal Country, where this work was written.

2. DISTRIBUTIVE LAWS AND ABSTRACT NERVE THEORY

The main result of this work is the construction of a monad (Equation (7.9)) and a graphical category (Section 8.3) and nerve theorem (Equation (8.4)) for circuit algebras. Equation (8.4) proved the machinery of abstract nerve theory [6], in combination with iterated distributive laws [5, 10].

Given the quantum topology origins of circuit algebras [3, 14] and their connections with Brauer categories and representations of classical groups [15, 34, 42, 44], this overview is aimed at a diverse audience including topologists, representation theorists and category theorists, and is intended to provide an accessible explanation of these formal concepts and how they interact in the proof of Equation (8.4), as well as to establish notation and terminology for what follows.

More technical details are included as remarks, and I refer those interested in further details on the general theory to [6, 51], for abstract nerve theory, and [5, 10], for (iterated) distributive laws.

2.1. Iterated distributive laws. This paper applies the general philosophy that, to understand something complicated, it is often helpful to break it up into pieces and instead study these simpler components and their local interactions. Monads are categorical devices that often serve to encode algebraic structures. In general, monads do not compose. But, in some cases, it may be possible to study a complicated algebraic structure via interactions between monads for simpler structures.

A monad $\mathbb{M} = (M, \mu^{\mathbb{M}}, \eta^{\mathbb{M}})$ on a category \mathbf{C} is given by an endofunctor $M: \mathbf{C} \rightarrow \mathbf{C}$, together with natural transformations $\mu^{\mathbb{M}}: M^2 \Rightarrow M$ (the *monadic multiplication*) and $\eta^{\mathbb{M}}: \text{id}_{\mathbf{C}} \Rightarrow M$ (the *monadic unit*) that satisfy axioms making \mathbb{M} into an associative monoid in the category of endofunctors on \mathbf{C} . (See, e.g., [35, Chapter VI], for detailed review of the general theory of monads and their algebras.)

An algebra (c, θ) for \mathbb{M} consists of an object c of \mathbf{C} , and a morphism $\theta: Mc \rightarrow c$ satisfying two axioms that assert the compatibility of θ with $\mu^{\mathbb{M}}$ and $\eta^{\mathbb{M}}$. Algebras are the objects of the *Eilenberg-Moore (EM) category* $\mathbf{C}^{\mathbb{M}}$ of algebras for \mathbb{M} , whose morphisms $(c, \theta) \rightarrow (d, \phi)$ are given by morphisms $f \in \mathbf{C}(c, d)$ such that $\phi \circ Mf = f \circ \theta$. The monad \mathbb{M} induces free-forgetful adjoint functors $\mathbf{C} \rightleftarrows \mathbf{C}^{\mathbb{M}}$ where the left adjoint $\mathbf{C} \rightarrow \mathbf{C}^{\mathbb{M}}$ is given by $c \mapsto (Mc, \mu^{\mathbb{M}}(c))$.

Let $\mathbb{M} = (M, \mu^{\mathbb{M}}, \eta^{\mathbb{M}})$ and $\mathbb{M}' = (M', \mu^{\mathbb{M}'}, \eta^{\mathbb{M}'})$ be monads on a category \mathbf{C} . In general, the composite functor MM' does not carry a monad structure, since the data of \mathbb{M} and \mathbb{M}' is not sufficient to define a multiplication $(MM')^2 = MM'MM' \Rightarrow MM'$. Observe, however, that any natural transformation $\lambda: M'M \Rightarrow MM'$ induces a natural transformation $\mu_\lambda \stackrel{\text{def}}{=} (\mu^{\mathbb{M}}\mu^{\mathbb{M}'}) \circ (M\lambda M'): (MM')^2 \Rightarrow MM'$.

Definition 2.1. A distributive law for \mathbb{M} and \mathbb{M}' is a natural transformation $\lambda: M'M \Rightarrow MM'$ such that the triple $(MM', \mu_\lambda, \eta^{\mathbb{M}}\eta^{\mathbb{M}'})$ defines a monad $\mathbb{M}\mathbb{M}'$ on \mathbf{C} .

Distributive laws were introduced by Beck [5] in terms of four axioms encoding their interactions with the monadic multiplications and units.

Let $\lambda: M'M \Rightarrow MM'$ be a distributive law between monads \mathbb{M} and \mathbb{M}' on \mathbf{C} . If (c, θ) is an $\mathbb{M}\mathbb{M}'$ algebra, then c admits an \mathbb{M}' algebra structure $\theta \circ \eta^{\mathbb{M}}M': M'c \rightarrow MM'c \rightarrow c$ and an \mathbb{M} algebra structure $\theta \circ \lambda \circ M\eta^{\mathbb{M}'}: Mc \rightarrow M'Mc \rightarrow MM'c \rightarrow c$. In particular, the induced functor $\mathbf{C}^{\mathbb{M}\mathbb{M}'} \rightarrow \mathbf{C}^{\mathbb{M}'}$ has a left adjoint and the monad \mathbb{M} *lifts* to a monad on $\mathbf{C}^{\mathbb{M}'}$ whose EM category of algebras is isomorphic to $\mathbf{C}^{\mathbb{M}\mathbb{M}'}$ (see [5, Section 3]). Indeed, the existence of a functorial lift of \mathbb{M} to \mathbb{M}' is equivalent to the existence of a distributive law $\lambda: M'M \Rightarrow MM'$ (see e.g. [10]).

In general, the functor $\mathbf{C}^{\mathbb{M}\mathbb{M}'} \rightarrow \mathbf{C}^{\mathbb{M}}$ does not have a left adjoint and so the monad \mathbb{M}' does not extend to a monad on $\mathbf{C}^{\mathbb{M}}$. However, in case the left adjoint does exist (in which case it is described by a certain right Kan extension [16, 48]) and the counit of the adjunction is an isomorphism, then \mathbb{M}' may be extended to a monad \mathbb{M}'_* on $\mathbf{C}^{\mathbb{M}}$ such that $(\mathbf{C}^{\mathbb{M}})^{\mathbb{M}'_*} \cong \mathbf{C}^{\mathbb{M}\mathbb{M}'}$ and the following square of adjunctions commutes:

$$(2.2) \quad \begin{array}{ccc} \mathbf{C}^{\mathbb{M}'} & \xleftarrow{\quad \top \quad} & \mathbf{C}^{\mathbb{M}\mathbb{M}'} \\ \uparrow \scriptstyle \vdash & \xrightarrow{\quad \top \quad} & \uparrow \scriptstyle \vdash \\ \mathbf{C} & \xleftarrow{\quad \top \quad} & \mathbf{C}^{\mathbb{M}} \end{array}$$

Example 2.3. The category of small categories is equivalent to the EM category of algebras for the category monad on the category of directed graphs. This sends a directed graph G to the directed graph \widetilde{G} with the same vertices, and – for each directed path (of length $n \geq 0$) from v to w in G – a directed edge $v \rightsquigarrow w$ in \widetilde{G} .

This monad may be obtained as a composite of the *semi-category monad* – which governs associative composition and assigns to each directed graph G the graph with the same vertices as G and edges

corresponding to non-trivial directed paths (of length $n > 0$) in G – and the *reflexive graph monad* that adjoins a distinguished loop at each vertex of a directed graph. The corresponding distributive law encodes the property that the adjoined loops provide identities for the (semi-categorical) composition.

Example 2.4. Similarly, the monad for modular operads on the category \mathbf{GS} of graphical species (see Equation (3.2)) is obtained in [41], as a composite \mathbb{DT} : the monad \mathbb{T} governs the composition structure (multiplication and contraction), and the monad \mathbb{D} adjoins distinguished elements that encode the combinatorics of the multiplicative unit. The distributive law ensures that the distinguished elements provide units for the multiplication (see also Section 6).

Let $\mathbb{M}_1, \dots, \mathbb{M}_n$ be an n -tuple of monads on a category \mathbf{C} . If there are distributive laws $\lambda_{i,j} : M_i M_j \Rightarrow M_j M_i$ (defined for all pairs $1 \leq i < j \leq n$) such that, for each triple $1 \leq i < j < k \leq n$, the corresponding triple of monads and distributive laws satisfies the so-called *Yang-Baxter* conditions

$$\lambda_{j,k} M_i \circ M_j \lambda_{i,k} \circ \lambda_{i,j} M_k = M_k \lambda_{i,j} \circ \lambda_{i,k} M_j \circ M_i \lambda_{j,k} : M_i M_j M_k \Rightarrow M_k M_j M_i,$$

then, by [10], $\mathbb{M}_1, \dots, \mathbb{M}_n$ may be composed to obtain a monad $\mathbb{M}_n \dots \mathbb{M}_1$ on \mathbf{C} . It follows that, for each $1 \leq i < n$, there are composable monads $\mathbb{M}_n \dots \mathbb{M}_{i+1}$ and $\mathbb{M}_i \dots \mathbb{M}_1$ on \mathbf{C} , and, in particular, that $\mathbb{M}_n \dots \mathbb{M}_{i+1}$ lifts to a monad $\widetilde{\mathbb{M}}_{(i+1)}$ on the EM category $\mathbf{C}^{\mathbb{M}_i \dots \mathbb{M}_{i+1}}$.

2.2. Presheaves, slice categories and nerve functors. Let \mathbf{V} be a category and let \mathbf{C} be an essentially small category. An \mathbf{V} -valued \mathbf{C} -*presheaf* is a functor $S : \mathbf{C}^{\text{op}} \rightarrow \mathbf{V}$. The corresponding functor category is denoted by $\mathbf{psh}_{\mathbf{V}}(\mathbf{C})$. If $\mathbf{V} = \mathbf{Set}$ is the category of sets and all set maps, then we will write $\mathbf{psh}(\mathbf{C}) \stackrel{\text{def}}{=} \mathbf{psh}_{\mathbf{V}}(\mathbf{C})$.

The category of elements $\text{el}_{\mathbf{C}}(P)$ of a \mathbf{Set} -valued presheaf $P : \mathbf{C}^{\text{op}} \rightarrow \mathbf{Set}$ is defined as follows:

Definition 2.5. *Objects of $\text{el}_{\mathbf{C}}(P)$ are pairs (c, x) – called elements of P – where c is an object of \mathbf{C} and $x \in P(c)$. Morphisms $(c, x) \rightarrow (d, y)$ are given by morphisms $f \in \mathbf{C}(c, d)$ such that $P(f)(y) = x$.*

A functor $\iota : \mathbf{D} \rightarrow \mathbf{C}$ induces a presheaf $\iota^* \mathbf{C}(-, c)$, $d \mapsto \mathbf{C}(\iota(d), c)$ on \mathbf{D} . For all $c \in \mathbf{C}$, the category $\text{el}_{\mathbf{D}}(\iota^* \mathbf{C}(-, c))$ is called the *slice category of \mathbf{D} over c* and is denoted by ι/c , or simply \mathbf{D}/c .

The *slice category of \mathbf{D} under c* , denoted by c/\mathbf{D} (or c/ι) is the category ι^{op}/c whose objects are morphisms $f \in \mathbf{C}(c, \iota(-))$ and whose morphisms $(d, f) \rightarrow (d', f')$ are given by $g \in \mathbf{D}(d, d')$ such that $\iota(g) \circ f = f' \in \mathbf{C}(c, \iota(d'))$.

A presheaf $P : \mathbf{C}^{\text{op}} \rightarrow \mathbf{Set}$ is *represented* by some $c \in \mathbf{C}$ if $P = \mathbf{C}(-, c)$. In this case, $\text{el}_{\mathbf{C}}(P)$ is precisely the *slice category \mathbf{C}/c* whose objects are pairs (d, f) where $f \in \mathbf{C}(d, c)$, and morphisms $(d, f) \rightarrow (d', f')$ are commuting triangles in \mathbf{C} :

$$\begin{array}{ccc} d & \xrightarrow{g} & d' \\ & \searrow f & \nearrow f' \\ & c & \end{array}$$

In particular, the *Yoneda embedding* $\mathbf{C} \rightarrow \mathbf{psh}(\mathbf{C}), c \mapsto \mathbf{C}(-, c)$ induces a canonical isomorphism $\text{el}_{\mathbf{C}}(P) \cong \mathbf{C}/P$ for all presheaves $P : \mathbf{C}^{\text{op}} \rightarrow \mathbf{Set}$. These categories will be identified in this work.

Definition 2.6. *The nerve functor $N = N(F) : \mathbf{C} \rightarrow \mathbf{psh}(\mathbf{D})$ associated to a functor $F : \mathbf{D} \rightarrow \mathbf{C}$ is given by $N(c)(d) = \mathbf{C}(Fd, c)$ for all $c \in \mathbf{C}$ and $d \in \mathbf{D}$.*

Let $\iota : \mathbf{D} \hookrightarrow \mathbf{C}$ be an inclusion of small categories. By [33, Proposition 5.1], the induced nerve $N : \mathbf{C} \rightarrow \mathbf{psh}(\mathbf{D})$ is fully faithful if and only if every object $c \in \mathbf{C}$ is canonically the colimit of the forgetful functor $\mathbf{D}/c \rightarrow \mathbf{C}$, $(f : \iota(d) \rightarrow c) \mapsto \iota(d)$. In this case, the inclusion $\iota : \mathbf{D} \hookrightarrow \mathbf{C}$ is called *dense*.

Example 2.7. The nerve of a small category \mathbf{C} is described by the image of \mathbf{C} under the nerve functor induced by the dense inclusion $\Delta \hookrightarrow \mathbf{Cat}$, where \mathbf{Cat} is the category of small categories, and Δ is the *simplex category* of non-empty finite ordinals and order preserving morphisms.

2.3. Weber’s abstract nerve theory. Every functor admits an (up to isomorphism) unique *bo-ff factorisation* as a bijective on objects functor followed by a fully faithful functor. For example, if $\mathbb{M} = (M, \mu, \eta)$ is a monad on a category \mathbf{C} , and $\mathbf{C}^{\mathbb{M}}$ is the EM category of algebras for \mathbb{M} , then the free functor $\mathbf{C} \rightarrow \mathbf{C}^{\mathbb{M}}$ has bo-ff factorisation $\mathbf{C} \rightarrow \mathbf{C}_{\mathbb{M}} \rightarrow \mathbf{C}^{\mathbb{M}}$, where $\mathbf{C}_{\mathbb{M}}$ is the Kleisli category whose objects are the *free \mathbb{M} -algebras* of the form $(Mc, \mu^{\mathbb{M}}(c))$ for $c \in \mathbf{C}$. See e.g. [35, Section VI.5].

Let $\mathbf{D} \subset \mathbf{C}$ be a dense subcategory and $\Theta_{\mathbb{M}, \mathbf{D}}$ the full subcategory of $\mathbf{C}_{\mathbb{M}}$ obtained in the bo-ff factorisation of the functor $\mathbf{D} \hookrightarrow \mathbf{C} \xrightarrow{\text{free}} \mathbf{C}^{\mathbb{M}}$. Consider the following diagram:

$$(2.8) \quad \begin{array}{ccccc} \Theta_{\mathbb{M}, \mathbf{D}} & \xrightarrow{\text{f.f.}} & \mathbf{C}^{\mathbb{M}} & \xrightarrow{N_{\mathbb{M}, \mathbf{D}}} & \mathbf{psh}(\Theta_{\mathbb{M}, \mathbf{D}}) \\ \text{b.o.} \uparrow j & & \uparrow \text{free} & \downarrow \text{forget} & \downarrow j^* \\ \mathbf{D} & \hookrightarrow & \mathbf{C} & \xrightarrow[N_{\mathbf{D}}]{\text{f.f.}} & \mathbf{psh}(\mathbf{D}). \end{array}$$

where j^* is the pullback of the bijective on objects functor $j: \mathbf{D} \rightarrow \Theta_{\mathbb{M}, \mathbf{D}}$. The left square of (2.8) commutes by definition, and the right square commutes up to natural isomorphism (c.f., [6, Section 1.6]).

If the fully faithful functor $\Theta_{\mathbb{M}, \mathbf{D}} \rightarrow \mathbf{C}^{\mathbb{M}}$ is also dense, the induced nerve $N_{\mathbb{M}, \mathbf{D}}: \mathbf{C}^{\mathbb{M}} \rightarrow \mathbf{psh}(\Theta_{\mathbb{M}, \mathbf{D}})$ is fully faithful. Moreover, if \mathbf{D} and \mathbb{M} satisfy certain conditions as described in [6, 51] – in which case \mathbf{D} is said to *provide arities for \mathbb{M}* – then, by [51, Section 4], the image of $N_{\mathbb{M}, \mathbf{D}}$ in $\mathbf{psh}(\Theta_{\mathbb{M}, \mathbf{D}})$ consists of precisely those objects that are mapped, up to equivalence, by j^* to the image of $N_{\mathbf{D}}$ in $\mathbf{psh}(\mathbf{D})$. This is the *Segal condition* for the nerve theorem.

More generally, if $\mathbf{D} \subset \mathbf{C}$ is a dense subcategory and \mathbb{M} is a monad on \mathbf{C} such that the induced nerve $N_{\mathbb{M}, \mathbf{D}}$ in (2.8) is fully faithful and the Segal condition is satisfied, then \mathbb{M} is said to be *D-nervous* [7]. Having arities is a sufficient, but not necessary condition for a monad to be nervous. Indeed, neither the modular operad monad of [41] nor the circuit algebra monad constructed in this work (Section 7.2) have arities, but they are both nervous (c.f., Section 8 and [41, Section 8]).

Remark 2.9. In many cases of interest, $\mathbf{C} = \mathbf{psh}(\mathcal{E})$ is the category of presheaves on some category \mathcal{E} . As the Yoneda embedding $\mathcal{E} \rightarrow \mathbf{C}$ is fully faithful and dense, any full subcategory $\mathbf{D} \subset \mathbf{C}$ that contains the representables is also dense in \mathbf{C} (and $\mathcal{E} \hookrightarrow \mathbf{D}$ embeds as a full dense subcategory) so that the induced nerve $\mathbf{C} \rightarrow \mathbf{psh}(\mathbf{D})$ is fully faithful. Bourke and Garner [7] call this situation *the presheaf context*.

If $\mathbf{C} = \mathbf{psh}(\mathcal{E})$ and \mathbb{M} is a D-nervous monad on \mathbf{C} , then the Segal condition has a particularly nice form: A presheaf P on $\Theta_{\mathbb{M}, \mathbf{D}}$ is equivalent to the nerve of an \mathbb{M} -algebra if and only if, for all $d \in \mathbf{D}$,

$$(2.10) \quad P(jd) \cong \lim_{(x, f) \in \mathcal{E}/d} j^* P(x).$$

Both the original Segal condition for the simplicial nerve theorem for small categories, and a Segal condition for the dendroidal nerve of an operad, may be obtained from monads with arities in the presheaf context (see e.g., [51, Section 4] or [27, Section 2] for details). The nerve theorems for modular operads [41, Theorem 8.2] and circuit algebras (Equation (8.4)) are also described in the presheaf context, though the corresponding monads on GS do not have arities in these cases.

Remark 2.11. (See also Remarks 2.9, 8.17 and 8.18.) A special class of monad with arities is described by *strongly cartesian monads* [6] (called *parametric right adjoint monads* in [51] and *polynomial monads* in [11]). These are *cartesian* monads (they satisfy certain pullback conditions) that are *locally*

right adjoint (certain functors of slice categories are right adjoints). For a strongly cartesian monad \mathbb{M} on \mathbf{C} , there is a canonical choice of arities $\mathbf{D}_{\mathbb{M}} \subset \mathbf{C}$ (see e.g., [6, section 2]). In the presheaf context, strongly cartesian monads and their associated Segal conditions, are central to the theory of *algebraic patterns* developed in [11].

Consider the diagram (2.8) for the case that \mathbb{M} is a strongly cartesian monad on $\mathbf{C} = \mathbf{psh}(\mathcal{E})$. Denote the canonical arities for \mathbb{M} by $\mathbf{D} \subset \mathbf{C}$. So, $\Theta_{\mathbb{M}, \mathbf{D}}$ is dense in $\mathbf{C}^{\mathbb{M}}$, the induced nerve is fully faithful, and $P \in \mathbf{psh}(\Theta_{\mathbb{M}, \mathbf{D}})$ is, up to equivalence, in the image of the nerve $N_{\mathbb{M}, \mathbf{D}}$ if and only if P satisfies (2.10). Since \mathbb{M} is strongly cartesian, there is, by [51, Proposition 2.6], a wide subcategory $\Theta_{\mathbb{M}, \mathbf{D}}^{\text{ref}} \subset \Theta_{\mathbb{M}, \mathbf{D}}$ of so-called *generic morphisms* or *refinements* [27] in $\Theta_{\mathbb{M}, \mathbf{D}}$ such that every morphism in $\Theta_{\mathbb{M}, \mathbf{D}}$ factors as a refinement followed by a morphism in the image of \mathbf{D} .

The strongly cartesian monad \mathbb{M} on $\mathbf{C} = \mathbf{psh}(\mathcal{E})$ then has the form

$$(2.12) \quad M(P)(x) \cong \text{colim}_{(y, f) \in x / \Theta_{\mathbb{M}, \mathbf{D}}^{\text{ref}}} \lim_{(e, \alpha) \in \mathcal{E} / \mathbf{D} y} P(e), \quad \text{for all } P \in \mathbf{C}, x \in \mathcal{E}.$$

In the terminology of [11], the factorisation system on $\Theta_{\mathbb{M}, \mathbf{D}}$, together with the underlying category \mathcal{E} , then describes an *extendable algebraic pattern* with corresponding *free Segal $\Theta_{\mathbb{M}, \mathbf{D}}$ -space monad* \mathbb{M} .

2.4. Applying distributive laws to prove nerve theorems. The proof of the nerve theorem Equation (8.4) for circuit algebras, like the proof of the modular operad nerve theorem of [41], exploits the distributive law decompositions of the corresponding composite monad.

Let \mathbb{M} be a monad on an essentially small category \mathbf{C} with subcategory $\mathbf{D} \subset \mathbf{C}$, and let $\Theta_{\mathbb{M}, \mathbf{D}} \subset \mathbf{C}^{\mathbb{M}}$ be the subcategory obtained in the bo-ff factorisation of $\mathbf{D} \hookrightarrow \mathbf{C} \rightarrow \mathbf{C}^{\mathbb{M}}$. Even if \mathbf{D} itself does not provide arities for \mathbb{M} , if $\mathbb{M} = \mathbb{M}_n \dots \mathbb{M}_1$ is a composite monad, then the decomposition of \mathbb{M} may provide other opportunities to apply the results of [6] in order to prove that the induced nerve $N_{\mathbb{M}, \mathbf{D}}: \mathbf{C}^{\mathbb{M}} \rightarrow \mathbf{psh}(\Theta_{\mathbb{M}, \mathbf{D}})$ is fully faithful:

Example 2.13. In the simplest case, let $\mathbb{M} = \mathbb{M}_2 \mathbb{M}_1$ be a composite monad on \mathbf{C} induced by a distributive law $\lambda: M_1 M_2 \Rightarrow M_2 M_1$, and let $\widetilde{\mathbb{M}}_2$ be the lift of \mathbb{M}_2 to $\mathbf{C}^{\mathbb{M}_1}$. Let $\mathbf{D} \subset \mathbf{C}$ be a full subcategory, $\Theta_{\mathbb{M}, \mathbf{D}}$ be the category obtained in the bo-ff factorisation of the functor $\mathbf{D} \rightarrow \mathbf{C}^{\mathbb{M}}$ and let $\mathbf{D}_{\mathbb{M}_1} \subset \mathbf{C}^{\mathbb{M}_1}$ be the full subcategory such that the bijective on objects functor $j: \mathbf{D} \rightarrow \Theta_{\mathbb{M}, \mathbf{D}}$ factors through $\mathbf{D}_{\mathbb{M}_1}$. This is illustrated in Diagram (2.14), with $\mathbb{M}' = \mathbb{M}_1$ and the free and forgetful functors $\mathbf{C}^{\mathbb{M}'} = \mathbf{C}^{\mathbb{M}_1} \rightleftarrows \mathbf{C}^{\mathbb{M}}$ corresponding to the monadic adjunction associated to the lift $\widetilde{\mathbb{M}}_2$ on $\mathbf{C}^{\mathbb{M}_1}$. The functor $\mathbf{D}_{\mathbb{M}_1} \rightarrow \mathbf{C}^{\mathbb{M}}$ induced by the lifted monad $\widetilde{\mathbb{M}}_2$ also has bo-ff factorisation through $\Theta_{\mathbb{M}, \mathbf{D}}$. So, if $\mathbf{D}_{\mathbb{M}_1}$ provides arities for $\widetilde{\mathbb{M}}_2$, then the nerve $N_{\mathbb{M}, \mathbf{D}}$ is fully faithful.

$$(2.14) \quad \begin{array}{ccccc} \Theta_{\mathbb{M}, \mathbf{D}} & \xrightarrow{\text{f.f.}} & \mathbf{C}^{\mathbb{M}} & \xrightarrow{N_{\mathbb{M}, \mathbf{D}}} & \mathbf{psh}(\Theta_{\mathbb{M}, \mathbf{D}}) \\ \uparrow \text{b.o.} & & \updownarrow \begin{array}{l} \text{free} \\ \text{forget} \end{array} & & \downarrow \\ \mathbf{D}_{\mathbb{M}'} & \xrightarrow{\text{f.f.}} & \mathbf{C}^{\mathbb{M}'} & \xrightarrow{N_{\mathbb{M}', \mathbf{D}}} & \mathbf{psh}(\mathbf{D}_{\mathbb{M}'}) \\ \uparrow \text{b.o.} & & \updownarrow \begin{array}{l} \text{free} \\ \text{forget} \end{array} & & \downarrow \\ \mathbf{D} & \xrightarrow{\text{f.f.}} & \mathbf{C} & \xrightarrow{N_{\mathbf{D}}} & \mathbf{psh}(\mathbf{D}). \end{array}$$

More generally, let $\mathbb{M} = \mathbb{M}_n \dots \mathbb{M}_1$ be a composite monad on \mathbf{C} and let $\mathbf{D} \subset \mathbf{C}$ be a full subcategory. For each $1 \leq i < n$, the monads $\mathbb{M}_n \dots \mathbb{M}_{i+1}$ and $\mathbb{M}_i \dots \mathbb{M}_1$ on \mathbf{C} are composable. In particular, $\mathbb{M}_n \dots \mathbb{M}_{i+1}$ lifts to a monad $\widetilde{\mathbb{M}}_{(i+1)}$ on the EM category $\mathbf{C}^{\mathbb{M}_i \dots \mathbb{M}_{i+1}}$.

There is a factorisation of the diagram (2.8) as

$$(2.15) \quad \begin{array}{ccccc} \Theta_{\mathbb{M},D} & \xrightarrow{\text{f.f.}} & \mathbb{C}^{\mathbb{M}} & \xrightarrow{N_{\mathbb{M},D}} & \text{psh}(\Theta_{\mathbb{M},D}) \\ \parallel & & \parallel & & \parallel \\ \Theta_{\mathbb{M}_n \dots \mathbb{M}_1, D} & \xrightarrow{\text{f.f.}} & \mathbb{C}^{\mathbb{M}_n \dots \mathbb{M}_1} & \xrightarrow{N_{\mathbb{M}_n \dots \mathbb{M}_1, D}} & \text{psh}(\Theta_{\mathbb{M}_n \dots \mathbb{M}_1, D}) \\ \uparrow \text{b.o.} \quad j_n & & \uparrow \text{free} \quad \downarrow \text{forget} & & \downarrow j_n^* \\ \vdots & & \vdots & & \vdots \\ \Theta_{\mathbb{M}_1, D} & \xrightarrow{\text{f.f.}} & \mathbb{C}^{\mathbb{M}_1} & \xrightarrow{N_{\mathbb{M}_1, D}} & \text{psh}(\Theta_{\mathbb{M}_1, D}) \\ \uparrow \text{b.o.} \quad j_1 & & \uparrow \text{free} \quad \downarrow \text{forget} & & \downarrow j_1^* \\ D & \xrightarrow{\text{f.f.}} & \mathbb{C} & \xrightarrow{N_D} & \text{psh}(D). \end{array}$$

The induced nerve $N_{\mathbb{M},D}$ is fully faithful if, for some $1 \leq i < n$, the lift of $\mathbb{M}_n \dots \mathbb{M}_{(i+1)}$ to $\mathbb{C}^{\mathbb{M}_i \dots \mathbb{M}_1}$ has arities $\Theta_{\mathbb{M}_i \dots \mathbb{M}_1, D}$ (as illustrated in (2.15)).

Example 2.16. (See also Equation (2.4).) Once again, let $\mathbb{M} = \mathbb{M}_2 \mathbb{M}_1$ be a composite monad on \mathbb{C} induced by a distributive law $\lambda: M_1 M_2 \Rightarrow M_2 M_1$. If the monad \mathbb{M}_1 extends to a monad \mathbb{M}_{1*} on $\mathbb{C}^{\mathbb{M}_2}$ such that $(\mathbb{C}^{\mathbb{M}_2})^{\mathbb{M}_{1*}} \cong \mathbb{C}^{\mathbb{M}_2 \mathbb{M}_1}$, then, by letting $\mathbb{M}' = \mathbb{M}_2$ in Diagram (2.14), the same technique as in Equation (2.13) may be applied.

For example, in [41, Theorem 7.46], the modular operad monad on \mathbf{GS} is obtained, via a distributive law, as a composite \mathbb{DT} . The proof of the nerve theorem and Segal condition ([41, Theorem 8.2]) follows from the fact that the monad \mathbb{T} extends to a monad \mathbb{T}_* , on the category \mathbf{GS}_* of \mathbb{D} -algebras, whose EM category of algebras coincides with the EM category of algebras for \mathbb{DT} on \mathbf{GS} . Whilst \mathbb{DT} does not have arities, \mathbb{T}_* does.

In Section 8, the proof of Equation (8.4) extends the proof of the modular operadic nerve theorem (Equation (2.16)) but requires a composite \mathbb{LDT} , obtained via iterated distributive laws, of three monads on \mathbf{GS} , and uses both lifts and extensions of monads. The general case is outlined in the following example.

Example 2.17. Examples 2.13 and 2.16 may be combined: Let $\mathbb{M} = \mathbb{M}_3 \mathbb{M}_2 \mathbb{M}_1$ be a monad on \mathbb{C} and let $D \subset \mathbb{C}$ be a full and dense subcategory. Assume that \mathbb{M}_1 extends to a monad \mathbb{M}_{1*} on the EM category $\mathbb{C}^{\mathbb{M}_2}$ of algebras for \mathbb{M}_2 and that $\mathbb{C}^{\mathbb{M}_2 \mathbb{M}_1} \cong (\mathbb{C}^{\mathbb{M}_2})^{\mathbb{M}_{1*}}$. By the theory of iterated distributive laws [10], \mathbb{M}_3 lifts functorially to both $\mathbb{C}^{\mathbb{M}_1}$ and $\mathbb{C}^{\mathbb{M}_2 \mathbb{M}_1}$. Hence, since $\mathbb{C}^{\mathbb{M}_2 \mathbb{M}_1} \cong (\mathbb{C}^{\mathbb{M}_2})^{\mathbb{M}_{1*}}$, the distributive law $M_1 M_3 \Rightarrow M_3 M_1$ between the monads \mathbb{M}_1 and \mathbb{M}_3 on \mathbb{C} lifts to a distributive law between the extension \mathbb{M}_{1*} of \mathbb{M}_1 , and the lift $\tilde{\mathbb{M}}_3$ of \mathbb{M}_3 to $\mathbb{C}^{\mathbb{M}_2}$. Moreover, algebras for $\tilde{\mathbb{M}}_3 \mathbb{M}_{1*}$ on $\mathbb{C}^{\mathbb{M}_2}$ correspond to algebras for \mathbb{M} on \mathbb{C} .

It follows that the nerve induced by the bo-ff factorisation $D \rightarrow \Theta_{\mathbb{M},D} \rightarrow \mathbb{C}^{\mathbb{M}}$ is fully faithful if $\tilde{\mathbb{M}}_3 \mathbb{M}_{1*}$ has arities obtained from the full subcategory $D_{\mathbb{M}_2} \subset \mathbb{C}^{\mathbb{M}_2}$ with the same objects as D (as in Diagram (2.14) with $\mathbb{M}' = \mathbb{M}_2$).

Remark 2.18. It is important to note that the conditions in each of the Examples 2.13-2.17 are not sufficient to establish a Segal condition in terms of presheaves on the category D . The Segal condition in [41, Theorem 8.2] and Equation (8.4), follows from the observation that the embedding $\mathbf{B}^{\mathfrak{s}} \hookrightarrow \mathbf{Gr}_*$ (described in Section 6.3) is dense.

Following Remarks 2.9 and 2.11, I conclude this section with some remarks on related work, specific to the presheaf context. These are used in Section 8, Remarks 8.17 and 8.18, to outline additional proofs of the nerve theorem 8.4.

Remark 2.19. Let $k: \mathcal{E}_1 \rightarrow \mathcal{E}_2$ be a bijective on objects functor and, for $i = 1, 2$, let \mathbb{M}_i be a monad on $\mathcal{C}_i = \mathbf{psh}(\mathcal{E}_i)$ such that the pullback $k^*: \mathcal{C}_2 \rightarrow \mathcal{C}_1$ describes a monad morphism $(\mathcal{C}_1, \mathbb{M}_1) \rightarrow (\mathcal{C}_1, \mathbb{M}_1)$ by $k^*\mathbb{M}_2 = \mathbb{M}_1 k^*$. By [6, Proposition 4.4], if \mathbb{M}_1 is strongly cartesian, then so is \mathbb{M}_2 and the canonical arities for \mathbb{M}_2 may be obtained from those of \mathbb{M}_2 by extension along k .

In the case of the circuit algebra and modular operad monads \mathbb{LDT} and \mathbb{DT} on \mathbf{GS} , the forgetful functor $\mathbf{GS}_* \rightarrow \mathbf{GS}$ associated to the monad \mathbb{D} on \mathbf{GS} is the pullback of the bijective on objects inclusion $k: \mathbf{B}^{\mathfrak{s}} \hookrightarrow \mathbf{B}_*^{\mathfrak{s}}$ described in Section 6.2. The corresponding nerve theorems then follow, as in Equation (8.18), by showing that the monads \mathbb{LT} and \mathbb{T} are strongly cartesian on \mathbf{GS} and admit extension, along k , to \mathbf{GS}_* .

Remark 2.20. Recall Remarks 2.9 and 2.11. Let $\mathcal{C} = \mathbf{psh}(\mathcal{E})$ and \mathcal{D} be a full subcategory containing the representables. Further, let \mathbb{M} be a \mathcal{D} -nervous monad on \mathcal{C} . So, the nerve $N_{\mathbb{M}, \mathcal{D}}: \mathcal{C}^{\mathbb{M}} \rightarrow \mathbf{psh}(\Theta_{\mathbb{M}, \mathcal{D}})$ in Diagram (2.8) is fully faithful and presheaves in its essential image satisfy the presheaf Segal condition (2.10). Assume also that \mathcal{D} is the smallest full subcategory of \mathcal{C} with respect to which \mathbb{M} is nervous.

Now, let $\Theta_{\mathbb{M}, \mathcal{D}}$ admit an algebraic pattern structure – with elementary objects in \mathcal{E} – as in [11]. This means that there is a factorisation system $(\Theta_{\mathbb{M}, \mathcal{D}}^{\text{ref}}, D')$ on $\Theta_{\mathbb{M}, \mathcal{D}}$ such that presheaves in the essential image of $N_{\mathbb{M}, \mathcal{D}}$ must also satisfy a second Segal condition

$$(2.21) \quad P(j'd) \cong \lim_{(x, f') \in \mathcal{E}'/d} j'^* P(x),$$

where $j': D' \hookrightarrow \Theta_{\mathbb{M}, \mathcal{D}}$ is the bijective on objects inclusion and $\mathcal{E}' \subset D'$ is the full subcategory on the objects from \mathcal{E} . Moreover, if the factorisation of $\Theta_{\mathbb{M}, \mathcal{D}}$ describes an extendable algebraic pattern as in [11, Proposition 8.8], then there is a strongly cartesian monad \mathbb{M}' on $\mathcal{C}' \stackrel{\text{def}}{=} \mathbf{psh}(\mathcal{E}')$, given by

$$M'(P)(x) \cong \text{colim}_{(y, f) \in x / \Theta_{\mathbb{M}, \mathcal{D}}^{\text{ref}}} \lim_{(e, \alpha') \in \mathcal{E}' / y} P(e), \quad \text{for all } P \in \mathcal{C}, x \in \mathcal{E},$$

whose EM category of algebras is precisely $\mathcal{C}^{\mathbb{M}}$.

This is summarised in the following diagram, where – as before (2.8) – the left squares commute and the right squares commute up to natural isomorphism:

$$(2.22) \quad \begin{array}{ccccccc} \mathcal{E}' & \xrightarrow[\text{dense}]{\text{f.f.}} & D' \subset \Theta_{\mathbb{M}, \mathcal{D}} & \xrightarrow[\text{dense}]{\text{f.f.}} & \mathcal{C}' & \xrightarrow[\text{f.f.}]{N_{D'}} & \mathbf{psh}(D') \\ & \searrow \text{b.o.} j' & \downarrow j' & \searrow \text{free}^{\mathbb{M}'} & \uparrow \text{forget}^{\mathbb{M}'} & \uparrow j'^* & \\ & & \Theta_{\mathbb{M}, \mathcal{D}} & \xrightarrow[\text{f.f.}]{} & \mathcal{C}^{\mathbb{M}} & \xrightarrow[\text{f.f.}]{N_{\mathbb{M}, \mathcal{D}}} & \mathbf{psh}(\Theta_{\mathbb{M}, \mathcal{D}}) \\ & \swarrow \text{b.o.} j & \uparrow j & \swarrow \text{free}^{\mathbb{M}} & \downarrow \text{forget}^{\mathbb{M}} & \downarrow j^* & \\ \mathcal{E} & \xrightarrow[\text{dense}]{\text{f.f.}} & \mathcal{D} \subset \mathcal{C} & \xrightarrow[\text{dense}]{\text{f.f.}} & \mathcal{C} & \xrightarrow[\text{f.f.}]{N_{\mathcal{D}}} & \mathbf{psh}(\mathcal{D}). \end{array}$$

For example, the situation of Equation (2.19) is obtained if there is a functor $i: \mathcal{D} \rightarrow D'$ such that $j' \circ i = j$, and \mathbb{M}' is the extension – along the restricted pullback $i^*: \mathcal{C}' \rightarrow \mathcal{C}$ – of a strongly cartesian monad \mathbb{M}_0 on \mathcal{C} .

It is natural to ask the conditions under which such a functor $i: \mathcal{D} \rightarrow D'$ exists, and, furthermore whether it is possible to classify the nervous monads \mathbb{M} for which (as with the circuit algebra and modular operad nerve constructions of Equation (8.4) and [41, Theorem 8.2]) there exists an iterated distributive law describing \mathbb{M} from which the strongly cartesian monad \mathbb{M}' on \mathcal{C}' may be obtained as an extension (and/or lift) as in Examples 2.13-2.17.

3. GRAPHICAL SPECIES, MODULAR OPERADS, AND CIRCUIT ALGEBRAS

In [42], circuit algebras and modular operads enriched in a symmetric monoidal category (\mathbf{X}, \otimes, I) were characterised axiomatically, as symmetric graded objects in \mathbf{X} equipped with operations satisfying certain axioms (see [42, Proposition 5.4] and [42, Definition 5.7]).

This section considers structures that satisfy the same axioms, but now defined *internally* to a category \mathbf{V} with sufficient (co)limits. When $\mathbf{X} = \mathbf{V} = \mathbf{Set}$, the enriched and internal definitions coincide.

3.1. Symmetric species. Let \mathbf{Set}_f denote the category of finite sets and functions and let $\mathbf{B} \subset \mathbf{Set}_f$ be the groupoid of finite sets and bijections. For each $n \in \mathbb{N}$, let \mathbf{n} denote the set $\{1, 2, \dots, n\}$ (so $\mathbf{0} = \emptyset$), and let Σ_n denote the group of permutations on \mathbf{n} . Let $\Sigma = \bigoplus_n \Sigma_n$ be the *symmetric groupoid* (with $\Sigma(n, n) = \Sigma_n$).

As Σ is a skeleton for \mathbf{B} , for any category \mathbf{V} , a \mathbf{V} -valued presheaf $P: \mathbf{B}^{\text{op}} \rightarrow \mathbf{V}$, also called a (*monochrome* or *single-sorted*) *species* [28] in \mathbf{V} , determines a presheaf $\Sigma^{\text{op}} \rightarrow \mathbf{V}$ by restriction. And conversely, a Σ -presheaf Q may always be extended to a \mathbf{B} -presheaf $Q_{\mathbf{B}}$, by setting

$$(3.1) \quad Q_{\mathbf{B}}(X) \stackrel{\text{def}}{=} \lim_{(\mathbf{n}, f) \in \Sigma/X} Q(n) \quad \text{for all } n \in \mathbb{N}.$$

Hence, a species $P: \mathbf{B}^{\text{op}} \rightarrow \mathbf{V}$ is equivalently a sequence $(P(n))_n$ of objects of \mathbf{V} such that Σ_n acts on $P(n)$ for all n .

For a set \mathfrak{D} , let $\text{list}(\mathfrak{D}) = \coprod_{n \in \mathbb{N}} \mathfrak{D}^n$ be the set of finite ordered sets $\mathbf{c} = (c_1, \dots, c_n)$ of elements of \mathfrak{D} that underlies the free associative monoid on \mathfrak{D} with (concatenation) product $\mathbf{cd} = \mathbf{c} \oplus \mathbf{d}$ given by

$$\mathbf{cd} \stackrel{\text{def}}{=} (c_1, \dots, c_m, d_1, \dots, d_n) \text{ for } \mathbf{c} = (c_1, \dots, c_m), \mathbf{d} = (d_1, \dots, d_n) \text{ in } \text{list}(\mathfrak{D}).$$

The empty list is the unit for \oplus and is denoted by \emptyset (or $\emptyset_{\mathfrak{D}}$).

The symmetric groupoid Σ acts on $\text{list}(\mathfrak{D})$ from the right by $\sigma: (\mathbf{c}\sigma) \stackrel{\text{def}}{=} (c_{\sigma 1}, \dots, c_{\sigma n}) \mapsto \mathbf{c}$, for all $\mathbf{c} = (c_1, \dots, c_n)$ and $\sigma \in \Sigma_n$. The monoidal groupoid so obtained is the *free symmetric groupoid on* \mathfrak{D} denoted by $\Sigma^{\mathfrak{D}}$. In particular, if $\mathfrak{D} = \emptyset$, then $\Sigma^{\mathfrak{D}} = \{\emptyset\}$.

An \mathfrak{D} -coloured species in \mathbf{V} is a \mathbf{V} -valued presheaf on $\Sigma^{\mathfrak{D}}$. This is equivalently described by a collection $(P(\mathbf{c}))_{\mathbf{c} \in \text{list}(\mathfrak{D})}$ of objects of \mathbf{V} such that Σ_n (right) acts on $P(n) = \coprod_{\mathbf{c} \in \mathfrak{D}^n} P(\mathbf{c})$ for all n .

3.2. Graphical species. Graphical species in \mathbf{Set} , introduced in [27] to describe various generalisations of (coloured) operads, generalise the notion of coloured species from Section 3.1 and have been used to describe coloured modular operads (called *compact closed categories* in [27]) in [23, 24, 27, 41]. For more details on graphical species in \mathbf{Set} , particularly regarding the involutive palettes (\mathfrak{C}, ω) that encode their expressive power, the reader is referred to [41, Section 1].

This section provides a short discussion on graphical species taking values in an arbitrary category \mathbf{V} . For convenience, \mathbf{V} will always be assumed to have sufficient limits and colimits. The terminal object of \mathbf{V} is denoted by $*$ and the initial object is denoted by $0 = 0_{\mathbf{V}}$.

As before, denote the groupoid of finite sets and bijections by \mathbf{B} . The category \mathbf{B}^{\S} (called *elGr* in [27]) is obtained from \mathbf{B} by adjoining a distinguished object \S that satisfies

- $\mathbf{B}^{\S}(\S, \S) \cong \mathbb{Z}/2\mathbb{Z}$ with generator τ ;
- for each finite set X and each element $x \in X$, there is a morphism $ch_x \in \mathbf{B}^{\S}(\S, X)$ that “chooses” x , and $\mathbf{B}^{\S}(\S, X) = \{ch_x, ch_x \circ \tau\}_{x \in X} \cong X \amalg X$;
- for all finite sets X and Y , $\mathbf{B}^{\S}(X, Y) = \mathbf{B}(X, Y)$;

- morphisms are equivariant with respect to the action of \mathbf{B} . That is, $ch_{f(x)} = f \circ ch_x \in \mathbf{B}^{\mathfrak{s}}(\mathfrak{s}, Y)$ for all $x \in X$ and all bijections $f: X \xrightarrow{\cong} Y$.

Definition 3.2. A graphical species S in a category \mathbf{V} is a presheaf $S: \mathbf{B}^{\mathfrak{s}\text{op}} \rightarrow \mathbf{V}$. The category of graphical species in \mathbf{V} is denoted by $\mathbf{GS}_{\mathbf{V}} \stackrel{\text{def}}{=} \mathbf{psh}_{\mathbf{V}}(\mathbf{B}^{\mathfrak{s}})$. When $\mathbf{V} = \mathbf{Set}$, we write $\mathbf{GS} \stackrel{\text{def}}{=} \mathbf{GS}_{\mathbf{Set}}$.

So, a graphical species S is described by a \mathbf{V} -valued species $(S_X)_{X \in \mathbf{B}}$, and an object $S_{\mathfrak{s}} \in \mathbf{V}$ with involution $S_{\tau} \in \mathbf{V}(S_{\mathfrak{s}}, S_{\mathfrak{s}})$, together with a family of projections $S(ch_x): S_X \rightarrow S_{\mathfrak{s}}$, defined for all finite sets X , and all $x \in X$, and equivariant with respect to the actions of \mathbf{B} and τ . In the definitions of circuit algebras and modular operads (Sections 3.3 & 3.4), the objects S_X describe the (modular operad) operations, or morphisms, whereas the object $S_{\mathfrak{s}}$ can be thought of as describing the “colours”, or generating the “object object”. As in [41, 42], a pair (\mathfrak{C}, ω) of a set \mathfrak{C} with involution $\omega: \mathfrak{C} \rightarrow \mathfrak{C}$ will be called an *(involutive) palette*.

Example 3.3. The terminal graphical species $K^{\mathbf{V}}$ is the constant (monochrome) graphical species that sends \mathfrak{s} and all finite sets X to the terminal object $*$ in \mathbf{V} .

Definition 3.4. A morphism $\gamma \in \mathbf{GS}_{\mathbf{V}}(S, S')$ is colour- or palette-preserving if its component at \mathfrak{s} is the identity on $(S_{\mathfrak{s}}, S_{\tau})$. A graphical species $S \in \mathbf{GS}_{\mathbf{V}}$ is called *monochrome* if $S_{\mathfrak{s}} = *$.

If (\mathfrak{C}, ω) is an involutive palette in \mathbf{Set} , objects of the subcategory $\mathbf{GS}^{(\mathfrak{C}, \omega)} \subset \mathbf{GS}$ of (\mathfrak{C}, ω) -coloured graphical species in \mathbf{Set} are graphical species $S \in \mathbf{GS}$ such that $(S_{\mathfrak{s}}, S_{\tau}) = (\mathfrak{C}, \omega)$. Morphisms are palette-preserving morphisms.

Definition 3.5. Let $S \in \mathbf{GS}^{(\mathfrak{C}, \omega)}$. For a finite set X , $S_X = \coprod_{\mathbf{c} \in \mathfrak{C}^X} S_{\mathbf{c}}$, where, for each $\mathbf{c} \in \mathfrak{C}^X$, the set $S_{\mathbf{c}}$ is the \mathbf{c} -(coloured) arity of S defined as the fibre above \mathbf{c} of the map $(S(ch_x))_{x \in X}: S_X \rightarrow \mathfrak{C}^X$.

It follows that S defines a \mathfrak{C} -coloured species $S': (\Sigma^{\mathfrak{C}})^{\text{op}} \rightarrow \mathbf{Set}$ by restriction. Conversely, for any involutive palette (\mathfrak{C}, ω) , a \mathfrak{C} -coloured species $P: (\Sigma^{\mathfrak{C}})^{\text{op}} \rightarrow \mathbf{Set}$ extends to a graphical species $\hat{P} \in \mathbf{GS}^{(\mathfrak{C}, \omega)}$ by (3.1).

Example 3.6. For any palette (\mathfrak{C}, ω) , the (\mathfrak{C}, ω) -coloured graphical species $K^{(\mathfrak{C}, \omega)}$ given by $K_{\mathbf{c}}^{(\mathfrak{C}, \omega)} = \{*\}$ for all $\mathbf{c} \in \text{list}(\mathfrak{C})$ is terminal in $\mathbf{GS}^{(\mathfrak{C}, \omega)}$. If $(\mathfrak{C}, \omega) = \emptyset$, then $K_{\mathbf{0}}^{\emptyset} = \{*\}$ and $K_X^{\emptyset} = \emptyset$ for $X \neq \mathbf{0}$.

The element category (Equation (2.5)) of a \mathbf{Set} -valued graphical species $S \in \mathbf{GS}$ is denoted by $\text{el}(S) \stackrel{\text{def}}{=} \text{el}_{\mathbf{B}\mathfrak{s}}(S)$.

Example 3.7. Let $(\mathfrak{Di}, \sigma_{\mathfrak{Di}})$ (called $\{\uparrow, \downarrow\}$ in [42]) be the involutive palette in \mathbf{Set} described by the unique non-trivial involution $\sigma_{\mathfrak{Di}}$ on the two-element set $\mathfrak{Di} = \{\text{in}, \text{out}\}$. This was used in [42, Section 3.2] to describe (monochrome) oriented circuit algebras (monochrome wheeled props), and in [41, Section 1.1] to describe props and wheeled operads.

If Di is the terminal $(\mathfrak{Di}, \sigma_{\mathfrak{Di}})$ -coloured graphical species in \mathbf{Set} , then, for all finite sets X , $Di_X = \mathfrak{Di}^X$ is the set of partitions of $X = X_{\text{in}} \amalg X_{\text{out}}$ into *input* and *output* sets and morphisms $Di_X \rightarrow Di_Y$ are bijections $X \xrightarrow{\cong} Y$ that preserve the partitions. Objects of the category $\mathbf{GS}^{(\mathfrak{Di}, \sigma_{\mathfrak{Di}})}$ are called *monochrome oriented graphical species* in \mathbf{Set} .

Since $(\text{in}) \cong (\text{out})$ in $\text{el}(Di)$, the element category $\text{el}(Di)$ of Di is equivalent to the category $(\mathbf{B} \times \mathbf{B}^{\text{op}})^+$ obtained from $\mathbf{B} \times \mathbf{B}^{\text{op}}$ by adjoining a distinguished object \downarrow (the *directed exceptional edge*), and morphisms $i_x, o_y: (X_{\text{in}}, X_{\text{out}}) \rightarrow (\downarrow)$ defined for each $x \in X_{\text{in}}$ and $y \in X_{\text{out}}$.

Definition 3.8. The category $\mathbf{OGS}_{\mathbf{V}}$ of oriented graphical species in \mathbf{V} is the presheaf category $\mathbf{psh}_{\mathbf{V}}(\text{el}(Di))$.

For all **Set**-valued graphical species $S' \in \mathbf{GS}$, there is a canonical equivalence $\mathbf{GS}/S' \cong \mathbf{psh}(\mathbf{el}(S'))$ such that $\gamma \in \mathbf{GS}(S, S')$ describes a presheaf $\tilde{S} \in \mathbf{psh}(\mathbf{el}(S'))$ by $\tilde{S}(\phi) = \gamma^{-1}(\phi)$ for all $\phi \in \mathbf{el}(S')$.

Example 3.9. (See also [41, Example 1.10].) Assume that \mathbf{V} has finite products and coproducts. Let $Di^{\mathbf{V}} \in \mathbf{GS}_{\mathbf{V}}$ be the graphical species given by $Di^{\mathbf{V}}_{\S} = * \amalg *$ (and $Di^{\mathbf{V}}_{\tau}$ acts by swapping the components), and, for each finite set \mathcal{X} , $Di^{\mathbf{V}}_{\mathcal{X}} = \coprod_{(X_{\text{in}}, X_{\text{out}}) \in Di_{\mathcal{X}}(*)} (*)$.

There is a canonical equivalence of categories $\mathbf{OGS}_{\mathbf{V}} \simeq \mathbf{GS}_{\mathbf{V}}/Di^{\mathbf{V}}$. If $(S, \gamma) \in \mathbf{GS}_{\mathbf{V}}/Di^{\mathbf{V}}$, then the corresponding presheaf $\tilde{S} \in \mathbf{psh}_{\mathbf{V}}(\mathbf{el}(Di)) = \mathbf{OGS}_{\mathbf{V}}$ is such that, for all $\mathcal{C} \in \mathbf{B}^{\S}$ and all $\phi \in S_{\mathcal{C}}$ (so ϕ is either a pair of finite sets or an element of $\{\text{in}, \text{out}\}$), $\tilde{S}(\phi)$ is given by the pullback

$$\begin{array}{ccc} \tilde{S}(\phi) & \xrightarrow{\quad} & S_{\mathcal{C}} \\ \downarrow & & \downarrow \gamma_{\mathcal{C}} \\ * & \xrightarrow{\quad inc_{\phi} \quad} & Di^{\mathbf{V}}_{\mathcal{C}} * \end{array}$$

where inc_{ϕ} is the obvious map that picks out the component indexed by ϕ in $Di^{\mathbf{V}}_{\mathcal{C}} = \coprod_{Di_{\mathcal{C}}} *$.

Since \mathbf{V} is always assumed to have sufficient (co)limits, these two descriptions of oriented graphical species – as \mathbf{V} -valued presheaves on $\mathbf{el}(Di)$ or as objects of $\mathbf{GS}_{\mathbf{V}}$ with morphisms to $Di^{\mathbf{V}}$ – will be used interchangeably in what follows.

Remark 3.10. When \mathbf{V} has finite products and coproducts, addition of natural numbers in Σ induces a symmetric monoidal product on the category of species in \mathbf{V} by

$$(3.11) \quad (S \otimes T)_X \stackrel{\text{def}}{=} \coprod_{(Y, Z), Y \amalg Z = X} S_Y \times T_Z, \text{ for all } X \in \mathbf{B}$$

for species S, T and all finite sets X . It is easily checked (see also [42, Section 2.1]) that monoids for this product correspond to (lax) monoidal presheaves on Σ .

This extends to a symmetric monoidal structure on $\mathbf{GS}_{\mathbf{V}}$ (and also $\mathbf{OGS}_{\mathbf{V}}$) whereby, given $S, T \in \mathbf{GS}_{\mathbf{V}}$, the graphical species $S \otimes T \in \mathbf{GS}_{\mathbf{V}}$, has $(S \otimes T)_X = \coprod_{(Y, Z), Y \amalg Z = X} S_Y \times T_Z$ as in (3.11) and $(S \otimes T)_{\S} \stackrel{\text{def}}{=} S_{\S} \amalg T_{\S}$.

By the universal property of the coproduct, the involutions on S_{\S} and T_{\S} induce an involution on $(S \otimes T)_{\S}$ and the \mathbf{B} -actions on S and T separately induce a \mathbf{B} -action on the collection $((S \otimes T)_X)_{X \in \mathbf{B}}$. For $x \in X$, the morphism $(S \otimes T)(ch_x^X): (S \otimes T)_X \rightarrow (S \otimes T)_{\S}$ is induced by the maps

$$(S \otimes T)(ch_x^{Y \amalg Z}) = \begin{cases} \iota_S \circ S(ch_x^Y) \circ p_{S_Y}: S_Y \times T_Z \rightarrow S_{\S} & \text{when } x \in Y \\ \iota_T \circ T(ch_x^Z) \circ p_{T_Z}: S_Y \times T_Z \rightarrow T_{\S} & \text{when } x \in Z. \end{cases}$$

Here, the maps $p_{(-)}$ from $S_Y \times T_Z$ are the projections and $\iota_{(-)}$ are the maps to the coproduct $(S \otimes T)_{\S}$.

It is straightforward to check that $S \otimes T \in \mathbf{GS}_{\mathbf{V}}$ and that \otimes is associative and the graphical species I defined by $I_{\S} = 0$, $I_{\emptyset} = *$ and $I_X = 0$ for $X \neq \emptyset$ (so $I = I_{\mathbf{V}} = K^{\emptyset}$ when $\mathbf{V} = \mathbf{Set}$) defines a unit for \otimes , making $(\mathbf{GS}_{\mathbf{V}}, \otimes, I)$ into a symmetric monoidal category.

3.3. Modular operads. This section generalises the definition of modular operads in **Set** from [41, Section 1.3] to define modular operads *internal to* arbitrary categories with finite limits. Modular operads internal to a category \mathbf{V} are defined as graphical species in \mathbf{V} equipped with a binary (unital) multiplication operation and unary contraction operation satisfying coherence axioms.

Let S be a graphical species in a category \mathbf{V} with finite limits. For finite sets X and Y with elements $x \in X$ and $y \in Y$, let $(S_X \times S_Y)^{x \ddagger y} \rightarrow S_X \times S_Y$ denote the pullback

$$(3.12) \quad \begin{array}{ccc} (S_X \times S_Y)^{x \ddagger y} & \longrightarrow & S_X \\ \downarrow & & \downarrow S(ch_x) \\ S_Y & \xrightarrow{S(ch_y \circ \tau)} & S_{\S}. \end{array}$$

More generally, given distinct elements $x_1, \dots, x_k \in X$ and $y_1, \dots, y_k \in Y$, let $(S_X \times S_Y)^{x_1 \ddagger y_1, \dots, x_k \ddagger y_k}$ be the limit of the collection $S_X \xrightarrow{S(ch_{x_i})} S_{\S} \xleftarrow{S(ch_{y_i} \circ \tau)} S_Y$, $1 \leq i \leq k$.

Let σ_2 denote the unique non-identity involution on $\mathbf{2}$. A morphism $\epsilon \in \mathbf{V}(S_{\S}, S_2)$ is *unit-like* for S if

$$(3.13) \quad \epsilon \circ \omega = S(\sigma_2) \circ \epsilon: S_{\S} \rightarrow S_2, \text{ and } S(ch_1) \circ \epsilon = id_{S_{\S}}.$$

Definition 3.14. A multiplication \diamond on S is a family of morphisms

$$\diamond_{X,Y}^{x \ddagger y}: (S_X \times S_Y)^{x \ddagger y} \rightarrow S_{(X \amalg Y) \setminus \{x,y\}},$$

in \mathbf{V} , defined for all pairs of finite sets X and Y , with elements $x \in X$ and $y \in Y$, and such that

(1) the obvious diagram

$$\begin{array}{ccc} (S_X \times S_Y)^{x \ddagger y} & \xrightarrow{\diamond_{X,Y}^{x \ddagger y}} & S_{(X \amalg Y) \setminus \{x,y\}} \\ \cong \downarrow & & \uparrow \diamond_{Y,X}^{y \ddagger x} \\ (S_Y \times S_X)^{y \ddagger x} & \xrightarrow{\diamond_{Y,X}^{y \ddagger x}} & S_{(X \amalg Y) \setminus \{x,y\}} \end{array}$$

commutes in \mathbf{V} ,

(2) \diamond is equivariant with respect to the \mathbf{B}^{\S} -action on S : if $\hat{\sigma}: X \setminus \{x\} \xrightarrow{\cong} W \setminus \{w\}$ and $\hat{\rho}: Y \setminus \{y\} \xrightarrow{\cong} Z \setminus \{z\}$ are restrictions of bijections $\sigma: X \xrightarrow{\cong} W$ and $\rho: Y \xrightarrow{\cong} Z$ such that $\sigma(x) = w$ and $\rho(y) = z$, then

$$S(\hat{\sigma} \sqcup \hat{\rho}) \diamond_{W,Z}^{w \ddagger z} = \diamond_{X,Y}^{x \ddagger y} S(\sigma \sqcup \rho),$$

(where $\sigma \sqcup \rho: X \amalg Y \xrightarrow{\cong} W \amalg Z$ is the block-wise permutation).

A multiplication \diamond on S is *unital* if S there is a unit-like morphism $\epsilon \in \mathbf{V}(S_{\S}, S_2)$ such that, for all finite sets X and all $x \in X$, the composite

$$(3.15) \quad S_X \xrightarrow{(id \times ch_x) \circ \Delta} S_X \times S_{\S} \xrightarrow{id \times \epsilon} S_X \times S_2 \xrightarrow{\diamond_{X,2}^{x \ddagger 2}} S_X$$

is the identity on S_X .

It is immediate from the definition (3.13) that any unit-like morphism in $\epsilon \in \mathbf{V}(S_{\S}, S_2)$ is monomorphic and, by (3.15) and condition (1) of Equation (3.14), if a multiplication \diamond on S admits a unit ϵ , then it is unique.

Now, let S be a graphical species in \mathbf{V} , and let $x \neq y$ be distinct elements of a finite set Z . Let $(S_Z)^{x \ddagger y} = S_Z^{y \ddagger x}$ denote the equaliser

$$(3.16) \quad S_Z^{x \ddagger y} \longrightarrow S_Z \xrightarrow[S(ch_y \circ \tau)]{S(ch_x)} S_{\S}.$$

More generally, for any morphism $f: C \rightarrow S_Z$ in \mathbf{V} , let $C^{x \dagger y} = C^{y \dagger x}$ denote the pullback of f along the universal map $S_Z^{x \dagger y} \rightarrow S_Z$. Given distinct elements $x_1, y_1, \dots, x_k, y_k$ of S_Z , and a morphism $f: C \rightarrow S_Z$ in \mathbf{V} , then $C^{x_1 \dagger y_1, \dots, x_k \dagger y_k} \stackrel{\text{def}}{=} (C^{x_1 \dagger y_1, \dots, x_{k-1} \dagger y_{k-1}})^{x_k \dagger y_k}$ is the obvious limit.

Invariance of $C^{x_1 \dagger y_1, \dots, x_k \dagger y_k}$ under permutations $(x_i, y_i) \mapsto (x_{\sigma i}, y_{\sigma i})$, $\sigma \in \Sigma_k$ follows from invariance of the defining morphisms

$$C^{x_1 \dagger y_1, \dots, x_k \dagger y_k} \longrightarrow S_Z^{x_1 \dagger y_1, \dots, x_k \dagger y_k} \longrightarrow S_Z.$$

Definition 3.17. A (graphical species) contraction ζ on S in \mathbf{V} is a family of maps $\zeta_X^{x \dagger y}: S_X^{x \dagger y} \rightarrow S_{X \setminus \{x, y\}}$ defined for each finite set X and pair of distinct elements $x, y \in X$.

The contraction ζ is equivariant with respect to the action of $\mathbf{B}^{\mathfrak{s}}$ on S : If $\hat{\sigma}: X \setminus \{x, y\} \xrightarrow{\cong} Z \setminus \{w, z\}$ is the restriction of a bijection $\sigma: X \xrightarrow{\cong} Z$ with $\sigma(x) = w$ and $\sigma(y) = z$. Then

$$(3.18) \quad S(\hat{\sigma}) \circ \zeta_Z^{w \dagger z} = \zeta_X^{x \dagger y} \circ S(\sigma): S_Z^{w \dagger z} \rightarrow S_{X \setminus \{x, y\}}.$$

In particular, by (3.18), if ζ is a contraction on S , then $\zeta_X^{x \dagger y} = \zeta_X^{y \dagger x}$ for all finite sets X and all pairs of distinct elements $x, y \in X$.

Definition 3.19. A nonunital modular operad (S, \diamond, ζ) in \mathbf{V} is a graphical species $S \in \mathbf{GS}_{\mathbf{V}}$ equipped with a multiplication \diamond and a contraction ζ such that the following four coherence axioms are satisfied:

(M1) Multiplication is associative (see Figure 1(M1)):

For all finite set X, Y, Z and elements $x \in X$, $z \in Z$, and distinct $y_1, y_2 \in Y$,

$$\begin{array}{ccc} (S_X \times S_Y \times S_Z)^{x \dagger y_1, y_2 \dagger z} & \xrightarrow{\diamond_{X, Y}^{x \dagger y_1} \times id_{S_Z}} & (S_{(X \amalg Y) \setminus \{x, y_1\}} \times S_Z)^{y_2 \dagger z} \\ \downarrow id_{S_X} \times \diamond_{Y, Z}^{y_2 \dagger z} & = & \downarrow \diamond_{X \amalg Y \setminus \{x, y_1\}, Z}^{y_2 \dagger z} \\ (S_X \times S_{(Y \amalg Z) \setminus \{y_2, z\}})^{x \dagger y_1} & \xrightarrow[\diamond_{X, Y \amalg Z \setminus \{y_2, z\}}^{x \dagger y_1}]{} & S_{(X \amalg Y \amalg Z) \setminus \{x, y_1, y_2, z\}}. \end{array}$$

(M2) Contractions commute (see Figure 1(M2)):

For all finite sets X with distinct elements w, x, y, z , the following square commutes:

$$\begin{array}{ccc} S_X^{w \dagger x, y \dagger z} & \xrightarrow{\zeta_X^{w \dagger x}} & S_{X \setminus \{w, x\}}^{y \dagger z} \\ \downarrow \zeta_X^{y \dagger z} & = & \downarrow \zeta_{X \setminus \{w, x\}}^{y \dagger z} \\ S_{X \setminus \{y, z\}}^{w \dagger x} & \xrightarrow[\zeta_{X \setminus \{y, z\}}^{w \dagger x}]{} & S_{X \setminus \{w, x, y, z\}} \end{array}.$$

(M3) Multiplication and contraction commute (see Figure 1(M2)):

For finite sets X, Y , distinct x_1, x_2 and x_3 in X , and $y \in Y$, the following diagram commutes:

$$\begin{array}{ccc} (S_X^{x_1 \dagger x_2} \times S_Y)^{x_3 \dagger y} & \xrightarrow{\zeta_X^{x_1 \dagger x_2} \times id_Y} & (S_{X \setminus \{x_1, x_2\}} \times S_Y)^{x_3 \dagger y} \\ \downarrow \diamond_{X, Y}^{x_3 \dagger y} & = & \downarrow \diamond_{X \setminus \{x_1, x_2\}, Y}^{x_3 \dagger y} \\ S_{(X \amalg Y) \setminus \{x_3, y\}}^{x_1 \dagger x_2} & \xrightarrow[\zeta_{X \amalg Y \setminus \{x_3, y\}}^{x_1 \dagger x_2}]{} & S_{(X \amalg Y) \setminus \{x_1, x_2, x_3, y\}}. \end{array}$$

(M4) “Parallel multiplication” of pairs (see Figure 1(M4)):

For finite sets X and Y , and distinct elements $x_1, x_2 \in X$ and $y_1, y_2 \in Y$, the following diagram commutes:

$$\begin{array}{ccc}
 (S_X \times S_Y)^{x_1 \ddagger y_1, x_2 \ddagger y_2} & \xrightarrow{\diamond_{X,Y}^{x_1 \ddagger y_1}} & S_{(X \amalg Y) \setminus \{x_1, y_1\}}^{x_2 \ddagger y_2} \\
 \downarrow \diamond_{x_2 \ddagger y_2, X,Y} & = & \downarrow \zeta_{X \amalg Y \setminus \{x_1, y_1\}}^{x_2 \ddagger y_2} \\
 S_{(X \amalg Y) \setminus \{x_2, y_2\}}^{x_1 \ddagger y_1} & \xrightarrow{\zeta_{x_1 \ddagger y_1, X \amalg Y \setminus \{x_2, y_2\}}} & S_{(X \amalg Y) \setminus \{x_1, x_2, y_1, y_2\}}
 \end{array}$$




FIGURE 1. Modular operad axioms (M1)-(M4).

(Axioms (M1)-(M4) are illustrated in Figure 1.)

The category of nonunital, modular operads in \mathbf{V} is denoted by $\mathfrak{MO}_{\mathbf{V}}^-$.

A (unital) modular operad $(S, \diamond, \zeta, \epsilon)$ in \mathbf{V} is a graphical species $S \in \mathbf{GS}_{\mathbf{V}}$ equipped with a unital multiplication (\diamond, ϵ) and contraction ζ such that $(S, \diamond, \zeta) \in \mathfrak{MO}_{\mathbf{V}}^-$. The category of modular operads in \mathbf{V} is denoted by $\mathfrak{MO}_{\mathbf{V}}$.

Remark 3.20. As discussed in Section 3.2, $\mathbf{GS}_{\mathbf{V}}$ has a monoidal category structure when \mathbf{V} admits finite coproducts. However, as will be explained in Equation (6.29), the categories \mathfrak{MO} and \mathfrak{MO}^- are not closed under the monoidal product in $\mathbf{GS}_{\mathbf{V}}$.

3.4. Circuit algebras. A *circuit algebra internal to a category* \mathbf{V} with finite limits (see Equation (3.25)) is a graphical species A in \mathbf{V} that is equipped with an equivariant contraction and such that $(A(\mathbf{n}))_{\mathbf{n}}$ describes a graded associative monoid. The monoid structure and contraction satisfy coherence axioms analogous to the (enriched) circuit algebra axioms (c1) – (c3), (e1) of [42, Proposition 5.4].

Definition 3.21. A (commutative) external product on a graphical species S in \mathbf{V} is given by a $\mathbf{B} \times \mathbf{B}$ -indexed collection of equivariant morphisms $\boxtimes_{X,Y} : S_X \times S_Y \rightarrow S_{X \amalg Y}$, in \mathbf{V} such that $\boxtimes_{Y,X} = \boxtimes_{X,Y} \circ \sigma_{Y,X}$ (where $\sigma_{Y,X} : S_Y \times S_X \xrightarrow{\cong} S_X \times S_Y$ is the swap isomorphism) and, for all elements $x \in X$, the following diagram commutes in \mathbf{V} :

$$\begin{array}{ccc} S_X \times S_Y & \xrightarrow{\boxtimes_{X,Y}} & S_{X \amalg Y} \\ \downarrow & & \downarrow S(ch_x^{X \amalg Y}) \\ S_X & \xrightarrow{S(ch_x^X)} & S_{\S} \end{array}$$

An external unit for the external product \boxtimes is a distinguished morphism $\ominus : * \rightarrow S_{\mathbf{0}}$ such that, for all finite sets X , the composite

$$S_X \xrightarrow{\cong} S_X \times * \xrightarrow{id \times \ominus} S_X \times S_{\mathbf{0}} \xrightarrow{\boxtimes} S_{(X \amalg \mathbf{0})} = S_X$$

is the identity on S_X for all finite sets X .

Remark 3.22. Henceforth, the monoidal unit \ominus for \boxtimes will be suppressed in the notation since most of the constructions in Sections 3-8 of this paper carry through without it. To agree with the circuit algebra literature, external products in this paper are always assumed to have a unit. See also Equation (5.17), and [31, Paragraph 4.7].

Recall that, if \mathbf{V} has finite coproducts, then $\mathbf{GS}_{\mathbf{V}}$ is a symmetric monoidal category. It follows immediately from Equation (3.21) that:

Lemma 3.23. A graphical species S in \mathbf{V} admits an external product (\boxtimes, \ominus) if and only if (S, \boxtimes, \ominus) is a monoid for the monoidal product \otimes on $\mathbf{GS}_{\mathbf{V}}$.

An external product \boxtimes and contraction ζ on a graphical species S induce an equivariant multiplication \diamond on S : for all finite sets X and Y , all $x \in X$ and $y \in Y$, the multiplication $\diamond_{X,Y}^{x \ddagger y}$ is defined by the composition

$$(3.24) \quad \diamond_{X,Y}^{x \ddagger y} : (S_X \times S_Y)^{x \ddagger y} \xrightarrow{\boxtimes_{X,Y}} (S_{X \amalg Y})^{x \ddagger y} \xrightarrow{\zeta^{x \ddagger y}} S_{(X \amalg Y) \setminus \{x,y\}}.$$

Definition 3.25. A nonunital circuit algebra in \mathbf{V} is a graphical species S in \mathbf{V} , equipped with an external product \boxtimes and a contraction ζ , such that the following three axioms (illustrated in Figure 2) are satisfied:

(C1) External product is associative (see Figure 2 (C1)):

For all finite sets X, Y, Z , the following square commutes:

$$\begin{array}{ccc} S_X \times S_Y \times S_Z & \xrightarrow{\boxtimes_{X,Y} \times id_{S_Z}} & S_{X \amalg Y} \times S_Z \\ \downarrow id_{S_X} \times \boxtimes_{Y,Z} & = & \downarrow \boxtimes_{(X \amalg Y), Z} \\ S_X \times S_{Y \amalg Z} & \xrightarrow{\boxtimes_{X, (Y \amalg Z)}} & S_{X \amalg Y \amalg Z}. \end{array}$$

(C2) Order of contraction does not matter. *This corresponds with Equation (3.19) (M2) (see also Figure 1 (M2)).*

(C3) Contraction and external product commute (see Figure 2 (C1)):

For all finite sets X and Y , and distinct elements $x_1, x_2 \in X$, the following square commutes:

$$\begin{array}{ccc}
 S_X^{x_1 \sharp x_2} \times S_Y & \xrightarrow{\boxtimes_{X,Y}} & S_{X \amalg Y}^{x_1 \sharp x_2} \\
 \zeta_X^{x_1 \sharp x_2} \times id_{S_Y} \downarrow & = & \downarrow \zeta_{X \amalg Y}^{x_1 \sharp x_2} \\
 S_{X \setminus \{x_1, x_2\}} \times S_Y & \xrightarrow{\boxtimes_{X \setminus \{x_1, x_2\}, Y}} & S_{X \setminus \{x_1, x_2\} \amalg Y}
 \end{array}$$

FIGURE 2. Circuit algebra axioms (C1) and (C3). For axiom (C2), see Figure 1(M2).

Let $\mathfrak{CA}_{\mathbf{V}}$ denote the category of nonunital circuit algebras in \mathbf{V} and morphisms of graphical species that preserve the external product and contraction.

A unital circuit algebra (in \mathbf{V}) is a nonunital circuit algebra (S, \boxtimes, ζ) together with a unit-like morphism $\epsilon: S_{\S} \rightarrow S_2$ (Equation (3.14)), such that, for all finite sets X and $x \in X$, the composite

$$(3.26) \quad S_X \xrightarrow{(id \times ch_x) \circ \Delta} (S_X \times S_{\S})^{x \sharp 2} \xrightarrow{id \times \epsilon} (S_X \times S_2)^{x \sharp 2} \xrightarrow{\boxtimes} (S_{X \amalg 2})^{x \sharp 2} \xrightarrow{\zeta_{X \amalg 2}^{x \sharp 2}} S_X$$

is the identity on S_X . (Here $\Delta: S_X \rightarrow S_X \times S_X$ is the diagonal and the last map makes use of the canonical isomorphism $(X \amalg \{1\}) \setminus \{x\} \cong X$.)

Morphisms in the category $\mathfrak{CA}_{\mathbf{V}}$ of circuit algebras in \mathbf{V} are morphisms in $\mathfrak{CA}_{\mathbf{V}}$ that preserve units.

For $\mathbf{V} = \mathbf{Set}$, we will write $\mathfrak{CA} \stackrel{\text{def}}{=} \mathfrak{CA}_{\mathbf{V}}$ and $\mathfrak{MO} \stackrel{\text{def}}{=} \mathfrak{MO}_{\mathbf{V}}$. Let \mathbf{CA} be the category, defined in [42, Section 5.2], of circuit algebras (of all colours) enriched in \mathbf{Set} . The following proposition follows immediately from [42, Proposition 5.4] and by unpacking Equation (3.25):

Proposition 3.27. *The inclusion $\Sigma \hookrightarrow \mathbf{B}$ induces a canonical equivalence of categories $\mathbf{CA} \simeq \mathfrak{CA}$.*

Remark 3.28. Henceforth, unless explicitly stated otherwise, we will not consider enriched circuit algebras: circuit algebras and modular operads will be assumed to be objects of the categories $\mathfrak{CA}_{\mathbf{V}}$ and $\mathfrak{MO}_{\mathbf{V}}$ (and $\mathfrak{CA}_{\mathbf{V}}^{\neg}$, $\mathfrak{MO}_{\mathbf{V}}^{\neg}$) so-defined, internally to a category \mathbf{V} with sufficient limits.

By (3.24), a (nonunital) circuit algebra (S, \boxtimes, ζ) admits a (nonunital) multiplication \diamond_S induced by \boxtimes and ζ . It follows immediately from (3.24) and axioms (C1)-(C3), that (S, \diamond_S, ζ) satisfy (M1)-(M4). Hence, circuit algebras in \mathbf{V} canonically admit a modular operad structure. This will be used

in Sections 4-7.2, to construct the composite monad for circuit algebras by modifying the modular operad monad of [41].

In fact, we have the following proposition, analogous to [42, Proposition 5.11], that will follow from Equation (7.9), Equation (5.14) and Equation (7.4), together with the classical theory of distributive laws [5]:

Proposition 3.29. *There are canonical monadic adjunctions*

$$(3.30) \quad \begin{array}{ccc} \mathfrak{CA}_V^- & \xrightleftharpoons{\top} & \mathfrak{CA}_V \\ \downarrow \vdash & & \downarrow \vdash \\ \mathfrak{MO}_V^- & \xrightleftharpoons{\top} & \mathfrak{MO}_V \end{array}$$

between the categories of (nonunital) modular and product modular operads in \mathbf{V} .

Proof. By Equation (7.9), \mathfrak{CA}_V is the EM category of algebras $\mathbf{GS}_V^{\mathbb{L}\mathbb{D}\mathbb{T}}$ for the composite monad $\mathbb{L}\mathbb{D}\mathbb{T}$ on \mathbf{GS}_V , constructed using iterated distributive laws. Now, $\mathfrak{MO}_V^- = \mathbf{GS}_V^{\mathbb{T}}$ by Equation (6.2), $\mathfrak{MO}_V = \mathbf{GS}_V^{\mathbb{D}\mathbb{T}}$ by Equation (6.27) (see also [41]), and $\mathfrak{CA}_V^- = \mathbf{GS}_V^{\mathbb{L}\mathbb{T}}$ by Equation (5.14) and Equation (7.4). Hence, the vertical adjunctions exist by the classical theory of distributive laws [5].

The horizontal adjunctions – where the left adjoints correspond to freely adjoining modular operadic units – exist by Equation (6.25). \square

The right adjoints of the pairs $\mathfrak{MO}_V \rightleftharpoons \mathfrak{CA}_V$ (and $\mathfrak{MO}_V^- \rightleftharpoons \mathfrak{CA}_V^-$) in Equation (3.29) correspond to the forgetful functor $(S, \boxtimes, \zeta) \mapsto (S, \diamond, \zeta)$ where the multiplication \diamond is defined by suitable compositions $\zeta \circ \boxtimes$ as in (3.24). The left adjoint is induced by the free graded monoid monad:

Let $\text{Core}(X/\text{Set}_f)$ be the maximal subgroupoid of the slice category X/Set_f whose objects are morphisms of finite sets $f: X \rightarrow Y$, and whose morphisms $(Y, f) \rightarrow (Y', f')$ are isomorphisms $g \cong Y \rightarrow Y'$ such that $gf = f'$. If S is a graphical species, then there is a graphical species LS defined by

$$LS_X = \text{colim}_{(Y, f) \in \text{Core}(X/\text{Set}_f)} \prod_{y \in Y} S_{f^{-1}(y)}.$$

The free monoid structure induces the external product on LS , and, if (S, \diamond, ζ) is a (nonunital) modular operad, then the modular operad axioms imply that the pair of operations ζ and \diamond , define a contraction on LS (see Section 7.2).

In particular, since units for the modular operadic multiplication are unique:

Corollary 3.31. *The canonical forgetful functors $\mathfrak{CA}_V \rightarrow \mathfrak{CA}_V^-$ and $\mathfrak{MO}_V \rightarrow \mathfrak{MO}_V^-$ are inclusions of categories.*

Example 3.32. By [13, 42], oriented circuit algebras enriched in a symmetric monoidal category (\mathbf{X}, \otimes, I) are equivalent to (coloured) wheeled props enriched in \mathbf{X} . The image of a (nonunital) wheeled prop under the forgetful functor $\mathfrak{CA} \rightarrow \mathfrak{MO}$ is its underlying (nonunital) *wheeled properad* [22, 52]. (See also Equation (3.6), Equation (6.28) and Equation (8.6) and Proposition 3.26 and Example 5.10 of [42].)

The graphical species Di^V (Examples 3.7 & 3.9) trivially admits a circuit algebra structure. Hence, Equation (3.32) motivates the following definition:

Definition 3.33. *The category \mathfrak{WP}_V of wheeled props in \mathbf{V} is defined as the slice category \mathfrak{CA}_V/Di^V . When $\mathbf{V} = \text{Set}$, we write $\mathfrak{WP} \stackrel{\text{def}}{=} \mathfrak{WP}_{\text{Set}}$.*

The category of nonunital wheeled props in \mathbf{V} is defined similarly by $\mathfrak{WP}_V^- \stackrel{\text{def}}{=} \mathfrak{CA}_V^-/Di^V$.

Remark 3.34. By Equation (3.29), a circuit algebra is a modular operad equipped with an external product that is compatible with the modular operadic multiplication in the sense of (3.24). By Equation (3.23), a graphical species with external product (S, \boxtimes, \ominus) is just a monoid object in \mathbf{GS}_V , so it is natural to ask whether there is a monoidal structure on \mathbf{MO}_V such that circuit algebras are precisely the monoid objects in \mathbf{MO}_V . In fact, this is not the case since, as will be explained in Equation (6.29), \mathbf{MO}_V is not closed under the monoidal product on \mathbf{GS}_V .

4. BACKGROUND ON GRAPHS

The combinatorics of modular operads have been described in [41, Section 8] in terms of a category Ξ of connected graphs. The remainder of the present paper is concerned with an analogous construction and graphical category Ξ^\times for circuit algebras.

Sections 4.1 and 4.2 provide a review of the category (Feynman) graphs and étale morphisms, first introduced in [27], that will underpin the rest of the paper. The interested reader is referred to [41, Sections 3 & 4] for details and explicit proofs related to the graphical formalism.

Section 4.3 gives a brief description of the operads of wiring diagrams and Brauer diagram category used to describe enriched circuit algebras (see e.g., [3, 13, 14] and [42]), and explains their relationship with Feynman graphs. This is purely for context and may be omitted.

4.1. Graphs and étale morphisms. The following definition is due to Joyal and Kock [27] (see also [23, 24, 30, 41]):

Definition 4.1. A (Feynman) graph \mathcal{G} is a diagram of finite sets

$$(4.2) \quad \mathcal{G} = \tau \bigcirc E \xleftarrow{s} H \xrightarrow{t} V ,$$

such that $s: H \rightarrow E$ is injective, and $\tau: E \rightarrow E$ is an involution without fixed points.

A subgraph $\mathcal{H} \subset \mathcal{G}$ is a triple of subsets $E' \subset E, H' \subset H, V' \subset V$ that inherit a graph structure from \mathcal{G} .

Elements of V are *vertices* of \mathcal{G} and elements of E are called *edges* of \mathcal{G} . The set of τ -orbits in E is denoted by \tilde{E} . Elements h of the set H are *half-edges* of \mathcal{G} .

Definition 4.3. The set $E_\bullet \subset E$ of inner edges of \mathcal{G} is the maximal subset of $\text{im}(s) \subset E$ that is closed under τ . The set of inner τ -orbits $\tilde{e} \in \tilde{E}$ with $e \in E_\bullet$ is denoted by \tilde{E}_\bullet . Elements of the set $E_0 \stackrel{\text{def}}{=} E \setminus \text{im}(s)$ are ports of \mathcal{G} .

A stick component of a graph \mathcal{G} is a pair $\{e, \tau e\}$ of edges of \mathcal{G} such that e and τe are both ports.

Definition 4.4. A graph is connected if it cannot be written as a disjoint union of non-empty graphs. A connected component \mathcal{H} of \mathcal{G} is a maximal connected subgraph of \mathcal{G} .

In particular, any graph \mathcal{G} is the disjoint union of its connected components, and the inclusion $\mathcal{H} \hookrightarrow \mathcal{G}$ of a connected component \mathcal{H} of \mathcal{G} describes a pointwise injective étale morphism.

Example 4.5. A subgraph $\mathcal{H} \subset \mathcal{G}$ of the form $\{e, \tau e\} \longleftarrow \emptyset \longrightarrow \emptyset$ describes a connected component of \mathcal{G} if and only if $\{e, \tau e\}$ is a stick component of \mathcal{G} .

Remark 4.6. A graph \mathcal{G} can be thought of as a rule for gluing closed intervals along their boundaries and may be realised geometrically as the one-dimensional space $|\mathcal{G}|$, obtained by taking the discrete space $\{*_v\}_{v \in V}$, and, for each $e \in E$, a copy $[0, \frac{1}{2}]_e$ of the interval $[0, \frac{1}{2}]$ subject to the identifications $0_{s(h)} \sim *_t(h)$ for $h \in H$ and $(\frac{1}{2})_e \sim (\frac{1}{2})_{\tau e}$ for all $e \in E$.

Disjoint union commutes with geometric realisation: for all graphs \mathcal{G} and \mathcal{H} , $|\mathcal{G} \amalg \mathcal{H}| = |\mathcal{G}| \amalg |\mathcal{H}|$. So, a graph \mathcal{G} is connected if and only if its realisation $|\mathcal{G}|$ is a connected space.

Example 4.7. (See also Equation (4.42).) Up to isomorphism, there are four distinct connected graphs with a single edge orbit:

(a) the **1-corolla** \mathcal{C}_1 has one vertex, no inner edges, and one port (see Figure 3(a))

$$\mathcal{C}_1 \stackrel{\text{def}}{=} \bigcirc \mathbf{2} \longleftarrow \mathbf{1} \longrightarrow \mathbf{1}$$

(b) the **stick graph** (\mathfrak{i}) has no vertices, no inner edges and two ports (see Figure 3(b))

$$(\mathfrak{i}) \stackrel{\text{def}}{=} \bigcirc \mathbf{2} \longleftarrow \mathbf{0} \longrightarrow \mathbf{0}$$

(c) the **wheel graph** \mathcal{W} has one vertex, one inner edge orbit and no ports (see Figure 3(c));

$$\mathcal{W} \stackrel{\text{def}}{=} \bigcirc \mathbf{2} \xleftarrow{id} \mathbf{2} \longrightarrow \mathbf{1}$$

(d) finally, there is a graph with two vertices, one inner edge orbit and no ports (Figure 3(d)):

$$\bigcirc \mathbf{2} \xleftarrow{id} \mathbf{2} \xrightarrow{id} \mathbf{2}.$$

The stick and wheel graphs (\mathfrak{i}) and \mathcal{W} are particularly important in what follows.




FIGURE 3. Realisations of connected graphs with one edge orbit: (a) the 1-corolla \mathcal{C}_1 ; (b) the stick graph (\mathfrak{i}) ; (c) the wheel graph \mathcal{W} consists of a single inner edge orbit with one end vertex; (d) a single inner edge orbit with distinct end vertices.

For any set X , let $X^\dagger \cong X$ denote its formal involution.

Example 4.8. (See also Figure 3(d).) The X -corolla \mathcal{C}_X associated to a finite set X has the form

$$\mathcal{C}_X : \quad \dagger \bigcirc X \amalg X^\dagger \xleftarrow{\text{inc}} X^\dagger \longrightarrow \{*\}.$$

Let \mathcal{G} be a graph with vertex and edge sets V and E respectively. For each vertex v , define $H/v \stackrel{\text{def}}{=} t^{-1}(v) \subset H$ to be the fibre of t at v , and let $E/v \stackrel{\text{def}}{=} s(H/v) \subset E$.

Definition 4.9. Edges in the set E/v are said to be incident on v .

The valency $|v|$ of a vertex $v \in V$ is given by $|v| \stackrel{\text{def}}{=} |H/v|$ and $V_n \subset V$ is the set of n -valent vertices of \mathcal{G} . A vertex v is bivalent if $|v| = 2$ and a bivalent graph is a graph \mathcal{G} with $V = V_2$. A vertex $v \in V$ with $|v| = 0$ is called an isolated vertex of \mathcal{G} .

Bivalent and isolated vertices are particularly important in Section 6.3.

Vertex valency induces an \mathbb{N} -grading on the edge set E of \mathcal{G} : For $n \geq 1$, define $H_n \stackrel{\text{def}}{=} t^{-1}(V_n)$ and $E_n \stackrel{\text{def}}{=} s(H_n)$. Then $s(H) = \coprod_{n \geq 1} E_n = E \setminus E_0$, so $E = \coprod_{n \in \mathbb{N}} E_n$.

Example 4.10. The stick graph (\mathfrak{i}) (Equation (4.7)) has $E_0(\mathfrak{i}) = E(\mathfrak{i}) = \mathbf{2}$, and $E_\bullet(\mathfrak{i}) = \emptyset$. Conversely, the wheel graph \mathcal{W} has $E_0(\mathcal{W}) = \emptyset$ and $E_\bullet(\mathcal{W}) = E_2(\mathcal{W}) = E(\mathcal{W}) \cong \mathbf{2}$.

The X -corolla \mathcal{C}_X (see Equation (4.8)) with vertex $*$ has set of ports $E_0(\mathcal{C}_X) = X$ and $E/* = X^\dagger$. So, if $X \cong \mathbf{n}$, then $|*| = n$, $V = V_n$, and $E = E_n \amalg E_0$ where $E_i \cong X$ for $i = 0, n$.

Definition 4.11. An étale morphism $f: \mathcal{G} \rightarrow \mathcal{G}'$ of graphs is a commuting diagram of finite sets

$$(4.12) \quad \begin{array}{ccccccc} \mathcal{G} & & E & \xleftarrow{\tau} & E & \xleftarrow{s} & H & \xrightarrow{t} & V \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \mathcal{G}' & & E' & \xleftarrow{\tau'} & E' & \xleftarrow{s'} & H' & \xrightarrow{t'} & V' \end{array}$$

such that the right-hand square is a pullback. The category of graphs and étale morphisms is denoted by \mathbf{Gr} .

It is straightforward to verify that the right hand square in (4.12) is a pullback if and only if, for all $v \in V$, the map $E/v \rightarrow E'/f(v)$ induced by the restriction of the map $V \rightarrow V'$ is bijective. Hence, as the name suggests, étale morphisms describe local isomorphisms.

Example 4.13. Let \mathcal{G} be a graph with edge set E . Then $\mathbf{Gr}(1, \mathcal{G}) = \{ch_e\}_{e \in E} \cong E$ where for each edge $e \in V$, $ch_e: (1) \rightarrow \mathcal{G}$ is the morphism $1 \mapsto e$ that chooses e .

Pointwise disjoint union defines a symmetric strict monoidal structure on \mathbf{Gr} with unit given by the empty graph $\emptyset \stackrel{\text{def}}{=} \emptyset \longleftarrow \emptyset \longrightarrow \emptyset$.

Example 4.14. Let X and Y be finite sets. Write $\mathcal{D}_{X,Y} \stackrel{\text{def}}{=} \mathcal{C}_X \amalg \mathcal{C}_Y$ for the disjoint union of X and Y -corollas. This has ports $E_0(\mathcal{D}_{X,Y}) = X \amalg Y = E_0(\mathcal{C}_X) \amalg E_0(\mathcal{C}_Y)$ and no internal edges. The canonical inclusions $\iota_X: \mathcal{C}_X \hookrightarrow \mathcal{D}_{X,Y} \hookleftarrow \mathcal{C}_Y: \iota_Y$ are étale.

Example 4.15. Up to isomorphism, (1) is the only connected graph with no vertices. As in [41], and following [30], a *shrub* \mathcal{S} is a disjoint union of stick graphs. Any commuting diagram of the form (4.12), where $\mathcal{G} = \mathcal{S}$ is a shrub, is trivially étale and hence defines a morphism in \mathbf{Gr} .

Example 4.16. (C.f. [42, Remark 3.12].) Let \mathbf{D} be the small category $\begin{array}{c} \bullet \\ \curvearrowright \end{array} \bullet \longrightarrow \bullet \longleftarrow \bullet$. So \mathbf{Gr} is a subcategory of $\mathbf{psh}(\mathbf{D})$. The endomorphisms $id, \tau: (1) \rightrightarrows (1)$ of (1) in \mathbf{Gr} are coequalised in $\mathbf{psh}(\mathbf{D})$ by $* \leftarrow \emptyset \rightarrow \emptyset$, which is not a graph. So, $id_{(1)}$ and τ do not admit a coequaliser in \mathbf{Gr} . This provided a motivating example for [41].

The two endomorphisms $id_{\mathcal{W}}, \tau_{\mathcal{W}}: \mathcal{W} \rightarrow \mathcal{W}$, viewed as morphisms of presheaves on \mathbf{D} , have coequaliser $* \leftarrow * \rightarrow *$ which is terminal in $\mathbf{psh}(\mathbf{D})$, and is not a graph. In particular, \mathbf{Gr} does not have a terminal object and is therefore not closed under (finite) limits.

As Equation (4.16) shows, \mathbf{Gr} does not, in general, admit finite colimits. However, if \mathcal{G} is a graph and $e_1, e_2 \in E_0$ are ports of \mathcal{G} with $e_1 \neq \tau e_2$, then the colimit $\mathcal{G}^{e_1 \dagger e_2}$ of the parallel morphisms

$$(4.17) \quad ch_{e_1}, ch_{e_2} \circ \tau: (1) \rightrightarrows \mathcal{G}$$

is the graph described by identifying $e_1 \sim \tau e_2$ and $e_2 \sim \tau e_1$ in the diagram (4.2) defining \mathcal{G} .

Example 4.18. The graph $\mathcal{C}_2^{e_1 \dagger \tau e}$ obtained by identifying the ports of the **2**-corolla is isomorphic to the wheel graph \mathcal{W} .

Example 4.19. More generally, for X a finite set with distinct elements x and y , the graph $\mathcal{C}_X^{x \dagger y}$ has ports in $E_0 = X \setminus \{x, y\}$, one inner τ -orbit $\{x, y\}$ (bold-face in Figure 4), and one vertex v :

$$\tau \begin{array}{c} \curvearrowright \end{array} (X \setminus \{x, y\}) \amalg X^\dagger \xleftarrow{s} X^\dagger \xrightarrow{t} \{v\},$$

where $\tau x^\dagger = y^\dagger$ and $\tau z = z^\dagger$ for $z \in X \setminus \{x, y\}$. Graphs of the form $\mathcal{C}_X^{x \dagger y}$ encode formal contractions in graphical species.

Since the monoidal structure on \mathbf{Gr} is induced by the cocartesian monoidal structure on diagrams of finite sets, colimits in \mathbf{Gr} , when they exist, commute with \amalg . In particular, for any graphs \mathcal{G} and \mathcal{H} and distinct ports e_1, e_2 of \mathcal{G} ,

$$(4.20) \quad \mathcal{G}^{e_1 \ddagger e_2} \amalg \mathcal{H} = (\mathcal{G} \amalg \mathcal{H})^{e_1 \ddagger e_2}.$$

Example 4.21. Recall (Equation (4.14)) that $\mathcal{D}_{X,Y}$ is the disjoint union $\mathcal{C}_X \amalg \mathcal{C}_Y$. Let $x \in X$, and $y \in Y$. By identifying the edges $x \sim \tau y, y \sim \tau x$, we obtain a graph $\mathcal{D}_{X,Y}^{x \ddagger y} \stackrel{\text{def}}{=} (\mathcal{C}_X \amalg \mathcal{C}_Y)^{x \ddagger y}$, that has two vertices and one inner edge orbit $\{x, y\}$, highlighted in bold-face in Figure 4.

$$\tau \bigcirc \left((X \amalg Y) \setminus \{x, y\} \amalg (X \amalg Y)^\dagger \right) \xleftarrow{s} \left((X \amalg Y)^\dagger \right) \xrightarrow{t} \{v_X, v_Y\}$$

with s the obvious inclusion and the involution τ described by $z \leftrightarrow z^\dagger$ for $z \in (X \amalg Y) \setminus \{x, y\}$ and $x^\dagger \leftrightarrow y^\dagger$. The map t is described by $t^{-1}(v_X) = (X \setminus \{x\})^\dagger \amalg \{y\}$ and $t^{-1}(v_Y) = (Y \setminus \{y\})^\dagger \amalg \{x\}$.

In the construction of modular operads, graphs of the form $\mathcal{D}_{X,Y}^{x \ddagger y}$ are used to encode formal multiplications in graphical species.




FIGURE 4. Realisations of $\mathcal{D}_{X,Y}^{x \ddagger y}$ and $\mathcal{C}_X^{x \ddagger y}$ for $X \cong 4$, $Y \cong 3$.

Example 4.22. For $k \geq 0$, the *line graph* \mathcal{L}^k is the connected bivalent graph (illustrated in Figure 5) with ports in $E_0 = \{1_{\mathcal{L}^k}, 2_{\mathcal{L}^k}\}$, and

- ordered set of edges $E(\mathcal{L}^k) = (l_j)_{j=0}^{2k+1}$ where $l_0 = 1_{\mathcal{L}^k} \in E_0$ and $l_{2k+1} = 2_{\mathcal{L}^k} \in E_0$, and the involution is given by $\tau(l_{2i}) = l_{2i+1}$, for $0 \leq i \leq k$,
- ordered set of k vertices $V(\mathcal{L}^k) = (v_i)_{i=1}^k$, such that $E/v_i = \{l_{2i-1}, l_{2i}\}$ for $1 \leq i \leq k$.

So, \mathcal{L}^k is described by a diagram of the form $\bigcirc \mathbf{2} \amalg 2(\mathbf{k}) \xleftarrow{\quad} 2(\mathbf{k}) \xrightarrow{\quad} \mathbf{k}$.

The line graphs $(\mathcal{L}^k)_{k \in \mathbb{N}}$ may be defined inductively by gluing copies of \mathcal{C}_2 :

$$\begin{aligned} \mathcal{L}^0 &= (i) \\ \mathcal{L}^{k+1} &\cong (\mathcal{L}^k \amalg \mathcal{C}_2)^{2_{\mathcal{L}^k} \ddagger 1_{\mathcal{C}_2}}, \quad k \geq 0. \end{aligned}$$

Example 4.23. For $m \geq 1$, the wheel graph \mathcal{W}^m (illustrated in Figure 5) is the connected bivalent graph with no ports, obtained as the coequaliser in $\mathbf{Grpsh}_f(\mathcal{D})$ of the morphisms $ch_{1_{\mathcal{L}^m}}, ch_{2_{\mathcal{L}^m}} \circ \tau: (i) \rightrightarrows \mathcal{L}^m$. So, $\mathcal{W}^m = \mathcal{L}^{m(1 \ddagger 2)}$ has

- $2m$ cyclically ordered edges $E(\mathcal{W}^m) = (a_j)_{j=1}^{2m}$, such that the involution satisfies $\tau(a_8) = a_1$ and $\tau(a_{2i}) = a_{2i+1}$ for $1 \leq i \leq m-1$,
- m cyclically ordered vertices $V(\mathcal{W}^m) = (v_i)_{i=1}^m$, that $E/v_i = \{a_{2i-1}, a_{2i}\}$ for $1 \leq i \leq m$,

and is described by a diagram of the form $\bigcirc 2(\mathbf{m}) \xleftarrow{\quad} 2(\mathbf{m}) \xrightarrow{\quad} \mathbf{m}$.




FIGURE 5. Line and wheel graphs.

By [41, Proposition 4.23], a connected bivalent graph is isomorphic to \mathcal{L}^k or \mathcal{W}^m for some k, m .

In general, given ports e_1, e_2 of a graph \mathcal{G} such that $e_1 \neq \tau e_2$, then $\mathcal{G}^{e_1 \sharp e_2}$ is formed by “gluing” the ports e_1 and e_2 (see Figure 4). As the following examples shows, the result of forming $\mathcal{G}^{e_1 \sharp e_2}$ is somewhat surprising when $e_1 = \tau e_2$, so e_1, e_2 are the edges of the same stick component:

Example 4.24. Observe that $ch_1^{(i)}: (i) \rightarrow (i)$, $1 \mapsto 1$ is the identity on (i) and $ch_2^{(i)}: (i) \rightarrow (i)$, $1 \mapsto 2$ is precisely τ . Hence $ch_2^{(i)} \circ \tau = \tau^2 = id_i$, and therefore

$$(i)^{1 \sharp 2} = (i).$$

It follows that, if a pair of ports $e_1, e_2 = \tau e_1 \in E_0(\mathcal{G})$ form a stick component of a graph \mathcal{G} (Equation (4.3)), then $\mathcal{G}^{e_1 \sharp e_2} \cong \mathcal{G}$.

Definition 4.25. Following [23], a graph embedding is a morphism $f \in \text{Gr}(\mathcal{G}, \mathcal{H})$ that factors as

$$\mathcal{G} \rightarrow \mathcal{G}^{e_1 \sharp e'_1, \dots, e_k \sharp e'_k} \hookrightarrow \mathcal{H}$$

where the first morphism describes a colimit $\mathcal{G} \rightarrow \mathcal{G}^{e_1 \sharp e'_1, \dots, e_k \sharp e'_k}$ of pairs of parallel maps

$$ch_{e_i}, ch_{e'_i} \circ \tau: (i) \rightrightarrows \mathcal{G},$$

(with $e_1, e'_1, \dots, e_k, e'_k$ a possibly empty set of distinct ports of \mathcal{G}), and the second morphism is a pointwise injection in Gr .

Embeddings form a subcategory of Gr that includes all isomorphisms. They may be thought of as analogous to inclusions of open subsets in topology. The category Gr also admits constructions that play the role of neighbourhoods of points in topology. (See Equation (4.37) for a description of the Grothendieck topology on Gr . This is discussed in more detail in [27, 41].)

Definition 4.26. A neighbourhood of a vertex $v \in V$ (respectively, an edge $e \in E$) is an embedding $u: \mathcal{H} \rightarrow \mathcal{G}$ such that v (respectively e) is in the image of u .

There is a canonical choice of neighbourhood for each vertex v and edge e of a graph \mathcal{G} :

Example 4.27. Let \mathcal{G} be a graph. For each orbit $\tilde{e} \in \tilde{E}$, the inclusion of the stick graph ${}_{\tilde{e}} \stackrel{\text{def}}{=} \{e, \tau e\} \longleftarrow \emptyset \longrightarrow \emptyset$ canonically induces an embedding $\iota_{\tilde{e}}: ({}_{\tilde{e}}) \hookrightarrow \mathcal{G}$.

Recall that, for each $v \in V$, $E/v \stackrel{\text{def}}{=} s(t^{-1}(v))$ is the set of edges incident on v . Let $\mathbf{v} = (E/v)^\dagger$ denote its formal involution.

Then the corolla $\mathcal{C}_{\mathbf{v}}$ is given by

$$\mathcal{C}_{\mathbf{v}} = \bigcirc_{\rightarrow} (E/v \amalg (E/v)^\dagger) \xleftarrow{s} H/v \xrightarrow{t} \{v\}.$$

The inclusion $E/v \hookrightarrow E$ induces an embedding $\iota_v^{\mathcal{G}}$ or $\iota_v: \mathcal{C}_{\mathbf{v}} \rightarrow \mathcal{G}$. Observe that, whenever there is an edge e such that e and τe are both incident on the same vertex v – so $\{e, \tau e\}$ forms a *loop* at v – then ι_v is not injective on edges.

If E/v is empty – so \mathcal{C}_v is an isolated vertex – then $\mathcal{C}_v \hookrightarrow \mathcal{G}$ is a connected component of \mathcal{G} .

For each $v \in V$ (respectively, $e \in E$), an embedding $u: \mathcal{H} \rightarrow \mathcal{G}$ is a neighbourhood of v (respectively e) precisely when $\iota_v: \mathcal{C}_v \rightarrow \mathcal{G}$ (respectively $\iota_{\bar{e}}: (\iota_{\bar{e}}) \hookrightarrow \mathcal{G}$) factors through u .

4.2. Evaluating graphical species on graphs. The assignments $\S \mapsto (\cdot)$, $X \mapsto \mathcal{C}_X$ (for finite sets X) define a full inclusion $\Phi: \mathbf{B}^\S \hookrightarrow \mathbf{Gr}$. Let $\Upsilon: \mathbf{Gr} \rightarrow \mathbf{GS}$ be the induced nerve, given by $\Upsilon\mathcal{G}: \mathcal{C} \mapsto \mathbf{Gr}(\mathcal{C}, \mathcal{G})$ for $\mathcal{C} \in \mathbf{B}^\S$.

Definition 4.28. For $\mathcal{G} \in \mathbf{Gr}$, the category $\mathbf{el}(\Upsilon\mathcal{G}) \cong \mathbf{B}^\S/\mathcal{G}$ is called the category of elements of \mathcal{G} and denoted simply by $\mathbf{el}(\mathcal{G})$.

The following lemma is proved in [41, Lemmas 4.16 & 4.24] and makes precise the intuitive idea that any graph \mathcal{G} is obtained by gluing together edges and corollas (the elements of \mathcal{G}):

Lemma 4.29. For all graphs \mathcal{G} , $\mathcal{G} = \text{colim}_{(\mathcal{C}, b) \in \mathbf{el}(\mathcal{G})} \mathcal{C}$ canonically. In other words, \mathbf{B}^\S is dense in \mathbf{Gr} .

If \mathbf{V} is a category with all small limits, then the pullback functor $\Phi^*: \mathbf{psh}_{\mathbf{V}}(\mathbf{Gr}) \rightarrow \mathbf{GS}_{\mathbf{V}}$ along $\Phi: \mathbf{B}^\S \hookrightarrow \mathbf{Gr}$ has a right adjoint $\Phi_*: \mathbf{GS}_{\mathbf{V}} \hookrightarrow \mathbf{psh}_{\mathbf{V}}(\mathbf{Gr})$ given by

$$(4.30) \quad S \longmapsto (\mathcal{G} \mapsto \lim_{(\mathcal{C}, b) \in \mathbf{el}(\mathcal{G})} S(\mathcal{C})).$$

Since Φ is fully faithful, so is Φ_* [1, Exposé ii]. The same notation will be used for graphical species viewed as (\mathbf{V} -valued) presheaves on \mathbf{B}^\S and for their image as presheaves on \mathbf{Gr} .

Example 4.31. By (4.30), $S(\mathcal{G}) \times S(\mathcal{H}) = S(\mathcal{G} \amalg \mathcal{H})$ for all graphs \mathcal{G} and \mathcal{H} . In particular,

$$(4.32) \quad S_X \times S_Y = S(\mathcal{C}_X) \times S(\mathcal{C}_Y) = S(\mathcal{C}_X \amalg \mathcal{C}_Y) \text{ for all finite sets } X \text{ and } Y.$$

So, an external product \boxtimes on S (Equation (3.21)) may be defined by a collection of equivariant maps $S(\mathcal{C}_X \amalg \mathcal{C}_Y) \rightarrow S_{X \amalg Y}$.

Example 4.33. Let X be a finite set with distinct elements x and y . By (3.16), $S_X^{x \dagger y} \in \mathbf{V}$ is the equaliser of the morphisms $S(ch_x), S(ch_y \circ \tau): S_X \rightrightarrows S_\S$. Since the graph $\mathcal{C}_X^{x \dagger y}$ is defined as the coequaliser of the morphisms $ch_x, ch_y \circ \tau: (\cdot) \rightarrow \mathcal{C}_X$ (see Equation (4.19)), it follows from (4.30) that

$$(4.34) \quad S_X^{x \dagger y} = S(\mathcal{C}_X^{x \dagger y}).$$

Hence, a contraction ζ on a graphical species S in \mathbf{V} (Equation (3.17)) is defined by a collection of maps $S(\mathcal{C}_X^{x \dagger y}) \rightarrow S_{X \setminus \{x, y\}}$ where X is a finite set with distinct elements $x, y \in X$.

Likewise, if X, Y are finite sets with elements $x \in X, y \in Y$, then by (3.12),

$$(4.35) \quad (S_X \times S_Y)^{x \dagger y} = S(\mathcal{D}_{X, Y}^{x \dagger y}) = S(\mathcal{D}_{X, Y}^{x \dagger y}),$$

since $\mathcal{D}_{X, Y}^{x \dagger y}$ is the coequaliser of the morphisms $ch_x, ch_y \circ \tau: (\cdot) \rightarrow \mathcal{C}_X \amalg \mathcal{C}_Y$.

So, a multiplication \diamond on S (Equation (3.14)) is defined by a collection of maps $S(\mathcal{D}_{X, Y}^{x \dagger y}) \rightarrow S_{X \amalg Y \setminus \{x, y\}}$ where X, Y are finite sets and $x \in X, y \in Y$.

Definition 4.36. When $\mathbf{V} = \mathbf{Set}$ and $S \in \mathbf{GS}$, an element $\alpha \in S(\mathcal{G})$ is called an S -structure on \mathcal{G} . Objects of the category $\mathbf{Gr}(S)$, of S -structured graphs, are pairs (\mathcal{G}, α) with $\alpha \in S(\mathcal{G})$, and morphisms $(\mathcal{G}, \alpha) \rightarrow (\mathcal{G}', \alpha')$ in $\mathbf{Gr}(S)$ are given by morphisms $f \in \mathbf{Gr}(\mathcal{G}, \mathcal{G}')$ such that $S(f)(\alpha') = \alpha$.

By Equation (4.29), \mathbf{Gr} may be viewed as a full subcategory of \mathbf{GS} and for graphs $\mathcal{G} \in \mathbf{Gr}$, I will write $\mathcal{G} \in \mathbf{GS}$, rather than $\Upsilon\mathcal{G} \in \mathbf{GS}$ where there is no risk of confusion.

Remark 4.37. Let \mathcal{G} be a graph. Families of jointly surjective morphisms define the covers at \mathcal{G} for a canonical *étale topology* J on \mathbf{Gr} . Since $\mathrm{el}(\mathcal{G})$ refines every cover at \mathcal{G} , the category \mathbf{GS} of graphical species in \mathbf{Set} is equivalent to the category $\mathbf{sh}(\mathbf{Gr}, J)$ of étale sheaves on \mathbf{Gr} . It follows, in particular, that there is a canonical isomorphism $S(\mathcal{G}) \cong \mathbf{GS}(\Upsilon\mathcal{G}, S)$. Hence, $\mathbf{Gr}(S)$ is canonically isomorphic to the slice category \mathbf{Gr}/S .

Example 4.38. The terminal oriented graphical species $Di \in \mathbf{GS}$ was defined in Equation (3.7). By Equation (4.37), a Di -structure $\phi \in \mathbf{GS}(\mathcal{G}, Di)$ on a graph \mathcal{G} is given by a bijection $\{e, \tau e\} \xrightarrow{\cong} \mathfrak{D}i$ for each edge orbit $\{e, \tau e\}$ of \mathcal{G} . In other words, ϕ defines an orientation on \mathcal{G} . Hence, the *category* $\mathbf{OGr} \cong \mathbf{Gr}(Di)$ of *directed graphs* is defined as the slice category \mathbf{Gr}/Di . This is a full subcategory of $\mathbf{OGS} = \mathbf{GS}/Di$ by Equation (4.29) and each directed graph $\tilde{\mathcal{G}} = (\mathcal{G}, \phi) \in \mathbf{OGr}$ is canonically the colimit of its induced oriented element category $\mathrm{el}_{Di}(\tilde{\mathcal{G}})$ obtained by pulling back the orientation ϕ on \mathcal{G} along each element $(\mathcal{C}, b) \in \mathrm{el}(\mathcal{G})$. (See also [41, Section 4.5].)

In particular, if $\tilde{S}: \mathrm{el}(Di)^{\mathrm{op}} \rightarrow \mathbf{V}$ is an oriented graphical species in \mathbf{V} corresponding to a pair $(S, \gamma) \in \mathbf{GS}_{\mathbf{V}}/Di^{\mathbf{V}}$, then \tilde{S} extends to a presheaf on \mathbf{OGr} such that, for all graphs $\mathcal{G} \in \mathbf{Gr}$,

$$S(\mathcal{G}) \cong \coprod_{\phi \in \mathfrak{D}i(\mathcal{G})} \tilde{S}(\mathcal{G}, \phi).$$

4.3. Relationship between graphs and Brauer diagrams and wiring diagrams. In Equation (7.9), circuit algebras internal to a category \mathbf{V} will be constructed as algebras for a composite monad \mathbb{LDT} on $\mathbf{GS}_{\mathbf{V}}$ that is described in terms of graphs in \mathbf{Gr} and, by Equation (8.4), circuit algebras in \mathbf{Set} are presheaves over \mathbf{Gr} satisfying additional properties. However, (enriched) circuit algebras are usually defined as algebras over an operad of wiring diagrams (see e.g., [3, 12, 14, 25, 49] and [42]).

In [42, Section 4], the operad \mathbf{WD} of (monochrome unoriented) wiring diagrams is defined using the symmetric monoidal category \mathbf{BD} of Brauer diagrams, familiar in the representation theory of classical groups [34, 42, 44]. In fact, (enriched) circuit algebras may equivalently be described by symmetric lax monoidal functors from categories of (coloured) Brauer diagrams.

A partial overview of how different variations of circuit algebras are associated to classical groups, and how they relate to the composite circuit algebra monad \mathbb{LDT} is provided in [42, Introduction]. This short section reviews the definitions of \mathbf{BD} and \mathbf{WD} and describes how wiring diagrams are related to graphs as in Equation (4.1).

Objects of \mathbf{BD} are natural numbers. Morphisms are (*monochrome, unoriented*) *Brauer diagrams*, where a Brauer diagram in $f: m \rightarrow n$ is a pair $f = (\tau, \mathfrak{k})$ such that $\mathfrak{k} \in \mathbb{N}_{\geq 0}$ is a natural number and τ is a fixed-point free involution (or *matching*) on $\mathbf{m} \amalg \mathbf{n}$ (as usual, $\mathbf{n} = \{1, \dots, n\}$ for all $n \geq 1$ and $\mathbf{0} = \emptyset$).

A Brauer diagram $(\tau, \mathfrak{k}): m \rightarrow n$ may be visualised, as in Figure 6 (b) by \mathfrak{k} closed loops next to a univalent graph that comprises of a row of n vertices bijectively labelled by \mathbf{n} above a row of m vertices labelled by \mathbf{m} , such that there is an edge connecting vertices x and y in $\mathbf{m} \amalg \mathbf{n}$ if and only if $x = \tau y$. The monoidal product \oplus on \mathbf{BD} is given by $f_1 \oplus f_2 = (\tau_1 \amalg \tau_2, \mathfrak{k}_1 + \mathfrak{k}_2) \in \mathbf{BD}(m_1 + m_2, n_1 + n_2)$ for $f_1 = (\tau_1, \mathfrak{k}_1) \in \mathbf{BD}(m_1, n_1)$ and $f_2 = (\tau_2, \mathfrak{k}_2) \in \mathbf{BD}(m_2, n_2)$. It is represented graphically by juxtaposition. Categorical composition is described by “diagram stacking” as in Figure 6 (see also [42, Section 3]).

The operad \mathbf{WD} is the \mathbb{N} -coloured *operad underlying* \mathbf{BD} . This means that, for each tuple of natural numbers $(m_1, \dots, m_k; n)$, the set of (*monochrome*) *wiring diagrams of type* $(m_1, \dots, m_k; n)$ is given by $\mathbf{WD}(m_1, \dots, m_k; n) \stackrel{\mathrm{def}}{=} \mathbf{BD}(\sum_{i=1}^k m_i; n)$. For $f \in \mathbf{BD}(\sum_{i=1}^k m_i; n)$, the corresponding wiring diagram is denoted by $\overline{f} \in \mathbf{WD}(m_1, \dots, m_k; n)$. The (strict) symmetric monoidal category structure on \mathbf{BD}

makes WD into an \mathbf{N} -coloured operad (see [42, Section 4]) with composition

$$\gamma: \text{WD}(m_1, \dots, m_k; n) \otimes \bigotimes_{i=1}^k \text{WD}(l_{i,1}, \dots, l_{i,k_i}; m_i) \rightarrow \text{WD}(l_{1,1}, \dots, l_{k,k_i}; n)$$

given by

$$\gamma(\bar{g}, (\bar{f}_i)_i) \stackrel{\text{def}}{=} \overline{(g \circ (f_1 \oplus \dots \oplus f_n))}.$$

It follows, in particular, that an algebra over WD (a monochrome circuit algebra) is, equivalently, a lax monoidal functor from BD (see [42, Sections 3 & 4]).

A monochrome wiring diagram $\bar{f} \in \text{WD}(m_1, \dots, m_k; n)$ induced by an Brauer diagram $f \in \text{BD}(\sum_{i=1}^k m_i, n)$ of the form $f = (\tau_f, 0)$ describes a graph:

$$(4.39) \quad \tau_f \bigcirc_{\uparrow} (\mathbf{n} \amalg (\coprod_{i=1}^k \mathbf{m}_i)) \xleftarrow{\text{inc}} \coprod_{i=1}^k \mathbf{m}_i \xrightarrow{p} \mathbf{k}$$

where $p: \coprod_{i=1}^k \mathbf{m}_i \rightarrow \mathbf{k}$ is the canonical projection.




FIGURE 6. (a) Composition of Brauer diagrams $f = (\tau_f, 0) \in \text{BD}(5, 7)$ and $g = (\tau_g, 0) \in \text{BD}(7, 3)$ (b) the resulting Brauer diagram $gf = (\tau_g \tau_f, 1) \in \text{BD}(5, 3)$ with a single loop formed in the composition.

Remark 4.40. Wiring diagrams are, essentially, symmetric versions of Jones's *planar diagrams* [26]. As such, they are commonly represented, not by Brauer diagrams, but in terms of immersions of compact 1-manifolds in punctured discs as in Figure 7 (see e.g., [13] and [42, Section 4.2] for details).

Consider the punctured disc representation of a wiring diagram \bar{f} as in Figure 7. It is clear that, by shrinking the punctures to points, and deleting the outer disc, we obtain a representation of a graph with k vertices of degree m_1, \dots, m_k and n loose edges. This is homeomorphic to the geometric realisation (Equation (4.6)) of the Feynman graph (4.39) corresponding to \bar{f} .

Example 4.41. When $X = \mathbf{n}$, the corolla \mathcal{C}_X defined in Equation (4.8) corresponds by (4.39) to the identity wiring diagram $\overline{id_n} \in \text{WD}(\mathbf{n}; \mathbf{n})$.

By [34, Theorem 2.6] or [2, Proposition 2.15], the category BD is generated, under composition and monoidal product, by the Brauer diagrams $\sigma_2 \in \text{BD}(2, 2)$ (the non-identity bijection between source and target sets that generates the symmetry in BD) and $id_1 \in \text{BD}(1, 1)$, $\cap \in \text{BD}(2, 0)$ and $\cup \in \text{BD}(0, 2)$.

Example 4.42. With the exception of σ_2 , each of the generating morphisms of BD has the form $(\tau_2, 0)$, where τ_2 is the unique matching on the two-element set. Hence τ_2 induces distinct wiring diagrams $\overline{id_1} \in \text{WD}(1; 1)$, $\cup \in \text{WD}(-; 2)$ and $\cap_{(2; 0)} \in \text{WD}(2; 0)$ and $\cap_{(1, 1; 0)} \in \text{WD}(1, 1; 0)$. By (4.39), these describe the four graphs with a single edge orbit from Equation (4.7):

- (a) for $\overline{id_1} \in \text{WD}(1; 1)$, the corresponding graph is the **1-corolla** \mathcal{C}_1 (Figure 3(a));
- (b) for $\cup \in \text{WD}(-; 2)$, the corresponding graph is the *stick graph* (i) (Figure 3(b));
- (c) for $\cap_{(2; 0)} \in \text{WD}(2; 0)$, the corresponding graph is the *wheel graph* \mathcal{W} (Figure 3(c));
- (d) for $\cap_{(1, 1; 0)} \in \text{WD}(1, 1; 0)$, the corresponding graph is given by Equation (4.7)(d) (Figure 3(d)).




FIGURE 7. Brauer diagram and punctured disc representations of the same composition of wiring diagrams.

Inclusions of connected components in \mathbf{Gr} describe factors of monoidal products in \mathbf{BD} . But, in general, morphisms in \mathbf{Gr} do not admit a nice description in terms of \mathbf{BD} (or \mathbf{WD}). Nonetheless, since the category \mathbf{CA} of circuit algebras in \mathbf{Set} is equivalent to the category \mathbf{CA} described in [42], and objects of \mathbf{CA} are algebras over operads of wiring diagrams [42, Section 4], there must be constructions on graphs that play the role of operadic composition in \mathbf{WD} . These will be described in the next section (Section 5.1).

Remark 4.43. The (internal) construction of circuit algebras discussed in this work characterises *all* circuit algebras as algebras for a single monad on \mathbf{GS}_V , with special cases, such as wheeled props (oriented circuit algebras) obtained as subcategories or via slice constructions. By contrast, different versions of (enriched) circuit algebras – such as oriented or nonoriented – are algebras over distinct (different coloured) operads of wiring diagrams.

New categories of Brauer diagrams – and hence new operads of wiring diagrams – may be obtained by adding colours to the connected components of the underlying graphs as in [42, Section 3.2]. In particular, the category \mathbf{OBD} of *oriented monochrome diagrams*, whose objects are elements of the monoid $\text{list}\{\text{in}, \text{out}\}$, may be defined similarly to \mathbf{BD} by orienting the edges of the Brauer diagrams. (See [42, Section 3.2] for a precise definition in terms of the palette $(\mathfrak{Di}, \sigma_{\mathfrak{Di}})$ defined in Equation (3.7).) This category is important in the representation theory of general linear groups [34, 42, 44].

Remark 4.44. Extending the Brauer’s 1937 work [8], categories of (oriented) Brauer diagrams have been used to study the representation theory of the (general linear) orthogonal and symplectic groups (see, e.g., [34, 43, 44]).

Building on the wheeled prop characterisation of algebras for the general linear groups in [15], [42, Theorem 6.3] provides characterises algebras over the finite dimensional orthogonal and symplectic groups as (enriched) circuit algebras satisfying two simple relations.

Remark 4.45. (See also Equation (5.3)) The subcategory $\mathbf{dBD} \subset \mathbf{BD}$ of *downward Brauer diagrams* is generated by $\sigma_2 = (\{s_1 \mapsto t_2, s_2 \mapsto t_1\}, 0) \in \mathbf{BD}(2, 2)$, $id_1 \in \mathbf{BD}(1, 1)$ and $\cap \in \mathbf{BD}(2, 0)$ (but not $\cup \in \mathbf{BD}(0, 2)$). Morphisms in $\mathbf{dBD}(m, n)$ have the form $f = (\tau, 0)$ where, for all $t_i \in \mathfrak{T}(f)$, $\tau(t_i) \in Sf$. In particular, Brauer diagrams in \mathbf{dBD} are “loop-free”.

By [44] finite-length (vector-space valued) functors from \mathbf{dBD} are equivalent to representations of the infinite dimensional orthogonal (or symplectic) group, and representations of the infinite dimensional general linear group are described by functors from the category of downward *oriented* Brauer diagrams.

In [42, Section 4], nonunital circuit algebras (i.e., $\mathbb{L}\mathbb{T}$ -algebras) are defined as lax symmetric monoidal functors from (coloured versions of) \mathbf{dBD} . In particular, the categories of algebras for the infinite orthogonal, symplectic and general linear groups admit descriptions in terms of nonunital circuit algebras. (See the discussions in [42, Introduction & Section 6.2], and [42, Theorem 6.13] in particular.)

5. A MONAD FOR NONUNITAL CIRCUIT ALGEBRAS

This section gives a short outline, closely related to [27] and [41, Section 5], of the construction of the nonunital circuit algebra monad \mathbb{T}^\times on \mathbf{GS}_V where, as usual, the category V is assumed to have enough limits and colimits for all the given constructions.

5.1. Gluing constructions and labelled graphs. As shown in Equation (4.16), the graph category \mathbf{Gr} is not closed under finite colimits. However, there is a class of diagrams in \mathbf{Gr} that always admit colimits, and are central to the definition of the nonunital circuit algebra monad \mathbb{T}^\times in Section 5.2.

The following terminology is based on [30]:

Definition 5.1. Let \mathcal{G} be a graph. A \mathcal{G} -shaped graph of graphs is a functor $\Gamma: \mathbf{el}(\mathcal{G}) \rightarrow \mathbf{Gr}$ (or $\Gamma^\mathcal{G}$) such that

$$\begin{aligned} \Gamma(a) &= ({}_1) && \text{for all } ({}_1, a) \in \mathbf{el}(\mathcal{G}), \\ E_0(\Gamma(b)) &= X && \text{for all } (C_X, b) \in \mathbf{el}(\mathcal{G}), \end{aligned}$$

and, for all $(C_{X_b}, b) \in \mathbf{el}(\mathcal{G})$ and all $x \in X_b$,

$$\Gamma(ch_x) = ch_x^{\Gamma(b)} \in \mathbf{Gr}({}_1, \Gamma(b)).$$

A \mathcal{G} -shaped graph of graphs $\Gamma: \mathbf{el}(\mathcal{G}) \rightarrow \mathbf{Gr}$ is *nondegenerate* if, for all $v \in V$, $\Gamma(\iota_v)$ has no stick components. Otherwise, Γ is called *degenerate*.

In particular, the empty \mathcal{C}_0 -indexed graph of graphs $\mathcal{C}_0 \mapsto \emptyset$ is nondegenerate.

A nondegenerate \mathcal{G} -shaped graph of graphs may be viewed informally as a rule for substituting graphs into vertices of \mathcal{G} as in Figure 8.

Proposition 5.2. A \mathcal{G} -shaped graph of graphs Γ admits a colimit $\Gamma(\mathcal{G})$ in \mathbf{Gr} , and, for each $(C, b) \in \mathbf{el}(\mathcal{G})$, the universal map $b_\Gamma: \Gamma(b) \rightarrow \Gamma(\mathcal{G})$ is an embedding in \mathbf{Gr} .

If Γ is nondegenerate and $\Gamma(b) \neq \emptyset$ for all $(C, b) \in \mathbf{el}(\mathcal{G})$, then

- Γ describes a bijection $E(\Gamma(\mathcal{G})) \cong E(\mathcal{G}) \amalg \coprod_{v \in V} E_\bullet(\Gamma(\iota_v))$ of sets;
- Γ induces an identity $E_0(\mathcal{G}) \xrightarrow{=} E_0(\Gamma(\mathcal{G}))$ of ports;
- there is a canonical surjective map $V(\Gamma(\mathcal{G})) \twoheadrightarrow V(\mathcal{G})$ induced by the embeddings $\Gamma(\iota_v): \Gamma(\iota_v) \rightarrow \Gamma(\mathcal{G})$ for each vertex v of \mathcal{G} .




FIGURE 8. A \mathcal{G} shaped graph of graphs $\mathbf{\Gamma}$ describes *graph substitution* in which each vertex v of \mathcal{G} is replaced by a graph \mathcal{G}_v according to a bijection $E_0(\mathcal{G}_v) \xrightarrow{\cong} (E/v)^\dagger$. If $\mathbf{\Gamma}$ is nondegenerate, then taking its colimit corresponds to erasing the inner (blue) nesting.

Proof. If $\mathbf{\Gamma}$ is nondegenerate, then $\mathbf{\Gamma}$ admits a colimit $\mathbf{\Gamma}(\mathcal{G})$ in \mathbf{Gr} by [41, Proposition 5.16]. In case $\mathbf{\Gamma}$ is degenerate, the existence of a colimit in \mathbf{Gr} follows from [41, Proposition 7.15].

The other statements follow directly from [41, Corollary 5.19]. \square

Remark 5.3. (See also Equation (4.45).) By Equation (3.27), there is an equivalence of categories between (internal) circuit algebras in \mathbf{Set} and algebras for operads of wiring diagrams. Under this equivalence, graphs of graphs play the role of operadic composition of wiring diagrams.

Let \mathcal{G} be a graph with n ports, k vertices, and such that each vertex $v_i \in V$ has $m_i = |E/v_i|$ adjacent edges. Up to orderings, \mathcal{G} describes a wiring diagram $\bar{g} \in \mathbf{WD}(\mathbf{m}_1, \dots, \mathbf{m}_k; \mathbf{n})$ with no closed components. A \mathcal{G} -shaped graph of graphs $\mathbf{\Gamma}$ such that $\mathbf{\Gamma}(b) \neq \emptyset$ for all $(\mathcal{C}, b) \in \mathbf{el}(\mathcal{G})$ describes a choice of k wiring diagrams $\bar{f}^i \in \mathbf{WD}(l_1, \dots, l_{k_i}; \mathbf{m}_i)$ for $(l_1, \dots, l_{k_i}) \in \mathbf{list}(\mathbb{N})$ and $1 \leq i \leq k$.

Non-degeneracy of $\mathbf{\Gamma}$ corresponds to the condition, discussed in [42, Remark 3.12], that, for each \bar{f}^i in the composition, the underlying morphism f^i of Brauer diagrams is a morphism of the downward Brauer category $f^i \in \mathbf{dBD}(l_1 + \dots + l_{k_i}, m_i)$. If $\mathbf{\Gamma}$ is nondegenerate, then its colimit $\mathbf{\Gamma}(\mathcal{G})$ is described by the composition $\gamma(\bar{g}, (\bar{f}_i)_i)$ in \mathbf{WD} . (Colimits of degenerate graphs of graphs are discussed in Section 6.3.)

Example 5.4. By Equation (4.29), every graph \mathcal{G} is the colimit of the (nondegenerate) *identity \mathcal{G} -shaped graph of graphs* $\mathbf{I}^{\mathcal{G}}$ given by the forgetful functor $\mathbf{el}(\mathcal{G}) \rightarrow \mathbf{Gr}$, $(\mathcal{C}, b) \mapsto \mathcal{C}$.

The identity \mathcal{G} -shaped graph of graphs corresponds, as in Equation (5.3) to a composition of wiring diagrams of the form $\gamma(\bar{g}, (\bar{id}_{m_i})_{i=1}^k)$.

Definition 5.5. The category $\mathbf{Gr}^{(\mathcal{G})}$ of nondegenerate \mathcal{G} -shaped graphs of graphs is the full subcategory of nondegenerate \mathcal{G} -shaped graphs of graphs $\mathbf{\Gamma}$ in the category of functors $\mathbf{el}(\mathcal{G}) \rightarrow \mathbf{Gr}$.

Definition 5.6. Let X be a finite set. A generalised X -graph (henceforth X -graph) is a \mathcal{C}_X -shaped graph of graphs. An X -graph is admissible if it is nondegenerate.

The maximal subgroupoid $\mathbf{Core}(\mathbf{Gr}^{(\mathcal{C}_X)}) \subset \mathbf{Gr}^{(\mathcal{C}_X)}$ is called the groupoid of (admissible) X -graphs and X -isomorphisms and is denoted by $X\text{-Gr}_{\text{iso}}$.

More generally, for any graph \mathcal{G} and any $(\mathcal{C}, b) \in \mathbf{el}(\mathcal{G})$, $[b]\text{-Gr}_{\text{iso}}$ denotes the groupoid of nondegenerate \mathcal{C} -shaped graphs of graphs and isomorphisms.

Hence, a generalised X -graph is given by a pair $\mathcal{X} = (\mathcal{G}, \rho)$, where \mathcal{G} is a graph and $\rho: E_0 \xrightarrow{\cong} X$. An X -graph is *admissible* if it has no stick components, and an X -isomorphism in $X\text{-Gr}_{\text{iso}}(\mathcal{X}, \mathcal{X}')$ is an isomorphism $g \in \text{CGr}(\mathcal{G}, \mathcal{G}')$ that preserves the X -labelling: $\rho' \circ g_{E_0} = \rho: E_0 \rightarrow X$.

Example 5.7. For any graph \mathcal{G} , a (nondegenerate) \mathcal{G} -shaped graph of graphs $\mathbf{\Gamma}: \text{el}(\mathcal{G}) \rightarrow \text{Gr}$ is precisely a choice of (admissible) X_b -graph \mathcal{X}_b for each $(\mathcal{C}_{X_b}, b) \in \text{el}(\mathcal{G})$.

Let S be a graphical species in \mathbf{V} , and \mathcal{G} a graph. If $\mathbf{\Gamma}: \text{el}(\mathcal{G}) \rightarrow \text{Gr}$ is a nondegenerate \mathcal{G} -shaped graph of graphs with colimit $\mathbf{\Gamma}(\mathcal{G})$ in Gr , then the composition

$$\text{el}(\mathcal{G})^{\text{op}} \xrightarrow{\mathbf{\Gamma}^{\text{op}}} \text{Gr}^{\text{op}} \xrightarrow{S} \mathbf{V}$$

defines a functor $\text{el}(\mathcal{G})^{\text{op}} \rightarrow \mathbf{V}$ with limit $S(\mathbf{\Gamma}(\mathcal{G}))$.

Example 5.8. Recall Equation (4.36). If S is a graphical species in Set and \mathcal{G} is any graph, then a nondegenerate \mathcal{G} -shaped graph of graphs $\mathbf{\Gamma}$ and choice of S -structure $\alpha \in S(\mathbf{\Gamma}(\mathcal{G}))$ determines a functor $\text{el}(\mathcal{G}) \rightarrow \text{Gr}(S)$ – called a *\mathcal{G} -shaped graph of S -structured graphs* – by $b \mapsto S(b_{\mathbf{\Gamma}})(\alpha)$ (where, for all $(\mathcal{C}, b) \in \text{el}(\mathcal{G})$, $b_{\mathbf{\Gamma}}$ is the universal map described in Equation (5.2)).

5.2. Nonunital circuit algebras. It is now straightforward to construct a monad $\mathbb{T}^\times = (T^\times, \mu^{\mathbb{T}^\times}, \eta^{\mathbb{T}^\times})$ on $\text{GS}_{\mathbf{V}}$, whose algebras are nonunital circuit algebras in \mathbf{V} .

The underlying endofunctor $T^\times = T^\times_{\mathbf{V}}: \text{GS}_{\mathbf{V}} \rightarrow \text{GS}_{\mathbf{V}}$ is defined, for all $S \in \text{GS}_{\mathbf{V}}$, by

$$(5.9) \quad \begin{aligned} T^\times S_{\S} &= S_{\S}, \\ T^\times S_X &= \text{colim}_{\mathcal{X} \in X\text{-Gr}_{\text{iso}}} S(\mathcal{X}) \quad \text{for all finite sets } X. \end{aligned}$$

This acts on isomorphisms in \mathbf{B}^{\S} , $T^\times S$ acts by relabelling graph ports. Let $g: \mathcal{X}' \rightarrow \mathcal{X}$ be a port-preserving isomorphism in $X\text{-Gr}_{\text{iso}}$, and for each $x \in X$, let $ch_x^{\mathcal{X}} \in \text{CGr}(\mathbf{1}, \mathcal{G})$ be the map $ch_{\rho^{-1}(x)}$ defined by $1 \mapsto \rho^{-1}(x) \in E_0(\mathcal{G})$. Then,

$$S(g \circ ch_x^{\mathcal{X}}) = S(ch_x^{\mathcal{X}})S(g) = S(ch_x^{\mathcal{X}'}).$$

Hence, the projections $T^\times S(ch_x)$ induced by $S(ch_x^{\mathcal{X}}): S(\mathcal{X}) \rightarrow S(\mathbf{1}) = S_{\S}$, are well-defined. So $T^\times S$ extends to all morphisms in \mathbf{B}^{\S} and therefore defines a graphical species in \mathbf{V} . The assignment $S \mapsto T^\times S$ is clearly natural on $S \in \text{GS}_{\mathbf{V}}$, and so T^\times defines an endofunctor on $\text{GS}_{\mathbf{V}}$.

As in Equation (5.2), for an X -graph \mathcal{X} and nondegenerate \mathcal{X} -shaped graph of graphs $\mathbf{\Gamma}$ in $\text{Gr}^{(\mathcal{X})}$, let $\mathbf{\Gamma}(\mathcal{X})$ denote the colimit of $\mathbf{\Gamma}$ in $\text{Gr}^{(\mathcal{X})}$.

Lemma 5.10. *For any graphical species S in \mathbf{V} and any finite set X , there is a canonical isomorphism*

$$(T^\times)^2 S_X \xrightarrow{\cong} \text{colim}_{\mathcal{X} \in X\text{-Gr}_{\text{iso}}} \left(\lim_{\mathbf{\Gamma} \in \text{Gr}^{(\mathcal{X})}} S(\mathbf{\Gamma}(\mathcal{X})) \right).$$

Proof. By definition, for all finite sets X ,

$$\begin{aligned} (T^\times)^2 S_X &= \text{colim}_{\mathcal{X} \in X\text{-Gr}_{\text{iso}}} T^\times S(\mathcal{X}) \\ &= \text{colim}_{\mathcal{X} \in X\text{-Gr}_{\text{iso}}} \lim_{(\mathcal{C}, b) \in \text{el}(\mathcal{X})} T^\times S(\mathcal{C}) \\ &= \text{colim}_{\mathcal{X} \in X\text{-Gr}_{\text{iso}}} \lim_{(\mathcal{C}, b) \in \text{el}(\mathcal{X})} \text{colim}_{\mathcal{Y} \in [b]\text{-Gr}_{\text{iso}}} S(\mathcal{Y}). \end{aligned}$$

Since $[b]\text{-Gr}_{\text{iso}}$ is a groupoid by definition, $\text{colim}_{\mathcal{Y} \in [b]\text{-Gr}_{\text{iso}}} S(\mathcal{Y}) = \lim_{\mathcal{Y} \in [b]\text{-Gr}_{\text{iso}}} S(\mathcal{Y})$ for all $(\mathcal{C}, b) \in \text{el}(\mathcal{X})$. By Equation (5.5),

$$\lim_{\mathbf{\Gamma} \in \text{Gr}^{(\mathcal{X})}} S(\mathbf{\Gamma}(\mathcal{X})) \cong \lim_{(\mathcal{C}, b) \in \text{el}(\mathcal{X})} \left(\lim_{\mathcal{Y} \in [b]\text{-Gr}_{\text{iso}}} S(\mathcal{Y}) \right)$$

canonically, and therefore, by the universal property of (co)limits, there is a unique isomorphism

$$(5.11) \quad (T^\times)^2 S_X \cong \text{colim}_{\mathcal{X} \in X\text{-Gr}_{\text{iso}}} \left(\lim_{\mathbf{\Gamma} \in \text{Gr}^{(\mathcal{X})}} S(\mathbf{\Gamma}(\mathcal{X})) \right).$$

□

By Equation (5.2), each $\Gamma(\mathcal{X})$ inherits the structure of an X -graph from \mathcal{X} . Hence, by an application of Equation (5.10), the assignment $(\mathcal{X}, \Gamma) \rightarrow \Gamma(\mathcal{X})$ describes a \mathbf{V} -morphism $\mu^{\mathbb{T}^\times} S_X \in E((T^\times)^2 S_X, T^\times S_X)$ in \mathbf{V} given by

$$(5.12) \quad \mu^{\mathbb{T}^\times} S_X : (T^\times)^2 S_X = \text{colim}_{\mathcal{X} \in X\text{-Gr}_{\text{iso}}} (\lim_{\Gamma \in \text{Gr}(\mathcal{X})} S(\Gamma(\mathcal{X}))) \rightarrow \text{colim}_{\mathcal{X}' \in X\text{-Gr}_{\text{iso}}} S(\mathcal{X}') = T^\times S_X.$$

This is natural in X and in S and hence defines a natural transformation $\mu^{\mathbb{T}^\times} : (T^\times)^2 \Rightarrow T^\times$.

The inclusion $\mathbf{B}^{\mathfrak{s}} \hookrightarrow \mathbf{Gr}$ induces a natural transformation $\eta^{\mathbb{T}^\times} : id_{\mathbf{GS}} \Rightarrow T^\times$, $S_X \xrightarrow{\eta^{\mathbb{T}^\times}} S(\mathcal{C}_X) \rightarrow T^\times S_X$ that provides a unit for the endofunctor T^\times on $\mathbf{GS}_{\mathbf{V}}$. It is straightforward to verify that the triple $\mathbb{T}^\times = (T^\times, \mu^{\mathbb{T}^\times}, \eta^{\mathbb{T}^\times})$ defines a monad.

Example 5.13. For $S \in \mathbf{GS}_{\mathbf{V}}$, $T^\times S$ admits a canonical nonunital circuit algebra structure:

Disjoint union of graphs induces an external product \boxtimes on $T^\times S$. For, let X, Y be finite sets and let \mathcal{X} be an X -graph and \mathcal{Y} a Y -graph. Then, the following diagram – in which the unlabelled maps are the defining morphisms for T^\times – commutes in \mathbf{V} and describes an external product \boxtimes on $T^\times S$:

$$\begin{array}{ccccccc} S(\mathcal{X} \amalg \mathcal{Y}) & \xlongequal{\quad} & S(\mathcal{X}) \times S(\mathcal{Y}) & \longrightarrow & T^\times S_X \times T^\times S_Y & \xlongequal{\quad} & T^\times S(\mathcal{C}_X \amalg \mathcal{C}_Y) \longrightarrow (T^\times)^2 S_{X \amalg Y} \\ & & & & \downarrow \boxtimes & & \downarrow \mu^{\mathbb{T}^\times} \\ & & & & T^\times S_{X \amalg Y} & & \end{array}$$

Since $\emptyset \in \mathbf{0}\text{-Gr}_{\text{iso}}$, there is a morphism $* = S(\emptyset) \rightarrow T^\times S_{\mathbf{0}}$ from the terminal object $*$ in \mathbf{V} . This provides an external unit for \boxtimes .

Similarly, the contraction ζ on $T^\times S$ is induced by commuting diagrams of the form

$$\begin{array}{ccccc} S(\mathcal{X}^{x \ddagger y}) & \longrightarrow & T^\times S(\mathcal{C}_X^{x \ddagger y}) & \longrightarrow & (T^\times)^2 S_{X \setminus \{x, y\}} \\ & & \downarrow \zeta_X^{x \ddagger y} & & \downarrow \mu^{\mathbb{T}^\times} \\ & & T^\times S_{X \setminus \{x, y\}} & & \end{array}$$

where X is a finite set with distinct elements $x, y \in X$, and \mathcal{X} an X -graph.

It follows immediately from the definitions (and the diagrams in the proof of Equation (5.14) below) that \boxtimes and ζ satisfy (C1)-(C3) of Equation (3.25).

Proposition 5.14. *There is a canonical isomorphism of categories $\mathbf{GS}_{\mathbf{V}}^{\mathbb{T}^\times} \cong \mathfrak{CA}_{\mathbf{V}}^-$.*

Proof. Let (A, h) be an algebra for \mathbb{T}^\times .

For finite sets X and Y , the morphism $\boxtimes_{X,Y} : A_X \times A_Y \rightarrow A_{X \amalg Y}$ in \mathbf{V} is obtained by composing the structure morphism $A_X \times A_Y = A(\mathcal{C}_X \amalg \mathcal{C}_Y) \rightarrow T^\times A_{X \amalg Y}$ with $h : T^\times A_{X \amalg Y} \rightarrow A_{X \amalg Y}$. Moreover, the composition $* = A(\emptyset) \rightarrow T^\times A_{\mathbf{0}} \xrightarrow{h} A_{\mathbf{0}}$ provides a unit for the external product \boxtimes so defined.

If a finite set X has distinct elements x and y , then we may define a morphism $\zeta_X^{x \ddagger y} : A_X \rightarrow A_{X \setminus \{x, y\}}$ in \mathbf{V} by composing $A_X^{x \ddagger y} = A(\mathcal{C}_X^{x \ddagger y}) \rightarrow T^\times A_{X \setminus \{x, y\}}$ with h .

The proof that (A, \boxtimes, ζ) satisfies the axioms for a circuit algebra proceeds by representing each side of the equations in (C1)-(C3) of Equation (3.25) in terms of graph of graphs, and then showing that these have the same colimit in \mathbf{Gr} .

For (C1), the external product $\boxtimes_{X,Y}$ is represented by the graph $\mathcal{C}_X \amalg \mathcal{C}_Y$, and so $\boxtimes_{X \amalg Y, Z} \circ (\boxtimes_{X,Y} \times id_{A_Z})$ is represented by the $\mathcal{C}_{X \amalg Y} \amalg \mathcal{C}_Z$ -shaped graphs of graphs $(\iota_{X \amalg Y} \mapsto \mathcal{C}_X \amalg \mathcal{C}_Y, \iota_Z \mapsto \mathcal{C}_Z)$. This has colimit $\mathcal{C}_X \amalg \mathcal{C}_Y \amalg \mathcal{C}_Z$ which is also the colimit of the $\mathcal{C}_X \amalg \mathcal{C}_{Y \amalg Z}$ -shaped graph of graphs $(\iota_X \mapsto \mathcal{C}_X, \iota_{Y \amalg Z} \mapsto \mathcal{C}_Y \amalg \mathcal{C}_Z)$.

More precisely, it follows from the monad algebra axioms that, for all finite sets X, Y and Z , the following diagram, in which the maps $A_{X \amalg Y} \times A_Z \rightarrow A_{X \amalg Y \amalg Z} \leftarrow A_X \times A_{Y \amalg Z}$ are just $\boxtimes_{X \amalg Y, Z}$ and $\boxtimes_{X, Y \amalg Z}$, commutes:

$$\begin{array}{c}
 A_X \times A_Y \times A_Z \\
 \swarrow = \quad \downarrow = \quad \searrow = \\
 A_{X \amalg Y} \times A_Z \xleftarrow{(\boxtimes, id)} A(C_X \amalg C_Y) \times A(C_Z) \xrightarrow{=} A(C_X \amalg C_Y \amalg C_Z) \xleftarrow{=} A(C_X) \times A(C_Y \amalg C_Z) \xrightarrow{(id, \boxtimes)} A_X \times A_{Y \amalg Z} \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 A(C_{X \amalg Y} \amalg C_Z) \xleftarrow{h} T^\times A(C_{X \amalg Y} \amalg C_Z) \quad T^\times A(C_X \amalg C_{Y \amalg Z}) \xrightarrow{h} A(C_X \amalg C_{Y \amalg Z}) \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 T^\times A_{X \amalg Y \amalg Z} \xleftarrow{T^\times h} T^{\times 2} A_{X \amalg Y \amalg Z} \xrightarrow{\mu^{T^\times} kA} T^\times A_{X \amalg Y \amalg Z} \xleftarrow{\mu^{T^\times} kA} T^{\times 2} A_{X \amalg Y \amalg Z} \xrightarrow{T^\times h} T^\times A_{X \amalg Y \amalg Z} \\
 \searrow h \quad \downarrow h \quad \swarrow h \\
 A_{X \amalg Y \amalg Z}
 \end{array}$$

Hence, \boxtimes satisfies (C1) of Equation (3.25).

For (C2), if w, x, y, z are mutually distinct elements of X , then, by the monad algebra axioms, the following diagram commutes in \mathbb{V} , and hence ζ satisfies (C2):

$$\begin{array}{c}
 (A_X)^{w \ddagger x, y \ddagger z} \\
 \swarrow = \quad \downarrow = \quad \searrow = \\
 (A_{X \setminus \{w, x\}})^{y \ddagger z} \xleftarrow{\zeta^{w \ddagger x}} (A(C_X^{w \ddagger x}))^{y \ddagger z} \xrightarrow{=} A((C_X \amalg C_Y)^{w \ddagger x, y \ddagger z}) \xleftarrow{=} (A(C_X^{y \ddagger z}))^{x \ddagger y} \xrightarrow{\zeta^{y \ddagger z}} (A_{X \setminus \{y, z\}})^{w \ddagger x} \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 A(C_{X \setminus \{w, x\}}^{y \ddagger z}) \xleftarrow{h} T^\times A(C_{X \setminus \{w, x\}}^{y \ddagger z}) \quad T^\times A(C_{X \setminus \{y, z\}}^{w \ddagger x}) \xrightarrow{h} A(C_{X \setminus \{y, z\}}^{w \ddagger x}) \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 T^\times A_{X \setminus \{w, x, y, z\}} \xleftarrow{T^\times h} T^{\times 2} A_{X \setminus \{w, x, y, z\}} \xrightarrow{\mu^{T^\times} kA} T^\times A_{X \setminus \{w, x, y, z\}} \xleftarrow{\mu^{T^\times} kA} T^{\times 2} A_{X \setminus \{w, x, y, z\}} \xrightarrow{T^\times h} T^\times A_{X \setminus \{w, x, y, z\}} \\
 \searrow h \quad \downarrow h \quad \swarrow h \\
 A_{X \setminus \{w, x, y, z\}}
 \end{array}$$

Finally, if X and Y are finite sets and x_1 and x_2 are distinct elements of X , then the diagram

$$\begin{array}{c}
 (A_X \times A_Y)^{x_1 \ddagger x_2} \\
 \swarrow = \quad \searrow = \\
 (A_{X \amalg Y})^{x_1 \ddagger x_2} \xleftarrow{\boxtimes} (A(C_X \amalg C_Y))^{x_1 \ddagger x_2} \xrightarrow{=} A(C_X^{x_1 \ddagger x_2}) \times A_Y \xrightarrow{\zeta^{x_1 \ddagger x_2}} A_{X \setminus \{x_1, x_2\}} \times A_Y \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 A(C_{X \amalg Y}^{x_1 \ddagger x_2}) \xleftarrow{h} T^\times A(C_{X \amalg Y}^{x_1 \ddagger x_2}) \quad T^\times A(C_{X \setminus \{x_1, x_2\}} \amalg C_Y) \xrightarrow{h} A(C_{X \setminus \{x_1, x_2\}} \amalg C_Y) \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 T^\times A_{(X \amalg Y) \setminus \{x_1, x_2\}} \xleftarrow{T^\times h} T^{\times 2} A_{(X \amalg Y) \setminus \{x_1, x_2\}} \xrightarrow{\mu^{T^\times} kA} T^\times A_{(X \amalg Y) \setminus \{x_1, x_2\}} \xleftarrow{\mu^{T^\times} kA} T^{\times 2} A_{(X \amalg Y) \setminus \{x_1, x_2\}} \xrightarrow{T^\times h} T^\times A_{(X \amalg Y) \setminus \{x_1, x_2\}} \\
 \searrow h \quad \downarrow h \quad \swarrow h \\
 A_{(X \amalg Y) \setminus \{x_1, x_2\}}
 \end{array}$$

commutes. Therefore, (C3) is satisfied and (A, \boxtimes, ζ) is a nonunital circuit algebra.

Conversely, let (S, \boxtimes, ζ) be a nonunital circuit algebra. For finite sets X_1, \dots, X_n , let $\boxtimes_{X_1, \dots, X_n} : \prod_{i=1}^n S_{X_i} \rightarrow S_{\coprod_i X_i}$ be the obvious morphism induced by \boxtimes . This is well defined since (S, \boxtimes, ζ) satisfies (C1)-(C3). Recall from Section 3.4 that, if $x_1, y_1, \dots, x_k, y_k$ are mutually distinct elements of $\coprod_i X_i$, then $(\prod_i S_{X_i})^{x_1 \ddagger y_1, \dots, x_k \ddagger y_k}$ is the limit of the diagram of parallel morphisms

$$\boxtimes_{X_1, \dots, X_n} \circ S(ch_{x_i}), \boxtimes_{X_1, \dots, X_n} \circ S(ch_{y_i} \circ \tau) : \prod_{i=1}^n S_{X_i} \rightarrow S_{\coprod_i X_i} \rightarrow S_{\S},$$

and that this is independent of the ordering of the pairs (x_i, y_i) .

Let X be a finite set and let \mathcal{X} be an admissible X -graph. For each $v \in V$, let $(C_{Z_v}, b_v) \in \text{el}(\mathcal{X})$ be a neighbourhood of v , and let $Z_v \cong |v|$ be such that $\coprod_v Z_v = X \amalg \{y_1, z_1, \dots, y_k, z_k\}$.

Since \mathcal{X} has no stick components,

$$(5.15) \quad S(\mathcal{X}) = \left(\prod_{v \in V(\mathcal{X})} S_{Z_v} \right)^{y_1 \ddagger z_1, \dots, y_k \ddagger z_k}.$$

In particular, the morphism $\prod_v S_{Z_v} \rightarrow S_{(\coprod_v Z_v)}$ induces a morphism

$$\boxtimes^V : S(\mathcal{X}) \rightarrow S_{\coprod_{v \in V} Z_v}^{y_1 \ddagger z_1, \dots, y_k \ddagger z_k}.$$

Let $\hat{h}_{\mathcal{X}} : S(\mathcal{X}) \rightarrow S(C_X) = S_X$ be the map (illustrated in Figure 9) defined, using (5.15), by the composition

$$(5.16) \quad S(\mathcal{X}) \xrightarrow{\boxtimes^V} S_{\coprod_{v \in V} Z_v}^{y_1 \ddagger z_1, \dots, y_k \ddagger z_k} \xrightarrow{\zeta^{(k-1)}} S_X$$

where $\zeta^{(k-1)}$ is obtained by the obvious iterated contractions. Since (S, \boxtimes, ζ) satisfies (C1) and (C2), these are independent of choices of ordering.




FIGURE 9. A representation of the structure map $\hat{h}_{\mathcal{X}} : S(\mathcal{X}) \rightarrow S_X$.

The maps $\hat{h}_{\mathcal{X}}$ so defined are the components of a morphism $\hat{h} : T^{\times} S \rightarrow S$ in $\text{GS}_{\mathbf{V}}$.

Moreover, for any \mathcal{X} -shaped graph of graphs $\mathbf{\Gamma}$, $\mathbf{\Gamma}(b_v)$ is a Z_v -graph. So, by (C3), we obtain morphisms $\hat{h}_{\mathbf{\Gamma}(b_v)} : S(\mathbf{\Gamma}(b_v)) \rightarrow S_{Z_v}$, and by iterating (5.16), it is straightforward to show that S , together with $\hat{h} : T^{\times} S \rightarrow S$ satisfy the monad algebra axioms. (See Figure 10 for an illustration of coherence of \hat{h} and $\mu^{\text{T}^{\times}}$.) \square




FIGURE 10. Compatibility of the structure map $\hat{h}: T^\times S \rightarrow S$ with the monad multiplication $\mu^{T^\times} S: (T^\times)^2 S \rightarrow T^\times S$.

Remark 5.17. The monad in [31] is defined identically to \mathbb{T}^\times on \mathbf{GS} except that, in [31], the empty graph \emptyset is not included as a $\mathbf{0}$ -graph. Consequently, algebras for the monad in [31] are equipped with a nonunital external product. See also Equation (3.22).

Remark 5.18. The monad \mathbb{T}^\times has arities $\mathbf{Gr} \subset \mathbf{GS}$ (in fact, \mathbb{T} is strongly cartesian) and hence there is an abstract nerve theorem in the style of [6] for \mathbf{Set} -valued nonunital circuit algebras. However, this construction is not described here since it is straightforward to derive from the – more interesting (and challenging) – nerve theorem for unital circuit algebras, Equation (8.4).

Example 5.19. Recall that the oriented graphical species $Di^\vee \in \mathbf{GS}_\vee$ defined in Equation (3.9) has a (unique) trivial circuit algebra structure and hence, there is a monad \mathbf{OT}^\times on the category \mathbf{GS}_\vee/Di^\vee of oriented graphical species in \mathbf{V} whose EM category of algebras is isomorphic to $\mathbf{GS}_\vee^{T^\times}/Di^\vee$. By Equation (5.14) and Equation (3.33), this is the category $\mathfrak{W}\mathfrak{P}_\vee^-$ of nonunital wheeled props in \mathbf{V} .

The monad \mathbf{OT}^\times is easy to describe: Following Equation (4.38), the endofunctor \mathbf{OT}^\times on oriented graphical species in \mathbf{V} is obtained by replacing the categories $X\text{-}\mathbf{Gr}_{\text{iso}}$ in the definition of T^\times with the categories of directed graphs with labelled input and output ports and port-preserving morphisms in $\mathbf{OGr} = \mathbf{Gr}/Di$, and applying the equivalence $\mathbf{GS}_\vee/Di^\vee \simeq \mathbf{psh}_\vee(\text{el}(Di))$.

Remark 5.20. In [46], a *biased* definition (following [52]) of (enriched) wheeled props is given and a groupoid-coloured operad \mathbb{W} for nonunital wheeled props is described and shown to be Koszul. The wheeled prop definition is essentially an oriented version of the description of enriched circuit algebras in [42, Proposition 5.4]. The operad \mathbb{W} is obtained from the operad of oriented downward wiring diagrams (see [42, Section 5.1]) by including isomorphisms in the (groupoid) colouring.

Equation (5.14) uses the idea that a graph is obtained by iterated “contraction” (or gluing ports) of a disjoint union of corollas. This technique is very similar to the *normal form of a graph* used in [46, Section 5] to prove that the (enriched) nonunital wheeled prop operad \mathbb{W} is Koszul.

6. REVIEW OF MODULAR OPERADS AND POINTED GRAPHS

In [41], modular operads in \mathbf{Set} are constructed as algebras for a composite monad \mathbb{DT} on \mathbf{GS} . The monads \mathbb{D} and \mathbb{T} will be used in Section 7.2 to construct the circuit algebra monad. Therefore, this

section provides a review of their definitions, and the distributive law between them, in the more general setting of \mathbf{GS}_V (where, as usual, V is assumed to have sufficient limits and colimits).

6.1. The monad \mathbb{T} and nonunital modular operads. The nonunital modular operad monad $\mathbb{T} = (T, \mu^{\mathbb{T}}, \eta^{\mathbb{T}})$ on \mathbf{GS} was described in [41, Section 5]. In this paper, the nonunital V -valued modular operad monad on \mathbf{GS}_V is also denoted by $\mathbb{T} = \mathbb{T}_V$.

For all finite sets X , let $X\text{-CGr}_{\text{iso}} \subset X\text{-Gr}_{\text{iso}}$ be the full subgroupoid of connected X -graphs. The endofunctor $T: \mathbf{GS}_V \rightarrow \mathbf{GS}_V$ is defined, for all $S \in \mathbf{GS}_V$, by

$$(6.1) \quad \begin{aligned} T^\times S_{\S} &= S_{\S}, \\ T^\times S_X &= \text{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{iso}}} S(\mathcal{X}), \quad \text{for all finite sets } X. \end{aligned}$$

By [41, Corollary 5.19] (which follows [30, Lemma 1.5.12]), if \mathcal{G} is a connected graph, and $\mathbf{\Gamma}: \text{el}(\mathcal{G}) \rightarrow \mathbf{CGr}$ is a nondegenerate \mathcal{G} -shaped graph of graphs such that $\mathbf{\Gamma}(b)$ is connected for all $(\mathcal{C}, b) \in \text{el}(\mathcal{G})$, then the colimit $\mathbf{\Gamma}(\mathcal{G})$ is also connected.

Hence, $\mathbb{T} = (T, \mu^{\mathbb{T}}, \eta^{\mathbb{T}})$, where $\mu^{\mathbb{T}}, \eta^{\mathbb{T}}$ are the obvious (co)restrictions to connected graphs, describes a monad on \mathbf{GS}_V .

Proposition 6.2. *The EM category $\mathbf{GS}_V^{\mathbb{T}}$ of algebras for \mathbb{T} on \mathbf{GS}_V is canonically equivalent to \mathfrak{MD}_V^- .*

Proof. This is proved in [41, Proposition 5.29] for the case $V = \mathbf{Set}$. The proof for general V is similar.

The inclusion $X\text{-CGr}_{\text{iso}} \hookrightarrow X\text{-Gr}_{\text{iso}}$ induces a monomorphism $TS_X \rightarrow T^\times S_X$. In particular, since $\mathcal{X}^{x \ddagger y}$ is connected for all connected X -graphs \mathcal{X} and all pairs x, y of distinct elements of X , if (A, h) is an algebra for \mathbb{T} , then A admits a contraction ζ given by $\zeta_X^{x \ddagger y}: A_X^{x \ddagger y} = A(\mathcal{C}_X^{x \ddagger y}) \xrightarrow{h} TA_{X \setminus \{x, y\}}$. This satisfies (M2) by the proof of Equation (5.14).

Moreover, there is a multiplication \diamond on A such that, for all finite sets X and Y and all elements $x \in X$, and $y \in Y$, $\diamond_{X, Y}^{x \ddagger y}$ is induced by $A(\mathcal{D}_{X, Y}^{x \ddagger y}) \rightarrow TA_{(X \amalg Y) \setminus \{x, y\}} \xrightarrow{h} A_{(X \amalg Y) \setminus \{x, y\}}$.

So, to show that (A, \diamond, ζ) satisfy the remaining axioms (M1)-(M3), we may construct (as in the proof of Equation (5.14) and [41, Proposition 5.29]), for each axiom, a pair of graphs of graphs that have the same colimit in \mathbf{CGr} .

The converse follows exactly the same method as the proof of [41, Proposition 5.29] and Equation (5.14). \square

6.2. The unit monad \mathbb{D} . The modular operad monad on \mathbf{GS}_V will be described in terms of a distributive law for composing \mathbb{T} with the *pointed graphical species monad* $\mathbb{D} = (D, \mu^{\mathbb{D}}, \eta^{\mathbb{D}})$. This was defined on graphical species in \mathbf{Set} in [41], and is generalised here to a monad, also denoted by $\mathbb{D} = \mathbb{D}_V$, on \mathbf{GS}_V for V with sufficient limits and colimits.

The endofunctor D on \mathbf{GS}_V is given by $DS_{\S} = S_{\S}$, $DS_{\tau} = S_{\tau}$ and, for finite sets X :

$$DS_X = \begin{cases} S_X & X \not\cong \mathbf{2}, \text{ and } X \not\cong \mathbf{0}, \\ S_{\mathbf{2}} \amalg S_{\S} & X = \mathbf{2}, \\ S_{\mathbf{0}} \amalg \tilde{S}_{\S} & X = \mathbf{0}, \end{cases}$$

where \tilde{S}_{\S} is the coequaliser of the identity and S_{τ} on S_{\S}

There are canonical natural transformations $\eta^{\mathbb{D}}: 1_{\mathbf{GS}_V} \Rightarrow D$, provided by the induced morphism $S \hookrightarrow DS$, and $\mu^{\mathbb{D}}: D^2 \Rightarrow D$ induced by the canonical projections $D^2 S_{\mathbf{2}} \rightarrow DS_{\mathbf{2}}$, so that $\mathbb{D} = (D, \mu^{\mathbb{D}}, \eta^{\mathbb{D}})$ defines the pointed graphical species monad on \mathbf{GS}_V .

Algebras for \mathbb{D} – called *pointed graphical species in \mathbf{V}* – are graphical species S in \mathbf{V} equipped with a distinguished unit-like morphism (Equation (3.14)) $\epsilon: S_{\S} \rightarrow S_{\mathbf{2}}$, and a distinguished morphism $o: S_{\S} \rightarrow S_{\mathbf{0}}$ that factors through \tilde{S}_{\S} .

Let \mathbf{B}_{*}^{\S} be category obtained from \mathbf{B}^{\S} by formally adjoining morphisms $u: \mathbf{2} \rightarrow \S$ and $z: \mathbf{0} \rightarrow \S$, subject to the relations:

- (i) $u \circ ch_1 = id \in \mathbf{B}^{\S}(\S, \S)$ and $u \circ ch_2 = \tau \in \mathbf{B}^{\S}(\S, \S)$;
- (ii) $\tau \circ u = u \circ \sigma_{\mathbf{2}} \in \mathbf{B}^{\S}(\mathbf{2}, \S)$;
- (iii) $z = \tau \circ z \in \mathbf{B}^{\S}(\mathbf{0}, \S)$.

It is straightforward to verify that \mathbf{B}_{*}^{\S} is completely described by:

- (i) $\mathbf{B}_{*}^{\S}(\S, \S) = \mathbf{B}^{\S}(\S, \S)$ and $\mathbf{B}_{*}^{\S}(Y, X) = \mathbf{B}^{\S}(Y, X)$ whenever $Y \not\cong \mathbf{0}$ and $Y \not\cong \mathbf{2}$;
- (ii) $\mathbf{B}_{*}^{\S}(\mathbf{0}, \S) = \{z\}$, and $\mathbf{B}_{*}^{\S}(\mathbf{0}, X) = \mathbf{B}^{\S}(\mathbf{0}, X) \amalg \{ch_x \circ z\}_{x \in X}$;
- (iii) $\mathbf{B}_{*}^{\S}(\mathbf{2}, \S) = \{u, \tau \circ u\}$, and $\mathbf{B}_{*}^{\S}(\mathbf{2}, X) = \mathbf{B}^{\S}(\mathbf{0}, X) \amalg \{ch_x \circ u, ch_x \circ \tau \circ u\}_{x \in X}$.

It follows that:

Lemma 6.3. *The EM category $\mathbf{GS}_{*\mathbf{V}}$ of algebras for \mathbb{D} is the category of \mathbf{V} -presheaves on \mathbf{B}_{*}^{\S} . In other words, the following are equivalent:*

- (1) S_{*} is a presheaf on \mathbf{B}_{*}^{\S} that restricts to a graphical species S on \mathbf{B}^{\S} ;
- (2) (S, ϵ, o) , with $\epsilon = S_{*}(u)$ and $o = S_{*}(z)$ is a pointed graphical species.

As a consequence of Equation (6.3), the notation S_{*} and (S, ϵ, o) will be used interchangeably to denote the same pointed graphical species.

Remark 6.4. It is convenient to think of ϵ as a *unit morphism* for \S , and of o as a *contracted unit morphism* for S . Likewise, the monad D may be viewed as adjoining formal units to $S_{\mathbf{2}}$ and formal contracted units to $S_{\mathbf{0}}$ as will now be explained.

6.3. Pointed graphs. The category \mathbf{CGr}_{*} of connected graphs and *pointed étale morphisms*, obtained in the bo-ff factorisation of the functor $\mathbf{CGr} \rightarrow \mathbf{GS}_{*}$ is described in detail in [41, Section 7.2]. Here, \mathbf{CGr}_{*} is extended to the category \mathbf{Gr}_{*} of *all* graphs and pointed étale morphisms. In other words, \mathbf{Gr}_{*} is the category defined by the bo-ff factorisation of the functor $\mathbf{Gr} \xrightarrow{\Upsilon} \mathbf{GS} \rightarrow \mathbf{GS}_{*}$, as in the following commuting diagram of functors:

$$(6.5) \quad \begin{array}{ccccc} & & \text{f.f.} & & \\ & \text{B}_{*}^{\S} & \xrightarrow{\quad \text{f.f.} \quad} & \mathbf{Gr}_{*} & \xrightarrow{\quad \Upsilon_{*} \quad} & \mathbf{GS}_{*} \\ & \uparrow \text{b.o.} & & \uparrow \text{b.o.} & & \uparrow \text{free}^{\mathbb{D}} \\ \text{B}^{\S} & \xrightarrow{\quad \quad} & \mathbf{Gr} & \xrightarrow{\quad \Upsilon \quad} & \mathbf{GS} \\ & & & & \downarrow \text{forget}^{\mathbb{D}} \end{array}$$

The inclusion $\mathbf{B}_{*}^{\S} \rightarrow \mathbf{Gr}_{*}$ is fully faithful (by uniqueness of bo-ff factorisation), and the induced nerve $\Upsilon_{*}: \mathbf{Gr}_{*} \rightarrow \mathbf{GS}_{*}$ is fully faithful by construction.

Let $\mathcal{G} \in \mathbf{Gr}$ be a graph. By Section 6.2, for each edge $e \in E$, the ch_e -coloured unit for $\Upsilon_{*}\mathcal{G}$ is given by $\epsilon_e^{\mathcal{G}} = ch_e \circ u \in \mathbf{Gr}_{*}(\mathbf{C}_2, \mathcal{G})$, and the corresponding contracted unit is given by $o_e^{\mathcal{G}} = ch_e \circ z \in \mathbf{Gr}_{*}(\mathbf{C}_0, \mathcal{G})$.

Since the functor Υ_{*} embeds \mathbf{Gr}_{*} as a full subcategory of \mathbf{GS}_{*} , I will write \mathcal{G} , rather than $\Upsilon_{*}\mathcal{G}$, where there is no risk of confusion. In particular, the element category $\mathbf{el}_{*}(\Upsilon_{*}\mathcal{G})$ is denoted simply by $\mathbf{el}_{*}(\mathcal{G})$ and called the *category of pointed elements of a graph \mathcal{G}* .

For all pointed graphical species S_* , the forgetful functor $\mathbf{GS}_* \rightarrow \mathbf{GS}$ induces an inclusion $\mathbf{el}(S) \hookrightarrow \mathbf{el}_*(S_*)$, and, for all graphs \mathcal{G} , the inclusion $\mathbf{el}(\mathcal{G}) \hookrightarrow \mathbf{el}_*(\mathcal{G})$ is final (see [35, Section IX.3]) by [41, Lemma 7.8]. Hence, for all pointed graphical species $S_* = (S, \epsilon, o)$ in \mathbf{V} ,

$$(6.6) \quad \lim_{(\mathcal{C}, b) \in \mathbf{el}_*(\mathcal{G})} S(\mathcal{C}) = \lim_{(\mathcal{C}, b) \in \mathbf{el}(\mathcal{G})} S(\mathcal{C}) = S(\mathcal{G}).$$

Moreover, by [41, Section 7.3], the induced inclusion $\mathbf{GS}_* \hookrightarrow \mathbf{psh}_{\mathbf{V}}(\mathbf{Gr}_*)$ is fully faithful.

To better understand morphisms in \mathbf{Gr}_* , let \mathcal{G} be a graph and let $W \subset V_0 \amalg V_2$ be a subset of isolated and bivalent vertices of \mathcal{G} .

The *vertex deletion functor* (for W) is the \mathcal{G} -shaped graph of graphs $\Lambda_{\setminus W}^{\mathcal{G}}: \mathbf{el}(\mathcal{G}) \rightarrow \mathbf{Gr}$ given by

$$\Lambda_{\setminus W}^{\mathcal{G}}(\mathcal{C}_X, b) = \begin{cases} (i) & \text{if } (\mathcal{C}_X, b) \text{ is a neighbourhood of } v \in W \text{ (so } X \cong \mathbf{0} \text{ or } X \cong \mathbf{2}), \\ \mathcal{C}_X & \text{otherwise.} \end{cases}$$

By Equation (5.2), this has a colimit $\mathcal{G}_{\setminus W}$ in \mathbf{Gr} , and, for all $(\mathcal{C}, b) \in \mathbf{el}(\mathcal{G})$, $\Lambda_{\setminus W}^{\mathcal{G}}$ describes a morphism $\mathcal{C} \rightarrow \Lambda_{\setminus W}^{\mathcal{G}}(b)$ in \mathbf{B}_*^{\S} . Hence, $\Lambda_{\setminus W}^{\mathcal{G}}$ induces a morphism $\mathbf{del}_{\setminus W} \in \mathbf{Gr}_*(\mathcal{G}, \mathcal{G}_{\setminus W})$.

Definition 6.7. *The morphism $\mathbf{del}_{\setminus W} \in \mathbf{Gr}_*(\mathcal{G}, \mathcal{G}_{\setminus W})$ induced by $\Lambda_{\setminus W}^{\mathcal{G}}: \mathbf{el}(\mathcal{G}) \rightarrow \mathbf{Gr}$ is called the vertex deletion morphism corresponding to $W \subset V_0 \cong V_2$.*

Example 6.8. Assume that $\mathcal{G} \not\cong \mathcal{C}_0$ is connected (so V_0 is empty). Let $W \subset V_2$ be a subset of bivalent vertices of \mathcal{G} . Unless $W = V(\mathcal{G})$ and $\mathcal{G} = \mathcal{W}^m$ for some $m \geq 1$, the graph $\mathcal{G}_{\setminus W}$ may be intuitively described as “ \mathcal{G} with a subset of bivalent vertices deleted” as in Figure 11.

In this case, the graph $\mathcal{G}_{\setminus W}$ is described explicitly by:

$$\mathcal{G}_{\setminus W} = \tau_{\setminus W} \bigcirc E_{\setminus W} \xleftarrow{s_{\setminus W}} H_{\setminus W} \xrightarrow{t_{\setminus W}} V_{\setminus W},$$

where

$$\begin{aligned} V_{\setminus W} &= V \setminus W, \\ H_{\setminus W} &= H \setminus \left(\coprod_{v \in W} H/v \right), \\ E_{\setminus W} &= E \setminus \left(\coprod_{v \in W} E/v \right), \end{aligned}$$

and $s_{\setminus W}, t_{\setminus W}$ are just the restrictions of s and t . The description of the involution $\tau_{\setminus W}$ is more complicated than needed here. (The interested reader is referred to [41, Section 7.2] which contains full descriptions of the vertex deletion morphisms).




FIGURE 11. Vertex deletion $\mathbf{del}_{\setminus W}: \mathcal{G} \rightarrow \mathcal{G}_{\setminus W}$.

Example 6.9. If $\mathcal{G} = \mathcal{C}_0$ is an isolated vertex $*$ and $W = \{*\}$, then $\mathbf{del}_{\setminus W}$ is precisely $z: \mathcal{C}_0 \rightarrow (i)$.

Example 6.10. If $\mathcal{G} = \mathcal{L}^k$, and $W = V$, then $\mathbf{\Lambda}_{\setminus W}^{\mathcal{G}}$ is the constant functor to (\mathfrak{l}) and hence $\mathcal{G}_{\setminus W} = (\mathfrak{l})$. The induced morphism is denoted by $u^k \stackrel{\text{def}}{=} \text{del}_{\setminus W}: \mathcal{L}^k \rightarrow \text{Gr}_*$ and described by the following diagram in Gr_* :

$$\begin{array}{ccccccc} (\mathfrak{l}) & \xrightarrow{ch_1} & \mathcal{C}_2 & \xleftarrow{ch_2 \circ \tau} & \dots & \xrightarrow{ch_1} & \mathcal{C}_2 & \xleftarrow{ch_2 \circ \tau} & (\mathfrak{l}) \\ & \searrow id_{(\mathfrak{l})} & \downarrow u & & & & \downarrow u & & \swarrow id_{(\mathfrak{l})} \\ & & (\mathfrak{l}) & = & \dots & = & (\mathfrak{l}) & & \end{array}$$

In particular, $u^1 = u: \mathcal{C}_2 \rightarrow (\mathfrak{l})$ and u^0 is just the identity on (\mathfrak{l}) .

Example 6.11. Let $\mathcal{G} = \mathcal{W}$ be the wheel graph with one vertex v . Then $\mathcal{W}_{\setminus \{v\}} = \text{colim}_{\text{el}(\mathcal{W})} \mathbf{\Lambda}_{\setminus \{v\}}^{\mathcal{W}}$ exists and is isomorphic to (\mathfrak{l}) in Gr . The induced morphism $\kappa \stackrel{\text{def}}{=} \text{del}_{\setminus \{v\}}: \mathcal{W} \rightarrow (\mathfrak{l})$ is described in (6.12). Hence, there are precisely two morphisms κ and $\tau \circ \kappa$ in $\text{Gr}_*(\mathcal{W}, \mathfrak{l})$:

(6.12)

$$\begin{array}{ccc} & \mathcal{W} & \\ ch_a \nearrow & & \nwarrow 1_{\mathcal{C}_2} \mapsto a \\ (\mathfrak{l}) & \xrightleftharpoons[ch_2 \circ \tau]{ch_1} & \mathcal{C}_2 \\ & \searrow u & \\ & (\mathfrak{l}) & \end{array}$$

(6.13)

$$\begin{array}{ccc} & \mathcal{W} & \\ ch_a \nearrow & & \nwarrow 1_{\mathcal{C}_2} \mapsto a \\ (\mathfrak{l}) & \xrightleftharpoons[ch_2 \circ \tau]{ch_1} & \mathcal{C}_2 \\ & \parallel & \downarrow \sigma_2 \\ (\mathfrak{l}) & \xrightleftharpoons[ch_1 \circ \tau]{ch_2} & \mathcal{C}_2 \\ & \searrow u & \\ & (\mathfrak{l}) & \end{array}$$

More generally, let \mathcal{W}^m be the wheel graph with m vertices $(v_i)_{i=1}^m$, and let $\iota \in \text{CGr}(\mathcal{L}^{m-1}, \mathcal{W}^m)$ be an étale inclusion. If W is the image of $V(\mathcal{L}^{m-1})$ in $V(\mathcal{W}^m)$, then $V(\mathcal{W}^m) \cong W \amalg \{*\}$, and by (6.12) and Equation (6.10), there are two distinct pointed morphisms, κ^m and $\tau \circ \kappa^m$, in $\text{Gr}_*(\mathcal{W}^m, \mathfrak{l})$. Hence, for all \mathcal{G} ,

$$\text{Gr}_*(\mathcal{W}^m, \mathcal{G}) = \text{Gr}(\mathcal{W}^m, \mathcal{G}) \amalg \{ch_e \circ \kappa^m\}_{e \in E(\mathcal{G})} \cong \text{Gr}_*(\mathcal{W}, \mathcal{G}).$$

For convenience, let us define $\mathcal{W}^0 \stackrel{\text{def}}{=} (\mathfrak{l}_{\bar{e}}) = \mathcal{W}_{\setminus V(\mathcal{W}^m)}^m$ ($m \geq 1$). Then for all $k \geq 0$, $u^k: \mathcal{L}^k \rightarrow (\mathfrak{l})$ factors canonically as $\mathcal{L}^k \rightarrow \mathcal{W}^k \xrightarrow{\kappa^k} (\mathfrak{l})$.

Remark 6.14. (See Section 4.3 and [42, Section 3].) In [42, Remark 3.2], it was explained that the composition $\bigcirc = \bigcap \circ \bigcup$ of Brauer diagrams is not described by a pushout of cospans as in [42, Diagram (3.13)]. Instead, the pushout for $(\bigcap \circ \bigcup)$ is induced by the trivial diagram $(\mathfrak{l}) \xrightarrow{id} (\mathfrak{l}) \xleftarrow{id} (\mathfrak{l})$, and therefore agrees with $\kappa: \mathcal{W} \rightarrow (\mathfrak{l})$ described in Equation (6.11). (See [41, Section 6] for more details on the combinatorics of contracted units.)

Definition 6.15. The similarity category $\text{Gr}_{\text{sim}} \hookrightarrow \text{Gr}_*$ is the identity-on-objects subcategory of Gr_* whose morphisms are generated under composition by vertex deletion morphisms and graph isomorphisms. Morphisms in Gr_{sim} are called similarity morphisms, and connected components of Gr_{sim} are similarity classes. Graphs are similar if they are in the same connected component of Gr_{sim} .

The following proposition summarises some important basic properties of vertex deletion morphisms. Detailed proofs may be found in [41, Section 7.2].

Proposition 6.16. For all graphs \mathcal{G} and all subsets $W \subset V_2$ of bivalent vertices of \mathcal{G} ,

- (1) $\text{del}_{\setminus W}: \mathcal{G} \rightarrow \mathcal{G}_{\setminus W}$ preserves connected components, and may be defined componentwise;

- (2) if \mathcal{G} is connected and $W = V$, then $\mathcal{G} \cong \mathcal{L}^k$ or $\mathcal{G} \cong \mathcal{W}^m$ for $k, m \geq 0$ and $\mathcal{G}_{\setminus W} \cong (1)$;
(3) if \mathcal{G} is connected, then, unless $\mathcal{G} \cong \mathcal{W}^m$ or $\mathcal{G} \cong \mathcal{C}_0$ and $W = V$ for $m \geq 1$, then $\text{del}_{\setminus W}$ induces an identity on ports: $\text{del}_{\setminus W}: E_0(\mathcal{G}) \xrightarrow{\cong} E_0(\mathcal{G}_{\setminus W})$.

The pair $(\text{Gr}_{\text{sim}}, \text{Gr})$ of subcategories of Gr_* defines an orthogonal factorisation system on Gr_* .

Example 6.17. For all graphical species S in \mathbf{V} , and all graphs \mathcal{G} with no isolated vertices,

$$(6.18) \quad DS(\mathcal{G}) = \coprod_{W \subset V_2} S(\mathcal{G}_{\setminus W}).$$

More generally, since $DS_0 = DS(\mathcal{C}_0) = S_0 \amalg \tilde{S}_8$:

$$(6.19) \quad DS(\mathcal{G}) \cong \text{colim}_{(\mathcal{H}, f) \in \mathcal{G}/\text{Gr}_{\text{sim}}} S(\mathcal{H}), \quad \text{for all graphs } \mathcal{G} \in \text{Gr}.$$

Example 6.20. Let $S_* = (S, \epsilon, o)$ be a pointed graphical species in \mathbf{V} . For $k, m \geq 0$, there are distinguished monomorphisms in \mathbf{V} :

$$(6.21) \quad S_*(u^k): S_8 \rightarrow S(\mathcal{L}^k), \quad \text{and } S_*(\kappa^m): S_8 \rightarrow S(\mathcal{W}^m).$$

6.4. The distributive law for modular operads. For the remainder of this section, all graphs will be assumed to be connected, unless stated otherwise.

Recall from Equation (5.6) that an X -graph \mathcal{X} is said to be admissible if it has no stick components. For any finite set X , we may enlarge the category $X\text{-CGr}_{\text{iso}}$ of admissible connected X -graphs and port-preserving isomorphisms, to form a category $X\text{-CGr}_{\text{sim}}$ that includes vertex deletion morphisms out of X -graphs together with any labelling of ports from the domain.

- If $X \not\cong 0$, $X \not\cong 2$, then $X\text{-CGr}_{\text{sim}}$ is the category whose objects are connected X -graphs and whose morphisms are similarity morphisms that preserve the labelling of the ports.
- For $X = 2$, $2\text{-CGr}_{\text{sim}}$ contains the morphisms $\text{del}_{\setminus V}: \mathcal{L}^k \rightarrow (1)$, and hence the labelled stick graphs $(1, id)$ and $(1, \tau)$. There are no non-trivial morphisms out of these special graphs, and \mathcal{X} is in the same connected component as $(1, id)$ if and only if $\mathcal{X} = \mathcal{L}^k$ (with the identity labelling) for some $k \in \mathbb{N}$. In particular, $\tau: (1) \rightarrow (1)$ does not induce a morphism in $2\text{-CGr}_{\text{sim}}$.
- Finally, when $X = 0$, the morphisms $\text{del}_{\setminus V}: \mathcal{W}^m \rightarrow (1)$, and $z: \mathcal{C}_0 \rightarrow (1)$ are not boundary-preserving, and, in particular, do not equip (1) with any labelling of its ports. So, the objects of $0\text{-CGr}_{\text{sim}}$ are the admissible 0 -graphs, and (1) .

In particular, $\mathcal{W}^m, \mathcal{C}_0$ and (1) are in the same connected component of $0X\text{-CGr}_{\text{sim}}$. Since (1) is not admissible, there are no non-trivial morphisms in $0\text{-CGr}_{\text{sim}}$ with (1) as domain.

To simplify notation in what follows, we write $\mathcal{C}_{0 \setminus V} \stackrel{\text{def}}{=} (1)$ and $\text{del}_{\setminus V} = z: \mathcal{C}_0 \rightarrow (1)$.

The following lemma is immediate from the definitions:

Lemma 6.22. *For all finite sets X and all connected X -graphs \mathcal{X} , $\mathcal{X}^\perp \stackrel{\text{def}}{=} \mathcal{X}_{/(V_2 \amalg V_0)}$ is terminal in the connected component of $X\text{-CGr}_{\text{sim}}$ containing \mathcal{X} .*

If $\mathcal{X} \cong \mathcal{L}^k$ or $\mathcal{X} \cong \mathcal{W}^m$, $\mathcal{X} \cong \mathcal{C}_0$ for some $k, m \geq 0$, then $\mathcal{X}^\perp \cong (1)$. Otherwise, \mathcal{X}^\perp is an admissible, connected X -graph (without bivalent or isolated vertices).

Let S be a graphical species in \mathbf{V} . By (6.19), if \mathcal{X} is a connected X -graph and $X\text{-CGr}_{\text{sim}}^{\mathcal{X}}$ is the connected component of \mathcal{X} in $X\text{-CGr}_{\text{sim}}$, then there is a canonical isomorphism

$$DS(\mathcal{X}) \cong \text{colim}_{\mathcal{X}' \in X\text{-CGr}_{\text{sim}}^{\mathcal{X}}} S(\mathcal{X}').$$

By (6.18), for $X \not\cong \mathbf{0}$,

$$TDS_X = \operatorname{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{iso}}} DS(\mathcal{X}) = \operatorname{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{iso}}} \coprod_{W \subset V_2(\mathcal{X})} S(\mathcal{X}_{\setminus W}).$$

And, since vertex deletion morphisms preserve ports when $X \neq \mathbf{0}$, each $\mathcal{X}_{\setminus W} \in X\text{-CGr}_{\text{iso}}$ is either an admissible X -graph or $\mathcal{X}_{\setminus W} \cong (\mathbf{i})$, and hence there are canonical morphisms

$$TDS_X \rightarrow \begin{cases} \operatorname{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{iso}}} S(\mathcal{X}) & = TS_X & = DTS_X & \text{when } X \not\cong \mathbf{0}, X \not\cong \mathbf{2}, \\ (\operatorname{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{iso}}} S(\mathcal{X})) \amalg S_{\S} & = TS_{\mathbf{2}} \amalg S_{\S} & = DTS_{\mathbf{2}} & \text{when } X = \mathbf{2} \end{cases}$$

(since (\mathbf{i}) is not a $\mathbf{2}$ -graph by convention). These describe the morphisms $\lambda_{\mathbb{DT}} S_X : TDS_X \rightarrow DTS_X$ in \mathbf{V} for $X \not\cong \mathbf{0}$.

When $X = \mathbf{0}$,

$$\begin{aligned} TDS_{\mathbf{0}} &= \operatorname{colim}_{\mathcal{X} \in \mathbf{0}\text{-CGr}_{\text{iso}}} DS(\mathcal{X}) \\ &= \left(\operatorname{colim}_{\substack{\mathcal{X} \in \mathbf{0}\text{-CGr}_{\text{iso}} \\ \mathcal{X} \not\cong \mathcal{C}_{\mathbf{0}}}} \coprod_{W \subset V_2(\mathcal{X})} S(\mathcal{X}_{\setminus W}) \right) \amalg DS(\mathcal{C}_{\mathbf{0}}) \\ &= \left(\operatorname{colim}_{\substack{\mathcal{X} \in \mathbf{0}\text{-CGr}_{\text{iso}} \\ \mathcal{X} \not\cong \mathcal{C}_{\mathbf{0}}}} \coprod_{W \subset V_2(\mathcal{X})} S(\mathcal{X}_{\setminus W}) \right) \amalg S_{\mathbf{0}} \amalg \tilde{S}_{\S}. \end{aligned}$$

Vertex deletion maps $\operatorname{del}_{\setminus W} : \mathcal{G} \rightarrow \mathcal{G}_{\setminus W}$ from $\mathbf{0}$ -graphs preserve ports unless $\operatorname{del}_{\setminus W} \cong z : \mathcal{C}_{\mathbf{0}} \rightarrow (\mathbf{i})$ or $\operatorname{del}_{\setminus W} \cong \kappa^m : \mathcal{W}^m \rightarrow (\mathbf{i})$. So, $\lambda_{\mathbb{DT}} S_{\mathbf{0}} : TDS_{\mathbf{0}} \rightarrow DTS_{\mathbf{0}}$ is given by the canonical morphism

$$TDS_{\mathbf{0}} = \left(\operatorname{colim}_{\substack{\mathcal{X} \in \mathbf{0}\text{-CGr}_{\text{iso}} \\ \mathcal{X} \not\cong \mathcal{C}_{\mathbf{0}}}} \coprod_{W \subset V_2(\mathcal{X})} S(\mathcal{X}_{\setminus W}) \right) \amalg S_{\mathbf{0}} \amalg \tilde{S}_{\S} \rightarrow \operatorname{colim}_{\mathcal{X} \in \mathbf{0}\text{-CGr}_{\text{iso}}} S(\mathcal{X}) \amalg \tilde{S}_{\S} = DTS_{\mathbf{0}}.$$

The verification that $\lambda_{\mathbb{DT}}$ satisfies the four distributive law axioms [5] follows by a straightforward application of the definitions. To help the reader gain familiarity with the constructions, I describe just one here, namely that the following diagram of endofunctors on $\mathbf{GS}_{\mathbf{V}}$ commutes:

$$(6.23) \quad \begin{array}{ccccc} T^2 D & \xrightarrow{T\lambda_{\mathbb{DT}}} & TDT & \xrightarrow{\lambda_{\mathbb{DT}} T} & DT^2 \\ \mu^T D \downarrow & & & & \downarrow D\mu^T \\ TD & \xrightarrow{\lambda_{\mathbb{DT}}} & & & DT. \end{array}$$

Recall, from (5.12) and (6.18), that, for all graphical species S in \mathbf{V} and all finite sets X ,

$$\begin{aligned} T^2 DS_X &= \operatorname{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{iso}}} \lim_{(\mathcal{C}, b) \in \operatorname{el}(\mathcal{X})} \operatorname{colim}_{\mathcal{Y}^b \in [b]\text{-CGr}_{\text{iso}}} \coprod_{W^b \subset V_2} S(\mathcal{Y}_{\setminus W^b}^b) \\ &= \operatorname{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{iso}}} \lim_{\Gamma \in \operatorname{CGr}(\mathcal{X})} \coprod_{W^{\Gamma} \subset V_2(\Gamma(\mathcal{X}))} S(\Gamma(\mathcal{X})_{\setminus W^{\Gamma}}), \end{aligned}$$

where, as usual, $\iota_v : \mathcal{C}_{\mathbf{v}} \rightarrow \mathcal{X}$ is the canonical morphism. So, to describe the two paths in (6.23), let \mathcal{X} be a connected X -graph, Γ a nondegenerate \mathcal{X} -shaped graph of connected graphs, and let $W^{\Gamma} \subset (V_0 \amalg V_2)(\Gamma(\mathcal{X}))$ be a subset of (bivalent or isolated) vertices of the colimit $\Gamma(\mathcal{X})$ of Γ in \mathbf{CGr} .

For each $(\mathcal{C}, b) \in \operatorname{el}(\mathcal{X})$, let

$$W^b \stackrel{\text{def}}{=} V(\Gamma(b)) \cap W^{\Gamma}, \quad \text{so } W^{\Gamma} \cong \coprod_{v \in V(\mathcal{X})} W^{\iota_v}.$$

If $(\mathcal{X}, \Gamma, W) = (\mathcal{C}_{\mathbf{0}}, \mathbf{I}^{\mathcal{C}_{\mathbf{0}}}, V(\mathcal{C}_{\mathbf{0}}))$, then $\Gamma(\mathcal{X}) = \mathcal{C}_{\mathbf{0}}$, and $(\mathcal{C}_{\mathbf{0}}, b)$ is the unique object of $\operatorname{el}(\mathcal{X})$.

Otherwise, since \mathcal{X} is connected, it must be the case that $W^{\Gamma} \subset V_2(\Gamma(\mathcal{X}))$.

In each case we may define $W \subset V(\mathcal{X})$ to be the set of vertices v of \mathcal{X} such that $W^{\iota_v} = V(\Gamma(\iota_v))$. Then, there is a unique nondegenerate $\mathcal{X}_{\setminus W}$ -shaped graph of connected graphs $\Gamma_{\setminus W}$ given by $\Gamma_{\setminus W}(\operatorname{del}_{\setminus W} \circ b) = \Gamma(b)_{\setminus W^b}$, and with colimit $\Gamma_{\setminus W}(\mathcal{X}_{\setminus W})$ in \mathbf{Gr}_{*} :

$$(6.24) \quad \Gamma_{\setminus W}(\mathcal{X}_{\setminus W}) = \Gamma(\mathcal{X})_{\setminus W^{\Gamma}}.$$




FIGURE 12. Diagram (6.23) commutes. Here the vertices in W are marked in green.

Then, the two paths (illustrated in Figure 12) described by the diagram (6.23) are induced by the following canonical maps (where, in each case, the vertical arrows are the defining universal morphisms):

Top-right

$$\begin{array}{ccccc}
 & \text{delete } W^b \text{ in } \Gamma(b) \rightarrow (\mathcal{X}, (\Gamma(b) \setminus W^b)_b, W) & \xrightarrow{\text{delete } W \text{ in } \mathcal{X}} & (\mathcal{X} \setminus W, \Gamma \setminus W) & \xrightarrow{\text{evaluate colimit } \Gamma \setminus W} \\
 & \downarrow & & \downarrow & \downarrow \\
 (\mathcal{X}, (\Gamma(b))_b, (W^b)_b) & & & & \Gamma \setminus W(\mathcal{X} \setminus W) \\
 & \searrow & \lim_{\text{el}(\mathcal{X})} S(\Gamma(b) \setminus W^b) & \xrightarrow{\quad} & S(\Gamma \setminus W(\mathcal{X} \setminus W)) \\
 \downarrow & & \downarrow & & \downarrow \\
 \lim_{\text{el}(\mathcal{X})} S(\Gamma(b) \setminus W^b) & & & & S(\Gamma \setminus W(\mathcal{X} \setminus W)) \\
 \downarrow & \xrightarrow{T\lambda_{\mathbb{DT}} S_X} & T D T S_X & \xrightarrow{\lambda_{\mathbb{DT}} T S_X} & D T^2 S_X \\
 T^2 D S_X & & & & \downarrow D\mu^T S_X \\
 & & & & D T S_X
 \end{array}$$

Left-bottom

$$\begin{array}{ccccc}
 (\mathcal{X}, \Gamma, W^\Gamma) & \xrightarrow{\text{evaluate colimit } \Gamma} & (\Gamma(\mathcal{X}), W^\Gamma) & \xrightarrow{\text{delete } W^\Gamma \text{ in } \Gamma(\mathcal{X})} & \Gamma(\mathcal{X}) \setminus W^\Gamma \\
 \downarrow & & \downarrow & & \downarrow \\
 S(\Gamma(\mathcal{X}) \setminus W^\Gamma) & \xlongequal{\quad} & S(\Gamma(\mathcal{X}) \setminus W^\Gamma) & \xlongequal{\quad} & S(\Gamma(\mathcal{X}) \setminus W^\Gamma) \\
 \downarrow & & \downarrow & & \downarrow \\
 T^2 D S_X & \xrightarrow{\mu^T D S_X} & T D S_X & \xrightarrow{\lambda_{\mathbb{DT}} S_X} & D T S_X
 \end{array}$$

These are equal since $S(\Gamma \setminus W(\mathcal{X} \setminus W)) = S(\Gamma(\mathcal{X}) \setminus W^\Gamma)$ by construction, and hence (6.23) commutes.

The proofs that $\lambda_{\mathbb{DT}}$ satisfies the remaining three distributive law axioms follow similarly (and are somewhat simpler than the proof that (6.23) commutes).

Proposition 6.25. *The monad \mathbb{T} on $\mathbf{GS}_{\mathbf{V}}$ extends to a monad $\mathbb{T}_* = (T_*, \mu^{\mathbb{T}_*}, \eta^{\mathbb{T}_*})$ on $\mathbf{GS}_{*\mathbf{V}}$ such that $\mathbf{GS}_{*\mathbf{V}}^{\mathbb{T}_*} \cong \mathbf{GS}_{\mathbf{V}}^{\mathbb{DT}}$.*

Proof. For $\mathbf{V} = \mathbf{Set}$, \mathbb{T}_* has been described in detail in [41, Section 7].

For graphical species in a general category \mathbf{V} with sufficient (co)limits, the endofunctor T_* is the quotient of T given, for all pointed graphical species $S_* = (S, \epsilon, o)$ in \mathbf{V} , by

$$(6.26) \quad T_* S_{\S} = S_{\S}, \text{ and } T_* S_X = \operatorname{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{sim}}} S(\mathcal{X}).$$

The unit $\epsilon^{T_* S} \stackrel{\text{def}}{=} T_*(\epsilon): S_{\S} \rightarrow T_* S_2$ is described by the obvious composite

$$S_{\S} \xrightarrow{\epsilon} S_2 \xrightarrow{\eta^{\mathbb{T}} S} T S_2 \longrightarrow T_* S_2,$$

and the contracted unit $o^{T_* S} \stackrel{\text{def}}{=} T_*(o): S_{\S} \rightarrow T_* S_0$ is described by

$$S_{\S} \xrightarrow{o} S_0 \xrightarrow{\eta^{\mathbb{T}} S} T S_0 \longrightarrow T_* S_0.$$

In particular, $\epsilon^{T_* S}$ is defined by maps $S(u^k): S_{\S} \rightarrow S(\mathcal{L}^k)$ (for $k \geq 1$), and $o^{T_* S}$ by maps $S(\kappa^m): S_{\S} \rightarrow S(\mathcal{W}^m)$, $m \geq 1$ and $S(o): S_{\S} \rightarrow S(\mathcal{C}_0)$ such that $o \circ S_{\tau} = o$.

The multiplication $\mu^{\mathbb{T}_*}$ for \mathbb{T}_* is the one induced by $\mu^{\mathbb{T}}$, that forgets the pair (\mathcal{X}, Γ) – of an X -graph \mathcal{X} , and a nondegenerate \mathcal{X} -shaped graph of connected graphs Γ – and replaces it with the colimit $\Gamma(\mathcal{X})$. The unit $\eta^{\mathbb{T}_*}$ for \mathbb{T}_* is defined by the maps $S_X = S(\mathcal{C}_X) \mapsto \operatorname{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{sim}}} S(\mathcal{X})$.

Let (A_*, h_*) be a \mathbb{T}_* algebra in $\mathbf{GS}_{*\mathbf{V}}$ such that $A_* = (A, \epsilon, o)$. The unique map $h: DTA \rightarrow A$ that preserves distinguished (contracted) units and factors through the quotient $TA \twoheadrightarrow T_* A$ on TA describes a \mathbb{DT} algebra structure on A .

Conversely, if (A, h) is a \mathbb{DT} algebra, then in particular, A has a \mathbb{D} -algebra structure (A, ϵ, o) and h , together with $\lambda_{\mathbb{DT}}$, induce a \mathbb{T}_* algebra structure on (A, ϵ, o) . (See the proof of [41, Proposition 7.39]).

It is straightforward (though somewhat laborious) to use the definition of $\lambda_{\mathbb{DT}}$ – as in the proof of [41, Proposition 7.39] – to check that these assignments extend to inverse functors $\mathbf{GS}_{*\mathbf{V}}^{\mathbb{T}_*} \rightleftarrows \mathbf{GS}_{\mathbf{V}}^{\mathbb{DT}}$. \square

It is then straightforward to derive the following theorem (that, in the case $\mathbf{V} = \mathbf{Set}$, was the main result, Theorem 7.46, of [41]) for general \mathbf{V} .

Theorem 6.27. *The EM category $\mathbf{GS}_{\mathbf{V}}^{\mathbb{DT}}$ of algebras for \mathbb{DT} is canonically isomorphic to $\mathfrak{MD}_{\mathbf{V}}$.*

Proof. Let (A, h) be a \mathbb{DT} -algebra in \mathbf{V} . Then h induces a \mathbb{T} -algebra structure $h^{\mathbb{T}}: TA \rightarrow A$ by restriction, and a \mathbb{D} -algebra structure $h^{\mathbb{D}} = h \circ D\eta^{\mathbb{T}} A: DA \rightarrow A$. In particular, $h^{\mathbb{T}}$ equips A with the a multiplication \diamond and contraction ζ such that (A, \diamond, ζ) is a nonunital modular operad in \mathbf{V} , and $h^{\mathbb{D}}$ equips A with the structure of a pointed graphical species $A_* = (A, \epsilon, o)$ in \mathbf{V} .

To prove that (A, h) describes a modular operad, it remains to show that ϵ is a unit for \diamond . But, this is immediate from the definition of the distributive law $\lambda_{\mathbb{DT}}$ or, equivalently, from Equation (6.25) and (6.26).

Conversely, if $(S, \diamond, \zeta, \epsilon)$ is a modular operad in \mathbf{V} , then (S, \diamond, ζ) defines a \mathbb{T} -algebra $p^{\mathbb{T}}: TS \rightarrow S$ by Equation (6.2), and $(S, \epsilon, \zeta\epsilon)$ defines a \mathbb{D} -algebra by Section 6.2. In particular, since ϵ is a unit for \diamond , and $p^{\mathbb{T}}$ is induced by iterations of \diamond and ζ , $p^{\mathbb{T}}$ induces a \mathbb{T}_* -algebra structure on $(S, \epsilon, \zeta\epsilon)$ by (6.26). Hence, $(S, \diamond, \zeta, \epsilon)$ describes an algebra for \mathbb{DT} . \square

Example 6.28. (See also Examples 3.7, 3.32 and 5.19, and Equation (3.33).) The results of this section may be modified for oriented graphical species (see Equation (3.8)). By restricting to connected directed graphs, we may obtain an oriented version of \mathbb{T} as in Equation (5.19). Since there are no

orientation preserving automorphisms of the directed exceptional edge \downarrow in \mathbf{OGr} , the endofunctor OD underlying the monad D on \mathbf{OGS}_V such that $\mathbf{OGS}_V^D \simeq \mathbf{GS}_V^{\mathbb{D}}/Di^V$ is defined, for all oriented graphical species $\tilde{S}: \mathbf{el}(Di)^{\text{op}} \rightarrow V$, by $OD\tilde{S}(\downarrow) = \tilde{S}(\downarrow)$ and


$$OD\tilde{S}(X_{\text{in}}, X_{\text{out}}) = \begin{cases} \tilde{S}(X_{\text{in}}, X_{\text{out}}) & \text{for } (X_{\text{in}}, X_{\text{out}}) \not\cong (\mathbf{0}, \mathbf{0}), \text{ and } (X_{\text{in}}, X_{\text{out}}) \not\cong (\mathbf{1}, \mathbf{1}), \\ \tilde{S}(\mathbf{1}, \mathbf{1}) \amalg \tilde{S}(\downarrow) & \text{for } (X_{\text{in}}, X_{\text{out}}) \cong (\mathbf{1}, \mathbf{1}), \\ \tilde{S}(\mathbf{0}, \mathbf{0}) \amalg \tilde{S}(\downarrow) & \text{for } (X_{\text{in}}, X_{\text{out}}) \cong (\mathbf{0}, \mathbf{0}) \end{cases}$$

A category \mathbf{OGr}_* of pointed directed graphs and a distributive law are obtained exactly as for the undirected case. This is equipped with distinguished morphisms $\hat{z}: \mathcal{C}_{\mathbf{0}, \mathbf{0}} \rightarrow (\downarrow)$ and $\hat{u}: \mathcal{C}_{\mathbf{1}, \mathbf{1}} \rightarrow (\downarrow)$ satisfying $\hat{u} \circ i_1 = \hat{u} \circ o_1 = id_{\downarrow}$.

Algebras for the composite monad on \mathbf{OGS}_V so-obtained are wheeled properads in V .


Remark 6.29. By Equation (3.34), circuit algebras are, in particular monoid objects in \mathbf{GS}_V . However, they do not describe monoids in the category \mathbf{MO}_V of modular operads since $\mathbf{MO}_V \simeq \mathbf{GS}_V^{\mathbb{D}^T}$ is not closed under the monoidal product \otimes in \mathbf{GS}_V : For, given $S, S' \in \mathbf{GS}_V$, there is, in general, no map $DT(S \otimes S') \rightarrow DT(S) \otimes DT(S')$. And hence, if $(A, h), (A', h')$ are algebras for \mathbb{D}^T it is not possible to construct a map $DT(A \otimes A') \rightarrow A \otimes A'$.

To see this, consider an element of $T(A \otimes A')_2 \subset T(A \otimes A')_2$ represented by an $A \otimes A'$ structure on \mathcal{L}^2 , where decoration by A is coloured in filled red, and decoration by A' is unfilled blue:



Since the TA and TA' components are both connected, this represents an element of $TA_1 \otimes TA'_1$ (“delete” the black circle) and hence, applying h, h' , we obtain an element of $A \otimes A'$.

However, in the following $A \otimes A'$ structure on \mathcal{L}^3 , the TA (red, filled) and TA' (blue, unfilled) components are no longer connected:



Hence this does not represent an element of $(DTA \otimes DTA')_2$ and so $\mathbf{GS}_V^{\mathbb{D}^T} \simeq \mathbf{MO}_V$ is not closed under the monoidal product \otimes on \mathbf{GS}_V .

7. A MONAD FOR CIRCUIT ALGEBRAS

Despite the similarity between the definitions of the monads \mathbb{T} and \mathbb{T}^\times on \mathbf{GS}_V , it is not possible to modify $\lambda_{\mathbb{D}^T}$ by replacing \mathbb{T} with \mathbb{T}^\times to obtain a distributive law for \mathbb{D} and \mathbb{T}^\times :

Namely, for all graphical species S , since $DS_2 \cong S_2 \amalg S_\S$ by definition, there is a canonical morphism

$$S(\amalg \amalg) = S_\S \times S_\S \rightarrow DS_2 \times DS_2 = DS(\mathcal{C}_2 \amalg \mathcal{C}_2) \rightarrow TDS_{2 \amalg 2}.$$

However, there is, in general, no morphism

$$(S_2 \amalg S_\S) \times (S_2 \amalg S_\S) \rightarrow TS_{2 \amalg 2} = DTS_{2 \amalg 2},$$

and therefore $\lambda_{\mathbb{D}^T}: TD \Rightarrow DT$ does not extend to a distributive law $T^\times D \Rightarrow DT^\times$.

The problem of finding the circuit algebra monad on \mathbf{GS}_V is easily solved by observing that the monad \mathbb{T}^\times is, itself a composite: the free graded monoid monad on $\mathbf{psh}(\Sigma)$ (see Section 3.4) induces a monad $\mathbb{L} = (L, \mu^{\mathbb{L}}, \eta^{\mathbb{L}})$ on \mathbf{GS}_V whose algebras are graphical species in V that are equipped with

an external product. Moreover, there is a distributive law $\lambda_{\mathbb{L}\mathbb{T}}: TL \Rightarrow LT$ such that $\mathbb{T}^\times = \mathbb{L}\mathbb{T}$ is the induced composite monad.

It will then be straightforward to prove (Equation (7.9)) that, for all (unpointed) graphical species S in \mathbb{V} , $LDT S$ describes a circuit algebra in \mathbb{V} .

By [10], there is a distributive law $\lambda_{\mathbb{L}(\mathbb{D}\mathbb{T})}: (DT)L \Rightarrow L(DT)$, and hence a composite monad $\mathbb{L}\mathbb{D}\mathbb{T}$ on $\mathbb{G}\mathbb{S}_{\mathbb{V}}$, if there are pairwise distributive laws $\lambda_{\mathbb{L}\mathbb{T}}: TL \Rightarrow LT$, and $\lambda_{\mathbb{L}\mathbb{D}}: DL \Rightarrow LD$, such that the following *Yang-Baxter diagram* of endofunctors on $\mathbb{G}\mathbb{S}_{\mathbb{V}}$ commutes:

$$(7.1) \quad \begin{array}{ccccc} & & DTL & \xrightarrow{D\lambda_{\mathbb{L}\mathbb{T}}} & DLT \\ & \nearrow \lambda_{\mathbb{D}\mathbb{T}}L & & & \searrow \lambda_{\mathbb{L}\mathbb{D}}T \\ TDL & & & & LDT \\ & \searrow T\lambda_{\mathbb{L}\mathbb{D}} & & & \nearrow L\lambda_{\mathbb{D}\mathbb{T}} \\ & & TLD & \xrightarrow{\lambda_{\mathbb{L}\mathbb{T}}D} & LTD \end{array}$$

In Section 7.1, the monad \mathbb{L} is defined and the distributive laws $\lambda_{\mathbb{L}\mathbb{T}}: TL \Rightarrow LT$ and $\lambda_{\mathbb{L}\mathbb{D}}: DL \Rightarrow LD$ are constructed. In Section 7.2, it is proved that the diagram (7.1) commutes, and the EM category of algebras for the composite monad $\mathbb{L}\mathbb{D}\mathbb{T}$ on $\mathbb{G}\mathbb{S}_{\mathbb{V}}$ is canonically equivalent to the category $\mathfrak{CA}_{\mathbb{V}}$ of circuit algebras in \mathbb{V} .

7.1. The free external product monad \mathbb{L} . Recall from Section 2.2 that, for any finite set X , $\text{Core}(X/\text{Set}_{\mathfrak{f}})$ is the maximal subgroupoid of $X/\text{Set}_{\mathfrak{f}}$ whose objects are pairs (Y, f) with $f \in \text{Set}_{\mathfrak{f}}(X, Y)$ a morphism of finite sets.

As usual, let \mathbb{V} be a category with sufficient (co)limits. The free graded monoid endofunctor L on $\mathbb{G}\mathbb{S}_{\mathbb{V}}$ is given by

$$(7.2) \quad \begin{aligned} LS_{\S} &= S_{\S} \\ LS_X &= \text{colim}_{(Y,f) \in \text{Core}(X/\text{Set}_{\mathfrak{f}})} \prod_{y \in Y} S_{f^{-1}(y)} \end{aligned}$$

for all finite sets X . This underlies a monad $\mathbb{L} = (L, \mu^{\mathbb{L}}, \eta^{\mathbb{L}})$ with monadic multiplication $\mu^{\mathbb{L}}$ induced by disjoint union of partitions, and monadic unit $\eta^{\mathbb{L}}$ induced by inclusion of trivial partitions.

The following lemma is immediate from the definitions (see also Equation (3.23)):

Lemma 7.3. *The EM category of algebras for the monad \mathbb{L} on $\mathbb{G}\mathbb{S}_{\mathbb{V}}$ is equivalent to the category of graphical species with unital external product in \mathbb{V} . In particular $\mathbb{G}\mathbb{S}_{\mathbb{V}}^{\mathbb{L}}$ is equivalent to the monoid subcategory of $\mathbb{G}\mathbb{S}_{\mathbb{V}}$.*

In order to describe a distributive law $\lambda_{\mathbb{L}\mathbb{T}}: TL \Rightarrow LT$ for \mathbb{L} and \mathbb{T} , observe first that $L \subset T^\times$ is a subfunctor since, if S is a graphical species in \mathbb{V} , and X a finite set, then

$$\begin{aligned} LS_X &= \text{colim}_{(Y,f) \in \text{Core}(X/\text{Set}_{\mathfrak{f}})} \prod_{y \in Y} S_{f^{-1}(y)} \\ &= \text{colim}_{\mathcal{X} \in X\text{-Cor}_{\boxtimes \text{iso}}} S(\mathcal{X}) \end{aligned}$$

where $X\text{-Cor}_{\boxtimes \text{iso}} \subset X\text{-Gr}_{\text{iso}}$ is the groupoid of X -graphs \mathcal{X}_{cor} that are disjoint unions of corollas $\mathcal{X}_{\text{cor}} = \coprod_{i \in I} \mathcal{C}_{X_i}$ (with $\coprod_{i \in I} X_i = X$). In particular, from this point of view, the multiplication $\mu^{\mathbb{L}}$ for \mathbb{L} is just the restriction of the multiplication $\mu^{\mathbb{T}^\times}$ for \mathbb{T}^\times (Equation (5.14)) to \mathcal{X}_{cor} -shaped graphs of disjoint unions of corollas, and the unit $\eta^{\mathbb{L}}$ for \mathbb{L} , is the obvious corestriction.

Given a finite set X , and a connected X -graph $\mathcal{X} \in X\text{-CGr}_{\text{iso}}$, let $\text{Cor}_{\boxtimes}^{(\mathcal{X})}$ denote the full subgroupoid of $\text{Gr}^{(\mathcal{X})}$ whose objects are nondegenerate \mathcal{X} -shaped graphs of graphs $\mathbf{\Gamma}$ such that, for all $(\mathcal{C}_{X_b}, b) \in$

$\text{el}(\mathcal{X}), \mathbf{\Gamma}(\mathcal{C}_{X_b}, b) \in X\text{-Cor}_{\boxtimes \text{iso}}$ is a disjoint union of corollas. Then,

$$\begin{aligned} TLS_X &= \text{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{iso}}} \lim_{(\mathcal{C}, b) \in \text{el}(\mathcal{X})} LS(\mathcal{C}) \\ &= \text{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{iso}}} \lim_{\mathbf{\Gamma} \in \text{Cor}_{\boxtimes}^{(\mathcal{X})}} S(\mathbf{\Gamma}(\mathcal{X})). \end{aligned}$$

Since, for all $\mathbf{\Gamma} \in \text{Cor}_{\boxtimes}^{(\mathcal{X})}$, the colimit $\mathbf{\Gamma}(\mathcal{X})$ has the structure of an admissible (in general, not connected) X -graph, there is a canonical morphism $TLS_X \rightarrow \text{colim}_{\mathcal{X}' \in X\text{-Gr}_{\text{iso}}} S(\mathcal{X}')$. And, therefore, since every graph is the coproduct in Gr of its connected components, $S(\mathcal{X}') = \prod_{k \in K} S(\mathcal{X}'_k)$ describes an element of TLS_X for all X -graphs $\mathcal{X}' = \coprod_{k \in K} \mathcal{X}'_k$.

Let $\lambda_{\mathbb{L}\mathbb{T}}: TL \Rightarrow LT$ be the natural transformation so obtained. In other words, $\lambda_{\mathbb{L}\mathbb{T}}$ is induced by the canonical morphisms,

$$\text{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{iso}}} \left(\text{colim}_{\mathbf{\Gamma} \in \text{Cor}_{\boxtimes}^{(\mathcal{X})}} S(\mathbf{\Gamma}(\mathcal{X})) \right) \rightarrow \text{colim}_{\mathcal{X}' \in X\text{-CGr}_{\text{iso}}} S(\mathcal{X}') \text{ (where } S \in \text{GS}_{\mathbf{V}}, \text{ and } X \in \mathbf{B}),$$

that forgets pairs $(\mathcal{X}, \mathbf{\Gamma})$ where \mathcal{X} is a connected admissible X -graph and $\mathbf{\Gamma} \in \text{Cor}_{\boxtimes}^{(\mathcal{X})}$, and remembers only the X -graphs given by the colimit $\mathbf{\Gamma}(\mathcal{X}) = \coprod_{k \in K} \mathbf{\Gamma}(\mathcal{X})_k$, in the evaluation of $S(\mathbf{\Gamma}(\mathcal{X}))$.

Lemma 7.4. *The natural transformation $\lambda_{\mathbb{L}\mathbb{T}}: TL \Rightarrow LT$ satisfies the four axioms of a distributive law, and $\mathbb{L}\mathbb{T} = \mathbb{T}^\times$ on $\text{GS}_{\mathbf{V}}$. Hence, algebras for $\mathbb{L}\mathbb{T}$ are nonunital circuit algebras.*

Proof. Since a graph is the disjoint union of its connected components, $LT = T^\times: \text{GS}_{\mathbf{V}} \rightarrow \text{GS}_{\mathbf{V}}$.

Moreover, $L \subset T^\times$ and $T \subset T^\times$ so $TL \subset (T^\times)^2$ is a subfunctor, and $\lambda_{\mathbb{L}\mathbb{T}}$ is just the restriction to TL of $\mu^{T^\times}: (T^\times)^2 \Rightarrow T^\times = LT$ on $\text{GS}_{\mathbf{V}}$. The four distributive law axioms then follow immediately from the monad axioms. \square

Observe also that there is an obvious choice of natural transformation $\lambda_{\mathbb{L}\mathbb{D}}: DL \Rightarrow LD$ that defines the distributive law for \mathbb{L} and \mathbb{D} :

Lemma 7.5. *The natural transformation $\lambda_{\mathbb{L}\mathbb{D}}: DL \Rightarrow LD$ obtained by extending the natural transformation $L\eta^{\mathbb{D}}: L \Rightarrow LD$ to a DL by*

$$S_{\S} \rightarrow DS_{\mathbf{2}} \xrightarrow{\eta^{\mathbb{L}} DS} LDS_{\mathbf{2}}, \text{ and } S_{\S} \rightarrow DS_{\mathbf{0}} \xrightarrow{\eta^{\mathbb{L}} DS} LDS_{\mathbf{0}}, \text{ for all } S \in \text{GS}_{\mathbf{V}}$$

describes a distributive law.

Proof. Informally, $\lambda_{\mathbb{L}\mathbb{D}}$ takes a tuple $(S_{X_i})_{i=1}^n \in LS_{\prod_i X_i}$ of objects in the image of S , and regards it as a tuple in the image of DS , and takes the image of S_{\S} in $DS_{\mathbf{0}}$ or $DS_{\mathbf{2}}$ and regards it as a 1-tuple in the image of $DS \hookrightarrow LDS$. This construction is natural in S and it is simple to show, using the monad unit axiom for $\eta^{\mathbb{D}}$, that $\lambda_{\mathbb{L}\mathbb{D}}$ so obtained satisfies the axioms of a distributive law. \square

7.2. An iterated distributive law for circuit algebras. So far, we have defined three monads \mathbb{T}, \mathbb{D} and \mathbb{L} on $\text{GS}_{\mathbf{V}}$, and three distributive laws $\lambda_{\mathbb{D}\mathbb{T}}: TD \Rightarrow DT$, $\lambda_{\mathbb{L}\mathbb{T}}: TL \Rightarrow LT$ and $\lambda_{\mathbb{L}\mathbb{D}}: DL \Rightarrow LD$. It remains to show that the Yang-Baxter diagram (7.1) commutes and therefore there is an iterated distributive law for the three monads such that the resulting composite monad $\mathbb{L}\mathbb{D}\mathbb{T}$ on $\text{GS}_{\mathbf{V}}$ is the desired circuit algebra monad.

Let \mathcal{G} be any graph and $W \subset V_2$ any subset of bivalent vertices of \mathcal{G} . As in Section 6.3, let $\mathcal{G}_{\setminus W}$ be the graph obtained from \mathcal{G} by deleting the vertices in W . If $\mathbf{\Gamma}_{\setminus W}: \text{el}(\mathcal{G}_{\setminus W}) \rightarrow \text{Gr}$ is a

nondegenerate $\mathcal{G}_{\setminus W}$ -shaped graph of graphs, then there is an induced nondegenerate \mathcal{G} -shaped graph of graphs $\mathbf{\Gamma}^W : \text{el}(\mathcal{G}) \rightarrow \text{Gr}$ defined by

$$(7.6) \quad (\mathcal{C}, b) \mapsto \begin{cases} \mathcal{C}_2 & (\mathcal{C}, b) \text{ is a neighbourhood of } v \in W \\ \mathbf{\Gamma}_{\mathcal{W}}(\text{del}_{\setminus W} \circ b) & \text{otherwise.} \end{cases}$$

Moreover, $\mathbf{\Gamma}^W$ induces an inclusion $W \hookrightarrow V_2(\mathbf{\Gamma}^W(\mathcal{X}))$ (see Figure 13), and

$$(7.7) \quad \mathbf{\Gamma}^W(\mathcal{X})_{\setminus W} = \mathbf{\Gamma}_{\setminus W}(\mathcal{X}_{\setminus W}).$$

Hence, the assignment $\mathbf{\Gamma}_{\setminus W} \rightarrow \mathbf{\Gamma}^W$ defines a canonical full embedding $\text{Gr}^{(\mathcal{G}_{\setminus W})} \hookrightarrow \text{Gr}^{(\mathcal{G})}$ of categories that preserves similarity classes of colimits: In particular, the embedding $\text{Gr}^{(\mathcal{G}_{\setminus W})} \hookrightarrow \text{Gr}^{(\mathcal{G})}$ restricts




FIGURE 13. The colimits of the \mathcal{G} -shaped graph of graphs $\mathbf{\Gamma}^W$ described in (7.6), and the $\mathcal{G}_{\setminus W}$ -shaped graph of graphs $\mathbf{\Gamma}_{\setminus W}$ are similar in Gr_* .

to an inclusion $\text{Cor}_{\boxtimes}^{(\mathcal{G}_{\setminus W})} \hookrightarrow \text{Cor}_{\boxtimes}^{(\mathcal{G})}$.

Proposition 7.8. *The triple of distributive laws $\lambda_{\mathbb{D}\mathbb{T}} : TD \Rightarrow DT$, $\lambda_{\mathbb{L}\mathbb{T}} : TL \Rightarrow LT$, and $\lambda_{\mathbb{L}\mathbb{D}} : DL \Rightarrow LD$ on GS , describe a composite monad \mathbb{LDT} on $\text{GS}_{\mathbb{V}}$.*

Proof. By [10, Theorem 1.6], we must check that the diagram (7.1) commutes for $\lambda_{\mathbb{D}\mathbb{T}}$, $\lambda_{\mathbb{L}\mathbb{T}}$, and $\lambda_{\mathbb{L}\mathbb{D}}$:

For all graphical species S in \mathbb{V} and all finite sets X ,


$$TDLS_X = \text{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{iso}}} \coprod_{W \subset (V_0 \amalg V_2)(\mathcal{X})} \lim_{\mathbf{\Gamma}_{\setminus W} \in \text{Cor}_{\boxtimes}^{(\mathcal{X}_{\setminus W})}} S(\mathbf{\Gamma}_{\setminus W}(\mathcal{X}_{\setminus W}))$$

is indexed by triples $(\mathcal{X}, W, \mathbf{\Gamma}_{\setminus W})$, where \mathcal{X} is a connected X -graph, $W \subset V_2(\mathcal{X})$ (or $\mathcal{X} \cong \mathcal{C}_0$ and $W = V(\mathcal{C}_0)$) and $\mathbf{\Gamma}_{\setminus W}$ is a nondegenerate $\mathcal{X}_{\setminus W}$ -shaped graph of graphs such that $\mathbf{\Gamma}_{\setminus W}(b)$ is a disjoint union of corollas for each $(\mathcal{C}_{X_b}, b) \in \text{el}(\mathcal{X}_{\setminus W})$.

Observe first that, if $W = V(\mathcal{X})$, then either $\mathcal{X} \cong \mathcal{C}_0$ or $\mathcal{X} \cong \mathcal{W}^m$ in $\mathbf{0}\text{-CGr}_{\text{iso}}$, or $\mathcal{X} \cong \mathcal{L}^k$ in $\mathbf{2}\text{-CGr}_{\text{iso}}$. In each case $\mathcal{X}_{\setminus W} \cong (1)$ and hence $\mathbf{\Gamma}_{\setminus W}$ is the trivial graph of graphs $id \mapsto (1)$. In particular, the maps $S(\mathbf{\Gamma}_{\setminus W}(\mathcal{X}_{\setminus W})) \rightarrow TDLS_X \rightarrow LDTS_X$ induced by each path $TDL \Rightarrow LDT$ in (7.1) coincide, since both factor through S_{\S} .


Assume therefore that $W \neq V(\mathcal{X})$. The two paths in (7.1) are then described by the following data:

Top:



where, in each case, we evaluate $S(\Gamma_{\setminus W}(\mathcal{X}_{\setminus W}))$ and take the appropriate colimit of such objects.

Bottom: The bottom path of (7.1) gives



where Γ^W is the nondegenerate \mathcal{X} -shaped graph of graphs in $\text{Cor}_{\boxtimes}^{(\mathcal{X})}$ obtained from $\Gamma_{\setminus W}$ as in (7.6).

So, $\Gamma_{\setminus W}(\mathcal{X}_{\setminus W}) = \Gamma^W(\mathcal{X})_{\setminus W}$ by (7.7). Hence, the diagram (7.1) commutes, and $(\lambda_{\mathbb{L}\mathbb{D}}, \lambda_{\mathbb{L}\mathbb{T}}, \lambda_{\mathbb{D}\mathbb{T}})$ describe an iterated distributive law for \mathbb{L}, \mathbb{D} and \mathbb{T} on $\text{GS}_{\mathbb{V}}$. \square

It therefore remains to prove:

Theorem 7.9. *The EM category of algebras for the composite monad $\mathbb{L}\mathbb{D}\mathbb{T}$ on $\text{GS}_{\mathbb{V}}$ is isomorphic to the category $\mathfrak{CA}_{\mathbb{V}}$ of circuit algebras in \mathbb{V} .*

Proof. By [5], for a composite monad is equipped with an algebra structure for each of the component monads. Hence, an algebra (A, h) for the composite monad $\mathbb{L}\mathbb{D}\mathbb{T}$ on $\text{GS}_{\mathbb{V}}$ has an $\mathbb{L}\mathbb{T}$ -algebra structure $h^{\mathbb{L}\mathbb{T}}: LTA \rightarrow A$, and also a $\mathbb{D}\mathbb{T}$ algebra structure $h^{\mathbb{D}\mathbb{T}}: DTA \rightarrow A$. Each structure is given by the corresponding restriction $LTA \rightarrow LDTA \xrightarrow{h} A$, respectively $DTA \rightarrow LDTA \xrightarrow{h} A$. Therefore, by Equation (7.4), h induces an external product \boxtimes , and contraction ζ on A such that (A, \boxtimes, ζ) is a nonunital circuit algebra. And, by Equation (6.27), h induces a unital multiplication (\diamond, ϵ) on A , such that

$$\diamond_{X,Y}^{x\ddagger y} = \zeta_{X\amalg Y}^{x\ddagger y} \boxtimes_{X,Y}: (A_X \times A_Y)^{x\ddagger y} \rightarrow A_{X\amalg Y \setminus \{x,y\}}.$$

Hence

$$A_X \xrightarrow{(id \times ch_x) \circ \Delta} (A_X \times A_{\S})^{x\ddagger 2} \xrightarrow{id \times \epsilon} (A_X \times A_2)^{x\ddagger 2} \xrightarrow{\zeta_{X \amalg 2}^{x\ddagger 2} \circ \boxtimes} A_X$$

is the identity on A_X . So, (A, h) has a canonical circuit algebra structure.

Conversely, a circuit algebra $(S, \boxtimes, \zeta, \epsilon)$ in \mathbf{V} has an underlying modular operad structure $(S, \diamond, \zeta, \epsilon)$, where $\diamond_{X,Y}^{x\ddagger y} = \zeta_{X \amalg Y}^{x\ddagger y} \boxtimes_{X,Y}$ and hence $(S, \boxtimes, \zeta, \epsilon)$ defines a \mathbb{DT} -algebra structure $h^{\mathbb{DT}}: DTS \rightarrow S$ by Equation (6.27). By Equation (7.3), (S, \boxtimes) has an \mathbb{L} -algebra structure where $h^{\mathbb{L}}: LS \rightarrow S$ is given by $\boxtimes_{X,Y}: S_X \times S_Y \rightarrow S_{X \amalg Y}$. It follows immediately from the circuit algebra axioms that $h^{\mathbb{L}} \circ Lh^{\mathbb{DT}}: L DTS \rightarrow S$ gives S the structure of an \mathbb{LDT} -algebra.

These maps are functorial by construction and extend to an equivalence $\mathfrak{CA}_{\mathbf{V}} \simeq \mathbf{GS}_{\mathbf{V}}^{\mathbb{LDT}}$. \square

Corollary 7.10. *If $\mathbf{V} = \mathbf{Set}$, then for each palette (\mathfrak{C}, ω) in \mathbf{Set} , the subcategory $\mathfrak{CA}^{(\mathfrak{C}, \omega)}$ of (\mathfrak{C}, ω) -coloured circuit algebras is the EM-category of algebras for the monad \mathbb{LDT} restricted to $\mathbf{GS}^{(\mathfrak{C}, \omega)}$. By Equation (3.27), this is equivalent to the category $\mathbf{Alg}_{\mathbf{Set}}(\mathbf{WD}^{(\mathfrak{C}, \omega)}) = \mathbf{CA}^{(\mathfrak{C}, \omega)}$ of \mathbf{Set} -valued algebras for the operad $\mathbf{WD}^{(\mathfrak{C}, \omega)}$ of (\mathfrak{C}, ω) -coloured wiring diagrams defined in [42, Section 5.2].*

Example 7.11. By Equation (3.33) and Equation (7.9), the category $\mathfrak{WP}_{\mathbf{V}}$ of wheeled props in \mathbf{V} is isomorphic to the slice category $\mathbf{GS}_{\mathbf{V}}^{\mathbb{LDT}}/Di^{\mathbf{V}}$. In particular – as in Examples 5.19 and 6.28 – the constructions of Sections 7.1 and 7.2 may be modified, in terms of directed graphs, to obtain oriented versions of the monads and distributive laws and a composite wheeled prop monad on $\mathbf{OGS}_{\mathbf{V}}$.

Remark 7.12. Like monads, operads encode algebraic structure. Indeed, for any operad in \mathbf{Set} , one may define a monad with the same algebras. Markl [37] defined distributive laws for operads and proved that the operad obtained by composing Koszul operads via an *operad distributive law* is Koszul. This result was used in [29] to compose an operad (of downward wiring diagrams [42, Section 5.1]) associated to the restriction of \mathbb{T} to $\mathbf{GS}^{\{*\}}$ and an operad associated to the restriction of \mathbb{L} to $\mathbf{GS}^{\{*\}}$. So, to determine whether the operad (of wiring diagrams) for monochrome unital circuit algebras obtained from \mathbb{LDT} is Koszul, it should be sufficient to explore whether the operad obtained by restricting \mathbb{D} is. This is not expected to be the case, and the question is not further explored here.

8. A NERVE THEOREM FOR CIRCUIT ALGEBRAS

This section recalls the nerve theorem for \mathbf{Set} -valued modular operads [41, Theorem 8.2] and gives a proof, by the same methods, of an analogous result for \mathbf{Set} -valued circuit algebras.

8.1. Overview of the method. Let $\mathbf{CGr}_* \hookrightarrow \mathbf{GS}_*$ be the full subcategory of connected graphs and pointed étale morphisms from Section 6.3, that is obtained in the bo-ff factorisation of the functor $\mathbf{CGr} \rightarrow \mathbf{GS} \rightarrow \mathbf{GS}_*$, and, as in Section 6.4, let \mathbb{T}_* be the monad on \mathbf{GS}_* whose EM category of algebras is equivalent to the category \mathfrak{MO} of modular operads (in \mathbf{Set}).

So, there is a commuting diagram of functors

$$(8.1) \quad \begin{array}{ccccccc} & & \Xi^{\mathbb{C}} & \xrightarrow{\text{f.f.}} & \mathfrak{MO} & \xrightarrow{N} & \mathbf{psh}(\Xi) \\ & & \uparrow j \text{ b.o.} & & \uparrow \text{free}^{\mathbb{T}*} \downarrow \text{forget}^{\mathbb{T}*} & & \downarrow j^* \\ \mathbf{B}_*^{\S} & \xrightarrow{\text{f.f.}} & \mathbf{CGr}_* & \xrightarrow{\text{f.f.}} & \mathbf{GS}_* & \xrightarrow{\text{f.f.}} & \mathbf{psh}(\mathbf{CGr}_*) \\ & \uparrow \text{b.o.} & \uparrow \text{b.o.} & & \uparrow \text{free}^{\mathbb{D}} \downarrow \text{forget}^{\mathbb{D}} & & \downarrow \\ \mathbf{B}^{\S} & \xrightarrow{\text{f.f.}} & \mathbf{CGr} & \xrightarrow{\text{f.f.}} & \mathbf{GS} & \xrightarrow{\text{f.f.}} & \mathbf{psh}(\mathbf{CGr}) \end{array}$$

(Note: The diagram above is a simplified representation of the complex commutative diagram in the image. The original image shows a more detailed structure with additional arrows and labels like 'dense' and 'bo-ff'.)

where Ξ is the category, whose objects are indexed by connected graphs, obtained in bo-ff factorisation of the functor $\mathbf{CGr} \rightarrow \mathbf{GS} \rightarrow \mathbf{M}\mathbf{D}$.

In [41, Section 8], it was shown (for $\mathbf{V} = \mathbf{Set}$) that, though the modular operad monad $\mathbb{D}\mathbb{T}$ on \mathbf{GS} does not have arities, the monad \mathbb{T}_* on \mathbf{GS}_* does, whereby the following theorem holds:

Theorem 8.2. [41, Theorem 8.2] *The monad \mathbb{T}_* , and the category $\mathbf{CGr}_* \hookrightarrow \mathbf{GS}_*$, satisfy the conditions of [6] (\mathbb{T}_* has arities \mathbf{CGr}_*). Hence, the nerve $N: \mathbf{M}\mathbf{D} \rightarrow \mathbf{psh}(\Xi)$ is fully faithful. Its essential image consists of those presheaves P on Ξ such that $P(\mathcal{G}) = \lim_{(\mathcal{C}, b) \in \text{el}(\mathcal{G})} P(\mathcal{C})$ for all graphs \mathcal{G} .*

The circuit algebra monad $\mathbb{L}\mathbb{D}\mathbb{T}$ on \mathbf{GS} also does not have arities. However, since $\mathbf{GS}^{\mathbb{T}_*} \cong \mathbf{GS}^{\mathbb{D}\mathbb{T}}$ (Equation (6.25)) and \mathbb{L} lifts functorially to $\mathbf{GS}^{\mathbb{D}\mathbb{T}}$, the distributive law $\lambda_{\mathbb{L}\mathbb{T}}: \mathbb{T}\mathbb{L} \Rightarrow \mathbb{L}\mathbb{T}$ on \mathbf{GS} extends to a distributive law $T_*\tilde{\mathbb{L}} \Rightarrow \tilde{\mathbb{L}}T_*$ for \mathbb{T}_* and the lift $\tilde{\mathbb{L}}$ of \mathbb{L} to \mathbf{GS}_* . By Equation (7.9), the EM category of algebras for this composite monad $\tilde{\mathbb{L}}\mathbb{T}_*$ on \mathbf{GS}_* is precisely the category \mathbf{CA} of circuit algebras in \mathbf{Set} .

By Section 6.3 (see also [41, Section 7.3]), the full subcategory $\mathbf{Gr}_* \subset \mathbf{GS}_*$ of all graphs and pointed morphisms induces a fully faithful inclusion $\mathbf{GS}_* \hookrightarrow \mathbf{psh}(\mathbf{Gr}_*)$. Hence, if Ξ^\times is the category obtained in the bo-ff factorisation of $\mathbf{Gr} \rightarrow \mathbf{GS} \rightarrow \mathbf{CA}$, we may consider the following diagram of functors in which the left and middle squares commute, and the right squares commute up to natural isomorphism:

$$(8.3) \quad \begin{array}{ccccccc} & & \Xi^\times & \hookrightarrow & \mathbf{CA} & \xrightarrow{N_{\Xi^\times}} & \mathbf{psh}(\Xi^\times) \\ & & \uparrow j \text{ b.o.} & & \uparrow \text{free } \tilde{\mathbb{L}}\mathbb{T}_* & & \downarrow j^* \\ \mathbf{B}_*^{\S} & \hookrightarrow & \mathbf{Gr}_* & \hookrightarrow & \mathbf{GS}_* & \hookrightarrow & \mathbf{psh}(\mathbf{Gr}_*) \\ \uparrow \text{b.o.} & \nearrow \text{dense} & \uparrow \text{b.o.} & & \uparrow \text{free } \mathbb{D} & & \downarrow \\ \mathbf{B}^{\S} & \hookrightarrow & \mathbf{Gr} & \hookrightarrow & \mathbf{GS} & \hookrightarrow & \mathbf{psh}(\mathbf{Gr}) \end{array}$$

(Horizontal arrows are fully faithful (f.f.). Vertical arrows are b.o. (bifibration over objects).)

The remainder of this paper is dedicated to proving the following theorem using a modification of the proof of Equation (8.2) ([41, Theorem 8.2]) to show that \mathbf{Gr}_* provides arities for $\tilde{\mathbb{L}}\mathbb{T}_*$:

Theorem 8.4. *The functor $N: \mathbf{CA} \rightarrow \mathbf{psh}(\Xi^\times)$ is full and faithful. Its essential image consists of precisely those presheaves P on Ξ^\times such that*

$$(8.5) \quad P(\mathcal{G}) = \lim_{(\mathcal{C}, b) \in \text{el}(\mathcal{G})} P(\mathcal{C}) \text{ for all graphs } \mathcal{G}.$$

Recall that the category $\mathbf{W}\mathbf{P} = \mathbf{GS}^{\mathbb{L}\mathbb{D}\mathbb{T}}/Di \cong \mathbf{OGS}^{\mathbb{L}\mathbb{D}\mathbb{T}}$ is the category of wheeled props in \mathbf{Set} . Hence, Equation (8.4) implies an abstract nerve theorem for wheeled props. For, let $\mathbf{O}\Xi^\times \stackrel{\text{def}}{=} \Xi^\times/Di$ be the induced oriented graphical category. Since $\mathbf{psh}(\mathbf{G})/Di \simeq \mathbf{psh}(\mathbf{G}/Di)$ for $\mathbf{G} = \mathbf{Gr}, \mathbf{Gr}_*, \Xi^\times$, it follows that:

Corollary 8.6. *The induced functor $\mathbf{ON}: \mathbf{W}\mathbf{P} \rightarrow \mathbf{psh}(\mathbf{O}\Xi^\times)$ is full and faithful. Its essential image consists of precisely those presheaves P on $\mathbf{O}\Xi^\times$ that restrict to graphical species.*

Sections 8.2-8.4 are devoted to constructing the graphical category Ξ^\times and giving a direct proof of Equation (8.4), in the style of Equation (8.2) (see also, [41, Theorem 8.2]). Further proofs, using results from [6] and [11] on strongly cartesian monads, are outlined in Remarks 8.18 and 8.17.

8.2. The monad $\tilde{\mathbb{L}}\mathbb{T}_*$ on $\mathbf{GS}_{*\mathbf{V}}$. The first step is to understand the monad $\tilde{\mathbb{L}}\mathbb{T}_*$ on $\mathbf{GS}_{*\mathbf{V}}$. Here \mathbf{V} is, as usual, an arbitrary category with sufficient limits and colimits.

The lift $\tilde{\mathbb{L}}$ of \mathbb{L} to $\mathbf{GS}_{*\mathbf{V}}$ is given by $\tilde{\mathbb{L}}(S_*) = (LS, \eta^{\mathbb{L}}\epsilon, \eta^{\mathbb{L}}o)$, for all pointed graphical species $S_* = (S, \epsilon, o)$ in \mathbf{V} (where $\mu^{\tilde{\mathbb{L}}}, \eta^{\tilde{\mathbb{L}}}$ are induced by $\mu^{\mathbb{L}}$ and $\eta^{\mathbb{L}}$ in the obvious way.)

Let X be a finite set. The category $X\text{-CGr}_{\text{sim}}$ of similar connected X -graphs was defined in Section 6.4 and the monad $\mathbb{T}_* = (T_*, \mu^{\mathbb{T}_*}, \eta^{\mathbb{T}_*})$ with $T_*S_X = \text{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{sim}}} S(\mathcal{X})$ was discussed in Section 6.4.

Similarity morphisms preserve connected components. Hence the category $X\text{-Gr}_{\text{sim}}$ of *all* X -graphs and similarity morphisms, may be defined by

$$(8.7) \quad X\text{-Gr}_{\text{sim}} \stackrel{\text{def}}{=} \text{colim}_{(Y, f) \in \text{Core}(X/\text{Set}_f)} \prod_{y \in Y} (f^{-1}(y)) X\text{-CGr}_{\text{sim}}.$$

The endofunctor $\tilde{L}T_*$ is described by the obvious quotient of T^\times : Let $S_* = (S, \epsilon, o)$ be a pointed graphical species in \mathbf{V} and let X be a finite set, then

$$(8.8) \quad \begin{aligned} \tilde{L}T_*S_X &= \text{colim}_{(Y, f) \in \text{Core}(X/\text{Set}_f)} \left(\prod_{y \in Y} T_*S_{f^{-1}(y)} \right) \\ &= \text{colim}_{(Y, f) \in \text{Core}(X/\text{Set}_f)} \left(\prod_{y \in Y} \left(\text{colim}_{(\mathcal{G}, \rho) \in (f^{-1}(y))\text{-CGr}_{\text{sim}}} S(\mathcal{G}) \right) \right) \\ &= \text{colim}_{\mathcal{X} \in X\text{-Gr}_{\text{sim}}} S(\mathcal{X}). \end{aligned}$$

And,

$$\begin{aligned} T_*\tilde{L}S_X &= \text{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{sim}}} \lim_{(\mathcal{C}, b) \in \text{el}(\mathcal{X})} LS(\mathcal{C}) \\ &= \text{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{sim}}} \lim_{\mathbf{\Gamma} \in \text{Cor}_{\boxtimes}^{(\mathcal{X})}} S(\mathbf{\Gamma}(\mathcal{X})). \end{aligned}$$

As $\text{GS}_*^{\mathbb{T}_*} \cong \text{GS}_*^{\mathbb{DT}}$ and \mathbb{L} lifts to $\text{GS}_*^{\mathbb{DT}}$, there is a distributive law $T_*\tilde{L} \Rightarrow \tilde{L}T_*$, induced by the distributive law $\lambda_{\mathbb{L}\mathbb{T}}: TL \Rightarrow LT = T^\times$, which is compatible with similarity morphisms by construction: for each finite set X , a colimit indexed by pairs $(\mathcal{X}, \mathbf{\Gamma})$ of a connected admissible X -graph \mathcal{X} , and $\mathbf{\Gamma} \in \text{Cor}_{\boxtimes}^{(\mathcal{X})}$, is replaced by a colimit indexed over (in general not connected) admissible X -graphs $\mathbf{\Gamma}(\mathcal{X})$.

8.3. A graphical category for circuit algebras. For the remainder of the paper, $\mathbf{V} = \text{Set}$.

The category Gr_* obtained in the bo-ff factorisation of $\text{Gr} \rightarrow \text{GS} \rightarrow \text{GS}_*$ has been discussed in Section 6.3. Let Ξ^\times be the category, whose objects are graphs, obtained in the bo-ff factorisation of $\text{Gr} \rightarrow \text{GS} \rightarrow \mathfrak{CA}$ (and also of $\text{Gr}_* \rightarrow \text{GS}_* \rightarrow \mathfrak{CA}$). This is the full subcategory of the Kleisli category $\text{GS}_*^{\tilde{\mathbb{L}}T_*}$ of $\tilde{\mathbb{L}}T_*$ on free circuit algebras of the form $\tilde{L}T_*\Upsilon_*\mathcal{G} = LDT\Upsilon\mathcal{G}$ for graphs $\mathcal{G} \in \text{Gr}$. By definition, morphisms $\beta \in \Xi^\times(\mathcal{G}, \mathcal{H})$ are given by morphisms in $\text{GS}_*(\Upsilon_*\mathcal{G}, \tilde{L}T_*\Upsilon_*\mathcal{H})$.

To understand Ξ^\times , the first step is therefore to describe $\tilde{L}T_*\Upsilon_*\mathcal{H}$ for all graphs \mathcal{H} .

Since each connected component of $X\text{-Gr}_{\text{sim}}$ contains an (in general, not connected) admissible X -graph \mathcal{X} , it follows from (8.8) that, for all finite sets X , elements of $(\tilde{L}T_*\Upsilon_*\mathcal{H})_X \cong \Xi^\times(\mathcal{C}_X, \mathcal{H})$ are represented by pairs (\mathcal{X}, f) where \mathcal{X} is an admissible X -graph and $f \in \text{Gr}_*(\mathcal{X}, \mathcal{H})$. In particular, f factors uniquely as a morphism $\mathcal{X} \rightarrow \mathcal{X}_{W_f}$ in Gr_{sim} followed by a morphism $f^\perp \in \text{Gr}(\mathcal{X}_{W_f}, \mathcal{H})$. Pairs (\mathcal{X}^1, f^1) and (\mathcal{X}^1, f^2) represent the same element $[\mathcal{X}, f]_* \in (\tilde{L}T_*\Upsilon_*\mathcal{H})_X$ if and only if there is an object $\mathcal{X}_f \in X\text{-Gr}_{\text{sim}}$ and a commuting diagram

$$(8.9) \quad \begin{array}{ccccc} \mathcal{X}^1 & \xrightarrow{g^1} & \mathcal{X}^f & \xleftarrow{g^2} & \mathcal{X}^2 \\ & \searrow f^1 & \downarrow f^\perp & \swarrow f^2 & \\ & & \mathcal{H} & & \end{array}$$

in Gr_* such that g^j are similarity morphisms in $X\text{-Gr}_{\text{sim}}$ for $j = 1, 2$, and $f^\perp \in \text{Gr}(\mathcal{X}^f, \mathcal{H})$ is an (unpointed) étale morphism.

In particular, for all \mathcal{H} , $e \in E(\mathcal{H})$ and $m \geq 1$, the following special case of (8.9) commutes in \mathbf{Gr}_* :

$$(8.10) \quad \begin{array}{ccccc} \mathcal{C}_0 & \xrightarrow{z} & (1) & \xleftarrow{\kappa} & \mathcal{W}^m \\ & \searrow \text{ch}_e \circ z & \downarrow \text{ch}_e & \swarrow \text{ch}_e \circ \kappa & \\ & & \mathcal{H} & & \end{array}$$

Since $\Xi^\times(\mathcal{G}, \mathcal{H}) \cong \tilde{L}T_*\Upsilon_*\mathcal{H}(\mathcal{G}) = \lim_{(\mathcal{C}, b) \in \text{el}(\mathcal{G})} \tilde{L}T_*\Upsilon_*\mathcal{H}(\mathcal{C})$, morphisms $\gamma \in \Xi^\times(\mathcal{G}, \mathcal{H})$ are represented by pairs $(\mathbf{\Gamma}, f)$, where $\mathbf{\Gamma} : \text{el}(\mathcal{G}) \rightarrow \mathbf{Gr}$ is a nondegenerate \mathcal{G} -shaped graph of graphs, and $f \in \mathbf{Gr}_*(\mathbf{\Gamma}(\mathcal{G}), \mathcal{H})$ is a morphism in \mathbf{Gr}_* (see [41, Section 8.1]).

The following lemma generalises [41, Lemma 8.9], and is proved in exactly the same manner.

Lemma 8.11. *Let \mathcal{G} and \mathcal{H} be graphs and, for $i = 1, 2$, let $\mathbf{\Gamma}^i$ be a nondegenerate \mathcal{G} -shaped graph of graphs with colimit $\mathbf{\Gamma}^i(\mathcal{G})$, and $f^i \in \mathbf{Gr}_*(\mathbf{\Gamma}^i(\mathcal{G}), \mathcal{H})$. Then $(\mathbf{\Gamma}^1, f^1), (\mathbf{\Gamma}^2, f^2)$ represent the same element γ of $\Xi^\times(\mathcal{G}, \mathcal{H})$ if and only if there is a representative $(\mathbf{\Gamma}, f)$ of γ , and a commuting diagram in \mathbf{Gr}_* of the following form:*

$$(8.12) \quad \begin{array}{ccccc} \mathbf{\Gamma}^1(\mathcal{G}) & \xrightarrow{\quad} & \mathbf{\Gamma}(\mathcal{G}) & \xleftarrow{\quad} & \mathbf{\Gamma}^2(\mathcal{G}) \\ & \searrow f^1 & \downarrow f & \swarrow f^2 & \\ & & \mathcal{H} & & \end{array}$$

where the morphisms in the top row are vertex deletion morphisms (including $z : \mathcal{C}_0 \rightarrow (1)$) in \mathbf{Gr}_{sim} , and $f \in \mathbf{Gr}(\mathbf{\Gamma}(\mathcal{G}), \mathcal{H})$.

The following terminology is from [27].

Definition 8.13. *A (pointed) free morphism in Ξ^\times is a morphism in the image of the inclusion $\mathbf{Gr}_* \hookrightarrow \Xi^\times$. An unpointed free morphism in Ξ^\times is a morphism in the image of $\mathbf{Gr} \hookrightarrow \mathbf{Gr}_* \hookrightarrow \Xi^\times$. A refinement morphism $[\mathbf{\Gamma}]$ in Ξ^\times is a morphism in $\Xi^\times(\mathcal{G}, \mathcal{H})$ with a representative of the form $(\mathbf{\Gamma}, \text{id}_{\mathcal{H}})$.*

So, a free morphism in $\Xi^\times(\mathcal{G}, \mathcal{H})$ has a representative of the form (id, f) where $\text{id} : \text{el}(\mathcal{G}) \rightarrow \mathbf{Gr}$ is the trivial identity \mathcal{G} -shaped graph of graphs with colimit \mathcal{G} and $f \in \mathbf{Gr}_*(\mathcal{G}, \mathcal{H})$. Refinements $[\mathbf{\Gamma}]$ in $\Xi^\times(\mathcal{G}, \mathcal{H})$ are described by nondegenerate \mathcal{G} -shaped graph of graphs $\mathbf{\Gamma}$ with colimit \mathcal{H} , and hence induce isomorphisms $E_0(\mathcal{G}) \xrightarrow{\cong} E_0(\mathcal{H})$ on boundaries.

In particular, morphisms in $\Xi^\times(\mathcal{G}, \mathcal{H})$ factor as $\mathcal{G} \xrightarrow{[\mathbf{\Gamma}]} \mathbf{\Gamma}(\mathcal{G}) \xrightarrow{\text{del}_{\setminus W}} \mathbf{\Gamma}(\mathcal{G})_{\setminus W} \xrightarrow{f} \mathcal{H}$. Here $\mathbf{\Gamma} : \text{el}(\mathcal{G}) \rightarrow \mathbf{Gr}$ is a nondegenerate \mathcal{G} -shaped graph of graphs with colimit $\mathbf{\Gamma}(\mathcal{G}) \in \mathbf{Gr}$, and $f \in \mathbf{Gr}(\mathbf{\Gamma}(\mathcal{G})_{\setminus W}, \mathcal{H})$ is an (unpointed) étale morphism \mathbf{Gr} .

8.4. Factorisation categories and the nerve theorem. Let $\mathbf{GS}_{*\tilde{\mathbb{L}}T_*}$ be the Kleisli category of $\tilde{\mathbb{L}}T_*$ on \mathbf{GS}_* . By [6, Proposition 2.5], to show that the monad $\tilde{\mathbb{L}}T_*$, together with the category \mathbf{Gr}_* satisfy the conditions of the nerve theorem [6, Theorem 1.10] (that is, $\tilde{\mathbb{L}}T_*$ has arities \mathbf{Gr}_*) and therefore the nerve functor $N : \mathfrak{CA} \rightarrow \mathbf{psh}(\Xi^\times)$ is fully faithful, it is sufficient to show that certain categories, obtained by factoring morphisms in $\mathbf{GS}_{*\tilde{\mathbb{L}}T_*}$, are connected.

To construct these categories, let S_* be a pointed graphical species in \mathbf{Set} , \mathcal{G} a graph in \mathbf{Gr} , and let $\beta \in \mathbf{GS}_{*\tilde{\mathbb{L}}T_*}(\mathcal{G}, S_*)$ be a morphism in the Kleisli category of $\tilde{\mathbb{L}}T_*$. By the Yoneda lemma,

$$\mathbf{GS}_{*\tilde{\mathbb{L}}T_*}(\mathcal{G}, S_*) \cong \mathbf{GS}_*(\mathcal{G}, \tilde{L}T_*S_*) \cong \tilde{L}T_*S_*(\mathcal{G}) = \text{colim}_{(\mathcal{C}, b) \in \text{el}(\mathcal{G})} \tilde{L}T_*S_*(\mathcal{C})$$

where

$$\tilde{L}T_*S_*(\mathcal{C}) \cong \begin{cases} S_{\S} & \mathcal{C} \cong (1), \\ \text{colim}_{\mathcal{X} \in X - \mathbf{Gr}_{\text{sim}}} S(\mathcal{X}) & \mathcal{C} \cong \mathcal{C}_X. \end{cases}$$

So, $\beta \in \mathbf{GS}_{*\mathbb{T}_*}(\mathcal{G}, S_*)$ is represented by pairs $(\mathbf{\Gamma}, \alpha)$ of

- a nondegenerate \mathcal{G} -shaped graph of graphs $\mathbf{\Gamma}$ with colimit $\mathbf{\Gamma}(\mathcal{G})$, viewed as a refinement $[\mathbf{\Gamma}] \in \Xi^\times(\mathcal{G}, \mathbf{\Gamma}(\mathcal{G})) = \mathbf{GS}_{*\mathbb{T}_*}(\mathcal{G}, \mathbf{\Gamma}(\mathcal{G}))$,
- an element $\alpha \in S(\mathbf{\Gamma}(\mathcal{G}))$ viewed, by the Yoneda lemma, as a morphism $\alpha \in \mathbf{GS}_{*\mathbb{T}_*}(\Upsilon_*\mathbf{\Gamma}(\mathcal{G}), S_*)$

such that the induced composition $\mathcal{G} \xrightarrow{[\mathbf{\Gamma}]} \mathbf{\Gamma}(\mathcal{G}) \xrightarrow{\alpha} S(\mathbf{\Gamma}(\mathcal{G})) \rightarrow (\tilde{L}T_*S_*)(\mathcal{G})$ is precisely β .

Definition 8.14. Objects of the factorisation category $\mathbf{fact}_*(\beta)$ of β are pairs $(\mathbf{\Gamma}, \alpha)$ that represent β as above, and morphisms in $\mathbf{fact}_*(\beta)((\mathbf{\Gamma}^1, \alpha^1), (\mathbf{\Gamma}^2, \alpha^2))$ are commuting diagrams in $\mathbf{GS}_{*\tilde{L}T_*}$

$$(8.15) \quad \begin{array}{ccccc} & & [\mathbf{\Gamma}^1](\mathcal{G}) & & \\ & \nearrow & \downarrow g & \searrow \alpha^1 & \\ \mathcal{G} & & & & S_* \\ & \searrow & \downarrow g & \nearrow \alpha^2 & \\ & & [\mathbf{\Gamma}^2](\mathcal{G}) & & \end{array}$$

such that g is a morphism in $\mathbf{Gr}_* \hookrightarrow \mathbf{GS}_{*\tilde{L}T_*}$.

Lemma 8.16. For all pointed graphical species S_* in \mathbf{Set} , all graphs $\mathcal{G} \in \mathbf{Gr}$, and all $\beta \in \mathbf{GS}_*(\mathcal{G}, \tilde{L}T_*S)$, the category $\mathbf{fact}_*(\beta)$ is connected.

Proof. If $\mathcal{G} \cong (1)$, then $\mathbf{\Gamma}$ is isomorphic to the identity (1) -shaped graph of graphs. If $\mathcal{G} \cong \mathcal{C}_X$ for some finite set X , and $\beta \in \tilde{L}T_*S_X \cong \mathbf{GS}_{*\mathbb{T}_*}(\mathcal{G}, S_*)$, then objects of $\mathbf{fact}_*(\beta)$ have the form (\mathcal{X}, α) where $\mathcal{X} \in X\text{-Gr}_{\text{iso}}$ is an admissible (in general not connected) X -graph, $\alpha \in S_*(\mathcal{X})$ and β is the image of α under the universal morphism $S_*(\mathcal{X}) \rightarrow (\tilde{L}T_*S_*)_X$. Since $(\tilde{L}T_*S_*)_X = \text{colim}_{\mathcal{X}' \in X\text{-Gr}_{\text{sim}}} S(\mathcal{X}')$, if (\mathcal{X}', α') also represents β , then \mathcal{X} and \mathcal{X}' are connected in $X\text{-Gr}_{\text{sim}}$ and hence they are similar in \mathbf{Gr}_* . Hence, if $\mathcal{G} = \mathcal{C}$ is in the image of $\mathbf{B}_*^{\mathfrak{S}} \subset \mathbf{Gr}_*$, then $\mathbf{fact}_*(\beta)$ is connected for all $\beta \in \mathbf{GS}_{*\mathbb{T}_*}(\mathcal{C}, S_*) \cong \tilde{L}T_*S(\mathcal{C})$.

For general $\mathcal{G} \in \mathbf{Gr}_*$, and $\beta \in \mathbf{GS}_{*\mathbb{T}_*}(\mathcal{G}, S_*)$, observe first that, since colimits of graphs of graphs are computed componentwise, we may assume that \mathcal{G} is connected. Assume also that $\mathcal{G} \not\cong \mathcal{C}_0$. Let $(\mathbf{\Gamma}^i, \alpha^i)$, $i = 1, 2$ be objects of $\mathbf{fact}_*(\beta)$. Then, for each $(\mathcal{C}_X, b) \in \text{el}(\mathcal{G})$, $\mathbf{\Gamma}^1(b)$ and $\mathbf{\Gamma}^2(b)$ are in the same connected component of $X\text{-Gr}_{\text{sim}}$, since the lemma holds when $\mathcal{G} \cong \mathcal{C}_X$. Therefore, $\mathbf{\Gamma}^1(\mathcal{G})$ and $\mathbf{\Gamma}^2(\mathcal{G})$ are similar in \mathbf{Gr}_* . \square

Equation (8.4) now follows from [6, Sections 1 & 2].

Proof of Equation (8.4). The induced nerve functor $N: \mathfrak{CA} \rightarrow \mathbf{psh}(\Xi^\times)$ is fully faithful by Equation (8.16), [6, Proposition 2.5], and [6, Propositions 1.5 & 1.9].

Moreover, by [6, Theorem 1.10], its essential image is the subcategory of those presheaves on Ξ^\times whose restriction to \mathbf{Gr}_* are in the image of the fully faithful embedding $\mathbf{GS}_* \hookrightarrow \mathbf{psh}(\mathbf{Gr}_*)$. So, a presheaf P on Ξ^\times described the nerve of a circuit algebra if and only if $P(\mathcal{G}) \cong \lim_{(\mathcal{C}, b) \in \text{el}_*(\mathcal{G})} P(\mathcal{C})$, and hence, by finality of $\text{el}(\mathcal{G}) \subset \text{el}_*(\mathcal{G})$, $P(\mathcal{G}) = \lim_{(\mathcal{C}, b) \in \text{el}(\mathcal{G})} P(\mathcal{C})$. Therefore, the Segal condition (8.5) is satisfied and Equation (8.4) is proved. \square

Following from Remarks 2.20 and 2.19, Equation (8.4) also admits proofs using extendable algebraic patterns [11], and extensions of strongly cartesian monads as described in [6, Proposition 4.4]:

Remark 8.17. Recall Remarks 2.11 and 2.20. By (8.8), for all finite sets X ,

$$\begin{aligned} \tilde{L}T_*S_X &= \text{colim}_{\mathcal{X} \in X\text{-Gr}_{\text{sim}}} S(\mathcal{X}) \\ &= \text{colim}_{\mathcal{X} \in X\text{-Gr}_{\text{sim}}} \lim_{(\mathcal{C}, b) \in \text{el}(\mathcal{X})} S(\mathcal{C}) \end{aligned}$$

And, by the previous discussion, a morphism in Ξ^\times factor as a refinement (induced by a graphs of graphs) followed by a Gr_* morphism.

Hence (e.g., by [11, Section 1]) Ξ^\times describes an extendable algebraic pattern, and $\tilde{\mathbb{L}}\mathbb{T}_*$ is the associated strongly cartesian monad, with canonical arities Gr_* . Equation (8.4) then follows immediately from [6, 51].

Remark 8.18. Recall Equation (2.19). By (6.1), for all finite sets X ,

$$T^\times S_X = \text{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{iso}}} S(\mathcal{X}) = \text{colim}_{\mathcal{X} \in X\text{-CGr}_{\text{iso}}} \lim_{(\mathcal{C}, b) \in \text{el}(\mathcal{X})} S(\mathcal{C}).$$

So (e.g., by [11, Section 1]) \mathbb{T}^\times is strongly cartesian with canonical arities $\text{Gr} \subset \text{GS}$.

By Section 6.2, the right adjoint functor $i^*: \text{GS}_* \rightarrow \text{GS}$ associated to the monad \mathbb{D} on GS is just the pullback to presheaves of the bijective on objects inclusion $i: \mathbf{B}^{\mathfrak{s}} \hookrightarrow \mathbf{B}_*^{\mathfrak{s}}$. And, by construction, $(LT)i^* = i^*(\tilde{L}T_*)$: $\text{GS}_* \rightarrow \text{GS}$ so the monad $\tilde{\mathbb{L}}\mathbb{T}_*$ on GS_* is the pushforward along i of $\mathbb{L}\mathbb{T} = \mathbb{T}^\times$ on GS . Hence, by [6, Proposition 4.4], $\tilde{\mathbb{L}}\mathbb{T}_*$ on GS_* is strongly cartesian with canonical arities Gr_* .

It remains, finally to remark on the homotopy theory of circuit algebras (see also the Introduction).

Remark 8.19. Both the nerve theorem [41, Theorem 8.2] for modular operads on which Equation (8.4) is built, as well as Hackney, Robertson and Yau's nerve theorem [24, 3.6], lead to well-defined models of $(\infty, 1)$ -modular operads in which the fibrant objects in the appropriate category of presheaves are precisely those that satisfy the weak Segal condition: in the case of [41] it followed from the results of [9] that these were given by $P: \Xi^{\text{op}} \rightarrow \mathbf{sSet}$ such that, for all (connected) graphs $\mathcal{G} \in \Xi$,

$$P(\mathcal{G}) \simeq \lim_{(\mathcal{C}, b) \in \text{el}(\mathcal{G})} P(\mathcal{C}).$$

However, describing a similar model for circuit algebras is more challenging. The presence of unconnected graphs means that, neither the method of [24], nor that of [41, Corollary 8.14] may be directly extended to a model structure on $\mathbf{psh}_{\mathbf{sSet}}(\Xi^\times)$ (or some subcategory thereof) that gives a good description of weak circuit algebras. It is for later work to investigate the existence of a such a model.

REFERENCES

- [1] Michael Artin, Alexander Grothendieck, J. L. Verdier, N. Bourbaki, P. Deligne, and Bernard Saint-Donat, editors. *Séminaire de géométrie algébrique du Bois-Marie 1963–1964. Théorie des topos et cohomologie étale des schémas. (SGA 4). Un séminaire dirigé par M. Artin, A. Grothendieck, J. L. Verdier. Avec la collaboration de N. Bourbaki, P. Deligne, B. Saint-Donat. Tome 1: Théorie des topos. Exposés I à IV. 2e éd.*, volume 269 of *Lect. Notes Math.* Springer, Cham, 1972.
- [2] Markus Banagl. High-Dimensional Topological Field Theory, Positivity, and Exotic Smooth Spheres. Preprint, arXiv:1508.01337 [math.AT], 2015.
- [3] Dror Bar-Natan and Zsuzsanna Dancso. Finite type invariants of w-knotted objects II: tangles, foams and the Kashiwara-Vergne problem. *Math. Ann.*, 367(3-4):1517–1586, 2017.
- [4] M. A. Batanin and C. Berger. Homotopy theory for algebras over polynomial monads. *Theory Appl. Categ.*, 32:148–253, 2017.
- [5] Jon Beck. Distributive laws. In *Sem. on Triples and Categorical Homology Theory (ETH, Zürich, 1966/67)*, pages 119–140. Springer, Berlin, 1969.
- [6] Clemens Berger, Paul-André Melliès, and Mark Weber. Monads with arities and their associated theories. *J. Pure Appl. Algebra*, 216(8-9):2029–2048, 2012.
- [7] John Bourke and Richard Garner. Monads and theories. *Adv. Math.*, 351:1024–1071, 2019.
- [8] Richard Brauer. On algebras which are connected with the semisimple continuous groups. *Ann. of Math. (2)*, 38(4):857–872, 1937.
- [9] Giovanni Caviglia and Geoffroy Horel. Rigidification of higher categorical structures. *Algebr. Geom. Topol.*, 16(6):3533–3562, 2016.
- [10] Eugenia Cheng. Iterated distributive laws. *Math. Proc. Cambridge Philos. Soc.*, 150(3):459–487, 2011.

- [11] Hongyi Chu and Rune Haugseng. Homotopy-coherent algebra via Segal conditions. *Adv. Math.*, 385:95, 2021. Id/No 107733.
- [12] Celeste Damiani and Vincent Florens. Alexander invariants of ribbon tangles and planar algebras. *J. Math. Soc. Japan*, 70(3):1063–1084, 2018.
- [13] Zsuzsanna Dancso, Iva Halacheva, and Marcy Robertson. Circuit algebras are wheeled props. *J. Pure Appl. Algebra*, 225(12):106767, 33, 2021.
- [14] Zsuzsanna Dancso, Iva Halacheva, and Marcy Robertson. A topological characterisation of the Kashiwara-Vergne groups. *Trans. Am. Math. Soc.*, 376(5):3265–3317, 2023.
- [15] Harm Derksen and Visu Makam. Invariant theory and wheeled PROPs. *J. Pure Appl. Algebra*, 227(9):30, 2023. Id/No 107302.
- [16] Adrián Doña Mateo. Pushforward monads. *Theory Appl. Categ.*, 44:927–963, 2025.
- [17] Gabriel C. Drummond-Cole and Philip Hackney. Dwyer-Kan homotopy theory for cyclic operads. *Proc. Edinb. Math. Soc., II. Ser.*, 64(1):29–58, 2021.
- [18] Jacques Gignoux, Guillaume Chérel, Ian D. Davies, Shayne R. Flint, and Eric Lateltin. Emergence and complex systems: The contribution of dynamic graph theory. *Ecological Complexity*, 31:34–49, 2017.
- [19] David G. Green. Emergence in complex networks of simple agents. *Journal of Economic Interaction and Coordination*, 18(3):419–462, July 2023.
- [20] Philip Hackney. Categories of graphs for operadic structures. *Math. Proc. Camb. Philos. Soc.*, 176(1):155–212, 2024.
- [21] Philip Hackney. Segal conditions for generalized operads. In *Higher structures in topology, geometry, and physics. AMS special session, virtual, March 26–27, 2022*, pages 161–194. Providence, RI: American Mathematical Society (AMS), 2024.
- [22] Philip Hackney, Marcy Robertson, and Donald Yau. *Infinity properads and infinity wheeled properads*, volume 2147 of *Lecture Notes in Mathematics*. Springer, Cham, 2015.
- [23] Philip Hackney, Marcy Robertson, and Donald Yau. A graphical category for higher modular operads. *Adv. Math.*, 365:107044, 2020.
- [24] Philip Hackney, Marcy Robertson, and Donald Yau. Modular operads and the nerve theorem. *Adv. Math.*, 370:107206, 39, 2020.
- [25] Iva Halacheva. Alexander type invariants of tangles, 2016. arXiv: 1611.09280.
- [26] V. F. R. Jones. The Potts model and the symmetric group. In *Subfactors (Kyuzeso, 1993)*, pages 259–267. World Sci. Publ., River Edge, NJ, 1994.
- [27] A. Joyal and J. Kock. Feynman graphs, and nerve theorem for compact symmetric multicategories (extended abstract). *Electronic Note in Theoretical Computer Science*, 270(2):105 – 113, 2011.
- [28] André Joyal. Une théorie combinatoire des séries formelles. *Adv. in Math.*, 42(1):1–82, 1981.
- [29] Ralph M. Kaufmann and Benjamin C. Ward. Schwarz Modular Operads Revisited. Preprint, arXiv:2404.17540 [math.AT], 2024.
- [30] Joachim Kock. Graphs, hypergraphs, and properads. *Collect. Math.*, 67(2):155–190, 2016.
- [31] Joachim Kock. Cospan construction of the graph category of Borisov and Manin. *Publ. Mat.*, 62(2):331–353, 2018.
- [32] A. Kupers and O. Randal-Williams. On the Torelli Lie algebra. *Forum Math. Pi*, 11:47, 2023. Id/No e13.
- [33] Joachim Lambek. *Completions of categories*. Seminar lectures given in 1966 in Zürich. Lecture Notes in Mathematics, No. 24. Springer-Verlag, Berlin-New York, 1966.
- [34] G. I. Lehrer and R. B. Zhang. The Brauer category and invariant theory. *J. Eur. Math. Soc. (JEMS)*, 17(9):2311–2351, 2015.
- [35] Saunders Mac Lane. *Categories for the working mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1998.
- [36] M. Markl, S. Merkulov, and S. Shadrin. Wheeled PROPs, graph complexes and the master equation. *J. Pure Appl. Algebra*, 213(4):496–535, 2009.
- [37] Martin Markl. Distributive laws and Koszulness. *Ann. Inst. Fourier*, 46(2):307–323, 1996.
- [38] Sergei A. Merkulov. Wheeled props in algebra, geometry and quantization. In *European Congress of Mathematics*, pages 83–114. Eur. Math. Soc., Zürich, 2010.
- [39] Ieke Moerdijk and Ittay Weiss. Dendroidal sets. *Algebr. Geom. Topol.*, 7:1441–1470, 2007.
- [40] Evan Patterson, Andrew Baas, Timothy Hosgood, and James Fairbanks. A diagrammatic view of differential equations in physics. *Mathematics in Engineering*, 5(2):1–59, 2023.
- [41] Sophie Raynor. Graphical combinatorics and a distributive law for modular operads. *Adv. Math.*, 392:Paper No. 108011, 87, 2021.
- [42] Sophie Raynor. Functorial, operadic and modular operadic combinatorics of circuit algebras. *J. Pure Appl. Algebra*, 229(11):37, 2025. Id/No 108105.

- [43] Hebing Rui and Linliang Song. Representations of Brauer category and categorification. *J. Algebra*, 557:1–36, 2020.
- [44] Steven V. Sam and Andrew Snowden. Stability patterns in representation theory. *Forum Math. Sigma*, 3:Paper No. e11, 108, 2015.
- [45] Albert Schwarz. Grassmannian and string theory. *Communications in Mathematical Physics*, 199:1–24, 1996.
- [46] Kurt Stoeckl. Koszul operads governing props and wheeled props. *Adv. Math.*, 454:80, 2024. Id/No 109869.
- [47] Robin Stoll. Modular operads as modules over the Brauer properad. *Theor. Appl. Categor.*, 38:1538–1607, 2022.
- [48] Ross Street. The formal theory of monads. *J. Pure Appl. Algebra*, 2:149–168, 1972.
- [49] Daniel Tubbenhauer. Virtual Khovanov homology using cobordisms. *J. Knot Theory Ramifications*, 23(9):1450046, 91, 2014.
- [50] Benjamin C. Ward. Massey products for graph homology. *Int. Math. Res. Not.*, 2022(11):8086–8161, 2022.
- [51] Mark Weber. Familial 2-functors and parametric right adjoints. *Theory Appl. Categ.*, 18:No. 22, 665–732, 2007.
- [52] Donald Yau and Mark W. Johnson. *A foundation for PROPs, algebras, and modules*, volume 203 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2015.