

# Higher rank Nichols algebras of diagonal type over fields of positive characteristic

Lijie Lei, Chen Yuan, Chen Qian, Jing Wang <sup>\*†</sup>

## Abstract

The classification of Nichols algebras is an essential step in the classification theory of pointed Hopf algebras by lifting method of N. Andruskiewitsch and H.-J. Schneider. Arithmetic root systems are invariants of Nichols algebras of diagonal type with a certain finiteness property. In this paper, all rank  $r \geq 5$  Nichols algebras of diagonal type with a finite irreducible root system over fields of positive characteristic are classified. Weyl groupoids and finite arithmetic root systems are crucial tools for our classification.

**Key Words:** Nichols algebra, Cartan graph, Weyl groupoid, Arithmetic root system

## 1 Introduction

The theory of Nichols algebras is motivated by the Hopf algebra theory and plays an important role in Hopf algebra. Besides, it has applications to other research fields such as Kac-Moody Lie superalgebras and conformal field theory [2, 19, 35, 40–42].

In Hopf algebra theory, the classification of all finite dimensional Hopf algebras is a tough question [1]. The classification of finite dimensional Nichols algebras is an essential step in the N. Andruskiewitsch and H.-J. Schneider lifting method for classifying finite dimensional pointed Hopf algebras [7–10]. The structure of Nichols algebras appears naturally in the classification of finite dimensional pointed Hopf algebras, as follows. Given a Hopf algebra  $H$ , consider its coradical filtration

$$H_0 \subset H_1 \subset \dots$$

such that  $H_0$  is a Hopf subalgebra of  $H$  and the associated graded coalgebra

$$\mathrm{gr}H = \bigoplus_{n \geq 0} H_n/H_{n-1},$$

where  $H_{-1} = 0$ . Then  $\mathrm{gr}H$  is a graded Hopf algebra, since the coradical  $H_0$  of  $H$  is a Hopf subalgebra. In addition, consider the projection  $\pi : \mathrm{gr}H \rightarrow H_0$ ; let  $R$  be the algebra of

---

<sup>\*</sup>Corresponding author: wang\_jing619@163.com

<sup>†</sup>The author is partially supported by National Natural Science Foundation of China(No.12471038).

coinvariants of  $\pi$ . Then, by a result of Radford and Majid,  $R$  is a braided Hopf algebra and  $\text{gr}H$  is the bosonization (or biproduct) of  $R$  and  $H_0$ :  $\text{gr}H \simeq R \# H_0$ . The principle of the “Lifting method” is first introduced to study  $R$ , then to transfer the information to  $\text{gr}H$  via bosonization, and finally to lift to  $H$  [7, 9]. The braided Hopf algebra  $R$  is generated by the vector space  $V$  of  $H_0$ -coinvariants of  $H_1/H_0$ , namely Nichols algebra  $\mathcal{B}(V)$  which is generated by  $V$  in commemoration of W. Nichols who started to study these objects as bialgebras of type one [7, 38]. Indeed, the conjecture on generation in degree one by N. Andruskiewitsch and H.-J. Schneider fails in positive characteristic. This is an important task for future work. Regarding the description of Nichols algebras, there are many different but alternative ways, see for example [36, 37, 39, 46, 47].

The crucial step to classify pointed Hopf algebras is to determine all Nichols algebras  $\mathcal{B}(V)$  that are finite dimensional. N. Andruskiewitsch stated the following question.

**Question 1.1. (N. Andruskiewitsch [1])** *Given a braiding matrix  $(q_{ij})_{1 \leq i, j \leq \theta}$  whose entries are roots of 1, when  $\mathcal{B}(V)$  is finite dimensional, where  $V$  is a vector space with basis  $x_1, \dots, x_\theta$  and braiding  $c(x_i \otimes x_j) = q_{ij}(x_j \otimes x_i)$ ? If so, compute  $\dim_{\mathbb{k}} \mathcal{B}(V)$ , and give a “nice” presentation by generators and relations.*

Several authors have obtained the classification results for infinite and finite dimensional Nichols algebras of Cartan type, see [8, 22, 39]. I. Heckenberger has classified all finite dimensional Nichols algebras of diagonal type, see [21, 23, 24, 26]. The generators and relations of such Nichols algebras are also given [11, 12]. With the classification results, N. Andruskiewitsch and H.-J. Schneider [10] have obtained a classification theorem about finite dimensional pointed Hopf algebras under some technical assumptions. Then the classification of finite dimensional pointed Hopf algebras with abelian coradical is established in a series of papers [3, 13, 14, 20]. Based on such successful applications, the analysis to the structure of Nichols algebras over arbitrary fields is crucial and has potential applications. However, significant differences emerge in positive characteristic  $p > 0$  due to the existence of  $p$ -roots: for any  $q \in \mathbb{k} \setminus \{0\}$  and  $n \in \mathbb{N}$ , the quantum integer  $(n)_q := 1 + q + \dots + q^{n-1}$  vanishes if and only if  $q^n = 1$  or  $p|n$ , where  $\mathbb{k}$  is a field of characteristic  $p$ . Over fields of positive characteristic, finite dimensional rank 2, rank 3, and rank 4 Nichols algebras of diagonal type are classified in [31, 44, 45]. Those classification results yield some important applications in pointed Hopf algebras and Nichols algebras [6, 27]. These low-rank results have already also demonstrated that the classification framework for characteristic zero cannot be directly extended—new concepts and tools are needed to capture the unique structures in positive characteristic. The methods used are the same as in rank 3 and 4. The present work focuses on the classification of rank  $r \geq 5$  Nichols algebras of diagonal type with finite irreducible root systems in positive characteristic.

The main theoretical tools for the classification of Nichols algebra  $\mathcal{B}(V)$  are the root system and the Weyl groupoid of  $\mathcal{B}(V)$ . I. Heckenberger in [22] defines the arithmetic root system and the Weyl groupoid of Nichols algebra of diagonal type based on the ideas of V. Kharchenko on Poincaré–Birkhoff–Witt basis of Nichols algebra of diagonal type [34]. Later, M. Cuntz, I. Heckenberger and H. Yamane present a combinatorial theory of these two structures [16, 32]. Then the theory of root systems and Weyl groupoids is developed for more general Nichols algebras [5, 28, 29]. Further, to more clearly see the extent of generality, M. Cuntz and I. Heckenberger achieve a classification of finite Weyl groupoids in [17, 18].

The authors focus on connected simply connected Cartan graphs such that the real roots form a finite irreducible root system. Such Cartan graphs are called *finite Cartan graphs*. In specific, they state the following theorem in [18]. This result provides the framework for our study.

**Theorem 1.2.** *There are exactly three families of connected simply connected Cartan graphs for which the real roots form a finite irreducible root system:*

- (1) *The family of Cartan graphs of rank two parametrized by triangulations of a convex  $n$ -gon by non-intersecting diagonals.*
- (2) *For each rank  $r > 2$ , the standard Cartan graphs of type  $A_r, B_r, C_r$  and  $D_r$ , and a series of  $r - 1$  further Cartan graphs described explicitly in Theorem 3.8.*
- (3) *A family consisting of 74 further “sporadic” Cartan graphs (including those of type  $F_4, E_6, E_7$  and  $E_8$ ).*

The paper is organized as follows. In Section 2, the main notations and some general results are recalled. One general result implies that we can attach a connected finite Cartan graph of rank  $r$  to a tuple  $M = (M_1, \dots, M_r)$  of one-dimensional Yetter–Drinfel’d over an abelian group. If  $r = 1$ , the Cartan graph contains no information about  $\mathcal{B}(V)$ . Therefore, we give restriction  $r \geq 2$ . We refer to a Cartan graph as *non-sporadic* if it is not sporadic. In Section 3, we provide an explicit characterization of finite Cartan graphs of rank  $r \geq 5$ . To do so, we divide into two subsections: non-sporadic finite Cartan graphs and sporadic Cartan finite graphs. In the first one we briefly recall the construction of root systems of finite Cartan graphs of type  $A_r, B_r, C_r, D_r$  and a series of  $r - 1$  further Cartan graphs of type  $D'(r, s)$  for  $s \in \{1, \dots, r - 1\}$ . In order to cover all such Cartan graphs of type  $D'(r, s)$  for  $s \in \{1, \dots, r - 1\}$ , we define “good  $D'_r$  neighborhood” by GAP software. In Theorem 3.11, we prove that every non-sporadic finite Cartan graph of rank  $r \geq 5$  is either standard or with a point which has the good  $D'_r$  neighborhood. In the second subsection we introduce the concepts of good  $A_r$  neighborhoods to explicitly characterize the sporadic finite Cartan graphs. And we formulate such properties in Theorem 3.15. Finally, in Section 4 we formulate the classification Theorem 4.1. As a corollary of Theorem 4.1, all rank  $r \geq 5$  finite dimensional Nichols algebras of diagonal type in positive characteristic are given, see Corollary 4.2. To conclude, Table 1 consists of all corresponding generalized Dynkin diagrams and in Table 2 we present the corresponding exchange graphs for the sporadic case of  $\mathcal{C}(M)$ .

Throughout this paper,  $\mathbb{k}$  always denotes a field of characteristic  $p > 0$ . Assume  $\mathbb{k}^* = \mathbb{k} \setminus \{0\}$ ,  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ ,  $G'_n = \{q \in \mathbb{k}^* \mid q^n = 1, q^k \neq 1 \text{ for all } 1 \leq k < n\}$  for  $n \in \mathbb{N}$ ,  $2 \leq r \in \mathbb{N}_0$ ,  $I = \{1, 2, 3, \dots, r\}$  and  $(\alpha_i)_{i \in I}$  be the standard basis of  $\mathbb{Z}^I$ .

## 2 Preliminaries

We recall the following standard definitions [15, 26, 31, 32, 44, 45] and refer to [30] for details.

## 2.1 Yetter–Drinfel’d modules and braided vector space

**Definition 2.1.** Let  $V$  be a  $r$ -dimensional vector space over  $\mathbb{k}$ . The pair  $(V, c)$  is called a braided vector space, if  $c \in \text{Aut}(V \otimes V)$  is a solution of the braid equation, that is

$$(c \otimes \text{id})(\text{id} \otimes c)(c \otimes \text{id}) = (\text{id} \otimes c)(c \otimes \text{id})(\text{id} \otimes c).$$

A braided vector space  $(V, c)$  is of diagonal type if  $V$  admits a basis  $\{x_i | i \in I\}$  such that for all  $i, j \in I$  one has

$$c(x_i \otimes x_j) = q_{ij} x_j \otimes x_i \quad \text{for some } q_{ij} \in \mathbb{k}^*.$$

**Definition 2.2.** The matrix  $(q_{ij})_{i,j \in I}$  is termed of the braiding matrix of  $V$ . The braiding matrix  $(q_{ij})_{i,j \in I}$  is indecomposable if for any  $i \neq j$  there exists a sequence  $i_1 = i, i_2, \dots, i_t = j$  of elements of  $I$  such that  $q_{i_s i_{s+1}} q_{i_{s+1} i_s} \neq 1$ , where  $1 \leq s \leq t-1$ .

This paper studies the braided vector spaces of diagonal type with an indecomposable braiding matrix.

**Definition 2.3.** Let  $H$  be a Hopf algebra. A Yetter–Drinfel’d module  $V$  over  $H$  is a left  $H$ -module with left action  $\lambda_L : H \otimes V \longrightarrow V$  and a left  $H$ -comodule with left coaction  $\delta_L : V \longrightarrow H \otimes V$  satisfying the compatibility condition

$$\delta_L(h.v) = h_{(1)} v_{(-1)} \kappa(h_{(3)}) \otimes h_{(2)}.v_{(0)}, \quad h \in H, v \in V.$$

We say that  $V$  is of diagonal type if  $H = \mathbb{k}G$  and  $V$  is a direct sum of one-dimensional Yetter–Drinfel’d modules over the group algebra  $\mathbb{k}G$ , where  $G$  is abelian.

We denote by  ${}^H_H\mathcal{YD}$  the category of Yetter–Drinfel’d modules over  $H$ , where morphisms preserve both the action and the coaction of  $H$ . The category  ${}^H_H\mathcal{YD}$  is braided with braiding

$$c_{V,W} : V \otimes W \longrightarrow W \otimes V, \quad v \otimes w \longrightarrow v_{(-1)}.w \otimes v_{(0)} \quad (1)$$

for all  $V, W \in {}^H_H\mathcal{YD}$ ,  $v \in V$ , and  $w \in W$ . Actually, the category  ${}^H_H\mathcal{YD}$  is a braided monoidal category, where the monoidal structure is given by the tensor product over  $\mathbb{k}$ . Then any Yetter–Drinfel’d module  $V \in {}^H_H\mathcal{YD}$  over  $H$  admits a braiding  $c_{V,V}$  and hence  $(V, c_{V,V})$  is a braided vector space. Conversely, any braided vector space can be realized as a Yetter–Drinfel’d module over some Hopf algebras if and only if the braiding is rigid [43, Section 2]. Notice that Yetter–Drinfel’d module structures on  $V$  with different Hopf algebras can give the same braiding and not all braidings of  $V$  are induced by the above Equation (1).

Let  $H = \mathbb{k}G$ . We write  ${}^G_G\mathcal{YD}$  for the category of Yetter–Drinfel’d modules over  $\mathbb{k}G$  and say that  $V \in {}^G_G\mathcal{YD}$  is a Yetter–Drinfel’d module over  $G$ . Notice that if  $V$  is of diagonal type, then  $(V, c_{V,V})$  is a braided vector space of diagonal type. Any braided vector space of diagonal type is also a Yetter–Drinfel’d module of diagonal type. Indeed, assume that  $(V, c)$  is a braided vector space of diagonal type with an indecomposable braiding matrix  $(q_{ij})_{i,j \in I}$  of a basis  $\{x_i | i \in I\}$ . Let  $G_0$  be a free abelian group generated by elements  $\{g_i | i \in I\}$ . Define the left coaction and left action by

$$\delta_L : V \longrightarrow \mathbb{k}G_0 \otimes V, \quad x_i \longrightarrow g_i \otimes x_i, \quad g_i.x_j = q_{ij}x_j \in V.$$

Then  $V = \bigoplus_{i \in I} \mathbb{k}x_i$  and each  $\mathbb{k}x_i$  is one-dimensional Yetter–Drinfel’d modules over  $G_0$ . Hence  $V$  is a Yetter–Drinfel’d module of diagonal type over  $G_0$ .

## 2.2 Nichols Algebras of Diagonal Type

In this section, we recall a definition of the Nichols algebra  $\mathcal{B}(V)$  generated by  $(V, c)$ .

**Definition 2.4.** *Let  $H$  be a Hopf algebra. A braided Hopf algebra in  ${}^H_H\mathcal{YD}$  is a 6-tuple  $\mathcal{B} = (\mathcal{B}, \mu, 1, \delta, \epsilon, \mathcal{S})$ , where  $(\mathcal{B}, \mu, 1)$  is an algebra in  ${}^H_H\mathcal{YD}$  and a coalgebra  $(\mathcal{B}, \delta, \epsilon)$  in  ${}^H_H\mathcal{YD}$ , and  $\mathcal{S} : \mathcal{B} \rightarrow \mathcal{B}$  is a morphism in  ${}^H_H\mathcal{YD}$  such that  $\delta, \epsilon$  and  $\mathcal{S}$  satisfy  $\mathcal{S}(b^{(1)})b^{(2)} = b^{(1)}\mathcal{S}(b^{(2)}) = \epsilon(b)1$ , where we define  $\delta(b) = b^{(1)} \otimes b^{(2)}$  as the coproduct of  $\mathcal{B}$  to avoid the confusion.*

Let  $(V, c)$  be a  $r$ -dimensional braided vector space of diagonal type. The Nichols algebras generated by  $(V, c)$  is defined as follows.

**Definition 2.5.** *The Nichols algebra generated by  $V \in {}^H_H\mathcal{YD}$  is defined as the quotient*

$$\mathcal{B}(V) = T(V)/\mathcal{I}(V) = (\oplus_{n=0}^{\infty} V^{\otimes n})/\mathcal{I}(V)$$

where  $\mathcal{I}(V)$  is the unique maximal coideal among all coideals of  $T(V)$  which are contained in  $\oplus_{n \geq 2} V^{\otimes n}$ . Nichols algebra  $\mathcal{B}(V)$  is said to be of diagonal type if  $V$  is a Yetter–Drinfel’d module of diagonal type. The dimension of  $V$  is the rank of Nichols algebra  $\mathcal{B}(V)$ .

Note that if  $B \in {}^H_H\mathcal{YD}$  and  $B$  is an algebra in  ${}^H_H\mathcal{YD}$  then  $B \otimes B$  is an algebra in  ${}^H_H\mathcal{YD}$  with the product given by

$$(a \otimes b)(c \otimes d) = a(b_{(-1)} \cdot c) \otimes b_{(0)}d, \quad (2)$$

for all  $a, b, c, d \in B$ , where  $\cdot$  denotes the left action of  $H$  on  $B$ .

The tensor algebra  $T(V)$  admits a natural structure of a Yetter–Drinfel’d module and an algebra structure in  ${}^H_H\mathcal{YD}$ . It is an  $\mathbb{N}_0$ -graded braided Hopf algebra in  ${}^H_H\mathcal{YD}$  with coproduct  $\delta(v)$  and counit  $\epsilon$  given by  $\delta(v) = 1 \otimes v + v \otimes 1$  and  $\epsilon(v) = 0$  for all  $v \in V$  such that  $\delta$  and  $\epsilon$  are the algebra morphisms. The antipode of  $T(V)$  exists [9, Section 2.1]. Notice that the product defined by Equation (2) on  $T(V)$  is the unique algebra structure such that  $\delta(v) = 1 \otimes v + v \otimes 1 \in T(V) \otimes T(V)$  for all  $v \in V$ . And the coproduct can be extended from  $V$  to  $T(V)$ . For example, for all  $v, w \in V$  one gets (we write the elements of  $T(V)$  without the tensor product sign for brevity)

$$\begin{aligned} \delta(vw) &= \delta(v)\delta(w) \\ &= (1 \otimes v + v \otimes 1)(1 \otimes w + w \otimes 1) \\ &= 1 \otimes vw + v_{(-1)} \cdot w \otimes v_{(0)} + v \otimes w + vw \otimes 1. \end{aligned}$$

Let  $(I_i)_{i \in I}$  be the family of all ideals of  $T(V)$  contained in  $\oplus_{n \geq 2} V^{\otimes n}$ , i.e.

$$\delta(I_i) \subset I_i \otimes T(V) + T(V) \otimes I_i.$$

Hence  $\mathcal{I}(V)$  is also an ideal, so  $\mathcal{B}(V)$  is a braided Hopf algebra in  ${}^H_H\mathcal{YD}$ . By [4, Proposition 3.2.12], Nichols algebra  $\mathcal{B}(V)$  is the unique  $\mathbb{N}_0$ -graded braided Hopf algebra generated by  $V$  in  ${}^H_H\mathcal{YD}$  with homogenous components  $\mathcal{B}(V)(0) = \mathbb{k}$ ,  $\mathcal{B}(V)(1) = V$ , and  $P(\mathcal{B}(V)) = V$ , where  $P(\mathcal{B}(V))$  is the space of primitive elements of  $\mathcal{B}(V)$ .

## 2.3 Cartan graphs, root systems, and Weyl groupoids

We start by the notion of a generalized Cartan matrix.

**Definition 2.6.** A generalized Cartan matrix is a matrix  $A = (a_{ij})_{i,j \in I}$  with integer entries such that

- (1)  $a_{ii} = 2$  and  $a_{jk} \leq 0$  for any  $i, j, k \in I$  with  $j \neq k$ ,
- (2) if  $i, j \in I$  and  $a_{ij} = 0$ , then  $a_{ji} = 0$ .

We say that a generalized Cartan matrix  $A \in \mathbb{Z}^{I \times I}$  is *decomposable* if there exists a nonempty proper subset  $I_1 \subset I$  such that  $a_{ij} = 0$  for any  $i \in I_1$  and  $j \in I \setminus I_1$ . And  $A$  is called *indecomposable* if it is not decomposable.

**Definition 2.7.** Let  $\mathcal{X}$  be a non-empty set and  $A^X = (a_{ij}^X)_{i,j \in I}$  be a generalized Cartan matrix for all  $X \in \mathcal{X}$ . For any  $i \in I$  let  $r_i : \mathcal{X} \rightarrow \mathcal{X}$ ,  $X \mapsto r(i, X)$ , where  $r : I \times \mathcal{X} \rightarrow \mathcal{X}$  is a map. The quadruple

$$\mathcal{C} = \mathcal{C}(I, \mathcal{X}, r, (A^X)_{X \in \mathcal{X}})$$

is called a *semi-Cartan graph* if  $r_i^2 = \text{id}_{\mathcal{X}}$  for all  $i \in I$ , and  $a_{ij}^X = a_{ij}^{r_i(X)}$  for all  $X \in \mathcal{X}$  and  $i, j \in I$ . We say that a semi-Cartan graph  $\mathcal{C}$  is *indecomposable* if  $A^X$  is indecomposable for all  $X \in \mathcal{X}$ .

The elements of the set  $\{r_i(X), i \in I\}$  are called the *neighbors* of  $X$  for all  $X \in \mathcal{X}$ . The cardinality of  $I$  is called the *rank* of  $\mathcal{C}$  and the elements of  $\mathcal{X}$  are called the *points* of  $\mathcal{C}$ .

**Definition 2.8.** The exchange graph of  $\mathcal{C}(I, \mathcal{X}, r, (A^X)_{X \in \mathcal{X}})$  is a labeled non-oriented graph with vertices corresponding to the elements of  $\mathcal{X}$ , and edges marked by elements  $i \in I$ , where two vertices  $X, Y$  are connected by an edge  $i$  if and only if  $X \neq Y$  and  $r_i(X) = Y$  (and  $r_i(Y) = X$ ).

Note that the exchange graph of  $\mathcal{C}$  may have multiple edges with different labels. For simplicity, we display only one edge with several labels instead of several edges.

**Definition 2.9.** Let  $\mathcal{C}$  be a semi-Cartan graph.  $\mathcal{C}$  is said to be *connected* if there is no proper non-empty subset  $\mathcal{Y} \subseteq \mathcal{X}$  such that  $r_i(Y) \in \mathcal{Y}$  for all  $i \in I$ ,  $Y \in \mathcal{Y}$ .

Let  $\mathcal{C} = \mathcal{C}(I, \mathcal{X}, r, (A^X)_{X \in \mathcal{X}})$  be a connected semi-Cartan graph. For any  $X \in \mathcal{X}$  and any  $i \in I$  let

$$s_i^X \in \text{Aut}(\mathbb{Z}^I), \quad s_i^X \alpha_j = \alpha_j - a_{ij}^X \alpha_i, \quad (3)$$

for all  $j \in I$ . Let  $\mathcal{D}(\mathcal{X}, I)$  be the category such that  $\text{Ob} \mathcal{D}(\mathcal{X}, I) = \mathcal{X}$  and morphisms  $\text{Hom}(X, Y) = \{(Y, f, X) | f \in \text{End}(\mathbb{Z}^I)\}$  for  $X, Y \in \mathcal{X}$ , where the composition of morphisms is defined by  $(Z, g, Y) \circ (Y, f, X) = (Z, gf, X)$  for all  $X, Y, Z \in \mathcal{X}$ ,  $f, g \in \text{End}(\mathbb{Z}^I)$ . Let  $\mathcal{W}(\mathcal{C})$  be the smallest subcategory of  $\mathcal{D}(\mathcal{X}, I)$  which contains all morphisms  $(r_i(X), s_i^X, X)$  with  $i \in I$ ,  $X \in \mathcal{X}$ . For the sake of simplicity, we write  $s_i^X$  instead of  $(r_i(X), s_i^X, X)$  if no confusion is possible. Notice that  $\mathcal{W}(\mathcal{C})$  is a groupoid since all generators are invertible.

**Definition 2.10.** For all  $X \in \mathcal{X}$ , the set

$$\Delta^{X \text{ re}} = \{\omega \alpha_i \in \mathbb{Z}^I \mid \omega \in \text{Hom}(\mathcal{W}(\mathcal{C}), X)\} \quad (4)$$

is called the set of real roots of  $\mathcal{C}$  at  $X$ . The real roots  $\alpha_i$ ,  $i \in I$ , are called simple. The elements of  $\Delta_+^{X \text{ re}} = \Delta^{X \text{ re}} \cap \mathbb{N}_0^I$  and  $\Delta_-^{X \text{ re}} = \Delta^{X \text{ re}} \cap -\mathbb{N}_0^I$  are called positive roots and negative roots, respectively. We say that  $\mathcal{C}$  is finite if the set  $\Delta^{X \text{ re}}$  is finite for all  $X \in \mathcal{X}$ .

**Definition 2.11.** The semi-Cartan graph  $\mathcal{C}$  is called a Cartan graph if the following hold :

- (1) For all  $X \in \mathcal{X}$ , the set  $\Delta^{X \text{ re}}$  consists of positive and negative roots.
- (2) If  $m_{ij}^X := |\Delta^{X \text{ re}} \cap (\mathbb{N}_0 \alpha_i + \mathbb{N}_0 \alpha_j)|$  is finite, then  $(r_i r_j)^{m_{ij}^X}(X) = X$ , where  $i, j \in I$  and  $X \in \mathcal{X}$ .

In this case,  $\mathcal{W}(\mathcal{C})$  is called the Weyl groupoid of  $\mathcal{C}$ .

**Definition 2.12.**  $\mathcal{R} = \mathcal{R}(\mathcal{C}, (\Delta^X)_{X \in \mathcal{X}})$  is a root system of type  $\mathcal{C}$  if for all  $X \in \mathcal{X}$ , the sets  $\Delta^X$  are the subsets of  $\mathbb{Z}^I$  such that

- (1)  $\Delta^X = (\Delta^X \cap \mathbb{N}_0^I) \cup -(\Delta^X \cap \mathbb{N}_0^I)$ .
- (2)  $\Delta^X \cap \mathbb{Z} \alpha_i = \{\alpha_i, -\alpha_i\}$  for all  $i \in I$ .
- (3)  $s_i^X(\Delta^X) = \Delta^{r_i(X)}$  for all  $i \in I$ .
- (4)  $(r_i r_j)^{m_{ij}^X}(X) = X$  for all  $i \neq j \in I$  where  $m_{ij}^X = |\Delta^X \cap (\mathbb{N}_0 \alpha_i + \mathbb{N}_0 \alpha_j)|$  is finite.

We say that  $\mathcal{R}$  is finite if  $\Delta^X$  is finite for all  $X \in \mathcal{X}$ .

We call  $\mathcal{W}(\mathcal{R}) := \mathcal{W}(\mathcal{C})$  is the groupoid of  $\mathcal{R}$ . As in [16, Definition 4.3] we say that  $\mathcal{R}$  is *reducible* if there exist non-empty disjoint subsets of  $I'$ ,  $I'' \subset I$  such that  $I = I' \cup I''$  and  $a_{ij} = 0$  for all  $i \in I'$ ,  $j \in I''$  and

$$\Delta^X = \left( \Delta^X \cap \sum_{i \in I'} \mathbb{Z} \alpha_i \right) \cup \left( \Delta^X \cap \sum_{j \in I''} \mathbb{Z} \alpha_j \right) \quad \text{for all } X \in \mathcal{X}.$$

In this case, we write  $\mathcal{R} = \mathcal{R}|_{I_1} \oplus \mathcal{R}|_{I_2}$ . Conversely, we say that  $\mathcal{R}$  is *irreducible* if it is not reducible.

Let  $\mathcal{R} = \mathcal{R}(\mathcal{C}, (\Delta^X)_{X \in \mathcal{X}})$  be a root system of type  $\mathcal{C}$ . Next we recall the following properties of  $\mathcal{R}$  from [16] and [17].

**Lemma 2.13.** Let  $X \in \mathcal{X}$ ,  $k \in \mathbb{Z}$ , and  $i, j \in I$  such that  $i \neq j$ . Then  $\alpha_j + k \alpha_i \in \Delta^{X \text{ re}}$  if and only if  $0 \leq k \leq -a_{ij}^X$ .

**Lemma 2.14.** Let  $\mathcal{C}$  be a connected semi-Cartan graph and  $\mathcal{R} = \mathcal{R}(\mathcal{C}, (\Delta^X)_{X \in \mathcal{X}})$  be a root system of type  $\mathcal{C}$ . Then the following are equivalent.

- (1)  $\mathcal{R}$  is finite.
- (2)  $\Delta^X$  is finite for some  $X \in \mathcal{X}$ .
- (3)  $\mathcal{C}$  is finite.

(4)  $\mathcal{W}(\mathcal{R})$  is finite.

**Proposition 2.15.** *Let  $\mathcal{C}$  be a connected semi-Cartan graph and  $\mathcal{R} = \mathcal{R}(\mathcal{C}, (\Delta^X)_{X \in \mathcal{X}})$  be a root system of type  $\mathcal{C}$ . Then*

- (1) *there exists  $X \in \mathcal{X}$  such that  $A^X$  is indecomposable if and only if the semi-Cartan graph  $\mathcal{C}$  is indecomposable.*
- (2) *if  $\mathcal{R}$  is finite then the semi-Cartan graph  $\mathcal{C}$  is indecomposable if and only if the root system  $\mathcal{R}$  is irreducible.*

The following proposition implies that if  $\mathcal{R}$  is a finite root system of type  $\mathcal{C}$  then all roots are real and  $\mathcal{R}$  is uniquely determined by  $\mathcal{C}$ .

**Proposition 2.16.** *Let  $\mathcal{R} = \mathcal{R}(\mathcal{C}, (\Delta^X)_{X \in \mathcal{X}})$  be a root system of type  $\mathcal{C}$ . Let  $X \in \mathcal{X}$ ,  $m \in \mathbb{N}_0$ , and  $i_1, \dots, i_m \in I$  such that*

$$\omega = \text{id}_X s_{i_1} s_{i_2} \cdots s_{i_m} \in \text{Hom}(\mathcal{W}(\mathcal{C}), X)$$

*and  $\ell(\omega) = m$ . Then the elements*

$$\beta_n = \text{id}_X s_{i_1} s_{i_2} \cdots s_{i_{n-1}}(\alpha_{i_n}) \in \Delta^X \cap \mathbb{N}_0^I,$$

*are pairwise different, where  $n \in \{1, 2, \dots, m\}$  (and  $\beta_1 = \alpha_{i_1}$ ). Here,*

$$\ell(\omega) = \min\{m \in \mathbb{N}_0 \mid \omega = \text{id}_X s_{i_1} s_{i_2} \cdots s_{i_m}, i_1, i_2, \dots, i_m \in I\}$$

*is the length of  $\omega \in \text{Hom}(\mathcal{W}(\mathcal{C}), X)$ . In particular, if  $\mathcal{R}$  is finite and  $\omega \in \text{Hom}(\mathcal{W}(\mathcal{C}))$  is the longest element, then*

$$\{\beta_n \mid 1 \leq n \leq \ell(\omega) = |\Delta^X|/2\} = \Delta^X \cap \mathbb{N}_0^I.$$

**Remark 2.17.** *If  $\mathcal{C}$  is a finite Cartan graph then  $\mathcal{R}$  is finite and hence*

$$\mathcal{R}^{re} = \mathcal{R}^{re}(\mathcal{C}, (\Delta^{X^{re}})_{X \in \mathcal{X}})$$

*is the unique root system of type  $\mathcal{C}$  by Proposition 2.16, that is,  $\mathcal{R}$  is uniquely determined by  $\mathcal{C}$ . And we say that  $(\mathcal{R}^{re})_+^X$  is the set of positive real roots at  $X \in \mathcal{X}$ .*

## 2.4 The semi-Cartan Graph of a Nichols Algebra of Diagonal Type

In this section, we attach a semi-Cartan graph to a tuple of finite dimensional Yetter–Drinfel’d modules under some finiteness conditions. Assume that  $G$  is an abelian group and  $\mathbb{k}G$  is the group algebra over  $G$ . Let  ${}^G_G\mathcal{YD}$  be the category of Yetter–Drinfel’d modules over  $\mathbb{k}G$ . Let  $\mathcal{F}_r^G$  be the set of  $r$ -tuples of finite dimensional irreducible objects in  ${}^G_G\mathcal{YD}$  and  $\mathcal{X}_r^G$  be the set of  $r$ -tuples of isomorphism classes of finite dimensional irreducible objects in  ${}^G_G\mathcal{YD}$ . For any  $(M_1, \dots, M_r) \in \mathcal{F}_r^G$ , write  $[M] := ([M_1], \dots, [M_r]) \in \mathcal{X}_r^G$  the corresponding isomorphism class of  $(M_1, \dots, M_r)$ .



We fix that  $M = (\mathbb{k}x_1, \mathbb{k}x_2, \dots, \mathbb{k}x_r) \in \mathcal{F}_r^G$  is a tuple of one-dimensional Yetter–Drinfel’d over  $G$  and  $[M] \in \mathcal{X}_r^G$ . Let  $V_M = \bigoplus_{i \in I} \mathbb{k}x_i \in {}_G^G \mathcal{YD}$  be a Yetter–Drinfel’d module of diagonal type over  $G$ , where  $\{x_i | i \in I\}$  is a basis of  $V$ . Then there exist a matrix  $(q_{ij})_{i,j \in I}$  and elements  $\{g_i | i \in I\} \subset G$  such that  $\delta(x_i) = g_i \otimes x_i$  and  $g_i \cdot x_j = q_{ij}x_j$  for all  $i, j \in I$ . We say that the matrix  $(q_{ij})_{i,j \in I}$  is the braiding matrix of  $M$ . Recall that the matrix does not depend on the basis  $\{x_i | i \in I\}$  up to permutation of  $I$ . We say that  $\mathcal{B}(V_M) = \mathcal{B}(\bigoplus_{i=1}^n \mathbb{k}x_i)$  is the Nichols algebra of the tuple  $M$ , denoted by  $\mathcal{B}(M)$ .

Recall that the adjoint action  $\text{ad}$  of a Nichols algebra  $\mathcal{B}(M)$  with antipode  $S$  from [30] is the linear map  $\text{ad} : \mathcal{B}(M) \otimes \mathcal{B}(M) \rightarrow \mathcal{B}(M)$

$$x \otimes y \rightarrow (\text{ad}x)(y) := x_{(1)}yS(x_{(2)})$$

for all  $x, y \in \mathcal{B}(M)$ .

In order to construct a semi-Cartan graph to  $M$ , we recall some finiteness conditions from [9] and [28].

**Definition 2.18.** *Let  $i \in I$ .  $M$  is  $i$ -finite if for any  $j \in I \setminus \{i\}$ ,  $(\text{ad}_c x_i)^m(x_j) = 0$  for some  $m \in \mathbb{N}$ .*

**Lemma 2.19.** *For any  $i, j \in I$  with  $i \neq j$ , the following are equivalent.*

- (1)  $(m+1)_{q_{ii}}(q_{ii}^m q_{ij} q_{ji} - 1) = 0$  and  $(k+1)_{q_{ii}}(q_{ii}^k q_{ij} q_{ji} - 1) \neq 0$  for all  $0 \leq k < m$ .
- (2)  $(\text{ad}_c x_i)^{m+1}(x_j) = 0$  and  $(\text{ad}_c x_i)^m(x_j) \neq 0$  in  $\mathcal{B}(V)$ .

Here  $(n)_q := 1 + q + \dots + q^{n-1}$ , which is 0 if and only if  $q^n = 1$  for  $q \neq 1$  or  $p|n$  for  $q = 1$ . Notice that  $(1)_q \neq 0$  for any  $q \in \mathbb{N}$ .

The lemma below follows directly from Lemma 2.19.

**Lemma 2.20.** *Let  $i \in I$ . Then  $M = (\mathbb{k}x_j)_{j \in I}$  is  $i$ -finite if and only if for any  $j \in I \setminus \{i\}$  there is a non-negative integer  $m$  satisfying  $(m+1)_{q_{ii}}(q_{ii}^m q_{ij} q_{ji} - 1) = 0$ .*

Let  $i \in I$ . Assume that  $M$  is  $i$ -finite. Let  $(a_{ij}^M)_{j \in I} \in \mathbb{Z}^I$  and  $R_i(M) = (R_i(M)_j)_{j \in I}$ , where

$$a_{ij}^M = \begin{cases} 2 & \text{if } j = i, \\ -\max\{m \in \mathbb{N}_0 \mid (\text{ad}_c x_i)^m(x_j) \neq 0\} & \text{if } j \neq i. \end{cases}$$

$$R_i(M)_i = \mathbb{k}y_i, \quad R_i(M)_j = \mathbb{k}(\text{ad}_c x_i)^{-a_{ij}^M}(x_j), \quad (5)$$

where  $y_i \in (\mathbb{k}x_i)^* \setminus \{0\}$ . If  $M$  is not  $i$ -finite, then let  $R_i(M) = M$ . Then  $R_i(M)$  is a  $r$ -tuple of one-dimensional Yetter–Drinfel’d modules over  $G$ .

Let

$$\mathcal{F}_r^G(M) = \{R_{i_1} \cdots R_{i_n}(M) \in \mathcal{F}_r^G \mid n \in \mathbb{N}_0, i_1, \dots, i_n \in I\}$$

and

$$\mathcal{X}_r^G(M) = \{[R_{i_1} \cdots R_{i_n}(M)] \in \mathcal{X}_r^G \mid n \in \mathbb{N}_0, i_1, \dots, i_n \in I\}.$$

**Definition 2.21.** *We say that  $M$  admits all reflections if  $N$  is  $i$ -finite for all  $N \in \mathcal{F}_r^G(M)$ .*

Notice that the reflections depend only on the braiding matrix  $(q_{ij})_{i,j \in I}$ . Next we recall the notion of generalized Dynkin diagram [25].

**Definition 2.22.** Let  $V$  be a  $r$ -dimensional braided vector space of diagonal type with the braiding matrix  $(q_{ij})_{i,j \in I}$ . The generalized Dynkin diagram of  $V$  is a non-directed graph  $\mathcal{D}$  with the following properties:

- (1) there is a bijective map  $\phi$  from  $I$  to the vertices of  $\mathcal{D}$ ,
- (2) for all  $i \in I$  the vertex  $\phi(i)$  is labeled by  $q_{ii}$ ,
- (3) for all  $i, j \in I$  with  $i \neq j$ , the number  $n_{ij}$  of edges between  $\phi(i)$  and  $\phi(j)$  is either 0 or 1. If  $q_{ij}q_{ji} = 1$  then  $n_{ij} = 0$ , otherwise  $n_{ij} = 1$  and the edge is labeled by  $q_{ij}q_{ji}$ .

We say that the generalized Dynkin diagram of  $M$  is the generalized Dynkin diagram of the braided vector space  $\oplus_{i \in I} M_i$ . The generalized Dynkin diagram of  $M$  is connected if the braiding matrix of  $M$  is indecomposable.

To compute the labels of the generalized Dynkin diagram of  $R_i(M) = (R_i(M)_j)_{j \in I}$ , we present the following theory.

**Lemma 2.23.** Let  $i \in I$ . Assume that  $M$  is  $i$ -finite and let  $a_{ij} := a_{ij}^M$  for all  $j \in I$ . Let  $(q'_{jk})_{j,k \in I}$  be the braiding matrix of  $R_i(M)$  with respect to  $(y_j)_{j \in I}$ . Then

$$q'_{jj} = \begin{cases} q_{ii} & \text{if } j = i, \\ q_{jj} & \text{if } j \neq i, q_{ij}q_{ji} = q_{ii}^{a_{ij}}, \\ q_{ii}q_{jj}(q_{ij}q_{ji})^{-a_{ij}} & \text{if } j \neq i, q_{ii} \in G'_{1-a_{ij}}, \\ q_{jj}(q_{ij}q_{ji})^{-a_{ij}} & \text{if } j \neq i, q_{ii} = 1, \end{cases}$$

$$q'_{ij}q'_{ji} = \begin{cases} q_{ij}q_{ji} & \text{if } j \neq i, q_{ij}q_{ji} = q_{ii}^{a_{ij}}, \\ q_{ii}^2(q_{ij}q_{ji})^{-1} & \text{if } j \neq i, q_{ii} \in G'_{1-a_{ij}}, \\ (q_{ij}q_{ji})^{-1} & \text{if } j \neq i, q_{ii} = 1, \end{cases}$$

and

$$q'_{jk}q'_{kj} = \begin{cases} q_{jk}q_{kj} & \text{if } q_{ir}q_{ri} = q_{ii}^{a_{ir}}, r \in \{j, k\}, \\ q_{jk}q_{kj}(q_{ik}q_{ki}q_{ii}^{-1})^{-a_{ij}} & \text{if } q_{ij}q_{ji} = q_{ii}^{a_{ij}}, q_{ii} \in G'_{1-a_{ik}}, \\ q_{jk}q_{kj}(q_{ij}q_{ji})^{-a_{ik}}(q_{ik}q_{ki})^{-a_{ij}} & \text{if } q_{ii} = 1, \\ q_{jk}q_{kj}q_{ii}^2(q_{ij}q_{ji}q_{ik}q_{ki})^{-a_{ij}} & \text{if } q_{ii} \in G'_{1-a_{ik}}, q_{ii} \in G'_{1-a_{ij}}. \end{cases}$$

for  $j \neq k$ ,  $|j - i| = 1$ ,  $|k - i| = 1$ . For  $|j - i| > 1$ ,  $|k - i| > 1$ ,  $q'_{jk}q'_{kj} = q_{jk}q_{kj}$ . Here,  $G'_n$  denotes the set of primitive  $n$ -th roots of unity in  $\mathbb{K}$ , that is,  $G'_n = \{q \in \mathbb{K}^* \mid q^n = 1, q^k \neq 1 \text{ for all } 1 \leq k < n\}$  for  $n \in \mathbb{N}$ .

If  $M$  admits all reflections, then we are able to construct a semi-Cartan graph  $\mathcal{C}(M)$  of  $M$  by [44, Proposition 1.5].

**Theorem 2.24.** Assume that  $M$  admits all reflections. For all  $X \in \mathcal{X}_r^G(M)$  let

$$[X]_r = \{Y \in \mathcal{X}_r^G(M) \mid Y \text{ and } X \text{ have the same generalized Dynkin diagram}\}.$$

Let  $\mathcal{Y}_r(M) = \{[X]_r \mid X \in \mathcal{X}_r^G(M)\}$  and  $A^{[X]_r} = A^X$  for all  $X \in \mathcal{X}_r^G(M)$ . Let  $t : I \times \mathcal{Y}_r(M) \rightarrow \mathcal{Y}_r(M)$ ,  $(i, [X]_r) \mapsto [R_i(X)]_r$ . Then the tuple

$$\mathcal{C}(M) = \{I, \mathcal{Y}_r(M), t, (A^Y)_{Y \in \mathcal{Y}_r(M)}\}$$

is a connected semi-Cartan graph. We say that  $\mathcal{C}(M)$  is the semi-Cartan graph attached to  $M$ .

Furthermore, we can attach a groupoid  $\mathcal{W}(M) := \mathcal{W}(\mathcal{C}(M))$  to  $M$  if  $M$  admits all reflections.

Notice that Nichols algebra  $\mathcal{B}(M)$  is  $\mathbb{N}_0^r$ -graded with  $\deg M_i = \alpha_i$  for all  $i \in I$ . Following the terminology in [28], we say that the Nichols algebra  $\mathcal{B}(M)$  is *decomposable* if there exists a totally ordered index set  $(L, \leq)$  and a sequence  $(W_l)_{l \in L}$  of finite dimensional irreducible  $\mathbb{N}_0^r$ -graded objects in  ${}^G\mathcal{YD}$  such that

$$\mathcal{B}(M) \simeq \bigotimes_{l \in L} \mathcal{B}(W_l). \quad (6)$$

For each decomposition (6), we define the set of *positive roots*  $\Delta_+^{[M]} \subset \mathbb{Z}^I$  and the set of *roots*  $\Delta^{[M]} \subset \mathbb{Z}^I$  of  $[M]$  by

$$\Delta_+^{[M]} = \{\deg(W_l) \mid l \in L\}, \quad \Delta^{[M]} = \Delta_+^{[M]} \cup -\Delta_+^{[M]}.$$

By [28, Theorem 4.5] we obtain that the set of roots  $\Delta^{[M]}$  of  $[M]$  does not depend on the choice of the decomposition.

**Remark 2.25.** If  $\dim M_i = 1$  for all  $i \in I$ , then the Nichols algebra  $\mathcal{B}(M)$  is decomposable based on the theorem of V. Kharchenko [34, Theorem 2]. The set of roots of Nichols algebra  $\mathcal{B}(M)$  can be always defined and it is denoted by  $\Delta^{[M]}$ . If the set of roots  $\Delta^{[M]}$  is finite then we can check that  $M$  admits all reflections by [28, Corollary 6.12].

If  $M$  admits all reflections and  $\Delta^{[M]}$  is finite, then we can define a finite root system  $\mathcal{R}(M)(\mathcal{C}(M), (\Delta^{[N]})_{N \in \mathcal{F}_r^G(M)})$  of type  $\mathcal{C}(M)$  [45, Theorem 1.25].

**Theorem 2.26.** Assume that  $M$  admits all reflections. Then the following are equivalent.

- (1)  $\Delta^{[M]}$  is finite.
- (2)  $\mathcal{C}(M)$  is a finite Cartan graph.
- (3)  $\mathcal{W}(M)$  is finite.
- (4)  $\mathcal{R}(M) := \mathcal{R}(M)(\mathcal{C}(M), (\Delta^{[N]})_{N \in \mathcal{F}_r^G(M)})$  is finite.

In all cases,  $\mathcal{R}(M)$  is the unique root system of type  $\mathcal{C}(M)$ .

To simplify the presentation of the generalized Dynkin diagrams, we recall the notion of simple chains [26]. Recall that we denote  $E = (\alpha_i)_{i \in I}$  the standard basis of  $\mathbb{Z}^I$ .

**Definition 2.27.** Assume that  $(\Delta, \chi, E)$  is an arithmetic root system of rank  $r > 2$  and  $D_{\chi, E}$  is a labeled path graph. Call this graph a simple chain (of length  $r$ ) if

$$(q_{11}q_{12}q_{21} - 1)(q_{11} + 1) = 0, \quad (q_{rr}q_{r,r-1}q_{r-1,r} - 1)(q_{rr} + 1) = 0$$

$$q_{ii}^2 q_{i-1,i} q_{i,i-1} q_{i,i+1} q_{i+1,i} = 1 \quad \text{for } 1 < i < r$$

The latter equations hold if and only if

$$q_{ii} + 1 = q_{i-1,i} q_{i,i-1} q_{i,i+1} q_{i+1,i} - 1 = 0 \quad \text{or} \quad q_{ii} q_{i-1,i} q_{i,i-1} = q_{ii} q_{i,i+1} q_{i+1,i} = 1.$$

Let  $\mathcal{D}_{\chi, E}$  be a simple chain and set  $q := q_{rr}^2 q_{r-1,r} q_{r,r-1}$ . By Definition 2.27, we obtain

$$q = \begin{cases} q_{rr} & \text{if } q_{rr} \neq -1, \\ q_{r-1,r} q_{r,r-1} & \text{if } q_{rr} = -1. \end{cases}$$

Then  $q_{11}^2 q_{12} q_{21} \in \{q, q^{-1}\}$  and  $q_{i-1,i} q_{i,i-1} \in \{q, q^{-1}\}$  for all  $i \in \{2, \dots, r\}$ . Moreover, we observe that the simple chain  $\mathcal{D}_{\chi, E}$  is uniquely determined by the value of  $q$  and the indices  $i$  ( $1 \leq i \leq r$ ) satisfying  $q_{i-1,i} q_{i,i-1} = q$ , with  $q_{01} q_{10}$  defined as  $1/(q_{11}^2 q_{12} q_{21})$ . In this case, we use the symbol

$$\boxed{C(r, q; i_1, \dots, i_j)}$$

to denote  $\mathcal{D}_{\chi, E}$ , where  $q_{i-1,i} q_{i,i-1} = q$  if and only if  $i \in \{i_1, i_2, \dots, i_j\}$ .

### 3 Finite Cartan graphs

In this section, we illustrate the properties of finite connected indecomposable Cartan graphs of rank  $r \geq 5$  in Theorem 3.11 and Theorem 3.15, which are essential for our classification in the next section.

Let  $M := (\mathbb{k}x_1, \dots, \mathbb{k}x_r)$  be a tuple of one-dimensional Yetter-Drinfel'd modules over  $\mathbb{k}G$ . Assume that  $M$  admits all reflections and  $\mathcal{C}(M) = \mathcal{C}(I, \mathcal{X}, r, (A^X)_{X \in \mathcal{X}})$  be the attached indecomposable semi-Cartan graph to  $M$ . Recall that a semi-Cartan graph  $\mathcal{C}$  is standard, if  $A^X = A^Y$  for all  $X, Y \in \mathcal{X}$ .

$$\text{Set } A_r = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \cdots & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & \cdots & 0 & -1 & 2 \end{pmatrix},$$

$$B_r = \begin{pmatrix} 2 & -2 & 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \cdots & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & \cdots & 0 & -1 & 2 \end{pmatrix},$$

$$\begin{aligned}
C_r = B_r^T &= \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ -2 & 2 & -1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \cdots & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & \cdots & 0 & -1 & 2 \end{pmatrix}, \\
D_r &= \begin{pmatrix} 2 & 0 & -1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 2 & -1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ -1 & -1 & 2 & -1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \cdots & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & \cdots & 0 & -1 & 2 \end{pmatrix}, \\
\text{and } E_r &= \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ -1 & 2 & 0 & -1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 2 & -1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & -1 & -1 & 2 & -1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & 0 & \cdots & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & \cdots & 0 & -1 & 2 \end{pmatrix}.
\end{aligned}$$

### 3.1 Non-sporadic finite Cartan graphs

We give the properties of non-sporadic finite Cartan graphs of rank  $r \geq 5$  in Theorem 3.11, which are crucial for our classification in Theorem 4.5. In this section, we first recall the arithmetic root systems of non-sporadic finite Cartan graphs of rank  $r \geq 5$ . Based on the analysis of these root systems, we then define the good  $D'_r$  neighborhood to cover all such Cartan graphs.

#### 3.1.1 The root systems

We focus mainly on the construction of root systems of type  $D'(r, s)$  for  $s \in \{1, \dots, r-1\}$  from [18]. For details on the construction of root systems of standard type  $A_r, B_r, C_r, D_r$ , we refer to the book [33, Chapter III, § 12.1]—the classic results on root systems of Lie algebras by J.E. Humphreys provide a benchmark for the construction of standard root systems.

For  $1 \leq i, j \leq r$ , let

$$\eta_{i,j} := \begin{cases} \sum_{k=i}^j \alpha_k & i \leq j, \\ 0 & i > j. \end{cases}$$

**Definition 3.1.** Let  $Z \subseteq \{1, \dots, r-1\}$ . Let  $\Phi_{r,Z}$  denote the set of roots

$$\begin{aligned} \eta_{i,j-1}, & \quad 1 \leq i < j \leq r, \\ \eta_{i,r-2} + \alpha_r, & \quad 1 \leq i < r, \\ \eta_{i,r} + \eta_{j,r-2}, & \quad 1 \leq i < j < r, \\ \eta_{j,r} + \eta_{j,r-2}, & \quad j \in Z. \end{aligned}$$

Let  $Y \subseteq \{1, \dots, r-1\}$ . Let  $\Psi_{r,Y}$  denote the set of roots

$$\begin{aligned} \eta_{i,j}, & \quad 1 \leq i \leq j \leq r, \\ \eta_{i,r} + \eta_{j,r-1}, & \quad 1 \leq i < j < r, \\ \eta_{j,r} + \eta_{j,r-1}, & \quad j \in Y. \end{aligned}$$

Further, denote  $\Psi'_{r,Y}$  the set obtained from  $\Psi_{r,Y}$  by exchanging  $\alpha_{r-1}$  and  $\alpha_r$ .

**Remark 3.2.** The sets  $\Phi_{r,\emptyset}$  and  $\Psi_{r,\{1,\dots,r-1\}}$  are the sets of positive real roots of the Weyl groups of type  $D_r$  and  $C_r$ , respectively, as constructed for the root systems in [33].

**Proposition 3.3.** Let  $Z_1, Z_2, Y_1, Y_2 \subseteq \{1, \dots, r-1\}$  with

$$|Z_1| = |Z_2| = |Y_1| + 1 = |Y_2| + 1.$$

Then there exists a finite Cartan graph  $\mathcal{C}$  with objects  $a, b, c, d$  such that

$$(R^{\text{re}})_+^a = \Phi_{r,Z_1}, \quad (R^{\text{re}})_+^b = \Phi_{r,Z_2}, \quad (R^{\text{re}})_+^c = \Psi_{r,Y_1} \quad \text{and} \quad (R^{\text{re}})_+^d = \Psi'_{r,Y_2}.$$

**Remark 3.4.** Let  $Z, Y \subseteq \{1, \dots, r-1\}$  with  $|Z| = |Y| + 1$ . Then  $\Phi_{r,Z}$ ,  $\Psi_{r,Y}$  and  $\Psi'_{r,Y}$  are isomorphic by Proposition 3.3.

**Definition 3.5.** Let  $\mathcal{C}$  be a finite Cartan graph of rank  $r \geq 3$ . If there exists a subset  $Z \subseteq \{1, \dots, r-1\}$  such that  $\Phi_{r,Z} = (R^{\text{re}})_+^a$  for some object  $a$ , then we say that  $\mathcal{C}$  is of type  $D'(r, |Z|)$ . If there exists a subset  $Y \subseteq \{1, \dots, r-1\}$  such that  $\Psi_{r,Y} = (R^{\text{re}})_+^a$  for some object  $a$ , then we say that  $\mathcal{C}$  is of type  $D'(r, |Y| + 1)$ .

Notice that if  $\mathcal{C}$  is of type  $D'(r, 0)$ , then it is standard of type  $D$ , and if  $\mathcal{C}$  is of type  $D'(r, r)$ , then it is standard of type  $C$ .

**Proposition 3.6.** Let  $\mathcal{C}$  be a non-sporadic finite Cartan graph of rank  $r \geq 3$ . Let  $a$  be an object of  $\mathcal{C}$  and assume that the Dynkin diagram of the object  $a$  is of type  $D'_r$ . Then  $\Phi_{r,\{r-1\}} \subseteq (R^{\text{re}})_+^a$ .

**Remark 3.7.** Let  $\mathcal{C}$  be a non-sporadic finite Cartan graph of rank  $r \geq 3$ . Then either  $\mathcal{C}$  is a standard Cartan graph of type  $A_r, B_r, C_r, D_r$ , or there is an object of Dynkin diagram type  $D'_r$  [18, Proposition 3.8]

**Theorem 3.8.** Let  $\mathcal{C}$  be a non-sporadic finite Cartan graph of rank  $r \geq 3$  and let

$$\mathcal{R}_+ := \{(R^{\text{re}})_+^a \mid a \in \mathcal{C}\}.$$

Then there are two possibilities:

- (1) The Cartan graph  $\mathcal{C}$  is standard ( $|\mathcal{R}_+| = 1$ ) of type  $A_r, B_r, C_r, D_r$ .
- (2) Up to equivalence, the root sets of  $\mathcal{C}$  are given by

$$\mathcal{R}_+ = \{\Phi_{r,Z}, \Psi_{r,Y}, \Psi'_{r,Y} \mid Z, Y \subseteq \{1, \dots, r-1\}, |Z| = s, |Y| = s-1\}$$

for some  $s \in \{1, \dots, r-1\}$ .

**Remark 3.9.** Let  $\mathcal{C}$  be a finite Cartan graph of rank  $r \geq 3$ . We can construct all root systems of  $\mathcal{C}$  by Theorem 3.8. Especially, we give a list of root systems of rank  $3 \leq r \leq 9$  that comprises the positive real roots of one object for each type of non-sporadic finite Cartan graph in Appendix 6. The rank 3 ones are taken from [17] and there are only five finite Cartan graphs are non-sporadic because of  $A_3 = D_3$ .

### 3.1.2 Good $D'_r$ neighborhood

In general, each point of a finite Cartan graph  $\mathcal{C}$  has its own neighborhood. We analyze all the positive real roots  $\Delta^{[M]}$  of non-sporadic finite Cartan graph by GAP. Depending on the output, we construct the following definition “good  $D'_r$  neighborhood” in order to cover all the finite connected indecomposable Cartan graphs in such way that at least one point of  $\mathcal{C}$  has a good  $D'_r$  neighborhood.

**Definition 3.10.** We say that  $X$  has a **good  $D'_r$  neighborhood** if there exists a permutation of  $I$  and an integer  $a \in \mathbb{N}$  such that  $A^X = A^{r_i(X)}$ ,  $i \in \{1, 4, \dots, r\}$ ,

$$A^{r_2(X)} = \begin{pmatrix} 2 & -1 & -1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ -1 & -1 & 2 & -1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \cdots & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & \cdots & 0 & -1 & 2 \end{pmatrix},$$

$$A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ -a & 2 & -1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \cdots & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & \cdots & 0 & -1 & 2 \end{pmatrix},$$

where  $a \in \{1, 2\}$  and  $a_{14}^{r_3 r_2(X)} = 0$ .

The following theorem illustrates an important property of the non-sporadic finite Cartan graphs.

**Theorem 3.11.** Assume that  $\mathcal{C}$  is a non-sporadic finite Cartan graph of rank  $r \geq 5$ . Then one of the following is true.

- (1) The Cartan graph  $\mathcal{C}(M)$  is standard of type  $A_r, B_r, C_r, D_r$ .

(2) Up to equivalence, there exists a point  $Y \in \mathcal{X}$  such that  $Y$  has a good  $D'_r$  neighborhood.

*Proof.* Let  $\mathcal{R} = \mathcal{R}(\mathcal{C}, (\Delta^X)_{X \in \mathcal{X}})$  be a root system of  $\mathcal{C}(M)$ . For any  $X \in \mathcal{X}$ , let  $\Delta_+^{X \text{ re}}$  be the positive roots of  $X$ . To a permutation of  $I$ , by Theorem 3.8 there exists a point  $X \in \mathcal{X}$  satisfying that the set  $\Delta_+^{X \text{ re}}$  is the type  $A_r, B_r, C_r, D_r$ , or type  $D'(r, s)$  for  $s \in \{1, \dots, r-1\}$ . There are precisely  $r-1+4$  such possible sets of real roots. We analyze all  $r+3$  sets in the following way. For point  $Y$ , we assume that  $\Delta_+^{Y \text{ re}}$  is one of the sets. Since the reflection  $s_i^Y$  maps  $\Delta_+^{Y \text{ re}} \setminus \{\alpha_i\}$  bijectively to  $\Delta_+^{r_i(Y) \text{ re}} \setminus \{\alpha_i\}$  for any  $i \in I$ , the Cartan matrices of all neighbors of  $Y$  can be obtained from  $\Delta_+^{Y \text{ re}}$  by Lemma 2.13. If the Cartan graph  $\mathcal{C}(M)$  is standard or  $Y$  has a good  $D'_r$  neighborhood, then the claim is true. Otherwise repeat the previous step to the neighbours of  $Y$ . Since  $\mathcal{X}$  is finite, this algorithm terminates. The elementary calculations by GAP and are skipped here.  $\square$

## 3.2 Sporadic finite Cartan graphs

In this section, we formulate the properties of sporadic finite Cartan graphs of rank  $r \geq 5$  in Theorem 3.15.

We analyze the positive real roots of sporadic finite Cartan graphs by the same way in Section 3.1.2. All such roots are listed in [18, Appendix B.2.]. Here, we define “good  $A_5$  neighborhood”, “good  $A_6$  neighborhood”, and “good  $A_7$  neighborhood” for ranks 5, 6, and 7, respectively.

**Definition 3.12.** We say that  $X$  has a **good  $A_5$  neighborhood** if there exists a permutation of  $I$  and integers  $a, b, c, d, e \in \mathbb{N}$  such that

$$A^X = A^{r_1(X)} = A^{r_2(X)} = A_5, \quad A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & -a & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & -b & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix},$$

$$A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -c \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -d & -1 & 2 \end{pmatrix}, \quad A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & -e & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix},$$

$$a_{15}^{r_2 r_3 r_4(X)} = 0, \quad a_{45}^{r_3 r_2 r_1(X)} \neq -3, \quad \text{and satisfies one of the following.}$$

$$(A_{5_1}) \quad (a, b, c, d, e) = (1, 1, 0, 0, 1), \quad (a_{41}^{r_2 r_3(X)}, a_{45}^{r_2 r_3(X)}) = (0, -1), \quad (a_{32}^{r_4 r_5(X)}, a_{34}^{r_2 r_1(X)}) \neq (-2, -2);$$

$$(A_{5_2}) \quad (a, b, c, d, e) = (0, 0, 1, 1, 2), \quad a_{25}^{r_3 r_4(X)} \in \{0, -1\}, \quad (a_{45}^{r_3 r_2 r_1(X)}, a_{45}^{r_3 r_2(X)}) \neq (-2, -2);$$

$$(A_{5_3}) \quad (a, b, c, d, e) = (0, 0, 1, 1, 1), \quad (a_{52}^{r_3 r_4(X)}, a_{32}^{r_5 r_4(X)}) = (-1, -1).$$

**Definition 3.13.** We say that  $X$  has a **good  $A_6$  neighborhood** if there exists a permutation of  $I$  and an integer  $a \in \mathbb{N}$  such that  $A^X = A^{r_1(X)} = A^{r_2(X)} = A^{r_3(X)} = A^{r_6(X)} = A_6$ ,



$$A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 & 0 \\ 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & -1 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}, A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & -a \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & -a & -1 & 2 \end{pmatrix},$$

$a_{36}^{r_5 r_4(X)} = 0$ , and  $a_{25}^{r_3 r_4(X)} = 0$ , where  $a$  satisfies one of the following.

( $A_{6_1}$ )  $a = 0$ ,  $a_{32}^{r_5 r_4(X)} = -1$ ,  $a_{43}^{r_5 r_6(X)} = -1$ ,  $(a_{45}^{r_3 r_2 r_1(X)}, a_{45}^{r_3 r_2(X)}) \neq (-2, -2)$ ;

( $A_{6_2}$ )  $a = 1$ ,  $a_{45}^{r_3 r_2(x)} = -1$ ,  $a_{45}^{r_3 r_2 r_1(X)} = -1$ .

**Definition 3.14.** We say that  $X$  has a **good  $A_7$  neighborhood** if there exists a permutation of  $I$  such that  $A^X = A^{r_1(X)} = A^{r_2(X)} = A^{r_4(X)} = A^{r_5(X)} = A^{r_6(X)} = A^{r_7(X)} = A_7$ ,

$$A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}, (a_{25}^{r_4 r_3(X)}, a_{21}^{r_4 r_3(X)}) = (0, -1),$$

$a_{45}^{r_2 r_3(x)} = -1$ ,  $a_{34}^{r_2 r_1(X)} = -1$ ,  $a_{32}^{r_4 r_5(X)} = -1$ ,  $a_{32}^{r_4 r_5 r_6(x)} = -1$ , and  $a_{32}^{r_4 r_5 r_6 r_7(X)} = -1$ .

We formulate the following property of the sporadic finite Cartan graphs.

**Theorem 3.15.** Assume that  $\mathcal{C}$  is a sporadic finite Cartan graph of rank  $r \geq 5$ . Then one of the following is true.

- (1) The Cartan graph  $\mathcal{C}(M)$  is standard of type  $E_6, E_7, E_8$ .
- (2) Up to equivalence, there exists a point  $Y \in \mathcal{X}$  such that  $Y$  has a good  $A_\theta$  neighborhood, where  $\theta = 5, 6, 7$ .

*Proof.* For any  $X \in \mathcal{X}$ , let  $\Delta_+^{X \text{ re}}$  be the positive real roots of  $X$ . By [18, Theorem 4.1] there exists a point  $X \in \mathcal{X}$  satisfying that the set  $\Delta_+^{X \text{ re}}$  is in the list of [18, Appendix B.2.] up to a permutation of  $I$ . There are precisely 6 sets, 4 sets, 2 sets and 1 set of real roots for rank 5, 6, 7, 8, respectively. We analyze each set of the real roots in the list. Since the reflection  $s_i^X$  maps  $\Delta_+^{X \text{ re}} \setminus \{\alpha_i\}$  bijectively to  $\Delta_+^{r_i(X) \text{ re}} \setminus \{\alpha_i\}$  for any  $i \in I$ , the Cartan matrices of all neighbors of  $X$  can be obtained from  $\Delta_+^{X \text{ re}}$  by Lemma 2.13. If the Cartan graph  $\mathcal{C}(M)$  is standard or  $Y$  has a good  $A_\theta$  neighborhood, then the claim is true. Otherwise repeat the previous step to the neighbours of  $Y$ . Since  $\mathcal{X}$  is finite, this algorithm terminates.  $\square$

## 4 The classification theorem

In this section, all rank  $r \geq 5$  Nichols algebras of diagonal type with a finite irreducible root system are determined. We formulate the main result in Theorem 4.1 and present the corresponding generalized Dynkin diagrams in Table 1.

**Theorem 4.1.** Let  $\mathbb{k}$  is a field of characteristic  $p > 0$ . Let  $r \geq 5$  and  $I = \{1, 2, \dots, r\}$ . Let  $(V, c)$  be a braided vector space of diagonal type over  $\mathbb{k}$  with basis  $\{x_k | k \in I\}$  satisfying

$$c(x_i \otimes x_j) = q_{ij} x_j \otimes x_i, \quad q_{ij} \in \mathbb{k}^*.$$

Assume that the  $(q_{ij})_{i,j \in I}$  is an indecomposable braiding matrix. Let  $M := (\mathbb{k}x_i)_{i \in I}$ . Then the Nichols algebra  $\mathcal{B}(V)$  generated by  $(V, c)$  has a finite set of roots  $\Delta^{[M]}$  if and only if the generalized Dynkin diagram  $\mathcal{D}$  of  $V$  appears in Table 1. In this case, the row of Table 1 containing  $\mathcal{D}$  consists precisely of the generalized Dynkin diagrams of all the points of  $\mathcal{C}(M)$ . In Table 2 we present the corresponding exchange graphs for the sporadic case of  $\mathcal{C}(M)$ .

Theorem 4.1 is enough to classify finite dimensional Nichols algebra of diagonal type by [31, Corollary 6].

**Corollary 4.2.** Assume that  $(V, c)$  be a braided vector space of diagonal type. Then the Nichols algebra  $\mathcal{B}(V)$  is finite dimensional if and only if the generalized Dynkin diagram  $\mathcal{D}$  of  $V$  appears in Table 1 and the labels of the vertices of  $\mathcal{D}$  are roots of unity (including 1).

**Remark 4.3.** To avoid confusion, we give the following statements.




Figure 1: Dynkin diagram of type  $D'_r$




Figure 2: generalized Dynkin diagram of type  $A_r$

- (1) Label the vertices of the generalized Dynkin diagrams from left to right and then top to bottom by  $1, \dots, r$  (for example: label  $D_r$  as in Fig. 1).
- (2) For a generalized Dynkin diagram  $\mathcal{D}$  and  $i_1, i_2, \dots, i_r \in I$ , write  $\tau_{i_1 i_2 \dots i_r} \mathcal{D}$  for the graph  $\mathcal{D}$  where the vertices of  $\mathcal{D}$  change to  $i_1, i_2, \dots, i_r$  respectively.
- (3) For a generalized Dynkin diagram  $\mathcal{D}$  and  $k \in I$ , the integer  $k$  denotes the label of the  $k$ -th vertex of the generalized Dynkin diagrams (for example: label  $A_r$  as in Fig. 2).
- (4) For a generalized Dynkin diagram  $\mathcal{D}$  and  $i, j \in \mathbb{N}$ , denote by  $\mathcal{D}_{i,j}^r$  the diagram  $\mathcal{D}$  of rank  $r$  located in the  $j$ -th position of row  $i$ .

The proof of the theorem is preceded by several special cases as follows.

Let  $X = [M]$ ,  $A^X := (a_{ij})_{i,j \in I}$  be the Cartan matrix of  $X$ ,  $(q_{i,j})_{i,j \in I}$  be the braiding matrix of  $X$  and  $(q_{i,j}^{r_i(X)})_{i,j \in I}$  be the braiding matrix of  $r_i(X)$ . To simplify the labels, we write  $\widetilde{q_{ij}} := q_{ij}q_{ji}$  for  $1 \leq i, j \leq r$ .

The following lemma will be helpful to simplify the proof in what follows.

**Lemma 4.4.** *If  $\mathcal{D}_{\chi,E}$  is of the form*

$$\begin{array}{ccccccccccccccc} q_{11} & r & q_{22} & q & q_{33} & s & q_{44} & t & q_{55} & q_{r-1,r-1} & u & q_{rr} \\ \circ & & \circ & & \circ & & \circ & & \circ & \cdots & \circ & & \circ \end{array}$$

and has a good  $D'_r$  neighborhood, then  $q_{22} = -1$  and  $qr \neq 1$ .

*Proof.* If  $q_{22} \neq -1$ , then  $A^{r_2(X)} = A_r$  by Lemma 2.23, which contradicts Definition 3.10. Then we obtain  $qr \neq 1$  by  $a_{13}^{r_2(X)} = -1$ .  $\square$

**Theorem 4.5.** *If  $\Delta^{[M]}$  is a finite set of roots of non-sporadic finite Cartan graphs of rank  $r \geq 5$  then all possible Dynkin diagrams of  $V$  appear in rows 1 to 10 of Table 1.*

*Proof.* By Theorem 3.11, we know that either  $\mathcal{C}(M)$  is standard or there exists a point  $X$  such that  $A^X$  has a good  $D'_r$  neighborhood.

Case *a.* Assume that  $\mathcal{C}(M)$  is standard of type  $A_r$ . Then  $A^X = A_r$  implies that  $(2)_{q_{ii}}(\widetilde{q_{ii}q_{i,i+1}} - 1) = (2)_{q_{jj}}(\widetilde{q_{jj}q_{j-1,j}} - 1) = 0$ , for all  $i \in \{1, 2, 3, \dots, r-1\}$  and  $j \in \{2, 3, 4, \dots, r\}$ . Applying Lemma 2.23 we distinguish the following subcases:  $a_1, a_2, a_3$ .

Subcase  $a_1$ . Consider that  $\widetilde{q_{ii}q_{i,i+1}} - 1 = \widetilde{q_{jj}q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 2, 3, \dots, r-1\}$  and  $j \in \{2, 3, 4, \dots, r\}$ . To avoid repetition, let  $q := q_{11} \neq -1$ . Then  $\mathcal{D}$ :

$$\begin{array}{ccccccccccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & q & q^{-1} & q & \cdots & q & q^{-1} & q \\ \circ & & \circ & & \circ & & \circ & & \circ & \cdots & \circ & & \circ \end{array}$$

is represented by a simple chain

$$\boxed{C(r, q; )}$$

which appears in row 1 of Table 1.

Subcase  $a_2$ . Consider that  $(2)_{q_{kk}} = 0$  for  $1 \leq k \leq (r+1)/2$  (Avoid symmetrical repetition) and  $\widetilde{q_{ii}q_{i,i+1}} - 1 = \widetilde{q_{jj}q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 2, 3, \dots, r-1\} \setminus \{k\}$  and  $j \in \{2, 3, 4, \dots, r\} \setminus \{k\}$ . Assume that  $r := \widetilde{q_{k-1,k}} \neq -1$  and  $q := \widetilde{q_{k,k+1}} \neq -1$  to avoid the repetition. If  $k \neq 1$ , then  $qr = 1$  by  $a_{k-1,k+1}^{r_k(X)} = 0$ . Hence  $\mathcal{D}$ :

$$\begin{array}{ccccccccccccccc} q & q^{-1} & q & q^{-1} & q & \cdots & q & q^{-1} & -1 & q & q^{-1} & q^{-1} & q & q^{-1} \\ \circ & & \circ & & \circ & \cdots & \circ & & \circ & & \circ & & \circ & & \circ \\ & & & & & & & & k & & & & & & \end{array}$$

is represented by the simple chain

$$\boxed{C(r, q^{-1}; 1, \dots, k)}$$

which appears in row 2 of Table 1. If  $k = 1$ , then  $\mathcal{D}$ :

$$\begin{array}{ccccccccccccccc} -1 & q & q^{-1} & q & q^{-1} & q & q^{-1} & q & q^{-1} & q^{-1} & q & q^{-1} \\ \circ & & \circ & & \circ & & \circ & & \circ & \cdots & \circ & & \circ \end{array}$$

is represented by the simple chain

$$\boxed{C(r, q^{-1}; 1)}$$

which appears in row 2 of Table 1.

Similar reflections and conditions occur in cases  $q_{i_1, i_1} = q_{i_2, i_2} = \dots = q_{i_k, i_k} = -1$  for  $i_1, i_2, \dots, i_k \in \{1, 2, \dots, r\}$ ,  $2 \leq k < r$ , and  $q_{ii} \widetilde{q_{i, i+1}} - 1 = q_{jj} \widetilde{q_{j-1, j}} - 1 = 0$  for all  $i \in \{1, 2, 3, \dots, r-1\} \setminus \{i_1, i_2, \dots, i_k\}$  and  $j \in \{2, 3, 4, \dots, r\} \setminus \{i_1, i_2, \dots, i_k\}$ . In these cases, all the generalized Dynkin diagrams  $\mathcal{D}$  appear in row 2 of Table 1.

Subcase  $a_3$ . Consider that  $(2)_{q_{ii}} = 0$  for all  $i \in \{1, 2, \dots, r\}$ . Then  $\mathcal{D}$  is the case of  $q = -1$  and  $p = 3$  in row 1 of Table 1.

Case  $b$ . Assume that  $\mathcal{C}(M)$  is standard of type  $B_r$ . Then  $A^X = B_r$  implies that  $(3)_{q_{11}}(q_{11}^2 \widetilde{q_{12}} - 1) = (2)_{q_{ii}}(q_{ii} \widetilde{q_{i, i+1}} - 1) = (2)_{q_{jj}}(q_{jj} \widetilde{q_{j-1, j}} - 1) = 0$  for all  $i \in \{2, 3, \dots, r-1\}$  and  $j \in \{2, 3, \dots, r\}$ . Hence we distinguish the following subcases:  $b_1, b_2, b_3, b_4$ .

Subcase  $b_1$ . Consider that  $q_{11}^2 \widetilde{q_{12}} - 1 = q_{ii} \widetilde{q_{i, i+1}} - 1 = q_{jj} \widetilde{q_{j-1, j}} - 1 = 0$  for all  $i \in \{2, 3, \dots, r-1\}$  and  $j \in \{2, 3, \dots, r\}$ . Let  $q := q_{11}$ . Then  $q \neq -1$  and  $\mathcal{D}$ :

$$\begin{array}{cccccccccccccccc} q & q^{-2} & q^2 & q^{-2} & q^2 & q^{-2} & q^2 & q^{-2} & q^2 & & q^2 & q^{-2} & q^2 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \cdots & \circ & \circ & \circ \end{array}$$

is represented by the simple chain

$$\tau_{r \dots 321} \left( \boxed{C(r-1, q^2; )} \right) \xrightarrow{q^{-2}} \overset{q}{\circ}$$

which appears in row 3 of Table 1.

Subcase  $b_2$ . Consider that  $q_{22} = -1$  and  $q_{11}^2 \widetilde{q_{12}} - 1 = q_{ii} \widetilde{q_{i, i+1}} - 1 = q_{jj} \widetilde{q_{j-1, j}} - 1 = 0$  for all  $i \in \{3, 4, \dots, r-1\}$  and  $j \in \{3, 4, \dots, r\}$ . In this case we assume that  $\widetilde{q_{23}} \neq -1$  to avoid the repetition. Let  $q_{11} := q$ ,  $\widetilde{q_{23}} := r$ . Then  $q \neq -1$ ,  $\widetilde{q_{12}} = q^{-2}$ . Hence we have  $\widetilde{q_{12}} \widetilde{q_{23}} = q^{-2} r = 1$  by  $a_{13}^{r_2(X)} = 0$ . Then  $\mathcal{D}$ :

$$\begin{array}{cccccccccccccccc} q & q^{-2} & -1 & q^2 & q^{-2} & q^2 & q^{-2} & q^2 & q^{-2} & & q^{-2} & q^2 & q^{-2} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \cdots & \circ & \circ & \circ \end{array}$$

is represented by the simple chain

$$\tau_{r \dots 321} \left( \boxed{C(r-1, q^2; 1, \dots, r-1)} \right) \xrightarrow{q^{-2}} \overset{q}{\circ}$$

which appears in row 4 of Table 1.

Similar reflections and conditions occur in cases  $q_{kk} = -1$  for  $k \in \{3, \dots, r\}$  and  $q_{11}^2 \widetilde{q_{12}} - 1 = q_{ii} \widetilde{q_{i, i+1}} - 1 = q_{jj} \widetilde{q_{j-1, j}} - 1 = 0$  for all  $i \in \{3, 4, \dots, r-1\} \setminus \{k\}$  and  $j \in \{3, 4, \dots, r\} \setminus \{k\}$ . It happens similarly for subcases with more vertices satisfying  $q_{i_1, i_1} = q_{i_2, i_2} = \dots = q_{i_k, i_k} = -1$  for  $i_1, i_2, \dots, i_k \in \{2, 3, \dots, r\}$  and  $1 < k < r$ . In these cases, all the generalized Dynkin diagrams  $\mathcal{D}$  appear in row 4 of Table 1.

Subcase  $b_3$ . Consider that  $(3)_{q_{11}} = 0$  and  $q_{ii} \widetilde{q_{i, i+1}} - 1 = q_{jj} \widetilde{q_{j-1, j}} - 1 = 0$  for all  $i \in \{2, 3, \dots, r-1\}$  and  $j \in \{2, 3, \dots, r\}$ . Assume that  $q_{11}^2 \widetilde{q_{12}} - 1 \neq 0$  and  $q_{22} \neq -1$  to avoid repetition. Let  $q_{11} := \zeta$  and  $q_{22} := q$ . The reflection of  $X$  is

$$X : \begin{array}{cccccccccccccccc} \zeta & q^{-1} & q & q^{-1} & q & q^{-1} & q & q^{-1} & q & & q & q^{-1} & q \\ \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \cdots & \circ & \circ & \circ \end{array}$$

$$\Rightarrow r_1(X) : \begin{array}{c} \zeta \quad \zeta^{-1}q \quad \zeta q^{-1}q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad \dots \quad q \quad q^{-1} \quad q \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \dots \quad \circ \quad \circ \quad \circ \end{array}$$

with  $(3)_\zeta = 0$  and  $q \notin \{-1, \zeta, \zeta^{-1}\}$  and applying Lemma 2.23 to  $A^{r_1(X)} = B_r$ , we obtain  $\zeta q^{-2} = 1$  or  $\zeta q^{-1} = -1$ . Hence  $q = -\zeta^{-1}$ ,  $p \neq 2$  or  $q = -\zeta$ ,  $p \neq 2$ . If  $p \neq 2, 3$ , then one has  $q = -\zeta^{-1} = -\zeta$ . Hence  $\mathcal{D}$ :

$$\begin{array}{c} \zeta \quad -\zeta \quad -\zeta^{-1} \quad -\zeta \quad -\zeta^{-1} \quad -\zeta \quad -\zeta^{-1} \quad -\zeta \quad -\zeta^{-1} \quad -\zeta^{-1} \quad -\zeta \quad -\zeta^{-1} \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \dots \quad \circ \quad \circ \end{array}$$

is represented by the simple chain

$$\tau_{r \dots 321} \left( \begin{array}{c} \zeta \\ \circ \end{array} \right) \text{---} \left( C(r-1, -\zeta^{-1};) \right) \text{---} \begin{array}{c} -\zeta \\ \circ \end{array}$$

which appears in row 5 of Table 1. If  $p = 3$ , then one has  $q = -1$  for  $q = -\zeta^{-1}$  and  $q = -\zeta$ . Hence  $\mathcal{D}$  appears in row 5' of Table 1.

Subcase  $b_4$ . Consider that  $q_{22} = -1$ ,  $(3)_{q_{11}} = 0$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{3, 4, \dots, r-1\}$  and  $j \in \{3, 4, \dots, r\}$ . Assume that the conditions  $q_{11}^2\widetilde{q_{12}} - 1 = 0$  and  $\widetilde{q_{23}} = -1$  don't hold to avoid repetition. Let  $q_{11} := \zeta$ ,  $q := \widetilde{q_{12}}$  and  $r := \widetilde{q_{23}}$ . Then  $r \neq -1$ ,  $qr = 1$  by  $A^{r_2(X)} = B_r$ . And one has  $(3)_\zeta = 0$  and  $q \notin \{-1, \zeta, \zeta^{-1}\}$ . The reflection of  $X$

$$X : \begin{array}{c} \zeta \quad q \quad -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad \dots \quad q \quad q^{-1} \quad q \\ \bullet \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \dots \quad \circ \quad \circ \quad \circ \end{array}$$

$$\Rightarrow r_1(X) : \begin{array}{c} \zeta \quad (\zeta q)^{-1} \quad -\zeta q^2 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad \dots \quad q \quad q^{-1} \quad q \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \dots \quad \circ \quad \circ \quad \circ \end{array}$$

imply that  $-\zeta q^2 = -1$  or  $-\zeta q^2(\zeta q)^{-1} = -\zeta q^2 q^{-1} = 1$  by  $A^{r_1(X)} = B_r$ . Then  $q = -\zeta$ ,  $p \neq 2, 3$ . Hence  $\mathcal{D}$ :

$$\begin{array}{c} \zeta \quad -\zeta \quad -1 \quad -\zeta^{-1} \quad -\zeta \quad -\zeta^{-1} \quad -\zeta \quad -\zeta^{-1} \quad -\zeta \quad \dots \quad -\zeta \quad -\zeta^{-1} \quad -\zeta \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \dots \quad \circ \quad \circ \quad \circ \end{array}$$

is represented by the simple chain

$$\tau_{r \dots 321} \left( \begin{array}{c} \zeta \\ \circ \end{array} \right) \text{---} \left( C(r-1, -\zeta^{-1}; 1, \dots, r-1) \right) \text{---} \begin{array}{c} -\zeta \\ \circ \end{array}$$

which appears in row 6 of Table 1.

Similar reflections and conditions occur in cases  $q_{kk} = -1$  for  $k \in \{3, 4, \dots, r\}$ ,  $(3)_{q_{11}} = 0$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{3, 4, \dots, r-1\} \setminus \{k\}$  and  $j \in \{3, 4, \dots, r\} \setminus \{k\}$ . It happens similarly for subcases with more vertices satisfying  $q_{i_1, i_1} = q_{i_2, i_2} = \dots = q_{i_k, i_k} = -1$  for  $i_1, i_2, \dots, i_k \in \{2, 3, \dots, r\}$  and  $1 < k < r$ . In these cases, all the generalized Dynkin diagrams  $\mathcal{D}$  appear in row 6 of Table 1.

Case  $c$ . Assume that  $\mathcal{C}(M)$  is standard of type  $C_r$ . By  $A^X = C_r$ , we obtain  $(3)_{q_{22}}(q_{22}^2\widetilde{q_{12}} - 1) = (2)_{q_{ii}}(q_{ii}\widetilde{q_{i,i+1}} - 1) = (2)_{q_{jj}}(q_{jj}\widetilde{q_{j-1,j}} - 1) = 0$  for all  $i \in \{1, 3, \dots, r-1\}$  and  $j \in \{3, 4, \dots, r\}$ . Hence we have the following subcases:  $c_1, c_2$ .

Subcase  $c_1$ . Consider that  $q_{22}^2\widetilde{q_{12}} - 1 = q_{22}\widetilde{q_{23}} - 1 = (2)_{q_{ii}}(q_{ii}\widetilde{q_{i,i+1}} - 1) = (2)_{q_{jj}}(q_{jj}\widetilde{q_{j-1,j}} - 1) = 0$  for all  $i \in \{1, 3, \dots, r-1\}$  and  $j \in \{3, 4, \dots, r\}$ . Let  $q := q_{22}$ . Then  $q \neq -1$ ,  $\widetilde{q_{12}} = q^{-2}$

and  $\widetilde{q_{23}} = q^{-1}$ . Hence  $q_{33}\widetilde{q_{23}} = 1$ . Otherwise, if  $q_{33}\widetilde{q_{23}} \neq 1$  then  $q_{33} = -1$  and hence  $a_{21}^{r_3} = -1$ , which is a contradiction. Then  $q_{33}\widetilde{q_{34}} = 1$  by  $a_{34} = -1$ . Hence  $q_{44}\widetilde{q_{34}} = 1$  and  $q_{44} \neq -1$ . Otherwise, if  $q_{44} = -1$  then  $a_{21}^{r_3 r_4(X)} = -1$  by Lemma 2.23, which is again a contradiction. In the same way, we obtain that  $q_{ii}\widetilde{q_{i,i+1}} = q_{jj}\widetilde{q_{j-1,j}} = 1$  for all  $i \in \{3, 4, \dots, r-1\}$  and  $j \in \{3, 4, \dots, r\}$ . If  $q_{11} = -1$  and  $\widetilde{q_{12}} \neq -1$ , then the reflection of  $X$

$$X : \overset{-1}{\bullet} \overset{q^{-2}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \dots \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ}$$

$$\Rightarrow r_1(X) : \overset{-1}{\circ} \overset{q^2}{\circ} \overset{-q^{-1}}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \dots \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ}$$

gives that  $-q^{-1}q^{-1} = 1$  by applying Lemma 2.23 on  $a_{23}^{r_1(X)} = -1$ . Then  $\widetilde{q_{12}} = q^{-2} = -1$ , which is a contradiction. If  $q_{11}\widetilde{q_{12}} - 1 = 0$ , then  $\mathcal{D}$ :

$$\overset{q^2}{\circ} \overset{q^{-2}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \dots \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ}$$

is represented by the simple chain

$$\tau_{r \dots 321} \left( C(r-2, q; ) \right) \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-2}}{\circ} \overset{q^2}{\circ}$$

which appears in row 7 of Table 1.

Subcase  $c_2$ . Consider that  $(3)_{q_{22}} = q_{22}\widetilde{q_{23}} - 1 = (2)_{ii}(q_{ii}\widetilde{q_{i,i+1}} - 1) = (2)_{jj}(q_{jj}\widetilde{q_{j-1,j}} - 1) = 0$  for all  $i \in \{1, 3, \dots, r-1\}$  and  $j \in \{3, 4, \dots, r\}$ . Assume that  $q_{22}^2\widetilde{q_{12}} - 1 \neq 0$  to avoid repetition. Let  $q_{22} := \zeta$ . Then  $\widetilde{q_{12}} \notin \{\zeta, \zeta^{-1}\}$ . By  $q_{22}\widetilde{q_{23}} - 1 = 0$  one obtains that  $\widetilde{q_{23}} = \zeta^{-1}$ , and  $a_{13}^{r_2(X)} = 0$  yields  $\widetilde{q_{12}}(q_{22})^{-1} = 1$ . Hence  $\widetilde{q_{12}} = \zeta$ , which is a contradiction.

Case  $d$ . Assume that  $\mathcal{C}(M)$  is standard of type  $D_r$ . Applying Lemma 2.23 to  $A^X = D_r$  shows that  $(2)_{q_{33}}(q_{33}\widetilde{q_{3i}} - 1) = (2)_{q_{ii}}(q_{ii}\widetilde{q_{3i}} - 1) = 0$  for all  $i \in \{1, 2, 4\}$  and  $(2)_{q_{jj}}(q_{jj}\widetilde{q_{j,j+1}} - 1) = (2)_{q_{kk}}(q_{kk}\widetilde{q_{k-1,k}} - 1) = 0$  for all  $j \in \{4, 5, \dots, r-1\}$  and  $k \in \{5, 6, \dots, r\}$ . This will be done in several subcases:  $d_1, d_2, d_3$ .

Subcase  $d_1$ . Consider that  $q_{33}\widetilde{q_{3i}} - 1 = q_{ii}\widetilde{q_{3i}} - 1 = q_{jj}\widetilde{q_{j,j+1}} - 1 = q_{kk}\widetilde{q_{k-1,k}} - 1 = 0$  for all  $i \in \{1, 2, 4\}$ ,  $j \in \{4, 5, \dots, r-1\}$  and  $k \in \{5, 6, \dots, r\}$ . To avoid repetition, let  $q := q_{33} \neq -1$ . Then  $\mathcal{D}$ :

$$\begin{array}{c} \overset{q}{\circ} \overset{q^{-1}}{\circ} \\ \diagdown \quad \diagup \\ \overset{q}{\circ} \overset{q^{-1}}{\circ} \end{array} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \dots \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ}$$

is represented by the simple chain

$$\tau_{r \dots 321} \left( C(r-2, q; ) \right) \begin{array}{c} \overset{q^{-1}}{\circ} \overset{q}{\circ} \\ \diagup \quad \diagdown \\ \overset{q^{-1}}{\circ} \overset{q}{\circ} \end{array}$$


which appears in row 8 of Table 1.

Subcase  $d_2$ . Consider that  $q_{33}\widetilde{q_{3i}} - 1 = 0$  for all  $i \in \{1, 2, 4\}$  and there exists  $l \in \{1, 2, 4, 5, \dots, r\}$  such that  $(2)_{q_{ll}} = (2)_{q_{mm}}q_{mm}\widetilde{q_{3m}} - 1 = (2)_{q_{jj}}q_{jj}\widetilde{q_{j,j+1}} - 1 = (2)_{q_{kk}}q_{kk}\widetilde{q_{k-1,k}} - 1 = 0$  for all  $m \in \{1, 2, 4\} \setminus \{l\}$ ,  $j \in \{4, 5, \dots, r-1\} \setminus \{l\}$ ,  $k \in \{5, 6, \dots, r\} \setminus \{l\}$ . Assume that  $q_{33} \neq -1$  in this case to avoid repetition. Let  $q := q_{33}$ . Then  $q \neq \pm 1$ . We obtain that  $q^{-2} = 1$  by applying Lemma 2.23 to  $A^{r_3 \dots r_l(X)} = D_r$ , which is a contradiction. It happens similarly for subcases with more vertices satisfying  $(2)_{q_{i_1, i_1}} = (2)_{q_{i_2, i_2}} = \dots = (2)_{q_{i_k, i_k}} = 0$  for  $i_1, i_2, \dots, i_k \in \{1, 2, 4, 5, \dots, r\}$  and  $1 < k < r$ , a contradiction can be obtained in the same way.


Subcase  $d_3$ . Consider  $(2)_{q_{33}} = (2)_{q_{ii}}(q_{ii}\widetilde{q_{3i}} - 1) = (2)_{q_{jj}}(q_{jj}\widetilde{q_{j,j+1}} - 1) = (2)_{q_{kk}}(q_{kk}\widetilde{q_{k-1,k}} - 1) = 0$  for all  $i \in \{1, 2, 4\}$ ,  $j \in \{4, 5, \dots, r-1\}$ ,  $k \in \{5, \dots, r\}$ . let  $q := \widetilde{q_{31}}$ ,  $r := \widetilde{q_{32}}$ ,  $s := \widetilde{q_{34}}$ . To avoid repetition, let  $q, r, s \neq -1$ . Since  $A^{r_3(X)} = D_r$ , we have  $qr = rs = qs = 1$ . Then  $q = r = s = -1$ , which is a contradiction.

Case  $d'$ . Assume that  $A^X$  has a good  $D'_r$  neighborhood. Let  $a := a_{21}^{r_3(X)}$ . By  $A^X = A_r$  we have  $(2)_{q_{ii}}(q_{ii}\widetilde{q_{i,i+1}} - 1) = (2)_{q_{jj}}(q_{jj}\widetilde{q_{j-1,j}} - 1) = 0$  for all  $i \in \{1, 2, 3, \dots, r-1\}$  and  $j \in \{2, 3, 4, \dots, r\}$ . From Lemma 4.4, we know  $q_{22} = -1$  and  $\widetilde{q_{12}}\widetilde{q_{23}} \neq 1$  for all cases. Now we have the following subcases:  $d'_1, d'_2, d'_3, d'_4$ .


Subcase  $d'_1$ . Consider that  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 3, 4, \dots, r-1\}$  and  $j \in \{3, 4, \dots, r\}$ . We assume that  $\widetilde{q_{23}} \neq -1$  to avoid repetition. From Lemma 2.23 one has  $a = 1$ . Let  $r := \widetilde{q_{12}}$  and  $q := \widetilde{q_{23}}$ . Then  $qr \neq 1$  and  $q \neq -1$ . The reflections of  $X$



imply that  $q^2r = 1$  by  $a_{14}^{r_3r_2(X)} = 0$ , and hence  $\mathcal{D}$ :



is represented by the simple chain



which appears in row 10 of Table 1.

Similar reflections and conditions also hold in cases  $q_{i_1, i_1} = q_{i_2, i_2} = \dots = q_{i_k, i_k} = -1$  for  $i_1, i_2, \dots, i_k \in \{4, 5, \dots, r\}$ ,  $1 \leq k < r-2$ , and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 3, 4, \dots, r-1\} \setminus \{i_1, i_2, \dots, i_k\}$  and  $j \in \{3, 4, \dots, r\} \setminus \{i_1, i_2, \dots, i_k\}$ . In these cases, all the generalized Dynkin diagrams  $\mathcal{D}$  appear in row 10 of Table 1.

Subcase  $d'_2$ . Consider that  $q_{11} = -1$ ,  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{3, 4, \dots, r-1\}$  and  $j \in \{3, 4, \dots, r\}$ . From Lemma 2.23, it follows that  $a = 1$ . Let  $r := \widetilde{q_{12}}$  and  $q := \widetilde{q_{23}}$ .

Then  $qr \neq 1$ . By applying Lemma 2.23 to  $a_{23}^{r_1(X)} = -1$ , we have  $r = -1, q \neq -1$ . The reflections of  $X$

$$\begin{aligned}
X : \begin{array}{c} -1 \quad -1 \quad -1 \quad q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad \dots \quad q^{-1} \quad q \quad q^{-1} \\ \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \dots \quad \circ \quad \circ \quad \circ \end{array} &\Rightarrow r_2(X) : \begin{array}{c} -1 \\ \circ \quad -q \quad -1 \\ \circ \quad q^{-1} \quad -1 \\ -1 \end{array} \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad \dots \quad q^{-1} \quad q \quad q^{-1} \\ \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \dots \quad \circ \quad \circ \quad \circ \end{array} \\
&\Rightarrow r_3 r_2(X) : \begin{array}{c} -q \\ \circ \quad (-q)^{-1} \quad -q^2 \\ \circ \quad q^{-1} \quad -1 \\ -1 \end{array} \begin{array}{c} q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad \dots \quad q^{-1} \quad q \quad q^{-1} \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \dots \quad \circ \quad \circ \quad \circ \end{array}
\end{aligned}$$

imply that  $q^2 = -1$  by  $a_{14}^{r_3 r_2(X)} = 0$ . Then  $q \in G'_4$  and  $p \neq 2$ . Hence  $\mathcal{D}$ :

$$\begin{array}{c} -1 \quad -1 \quad -1 \quad q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad \dots \quad q^{-1} \quad q \quad q^{-1} \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \dots \quad \circ \quad \circ \quad \circ \end{array}$$

is represented by the simple chain

$$\tau_{r \dots 321} \left( C(r-1, q; 1, \dots, r-1) \right) \begin{array}{c} -1 \quad -1 \\ \circ \quad \circ \end{array}$$

which is the case of  $q^2 = -1$  and  $p \neq 2$  in row 10 of Table 1.

We apply a similar proof as above in cases  $q_{11} = q_{i_1, i_1} = q_{i_2, i_2} = \dots = q_{i_k, i_k} = -1$  for  $i_1, i_2, \dots, i_k \in \{4, 5, \dots, r\}$ ,  $1 \leq k < r-2$ , and  $q_{ii} \widetilde{q_{i, i+1}} - 1 = q_{jj} \widetilde{q_{j-1, j}} - 1 = 0$  for all  $i \in \{3, 4, \dots, r-1\} \setminus \{i_1, i_2, \dots, i_k\}$  and  $j \in \{3, 4, \dots, r\} \setminus \{i_1, i_2, \dots, i_k\}$ . In these cases, all the generalized Dynkin diagrams  $\mathcal{D}$  are the case of  $q^2 = -1$  and  $p \neq 2$  in row 10 of Table 1.

Subcase  $d'_3$ . Consider that  $q_{33} = -1$ ,  $q_{ii} \widetilde{q_{i, i+1}} - 1 = q_{jj} \widetilde{q_{j-1, j}} - 1 = 0$  for all  $i \in \{1, 4, \dots, r-1\}$  and  $j \in \{4, 5, \dots, r\}$ . Assume that  $\widetilde{q_{34}} \neq -1$  to avoid repetition. Let  $r := \widetilde{q_{12}}$ ,  $q := \widetilde{q_{23}}$  and  $s := \widetilde{q_{34}}$ . Then  $qr \neq 1$ ,  $s \neq -1$ . We have  $qs = 1$  by  $a_{24}^{r_3(X)} = 0$ . The reflections of  $X$

$$\begin{aligned}
r_2(X) : \begin{array}{c} -1 \\ \circ \quad qr \\ \circ \quad q^{-1} \quad -1 \\ \circ \quad q^{-1} \quad -1 \end{array} \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad \dots \quad q \quad q^{-1} \quad q \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \dots \quad \circ \quad \circ \quad \circ \end{array} &\Leftarrow X : \begin{array}{c} r^{-1} \quad r \quad -1 \quad q \quad -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad \dots \quad q \quad q^{-1} \quad q \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \dots \quad \circ \quad \circ \quad \circ \end{array} \\
&\Rightarrow r_3(X) : \begin{array}{c} r^{-1} \quad r \quad q \quad q^{-1} \quad -1 \quad q \quad -1 \quad q^{-1} \quad q \quad \dots \quad q \quad q^{-1} \quad q \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \dots \quad \circ \quad \circ \quad \circ \end{array}
\end{aligned}$$

imply that  $q^2 r = 1$  by  $a_{13}^{r_2(X)} = -1$  and  $a = 2$ . Then  $\mathcal{D}$ :

$$\begin{array}{c} q^2 \quad q^{-2} \quad -1 \quad q \quad -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad \dots \quad q \quad q^{-1} \quad q \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \dots \quad \circ \quad \circ \quad \circ \end{array}$$

is represented by the simple chain

$$\tau_{r \dots 321} \left( C(r-2, q^{-1}; 1, \dots, r-2) \right) \begin{array}{c} q \quad -1 \quad q^{-2} \quad q^2 \\ \circ \quad \circ \quad \circ \quad \circ \end{array}$$



which appears in row 9 of Table 1.

Applying the aforementioned method in cases  $q_{33} = q_{i_1, i_1} = q_{i_2, i_2} = \dots = q_{i_k, i_k} = -1$  for  $i_1, i_2, \dots, i_k \in \{4, 5, \dots, r\}$ ,  $1 \leq k < r - 2$ , and  $\widetilde{q_{ii}q_{i, i+1}} - 1 = \widetilde{q_{jj}q_{j-1, j}} - 1 = 0$  for all  $i \in \{1, 4, \dots, r-1\} \setminus \{i_1, i_2, \dots, i_k\}$  and  $j \in \{4, \dots, r\} \setminus \{i_1, i_2, \dots, i_k\}$ , we obtain that all the generalized Dynkin diagrams  $\mathcal{D}$  appear in row 10 of Table 1.

Subcase  $d'_4$ . Consider that  $q_{11} = q_{33} = -1$ ,  $\widetilde{q_{ii}q_{i, i+1}} - 1 = \widetilde{q_{jj}q_{j-1, j}} - 1 = 0$  for all  $i \in \{4, \dots, r-1\}$  and  $j \in \{4, 5, \dots, r\}$ . Let  $r := q'_{12}$ ,  $q := \widetilde{q_{23}}$  and  $s := \widetilde{q_{34}}$ , then  $qr \neq 1$ . To avoid repetition, let  $s \neq -1$ . We have  $r = -1$ ,  $p \neq 2$ , and  $qs = 1$  by  $a_{23}^{r_1(X)} = -1$  and  $a_{24}^{r_3(X)} = 0$ . The reflections of  $X$

$$\begin{aligned}
 r_2(X) : & \begin{array}{c} -1 \\ \circ \\ \diagup \quad \diagdown \\ -1 \quad q \quad q^{-1} \\ \diagdown \quad \diagup \\ \circ \\ -1 \end{array} \begin{array}{c} q \\ q^{-1} \\ q \\ q^{-1} \\ q \\ \dots \\ q \\ q^{-1} \\ q \end{array} \Leftarrow X : \begin{array}{c} -1 \quad -1 \quad -1 \quad q \quad -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad \dots \quad q \quad q^{-1} \quad q \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \dots \quad \circ \quad \circ \quad \circ \end{array} \\
 \Rightarrow r_3(X) : & \begin{array}{c} -1 \quad -1 \quad q \quad q^{-1} \quad -1 \quad q \quad -1 \quad q^{-1} \quad q \quad \dots \quad q \quad q^{-1} \quad q \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \dots \quad \circ \quad \circ \quad \circ \end{array}
 \end{aligned}$$

imply that  $q^2 = -1$  by  $a_{31}^{r_2(X)} = -1$  and hence  $a = 2$ . Then  $\mathcal{D}$ :

$$\begin{array}{c} -1 \quad -1 \quad -1 \quad q \quad -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad \dots \quad q \quad q^{-1} \quad q \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \dots \quad \circ \quad \circ \quad \circ \end{array}$$

is represented by the simple chain

$$\tau_{r \dots 321} \left( \text{C}(r-2, q^{-1}; 1, \dots, r-2) \right) \begin{array}{c} q \quad -1 \quad -1 \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \end{array}$$

which is the case of  $q^2 = -1$  and  $p \neq 2$  in row 9 of Table 1.

By applying the aforementioned method in cases  $q_{11} = q_{33} = q_{i_1, i_1} = q_{i_2, i_2} = \dots = q_{i_k, i_k} = -1$  for  $i_1, i_2, \dots, i_k \in \{4, 5, \dots, r\}$ ,  $1 \leq k < r - 2$ , and  $\widetilde{q_{ii}q_{i, i+1}} - 1 = \widetilde{q_{jj}q_{j-1, j}} - 1 = 0$  for all  $i \in \{4, \dots, r-1\} \setminus \{i_1, i_2, \dots, i_k\}$  and  $j \in \{4, \dots, r\} \setminus \{i_1, i_2, \dots, i_k\}$ , we obtain that all the generalized Dynkin diagrams  $\mathcal{D}$  are the case of  $q^2 = -1$  and  $p \neq 2$  in row 10 of Table 1.  $\square$

Theorem 3.15 allows us to discuss case by case by hand. The following discussions cover the sporadic cases of ranks 5, 6, 7, and 8.

**Lemma 4.6.** *If  $\mathcal{D}_{\chi, E}$  is of the form*

$$\begin{array}{c} q_{11} \quad q \quad q_{22} \quad r \quad q_{33} \quad s \quad q_{44} \quad t \quad q_{55} \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}$$

*and has a good  $A_5$  neighborhood, then there exists  $i \in \{3, 4\}$  such that  $q_{ii} = -1$  and the following cases are impossible.*

$$(a_1) \quad \begin{array}{c} q^{-1} \quad q \quad q^{-1} \quad q \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}, \quad q \neq -1, p \neq 2$$

$$\begin{aligned}
(a_2) \quad & \begin{array}{c} -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad q \quad q^{-1} \quad q \quad q^{-1} \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}, \quad q \neq -1, p \neq 2 \\
(a_3) \quad & \begin{array}{c} -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad q \quad q^{-1} \quad q \quad -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}, \quad q \neq -1, p \neq 2 \\
(a_4) \quad & \begin{array}{c} -1 \quad q \quad q^{-1} \quad q \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}, \quad q \neq -1, p \neq 2 \\
(b_1) \quad & \begin{array}{c} -1 \quad q^{-2} \quad q^2 \quad q^{-2} \quad q^2 \quad q^{-2} \quad -1 \quad q \quad -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}, \quad q \notin G'_3, p \neq 3 \\
(b_2) \quad & \begin{array}{c} -1 \quad q^2 \quad -1 \quad q^{-2} \quad q^2 \quad q^{-2} \quad -1 \quad q \quad -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}, \quad q \notin G'_3, p \neq 3 \\
(b_3) \quad & \begin{array}{c} -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad q \quad -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}, \quad q \neq -1, p \neq 2 \\
(c_1) \quad & \begin{array}{c} q^{-1} \quad q \quad -1 \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad -1 \quad -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}, \quad q \neq -1, p \neq 2 \\
(c_2) \quad & \begin{array}{c} -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad -1 \quad -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}, \quad q \neq -1, p \neq 2 \\
(c_3) \quad & \begin{array}{c} -1 \quad q \quad -1 \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad -1 \quad -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}, \quad q \neq -1, p \neq 2 \\
(c_4) \quad & \begin{array}{c} -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad q^{-1} \quad q \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}, \quad q \neq -1, p \neq 2
\end{aligned}$$

*Proof.* If  $q_{33} \neq -1$  and  $q_{44} \neq -1$ , then  $A^{r_3(X)} = A^{r_4(X)} = A_5$  by Lemma 2.23, which contradicts Definition 3.12. In the following we analyze the cases  $(a_1) - (c_3)$ .

- (i) If  $X$  is of the type  $\begin{array}{c} q_{11} \quad q \quad q^{-1} \quad q \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$  then it satisfies the case  $(A_{5_1})$  of Definition 3.12. Then the transformation

$$\begin{array}{c} q_{11} \quad q \quad q^{-1} \quad q \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \\ \circ \text{---} \circ \text{---} \bullet \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array} \Rightarrow \begin{array}{c} \begin{array}{c} -1 \\ \circ \end{array} \\ \begin{array}{c} q^{-1} \quad -1 \\ \circ \text{---} \circ \end{array} \\ \begin{array}{c} -1 \quad -1 \quad -1 \\ \circ \text{---} \circ \text{---} \circ \end{array} \end{array}$$

implies  $q_{11} = q^{-1}$  or  $q_{11} = -1$ . By  $a_{41}^{r_2 r_3(X)} = 0$ , one has  $q \in G'_4$  and hence the further transformation

$$\begin{array}{c} \begin{array}{c} -1 \\ \circ \end{array} \\ \begin{array}{c} q^{-1} \quad -1 \\ \circ \text{---} \circ \end{array} \\ \begin{array}{c} -1 \quad -1 \quad -1 \\ \circ \text{---} \circ \text{---} \circ \end{array} \end{array} \Rightarrow \begin{array}{c} q^{-1} \quad q \\ \circ \text{---} \circ \end{array} \begin{array}{c} -1 \quad -q^{-1} \quad -q \quad -1 \quad -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$$

where

$$q'_{11} = \begin{cases} -1 & \text{if } q_{11} = q^{-1}, \\ q & \text{if } q_{11} = -1. \end{cases}$$

implies  $a_{45}^{r_2 r_3(X)} = -2$ , which contradicts  $a_{45}^{r_2 r_3(X)} = -1$ . Hence the cases  $(a_1)$  and  $(a_4)$  are impossible.

(ii) If  $X$  is of the type  $\begin{array}{ccccccc} -1 & -1 & -1 & -1 & -1 & q & q^{-1} & q & q_{55} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$  then the transformation

$$\begin{array}{ccccccc} -1 & -1 & -1 & -1 & -1 & q & q^{-1} & q & q_{55} \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow \begin{array}{ccccccc} & & & & -1 & & & & \\ & & & & \circ & & & & \\ -1 & -1 & -1 & -1 & q^{-1} & -q & -1 & q & q_{55} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $q_{55} = q^{-1}$  or  $q_{55} = -1$ . From  $a_{41}^{r_2 r_3(X)} = 0$ , we have  $q = 1$ , which is a contradiction. Hence the cases  $(a_2)$  and  $(a_3)$  are impossible.

(iii) If  $X$  is of the type  $\begin{array}{ccccccc} -1 & q^{-2} & q^2 & q^{-2} & q^2 & q^{-2} & -1 & q & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$  then it falls into case  $(A_{5_2})$  of Definition 3.12. The following transformations

$$\begin{array}{ccccccc} -1 & q^{-2} & q^2 & q^{-2} & q^2 & q^{-2} & -1 & q & -1 \\ \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow \begin{array}{ccccccc} -1 & q^2 & -1 & q^{-2} & q^2 & q^{-2} & -1 & q & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

$$\Rightarrow \begin{array}{ccccccc} -1 & q^{-2} & -1 & q^2 & -1 & q^{-2} & -1 & q & -1 \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow \begin{array}{ccccccc} -1 & q^{-2} & q^2 & q^{-2} & -1 & q^2 & q^{-2} & q & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

give  $a_{45}^{r_3 r_2 r_1(X)} = -3$ , which contradicts  $a_{45}^{r_3 r_2 r_1(X)} \neq -3$ . Hence the case  $(b_1)$  is impossible.

(iv) If  $X$  is of the type  $\begin{array}{ccccccc} -1 & q^2 & -1 & q^{-2} & q^2 & q^{-2} & -1 & q & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$  then it satisfies the conditions for case  $(A_{5_2})$  in Definition 3.12. Then we have

$$\begin{array}{ccccccc} -1 & q^2 & -1 & q^{-2} & q^2 & q^{-2} & -1 & q & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ \end{array} \Rightarrow \begin{array}{ccccccc} -1 & q^2 & -1 & q^{-2} & -1 & q^2 & q^{-2} & -1 & q \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow \begin{array}{ccccccc} & & & & q^2 & & & & \\ & & & & \circ & & & & \\ -1 & q^2 & -1 & q^{-2} & q^{-1} & -q & -1 & q & q \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow \tau_{12534} \begin{array}{ccccccc} & & & & -1 & & & & \\ & & & & \circ & & & & \\ -1 & q^2 & q^{-2} & q^2 & q & -1 & q^{-2} & q^2 & \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

Hence  $q = 1$  by  $a_{15}^{r_2 r_3 r_4(X)} = 0$ , which is a contradiction. Hence the case  $(b_2)$  is impossible.


(v) If  $X$  is of the type  $\begin{array}{ccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & q & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$  it satisfies the case  $(A_{5_2})$  in Definition 3.12. Then we have the reflections of  $X$

$$\begin{array}{ccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & q & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ \end{array} \Rightarrow \begin{array}{ccccccc} & & & & -1 & & & & \\ & & & & \circ & & & & \\ -1 & -1 & -1 & -1 & -1 & -1 & -q & q & \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$


$$\Rightarrow \tau_{12534} \begin{array}{ccccccc} & & & & -1 & & & & \\ & & & & \circ & & & & \\ -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 & \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow \tau_{51234} \begin{array}{ccccccc} -1 & & & & & & & & \\ \circ & & & & & & & & \\ q^{-1} & -1 & -1 & -1 & -1 & -1 & -1 & & \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

and obtain that  $q \in G'_4$  by  $a_{53}^{r_4(X)} = -1$  and  $a_{15}^{r_2 r_3 r_4(X)} = -1$ , which contradicts  $a_{15}^{r_2 r_3 r_4(X)} = 0$ . Hence the case  $(b_3)$  is impossible.

(vi) If  $X$  is of the type  $\begin{array}{ccccccc} q^{-1} & q & -1 & q^{-1} & q & q^{-1} & -1 & -1 & -1 \\ \circ & - & \circ & - & \circ & - & \circ & - & \circ \end{array}$  then it belongs to the case  $(A_{5_3})$  in Definition 3.12. On the one hand, we have




and  $q \in G'_3$  by  $a_{52}^{r_3 r_4(X)} = -1$ . On the other hand, the transformation



implies that  $a_{32}^{r_5 r_4(X)} = -2$  by  $q \in G'_3$ , which contradicts  $a_{32}^{r_5 r_4(X)} = -1$ . Hence the case  $(c_1)$  is impossible. Similarly, the same reasoning as in the proof of case  $(c_1)$  shows that cases  $(c_2)$  and  $(c_3)$  are also impossible.

(vii) If  $X$  is of the type  $\begin{array}{ccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & q^{-1} & q \\ \circ & - & \circ & - & \circ & - & \circ & - & \circ \end{array}$  then the reflections of  $X$



imply  $q = 1$  by  $a_{32}^{r_5 r_4(X)} = -1$ , which is a contradiction. Hence the case  $(c_4)$  is impossible.

□

Lemma 4.6 allows us to give a much simpler proof in what follows.

**Theorem 4.7.** *If  $\Delta^{[M]}$  is a finite set of roots of sporadic finite Cartan graphs of rank 5 then all possible Dynkin diagrams of  $V$  appear in rows 11 to 15 of Table 1.*

*Proof.* Let  $a := -a_{24}^{r_3(X)}$ ,  $b := -a_{42}^{r_3(X)}$ ,  $c := -a_{35}^{r_4(X)}$ ,  $d := -a_{53}^{r_4(X)}$ , and  $e := -a_{43}^{r_5(X)}$ . From Theorem 3.15, we have that  $A^X$  has a good  $A_5$  neighborhood. By Lemma 4.6, there exists  $i \in \{3, 4\}$  such that  $q_{ii} = -1$  for all cases. Then  $A^X = A_5$  implies  $(2)_{q_{ii}}(\widetilde{q_{ii} q_{i,i+1}} - 1) = (2)_{q_{jj}}(\widetilde{q_{jj} q_{j-1,j}} - 1) = 0$  for all  $i \in \{1, 2, 3, 4\}$  and  $j \in \{2, 3, 4, 5\}$ . Depending on the number of vertices that satisfy  $q_{ii} = -1$  for  $i \in \{1, 2, 3, 4, 5\}$ , we distinguish the following categories: Step 1 to Step 5.

Step 1. Assume that there is one vertex labeled by  $-1$ . We split in two cases.

Step 1.1. Consider that  $q_{33} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 2, 4\}$  and  $j \in \{2, 4, 5\}$ . Then  $e = 1$ . By Definition 3.12 one obtains that  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . Let  $q := q_{11}$  and  $r := q_{44}$ . From  $a_{24}^{r_3(X)} = -1$ , we have  $qr \neq 1$ . And the reflections of  $X$

$$\begin{aligned}
 X : & \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad r \quad r^{-1} \quad r \\ \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_3(X) : \begin{array}{c} \begin{array}{c} -1 \\ \circ \end{array} \\ q \quad q^{-1} \quad -1 \quad \begin{array}{c} q \quad r \\ \circ \quad \circ \end{array} \quad -1 \quad (qr)^{-1} \quad -1 \quad (r^{-1}) \quad r \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \\
 \Rightarrow r_2 r_3(X) : & \begin{array}{c} \begin{array}{c} -1 \\ \circ \end{array} \\ q \quad q^{-1} \quad -1 \quad \begin{array}{c} q \quad q^{-2} r^{-1} \\ \circ \quad \circ \end{array} \quad qr \quad (qr)^{-1} \quad r^{-1} \quad r \\ \tau_{32145} \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}
 \end{aligned}$$

imply  $q^2 r = 1$  by  $a_{41}^{r_2 r_3(X)} = 0$ ,  $qr = -1$  or  $qr^2 = 1$  by  $a_{45}^{r_2 r_3(X)} = -1$ . Then we obtain  $q = r \in G'_3$  and  $p \neq 3$ . Hence  $\mathcal{D} = \mathcal{D}_{11,1}^5$ .

Step 1.2. Consider that  $q_{44} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 2, 3\}$  and  $j \in \{2, 3, 5\}$ . Then  $(a, b, e) = (0, 0, 1)$ . Hence in this case  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . Let  $q := q_{11}$  and  $r := q_{55}$ . By  $a_{35}^{r_4(X)} = -1$ , we obtain  $qr \neq 1$ . On the one hand, the reflections of  $X$

$$\begin{aligned}
 X : & \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad r \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} \begin{array}{c} -1 \\ \circ \end{array} \\ q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad \begin{array}{c} q \quad r \\ \circ \quad \circ \end{array} \quad (qr)^{-1} \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \end{array} \\
 \Rightarrow r_5 r_4(X) : & \begin{array}{c} \tau_{12354} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad (qr)^{-1} \quad qr \quad -1 \quad r^{-1} \quad r \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}
 \end{aligned}$$

imply  $qr = -1$  or  $q^2 r = 1$  by  $a_{32}^{r_5 r_4(X)} = -1$ . On the other hand, the reflections of  $X$

$$\begin{aligned}
 X : & \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad r \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} \begin{array}{c} -1 \\ \circ \end{array} \\ q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad \begin{array}{c} q \quad r \\ \circ \quad \circ \end{array} \quad (qr)^{-1} \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \\
 \Rightarrow r_3 r_4(X) : & \begin{array}{c} \begin{array}{c} (qr)^{-1} \\ \circ \end{array} \\ q \quad q^{-1} \quad -1 \quad \begin{array}{c} q^{-2} r^{-1} \quad qr \\ \circ \quad \circ \end{array} \quad -1 \quad q^{-1} \quad q \\ \tau_{12534} \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}
 \end{aligned}$$

imply  $qr = -1$  or  $q^3 r^2 = 1$  by  $a_{52}^{r_3 r_4(X)} = -1$ . Summing up all proceeding conditions, we have  $qr = -1$  and  $p \neq 2$ . Then  $-q^{-2} = 1$  by  $a_{15}^{r_2 r_3 r_4(X)} = 0$ . Hence  $q \in G'_4$  and  $\mathcal{D} = \mathcal{D}_{14,4}^5$ ,  $p \neq 2$ .

Step 2. Assume that there are two vertices labeled by  $-1$ . Below we distinguish the following seven steps.

Step 2.1. Consider that  $q_{44} = q_{55} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 2, 3\}$  and  $j \in \{2, 3\}$ . Hence  $(a, b) = (0, 0)$ . From Definition 3.12, it follows that either  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$  or  $(a, b, c, d, e) = (0, 0, 1, 1, 2)$ . Let  $q := q_{11}$  and  $r^{-1} := \widetilde{q_{45}}$ . If

$$(a, b, c, d, e) = (0, 0, 1, 1, 1), \text{ that is } A^{r_3(X)} = A_5, A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix}, \text{ and }$$

$A^{r_5(X)} = A_5$ . Then  $qr \neq 1$ . The reflection of  $X$

$$X : \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \bullet \end{array} \Rightarrow r_5(X) : \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad r^{-1} \quad r \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . Then the reflections of  $X$

$$\begin{aligned}
 X : & \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad -1 \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \begin{array}{c} q \\ -1 \\ -q^{-1} \\ -1 \end{array} \\
 \Rightarrow r_5 r_4(X) : & \tau_{12354} \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad -q^{-1} \quad -q \quad -1 \quad -1 \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}
 \end{aligned}$$

imply  $q^2 = -1$  by  $a_{32}^{r_5 r_4(X)} = -1$ . Thus  $q \in G'_4$ . In addition, the reflection of  $X$

$$\begin{aligned}
 r_4(X) : & \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \\ \circ \quad \circ \quad \circ \quad \bullet \quad \circ \end{array} \begin{array}{c} q \\ -1 \\ -q^{-1} \\ -1 \end{array} \Rightarrow r_3 r_4(X) : \tau_{12534} \begin{array}{c} q \quad q^{-1} \quad -1 \\ \circ \quad \circ \quad \circ \end{array} \begin{array}{c} -q^{-1} \\ -q^{-2} \\ -q \\ -1 \\ q^{-1} \end{array} q
 \end{aligned}$$

implies  $q^3 = 1$  and  $p \neq 3$  by  $a_{52}^{r_3 r_4(X)} = -1$ . Thus  $q \in G'_3$ , which contradicts  $q \in G'_4$ .

$$\text{If } (a, b, c, d, e) = (0, 0, 1, 1, 2), \text{ it is } A^{r_3(X)} = A_5, A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & -0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix} \text{ and}$$

$$A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & -2 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix}. \text{ Then } qr \neq 1. \text{ The reflection of } X$$

$$\begin{aligned}
 X : & \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \begin{array}{c} q \\ r \\ (qr)^{-1} \\ r^{-1} \end{array}
 \end{aligned}$$

implies  $r = -1$  or  $qr^2 = 1$  by  $a_{53}^{r_4(X)} = -1$ . If  $r = -1$ , the reflection of  $X$  is

$$X : \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad -1 \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_5(X) : \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad -1 \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}$$

which contradicts  $e = 2$ . If  $qr^2 = 1$ , we have either  $a_{25}^{r_3 r_4(X)} = 0$  or  $a_{25}^{r_3 r_4(X)} = -1$  by Definition 3.12 ( $A_{5_2}$ ). When  $a_{25}^{r_3 r_4(X)} = 0$ , we have  $r^3 = 1$ . Then  $q = r \in G'_3$  and  $p \neq 3$ . Hence  $\mathcal{D} = \mathcal{D}_{12,2}^5$ . When  $a_{25}^{r_3 r_4(X)} = -1$ , the reflection of  $X$

$$\begin{aligned}
 r_4(X) : & \begin{array}{c} r^{-2} \quad r^2 \quad r^{-2} \quad r^2 \quad -1 \\ \circ \quad \circ \quad \circ \quad \bullet \quad \circ \end{array} \begin{array}{c} r^{-2} \\ r \\ r \\ -1 \end{array} \Rightarrow r_3 r_4(X) : \tau_{12534} \begin{array}{c} r^{-2} \quad r^2 \quad -1 \\ \circ \quad \circ \quad \circ \end{array} \begin{array}{c} -1 \\ r^3 \\ r^{-1} \\ r^{-2} \\ -1 \\ r^2 \end{array} r^{-2}
 \end{aligned}$$

implies  $r \in G'_5$ . Hence  $\mathcal{D} = \mathcal{D}_{15,6}^5$ ,  $p \neq 5$ .

Step 2.2. Consider that  $q_{11} = q_{33} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{2, 4\}$  and  $j \in \{2, 4, 5\}$ . Then  $(c, d, e) = (0, 0, 1)$ . Hence  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . Let  $q := q_{22}$  and  $r := q_{44}$ . We obtain  $qr \neq 1$  by  $a_{24}^{r_3(X)} = -1$ . The reflections of  $X$

$$\begin{aligned}
 X : \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array} \Rightarrow r_3(X) : \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{(qr)^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array} \\
 \Rightarrow r_2 r_3(X) : \begin{array}{c} q^{-1} \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{(qr)^{-1}} \begin{array}{c} q^{-2}r^{-1} \\ \circ \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array}
 \end{aligned}$$

imply  $q^2r = 1$  by  $a_{41}^{r_2r_3(X)} = 0$ . And  $a_{45}^{r_2r_3(X)} = -1$  gives  $qr^2 = 1$  or  $qr = -1$ . Then  $q = r \in G'_3$  and  $p \neq 3$ . Hence  $\mathcal{D} = \mathcal{D}_{12,13}^5$ .

Step 2.3. Consider that  $q_{33} = q_{55} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 2, 4\}$  and  $j \in \{2, 4\}$ . Then  $(c, d) = (0, 0)$ . Hence  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . This case is symmetric to Step 2.2. Hence  $\mathcal{D} = \tau_{54321}\mathcal{D}_{12,13}^5$ .

Step 2.4. Consider that  $q_{11} = q_{44} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{2, 3\}$  and  $j \in \{2, 3, 5\}$ . Then  $(a, b, e) = (0, 0, 1)$ . By Definition 3.12 one obtains  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . Let  $q := q_{33}$  and  $r := q_{55}$ . Lemma 2.23 gives  $qr \neq 1$ . On the one hand, the reflections of  $X$

$$\begin{aligned}
 X : \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{(qr)^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \\
 \Rightarrow r_3 r_4(X) : \tau_{12534} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{(qr)^{-1}} \begin{array}{c} q^{-2}r^{-1} \\ \circ \end{array} \xrightarrow{qr} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \quad (7)
 \end{aligned}$$

imply  $qr = -1$  or  $q^3r^2 = 1$  by  $a_{52}^{r_3r_4(X)} = -1$ . And we have  $q^2 = -1$  by  $a_{15}^{r_2r_3r_4(X)} = 0$ . Hence  $q \in G'_4$ . On the other hand, the reflections of  $X$

$$\begin{aligned}
 X : \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{(qr)^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \\
 \Rightarrow r_5 r_4(X) : \tau_{12354} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} (qr)^{-1}qr \\ \circ \end{array} \xrightarrow{-1} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array}
 \end{aligned}$$

imply  $qr = -1$  or  $q^2r = 1$  by  $a_{32}^{r_5r_4(X)} = -1$ . Hence one has to have  $qr = -1$  and  $p \neq 2$ . Then from the reflections of  $X$

$$\begin{aligned}
 X : \begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{-q^{-1}} \begin{array}{c} -q^{-1} \\ \circ \end{array} \Rightarrow r_1(X) : \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q} \begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{-q^{-1}} \begin{array}{c} -q^{-1} \\ \circ \end{array} \\
 \Rightarrow r_2 r_1(X) : \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q} \begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{-q^{-1}} \begin{array}{c} -q^{-1} \\ \circ \end{array} \Rightarrow r_3 r_2 r_1(X) : \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q} \begin{array}{c} q^{-1} \\ \circ \end{array} \xrightarrow{-q^{-1}} \begin{array}{c} -q^{-1} \\ \circ \end{array}
 \end{aligned}$$

we obtain  $a_{45}^{r_3 r_2 r_1(X)} = -3$ , which contradicts  $a_{45}^{r_3 r_2 r_1(X)} \neq -3$ .

Step 2.5.. Consider that  $q_{22} = q_{33} = -1$  and  $q_{ii} \widetilde{q_{i,i+1}} - 1 = q_{jj} \widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 4\}$  and  $j \in \{4, 5\}$ . Hence  $(c, d, e) = (0, 0, 1)$ . Definition 3.12 implies  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . Let  $q := q_{11}$ ,  $r^{-1} := \widetilde{q_{23}}$  and  $s := q_{44}$ . By  $A^{r_2(X)} = A_5$ , one obtains  $qr = 1$ . And  $a_{35}^{r_4(X)} = -1$  yields  $rs \neq 1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccccc} r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & s & s^{-1} & s \\ \circ & & \bullet & & \circ & & \circ & & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{ccccccccc} -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & s & s^{-1} & s \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}$$

gives  $r = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.6 ( $a_2$ ), this case is impossible.

Step 2.6. Consider that  $q_{33} = q_{44} = -1$  and  $q_{ii} q'_{i,i+1} - 1 = q_{jj} q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2\}$  and  $j \in \{2, 5\}$ . Then  $e = 1$ . From the definition of good  $A_5$  neighborhood, we have either  $(a, b, c, d) = (0, 0, 1, 1)$  or  $(a, b, c, d) = (1, 1, 0, 0)$ . Let  $q := q_{11}$ ,  $r^{-1} := q'_{34}$  and  $s := q_{55}$ . If

$$(a, b, c, d) = (0, 0, 1, 1), \text{ that is } A^{r_3(X)} = A_5 \text{ and } A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix}. \text{ Hence}$$

$qr = 1$  and  $rs \neq 1$ . The reflection of  $X$

$$X : \begin{array}{ccccccccc} r^{-1} & r & r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & s \\ \circ & & \circ & & \bullet & & \circ & & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccc} r^{-1} & r & -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & s \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.6 ( $c_4$ ), this case is impossible.

If  $(a, b, c, d) = (1, 1, 0, 0)$ , then  $A^{r_4(X)} = A_5$  and

$$A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & -1 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & -1 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix}. \text{ Hence } qr \neq 1 \text{ and } rs = 1. \text{ The reflection of } X$$

$$X : \begin{array}{ccccccccc} q & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & r & r^{-1} \\ \circ & & \circ & & \circ & & \bullet & & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccccc} q & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & r^{-1} & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.6 ( $a_1$ ), this case is impossible.

Step 2.7. Consider that  $q_{22} = q_{44} = -1$  and  $q_{ii} \widetilde{q_{i,i+1}} - 1 = q_{jj} \widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 3\}$  and  $j \in \{3, 5\}$ . Then  $(a, b, e) = (0, 0, 1)$ . Hence in this case  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . Let  $s := q_{11}$ ,  $q := q_{33}$  and  $r := q_{55}$ . Then  $rs \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . And  $A^{r_2(X)} = A_5$  yields  $qr = 1$ . Following the same method as in the proof of Step 2.4, one obtains  $a_{45}^{r_3 r_2 r_1(X)} = -3$ , which is a contradiction.

Step 3. Assume that there are three vertices labeled by  $-1$ . Next we distinguish the following categories: Step 3.1 to Step 3.9.

Step 3.1. Consider that  $q_{11} = q_{22} = q_{33} = -1$ ,  $q_{44} \widetilde{q_{45}} - 1 = 0$  and  $q_{jj} \widetilde{q_{j-1,j}} - 1 = 0$  for  $j \in \{4, 5\}$ . Then  $(c, d, e) = (0, 0, 1)$ . The definition of good  $A_5$  neighborhood implies  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . Let  $q^{-1} := \widetilde{q_{12}}$ ,  $r^{-1} := \widetilde{q_{23}}$  and  $s := q_{44}$ . Then  $a_{24}^{r_3(X)} = -1$  gives  $rs \neq 1$ . And since  $A^{r_2(X)} = A_5$ , we have  $qr = 1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccccc} -1 & r & -1 & r^{-1} & -1 & s^{-1} & s & s^{-1} & s \\ \circ & & \bullet & & \circ & & \circ & & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{ccccccccc} r & r^{-1} & -1 & r & r^{-1} & s^{-1} & s & s^{-1} & s \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}$$



implies  $r = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.6 ( $a_2$ ), this case is impossible.

Step 3.2. Consider that  $q_{33} = q_{44} = q_{55} = -1$ ,  $q_{22}\widetilde{q_{12}} - 1 = 0$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = 0$  for  $i \in \{1, 2\}$ . Definition 3.12 implies either  $(a, b, c, d) = (1, 1, 0, 0)$  or  $(a, b, c, d) = (0, 0, 1, 1)$ . For the same reason as in Step 2.6, one finds that the cases  $(a, b, c, d) = (1, 1, 0, 0)$  and  $(a, b, c, d) = (0, 0, 1, 1)$  are impossible.

Step 3.3. Consider that  $q_{11} = q_{22} = q_{44} = -1$ ,  $q_{33}\widetilde{q_{34}} - 1 = 0$  and  $q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for  $j \in \{3, 5\}$ . Then  $(a, b, e) = (0, 0, 1)$ . By the definition of good  $A_5$  neighborhood, we have  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . Let  $q^{-1} := \widetilde{q_{12}}$ ,  $r := q_{33}$  and  $s := q_{55}$ . Then  $rs \neq 1$ . And we obtain  $qr = 1$  from  $A^{r_2(X)} = A_5$ . On the one hand, the reflections of  $X$

$$\begin{aligned} X : \begin{array}{c} -1 \quad r \quad -1 \quad r^{-1} \quad r \quad r^{-1} \quad -1 \quad s^{-1} \quad s \\ \circ \quad \quad \circ \quad \quad \circ \quad \quad \bullet \quad \quad \circ \end{array} &\Rightarrow r_4(X) : \begin{array}{c} -1 \quad r \quad -1 \quad r^{-1} \quad -1 \quad \begin{array}{c} \circ \quad -1 \\ r \quad \quad s \\ (rs)^{-1} \quad \circ \quad -1 \end{array} \\ \circ \quad \quad \circ \quad \quad \bullet \quad \quad \circ \end{array} \\ &\Rightarrow r_3 r_4(X) : \tau_{12534} \begin{array}{c} \begin{array}{c} (rs)^{-1} \\ r^{-2}s^{-1} \quad \circ \quad rs \\ r \quad r^{-1} \quad r \quad -1 \quad r^{-1} \quad r \end{array} \\ \circ \quad \quad \circ \quad \quad \circ \quad \quad \circ \end{array} \end{aligned}$$

imply  $rs = -1$  or  $r^3 s^2 = 1$  by  $a_{52}^{r_3 r_4(X)} = -1$ . On the other hand, the reflections of  $X$

$$\begin{aligned} X : \begin{array}{c} -1 \quad r \quad -1 \quad r^{-1} \quad r \quad r^{-1} \quad -1 \quad s^{-1} \quad s \\ \circ \quad \quad \circ \quad \quad \circ \quad \quad \bullet \quad \quad \circ \end{array} &\Rightarrow r_4(X) : \begin{array}{c} -1 \quad r \quad -1 \quad r^{-1} \quad -1 \quad \begin{array}{c} \circ \quad -1 \\ r \quad \quad s \\ (rs)^{-1} \quad \circ \quad -1 \end{array} \\ \circ \quad \quad \circ \quad \quad \bullet \quad \quad \circ \end{array} \\ &\Rightarrow r_5 r_4(X) : \tau_{12354} \begin{array}{c} -1 \quad r \quad -1 \quad r^{-1} (rs)^{-1} r s \quad -1 \quad s^{-1} \quad s \\ \circ \quad \quad \circ \quad \quad \circ \quad \quad \circ \quad \quad \circ \quad \quad \circ \end{array} \end{aligned}$$

imply  $rs = -1$  or  $r^2 s = 1$  by  $a_{32}^{r_5 r_4(X)} = -1$ . Then  $rs = -1$  and  $p \neq 2$ . Since  $a_{15}^{r_2 r_3 r_4(X)} = 0$  yields  $rs = 1$ , which contradicts  $rs = -1$ .

Step 3.4. Consider that  $q_{22} = q_{44} = q_{55} = -1$ ,  $q_{33}\widetilde{q_{23}} - 1 = 0$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = 0$  for  $i \in \{1, 3\}$ . Then  $(a, b) = (0, 0)$ . Definition 3.12 implies that either  $(a, b, c, d, e) = (0, 0, 1, 1, 2)$  or  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . Let  $q := q_{11}$ ,  $r := q_{33}$  and  $s^{-1} := \widetilde{q_{45}}$ . If

$$(a, b, c, d, e) = (0, 0, 1, 1, 2), \text{ it is } A^{r_3(X)} = A_5, \quad A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix} \text{ and}$$

$$A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & -2 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix}. \text{ Then } qr = 1 \text{ and } rs \neq 1. \text{ The reflection of } X$$

$$X : \begin{array}{c} r^{-1} \quad r \quad -1 \quad r^{-1} \quad r \quad r^{-1} \quad -1 \quad s^{-1} \quad -1 \\ \circ \quad \quad \circ \quad \quad \circ \quad \quad \bullet \quad \quad \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} r^{-1} \quad r \quad -1 \quad r^{-1} \quad -1 \quad \begin{array}{c} \circ \quad -1 \\ r \quad \quad s \\ (rs)^{-1} \quad \circ \quad s^{-1} \end{array} \\ \circ \quad \quad \circ \quad \quad \bullet \quad \quad \circ \end{array} \quad (8)$$

gives  $s = -1$  or  $rs^2 = 1$  by  $a_{53}^{r_4(X)} = -1$ . Moreover, the reflection of  $X$

$$X : \begin{array}{ccccccc} r^{-1} & r & -1 & r^{-1} & r & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{ccccccc} r^{-1} & r & -1 & r^{-1} & r & r^{-1} & s^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \begin{array}{c} s \\ -1 \end{array}$$

implies  $s \neq -1$  by  $a_{43}^{r_5(X)} = -2$ . Hence  $rs^2 = 1$ . By Definition 3.12 ( $A_{5_2}$ ) one obtains that either  $a_{25}^{r_3r_4(X)} = 0$  or  $a_{25}^{r_3r_4(X)} = -1$ . When  $a_{25}^{r_3r_4(X)} = 0$ , the reflection (8) of  $X$  implies  $r^2s = 1$ . Hence  $r = s \in G'_3$  and  $p \neq 3$ . Then the reflections of  $X$

$$X : \begin{array}{ccccccc} r^{-1} & r & -1 & r^{-1} & r & r^{-1} & -1 \\ \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{ccccccc} -1 & r^{-1} & -1 & r & -1 & r^{-1} & -1 \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ \end{array}$$

$$\Rightarrow r_3r_2(X) : \begin{array}{ccccccc} -1 & r^{-1} & r & r^{-1} & -1 & r & r^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \begin{array}{c} -1 \\ -1 \end{array}$$

give  $(a_{45}^{r_3r_2r_1(X)}, a_{45}^{r_3r_2(X)}) = (-2, -2)$ , which contradicts  $(a_{45}^{r_3r_2r_1(X)}, a_{45}^{r_3r_2(X)}) \neq (-2, -2)$ . When  $a_{25}^{r_3r_4(X)} = -1$ , the reflection of  $X$

$$r_4(X) : \begin{array}{ccccccc} r^{-1} & r & -1 & r^{-1} & -1 & r & -1 \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ \end{array} \begin{array}{c} s \\ (rs)^{-1} \\ s^{-1} \end{array} \Rightarrow r_3r_4(X) : \begin{array}{ccccccc} r^{-1} & r & r^{-1} & r & -1 & r^{-1} & r \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \begin{array}{c} -r^{-1}s^{-2} \\ r^{-2}s^{-1} \\ rs \\ -1 \end{array}$$

implies  $r = -1$  or  $r^3s = 1$ . If  $r = -1$ , we have  $s = -1$  and  $p \neq 2$  by  $a_{15}^{r_2r_3r_4(X)} = 0$ , which is a contradiction. If  $r^3s = 1$ , we obtain  $r \in G'_5$  by  $rs^2 = 1$ . Similarly to the proof in Case  $a_{24}$ , we have  $a_{45}^{r_3r_2r_1(X)} = -3$ , which contradicts  $a_{45}^{r_3r_2r_1(X)} \neq -3$ .

$$\text{If } (a, b, c, d, e) = (0, 0, 1, 1, 1), \text{ then } A^{r_3(X)} = A_5, A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix} \text{ and}$$

$A^{r_5(X)} = A_5$ . Then  $qr = 1$  and  $rs \neq 1$ . The reflection of  $X$

$$X : \begin{array}{ccccccc} r^{-1} & r & -1 & r^{-1} & r & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{ccccccc} r^{-1} & r & -1 & r^{-1} & r & r^{-1} & s^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \begin{array}{c} s \\ -1 \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . By Lemma 4.6 ( $c_1$ ), this case is impossible.

Step 3.5. Consider that  $q_{11} = q_{44} = q_{55} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{2, 3\}$  and  $j \in \{2, 3\}$ . Hence  $(a, b) = (0, 0)$ . Then we have either  $(a, b, c, d, e) = (0, 0, 1, 1, 2)$  or  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . Let  $q := q_{22}$  and  $r^{-1} := \widetilde{q_{45}}$ . If  $(a, b, c, d, e) = (0, 0, 1, 1, 2)$ , then

$$A^{r_3(X)} = A_5, A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix} \text{ and } A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & -2 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix}.$$

Thus  $qr \neq 1$ . The reflection of  $X$

$$X : \begin{array}{ccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccc} -1 & q^{-1} & q & q^{-1} & -1 & q & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \begin{array}{c} q \\ (qr)^{-1} \\ r \\ r^{-1} \end{array} \quad (9)$$

gives  $r = -1$  or  $qr^2 = 1$  by  $a_{53}^{r_4(X)} = -1$ . Furthermore, the reflection of  $X$

$$X : \begin{array}{c} -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \end{array} \Rightarrow r_5(X) : \begin{array}{c} -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad r^{-1} \quad r \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}$$

implies  $r \neq -1$ . Hence  $qr^2 = 1$ . By Definition 3.12 ( $A_{52}$ ), we obtain  $a_{25}^{r_3 r_4(X)} \in \{0, 1\}$ . If  $a_{25}^{r_3 r_4(X)} = 0$ , the reflection (9) of  $X$  shows  $q^2 r = 1$ . Then  $r = q \in G'_3$  and  $p \neq 3$  by  $qr^2 = 1$ . Hence  $\mathcal{D} = \mathcal{D}_{13,3}^5$ . If  $a_{25}^{r_3 r_4(X)} = -1$ , the reflection of  $X$

$$r_4(X) : \begin{array}{c} -1 \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad q \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \end{array} \Rightarrow r_3 r_4(X) : \begin{array}{c} -1 \quad q^{-1} \quad -1 \quad q \quad -1 \quad q^{-1} \quad q \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}$$

implies  $q^3 r = 1$  by  $a_{15}^{r_2 r_3 r_4(X)} = 0$ . Then  $r \in G'_5$  by  $qr^2 = 1$ . Then  $\mathcal{D}$  is

$$\begin{array}{c} -1 \quad r^2 \quad r^{-2} \quad r^2 \quad r^{-2} \quad r^2 \quad -1 \quad r^{-1} \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}$$

which is impossible by Lemma 4.6 ( $b_1$ ).

$$\text{If } (a, b, c, d, e) = (0, 0, 1, 1, 1), \text{ then } A^{r_3(X)} = A_5, A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix} \text{ and}$$

$A^{r_5(X)} = A_5$ . Hence  $qr \neq 1$ . The reflection of  $X$

$$X : \begin{array}{c} -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \end{array} \Rightarrow r_5(X) : \begin{array}{c} -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad r^{-1} \quad r \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . By Lemma 4.6 ( $c_2$ ), this case is impossible.

Step 3.6. Consider that  $q_{11} = q_{33} = q_{44} = -1$ ,  $q_{22}\widetilde{q_{23}} - 1 = 0$  and  $q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for  $j \in \{2, 5\}$ . Then  $e = 1$ . Since  $X$  has a good  $A_5$  neighborhood, we have either  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$  or  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . Let  $q := q_{22}$ ,  $r^{-1} := \widetilde{q_{34}}$  and  $s := q_{55}$ . If  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ , that is

$$A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & -1 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & -1 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix} \text{ and } A^{r_4(X)} = A^{r_5(X)} = A_5. \text{ Then } qr \neq 1 \text{ and } rs = 1.$$

By  $a_{32}^{r_4(X)} = -1$ , we obtain  $r = -1$  and  $p \neq 2$ . By Lemma 4.6 ( $a_4$ ), this case is impossible.

$$\text{If } (a, b, c, d, e) = (0, 0, 1, 1, 1), \text{ that is } A^{r_3(X)} = A_5, A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix}$$

and  $A^{r_5(X)} = A_5$ . Given this condition, we have  $qr = 1$  and  $rs \neq 1$ . And the reflection of  $X$

$$X : \begin{array}{c} -1 \quad r \quad r^{-1} \quad r \quad -1 \quad r^{-1} \quad -1 \quad s^{-1} \quad s \\ \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_3(X) : \begin{array}{c} -1 \quad r \quad -1 \quad r^{-1} \quad -1 \quad r \quad r^{-1} \quad s^{-1} \quad s \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . Lemma 4.6 ( $c_4$ ) shows that this case is impossible.

Step 3.7. Consider that  $q_{22} = q_{33} = q_{55} = -1$ ,  $q_{44}\widetilde{q_{34}} - 1 = 0$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = 0$  for  $i \in \{1, 4\}$ . Then in this case  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . Let  $q := q_{11}$ ,  $r^{-1} := \widetilde{q_{23}}$  and  $s := q_{44}$ . By  $a_{24}^{r_3(X)} = -1$  one obtains  $rs \neq 1$ . The reflection of  $X$

$$X : \begin{array}{ccccccc} q & q^{-1} & -1 & r^{-1} & -1 & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{ccccccc} -1 & q & -1 & r & r^{-1} & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$  and  $r = -1$  by  $a_{34}^{r_2(X)} = -1$ . Then  $q = r = -1$  and  $p \neq 2$ . Thus by Lemma 4.6 ( $a_3$ ), this case is impossible.

Step 3.8. Consider that  $q_{11} = q_{33} = q_{55} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{2, 4\}$  and  $j \in \{2, 4\}$ . In this case  $(c, d) = (0, 0)$ . Hence  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . Let  $q := q_{22}$  and  $r := q_{44}$ . We obtain  $qr \neq 1$  by  $a_{24}^{r_3(X)} = -1$ . The reflection of  $X$

$$X : \begin{array}{ccccccc} -1 & q^{-1} & q & q^{-1} & -1 & r^{-1} & r & r^{-1} & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccc} -1 & q^{-1} & -1 & (qr)^{-1} & -1 & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $q^2r = 1$  by  $a_{41}^{r_2r_3(X)} = 0$ . And  $a_{45}^{r_2r_3(X)} = -1$  gives  $qr = -1$  or  $qr^2 = 1$ . Then we obtain  $q = r \in G'_3$  and  $p \neq 3$ . Hence we have  $a_{32}^{r_4r_5(X)} = -2$  by the reflections of  $X$

$$X : \begin{array}{ccccccc} -1 & q^{-1} & q & q^{-1} & -1 & r^{-1} & r & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{ccccccc} -1 & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & r & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array}$$

$$\Rightarrow r_4r_5(X) : \begin{array}{ccccccc} -1 & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & r^{-1} & r \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

And the reflections of  $X$

$$X : \begin{array}{ccccccc} -1 & q^{-1} & q & q^{-1} & -1 & r^{-1} & r & r^{-1} & -1 \\ \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_1(X) : \begin{array}{ccccccc} -1 & q & -1 & q^{-1} & -1 & r^{-1} & r & r^{-1} & -1 \\ \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

$$\Rightarrow r_2r_1(X) : \begin{array}{ccccccc} q & q^{-1} & -1 & q & q^{-1} & r^{-1} & r & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

imply  $a_{34}^{r_2r_1(X)} = -2$ , which contradicts  $(a_{32}^{r_4r_5(X)}, a_{34}^{r_2r_1(X)}) \neq (-2, -2)$ .

Step 3.9. Consider that  $q_{22} = q_{33} = q_{44} = -1$ ,  $q_{11}\widetilde{q_{12}} - 1 = 0$  and  $q_{55}\widetilde{q_{45}} - 1 = 0$ . We have  $e = 1$ . According to Definition 3.12, the tuple  $(a, b, c, d, e)$  must be either  $(1, 1, 0, 0, 1)$  or  $(0, 0, 1, 1, 1)$ . Let  $q := q_{11}$ ,  $r^{-1} := \widetilde{q_{23}}$ ,  $s^{-1} := \widetilde{q_{34}}$  and  $t := q_{55}$ . If  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ , that is

$$A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & -1 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & -1 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix} \text{ and } A^{r_4(X)} = A_5. \text{ Thus } qr = st = 1 \text{ and } rs \neq 1. \text{ The}$$

reflection of  $X$

$$X : \begin{array}{ccccccc} r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & -1 & s & s^{-1} \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{ccccccc} -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & -1 & s & s^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . And the reflection of  $X$

$$X : \begin{array}{ccccccc} -1 & -1 & -1 & -1 & -1 & s^{-1} & -1 & s & s^{-1} \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccc} -1 & -1 & -1 & -1 & s^{-1} & s & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . Hence this case is a contradiction to  $rs \neq 1$ .

$$\text{If } (a, b, c, d, e) = (0, 0, 1, 1, 1), \text{ then } A^{r_3(X)} = A_5 \text{ and } A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix}.$$

Hence we have  $qr = rs = 1$  and  $st \neq 1$ . From  $a_{45}^{r_3(X)} = -1$ , it follows that  $s = -1$  and  $p \neq 2$ . By Lemma 4.6 ( $c_4$ ), this case is impossible.

Step 4. Assume that there are four vertices labeled by  $-1$ . Now we distinguish the following five steps.

Step 4.1. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = -1$  and  $q_{55}\widetilde{q}_{45} - 1 = 0$ . Then  $e = 1$ . Hence, in this case either  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$  or  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . The same argument as in Step 3.9 shows that both  $(a, b, c, d) = (1, 1, 0, 0, 1)$  and  $(a, b, c, d) = (0, 0, 1, 1, 1)$  are impossible.

Step 4.2. Consider that  $q_{22} = q_{33} = q_{44} = q_{55} = -1$  and  $q_{11}\widetilde{q}_{12} - 1 = 0$ . Since  $X$  has a good  $A_5$  neighborhood, we have either  $(a, b, c, d) = (1, 1, 0, 0)$  or  $(a, b, c, d) = (0, 0, 1, 1)$ . Let  $q := q_{11}$ ,  $r^{-1} := \widetilde{q}_{23}$ ,  $s^{-1} := \widetilde{q}_{34}$  and  $t^{-1} := \widetilde{q}_{45}$ . If  $(a, b, c, d) = (1, 1, 0, 0)$ . Applying the same directional reflection as  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$  of Step 3.9 for the same reason leads to a contradiction.

$$\text{If } (a, b, c, d) = (0, 0, 1, 1), \text{ that is } A^{r_3(X)} = A_5 \text{ and } A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix}.$$

Hence  $qr = rs = 1$  and  $st \neq 1$ . What's more, the reflection of  $X$

$$X : \begin{array}{ccccccccc} s & s^{-1} & -1 & s & -1 & s^{-1} & -1 & t^{-1} & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccc} s & s^{-1} & s & s^{-1} & -1 & s & s^{-1} & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.6 ( $b_3$ ), this case is impossible.

Step 4.3. Consider that  $q_{11} = q_{22} = q_{33} = q_{55} = -1$ ,  $q_{44}\widetilde{q}_{45} - 1 = 0$  and  $q_{44}\widetilde{q}_{34} - 1 = 0$ . Then  $(c, d) = (0, 0)$ . Hence in this case  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . Let  $q^{-1} := \widetilde{q}_{12}$ ,  $r^{-1} := \widetilde{q}_{23}$  and  $s := q_{44}$ . Then one obtains  $rs \neq 1$  by  $a_{24}^{r_3(X)} = -1$ . The reflection of  $X$

$$X : \begin{array}{ccccccccc} -1 & q^{-1} & -1 & r^{-1} & -1 & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{ccccccccc} q^{-1} & q & -1 & r & r^{-1} & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$  and  $r = -1$  by  $a_{34}^{r_2(X)} = -1$ . Hence  $q = r = -1$  and  $p \neq 2$ . By Lemma 4.6 ( $a_3$ ), this case is impossible.

Step 4.4. Consider that  $q_{11} = q_{33} = q_{44} = q_{55} = -1$ ,  $q_{22}\widetilde{q}_{23} - 1 = 0$  and  $q_{22}\widetilde{q}_{12} - 1 = 0$ . From Definition 3.12, the tuple  $(a, b, c, d)$  can only be  $(1, 1, 0, 0)$  or  $(0, 0, 1, 1)$ . Let  $q := q_{22}$ ,  $r^{-1} := \widetilde{q}_{34}$  and  $s^{-1} := \widetilde{q}_{45}$ . If  $(a, b, c, d) = (1, 1, 0, 0)$ , that is

$$A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & -1 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & -1 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix} \text{ and } A^{r_4(X)} = A_5. \text{ Then } qr \neq 1 \text{ and } rs = 1. \text{ The}$$

reflection of  $X$

$$X : \begin{array}{c} -1 & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & r & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} -1 & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & r^{-1} & r \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.6 ( $a_4$ ), this case is impossible.

$$\text{If } (a, b, c, d) = (0, 0, 1, 1), \text{ then } A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix} \text{ and } A^{r_3(X)} = A_5. \text{ In}$$

this case  $qr = 1$  and  $rs \neq 1$ . The reflection of  $X$

$$X : \begin{array}{c} -1 & r & r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{c} -1 & r & -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.6 ( $b_3$ ), this case is impossible.

Step 4.5. Consider that  $q_{11} = q_{22} = q_{44} = q_{55} = -1$ ,  $q_{33}\widetilde{q}_{34} - 1 = 0$  and  $q_{33}\widetilde{q}_{23} - 1 = 0$ . Then  $(a, b) = (0, 0)$ . Thus, either  $(a, b, c, d, e) = (0, 0, 1, 1, 2)$  or  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . Let  $q^{-1} := \widetilde{q}_{12}$ ,  $r := q_{33}$  and  $s^{-1} := \widetilde{q}_{45}$ . If  $(a, b, c, d, e) = (0, 0, 1, 1, 2)$ , that is  $A^{r_3(X)} = A_5$ ,

$$A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix} \text{ and } A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & -2 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix}. \text{ Then } qr = 1$$

and  $rs \neq 1$ . On the one hand, the reflection of  $X$

$$X : \begin{array}{c} -1 & r & -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} -1 & r & -1 & r^{-1} & -1 & r & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \quad (10)$$

gives  $s = -1$  or  $rs^2 = 1$  by  $a_{53}^{r_4(X)} = -1$ . On the other hand, the reflection of  $X$

$$X : \begin{array}{c} -1 & r & -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{c} -1 & r & -1 & r^{-1} & r & r^{-1} & s^{-1} & s & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s \neq -1$ . Hence we obtain  $rs^2 = 1$ . From Definition 3.12 ( $A_{52}$ ),  $a_{25}^{r_3r_4(X)} \in \{0, 1\}$ . If  $a_{25}^{r_3r_4(X)} = 0$ , the reflection (10) of  $X$  implies  $r^2s = 1$ . Then  $r = s \in G_3'$  and  $p \neq 3$ . Hence  $\mathcal{D} = \mathcal{D}_{13,5}^5$ . If  $a_{25}^{r_3r_4(X)} = -1$ , the reflection of  $X$

$$r_4(X) : \begin{array}{c} -1 & r & -1 & r^{-1} & -1 & r & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3r_4(X) : \begin{array}{c} -1 & r & -1 & r^{-1} & -1 & r & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  or  $r^3s = 1$ . If  $r = -1$ , we have  $s = -1$  and  $p \neq 2$  by  $a_{15}^{r_2r_3r_4(X)} = 0$ , which is a contradiction. If  $r^3s = 1$ , we obtain  $s \in G_5'$  by  $rs^2 = 1$ . Then  $\mathcal{D}$  is

$$\begin{array}{c} -1 & s^{-2} & -1 & s^2 & s^{-2} & s^2 & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$


By Lemma 4.6 ( $b_2$ ), this case is impossible.

$$\text{If } (a, b, c, d, e) = (0, 0, 1, 1, 1), \text{ that is } A^{r_3(X)} = A_5, A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix}$$

and  $A^{r_5(X)} = A_5$ . Then  $qr = 1$  and  $rs \neq 1$ . Since  $a_{43}^{r_5(X)} = -1$ , we obtain  $s = -1$  and  $p \neq 2$ . By Lemma 4.6 ( $c_3$ ), this case is impossible.

Step 5. Assume that there are five vertices labeled by  $-1$ . Then that is  $q_{11} = q_{22} = q_{33} = q_{44} = q_{55} = -1$ . By the same reasoning used in Step 4.2, this case is impossible.  $\square$

**Lemma 4.8.** *If  $\mathcal{D}_{\chi, E}$  is of the form*




*and has a good  $A_6$  neighborhood, then  $q_{44} = -1$  and the following are impossible.*

- ( $a_1$ )  $\begin{array}{cccccccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & -1 & -1 & -1 & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- ( $a_2$ )  $\begin{array}{cccccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & -1 & -1 & -1 & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- ( $a_3$ )  $\begin{array}{cccccccccccc} q^{-1} & q & -1 & q^{-1} & q & q^{-1} & -1 & -1 & -1 & -1 & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- ( $a_4$ )  $\begin{array}{cccccccccccc} -1 & q & -1 & q^{-1} & q & q^{-1} & -1 & -1 & -1 & -1 & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- ( $a_5$ )  $\begin{array}{cccccccccccc} q & q^{-1} & -1 & q & -1 & q^{-1} & -1 & -1 & -1 & -1 & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- ( $b_1$ )  $\begin{array}{cccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & q^{-1} & -1 & -1 & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \in G'_4, p \neq 2$
- ( $b_2$ )  $\begin{array}{cccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & q^{-1} & q & q^{-1} & q \\ \circ & & \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- ( $b_3$ )  $\begin{array}{cccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & q^{-1} & q & -1 & q \\ \circ & & \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- ( $b_4$ )  $\begin{array}{cccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & q^{-1} & q & q^{-1} & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- ( $b_5$ )  $\begin{array}{cccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & q^{-1} & -1 & q & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- ( $b_6$ )  $\begin{array}{cccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & q & -1 & q^{-1} & q \\ \circ & & \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$


*Proof.* If  $q_{44} \neq -1$ , then  $A^{r_4(X)} = A_6$ , which contradicts Definition 3.13. We proceed by considering the following cases.

(i) If  $X$  is of type  $\begin{array}{ccccccccccc} q & q^{-1} & -1 & q & -1 & q^{-1} & -1 & -1 & -1 & -1 & -1 \end{array}$  where  $q \neq -1$ , then one has




and obtains  $q = -1$  and  $p \neq 2$  by  $a_{36}^{r_5 r_4} = 0$ , which is a contradiction to  $q \neq -1$ . Hence the case  $(a_1)$  is impossible.

(ii) If  $X$  is of type  $\begin{array}{ccccccccccc} q^{-1} & q & -1 & q^{-1} & q & q^{-1} & -1 & -1 & -1 & -1 & -1 \end{array}$  where  $q \neq -1$ . We have



Then  $q = 1$  by  $a_{36}^{r_5 r_4} = 0$ , which contradicts  $q \neq 1$ . Hence the case  $(a_2)$  is impossible. An analogous argument, based on similar reflections and conditions, applies to cases  $(a_3)$ ,  $(a_4)$  and  $(a_5)$ . Hence, the cases  $(a_3)$ ,  $(a_4)$  and  $(a_5)$  are impossible.

(iii) If  $X$  is of type  $\begin{array}{ccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & q^{-1} & -1 & -1 & -1 \end{array}$  where  $q \in G'_4$ , then we have



and obtain  $q = 1$  by  $a_{25}^{r_3 r_4(X)} = 0$ , which contradicts  $q \neq 1$ . Hence the case  $(b_1)$  is impossible. In view of the similar reflections and conditions present in cases  $(b_2)$ – $(b_6)$ , the same reasoning as in case  $(b_1)$  shows that these cases are likewise impossible.

□

**Theorem 4.9.** *If  $\Delta^{[M]}$  is a finite set of roots of sporadic finite Cartan graphs of rank 6 then all possible Dynkin diagrams of  $V$  appear in rows 16 to 19 of Table 1.*

*Proof.* Theorem 3.15 implies that either  $C(M)$  is standard of type  $E_6$  or there exists a point  $X$  such that  $A^X$  has a good  $A_6$  neighborhood.

Case  $e$ . Assume that  $C(M)$  is standard of type  $E_6$ . Since  $A^X = E_6$ , we have  $(2)_{q_{22}}(q_{22}\widetilde{q_{24}} - 1) = (2)_{q_{ii}}(q_{ii}\widetilde{q_{i,i+1}} - 1) = (2)_{q_{jj}}(q_{jj}\widetilde{q_{j-1,j}} - 1) = 0$  for all  $i \in \{1, 3, 4, 5\}$  and  $j \in \{2, 4, 5, 6\}$ . Let  $q := \widetilde{q_{12}}$ . If  $q \neq -1$ , then  $q^2 \neq 1$ . By Lemma 2.23, we obtain  $q_{ii} \neq -1$  for all  $i \in \{1, 2, 3, 4, 5, 6\}$  and  $q_{22}\widetilde{q_{24}} = q_{ii}\widetilde{q_{i,i+1}} = q_{jj}\widetilde{q_{j-1,j}} = 1$  for all  $i \in \{1, 3, 4, 5\}$  and  $j \in \{2, 4, 5, 6\}$ . Otherwise, if any vertex is  $-1$ , then  $q^2 = 1$  by  $A^X = E_6$ , which is a contradiction. Hence  $\mathcal{D} = \mathcal{D}_{16,1}^6$ . If  $q = -1$ , then  $q_{ii} = -1$  for all  $i \in \{1, 2, 3, 4, 5, 6\}$  by  $A^X = E_6$ . Hence  $\mathcal{D} = \mathcal{D}_{16',1}^6$ .

Case  $a$ . Assume that  $X$  has a good  $A_6$  neighborhood. Then  $A^X = A_6$  implies  $(2)_{q_{ii}}(q_{ii}\widetilde{q_{i,i+1}} - 1) = (2)_{q_{jj}}(q_{jj}\widetilde{q_{j-1,j}} - 1) = 0$ , for all  $i \in \{1, 2, 3, 4, 5\}$  and  $j \in \{2, 3, 4, 5, 6\}$ . From Lemma 4.8,



we know  $q_{44} = -1$  for all cases. Depending on the number of vertices that satisfy  $q_{ii} = -1$ , we divide the cases into the following categories: Step 1 to Step 6.

Step 1. Assume that there is one vertex labeled by  $-1$ . Since Lemma 4.8 implies  $q_{44} = -1$ , there is only one case:  $q_{44} = -1$  and  $\widetilde{q_{ii}q_{i,i+1} - 1} = \widetilde{q_{jj}q_{j-1,j} - 1} = 0$  for all  $i \in \{1, 2, 3, 5\}$  and  $j \in \{2, 3, 5, 6\}$ . Hence  $a = 0$ . Let  $q := q_{11}$  and  $r := q_{55}$ . We obtain  $qr \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . The reflections of  $X$

$$\begin{aligned}
 X : & \quad \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad r \quad r^{-1} \quad r \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} \begin{array}{c} -1 \\ \swarrow q \quad \searrow r \\ \circ \quad \circ \\ \nwarrow (qr)^{-1} \quad \nearrow -1 \\ \circ \quad \circ \end{array} \\ q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad r \end{array} \\
 \Rightarrow r_5 r_4(X) : & \quad \begin{array}{c} \begin{array}{c} r \\ \swarrow \quad \searrow \\ \circ \quad \circ \\ \nwarrow r^{-1} \quad \nearrow r \end{array} \\ \tau_{123564} \quad q \quad q^{-1} \quad q \quad q^{-1} (qr)^{-1} \quad qr \quad -1 \end{array}
 \end{aligned}$$

imply  $rq = -1$  or  $q^2 r = 1$  by  $a_{32}^{r_5 r_4(X)} = -1$ . And  $a_{36}^{r_5 r_4(X)} = 0$  gives  $qr^2 = 1$ . Then we obtain  $q = r \in G'_3$ . Hence  $\mathcal{D} = \mathcal{D}_{17,1}^6$  and  $p \neq 3$ .

Step 2. Assume that there are two vertices labeled by  $-1$ . By  $q_{44} = -1$ , we distinguish the following five steps.

Step 2.1. Consider that  $q_{11} = q_{44} = -1$  and  $\widetilde{q_{ii}q_{i,i+1} - 1} = \widetilde{q_{jj}q_{j-1,j} - 1} = 0$  for all  $i \in \{2, 3, 5\}$  and  $j \in \{2, 3, 5, 6\}$ . Hence  $a = 0$ . Let  $q := q_{22}$  and  $r := q_{55}$ . Then we have  $qr \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . The reflection of  $X$

$$\begin{aligned}
 X : & \quad \begin{array}{c} -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad r \quad r^{-1} \quad r \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} \begin{array}{c} -1 \\ \swarrow q \quad \searrow r \\ \circ \quad \circ \\ \nwarrow (qr)^{-1} \quad \nearrow -1 \\ \circ \quad \circ \end{array} \\ -1 \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad r \end{array}
 \end{aligned}$$

implies  $r = q^2$  by  $a_{25}^{r_3 r_4(X)} = 0$  and  $q = r^2$  by  $a_{36}^{r_5 r_4(X)} = 0$ . Then  $r = q \in G'_3$ . Hence  $\mathcal{D} = \mathcal{D}_{18,19}^6$  and  $p \neq 3$ .

Step 2.2. Consider that  $q_{22} = q_{44} = -1$  and  $\widetilde{q_{ii}q_{i,i+1} - 1} = \widetilde{q_{jj}q_{j-1,j} - 1} = 0$  for all  $i \in \{1, 3, 5\}$  and  $j \in \{3, 5, 6\}$ . Hence  $a = 0$ . Let  $q := q_{11}$ ,  $r := q_{33}$  and  $s := q_{55}$ . Then we obtain  $qr = 1$  and  $rs \neq 1$ . The reflection of  $X$

$$\begin{aligned}
 X : & \quad \begin{array}{c} q \quad q^{-1} \quad -1 \quad q \quad q^{-1} \quad q \quad -1 \quad s^{-1} \quad s \quad s^{-1} \quad s \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} \begin{array}{c} -1 \\ \swarrow q^{-1} \quad \searrow s \\ \circ \quad \circ \\ \nwarrow qs^{-1} \quad \nearrow -1 \\ \circ \quad \circ \end{array} \\ q \quad q^{-1} \quad -1 \quad q \quad -1 \quad s^{-1} \quad s \end{array}
 \end{aligned}$$

implies  $s = q^2$  by  $a_{25}^{r_3 r_4(X)} = 0$  and  $q = s^2$  by  $a_{36}^{r_5 r_4(X)} = 0$ . Hence  $q \in G'_3$  and  $p \neq 3$ . Then the reflections of  $X$

$$\begin{aligned}
 X : & \quad \begin{array}{c} q \quad q^{-1} \quad -1 \quad q \quad q^{-1} \quad q \quad -1 \quad s^{-1} \quad s \quad s^{-1} \quad s \\ \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_2(X) : \begin{array}{c} -1 \quad q \quad -1 \quad q^{-1} \quad -1 \quad q \quad -1 \quad s^{-1} \quad s \quad s^{-1} \quad s \\ \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \\
 \Rightarrow r_3 r_2(X) : & \quad \begin{array}{c} -1 \quad q \quad q^{-1} \quad q \quad -1 \quad q^{-1} \quad q \quad s^{-1} \quad s \quad s^{-1} \quad s \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}
 \end{aligned}$$

imply  $(a_{45}^{r_3 r_2 r_1(X)}, a_{45}^{r_3 r_2(X)}) = (-2, -2)$ , which contradicts  $(a_{45}^{r_3 r_2 r_1(X)}, a_{45}^{r_3 r_2(X)}) \neq (-2, -2)$ .

Step 2.3. Consider that  $q_{33} = q_{44} = -1$  and  $q_{ii} \widetilde{q_{i,i+1}} - 1 = q_{jj} \widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 2, 5\}$  and  $j \in \{2, 5, 6\}$ . Hence  $a = 0$ . Let  $q := q_{11}$ ,  $r^{-1} := \widetilde{q_{34}}$ ,  $s := q_{55}$ . By Lemma 2.23, we have  $qr = 1$  and  $rs \neq 1$ . Then the reflection of  $X$

$$X : \begin{array}{cccccccc} r^{-1} & r & r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & s & s^{-1} & s \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{cccccccc} r^{-1} & r & -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & s & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.8 (b<sub>2</sub>), this case is impossible.

Step 2.4. Consider that  $q_{44} = q_{55} = -1$  and  $q_{ii} \widetilde{q_{i,i+1}} - 1 = q_{jj} \widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 2, 3\}$  and  $j \in \{2, 3, 6\}$ . Let  $q := q_{11}$ ,  $r^{-1} := \widetilde{q_{45}}$  and  $s := q_{66}$ . Since  $X$  has a good  $A_6$  neighborhood, we have  $qr \neq 1$ . And we obtain  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . The reflection of  $X$

$$X : \begin{array}{cccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & -1 & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} q & q^{-1} & q & q^{-1} & -1 & -1 & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $q^2 = -1$  by  $a_{25}^{r_3 r_4(X)} = 0$  and  $qs = -1$  by  $a_{36}^{r_5 r_4(X)} = 0$ . Hence  $q = s \in G'_4$  and  $p \neq 2$ . Then  $\mathcal{D} = \mathcal{D}_{19,1}^6$  and  $p \neq 2$ .

Step 2.5. Consider that  $q_{44} = q_{66} = -1$  and  $q_{ii} \widetilde{q_{i,i+1}} - 1 = q_{jj} \widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 2, 3, 5\}$  and  $j \in \{2, 3, 5\}$ . Then  $a = 0$ . Let  $q := q_{11}$  and  $r := q_{55}$ . Then  $qr \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . Hence we obtain  $r = q^{-2}$  by  $a_{25}^{r_3 r_4(X)} = 0$  and  $q = r^{-2}$  by  $a_{36}^{r_5 r_4(X)} = -1$ . Thus  $r = q \in G'_3$  and  $p \neq 3$ . Then the reflections of  $X$

$$X : \begin{array}{cccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & q^{-1} & q & q^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_6(X) : \begin{array}{cccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & q^{-1} & -1 & q & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ \end{array}$$

$$\Rightarrow r_5 r_6(X) : \begin{array}{cccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & q^{-1} & q & -1 & q^{-1} & q \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

imply  $q = -1$  and  $p \neq 2$  by  $a_{43}^{r_5 r_6(X)} = -1$ , which contradicts  $q \in G'_3$ .

Step 3. Assume that there are three vertices labeled by  $-1$ . By  $q_{44} = -1$ , we distinguish the following eight steps.

Step 3.1. Consider that  $q_{11} = q_{22} = q_{44} = -1$  and  $q_{ii} \widetilde{q_{i,i+1}} - 1 = q_{jj} \widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{3, 5\}$  and  $j \in \{3, 5, 6\}$ . Then  $a = 0$ . Let  $q^{-1} := \widetilde{q_{12}}$ ,  $r := q_{33}$  and  $s := q_{55}$ . We obtain  $rs \neq 1$  by  $a_{35}^{r_4(X)} = -1$  and  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$ . Then the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & s & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & r & -1 & r^{-1} & -1 & r & s & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

gives  $s = r^{-2}$  by  $a_{25}^{r_3 r_4(X)} = 0$  and  $r = s^{-2}$  by  $a_{36}^{r_5 r_4(X)} = 0$ . Hence  $r = s \in G'_3$  and  $p \neq 3$ . Then  $\mathcal{D} = \mathcal{D}_{18,16}^6$ .

Step 3.2. Consider that  $q_{11} = q_{33} = q_{44} = -1$  and  $q_{ii} \widetilde{q_{i,i+1}} - 1 = q_{jj} \widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{2, 5\}$  and  $j \in \{2, 5, 6\}$ . Hence  $a = 0$ . By Definition 3.13 and Step 2.3, this case is impossible. Similarly, for the symmetric case with  $q_{33} = q_{44} = q_{66} = -1$  and

$q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 2, 5\}$  and  $j \in \{2, 5\}$ , by Lemma 4.10 ( $b_4$ ) one obtains that this case is impossible.

Step 3.3. Consider that  $q_{11} = q_{44} = q_{55} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{2, 3\}$  and  $j \in \{2, 3, 6\}$ . By Definition 3.13, we have  $a \in \{0, 1\}$ . Let  $q := q_{22}, r^{-1} := \widetilde{q_{45}}$  and  $s := q_{66}$ . By  $a_{35}^{r_4(X)} = -1$ , we obtain  $qr \neq 1$ . Below we split in two steps.

Step 3.3.1. If  $a = 0$ , then  $a_{46}^{r_5(X)} = 0$  implies  $rs = 1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & r & r^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{cccccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . By Lemma 4.8 ( $a_2$ ), this case is impossible.

Step 3.3.2. If  $a = 1$ , then  $a_{46}^{r_5(X)} = -1$ . Then  $rs \neq 1$ . And the reflection of  $X$

$$X : \begin{array}{ccccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{ccccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . Then the reflections of  $X$

$$X : \begin{array}{ccccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & -1 & -1 & s^{-1} & s \\ \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_1(X) : \begin{array}{ccccccccccc} -1 & q & -1 & q^{-1} & q & q^{-1} & -1 & -1 & -1 & s^{-1} & s \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

$$\Rightarrow r_2r_1(X) : \begin{array}{ccccccccccc} q & q^{-1} & -1 & q & -1 & q^{-1} & -1 & -1 & -1 & s^{-1} & s \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

$$\Rightarrow r_3r_2r_1(X) : \begin{array}{ccccccccccc} q & q^{-1} & q & q^{-1} & -1 & q & q^{-1} & -1 & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

imply  $q = -1$  and  $p \neq 2$  by  $a_{45}^{r_3r_2r_1(X)} = -1$ , which contradicts  $qr \neq 1$ .

Step 3.4. Consider that  $q_{11} = q_{44} = q_{66} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{2, 3, 5\}$  and  $j \in \{2, 3, 5\}$ . Hence  $a = 0$ . Let  $q := q_{22}$  and  $r := q_{55}$ . By  $a_{35}^{r_4(X)} = -1$ , we obtain  $qr \neq 1$ . Then the reflections of  $X$

$$X : \begin{array}{ccccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & r & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ \end{array} \Rightarrow r_6(X) : \begin{array}{ccccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & r & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ \end{array}$$

$$\Rightarrow r_5r_6(X) : \begin{array}{ccccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & r^{-1} & r \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

imply  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5r_6(X)} = -1$ . By Lemma 4.8 ( $a_2$ ), this case is impossible.

Step 3.5. Consider that  $q_{22} = q_{33} = q_{44} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 5\}$  and  $j \in \{5, 6\}$ . Hence  $a = 0$ . From Definition 3.13 and Step 2.3, we know that this case is impossible.

Consider the symmetric case that  $q_{33} = q_{44} = q_{55} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 2\}$  and  $j \in \{2, 6\}$ . Let  $q := q_{11}, r^{-1} := \widetilde{q_{34}}, s^{-1} := \widetilde{q_{45}}$  and  $t := q_{66}$ . In this case we have  $qr = 1$  and  $rs \neq 1$ . The reflection of  $X$

$$X : \begin{array}{ccccccccccc} r^{-1} & r & r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & -1 & t^{-1} & t \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccccc} r^{-1} & r & -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

$$X : \overset{r^{-1}}{\circ} \overset{r}{\text{---}} \overset{r^{-1}}{\circ} \overset{r}{\text{---}} \overset{-1}{\circ} \overset{r^{-1}}{\text{---}} \overset{-1}{\bullet} \overset{s^{-1}}{\text{---}} \overset{-1}{\circ} \overset{t^{-1}}{\text{---}} \overset{t}{\circ} \quad \Rightarrow \quad r_4(X) : \overset{r^{-1}}{\circ} \overset{r}{\text{---}} \overset{r^{-1}}{\circ} \overset{r}{\text{---}} \overset{r^{-1}}{\circ} \overset{(rs)^{-1}}{\text{---}} \overset{-1}{\circ} \overset{s^{-1}}{\text{---}} \overset{-1}{\circ} \overset{t^{-1}}{\text{---}} \overset{t}{\circ}$$

Step 3.6. Consider that  $q_{22} = q_{44} = q_{55} = -1$  and  $q_{ii}\widetilde{q_{i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 3\}$  and  $j \in \{3, 6\}$ . By Definition 3.13, we have either  $a = 0$  or  $a = 1$ . Let  $q := q_{11}$ ,  $r := q_{33}$ ,  $s^{-1} := \widetilde{q_{45}}$  and  $t := q_{66}$ . Then we obtain  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$  and  $rs \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . Below we split in two steps.

$$X : \overset{r^{-1}}{\circ} \overset{r}{\text{---}} \overset{-1}{\circ} \overset{r^{-1}}{\text{---}} \overset{r}{\text{---}} \overset{r^{-1}}{\text{---}} \overset{-1}{\circ} \overset{s^{-1}}{\text{---}} \overset{-1}{\bullet} \overset{s}{\text{---}} \overset{s^{-1}}{\circ} \Rightarrow r_5(X) : \overset{r^{-1}}{\circ} \overset{r}{\text{---}} \overset{-1}{\circ} \overset{r^{-1}}{\text{---}} \overset{r}{\text{---}} \overset{r^{-1}}{\text{---}} \overset{s^{-1}}{\text{---}} \overset{s}{\text{---}} \overset{-1}{\circ} \overset{s^{-1}}{\text{---}} \overset{-1}{\circ}$$
$$X : \overset{r^{-1}}{\circ} \overset{r}{\text{---}} \overset{-1}{\circ} \overset{r^{-1}}{\text{---}} \overset{r}{\text{---}} \overset{r^{-1}}{\text{---}} \overset{-1}{\text{---}} \overset{s^{-1}}{\text{---}} \overset{-1}{\bullet} \overset{t^{-1}}{\text{---}} \overset{t}{\circ} \Rightarrow r_5(X) : \overset{r^{-1}}{\circ} \overset{r}{\text{---}} \overset{-1}{\circ} \overset{r^{-1}}{\text{---}} \overset{r}{\text{---}} \overset{r^{-1}}{\text{---}} \overset{s^{-1}}{\text{---}} \overset{s}{\circ} \overset{t}{\text{---}} \overset{(st)^{-1}}{\circ} \overset{-1}{\text{---}} \overset{-1}{\circ}$$
$$\begin{array}{ccc}
X : \overset{r^{-1}}{\circ} \xrightarrow{r} \overset{-1}{\bullet} \xrightarrow{r^{-1}} \overset{r}{\circ} \xrightarrow{r^{-1}} \overset{-1}{\circ} \xrightarrow{-1} \overset{-1}{\circ} \xrightarrow{t^{-1}} \overset{t}{\circ} & \Rightarrow & r_2(X) : \overset{-1}{\circ} \xrightarrow{r^{-1}} \overset{-1}{\circ} \xrightarrow{r} \overset{-1}{\bullet} \xrightarrow{r^{-1}} \overset{-1}{\circ} \xrightarrow{-1} \overset{-1}{\circ} \xrightarrow{t^{-1}} \overset{t}{\circ} \\
\Rightarrow & & r_3 r_2(X) : \overset{-1}{\circ} \xrightarrow{r^{-1}} \overset{r}{\circ} \xrightarrow{r^{-1}} \overset{-1}{\circ} \xrightarrow{r} \overset{r^{-1}}{\circ} \xrightarrow{-1} \overset{-1}{\circ} \xrightarrow{t^{-1}} \overset{t}{\circ}
\end{array}$$

Step 3.7. Consider that  $q_{22} = q_{44} = q_{66} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 3, 5\}$  and  $j \in \{3, 5\}$ . In this case, we have  $a = 0$ . Let  $q := q_{11}$ ,  $r := q_{33}$  and  $s := q_{55}$ . By Lemma 2.23, we obtain  $qr = 1$  and  $rs \neq 1$ . Then the reflections of  $X$

$$\begin{array}{ccc}
X : \overset{r^{-1}}{\circ} \overset{r}{\text{---}} \overset{-1}{\circ} \overset{r^{-1}}{\text{---}} \overset{r}{\text{---}} \overset{r^{-1}}{\circ} \overset{-1}{\text{---}} \overset{s^{-1}}{\text{---}} \overset{s}{\text{---}} \overset{s^{-1}}{\text{---}} \bullet & \Rightarrow & r_6(X) : \overset{r^{-1}}{\circ} \overset{r}{\text{---}} \overset{-1}{\circ} \overset{r^{-1}}{\text{---}} \overset{r}{\text{---}} \overset{r^{-1}}{\text{---}} \overset{-1}{\circ} \overset{s^{-1}}{\text{---}} \bullet \overset{s}{\text{---}} \overset{-1}{\circ} \\
\\
\Rightarrow & r_5 r_6(X) : \overset{r^{-1}}{\circ} \overset{r}{\text{---}} \overset{-1}{\circ} \overset{r^{-1}}{\text{---}} \overset{r}{\text{---}} \overset{r^{-1}}{\text{---}} \overset{s^{-1}}{\text{---}} \overset{s}{\text{---}} \overset{-1}{\circ} \overset{s^{-1}}{\text{---}} \overset{s}{\text{---}} \circ
\end{array}$$

Step 3.8. Consider that  $q_{44} = q_{55} = q_{66} = -1$  and  $\widetilde{q_{ii}q_{i,i+1}} - 1 = \widetilde{q_{jj}q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 2, 3\}$  and  $j \in \{2, 3\}$ . Then  $a \in \{0, 1\}$ . Let  $q := q_{11}$ ,  $r^{-1} := \widetilde{q_{45}}$ ,  $s^{-1} := \widetilde{q_{56}}$ . By  $a_{35}^{r_4(X)} = -1$ , we obtain  $qr \neq 1$ . Below we split in two steps.

Step 3.8.1. If  $a = 0$ , then  $a_{46}^{r_5(X)} = 0$  implise  $rs = 1$ . In addition, the reflection of  $X$

$$X : \begin{array}{ccccccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{ccccccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & s & s^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . Hence  $r = s = -1$ . By Lemma 4.8 ( $a_1$ ), this case is impossible.

Step 3.8.2. If  $a = 1$ , then  $a_{46}^{r_5(X)} = -1$ . Hence  $rs \neq 1$ . The reflection of  $X$

$$X : \begin{array}{ccccccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{ccccccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & s & s^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{46}^{r_5(X)} = -1$ . Additionally, the reflection of  $X$

$$X : \begin{array}{ccccccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & -1 & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_6(X) : \begin{array}{ccccccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & -1 & s^{-1} & s & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{54}^{r_6(X)} = -1$ , which contradicts  $rs \neq 1$ .

Step 4. Assume that there are four vertices labeled by  $-1$ . By  $q_{44} = -1$ , this will be done in eight steps.

Step 4.1. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = -1$ ,  $q_{55}\widetilde{q_{56}} - 1 = 0$  and  $q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for  $j \in \{5, 6\}$ . Then  $a = 0$ . By Definition 3.13 and Step 2.3, this case is impossible.

Consider the symmetric case that  $q_{33} = q_{44} = q_{55} = q_{66} = -1$ ,  $q_{22}\widetilde{q_{21}} - 1 = 0$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = 0$  for  $i \in \{1, 2\}$ . Let  $q := q_{11}$ ,  $r^{-1} := \widetilde{q_{34}}$ ,  $s^{-1} := \widetilde{q_{45}}$  and  $t^{-1} := \widetilde{q_{56}}$ . It follows from Lemma 2.23 and Definition 3.13 that  $qr = 1$  and  $rs \neq 1$ . The reflection of  $X$

$$X : \begin{array}{ccccccccccc} r^{-1} & r & r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccccc} r^{-1} & r & -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

gives  $r = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & s^{-1} & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_6(X) : \begin{array}{ccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & s^{-1} & t^{-1} & t & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $st = 1$  or  $t = -1$  by  $a_{54}^{r_6(X)} = -1$ . If  $st = 1$ , this case is impossible by Lemma 4.8 ( $b_5$ ). If  $t = -1$ , then  $s \neq -1$  and the reflection of  $X$

$$X : \begin{array}{ccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & s^{-1} & -1 & -1 & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & s^{-1} & -1 & -1 & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s^2 = -1$  by  $a_{53}^{r_4(X)} = -1$ . Hence  $s \in G'_4$ . By Lemma 4.8 ( $b_1$ ), this case is impossible.

Step 4.2. Consider that  $q_{11} = q_{22} = q_{44} = q_{55} = -1$ ,  $q_{33}\widetilde{q_{34}} - 1 = 0$  and  $q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for  $j \in \{3, 6\}$ . Then  $a \in \{0, 1\}$ . Let  $q^{-1} := \widetilde{q_{12}}$ ,  $r := q_{33}$ ,  $s^{-1} := \widetilde{q_{45}}$  and  $t := q_{66}$ . We obtain  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$  and  $rs \neq 1$  by  $a_{35}^{r_4(X)} = -1$ .

Step 4.2.1. If  $a = 0$ . Then  $a_{46}^{r_5(X)} = 0$  yields  $st = 1$ . Hence the reflection of  $X$

$$X : \begin{array}{ccccccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & -1 & s & s^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{ccccccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & s^{-1} & s & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$



Step 4.6. Consider that  $q_{11} = q_{44} = q_{55} = q_{66} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{2, 3\}$  and  $j \in \{2, 3\}$ . By the definition of good  $A_6$  neighborhood, we have  $a \in \{0, 1\}$ . Let  $q := q_{22}$ ,  $r^{-1} := \widetilde{q_{45}}$  and  $s^{-1} := \widetilde{q_{56}}$ . By  $a_{35}^{r_4(X)} = -1$ , we have  $qr \neq 1$ .

Step 4.6.1. If  $a = 0$ , then  $a_{46}^{r_5(X)} = 0$ . Hence  $rs = 1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & r & -1 \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{ccccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & r^{-1} & r \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . By Lemma 4.8 ( $a_2$ ), this case is impossible.

Step 4.6.2. If  $a = 1$ , then  $a_{46}^{r_5(X)} = -1$ . Hence  $rs \neq 1$ . The same argument as in Step 3.8.2 shows that this step is a contradiction.

Step 4.7. Consider that  $q_{22} = q_{33} = q_{44} = q_{55} = -1$ ,  $q_{11}\widetilde{q_{12}} - 1 = 0$  and  $q_{66}\widetilde{q_{56}} - 1 = 0$ . Hence  $a \in \{0, 1\}$ . Let  $q := q_{11}$ ,  $r^{-1} := \widetilde{q_{23}}$ ,  $s^{-1} := \widetilde{q_{34}}$ ,  $t^{-1} := \widetilde{q_{45}}$  and  $u := q_{66}$ . By Lemma 2.23, we obtain  $rs = qr = 1$  and  $st \neq 1$ . Then  $qt \neq 1$ . Below we split in two steps.

Step 4.7.1. If  $a = 0$ , that is  $A^{r_5(X)} = A_6$ . Hence  $tu = 1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccccccc} q & q^{-1} & -1 & q & -1 & q^{-1} & -1 & t^{-1} & -1 & t & t^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{ccccccccccc} q & q^{-1} & -1 & q & -1 & q^{-1} & t^{-1} & t & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . Thus this case is impossible by Lemma 4.8 ( $a_5$ ).

Step 4.7.2. If  $a = 1$ , then  $a_{46}^{r_5(X)} = -1$ . Hence  $tu \neq 1$ . The reflection of  $X$

$$X : \begin{array}{ccccccccccc} q & q^{-1} & -1 & q & -1 & q^{-1} & -1 & t^{-1} & -1 & u^{-1} & u \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{ccccccccccc} q & q^{-1} & -1 & q & -1 & q^{-1} & t^{-1} & t & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . On the other hand, the reflection of  $X$

$$X : \begin{array}{ccccccccccc} q & q^{-1} & -1 & q & -1 & q^{-1} & -1 & -1 & u^{-1} & u \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccccc} q & q^{-1} & q & q^{-1} & -1 & q & q^{-1} & -1 & -1 & u^{-1} & u \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $q = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ , which contradicts  $qt \neq 1$ .

Step 4.8. Consider that  $q_{22} = q_{44} = q_{55} = q_{66} = -1$ ,  $q_{33}\widetilde{q_{32}} - 1 = 0$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = 0$  for  $i \in \{1, 3\}$ . Hence we have either  $a = 0$  or  $a = 1$ . Let  $q := q_{11}$ ,  $r := q_{33}$ ,  $s^{-1} := \widetilde{q_{45}}$  and  $t^{-1} := \widetilde{q_{56}}$ . Then  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$  and  $rs \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . The further discussion is similar to Step 4.6.

Step 5. Assume that there are five vertices labeled by  $-1$ . Since  $q_{44} = -1$ , we consider the following three situations separately.

Step 5.1. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{55} = -1$  and  $q_{66}\widetilde{q_{56}} - 1 = 0$ . Let  $q^{-1} := \widetilde{q_{12}}$ ,  $r^{-1} := \widetilde{q_{23}}$ ,  $s^{-1} := \widetilde{q_{34}}$ ,  $t^{-1} := \widetilde{q_{45}}$  and  $u := q_{66}$ . We have  $rs = 1$  by  $a_{24}^{r_3(X)} = 0$ ,  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$ , and  $st \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccccccc} -1 & s^{-1} & -1 & s & -1 & s^{-1} & -1 & t^{-1} & -1 & u^{-1} & u \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccccc} -1 & s^{-1} & s & s^{-1} & -1 & s & s^{-1} & t^{-1} & -1 & u^{-1} & u \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . Hence  $t \neq -1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & t^{-1} & -1 & u^{-1} & u \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & t^{-1} & -1 & u^{-1} & u \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $tu = 1$  by  $a_{56}^{r_4(X)} = -1$ . By Lemma 4.8 ( $b_6$ ), this case is impossible.

Consider the symmetric case that  $q_{22} = q_{33} = q_{44} = q_{55} = q_{66} = -1$  and  $q_{11}\widetilde{q_{12}} - 1 = 0$ . Let  $q := q_{11}$ ,  $r^{-1} := \widetilde{q_{23}}$ ,  $s^{-1} := \widetilde{q_{34}}$ ,  $t^{-1} := \widetilde{q_{45}}$ , and  $u^{-1} := \widetilde{q_{56}}$ . Then  $rs = 1$ ,  $qr = 1$  and  $st \neq 1$ . Employing an argument analogous to the one above, we find from  $a_{45}^{r_3(X)} = -1$  that  $q = -1$  and  $p \neq 2$ , and from  $a_{56}^{r_4(X)} = -1$  that  $tu = 1$ . By Lemma 4.8 ( $b_5$ ), this case is impossible.

Step 5.2. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{66} = -1$  and  $q_{55}\widetilde{q_{56}} - 1 = q_{55}\widetilde{q_{45}} - 1 = 0$ . Then  $a = 0$ . Let  $q^{-1} := \widetilde{q_{12}}$ ,  $r^{-1} := \widetilde{q_{23}}$ ,  $s^{-1} := \widetilde{q_{34}}$  and  $t := q_{55}$ . By Definition 3.13, we obtain  $qr = rs = 1$  and  $st \neq 1$ . Then the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & q^{-1} & -1 & q & -1 & q^{-1} & -1 & t^{-1} & t & t^{-1} & -1 \end{array} \Rightarrow r_3(X) : \begin{array}{cccccccc} -1 & q^{-1} & q & q^{-1} & -1 & q & q^{-1} & t^{-1} & t & t^{-1} & -1 \end{array}$$

implies  $q = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.8 ( $b_4$ ), this case is impossible.

Consider the symmetric case that  $q_{11} = q_{33} = q_{44} = q_{55} = q_{66} = -1$ , and  $q_{22}\widetilde{q_{23}} - 1 = q_{22}\widetilde{q_{12}} - 1 = 0$ . Let  $q := q_{22}$ ,  $r^{-1} := \widetilde{q_{34}}$ ,  $s^{-1} := \widetilde{q_{45}}$  and  $t^{-1} := \widetilde{q_{56}}$ . We obtain  $qr = 1$  by  $a_{24}^{r_3(X)} = 0$  and  $rs \neq 1$   $a_{35}^{r_4(X)} = -1$ . Using an argument similar to that in Step 5.1, we deduce from  $a_{45}^{r_3(X)} = -1$  that  $q = -1$  and  $p \neq 2$ , and from  $a_{56}^{r_4(X)} = -1$  that  $st = 1$ . By Lemma 4.8 ( $b_5$ ), this case is impossible.

Step 5.3. Consider that  $q_{11} = q_{22} = q_{44} = q_{55} = q_{66} = -1$  and  $q_{33}\widetilde{q_{34}} - 1 = q_{33}\widetilde{q_{23}} - 1 = 0$ . By Definition 3.13, it follows that  $a \in \{0, 1\}$ . Let  $q^{-1} := \widetilde{q_{12}}$ ,  $r := q_{33}$ ,  $s^{-1} := \widetilde{q_{45}}$  and  $t^{-1} := \widetilde{q_{56}}$ . By  $a_{13}^{r_2(X)} = 0$ , we obtain  $qr = 1$  and from  $a_{35}^{r_4(X)} = -1$ , we have  $rs \neq 1$ . The application of the same directional reflection as in Step 4.6 demonstrates the impossibility of this case.

Step 6. Assume that all vertices are labeled by  $-1$ . That is  $q_{11} = q_{22} = q_{33} = q_{44} = q_{55} = q_{66} = -1$ . Let  $q^{-1} := \widetilde{q_{12}}$ ,  $r^{-1} := \widetilde{q_{23}}$ ,  $s^{-1} := \widetilde{q_{34}}$ ,  $t^{-1} := \widetilde{q_{45}}$  and  $u^{-1} := \widetilde{q_{56}}$ . We have  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$ ,  $rs = 1$  by  $a_{24}^{r_3(X)} = 0$  and  $st \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . Then the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & s^{-1} & -1 & t^{-1} & -1 & u^{-1} & -1 \end{array} \Rightarrow r_3(X) : \begin{array}{cccccccc} -1 & s^{-1} & s & s^{-1} & -1 & s & s^{-1} & t^{-1} & -1 & u^{-1} & -1 \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . Hence  $t \neq -1$ . Then the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & t^{-1} & -1 & u^{-1} & -1 \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & t^{-1} & -1 & u^{-1} & -1 \end{array}$$

implies  $tu = 1$  by  $a_{56}^{r_4(X)} = -1$ . By Lemma 4.8 ( $b_5$ ), this case is impossible.  $\square$

**Lemma 4.10.** *If  $\mathcal{D}_{X,E}$  is of the form*

$$\begin{array}{cccccccccccc} q_{11} & r & q_{22} & s & q_{33} & t & q_{44} & u & q_{55} & v & q_{66} & w & q_{77} \\ \circ & & \circ & & \circ & & \circ & & \circ & & \circ & & \circ \end{array}$$

*and has a good  $A_7$  neighborhood, then  $q_{33} = -1$  and the following are impossible.*

$$(a_1) \quad \begin{array}{cccccccc} -1 & q^{-1} & -1 & q & -1 & -1 & -1 & -1 & -1 & -1 & -1 \end{array}, \quad q \neq -1, p \neq 2$$



$$(c_8) \quad \begin{array}{cccccccccccc} & -1 & -1 & -1 & -1 & q^{-1} & q & q^{-1} & -1 & q & q^{-1} & q & q^{-1} \\ \circ & & & & \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$$

*Proof.* If  $q_{33} \neq -1$ , then  $A^{r_3(X)} = A_7$ , which contradicts Definition 3.14. Next consider the following cases.

(i) Assume that  $X$  is of the type  $\begin{array}{cccccccccccc} q_{11} & q^{-1} & -1 & & q & & -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 \end{array}$  where  $q_{11} = -1$  or  $q_{11} = q$ . Since  $X$  has a good  $A_7$  neighborhood, the transformation

$$q_{11} \quad q^{-1} \quad -1 \quad q \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad \Rightarrow \quad q'_{11} \quad q \quad -1 \quad q^{-1} \quad q \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1$$

implies  $q = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ , where

$$q'_{11} = \begin{cases} q^{-1} & \text{if } q_{11} = -1, \\ -1 & \text{if } q_{11} = q. \end{cases}$$

which contradicts  $q \neq -1$ . Hence the cases  $(a_1)$  and  $(a_2)$  are impossible.

(ii) Assume that  $X$  is of the type  $q^{11} q^{-1} q q^{-1} -1 -1 -1 -1 -1 -1 -1$  where  $q_{11} = -1$  or  $q_{11} = q$ . We split in two cases  $q^2 = -1$  and  $q^2 \neq -1$ . If  $q^{-2} = -1$ , the transformations

$$\Rightarrow \tau_{3124567} \begin{array}{c} \circ q'_{11} \\ | q \\ \circ \begin{array}{c} q \quad q^{-1} \quad -1 \quad -q \quad -q^{-1} \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \end{array} \end{array}$$

imply that  $q = 1$  by  $a_{45}^{r_2 r_3(x)} = -1$ , where

$$q'_{11} = \begin{cases} q^{-1} & \text{if } q_{11} = -1, \\ -1 & \text{if } q_{11} = q. \end{cases}$$

which is a contradiction. If  $q^{-2} \neq -1$ , the transformations

$$\begin{array}{c} \begin{array}{c} q_{11} \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \end{array} \Rightarrow \begin{array}{c} \circ -1 \\ | q \\ \bullet \quad \circ \begin{array}{c} -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \end{array} \end{array} \\ \\ \Rightarrow \tau_{3214567} \begin{array}{c} \circ q'_{11} \\ | q \\ \circ \begin{array}{c} q \quad q^{-1} \quad -1 \quad -q \quad -q^{-1} \quad -1 \quad -1 \quad -1 \quad -1 \end{array} \end{array} \end{array}$$

imply that  $q = 1$  by  $a_{45}^{r_2 r_3(x)} = -1$ , where

$$q'_{11} = \begin{cases} q^{-1} & \text{if } q_{11} = -1 \\ -1 & \text{if } q_{11} = q \end{cases}$$



which is also a contradiction. Hence the cases  $(b_1)$  and  $(b_2)$  are impossible.

- (iii) If  $X$  is of the type  $\begin{array}{c} -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad q \end{array}$ , then the analysis splits depending on whether  $q^2 = -1$  or not. If  $q^{-2} = -1$ , then the transformations

$$\begin{array}{c} \begin{array}{c} -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad q \end{array} \Rightarrow \begin{array}{c} \circ -1 \\ | q \\ \bullet \quad \circ \begin{array}{c} -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \end{array} \end{array} \\ \\ \Rightarrow \begin{array}{c} \circ q \\ | q^{-1} \\ \circ \begin{array}{c} -1 \quad -1 \quad -q^{-1} \quad -q \quad -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \end{array} \end{array} \end{array}$$

imply  $q = 1$  by  $a_{21}^{r_4 r_3(x)} = -1$ , which is a contradiction. If  $q^{-2} \neq -1$ , then the transformations

$$\begin{array}{c} \begin{array}{c} -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad q \end{array} \Rightarrow \begin{array}{c} \circ -1 \\ | q \\ \bullet \quad \circ \begin{array}{c} -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \end{array} \end{array}$$



implies that  $q = -1$ ,  $p \neq 2$  or  $q \in G'_3$ ,  $p \neq 3$  by  $a_{21}^{r_4 r_3(X)} = -1$ . Since  $q = -1$  yields  $r = 1$ , it is a contradiction. Rewrite  $X$  as

$$\begin{array}{c} q & q^{-1} & q & q^{-1} & -1 & q^{-1} & q & q^{-1} & q & q^{-1} & q & q^{-1} & q \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

where  $q \in G'_3$ ,  $p \neq 3$ . Hence  $\mathcal{D} = \tau_{7654321}\mathcal{D}_{21,1}^7$  and  $p \neq 3$ .

Step 1.2. If  $rq = -1$ , then the reflection of  $X$

$$r_3(X) : \begin{array}{c} \begin{array}{c} -1 \\ \swarrow \quad \searrow \\ -q^{-1} \quad -q \quad -1 \end{array} \\ \begin{array}{c} \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \end{array} \Rightarrow r_4 r_3(X) : \begin{array}{c} \begin{array}{c} q \\ \swarrow \quad \searrow \\ -1 \quad -1 \end{array} \\ \begin{array}{c} \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \end{array}$$

implies  $q = -1$  and  $p \neq 2$  by  $a_{25}^{r_4 r_3(X)} = 0$ . Rewrite  $X$  as

$$\begin{array}{c} r & r^{-1} & r & r^{-1} & -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

By Lemma 4.10 ( $b_2$ ), this case is impossible.

Step 2. Assume that there are two vertices labeled by  $-1$ . By  $q_{33} = -1$ , we distinguish the following three steps.

Step 2.1. Consider that  $q_{11} = q_{33} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{2, 4, 5, 6\}$  and  $j \in \{2, 4, 5, 6, 7\}$ . Let  $r := q_{22}$  and  $q := q_{44}$ . Then  $rq \neq 1$  by  $a_{24}^{r_3(X)} = -1$ . The reflections of  $X$

$$\begin{array}{c} X : \begin{array}{c} -1 \quad r^{-1} \quad r \quad r^{-1} \quad -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad q \\ \bullet \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_1(X) : \begin{array}{c} -1 \quad r \quad -1 \quad r^{-1} \quad -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad q \\ \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \\ \Rightarrow r_2 r_1(X) : \begin{array}{c} r \quad r^{-1} \quad -1 \quad r \quad r^{-1} \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad q \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \end{array}$$

imply that  $r = -1$  and  $p \neq 2$  by  $a_{34}^{r_2 r_1(X)} = -1$ . By Lemma 4.10 ( $c_1$ ), this case is impossible.

Step 2.2. Consider that  $q_{22} = q_{33} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 4, 5, 6\}$  and  $j \in \{4, 5, 6, 7\}$ . Let  $r := q_{11}$ ,  $q := \widetilde{q_{23}}$  and  $s := q_{44}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $r^{-1}q = 1$  and  $qs^{-1} \neq 1$ . The reflection of  $X$

$$\begin{array}{c} X : \begin{array}{c} q \quad q^{-1} \quad -1 \quad q \quad -1 \quad s^{-1} \quad s \quad s^{-1} \quad s \quad s^{-1} \quad s \quad s^{-1} \quad s \\ \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_2(X) : \begin{array}{c} -1 \quad q \quad -1 \quad q^{-1} \quad q \quad s^{-1} \quad s \quad s^{-1} \quad s \quad s^{-1} \quad s \quad s^{-1} \quad s \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \end{array}$$

implies  $q = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.10 ( $c_1$ ), this case is impossible.

Step 2.3. Consider that  $q_{33} = q_{kk} = -1$  for  $k \in \{4, 5, 6, 7\}$  and  $q_{ii}\widetilde{q'_{i,i+1}} - 1 = q_{jj}\widetilde{q'_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 2, 4, 5, 6\} \setminus \{k\}$  and  $j \in \{2, 4, 5, 6, 7\} \setminus \{k\}$ . Let  $r := q_{11}$ ,  $q := q'_{34}$  and  $s := q_{k-1,k}$ . Then we have  $r^{-1}q \neq 1$  by  $a_{24}^{r_3(X)} = -1$  and  $qs = 1$  by  $A^{r_k(X)} = A_7$ . Hence we obtain  $q = -1$  and  $p \neq 2$  by  $a_{32}^{r_4 \cdots r_k(X)} = -1$ . By Lemma 4.10 ( $b_2$ ), these cases are impossible.

Step 3. Assume that there are three vertices labeled by  $-1$ . By  $q_{33} = -1$ , this will be done in seven steps.

Step 3.1. Consider that  $q_{11} = q_{22} = q_{33} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{4, 5, 6\}$  and  $j \in \{4, 5, 6, 7\}$ . Let  $s := \widetilde{q_{12}}$ ,  $r := \widetilde{q_{23}}$  and  $q := q_{44}$ . We have  $rq^{-1} \neq 1$  by  $a_{24}^{r_3(X)} = -1$  and  $sr = 1$  by  $a_{13}^{r_2(X)} = 0$ . Then the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & r^{-1} & -1 & & r & & -1 & q^{-1} & q & q^{-1} & q & q^{-1} & q & q^{-1} & q \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccc} r^{-1} & r & -1 & r^{-1} & r & q^{-1} & q & q^{-1} & q & q^{-1} & q & q^{-1} & q \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.10 ( $c_1$ ), this case is impossible.

Step 3.2. Consider that  $q_{11} = q_{33} = q_{kk} = -1$  for  $k \in \{4, 5, 6, 7\}$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{2, 4, 5, 6\} \setminus \{k\}$  and  $j \in \{2, 4, 5, 6, 7\} \setminus \{k\}$ . Let  $r := q_{22}$ ,  $s := \widetilde{q_{34}}$ , and  $q := \widetilde{q_{k-1,k}}$ . By  $a_{24}^{r_3(X)} = -1$ , we have  $rs \neq 1$ . And for  $k \neq 7$ ,  $A^{r_k(X)} = A_7$  implies  $\widetilde{q_{k,k+1}} = q^{-1}$ . Then we obtain  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.10 ( $b_1$ ), these cases are impossible.

Step 3.3. Consider that  $q_{22} = q_{33} = q_{44} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 5, 6\}$  and  $j \in \{5, 6, 7\}$ . Let  $q := q_{11}$ ,  $s := \widetilde{q_{23}}$ ,  $t := \widetilde{q_{34}}$  and  $r := q_{55}$ . We have  $r^{-1}s = 1$  by  $a_{13}^{r_2(X)} = 0$ ,  $st \neq 1$  by  $a_{24}^{r_3(X)} = 0$ , and  $tq^{-1} = 1$  by  $a_{35}^{r_4(X)} = 0$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} s & s^{-1} & -1 & & s & & -1 & t & -1 & t^{-1} & t & t^{-1} & t \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} s & s^{-1} & -1 & & s & & t & t^{-1} & -1 & t & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.10 ( $a_2$ ), this case is impossible.

Step 3.4. Consider that  $q_{22} = q_{33} = q_{55} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 4, 6\}$  and  $j \in \{4, 6, 7\}$ . Let  $r := q_{11}$ ,  $s := \widetilde{q_{23}}$ ,  $q := q_{44}$  and  $t := q_{66}$ . We have  $r^{-1}s = 1$  by  $a_{13}^{r_2(X)} = 0$ ,  $sq^{-1} \neq 1$  by  $a_{24}^{r_3(X)} = 0$ , and  $qt = 1$  by  $a_{46}^{r_5(X)} = 0$ . Then the reflection of  $X$

$$X : \begin{array}{cccccccc} s & s^{-1} & -1 & & s & & -1 & q^{-1} & q & q^{-1} & -1 & q & q^{-1} & q & q^{-1} \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccc} -1 & s & -1 & s^{-1} & s & q^{-1} & q & q^{-1} & -1 & q & q^{-1} & q & q^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.10 ( $c_8$ ), this case is impossible.

Step 3.5. Consider that  $q_{22} = q_{33} = q_{66} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 4, 5\}$  and  $j \in \{4, 5, 7\}$ . Let  $r := q_{11}$ ,  $s := \widetilde{q_{23}}$ ,  $q := q_{44}$  and  $t := q_{77}$ . From Definition 3.14 and Lemma 2.23, we have  $r^{-1}s = qt = 1$  and  $sq^{-1} \neq 1$ . Furthermore, by  $a_{34}^{r_2(X)} = -1$ , we obtain  $s = -1$  and  $p \neq 2$ . By Lemma 4.10 ( $c_3$ ), this case is impossible.

Step 3.6. Consider that  $q_{22} = q_{33} = q_{77} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 4, 5, 6\}$  and  $j \in \{4, 5, 6\}$ . Let  $r := q_{11}$ ,  $s := \widetilde{q_{23}}$  and  $q := q_{44}$ . We obtain  $r^{-1}s = 1$  by  $a_{13}^{r_2(X)} = 0$  and  $sq^{-1} \neq 1$  by  $a_{24}^{r_3(X)} = 0$ . As in the preceding step, we obtain  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.10 ( $c_2$ ), this case is impossible.

Step 3.7. Consider that  $q_{33} = q_{i_1,i_1} = q_{i_2,i_2} = -1$  for  $i_1, i_2 \in \{4, 5, 6, 7\}$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 2, 4, 5, 6\} \setminus \{i_1, i_2\}$  and  $j \in \{2, 4, 5, 6, 7\} \setminus \{i_1, i_2\}$ . Let  $s := \widetilde{q_{23}}$ ,  $q := \widetilde{q_{i_1-1,i_1}}$  and  $t := \widetilde{q_{i_2-1,i_2}}$ . By Definition 3.14, we have  $qt = 1$  and  $sq \neq 1$ . And  $q_{77} = t$  if  $i_2 \neq 7$ . Then we obtain  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4 \cdots r_{i_1}} = -1$ . By Lemma 4.10 ( $b_2$ ), these cases are impossible.

Step 4. Assume that there are four vertices labeled by  $-1$ . By  $q_{33} = -1$ , we distinguish the following ten steps.

Step 4.1. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{5, 6\}$  and  $j \in \{5, 6, 7\}$ . Let  $r := \widetilde{q_{12}}$ ,  $s := \widetilde{q_{23}}$ ,  $t := \widetilde{q_{34}}$  and  $q := q_{55}$ . Then we

have  $rs = 1$  by  $a_{13}^{r_2(X)} = 0$ ,  $tq^{-1} = 1$  by  $a_{35}^{r_4(X)} = 0$  and  $st \neq 1$  by  $a_{24}^{r_3(X)} = -1$ . Hence, by  $a_{32}^{r_4(X)} = -1$ , we have  $t = -1$  and  $p \neq 2$ . By Lemma 4.10 ( $a_1$ ), this case is impossible.

Step 4.2. Consider that  $q_{11} = q_{22} = q_{33} = q_{55} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{4, 6\}$  and  $j \in \{4, 6, 7\}$ . Let  $r := \widetilde{q_{12}}$ ,  $s := \widetilde{q_{23}}$ ,  $t := q_{44}$  and  $q := q_{66}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs = tq = 1$  and  $st^{-1} \neq 1$ . Hence, it follows from  $a_{34}^{r_2(X)} = -1$  that  $s = -1$  and  $p \neq 2$ . By Lemma 4.10 ( $c_8$ ), this case is impossible.

Step 4.3. Consider that  $q_{11} = q_{22} = q_{33} = q_{66} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{4, 5\}$  and  $j \in \{4, 5, 7\}$ . Let  $r := \widetilde{q_{12}}$ ,  $s := \widetilde{q_{23}}$ ,  $t := q_{44}$  and  $q := q_{77}$ . By Lemma 2.23 and Definition 3.14, we have  $rs = 1$ ,  $tq = 1$  and  $st^{-1} \neq 1$ . Then the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & t^{-1} & t & t^{-1} & t & t^{-1} & -1 & t & t^{-1} \\ \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccc} s^{-1} & s & -1 & s^{-1} & s & t^{-1} & t & t^{-1} & t & t^{-1} & -1 & t & t^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.10 ( $c_3$ ), this case is impossible.

Step 4.4. Consider that  $q_{11} = q_{22} = q_{33} = q_{77} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{4, 5, 6\}$  and  $j \in \{4, 5, 6\}$ . As in the previous Step 4.3, we conclude that this case is impossible by Lemma 4.10 ( $c_2$ ).

Step 4.5. Consider that  $q_{11} = q_{33} = q_{i_1,i_1} = q_{i_2,i_2} = -1$  for  $i_1, i_2 \in \{4, 5, 6, 7\}$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{2, 4, 5, 6\} \setminus \{i_1, i_2\}$  and  $j \in \{2, 4, 5, 6, 7\} \setminus \{i_1, i_2\}$ . Let  $r := q_{22}$ ,  $s := \widetilde{q_{34}}$  and  $t := \widetilde{q_{i_1,i_1+1}}$ . Then  $rs \neq 1$  and  $st = 1$  by Definition 3.14. And  $q_{i_2,i_2+1} = q$  if  $i_2 \neq 7$ . Hence, we obtain  $s = -1$  and  $q \neq 2$  by  $a_{32}^{r_4 \cdots r_{i_1}(X)} = -1$ . By Lemma 4.10 ( $b_1$ ), these cases are impossible.

Step 4.6. Consider that  $q_{22} = q_{33} = q_{55} = q_{77} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 4, 6\}$  and  $j \in \{4, 6\}$ . Let  $r := q_{11}$ ,  $s := \widetilde{q_{23}}$ ,  $t := q_{44}$  and  $q := q_{66}$ . In this case, we have  $r^{-1}s = tq = 1$  and  $st^{-1} \neq 1$ . The further discussion is similar to Step 4.3. By Lemma 4.10 ( $c_7$ ), this case is impossible.

Step 4.7. Consider that  $q_{22} = q_{33} = q_{44} = q_{kk} = -1$  for  $k \in \{5, 6, 7\}$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 5, 6\} \setminus \{k\}$  and  $j \in \{5, 6, 7\} \setminus \{k\}$ . Let  $r := q_{11}$ ,  $s := \widetilde{q_{23}}$ ,  $t := \widetilde{q_{34}}$ , and  $u := \widetilde{q_{45}}$ . By Definition 3.14, we obtain  $r^{-1}s = tu = 1$  and  $st \neq 1$ . If  $k \neq 7$ , then  $q_{k,k+1} = u$  from  $A^{r_k(X)} = A_7$ . Hence we obtain  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.10 ( $a_2$ ), these cases are impossible.

Step 4.8. Consider that  $q_{22} = q_{33} = q_{55} = q_{66} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 4\}$  and  $j \in \{4, 7\}$ . Let  $r := q_{11}$ ,  $s := \widetilde{q_{23}}$ ,  $t := q_{44}$  and  $q = q_{77}$ . By Definition 3.14, we obtain  $r^{-1}s = t^{-1}u = uq^{-1} = 1$  and  $st^{-1} \neq 1$ . The subsequent discussion is analogous to that in Step 4.3. By Lemma 4.10 ( $c_6$ ), this case is impossible.

Step 4.9. Consider that  $q_{22} = q_{33} = q_{66} = q_{77} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 4, 5\}$  and  $j \in \{4, 5\}$ . Let  $r := q_{11}$ ,  $s := \widetilde{q_{23}}$ ,  $t := q_{44}$  and  $q := \widetilde{q_{67}}$ . By Definition 3.14, we have  $r^{-1}s = t^{-1}q = 1$  and  $st^{-1} \neq 1$ . As in Step 4.3, we obtain  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.10 ( $c_4$ ), this case is impossible.

Step 4.10. Consider that  $q_{33} = q_{i_1,i_1} = q_{i_2,i_2} = q_{i_3,i_3} = -1$  for  $i_1, i_2, i_3 \in \{4, 5, 6, 7\}$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{1, 2, 4, 5, 6\} \setminus \{i_1, i_2, i_3\}$  and  $j \in \{2, 4, 5, 6, 7\} \setminus \{i_1, i_2, i_3\}$ . Let  $r := q_{11}$ ,  $s := \widetilde{q_{34}}$ ,  $t := \widetilde{q_{i_1,i_1+1}}$  and  $q := \widetilde{q_{i_2,i_2+1}}$ . By Lemma 2.23, we obtain that  $st = tq = 1$  and  $r^{-1}s \neq 1$ . And  $q_{77} = q$  if  $i_3 \neq 7$ . Then we obtain  $s = -1$  and  $p \neq 2$  from  $a_{32}^{r_4 \cdots r_{i_1}(X)} = -1$ . By Lemma 4.10 ( $b_2$ ), these cases are impossible.

Step 5. Assume that there are five vertices labeled by  $-1$ . By  $q_{33} = -1$ , the analysis is divided into the following seven steps.

Step 5.1. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{kk} = -1$ ,  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for  $i \in \{5, 6\} \setminus \{k\}$  and  $j \in \{6, 7\} \setminus \{k\}$ . Let  $r := \widetilde{q_{12}}$ ,  $s := \widetilde{q_{23}}$ ,  $t := \widetilde{q_{34}}$ , and  $u := \widetilde{q_{45}}$ . Then  $rs = 1$  by  $a_{13}^{r_2(X)} = 0$ ,  $st \neq 1$  by  $a_{24}^{r_3(X)} = -1$ , and  $tu = 1$  by  $a_{35}^{r_4(X)} = 0$ . And for  $k \neq 7$ ,  $A^{r_k(X)} = A_7$  yields  $\widetilde{q_{k,k+1}} = u^{-1}$ . Hence, we obtain that  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.10 ( $a_1$ ), these cases are impossible.

By Lemma 4.10 ( $b_2$ ), and following an argument similar to the one above, we conclude that the case with  $q_{33} = q_{44} = q_{55} = q_{66} = q_{77} = -1$ ,  $q_{22}\widetilde{q_{12}} - 1 = 0$ , and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = 0$  for  $i \in \{2, 4, 5\}$  is also impossible.

Step 5.2. Consider that  $q_{11} = q_{22} = q_{33} = q_{55} = q_{66} = -1$ ,  $q_{44}\widetilde{q_{45}} - 1 = 0$  and  $q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for  $j \in \{4, 7\}$ . Let  $r := \widetilde{q_{12}}$ ,  $s := \widetilde{q_{23}}$ ,  $t := q_{44}$ ,  $u := \widetilde{q_{56}}$  and  $q := q_{77}$ . By Definition 3.14, we have  $rs = t^{-1}u = uq^{-1} = 1$  and  $st^{-1} \neq 1$ . Then the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & t^{-1} & t & t^{-1} & -1 & t & -1 & t^{-1} & t \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccc} s^{-1} & s & -1 & s^{-1} & s & t^{-1} & t & t^{-1} & -1 & t & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.10 ( $c_6$ ), this case is impossible.

Consider the symmetric case that  $q_{22} = q_{33} = q_{55} = q_{66} = q_{77} = -1$ ,  $q_{44}\widetilde{q_{34}} - 1 = 0$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = 0$  for  $i \in \{1, 4\}$ . Similarly to the proof above, this case is impossible by Lemma 4.10 ( $c_5$ ).

Step 5.3. Consider that  $q_{11} = q_{22} = q_{33} = q_{55} = q_{77} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{4, 6\}$  and  $j \in \{4, 6\}$ . Apply the same reflection conditions as in the preceding step. By Lemma 4.10 ( $c_7$ ), this case is impossible.

Step 5.4. Consider that  $q_{11} = q_{22} = q_{33} = q_{66} = q_{77} = -1$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{4, 5\}$  and  $j \in \{4, 5\}$ . Using the same calculation method as in Step 5.3, we find that this case is impossible by Lemma 4.10 ( $c_4$ ).

Step 5.5. Consider that  $q_{11} = q_{33} = q_{i_1,i_1} = q_{i_2,i_2} = q_{i_3,i_3} = -1$  for  $i_1, i_2, i_3 \in \{4, 5, 6, 7\}$  and  $q_{ii}\widetilde{q_{i+1,i}} - 1 = q_{jj}\widetilde{q_{j-1,j}} - 1 = 0$  for all  $i \in \{2, 4, 5, 6\} \setminus \{i_1, i_2, i_3\}$  and  $j \in \{2, 4, 5, 6, 7\} \setminus \{i_1, i_2, i_3\}$ . Let  $r := q_{22}$ ,  $s := \widetilde{q_{34}}$ ,  $t := \widetilde{q_{45}}$ ,  $u := \widetilde{q_{56}}$  and  $q := q_{77}$ . By Definition 3.14, we have  $st = tu = uq^{-1} = 1$  and  $r^{-1}s \neq 1$ . Then the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & -1 & s & -1 & s^{-1} & -1 & s & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & s & s^{-1} & -1 & s & s^{-1} & s & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.10 ( $b_1$ ), these cases are impossible.

Step 5.6. Consider that  $q_{22} = q_{33} = q_{44} = q_{55} = q_{66} = -1$  and  $q_{11}\widetilde{q_{12}} - 1 = q_{77}\widetilde{q_{67}} - 1 = 0$ . Let  $r := q_{11}$ ,  $s := \widetilde{q_{23}}$ ,  $t := \widetilde{q_{34}}$ ,  $u := \widetilde{q_{45}}$ ,  $v := \widetilde{q_{56}}$  and  $q := q_{77}$ . By Definition 3.14, we have  $r^{-1}s = tu = uv = vq^{-1} = 1$  and  $st \neq 1$ . Then the reflection of  $X$

$$X : \begin{array}{cccccccc} s & s^{-1} & -1 & s & -1 & t & -1 & t^{-1} & -1 & t & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} s & s^{-1} & -1 & s & t & t^{-1} & -1 & t & t^{-1} & t & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.10 ( $a_2$ ), this case is impossible.

Step 5.7. Consider that  $q_{22} = q_{33} = q_{44} = q_{66} = q_{77} = -1$ ,  $q_{55}\widetilde{q_{45}} - 1 = 0$  and  $q_{ii}\widetilde{q_{i,i+1}} - 1 = 0$  for  $i \in \{1, 5\}$ . Let  $r := q_{11}$ ,  $s := \widetilde{q_{23}}$ ,  $t := \widetilde{q_{34}}$ ,  $u := q_{55}$  and  $q := \widetilde{q_{67}}$ . By Lemma 2.23, we obtain  $r^{-1}s = tu^{-1} = u^{-1}q = 1$  and  $st \neq 1$ . An analogous argument to that in Step 5.6, based on the same reflection conditions and results, rules out this case.

Step 6. Assume that there are six vertices labeled by  $-1$ . By  $q_{33} = -1$ , the following four steps are distinguished.

Step 6.1. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{55} = q_{66} = -1$  and  $q_{77}\widetilde{q_{67}} - 1 = 0$ . Let  $r := \widetilde{q_{12}}$ ,  $s := \widetilde{q_{23}}$ ,  $t := \widetilde{q_{34}}$ ,  $u := \widetilde{q_{45}}$ ,  $v := \widetilde{q_{56}}$  and  $q := q_{77}$ . By Lemma 2.23 and Definition 3.14, we have  $rs = tu = uv = vq^{-1} = 1$  and  $st \neq 1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccc} -1 & s^{-1} & -1 & s & -1 & t & -1 \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccc} -1 & s^{-1} & -1 & s & t & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.10 ( $a_1$ ), this case is impossible.

Consider the symmetric case that  $q_{22} = q_{33} = q_{44} = q_{55} = q_{66} = q_{77} = -1$  and  $q_{11}\widetilde{q_{12}} - 1 = 0$ . Similarly to the proof above, this case is impossible by Lemma 4.10 ( $a_2$ ).

Step 6.2. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{55} = q_{77} = -1$  and  $q_{66}\widetilde{q_{56}} - 1 = q_{66}\widetilde{q_{67}} - 1 = 0$ . Let  $r := \widetilde{q_{12}}$ ,  $s := \widetilde{q_{23}}$ ,  $t := \widetilde{q_{34}}$ ,  $u := \widetilde{q_{45}}$  and  $q := q_{66}$ . Then  $rs = tu = uq^{-1} = 1$  and  $st \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{ccccccc} -1 & s^{-1} & -1 & s & -1 & t & -1 \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccc} -1 & s^{-1} & -1 & s & t & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.10 ( $a_1$ ), this case is impossible.

Consider the symmetric case that  $q_{11} = q_{33} = q_{44} = q_{55} = q_{66} = q_{77} = -1$  and  $q_{22}\widetilde{q_{12}} - 1 = q_{22}\widetilde{q_{23}} - 1 = 0$ . By an argument analogous to the one above, Lemma 4.10 ( $b_1$ ) implies that this case is impossible.

Step 6.3. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{66} = q_{77} = -1$  and  $q_{55}\widetilde{q_{45}} - 1 = q_{55}\widetilde{q_{56}} - 1 = 0$ . Let  $r := \widetilde{q_{12}}$ ,  $s := \widetilde{q_{23}}$ ,  $t := \widetilde{q_{34}}$ ,  $u := q_{55}$  and  $q := \widetilde{q_{67}}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs = tu^{-1} = u^{-1}q = 1$  and  $st \neq 1$ . Hence  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . The same reflection conditions and results from step 6.2 lead to the impossibility of this case.

Step 6.4. Consider that  $q_{11} = q_{22} = q_{33} = q_{55} = q_{66} = q_{77} = -1$  and  $q_{44}\widetilde{q_{34}} - 1 = q_{44}\widetilde{q_{45}} - 1 = 0$ . Let  $r := \widetilde{q_{12}}$ ,  $s := \widetilde{q_{23}}$ ,  $t := q_{44}$ ,  $u := \widetilde{q_{56}}$  and  $q := \widetilde{q_{67}}$ . By Definition 3.14, we obtain  $rs = t^{-1}u = uq = 1$  and  $st^{-1} \neq 1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccc} -1 & s^{-1} & -1 & s & -1 & t^{-1} & t \\ \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{ccccccc} s^{-1} & s & -1 & s^{-1} & s & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.10 ( $c_5$ ), this case is impossible.

Step 7. Assume that all vertices are labeled by  $-1$ . That is  $q_{11} = q_{22} = q_{33} = q_{44} = q_{55} = q_{66} = q_{77} = -1$ . Let  $r := \widetilde{q_{12}}$ ,  $s := \widetilde{q_{23}}$ ,  $t := \widetilde{q_{34}}$ ,  $u := \widetilde{q_{45}}$ ,  $v := \widetilde{q_{56}}$ , and  $q := \widetilde{q_{67}}$ . Then in this case, we have  $rs = tu = uv = vq = 1$  and  $st \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{ccccccc} -1 & s^{-1} & -1 & s & -1 & t & -1 \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccc} -1 & s^{-1} & -1 & s & t & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.10 ( $a_1$ ), this case is impossible.  $\square$

**Theorem 4.12.** *If  $\Delta^{[M]}$  is a finite set of roots of sporadic finite Cartan graphs of rank 8 then all possible Dynkin diagrams of  $V$  appear in row 22 of Table 1.*

*Proof.* By Theorem 3.15,  $C(M)$  is standard of type  $E_8$ . Then  $A^X = E_8$  yields  $(2)_{q_{22}}(q_{22}\widetilde{q_{24}} - 1) = (2)_{q_{ii}}(q_{ii}\widetilde{q_{i,i+1}} - 1) = (2)_{q_{jj}}(q_{jj}\widetilde{q_{j-1,j}} - 1) = 0$  for all  $i \in \{1, 3, 4, 5, 6, 7\}$  and  $j \in \{2, 4, 5, 6, 7, 8\}$ . Let  $q := \widetilde{q_{12}}$ . By  $A^X = E_8$ , one obtains  $\mathcal{D} = \mathcal{D}_{22,1}^8$  if  $q \neq -1$  and  $\mathcal{D} = \mathcal{D}_{22,1}^8$  if  $q = -1$ .  $\square$



*Proof of Theorem 4.1.* The if part is clear. Indeed, we assume that the generalized Dynkin diagram  $\mathcal{D}$  of  $V$  appears in row  $s$  or  $s'$  of any of Table 1. Then  $M$  admits all reflections by Lemmas 2.20 and 2.23 and we can attach a Cartan graph  $\mathcal{C}(M)$  to  $V$  by Theorem 2.24. Notice that the Cartan graph  $\mathcal{C}(M)$  is the same with the Cartan graph obtained from the Dynkin diagrams listed in row  $s$  of Table 4 in [26], which appears in the third column of Table 2. Moreover, the arithmetic root systems of the above Cartan graph are finite, see [26, Theorem 22]. Hence  $\mathcal{W}(\mathcal{C}(M))$  is finite. Then  $\mathcal{B}(V)$  has a finite set of roots  $\Delta^{[M]}$  by Theorem 2.26.

Next we prove that if  $\mathcal{B}(V)$  has a finite set of roots then the generalized Dynkin diagram of  $V$  appears in Table 1. Assume that  $\mathcal{B}(V)$  has a finite set of roots  $\Delta^{[M]}$ . By Theorem 2.26, we obtain that  $\mathcal{C}(M)$  is a finite Cartan graph. From Theorem 1.2, the finite Cartan graph  $\mathcal{C}(M)$  is either non-sporadic or sporadic. If  $\mathcal{C}(M)$  is a non-sporadic finite Cartan graph, Theorem 4.5 gives that the generalized Dynkin diagram  $\mathcal{D}$  of  $V$  appears in rows 1 to 10 of Table 1. If  $\mathcal{C}(M)$  is a sporadic finite Cartan graph, then the generalized Dynkin diagram  $\mathcal{D}$  of  $V$  appears in rows 11 to 22 of Table 1 by Theorem 4.7, 4.9, 4.11, and 4.12.  $\square$

**Remark 4.13.** Notice that the row of Table 1 containing  $\mathcal{D}$  consists precisely of the generalized Dynkin diagrams of all points of  $\mathcal{C}(M)$  and we present the corresponding exchange graphs for the sporadic case of  $\mathcal{C}(M)$  in Table 2.




## 5 Generalized Dynkin diagrams and exchanges graphs

### 5.1 Generalized Dynkin diagrams

row	gener. Dynkin diagrams	fixed parameters	char $\mathbb{k}$
1	$C(r, q; )$	$q \in \mathbb{k}^* \setminus \{1\}$	$p > 0$
2	$C(r, q; i_1, \dots, i_j)$ $C(r, q^{-1}; i_1, \dots, i_{r+1-j})$	$q \in \mathbb{k}^*, q^2 \neq 1$ $1 \leq j \leq (r+1)/2$	$p > 0$
3	$C(r-1, q^2; )$	$q \in \mathbb{k}^*, q^2 \neq 1$	$p > 0$
4	$C(r-1, q^2; i_1, \dots, i_j)$ $C(r-1, q^{-2}; i_1, \dots, i_{r-j})$	$q \in \mathbb{k}^*, q^4 \neq 1$ $1 \leq j \leq r-1$	$p > 0$
5	$C(r-1, -\zeta^{-1}; )$	$\zeta \in G'_3$	$p \neq 2, 3$
5'			$p = 3$
6	$C(r-1, -\zeta^{-1}; i_1, \dots, i_j)$ $C(r-1, -\zeta; i_1, \dots, i_{r-j})$	$\zeta \in G'_3$ $1 \leq j \leq r-1$	$p \neq 2, 3$
7	$C(r-2, q; )$	$q \in \mathbb{k}^*, q^2 \neq 1$	$p > 0$
8	$C(r-2, q; )$	$q \in \mathbb{k}^* \setminus \{1\}$	$p > 0$
9	$C(r-2, q; i_1)$ $C(r-2, q^{-1}; 1, \dots, r-2)$ $C(r-2, q; )$	$q \in \mathbb{k}^*, q^2 \neq 1$	$p > 0$
10	$C(r-1, q; i_1, \dots, i_j)$ $C(r-2, q; i_1, \dots, i_{j-1})$ $C(r-2, q^{-1}; i_1, \dots, i_{r-j})$	$q \in \mathbb{k}^*, q^2 \neq 1$ $2 \leq j \leq r-1$	$p > 0$

row	gener. Dynkin diagrams	fixed parameters	char $\mathbb{k}$
11		$\zeta \in G'_3$	$p \neq 3$
12		$\zeta \in G'_3$	$p \neq 3$

row	gener. Dynkin diagrams	fixed parameters	char $\mathbb{k}$
13		$\zeta \in G'_3$	$p \neq 3$

row	gener. Dynkin diagrams	fixed parameters	char $\mathbb{k}$
	 		
14		$\zeta \in G'_4$	$p \neq 2$

row	gener. Dynkin diagrams	fixed parameters	char $\mathbb{k}$
15		$\zeta \in G'_5$	$p \neq 5$
16		$q \in \mathbb{k}^* \setminus \{1\}$	$p > 0$
16'			$p > 0$
17		$\zeta \in G'_3$	$p \neq 3$

row	gener. Dynkin diagrams	fixed parameters	char $\mathbb{k}$
18	         	$\zeta \in G'_3$	$p \neq 3$

row	gener. Dynkin diagrams	fixed parameters	char $\mathbb{k}$
19		$\zeta \in G'_4$	$p \neq 2$



row	gener. Dynkin diagrams	fixed parameters	char $\mathbb{k}$
20		$q \in \mathbb{k}^* \setminus \{1\}$	$p > 0$
20'			$p > 0$
21		$\zeta \in G_3$	$p \neq 3$
22		$q \in \mathbb{k}^* \setminus \{1\}$	$p > 0$
22'			$p > 0$

Table 1: Dynkin diagrams in all positive characteristic  $p > 0$

## 5.2 Exchange graphs

	exchange graph	row	char $\mathbb{k}$
11	$  \begin{array}{c}  \mathcal{D}_{11,1}^5 \xrightarrow{3} \mathcal{D}_{11,2}^5 \xrightarrow{2} \tau_{54321} \mathcal{D}_{11,3}^5 \xrightarrow{1} \tau_{54321} \mathcal{D}_{11,4}^5 \\    \quad 4 \\  \tau_{12435} \mathcal{D}_{11,3}^5 \xrightarrow{5} \tau_{12435} \mathcal{D}_{11,4}^5  \end{array}  $	11	$p \neq 3$
12	$  \begin{array}{c}  \tau_{13245} \mathcal{D}_{12,14}^5 \xrightarrow{2} \mathcal{D}_{12,11}^5 \xrightarrow{3} \mathcal{D}_{12,13}^5 \\  \quad \quad \quad   \quad 1 \quad \quad \quad   \quad 1 \\  \quad \quad \quad \mathcal{D}_{12,9}^5 \xrightarrow{3} \mathcal{D}_{12,12}^5 \xrightarrow{2} \mathcal{D}_{12,15}^5 \\  \quad \quad \quad   \quad 4 \\  \tau_{12435} \mathcal{D}_{12,7}^5 \xrightarrow{1} \tau_{12435} \mathcal{D}_{12,6}^5 \xrightarrow{2} \tau_{12435} \mathcal{D}_{12,5}^5 \\    \quad 5 \quad \quad \quad   \quad 5 \quad \quad \quad   \quad 5 \\  \tau_{12435} \mathcal{D}_{12,10}^5 \xrightarrow{1} \tau_{12435} \mathcal{D}_{12,8}^5 \xrightarrow{2} \tau_{12435} \mathcal{D}_{12,4}^5 \\  \quad \quad \quad   \quad 4 \\  \quad \quad \quad \tau_{12435} \mathcal{D}_{12,3}^5 \xrightarrow{3} \tau_{12435} \mathcal{D}_{12,2}^5 \xrightarrow{5} \tau_{12435} \mathcal{D}_{12,1}^5  \end{array}  $	12	$p \neq 3$
13	$  \begin{array}{c}  \mathcal{D}_{13,21}^5 \xrightarrow{1} \mathcal{D}_{13,20}^5 \xrightarrow{2} \mathcal{D}_{13,19}^5 \xrightarrow{3} \mathcal{D}_{13,18}^5 \\  \quad \quad \quad   \quad 4 \\  \tau_{12435} \mathcal{D}_{13,16}^5 \xrightarrow{1} \tau_{12435} \mathcal{D}_{13,15}^5 \xrightarrow{2} \tau_{12435} \mathcal{D}_{13,17}^5 \\    \quad 5 \quad \quad \quad   \quad 5 \quad \quad \quad   \quad 5 \\  \tau_{12435} \mathcal{D}_{13,13}^5 \xrightarrow{1} \tau_{12435} \mathcal{D}_{13,10}^5 \xrightarrow{2} \tau_{12435} \mathcal{D}_{13,14}^5 \\    \quad 4 \quad \quad \quad   \quad 4 \\  \tau_{12435} \mathcal{D}_{13,12}^5 \xrightarrow{2} \tau_{12435} \mathcal{D}_{13,9}^5 \xrightarrow{1} \tau_{12435} \mathcal{D}_{13,6}^5 \\    \quad 3 \quad \quad \quad   \quad 3 \quad \quad \quad   \quad 3 \\  \tau_{12435} \mathcal{D}_{13,11}^5 \xrightarrow{4} \tau_{12435} \mathcal{D}_{13,8}^5 \xrightarrow{2} \tau_{12435} \mathcal{D}_{13,5}^5 \xrightarrow{1} \tau_{12435} \mathcal{D}_{13,3}^5 \\    \quad 5 \quad \quad \quad   \quad 5 \quad \quad \quad   \quad 5 \\  \tau_{12435} \mathcal{D}_{13,7}^5 \xrightarrow{4} \tau_{12435} \mathcal{D}_{13,4}^5 \xrightarrow{2} \tau_{12435} \mathcal{D}_{13,2}^5 \xrightarrow{1} \tau_{12435} \mathcal{D}_{13,1}^5  \end{array}  $	13	$p \neq 3$
14	$  \begin{array}{c}  \tau_{12354} \mathcal{D}_{14,6}^5 \xrightarrow{1} \tau_{12354} \mathcal{D}_{14,5}^5 \\    \quad 2 \\  \mathcal{D}_{14,4}^5 \xrightarrow{4} \tau_{12354} \mathcal{D}_{14,3}^5 \xrightarrow{5} \tau_{12354} \mathcal{D}_{14,1}^5 \xrightarrow{3} \tau_{12354} \mathcal{D}_{14,2}^5 \\    \quad 3 \quad \quad \quad   \quad 5 \\  \tau_{12534} \mathcal{D}_{14,2}^5 \xrightarrow{5} \tau_{12534} \mathcal{D}_{14,1}^5 \xrightarrow{3} \tau_{12534} \mathcal{D}_{14,3}^5 \\    \quad 2 \quad \quad \quad   \quad 4 \\  \tau_{12534} \mathcal{D}_{14,5}^6 \xrightarrow{1} \tau_{12534} \mathcal{D}_{14,6}^6 \quad \tau_{12543} \mathcal{D}_{14,4}^5  \end{array}  $	14	$p \neq 2$
15	$  \begin{array}{c}  \mathcal{D}_{15,1}^5 \xrightarrow{3} \mathcal{D}_{15,2}^5 \xrightarrow{2} \mathcal{D}_{15,4}^5 \xrightarrow{1} \mathcal{D}_{15,5}^5 \\    \quad 4 \\  \tau_{12435} \mathcal{D}_{15,3}^5 \xrightarrow{5} \tau_{12453} \mathcal{D}_{15,6}^5 \xrightarrow{3} \tau_{12453} \mathcal{D}_{15,7}^5  \end{array}  $	15	$p \neq 5$
16	$\mathcal{D}_{16,1}^6$	16	$p > 0$
16'	$\mathcal{D}_{16',1}^6$	16	$p > 0$

(continued)	exchange graph	row	char $\mathbb{k}$
17	$\begin{array}{c} \mathcal{D}_{17,7}^6 \xrightarrow{1} \mathcal{D}_{17,6}^6 \xrightarrow{2} \mathcal{D}_{17,4}^6 \xrightarrow{3} \mathcal{D}_{17,2}^6 \xrightarrow{4} \mathcal{D}_{17,1}^6 \\   \\ \tau_{123546} \mathcal{D}_{17,3}^6 \xrightarrow{6} \tau_{123546} \mathcal{D}_{17,5}^6 \end{array}$	17	$p \neq 3$
18	$\begin{array}{ccccccc} & & \mathcal{D}_{18,9}^6 & \xrightarrow{1} & \mathcal{D}_{18,7}^6 & \xrightarrow{2} & \mathcal{D}_{18,5}^6 & \xrightarrow{3} & \mathcal{D}_{18,4}^6 & \xrightarrow{4} & \mathcal{D}_{18,3}^6 \\ & &   & &   & &   & &   & &   \\ \tau_{125346} \mathcal{D}_{18,21}^6 & & \mathcal{D}_{18,12}^6 & \xrightarrow{1} & \mathcal{D}_{18,10}^6 & \xrightarrow{2} & \mathcal{D}_{18,8}^6 & \xrightarrow{3} & \mathcal{D}_{18,6}^6 & & \mathcal{D}_{18,2}^6 \\ & &   & &   & &   & &   & &   \\ \tau_{125346} \mathcal{D}_{18,20}^6 & \xrightarrow{3} & \tau_{123546} \mathcal{D}_{18,15}^6 & \xrightarrow{1} & \tau_{123546} \mathcal{D}_{18,13}^6 & \xrightarrow{3} & \tau_{125346} \mathcal{D}_{18,17}^6 & & \mathcal{D}_{18,1}^6 \\ & &   & &   & &   & &   \\ \tau_{125346} \mathcal{D}_{18,17}^6 & & \tau_{123546} \mathcal{D}_{18,19}^6 & \xrightarrow{1} & \tau_{123546} \mathcal{D}_{18,16}^6 & & \tau_{123546} \mathcal{D}_{18,11}^6 & \xrightarrow{4} & \mathcal{D}_{18,8}^6 \\ & &   & &   & &   & &   \\ & & \tau_{123546} \mathcal{D}_{18,18}^6 & \xrightarrow{3} & \tau_{123546} \mathcal{D}_{18,14}^6 & \xrightarrow{5} & \tau_{123546} \mathcal{D}_{18,11}^6 & \xrightarrow{4} & \mathcal{D}_{18,8}^6 \end{array}$	18	$p \neq 3$
19	$\begin{array}{ccccccccc} & & \mathcal{D}_{19,1}^6 & \xrightarrow{4} & \mathcal{D}_{19,2}^6 & \xrightarrow{3} & \mathcal{D}_{19,5}^6 & \xrightarrow{2} & \mathcal{D}_{19,6}^6 & \xrightarrow{1} & \mathcal{D}_{19,7}^6 \\ & &   & &   & &   & &   & &   \\ \tau_{123465} \mathcal{D}_{19,4}^6 & \xrightarrow{6} & \mathcal{D}_{19,3}^6 & & \tau_{123546} \mathcal{D}_{19,3}^6 & \xrightarrow{6} & \tau_{123564} \mathcal{D}_{19,4}^6 & & & & \\ & &   & &   & &   & &   & &   \\ & & \tau_{123546} \mathcal{D}_{19,2}^6 & \xrightarrow{5} & \tau_{123546} \mathcal{D}_{19,1}^6 & & & & & & \\ & &   & &   & &   & &   & &   \\ & & \tau_{125346} \mathcal{D}_{19,5}^6 & \xrightarrow{2} & \tau_{125346} \mathcal{D}_{19,6}^6 & \xrightarrow{1} & \tau_{125346} \mathcal{D}_{19,7}^6 & & & & \end{array}$	19	$p \neq 2$
20	$\mathcal{D}_{20,1}^7$	20	$p > 0$
20'	$\mathcal{D}_{20',1}^7$	20	$p > 0$
21	$\begin{array}{ccccc} \tau_{7654321} \mathcal{D}_{21,4}^7 & & \tau_{7654321} \mathcal{D}_{21,1}^7 & & \tau_{7654321} \mathcal{D}_{21,8}^7 \xrightarrow{7} \tau_{7654321} \mathcal{D}_{21,7}^7 \\   & &   & &   \\ \tau_{7654321} \mathcal{D}_{21,3}^7 & \xrightarrow{2} & \tau_{7654321} \mathcal{D}_{21,2}^7 & \xrightarrow{4} & \tau_{7654321} \mathcal{D}_{21,5}^7 \xrightarrow{5} \tau_{7654321} \mathcal{D}_{21,6}^7 \end{array}$	21	$p \neq 3$
22	$\mathcal{D}_{22,1}^8$	22	$p > 0$
22'	$\mathcal{D}_{22',1}^8$	22	$p > 0$

Table 2: The exchange graphs of  $\mathcal{C}(M)$ 

## 6 Appendix

For brevity, we adopt the product notation  $\prod_{i=1}^r i^{x_i}$  as a shorthand for the sum  $\sum_{i=1}^r x_i \alpha_{r+1-i}$ .

**Rank 3:** Nr. 1 with 6 positive roots (type  $A_3$ ): 1, 2, 3, 12, 13, 123

Nr. 2 with 7 positive roots (type  $D'(3, 1)$ ): 1, 2, 3, 12, 13, 23, 123

Nr. 3 with 8 positive roots (type  $D'(3, 2)$ ): 1, 2, 3, 12, 13,  $1^2 2$ , 123,  $1^2 23$

Nr. 4 with 9 positive roots (type  $C_3$ ): 1, 2, 3, 12, 13,  $1^2 2$ , 123,  $1^2 23$ ,  $1^2 23^2$

Nr. 5 with 9 positive roots (type  $B_3$ ): 1, 2, 3, 12, 23,  $1^2 2$ , 123,  $1^2 23$ ,  $1^2 2^2 3$

**Rank 4:** Nr. 1 with 10 positive roots (type  $A_4$ ): 1, 2, 3, 4, 12, 23, 34, 123, 234, 1234

Nr. 2 with 12 positive roots (type  $D_4$ ): 1, 2, 3, 4, 13, 23, 34, 123, 134, 234, 1234,  $123^2 4$

Nr. 3 with 13 positive roots (type  $D'(4, 1)$ ): 1, 2, 3, 4, 12, 13, 23, 34, 123, 134, 234, 1234,  $123^2 4$

Nr. 4 with 14 positive roots (type  $D'(4, 2)$ ): 1, 2, 3, 4, 12, 13, 23, 34, 123, 134, 234,  $123^2$ , 1234,  $123^2 4$

Nr. 5 with 15 positive roots (type  $D'(4, 3)$ ): 1, 2, 3, 4, 12, 13, 23, 34, 123, 134, 234,  $123^2$ ,  $1234$ ,  $123^24$ ,  $123^24^2$

Nr. 6 with 16 positive roots (type  $C_4$ ): 1, 2, 3, 4, 12, 13, 34,  $12^2$ , 123, 134,  $12^2 3$ , 1234,  $12^2 3^2$ ,  $12^2 34$ ,  $12^2 3^2 4$ ,  $12^2 3^2 4^2$

Nr. 7 with 16 positive roots (type  $B_4$ ): 1, 2, 3, 4, 12, 23, 34,  $1^2 2$ , 123, 234,  $1^2 23$ , 1234,  $1^2 2^2 3$ ,  $1^2 234$ ,  $1^2 2^2 34$ ,  $1^2 2^2 3^2 4$

**Rank 6:** Nr. 1 with 21 positive roots (type  $A_6$ ): 1, 2, 3, 4, 5, 6, 12, 23, 34, 45, 56, 123, 234, 345, 456, 1234, 2345, 3456, 12345, 23456, 123456  
Nr. 2 with 30 positive roots (type  $D_6$ ): 1, 2, 3, 4, 5, 6, 13, 23, 34, 45, 56, 123, 134, 234, 345, 456, 1234, 1345, 2345, 3456,  $123^2 4$ , 12345, 13456, 23456,  $123^2 45$ , 123456,  $123^2 4^2 5$ ,  $123^2 456$ ,  $123^2 4^2 56$ ,  $123^2 4^2 5^2 6$   
Nr. 3 with 31 positive roots (type  $D'(6, 1)$ ): 1, 2, 3, 4, 5, 6, 12, 13, 23, 34, 45, 56, 123, 134, 234, 345, 456, 1234, 1345, 2345, 3456,  $123^2 4$ , 12345, 13456, 23456,  $123^2 45$ , 123456,  $123^2 4^2 5$ ,  $123^2 456$ ,  $123^2 4^2 56$ ,  $123^2 4^2 5^2 6$   
Nr. 4 with 32 positive roots (type  $D'(6, 2)$ ): 1, 2, 3, 4, 5, 6, 12, 13, 23, 34, 45, 56, 123, 134, 234, 345, 456,  $123^2$ , 1234, 1345, 2345, 3456,  $123^2 4$ , 12345, 13456, 23456,  $123^2 45$ , 123456,  $123^2 4^2 5$ ,  $123^2 456$ ,  $123^2 4^2 56$ ,  $123^2 4^2 5^2 6$   
Nr. 5 with 33 positive roots (type  $D'(6, 3)$ ): 1, 2, 3, 4, 5, 6, 12, 13, 23, 34, 45, 56, 123, 134, 234, 345, 456,  $123^2$ , 1234, 1345, 2345, 3456,  $123^2 4$ , 12345, 13456, 23456,  $123^2 4^2$ ,  $123^2 45$ , 123456,  $123^2 4^2 5$ ,  $123^2 456$ ,  $123^2 4^2 56$ ,  $123^2 4^2 5^2 6$   
Nr. 6 with 34 positive roots (type  $D'(6, 4)$ ): 1, 2, 3, 4, 5, 6, 12, 13, 23, 34, 45, 56, 123, 134, 234, 345, 456,  $123^2$ , 1234, 1345, 2345, 3456,  $123^2 4$ , 12345, 13456, 23456,  $123^2 4^2$ ,  $123^2 45$ , 123456,  $123^2 4^2 5$ ,  $123^2 456$ ,  $123^2 4^2 5^2$ ,  $123^2 4^2 56$ ,  $123^2 4^2 5^2 6$   
Nr. 7 with 35 positive roots (type  $D'(6, 5)$ ): 1, 2, 3, 4, 5, 6, 12, 13, 23, 34, 45, 56, 123, 134, 234, 345, 456,  $123^2$ , 1234, 1345, 2345, 3456,  $123^2 4$ , 12345, 13456, 23456,  $123^2 4^2$ ,  $123^2 45$ , 123456,  $123^2 4^2 5$ ,  $123^2 456$ ,  $123^2 4^2 5^2$ ,  $123^2 4^2 56$ ,  $123^2 4^2 5^2 6$ ,  $123^2 4^2 5^2 6^2$   
Nr. 8 with 36 positive roots (type  $C_6$ ): 1, 2, 3, 4, 5, 6, 12, 23, 34, 45, 56,  $12^2$ , 123, 234, 345, 456,  $12^2 3$ , 1234, 2345, 3456,  $12^2 3^2$ ,  $12^2 34$ , 12345, 23456,  $12^2 3^2 4$ ,  $12^2 345$ , 123456,  $12^2 3^2 4^2$ ,  $12^2 3^2 45$ ,  $12^2 3456$ ,  $12^2 3^2 4^2 5$ ,  $12^2 3^2 4^2 56$ ,  $12^2 3^2 4^2 5^2 6$ ,  $12^2 3^2 4^2 5^2 6^2$   
Nr. 9 with 36 positive roots (type  $B_6$ ): 1, 2, 3, 4, 5, 6, 12, 23, 34, 45, 56,  $12^2$ , 123, 234, 345, 456,  $12^2 23$ , 1234, 2345, 3456,  $12^2 2^3$ ,  $12^2 34$ , 12345, 23456,  $12^2 2^2 34$ ,  $12^2 345$ , 123456,  $12^2 2^2 3^2 4$ ,  $12^2 2^2 345$ , 123456,  $12^2 2^2 3^2 4^2$ ,  $12^2 2^2 3^2 45$ ,  $12^2 2^2 3^2 456$ ,  $12^2 2^2 3^2 4^2 56$ ,  $12^2 2^2 3^2 4^2 5^2 6$

**Rank 8:** Nr. 1 with 36 positive roots (type  $A_8$ ): 1, 2, 3, 4, 5, 6, 7, 8, 12, 23, 34, 45, 56, 67, 78, 123, 234, 345, 456, 567, 678, 1234, 2345, 3456, 4567,

Nr. 2 with 56 positive roots (type  $D_8$ ): 1, 2, 3, 4, 5, 6, 7, 8, 13, 23, 34, 45, 56, 67, 78, 123, 134, 234, 345, 456, 567, 678, 1234, 1345, 2345, 3456, 4567, 5678,  $123^2$  4, 12345, 13456, 23456, 34567, 45678,  $123^2$  45, 123456, 134567, 234567, 345678,  $123^2$  4<sup>2</sup> 5,  $123^2$  456, 1234567, 1345678, 2345678,  $123^2$  4<sup>2</sup> 56,  $123^2$  4567, 12345678,  $123^2$  4<sup>2</sup> 5<sup>2</sup> 6,  $123^2$  4<sup>2</sup> 567,  $123^2$  45678,  $123^2$  4<sup>2</sup> 5<sup>2</sup> 6<sup>2</sup> 7,  $123^2$  4<sup>2</sup> 5678,  $123^2$  4<sup>2</sup> 5<sup>2</sup> 6<sup>2</sup> 78,  $123^2$  4<sup>2</sup> 5<sup>2</sup> 6<sup>2</sup> 7<sup>2</sup> 8

Nr. 4 with 58 positive roots (type  $D(8, 2)$ ): 1, 2, 3, 4, 5, 6, 7, 8, 12, 13, 23, 34, 45, 56, 67, 78, 123, 134, 234, 345, 456, 567, 678,  $123^2$ , 1234, 1345, 2345, 3456, 4567, 5678,  $123^2 4$ , 12345, 13456, 23456, 34567, 45678,  $123^2 45$ , 123456, 134567, 234567, 345678,  $123^2 4^2 5$ ,  $123^2 456$ , 1234567, 12345678,  $123^2 4^2 56$ ,  $123^2 4567$ , 12345678,  $123^2 4^2 5^2 6$ ,  $123^2 4^2 567$ ,  $123^2 45678$ ,  $123^2 4^2 5^2 67$ ,  $123^2 4^2 5678$ ,  $123^2 4^2 5^2 6^2 7$ ,  $123^2 4^2 5^2 678$ ,  $123^2 4^2 5^2 6^2 78$ ,  $123^2 4^2 5^2 6^2 7^2 8$

Nr. 6 with 60 positive roots (type  $D'(8,4)$ ): 1, 2, 3, 4, 5, 6, 7, 8, 12, 13, 23, 34, 45, 56, 67, 78, 123, 134, 234, 345, 456, 567, 678,  $123^2$ , 1234, 1345, 2345, 3456, 4567, 5678,  $123^2 4$ , 12345, 13456, 23456, 34567, 45678,  $123^2 4^2$ ,  $123^2 45$ , 123456, 134567, 234567, 345678,  $123^2 4^2 5$ ,  $123^2 456$ , 1234567, 1345678, 2345678,  $123^2 4^2 5^2$ ,  $123^2 4^2 56$ ,  $123^2 4567$ , 12345678,  $123^2 4^2 5^2 6$ ,  $123^2 4^2 567$ ,  $123^2 45678$ ,  $123^2 4^2 5^2 67$ ,  $123^2 4^2 5678$ ,  $123^2 4^2 5^2 6^2 7$ ,  $123^2 4^2 5^2 6^2 78$ ,  $123^2 4^2 5^2 6^2 7^2 8$

Nr. 8 with 62 positive roots (type  $D'(8,6)$ ): 1, 2, 3, 4, 5, 6, 7, 8, 12, 13, 23, 34, 45, 56, 67, 78, 123, 134, 234, 345, 456, 567, 678,  $123^2$ , 1234, 1345, 2345, 3456, 4567, 5678,  $123^2 4$ , 12345, 13456, 23456, 34567, 45678,  $123^2 4^2$ ,  $123^2 45$ , 123456, 134567, 234567, 345678,  $123^2 4^2 5$ ,  $123^2 456$ , 1234567, 1345678, 2345678,  $123^2 4^2 5^2$ ,  $123^2 4^2 56$ ,  $123^2 4567$ ,  $123^2 4^2 5^2 6$ ,  $123^2 4^2 567$ ,  $123^2 45678$ ,  $123^2 4^2 5^2 6^2$ ,  $123^2 4^2 5^2 67$ ,  $123^2 4^2 5678$ ,  $123^2 4^2 5^2 6^2 7$ ,  $123^2 4^2 5^2 6^2 78$ ,  $123^2 4^2 5^2 6^2 7^2 8$

Nr. 10 with 64 positive roots (type  $C_8$ ): 1, 2, 3, 4, 5, 6, 7, 8, 12, 23, 34, 45, 56, 67, 78,  $12^2$ , 123, 234, 345, 456, 567, 678,  $12^2 3$ , 1234, 2345, 3456, 4567, 5678,  $12^2 3^2$ ,  $12^2 34$ , 12345, 23456, 34567, 45678,  $12^2 3^2 4$ ,  $12^2 345$ , 123456, 234567, 345678,  $12^2 3^2 4^2$ ,  $12^2 3^2 45$ ,  $12^2 3456$ , 1234567, 12345678,  $12^2 3^2 4^2 5$ ,  $12^2 3^2 4^2 56$ ,  $12^2 3^2 4567$ ,  $12^2 3^2 4^2 567$ ,  $12^2 3^2 45678$ ,  $12^2 3^2 4^2 5678$ ,  $12^2 3^2 4^2 5678^2$ ,  $12^2 3^2 4^2 5678^2 7$ ,  $12^2 3^2 4^2 5678^2 7^2$ ,  $12^2 3^2 4^2 5678^2 7^2 8$ ,  $12^2 3^2 4^2 5678^2 7^2 8^2$ ,  $12^2 3^2 4^2 5678^2 7^2 8^2 7$ ,  $12^2 3^2 4^2 5678^2 7^2 8^2 7^2$ ,  $12^2 3^2 4^2 5678^2 7^2 8^2 7^2 8$ ,  $12^2 3^2 4^2 5678^2 7^2 8^2 7^2 8^2$ .

**Rank 9:** Nr. 1 with 45 positive roots (type  $A_9$ ): 1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 23, 34, 45, 56, 67, 78, 89, 123, 234, 345, 456, 567, 678, 789, 1234, 2345, 3456, 4567, 5678, 6789, 12345, 23456, 34567, 45678, 56789, 123456, 234567, 345678, 456789, 1234567, 2345678, 3456789, 123456789

Nr. 3 with 73 positive roots (type  $D(9,1)$ ): 1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 13, 23, 34, 45, 56, 67, 78, 89, 123, 134, 234, 345, 456, 567, 678, 789, 1234, 1345, 2345, 3456, 4567, 5678, 6789,  $123^24$ ,  $12345$ ,  $13456$ ,  $23456$ ,  $34567$ ,  $45678$ ,  $56789$ ,  $123^245$ ,  $123456$ ,  $134567$ ,  $234567$ ,  $345678$ ,  $456789$ ,  $123^24^25$ ,  $123^2456$ ,  $1234567$ ,  $1345678$ ,  $2345678$ ,  $3456789$ ,  $123^24^256$ ,  $123^24567$ ,  $12345678$ ,  $13456789$ ,  $23456789$ ,  $123^24^2567$ ,  $123^24^25678$ ,  $123^24^256789$ ,  $123^24^25678^2$ ,  $123^24^256789^2$ ,  $123^24^256^278$ ,  $123^24^256^2789$ ,  $123^24^256^2789^2$ ,  $123^24^256^2789^28$ ,  $123^24^256^2789^289$ ,  $123^24^256^2789^289^2$ .

Nr. 4 with 74 positive roots (type  $D'(9, 2)$ ): 1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 13, 23, 34, 45, 56, 67, 78, 89, 123, 134, 234, 345, 456, 567, 678, 789,  $123^2$  1234, 1345, 2345, 3456, 4567, 5678, 6789,  $123^2$  4, 12345, 13456, 23456, 34567, 45678, 56789,  $123^2$  45, 123456, 134567, 234567, 345678, 456789,  $123^2$  4<sup>2</sup>5,  $123^2$  456, 1234567, 1345678, 2345678, 3456789,  $123^2$  4<sup>2</sup>56,  $123^2$  4567, 12345678, 13456789, 23456789,  $123^2$  4<sup>2</sup>567,  $123^2$  4<sup>2</sup>56<sup>2</sup>,  $123^2$  45678, 123456789,  $123^2$  4<sup>2</sup>5678,  $123^2$  4<sup>2</sup>56<sup>2</sup>7,  $123^2$  4<sup>2</sup>56<sup>2</sup>78,  $123^2$  4<sup>2</sup>56<sup>2</sup>789,  $123^2$  4<sup>2</sup>56<sup>2</sup>7<sup>2</sup>8,  $123^2$  4<sup>2</sup>56<sup>2</sup>7<sup>2</sup>89,  $123^2$  4<sup>2</sup>56<sup>2</sup>7<sup>2</sup>8<sup>2</sup>9

Nr. 5 with 75 positive roots (type  $D'(9, 3)$ ): 1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 13, 23, 34, 45, 56, 67, 78, 89, 123, 134, 234, 345, 456, 567, 678, 789,  $123^2$  1234, 1345, 2345, 3456, 4567, 5678, 6789,  $123^2$  4, 12345, 13456, 23456, 34567, 45678, 56789,  $123^2$  4<sup>2</sup>,  $123^2$  45, 123456, 134567, 234567, 345678, 456789,  $123^2$  4<sup>2</sup> 5,  $123^2$  456, 1234567, 1345678, 2345678, 3456789,  $123^2$  4<sup>2</sup> 56,  $123^2$  4567,  $123^2$  4<sup>2</sup> 56<sup>2</sup>,  $123^2$  45678,  $123^2$  4<sup>2</sup> 5<sup>2</sup> 67,  $123^2$  4<sup>2</sup> 5678,  $123^2$  4<sup>2</sup> 5<sup>2</sup> 67<sup>2</sup>,  $123^2$  4<sup>2</sup> 5<sup>2</sup> 678,  $123^2$  4<sup>2</sup> 5<sup>2</sup> 678<sup>2</sup>,  $123^2$  4<sup>2</sup> 5<sup>2</sup> 6<sup>2</sup> 78,  $123^2$  4<sup>2</sup> 5<sup>2</sup> 6<sup>2</sup> 78<sup>2</sup>,  $123^2$  4<sup>2</sup> 5<sup>2</sup> 6<sup>2</sup> 78<sup>2</sup> 9,  $123^2$  4<sup>2</sup> 5<sup>2</sup> 6<sup>2</sup> 7<sup>2</sup> 89,  $123^2$  4<sup>2</sup> 5<sup>2</sup> 6<sup>2</sup> 7<sup>2</sup> 8<sup>2</sup> 9

Nr. 6 with 76 positive roots (type  $D'(9,4)$ ): 1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 13, 23, 34, 45, 56, 67, 78, 89, 123, 134, 234, 345, 456, 567, 678, 789,  $123^2$  1234, 1345, 2345, 3456, 4567, 5678, 6789,  $123^2 4$ , 12345, 13456, 23456, 34567, 45678, 56789,  $123^2 4^2$ ,  $123^2 45$ , 123456, 134567, 234567, 345678, 456789,  $123^2 4^2 5$ ,  $123^2 456$ , 1234567, 1345678, 2345678, 3456789,  $123^2 4^2 5^2$ ,  $123^2 4^2 56$ ,  $123^2 4567$ , 12345678, 13456789, 23456789,  $123^2 4^2 567$ ,  $123^2 4^2 5^2 6$ ,  $123^2 45678$ , 123456789,  $123^2 4^2 5^2 67$ ,  $123^2 4^2 5678$ ,  $123^2 456789$ ,  $123^2 4^2 5^2 6^2 7$ ,  $123^2 4^2 56789$ ,  $123^2 4^2 5^2 678$ ,  $123^2 4^2 5^2 6^2 78$ ,  $123^2 4^2 5^2 6^2 7^2 8$ ,  $123^2 4^2 5^2 6^2 7^2 89$ ,  $123^2 4^2 5^2 6^2 7^2 8^2 9$

Nr. 7 with 77 positive roots (type  $D'(9,5)$ ): 1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 13, 23, 34, 45, 56, 67, 78, 89, 123, 134, 234, 345, 456, 567, 678, 789,  $123^2$  1234, 1345, 2345, 3456, 4567, 5678, 6789,  $123^2 4$ , 12345, 13456, 23456, 34567, 45678, 56789,  $123^2 4^2$ ,  $123^2 45$ , 123456, 134567, 234567, 345678, 456789,  $123^2 4^2 5$ ,  $123^2 456$ , 1234567, 1345678, 2345678, 3456789,  $123^2 4^2 5^2$ ,  $123^2 4^2 56$ ,  $123^2 4567$ , 12345678, 13456789, 23456789,  $123^2 4^2 567$ ,  $123^2 4^2 5^2 6$ ,  $123^2 45678$ , 123456789,  $123^2 4^2 5^2 6^2$ ,  $123^2 4^2 5^2 67$ ,  $123^2 4^2 5678$ ,  $123^2 456789$ ,  $123^2 4^2 5^2 6^2 7$ ,  $123^2 4^2 56789$ ,  $123^2 4^2 5^2 678$ ,  $123^2 4^2 5^2 6^2 78$ ,  $123^2 4^2 5^2 6^2 7^2 8$ ,  $123^2 4^2 5^2 6^2 7^2 89$ ,  $123^2 4^2 5^2 6^2 7^2 8^2 9$

Nr. 8 with 78 positive roots (type  $D'(9,6)$ ): 1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 13, 23, 34, 45, 56, 67, 78, 89, 123, 134, 234, 345, 456, 567, 678, 789,  $123^2$  1234, 1345, 2345, 3456, 4567, 5678, 6789,  $123^2 4$ , 12345, 13456, 23456, 34567, 45678, 56789,  $123^2 4^2$ ,  $123^2 45$ , 123456, 134567, 234567, 345678, 456789,  $123^2 4^2 5$ ,  $123^2 456$ , 1234567, 1345678, 2345678, 3456789,  $123^2 4^2 5^2$ ,  $123^2 4^2 56$ ,  $123^2 4567$ , 12345678, 13456789, 23456789,  $123^2 4^2 567$ ,  $123^2 4^2 5^2 6$ ,  $123^2 45678$ , 123456789,  $123^2 4^2 5^2 6^2$ ,  $123^2 4^2 5^2 67$ ,  $123^2 4^2 5678$ ,  $123^2 456789$ ,  $123^2 4^2 5^2 6^2 7$ ,  $123^2 4^2 56789$ ,  $123^2 4^2 5^2 678$ ,  $123^2 4^2 5^2 6^2 78$ ,  $123^2 4^2 5^2 6^2 7^2 8$ ,  $123^2 4^2 5^2 6^2 7^2 89$ ,  $123^2 4^2 5^2 6^2 7^2 8^2 9$

Nr. 9 with 79 positive roots (type  $D'(9,7)$ ): 1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 13, 23, 34, 45, 56, 67, 78, 89, 123, 134, 234, 345, 456, 567, 678, 789,  $123^2$  1234, 1345, 2345, 3456, 4567, 5678, 6789,  $123^2 4$ , 12345, 13456, 23456, 34567, 45678, 56789,  $123^2 4^2$ ,  $123^2 45$ , 123456, 134567, 234567, 345678, 456789,  $123^2 4^2 5$ ,  $123^2 456$ , 1234567, 1345678, 2345678, 3456789,  $123^2 4^2 5^2$ ,  $123^2 4^2 56$ ,  $123^2 4567$ , 12345678, 13456789, 23456789,  $123^2 4^2 567$ ,  $123^2 4^2 5^2 6$ ,  $123^2 45678$ , 123456789,  $123^2 4^2 5^2 6^2$ ,  $123^2 4^2 5^2 67$ ,  $123^2 4^2 5678$ ,  $123^2 456789$ ,  $123^2 4^2 5^2 6^2 7$ ,  $123^2 4^2 56789$ ,  $123^2 4^2 5^2 678$ ,  $123^2 4^2 5^2 6^2 78$ ,  $123^2 4^2 5^2 6^2 7^2 8$ ,  $123^2 4^2 5^2 6^2 7^2 89$ ,  $123^2 4^2 5^2 6^2 7^2 8^2 9$

Nr. 10 with 80 positive roots (type  $D'(9,8)$ ): 1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 13, 23, 34, 45, 56, 67, 78, 89, 123, 134, 234, 345, 456, 567, 678, 789,  $123^2$  1234, 1345, 2345, 3456, 4567, 5678, 6789,  $123^2 4$ , 12345, 13456, 23456, 34567, 45678, 56789,  $123^2 4^2$ ,  $123^2 45$ , 123456, 134567, 234567, 345678, 456789,  $123^2 4^2 5$ ,  $123^2 456$ , 1234567, 1345678, 2345678, 3456789,  $123^2 4^2 5^2$ ,  $123^2 4^2 56$ ,  $123^2 4567$ , 12345678, 13456789, 23456789,  $123^2 4^2 567$ ,  $123^2 4^2 5^2 6$ ,  $123^2 45678$ , 123456789,  $123^2 4^2 5^2 6^2$ ,  $123^2 4^2 5^2 67$ ,  $123^2 4^2 5678$ ,  $123^2 456789$ ,  $123^2 4^2 5^2 6^2 7$ ,  $123^2 4^2 56789$ ,  $123^2 4^2 5^2 678$ ,  $123^2 4^2 5^2 6^2 78$ ,  $123^2 4^2 5^2 6^2 7^2 8$ ,  $123^2 4^2 5^2 6^2 7^2 89$ ,  $123^2 4^2 5^2 6^2 7^2 8^2 9$

Nr. 11 with 81 positive roots (type  $C_9$ ): 1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 23, 34, 45, 56, 67, 78, 89,  $12^2$ , 123, 234, 345, 456, 567, 678, 789,  $12^2 3$ , 1234, 2345, 3456, 4567, 5678, 6789,  $12^2 3^2$ ,  $12^2 34$ , 12345, 23456, 34567, 45678, 56789,  $12^2 3^2 4$ ,  $12^2 345$ , 123456, 234567, 345678, 456789,  $12^2 3^2 4^2$ ,  $12^2 3^2 45$ ,  $12^2 3456$ , 1234567, 2345678, 3456789,  $12^2 3^2 4^2 5$ ,  $12^2 3^2 456$ ,  $12^2 34567$ , 12345678, 23456789,  $12^2 3^2 4^2 5^2$ ,  $12^2 3^2 4^2 56$ ,  $12^2 3^2 4567$ ,  $12^2 345678$ , 123456789,  $12^2 3^2 4^2 5^2 6$ ,  $12^2 3^2 4^2 567$ ,  $12^2 3^2 45678$ ,  $12^2 3456789$ ,  $12^2 3^2 4^2 5^2 6^2$ ,  $12^2 3^2 4^2 5^2 67$ ,  $12^2 3^2 4^2 5678$ ,  $12^2 3^2 4^2 5^2 678$ ,  $12^2 3^2 4^2 5^2 6^2 7$ ,  $12^2 3^2 4^2 5^2 6789$ ,  $12^2 3^2 4^2 5^2 6^2 7^2$ ,  $12^2 3^2 4^2 5^2 6^2 7^2 8$ ,  $12^2 3^2 4^2 5^2 6^2 7^2 89$ ,  $12^2 3^2 4^2 5^2 6^2 7^2 8^2 9$

Nr. 12 with 81 positive roots (type  $B_9$ ): 1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 23, 34, 45, 56, 67, 78, 89,  $1^2 2$ , 123, 234, 345, 456, 567, 678, 789,  $1^2 23$ , 1234, 2345, 3456, 4567, 5678, 6789,  $1^2 2^2 3$ ,  $1^2 234$ , 12345, 23456, 34567, 45678, 56789,  $1^2 2^2 34$ ,  $1^2 2345$ , 123456, 234567, 345678, 456789,  $1^2 2^2 3^2 4$ ,  $1^2 2^2 345$ ,  $1^2 23456$ , 1234567, 2345678, 3456789,  $1^2 2^2 3^2 4^2$ ,  $1^2 2^2 3^2 45$ ,  $1^2 2^2 3456$ , 12345678, 23456789,  $1^2 2^2 3^2 4^2 5$ ,  $1^2 2^2 3^2 4^2 56$ ,  $1^2 2^2 3^2 4567$ ,  $1^2 2345678$ , 123456789,  $1^2 2^2 3^2 4^2 5^2 6$ ,  $1^2 2^2 3^2 4^2 5^2 67$ ,  $1^2 2^2 3^2 4^2 5678$ ,  $1^2 2^2 3^2 4^2 5^2 678$ ,  $1^2 2^2 3^2 4^2 5^2 6^2 7$ ,  $1^2 2^2 3^2 4^2 5^2 6^2 78$ ,  $1^2 2^2 3^2 4^2 5^2 6^2 7^2 8$ ,  $1^2 2^2 3^2 4^2 5^2 6^2 7^2 89$ ,  $1^2 2^2 3^2 4^2 5^2 6^2 7^2 8^2 9$

## Acknowledgment

The authors thank N. Andruskiewitsch and H. Yamane for the invaluable suggestions to the references. Furthermore, the author J. Wang would like to thank her PhD advisor Prof. I. Heckenberger for the fruitful discussions during her PhD study in Germany, which have been crucial to the entire project.

## References

- [1] N. Andruskiewitsch. About finite dimensional Hopf algebras. *Contemp. Math.*, 294:1–57, 01 2002.
- [2] N. Andruskiewitsch. On finite-dimensional Hopf algebras. *Proc. Int. Cong. Math.*, 2:117–141, 2014.
- [3] N. Andruskiewitsch, I. Angiono, and A. G. Iglesias. Liftings of Nichols algebras of diagonal type I. Cartan type A. *Int. Math. Res. Not.*, 2017(9):2793–2884, 2017.
- [4] N. Andruskiewitsch and M. Graña. Braided Hopf algebras over non abelian finite groups. *Bol. Acad. Nac. Cienc. (Córdoba)*, 63:45–78, 1998.
- [5] N. Andruskiewitsch, I. Heckenberger, and H.-J. Schneider. The Nichols algebra of a semisimple Yetter-Drinfeld module. *Amer. J. Math.*, 132(6):1493–1547, 2010.
- [6] N. Andruskiewitsch, I. Heckenberger, and L. Vendramin. Pointed hopf algebras of odd dimension and nichols algebras over solvable groups. *arXiv preprint 2411.02304*, 2024.
- [7] N. Andruskiewitsch and H.-J. Schneider. Lifting of quantum linear spaces and pointed Hopf algebras of order  $p^3$ . *J. Algebra*, 209:658–691, 1998.
- [8] N. Andruskiewitsch and H.-J. Schneider. Finite Quantum Groups and Cartan Matrices. *Adv. Math.*, 154(1):1–45, 2000.
- [9] N. Andruskiewitsch and H.-J. Schneider. Pointed Hopf algebras. *Math. Sci. Res. Inst. Publ.*, 43, 10 2001.
- [10] N. Andruskiewitsch and H.-J. Schneider. On the classification of finite-dimensional pointed Hopf algebras. *Ann. Math.*, 171(1):375–417, 2010.
- [11] I. Angiono. On Nichols algebras of diagonal type. *J. Reine Angew. Math.*, 2013(683):189–251, 2013.
- [12] I. Angiono. A presentation by generators and relations of Nichols algebras of diagonal type and convex orders on root systems. *J. Europ. Math. Soc.*, 17(10):2643–2671, 2015.
- [13] I. Angiono and A. G. Iglesias. Liftings of Nichols algebras of diagonal type II: All liftings are cocycle deformations. *Sel. Math., New Ser.*, 25(1):95, 2019.
- [14] I. Angiono and A. G. Iglesias. Pointed Hopf algebras: a guided tour to the liftings. *Rev. Colomb. Mat.*, 53:1–44, 2019.
- [15] M. Cuntz and I. Heckenberger. Weyl groupoids of rank two and continued fractions. *Algebra Number Theory*, 3(3):317–340, 2009.
- [16] M. Cuntz and I. Heckenberger. Weyl groupoids with at most three objects. *J. Pure Appl. Algebra*, 213(6):1112–1128, 2009.

- [17] M. Cuntz and I. Heckenberger. Finite Weyl groupoids of rank three. *Trans. Amer. Math. Soc.*, 364(3):1369–1393, 2012.
- [18] M. Cuntz and I. Heckenberger. Finite Weyl groupoids. *Reine Angew. Math.*, 2015(702):77–108, 2015.
- [19] I. Flandoli and S.D. Lentner. Algebras of non-local screenings and diagonal Nichols algebras. *SIGMA, Symmetry Integrability Geom. Methods Appl.*, 18:paper 018, 81, 2022.
- [20] A. G.Iglesias and J.M.J. Giralaldi. Liftings of Nichols algebras of diagonal type III. Cartan type  $G_2$ . *J. Algebra*, 478:506–568, 2017.
- [21] I. Heckenberger. Weyl equivalence for rank 2 Nichols algebras of diagonal type. *Ann. Uni. Ferrara*, 51(1):281–289, 2005.
- [22] I. Heckenberger. The Weyl groupoid of a Nichols algebra of diagonal type. *Invent. Math.*, 164(1):175–188, 2006.
- [23] I. Heckenberger. Classification of arithmetic root systems of rank 3. In *Actas del XVI coloquio Latinoamericano de álgebra.*, pages 227–252. Madrid: Revista Matemática Iberoamericana, 2007.
- [24] I. Heckenberger. Examples of finite-dimensional rank 2 Nichols algebras of diagonal type. *Compos. Math.*, 143(1):165–190, 2007.
- [25] I. Heckenberger. Rank 2 Nichols algebras with finite arithmetic root system. *Algebr. Represent. Theory*, 11(2):115–132, 2008.
- [26] I. Heckenberger. Classification of arithmetic root systems. *Adv. Math.*, 220(1):59–124, 2009.
- [27] I. Heckenberger, E. Meir, and L. Vendramin. Finite-dimensional nichols algebras of simple yetter–drinfeld modules (over groups) of prime dimension. *Advances in Mathematics*, 444:109637, 2024.
- [28] I. Heckenberger and H.-J. Schneider. Root systems and Weyl groupoids for Nichols algebras. *Proc. Lond. Math. Soc.*, 101(3):623–654, 2010.
- [29] I. Heckenberger and H.-J. Schneider. Right coideal subalgebras of Nichols algebras and the duflo order on the Weyl groupoid. *Israel J. Math.*, 197(1):139–187, 2013.
- [30] I. Heckenberger and H.-J. Schneider. *Hopf algebras and root systems*, volume 247. American Mathematical Soc., 2020.
- [31] I. Heckenberger and J. Wang. Rank 2 Nichols algebras of diagonal type over fields of positive characteristic. *SIGMA Symmetry. Integr. Geom.*, 11:011, 2015.
- [32] I. Heckenberger and H. Yamane. A generalization of Coxeter groups, root systems, and Matsumoto’s theorem. *Math. Z.*, 259(2):255–276, 2008.



- [33] J.E. Humphreys. *Introduction to Lie Algebras and Representation Theory*, volume 9 of *Graduate Texts in Mathematics*. Springer Science & Business Media, 1972.
- [34] V. Kharchenko. A quantum analog of the Poincare-Birkhoff-Witt theorem. *Algebra Logic*, 38(4):259–276, 1999.
- [35] S.D. Lentner. Quantum groups and nichols algebras acting on conformal field theories. *Advances in Mathematics*, 378:107517, 2021.
- [36] G. Lusztig. Noncommutative Differentials and Yang-Mills on Permutation Groups  $s_n$ . *Lect. Notes Pure Appl. Math.*, 239, 2001.
- [37] G. Lusztig. *Introduction to Quantum Groups*. Modern Birkhäuser Classics (MBC). Springer Science & Business Media, 1st ed. 1993, corr. 3rd printing edition, 2010.
- [38] W.D. Nichols. Bialgebras of type one. *Commun. Alg.*, 6(15):1521–1552, 1978.
- [39] M. Rosso. Quantum groups and quantum shuffles. *Invent. Math.*, 133(2):399–416, 1998.
- [40] A.M Semikhatov. Virasoro central charges for Nichols algebras. In *Conformal Field Theories and Tensor Categories*, pages 67–92. Springer, 2014.
- [41] A.M. Semikhatov and I.Yu. Tipunin. The Nichols algebra of screenings. *Commun. Contemp. Math.*, 14(04):1250029, 2012.
- [42] A.M. Semikhatov and I.Yu. Tipunin. Logarithmic  $sl_2$  CFT models from Nichols algebras: I. *J. Phys.*, 46(49):494011, 2013.
- [43] M. Takeuchi. Survey of braided hopf algebras. *Contemp. Math.*, 267:301–324, 2000.
- [44] J. Wang. Rank three Nichols algebras of diagonal type over arbitrary fields. *Israel J. Math.*, 218(1):1–26, 2017.
- [45] J. Wang. Rank 4 finite-dimensional Nichols algebras of diagonal type in positive characteristic. *J. Algebra*, 559:547–579, 2020.
- [46] S.L. Woronowicz. Compact matrix pseudogroups. *Comm. Math. Phys.*, 111(4):613–665, 1987.
- [47] S.L. Woronowicz. Differential calculus on compact matrix pseudogroups (quantum groups). *Comm. Math. Phys.*, 122(1):125–170, 1989.