

On Fundamental Theorems of Invariant Theory for the Special Linear Supergroup

J. Razzaq

*Mathematical Institute of Charles University.
Sokolovská 83, Prague, Czech Republic.
e-mail: junaid.razzaq@matfyz.cuni.cz*

R. Fiorese

*Fabit, Università di Bologna.
Via San Donato, 15. 40126 Bologna, Italy.
and INFN, Sezione di Bologna, Italy
e-mail: rita.fioresi@unibo.it*

M. A. Lledó

*Departament de Física Teòrica, Universitat de València and
IFIC (CSIC-UVEG).
C/Dr. Moliner, 50, E-46100 Burjassot (València), Spain.
e-mail: maria.lledo@ific.uv.es*

Abstract

The purpose of this paper is to prove the First and Second Fundamental Theorems of invariant theory for the complex special linear supergroup and discuss the superalgebra of invariants, via the super Plücker relations.

1 Introduction

Since its origins, in the XIXth century [6], classical invariant theory rapidly provided important results, whose applications involve both, mathematics and physics (see [3] for an historical review of the theory).

One of the classical invariant theory main questions, which led to the spectacular result by Hilbert [16], is the search for invariants, with respect to a given group action, in the ring of polynomial functions ([22, 23]).

We can look, for example, at the polynomial functions on the set of $r \times p$ ($r \leq p$) complex matrices $M_{r \times p}$, that are invariant under the action of the complex special linear group $SL_r(\mathbb{C})$:

$$\begin{aligned} \mathbb{C}[M_{r \times p}] \times SL_r(\mathbb{C}) &\longrightarrow \mathbb{C}[M_{r \times p}] \\ (f, g) &\longrightarrow f \cdot g, \end{aligned}$$

where $(f \cdot g)(M) := f(gM)$ and $\mathbb{C}[M_{r \times p}]$ denotes the algebra of polynomials functions on the entries of $M_{r \times p}$.

Let $X_{i_1 \dots i_r}$ denote the $r \times r$ minor formed with the columns (i_1, \dots, i_r) of a matrix in $M_{r \times p}$. The First and Second fundamental Theorems of invariant theory for $SL_r(\mathbb{C})$ state the following (see [13, 12, 11] and Refs. therein for a full account).

Theorem 1.1. *Let the notation be as above.*

1. **First Fundamental Theorem (FFT) of invariant theory.** *The ring of invariants $\mathbb{C}[M_{r \times p}]^{SL_r(\mathbb{C})}$ is generated by the minors $X_{i_1 \dots i_r}$ of the $r \times r$ submatrices in $M_{r \times p}$.*
2. **Second Fundamental Theorem (SFT) of invariant theory.** *We have a presentation of the ring of invariants via the ideal I of the Plücker relations:*

$$\begin{aligned} \mathbb{C}[M_{r \times p}]^{SL_r(\mathbb{C})} &\cong \mathbb{C}[X_{i_1 \dots i_r}] / I, \quad \text{with} \\ I &:= \left(\sum_{k=1}^{r+1} (-1)^k X_{i_1 \dots i_{r-1} j_k} X_{j_1 \dots \tilde{j}_k \dots j_{r+1}} \right) \end{aligned} \quad (1)$$

where

$$\begin{aligned} 1 &\leq i_1 < \dots < i_{r-1} \leq p \\ 1 &\leq j_1 < \dots < \tilde{j}_k < \dots < j_{r+1} \leq p \end{aligned}$$

and \tilde{j}_k means that the index is removed.

In this work we want to provide a generalization of this result to super-geometry. The generalization of some questions of invariant theory to the super setting appeared early in the literature: for example, the Young super tableaux and techniques regarding their manipulation, appeared first in [18]

and at the same time in [1], to address questions of representation theory of Lie superalgebras and the double commutant theorem, a fundamental result, originally proven, in the ordinary setting, by Schur [22].

In particular, in [1], the authors prove also a version of the straightening algorithm for Young super tableaux, of uttermost importance in representation theory and directly linked to the Plücker relations [11]. Later on, in the papers [15, 4], appeared such algorithm in the super setting. Their idea, involving the so called *virtual variables method*, provides an elegant approach which encompasses both, the ordinary and the super setting at once. More recently in [24, 9] appeared generalizations to other supergroups as the orthosymplectic one and in [7] a categorical approach to super invariant theory allowed to go beyond the characteristic zero setting (see also [8]).

In the present work, however, we are interested in a related, and yet different topic. It has its significance rooted into the geometric meaning of these combinatorial questions. In the ordinary setting Theorem 1.1 provides with a presentation of the ring of invariants via the Plücker relations, which give an embedding of the Grassmannian manifold into a suitable projective space. In supergeometry, super Grassmannians do not admit, in general, projective embeddings (see [19], Chapter 4). However, in [20], the authors define a new graded version of super projective space, with negative grading, which allows for such embedding. Indeed, the Berezinians of sub-supermatrices replace truly the notion of minors, which give the projective coordinates for the classical Plücker embedding. Then, they allow to proceed and give the correct supergeometric counterpart of the Plücker embedding and Plücker relations [20]. Furthermore, in [21], these super Plücker relations has been studied from the perspective of constructing super cluster algebras. However, due to the very nature of Berezinians, which are defined only when the supermatrix is invertible, one is forced to restrict the analysis to an open set. Once this geometric constraint is set in place, one can state and prove the super version of the First Fundamental Theorem and progress towards the full proof of the more elusive Second Fundamental Theorem.

The paper is organized as follows.

In Section 2, we introduce the notation and a few facts concerning the Berezinians. We have included in Appendix A some very basic notions of supergeometry. We also state the super Cramer rule, first proved in [2] with different methods, which is instrumental for the proofs of the results in the next sections. Due to its importance to the present work, and in order to

be self contained, we have included our new proof of super Cramer's rule in Appendix B, which is very much in line with the reasoning of the rest of the paper.

In Section 3, we state and prove the *First Fundamental Theorem of super invariant theory* for the special linear supergroup, Theorem 3.1.

In Section 4, we introduce and prove a super version of the Jacobi identity (a classical determinant identity, for details see Appendix C), Theorem 4.1, one of our main results, which is instrumental for the following sections.

In Sections 5, 6 we derive the *super Plücker relations* from the super Jacobi identity in a special, yet illustrative, case (i.e. of $SL(1|1)$). We also prove *Second Fundamental Theorem of super invariant theory* for $SL(1|1)$.

In Section 7 we derive the *super Plücker relations* for the case of $SL(r|s)$ and conjectured *Second Fundamental Theorem of super invariant theory* for $SL(r|s)$. It is important to note that the super Plücker relations we derived are same as the one given in [20], however, we used a different construction based on super Jacobi identity.

2 Preliminaries

In this section we establish our notation on supermatrices. The reader can go to Appendix A or to references [5, 10, 20] for terminology.

Let A be a fixed commutative superalgebra. All entries of supermatrices are intended to be in A . An *even supermatrix*

$$B = \left(\begin{array}{c|c} B_1 & B_2 \\ \hline B_3 & B_4 \end{array} \right) \quad (2)$$

of size $p|q \times r|s$ is a block matrix whose first r columns are even vectors and last s columns are odd vectors. Consequently, B_1 and B_4 contain even entries, while B_2 and B_3 contain odd entries.

We also need to introduce the concept of *fake supermatrices*¹.

Definition 2.1. A *fake supermatrix of type-I* is a matrix obtained by inserting an odd vector at an even vector's position of an even supermatrix. A *fake supermatrix of type-II* is a matrix obtained by inserting an even vector at an odd vector's position of an even supermatrix.

¹These are called 'wrong supermatrices' in [20].

As usual, for any even supermatrix or fake supermatrix, as in (2), we define its *parity reversed matrix*, denoted by B^Π by swapping B_1 with B_4 and B_2 with B_3 , i.e.

$$\left(\begin{array}{c|c} B_1 & B_2 \\ \hline B_3 & B_4 \end{array} \right)^\Pi = \left(\begin{array}{c|c} B_4 & B_3 \\ \hline B_2 & B_1 \end{array} \right).$$

Notice that if B is a fake supermatrix of type I, then B^Π is a fake supermatrix of type II and viceversa.

Definition 2.2. Let B be an even supermatrix or a fake supermatrix of type-I:

$$B = \left(\begin{array}{c|c} B_1 & B_2 \\ \hline B_3 & B_4 \end{array} \right). \quad (3)$$

Assume that B_4 is invertible. We define the *Berezinian* of B as:

$$\text{Ber} B := \det B_4^{-1} \det(B_1 - B_2 B_4^{-1} B_3)$$

Notice that, although this definition is the same as the first line of (47), in Appendix A, here we use it in a more general context, when B can be a fake supermatrix of type-I.

Definition 2.3. Let B be an even supermatrix or a fake supermatrix of type-II:

$$B = \left(\begin{array}{c|c} B_1 & B_2 \\ \hline B_3 & B_4 \end{array} \right) \quad (4)$$

Assume that B_1 is invertible. We define the *inverted berezinian* of B :

$$\text{Ber}^* B := \text{Ber} B^\Pi.$$

It is clear that for an invertible supermatrix B :

$$\text{Ber}^* B = \text{Ber} B^{-1}. \quad (5)$$

Remark 2.4. For an even supermatrix B , we denote with i the even indices and with \hat{i} the odd ones, as we exemplify below:

$$B = \left(\begin{array}{ccc|ccc} b_{11} & \dots & b_{1p} & b_{1\hat{1}} & \dots & b_{1\hat{q}} \\ \cdot & \dots & \cdot & \cdot & \dots & \cdot \\ \cdot & \dots & \cdot & \cdot & \dots & \cdot \\ b_{r1} & \dots & b_{rp} & b_{r\hat{1}} & \dots & b_{r\hat{q}} \\ \hline b_{\hat{1}1} & \dots & b_{\hat{1}p} & b_{\hat{1}\hat{1}} & \dots & b_{\hat{1}\hat{q}} \\ \cdot & \dots & \cdot & \cdot & \dots & \cdot \\ \cdot & \dots & \cdot & \cdot & \dots & \cdot \\ b_{\hat{s}1} & \dots & b_{\hat{s}p} & b_{\hat{s}\hat{1}} & \dots & b_{\hat{s}\hat{q}} \end{array} \right).$$

We state now the super version of Cramer's rule. Recall that, in ordinary linear algebra, for an invertible matrix T of size $n \times n$ and a column vector \mathbf{b} of size n , the solution to the equation:

$$T\mathbf{x} = \mathbf{b}$$

is a column vector \mathbf{x} whose entries are given by:

$$x_i = \frac{\det T_i(\mathbf{b})}{\det T}$$

where $T_i(\mathbf{b})$ denotes the matrix obtained by replacing the i -th column of T with \mathbf{b} .

In super linear algebra, we have the following theorem:

Theorem 2.5. *Super Cramer's rule. Let $M \in \text{GL}(r|s)$ be an invertible even supermatrix and let $\mathbf{b} \in \mathbf{k}^{r|s}$ be an even vector of size $r|s$. Then, the solution to the equation*

$$M\mathbf{x} = \mathbf{b}$$

is given by:

$$x_i = \frac{\text{Ber} M_i(\mathbf{b})}{\text{Ber} M} \quad \text{for } i = 1, \dots, r$$

$$x_{\hat{j}} = \frac{\text{Ber}^* M_{\hat{j}}(\mathbf{b})}{\text{Ber}^* M} \quad \text{for } \hat{j} = \hat{1} \cdots \hat{s}$$

where $M_i(\mathbf{b})$ is the supermatrix obtained by replacing i -th even column of M with \mathbf{b} and $M_{\hat{j}}(\mathbf{b})$ is the fake supermatrix of type-II obtained by replacing \hat{j} -th odd column of M with \mathbf{b} .

This is a known result, see [2]. However, for completeness, we have included in Appendix B a proof of it, obtained with different, elementary methods.

As stated in Remark B.1, the same expressions hold for the equation $M\mathbf{x} = \mathbf{b}$ if \mathbf{b} is an odd vector of size $r|s$.

Example 2.6. Let $g = \left(\begin{array}{c|c} a & \alpha \\ \hline \beta & b \end{array} \right) \in \text{GL}(1|1)$. Suppose we need to find $\begin{pmatrix} x \\ \eta \end{pmatrix}$ such that:

$$g \begin{pmatrix} x \\ \eta \end{pmatrix} = \begin{pmatrix} c \\ \gamma \end{pmatrix}.$$

Then, using super Cramer's rule, one gets:

$$x = \frac{\text{Ber} \left(\begin{array}{c|c} c & \alpha \\ \hline \gamma & b \end{array} \right)}{\text{Ber} \left(\begin{array}{c|c} a & \alpha \\ \hline \beta & b \end{array} \right)} = \frac{c - \alpha b^{-1} \gamma}{a - \alpha b^{-1} \beta}$$

and

$$\eta = \frac{\text{Ber}^* \left(\begin{array}{c|c} a & c \\ \hline \beta & \gamma \end{array} \right)}{\text{Ber}^* \left(\begin{array}{c|c} a & \alpha \\ \hline \beta & b \end{array} \right)} = \frac{\gamma - \beta a^{-1} c}{b - \beta a^{-1} \alpha}.$$

3 First Fundamental Theorem of invariant theory for $\text{SL}(r|s)$

In ordinary geometry, we have a natural right action of the complex special linear group $\text{SL}_r(\mathbb{C})$ on the algebra of functions $\mathbb{C}[\text{M}_{r \times p}]$ of the $r \times p$ complex matrices $\text{M}_{r \times p}$:

$$\begin{aligned} \mathbb{C}[\text{M}_{r \times p}] \times \text{SL}_r(\mathbb{C}) &\longrightarrow \mathbb{C}[\text{M}_{r \times p}] \\ (f, g) &\longrightarrow f \cdot g, \end{aligned} \tag{6}$$

where $(f \cdot g)(M) := f(gM)$, for all $M \in \text{M}_{r \times p}(\mathbb{C})$.

The First Fundamental Theorem of ordinary invariant theory realizes the ring of invariants with respect to this action, as the subring of $\mathbb{C}[\text{M}_{r \times p}]$ generated by the determinants of the $r \times r$ minors in $\text{M}_{r \times p}(\mathbb{C})$.

We now extend the action (6) to the super setting and compute the superalgebra of invariants. Let $\text{M}_{r|s \times p|q}$ be the super vector space of supermatrices and let \mathcal{O} be the superalgebra of 'functions' on $\text{M}_{r|s \times p|q}$ (see Observation A.1). For notational purposes, let us organize the generators of \mathcal{O} (or 'coordinates') in a supermatrix, as follows:

$$A = \left(\begin{array}{ccc|ccc} x_{11} & \dots & x_{1p} & \alpha_{1\hat{1}} & \dots & \alpha_{1\hat{q}} \\ \cdot & \dots & \cdot & \cdot & \dots & \cdot \\ \cdot & \dots & \cdot & \cdot & \dots & \cdot \\ x_{r1} & \dots & x_{rp} & \alpha_{r\hat{1}} & \dots & \alpha_{r\hat{q}} \\ \hline \beta_{\hat{1}1} & \dots & \beta_{\hat{1}p} & y_{\hat{1}\hat{1}} & \dots & y_{\hat{1}\hat{q}} \\ \cdot & \dots & \cdot & \cdot & \dots & \cdot \\ \cdot & \dots & \cdot & \cdot & \dots & \cdot \\ \beta_{\hat{s}1} & \dots & \beta_{\hat{s}p} & y_{\hat{s}\hat{1}} & \dots & y_{\hat{s}\hat{q}} \end{array} \right), \quad (7)$$

where the hat denotes the odd indices. Then,

$$\mathcal{O} = k[x_{ij}, y_{\hat{k}\hat{l}}, \alpha_{i\hat{k}}, \beta_{\hat{l}j}].$$

Let \mathcal{A} be a commutative superalgebra. For every supermatrix in the set of \mathcal{A} -points of $M_{r|s \times p|q}$ (see Appendix A), $M \in M_{r|s \times p|q}(\mathcal{A})$ and $f \in \mathcal{O}$ one can compute an element of \mathcal{A} . This is denoted, in the terminology of the \mathcal{A} -points, as $f(M)$ instead of the, perhaps more appropriate, $M(f)$.

Let $SL(r|s)$ be the special linear supergroup (see, for example, [5], Chapter 1). Its \mathcal{A} -points are the matrices in $GL(r|s)(\mathcal{A})$ with Berezinian equal to 1. In terms of \mathcal{A} -points, one can write a left action of the special linear supergroup on the supermatrices:

$$\begin{aligned} SL(r|s)(\mathcal{A}) \times M_{r|s \times p|q}(\mathcal{A}) &\longrightarrow M_{r|s \times p|q}(\mathcal{A}) \\ (g, M) &\longrightarrow gM. \end{aligned}$$

One can also define a right action of $SL(r|s)$ on \mathcal{O} , exactly as in the ordinary case (6). For $g \in SL(r|s)(\mathcal{A})$ and $f \in \mathcal{O}$ one has

$$(f \cdot g)(M) := f(gM), \quad M \in M_{r|s \times p|q}(\mathcal{A}).$$

We want to be able to define the Berezinians of all $r|s \times r|s$ minors appearing in the matrix (7). We then have to localize \mathcal{O} at the determinants

$$D_{i_1, \dots, i_r} := \det \begin{pmatrix} x_{1i_1} & \dots & x_{1i_r} \\ \cdot & \dots & \cdot \\ \cdot & \dots & \cdot \\ x_{ri_1} & \dots & x_{ri_r} \end{pmatrix}, \quad D_{\hat{j}_1, \dots, \hat{j}_s} := \det \begin{pmatrix} y_{\hat{1}\hat{j}_1} & \dots & y_{\hat{1}\hat{j}_s} \\ \cdot & \dots & \cdot \\ \cdot & \dots & \cdot \\ y_{\hat{s}\hat{j}_1} & \dots & y_{\hat{s}\hat{j}_s} \end{pmatrix},$$

in the diagonal blocks, in the expression (7). We can now look at the action of the supergroup $\mathrm{SL}(r|s)$ on the localization

$$\tilde{\mathcal{O}} := \mathcal{O}[D_{i_1 \dots i_r}^{-1}, D_{\hat{j}_1 \dots \hat{j}_s}^{-1}]. \quad (8)$$

This corresponds to take the subset $\tilde{M}_{r|s \times p|q}(\mathcal{A}) \subset M_{r|s \times p|q}(\mathcal{A})$ where all the corresponding minors are invertible.

In this way we can formulate the super version of the First Fundamental Theorem of invariant theory. As we shall see in the next section, it is related with the Grassmannian and its super Plücker embedding, and we adopted a very similar notation for generators as in [20].

Theorem 3.1. *Let $A_{i_1 \dots i_r | \hat{j}_1 \dots \hat{j}_s}$ be the supermatrix formed with the columns $(i_1, \dots, i_r | \hat{j}_1, \dots, \hat{j}_s)$ of the supermatrix A in (7). Then, the ring of invariants $\tilde{\mathcal{O}}^{\mathrm{SL}(r|s)}$ is generated by the super minors*

$$\begin{aligned} X_{i_1 \dots i_r | \hat{j}_1 \dots \hat{j}_s} &:= \mathrm{Ber} A_{i_1 \dots i_r | \hat{j}_1 \dots \hat{j}_s} \\ X_{i_1 \dots i_r | \hat{j}_1 \dots \hat{j}_s}^* &:= \mathrm{Ber}^* A_{i_1 \dots i_r | \hat{j}_1 \dots \hat{j}_s} \end{aligned} \quad (9)$$

and the ‘fake’ super minors,

$$X_{i_1 \dots i_{k-1} \hat{j}_k i_{k+1} \dots i_r | \hat{j}_1 \dots \hat{j}_s} := \mathrm{Ber} A_{i_1 \dots i_{k-1} \hat{j}_k i_{k+1} \dots i_r | \hat{j}_1 \dots \hat{j}_s} \quad (10)$$

$$X_{i_1 \dots i_r | \hat{j}_1 \dots \hat{j}_{l-1} i_l \hat{j}_{l+1} \dots \hat{j}_s} := \mathrm{Ber}^* A_{i_1 \dots i_r | \hat{j}_1 \dots \hat{j}_{l-1} i_l \hat{j}_{l+1} \dots \hat{j}_s} \quad (11)$$

where $1 \leq i_1 \leq \dots \leq i_r \leq p$ and $1 \leq \hat{j}_1 \leq \dots \leq \hat{j}_s \leq q$.

Proof. Firstly, it is clear, from the multiplicative property of Berezinian, that super-minors and fake super-minors are invariants.

To show that they generate the ring $\tilde{\mathcal{O}}^{\mathrm{SL}(r|s)}$, we decompose the matrix of coordinate functions (7) as $A = \tilde{A}_{1 \dots r | \hat{1} \dots \hat{s}} B$, where

$$\tilde{A}_{1\dots r|\hat{1}\dots\hat{s}} = \left(\begin{array}{ccc|ccc} \frac{x_{11}}{X_{1\dots r|\hat{1}\dots\hat{s}}} & \cdots & x_{1r} & \alpha_{1\hat{1}} & \cdots & \alpha_{1\hat{s}} \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ \frac{x_{r1}}{X_{1\dots r|\hat{1}\dots\hat{s}}} & \cdots & x_{rr} & \alpha_{r\hat{1}} & \cdots & \alpha_{r\hat{s}} \\ \hline \frac{\beta_{\hat{1}1}}{X_{1\dots r|\hat{1}\dots\hat{s}}} & \cdots & \beta_{\hat{1}r} & y_{\hat{1}\hat{1}} & \cdots & y_{\hat{1}\hat{s}} \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ \frac{\beta_{\hat{s}1}}{X_{1\dots r|\hat{1}\dots\hat{s}}} & \cdots & \beta_{\hat{s}r} & y_{\hat{s}\hat{1}} & \cdots & y_{\hat{s}\hat{s}} \end{array} \right).$$

Therefore, we have,

$$B = \tilde{A}_{1\dots r|\hat{1}\dots\hat{s}}^{-1} A.$$

Notice that, we are working over $\mathcal{O}[D_{i_1\dots i_r}^{-1}, D_{\hat{j}_1\dots\hat{j}_s}^{-1}]$, hence, $\tilde{A}_{1\dots r|\hat{1}\dots\hat{s}}^{-1}$ is well-defined. Using super Cramer's rule we can find the entries of $B = \left(\begin{array}{c|c} U & V \\ \hline W & Z \end{array} \right)$:

$$U_{ij} = \begin{cases} X_{j\dots r|\hat{1}\dots\hat{s}} & \text{for } i = 1, 1 \leq j \leq p, \\ X_{1\dots r|\hat{1}\dots\hat{s}}^* X_{1\dots(i-1)j(i+1)\dots r|\hat{1}\dots\hat{s}} & \text{for } 1 < i \leq r, 1 \leq j \leq p \end{cases} \quad (12)$$

$$W_{\hat{k}j} = X_{1\dots r|\hat{1}\dots\hat{s}} X_{1\dots r|\hat{1}\dots(\hat{k}-\hat{1})j(\hat{k}+\hat{1})\dots\hat{s}}^* \quad \text{for } 1 \leq k \leq s, 1 \leq j \leq p. \quad (13)$$

$$V_{i\hat{l}} = \begin{cases} X_{\hat{l}\dots r|\hat{1}\dots\hat{s}} & \text{for } i = 1, 1 \leq l \leq q, \\ X_{1\dots r|\hat{1}\dots\hat{s}}^* X_{1\dots(i-1)\hat{l}(i+1)\dots r|\hat{1}\dots\hat{s}} & \text{for } 1 < i \leq r, 1 \leq l \leq q. \end{cases} \quad (14)$$

$$Z_{\hat{k}\hat{l}} = X_{1\dots r|\hat{1}\dots\hat{s}} X_{1\dots r|\hat{1}\dots(\hat{k}-\hat{1})\hat{l}(\hat{k}+\hat{1})\dots\hat{s}}^* \quad \text{for } 1 \leq k \leq s, 1 \leq l \leq q. \quad (15)$$

Let \mathcal{A} be any superalgebra and consider an arbitrary element $T \in \tilde{M}_{r|s \times p|q}(\mathcal{A})$. We can write T as a product,

$$T = \tilde{T}_{1\dots r|\hat{1}\dots\hat{s}} R \quad (16)$$

where, $\tilde{T}_{1\dots r|\hat{1}\dots\hat{s}}$ is an element of $\mathrm{SL}(r|s)(\mathcal{A})$ and $R \in \tilde{M}_{r|s \times p|q}(\mathcal{A})$ is computed in terms of superminors and fake superminors. Explicitly, each entry of R can be computed by evaluating on T the corresponding superminor or fake superminor as in B above.

Let $f \in \tilde{\mathcal{O}}^{\mathrm{SL}(r|s)}$ be an invariant. Then, by definition, we have:

$$f.g(R) = f(gR) = f(R), \quad \text{for all } R \in \tilde{M}_{r|s \times p|q}(\mathcal{A}) \text{ and } g \in \mathrm{SL}(r|s)(\mathcal{A}).$$

But then, for any $T \in \tilde{M}_{r|s \times p|q}(\mathcal{A})$,

$$f(T) = f(\tilde{T}_{1\dots r|\hat{1}\dots\hat{s}}R) = f(R),$$

where R is as in (16). Therefore, f is a function of superminors and fake superminors, as we wanted to prove. \square

4 The super Jacobi identity

In this section we want to obtain a super Jacobi identity, a generalization of the Jacobi identity, Theorem C.1 in the Appendix C, to the super setting. We first introduce some notation.

Notation. Let A be an invertible supermatrix of size $p|q \times p|q$. Let $u = (p - r + 1, \dots, p|\hat{q} - \hat{s} + \hat{1}, \dots, \hat{q})$ and $v = (1, \dots, r|\hat{1}, \dots, \hat{s})$ be two sets of indices, $r < p$, $\hat{s} < \hat{q}$. Let A_v^u denote the matrix obtained from A by deleting the rows and columns contained in u and v , respectively, *i.e.* by deleting the last r even rows, the last \hat{s} odd rows, first r even columns and first \hat{s} odd columns. Also, set $\tilde{u} = (1, \dots, p - r|\hat{1}, \dots, \hat{q} - \hat{s})$ and $\tilde{v} = (r + 1, \dots, p|\hat{s} + \hat{1}, \dots, \hat{q})$ as the complements of u and v , respectively.

Theorem 4.1. *The super Jacobi Identity. Let the notation be as above. If A_v^u and $(A^{-1})_{\tilde{u}}^{\tilde{v}}$ are invertible, then*

$$\mathrm{Ber} A \mathrm{Ber} (A^{-1})_{\tilde{u}}^{\tilde{v}} = (-1)^t \mathrm{Ber} A_v^u \quad (17)$$

where $t = r(p + 1) + s(q + 1)$.

We call (17) the *super Jacobi identity*.

Proof. A similar strategy to what we used in the proof of Theorem C.1 (see Appendix C) will work here. Let

$$A = \left(\begin{array}{c|c} [a_{ij}] & [a_{i\hat{k}}] \\ \hline [a_{\hat{l}j}] & [a_{\hat{l}\hat{k}}] \end{array} \right) \quad \text{and} \quad A^{-1} = \left(\begin{array}{c|c} [b_{ij}] & [b_{i\hat{k}}] \\ \hline [b_{\hat{l}j}] & [b_{\hat{l}\hat{k}}] \end{array} \right).$$

According to our notation,

$$(A^{-1})_{\tilde{u}}^{\tilde{v}} = \left(\begin{array}{ccc|ccc} b_{1(p-r+1)} & \cdots & b_{1p} & b_{1(\hat{q}-\hat{s}+\hat{1})} & \cdots & b_{1\hat{q}} \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ \hline b_{r(p-r+1)} & \cdots & b_{rp} & b_{r(\hat{q}-\hat{s}+\hat{1})} & \cdots & b_{r\hat{q}} \\ b_{\hat{1}(p-r+1)} & \cdots & b_{\hat{1}p} & b_{\hat{1}(\hat{q}-\hat{s}+\hat{1})} & \cdots & b_{\hat{1}\hat{q}} \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ \hline b_{\hat{s}(p-r+1)} & \cdots & b_{\hat{s}p} & b_{\hat{s}(\hat{q}-\hat{s}+\hat{1})} & \cdots & b_{\hat{s}\hat{q}} \end{array} \right)$$

One can see that

$$\text{Ber}(A^{-1})_{\tilde{u}}^{\tilde{v}} = \text{Ber}T \quad (18)$$

where T is the super matrix of size $p|q \times p|q$ described as follows:

$$T_j = \begin{cases} (A^{-1})_{p-r+j} & \text{for } 1 \leq j \leq r \\ e_j & \text{for } r+1 \leq j \leq p \end{cases}$$

where T_j denotes the j -th even column of T , $(A^{-1})_k$ is the k -th even column of A^{-1} and e_j denotes an even vector of size $p|q$, whose entries are all 0 except at the j -th even position, where the entry is 1. On the other hand,

$$T_{\hat{l}} = \begin{cases} (A^{-1})_{\hat{q}-\hat{s}+\hat{l}} & \text{for } \hat{1} \leq \hat{l} \leq \hat{s} \\ e_{\hat{l}} & \text{for } \hat{s} + \hat{1} \leq \hat{l} \leq \hat{q}, \end{cases}$$

where $T_{\hat{l}}$ denotes the \hat{l} -th odd column of T , $(A^{-1})_{\hat{k}}$ is the \hat{k} -th-odd column of A^{-1} and $e_{\hat{l}}$ denotes an odd vector of size $p|q$ whose entries are all 0 except at the \hat{l} -th odd position, where the entry is 1.

In other words, T is a supermatrix whose first r even columns coincide with the last r even columns of A^{-1} , the first \hat{s} odd columns coincide with the last \hat{s} odd columns of A^{-1} and the remaining even and odd columns are e_j and $e_{\hat{l}}$, respectively.

Therefore, using Equation (18) and the multiplicative property of Berezinian, we have:

$$\text{Ber} A \text{Ber}(A^{-1})_{\tilde{u}}^{\tilde{v}} = \text{Ber} A \text{Ber} T = \text{Ber}(AT). \quad (19)$$

Let us denote $C = AT$; then, using above notation:

$$C_j = \begin{cases} e_{p-r+j} & \text{for } 1 \leq j \leq r, \\ A_j & \text{for } r+1 \leq j \leq p, \end{cases}$$

and

$$C_{\hat{l}} = \begin{cases} e_{\hat{q}-\hat{s}+\hat{l}} & \text{for } \hat{1} \leq \hat{l} \leq \hat{s} \\ A_{\hat{l}} & \text{for } \hat{s} + \hat{1} \leq \hat{l} \leq \hat{q}. \end{cases}$$

Then, one can compute:

$$\text{Ber}(AT) = \text{Ber} C = (-1)^{r(p+1)+s(q+1)} \text{Ber} A_v^u \quad (20)$$

and by putting this into Equation (19) the desired identity is obtained. \square

5 The Super Plücker Relations for $\text{SL}(1|1)$

We want to use the super Jacobi identity as in Theorem (4.1) to derive the relations among the generators in the superalgebra $\tilde{\mathcal{O}}$ defined in (8). This approach for the construction of classical Plücker relations has been explained in the Appendix C, Proposition C.2.

We illustrate in detail in this section the case of $r = s = 1$, the general case will be discussed shortly, in the next section, following the same strategy.

Theorem 5.1. *The super Plücker relations for $\text{SL}(1|1)$. Let the notation be*

as in (9), (10) and (11) for $r = s = 1$. Then:

$$X_{i|\hat{\mu}}X_{i|\hat{\mu}}^* = 1, \quad (21)$$

$$X_{i|\hat{\mu}}X_{j|\hat{\nu}} = \text{Ber} \left(\frac{X_{j|\hat{\mu}}}{X_{i|j}^*} \middle| \frac{X_{\hat{\nu}|\hat{\mu}}}{X_{i|\hat{\nu}}^*} \right), \quad (22)$$

$$X_{i|\hat{\mu}}X_{\hat{\lambda}|\hat{\nu}} = \text{Ber} \left(\frac{X_{\hat{\lambda}|\hat{\mu}}}{X_{i|\hat{\lambda}}^*} \middle| \frac{X_{\hat{\nu}|\hat{\mu}}}{X_{i|\hat{\nu}}^*} \right), \quad (23)$$

$$X_{i|\hat{\mu}}^*X_{j|k}^* = \text{Ber}^* \left(\frac{X_{j|\hat{\mu}}}{X_{i|j}^*} \middle| \frac{X_{k|\hat{\mu}}}{X_{i|k}^*} \right), \quad (24)$$

where the Latin (even) indices run from 1 to p and the Greek (odd) indices run from 1 to q .

We call these the *super Plücker relations* for $\text{SL}(1|1)$. Moreover, it is important to note that these super Plücker relations coincides with the super Plücker relations appeared in [20, Th. 8] obtained with a different approach.

Proof. The relations (21) follow from the Equation (5).

Now, for any indices $(i|\hat{\mu})$ and $(j|\hat{\nu})$, fix the following matrix:

$$B = \left(\begin{array}{cc|cc} x_{1i} & x_{1j} & x_{1\hat{\mu}} & x_{1\hat{\nu}} \\ 0 & 1 & 0 & 0 \\ \hline x_{\hat{1}i} & x_{\hat{1}j} & x_{\hat{1}\hat{\mu}} & x_{\hat{1}\hat{\nu}} \\ 0 & 0 & 0 & 1 \end{array} \right).$$

Clearly,

$$\text{Ber} B = X_{i|\hat{\mu}} \quad (25)$$

By setting $u = (2, \hat{2})$ and $v = (1, \hat{1})$ we get:

$$\text{Ber} B_v^u = \text{Ber} \left(\frac{x_{1j}}{x_{\hat{1}j}} \middle| \frac{x_{1\hat{\nu}}}{x_{\hat{1}\hat{\nu}}} \right) = X_{j|\hat{\nu}}. \quad (26)$$

Moreover, since $\tilde{u} = (1|\hat{1})$ and $\tilde{v} = (2|\hat{2})$, using the super Cramer rule one can compute:

$$(B^{-1})_{\tilde{u}}^{\tilde{v}} = \left(\frac{(B^{-1})_{12}}{(B^{-1})_{\hat{1}2}} \middle| \frac{(B^{-1})_{1\hat{2}}}{(B^{-1})_{\hat{1}\hat{2}}} \right) = \left(\frac{-X_{j|\hat{\mu}}X_{i|\hat{\mu}}^*}{-X_{i|j}^*X_{i|\hat{\mu}}} \middle| \frac{-X_{\hat{\nu}|\hat{\mu}}X_{i|\hat{\mu}}^*}{-X_{i|\hat{\nu}}^*X_{i|\hat{\mu}}} \right).$$

Hence,

$$\text{Ber}(B^{-1})_{\hat{u}}^{\hat{v}} = (X_{i|\hat{\mu}}^*)^2 \text{Ber} \left(\begin{array}{c|c} X_{j|\hat{\mu}} & X_{\hat{\nu}|\hat{\mu}} \\ \hline X_{i|j}^* & X_{i|\hat{\nu}}^* \end{array} \right). \quad (27)$$

Substituting Equations (25), (26) and (27) in the super Jacobi identity, one arrives to the relations (22).

For the other relations, we would need to generalize the super Jacobi identity to include some fake supermatrices. We instead use an argument, similar to the one used in the proof of the super Jacobi identity. Consider the matrix

$$B = \left(\begin{array}{cc|cc} x_{1i} & 0 & x_{1\hat{\mu}} & x_{1\hat{\nu}} \\ 0 & 1 & 0 & 0 \\ \hline x_{\hat{1}i} & 0 & x_{\hat{1}\hat{\mu}} & x_{\hat{1}\hat{\nu}} \\ 0 & 0 & 0 & 1 \end{array} \right) \quad (28)$$

with $\text{Ber} B = X_{i|\hat{\mu}}$. We construct a fake supermatrix T in the following way: the first even column is the second even column of B^{-1} , the first odd column is the second odd column of B^{-1} ; at the second even column position we have $B^{-1}t$ where

$$t := \left(\begin{array}{c} x_{1\hat{\lambda}} \\ 0 \\ \hline x_{\hat{1}\hat{\lambda}} \\ 0 \end{array} \right).$$

and at the second odd column position we have $e_{\hat{2}}$. Then, clearly:

$$\text{Ber}(BT) = \text{Ber} \left(\begin{array}{cc|cc} 0 & x_{1\hat{\lambda}} & 0 & x_{1\hat{\nu}} \\ 1 & 0 & 0 & 0 \\ \hline 0 & x_{\hat{1}\hat{\lambda}} & 0 & x_{\hat{1}\hat{\nu}} \\ 0 & 0 & 1 & 1 \end{array} \right) = X_{\hat{\lambda}|\hat{\nu}}. \quad (29)$$

On the other hand, one can explicitly compute, using super Cramer's rule:

$$T = \left(\begin{array}{cc|cc} 0 & X_{\hat{\lambda}|\hat{\mu}} X_{i|\hat{\mu}}^* & -X_{\hat{\nu}|\hat{\mu}} X_{i|\hat{\mu}}^* & 0 \\ 1 & 0 & 0 & 0 \\ \hline 0 & X_{i|\hat{\lambda}}^* X_{i|\hat{\mu}} & -X_{i|\hat{\nu}}^* X_{i|\hat{\mu}} & 0 \\ 0 & 0 & 1 & 1 \end{array} \right) \quad (30)$$

Therefore,

$$\text{Ber}T = (X_{i|\hat{\mu}}^*)^2 \text{Ber} \left(\begin{array}{c|c} X_{\hat{\lambda}|\hat{\mu}} & X_{\hat{\nu}|\hat{\mu}} \\ \hline X_{i|\hat{\lambda}}^* & X_{i|\hat{\nu}}^* \end{array} \right). \quad (31)$$

Using the multiplicative property of the Berezinian and equation (29) one arrives to the relations (23).

Similarly, the relations (24) follow. Consider the supermatrix

$$B = \left(\begin{array}{cc|cc} x_{\hat{1}\hat{\mu}} & 0 & x_{\hat{1}i} & x_{\hat{1}j} \\ 0 & 1 & 0 & 0 \\ \hline x_{1\hat{\mu}} & 0 & x_{1i} & x_{1j} \\ 0 & 0 & 0 & 1 \end{array} \right),$$

whose Berezinian is $\text{Ber}B = X_{i|\hat{\mu}}^*$. Define a the supermatrix T whose first even column is the second even column of B^{-1} and whose second even column is $B^{-1}t$ with

$$t = \left(\begin{array}{c} x_{\hat{1}k} \\ 0 \\ x_{1k} \\ 0 \end{array} \right);$$

on the other hand, its first odd column is the second odd column of B^{-1} and its second odd column is $e_{\hat{2}}$. Then

$$BT = \left(\begin{array}{cc|cc} 0 & x_{\hat{1}k} & 0 & x_{\hat{1}j} \\ 1 & 0 & 0 & 0 \\ \hline 0 & x_{1k} & 0 & x_{1j} \\ 0 & 0 & 1 & 1 \end{array} \right).$$

The Berezinian $\text{Ber}(BT) = X_{j|k}^*$. As before, we can compute T explicitly with the help of the super Cramer rule:

$$T = \left(\begin{array}{cc|cc} 0 & X_{i|k}^* X_{i|\hat{\mu}} & -X_{i|j}^* X_{i|\hat{\mu}} & 0 \\ 1 & 0 & 0 & 0 \\ \hline 0 & X_{k|\hat{\mu}} X_{i|\mu}^* & -X_{j|\hat{\mu}} X_{i|\mu}^* & 0 \\ 0 & 0 & 0 & 1 \end{array} \right).$$

Then

$$\text{Ber}T = X_{i|\hat{\mu}}^2 \text{Ber} \left(\begin{array}{c|c} X_{i|\hat{\mu}}^* & X_{i|j}^* \\ \hline X_{k|\hat{\mu}} & X_{j|\hat{\mu}} \end{array} \right),$$

so

$$\text{Ber}(BT) = \text{Ber}B \text{Ber}T = X_{i|\hat{\mu}}^* X_{i|\hat{\mu}}^2 \text{Ber}^* \left(\begin{array}{c|c} X_{j|\hat{\mu}} & X_{k|\hat{\mu}}^* \\ \hline X_{i|j} & X_{i|k}^* \end{array} \right).$$

We then arrive to (24), as we wanted to show. \square

6 Second Fundamental Theorem of super invariant theory for $\text{SL}(1|1)$

Now, we want to prove that the relations that appeared in Equations (21), (22), (23), and (24) completely characterize the super ring of invariants $\tilde{\mathcal{O}}^{\text{SL}(1|1)}$. In this way, we prove the Second Fundamental Theorem of super invariant theory for $\text{SL}(1|1)$.

Let us recall the following basic fact from linear algebra.

Let $\{v_1, v_2, \dots\}$ be an ordered (countable) basis of a vector space V . For any non-zero vector $v = \sum_i a_i v_i$, define the *leading term* of v as $\text{Lt}(v) := v_{i_0}$, where $a_{i_0} \neq 0$, and $a_i = 0$ for all $v_i < v_{i_0}$.

Lemma 6.1. *Any collection of vectors $\{v_1, v_2, \dots\}$ in V , with distinct leading terms with respect to an ordered basis, as above, is linearly independent.*

The superalgebra $\tilde{\mathcal{O}}^{\text{SL}(1|1)}$ is generated by $X_{i|\hat{\mu}}$, $X_{\hat{\mu}|\hat{\nu}}$, $X_{j|\hat{\nu}}^*$, and $X_{i|j}^*$. Each element of $\tilde{\mathcal{O}}^{\text{SL}(1|1)}$ is a linear combination of expressions of the following form:

$$\prod_{i|\hat{\mu}, j|\hat{\lambda}, \hat{\eta}|\hat{\nu}, k|l} X_{i|\hat{\mu}}^{d_{i|\hat{\mu}}} X_{\hat{\eta}|\hat{\nu}}^{d_{\hat{\eta}|\hat{\nu}}} X_{j|\hat{\lambda}}^{*d_{j|\hat{\lambda}}} X_{k|l}^{*d_{k|l}}, \quad (32)$$

where $d_{i|\hat{\mu}}, d_{j|\hat{\lambda}} \in \{0, 1, 2, \dots\}$ and $d_{\hat{\eta}|\hat{\nu}}, d_{k|l} \in \{0, 1\}$. However, due to super Plücker relations (21), (22), (23) and (24), we can express:

$$X_{i|\hat{\mu}} = X_{1|\hat{1}}^* X_{1|\hat{\mu}} X_{i|\hat{1}} - X_{1|\hat{1}}^* X_{\hat{\mu}|\hat{1}} X_{1|\hat{\mu}}^2 X_{1|i}^*, \quad (33)$$

$$X_{j|\hat{\lambda}}^* = X_{1|\hat{1}} X_{j|\hat{1}}^* X_{1|\hat{\lambda}}^* - X_{1|\hat{1}} X_{1|j}^* X_{j|\hat{1}}^2 X_{\hat{\lambda}|\hat{1}}, \quad (34)$$

$$X_{\hat{\eta}|\hat{\nu}} = X_{1|\hat{1}}^* X_{1|\hat{\nu}} X_{\hat{\eta}|\hat{1}} - X_{1|\hat{1}}^* X_{\hat{\nu}|\hat{1}} X_{1|\hat{\nu}}^2 X_{1|\hat{\eta}}^*, \quad (35)$$

$$X_{k|l}^* = X_{1|\hat{1}} X_{k|\hat{1}}^* X_{1|l}^* - X_{1|\hat{1}} X_{1|k}^* X_{k|\hat{1}}^2 X_{l|\hat{1}}. \quad (36)$$

for $1 \leq i, j, k, l \leq p$ and $1 \leq \mu, \lambda, \nu, \eta \leq q$.

Therefore, every expression in (32) can be written and arranged as a linear combination of expressions of the following form:

$$P = P_{1|\underline{\hat{\mu}}} P_{i|\hat{1}} P_{\hat{\eta}|\hat{1}} P_{1|\hat{\lambda}}^* P_{j|\hat{1}}^* P_{1|\underline{\hat{l}}}, \quad (37)$$

(the underlined indices mean multiindices) where:

$$\begin{aligned} P_{1|\underline{\hat{\mu}}} &:= X_{1|\hat{\mu}_1}^{d_{\hat{\mu}_1}} \cdots X_{1|\hat{\mu}_\alpha}^{d_{\hat{\mu}_\alpha}}, & P_{i|\hat{1}} &:= X_{i_1|\hat{1}}^{d_{i_1}} \cdots X_{i_a|\hat{1}}^{d_{i_a}}, & P_{\hat{\eta}|\hat{1}} &:= X_{\hat{\eta}_1|\hat{1}} \cdots X_{\hat{\eta}_\beta|\hat{1}}, \\ P_{1|\hat{\lambda}}^* &:= X_{1|\hat{\lambda}_1}^{*d_{\hat{\lambda}_1}} \cdots X_{1|\hat{\lambda}_\gamma}^{*d_{\hat{\lambda}_\gamma}}, & P_{j|\hat{1}}^* &:= X_{j_1|\hat{1}}^{*d_{j_1}} \cdots X_{j_b|\hat{1}}^{*d_{j_b}}, & P_{1|\underline{\hat{l}}} &:= X_{1|\underline{l}_1}^* \cdots X_{1|\underline{l}_c}^*, \end{aligned}$$

and

$$\begin{aligned} 1 \leq \mu_1 < \cdots < \mu_\alpha \leq q, & \quad 1 < i_1 < \cdots < i_a \leq p, \\ 1 < \eta_1 < \cdots < \eta_\beta \leq q, & \quad 1 \leq \lambda_1 < \cdots < \lambda_\gamma \leq q, \\ 1 < j_1 < \cdots < j_b \leq p, & \quad 1 < l_1 < \cdots < l_c \leq p, \end{aligned}$$

with the conditions

- $\hat{\mu}_\epsilon \neq \hat{\lambda}_\delta$ for any $\epsilon, \delta \in \{1, \dots, q\}$,
- $i_m \neq j_n$ for any $m, n \in \{2, \dots, p\}$.

Definition 6.2. We call the elements of the set:

$$\{X_{1|\hat{\mu}}, X_{i|\hat{1}}, X_{1|\hat{\mu}}^*, X_{i|\hat{1}}^*, X_{\hat{\lambda}|\hat{1}}, X_{1|\hat{j}}^* | 1 \leq i, j \leq p, 2 \leq \hat{\mu}, \hat{\lambda} \leq q\}$$

standard expressions. The products of the standard expressions that appear in P , as in (37), are *standard products*.

Lemma 6.3. *The standard products are linearly independent and form a vector space basis of $\tilde{\mathcal{O}}^{\text{SL}(1|1)}$.*

Proof. The standard products generate $\tilde{\mathcal{O}}^{\text{SL}(1|1)}$, hence we only need to show that they are linearly independent.

Let \mathcal{B} be the monomial basis of $\tilde{\mathcal{O}}$, with expressions of the following form:

$$\Pi_{(i,j)} x_{ij}^{a_{ij}} (x_{ij}^{-1})^{b_{ij}} \Pi_{(k,l)} x_{kl}^{c_{kl}},$$

where:

- x_{ij} and x_{kl} denotes even and odd generators respectively,
- $a_{ij}, b_{ij} \in \{0, 1, 2, \dots\}$, $c_{kl} \in \{0, 1\}$,
- for a fixed (i, j) at least one of a_{ij} or b_{ij} is zero.

Now, consider the following ordering of tuples,

$$\begin{aligned} (1, 1) &< \dots < (1, p) < (\hat{1}, \hat{1}) < \dots < (\hat{1}, \hat{q}) < \\ (1, 1)^{-1} &< \dots < (1, p)^{-1} < (\hat{1}, \hat{1})^{-1} < \dots < (\hat{1}, \hat{q})^{-1} < \\ (1, \hat{1}) &< \dots < (1, \hat{q}) < (\hat{1}, 1) < \dots < (\hat{1}, p), \end{aligned}$$

where $(i, j)^{-1}$ is just a notation for denoting the generators x_{ij}^{-1} . This gives us a total ordering on the basis \mathcal{B} of $\tilde{\mathcal{O}}$ as follows:

$$\Pi_{(i,j)} x_{ij}^{a_{ij}} (x_{ij}^{-1})^{b_{ij}} \Pi_{(k,l)} x_{kl}^{c_{kl}} < \Pi_{(i,j)} x_{ij}^{a'_{ij}} (x_{ij}^{-1})^{b'_{ij}} \Pi_{(k,l)} x_{kl}^{c'_{kl}}$$

if one of the following holds:

(i) \exists an (i, j) for which $a_{ij} \neq a'_{ij}$ and (i, j) is smallest such tuple, then $a_{ij} > a'_{ij}$.

(ii) $a_{ij} = a'_{ij}$ for all (i, j) , and \exists an (i, j) for which $b_{ij} \neq b'_{ij}$ and (i, j) is smallest such tuple then, $b_{ij} < b'_{ij}$.

(iii) $a_{ij} = a'_{ij}$ and $b_{ij} = b'_{ij}$ for all (i, j) , and (k, l) is smallest tuple for which $c_{kl} \neq c'_{kl}$, then $c_{kl} > c'_{kl}$.

With this ordering the leading terms of $P_{1|\underline{\mu}}$, $P_{\underline{i}|\hat{1}}$, $P_{\underline{\hat{q}}|\hat{1}}$, $P_{1|\underline{\hat{\lambda}}}^*$, $P_{\underline{j}|\hat{1}}^*$, $P_{1|\underline{l}}^*$ turns out to be:

$$\begin{aligned} \text{Lt}(P_{1|\underline{\mu}}) &= x_{11}^{d_{\underline{\mu}}} (x_{\hat{1}\hat{\mu}_1}^{-1})^{d_{\hat{\mu}_1}} \dots (x_{\hat{1}\hat{\mu}_\alpha}^{-1})^{d_{\hat{\mu}_\alpha}}, & \text{Lt}(P_{\underline{i}|\hat{1}}) &= x_{1i_1}^{d_{i_1}} \dots x_{1i_a}^{d_{i_a}} (x_{\hat{1}\hat{1}}^{-1})^{d_{\hat{1}}}, \\ \text{Lt}(P_{\underline{\hat{q}}|\hat{1}}) &= (x_{\hat{1}\hat{1}}^{-1})^\beta x_{1\hat{q}_1} \dots x_{1\hat{q}_\beta}, & \text{Lt}(P_{1|\underline{\hat{\lambda}}}^*) &= x_{\hat{1}\hat{\lambda}_1}^{d_{\hat{\lambda}_1}} \dots x_{\hat{1}\hat{\lambda}_\gamma}^{d_{\hat{\lambda}_\gamma}} (x_{11}^{-1})^{d_{\hat{\lambda}}}, \\ \text{Lt}(P_{\underline{j}|\hat{1}}^*) &= x_{11}^{d_j} (x_{1j_1}^{-1})^{d_{j_1}} \dots (x_{1j_1}^{-1})^{d_{j_b}}, & \text{Lt}(P_{1|\underline{l}}^*) &= (x_{11}^{-1})^c x_{1l_1} \dots x_{1l_c}, \end{aligned}$$

where $d_{\underline{\mu}} := d_{\mu_1} + \cdots d_{\mu_\alpha}$ and similarly others. Moreover, the leading term $\text{Lt}(P)$ of the standard generator $P = P_{1|\underline{\mu}} P_{i|\hat{1}} P_{\hat{1}|\hat{1}} P_{1|\hat{\Delta}}^* P_{j|\hat{1}}^* P_{1|\hat{l}}^*$ is the product of these leading terms. Therefore, distinct standard generators have distinct leading terms, and, by Lemma 6.1, they are linearly independent. \square

The Lemma 6.3 implies that the super Plücker relations are all the algebraic relations among the generators. We state this precisely in the theorem below.

Theorem 6.4. *Second Fundamental Theorem of super invariant theory for $\text{SL}(1|1)$. We have a presentation of the ring of invariants,*

$$\tilde{\mathcal{O}}^{\text{SL}(1|1)} \cong \mathbb{C}[Y_{i|\hat{\mu}}, Y_{\hat{\mu}|\hat{\nu}}, Y_{j|\hat{\nu}}^*, Y_{i|\hat{j}}^*] / \text{I}_{sp}$$

where $Y_{i|\hat{\mu}}$ and $Y_{j|\hat{\nu}}^*$ are even variables, $Y_{\hat{\mu}|\hat{\nu}}$ and $Y_{i|\hat{j}}^*$ are odd variables, and I_{sp} is the ideal generated by super Plücker relations given in Equations (21), (22), (23) and (24).

Proof. Define,

$$\begin{aligned} \pi : \mathbb{C}[Y_{i|\hat{\mu}}, Y_{\hat{\mu}|\hat{\nu}}, Y_{j|\hat{\nu}}^*, Y_{i|\hat{j}}^*] &\longrightarrow \tilde{\mathcal{O}}^{\text{SL}(1|1)} \\ Y_{i|\hat{\mu}}, Y_{\hat{\mu}|\hat{\nu}}, Y_{j|\hat{\nu}}^*, Y_{i|\hat{j}}^* &\mapsto X_{i|\hat{\mu}}, X_{\hat{\mu}|\hat{\nu}}, X_{j|\hat{\nu}}^*, X_{i|\hat{j}}^*. \end{aligned} \quad (38)$$

Then, clearly, $\text{I}_{sp} \subset \ker(\pi)$. We need to show that, in fact, $\text{I}_{sp} = \ker(\pi)$.

Assume $f \in \ker(\pi)$. By Lemma 6.3 we can write:

$$f = \sum_i a_i P_i(Y) + Q$$

where the $P_i(Y)$ are standard products in Y 's and $Q \in \text{I}_{sp}$. We have

$$\pi(f) = \sum_i a_i P_i(X) = 0$$

where $P_i(X)$ are standard products in X 's. By Lemma 6.3, we have $a_i = 0$ for all i , so that $f = Q \in \text{I}_{sp}$. \square

7 Super Plücker relations for $\text{SL}(r|s)$

In this section, we prove one of the main results of our paper: the generalization to the super setting of the Plücker relations for $\text{SL}(r|s)$, expressed for the ordinary setting in Thm. 1.1. Again, it is important to note that these super Plücker relations in the general case also coincides with the relations appeared in [20, Th. 9] obtained with a different approach.

Theorem 7.1. *Let the notation be as in (9), (10) and (11). Then,*

$$X_{\underline{i}|\underline{\hat{\mu}}} X_{\underline{i}|\underline{\hat{\mu}}}^* = 1 \quad (39)$$

$$(X_{\underline{i}|\underline{\hat{\mu}}})^{r+s-1} X_{\underline{j}|\underline{\hat{\nu}}} = \text{Ber} \left(\begin{array}{c|c} X_{\underline{i}_a(j_t)|\underline{\hat{\mu}}} & X_{\underline{i}_a(\nu_\beta)|\underline{\hat{\mu}}} \\ \hline X_{\underline{i}|\underline{\hat{\mu}}_\alpha}^*(j_t) & X_{\underline{i}|\underline{\hat{\mu}}_\alpha}^*(\nu_\beta) \end{array} \right)_{\substack{a,t=1,\dots,r \\ \alpha,\beta=1,\dots,s}} \quad (40)$$

$$(X_{\underline{i}|\underline{\hat{\mu}}})^{r+s-1} X_{j_1 \dots j_{r-1} \hat{\lambda}|\underline{\hat{\nu}}} = \text{Ber} \left(\begin{array}{cc|c} X_{\underline{i}_a(j_t)|\underline{\hat{\mu}}} & X_{\underline{i}_a(\hat{\lambda})|\underline{\hat{\mu}}} & X_{\underline{i}_a(\nu_\beta)|\underline{\hat{\mu}}} \\ \hline X_{\underline{i}|\underline{\hat{\mu}}_\alpha}^*(j_t) & X_{\underline{i}|\underline{\hat{\mu}}_\alpha}^*(\hat{\lambda}) & X_{\underline{i}|\underline{\hat{\mu}}_\alpha}^*(\nu_\beta) \end{array} \right)_{\substack{a=1,\dots,r \\ t=1,\dots,r-1 \\ \alpha,\beta=1,\dots,s}} \quad (41)$$

$$(X_{\underline{i}|\underline{\hat{\mu}}})^{-r-s+1} X_{\underline{j}|\nu_1 \dots \nu_{s-1} y}^* = \text{Ber}^* \left(\begin{array}{c|cc} X_{\underline{i}_a(j_t)|\underline{\hat{\mu}}} & X_{\underline{i}_a(\nu_\beta)|\underline{\hat{\mu}}} & X_{\underline{i}_a(y)|\underline{\hat{\mu}}} \\ \hline X_{\underline{i}|\underline{\hat{\mu}}_\alpha}^*(j_t) & X_{\underline{i}|\underline{\hat{\mu}}_\alpha}^*(\nu_\beta) & X_{\underline{i}|\underline{\hat{\mu}}_\alpha}^*(y) \end{array} \right)_{\substack{a,t=1,\dots,r \\ \alpha=1,\dots,s \\ \beta=1,\dots,s-1}} \quad (42)$$

where $\underline{i}_a(j_t)|\underline{\hat{\mu}}$ denotes replacing the a -th index of $\underline{i} = i_1, \dots, i_r$ with j_t and similarly others. We call these the super Plücker relations for $\text{SL}(r|s)$.

Proof. Consider the ordered multi-indices $\underline{i}|\underline{\hat{\mu}} = (i_1, \dots, i_r|\hat{\mu}_1, \dots, \hat{\mu}_s)$ and $\underline{j}|\underline{\hat{\nu}} = (j_1, \dots, j_r|\hat{\nu}_1, \dots, \hat{\nu}_s)$ taken from $(1, \dots, p|\hat{1}, \dots, \hat{q})$. We fix the following matrix of size $(2r|2s) \times (2r|2s)$:

$$A = \left(\begin{array}{cc|cc} A_1 & A_2 & A_3 & A_4 \\ O & I_r & O & O \\ \hline A_5 & A_6 & A_7 & A_8 \\ O & O & O & I_s \end{array} \right).$$

where O and I denote the null and identity matrices, respectively, and

$$\begin{array}{llll} A_1 = [x_{ki_l}] & A_2 = [x_{kj_l}] & A_3 = [x_{k\hat{\mu}_n}] & A_4 = [x_{k\hat{\nu}_n}] \\ A_5 = [x_{\hat{m}i_l}] & A_6 = [x_{\hat{m}j_l}] & A_7 = [x_{\hat{m}\hat{\mu}_n}] & A_8 = [x_{\hat{m}\hat{\nu}_n}] \end{array}$$

where $k, l = 1, \dots, r$ and $m, n = 1, \dots, s$. Then, clearly,

$$\text{Ber}(A) = X_{\underline{i}|\underline{\hat{\mu}}}. \quad (43)$$

Moreover, by setting,

$$u = (r+1, \dots, 2r | s+1, \dots, 2s) \quad \text{and} \quad v = (1, \dots, r | \hat{1}, \dots, \hat{s})$$

we get:

$$\text{Ber}(A_v^u) = \text{Ber} \left(\begin{array}{c|c} A_2 & A_4 \\ \hline A_6 & A_8 \end{array} \right) = X_{\underline{j}|\underline{\hat{\nu}}}. \quad (44)$$

On the other hand:

$$(A^{-1})_{\underline{u}}^{\underline{\hat{v}}} = \left(\begin{array}{c|c} (A^{-1})_{ab} & (A^{-1})_{a\hat{d}} \\ \hline (A^{-1})_{\hat{c}b} & (A^{-1})_{\hat{c}\hat{d}} \end{array} \right)$$

where $a = 1, \dots, r$, $b = r+1, \dots, 2r$, $c = 1, \dots, s$ and $d = s+1, \dots, 2s$.

These entries of A^{-1} can be computed using the super Cramer rule. For instance,

$$(A^{-1})_{ab} = \frac{\text{Ber}(A_a(e_b))}{\text{Ber}(A)} = -X_{\underline{i}_a(j_t)|\underline{\hat{\mu}}} X_{\underline{i}|\underline{\hat{\mu}}}^*$$

where $t = b - r$.

Similarly, the other entries are obtained. We get:

$$(A^{-1})_{\underline{u}}^{\underline{\hat{v}}} = \left(\begin{array}{c|c} -X_{\underline{i}_a(j_t)|\underline{\hat{\mu}}} X_{\underline{i}|\underline{\hat{\mu}}}^* & -X_{\underline{i}_a(\nu_\beta)|\underline{\hat{\mu}}} X_{\underline{i}|\underline{\hat{\mu}}}^* \\ \hline -X_{\underline{i}|\underline{\hat{\mu}}_\alpha}^*(j_t) X_{\underline{i}|\underline{\hat{\mu}}} & -X_{\underline{i}|\underline{\hat{\mu}}_\alpha}^*(\nu_\beta) X_{\underline{i}|\underline{\hat{\mu}}} \end{array} \right)_{\substack{a,t=1,\dots,r \\ \alpha,\beta=1,\dots,s}} \quad (45)$$

Substituting Equations. (43), (44) and (45) into the super Jacobi identity, we get relations in Equation (40). Similarly, the other super Plücker relations in Eqs. (41) and (42) are obtained by modifying the strategy used in the proof of super Jacobi identity, as we explicitly presented in the previous section for the case of $\text{SL}(1|1)$. \square

We end this section by stating a conjecture.

Conjecture 7.2. $\tilde{\mathcal{O}}^{\text{SL}(r|s)}$ is isomorphic to the superalgebra generated by $Y_{\underline{i}|\underline{\hat{\mu}}}$, $Y_{\underline{i}_a(\hat{\nu})|\underline{\hat{\mu}}}$, $Y_{\underline{i}|\underline{\hat{\mu}}_\alpha}^*(j)$ and $Y_{\underline{i}|\underline{\hat{\mu}}}^*$ subject to the super Plücker relations given in Equations (39), (40), (41) and (42).

Acknowledgments

We thank Prof. Shemyakova and Prof. Voronov for helpful discussions. R. Fioresi and J. Razzaq want to thank the Departament de Física Teòrica, Universitat de València and the Department of Mathematics, LUMS for the hospitality while part of the work was done. M. A. Lledó wants to thank Fabit, Università di Bologna, for its kind hospitality. This research was supported by GNSAGA-Indam, PNRR MNESYS, PNRR National Center for HPC, Big Data and Quantum Computing CUP J33C22001170001, PNRR SIMQuSEC CUP J13C22000680006. This work is also supported by the Spanish Grants PID2020-116567GB-C21, PID2023-001292-S and CEX2023-001292-S funded by MCIU/AEI/10.13039/501100011033. This work was supported by Horizon Europe EU projects MSCA-SE CaLIGOLA, Project ID: 101086123, MSCA-DN CaLiForNIA, Project ID: 101119552.

This article is based upon work from COST Action CaLISTA CA21109 supported by COST (European Cooperation in Science and Technology), www.cost.eu.

Declaration

This work is pure mathematical and does not have associated data. Moreover, the authors declare no conflict of interest.

A Supergeometry

We briefly recall some notions regarding supergroups, superspaces and their \mathcal{A} -points. For more details see, for example, Ref. [5], Chapters 1 and 3 and [10], Chapter 1.

Let \mathbf{k} denote the ground field ($\mathbf{k} = \mathbb{R}, \mathbb{C}$). A *super vector space* is a \mathbb{Z}_2 -graded vector space over \mathbf{k} $V = V_0 \oplus V_1$. We call the pair of numbers $\dim V_0 | \dim V_1$ the *super dimension* of V and we define $\mathbf{k}^{m|n} := \mathbf{k}^m \oplus \mathbf{k}^n$.

We say that \mathcal{A} is a *commutative superalgebra* if it is a \mathbb{Z}_2 -graded algebra, with grading preserving multiplication and $ab = (-1)^{p(a)p(b)}ba$, for all (homogeneous) $a, b \in \mathcal{A}$, being $p(a), p(b) = 0, 1$ according to the grading ($p(a)$ is the *parity* of an element $a \in \mathcal{A}$). An important example of commutative

superalgebra is the *superalgebra of polynomials*:

$$\mathbf{k}[x_1, \dots, x_m, \xi_1, \dots, \xi_n] := k[x_1, \dots, x_m] \otimes \wedge(\xi_1, \dots, \xi_n)$$

where Latin letters denote the *even* (commuting) variables, while Greek letters denote the *odd* (anticommuting) ones.

For a commutative superalgebra \mathcal{A} , we define the free \mathcal{A} -module $\mathcal{A}^{m|n} = \mathcal{A} \otimes k^{m|n}$, where

$$\mathcal{A}_0^{m|n} := \mathcal{A}_0 \otimes k^m \oplus \mathcal{A}_1 \otimes k^n \quad \text{and} \quad \mathcal{A}_1^{m|n} := \mathcal{A}_0 \otimes k^n \oplus \mathcal{A}_1 \otimes k^m$$

are, respectively, its even and odd parts.

Given a commutative superalgebra \mathcal{A} , the \mathcal{A} -points of the supervector space $V = V_0 \oplus V_1$ are:

$$V(\mathcal{A}) = (\mathcal{A} \otimes V)_0 = \mathcal{A}_0 \otimes V_0 + \mathcal{A}_1 \otimes V_1$$

We call the elements in $(\mathcal{A} \otimes V)_0$ *even vectors*, while the elements in $(\mathcal{A} \otimes V)_1$ are *odd vectors*.

For example, the \mathcal{A} -points of the supervector space $k^{m|n}$ are:

$$k^{m|n}(\mathcal{A}) := \mathcal{A}_0 \otimes k^m \oplus \mathcal{A}_1 \otimes k^n = \{(a_1, \dots, a_m, \alpha_1, \dots, \alpha_n)\} \quad (46)$$

Hence $(a_1, \dots, a_m, \alpha_1, \dots, \alpha_n)$ is an even vector while $(\beta_1, \dots, \beta_m, b_1, \dots, b_n)$ is an odd one.

Observation A.1. We equivalently interpret the \mathcal{A} -points as follows:

$$\mathbf{k}^{m|n}(\mathcal{A}) = \text{Hom}_{(\text{salg})}(\mathbf{k}[x_1, \dots, x_m, \xi_1, \dots, \xi_n], \mathcal{A})$$

where (salg) denotes the category of commutative superalgebras. In fact, an expression as (46), gives a morphism f by specifying the images of the generators x_i 's and ξ_j 's: $f(x_i) = a_i$, $f(\xi_j) = \alpha_j$ and vice-versa. We say, with an abuse of language, that $\mathbf{k}[x_1, \dots, x_m, \xi_1, \dots, \xi_n]$ is the superalgebra of ‘functions’ on the vector superspace $k^{m|n}$. To be precise, $\mathbf{k}[x_1, \dots, x_m, \xi_1, \dots, \xi_n]$ represents $\mathbf{k}^{m|n}$ when we adopt the language of the *functor of points*, that we shall not pursue here. The above superalgebra, $\mathbf{k}[x_1, \dots, x_m, \xi_1, \dots, \xi_n]$, can be seen also as the superalgebra of global sections of the structural sheaf on the supervariety $\mathbf{k}^{m|n}$. We shall not go further into the theory, the reader can see, for example, [5], Chapter 3.

We now take into exam a special case, important in our treatment. We define the super vector space of *supermatrices* as:

$$\begin{aligned} M_{r|s \times p|q} &= \left\{ \left(\begin{array}{c|c} X & Y \\ \hline Z & W \end{array} \right) \right\}, \quad \text{with} \\ (M_{r|s \times p|q})_0 &= \left\{ \left(\begin{array}{c|c} X & 0 \\ \hline 0 & W \end{array} \right) \right\}, \quad (M_{r|s \times p|q})_1 = \left\{ \left(\begin{array}{c|c} 0 & Y \\ \hline Z & 0 \end{array} \right) \right\} \end{aligned}$$

where X , Y , Z , W are respectively $r \times p$, $r \times q$, $s \times p$ and $s \times q$ matrices with entries in k . Whenever $p|q = r|s$, we write just $M(r|s)$.

For a commutative superalgebra \mathcal{A} , $(\mathcal{A} \otimes M_{r|s \times p|q})_0$ are the *even supermatrices*, and $(\mathcal{A} \otimes M_{r|s \times p|q})_1$ are the *odd supermatrices*:

$$\begin{aligned} (\mathcal{A} \otimes M_{r|s \times p|q})_0 &= \left\{ \left(\begin{array}{c|c} [a_{ij}] & [\alpha_{il}] \\ \hline [\beta_{kj}] & [b_{kl}] \end{array} \right) \right\} \\ (\mathcal{A} \otimes M_{r|s \times p|q})_1 &= \left\{ \left(\begin{array}{c|c} [\sigma_{ij}] & [c_{il}] \\ \hline [d_{kj}] & [\delta_{kl}] \end{array} \right) \right\} \end{aligned}$$

Notice that the vectors forming the first p columns of a $r|s \times p|q$ even supermatrix are even, while the vectors forming the last q columns are odd. Vice-versa for odd supermatrices the first p columns consist of odd vectors and the last q of even ones.

The set of \mathcal{A} -points of the *general linear supergroup*, denoted, as usual, $GL(r|s)(\mathcal{A})$, is the group of automorphisms of $\mathcal{A}^{r|s}$. It consists of all invertible elements in $M(r|s)(\mathcal{A})$. They are characterized by having invertible diagonal blocks (see, for example, [5], Chapter 1). One can then define the *Berezinian*:

$$\begin{aligned} \text{Ber} \left(\begin{array}{c|c} X & Y \\ \hline Z & W \end{array} \right) &= \det(X - YW^{-1}Z) \det(W)^{-1} = \\ &= \det(X) \det(W - ZX^{-1}Y)^{-1}, \end{aligned} \quad (47)$$

where ‘det’ is the usual determinant.

B The super Cramer rule

We give here a proof of the super Cramer rule, Theorem 2.5.

We need first some notation: for any even vector \mathbf{b} of size $r|s$, we denote by $\mathbf{b}^{(e)}$ and $\mathbf{b}^{(o)}$ the column vectors consisting of even and odd coefficients of \mathbf{b} , respectively:

$$\mathbf{b} = \begin{pmatrix} b_1 \\ \cdot \\ \cdot \\ \cdot \\ b_r \\ b_{\hat{1}} \\ \cdot \\ \cdot \\ \cdot \\ b_{\hat{s}} \end{pmatrix}, \quad \mathbf{b}^{(e)} = \begin{pmatrix} b_1 \\ \cdot \\ \cdot \\ \cdot \\ b_r \end{pmatrix}, \quad \mathbf{b}^{(o)} = \begin{pmatrix} b_{\hat{1}} \\ \cdot \\ \cdot \\ \cdot \\ b_{\hat{s}} \end{pmatrix}$$

Theorem 2.5 states:

Let $M \in \text{GL}(r|s)$ be an invertible even supermatrix and let $\mathbf{b} \in \mathbf{k}^{r|s}$ be an even vector of size $r|s$, then the solution to the equation

$$M\mathbf{x} = \mathbf{b}$$

is:

$$x_i = \frac{\text{Ber} M_i(\mathbf{b})}{\text{Ber} M} \quad \text{for } i = 1, \dots, r$$

$$x_{\hat{j}} = \frac{\text{Ber}^* M_{\hat{j}}(\mathbf{b})}{\text{Ber}^* M} \quad \text{for } \hat{j} = \hat{1}, \dots, \hat{s}$$

where $M_i(\mathbf{b})$ is the supermatrix obtained by replacing i -th even column of M with \mathbf{b} and $M_{\hat{j}}(\mathbf{b})$ is the fake supermatrix of type-II obtained by replacing \hat{j} -th odd column of M with \mathbf{b} .

Proof. We have to show that the above statement holds for all $M \in \text{GL}(r|s)$. We will do it in three steps.

1. One can decompose any supermatrix $M \in \text{GL}(r|s)$ as a product $M = M_+ M_0 M_-$ where M_+ , M_0 and M_- are supermatrices of the following types respectively:

$$\left(\begin{array}{c|c} I & X \\ \hline O & I \end{array} \right), \quad \left(\begin{array}{c|c} V & O \\ \hline O & W \end{array} \right), \quad \left(\begin{array}{c|c} I & O \\ \hline Z & I \end{array} \right),$$

where I denotes the identity matrix and O denotes the null matrix.

For $M = \left(\begin{array}{c|c} M_1 & M_2 \\ \hline M_3 & M_4 \end{array} \right)$, one gets:

$$\begin{aligned} X &= M_2 M_4^{-1} \\ V &= M_1 - M_2 M_4^{-1} M_3 \\ W &= M_4 \\ Z &= M_4^{-1} M_3. \end{aligned}$$

2. The statement of the theorem holds for supermatrices of the form $C = AB$, where A and B are of the form M_0 and M_- respectively.

Let $A = \left(\begin{array}{c|c} V & O \\ \hline O & W \end{array} \right)$ and $B = \left(\begin{array}{c|c} I & O \\ \hline Z & I \end{array} \right)$ so,

$$C = \left(\begin{array}{c|c} V & O \\ \hline WZ & W \end{array} \right), \quad \text{and}$$

$$C^{-1} = \left(\begin{array}{c|c} V^{-1} & O \\ \hline -ZV^{-1} & W^{-1} \end{array} \right) = \left(\begin{array}{c|c} \left[\frac{\det V_i(e_j)}{\det V} \right] & O \\ \hline \left[\frac{-\sum_{i=1}^r z_{ki} \det V_i(e_j)}{\det V} \right] & \left[\frac{\det W_k(e_l)}{\det W} \right] \end{array} \right)$$

where e_j denotes the column vector having 1 at j -th place and 0 at all other places. Therefore,

$$C^{-1}\mathbf{b} = \left(\begin{array}{c} \frac{\sum_{j=1}^r \det V_1(e_j)b_j}{\det V} \\ \cdot \\ \cdot \\ \frac{\sum_{j=1}^r \det V_r(e_j)b_j}{\det V} \\ \hline -\frac{\sum_{i,j=1}^r z_{\hat{1}i} \det V_i(e_j)b_j}{\det V} + \frac{\sum_{l=\hat{1}}^{\hat{s}} \det W_1(e_l)b_l}{\det W} \\ \cdot \\ \cdot \\ -\frac{\sum_{i,j=1}^r z_{\hat{s}i} \det V_i(e_j)b_j}{\det V} + \frac{\sum_{l=\hat{1}}^{\hat{s}} \det W_s(e_l)b_l}{\det W} \end{array} \right). \quad (48)$$

On the other hand, using the linearity of determinant, we have:

$$\begin{aligned} \frac{\text{Ber} C_i(\mathbf{b})}{\text{Ber} C} &= \frac{\det W^{-1} \det V_i(\mathbf{b}^{(e)})}{\det W^{-1} \det V} = \frac{\det V_i(\mathbf{b}^{(e)})}{\det V} = \\ &= \frac{\sum_{j=1}^r \det V_i(e_j)b_j}{\det V} \end{aligned} \quad (49)$$

and

$$\begin{aligned} \frac{\text{Ber}^* C_{\hat{k}}(\mathbf{b})}{\text{Ber}^* C} &= \frac{\det V^{-1} \det (W_k(\mathbf{b}^{(o)}) - WZV^{-1}O_k(\mathbf{b}^{(e)}))}{\det V^{-1} \det W} = \\ &= \frac{\det W_k(\mathbf{b}^{(o)}) - \det W_k(WZV^{-1}\mathbf{b}^{(e)})}{\det W} = \\ &= \frac{\sum_{l=\hat{1}}^{\hat{s}} \det W_k(e_l)b_l}{\det W} - \frac{\sum_{i,j=1}^r z_{\hat{k}i} \det V_i(e_j)b_j}{\det V}. \end{aligned} \quad (50)$$

The last two equalities follows from the multi-linearity property of determinant and ordinary the Cramer rule, respectively. Hence, comparing Eqs. (48), (49) and (50) one realizes that the theorem holds for supermatrices of type C .

3. The theorem holds for supermatrices of type $D = EF$ where $F = \left(\begin{array}{c|c} U & Y \\ \hline V & W \end{array} \right)$ is an even supermatrix which already satisfies the theorem and $E = \left(\begin{array}{c|c} I & X \\ \hline O & I \end{array} \right)$ with X having only one non-zero entry. Without loss of generality, we suppose that the top-left entry $x_{1\hat{1}} \neq 0$.

Therefore,

$$D = \left(\begin{array}{ccc|ccc} u_{11} + x_{1\hat{1}}v_{\hat{1}1} & \cdots & u_{1r} + x_{1\hat{1}}v_{\hat{1}r} & y_{1\hat{1}} + x_{1\hat{1}}w_{\hat{1}\hat{1}} & \cdots & y_{1\hat{s}} + x_{1\hat{1}}w_{\hat{1}\hat{s}} \\ u_{21} & \cdots & u_{2r} & y_{2\hat{1}} & \cdots & y_{2\hat{s}} \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ u_{r1} & \cdots & u_{rr} & y_{r\hat{1}} & \cdots & y_{r\hat{s}} \\ \hline & & V & & & W \end{array} \right)$$

Note that,

$$\mathbf{x} = D^{-1}\mathbf{b} = F^{-1}(E^{-1}\mathbf{b}). \quad (51)$$

Fix

$$\mathbf{c} = E^{-1}\mathbf{b} = \left(\begin{array}{c} b_1 - x_{1\hat{1}}b_{\hat{1}} \\ \cdot \\ \cdot \\ b_r \\ \hline b_{\hat{1}} \\ \cdot \\ \cdot \\ b_{\hat{s}} \end{array} \right).$$

Since F satisfies the theorem, equation (51) implies

$$x_i = \frac{\text{Ber} F_i(\mathbf{c})}{\text{Ber} F} = \frac{\det(U_i(\mathbf{c}^{(e)}) - YW^{-1}V_i(\mathbf{b}^{(o)}))}{\det(U - YW^{-1}V)} \quad (52)$$

and

$$x_{\hat{j}} = \frac{\text{Ber}^* F_{\hat{j}}(\mathbf{c})}{\text{Ber}^* F} = \frac{\det(W_{\hat{j}}(\mathbf{c}^{(o)}) - VU^{-1}Y_{\hat{j}}(\mathbf{c}^{(e)}))}{\det(W - VU^{-1}Y)}. \quad (53)$$

On the other hand,

$$\begin{aligned} \frac{\text{Ber} D_i(\mathbf{b})}{\text{Ber} D} &= \\ \frac{\det((U + XV)_i(\mathbf{b}^{(e)}) - (Y + XW)W^{-1}V_i(\mathbf{b}^{(o)}))}{\det(U - YW^{-1}V)} &= \\ \frac{\det((U + XV)_i(\mathbf{b}^{(e)}) - XV_i(\mathbf{b}^{(o)}) - YW^{-1}V_i(\mathbf{b}^{(o)}))}{\det(U - YW^{-1}V)} &= \\ \frac{\det(U_i(\mathbf{c}^{(e)}) - YW^{-1}V_i(\mathbf{b}^{(o)}))}{\det(U - YW^{-1}V)} &= \end{aligned} \quad (54)$$

and

$$\begin{aligned} \frac{\text{Ber}^* D_{\hat{j}}(\mathbf{b})}{\text{Ber}^* D} &= \frac{\det(W_{\hat{j}}(\mathbf{b}^{(o)}) - V(U + XV)^{-1}(Y + XW)_{\hat{j}}(\mathbf{b}^{(e)}))}{\det(W - V(U + XV)^{-1}(Y + XW))} = \\ \frac{\det(W_{\hat{j}}(\mathbf{b}^{(o)}) - V(U^{-1} - U^{-1}XVU^{-1})(Y_{\hat{j}}(\mathbf{c}^{(e)}) + XW_{\hat{j}}(\mathbf{b}^{(o)})))}{\det(W - V(U^{-1} - U^{-1}XVU^{-1})(Y + XW))} &= \\ \frac{\det(W_{\hat{j}}(\mathbf{b}^{(o)}) - VU^{-1}Y_{\hat{j}}(\mathbf{c}^{(e)}) - VU^{-1}XW_{\hat{j}}(\mathbf{b}^{(o)}) - VU^{-1}XVU^{-1}Y_{\hat{j}}(\mathbf{c}^{(e)}))}{\det(W - VU^{-1}Y - VU^{-1}XW + VU^{-1}XVU^{-1}Y)} &= \\ \frac{\det(I - VU^{-1}X) \det(W_{\hat{j}}(\mathbf{c}^{(o)}) - VU^{-1}Y_{\hat{j}}(\mathbf{c}^{(e)}))}{\det(I - VU^{-1}X) \det(W - VU^{-1}Y)} &= \\ \frac{\det(W_{\hat{j}}(\mathbf{c}^{(o)}) - VU^{-1}Y_{\hat{j}}(\mathbf{c}^{(e)}))}{\det(W - VU^{-1}Y)}. & \end{aligned} \quad (55)$$

By comparing Eqs. (52), (53), (54) and (55), one concludes that the matrices of type F satisfy the Theorem.

One can easily observe that every even supermatrix $M \in \text{GL}(r|s)$ is of type F and this completes the proof. \square

Remark B.1. It is important to note that, in the above proof, we never use the fact that \mathbf{b} is even. The same expressions work for the equation $M\mathbf{x} = \mathbf{b}$ if \mathbf{b} is an odd vector of size $r|s$.

C The Jacobi identity and the Plücker relations

We start by recalling a determinant identity due to Jacobi, the *Jacobi complementary minor theorem* [17] or the *Jacobi identity* for short.

Theorem C.1. *The Jacobi complementary minor theorem or Jacobi identity. Let A be an invertible $n \times n$ matrix. Fix the two sets of indices:*

$$u = (n - r + 1, \dots, n) \quad \text{and} \quad v = (1, \dots, r)$$

Let \tilde{u} and \tilde{v} denote the complements of u and v respectively in $(1, \dots, n)$. Then:

$$\det A \det(A^{-1})_{\tilde{u}}^{\tilde{v}} = (-1)^{r(n+1)} \det A_v^u \quad (56)$$

where A_v^u denotes the matrix obtained from A by deleting its rows and columns whose indices are contained in u and v , respectively.

Proof. We denote $A = [a_{ij}]_{n \times n}$ and, $A^{-1} = [b_{kl}]_{n \times n}$. Therefore,

$$(A^{-1})_{\tilde{u}}^{\tilde{v}} = \begin{pmatrix} b_{1(n-r+1)} & \cdots & b_{1n} \\ \vdots & \cdots & \vdots \\ \vdots & \cdots & \vdots \\ b_{r(n-r+1)} & \cdots & b_{rn} \end{pmatrix}.$$

Notice that,

$$\det(A^{-1})_{\tilde{u}}^{\tilde{v}} = \det \begin{pmatrix} b_{1(n-r+1)} & \cdots & b_{1n} \\ \cdot & \cdots & \cdot \\ \cdot & \cdots & \cdot \\ b_{r(n-r+1)} & \cdots & b_{rn} \end{pmatrix} =$$

$$\det \begin{pmatrix} b_{1(n-r+1)} & \cdots & b_{1n} & 0 & \cdots & 0 \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ b_{r(n-r+1)} & \cdots & b_{rn} & 0 & \cdots & 0 \\ b_{(r+1)(n-r+1)} & \cdots & b_{(r+1)n} & 1 & \cdots & 0 \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ b_{n(n-r+1)} & \cdots & b_{nn} & 0 & \cdots & 1 \end{pmatrix}_{n \times n} \quad (57)$$

Let us denote the $n \times n$ matrix that appeared in Equation (57) by T . Hence, using the multiplicative property of the determinant we get:

$$\det A \det(A^{-1})_{\tilde{u}}^{\tilde{v}} = \det A \det T = \det(AT) =$$

$$\det \begin{pmatrix} 0 & \cdots & 0 & a_{1(r+1)} & \cdots & a_{1n} \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ 0 & \cdots & 0 & a_{(n-r)(r+1)} & \cdots & a_{(n-r)n} \\ 1 & \cdots & 0 & a_{(n-r+1)r+1} & \cdots & a_{(n-r+1)n} \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ 0 & \cdots & 1 & a_{n(r+1)} & \cdots & a_{nn} \end{pmatrix} =$$

$$(-1)^{r(n+1)} \det \begin{pmatrix} a_{1(r+1)} & \cdots & a_{1n} \\ \cdot & \cdots & \cdot \\ \cdot & \cdots & \cdot \\ a_{(n-r)(r+1)} & \cdots & a_{(n-r)n} \end{pmatrix} =$$

$$(-1)^{r(n+1)} \det(A_v^u).$$

as desired. □

There are several and more general versions of the Jacobi identity, (see [14], Section 6 and references therein). However, we show now that the Jacobi identity, as in (56), is enough to recover the ordinary Plücker relations as in Equation (1) of Theorem 1.1.

Proposition C.2. *Let $X_{i_1 \dots i_r}$ denote the minor of $M \in M_{r \times p}$ ($r \leq p$) corresponding to the submatrix formed by the columns (i_1, \dots, i_r) of M . Then, the relations*

$$\sum_{k=1}^{r+1} (-1)^k X_{i_1 \dots i_{r-1} j_k} X_{j_1 \dots \tilde{j}_k \dots j_{r+1}} = 0$$

come directly from the Jacobi identity (56).

Proof. Fix an $r \times p$ matrix:

$$C = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1p} \\ \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdots & \cdot \\ a_{r1} & a_{r2} & \cdots & a_{rp} \end{pmatrix}$$

Set any two sets $\{i_1, \dots, i_r\}$ and $\{j_1, \dots, j_r\}$ of ordered indices from $\{1, \dots, p\}$ and define the following $2r \times 2r$ matrix:

$$A = \begin{pmatrix} a_{1i_1} & \cdots & a_{1i_r} & a_{1j_1} & \cdots & a_{1j_r} \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ a_{ri_1} & \cdots & a_{ri_r} & a_{rj_1} & \cdots & a_{rj_r} \\ 0 & \cdots & 0 & 1 & \cdots & 0 \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdots & \cdot & \cdot & \cdots & \cdot \\ 0 & \cdots & 0 & 0 & \cdots & 1 \end{pmatrix} \quad \text{which satisfies} \quad \det A = X_{i_1 \dots i_r}.$$

Suppose that $X_{i_1 \dots i_r}$ is invertible. Now, fix

$$u = (r+1, \dots, 2r) \quad \text{and} \quad v = (1, \dots, r),$$

which implies,

$$\tilde{u} = (1, \dots, r) \quad \text{and} \quad \tilde{v} = (r+1, \dots, 2r).$$

Therefore,

$$A_v^u = \begin{pmatrix} a_{1j_1} & \cdots & a_{1j_r} \\ \vdots & \cdots & \vdots \\ \vdots & \cdots & \vdots \\ a_{rj_1} & \cdots & a_{rj_r} \end{pmatrix} \quad \text{and} \quad \det A_v^u = X_{j_1 \cdots j_r}.$$

Putting this in Equation (56) one gets

$$X_{j_1 \cdots j_r} = (-1)^r X_{i_1 \cdots i_r} \det(A^{-1})_{\tilde{u}}^{\tilde{v}}. \quad (58)$$

Now,

$$(A^{-1})_{\tilde{u}}^{\tilde{v}} = [(A^{-1})_{kl}]_{\substack{k=1, \dots, r \\ l=r+1, \dots, 2r}}$$

Using Cramer's rule, we know that:

$$(A^{-1})_{kl} = \frac{\det A_k(e_l)}{\det(A)}.$$

Therefore, one can easily compute,

$$(A^{-1})_{\tilde{u}}^{\tilde{v}} = \begin{pmatrix} -X_{j_1 i_2 \cdots i_r} X_{i_1 \cdots i_r}^{-1} & -X_{j_2 i_2 \cdots i_r} X_{i_1 \cdots i_r}^{-1} & \cdots & -X_{j_r i_2 \cdots i_r} X_{i_1 \cdots i_r}^{-1} \\ -X_{i_1 j_1 i_3 \cdots i_r} X_{i_1 \cdots i_r}^{-1} & -X_{i_1 j_2 i_3 \cdots i_r} X_{i_1 \cdots i_r}^{-1} & \cdots & -X_{i_1 j_r i_3 \cdots i_r} X_{i_1 \cdots i_r}^{-1} \\ \vdots & \vdots & \cdots & \vdots \\ \vdots & \vdots & \cdots & \vdots \\ -X_{i_1 i_2 \cdots j_1 i_r} X_{i_1 \cdots i_r}^{-1} & -X_{i_1 i_2 \cdots j_2 i_r} X_{i_1 \cdots i_r}^{-1} & \cdots & -X_{i_1 i_2 \cdots j_r i_r} X_{i_1 \cdots i_r}^{-1} \\ -X_{i_1 i_2 \cdots i_{r-1} j_1} X_{i_1 \cdots i_r}^{-1} & -X_{i_1 i_2 \cdots i_{r-1} j_2} X_{i_1 \cdots i_r}^{-1} & \cdots & -X_{i_1 i_2 \cdots i_{r-1} j_r} X_{i_1 \cdots i_r}^{-1} \end{pmatrix}$$

The Laplace expansion of the determinant, using the first row, gives us:

$$\det(A^{-1})_{\tilde{u}}^{\tilde{v}} = (X_{i_1 \cdots i_r})^{-1} \sum_{t=1}^r -X_{j_t i_2 \cdots i_r} c_{1,t} \quad (59)$$

where $c_{1,t}$ is the cofactor of the entry $((A^{-1})_{\tilde{u}}^{\tilde{v}})_{1t}$. Therefore, substituting these in Equation (58) we get:

$$X_{j_1 \cdots j_r} = (-1)^{r+1} \sum_{t=1}^r X_{j_t i_2 \cdots i_r} c_{1,t}. \quad (60)$$

Now, to calculate $c_{1,1}$, consider the same calculations with a matrix A' where we fix $j_1 = i_1$. This gives us:

$$(A'^{-1})_{\tilde{u}}^{\tilde{v}} = \begin{pmatrix} -1 & -X_{j_2 i_2 \dots i_r} X_{i_1 \dots i_r}^{-1} & \dots & -X_{j_r i_2 \dots i_r} X_{i_1 \dots i_r}^{-1} \\ 0 & -X_{i_1 j_2 i_3 \dots i_r} X_{i_1 \dots i_r}^{-1} & \dots & -X_{i_1 j_r i_3 \dots i_r} X_{i_1 \dots i_r}^{-1} \\ \cdot & \cdot & \dots & \cdot \\ \cdot & \cdot & \dots & \cdot \\ 0 & -X_{i_1 i_2 \dots j_2 i_r} X_{i_1 \dots i_r}^{-1} & \dots & -X_{i_1 \dots j_r i_r} X_{i_1 \dots i_r}^{-1} \\ 0 & -X_{i_1 i_2 \dots i_{r-1} j_2} X_{i_1 \dots i_r}^{-1} & \dots & -X_{i_1 i_2 \dots i_{r-1} j_r} X_{i_1 \dots i_r}^{-1} \end{pmatrix}$$

and from Equation (56) it follows that:

$$\begin{aligned} X_{i_1 j_2 \dots j_r} &= (-1)^r X_{i_1 \dots i_r} \det(A'^{-1})_{\tilde{u}}^{\tilde{v}} \\ &= (-1)^{r+1} X_{i_1 \dots i_r} c_{1,1} \end{aligned}$$

Hence, we get:

$$c_{1,1} = (-1)^{r+1} X_{i_1 j_2 \dots j_r} (X_{i_1 \dots i_r})^{-1}.$$

Similarly, one can get:

$$c_{1,t} = (-1)^{r+1} X_{j_1 \dots j_{t-1} i_1 j_{t+1} \dots j_r} (X_{i_1 \dots i_r})^{-1}.$$

By putting this in Equation (60) we get:

$$X_{i_1 \dots i_r} X_{j_1 \dots j_r} = \sum_{t=1}^r X_{j_t i_2 \dots i_r} X_{j_1 \dots j_{t-1} i_1 j_{t+1} \dots j_r} \quad (61)$$

which are exactly the Plücker relations in (1), up to renaming some indices. \square

We illustrate this result with an example given below.

Example C.3. Fix a 2×4 matrix

$$C = \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \end{pmatrix}$$

and set $X_{ij} := \det \begin{pmatrix} a_{1i} & a_{1j} \\ a_{2i} & a_{2j} \end{pmatrix}$. Suppose that X_{12} is invertible. Now, consider the following matrix:

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \text{which satisfies} \quad \det(A) = X_{12}.$$

By fixing $u = (3, 4)$ and $v = (1, 2)$, we get:

$$\det A_v^u = \det \begin{pmatrix} a_{13} & a_{14} \\ a_{23} & a_{24} \end{pmatrix} = X_{34}.$$

Moreover, using Cramer's rule, one can easily compute:

$$(A^{-1})_{\tilde{u}}^{\tilde{v}} = \begin{pmatrix} (A^{-1})_{13} & (A^{-1})_{14} \\ (A^{-1})_{23} & (A^{-1})_{24} \end{pmatrix} = \begin{pmatrix} X_{23}X_{12}^{-1} & X_{24}X_{12}^{-1} \\ -X_{13}X_{12}^{-1} & -X_{14}X_{12}^{-1} \end{pmatrix}.$$

Hence, substituting these values in Jacobi's identity,

$$\det(A_v^u) = \det(A) \det(A^{-1})_{\tilde{u}}^{\tilde{v}}$$

we get,

$$\begin{aligned} X_{34} &= X_{12} \det \begin{pmatrix} X_{23}X_{12}^{-1} & X_{24}X_{12}^{-1} \\ -X_{13}X_{12}^{-1} & -X_{14}X_{12}^{-1} \end{pmatrix} \\ \Rightarrow X_{12}X_{34} - X_{13}X_{24} + X_{14}X_{23} &= 0 \end{aligned}$$

which is exactly the Plücker relation in this case.

References

- [1] A. Berele and A. Regev, *Hook Young diagrams with applications to combinatorics and to representations of Lie superalgebras*, Adv. Math. **64** 118-175, (1987).
- [2] M. J. Bergvelt and J. M. Rabin *Supercurves, their Jacobians, and super KP equations*. Duke Math. J. **98** 1, 1–57, (1999).
- [3] A. Borel, *Essays in the history of Lie groups and algebraic groups*. History of Mathematics, **21**. American Mathematical Society and London Mathematical Society, (2001).
- [4] A. Brini, R. Q. Huang and A. Teolis, *The umbral symbolic method for supersymmetric tensors*, Adv. Math. **96** 123–193 (1992).
- [5] C. Carmeli, L. Caston, R. Fioresi, with an Appendix by I. Dimitrov, *Mathematical Foundations of Supersymmetry*, EMS Ser. Lect. Math., European Math. Soc., Zurich, (2011).

- [6] A. Cayley, *1889-1898. The collected mathematical papers of Arthur Cayley*, Cambridge Univ. Press. Cambridge.
- [7] K. Coulembier, P. Etingof, A. Kleshchev, V. Ostrik, *Super invariant theory in positive characteristic*. Eur. J. Math. **9**, 94 (2023).
- [8] K. Coulembier, P. Etingof, V. Ostrik, *On Frobenius exact symmetric tensor categories* (with Appendix A by Alexander Kleshchev) Ann. Math. **197** 3 1235-1279 (2023).
- [9] P. Deligne, G. I. Lehrer, R. B. Zhang *The first fundamental theorem of invariant theory for the orthosymplectic super group*. Advances in Math. **327** (2018).
- [10] R. Fiorese, M. A. Lledó, *The Minkowski and conformal superspaces. The classical and quantum descriptions*. World Scientific Publishing, Hackensack, NJ (2015).
- [11] W. Fulton, *Young Tableaux: With Applications to Representation theory and Geometry*. Cambridge University Press, Cambridge, (1997).
- [12] W. Fulton, J. Harris. Representation theory. A first course. Graduate Texts in Mathematics, Vol. 129. Springer-Verlag, New York (1991).
- [13] R. Goodman, N. R. Wallach. *Representations and Invariants of the Classical Groups*, Cambridge University Press, Cambridge (1998).
- [14] D. Grinberg *Notes on the combinatorial fundamentals of algebra*, <https://arxiv.org/abs/2008.09862>.
- [15] F. D. Grosshans, G. C. Rota, *Invariant theory, Young bitableaux and combinatorics*, Adv. in Math. **27** 63-92 (1978).
- [16] Hilbert, D. *Ueber die Theorie der algebraischen Formen*, Mathematische Annalen, **36** 4. 473–534 (1890).
- [17] C. Jacobi *De formatione et proprietatibus Determinantium* Journal fur die reine und angewandte Mathematik, **22** (1841).
- [18] P. H. Dondi, P. D. Jarvis, *Diagram and superfield techniques in the classical superalgebras*, J. Phys. A. Math. Gen. **14**, 547-563 (1981).

- [19] Y. I. Manin, *Gauge Field Theory and Complex Geometry*. Springer, Berlin (1997).
- [20] E. Shemyakova, T. Voronov, *On Super Plucker Embedding and Cluster Algebras*, *Selecta Mathematica*, **28**, Article number: 39 (2022).
- [21] E. Shemyakova, *On super cluster algebras based on super Plücker and super Ptolemy relations*, *Journal of Geometry and Physics*, 188, 104776 (2023).
- [22] I. Schur, *Über die rationalen Darstellungen der allgemeinen linearen Gruppe* (1927) in “I. Schur, *Gesammelte Abhandlungen III*,” pp. 68-85, Springer-Verlag, Berlin, (1973).
- [23] H. Weyl, *The Classical Groups*, Princeton Math. Series, No. 1, Princeton Univ. Press, Princeton, N.J. (1939).
- [24] Y. Zhang, *On the second fundamental theorem of invariant theory for the orthosymplectic super- group*. *J. Algebra* **501**, 394–434 (2018).