

# Herglotz's formalism, Eisenhart lift and Killing vectors

Krystian Bartczak\* and Piotr Kosiński†

*Faculty of Physics and Applied Informatics, University of Lodz, Poland*

## Abstract

The Eisenhart lift is extended to the case of dynamics described by action-dependent Lagrangians. The resulting Brinkmann metric depends on all coordinates. It is shown that the symmetries of the initial dynamics result in the existence of (conformal) Killing vectors of the Brinkmann metric. An example is given of equivalent time- and action-dependent descriptions which result in conformally equivalent metrics. It is also shown how the scaling invariance fits naturally into this scheme.

---

\*krystian.bartczak@edu.uni.lodz.pl

†piotr.kosinski@uni.lodz.pl

# Contents

|          |  |           |
|----------|--|-----------|
| <b>1</b> | <b>Introduction</b>                                      | <b>2</b>  |
| <b>2</b> | <b>Dimensional reduction of dynamics</b>                 | <b>5</b>  |
| <b>3</b> | <b>Eisenhart lift</b>                                    | <b>6</b>  |
| <b>4</b> | <b>Eisenhart lift - the general case</b>                 | <b>7</b>  |
| <b>5</b> | <b>A note on point transformations</b>                   | <b>9</b>  |
| <b>6</b> | <b>Symmetries, Noether's theorem and Killing vectors</b> | <b>10</b> |
| <b>7</b> | <b>Time versus action dependence</b>                     | <b>13</b> |
| <b>8</b> | <b>Scaling symmetry</b>                                  | <b>15</b> |
| <b>9</b> | <b>Summary</b>   | <b>17</b> |
|          | <b>Appendix</b>  | <b>18</b> |
| <b>A</b> | <b>Herglotz variational principle</b>                    | <b>18</b> |
| <b>B</b> | <b>Reparametrization invariance</b>                      | <b>20</b> |

## 1 Introduction

A geometric approach to classical dynamics, currently known as Eisenhart lift, had been originally proposed by Eisenhart in 1928 [1] and then rediscovered by a number of authors [2, 3, 4, 5, 6] (for a review see [7]).

Eisenhart lift relates the Newtonian dynamics in  $n$  dimensions to the geodesic motion in  $(n + 2)$  dimensional (pseudo-)Riemannian manifold equipped with the Brinkmann metric [8, 5]. Additional two variables are identified with time and action. Originally, the Eisenhart lift concerned conservative systems; however, the generalization to the time-dependent Lagrangians is quite straightforward [9]. In the latter case the components of the Brinkmann metric tensor depend, in general, on  $(n + 1)$  coordinates. The most general case

of the metric depending on  $(n + 2)$  coordinates is more problematic. This is because the last coordinate is identified, as mentioned above, with the action equal to the time integral of the Lagrange function. However, there exists a generalization of the Lagrangian (and Hamiltonian) formalism which admits the Lagrange functions depending on additional variable which can be viewed as a generalization of the action. This formalism is called Herglotz’s variational principle [10, 11, 12, 13]. It appears that the Herglotz dynamical systems occur naturally when the Lagrangian dynamics is reduced to lower-dimensional configuration space. Namely, the Lagrangian dynamics defined by the Lagrangian homogeneous of degree two in velocities, when reduced to the configuration space one dimension lower, is described, in general, by action-dependent Lagrangians [14]. Guided by this result we consider here the Eisenhart lift procedure for the metric depending on all coordinates. We show that the general Brinkmann metric in  $(n + 2)$ -dimensional space-time can be obtained by lifting the dynamics in  $n$  dimensions provided the latter is described by an action-dependent Lagrangians. Inherent in Eisenhart lift relation between symmetries, conservation laws and (conformal) Killing vectors is extended to the most general Brinkmann metric. It is also shown on the simple example how the equivalence of the descriptions of dissipative dynamics in terms of time-dependent and action-dependent Lagrangians is reflected in conformal equivalence of the relevant Brinkmann metrics.

One should admit that the Herglotz formalism is not particularly popular amongst physicists. However, it is closely related to the theory of contact transformations, a natural generalization of symplectic ones. Therefore, it finds applications in thermodynamics [15, 16, 17, 18], electromagnetic dissipative systems [19], hydrodynamics [20], generalization of Einstein’s equations in GR [21, 22], optimal control theory [23], dark energy [24], cosmology [25] or light propagation in general gravitational field [14]. It should be also mentioned that making the relevant metric action-dependent destroys the property that it provides the solution to vacuum Einstein’s equations. However, the latter property is by no means necessary in the context of Eisenhart lift the essence of which is the geometrization of dynamics.

The paper is organized as follows. In Sec. 2 we briefly review the general result concerning the “dimensional reduction” of dynamics described by the Lagrangians homogeneous in velocities. Then, in Sec. 3, the standard Eisenhart lift is described. Sec. 4 is devoted to the extension of the Eisenhart formalism to the case of general Brinkmann metrics. It is shown that the latter emerges from lifting the dynamics described by Herglotz’s variational

principle. Sec. 5 contains small remark concerning the role of point transformations in Brinkmann manifolds. It is shown further in Sec. 6 that the relation between symmetry transformations, Noether's theorem and Killing vectors, established in the framework of standard Eisenhart lift, extends naturally to the general case described here. In Sec. 7 a simple example is given how the equivalence between time-dependent and action-dependent descriptions manifests itself on the level of Brinkmann metrics. In Sec. 8 we consider the scaling symmetry showing that the Herglotz formalism provides a simple and natural framework for this kind of symmetry. The Eisenhart lift leads in this case to conformal transformations of Brinkmann metrics. Finally, Sec. 9 is devoted to some summary. In Appendix the Herglotz formalism is briefly described and some remarks concerning the reparametrization invariance are provided.

## 2 Dimensional reduction of dynamics

Eisenhart lift allows us to describe the dynamics of  $n$  degrees of freedom in terms of geodesic motion in appropriately chosen (pseudo-)Riemannian metric. In the original version, one starts with Lagrangian (or Hamiltonian) dynamics; the lifted dynamics is also described by the Lagrangian which is homogeneous of degree two in generalized velocities. However, in general it cannot be taken for granted that the reduction of geodesic motion to lower dimension results in Lagrangian dynamics. Still, the reduction procedure makes sense provided one admits a more general notion of Lagrangian dynamics; namely, one can consider Lagrangians depending on the action variable which results in generalization of variational principle known as Herglotz's variational principle [10, 11, 12, 13] (see Appendix for a brief description). Once such a generalization is admitted the following general theorem can be proven [14]. Consider a system with  $(n + 1)$  degrees of freedom  $q^\mu$ ,  $\mu = 0, 1, \dots, n$ , described by the Lagrangian  $L = L(\underline{q}, \underline{\dot{q}})$ ; here dot denotes the differentiation with respect to some evolution parameter  $\sigma$ . One assumes that:

- (i)  $L$  is nondegenerate,  $\det \left( \frac{\partial^2 L}{\partial \dot{q}^\mu \partial \dot{q}^\nu} \right) \neq 0$ ;
- (ii)  $L$  is a homogeneous function of generalized velocities  $\dot{q}^\mu$  of degree  $N \neq 1$ ;
- (iii)  $L$  does not depend explicitly on the evolution parameter  $\sigma$ .

It follows then that the value of the Lagrangian is a constant of motion. Let us fix the value  $C$  of  $L$  and consider the subset of trajectories obeying

$$L(\underline{q}, \underline{\dot{q}}) = C \tag{1}$$

Equation (1) can be solved, at least locally, with respect to  $\dot{q}^0$ :

$$\frac{dq^0}{d\sigma} = \mathcal{L}(q^0, q^1, \dots, q^n, \dot{q}^1, \dots, \dot{q}^n; C) \tag{2}$$

Then the projection of initial dynamics on the generalized coordinates  $q^1, \dots, q^n$  is described by the Herglotz equations:

$$\frac{\partial \mathcal{L}}{\partial q^i} - \frac{d}{d\sigma} \left( \frac{\partial \mathcal{L}}{\partial \dot{q}^i} \right) + \frac{\partial \mathcal{L}}{\partial q^0} \frac{\partial \mathcal{L}}{\partial \dot{q}^i} = 0, \quad i = 1, \dots, n \tag{3}$$

together with eq. (2). We conclude that the reduced configuration space is spanned by  $q^1, \dots, q^n$  while  $q^0$  plays the role of action.

The particularly interesting case concerns the Lagrangians  $L$  being the polynomials of degree 2 ( $N = 2$ ) in velocities:

$$L = \frac{1}{2} g_{\mu\nu}(\underline{q}) \dot{q}^\mu \dot{q}^\nu \quad (4)$$

Assume that  $g_{\mu\nu}(\underline{q})$  has the Lorentz signature. Consider the set of trajectories corresponding to  $L = 0$ . Then  $\mathcal{L}$ , defined by eq. (2), is homogeneous of first degree and defines reparametrization invariant dynamics. Therefore, one can choose the evolution parameter  $\sigma$  at will; for example, one can set  $\sigma = q^n$ , thereby reducing the number of degrees of freedom by two.

Finally, let us note that  $\mathcal{L}$  defines standard Lagrangian mechanics if  $L$  does not depend on  $q^0$ .

### 3 Eisenhart lift

The standard Eisenhart lift deals with the  $(n + 2)$ -dimensional manifold parametrized by the coordinates  $x^1, \dots, x^n, u, w$ , equipped with the so-called Brinkmann metric [8, 5]:

$$ds^2 = g_{\mu\nu}(\underline{x}) dx^\mu dx^\nu = h_{ij}(\underline{x}) dx^i dx^j + 2A_i(\underline{x}) dx^i du - 2V(\underline{x}) du^2 - 2du dw \quad (5)$$

Assuming  $\sigma$  to be an affine parameter one concludes that the geodesics are described by the Lagrangian:

$$L = \frac{1}{2} h_{ij}(\underline{x}) \dot{x}^i \dot{x}^j + A_i(\underline{x}) \dot{x}^i \dot{u} - V(\underline{x}) \dot{u}^2 - \dot{u} \dot{w} \quad (6)$$

Lagrange equations resulting from (6) are equivalent to the geodesic equations for Brinkmann metric:

$$\ddot{x}^i + \Gamma_{jk}^i \dot{x}^j \dot{x}^k + h^{ik} F_{jk} \dot{x}^j \dot{u} + h^{ik} \partial_k V \dot{u}^2 = 0 \quad (7)$$

$$\ddot{u} = 0 \quad (8)$$

$$\ddot{w} + \left( A_k \Gamma_{ij}^k - \frac{1}{2} (\partial_i A_j + \partial_j A_i) \right) \dot{x}^i \dot{x}^j + \left( A_j h^{jk} F_{ik} + 2 \partial_i V \right) \dot{x}^i \dot{u} + A_j h^{ij} \partial_i V \dot{u}^2 = 0 \quad (9)$$

here  $\Gamma_{jk}^i$  are Christoffel symbols for the  $h_{ij}$  metric,  $h^{ij}$  is the inverse of  $h_{ij}$  and  $F_{ij} = \partial_i A_j - \partial_j A_i$ .

Eq. (8) implies that  $u$  may be viewed as the affine parameter (the latter is defined up to an affine transformation). Choosing additionally the trajectories with  $L = 0$  one concludes that

$$\dot{w} = \dot{u} \left( \frac{1}{2} h_{ij} \frac{\dot{x}^i \dot{x}^j}{\dot{u}^2} + A_i \frac{\dot{x}^i}{\dot{u}} - V \right) \quad (10)$$

Eqs. (7) and (10) imply finally

$$\frac{d^2 x^i}{du^2} + \Gamma_{jk}^i \frac{dx^j}{du} \frac{dx^k}{du} + h^{ik} F_{jk} \frac{dx^j}{du} + h^{ik} \partial_k V = 0 \quad (11)$$

$$\frac{dw}{du} = \frac{1}{2} h_{ij} \frac{dx^i}{du} \frac{dx^j}{du} + A_i \frac{dx^i}{du} - V = \mathcal{L} \left( \underline{x}, \frac{d\underline{x}}{du} \right) \quad (12)$$

By noting that (11) are the Lagrange equations resulting from  $\mathcal{L} \left( \underline{x}, \frac{d\underline{x}}{du} \right)$  we conclude that  $u$  may be identified with time while  $w$  with the action for particle moving in the curved space (with the metrics  $h_{ij}$ ) under the influence of the scalar and vector potentials  $V$  and  $A_i$ , respectively.

It is not difficult to generalize the above reasoning to the case of  $u$ -dependent (i.e.  $t$ -dependent) metric and potentials.

## 4 Eisenhart lift - the general case

Consider now the general Brinkmann metric depending on all coordinate  $x^\mu = (x^k, u, w)$ :

$$ds^2 = h_{ij}(\underline{x}, u, w) dx^i dx^j + 2A_i(\underline{x}, u, w) dx^i du - 2V(\underline{x}, u, w) du^2 - 2du dw \quad (13)$$

or

$$L = \frac{1}{2} h_{ij}(\underline{x}, u, w) \dot{x}^i \dot{x}^j + A_i(\underline{x}, u, w) \dot{x}^i \dot{u} - V(\underline{x}, u, w) \dot{u}^2 - \dot{u} \dot{w} \quad (14)$$

$L$  is again an ordinary Lagrangian describing dynamics of  $(n+2)$  degrees of freedom. We shall show that, in the spirit of the result described in Sec. 2,  $L$  can be viewed as describing the Eisenhart-like lift of  $n$ -dimensional Herlotz dynamics. The relevant Lagrange equations (geodesic equations for the

metric (13)) read:

$$h_{kj}\ddot{x}^j + \Gamma_{kij}\dot{x}^i\dot{x}^j + \dot{u}^2 \partial_k V + A_k \ddot{u} + F_{ik}\dot{x}^i\dot{u} + \partial_u h_{jk}\dot{x}^j\dot{u} + \partial_u A_k \dot{u}^2 + \partial_w h_{jk}\dot{x}^j\dot{w} + \partial_w A_k \dot{u}\dot{w} = 0 \quad (15)$$

$$\ddot{u} + \frac{1}{2} \partial_w h_{ij}\dot{x}^i\dot{x}^j + \partial_w A_i \dot{x}^i\dot{u} - \partial_w V \dot{u}^2 = 0 \quad (16)$$

$$\ddot{w} + \frac{1}{2} \partial_u h_{ij}\dot{x}^i\dot{x}^j + \partial_u A_i \dot{x}^i\dot{u} - \partial_u V \dot{u}^2 - \frac{d}{d\sigma} (A_i \dot{x}^i - 2V\dot{u}) = 0 \quad (17)$$

Again, let us impose the condition  $L = 0$ ;  
this yields

$$\dot{w} = \dot{u} \left( \frac{1}{2} h_{ij} \frac{\dot{x}^i \dot{x}^j}{\dot{u}^2} + A_i \frac{\dot{x}^i}{\dot{u}} - V \right) \quad (18)$$

Eq. (18) is reparametrization invariant. It can be rewritten as

$$\frac{dw}{du} = \frac{1}{2} h_{ij} \frac{dx^i}{du} \frac{dx^j}{du} + A_i \frac{dx^i}{du} - V \equiv \mathcal{L} \left( \underline{x}, \frac{d\underline{x}}{d\underline{u}}, u, w \right) \quad (19)$$

Eq. (19) suggests that, in the framework of Herglotz formalism,  $u$  can be identified with time while  $w$  – with action. In order to show that this interpretation is correct we consider eq. (15) and (16) (eq. (17) is redundant once we imposed the constraint (18)). Dividing them by  $\dot{u}^2$  one finds:

$$\frac{\ddot{u}}{\dot{u}^2} = -\partial_w \mathcal{L} \quad (20)$$

$$h_{kj} \frac{\ddot{x}^j}{\dot{u}^2} + \Gamma_{kij} \frac{\dot{x}^i \dot{x}^j}{\dot{u}^2} + \partial_k V + A_k \frac{\ddot{u}}{\dot{u}^2} + F_{ik} \frac{\dot{x}^i}{\dot{u}} + \partial_u \left( h_{jk} \frac{\dot{x}^j}{\dot{u}} + A_k \right) + \partial_w \left( h_{jk} \frac{\dot{x}^j}{\dot{u}^2} + \frac{A_k}{\dot{u}} \right) \dot{w} = 0 \quad (21)$$

Moreover,

$$\frac{d}{d\sigma} = \dot{u} \frac{d}{du} \quad (22)$$

$$\frac{1}{\dot{u}^2} \frac{d^2}{d\sigma^2} = \frac{\ddot{u}}{\dot{u}^2} \frac{d}{du} + \frac{d^2}{du^2} = -\partial_w \mathcal{L} \frac{d}{du} + \frac{d^2}{du^2} \quad (23)$$



Using (19), (22) and (23) one can rewrite eq. (21) in terms of  $u$ -derivatives:

$$\begin{aligned} & h_{kj} \frac{d^2 x^j}{du^2} + \Gamma_{kij} \frac{dx^i}{du} \frac{dx^j}{du} + \\ & + \partial_k V + F_{ik} \frac{dx^i}{du} + \partial_u \left( h_{ik} \frac{dx^i}{du} + A_k \right) + \\ & + \partial_w \left( h_{ik} \frac{dx^i}{du} + A_k \right) \mathcal{L} - \left( h_{ik} \frac{dx^i}{du} + A_k \right) \partial_w \mathcal{L} = 0 \end{aligned} \quad (24)$$

It is now easy to check (again using (19)) that eq. (24) is the Herglotz equations for  $\mathcal{L}$ :

$$\frac{\partial \mathcal{L}}{\partial x^k} - \frac{d}{d\sigma} \left( \frac{\partial \mathcal{L}}{\partial \left( \frac{dx^k}{du} \right)} \right) + \frac{\partial \mathcal{L}}{\partial w} \frac{\partial \mathcal{L}}{\partial \left( \frac{dx^k}{du} \right)} = 0 \quad (25)$$

which, together with (19), forms the basis of Herglotz variational principle.

Let us stress that the dynamics resulting from the constraint  $L = 0$  is reparametrization invariant. Therefore, we can keep  $\sigma$  as an evolution parameter when deriving the reduced equations of motion and put  $\sigma = u$  at the very end (this is briefly sketched in Appendix). Concluding, we have shown that the Eisenhart lift can be applied to the quadratic, action- and time-dependent Lagrangian. The relevant  $(n+2)$ -dimensional metric takes the Brinkmann form with the metric tensor components depending also on  $w$ .

## 5 A note on point transformations

The Eisenhart lift allows us to relate the point transformations of initial dynamics to coordinate transformations in  $(n+2)$ -dimensional manifold equipped with Brinkmann metric. Consider an arbitrary coordinate transformation in the latter

$$x'^{\mu} = x'^{\mu}(x^{\nu}), \quad \mu, \nu = 0, 1, \dots, n, n+1 \quad (26)$$

with  $x^0 = w$ ,  $x^{n+1} = u$ . In general, (26) does not preserve the Brinkmann form (13) of the metric. This is because in the Herglotz formalism the transformation rule for the Lagrangian takes, in general, the form of implicit relation<sup>1</sup>. As a result the new Lagrangian is no longer quadratic in velocities

---

<sup>1</sup>We thank Tomasz Ruciński for the enlightening discussion on this point.

(this is also the case in the standard Lagrangian theory if the new time variable depends on old coordinates). However, in view of the theorem described in Sec. 2, eq. (26) should imply the correct rule for the transformation of action-dependent Lagrangian. Eq. (26) yields

$$\frac{dx'^0}{d\sigma} = \frac{\partial x'^0}{\partial x^0} \frac{dx^0}{d\sigma} + \sum_{k=1}^n \frac{\partial x'^0}{\partial x^k} \frac{dx^k}{d\sigma} + \frac{\partial x'^0}{\partial x^{n+1}} \frac{dx^{n+1}}{d\sigma} \quad (27)$$

Multiplying by  $\left(\frac{dx^{n+1}}{d\sigma}\right)^{-1}$  one finds

$$\left(\frac{dx^{n+1}}{d\sigma}\right)^{-1} \frac{dx'^0}{d\sigma} = \frac{\partial x'^0}{\partial x^0} \frac{dx^0}{d\sigma} \left(\frac{dx^{n+1}}{d\sigma}\right)^{-1} + \sum_{k=1}^n \frac{\partial x'^0}{\partial x^k} \frac{dx^k}{d\sigma} \left(\frac{dx^{n+1}}{d\sigma}\right)^{-1} + \frac{\partial x'^0}{\partial x^{n+1}} \quad (28)$$

or

$$\frac{dx'^{n+1}}{dx^{n+1}} \frac{dx'^0}{dx'^{n+1}} = \frac{\partial x'^0}{\partial x^0} \frac{dx^0}{dx^{n+1}} + \sum_{k=1}^n \frac{\partial x'^0}{\partial x^k} \frac{dx^k}{dx^{n+1}} + \frac{\partial x'^0}{\partial x^{n+1}} \quad (29)$$

Taking into account that  $x^0$  and  $x^{n+1}$  are identified with the action and time variables respectively, we conclude that eq. (29) coincides with the transformation rule (A10).

## 6 Symmetries, Noether's theorem and Killing vectors

The famous Noether's theorem relates the symmetries to conservation laws. In the framework of Eisenhart formalism the trajectories are lifted to geodesics in  $(n+2)$ -dimensional (pseudo-)Riemannian manifold. Brinkmann metric is determined by the initial Lagrangian; therefore, the symmetries of the latter can be lifted to the symmetries of the metric which, in turn, are determined by the relevant Killing vectors. Actually, since we are considering null geodesics, it is sufficient to restrict ourselves to conformal Killing vectors, i.e. we are really interested in the classes of conformally equivalent metrics. One can single out uniquely one metric from each class by choosing the normalization condition  $g_{uw} = -1$ .

Let us now extend the reasoning sketched above to the case of action-dependent Lagrangians. Assume that the transformations

$$x^i \mapsto x^i + \delta x^i(\underline{x}, u, w) \quad (30)$$

$$u \mapsto u + \delta u(\underline{x}, u, w) \quad (31)$$

$$w \mapsto w + \delta w(\underline{x}, u, w) \quad (32)$$

define the symmetry of the Lagrangian (19). The symmetry condition (A11) reads here

$$\begin{aligned} & \left( \frac{1}{2} \partial_k h_{ij} \dot{x}^i \dot{x}^j + \partial_k A_i \dot{x}^i - \partial_k V \right) \delta x^k + \\ & + (h_{ki} \dot{x}^i + A_k) (\delta \dot{x}^k - \dot{x}^k \delta \dot{u}) + \left( \frac{1}{2} \partial_u h_{ij} \dot{x}^i \dot{x}^j + \partial_u A_i \dot{x}^i - \partial_u V \right) \delta u + \\ & + \left( \frac{1}{2} \partial_w h_{ij} \dot{x}^i \dot{x}^j + \partial_w A_i \dot{x}^i - \partial_w V \right) \delta w + \left( \frac{1}{2} h_{ij} \dot{x}^i \dot{x}^j + A_i \dot{x}^i - V \right) \delta \dot{u} - \delta \dot{w} = 0 \end{aligned} \quad (33)$$

where, for notational simplicity, dot denotes differentiation with respect to  $u$  and

$$\delta \dot{x}^k = \dot{x}^l \partial_l \delta x^k + \partial_u \delta x^k + \dot{w} \partial_w \delta x^k \quad (34)$$

$$\delta \dot{u} = \dot{x}^k \partial_k \delta u + \partial_u \delta u + \dot{w} \partial_w \delta u \quad (35)$$

$$\delta \dot{w} = \dot{x}^k \partial_k \delta w + \partial_u \delta w + \dot{w} \partial_w \delta w \quad (36)$$

and

$$\dot{w} = \frac{1}{2} h_{ij} \dot{x}^i \dot{x}^j + A_i \dot{x}^i - V \quad (37)$$

Eqs. (34)-(37), when inserted into eq. (33) lead to the identity with the left-hand side being fourth degree polynomial in velocities  $\dot{x}^i$ . Therefore, one obtains five equations for the coefficients in front of consecutive powers of velocities. Fourth degree term yields

$$(h_{ij} h_{kl} + h_{ik} h_{jl} + h_{il} h_{jk}) \partial_w \delta u = 0 \quad (38)$$

and, after multiplying by  $h^{ij}h^{kl}$ ,

$$\partial_w \delta u = 0 \quad (39)$$

Third degree term leads to the relation

$$\begin{aligned} & (h_{il}h_{jk} + h_{jl}h_{ik} + h_{ij}h_{kl}) \partial_w \delta x^l + \\ & - (h_{ik} \partial_j + h_{ij} \partial_k + h_{jk} \partial_i) \delta u = 0 \end{aligned} \quad (40)$$

where (39) has been already used. By multiplying eq. (40) by  $h^{jk}$  one finds

$$h_{il} \partial_w \delta x^l = \partial_i \delta u \quad (41)$$

Putting back (41) into (40) one finds that it is satisfied identically.

The remaining identities, corresponding to the terms of second, first and zero degree, read, respectively

$$\begin{aligned} & (\delta u \partial_u - \partial_u \delta u) h_{ij} + (\delta w \partial_w - \partial_w \delta w) h_{ij} + A_k \partial_w \delta x^k h_{ij} + \\ & + (\partial_i \delta u A_j + \partial_j \delta u A_i) + (h_{ik} \partial_j + h_{jk} \partial_i) \delta x^k + \delta x^k \partial_k h_{ij} = 0 \end{aligned} \quad (42)$$

$$\begin{aligned} & (\delta w \partial_w - \partial_w \delta w) A_i + \delta u \partial_u A_i - 2V \partial_i \delta u - \partial_i \delta w + \\ & + (\partial_k A_i + A_k \partial_i) \delta x^k + h_{ik} \partial_u \delta x^k + A_i A_k \partial_w \delta x^k = 0 \end{aligned} \quad (43)$$

$$-\delta x^k \partial_k V - \delta u \partial_u V - \partial_u \delta u V - \delta w \partial_w V - \partial_w \delta w V + V \partial_w \delta w + A_k \partial_u \delta x^k - A_k \partial_w \delta x^k V = 0 \quad (44)$$

As in the standard case one can now find the Killing vector corresponding to the symmetries under consideration. Namely, we put

$$\begin{aligned} K & \equiv K^\mu \partial_\mu = K^i \partial_i + K^u \partial_u + K^w \partial_w = \\ & = \delta x^i \partial_i + \delta u \partial_u + \delta w \partial_w \end{aligned} \quad (45)$$

Using the identities (39), (41)-(44) one finds that  $K$  is a conformal vector field:

$$\nabla_\mu K_\nu + \nabla_\nu K_\mu = \lambda g_{\mu\nu} \quad (46)$$

with

$$\lambda \equiv \partial_u \delta u + \partial_w \delta w - A_i \partial_w \delta x^i \quad (47)$$

The conserved quantity related to the Killing field  $K$  reads

$$K_\mu \frac{dx^\mu}{d\sigma} = g_{\mu\nu} K^\mu \frac{dx^\nu}{d\sigma} = \frac{du}{d\sigma} Q \quad (48)$$

$$Q \equiv \left( h_{ij} \dot{x}^j + A_i \right) \delta x^i - \left( \frac{1}{2} h_{ij} \dot{x}^i \dot{x}^j + V \right) \delta u - \delta w \quad (49)$$

where we have used eqs. (5), (12) and (45). In the standard case  $Q$  is the Noether charge ( $\delta w$  accounts for total time derivative which, in general, enters Noether identity). Due to eq. (8)  $\frac{du}{d\sigma}$  is a constant. However, in the general case, eq. (16), only the product  $\frac{du}{d\sigma} Q$  is a constant of motion. The  $u$  variable, which plays the role of time in reduced dynamics is not any longer an affine parameter. Given any parametrization of a geodesics one can find the affine parameter by solving an ordinary differential equation which, however, depends on the choice of particular geodesic. In this sense the charge (48) is nonlocal. Specifically, one has

$$\frac{d}{d\sigma} \left( K_\mu \frac{dx^\mu}{d\sigma} \right) = \frac{d^2 u}{d\sigma^2} Q + \frac{du}{d\sigma} \frac{dQ}{d\sigma} \quad (50)$$

Using (20) we obtain

$$\begin{aligned} \frac{d}{d\sigma} \left( K_\mu \frac{dx^\mu}{d\sigma} \right) &= - \left( \frac{du}{d\sigma} \right)^2 \frac{\partial \mathcal{L}}{\partial w} Q + \frac{du}{d\sigma} \frac{dQ}{d\sigma} = \\ &= \left( \frac{du}{d\sigma} \right)^2 \left( - \frac{\partial \mathcal{L}}{\partial w} Q + \frac{dQ}{du} \right) = \\ &= \left( \frac{du}{d\sigma} \right)^2 e^{\int^u \frac{\partial \mathcal{L}}{\partial w} du'} \frac{d}{du} \left( e^{-\int^u \frac{\partial \mathcal{L}}{\partial w} du'} Q \right) \end{aligned} \quad (51)$$

in agreement with the generalized Noether theorem (cf. Appendix, eq. (A15))

## 7 Time versus action dependence

It is well known that the dissipative systems, demanding time dependent Lagrangians for their description, can be also described with help of action

dependent Lagrangians [9]. As a simple example consider the Lagrangian

$$L = e^{\gamma t} \left( \frac{1}{2} \dot{\vec{x}}^2 - V(\vec{x}) \right) \quad (52)$$

Lagrange equations of motion read:

$$\ddot{\vec{x}} + \gamma \dot{\vec{x}} + \frac{\partial V}{\partial \vec{x}} = 0 \quad (53)$$

Equivalently, eq. (53) can be derived from the following Lagrangian

$$\tilde{L} = \frac{1}{2} \dot{\vec{x}}^2 - V(\vec{x}) - \gamma \tilde{S} \quad (54)$$

Solving  $\dot{S} = L$  with the initial condition  $S(t_0) = 0$  one finds

$$S(t) = \int_{t_0}^t e^{\gamma t'} \left( \frac{1}{2} \dot{\vec{x}}^2(t') - V(\vec{x}(t')) \right) dt' \quad (55)$$

Analogously, solving  $\dot{\tilde{S}} = \tilde{L}$  with the same initial conditions yields

$$\tilde{S}(t) = e^{-\gamma t} \int_{t_0}^t e^{\gamma t'} \left( \frac{1}{2} \dot{\vec{x}}^2(t') - V(\vec{x}(t')) \right) dt' \quad (56)$$

Comparing eqs. (55) and (56) we infer that the above descriptions are related by transformation

$$\tilde{\vec{x}} = \vec{x} \quad (57)$$

$$\tilde{t} = t \quad (58)$$

$$\tilde{S} = e^{-\gamma t} S \quad (59)$$

Indeed, one finds from (A10)

$$\tilde{L} = e^{-\gamma t} L - \gamma \tilde{S} \quad (60)$$

On the level of Brinkmann metrics one has

$$ds^2 = e^{\gamma t} d\vec{x}^2 - 2e^{\gamma t} V du^2 - 2 du dw \quad (61)$$

$$d\tilde{s}^2 = d\tilde{x}^2 - 2(V + \gamma\tilde{w}) d\tilde{u}^2 - 2 d\tilde{u} d\tilde{w} \quad (62)$$

Eqs. (57)-(59) suggest the following transformation:

$$\tilde{x} = \vec{x} \quad (63)$$

$$\tilde{u} = u \quad (64)$$

$$\tilde{w} = e^{-\gamma u} w \quad (65)$$

which implies the following relation

$$d\tilde{s}^2 = e^{-\gamma\tilde{u}} ds^2 \quad (66)$$

The metrics  $ds^2$  and  $d\tilde{s}^2$  are conformally equivalent so they share the same set of null geodesics.

## 8 Scaling symmetry

Another nice example of the application of Herglotz formalism is the scaling symmetry.

Let  $L(q, \frac{dq}{dt}, t)$  be standard (i.e. action-independent) Lagrangian. Consider the following generalized point transformations:

$$\begin{aligned} q' &= q'(q, t) \\ t' &= t'(q, t) \\ S' &= \Lambda S + f(q, t), \quad \Lambda \in \mathbb{R} \end{aligned} \quad (67)$$

Then the symmetry condition (A11) takes the form:

$$\mathcal{L} \left( q', \frac{dq'}{dt'}, t' \right) \frac{dt'}{dt} = \Lambda \mathcal{L} \left( q, \frac{dq}{dt}, t \right) + \frac{df(q, t)}{dt} \quad (68)$$

which coincides with eq. (II.1) from [26]. The infinitesimal form of eq. (67) reads:

$$\begin{aligned} t' &= t + \delta t(q, t) \\ q' &= q + \delta q(q, t) \\ S' &= S + \delta \Lambda S + \delta f(q, t) \end{aligned} \quad (69)$$

and the Noether identity (A15) reads ( $\frac{\partial \mathcal{L}}{\partial S} = 0$ !)

$$\frac{d}{dt} \left( \frac{\partial \mathcal{L}}{\partial \dot{q}} \delta q - H \delta t - \delta f - \delta \Lambda S \right) = 0 \quad (70)$$

Now, taking into account that  $S$  is the standard action,  $S = \int^t L d\tau$ , we arrive at the eq. (II.4) from [26].

Consider now the Eisenhart lift. Assume that the Lagrangian (19) is invariant under the transformation:

$$\begin{aligned} x'^i &= x'^i(\underline{x}, u) \\ u' &= u'(u) \end{aligned} \quad (71)$$

in the sense that eq. (68) is obeyed (note that, due to the quadratic dependence of  $\mathcal{L}$  on velocities, this can happen only provided  $u'$  is a function of  $u$  only):

$$\begin{aligned} & \left( \frac{1}{2} h_{ij}(\underline{x}', u') \frac{dx'^i}{du'} \frac{dx'^j}{du'} + A_i(\underline{x}', u') \frac{dx'^i}{du'} - V(\underline{x}', u') \right) \frac{du'}{du} = \\ &= \Lambda \left( \frac{1}{2} h_{ij}(\underline{x}, u) \frac{dx^i}{du} \frac{dx^j}{du} + A_i(\underline{x}, u) \frac{dx^i}{du} - V(\underline{x}, u) \right) + \frac{df}{du}(\underline{x}, u) \end{aligned} \quad (72)$$

which can be rewritten as:

$$\begin{aligned} & \frac{1}{2} h_{ij}(\underline{x}', u') dx'^i dx'^j + A_i(\underline{x}', u') dx'^i du' - V(\underline{x}', u') du'^2 = \\ &= \Lambda \frac{du'}{du} \left( \frac{1}{2} h_{ij}(\underline{x}, u) dx^i dx^j + A_i(\underline{x}, u) dx^i du - V(\underline{x}, u) du^2 \right) + \frac{du'}{du} du df \end{aligned} \quad (73)$$

Let us supply (71) with the transformation rule:

$$w' = \Lambda w + f(\underline{x}, u) \quad (74)$$



so that

$$dw' = \Lambda dw + df \quad (75)$$

and

$$du' dw' = \frac{du'}{du} \Lambda du dw + \frac{du'}{du} du df \quad (76)$$

It follows then from eqs. (13), (73) and (76) that, under the transformations (71), (74) the metric transforms conformally:

$$(ds')^2 = \Lambda \frac{du'}{du} ds^2 \quad (77)$$

## 9 Summary

We have shown that the idea of Eisenhart lift can be extended to the case of general Brinkmann metric with components depending on all coordinates. The price to be paid is that one has to start with Herglotz's formalism which is more general than the Lagrangian one. So, starting with an action-dependent Lagrangian and performing essentially the same steps as in the case of standard Eisenhart lift one arrives at the most general form of Brinkmann metric. The important difference is that the coordinate  $u$ , identified with the physical time, cannot any longer serve as an affine parameter for null geodesics in Brinkmann metric (cf. eqs. (8) and (16)). This has profound consequences. The conserved charge, eq. (48), involves the derivative of  $u$  with respect to an affine parameter which, when expressed in terms of  $u$ , results in nonlocal expression for the conserved quantity, in agreement with known results [27, 28, 29].

We have also shown on a simple example that the equivalence between time-dependent and action-dependent descriptions of dynamics translates into the conformal equivalence of relevant Brinkmann metrics. It would be interesting to find the Hamiltonian description of the above reduction procedure. This is important because the Noether theorem can be extended to Hamiltonian formalism: the conserved quantities result also from symmetries described by canonical transformations which do not reduce to the point ones. On the level of lifted dynamics this should correspond to the existence of (conformal) Killing tensors.

# Appendix

## A Herglotz variational principle

Standard Lagrangian formalism can be generalized by considering the Lagrange functions which depend on additional variable:

$$L = L(q, \dot{q}, t, S) \quad (\text{A1})$$

(for simplicity we consider one degree of freedom; the generalization to any finite number of degrees of freedom is straightforward). Let us choose a function  $q = q(t)$ , defined on the interval  $[t_0, t_1]$ , and consider the initial value problem

$$\frac{dS}{dt} = L(q(t), \dot{q}(t), t, S) \quad (\text{A2})$$

$$S(t_0) = S_0 \quad (\text{A3})$$

where  $S_0$  is some fixed constant. Herglotz's variational principle states that the physical trajectory is determined by the following condition:

$$\delta S(t_1) = 0 \quad (\text{A4})$$

for all variations  $q(t) \mapsto q(t) + \delta q(t)$  obeying

$$\delta q(t_0) = 0 = \delta q(t_1) \quad (\text{A5})$$

In particular, if  $L$  does not depend on  $S$  we obtain the standard Lagrangian formalism.

Herglotz formalism shares many properties of the Lagrangian one. Euler-Lagrange equations are replaced by eq. (A2) together with

$$\frac{\partial L}{\partial q} - \frac{d}{dt} \left( \frac{\partial L}{\partial \dot{q}} \right) + \frac{\partial L}{\partial S} \frac{\partial L}{\partial \dot{q}} = 0 \quad (\text{A6})$$

The generalized point transformations

$$t' = t'(t, q, S) \quad (\text{A7})$$

$$q' = q'(t, q, S) \quad (\text{A8})$$

$$S' = S'(t, q, S) \quad (\text{A9})$$

preserve the form of the equations of motion provided the Lagrangian transforms according to the formula:

$$L' \frac{dt'}{dt} = \frac{\partial S'}{\partial S} L + \frac{\partial S'}{\partial q} \frac{dq}{dt} + \frac{\partial S'}{\partial t} \quad (\text{A10})$$

Point transformations (A7)-(A9) define a symmetry if the new Lagrangian has the same functional form, i.e.

$$L \left( q', \frac{dq'}{dt'}, t', S' \right) \frac{dt'}{dt} = \frac{\partial S'}{\partial S} L \left( q, \frac{dq}{dt}, t, S \right) + \frac{\partial S'}{\partial q} \frac{dq}{dt} + \frac{\partial S'}{\partial t} \quad (\text{A11})$$

Continuous symmetry transformations generate, via generalized Noether's theorem, the corresponding conservation laws. Consider namely the infinitesimal version of (A7)-(A9):

$$t' = t + \delta t(t, q, S) \quad (\text{A12})$$

$$q' = q + \delta q(t, q, S) \quad (\text{A13})$$

$$S' = S + \delta S(t, q, S); \quad (\text{A14})$$

then, for physical trajectories, the following relation is fulfilled

$$\frac{d}{dt} \left( e^{-\int^t \frac{\partial L}{\partial S} dt'} \left( L \delta t + \frac{\partial L}{\partial \dot{q}} (\delta q - \dot{q} \delta t) - \delta S \right) \right) = 0 \quad (\text{A15})$$

Eq. (A15) reduces to the standard conservation law provided  $\frac{\partial L}{\partial S} = 0$ . In general, however, (A15) is more complicated since the exponential term depends on the whole past history of the system.

Therefore, it ceases to be a useful tool of integration the equations of motion. It can be shown that it is also possible to generalize appropriately the Hamiltonian formalism, Poisson brackets, canonical transformations etc. to the case of action dependent Lagrangians. We will not dwell on it here.

## B Reparametrization invariance

The Lagrangian defined by eq. (10) in the main text reads

$$\tilde{\mathcal{L}} = \frac{1}{2} h_{ij} \frac{\dot{x}^i \dot{x}^j}{\dot{u}} + A_i \dot{x}^i - V \dot{u} \quad (\text{B1})$$

We have shown that the final equations of motion can be written in terms of  $u$ -derivatives only. This can be also inferred from the homogeneity of  $\tilde{\mathcal{L}}$  in velocities. Indeed, (B1) implies:

$$\dot{x}^k \frac{\partial \tilde{\mathcal{L}}}{\partial \dot{x}^k} + \dot{u} \frac{\partial \tilde{\mathcal{L}}}{\partial \dot{u}} = \tilde{\mathcal{L}} \quad (\text{B2})$$

The equations of motion read:

$$\frac{\partial \tilde{\mathcal{L}}}{\partial x^k} - \frac{d}{d\sigma} \left( \frac{\partial \tilde{\mathcal{L}}}{\partial \dot{x}^k} \right) + \frac{\partial \tilde{\mathcal{L}}}{\partial w} \frac{\partial \tilde{\mathcal{L}}}{\partial \dot{x}^k} = 0 \quad (\text{B3})$$

$$\frac{\partial \tilde{\mathcal{L}}}{\partial u} - \frac{d}{d\sigma} \left( \frac{\partial \tilde{\mathcal{L}}}{\partial \dot{u}} \right) + \frac{\partial \tilde{\mathcal{L}}}{\partial w} \frac{\partial \tilde{\mathcal{L}}}{\partial \dot{u}} = 0 \quad (\text{B4})$$

Taking into account that  $\tilde{\mathcal{L}} = \tilde{\mathcal{L}}(\underline{x}, u, w, \underline{\dot{x}}, \dot{u})$  and  $\dot{w} = \tilde{\mathcal{L}}$  one easily finds from (B2):

$$\dot{x}^k (\text{B3})_k + \dot{u} (\text{B4}) \equiv 0 \quad (\text{B5})$$

as an identity. Therefore, (B3) implies (B4). Now,  $\tilde{\mathcal{L}}$  obeys:

$$\tilde{\mathcal{L}} \left( \underline{x}, u, w, \frac{dx}{d\sigma'}, \frac{du}{d\sigma'} \right) = \frac{d\sigma}{d\sigma'} \tilde{\mathcal{L}} \left( \underline{x}, u, w, \frac{dx}{d\sigma}, \frac{du}{d\sigma} \right) \quad (\text{B6})$$

and it is straightforward to check that (B3) is invariant under the reparametrization  $\sigma \mapsto \sigma'$ .

## References

- [1] L. Eisenhart. *Ann. Math.* 30 (1928), 591–606.
- [2] J. Gomis, J. Pons. *Phys. Lett. A* 66 (1978), 463–465.
- [3] C. Duval, G. Burdet, H. Künzle, M. Perrin. *Phys. Rev. D* 31 (1985), 1841–1853.
- [4] A. Balachandran, H. Gomm, R. Sorkin. *Nucl. Phys. B* 281 (1987), 573–612.
- [5] C. Duval, G. Gibbons, P. Horvathy. *Phys. Rev. D* 43 (1991), 3907–3922.
- [6] A. Fordy, A. Galajinsky. *The Euro. Phys. Journ. C* 79 (2019), 301.
- [7] M. Cariglia. *Rev. Mod. Phys.* 86 (2014), 1283–1336.
- [8] H. Brinkmann. *Math. Ann.* 94 (1925), 119–145.
- [9] M. Cariglia, C. Duval, G. Gibbons, P. Horvathy. *Ann. Phys.* 373 (2016), 631–654.
- [10] G. Herglotz. *Lectures at the University of Göttingen*. Göttingen, 1930.
- [11] R. Guenther, C. Guenther, J. Gottsch. *The Herglotz Lectures on Contact Transformations and Hamiltonian Systems*. Lecture Notes in Nonlinear Analysis. Juliusz Schauder Center for Nonlinear Studies. Nicholas Copernicus University, 1996.
- [12] M. Lazo, J. Paiva, J. Amaral, G. Frederico. *Journ. Math. Phys.* 59 (2018), 032902.
- [13] E. Massa, E. Pagani. *Journ. Math. Phys.* 64 (2023), 102902.
- [14] J. Piwnik, J. Gonera, P. Kosiński. *Nucl. Phys. B* 1010 (2025), 116744.
- [15] A. Simoes, M. de León, M. Valcázar, D. M. de Diego. *Proc. R. Soc. Lond. A* 476 (2020), 20200244.
- [16] R. Mrugała, J. Nulton, J. Schön, P. Salamon. *Rep. Math. Phys.* 29 (1991), 109.
- [17] M. Grmela. *Entropy* 16 (2014), 1652.
- [18] D. Eberard, B. Maschke, A. van der Schaft. *Rep. Math. Phys.* 60 (2007), 175.

- [19] J. Gaset, A. Marín-Salvador. *Int. J. Geom. Methods Mod. Phys.* 19 (2022), 2250156.
- [20] J. Etnyre, R. Ghrist. *Nonlinearity* 13 (2000), 441.
- [21] J. Gaset, A. Mas. *J. Geom. Mech.* 15 (2023), 357.
- [22] J. Paiva, M. Lazo, V. Zanchin. *Phys. Rev. D* 105 (2022), 124023.
- [23] M. de León, M. Lainz, M. Muñoz-Lecanda. *Journ. Nonlin. Sci.* 33 (2023), 9.
- [24] M. Lazo, J. Paiva, J. Amaral, G. Frederico. *Phys. Rev. D* 95 (2017), 101501.
- [25] D. Sloan. *Class. Quant. Grav.* 40 (2023), 115008.
- [26] P.-M. Zhang, M. Elbistan, P. Horvathy, P. Kosiński. *Eur. Phys. Journ. Plus* 135 (2020), 223.
- [27] B. Georgieva, R. Guenther. *Topological Methods in Nonlinear Analysis, Journ. of the Juliusz Schauder Center* 26 (2005), 307–314.
- [28] B. Georgieva, R. Guenther, T. Bodurov. *Journ. Math. Phys.* 44 (2003), 3911–3927.
- [29] M. Lazo, J. Paiva, G. Frederico. *Nonlinear Dynamics* 97 (2019), 1125–1136.