

Stable central limit theorems for discrete-time lag martingale difference arrays

Walter Dempsey*

Easton Huch†

December 5, 2025

Abstract

Recent work in dynamic causal inference introduced a class of discrete-time stochastic processes that generalize martingale difference sequences and arrays as follows: the random variates in each sequence have expectation zero given certain lagged filtrations but not given the natural filtration. We formalize this class of stochastic processes and prove a stable central limit theorem (CLT) via a Bernstein blocking scheme and an application of the classical martingale CLT. We generalize our limit theorem to vector-valued processes via the Cramér–Wold device and develop a simple form for the limiting variance. We demonstrate the application of these results to a problem in dynamic causal inference and present a simulation study supporting their validity.

Keywords: Causal inference; Cramér–Wold device; Lindeberg method; stochastic process.

1 Introduction

Recent work by Bojinov, Rambachan, and Shephard (hereafter, BRS) proposes randomization-based methods for estimating time-averaged effects of treatments on future outcomes (Bojinov et al., 2021; Bojinov and Shephard, 2019). As a simple example, consider a setting in which an experimenter sequentially randomizes a single individual to a binary treatment $A_t \in \{0, 1\}$ for $T \in \mathbb{N}$ time points. Following each treatment, the experimenter observes an outcome $Y_t \in \mathbb{R}$. The proposed methods enable the experimenter to estimate the time-averaged effect of A_t on Y_{t+p} for $p = 0, 1, 2, \dots$. With $p = 0$, the result is an estimate of the ‘contemporaneous’ or immediate effect of A_t on Y_t . For $p \geq 1$, the resulting estimate provides a measure of effect persistence over time, with significant effects at larger values of p indicating greater persistence. Assessing treatment persistence is a well-recognized goal in dynamic causal inference, and other authors have proposed similar definitions of persistence, usually in a superpopulation setting (Blackwell and Glynn, 2018; Boruvka et al., 2018; Robins et al., 1999).

BRS propose asymptotically Gaussian inference procedures justified by the martingale central limit theorem (CLT) (Hall and Heyde, 1980; Häusler and Luschgy, 2015); however, the martingale CLT does not apply for $p \geq 1$. We show in Section 7 that the theory fails for $p \geq 1$ because the supposed-martingale sequence, $(U_t)_{1 \leq t \leq T-p}$, generally does not satisfy the *martingale property*, $\mathbb{E}(U_t | \mathcal{H}_{t-1}) = 0$, where $(\mathcal{H}_t)_{-p \leq t \leq T-p}$ is a filtration with each U_t being \mathcal{H}_t -measurable. Rather, it satisfies a condition that we refer to as the *lag martingale property*: $\mathbb{E}(U_t | \mathcal{H}_{t-p-1}) = 0$. For $p \geq 1$, we show in (8) that the covariance between neighboring time points is nonnegligible; hence, the martingale CLT does not apply.

*Department of Biostatistics, University of Michigan, Ann Arbor, MI 48109, USA. wdem@umich.edu

†Johns Hopkins Carey Business School, 555 Pennsylvania Ave NW, Washington, DC 20001, USA. ehuch@jhu.edu

In spite of this theoretical issue, the proposed framework of BRS poses considerable merit because it produces unbiased estimates of dynamic causal effects under weak assumptions. In particular, the framework imposes only weak regularity conditions on the data-generating process for the outcomes. This stands in contrast to many competing methods in dynamic causal inference that impose restrictive conditions, such as ‘no carryover effects’ across time periods (Arkhangelsky et al., 2021; Athey and Imbens, 2022; de Chaisemartin and D’Haultfoeuille, 2020; Imai and Kim, 2019). In addition, the BRS framework does not require a superpopulation assumption or independent units, enabling its application to a finite set of experimental units. Consequently, addressing this theoretical issue would provide significant benefit to practitioners, enabling valid asymptotic inference for the case $p \geq 1$ under weak assumptions.

The primary contribution of this work is stable CLTs for lag martingale difference processes (defined in Section 2), such as those arising from the BRS framework. Similar to the classical martingale CLT, our results guarantee stable convergence to a scale mixture of Gaussian random variables. An intuitive characterization of stable convergence is (Jacod and Shiryaev, 2003, Proposition VIII.5.33)

$$X_n \rightarrow X \text{ } \mathcal{G}\text{-stably} \iff (X_n, Y) \xrightarrow{d} (X, Y) \text{ for all } \mathcal{G}\text{-measurable } Y,$$

where \xrightarrow{d} denotes (weak) convergence in distribution. Stability enables us to obtain a limiting Gaussian distribution (as opposed to a scale-mixture of Gaussian distributions) by applying the stable Cramér–Slutsky rule (Häusler and Luschgy, 2015, Theorem 3.18 (b)):

$$X_n \rightarrow X \text{ } \mathcal{G}\text{-stably}, \quad Y_n \xrightarrow{P} Y, \quad Y \text{ is } \mathcal{G}\text{-measurable} \implies X_n Y_n \xrightarrow{d} XY$$

where \xrightarrow{P} denotes convergence in probability. In the dynamic causal inference setting discussed above, $1/Y^2$ is the variance of the estimator and may be random.

From a technical standpoint, our results rely on a Bernstein blocking strategy, partitioning the rows of a triangular array into minor and major blocks (Bernstein, 1927; Dehling and Philipp, 2002). The minor blocks are constructed to be asymptotically negligible so that the limiting behavior is determined by partial sums arising from the major blocks, which obey the classical martingale CLT. Researchers have applied related blocking strategies to obtain CLTs for a wide variety of stochastic processes exhibiting forms of weak dependence across time, including mixingales, near-epoch dependent sequences, and mixing sequences (Bardet et al., 2008; de Jong, 1997; Jiang and Hahn, 2008; Merlevede and Peligrad, 2000; Merlevède, 2003).

Of the results in the literature, mixingale CLTs bear the greatest similarity to our CLTs because they apply to nonstationary data and guarantee stable convergence relative to a σ -field large enough to contain the asymptotic variance (Ikeda, 2017; McLeish, 1977; Wu, 2007). For a fixed lag length p , mixingale arrays are easily seen to nest lag martingale difference arrays as a special case; however, our specialized theory offers several benefits. First, our Theorem 3 allows the lag length to increase sublinearly with the sequence length, enabling application to non-mixingale arrays. Second, our results provide a convenient form for the asymptotic variance involving only variances and covariances within the lag window, facilitating asymptotic inference. Third, our approach permits a straightforward extension to multivariate processes. In fact, these benefits motivated this work and enable various improvements to the BRS framework; though, we leave the methodological details to future work.

The remainder of the paper is organized as follows. Section 2 provides the basic setup and definitions of scalar-valued lag- p martingale difference sequences and arrays. Section 3 gives a basic CLT for scalar-valued processes with a fixed, finite lag length. Section 4 extends the CLT to

multivariate processes. Section 5 provides the extension to diverging, sublinear lag lengths. Section 6 derives the simplified form of the asymptotic variance. Section 7 illustrates the application of our theoretical results to the BRS framework and includes a simulation study supporting their validity. Section 8 concludes with a brief discussion of the results.

2 Lag martingale difference sequences and arrays

Let $(k_n)_{n \in \mathbb{N}}$ be a nondecreasing sequence with $k_n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} k_n = \infty$. Let $(X_{nk})_{1 \leq k \leq k_n}$ be a sequence of real-valued random variables defined on the probability space (Ω, \mathcal{X}, P) for all $n \in \mathbb{N}$. Let $(\mathcal{X}_{nk})_{0 \leq k \leq k_n}$ be a filtration in \mathcal{X} so that $\mathcal{X}_{n0} \subset \mathcal{X}_{n1} \subset \dots \subset \mathcal{X}_{nk_n} \subset \mathcal{X}$. For all $n \in \mathbb{N}$, we assume that $(X_{nk})_{1 \leq k \leq k_n}$ is adapted to the filtration $(\mathcal{X}_{nk})_{0 \leq k \leq k_n}$, meaning that X_{nk} is \mathcal{X}_{nk} -measurable for all $1 \leq k \leq k_n$. In other words, the triangular array $(X_{nk})_{1 \leq k \leq k_n, n \in \mathbb{N}}$ of random variables is adapted to the triangular array $(\mathcal{X}_{nk})_{0 \leq k \leq k_n, n \in \mathbb{N}}$ of σ -fields. We say that $A \in \mathcal{L}^q(P)$ if $\mathbb{E}(|A|^q) < \infty$ in P -probability. With this notation in place, scalar-valued lag- p martingale difference sequences are defined as follows.

Definition 1 (Lag- p martingale difference sequence). Let $p \in \{0, 1, \dots\}$ and $n \in \mathbb{N}$ such that $k_n \geq p + 1$. The sequence $(X_{nk})_{p+1 \leq k \leq k_n}$ is a (scalar-valued) *lag- p martingale difference sequence* if $\mathbb{E}(X_{nk} | \mathcal{X}_{n, k-p-1}) = 0$ and $X_{nk} \in \mathcal{L}^1(P)$ for all $p + 1 \leq k \leq k_n$.

Lag- p martingale difference arrays are defined similarly, allowing n to range over \mathbb{N} . Definition 1 nests martingale difference sequences (and, by extension, arrays) as the special case $p = 0$. We can relate different lag lengths as follows.

Proposition 1 (Lag relationships). Let $p \in \{0, 1, \dots\}$, $q \in \{p + 1, p + 2, \dots\}$, and $n \in \mathbb{N}$ such that $k_n \geq q + 1$. If $(X_{nk})_{p+1 \leq k \leq k_n}$ is a lag- p martingale, then $(X_{nk})_{q+1 \leq k \leq k_n}$ is a lag- q martingale.

Proof. Apply the tower property of expectations. \square

In particular, Proposition 1 implies that a martingale difference sequence is a lag- p martingale difference sequence for any $p \in \mathbb{N}$; however, the converse is false in general.

3 Basic CLT

We now provide a stable CLT for scalar-valued lag- p martingale difference arrays.

3.1 Setup

Let $p \in \{0, 1, \dots\}$. We partition the elements of $(X_{nk})_{p+1 \leq k \leq k_n}$ into major and minor blocks. The starting indices (in k) of these major blocks are given by a sequence $(b_j)_{j \in \mathbb{N}}$ with $b_{j+1} - b_j \geq p + 1$ for all $j \in \mathbb{N}$; by construction, we set $b_1 = p + 1$. The minor blocks fall between the major blocks and are of length p . Their starting indices are given by the sequence $(a_j)_{j \in \mathbb{N}}$ with $a_j := b_{j+1} - p$ denoting the starting index of the minor block immediately following the j th major block. The major block sizes are then given by $d_j := a_j - b_j$. We define the minor and major block sums, respectively, as

$$A_{nj} := \sum_{k=a_j}^{\min(b_{j+1}-1, k_n)} X_{nk}, \quad B_{nj} := \sum_{k=b_j}^{\min(a_j-1, k_n)} X_{nk}.$$

We then define corresponding minor and major partial sums as $S_n^{(A)} := \sum_{j=1}^{j_n^{(A)}} A_{nj}$ and $S_n^{(B)} := \sum_{j=1}^{j_n^{(B)}} B_{nj}$, respectively, where $j_n^{(A)} := \max\{j \in \mathbb{N} : a_j \leq k_n\}$ is the number of minor blocks through index k_n and $j_n^{(B)} := \max\{j \in \mathbb{N} : a_j - 1 \leq k_n\}$ is the number of *complete* major blocks. We can then decompose the partial sum as $S_n := \sum_{k=p+1}^{k_n} X_{nk} = S_n^{(A)} + S_n^{(B)} + S_n^{(C)}$, where

$$S_n^{(C)} := \begin{cases} B_{n, j_n^{(B)}+1} & \text{if } k_n \in [b_{j_n^{(B)}+1}, a_{j_n^{(B)}+1} - 2], \\ 0 & \text{otherwise} \end{cases}$$

comprises the final block if it is an *incomplete* major block (i.e., if one or more terms are excluded because $k_n < a_{j_n^{(B)}+1} - 1$). We define the filtration array $(\mathcal{B}_{nj})_{0 \leq j \leq j_n^{(B)}, n \in \mathbb{N}}$ with each $\mathcal{B}_{nj} = \mathcal{X}_{n, a_j-1}$. Lastly, a triangular array is *square integrable* if $X_{nk} \in \mathcal{L}^2(P)$ for all $1 \leq k \leq k_n$ and $n \in \mathbb{N}$.

3.2 Theorem statement

Our basic CLT generalizes the classical martingale CLT (Häusler and Luschgy, 2015, Theorem 6.1).

Theorem 1 (CLT). *For some $p \in \{0, 1, \dots\}$, let $(X_{nk})_{p+1 \leq k \leq k_n, n \in \mathbb{N}}$ be a square-integrable lag- p martingale difference array adapted to $(\mathcal{X}_{nk})_{0 \leq k \leq k_n, n \in \mathbb{N}}$. Assume that there exists a blocking sequence $(b_j)_{j \in \mathbb{N}}$ as defined in Section 3.1 such that the following conditions hold:*

- (D) $d_j \rightarrow \infty$ as $j \rightarrow \infty$,
- (SP) $S_n^{(A)} + S_n^{(C)} \xrightarrow{P} 0$ as $n \rightarrow \infty$,
- (MP) $\max_{p+1 \leq k \leq k_n} |X_{nk}| \xrightarrow{P} 0$ as $n \rightarrow \infty$,
- (N) $\sum_{j=1}^{j_n^{(B)}} \mathbb{E}(B_{nj}^2 | \mathcal{B}_{n, j-1}) \xrightarrow{P} \psi$ as $n \rightarrow \infty$ for some \mathcal{G} -measurable real random variable $\psi \geq 0$, where $\mathcal{G}_{nj} := \cap_{m \geq n} \mathcal{B}_{mj}$ and $\mathcal{G} := \sigma(\cup_{n=1}^{\infty} \mathcal{G}_{n, j_n^{(B)}})$, and
- (CLB) $\sum_{j=1}^{j_n^{(B)}} \mathbb{E}(B_{nj}^2 \mathbf{1}_{\{|B_{nj}| \geq \epsilon\}} | \mathcal{B}_{n, j-1}) \xrightarrow{P} 0$ as $n \rightarrow \infty$ for every $\epsilon > 0$.

Then $S_n \rightarrow \sqrt{\psi}Z$ \mathcal{G} -stably as $n \rightarrow \infty$, where Z is a standard Gaussian random variable that is independent of \mathcal{G} .

In typical applications, we have a martingale difference sequence $(Y_k)_{k \in \mathbb{N}}$ and corresponding filtration $(\mathcal{Y}_k)_{k \in \{0\} \cup \mathbb{N}}$. In this case, we set $X_{nk} = n^{-1/2}Y_k$ and $\mathcal{X}_{nk} = \mathcal{Y}_k$. As a result, the filtration array $(\mathcal{X}_{nk})_{0 \leq k \leq k_n, n \in \mathbb{N}}$ is *nested* because $\mathcal{X}_{nk} = \mathcal{Y}_k = \mathcal{X}_{n+1, k}$ for all $n \in \mathbb{N}$ and $1 \leq k \leq k_n$. Consequently, we can set $\mathcal{G} = \mathcal{Y}_{\infty} := \sigma(\cup_{k=0}^{\infty} \mathcal{Y}_k)$, and Theorem 1 yields full stable convergence: the random variable ψ can depend on any information in \mathcal{Y}_{∞} , including all elements of $(Y_k)_{k \in \mathbb{N}}$.

3.3 Basic CLT proof

This section provides a proof of Theorem 1.

3.3.1 Convergence in distribution.

We first prove the following Lemma.

Lemma 1 (CLT). *Assume the conditions of Theorem 1. Then $S_n \xrightarrow{d} \sqrt{\psi}Z$ as $n \rightarrow \infty$, where Z is a standard Gaussian random variable.*

Proof. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a bounded function that is three times continuously differentiable with bounded derivatives, and define $C := \max\{\sup_{x \in \mathbb{R}} f(x), \sup_{x \in \mathbb{R}} f'(x)\}$. The conclusion of Lemma 1 will follow if we can show that $\mathbb{E}\{f(S_n) - f(\sqrt{\psi}Z)\} \rightarrow 0$ as $n \rightarrow \infty$. We decompose $\mathbb{E}\{f(S_n) - f(\sqrt{\psi}Z)\}$ as

$$\mathbb{E}\left\{f(S_n) - f\left(S_n^{(B)}\right)\right\} + \mathbb{E}\left\{f\left(S_n^{(B)}\right) - f(\sqrt{\psi}Z)\right\} = \mathbb{E}(I_n) + \mathbb{E}(II_n). \quad (1)$$

By Taylor's formula and the decomposition of S_n , we have

$$f(S_n) = f\left(S_n^{(B)}\right) + \left(S_n^{(A)} + S_n^{(C)}\right) f'(s) \quad \text{for some } s \text{ between } S_n^{(B)} \text{ and } S_n.$$

This formula gives $|I_n| = \left|S_n^{(A)} + S_n^{(C)}\right| \cdot |f'(s)| \leq 2C$ because C bounds f . But we also have $|f'(s)| \leq C$, implying that $|I_n| \xrightarrow{P} 0$ because $S_n^{(A)} + S_n^{(C)} \xrightarrow{P} 0$ by (SP); thus, dominated convergence gives $\mathbb{E}|I_n| \rightarrow 0$ as $n \rightarrow \infty$. As for $\mathbb{E}(II_n)$, the array $(B_{nj})_{1 \leq j \leq j_n^{(B)}, n \in \mathbb{N}}$ is easily shown to be a square-integrable martingale difference array adapted to $(\mathcal{B}_{nj})_{0 \leq j \leq j_n^{(B)}, n \in \mathbb{N}}$. Consequently, we can apply the martingale CLT (Häusler and Luschgy, 2015, Theorem 6.1) to show that $\mathbb{E}(II_n) \rightarrow 0$ as $n \rightarrow \infty$. \square

3.3.2 Proof of Theorem 1.

Using Lemma 1, we now adapt step 4 in the proof of Theorem 6.1 in (Häusler and Luschgy, 2015) to prove Theorem 1. Let $\mathcal{E} := \cup_{n=1}^{\infty} \mathcal{G}_{n, j_n^{(B)}}$, where \mathcal{G}_{nj} is defined in (N). The sequence $(\mathcal{G}_{nj})_{1 \leq j \leq j_n^{(B)}, n \in \mathbb{N}}$ is nondecreasing in both indices, so $\sigma(\mathcal{E}) = \mathcal{G}$, $\Omega \in \mathcal{E}$, and \mathcal{E} is closed under finite intersections. By (Häusler and Luschgy, 2015, Theorem 3.17, (i) \iff (iv)),

$$S_n \rightarrow \sqrt{\psi}Z \quad \mathcal{G}\text{-stably} \quad \iff \quad P_F^{S_n} \rightarrow P_F^{\sqrt{\psi}Z} \quad \text{weakly for every } F \in \mathcal{E}, P(F) > 0,$$

where P_F^Y denotes the conditional distribution of Y given F . We now let $F \in \mathcal{E}$ be arbitrary with $P(F) > 0$. By monotonicity of $(\mathcal{G}_{n, j_n^{(B)}})_{n \in \mathbb{N}}$, there exists $m \in \mathbb{N}$ such that $F \in \mathcal{G}_{m, j_m^{(B)}}$. Define $J := j_m^{(B)} + 1$ and $N := \min\{n \in \mathbb{N} : k_n \geq b_J\}$. Under P_F , the array $(X_{nk})_{b_J \leq k \leq k_n, n \geq N}$ is a lag- p martingale difference array adapted to $(\mathcal{X}_{nk})_{a_{J-1}-1 \leq k \leq k_n, n \geq N}$. We then verify the conditions of Theorem 1. First, (D) holds because we have removed only a finite number of terms. Second, (SP) yields

$$\sum_{j=J}^{j_n^{(B)}} A_{nj} + S_n^{(C)} \xrightarrow{P} 0 \quad \text{as } n \rightarrow \infty \text{ in } P\text{-probability}$$

because the omitted terms $(A_{nj})_{1 \leq j < J}$ are asymptotically negligible by (MP). Because $F \in \mathcal{E}$ is fixed with $P(F) > 0$, the above statement also holds in P_F -probability, so that (SP) continues to hold for the arrays $(X_{nk})_{b_J \leq k \leq k_n, n \geq N}$ and $(\mathcal{X}_{nk})_{a_{J-1}-1 \leq k \leq k_n, n \geq N}$ in P_F -probability. Third,

$\max_{p+1 \leq k \leq k_n} |X_{nk}| \xrightarrow{P} 0$ as $n \rightarrow \infty$ in P -probability by (MP), so this convergence also holds in P_F -probability. We then have $\max_{b_J \leq k \leq k_n} |X_{nk}| \xrightarrow{P} 0$ as $n \rightarrow \infty$ in P_F -probability because

$$\max_{b_J \leq k \leq k_n} |X_{nk}| \leq \max_{p+1 \leq k \leq k_n} |X_{nk}| \quad P_F\text{-almost surely,}$$

so (MP) continues to hold. Fourth, by conditions (MP) and (N), we have

$$\sum_{j=J}^{j_n^{(B)}} \mathbb{E}_P(B_{nj}^2 | \mathcal{B}_{n,j-1}) \xrightarrow{P} \psi \quad \text{in } P\text{-probability as } n \rightarrow \infty$$

because, again, the omitted terms are asymptotically negligible. We also have $\mathbb{E}_P(B_{nj}^2 | \mathcal{B}_{n,j-1}) = \mathbb{E}_{P_F}(B_{nj}^2 | \mathcal{B}_{n,j-1})$ P_F -almost surely for all $J \leq j \leq j_n^{(B)}$ and $n \geq N$, which gives

$$\sum_{j=J}^{j_n^{(B)}} \mathbb{E}_{P_F}(B_{nj}^2 | \mathcal{B}_{n,j-1}) \xrightarrow{P} \psi \quad \text{in } P_F\text{-probability as } n \rightarrow \infty,$$

so that (N) holds. Finally, (CLB) immediately implies that

$$\sum_{j=J}^{j_n^{(B)}} \mathbb{E}_{P_F}(B_{nj}^2 \mathbf{1}_{\{|B_{nj}| \geq \epsilon\}} | \mathcal{B}_{n,j-1}) \xrightarrow{P} 0 \quad \text{in } P_F\text{-probability as } n \rightarrow \infty$$

for all $\epsilon > 0$. Because these conditions are satisfied, we may apply Lemma 1 to obtain that $\sum_{k=b_J}^{k_n} X_{nk} \xrightarrow{d} \sqrt{\psi} Z$ as $n \rightarrow \infty$ under P_F . As for the omitted terms, we have $\sum_{k=p+1}^{b_J-1} X_{nk} \xrightarrow{P} 0$ by (MP); thus, we also have $S_n \xrightarrow{d} \sqrt{\psi} Z$ as $n \rightarrow \infty$ under P_F .

4 Multivariate extension

We now show that Theorem 1 permits a multivariate extension.

4.1 Multivariate definitions

In this section, we consider vector-valued arrays $(\mathbf{X}_{nk})_{1 \leq k \leq k_n, n \in \mathbb{N}}$ adapted to $(\mathcal{X}_{nk})_{0 \leq k \leq k_n, n \in \mathbb{N}}$. Each element \mathbf{X}_{nk} takes values in \mathbb{R}^q for some $q \geq 2$. Letting $\mathbf{0}_q$ denote the zero vector in \mathbb{R}^q , Definition 1 can then be generalized as follows.

Definition 2 (Multivariate lag- p martingale difference sequence). Let $p \in \{0, 1, \dots\}$ and $n \in \mathbb{N}$ such that $k_n \geq p+1$. The sequence $(\mathbf{X}_{nk})_{p+1 \leq k \leq k_n}$ is a *multivariate lag- p martingale difference sequence* if $\mathbb{E}(\mathbf{X}_{nk} | \mathcal{X}_{n,k-p-1}) = \mathbf{0}_q$ and $X_{nkj} \in \mathcal{L}^1(P)$ for all $p+1 \leq k \leq k_n$ and $1 \leq j \leq q$.

As above, the array version is similar, allowing n to range over \mathbb{N} .

4.2 Multivariate CLT

For $\mathbf{x}, \mathbf{y} \in \mathbb{R}^q$, let $\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^\top \mathbf{y}$ and $\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle}$. For a symmetric positive semidefinite matrix \mathbf{Q} , we let $\mathbf{Q}^{1/2}$ denote the (unique) symmetric, positive semidefinite square-root matrix such that $\mathbf{Q}^{1/2} \mathbf{Q}^{1/2} = \mathbf{Q}$. The array $(\mathbf{X}_{nk})_{1 \leq k \leq k_n, n \in \mathbb{N}}$ is square-integrable if $X_{nkj} \in \mathcal{L}^2(P)$ for all $n \in \mathbb{N}$, $1 \leq k \leq k_n$, and $1 \leq j \leq q$. We define \mathbf{S}_n , $\mathbf{S}_n^{(A)}$, $\mathbf{S}_n^{(B)}$, $\mathbf{S}_n^{(C)}$, \mathbf{A}_{nj} , and \mathbf{B}_{nj} as in Section 3.1, replacing X_{nk} with \mathbf{X}_{nk} . Theorem 1 can then be extended as follows to the multivariate setting.

Theorem 2 (Multivariate CLT). *For some $p \in \{0, 1, \dots\}$, let $(\mathbf{X}_{nk})_{p+1 \leq k \leq k_n, n \in \mathbb{N}}$ be a square-integrable lag- p martingale difference array adapted to $(\mathcal{X}_{nk})_{0 \leq k \leq k_n, n \in \mathbb{N}}$ with $\mathbf{X}_{nk} \in \mathbb{R}^q$ for some $q \geq 2$. Assume there exists a blocking sequence $(b_j)_{j \in \mathbb{N}}$ as defined in Section 3.1 such that (D) and the following conditions hold:*

$$(\text{SP}_q) \quad \|\mathbf{S}_n^{(A)} + \mathbf{S}_n^{(C)}\| \xrightarrow{\text{P}} 0 \text{ as } n \rightarrow \infty,$$

$$(\text{MP}_q) \quad \max_{1 \leq k \leq k_n} \|\mathbf{X}_{nk}\| \xrightarrow{\text{P}} 0 \text{ as } n \rightarrow \infty,$$

$$(\text{N}_q) \quad \sum_{j=1}^{j_n^{(B)}} \mathbb{E}(\mathbf{B}_{nj} \mathbf{B}_{nj}^\top | \mathcal{B}_{n,j-1}) \xrightarrow{\text{P}} \boldsymbol{\Psi} \text{ as } n \rightarrow \infty \text{ for some random } \mathcal{G}\text{-measurable symmetric, positive semidefinite matrix } \boldsymbol{\Psi} \in \mathbb{R}^{q \times q}, \text{ and}$$

$$(\text{CLB}_q) \quad \sum_{j=1}^{j_n^{(B)}} \mathbb{E}(\|\mathbf{B}_{nj}\|^2 \mathbf{1}_{\{\|\mathbf{B}_{nj}\| \geq \epsilon\}} | \mathcal{B}_{n,j-1}) \xrightarrow{\text{P}} 0 \text{ as } n \rightarrow \infty \text{ for every } \epsilon > 0.$$

Then $\mathbf{S}_n \rightarrow \boldsymbol{\Psi}^{1/2} \mathbf{Z}_q$ \mathcal{G} -stably as $n \rightarrow \infty$, where $\mathbf{Z}_q \in \mathbb{R}^q$ is a standard Gaussian random vector that is independent of \mathcal{G} .

The conditions in Theorem 2 are straightforward generalizations of those in Theorem 1. Conditions (N_q) and (CLB_q) appear in Häusler and Luschgy (2024).

4.3 Multivariate CLT proof

For $q = 1$, the conclusion follows from Theorem 1. Following the proof strategy in Häusler and Luschgy (2024), we employ a stable version of the Cramér–Wold device (Häusler and Luschgy, 2015, Corollary 3.19, (i) \iff (iii)). According to this device, it suffices to show that for any $q \in \mathbb{N}$ and an arbitrary nonrandom vector $\mathbf{u} \in \mathbb{R}^q \setminus \{\mathbf{0}_q\}$, that

$$\langle \mathbf{u}, \mathbf{S}_n \rangle \rightarrow \langle \mathbf{u}, \boldsymbol{\Psi}^{1/2} \mathbf{Z}_q \rangle \quad \mathcal{G}\text{-stably as } n \rightarrow \infty, \quad (2)$$

which can be shown via Theorem 1 because $\langle \mathbf{u}, \mathbf{S}_n \rangle$ is scalar-valued. Accordingly, we fix $\mathbf{u} \in \mathbb{R}^q \setminus \{\mathbf{0}_q\}$. For all $n \in \mathbb{N}$, we have $\langle \mathbf{u}, \mathbf{S}_n \rangle = \sum_{k=1}^{k_n} \langle \mathbf{u}, \mathbf{X}_{nk} \rangle = \sum_{k=1}^{k_n} Y_{nk}$, where $Y_{nk} := \langle \mathbf{u}, \mathbf{X}_{nk} \rangle$. The array $(Y_{nk})_{p+1 \leq k \leq k_n, n \in \mathbb{N}}$ is a scalar-valued lag- p martingale difference array adapted to the array $(\mathcal{X}_{nk})_{0 \leq k \leq k_n, n \in \mathbb{N}}$. We now verify the conditions of Theorem 1 making use of the inequality $|Y_{nk}| = |\langle \mathbf{u}, \mathbf{X}_{nk} \rangle| \leq \|\mathbf{u}\| \cdot \|\mathbf{X}_{nk}\|$. Below, we use redefine the scalar quantities $S_n^{(A)}$, $S_n^{(B)}$, $S_n^{(C)}$, and B_{nj} in terms of Y_{nk} . We have

$$|S_n^{(A)} + S_n^{(C)}| = |\langle \mathbf{u}, \mathbf{S}_n^{(A)} + \mathbf{S}_n^{(C)} \rangle| \leq \|\mathbf{u}\| \cdot \|\mathbf{S}_n^{(A)} + \mathbf{S}_n^{(C)}\| \xrightarrow{\text{P}} 0 \text{ as } n \rightarrow \infty$$

by (SP_q), verifying (SP) for the array $(Y_{nk})_{p+1 \leq k \leq k_n, n \in \mathbb{N}}$. Similar reasoning yields

$$\max_{p+1 \leq k \leq k_n} |Y_{nk}| \leq \|\mathbf{u}\| \max_{p+1 \leq k \leq k_n} \|\mathbf{X}_{nk}\| \rightarrow 0 \text{ as } n \rightarrow \infty,$$

verifying (MP). Condition (N_q) gives

$$\sum_{j=1}^{j_n^{(B)}} \mathbb{E}(B_{nj}^2 | \mathcal{B}_{n,j-1}) = \mathbf{u}^\top \left\{ \sum_{j=1}^{j_n^{(B)}} \mathbb{E}(\mathbf{B}_{nj} \mathbf{B}_{nj}^\top | \mathcal{B}_{n,j-1}) \right\} \mathbf{u} \xrightarrow{\text{P}} \mathbf{u}^\top \boldsymbol{\Psi} \mathbf{u} \text{ as } n \rightarrow \infty,$$

verifying (N). Lastly, we have

$$\sum_{j=1}^{j_n^{(B)}} \mathbb{E} \left(B_{nj}^2 \mathbf{1}_{\{\|B_{nj}\| \geq \epsilon\}} | \mathcal{B}_{n,j-1} \right) \leq \|\mathbf{u}\|^2 \sum_{j=1}^{j_n^{(B)}} \mathbb{E} \left(\|B_{nj}\|^2 \mathbf{1}_{\{\|B_{nj}\| \geq \sqrt{\epsilon}/\|\mathbf{u}\|\}} | \mathcal{B}_{n,j-1} \right) \xrightarrow{P} 0$$

as $n \rightarrow \infty$ for every $\epsilon > 0$ by (CLB_q), verifying (CLB). Applying Theorem 1, we obtain

$$\left\langle \mathbf{u}, \sum_{k=1}^{k_n} X_{nk} \right\rangle = \sum_{k=1}^{k_n} Y_{nk} \rightarrow (\mathbf{u}^\top \Psi \mathbf{u})^{1/2} Z \quad \mathcal{G}\text{-stably as } n \rightarrow \infty.$$

Because $(\mathbf{u}^\top \Psi \mathbf{u})^{1/2} Z \stackrel{d}{=} \langle \mathbf{u}, \Psi^{1/2} \mathbf{Z}_q \rangle$ and Ψ is \mathcal{G} -measurable, (2) holds as well. Because $\mathbf{u} \in \mathbb{R}^q \setminus \{\mathbf{0}\}_q$ and $q \in \mathbb{N}$ are arbitrary, the conclusion of Theorem 2 follows.

5 Diverging lags

We now consider a generalization of Definition 2 that allows for diverging lag lengths.

5.1 Diverging lag definition

We generalize Definition 2 as follows.

Definition 3 (Lag martingale difference sequence). Let $n, q, s \in \mathbb{N}$ with $k_n \geq s$, and let $(\mathbf{X}_{nk})_{s \leq k \leq k_n}$ be a sequence of random variables in \mathbb{R}^q . Then $(\mathbf{X}_{nk})_{s \leq k \leq k_n}$ is a *lag martingale difference sequence* if there exists a sequence $(p_k)_{s \leq k \leq k_n}$, $p_k \in \mathbb{N}$ such that $\mathbb{E}(\mathbf{X}_{nk} | \mathcal{X}_{n,k-p_k-1}) = \mathbf{0}_q$, $X_{nkj} \in \mathcal{L}^1(P)$, and $k - p_k - 1 \geq 0$ for all $s \leq k \leq k_n$ and $1 \leq j \leq q$.

As above, martingale difference arrays are defined similarly, allowing n to range over \mathbb{N} . We refer to the lag as *diverging* if $p_k \rightarrow \infty$ as $k \rightarrow \infty$. We say that $(\mathbf{X}_{nk})_{s \leq k \leq k_n, n \in \mathbb{N}}$ is a lag martingale difference sequence of *order* $\gamma \geq 0$ if there exists $C > 0$ such that $p_k \leq Ck^\gamma$ for all $k \in \mathbb{N}$. The lag is *sublinear* if $(\mathbf{X}_{nk})_{s \leq k \leq k_n, n \in \mathbb{N}}$ is of order $\gamma < 1$.

5.2 Blocking sequences

One challenge in generalizing Theorem 2 to diverging lags is developing a suitable blocking scheme. Proposition 2 demonstrates the existence of such a scheme for sublinear lags. In Proposition 2, condition (CN) ensures that the minor blocks comprise a vanishing proportion of the total number of terms, (DN) ensures that each major block is of negligible size compared to the total number of major-block terms, and (L) guarantees that the martingale property eventually holds for the major-block sums.

Proposition 2 (Diverging blocks). *Let $(\mathbf{X}_{nk})_{s \leq k \leq k_n, n \in \mathbb{N}}$ be a lag martingale difference array with sublinear lag. Then there exist sequences $(a_j)_{j \in \mathbb{N}}$ and $(b_j)_{j \in \mathbb{N}}$ defining the starting indices of minor and major blocks, respectively, such that (D) and the following conditions hold:*

$$(CN) \quad \sum_{j=1}^J c_j / \sum_{j=1}^J (c_j + d_j) \rightarrow 0 \text{ as } J \rightarrow \infty, \text{ where } c_j := b_{j+1} - a_j \text{ and } d_j := a_j - b_j,$$

$$(DN) \quad \max_{1 \leq j \leq J} (d_j) / \sum_{j=1}^J d_j \rightarrow 0 \text{ as } J \rightarrow \infty, \text{ and}$$

$$(L) \quad \text{There exists } J \in \mathbb{N} \text{ such that } \mathbb{E}(\mathbf{X}_{nk} | \mathcal{X}_{n,a_j-1}) = \mathbf{0}_q \text{ for all } b_j \leq k < a_j, j \geq J, \text{ and } n \in \mathbb{N}.$$

Proof. Let $\gamma \in [0, 1)$ be the order of $(\mathbf{X}_{nk})_{s \leq k \leq k_n, n \in \mathbb{N}}$. Let $A, B > 0$ be arbitrary and let $\alpha, \beta > 0$ such that $\gamma < \alpha/(1 + \beta)$ and $\alpha < \beta$. The following sequences are easily shown to satisfy the above conditions:

$$a_j := \left\lceil \sum_{l=1}^j (lB)^\beta + \sum_{l=1}^{j-1} (lA)^\alpha \right\rceil + s, \quad b_j := \left\lceil \sum_{l=1}^{j-1} \left\{ (lB)^\beta + (lA)^\alpha \right\} \right\rceil + s.$$

□

5.3 Diverging-lag CLT

We adopt the notation of Section 4.2 for $\mathbf{S}_n, \mathbf{S}_n^{(A)}, \mathbf{S}_n^{(B)}, \mathbf{S}_n^{(C)}, \mathbf{A}_{nj}, \mathbf{B}_{nj}$, and related quantities. We can then adapt Theorem 2 as follows.

Theorem 3 (Diverging-lag CLT). *Let $(\mathbf{X}_{nk})_{s \leq k \leq k_n, n \in \mathbb{N}}$ be a square-integrable lag martingale difference array of sublinear lag length adapted to $(\mathcal{X}_{nk})_{0 \leq k \leq k_n, n \in \mathbb{N}}$ with $\mathbf{X}_{nk} \in \mathbb{R}^q$ for some $q, s \in \mathbb{N}$. Assume there exist blocking sequences $(a_j)_{j \in \mathbb{N}}, (b_j)_{j \in \mathbb{N}}$ as defined in Section 5.2 such that (D), (L), and the conditions of Theorem 2 hold. Then $\mathbf{S}_n \rightarrow \Psi^{1/2} \mathbf{Z}_q$ \mathcal{G} -stably as $n \rightarrow \infty$, where $\mathbf{Z}_q \in \mathbb{R}^q$ is a standard Gaussian random vector that is independent of \mathcal{G} .*

The primary difference between Theorems 2 and 3 is that Theorem 3 replaces the highly structured blocking sequence defined in Section 3.1 with condition (L) from Proposition 2. We do not explicitly require conditions (CN) and (DN) from Proposition 2. For data-generating processes likely to arise in practice, however, we expect these conditions to serve as prerequisites for those of Theorem 2; otherwise, the minor blocks may not be asymptotically negligible, and some major block terms may dominate the summations in (\mathbf{N}_q) and (\mathbf{CLB}_q) .

5.4 Diverging-lag CLT Proof

To prove Theorem 3, we first provide a slight generalization of Lemma 1.

Lemma 2. *Assume the conditions of Theorem 3 with dimension $q = 1$. Further, assume that condition (L) of Proposition 2 holds with $J = 1$. Then $S_n \xrightarrow{d} \sqrt{\psi} Z$ as $n \rightarrow \infty$, where Z is a standard Gaussian random variable.*

Proof. Because condition (L) holds with $J = 1$, we have $\mathbb{E}(B_{nj} | \mathcal{B}_{n,j-1}) = 0$ for all $1 \leq j \leq j_n^{(B)}$ and $n \in \mathbb{N}$; thus $(B_{nj})_{1 \leq j \leq j_n^{(B)}, n \in \mathbb{N}}$ is a martingale difference array adapted to $(\mathcal{B}_{nj})_{1 \leq j \leq j_n^{(B)}, n \in \mathbb{N}}$, so we can apply the proof technique of Lemma 1. □

We now prove Theorem 3, beginning with the case $q = 1$. We proceed as in Section 3.3. Let $F \in \mathcal{E}$ be arbitrary with $P(F) > 0$. We can find $m \in N$ such that $F \in \mathcal{G}_{m, j_m^{(B)}}$. We then define $J_1 := j_m^{(B)} + 1$. Now let $J_2 \in \mathbb{N}$ be the constant J in Proposition 2 condition (L) for the given sequences $(a_j)_{j \in \mathbb{N}}$ and $(b_j)_{j \in \mathbb{N}}$. Set $J := \max(J_1, J_2)$ and $N := \min\{n \in N : k_n \geq b_J\}$. We then have that $(X_{nk})_{b_J \leq k \leq k_n, n \geq N}$ is a square-integrable lag martingale difference array with sublinear lag adapted to $(\mathcal{X}_{nk})_{a_{J-1}-1 \leq k \leq k_n, n \geq N}$. Using the same steps as in Section 3.3, the array $(X_{nk})_{b_J \leq k \leq k_n, n \geq N}$ satisfies conditions (SP), (MP), (N), and (CLB) in P_F -probability with respect to $(\mathcal{X}_{nk})_{a_{J-1}-1 \leq k \leq k_n, n \geq N}$. Further, $(b_j)_{j \in \mathbb{N}}$ satisfies (D) by assumption, and condition (L) is satisfied

for all major blocks as required by Lemma 2 because each block index j corresponding to the array $(X_{nk})_{b_J \leq k \leq k_n, n \geq N}$ is greater than or equal to J_2 ; thus, Lemma 2 gives

$$\sum_{k=b_J}^{k_n} X_{nk} \xrightarrow{d} \sqrt{\psi} Z \quad \text{as } n \rightarrow \infty \text{ under } P_F.$$

As in Section 3.3, condition (MP) guarantees that $\sum_{k=s}^{b_J-1} X_{nk} \xrightarrow{P} 0$, so we also have $S_n \xrightarrow{d} \sqrt{\psi} Z$ as $n \rightarrow \infty$ under P_F , proving that $S_n \rightarrow \sqrt{\psi} Z$ \mathcal{G} -stably as $n \rightarrow \infty$. For $q \geq 2$, apply the Cramér–Wold device as in the proof of Theorem 2 to deduce the multivariate statement from the univariate case.

6 Limiting variance

This section discusses the limiting variance appearing in Theorems 1, 2, and 3. For generality, we consider the setting of Theorem 3 with limiting variance Ψ . Condition (N_q) gives a summation that converges in probability to Ψ . Each term in this summation can be expressed as $\mathbb{E}(\Psi_{nj} | \mathcal{B}_{n,j-1})$, where

$$\begin{aligned} \Psi_{nj} := & \sum_{k=b_J}^{a_j-1} \left[\text{Var}(\mathbf{X}_{nk} | \mathcal{X}_{n,k-p_k-1}) + \sum_{l=1}^{\min(p_k, k-b_J)} \left\{ \text{Cov}(\mathbf{X}_{nk}, \mathbf{X}_{n,k-l} | \mathcal{X}_{n,k-p_k-1}) \right. \right. \\ & \left. \left. + \text{Cov}(\mathbf{X}_{n,k-l}, \mathbf{X}_{nk} | \mathcal{X}_{n,k-p_k-1}) \right\} \right]. \end{aligned} \quad (3)$$

In applications, it would be convenient to have an expression for the asymptotic variance that involves only expectations of a finite lag length. Accordingly, we define

$$\Psi_n := \sum_{j=1}^{j_n^{(B)}} \Psi_{nj}, \quad (4)$$

$$\begin{aligned} \bar{\Psi}_n := & \sum_{k=s}^{k_n} \left[\text{Var}(\mathbf{X}_{nk} | \mathcal{X}_{n,k-p_k-1}) + \sum_{l=1}^{\min(p_k, k-s)} \left\{ \text{Cov}(\mathbf{X}_{nk}, \mathbf{X}_{n,k-l} | \mathcal{X}_{n,k-p_k-1}) \right. \right. \\ & \left. \left. + \text{Cov}(\mathbf{X}_{n,k-l}, \mathbf{X}_{nk} | \mathcal{X}_{n,k-p_k-1}) \right\} \right], \end{aligned} \quad (5)$$

where the second expression, $\bar{\Psi}_n$, includes the minor-block terms omitted from Ψ_n . Theorem 4 provides conditions under which $\Psi_n \xrightarrow{P} \Psi$ as $n \rightarrow \infty$. Below, we let $\|\mathbf{Q}\|_F := \|\text{vec}(\mathbf{Q})\|$ denote the Frobenius norm of a matrix, \mathbf{Q} .

Theorem 4 (Asymptotic variance). *Assume that the following conditions hold for the array of matrices given as $(\Psi_{nj})_{1 \leq j \leq j_n^{(B)}, n \in \mathbb{N}}$:*

(CLB_F¹) $\sum_{j=1}^{j_n^{(B)}} \mathbb{E}(\|\Psi_{nj}\|_F \mathbf{1}_{\{\|\Psi_{nj}\|_F \geq \epsilon\}} | \mathcal{B}_{n,j-1}) \xrightarrow{P} 0$ as $n \rightarrow \infty$ for every $\epsilon > 0$ and

(SB_F) $\left(\sum_{j=1}^{j_n^{(B)}} \mathbb{E}(\|\Psi_{nj}\|_F^2 | \mathcal{B}_{n,j-1}) \right)_{n \in \mathbb{N}}$ is bounded in probability.

Then $\Psi_n \xrightarrow{P} \Psi$ as $n \rightarrow \infty$.

We provide a proof of Theorem 4 in Appendix A. Corollary 1 provides a condition under which we also have $\bar{\Psi}_n \xrightarrow{P} \Psi$ as $n \rightarrow \infty$.

Corollary 1 (Simple asymptotic variance). *Assume that the omitted terms $\bar{\Psi}_n - \Psi_n$ are asymptotically negligible in the sense of the following assumption:*

$$(VN) \quad \|\bar{\Psi}_n - \Psi_n\|_F \xrightarrow{P} 0 \text{ as } n \rightarrow \infty.$$

Then $\bar{\Psi}_n \xrightarrow{P} \Psi$ as $n \rightarrow \infty$.

Proof. Decompose the normed difference as

$$\|\bar{\Psi}_n - \Psi\|_F \leq \|\bar{\Psi}_n - \Psi_n\|_F + \|\Psi_n - \Psi\|_F = I_n + II_n.$$

As $n \rightarrow \infty$, we have $I_n \xrightarrow{P} 0$ by (VN) and $II_n \xrightarrow{P} 0$ by Theorem 4. \square

In applications, provided the variance and covariances within the minor blocks are similar in size to those of the major blocks, (VN) will typically hold because the minor blocks comprise a vanishing proportion of the total number of terms. We generally expect $\bar{\Psi}_n$ to outperform Ψ_n as an estimator of Ψ because Ψ_n omits the variance contribution of the minor blocks, but $\bar{\Psi}_n$ does not.

7 Application

We now apply our theoretical results to the BRS framework described in Section 1.

7.1 Setup

We consider a simplified setting with a single experimental unit representing, for example, a single individual. The individual is assigned a binary treatment $A_t \in \{0, 1\}$ at each time point $t \in \{1, 2, \dots, T\}$ for $T \in \mathbb{N}$. From $t = 2$ onward, we posit potential values (Rubin, 1974) for each of these treatment variables. At $t = 2$, we have $A_2(0)$ and $A_2(1)$, representing the treatment that would be assigned under $A_1 = 0$ or $A_1 = 1$, respectively. The observed treatment is $A_2 = A_2(A_1)$. For $t \geq 3$, these potential values depend on the full treatment path up to $t - 1$, denoted as $\mathbf{A}_{t-1} := (A_1, A_2, \dots, A_{t-1})$ or \mathbf{a}_{t-1} for a fixed (nonrandom) realization. In general, we have potential treatments $(A_t(\mathbf{a}_{t-1}))_{\mathbf{a}_{t-1} \in \{0,1\}^{t-1}}$ and an observed treatment $A_t = A_t(\mathbf{A}_{t-1})$.

Associated with each time point, we also have real-valued potential outcomes $(Y_t(\mathbf{a}_t))_{\mathbf{a}_t \in \{0,1\}^t}$ and an observed outcome $Y_t = Y_t(\mathbf{A}_t)$. We denote the outcomes up to time point t associated with \mathbf{a}_t as $\mathbf{Y}_t(\mathbf{a}_t) := (Y_1(a_1), Y_2(\mathbf{a}_2), \dots, Y_t(\mathbf{a}_t))$, and we denote the observed vector of outcomes up to time point t as $\mathbf{Y}_t := \mathbf{Y}_t(\mathbf{A}_t)$. We denote the full set of potential outcomes across all possible treatment paths as $\mathbf{Y} := (Y_t(\mathbf{a}_t))_{\mathbf{a}_t \in \{0,1\}^t, t \in \{1,2,\dots,T\}}$. As in Bojinov et al. (2021) and Bojinov and Shephard (2019), we assume that the treatments are sequentially randomized.

Assumption 1 (Sequential randomization). For all t , a_t , and \mathbf{a}_{t-1} , we have

$$\Pr(A_t = a_t | \mathbf{A}_{t-1} = \mathbf{a}_{t-1}, \mathbf{Y}) = \Pr\{A_t = a_t | \mathbf{A}_{t-1} = \mathbf{a}_{t-1}, \mathbf{Y}_{t-1}(\mathbf{a}_{t-1})\}.$$

Below, we use colon notation to denote the value of a variable between certain time points; for example, $\mathbf{A}_{4:6} := (A_4, A_5, A_6)$. With this notation, we denote the assignment probabilities as follows.

Definition 4 (Assignment probabilities). Let $p \in \{0, 1, \dots, T\}$ and $t \in \{1, 2, \dots, T - p\}$. For all $\mathbf{a}_{t:(t+p)}$, define

$$p_t(\mathbf{a}_{t:(t+p)}; \mathbf{A}_{t-1}, \mathbf{Y}_{t-1}(\mathbf{a}_{t-1})) := \Pr(\mathbf{A}_{t:(t+p)} = \mathbf{a}_{t:(t+p)} | \mathbf{A}_{t-1}, \mathbf{Y}_{t-1}(\mathbf{a}_{t-1})).$$

For brevity, we write $P_t(\mathbf{a}_{t:(t+p)}) := p_t(\mathbf{a}_{t:(t+p)}; \mathbf{A}_{t-1}, \mathbf{Y}_{t-1}(\mathbf{a}_{t-1}))$. We further assume that the assignment probabilities for the observed treatment path are known.

Assumption 2 (Known assignment probabilities). For all $p \in \{0, 1, \dots, T\}$ and $t \in \{1, 2, \dots, T - p\}$, the quantity $P_t(\mathbf{A}_{t:(t+p)})$ is known.

We define a filtration $(\mathcal{F}_t)_{0 \leq t \leq T}$, where $\mathcal{F}_t := \sigma(\mathbf{Y}, \mathbf{A}_t)$ is the σ -field generated by \mathbf{Y} and \mathbf{A}_t . Because \mathbf{Y} is used to generate every \mathcal{F}_t , the setup and corresponding statistical procedures effectively fix \mathbf{Y} at its realized value.

7.2 Causal estimand

Except where noted, the remainder of this section focuses on the special case of $p = 1$ and the following lag-1 causal estimand.

Definition 5 (Lag-1 causal effect). For $t \in \{2, 3, \dots, T\}$, the lag-1 causal effect is

$$\tau_t := \frac{Y_t(\mathbf{A}_{t-2}, 1, 1) - Y_t(\mathbf{A}_{t-2}, 0, 1)}{2} + \frac{Y_t(\mathbf{A}_{t-2}, 1, 0) - Y_t(\mathbf{A}_{t-2}, 0, 0)}{2}.$$

The lag-1 causal effect, τ_t , quantifies the effect of treatment A_{t-1} on the outcome at the next time point, Y_t , averaging over $A_t = 0$ and $A_t = 1$ with equal probability. Our statistical goal is to estimate an average of these lag-1 causal effects over time.

Definition 6 (Average lag-1 causal effect). The average lag-1 causal effect is

$$\tau := \frac{1}{T-1} \sum_{t=2}^T \tau_t.$$

Definitions 5 and 6 correspond with Definition 2 in [Bojinov and Shephard \(2019\)](#) with $p = 1$ and uniform weights. The estimand τ describes the degree to which treatment effects persist across two consecutive time periods, averaged over time. BRS show that τ_t and τ can be unbiasedly estimated as

$$\hat{\tau}_t := \frac{(-1)^{1-A_{t-1}} Y_t}{2P_{t-1}(\mathbf{A}_{(t-1):t})}, \quad \hat{\tau} := \frac{1}{T-1} \sum_{t=2}^T \hat{\tau}_t. \quad (6)$$

To study the properties of their general estimators for $p \geq 0$, BRS define a quantity $U_t := \hat{\tau}_{t+p} - \tau_{t+p}$ and show that $\mathbb{E}(U_t | \mathcal{F}_{t-1}) = 0$; however, this definition has the disadvantage that U_t is \mathcal{F}_{t+p} -measurable but not \mathcal{F}_t -measurable unless $p = 0$ (in Section 1, $\mathcal{H}_t = \mathcal{F}_{t+p}$). Hence, the classical martingale CLT applies only if $p = 0$ because $\mathbb{E}(U_t | \mathcal{F}_{t+p-1}) \neq 0$ in general. For clarity, we instead define $W_t := \hat{\tau}_t - \tau_t$, which is easily seen to be \mathcal{F}_t -measurable with $\mathbb{E}(W_t | \mathcal{F}_{t-p-1}) = 0$. BRS's assumption further imply that $W_t \in \mathcal{L}^1(P)$; thus, $(W_t)_{p+1 \leq t \leq T}$ is a lag- p martingale difference sequence adapted to $(\mathcal{F}_t)_{0 \leq t \leq T}$.

To apply Theorem 1 and Corollary 1 with $p = 1$, we define the arrays $(X_{nt})_{2 \leq t \leq n, n \in \mathbb{N}}$ and $(\mathcal{X}_{nt})_{0 \leq t \leq n, n \in \mathbb{N}}$ with $X_{nt} := n^{-1/2} W_t$ and $\mathcal{X}_{nt} := \mathcal{F}_t$, and we assume that \mathbf{Y} and \mathbf{A}_T are embedded in an asymptotic regime under which the assumptions of Theorem 1 and Corollary 1 are satisfied.

In particular, this will generally require that $\min\{P_t(A_t), 1 - P_t(A_t)\}$ is bounded away from zero (or approaches zero sufficiently slowly); otherwise, the conditional variances may increase without bound. Combining the \mathcal{G} -stable convergence of Theorem 1 with Corollary 1, we have

$$\frac{S_n}{\sqrt{\bar{\psi}_n}} \rightarrow Z \quad \mathcal{G}\text{-mixing as } n \rightarrow \infty, \quad (7)$$

where Z is standard Gaussian and $\bar{\psi}_n$ is a scalar version of $\bar{\Psi}_n$ defined in (5). This result enables asymptotic Gaussian inference for τ_n even though $\bar{\psi}_n$ converges in probability to a random variable, ψ .

In this application, $\bar{\psi}_n$ is not computable from the observed data because it depends on unobserved potential outcomes. In their work, BRS address this challenge by proposing an estimable variance bound. A similar strategy could be applied in our setting; though, we would need to bound the sum of variance and covariance terms defining $\bar{\psi}_n$. Because our primary goal is to validate the theoretical results of Sections 3–6, our simulation study will use the ‘oracle’ variance estimator, $\bar{\psi}$ (we omit the subscript n in the simulation results). Future methodological work in this area could address the additional statistical challenge of deriving an estimable bound for $\bar{\psi}$.

We proceed by deriving the terms needed to compute $\bar{\psi}$ in the simulation study. In the expressions that follow, $p_t(a_t; \mathbf{A}_{t-2}, a, \mathbf{Y}) := \Pr(A_t = a_t | \mathbf{A}_{t-2}, A_{t-1} = a, \mathbf{Y})$ as in Definition 4. The variance $\text{Var}(W_t | \mathcal{F}_{t-2})$ is

$$\begin{aligned} & \frac{P_{t-1}(0)P_{t-1}(1)}{4} \left\{ \frac{\sum_{a=0}^1 Y_t(\mathbf{A}_{t-2}, 1, a)}{P_{t-1}(1)} + \frac{\sum_{a=0}^1 Y_t(\mathbf{A}_{t-2}, 0, a)}{P_{t-1}(0)} \right\}^2 + \\ & \sum_{a=0}^1 \left\{ \frac{Y_t(\mathbf{A}_{t-2}, a, 1)}{p_t(1; \mathbf{A}_{t-2}, a, \mathbf{Y})} - \frac{Y_t(\mathbf{A}_{t-2}, a, 0)}{p_t(0; \mathbf{A}_{t-2}, a, \mathbf{Y})} \right\}^2 \frac{p_t(1; \mathbf{A}_{t-2}, a, \mathbf{Y}) p_t(0; \mathbf{A}_{t-2}, a, \mathbf{Y})}{4P_{t-1}(a)}. \end{aligned}$$

The required covariance term, $\text{Cov}(W_{t-1}, W_t | \mathcal{F}_{t-2})$, is

$$\begin{aligned} & \frac{(-1)^{1-A_{t-2}}}{4P_{t-2}(A_{t-2})} \sum_{a_{t-1}=0}^1 \frac{(-1)^{1-a_{t-1}} Y_{t-1}(\mathbf{A}_{t-2}, a_{t-1})}{P_{t-1}(a_{t-1})} \sum_{a_t=0}^1 Y_t(\mathbf{A}_{t-2}, a_{t-1}, a_t) \\ & - \tau_t \frac{(-1)^{1-A_{t-2}}}{2P_{t-2}(A_{t-2})} \sum_{a=0}^1 Y_{t-1}(\mathbf{A}_{t-2}, a). \end{aligned} \quad (8)$$

7.3 Simulation study

We now describe a simulation study we performed to evaluate our theoretical results.

7.3.1 Simulation setup.

We set $T=100,000$ and perform 10,000 independent replications. We implemented our simulation in Python, coding the main simulation loop in Cython (Behnel et al., 2010) to reduce computation time. The simulation completes in under one minute on a personal laptop with 48 GB of RAM and 14 CPUs. We initially set the assignment probabilities to 0.5. After observing $A_t = 1$ at 100 distinct time points, we alter the assignment probability as follows to ensure ψ is a nondegenerate random variable. If the 100th value of $A_t = 1$ occurs at an odd time point, we set

$$\Pr(A_t = 1 | \mathbf{A}_{t-1}, \mathbf{Y}) = \begin{cases} 0.1 & \text{if } A_{t-1} = 1, \\ 0.5 & \text{otherwise.} \end{cases}$$




Figure 1: Estimates from the simulation study. Panel (a) plots kernel density estimates (KDEs) of the sampled values of τ and the corresponding unbiased estimator $\hat{\tau}$. Panel (b) compares a KDE of the sampled values of $W := \hat{\tau} - \tau$ to a Gaussian distribution with $\sigma_W := \text{SD}(W)$, offering visual evidence of the non-Gaussianity of W . Panel (c) shows the estimates of $\bar{\psi}$ segmented by whether the 100th value of $A_t = 1$ is even or odd, highlighting how the limiting variance is a random variable.

On the other hand, if the 100th value of $A_t = 1$ occurs at an even time point, we set $\Pr(A_t = 1 | \mathbf{A}_{t-1}, \mathbf{Y}) = 0.5$ uniformly. We draw the outcomes as follows:

$$Y_t(\mathbf{A}_{t-1}, a) | \mathbf{A}_{t-1} \stackrel{iid}{\sim} \text{Gamma}\{\theta_t(A_{t-1}), 1\}, \quad \theta_t(A_{t-1}) = \begin{cases} 2 & \text{if } A_{t-1} = 0, \\ 3 & \text{if } A_{t-1} = 1 \end{cases}$$

for $a = 0, 1$. This sampling scheme results in a causal effect τ that differs slightly between replications with $\mathbb{E}(\tau) = 1$. Within each replication, $\hat{\tau}$ is conditionally unbiased for the sampled value τ .

7.3.2 Simulation results.

Figure 1 displays the estimates from the simulation study. Panel 1a plots kernel density estimates (KDEs) of τ and $\hat{\tau}$ across replications. Panel 1b plots a KDE for $W := \hat{\tau} - \tau$ compared to a Gaussian distribution with standard deviation $\sigma_W := \text{SD}(W)$, providing visual evidence that W is not (marginally) Gaussian. We also applied the Kolmogorov–Smirnov (KS) test (Kolmogorov, 1933; Smirnov, 1948) against a Gaussian distribution with standard deviation equal to σ_W , resulting in a p-value of 1.5×10^{-35} .

Theorem 1 implies that the asymptotic distribution of W is a scale mixture of Gaussian random variables with ψ playing the role of random scale parameter. Panel 1c plots the simulated values of $\bar{\psi}$ —estimates of ψ in the sense of Corollary 1—segmented by whether the 100th value of $A_t = 1$ occurs at an even or odd time point. These estimates concentrate around 32.5 and 245.4 for the even and odd groups, respectively, with each group comprising roughly 50% of the simulation replicates. Consequently, Theorem 1 implies that the asymptotic distribution of $\sqrt{T}W$ is an approximately equal mixture of Gaussian random variables with mean zero and variances corresponding to those of these two groups (estimated at 32.5 and 245.4). To further evaluate this theoretical implication, we fit a two-component Gaussian mixture model to the sampled values of $\sqrt{T}W$. The fitted model estimates one component with a mixture weight of 52%, mean of -0.2 , and variance of 241.4; the other component is estimated at 48%, -0.1 , and 30.8 for these quantities, respectively. These estimates closely match the theoretical values we expect based on Theorem 1 and Corollary 1, providing additional empirical support for their validity.

To further illustrate the implications of stability in Theorems 1 and 2, we define the quantity $Z := (T/\bar{\psi})^{1/2}W$: a realized value from the sequence $(\bar{\psi}_n^{-1/2}S_n)_{n \in \mathbb{N}}$. By (7), the asymptotic




Figure 2: Normalized estimates of W from the simulation study. Panel (a) plots a KDE of $Z := (T/\bar{\psi})^{1/2} W$ vs. a standard Gaussian distribution, showing close agreement as expected based on (7). Panel (b) performs a similar comparison but omitting the covariance terms in the definition of $\bar{\psi}$; we label the resulting quantity Z -naive. The comparison of the KDE for Z -naive to the standard Gaussian distribution shows significant disagreement, highlighting the need for the covariance terms to obtain valid asymptotic inference.

distribution of Z is standard Gaussian. Figure 2 plots the empirical distribution of Z and a related quantity, Z -naive, that we formed using a variant of $\bar{\psi}$ that omits the covariance terms defined in (8). Panel 2a shows that the empirical distribution of Z closely matches that of a standard Gaussian as expected. We applied the KS test and two-sided t-tests for $\mathbb{E}(Z) = 0$ and $\mathbb{E}(Z^2) = 1$, resulting in p-values of 0.34, 0.61, and 0.83, respectively, offering little to no evidence against the null hypothesis that Z is standard Gaussian. Panel 2b includes a similar plot for Z -naive, showing noticeable disagreement between the empirical distribution and the standard Gaussian distribution. We applied the hypothesis tests above, resulting in p-values of 0.29, 0.53, and 4.2×10^{-37} , respectively, providing conclusive evidence that the variance is not equal to one. This comparison highlights the need for the covariance terms in $\bar{\psi}$. For general $p \geq 1$, their omission results in miscalibrated asymptotic inference due to the possibility of short-term temporal correlation in lag martingales. In contrast, for $p = 0$, these covariance terms are zero, so they need not be included and inference requires only the classical martingale CLT.

8 Conclusion

This paper introduces lag martingales and proves several CLTs for their partial sums. Our CLTs guarantee stable convergence to a scale mixture of Gaussian random variables. Our most general CLT (Theorem 3) applies to vector-valued processes with sublinear lag lengths. In addition to the CLTs, we also derive a convenient form for the asymptotic variance in Corollary 1.

This work was motivated by the dynamic causal inference method of BRS (Bojinov et al., 2021; Bojinov and Shephard, 2019). Our theoretical results enable application of their framework with lag length $p \geq 1$. The stability of our CLTs explicitly allows for a random limiting variance—a possibility that is difficult to rule out in practice. We illustrate the application of our theoretical results to the BRS framework in Section 7 and provide a simulation study to empirically verify their implications. Future methodological work in this area could leverage our theoretical results to

generalize the BRS framework in various ways, including to estimate effect moderation and allow for dependence across units in the treatment assignment process.

The literature on martingale CLTs includes a wide range of sufficient conditions (Häusler and Luschgy, 2015, 2024). In the present work, we have employed various versions of the nesting (N) and conditional Lindeberg (CLB) assumptions because other authors have argued that these assumptions produce the “most common and useful” CLT (Häusler and Luschgy, 2024, p. 4) and, indeed, in our application setting, these assumptions lead to a convenient expression for the asymptotic variance. Future work could develop additional sufficient conditions for our CLTs and, perhaps, demonstrate their application to other stochastic processes satisfying the lag martingale property, such as moving-average processes.

Acknowledgements

We wish to thank Gabriel Durham and Inbal Billie Nahum–Shani for helpful feedback and discussions on early versions of this manuscript.

Funding

This work was supported by NIH grants P50DA054039 and R01GM152549 provided by the NIDDK and NIGMS, respectively. In addition, Dr. Huch gratefully acknowledges research funding from the Johns Hopkins Carey Business School, which supported him during the late stages of this project.

Research Data

The code and results for the simulation found in Section 7.3 can be found at <https://github.com/eastonhuch/lag-martingales>.

References

- Arkhangelsky, D., of Economic Research., N. B., and Imbens, G. W. (2021). *Double-Robust Identification for Causal Panel Data Models / Dmitry Arkhangelsky, Guido W. Imbens*. NBER working paper series no. w28364. National Bureau of Economic Research, Cambridge, Mass.
- Athey, S. and Imbens, G. W. (2022). Design-based analysis in difference-in-differences settings with staggered adoption. *Journal of Econometrics*, 226(1):62–79. Annals Issue in Honor of Gary Chamberlain.
- Bardet, J.-M., Doukhan, P., Lang, G., and Ragache, N. (2008). Dependent Lindeberg central limit theorem and some applications. *Probability and Statistics*, 12:154–172.
- Behnel, S., Bradshaw, R., Citro, C., Dalcin, L., Seljebotn, D. S., and Smith, K. (2010). Cython: The best of both worlds. *Computing in Science & Engineering*, 13(2):31–39.
- Bernstein, S. (1927). Sur l’extension du théorème limite du calcul des probabilités aux sommes de quantités dépendantes. *Mathematische Annalen*, 97(1):1–59.
- Blackwell, M. and Glynn, A. N. (2018). How to make causal inferences with time-series cross-sectional data under selection on observables. *The American Political Science Review*, 112(4):1067–1082.

- Bojinov, I., Rambachan, A., and Shephard, N. (2021). Panel experiments and dynamic causal effects: A finite population perspective. *Quantitative Economics*, 12(4):1171–1196.
- Bojinov, I. and Shephard, N. (2019). Time series experiments and causal estimands: Exact randomization tests and trading. *Journal of the American Statistical Association*, 114(528):1665–1682.
- Boruvka, A., Almirall, D., Witkiewitz, K., and Murphy, S. A. (2018). Assessing time-varying causal effect moderation in mobile health. *Journal of the American Statistical Association*, 113(523):1112–1121.
- de Chaisemartin, C. and D’Haultfoeuille, X. (2020). Two-way fixed effects estimators with heterogeneous treatment effects. *The American Economic Review*, 110(9):2964–2996.
- de Jong, R. M. (1997). Central limit theorems for dependent heterogeneous random variables. *Econometric theory*, 13(3):353–367.
- Dehling, H. and Philipp, W. (2002). Empirical process techniques for dependent data. In Dehling, H., Mikosch, T., and Sørensen, M., editors, *Empirical Process Techniques for Dependent Data*, pages 3–113. Birkhäuser Boston, Boston, MA, 1st edition.
- Hall, P. and Heyde, C. C. (1980). *Martingale Limit Theory and its Application*. Elsevier, New York.
- Häusler, E. and Luschgy, H. (2015). *Stable Convergence and Stable Limit Theorems*, volume 74 of *Probability Theory and Stochastic Modelling*. Springer International Publishing, Cham.
- Häusler, E. and Luschgy, H. (2024). On stable central limit theorems for multivariate discrete-time martingales. arXiv:2407.18633.
- Ikeda, S. S. (2017). A note on mixingale limit theorems and stable convergence in law. *Communications in Statistics – Theory and Methods*, 46(19):9377–9387.
- Imai, K. and Kim, I. S. (2019). When should we use unit fixed effects regression models for causal inference with longitudinal data? *American Journal of Political Science*, 63(2):467–490.
- Jacod, J. and Shiryaev, A. (2003). *Limit Theorems for Stochastic Processes*. Grundlehren der mathematischen Wissenschaften, A Series of Comprehensive Studies in Mathematics, 288. Springer Berlin Heidelberg, Berlin, Heidelberg, 2nd edition.
- Jiang, X. and Hahn, M. (2008). A self-normalized central limit theorem for ρ -mixing stationary sequences. *Statistics & Probability Letters*, 78(12):1541–1547.
- Kolmogorov, A. N. (1933). Sulla determinazione empirica di una legge di distribuzione. *Giornale dell’Istituto Italiano degli Attuari*, 4:83–91.
- McLeish, D. L. (1977). On the invariance principle for nonstationary mixingales. *The Annals of Probability*, 5(4):616–621.
- Merlevede, F. and Peligrad, M. (2000). The functional central limit theorem under the strong mixing condition. *The Annals of Probability*, 28(3):1336–1352.
- Merlevède, F. (2003). On the central limit theorem and its weak invariance principle for strongly mixing sequences with values in a Hilbert space via martingale approximation. *Journal of Theoretical Probability*, 16(3):625–653.

- Robins, J. M., Greenland, S., and Hu, F.-C. (1999). Estimation of the causal effect of a time-varying exposure on the marginal mean of a repeated binary outcome. *Journal of the American Statistical Association*, 94(447):687–700.
- Rubin, D. B. (1974). Estimating causal effects of treatments in randomized and nonrandomized studies. *Journal of Educational Psychology*, 66(5):688–701.
- Smirnov, N. (1948). Table for estimating the goodness of fit of empirical distributions. *Annals of Mathematical Statistics*, 19(2):279–281.
- Wu, W. B. (2007). Strong invariance principles for dependent random variables. *The Annals of Probability*, 35(6):2294–2320.

A Proof of Theorem 4

To prove Theorem 4, we first present a generalization of Proposition 6.16 in Häusler and Luschgy (2015).

Lemma 3. *Let $(X_{nk})_{1 \leq k \leq k_n, n \in \mathbb{N}}$ be an array of random variables with $X_{nk} \in \mathcal{L}^1(P)$ for all $1 \leq k \leq k_n$ and $n \in \mathbb{N}$, and let $(\mathcal{X}_{nk})_{0 \leq k \leq k_n, n \in \mathbb{N}}$ be an array of σ -fields. Assume*

$$(\text{CLB}^1) \quad \sum_{k=1}^{k_n} \mathbb{E}(|X_{nk}| \mathbf{1}_{\{|X_{nk}| \geq \epsilon\}} | \mathcal{X}_{n,k-1}) \xrightarrow{P} 0 \text{ as } n \rightarrow \infty \text{ for every } \epsilon > 0 \text{ and}$$

$$(\text{SB}) \quad \left(\sum_{k=1}^{k_n} \mathbb{E}(|X_{nk}| | \mathcal{X}_{n,k-1}) \right)_{n \in \mathbb{N}} \text{ is bounded in probability.}$$

Then

$$\sum_{k=1}^{k_n} \mathbb{E}(X_{nk} | \mathcal{X}_{n,k-1}) - \sum_{k=1}^{k_n} X_{nk} \xrightarrow{P} 0 \text{ as } n \rightarrow \infty.$$

Proof. The proof follows from a slight generalization of the proofs of Lemma 6.14 and Proposition 6.16 in Häusler and Luschgy (2015). \square

We extend this lemma to vector-valued processes via the Cramér–Wold device.

Lemma 4. *Let $(\mathbf{X}_{nk})_{1 \leq k \leq k_n, n \in \mathbb{N}}$ be an array of random q -vectors with $X_{nkj} \in \mathcal{L}^1(P)$ for all $n \in \mathbb{N}$, $1 \leq k \leq k_n$, and $j \in \{1, 2, \dots, q\}$. Let $(\mathcal{X}_{nk})_{0 \leq k \leq k_n, n \in \mathbb{N}}$ be an array of σ -fields. Assume*

$$(\text{CLB}_q^1) \quad \sum_{k=1}^{k_n} \mathbb{E}(\|\mathbf{X}_{nk}\| \mathbf{1}_{\{\|\mathbf{X}_{nk}\| \geq \epsilon\}} | \mathcal{X}_{n,k-1}) \xrightarrow{P} 0 \text{ as } n \rightarrow \infty \text{ for every } \epsilon > 0 \text{ and}$$

$$(\text{SB}_q) \quad \left(\sum_{k=1}^{k_n} \mathbb{E}(\|\mathbf{X}_{nk}\| | \mathcal{X}_{n,k-1}) \right)_{n \in \mathbb{N}} \text{ is bounded in probability.}$$

Then

$$\sum_{k=1}^{k_n} \mathbb{E}(\mathbf{X}_{nk} | \mathcal{X}_{n,k-1}) - \sum_{k=1}^{k_n} \mathbf{X}_{nk} \xrightarrow{P} \mathbf{0}_q \text{ as } n \rightarrow \infty.$$

Proof. The case $q = 1$ follows directly from Lemma 3. For $q > 1$, we apply the Cramér–Wold device as in the proof of Theorem 2. Accordingly, let $q \in \mathbb{N}$ and $\mathbf{u} \in \mathbb{R}^q \setminus \{\mathbf{0}_q\}$ be arbitrary. Our goal is to show that

$$\left\langle \mathbf{u}, \sum_{k=1}^{k_n} \mathbb{E}(\mathbf{X}_{nk} | \mathcal{X}_{n,k-1}) - \sum_{k=1}^{k_n} \mathbf{X}_{nk} \right\rangle \xrightarrow{d} \langle \mathbf{u}, \mathbf{0}_q \rangle \text{ as } n \rightarrow \infty, \quad (9)$$

which will then imply that

$$\sum_{k=1}^{k_n} \mathbb{E}(\mathbf{X}_{nk} | \mathcal{X}_{n,k-1}) - \sum_{k=1}^{k_n} \mathbf{X}_{nk} \xrightarrow{d} \mathbf{0}_q \quad \text{as } n \rightarrow \infty,$$

implying the conclusion of the lemma because the limits are constant. We define $Y_{nk} := \langle \mathbf{u}, \mathbf{X}_{nk} \rangle$ and show that the conditions of Lemma 3 are satisfied for $(Y_{nk})_{1 \leq k \leq k_n, n \in \mathbb{N}}$. First, we have $Y_{nk} \in \mathcal{L}^1(P)$ for all $n \in \mathbb{N}$ and $1 \leq k \leq k_n$ because $Y_{nkj} \in \mathcal{L}^1(P)$, $j \in \{1, \dots, q\}$. Second, we have

$$\sum_{k=1}^{k_n} \mathbb{E}(|Y_{nk}| \mathbf{1}_{\{|Y_{nk}| \geq \epsilon\}} | \mathcal{X}_{n,k-1}) \leq \|\mathbf{u}\| \sum_{k=1}^{k_n} \mathbb{E}(\|\mathbf{X}_{nk}\| \mathbf{1}_{\{\|\mathbf{X}_{nk}\| \geq \epsilon/\|\mathbf{u}\|\}} | \mathcal{X}_{n,k-1}) \xrightarrow{P} 0$$

as $n \rightarrow \infty$ for all $\epsilon > 0$ by (CLB_q¹) so that (CLB¹) holds. Third, we have

$$\sum_{k=1}^{k_n} \mathbb{E}(|Y_{nk}| | \mathcal{X}_{n,k-1}) \leq \|\mathbf{u}\| \sum_{k=1}^{k_n} \mathbb{E}(\|\mathbf{X}_{nk}\| | \mathcal{X}_{n,k-1}),$$

which is bounded in probability by (SB_q), implying (SB); thus, Lemma 3 gives

$$\sum_{k=1}^{k_n} \mathbb{E}(Y_{nk} | \mathcal{X}_{n,k-1}) - \sum_{k=1}^{k_n} Y_{nk} \xrightarrow{d} 0.$$

We also have

$$\left\langle \mathbf{u}, \sum_{k=1}^{k_n} \mathbb{E}(\mathbf{X}_{nk} | \mathcal{X}_{n,k-1}) - \sum_{k=1}^{k_n} \mathbf{X}_{nk} \right\rangle = \sum_{k=1}^{k_n} \mathbb{E}(Y_{nk} | \mathcal{X}_{n,k-1}) - \sum_{k=1}^{k_n} Y_{nk}$$

and $\langle \mathbf{u}, \mathbf{0}_q \rangle = 0$, so (9) also holds, implying the conclusion of the lemma. \square

We now prove Theorem 4. Note that $\Psi_n := \sum_{j=1}^{j_n^{(B)}} \Psi_{nj}$ and $\sum_{j=1}^{j_n^{(B)}} \mathbb{E}(\Psi_{nj} | \mathcal{B}_{n,j-1}) \xrightarrow{P} \Psi$ as $n \rightarrow \infty$. Then apply Lemma 4 to the array $(\text{vec}(\Psi_{nj}))_{1 \leq j \leq j_n^{(B)}, n \in \mathbb{N}}$.