

RESOLUTIONS OF TYPE A QUANTUM SURFACE SINGULARITIES

SIMON CRAWFORD AND SUSAN J. SIERRA

ABSTRACT. Let $B = \mathbb{k}_q[u, v]^{C_{n+1}}$ be a Type A_n quantum Kleinian singularity, which is an example of a noncommutative surface singularity. This singularity is known to have a noncommutative quasi-crepant resolution Λ , which is an “algebraic” resolution of B . We construct a category \mathcal{X} which serves as a “geometric” resolution of B by adapting techniques from quiver GIT and show that \mathcal{X} and $\text{mod-}\Lambda$ are derived equivalent. Furthermore, we show that the intersection arrangement of lines in the exceptional locus of \mathcal{X} corresponds to a Type A_n Dynkin diagram. This generalises the geometric McKay correspondence for classical Kleinian singularities.

CONTENTS

1. Introduction	1
2. Preliminaries	4
3. A noncommutative singularity	8
4. The representation ring	13
5. A geometric resolution: the ring	18
6. A geometric resolution: the category	25
7. A system for localising T -modules	34
8. A derived equivalence of resolutions	41
9. Intersection theory	49
References	54

1. INTRODUCTION

Throughout, let \mathbb{k} be an algebraically closed field of characteristic 0. Let $q \in \mathbb{k}^\times$, and let

$$A_q = \mathbb{k}_q[u, v] := \frac{\mathbb{k}\langle u, v \rangle}{\langle vu - quv \rangle}$$

be the q -quantum plane. The cyclic group C_{n+1} of order $n+1$ generated by

$$g := \begin{pmatrix} \zeta & 0 \\ 0 & \zeta^{-1} \end{pmatrix},$$

where $\zeta \in \mathbb{k}$ is a primitive $(n+1)$ th root of unity, acts on A_q via graded automorphisms. Let $B_q = A_q^{C_{n+1}}$ be the invariant ring of this action. This is a well-known *quantum Kleinian singularity*, as studied in [Cra18, CKWZ18, CKWZ19] and others. In particular, B_q is singular, i.e. it has infinite global dimension, and this singularity can be resolved, in an appropriate sense, by the smash product $\Lambda_q := A_q \# G$. Further, there is a noncommutative McKay correspondence linking the representation theory of G , of B_q , and of Λ_q ; see [CKWZ18, CKWZ19].

In this paper we extend the McKay correspondence for B_q by establishing a *(derived) geometric McKay correspondence*. That is, we construct an \mathbb{N} -graded ring T_q using methods inspired by geometric invariant theory, and a further construct a category \mathcal{X}_q from T_q by generalising

Date: December 8, 2025.

2020 Mathematics Subject Classification. Primary: 14A22, 14E15. Secondary: 16S38, 14E16.

Key words and phrases. Quantum Kleinian singularity, invariant ring, resolution of singularities, McKay correspondence.

standard techniques of noncommutative geometry. We show that \mathcal{X}_q is also a resolution of (the module category of) B_q , again in an appropriate sense and, further, that \mathcal{X}_q and $\text{mod-}\Lambda_q$ are derived equivalent. This establishes an analogue of the derived geometric McKay correspondence, as given in [BKR01], for the quantum Kleinian singularity B_q . Thus we have quantised the full classical McKay correspondence for type \mathbb{A} Kleinian surface singularities.

In order to describe our results in more detail, we first review the classical (commutative) McKay correspondence for Kleinian surface singularities. The finite subgroups of $\text{SL}_2(\mathbb{k})$ were famously classified by Klein in the case $\mathbb{k} = \mathbb{C}$. Any such subgroup G acts on \mathbb{A}^2 by construction, and the quotient singularity

$$X = \mathbb{A}^2/G := \text{Spec } \mathbb{k}[u, v]^G$$

is known as a *Kleinian (surface) singularity*. These were studied by Du Val [DV34], Artin [Art66], and many others from a geometric perspective, but in 1981 McKay [McK80] established a beautiful relationship between the geometry of X and its minimal resolution and the representation theory of G , later extended and deepened by many others. We shall now give an ahistorical survey of the McKay correspondence for Kleinian singularities.

A finite subgroup G of $\text{SL}_2(\mathbb{k})$ acts naturally on $V = \mathbb{k}^2$ and thus on $A := S(V) = \mathbb{k}[u, v]$. Set $B = A^G$, and let $X = \mathbb{A}^2/G = \text{Spec } B$ be the associated Kleinian singularity. Let W_0, \dots, W_n be the irreducible representations of G , up to isomorphism, where $W_0 = \mathbb{k}$ is the trivial representation. As \mathbb{k} has characteristic 0, there are non-negative integers m_{ij} so that

$$V \otimes W_j \cong \bigoplus_i W_i^{m_{ij}}.$$

The *McKay quiver* Q associated to the action of G on V has vertex set $Q_0 = \{0, \dots, n\}$ and m_{ij} arrows from vertex i to vertex j . (Since $G \leq \text{SL}_2(\mathbb{k})$, we in fact have $m_{ij} = m_{ji}$ [LW12, Lemma 6.25].) Auslander had observed [Aus62] that $A \# G \cong \text{End}_B(A)$, and Reiten and Van den Bergh later showed (see in particular the proof of [RVdB89, Lemma 2.13]) that $A \# G$ is Morita equivalent to the *preprojective algebra* $\Pi(Q)$ of the McKay quiver, which is the path algebra of Q factored by certain natural relations; see [CBH98] for the definition. In fact, if G is cyclic, then $A \# G \cong \Pi(Q)$. There are many relationships between the representation theory of G , of $\Lambda := \Pi(Q)$, and of $A \# G$; see [LW12, Chapter 5] for a discussion.

In 1981, McKay [McK80] observed that the McKay quiver can also be obtained from the geometry of X . Let $\phi : \tilde{X} \rightarrow X$ be the minimal resolution of X , which exists and is unique for surface singularities. The exceptional divisor of ϕ is a union of irreducible curves, each of which is a \mathbb{P}^1 of self-intersection -2 , and it was known that there are n of these (recall that G has $n + 1$ irreducible representations). McKay showed:

Theorem 1.1. ([McK80]) *Let L_1, \dots, L_n be the irreducible components of the exceptional locus of ϕ . Form the dual graph Γ of the exceptional divisor by replacing each curve by a vertex and drawing $\#(L_i \cap L_j)$ edges between the vertices corresponding to L_i and L_j if $i \neq j$. Then Γ is a Dynkin diagram of type \mathbb{A}_n , \mathbb{D}_n , or \mathbb{E}_n , and is equal to the graph obtained from the McKay quiver of G by removing the vertex 0 and replacing each pair of opposing arrows by an edge.*

McKay's result is essentially combinatorial, but it was later extended, originally by Kapranov and Vasserot, to a derived correspondence between \tilde{X} and Λ :

Theorem 1.2. ([KV00]) *Let \tilde{X} and Λ be as above. The categories $\text{coh } \tilde{X}$ and $\text{mod-}\Lambda$ are derived equivalent.*

It is well-known that the resolution $\phi : \tilde{X} \rightarrow X$ is *crepant*: that is, $\phi^* \omega_X \cong \omega_{\tilde{X}}$, or equivalently the relative dualising complex $\phi^! \mathcal{O}_X$ is trivial. Further, the algebra $A \# G$ is Van den Bergh's original example [VdB04a, Example 1.1] of a *noncommutative crepant resolution*. We do not give Van den Bergh's full definition here, which relies on the assumption that the singularity being resolved is commutative. Instead we note that $A \# G$ is 2-Calabi-Yau (see Remark 3.2.7) and A^G is Gorenstein. The dualising complexes of $A \# G$ and of A^G are both trivial and so one

expects the “relative dualising complex” to be trivial: in other words, it makes sense to regard $A \# G$ as a crepant resolution of A^G . See [VdB23] for more discussion.

We briefly mention one more aspect of the commutative McKay correspondence: it is possible to construct the minimal resolution \tilde{X} of X from the preprojective algebra Λ , as in [BKR01]. This is done by using quiver GIT to construct a graded ring T so that $\tilde{X} = \text{Proj } T$. See the beginning of Section 4 for details.

We now discuss how to quantise the above. The appropriate analogue of a finite subgroup of $\text{SL}_2(\mathbb{k})$ in the noncommutative context is a *quantum binary polyhedral group*: a pair (H, A) where A is a two-dimensional Artin-Schelter regular algebra and H is a semisimple finite-dimensional Hopf algebra, acting on A inner faithfully, homogeneously, and with trivial homological determinant. These pairs were classified in [CKWZ16]; the cyclic group C_{n+1} acting on the quantum plane A_q as above is an example. If (H, A) is a quantum binary polyhedral group, then it makes sense to call the invariant ring A^H a *quantum Kleinian singularity*.

It is known that many aspects of the classical McKay correspondence extend to quantum Kleinian singularities. Let (H, A) be a quantum binary polyhedral group. By [CKWZ18, Theorem 0.3] the noncommutative version of *Auslander’s theorem* holds: the natural map

$$\gamma : A \# H \rightarrow \text{End}_{A^H}(A), \quad \gamma(a \# h)(b) = a(h \cdot b)$$

is an isomorphism of rings. It makes sense to view the nonsingular ring $A \# H$ (or, equivalently, $\text{End}_{A^G}(A)$) as a resolution of singularities of A^G ; this is called a *noncommutative quasiresolution* in [QWZ19].

Furthermore, one may construct the McKay quiver Q of the pair (H, A) (see [CKWZ19]), and $A \# H$ is Morita equivalent to a quotient of Λ of $\mathbb{k}Q$ [Cra22]. The relationships between the representation theory of H and of the preprojective algebra in the commutative context carry over to our situation; see [CKWZ19].

In the case of the pair (C_{n+1}, A_q) , the McKay quiver Q is the quiver corresponding to an affine A_n Dynkin diagram. Further, $A_q \# C_{n+1}$ is itself a basic algebra and may be viewed as a quotient of $\mathbb{k}Q$. We set $\Lambda_q := A_q \# C_{n+1}$. We refer to this as an *algebraic resolution* of $B_q = A_q^{C_{n+1}}$, as it clearly parallels the (algebraic) NCCR of a commutative Kleinian singularity. We show in Section 3 that Λ_q is also a crepant resolution of B_q , suitably defined.

In the commutative case the Kleinian singularity $X = \text{Spec}(A^G)$ has two resolutions: the algebraic resolution Λ and the geometric resolution \tilde{X} . In this paper, for the case that $(H, A) = (C_{n+1}, A_q)$, we use techniques of noncommutative algebraic geometry to construct a category \mathcal{X}_q which gives a second resolution of $B_q = A_q^{C_{n+1}}$. That is, there is a pair of adjoint functors

$$\begin{array}{ccc} & \mathcal{X}_q & \\ \uparrow \phi^* & & \downarrow \phi_* \\ & \text{Mod-}B_q & \end{array} \tag{1.3}$$

(These functors are the appropriate noncommutative analogue of a morphism of schemes; see [Smi04] and Section 2.4.) We show:

Theorem 1.4. *The category \mathcal{X}_q is a resolution of $\text{mod-}B_q$. That is:*

- (1) (Theorem 7.2.7) \mathcal{X}_q has finite homological dimension;
- (2) (Theorem 6.3.1(1)) \mathcal{X}_q is proper over $\text{mod-}B_q$ in the sense that ϕ_* takes noetherian objects to noetherian objects; and
- (3) (Proposition 6.2.2) away from the singularity of $\text{mod-}B_q$ the functors (1.3) become inverse equivalences.

We also show that \mathcal{X}_q contains exceptional objects L_1, \dots, L_n whose noncommutative intersection theory gives an A_n Dynkin diagram, extending the classical McKay correspondence Theorem 1.1. See Theorem 9.3.

The construction of \mathcal{X}_q comes from a graded ring T_q which generalises (in fact deforms) the construction of \tilde{X} via quiver GIT. In particular, we also deform the representation variety of Λ to construct a “representation ring” of the algebraic resolution Λ_q , which we show (Theorem 4.2.4) is universal for a certain class of representations of Λ_q . However, to prove most of Theorem 1.4 we must approach \mathcal{X}_q in another way: we show that \mathcal{X}_q may also be viewed as a noncommutative toric variety (Theorem 7.1.11). This allows us to access \mathcal{X}_q via charts and local techniques and in particular is needed to prove Theorem 1.4(1): that \mathcal{X}_q is nonsingular.

Our other main theorem is an extension of Theorem 1.2 to show that the geometric and algebraic resolutions are derived equivalent.

Theorem 1.5. (Theorem 8.3.3) *The derived categories of \mathcal{X}_q and $\text{mod-}\Lambda_q$ are equivalent.*

When $q = 1$, the proof of Theorem 1.5 recovers the proof of Theorem 1.2 given in [Wem16].

Since Λ_q is a crepant resolution of B_q in an appropriate sense, as discussed above, and \mathcal{X}_q and Λ_q are derived equivalent, one expects that \mathcal{X}_q is also a crepant resolution of B_q . We prove this at the end of Section 8.3.

In [Kaw24], Kawamata considers a closely related family of noncommutative singularities, namely the deformations of type \mathbb{A} Kleinian singularities defined in [CBH98], and constructs noncommutative schemes which serve as resolutions of these singularities. See also [Kaw25]. We note that the singularities Kawamata considers, as well as the techniques employed, are distinct from those in this paper.

By adapting the methods in this paper, it should be possible to construct an analogue of the category \mathcal{X}_q for other quantum Kleinian singularities A^H ; in work in progress, we consider the case of an \mathbb{L}_1 singularity.

Organisation of the paper. In Section 2, we give some definitions and basic results, and recall the notion of a noncommutative crepant resolution and a generalisation to noncommutative singularities. Section 3 introduces the singularities B_q of interest, constructs algebraic and categorical resolutions of these singularities, and discusses their crepancy.

The remainder of the paper is dedicated to constructing and studying a geometric resolution of B_q . Motivated by quiver GIT, in Section 4 we define the *representation ring* of Λ_q and show that it is universal in an appropriate sense. We then use this in Section 5 to construct the ring T_q which gives rise to the geometric resolution, and outline some of its properties. In Section 6, we define the category \mathcal{X}_q which is our geometric resolution of B_q and calculate some related cohomology groups. To establish further properties of \mathcal{X}_q , in Section 7 we show that \mathcal{X}_q is equivalent to a noncommutative toric variety, i.e. the category of coherent sheaves over a noncommutative ringed space \mathfrak{X} with a suitable action of a quantum torus. This allows us to, in Section 8, establish a derived equivalence between our algebraic and geometric resolutions. Finally, in Section 9 we compute the intersection arrangement of the lines in the “exceptional locus” in \mathcal{X}_q .

Acknowledgements. The second named author is partially supported by the EPSRC grant EP/T018844/1, and thanks the EPSRC for their support.

We thank the Simons Laufer Mathematical Sciences Institute for hosting us both in the spring of 2024 as part of the program “Noncommutative Algebraic Geometry”. We thank Jenny August, Arend Bayer, Ellen Kirkman, Jon Pridham, Michael Wemyss, and James Zhang for helpful discussions. We also thank Colin Ingalls for sharing his unpublished manuscript [Ing16] on noncommutative toric varieties with us; it gave us helpful inspiration, although our construction of \mathcal{X}_q as a noncommutative toric variety uses a different method.

2. PRELIMINARIES

2.1. Conventions. Throughout \mathbb{k} will denote an algebraically closed field of characteristic 0. Let R be a \mathbb{Z} -graded ring. We write $\text{Gr-}R$ (respectively, $R\text{-Gr}$) for the category of \mathbb{Z} -graded right (respectively, left) R -modules with degree-preserving morphisms, and $\text{gr-}R$ (respectively, $R\text{-gr}$)

for the full subcategory of noetherian objects. More generally, if Xyz is an abelian category, then xyz is the full subcategory of noetherian objects. Given $M \in \text{Gr-}R$, we define $M(i)$ to be the graded module which is isomorphic to M as an ungraded module, but which satisfies $M(i)_n = M_{i+n}$. The categories $\text{Mod-}R$ and $R\text{-Mod}$ are the ungraded right and left module categories of R .

If $M, N \in \text{Gr-}R$, then we write $\text{Hom}_{\text{Gr-}R}(M, N)$ for the space of graded (i.e. degree-preserving) morphisms. By $\text{Hom}_R(M, N)$ we mean $\text{Hom}_{\text{Mod-}R}(M, N)$. We have

$$\underline{\text{Hom}}_R(M, N) = \bigoplus_{s \in \mathbb{Z}} \text{Hom}_{\text{Gr-}R}(M, N(s)).$$

If M_R is finitely generated, then $\text{Hom}_R(M, N) \cong \underline{\text{Hom}}_R(M, N)$.

We write \otimes for $\otimes_{\mathbb{k}}$. The set of non-negative integers is denoted by \mathbb{N} .

2.2. Definitions and basic results. We begin by recalling some definitions and basic results that we will use without mention throughout the paper.

Definition 2.2.1. Suppose that R is a \mathbb{k} -algebra. We say that R is *connected graded* if it is \mathbb{N} -graded with $R_0 = \mathbb{k}$. We say that $M \in \text{Gr-}R$ is *locally finite* if $\dim_{\mathbb{k}} M_i < \infty$ for all $i \in \mathbb{Z}$. The graded module M is called *left bounded* if $M_i = 0$ for $i \ll 0$, *right bounded* if $M_i = 0$ for $i \gg 0$, and *bounded* if it is left and right bounded. If $M \in \text{Gr-}R$ is locally finite, the *Hilbert series* of M is the formal Laurent series

$$\text{hilb } M = \sum_{i \in \mathbb{Z}} (\dim_{\mathbb{k}} M_i) t^i.$$

The following notion provides a suitable concept of dimension for a noncommutative \mathbb{k} -algebra:

Definition 2.2.2. Suppose that R is a \mathbb{k} -algebra and M is a right R -module. The *Gelfand-Kirillov (GK-) dimension* of M is

$$\text{GKdim } M := \sup_{M_0, V} \limsup_{n \rightarrow \infty} \log_n (\dim_{\mathbb{k}} M_0 V^n)$$

where the supremum is taken over all finite-dimensional subspaces M_0 of M and V of A .

We will freely use standard results about GK-dimension, proofs of which can be found in [KL00], for example. If R is a connected graded \mathbb{k} -algebra which is locally finite and $M \in \text{gr-}A$ (which forces M to be left bounded) then, by [KL00, Proposition 6.6], we have

$$\text{GKdim } M = \limsup_{n \rightarrow \infty} \log_n (\dim_{\mathbb{k}} M_n) + 1.$$

Definition 2.2.3. Let $q \in \mathbb{k}^\times$. The *quantum plane* is the ring

$$A_q = \mathbb{k}_q[u, v] := \frac{\mathbb{k}\langle u, v \rangle}{\langle vu - quv \rangle}.$$

This ring is a connected graded noetherian domain. We will sometimes suppress the q and let $A = A_q$. Throughout this paper, we will reserve the letter A for the quantum plane as presented above.

More generally, if $\mathbf{q} = (q_{ij}) \in M_n(\mathbb{k})$ is a multiplicatively antisymmetric matrix, then to this data we can associate a *quantum polynomial ring*, or *quantum n -space*, as follows:

$$\mathbb{k}_{\mathbf{q}}[x_1, \dots, x_n] := \frac{\mathbb{k}\langle x_1, \dots, x_n \rangle}{\langle x_j x_i - q_{ij} x_i x_j \mid 1 \leq i, j \leq n \rangle}.$$

As before, this ring is always a connected graded noetherian domain.

We will frequently work with rings which are (factor rings of) quantum polynomial rings. When performing computations in such rings, the following notation will be convenient: if $q \in \mathbb{k}^\times$, we will use q^\bullet to mean an unspecified power of q . For example, in $\mathbb{k}_q[u, v]$, the monomials u^i and v^j q^\bullet -commute. Moreover, we will sometimes write $\mathbb{k}_{q^\bullet}[x_1, \dots, x_n]$ for a quantum polynomial ring where the generators x_i, x_j commute up to some power of q (possibly

depending on i and j) which can be calculated in principle, but whose specific value does not affect the result.

We will frequently need to invert elements of noetherian domains; the next definition mainly establishes notation.

Definition 2.2.4. Suppose that R is a noetherian domain. Then the set $X = R \setminus \{0\}$ forms a left and right Ore set in R , so we can form the *quotient ring* $Q(R) := RX^{-1} = X^{-1}R$. If, in addition, R is graded by some semigroup Σ , then the set Y of nonzero Σ -homogeneous elements of R forms a left and right Ore set in R [GS00, Theorem 5(a)], so we can form the *graded quotient ring* $Q_{\text{gr}}^{\Sigma}(R) := RY^{-1} = Y^{-1}R$. When the grading is clear from context, we will simply write $Q_{\text{gr}}(R)$.

We remark that the hypotheses imposed on R in the above definition are stronger than required, but all rings to which we need to apply this construction will be noetherian domains.

We will be interested in rings which have various good homological properties. We recall the relevant definitions below:

Definition 2.2.5. Let R be a ring and M a right R -module. The *grade* of M is defined to be

$$j(M) := \inf\{i \mid \text{Ext}_R^i(M, R) \neq 0\} \in \mathbb{N} \cup \{\infty\},$$

with the definition for left modules being similar. We say that M satisfies the *Auslander condition* if for all $i \geq 0$ and all left submodules $N \subseteq \text{Ext}_R^i(M, R)$ we have $j(N) \geq i$.

We say that R is *Gorenstein* if R has finite injective dimension when viewed as a right or left R -module, and that R is *Auslander–Gorenstein* if it is Gorenstein and every right or left R -module satisfies the Auslander condition. If R is Auslander–Gorenstein and has finite global dimension, it is *Auslander regular*.

In many cases, the properties of rings of infinite global dimension are encoded by properties of their so-called maximal Cohen–Macaulay modules, whose definition we now recall:

Definition 2.2.6. Let R be a Gorenstein ring. We say that a finitely generated R -module M is *maximal Cohen–Macaulay (MCM)* if it satisfies $\text{Ext}_R^i(M, R) = 0$ for all $i \geq 1$.

In [CKWZ19], the authors give a more general definition which specialises to the above definition when R is Gorenstein.

We will assume that the reader is familiar with the language of sheaves and derived categories; [Har77] and [Wei94] are standard references for the definitions and results we will use. We will, however, set some notation. If \mathcal{C} is an abelian category, we will write $D(\mathcal{C})$ and $D^b(\mathcal{C})$ for its derived and bounded derived categories, respectively. If R is a noetherian ring, we write $D(R)$ for $D(\text{Mod-}R)$ and $\text{Perf}(R)$ for the full subcategory of perfect complexes. We write $\mathcal{D}(R)$ for $D^b(\text{mod-}R)$, following [VdB23].

2.3. Noncommutative crepant resolutions. Suppose that X is a normal variety with resolution of singularities $\pi : \tilde{X} \rightarrow X$. This resolution is said to be *crepant* if the canonical sheaves ω_X and $\omega_{\tilde{X}}$ satisfy $\pi^*\omega_X = \omega_{\tilde{X}}$. In some sense, crepant resolutions are the “best” nonsingular approximation of a singular variety. In particular, if X is Gorenstein and we want \tilde{X} to also be Calabi–Yau, then the resolution must be crepant.

In [VdB04b, Theorem 4.4.2], Van den Bergh proved the following result, which we paraphrase:

Theorem 2.3.1. *Let R be a (commutative) Gorenstein \mathbb{k} -algebra which is a normal domain, and let $\pi : \tilde{X} \rightarrow X = \text{Spec } R$ be a crepant resolution of singularities. Assume that the fibres of π have dimension at most 1. Then there exists an MCM R -module M such that $\Lambda := \text{End}_R(M)$ is MCM as an R -module and has finite global dimension. Moreover, \tilde{X} and Λ are derived equivalent.*

This motivates the following definition, which is equivalent to the original one due to Van den Bergh:

Definition 2.3.2. Let R be a commutative CM normal domain. A *noncommutative crepant resolution (NCCR)* of R (or of $\text{Spec } R$) is a ring of the form $\Lambda := \text{End}_R(M)$ for some reflexive R -module M , such that Λ has finite global dimension and is an MCM R -module.

For example, by [VdB04a, Example 1.1], if G is a finite subgroup of $\text{SL}_2(\mathbb{k})$ then $\mathbb{k}[x, y] \# G = \text{End}_{\mathbb{k}[x, y]^G}(\mathbb{k}[x, y])$ is an NCCR of the Kleinian singularity $\mathbb{k}[x, y]^G$. If R is a CM local normal domain of dimension ≤ 3 , then it is known that all crepant resolutions and NCCRs of R are derived equivalent [IW13, Theorem 1.5].

If we replace R by a noncommutative ring, then there are several ways to generalise the definition of an NCCR. One such way is the *noncommutative quasi-crepant resolution* defined in [QWZ19]. We begin by recalling some more definitions.

Definition 2.3.3. We say that a \mathbb{k} -algebra R is *GK-Cohen–Macaulay* if

$$\text{GKdim}(M) + j(M) = \text{GKdim}(R) < \infty,$$

for all finitely generated nonzero right R -modules M .

If R is commutative and noetherian, then [BM17, Theorem 4.8] shows that R is GK-Cohen–Macaulay if and only if it is Cohen–Macaulay and equicodimensional (i.e. all of its maximal ideals have the same height).

Definition 2.3.4. Suppose that R and S are locally finite noetherian \mathbb{N} -graded \mathbb{k} -algebras with finite GK-dimension. We say that two graded (R, S) -bimodules M and N are *n -isomorphic*, denoted $M \cong_n N$, if there exist a graded (R, S) -bimodule P and bimodule morphisms

$$f : M \rightarrow P \quad \text{and} \quad g : N \rightarrow P$$

such that the kernels and cokernels of f and g have GK-dimension at most n .

We are now in a position to define the notion of a *noncommutative quasi-crepant resolution*, as given in [QWZ19].

Definition 2.3.5. Suppose that R is a locally finite noetherian \mathbb{N} -graded \mathbb{k} -algebra and that $\text{GKdim } R = d < \infty$. A *noncommutative quasi-resolution (NQR)* of R is a triple (S, M, N) where $S \in \mathcal{A}$ is Auslander regular and GK-Cohen–Macaulay with $\text{GKdim } S = d$, and ${}_S M_R$ and ${}_R N_S$ are bimodules which satisfy

$$M \otimes_R N \cong_0 S \quad \text{and} \quad N \otimes_S M \cong_0 R.$$

If, in addition, R is Auslander–Gorenstein and ${}_S M_R$ and ${}_R N_S$ are reflexive on both sides, then we call this triple a *noncommutative quasi-crepant resolution (NQCR)* of R .

By [QWZ19, Example 8.5], if S is a (commutative) Type \mathbb{A}_n Kleinian singularity, then an NCCR of S is also an NQR.

There is also a notion of a *categorical resolution*, due to Kuznetsov [Kuz08], which may be weakly or strongly crepant. This is discussed in Section 3.3.

2.4. Maps of noncommutative spaces. It will be useful to be able to use the language of maps of noncommutative spaces to discuss functors between categories. Following [Smi04], let \mathbf{A}, \mathbf{B} be (possibly triangulated) categories. A *map* $\Phi : \mathbf{A} \rightarrow \mathbf{B}$ is a pair of (triangulated) functors

$$\begin{array}{c} \mathbf{A} \\ \uparrow \Phi^* \quad \downarrow \Phi_* \\ \mathbf{B} \\ 7 \end{array}$$

so that (Φ^*, Φ_*) is an adjoint pair. This is by analogy with the fact that, if $\pi : Y \rightarrow X$ is a morphism of schemes, then there is an adjoint pair

$$\begin{array}{ccc} & \text{Qcoh } Y & \\ \pi^* \uparrow & & \downarrow \pi_* \\ & \text{Qcoh } X & \end{array}$$

For example, if R, S are rings, then an (R, S) -bimodule M induces a map $\Phi : \text{Mod-}S \rightarrow \text{Mod-}R$ given by

$$\begin{array}{ccc} & \text{Mod-}S & \\ \Phi^* = - \otimes_R M \uparrow & & \downarrow \Phi_* = \text{Hom}_S(M, -) \\ & \text{Mod-}R & \end{array}$$

3. A NONCOMMUTATIVE SINGULARITY

3.1. The singularity of interest. We now introduce the main protagonist of this paper.

Notation 3.1.1. Fix $n \geq 2$ and $q \in \mathbb{k}^\times$. The cyclic group G of order $n+1$ generated by

$$g := \begin{pmatrix} \zeta & 0 \\ 0 & \zeta^{-1} \end{pmatrix},$$

where $\zeta \in \mathbb{k}$ is a primitive $(n+1)$ th root of unity, acts on $A_q = \mathbb{k}_q[u, v]$ as a group of graded automorphisms. Explicitly, we have

$$g \cdot u = \zeta u, \quad g \cdot v = \zeta^{-1} v.$$

To this data we can associate the *skew group ring* $A_q \# G$. As an abelian group this ring is equal to $A_q \otimes \mathbb{k}G$, and we write $a \# g$ for the simple tensor $a \otimes g$. On simple tensors, the multiplication in $A \# G$ is given by

$$(a \# g)(b \# h) := a(g \cdot b) \# gh,$$

which is then extended linearly to all of $A \# G$.

We write $B_q := A_q^G$ for the invariant ring of the action of G on A_q . By [CKWZ18, Table 3], B_q is generated by the elements

$$x := u^{n+1}, \quad y := v^{n+1}, \quad z := uv,$$

and can be presented as a quotient of a quantum polynomial ring by the ideal generated by a single homogeneous, normal, nonzerodivisor:

$$B_q \cong \frac{\mathbb{k}_{q^\bullet}[x, y, z]}{\langle xy - q^{-\frac{1}{2}n(n+1)}z^{n+1} \rangle}, \quad \text{where} \quad \mathbb{k}_{q^\bullet}[x, y, z] := \frac{\mathbb{k}\langle x, y, z \rangle}{\left\langle \begin{array}{c} yx - q^{(n+1)^2}xy \\ zx - q^{n+1}xz \\ zy - q^{-(n+1)}yz \end{array} \right\rangle}. \quad (3.1.2)$$

Following [CKWZ18], we call B_q a *quantum Kleinian singularity of type \mathbb{A}_n* or a *quantum type \mathbb{A} singularity*. The singularity category of B_q is that of a commutative type \mathbb{A} singularity by [Cra18, Theorem 8.2.7].

We now recall various properties of A_q , B_q , and related rings. We begin by noting that B_q , although singular, has good ring-theoretic and homological properties:

Lemma 3.1.3. *The ring $B_q = A_q^G$:*

- (1) *is a noetherian domain;*
- (2) *is Auslander–Gorenstein of injective dimension 2;*
- (3) *is GK–Cohen–Macaulay;*

- (4) has GK-dimension 2; and
(5) has infinite global dimension.

Proof. These are all established in [Cra18, Lemma 6.1.3, Proposition 6.1.6]. \square

The ring B_q has an important family of modules $M^{(i)}$, where $0 \leq i \leq n$, which we now define. We first define (left and right) B_q -modules $M^{i,G}$ for $0 \leq i \leq n$ as follows:

$$M^{i,G} := \{a \in A_q \mid g \cdot a = \zeta^{-i}a\}.$$

As right B_q -modules, these have the explicit form

$$M^{0,G} = B_q, \quad M^{i,G} = u^{n+1-i}B_q + v^iB_q, \quad 1 \leq i \leq n.$$

In terms of the isomorphism (3.1.2), we can write these modules as

$$M^{(0)} = B_q, \quad M^{(i)} = xB_q + z^iB_q, \quad 1 \leq i \leq n.$$

Under the grading inherited from A_q , there are graded isomorphisms

$$M^{(i)} \cong M^{i,G}(i) \quad 1 \leq i \leq n,$$

induced by multiplication by u^i . We are particularly interested in these modules because they are maximal Cohen–Macaulay and they form essentially a complete list of such modules:

Proposition 3.1.4. ([CKWZ19, Theorem C]) *The modules $M^{(i)}$ are indecomposable and maximal Cohen–Macaulay. Moreover, any indecomposable MCM B_q -module is isomorphic (in the ungraded category) to some $M^{(i)}$.*

We now determine the Hilbert series of B_q and the $M^{(i)}$. Note that these Hilbert series are independent of q so, in particular, we can restrict attention to the case $q = 1$ (i.e. when A_q and B_q are commutative). In this case, by [BH93, Theorem 6.4.8] we have

$$\text{hilb } M^{i,G} = \frac{1}{n+1} \sum_{j=0}^n \frac{\zeta^{ij}}{(1 - \zeta^j t)(1 - \zeta^{-j} t)}.$$

A tedious calculation, similar to [CGW25, Example 6.19], gives a closed form for this Hilbert series:

$$\text{hilb } M^{i,G} = \frac{t^i + t^{n+1-i}}{(1 - t^2)(1 - t^{n+1})},$$

and hence

$$\text{hilb } M^{(i)} = \frac{t^{2i} + t^{n+1}}{(1 - t^2)(1 - t^{n+1})}. \quad (3.1.5)$$

Lemma 3.1.6. *For $0 \leq i \leq n$, there is a short exact sequence of right B_q -modules given by*

$$0 \rightarrow M^{(n+1-i)}(-2i) \xrightarrow{\begin{pmatrix} x^{-1} & z^i \\ & -1 \end{pmatrix}} B_q(-(n+1)) \oplus B_q(-2i) \xrightarrow{\begin{pmatrix} x & z^i \end{pmatrix}} M^{(i)} \rightarrow 0,$$

where the maps are given by left multiplication by the indicated matrices.

Proof. It is straightforward to check that the given maps are well-defined and that the sequence forms a complex. For exactness, it suffices to verify that the Euler characteristic of the sequence is 0; indeed, by (3.1.5), the Euler characteristic is

$$-t^{2i} \frac{t^{2(n+1-i)} + t^{n+1}}{(1 - t^2)(1 - t^{n+1})} + (t^{n+1} + t^{2i}) \frac{1 + t^{n+1}}{(1 - t^2)(1 - t^{n+1})} - \frac{t^{2i} + t^{n+1}}{(1 - t^2)(1 - t^{n+1})} = 0,$$

as desired. \square

We will also require the following fact:

Lemma 3.1.7. *There is an isomorphism*

$$\mathrm{Hom}_{B_q}(M^{(i)}, B_q) \cong M^{(n+1-i)}$$

given by $\eta : M^{(n+1-i)} \rightarrow \mathrm{Hom}_{B_q}(M^{(i)}, B_q)$, $\eta(m)(m') = x^{-1}mm'$.

Proof. By the proof of [CKWZ19, Corollary 2.6], there is an equivalence of categories

$$\{\text{graded projective } A_q \# G\text{-modules}\} \rightarrow \mathrm{gr}\text{-}B_q, \quad N \mapsto N^G.$$

Using this, and letting V_i denote the $\mathbb{k}G$ -module with character χ_i satisfying $\chi_i(g) = \zeta^i$, we obtain a chain of isomorphisms

$$\begin{aligned} \mathrm{Hom}_{B_q}(M^{i,G}, B_q) &\cong \mathrm{Hom}_{A_q^G}((V_i \otimes A_q)^G, A_q^G) \\ &\cong \mathrm{Hom}_{A_q \# G}(V_i \otimes A_q, A_q) \\ &\cong \mathrm{Hom}_{A_q}(V_i \otimes A_q, A_q)^G \\ &\cong \mathrm{Hom}_{A_q}(A_q, V_i^* \otimes A_q)^G \\ &\cong \mathrm{Hom}_{A_q}(A_q, V_{n+1-i} \otimes A_q)^G \\ &\cong (V_{n+1-i} \otimes A_q)^G \\ &\cong M^{n+1-i,G}. \end{aligned}$$

Identifying $M^{i,G}$ with $M^{(i)}$ and chasing through the isomorphisms shows that the map giving rise to this isomorphism is as claimed. \square

3.2. An algebraic resolution of B . Write $M = \bigoplus_{i=0}^n M^{(i)}$ for the direct sum of the indecomposable MCM B_q -modules. We in fact have $M \cong A_q$ as objects of $\mathrm{mod}\text{-}B_q$; we obtain an isomorphism of *graded* modules if we replace each $M^{(i)}$ by $M^{i,G}$. The endomorphism ring of this module will play an important role. By [CKWZ18, Theorem 0.3], the *Auslander map*

$$\gamma : A_q \# G \rightarrow \mathrm{End}_{B_q}(A_q), \quad \gamma(a \# g)(b) = a(g \cdot b) \quad (3.2.1)$$

is an isomorphism, and there is a similar isomorphism when we view A_q as a left B_q -module. By [Cra22, Example 6.6], there is a ring isomorphism

$$A_q \# G \cong \Lambda_q \quad (3.2.2)$$

where Λ_q is the quotient of the path algebra of the following quiver by the given relations:

$$\begin{array}{ccc} & 0 & \xleftrightarrow{\bar{\alpha}_0} 1 \\ & \nearrow \bar{\alpha}_n \quad \searrow \alpha_n & \\ n & & 2 \\ & \nwarrow \bar{\alpha}_{n-1} \quad \nearrow \alpha_{n-1} & \\ & & \dots \end{array} \quad \begin{array}{ccc} & 1 & \nwarrow \bar{\alpha}_1 \quad \nearrow \alpha_1 \\ & & 2 \\ & \nwarrow \bar{\alpha}_2 \quad \nearrow \alpha_2 & \\ & & \dots \end{array} \quad \bar{\alpha}_i \alpha_i = q \alpha_{i+1} \bar{\alpha}_{i+1} \text{ for } 0 \leq i \leq n. \quad (3.2.3)$$

(Note that our convention is to compose arrows in quivers from left to right.)

Moreover, we have $e_0 \Lambda_q e_0 \cong B_q$ and, under this isomorphism, we can identify the right B_q -module $M^{i,G}$ with the right $e_0 \Lambda_q e_0$ -module $e_i \Lambda_q e_0$ and identify $M \cong A_q$ with $\Lambda_q e_0$. Under this identification, we can rewrite the isomorphism of (3.2.1) as

$$\gamma' : \Lambda_q \rightarrow \mathrm{End}_{e_0 \Lambda_q e_0}(\Lambda_q e_0), \quad \gamma'(x)(ye_0) = xye_0. \quad (3.2.4)$$

The algebra Λ_q is of particular interest to us because it is an NQCR of B_q .

Lemma 3.2.5. *The triple $(\Lambda_q, e_0\Lambda_q, \Lambda_q e_0)$ is an NQCR of $B_q = e_0\Lambda_q e_0$.*

Proof. That $(\Lambda_q, e_0\Lambda_q, \Lambda_q e_0)$ is an NQR of B_q is [QWZ19, Example 8.5] (where they show that the isomorphic algebra $A_q \# G$ is an NQR). Since B_q is Auslander–Gorenstein, it remains to show that $e_0\Lambda_q$ and $\Lambda_q e_0$ are reflexive on both sides; by symmetry, it suffices to show this for $\Lambda_q e_0$.

Reflexivity of $\Lambda_q e_0$ as a left Λ_q -module follows from the fact that $\Lambda_q e_0$ is a direct summand of Λ_q , hence projective. To prove that $\Lambda_q e_0$ is reflexive as a right $e_0\Lambda_q e_0$ -module, we use (3.2.4). Multiplying this isomorphism on the left by e_0 gives $\text{Hom}_{e_0\Lambda_q e_0}(\Lambda_q e_0, e_0\Lambda_q e_0) \cong e_0\Lambda_q$, and similarly we have $\text{Hom}_{e_0\Lambda_q e_0}(e_0\Lambda_q, e_0\Lambda_q e_0) \cong \Lambda_q e_0$. Therefore

$$\text{Hom}_{e_0\Lambda_q e_0}(\text{Hom}_{e_0\Lambda_q e_0}(\Lambda_q e_0, e_0\Lambda_q e_0), e_0\Lambda_q e_0) \cong \text{Hom}_{e_0\Lambda_q e_0}(e_0\Lambda_q, e_0\Lambda_q e_0) \cong \Lambda_q e_0,$$

so that $\Lambda_q e_0$ is reflexive as a right $e_0\Lambda_q e_0$ -module. \square

Another good property of Λ_q is that it is Morita equivalent to B_q “away from the singularity”. This term is imprecise for a noncommutative ring which has no associated geometry, but note that $B_q[x^{-1}]$, $B_q[y^{-1}]$, $B_q[z^{-1}]$ are all localisations of quantum planes and so have finite global dimension. On the other hand, B_q has infinite global dimension, and by the previous sentence it makes some sense to view the singularity of B as being concentrated where $x = y = z = 0$.

Lemma 3.2.6. *Let $w \in \{x, y, z\}$. Then the algebras $B_q[w^{-1}]$ and $\Lambda_q[w^{-1}]$ are Morita equivalent.*

Proof. First let $w = x$. It is clear that $M^{(i)}[x^{-1}] = B[x^{-1}]$ for all i , and hence $M[x^{-1}] \cong B[x^{-1}]^{n+1}$. Therefore, combining (3.2.2) with the fact that the Auslander map is an isomorphism and localising, we obtain

$$\Lambda_q[x^{-1}] = \text{End}_{B_q}(M)[x^{-1}] \cong \text{End}_{B_q[x^{-1}]}(M[x^{-1}]) \cong \text{End}_{B_q[x^{-1}]}(B_q[x^{-1}]^{n+1}) \cong M_{n+1}(B_q[x^{-1}]),$$

and hence $B_q[x^{-1}]$ and $\Lambda_q[x^{-1}]$ are Morita equivalent. The same proof will work to show that $B_q[y^{-1}]$ and $\Lambda_q[y^{-1}]$ are Morita equivalent, provided we can show that $M^{(i)}[y^{-1}] \cong B_q[y^{-1}]$ for $1 \leq i \leq n$ (the claim for $i = 0$ is clear). Indeed, this follows from the fact that the maps

$$\alpha_i : M^{(i)}[y^{-1}] \rightarrow B_q[y^{-1}], \quad \alpha_i(m) = z^{-1}ym$$

are easily seen to be well-defined maps of right $B_q[y^{-1}]$ -modules which are invertible. Finally, if $w = z$ then the relation $xy = z^{n+1}$ means that we have also inverted x and y , and so the previous arguments apply. \square

Remark 3.2.7. By [RRZ14, Theorem 4.1(c)], Λ_q is *twisted 2-Calabi-Yau* (sometimes called *skew 2-Calabi-Yau*). While we do not give a precise definition, in particular this says that Λ_q is homologically smooth, in the sense that Λ_q has a finite length projective resolution in the category of Λ_q -bimodules such that each term is finitely generated. By [YZ06, Lemma 4.3(b)], this implies that Λ_q has finite global dimension, although not all rings of finite global dimension are homologically smooth. Since Λ_q is twisted 2-CY, it is reasonable to view $\text{mod-}\Lambda_q$ as being “smooth”.

3.3. The algebraic resolution is categorical. Let $\Lambda = \Lambda_q$ and $B = B_q$. As in Section 2.4, the functors

$$\begin{array}{ccc} & \text{Mod-}\Lambda & \\ \psi^* = - \otimes_B e_0\Lambda & \uparrow \downarrow & \psi_* = \text{Hom}_\Lambda(e_0\Lambda, -) \\ & \text{Mod-}B & \end{array}$$

give a map of noncommutative spaces $\psi : \text{Mod-}\Lambda \rightarrow \text{Mod-}B$.

Since Λ has finite global dimension, we think of the category $\text{Mod-}\Lambda$ as being nonsingular, and $\psi : \text{Mod-}\Lambda \rightarrow \text{Mod-}B$ may be viewed as a resolution of singularities, in a sense we will

now make more precise. Let \tilde{D} be a triangulated category and suppose that there is a map of noncommutative spaces $\Psi : \tilde{D} \rightarrow \mathcal{D}(B)$, that is, an adjoint pair of triangulated functors

$$\Psi^* : \text{Perf}(B) \rightarrow \tilde{D}, \quad \Psi_* : \tilde{D} \rightarrow \mathcal{D}(B).$$

Following [Kuz08], \tilde{D} is a *categorical resolution* of $\mathcal{D}(B)$ if \tilde{D} is “smooth” and the natural morphism of functors $\text{id}_{\text{Perf}(B)} \rightarrow \Psi_* \Psi^*$ is an isomorphism. The term “smooth” is somewhat open to interpretation, but it is reasonable to consider the derived category of an abelian category of a homologically smooth algebra to be smooth. (Recall from Remark 3.2.7 that Λ is homologically smooth.) A categorical resolution is said to be *weakly crepant* if (Ψ_*, Ψ^*) also form an adjoint pair. If X is a variety with rational singularities then a resolution of singularities $\Psi : \tilde{X} \rightarrow X$ is crepant exactly when $D^b(\tilde{X})$ is a weakly crepant categorical resolution of $D^b(X)$.

We will see that $\mathcal{D}(\Lambda)$ is a weakly crepant categorical resolution of $\mathcal{D}(B)$. We need some preliminary results first.

Lemma 3.3.1. *Let R be a right noetherian ring, let \mathcal{D} be a triangulated category, and let $\alpha, \beta : \mathcal{D}(R) \rightarrow \mathcal{D}$ be triangulated functors. Suppose that*

- (a) *there is an isomorphism $\theta : \alpha(R) \rightarrow \beta(R)$; and*
- (b) *for all $r \in R$, the following diagram commutes:*

$$\begin{array}{ccc} \alpha(R) & \xrightarrow{\alpha(r \cdot -)} & \alpha(R) \\ \theta \downarrow & & \downarrow \theta \\ \beta(R) & \xrightarrow{\beta(r \cdot -)} & \beta(R) \end{array}$$

Then $\alpha \cong \beta$ as functors from $\text{Perf}(R) \rightarrow \mathcal{D}$.

Proof. Consider the full subcategory of $\mathcal{D}(R)$ with one object, R . The hypothesis is that α, β are naturally isomorphic as functors from this subcategory to \mathcal{D} . The result follows since $\text{Perf}(R)$ is the triangulated envelope of this subcategory. \square

Consider now the two other natural functors between $\text{Mod-}\Lambda$ and $\text{Mod-}B$, namely

$$\begin{array}{ccc} & \text{Mod-}\Lambda & \\ \psi_! = - \otimes_{\Lambda} \Lambda e_0 & \begin{array}{c} \downarrow \uparrow \\ \text{Mod-}B \end{array} & \psi^! = \text{Hom}_B(\Lambda e_0, -) \end{array}$$

given by the isomorphism $\Lambda \cong \text{End}_B(\Lambda e_0)$. Note that $(\psi_!, \psi^!)$ are also an adjoint pair.

Lemma 3.3.2. *There is a natural isomorphism $\psi_* \cong \psi_!$, and the functors ψ^* and $\psi^!$ are equivalent upon restriction to the full subcategory of projective B -modules.*

Proof. We first consider the claimed natural isomorphism $\psi_* \cong \psi_!$. A straightforward adaptation of [AF92, Proposition 20.11] shows that there is a natural isomorphism

$$\text{Hom}_{\Lambda}(\Lambda, -) \otimes_{\Lambda} \Lambda e_0 \cong \text{Hom}_{\Lambda}(\text{Hom}_{\Lambda}(\Lambda e_0, \Lambda), -).$$

However, the left hand functor is naturally isomorphic to $\psi_!$ and, since $\text{Hom}_{\Lambda}(\Lambda e_0, \Lambda) \cong e_0 \Lambda$, the right hand functor is naturally isomorphic to ψ_* , giving the claimed result. Explicitly, the natural isomorphism $\eta : \psi_! \rightarrow \psi_*$ arising this way is given by, for $N \in \text{Mod-}\Lambda$,

$$\eta_N : N \otimes_{\Lambda} \Lambda e_0 \rightarrow \text{Hom}_{\Lambda}(e_0 \Lambda, N), \quad \eta_N(n \otimes x e_0)(e_0 y) = n x e_0 y.$$

For the second claim, it is straightforward to verify that $\theta : \psi^* \rightarrow \psi^!$ defined on $K \in \text{Mod-}B$ via

$$\theta_K : K \otimes_B e_0 \Lambda \rightarrow \text{Hom}_B(\Lambda e_0, K), \quad \theta_K(k \otimes e_0 x)(y e_0) = k e_0 x y e_0,$$

is a natural transformation. To show that this is an equivalence upon restricting to projective modules, by [AF92, Lemma 20.9], it suffices to check that ψ^* and $\psi^!$ agree on $B = e_0\Lambda e_0$. Indeed,

$$\psi^*(B) = B \otimes_B e_0\Lambda \cong e_0\Lambda \cong \text{Hom}_B(\Lambda e_0, e_0\Lambda e_0) = \text{Hom}_B(\Lambda e_0, B) = \psi^!(B),$$

where the isomorphism $e_0\Lambda \cong \text{Hom}_B(\Lambda e_0, e_0\Lambda e_0)$ follows from (3.2.4). \square

Proposition 3.3.3. *Since ψ^* and ψ_* are left exact and right exact, respectively, we may derive them to obtain functors $\text{Perf}(B) \rightarrow D(\Lambda)$ and $D(\Lambda) \rightarrow \mathcal{D}(B)$, which we will also call ψ^* and ψ_* . This data gives a weakly crepant categorical resolution of $\mathcal{D}(B)$.*

Proof. The functors $\psi^* = - \otimes_B e_0\Lambda$ and $\psi_* = \text{Hom}_\Lambda(e_0\Lambda, -)$ form an adjoint pair, and the same is true for their derived analogues. There is an isomorphism

$$\psi_*\psi^*(B) = \text{Hom}_\Lambda(e_0\Lambda, B \otimes_B e_0\Lambda) \cong \text{End}_\Lambda(e_0\Lambda) \cong e_0\Lambda e_0 = B,$$

from which it follows that the natural morphism $\text{id}_{\text{Mod-}B} \rightarrow \psi_*\psi^*$ restricts to an equivalence on projective B -modules. Lemma 3.3.1 then shows that ψ^* and ψ_* give rise to a categorical resolution of $\mathcal{D}(B)$. Moreover, if $N \in \text{Mod-}\Lambda$ and P is a projective B -module then

$$\text{Hom}_B(\psi_*(N), P) \cong \text{Hom}_B(\psi_!(N), P) \cong \text{Hom}_\Lambda(N, \psi^!(P)) \cong \text{Hom}_\Lambda(N, \psi^*(P)),$$

where we have used Lemma 3.3.2 and the fact that $(\psi_!, \psi^!)$ form an adjoint pair. Thus (ψ_*, ψ^*) are an adjoint pair on the level of modules and hence the same is true for their derived counterparts, and so this categorical resolution is weakly crepant. \square

There is also a notion of a *strongly crepant* categorical resolution [VdB23], corresponding to triviality of the relative dualising complex; we believe this resolution to be strongly crepant, but have not been able to prove it. By [VdB23, Lemma 2.2.1], NCCRs of normal Gorenstein algebraic varieties are strongly crepant.

There is another sense in which $\Lambda = \text{End}_B(M)$ is a “crepant” resolution of B : the module M_B is reflexive and the inclusion $B = A^{C_{n+1}} \subseteq A \# C_{n+1} \cong \Lambda$ makes Λ an MCM left and right B -module. Indeed, as a (left or right) B -module we have $A \# C_{n+1} \cong A^{n+1}$, and A is an MCM B -module by [CKWZ16, Theorem A]. Thus Λ satisfies the definition of an NCCR if we allow the singular ring to be noncommutative.

4. THE REPRESENTATION RING

Lemmata 3.2.5 and 3.2.6 show that it is reasonable to think of Λ_q as an algebraic resolution of singularities of B_q . The rest of the paper will be devoted to constructing a more “geometric” resolution of B_q , studying it, and comparing this resolution to Λ_q .

We begin by reviewing the geometric resolution \tilde{X} of $X = \mathbb{A}^2/G = \text{Spec } B_1$. This can be constructed via quiver GIT (geometric invariant theory), since $G = C_{n+1} \leq \text{SL}_2(\mathbb{k})$ is cyclic. This is essentially due to [BKR01], but we follow [VdB23, Theorem 6.3.1], as explained in [Wem16, Section 3].

Let $q = 1$ and let $\Lambda = \Lambda_1$ be the ring defined by (3.2.3), which we have seen is an NCCR of the type \mathbb{A} singularity $B = B_1$. Let $Y = \text{Rep}(\Lambda, (1, 1, \dots, 1))$ be the variety parametrising representations of Λ with dimension vector $(1, 1, 1, \dots, 1)$. Such a representation is given by a choice of $2(n+1)$ scalars, by which the generators of Λ act, satisfying the relations of Λ . Equivalently we have

$$\mathbb{k}[Y] = \mathbb{k}[a_0, \dots, a_n, \bar{a}_0, \dots, \bar{a}_n] / \langle a_0\bar{a}_0 = a_1\bar{a}_1 = \dots a_n\bar{a}_n \rangle.$$

Let $R := \mathbb{k}[Y]$. There is a natural action of $\mathbb{T}^{n+1} = (\mathbb{k}^\times)^{n+1}$ on Y , given by scaling the coordinates in such a way as to preserve $a_0\bar{a}_0$, and two representations are isomorphic if and only if they are in the same \mathbb{T}^{n+1} -orbit. Equivalently, there is a natural \mathbb{Z}^{n+1} -grading on R given by setting $\deg a_i = e_i$, $\deg \bar{a}_i = -e_i$. Note that the invariant ring $R^{\mathbb{T}^{n+1}} = R_{\vec{0}}$ is isomorphic to the singularity $A^{C_{n+1}} = \mathbb{k}[X] = B_1$; that is, X is isomorphic to the categorical quotient

$Y//\mathbb{T}^{n+1}$. Moreover, the GIT quotient $Y/\chi \mathbb{T}^{n+1}$, for a suitable weight χ , gives the minimal resolution \tilde{X} of X . Explicitly, let $\chi = (-n, 1, \dots, 1)$ and define

$$\tilde{X} := Y/\chi \mathbb{T}^{n+1} = \text{Proj}\left(\bigoplus_{n \geq 0} R_{n\chi}\right).$$

Thus it is natural to attempt to construct an alternative resolution of our noncommutative singularity B_q is by quantising the representation variety Y , or at least its coordinate ring R .

4.1. Defining the representation ring. Let $Y = \text{Rep}(\Lambda_1) := \text{Rep}(\Lambda_1, (1, 1, \dots, 1))$ be the representation variety of Λ_1 , where we have suppressed the dimension vector in the notation. We refer to the coordinate ring of Y , that is

$$R_1 = \mathbb{k}[Y] = \mathbb{k}[\text{Rep}(\Lambda_1)] = \frac{\mathbb{k}[a_i, \bar{a}_i \mid 0 \leq i \leq n]}{\langle a_0 \bar{a}_0 = a_1 \bar{a}_1 = \dots = a_n \bar{a}_n \rangle}, \quad (4.1.1)$$

as the *representation ring* of Λ_1 .

Now let $q \in \mathbb{k}^\times$ be arbitrary. As usual, we cannot expect a geometric object to parametrise representations of Λ_q , but we can still define a useful deformation of the representation ring R_1 .

Definition 4.1.2. The *representation ring* of Λ_q is

$$R_q^{(n)} := \frac{\mathbb{k}\langle a_i, \bar{a}_i \mid 0 \leq i \leq n \rangle}{\left\langle \begin{array}{cc} a_j a_i = a_i a_j & \bar{a}_j \bar{a}_i = \bar{a}_i \bar{a}_j \\ \bar{a}_j a_i = q a_i \bar{a}_j & a_0 \bar{a}_0 = a_i \bar{a}_i \end{array} \mid 0 \leq i, j \leq n \right\rangle}.$$

When the values of n or q are clear from context or immaterial, we will simply write R (or sometimes R_q) for this ring.

We begin by recording some basic properties of $R_q^{(n)}$.

Lemma 4.1.3. Let $R^{(n)} = R_q^{(n)}$. Then $R^{(n)}$:

(1) *is connected graded with Hilbert series*

$$h_{R^{(n)}}(t) = \frac{(1 - t^2)^n}{(1 - t)^{2(n+1)}}$$

if we set $\deg a_i = 1 = \deg \bar{a}_i$ for all i ;

(2) *has GK-dimension $n + 2$;*

(3) *is a domain;*

(4) *is noetherian.*

Proof.

(1) Consider the monomials of $R^{(n)}$ under the graded lexicographic ordering with

$$a_0 < \bar{a}_0 < a_1 < \bar{a}_1 < \dots < a_n < \bar{a}_n.$$

It is then straightforward to check that the given generators for the ideal of relations defining $R^{(n)}$ form a Gröbner basis. Therefore a basis for the degree d piece of $R^{(n)}$ is

$$\{a_0^{i_0} \bar{a}_0^{j_0} x_1^{i_1} x_2^{i_2} \dots x_n^{i_n} \mid 0 \leq i_k \leq d, 0 \leq j_0 \leq d, i_0 + j_0 + i_1 + \dots + i_n = d, x_k \in \{a_k, \bar{a}_k\}\}.$$

In particular, it follows that a_0 is a nonzerodivisor which is also normal, and therefore we have

$$h_{R^{(n)}}(t) = \frac{h_{R^{(n)}/\langle a_0 \rangle}(t)}{1 - t}.$$

Now define an auxiliary algebra

$$S^{(n)} := \frac{\mathbb{k}\langle a_i, \bar{a}_i \mid 1 \leq i \leq n \rangle}{\left\langle \begin{array}{cc} a_j a_i = a_j a_i & \bar{a}_j \bar{a}_i = \bar{a}_i \bar{a}_j \\ \bar{a}_j a_i = q a_i \bar{a}_j & a_i \bar{a}_i = 0 \end{array} \mid 1 \leq i, j \leq n \right\rangle},$$

and note that

$$R^{(n)}/\langle a_0 \rangle \cong S^{(n)}[\bar{a}_0; \theta].$$

is a skew polynomial ring for an appropriate choice of automorphism $\theta : S^{(n)} \rightarrow S^{(n)}$. It follows that

$$h_{R^{(n)}}(t) = \frac{h_{S^{(n)}[\bar{a}_0]}(t)}{1-t} = \frac{h_{S^{(n)}}(t)}{(1-t)^2},$$

so it suffices to determine the Hilbert series of $S^{(n)}$.

By an argument similar to the one used for $R^{(n)}$, a basis for the degree d piece of $S^{(n)}$ is given by

$$\{x_1^{i_1} \dots x_n^{i_n} \mid 0 \leq i_k \leq d, i_1 + \dots + i_n = d, x_k \in \{a_k, \bar{a}_k\}\}.$$

We can decompose this set as

$$\bigsqcup_{\ell=0}^n \{x_{j_1}^{i_{j_1}} \dots x_{j_\ell}^{i_{j_\ell}} \mid 1 \leq i_{j_k} \leq d, i_{j_1} + \dots + i_{j_\ell} = d, x_k \in \{a_k, \bar{a}_k\}, 1 \leq j_1 < j_2 < \dots < j_\ell \leq n\}.$$

In each disjoint subset of this union, there are $\binom{n}{\ell}$ choices for (j_1, \dots, j_ℓ) and then 2^ℓ choices for $(x_{j_1}, \dots, x_{j_\ell})$. Therefore the generating function for the ℓ th stratum is

$$\frac{2^\ell \binom{n}{\ell} t^\ell}{(1-t)^\ell} = \frac{\binom{n}{\ell} (2t)^\ell}{(1-t)^\ell}.$$

It follows that

$$h_{S^{(n)}}(t) = \sum_{\ell=0}^n \frac{\binom{n}{\ell} (2t)^\ell}{(1-t)^\ell} = \frac{1}{(1-t)^n} \sum_{\ell=0}^n \binom{n}{\ell} (2t)^\ell (1-t)^{n-\ell} = \frac{\left((1-t) + (2t)\right)^n}{(1-t)^n} = \frac{(1-t^2)^n}{(1-t)^{2n}}.$$

Therefore,

$$h_{R^{(n)}}(t) = \frac{h_{S^{(n)}}(t)}{(1-t)^2} = \frac{(1-t^2)^n}{(1-t)^{2(n+1)}}.$$

(2) In reduced form, the Hilbert series of $R^{(n)}$ is

$$h_{R^{(n)}}(t) = \frac{(1+t)^n}{(1-t)^{n+2}}.$$

Since this has a pole at 1 of order $n+2$, we have $\text{GKdim } R^{(n)} = n+2$.

(3) The proof of part (1) shows that $z := a_0 \bar{a}_0$ is a normal nonzerodivisor in $R^{(n)}$, so we have an inclusion

$$R^{(n)} \hookrightarrow R^{(n)}[z^{-1}].$$

It is straightforward to see that there is an isomorphism

$$R^{(n)}[z^{-1}] \cong \frac{\mathbb{K}[a_0^{\pm 1}, \dots, a_n^{\pm 1}] \langle z^{\pm 1} \rangle}{\langle za_i = qa_i z \mid 0 \leq i \leq n \rangle},$$

which, being a localisation of a quantum plane, is a domain, and hence the same is true of $R^{(n)}$.

(4) Since $R^{(n)}$ is a quotient of a quantum polynomial ring, it is noetherian. \square

4.2. A universal property. Let $R = R_q^{(n)}$. If $q = 1$ then R is the coordinate ring of the representation variety of Λ_1 . It is thus reasonable to ask if R is “universal” for representations of Λ_q for arbitrary q . In this section we show that this is true at least when q is not a root of unity, or when we consider R_q as a family parametrised by q .

We first give a general definition.

Definition 4.2.1. Let \mathbb{K} be a commutative ring, let Q be a quiver, and let $\Lambda = \mathbb{K}Q/I$ be a basic \mathbb{K} -algebra. We assume here that $I \subseteq (\mathbb{K}Q)_{\geq 2}$ is a graded ideal with respect to the path length grading on $\mathbb{K}Q$. Let Γ be an \mathbb{N} -graded \mathbb{K} -algebra, and induce a grading on $M_{Q_0}(\Gamma)$ by

giving the matrix units degree 0. A *graded Γ -representation* of Λ is an \mathbb{N} -graded \mathbb{K} -algebra homomorphism

$$\iota = \iota_\Gamma : \Lambda \rightarrow M_{Q_0}(\Gamma) = \text{End}_\Gamma(\Gamma^{Q_0})$$

which sends, for each $i \in Q_0$, the vertex idempotent e_i to the matrix unit e_{ii} , and which does not send any arrows (length 1 paths in Λ) to 0.

Thus $\iota(e_i \Lambda e_j) \subseteq e_i M_{Q_0}(\Gamma) e_j = e_{ij} \Gamma$, which we identify with Γ . We will abuse notation and write $\iota(e_i \Lambda e_j) \subseteq \Gamma$.

Remark 4.2.2. Technically, this should be called a graded Γ -representation with dimension vector $(1, \dots, 1)$, but we will suppress the dimension vector in the notation.

Given a graded Γ -representation of Λ , then ι_Γ induces an action of Λ on Γ^{Q_0} , which is thus a representation of Λ in the usual sense: the arrows in Λ take us from one copy of Γ to another. This corresponds to a right action of Λ on Γ^{Q_0} , and so we regard Γ^{Q_0} as a row vector.

We now formally define the family of the R_q . Let $\mathbb{K} = \mathbb{k}[q, q^{-1}]$, where q is an indeterminate. Fix n and let $R_\mathbb{K} = R_\mathbb{K}^{(n)}$ be the \mathbb{K} -algebra defined by the presentation

$$R_\mathbb{K} := \frac{\mathbb{K}\langle a_i, \bar{a}_i \mid 0 \leq i \leq n \rangle}{\left\langle \begin{array}{cc} a_j a_i = a_i a_j & \bar{a}_j \bar{a}_i = \bar{a}_i \bar{a}_j \\ \bar{a}_j a_i = q a_i \bar{a}_j & a_0 \bar{a}_0 = a_i \bar{a}_i \end{array} \mid 0 \leq i, j \leq n \right\rangle}.$$

Let $\Lambda_\mathbb{K}$ be the path algebra over \mathbb{K} of the quiver Q with relations as in (3.2.3). Further, let $A_\mathbb{K} = \mathbb{K}\langle u, v \rangle / \langle vu - quv \rangle$.

Lemma 4.2.3. *The \mathbb{K} -algebras $A_\mathbb{K}$, $R_\mathbb{K}$, and $\Lambda_\mathbb{K}$ are flat over \mathbb{K} .*

Proof. For $A_\mathbb{K}$ the statement is obvious. For $R_\mathbb{K}$ and $\Lambda_\mathbb{K}$ it follows from Lemma 4.1.3(1) and [Mat89, Theorem 23.1] (see also [Har77, Theorem III.9.9]). \square

In this section we will prove that $R_\mathbb{K}$ gives a representation of $\Lambda_\mathbb{K}$ which is universal for graded $A_\mathbb{K}$ -representations, and that the same holds when we specialise to let $q \in \mathbb{k}^\times$ be any non-root of unity.

Theorem 4.2.4.

- (1) *There is a graded $R_\mathbb{K}$ -representation $\iota_{R_\mathbb{K}}$ of $\Lambda_\mathbb{K}$ given by sending $\alpha_i \rightarrow a_i$, $\bar{\alpha}_i \rightarrow \bar{a}_i$. This specialises to give a graded R_q -representation of Λ_q for any $q \in \mathbb{k}^\times$.*
- (2) *$R_\mathbb{K}$ is universal for graded $A_\mathbb{K}$ -representations of $\Lambda_\mathbb{K}$ in the sense that given a graded $A_\mathbb{K}$ -representation ι_A of $\Lambda_\mathbb{K}$, there is a unique \mathbb{K} -algebra homomorphism $\phi : R_\mathbb{K} \rightarrow A_\mathbb{K}$ so that the diagram*

$$\begin{array}{ccc} \Lambda_\mathbb{K} & \xrightarrow{\iota_{R_\mathbb{K}}} & M_{Q_0}(R_\mathbb{K}) \\ & \searrow \iota_A & \downarrow M_{Q_0}(\phi) \\ & & M_{Q_0}(A_\mathbb{K}) \end{array}$$

commutes.

- (3) *If q is not a root of unity, then R_q is universal for graded A_q -representations of Λ_q in the sense defined by the obvious modification of (2).*

Remark 4.2.5. As usual, it follows from the universal properties in Theorem 4.2.4 that $R_\mathbb{K}$ is unique up to isomorphism, and similarly for R_q when q is not a root of unity. This justifies calling R the representation ring of Λ .

Before giving the proof of Theorem 4.2.4, we give two quick lemmata about multiplication in $A_\mathbb{K}$ and A_q .

Lemma 4.2.6.

- (1) *Let f, g be nonzero elements of $(A_\mathbb{K})_{n+1}$. Then $gf = q^{(n+1)^2} fg$ if and only if $g \in \mathbb{K}v^{n+1}$, $f \in \mathbb{K}u^{n+1}$.*

- (2) Suppose that q is not a root of unity, let $A = A_q$, and let f, g be nonzero elements of A_{n+1} . Then $gf = q^{(n+1)^2}fg$ if and only if $g \in \mathbb{K}v^{n+1}$, $f \in \mathbb{K}u^{n+1}$.

Proof. The proofs are similar; we prove (2). We have

$$u^a v^b u^c v^d = q^{bc-ad} u^c v^d u^a v^b, \quad (4.2.7)$$

and sufficiency follows immediately. For necessity, first suppose that $g = u^{n+1-b}v^b$, $f = u^c v^{n+1-c}$ are monomials. From (4.2.7), $gf = q^{(n+1)^2}fg$, and the assumption that q is not a root of unity, we have

$$(n+1)^2 = bc - (n+1-b)(n+1-c) = (n+1)(b+c-(n+1));$$

in other words $b+c = 2(n+1)$. As $0 \leq b, c \leq n+1$, the only way this can happen is if $b = c = n+1$.

For general f, g , put a monomial order on monomials in A so that $u > v$, so v^{n+1} is the smallest monomial in A_{n+1} . The leading monomials of f, g must $q^{(n+1)^2}$ -commute, and the monomial case forces $g = \lambda v^{n+1}$. It is then easy to see that for f to $q^{(n+1)^2}$ -commute with g forces $f = \mu u^{n+1}$. \square

Lemma 4.2.8.

- (1) Let $x_0, \dots, x_n \in (A_{\mathbb{K}})_1$ be nonzero. Then $x_0 x_1 \dots x_n \in \mathbb{K}u^{n+1}$ if and only if $x_0, \dots, x_n \in \mathbb{K}u$, and $x_0 x_1 \dots x_n \in \mathbb{K}v^{n+1}$ if and only if $x_0, \dots, x_n \in \mathbb{K}v$.
(2) Let $q \in \mathbb{K}^\times$ be arbitrary and let $A = A_q$. Let $x_0, \dots, x_n \in A_1$ be nonzero. Then $x_0 x_1 \dots x_n \in \mathbb{K}u^{n+1}$ if and only if $x_0, \dots, x_n \in \mathbb{K}u$, and $x_0 x_1 \dots x_n \in \mathbb{K}v^{n+1}$ if and only if $x_0, \dots, x_n \in \mathbb{K}v$.

Proof. For the first claim, consider the leading term of $x_0 \dots x_n$ in a monomial order with $v > u$. For the second claim, use an order with $u > v$. \square

Proof of Theorem 4.2.4. (1) We wish to define a homomorphism $\iota_R : \Lambda_{\mathbb{K}} \rightarrow M_{Q_0}(R_{\mathbb{K}})$ by sending

$$\alpha_i \mapsto e_{i,i+1}a_i, \quad \bar{\alpha}_i \mapsto e_{i+1,i}\bar{a}_i,$$

where the subscripts are interpreted modulo $n+1$. By construction, ι_R respects the relations of $\mathbb{K}Q$. Moreover, we have

$$\iota_R(\bar{\alpha}_i)\iota_R(\alpha_i) = e_{i+1,i+1}\bar{a}_i a_i = q e_{i+1,i+1}a_{i+1}\bar{a}_{i+1} = q \iota_R(\alpha_{i+1})\iota_R(\bar{\alpha}_{i+1}),$$

so ι_R respects the relations of $\Lambda_{\mathbb{K}}$ and gives a well-defined homomorphism. It is clear that ι_R specialises as claimed.

Statements (2) and (3) have similar proofs; we prove (2). A graded $A_{\mathbb{K}}$ -representation of $\Lambda_{\mathbb{K}}$ is equivalent to a choice of nonzero $a_0, \dots, a_n, \bar{a}_0, \dots, \bar{a}_n \in (A_{\mathbb{K}})_1$ so that

$$\bar{a}_i a_i = q a_{i+1} \bar{a}_{i+1}$$

for $i \in \{0, \dots, n\}$, where the subscripts are understood modulo $n+1$. We will show (abusing notation) that $a_0, \dots, a_n, \bar{a}_0, \dots, \bar{a}_n$ satisfy the relations of $R_{\mathbb{K}}$.

Let $f = a_0 \dots a_n$, $g = \bar{a}_n \dots \bar{a}_0$. Then

$$\begin{aligned} gf &= q \bar{a}_n \dots \bar{a}_2 (\bar{a}_1 a_1)^2 a_2 \dots a_n \\ &= q q^2 \bar{a}_n \dots \bar{a}_3 (\bar{a}_2 a_2)^3 a_3 \dots a_n \\ &= \dots = q^{1+\dots+n} (\bar{a}_n a_n)^{n+1} = q^{\binom{n+1}{2}} q^{n+1} (a_0 \bar{a}_0)^{n+1}. \end{aligned}$$

Likewise,

$$fg = q^{-\binom{n+1}{2}} (a_0 \bar{a}_0)^{n+1},$$

so $gf = q^{(n+1)^2}fg$. Combining Lemmata 4.2.6 and 4.2.8, we see that $a_0, \dots, a_n \in \mathbb{K}u$, and $\bar{a}_0, \dots, \bar{a}_n \in \mathbb{K}v$. It follows immediately that the relations for $R_{\mathbb{K}}$ hold for $a_0, \dots, a_n, \bar{a}_0, \dots, \bar{a}_n$. \square

Remark 4.2.9. If $q = 1$, then the representation variety of Λ parametrises all (commutative) representations of Λ , and so R is universal for (ungraded) representations of Λ into any commutative ring, using the obvious slight generalisation of Definition 4.2.1. Since any graded A -representation of Λ is a commutative representation, it follows that R is also universal for graded A -representations. The argument here is quite different, however, than the proof of Theorem 4.2.4.

Example 4.2.10. If q is a root of unity then R_q need not be universal in the sense of Theorem 4.2.4 (3), even given the universality of $R_{\mathbb{K}}$. For example, fix n and suppose that q is a primitive $(n+1)$ th root of unity. Let $A = A_q$, $\Lambda = \Lambda_q$, and define an A -representation of Λ by

$$\iota_A : \Lambda \rightarrow M_{Q_0}(A), \quad a_i \mapsto v e_{i,i+1}, \quad \bar{a}_i \mapsto q^{-2i} u e_{i+1,i}.$$

This gives an A -representation of Λ since

$$\iota_A(\bar{a}_i a_i - q a_{i+1} \bar{a}_{i+1}) = \begin{cases} q^{-2i} uv - q^{-2(i+1)+1} vu = q^{-2i} uv - q^{-2(i+1)+2} uv = 0 & \text{if } i \neq n, \\ q^{-2n} uv - q v u = q^{-2n} uv - q^2 uv = 0 & \text{if } i = n. \end{cases}$$

However, it is easy to see that this representation does not factor through R_q .

4.3. An embedding of R . Fix n, q and let $R = R_q^{(n)}$. For later use, we note that R embeds in an iterated Laurent polynomial ring over A .

Lemma 4.3.1. *There is an embedding*

$$\varphi : R \hookrightarrow A[s_0^{\pm 1}, \dots, s_n^{\pm 1}], \quad a_i \mapsto u \frac{s_{i+1}}{s_i}, \quad \bar{a}_i \mapsto v \frac{s_i}{s_{i+1}}.$$

Proof. It is straightforward to verify that φ respects the relations of R . Let

$$R' = \mathbb{K}\langle u \frac{s_{i+1}}{s_i}, v \frac{s_i}{s_{i+1}} \mid 0 \leq i \leq n \rangle \subseteq A[s_0^{\pm 1}, \dots, s_n^{\pm 1}] \quad (4.3.2)$$

be the image of φ . Note that uv is a normal element of R' and that

$$R'[(uv)^{-1}] = \mathbb{K}_{\mathbf{q}}\left[\left(v \frac{s_1}{s_0}\right)^{\pm 1}, \left(v \frac{s_2}{s_1}\right)^{\pm 1}, \dots, \left(v \frac{s_n}{s_{n-1}}\right)^{\pm 1}, (uv)^{\pm 1}\right],$$

which is a localisation of some quantum polynomial ring in $n+2$ variables and hence has GK-dimension $n+2$. It follows that $\text{GKdim } R' = n+2$. We therefore have a surjection $R \twoheadrightarrow R'$ between domains of GK-dimension $n+2$. By [KL00, Proposition 3.15], this map must be an isomorphism. \square

5. A GEOMETRIC RESOLUTION: THE RING

In this section we construct an \mathbb{N} -graded ring T which will give our “geometric” resolution of $B = B_q$. Unsurprisingly, T will be a deformation of the semi-invariant ring used to define the minimal resolution of a commutative \mathbb{A}_n singularity.

5.1. The ring T . We first define T , or $T_q^{(n)}$ if we want to make the dependence on n and q explicit. Again, let $A = A_q$ and $R = R_q^{(n)}$. If we give $\mathbb{K}[s_0^{\pm 1}, \dots, s_n^{\pm 1}]$ the standard \mathbb{Z}^{n+1} -grading with $\deg s_i = e_i$, then there is an induced \mathbb{Z}^{n+1} -grading on $A[s_0^{\pm 1}, \dots, s_n^{\pm 1}]$ and on R via the embedding in Lemma 4.3.1. We will refer to this grading as the *weight grading* on R , and write $\text{wt}(a)$ for the degree of a weight-homogeneous element a .

Following the approach of quiver GIT, set

$$\chi := (-n, 1, 1, \dots, 1) \in \mathbb{Z}^{n+1}$$

and let $T = T_q^{(n)}$ be the ring of semi-invariants of R corresponding to χ :

$$T := \bigoplus_{k \in \mathbb{N}} R_{k\chi}.$$

That is, T is the \mathbb{N} -graded ring whose k th graded piece consists of elements of R of weight $k\chi$. We will refer to this as the *projective grading* on T , to avoid confusion with the other gradings that we can (and will) use. If $k \in \mathbb{N}$, then T_k is always the k th graded piece under the projective grading. Later we will use the projective grading on T to define a category $\text{qgr-}T$, as in noncommutative projective geometry.

Before doing this, though, we need to study the elementary algebraic properties of T . We can view $T = T_q^{(n)}$ as a subring of $R = R_q^{(n)}$ as given in (4.1.1), or as a subring of $A_q[s_i^{\pm 1}]$ via Lemma 4.3.1; both perspectives will be useful.

We first record a generating set for T .

Proposition 5.1.1. *Fix n and let $T = T_q^{(n)}$. Then T is generated by the elements*

$$\begin{aligned} x &:= a_0 a_1 \dots a_n, & y &:= \bar{a}_0 \bar{a}_1 \dots \bar{a}_n, & z &:= a_0 \bar{a}_0, \\ t_i &:= a_0^{n-i} a_1^{n-i-1} \dots a_{n-i-2}^2 a_{n-i-1} \bar{a}_{n-i+1} \bar{a}_{n-i+2}^2 \dots \bar{a}_{n-1}^{i-1} \bar{a}_n^i. \end{aligned}$$

Further, $T_0 = B$ and $\text{GKdim } T \geq 3$.

Remark 5.1.2. We will see later that $\text{GKdim } T = 3$.

Proof of Proposition 5.1.1. Observe that $x, y, z \in T_0$ and $t_0, \dots, t_n \in T_1$. A general element of R is homogeneous of projective degree d if and only if each monomial in the a_i and \bar{a}_j appearing in it with nonzero coefficient has projective degree d , so it suffices to show that the claimed generating set generates all monomials in T_d for all $d \in \mathbb{N}$. We will prove this by induction on d .

We first show that x, y , and z generate T_0 . Any monomial in R can be written in the form

$$a = a_0^{i_0} \dots a_n^{i_n} \bar{a}_0^{j_0} \dots \bar{a}_n^{j_n},$$

where all of the exponents are non-negative. So suppose a monomial a as above has projective degree 0. For $0 \leq k \leq n$, if i_k and j_k are both nonzero, then we can pull out a factor of $z = a_k \bar{a}_k$. Therefore we may assume that at least one of i_k and j_k is 0 for all $0 \leq k \leq n$. By considering the $n+1$ components comprising the weight of a , we obtain $n+1$ linear equations:

$$-i_k + i_{k-1} + j_k - j_{k-1} = 0, \quad 0 \leq k \leq n,$$

where the subscripts are interpreted modulo $n+1$. First assume $j_0 = 0$, so that $-i_0 + i_n - j_n = 0$. For contradiction, suppose $j_n > 0$. By our earlier assumption, this forces $i_n = 0$, and hence $i_0 + j_n = 0$, which is absurd. Therefore $j_n = 0$, which also gives $i_0 = i_n$. Repeating this with the other equations shows that $i_0 = i_1 = \dots = i_n =: c$ and $j_k = 0$ for $0 \leq k \leq n$. Therefore $a = a_0^c a_1^c \dots a_n^c$, which is a scalar multiple of x^c . If instead $i_0 = 0$, we find that a is a multiple of y^c .

This shows that T_0 is the subring of A generated by x, y, z . Under the embedding in Lemma 4.3.1, we can identify $x = u^{n+1}$, $y = v^{n+1}$, $z = uv$, and therefore $T_0 = A^G = B$. As R is a domain and t_0 clearly q^\bullet -commutes with x, y , the subring $\mathbb{k}\langle x, y, t_0 \rangle$ of T is isomorphic to a quantum 3-space and hence $\text{GKdim } T \geq 3$.

Now suppose that $m \in T$ is a monomial of projective degree $d \geq 1$. Write

$$m = a_0^{i_0} a_1^{i_1} \dots a_n^{i_n} \bar{a}_0^{j_0} \bar{a}_1^{j_1} \dots \bar{a}_n^{j_n},$$

for some $i_k, j_k \geq 0$. By factoring out a power of $z = a_k \bar{a}_k$, up to rescaling m we may assume that $i_k = 0$ or $j_k = 0$ for $0 \leq k \leq n$.

First assume that at least one i_k is nonzero, and fix

$$k := \max\{\ell \mid i_\ell \neq 0\}.$$

Therefore $i_{k+1} = \dots = i_n = 0$, $i_k \geq 1$, and $j_k = 0$. The k th coordinate of the equation

$$\text{wt}(m) = d(-n, 1, 1, \dots, 1), \tag{5.1.3}$$

(where here we 0-index this vector), tells us that

$$d = i_{k-1} - j_{k-1} - i_k,$$

where $i_{k-1} = 0$ or $j_{k-1} = 0$. If $i_{k-1} = 0$ then we obtain

$$d = -j_{k-1} - i_k \leq 0,$$

which is absurd, so we must have $j_{k-1} = 0$. It then follows that

$$i_{k-1} = d + i_k \geq 1 + 1 = 2.$$

Repeating this argument along the entries of (5.1.3) shows that

$$i_k \geq 1, \quad i_{k-1} \geq 2, \quad i_{k-2} \geq 3, \quad \dots, \quad i_0 \geq k + 1.$$

Similarly, if we look at the $(k + 2)$ nd coordinate of (5.1.3), we obtain the equation

$$-j_{k+1} + j_{k+2} = d.$$

If $j_{k+2} = 0$ then this equation cannot be satisfied, so we must have $j_{k+2} \geq 1$. An argument similar to the one used for the i_ℓ shows that

$$j_{k+2} \geq 1, \quad j_{k+3} \geq 2, \quad \dots, \quad j_n \geq n - k - 1.$$

This shows that $a_0^{k+1} a_1^k \dots a_{k-1}^2 a_k \bar{a}_{k+2} \bar{a}_{k+3}^2 \dots \bar{a}_{n-1}^{n-k-2} \bar{a}_n^{n-k-1} = t_{n-k-1}$ is a factor of m , so we can write $m = \lambda t_{n-k-1} a'$ for some $\lambda \in \mathbb{k}^\times$ and a monomial $a' \in R$. If instead all of the i_ℓ are 0, a similar argument shows that $m = \lambda t_n a'$.

By the above, we have $m = \lambda t_\ell a'$, where $a' \in R$ is a monomial and $\lambda \in \mathbb{k}^\times$. We have $\text{wt}(a') = \text{wt}(m) - \text{wt}(t_\ell) = (d-1)\chi$, so $a' \in T_{d-1} = R_{(d-1)\chi}$. The claim follows by induction. \square

5.2. A presentation for T . We now seek to give a presentation for T . Define

$$\mathbb{k}_{\mathbf{q}}[x, y, z, t_0, \dots, t_n] = \frac{\mathbb{k}\langle x, y, z, t_0, \dots, t_n \rangle}{\left\langle \begin{array}{lll} xy = q^{(n+1)^2} xy & zx = q^{n+1} xz & zy = q^{-(n+1)} yz \\ t_i x = q^{(n+1)\binom{i+1}{2}} x t_i & t_i y = q^{-(n+1)\binom{n-i+1}{2}} y t_i & t_i z = q^{\frac{1}{2}(n+1)(2i-n)} z t_i \\ t_j t_i = q^{\frac{1}{4}(n+1)(i-j)(2ij-n(i+j+1))} t_i t_j \end{array} \right\rangle}, \quad (5.2.1)$$

which is a skew polynomial ring, and hence AS regular [AS87]. We note that the exponent

$$\frac{1}{4}(n+1)(i-j)(2ij-n(i+j+1))$$

is always an integer, so the last relation is well-defined. Also set

$$T_{\text{pres}} := \frac{\mathbb{k}_{\mathbf{q}}[x, y, z, t_0, \dots, t_n]}{\left\langle \begin{array}{ll} xy = q^{-\binom{n+1}{2}} z^{n+1} & \\ xt_i = q^{\alpha_i} z^i t_{i-1} & \text{for } 1 \leq i \leq n \\ yt_i = q^{\beta_i} z^{n-i} t_{i+1} & \text{for } 0 \leq i \leq n-1 \\ t_{i-1} t_{j+1} = q^{\gamma_{ij}} z^{j-i+1} t_i t_j & \text{for } 1 \leq i \leq j \leq n-1 \end{array} \right\rangle},$$

where

$$\alpha_i = -\binom{i}{2} - i\binom{n-i+2}{2}, \quad \beta_i = (i+2)\binom{n-i+1}{2},$$

$$\gamma_{ij} = -(j+1)\binom{n-j+1}{2} - (n-j)\binom{i}{2} - (j-i+1)\binom{n-i+1}{2} - \binom{j-i+1}{2}.$$

As suggested by our choice of notation, we claim that T and T_{pres} are isomorphic. As a first step, we show that the generators of T satisfy the relations in T_{pres} :

Lemma 5.2.2. *The elements $x, y, z, t_i \in T$ satisfy the relations in T_{pres} .*

Proof. Let $x, y, z, t_0, \dots, t_n \in T$ be as in Proposition 5.1.1. We begin the proof by showing that these elements satisfy the skew-commutativity relations given in the presentation for $\mathbb{k}_{\mathbf{q}}[x, y, z, t_0, \dots, t_n]$. It is clear that any pair of generators q^\bullet -commute, for some power q^\bullet of q .

To determine the precise power of q , we will view the generators as elements of $A[s_0^{\pm 1}, \dots, s_n^{\pm 1}]$ via Lemma 4.3.1. If $a \in A[s_0^{\pm 1}, \dots, s_n^{\pm 1}]$, write $a \approx f(u, v)$ for the image of a in

$$A[s_0^{\pm 1}, \dots, s_n^{\pm 1}]/\langle s_i - 1 \mid 0 \leq i \leq n \rangle \cong A.$$

It suffices to work in this quotient ring to determine the power of q in the q^\bullet -commutativity relation between two generators.

With this notation, we have

$$x \approx u^{n+1}, \quad y \approx v^{n+1}, \quad z \approx uv, \quad t_i \approx u^{\binom{n-i+1}{2}} v^{\binom{i+1}{2}}.$$

Direct calculation now gives

$$\begin{aligned} yx &\approx v^{n+1} u^{n+1} = q^{(n+1)^2} u^{n+1} v^{n+1} \approx q^{(n+1)^2} xy, \\ zx &\approx uv \cdot u^{n+1} = q^{n+1} u^{n+1} \cdot uv \approx q^{n+1} xz, \\ zy &\approx uv \cdot v^{n+1} = q^{-(n+1)} v^{n+1} \cdot uv \approx q^{-(n+1)} yz. \end{aligned}$$

For the commutativity relations involving t_i and one of x, y , or z , we have

$$\begin{aligned} t_i x &\approx u^{\binom{n-i+1}{2}} v^{\binom{i+1}{2}} u^{n+1} = q^{(n+1)\binom{i+1}{2}} u^{n+1} u^{\binom{n-i+1}{2}} v^{\binom{i+1}{2}} \approx q^{(n+1)\binom{i+1}{2}} x t_i, \\ t_i y &\approx u^{\binom{n-i+1}{2}} v^{\binom{i+1}{2}} v^{n+1} = q^{-(n+1)\binom{n-i+1}{2}} v^{n+1} u^{\binom{n-i+1}{2}} v^{\binom{i+1}{2}} \approx q^{-(n+1)\binom{n-i+1}{2}} y t_i \\ t_i z &\approx u^{\binom{n-i+1}{2}} v^{\binom{i+1}{2}} uv = q^{\binom{i+1}{2} - \binom{n-i+1}{2}} uv u^{\binom{n-i+1}{2}} v^{\binom{i+1}{2}} = q^{\frac{1}{2}(n+1)(2i-n)} z t_i. \end{aligned}$$

The only remaining commutativity relation left to determine is that relating t_i and t_j :

$$\begin{aligned} t_j t_i &\approx u^{\binom{n-j+1}{2}} v^{\binom{j+1}{2}} u^{\binom{n-i+1}{2}} v^{\binom{i+1}{2}} \\ &= q^{\binom{j+1}{2}\binom{n-i+1}{2}} u^{\binom{n-j+1}{2}} u^{\binom{n-i+1}{2}} v^{\binom{i+1}{2}} v^{\binom{j+1}{2}} \\ &= q^{\binom{j+1}{2}\binom{n-i+1}{2} - \binom{i+1}{2}\binom{n-j+1}{2}} u^{\binom{n-j+1}{2}} v^{\binom{i+1}{2}} u^{\binom{n-i+1}{2}} v^{\binom{j+1}{2}} \\ &= q^{\frac{1}{4}(n+1)(i-j)(2ij-n(i+j+1))} t_i t_j \end{aligned}$$

where the equality in the exponent of q is an easy computation.

We now show that the generators satisfy the relations by which we factor $\mathbb{k}_{\mathbf{q}}[x, y, z, t_0, \dots, t_n]$ to obtain T_{pres} . The first three are relatively straightforward. First,

$$xy \approx u^{n+1} v^{n+1} = q^{-\binom{n+1}{2}} (uv)^{n+1} \approx q^{-\binom{n+1}{2}} z^{n+1}.$$

Since $xy, z^{n+1} \in T_0 \subseteq A$, this suffices to show that $xy = q^{-\binom{n+1}{2}} z^{n+1}$.

Likewise,

$$\begin{aligned} x t_i &\approx u^{n+1} \cdot u^{\binom{n-i+1}{2}} v^{\binom{i+1}{2}} \\ &= u^i \cdot u^{n-i+1} u^{\binom{n-i+1}{2}} v^{i+\binom{i}{2}} \\ &= q^{-i\binom{n-i+2}{2}} u^i v^i u^{\binom{n-i+2}{2}} v^{\binom{i}{2}} \\ &\approx q^{-\binom{i}{2} - i\binom{n-i+2}{2}} z^i t_{i-1}. \end{aligned}$$

Similar to the above, $x t_i, z^i t_{i-1} \in A s_0^{-n} s_1 \dots s_n$ and so the above suffices to show that the relation holds. Finally,

$$\begin{aligned} y t_i &\approx v^{n+1} \cdot u^{\binom{n-i+1}{2}} v^{\binom{i+1}{2}} \\ &= q^{(i+1)\binom{n-i+1}{2}} v^{n-i} u^{n-i+\binom{n-i}{2}} v^{i+1+\binom{i+1}{2}} \\ &= q^{(i+1)\binom{n-i+1}{2} + \binom{n-i+1}{2}} (uv)^{n-i} u^{\binom{n-i}{2}} v^{\binom{i+2}{2}} \\ &\approx q^{(i+2)\binom{n-i+1}{2}} z^{n-i} t_{i+1}, \end{aligned}$$

and again this is sufficient.

The final relation requires a little more care. Again, it suffices to check the relations are satisfied up to \approx . If $1 \leq i \leq j \leq n-1$, then

$$\begin{aligned}
t_{i-1}t_{j+1} &\approx u^{\binom{n-i+2}{2}}v^{\binom{i}{2}}u^{\binom{n-j}{2}}v^{\binom{j+2}{2}} \\
&= u^{n-i+1+\binom{n-i+1}{2}}v^{\binom{i}{2}}u^{\binom{n-j}{2}}v^{j+1+\binom{j+1}{2}} \\
&= q^{-(j+1)\binom{n-j}{2}}u^{j-i+1}u^{n-j}u^{\binom{n-i+1}{2}}v^{j+1}v^{\binom{i}{2}}u^{\binom{n-j}{2}}v^{\binom{j+1}{2}} \\
&= q^{-(j+1)\binom{n-j}{2}-(n-j)\binom{(j+1)+\binom{i}{2}}{2}}u^{j-i+1}u^{\binom{n-i+1}{2}}v^{j+1}v^{\binom{i}{2}}u^{n-j+\binom{n-j}{2}}v^{\binom{j+1}{2}} \\
&= q^{-(j+1)\binom{n-j+1}{2}-(n-j)\binom{i}{2}-(j-i+1)\binom{n-i+1}{2}}u^{j-i+1}v^{j-i+1}u^{\binom{n-i+1}{2}}v^{i+\binom{i}{2}}u^{n-j+\binom{n-j}{2}}v^{\binom{j+1}{2}} \\
&= q^{-(j+1)\binom{n-j+1}{2}-(n-j)\binom{i}{2}-(j-i+1)\binom{n-i+1}{2}-(\binom{j-i+1}{2})}z^{j-i+1}t_it_j.
\end{aligned}$$

Thus the generators of T satisfy all the relations in T_{pres} . \square

By Lemma 5.2.2, there is a surjection $T_{\text{pres}} \twoheadrightarrow T$. To show that there are no more relations between the elements x, y, z, t_i other than those given in T_{pres} , we first make the following useful observation.

Proposition 5.2.3. $R = R^{(n)}$ has a \mathbb{Z}^3 -grading given by

$$\begin{aligned}
\deg(a_i) &= (\delta_{i,n}, 0, \delta_{i,n-1} - \delta_{i,n}) \\
\deg(\bar{a}_i) &= (1 - \delta_{i,n}, 1, -\delta_{i,n-1} + \delta_{i,n}),
\end{aligned}$$

where $\delta_{i,j}$ is the Kronecker delta. Under this grading T is \mathbb{N}^3 -graded with

$$\begin{aligned}
\deg x &= (1, 0, 0), \quad \deg y = (n, n+1, 0), \quad \deg z = (1, 1, 0), \\
\deg t_i &= \left(\binom{i}{2}, \binom{i+1}{2}, 1 \right), \quad 0 \leq i \leq n.
\end{aligned}$$

Proof. The relations of R are homogeneous with respect to this grading on the generators, giving the first statement. The second statement is a consequence of Proposition 5.1.1. \square

We often refer to the grading on T given in Proposition 5.2.3 as the *toric grading* of T . Since the relations in T_{pres} are a subset of those of T (and ultimately we will show they satisfy precisely the same relations), this also gives an \mathbb{N}^3 -grading on T_{pres} which we also call the toric grading. One reason that this grading is useful is because the graded pieces of T_{pres} are at most one-dimensional.

Proposition 5.2.4. The graded pieces of T_{pres} in the toric grading satisfy $\dim(T_{\text{pres}})_{\vec{n}} \leq 1$ for all $\vec{n} \in \mathbb{N}^3$.

Proof. Let

$$\mathcal{S} = \{x^i z^k \mid i, k \geq 0\} \cup \{y^j z^k \mid j \geq 1, k \geq 0\} \quad (5.2.5)$$

$$\cup \{x^i z^k t_0^{\ell_0} \mid i, k, \ell_0 \geq 0\} \cup \{y^j z^k t_n^{\ell_n} \mid j, k, \ell_n \geq 0\} \quad (5.2.6)$$

$$\cup \bigcup_{m=1}^{n-1} \{z^k t_m^{\ell_m} \mid k \geq 0, \ell_m \geq 1\} \cup \bigcup_{m=0}^{n-1} \{z^k t_m^{\ell_m} t_{m+1}^{\ell_{m+1}} \mid k \geq 0, \ell_m, \ell_{m+1} \geq 1\}. \quad (5.2.7)$$

We first show that \mathcal{S} spans T_{pres} . By applying the relations in $\mathbb{k}_{\mathbf{q}}[x, y, z, t_0, \dots, t_n]$, a monomial in T_{pres} can be written as $x^i y^j z^k t_0^{\ell_0} \dots t_n^{\ell_n}$, up to multiplication by a scalar, where all the exponents are non-negative. Moreover, repeated application of the relation $t_{i-1}t_{j+1} = q^{\bullet} z^{j-i+1} t_i t_j$ to the flanking t_i terms in this monomial shows that, up to scaling, we can write it as

$$x^i y^j z^k t_m^{\ell_m} t_{m+1}^{\ell_{m+1}}$$

for some $0 \leq m \leq n-1$ and exponents i, j, k, ℓ , some of which are possibly 0. Additionally, the relation $xy = q^{\bullet} z^{n+1}$ allows us to assume $i = 0$ or $j = 0$.

If $i = 0 = j$ then this element is in \mathcal{S} , so assume that $j = 0$ and $i > 0$; the case $i = 0$ and $j > 0$ is similar. We prove by induction on i that the monomial $x^i z^k t_m^{\ell_m} t_{m+1}^{\ell_{m+1}}$ lies in $\text{span } \mathcal{S}$. If $\ell_{m+1} \geq 1$, then the relation $xt_{m+1} = q^{\alpha_{m+1}} z^{m+1} t_m$ gives

$$x^i z^k t_m^{\ell_m} t_{m+1}^{\ell_{m+1}} = q^\bullet x^{i-1} z^{k'} t_m^{\ell_m+1} t_{m+1}^{\ell_{m+1}-1}.$$

By induction, this element lies in $\text{span } \mathcal{S}$. Now suppose $\ell_{m+1} = 0$; we consider two subcases. If $m = 0$, or if $m \neq 0$ and $\ell_m = 0$, then this monomial is already an element of \mathcal{S} . Otherwise $m \neq 0$ and $\ell_m \geq 1$, and we have

$$x^i z^k t_m^{\ell_m} = q^\bullet x^{i-1} z^{k'} t_{m-1} t_m^{\ell_m-1},$$

and this element lies in $\text{span } \mathcal{S}$ by induction.

We now show that distinct elements of \mathcal{S} have different toric degrees. By considering the third coordinate, it is clear that the sets in (5.2.5) are disjoint from the other sets. Moreover,

$$\deg x^i z^k = (i + k, k, 0), \quad \deg y^j z^{k'} = (jn + k', j(n + 1) + k', 0),$$

and these are distinct for $i, k, k' \geq 0$ and $j \geq 1$.

It remains to show that the toric degrees of elements in the remaining four sets in (5.2.6) and (5.2.7) are distinct. Call these four sets (a), (b), (c), and (d), in the order given; in principle we need to show that two distinct elements chosen from these four sets (possibly chosen from the same set) have distinct degrees. Most of these are routine, so we give only some of the proofs.

For example, consider elements $x^i z^k t_0^\ell$ and $y^{j'} z^{k'} t_n^{\ell'}$ from sets (a) and (b), which have the same weight. Equating weights gives three equations:

$$i + k = j'n + k' + \ell' \binom{n}{2}, \quad k = j'(n + 1) + k' + \ell' \binom{n+1}{2}, \quad \ell = \ell'.$$

Subtracting the second equation from the first and using the third equation gives $i = j' - \ell n$, where the left hand side is non-negative and the right hand side is negative, a contradiction.

As another example, consider elements $z^k t_m^{\ell_m}$ and $z^{k'} t_{m'}^{\ell_{m'}} t_{m'+1}^{\ell_{m'+1}}$ from sets (c) and (d). Equating degrees gives three equations

$$\begin{aligned} k + \ell_m \binom{m}{2} &= k' + \ell_{m'} \binom{m'}{2} + \ell_{m'+1} \binom{m'+1}{2}, \\ k + \ell_m \binom{m+1}{2} &= k' + \ell_{m'} \binom{m'+1}{2} + \ell_{m'+1} \binom{m'+2}{2} \\ \ell_m &= \ell_{m'} + \ell_{m'+1}. \end{aligned}$$

Subtracting the first equation from the second gives

$$\ell_m m = \ell_{m'} m' + \ell_{m'+1} (m' + 1) = (\ell_{m'} + \ell_{m'+1}) m' + \ell_{m'+1} = \ell_m m' + \ell_{m'+1}.$$

Therefore

$$\ell_m (m - m') = \ell_{m'+1}.$$

Now, if $m' \geq m$ then $\ell_{m'+1} \leq 0$, which is a contradiction, so we must have $m > m'$. But then

$$\ell_{m'+1} = \ell_m (m - m') \geq \ell_m = \ell_{m'} + \ell_{m'+1} > \ell_{m'+1},$$

which is also a contradiction. The remaining cases are similar and left to the reader. \square

With this result in hand, we are now able to show that T_{pres} gives a presentation of T :

Theorem 5.2.8. *There is an isomorphism $T \cong T_{\text{pres}}$ of \mathbb{N}^3 -graded rings. Moreover, these rings are domains of GK-dimension 3.*

Proof. We first note that, as T is a subring of the domain R , it is also a domain. Now, temporarily denote the generators of T by $\mathbf{x}, \mathbf{y}, \mathbf{z}, \mathbf{t}_0, \dots, \mathbf{t}_n$. Let $\nu : T_{\text{pres}} \rightarrow T$ be the obvious “boldface” map sending $x \mapsto \mathbf{x}$, etc., which exists by Lemma 5.2.2.

Since we know that ν is surjective by Proposition 5.1.1, to show that it is an isomorphism it remains to show that it is injective. It suffices to show that its restriction to each (toric) graded piece $(T_{\text{pres}})_{\vec{n}}$ is injective; since each of these are at most one-dimensional, we only need to show that the restriction is not the zero map when $\dim(T_{\text{pres}})_{\vec{n}} = 1$. Indeed, if $s \in \mathcal{S}$ then $\nu(s)$ is nonzero, since it is a monomial in the (nonzero) generators of the domain T . Thus ν is injective, and hence an isomorphism.

We will now make the identification $T = T_{\text{pres}}$; in particular, we will drop the bold notations \mathbf{x} , etc., for generators of T . It remains to show that $\text{GKdim } T = 3$.

Let $\Sigma = \{\vec{n} \mid T_{\vec{n}} \neq 0\}$ be the sub-semigroup of \mathbb{N}^3 on which T is supported. If v_1, \dots, v_k are nonzero \mathbb{N}^3 -homogeneous elements of T and $V = \mathbb{k} \cdot (v_1, \dots, v_k)$, we write

$$\Sigma_V = \{\deg(v_1), \dots, \deg(v_k)\} \subseteq \Sigma.$$

Since $\vec{n} \in \Sigma$ if and only if $\dim T_{\vec{n}} = 1$, we see that $\dim V = \#\Sigma_V$.

Now let W be an arbitrary finite-dimensional subspace of T , and let V be a finite-dimensional \mathbb{N}^3 -graded subspace containing W . Then $\dim W^n \leq \dim V^n = \#(\Sigma_V)^n$, and so $\text{GKdim } T$ is bounded above by the growth of Σ . As Σ is a sub-semigroup of \mathbb{N}^3 , which has cubic growth, $\text{GKdim } T \leq 3$. But by Proposition 5.1.1, $\text{GKdim } T \geq 3$. Thus $\text{GKdim } T = 3$. \square

5.3. The toric embedding of T . We now construct a useful embedding of T into a quantum 3-torus which is compatible with the toric grading. Let

$$D = \frac{\mathbb{k}\langle \mathbf{a}^{\pm 1}, \mathbf{b}^{\pm 1}, \mathbf{s}^{\pm 1} \rangle}{\left\langle \begin{array}{l} \mathbf{ba} = q^{n+1} \mathbf{ab} \\ \mathbf{sa} = \mathbf{as} \\ \mathbf{sb} = q^{-\binom{n+1}{2}} \mathbf{bs} \end{array} \right\rangle} \quad (5.3.1)$$

This has a \mathbb{Z}^3 -grading given by declaring

$$\deg \mathbf{a} = (1, 0, 0), \quad \deg \mathbf{b} = (0, 1, 0), \quad \deg \mathbf{s} = (0, 0, 1).$$

Let $Q_{\text{gr}}^{\mathbb{Z}^3}(T)$ be the \mathbb{Z}^3 -graded quotient ring of T obtained by inverting all \mathbb{Z}^3 -homogeneous elements.

We claim that $Q_{\text{gr}}^{\mathbb{Z}^3}(T) \cong D$ as graded rings; we will use this to construct an embedding $T \hookrightarrow D$. We first require a lemma:

Lemma 5.3.2. *We have $\dim Q_{\text{gr}}^{\mathbb{Z}^3}(T)_{\vec{n}} = 1$ for all $\vec{n} \in \mathbb{Z}^3$.*

Proof. Let $D' = Q_{\text{gr}}^{\mathbb{Z}^3}(T)$. We first show that $\dim D'_{\vec{n}} \leq 1$. Suppose that $u, v \in D'_{\vec{n}}$ are nonzero. By multiplying by a suitable homogeneous element $d \in T_{\vec{m}}$ (essentially clearing denominators), we obtain $ud, vd \in T_{\vec{n}+\vec{m}}$. Since $T_{\vec{n}+\vec{m}}$ is one-dimensional by Proposition 5.2.4, it follows that $ud = \lambda vd$ for some $\lambda \in \mathbb{k}^\times$. Multiplying by d^{-1} shows that $u = \lambda v$, so that $\dim D'_{\vec{n}} \leq 1$. Since D' contains the elements x, t_0, yz^{-n} whose respective degrees are $(1, 0, 0)$, $(0, 0, 1)$, and $(0, 1, 0)$, the reverse inequality also holds. \square

Now define a map

$$\begin{aligned} \bar{\theta} : \mathbb{k}\langle \mathbf{a}, \mathbf{b}, \mathbf{s} \rangle &\rightarrow Q_{\text{gr}}^{\mathbb{Z}^3}(T), \\ \mathbf{a} &\mapsto x, \quad \mathbf{b} \mapsto yz^{-n}, \quad \mathbf{s} \mapsto t_0. \end{aligned}$$

The following calculations shows that this descends to a map $\theta : D \rightarrow Q_{\text{gr}}^{\mathbb{Z}^3}(T)$:

$$\begin{aligned} \bar{\theta}(\mathbf{ba}) &= yz^{-n}x = yq^{-(n+1)}xz^{-n} = q^{(n+1)^2-n(n+1)}xyz^{-n} = q^{n+1}\theta(\mathbf{ab}), \\ \bar{\theta}(\mathbf{sa}) &= t_0x = xt_0 = \bar{\theta}(\mathbf{as}), \\ \bar{\theta}(\mathbf{sb}) &= t_0yz^{-n} = q^{-\frac{1}{2}n(n+1)^2}yt_0z^{-n} = q^{-\frac{1}{2}n(n+1)^2+\frac{1}{2}n^2(n+1)}yz^{-n}t_0 = q^{-\frac{1}{2}n(n+1)}\bar{\theta}(\mathbf{bs}). \end{aligned}$$

Moreover, it is straightforward to check that $\theta : D \rightarrow Q_{\text{gr}}^{\mathbb{Z}^3}(T)$ is a graded map.

Proposition 5.3.3. *The map*

$$\theta : D \rightarrow Q_{\text{gr}}^{\mathbb{Z}^3}(T), \quad \mathbf{a} \mapsto x, \quad \mathbf{b} \mapsto yz^{-n}, \quad \mathbf{s} \mapsto t_0,$$

is an isomorphism of \mathbb{Z}^3 -graded rings.

Proof. We first show that θ surjects. Indeed, we have

$$\begin{aligned} \theta(\mathbf{a}) &= x, \\ \theta(\mathbf{a}^n \mathbf{b}^{n+1}) &= x^n (yz^{-n})^{n+1} = q^\bullet y (xyz^{-(n+1)})^n = q^\bullet y \\ \theta(\mathbf{ab}) &= xyz^{-n} = q^\bullet z (xyz^{-(n+1)}) = q^\bullet z \\ \theta(\mathbf{a}^{(i)} \mathbf{b}^{(i+1)} \mathbf{s}) &= x^{(i)} (yz^{-n})^{(i+1)} t_0 = q^\bullet (xy)^{(i)} z^{-n} y^{(i+1)} t_0 \\ &= q^\bullet z^{(n+1)} z^{-n} z^{in-(i)} t_i = q^\bullet t_i. \end{aligned} \tag{5.3.4}$$

Since D and $Q_{\text{gr}}^{\mathbb{Z}^3}(T)$ are 1-dimensional in each graded piece (the latter by Lemma 5.3.2), it follows that θ is in fact an isomorphism. \square

Now define $\eta : T \rightarrow D$ to be the map $\eta = (\theta^{-1})|_T$, which is necessarily an embedding of T into D . From (5.3.4), we have

$$\eta(x) = \mathbf{a}, \quad \eta(y) = q^\bullet \mathbf{a}^n \mathbf{b}^{n+1}, \quad \eta(z) = q^\bullet \mathbf{ab}, \quad \eta(t_i) = q^\bullet \mathbf{a}^{(i)} \mathbf{b}^{(i+1)} \mathbf{s}. \tag{5.3.5}$$

The precise powers of q in the above are not important for our needs, but can be calculated with a bit of care. We call the map $\eta : T \rightarrow D$ the *toric embedding* of T , and we write

$$T_{\text{toric}} := \mathbb{k}\langle \mathbf{a}, \mathbf{a}^n \mathbf{b}^{n+1}, \mathbf{ab}, \mathbf{a}^{(i)} \mathbf{b}^{(i+1)} \mathbf{s} \mid 0 \leq i \leq n \rangle$$

for the image of η . These will play a major role in later calculations.

6. A GEOMETRIC RESOLUTION: THE CATEGORY

We will now use the ring T and methods of noncommutative algebraic geometry to construct a category \mathcal{X}_q which we will see is also a resolution of B_q .

In this section we consider T with its projective grading; that is, T is \mathbb{N} -graded with $T_d = R_{dX}$. We will define a category $\text{qgr-}T$, a suitable localisation of the graded module category of T , and give many of its properties. Much of this is standard noncommutative geometry as in [AZ94], but we need to give some proofs as the standing assumption of [AZ94] is that the degree 0 part of the ring is a finite module over a *commutative* ground ring.

We begin, however, by recalling the basics of local cohomology.

6.1. Local cohomology. In this paper we will be primarily interested in local cohomology of graded T -modules with respect to $T_+ = \bigoplus_{s \geq 1} T_s$. However, we recall the basic definitions more generally. Thus, in this section let Ω be a ring, I an ideal of Ω , and M an Ω -module. Recall that the j th local cohomology of M with respect to I is

$$H_I^j(M) = \lim_{s \rightarrow \infty} \text{Ext}_{\Omega}^j(\Omega/I^s, M).$$

Equivalently, the $H_I^j(-)$ are the right derived functors of $H_I^0(-) = \lim_{s \rightarrow \infty} \text{Hom}_{\Omega}(\Omega/I^s, -)$. Since Ω/I^s is an Ω -bimodule, $H_I^j(M)$ is an Ω -module. Note that $H_I^0(M)$ is a submodule of M ; we refer to it as the *I -torsion submodule* of M .

Lemma 6.1.1. *Local cohomology modules are I -torsion. That is, for all $j \in \mathbb{N}$ and for any Ω -module M , we have*

$$H_I^0(H_I^j(M)) = H_I^j(M).$$

Proof. As $I^s(\Omega/I^s) = 0$, we have $\text{Hom}_{\Omega}(\Omega/I^s, M)I^s = 0$, and thus $H_I^0(H_I^0(M)) = H_I^0(M)$. The general result follows by dimension shifting. \square

If Ω is commutative then it is well-known that local cohomology may be computed via Čech cohomology. The same is true if Ω is noncommutative and I is normally generated. Indeed, suppose now that $I = \langle x_0, \dots, x_n \rangle$, where x_0, \dots, x_n are normal elements of Ω and let M be a right Ω -module. The Čech complex $\check{C}(x_0, \dots, x_n; M)$ of M with respect to I is defined as follows, where we use the (slightly nonstandard) notation that $[n] = \{0, \dots, n\}$:

$$\check{C}(x_0, \dots, x_n; M) : \quad 0 \rightarrow M \xrightarrow{d_0} \bigoplus_{i=0}^n M[x_i^{-1}] \xrightarrow{d_1} \dots \xrightarrow{d_{s-1}} \bigoplus_{\substack{J \subseteq [n] \\ |J|=s}} M[x_J^{-1}] \xrightarrow{d_s} \dots \xrightarrow{d_n} M[x_{[n]}^{-1}] \rightarrow 0.$$

Here, we define

$$M[x_J^{-1}] = M[x_j^{-1} \mid j \in J].$$

The differential is defined on $m_J \in M[x_J^{-1}]$ by

$$d_{|J|}(m_J) = \sum_{k \notin J} (-1)^{o_J(k)} m_{J \cup \{k\}},$$

where $o_J(k)$ denotes the number of elements of J which are less than k , and $m_{J \cup \{k\}}$ is the image of $m_J \in M[x_J^{-1}]$ in $M[x_{J \cup \{k\}}^{-1}] = M[x_J^{-1}][x_k^{-1}]$.

Proposition 6.1.2. *If x_0, \dots, x_n are normal elements of Ω and $I = \langle x_0, \dots, x_n \rangle$, then the cohomology of the Čech complex computes the local cohomology of M with respect to I : that is,*

$$H^j(\check{C}(x_0, \dots, x_n; M)) \cong H_I^j(M)$$

for all $j \geq 0$.

Proof. The proof given in [BS13, Theorem 5.1.19] for commutative rings also works in our situation. \square

6.2. Defining the category \mathcal{X}_q . We now consider T with its projective grading. Let M be a graded (right) T -module. The *torsion* submodule $\tau(M)$ of M is the T_+ -torsion submodule; that is, $\tau(M) = H_{T_+}^0(M)$. If $M = \tau(M)$ then M is *torsion*. As usual, let $\text{Tors-}T$ be the full subcategory of $\text{Gr-}T$ consisting of torsion modules, and let $\text{Qgr-}T = \text{Gr-}T / \text{Tors-}T$. Let $\pi : \text{Gr-}T \rightarrow \text{Qgr-}T$ be the quotient functor.

More explicitly, the objects of $\text{Qgr-}T$ are the same as the objects of $\text{Gr-}T$, and the maps are defined by

$$\text{Hom}_{\text{Qgr-}T}(\pi M, \pi N) = \varinjlim_{\substack{M' \subseteq M \\ M/M' \text{ is torsion}}} \text{Hom}_{\text{Gr-}T}(M', N/\tau(N)).$$

If M is finitely generated, then

$$\text{Hom}_{\text{Qgr-}T}(\pi M, \pi N) = \varinjlim_s \text{Hom}_{\text{Gr-}T}(M_{\geq s}, N).$$

By definition, the elements of $\text{Hom}_{\text{Gr-}T}(M, N)$ are degree-preserving, and thus the same is true for elements of $\text{Hom}_{\text{Qgr-}T}(\mathcal{M}, \mathcal{N})$ for $\mathcal{M}, \mathcal{N} \in \text{Qgr-}T$. We will also sometimes need

$$\underline{\text{Hom}}_{\text{Qgr-}T}(\mathcal{M}, \mathcal{N}) = \bigoplus_{s \in \mathbb{Z}} \text{Hom}_{\text{Qgr-}T}(\mathcal{M}, \mathcal{N}(s)).$$

Recall our convention that if Xyz is an abelian category, then xyz is the full subcategory of noetherian objects. Since T is noetherian, $\text{qgr-}T \simeq \text{gr-}T / \text{tors-}T$.

By construction, the functor $\pi : \text{Gr-}T \rightarrow \text{Qgr-}T$ is exact and preserves colimits. By the adjoint functor theorem, therefore, π has a right adjoint, which we denote ω . If $\mathcal{N} \in \text{Qgr-}T$, $s \in \mathbb{Z}$ then

$$(\omega \mathcal{N})_s = \text{Hom}_{\text{Gr-}T}(T, \omega \mathcal{N}(s)) = \text{Hom}_{\text{Qgr-}T}(\pi T, \mathcal{N}(s))$$

and so

$$\omega \mathcal{N} = \underline{\text{Hom}}_{\text{Qgr-}T}(\pi T, \mathcal{N}).$$

If $\mathcal{N} \in \text{Qgr-}T$ then the (B, T) -bimodule structure on T coming from the inclusion $B = T_0 \subseteq T$ gives $\text{Hom}_{\text{Qgr-}T}(\pi T, \mathcal{N})$ the structure of a B -module. We denote the functor

$$H_{\text{Qgr-}T}^0(-) := \text{Hom}_{\text{Qgr-}T}(\pi T, -) : \text{Qgr-}T \rightarrow \text{Mod-}B$$

by ϕ_* . As this functor is left exact and preserves limits, the adjoint functor theorem guarantees that it has a left adjoint ϕ^* . We thus have a map of noncommutative spaces $\phi : \text{Qgr-}T \rightarrow \text{Mod-}B$ given by

$$\begin{array}{c} \text{Qgr-}T \\ \uparrow \phi^* \quad \downarrow \phi_* \\ \text{Mod-}B \end{array}$$

We now show that $\phi^* = \pi(- \otimes_B T)$.

Proposition 6.2.1. *Let $\phi_* = H_{\text{Qgr-}T}^0(-)$ as defined above and let $\phi^* = \pi(- \otimes_B T)$. Then (ϕ^*, ϕ_*) are an adjoint pair.*

Proof. Let $M \in \text{Mod-}B$, $\mathcal{N} \in \text{Qgr-}T$. We have

$$\begin{aligned} \text{Hom}_{\text{Qgr-}T}(\phi^* M, \mathcal{N}) &= \text{Hom}_{\text{Qgr-}T}(\pi(M \otimes_B T), \mathcal{N}) && \text{by definition} \\ &\cong \text{Hom}_{\text{Gr-}T}(M \otimes_B T, \omega \mathcal{N}) && \text{as } (\pi, \omega) \text{ are an adjoint pair} \\ &= \underline{\text{Hom}}_T(M \otimes_B T, \omega \mathcal{N})_0 \\ &\cong \text{Hom}_B(M, (\omega \mathcal{N})_0) && \text{by hom-tensor adjointness.} \end{aligned}$$

Now, $(\omega \mathcal{N})_0 = \text{Hom}_{\text{Gr-}T}(T, \omega \mathcal{N}) \cong \text{Hom}_{\text{Qgr-}T}(\pi T, \mathcal{N})$, using adjointness of (π, ω) again. But this last group is $\phi_* \mathcal{N}$ by definition, proving that

$$\text{Hom}_{\text{Qgr-}T}(\phi^* M, \mathcal{N}) \cong \text{Hom}_B(M, \phi_* \mathcal{N}).$$

The isomorphisms used in the proof are all natural in \mathcal{N} and (where relevant) M . \square

It is clear from the definition that $\phi^*(\text{mod-}B) \subseteq \text{qgr-}T$. We will see later that ϕ is “proper”: that is, that $\phi_*(\text{qgr-}T) \subseteq \text{mod-}B$, and that the map $\phi = (\phi^*, \phi_*) : \text{qgr-}T \rightarrow \text{mod-}B$ is a resolution of the singularity of B .

Similarly to Lemma 3.2.6, we now show that the map $\phi : \text{Qgr-}T \rightarrow \text{Mod-}B$ is an isomorphism above the “nonsingular locus” of B .

Proposition 6.2.2. *Let $w \in \{x, y, z\}$. Then ϕ^*, ϕ_* restrict to inverse equivalences between $\text{Qgr-}T[w^{-1}]$ and $\text{Mod-}B[w^{-1}]$.*

Proof. We give the proof for $w = x$; the other cases are similar. In $T[x^{-1}]$ we have $t_i = q^\bullet x^{-1} z^i t_{i-1}$ for all $i \geq 1$, so

$$T[x^{-1}] = B[x^{-1}][t_0; \theta]$$

for a suitable automorphism θ of $B[x^{-1}]$.

Let $N \in \text{Gr-}T[x^{-1}]$ and let $\mathcal{N} = \pi N$. Then $\omega \mathcal{N} = N[t_0^{-1}]$, so

$$\phi_* \mathcal{N} = N[t_0^{-1}]_0. \tag{6.2.3}$$

As $N[t_0^{-1}]_0 \otimes_{B[x^{-1}]} T[x^{-1}] = N[t_0^{-1}]_{\geq 0}$, which is isomorphic to $N[t_0^{-1}]$ in $\text{Qgr-}T$, we see that $\phi^* \phi_* \mathcal{N} \cong \mathcal{N}$.

If $M \in \text{Mod-}B[x^{-1}]$, it follows from (6.2.3) that

$$\phi_* \phi^* M = (M \otimes_{B[x^{-1}]} T[x^{-1}])_0 = M. \quad \square$$

We now define cohomology in $\text{Qgr-}T$, that is, the derived functors of $\phi_* = H_{\text{Qgr-}T}^0(-)$. This is similar to the first few pages of [AZ94, Section 7], and we refer the reader there for proofs.

As $\text{Qgr-}T$ is a quotient category of $\text{Gr-}T$, it has enough injectives. In particular, we have:

Lemma 6.2.4. *Let I be a graded-injective T -module. Then πI is an injective object of $\text{Qgr-}T$. As a consequence, $\text{Qgr-}T$ has enough injectives.* \square

It follows that if $\mathcal{N} \in \text{Qgr-}T$ then we can define $\text{Ext}_{\text{Qgr-}T}^j(\mathcal{N}, -)$ to be the j th right derived functor of $\text{Hom}_{\text{Qgr-}T}(\mathcal{N}, -)$. We similarly define $\underline{\text{Ext}}_{\text{Qgr-}T}^j(\mathcal{N}, -)$ as right derived functors of $\underline{\text{Hom}}_{\text{qgr-}T}(\mathcal{N}, -)$, where we recall that $\underline{\text{Hom}}_{\text{qgr-}T}(\mathcal{N}, \mathcal{M}) = \bigoplus_{s \in \mathbb{Z}} \text{Hom}_{\text{Gr-}T}(\mathcal{M}, \mathcal{N}(s))$.

We define the j th cohomology of \mathcal{N} to be

$$H_{\text{Qgr-}T}^j(\mathcal{N}) = \text{Ext}_{\text{Qgr-}T}^j(\pi T, \mathcal{N}),$$

and we also define

$$\underline{H}_{\text{Qgr-}T}^j(\mathcal{N}) = \underline{\text{Ext}}_{\text{Qgr-}T}^j(\pi T, \mathcal{N}).$$

Thus $\underline{H}_{\text{Qgr-}T}^0(\mathcal{N}) = \omega \mathcal{N}$.

We can relate cohomology $\underline{H}_{\text{Qgr-}T}^\bullet(-)$ on $\text{Qgr-}T$ to local cohomology $H_{T_+}^{\bullet+1}(-)$, as usual.

Proposition 6.2.5. *Let M, N be graded right T -modules, with M finitely generated.*

(1) *If $j \geq 0$ then*

$$\underline{\text{Ext}}_{\text{Qgr-}T}^j(\pi M, \pi N) \cong \lim_{d \rightarrow \infty} \underline{\text{Ext}}_T^j(M_{\geq d}, N)$$

and

$$\underline{H}_{\text{Qgr-}T}^j(\pi N) \cong \lim_{d \rightarrow \infty} \underline{\text{Ext}}_T^j(T_{\geq d}, N).$$

(2) *There is an exact sequence*

$$0 \rightarrow \tau(N) \rightarrow N \rightarrow \underline{H}_{\text{Qgr-}T}^0(\pi N) \rightarrow H_{T_+}^1(N) \rightarrow 0,$$

and for $j \geq 1$,

$$\underline{H}_{\text{Qgr-}T}^j(\pi N) \cong H_{T_+}^{j+1}(N).$$

Proof. This is [AZ94, Proposition 7.2], and the proof there works in our situation. \square

Corollary 6.2.6. *Let N be a graded right T -module. Then N and $\omega(\pi N)$ are isomorphic in $\text{Qgr-}T$.*

Proof. This follows from Proposition 6.2.5 and Lemma 6.1.1. \square

6.3. Properness. The goal of this section is to prove the following “relative Serre finiteness” result for $\text{qgr-}T$.

Theorem 6.3.1. (cf. [Har77, Theorem III.5.2], [AZ94, Theorem 7.4]) *Let $\mathcal{N} \in \text{qgr-}T$. Then:*

- (1) *for every $i \geq 0$, $H_{\text{Qgr-}T}^i(\mathcal{N})$ is a finitely generated right B -module;*
- (2) *for every $i \geq 1$, $\underline{H}_{\text{Qgr-}T}^i(\mathcal{N})$ is right bounded: that is, there is $s_0 \in \mathbb{N}$ so that $H_{\text{Qgr-}T}^i(\mathcal{N}(s)) = 0$ for $s \geq s_0$.*

Theorem 6.3.1(1) may be thought of as establishing the properness of ϕ , as if $\psi : Y \rightarrow X$ is a proper morphism of noetherian schemes, then $\mathbb{R}^i \psi_*(\text{coh } Y) \subseteq \text{coh } X$ for all i by [Gro61, 3.2.1]. Theorem 6.3.1(2) shows that $T(1)$ is relatively ample (over B) and so we may think of $\text{qgr-}T$ as being projective over $\text{mod-}B$.

We will prove Theorem 6.3.1 by establishing the Artin–Zhang χ conditions for T . We give the relevant definitions here, generalised from [AZ94]. As in Definition 2.2.1, we say that a graded vector space V is *left bounded* if $V_s = 0$ for $s \ll 0$, *right bounded* if $V_s = 0$ for $s \gg 0$, and *bounded* if it is left and right bounded. Let Ω be a left and right noetherian \mathbb{N} -graded \mathbb{k} -algebra and let $i \in \mathbb{N}$. Then Ω satisfies χ_i° if $\underline{\text{Ext}}_\Omega^j(\Omega/\Omega_+, N)$ is bounded for all $j \leq i$ and for all finitely generated graded Ω -modules N . Further, Ω satisfies χ_i if for all finitely generated graded Ω -modules N , for all d , and for all $j \leq i$, there exists an integer s_0 so that $\underline{\text{Ext}}_\Omega^j(\Omega/\Omega_{\geq s}, N)_{\geq d}$ is a finitely generated Ω -module for all $s \geq s_0$. If Ω satisfies χ_i° for all i , we say that Ω satisfies χ° , and if Ω satisfies χ_i for all i , we say that Ω satisfies χ .

We will see in this section that T satisfies χ , and as a consequence that Theorem 6.3.1 holds. To do this, we generalise a number of results from [AZ94], as to use [AZ94] directly we would need T_0 to be module-finite over a commutative ring.

We first give some general facts about bounds on $\underline{\text{Ext}}$ groups. Note that if Ω is an \mathbb{N} -graded \mathbb{k} -algebra and N is a finitely generated graded Ω -module, then all $\underline{\text{Ext}}_{\Omega}^i(\Omega/\Omega_{\geq s}, N)$ are automatically left bounded (cf. [AZ94, Proposition 3.1(c)]). If Ω satisfies χ_i , then since the modules $\underline{\text{Ext}}_{\Omega}^j(\Omega/\Omega_{\geq s}, N)$ are automatically Ω_+ -torsion, we see that $\underline{\text{Ext}}_{\Omega}^j(\Omega/\Omega_{\geq s}, N)$ is bounded for $j \leq i$ and $s \gg 0$.

Lemma 6.3.2. (cf. [AZ94, Lemma 3.4]) *Let Ω be an \mathbb{N} -graded \mathbb{k} -algebra and let $i \in \mathbb{N}$. Let M, N be finitely generated graded Ω -modules, with M bounded with left bound ℓ and right bound r . Let λ be the minimum left bound and ρ the maximum right bound of $\underline{\text{Ext}}_{\Omega}^j(\Omega/\Omega_+, N)$ for $j \leq i$ (note that we may have $|\lambda|, |\rho| = \infty$). Then for $j \leq i$, the left bound of $\underline{\text{Ext}}_{\Omega}^j(M, N)$ is at least $\lambda - r$, and the right bound is at most $\rho - \ell$.*

Proof. The proof of [AZ94, Lemma 3.4] does not use the standing assumption in [AZ94] that Ω_0 is finite-over-commutative, so may be used unchanged. \square

Proposition 6.3.3. *Let Ω be a left and right noetherian \mathbb{N} -graded \mathbb{k} -algebra that satisfies χ_i . Then Ω satisfies χ_i° .*

Proof. We follow the proof of [AZ94, Corollary 3.6(2)]. Let N be a finitely generated graded Ω -module. By assumption, using the comment before Lemma 6.3.2, there exists s so that $\underline{\text{Ext}}_{\Omega}^j(\Omega/\Omega_{\geq s}, N)$ is bounded for $j \leq i$.

Fix $j \leq i$. By induction, $\underline{\text{Ext}}_{\Omega}^k(\Omega/\Omega_+, N)$ is bounded for $k < j$ and so by Lemma 6.3.2 $\underline{\text{Ext}}_{\Omega}^{j-1}(\Omega_+/\Omega_{\geq s}, N)$ is bounded. From the exact sequence

$$\underline{\text{Ext}}_{\Omega}^{j-1}(\Omega_+/\Omega_{\geq s}, N) \rightarrow \underline{\text{Ext}}_{\Omega}^j(\Omega/\Omega_+, N) \rightarrow \underline{\text{Ext}}_{\Omega}^j(\Omega/\Omega_{\geq s}, N),$$

we see that $\underline{\text{Ext}}_{\Omega}^j(\Omega/\Omega_+, N)$ is bounded. \square

We now consider the χ condition for the graded ring

$$S = \mathbb{k}_{\mathbf{q}}[x, y, z, t_0, \dots, t_n]$$

defined in (5.2.1). Here the grading on S is such that x, y, z are in degree 0 and $\deg t_i = 1$ for all i . As $T \cong T_{\text{pres}}$ by Theorem 5.2.8, the obvious map $S \rightarrow T$ is well-defined.

We define torsion S -modules as we did torsion T -modules.

Proposition 6.3.4. (cf. [AZ94, Theorem 8.1]) *Let M, N be finitely generated graded S -modules such that M is torsion. Then $\underline{\text{Ext}}_S^i(M, N)$ is bounded for all i . Further, $\underline{\text{Ext}}_S^i(S/S_{\geq s}, N)$ is a finitely generated S -module for all i and s , and in particular S satisfies χ .*

Proof. As S is noetherian, we have $MS_{\geq s} = 0$ for some s . From the long exact $\underline{\text{Ext}}$ sequence, it is enough to prove the first statement in the case that $MS_+ = 0$, i.e. that M is an S/S_+ -module.

First assume that $M = S/S_+$ and $N = S$. From the Koszul resolution of S/S_+ as an S -module we immediately compute that

$$\underline{\text{Ext}}_S^i(S/S_+, S) = \begin{cases} S_0(n+1) & \text{if } i = n+1, \\ 0 & \text{otherwise.} \end{cases}$$

It follows immediately that the proposition holds if M_{S/S_+} and N_S are projective.

Now let N be an arbitrary S -module and consider a partial resolution $0 \rightarrow N' \rightarrow P \rightarrow N \rightarrow 0$ of S -modules, where P is projective. By induction on $\text{pd}_S N$ and the long exact sequence in $\underline{\text{Ext}}$ we see that $\underline{\text{Ext}}_S^i(S/S_+, N)$ is bounded and finitely generated over S for all i .

Using the long exact $\underline{\text{Ext}}$ sequence and the conclusion of the previous paragraph, we see that $\underline{\text{Ext}}_S^i(S/S_{\geq s}, N)$ is finitely generated and bounded for all i, s , and that if M is an arbitrary finitely generated torsion module then $\underline{\text{Ext}}_S^i(M, N)$ is bounded for all i . \square

Before we can prove that T satisfies χ , we will need to consider bounds on the (S, T) -bimodules $\text{Tor}_i^S(S/S_{\geq s}, T)$, which have a \mathbb{Z} -grading induced from the gradings on S and T .

Proposition 6.3.5. (cf. [AZ94, Proposition 2.4]) *Let $i, s \in \mathbb{N}$. Then $\text{Tor}_i^S(S/S_{\geq s}, T)$ is a finitely generated and bounded T -module, and if $i \geq 1$ the left bound is at least s .*

Proof. That $\text{Tor}_i^S(S/S_{\geq s}, T)$ is finitely generated over T follows by resolving $S/S_{\geq s}$ by finitely generated graded S -modules. Resolving T as an S -module, we see that $\text{Tor}_i^S(S/S_{\geq s}, T)$ is torsion, and thus bounded since it is finitely generated. To compute the left bound, we follow [AZ94]. For $i \geq 2$, $\text{Tor}_i^S(S/S_{\geq s}, T) \cong \text{Tor}_{i-1}^S(S_{\geq s}, T)$. Taking a resolution of T by finitely generated graded S -modules, we can see the left bound of $\text{Tor}_{i-1}^S(S_{\geq s}, T)$ is at least s . Further, $\text{Tor}_1^S(S/S_{\geq s}, T) \subseteq S_{\geq s} \otimes_S T$, which has left bound s . \square

Proposition 6.3.6. (cf. [AZ94, Lemma 8.2])

- (1) T satisfies χ and χ° .
- (2) Let M, N be finitely generated graded T -modules with M torsion. Then all $\underline{\text{Ext}}_T^i(M, N)$ are bounded.

Proof. We prove (1) following the proof of [AZ94, Lemma 8.2]. By induction, assume T satisfies χ_{i-1}° . Let $N \in \text{gr-}T$ and $s \in \mathbb{N}$, and consider the spectral sequence

$$E_2^{p,r} = \underline{\text{Ext}}_T^p(\text{Tor}_r^S(S/S_{\geq s}, T), N) \Rightarrow \underline{\text{Ext}}_S^{p+r}(S/S_{\geq s}, N_S)$$

and the edge map

$$E_2^{i,0} = \underline{\text{Ext}}_T^i(T/T_{\geq s}, N) \rightarrow \underline{\text{Ext}}_S^i(S/S_{\geq s}, N_S). \quad (6.3.7)$$

Let $r \geq 1$. By Proposition 6.3.5, the left bound of $\text{Tor}_r^S(S/S_{\geq s}, T)$ goes to ∞ as $s \rightarrow \infty$. By the induction hypothesis, $\underline{\text{Ext}}_T^{p-r}(T/T_{\geq s}, N)$ is bounded and so by Lemma 6.3.2 the right bound of $E_2^{p-r,r} = \underline{\text{Ext}}_T^{p-r}(\text{Tor}_r^S(S/S_{\geq s}, T), N)$ goes to $-\infty$ as $s \rightarrow \infty$. For $d \in \mathbb{Z}$, therefore, by taking $s \gg 0$ we may assume that $(E_2^{i-r,r})_{\geq d} = 0$ and thus that (6.3.7) is an isomorphism in degrees $\geq d$.

By Proposition 6.3.4,

$$\underline{\text{Ext}}_T^i(T/T_{\geq s}, N)_{\geq d} \cong \underline{\text{Ext}}_S^i(S/S_{\geq s}, N_S)_{\geq d}$$

is a finitely generated S -module and thus a finitely generated T -module. Thus T satisfies χ_i , and by Proposition 6.3.3 it satisfies χ_i° .

Part (2) is a consequence of the fact that T satisfies χ° and Lemma 6.3.2. \square

We need two subsidiary results before proving Serre finiteness for T .

Proposition 6.3.8. (cf. [AZ94, Proposition 3.5]) *Let $M, N \in \text{gr-}T$.*

- (1) *For all $d \in \mathbb{Z}$ and $i \in \mathbb{N}$ there is $s_0 \in \mathbb{Z}$ so that for all $s \geq s_0$ we have*

$$\underline{\text{Ext}}_T^i(M/M_{\geq s_0}, N)_{\geq d} \cong \underline{\text{Ext}}_T^i(M/M_{\geq s}, N)_{\geq d}.$$

- (2) *For all $d \in \mathbb{Z}$ and $i \in \mathbb{N}$ there is $s_0 \in \mathbb{Z}$ so that for all $s \geq s_0$ we have*

$$\underline{\text{Ext}}_T^i(M_{\geq s_0}, N)_{\geq d} \cong \underline{\text{Ext}}_T^i(M_{\geq s}, N)_{\geq d}.$$

- (3) *For all $i \in \mathbb{N}$, there is $\ell \in \mathbb{Z}$ so that for all $s \in \mathbb{Z}$ we have*

$$\underline{\text{Ext}}_T^i(M, N)_{\geq \ell} \cong \underline{\text{Ext}}_T^i(M_{\geq s}, N)_{\geq \ell}.$$

Proof. By Proposition 6.3.6 T satisfies χ° . Then the proof of [AZ94, Proposition 3.5] goes through in our context, using Lemma 6.3.2 instead of [AZ94, Lemma 3.4]. \square

Proposition 6.3.9. (cf. [AZ94, Proposition 3.13]) *Let $N \in \text{gr-}T$.*

- (1) *For all $d \in \mathbb{Z}$ and for all $s \gg 0$, we have*

$$\underline{\text{Hom}}_T(T_{\geq s}, N)_{\geq d} \cong (\omega\pi N)_{\geq d}.$$

- (2) *There is $\ell \in \mathbb{Z}$ so that*

$$N_{\geq \ell} \cong (\omega\pi N)_{\geq \ell}.$$

Proof. We follow the proof in [AZ94]. By definition

$$\omega\pi N = \lim_{s \rightarrow \infty} \underline{\text{Hom}}_T(T_{\geq s}, N). \quad (6.3.10)$$

Then (1) is (6.3.10) combined with Proposition 6.3.8(2). For (2), by Proposition 6.3.8(3) there is ℓ so that $N_{\geq \ell} \cong \underline{\text{Hom}}_T(T_{\geq s}, N)_{\geq \ell}$ for all s . But by (1) this is $(\omega\pi N)_{\geq \ell}$. \square

We can now prove relative Serre finiteness for $\text{qgr-}T$.

Proof of Theorem 6.3.1. Let $N \in \text{gr-}T$ so that $\pi N = \mathcal{N}$, and let $i \in \mathbb{N}$. By Proposition 6.3.8(1) there is s_0 so that for all $j \leq i$ we have

$$\underline{H}_{\text{Qgr-}T}^j(\mathcal{N}) = \lim_{s \rightarrow \infty} \underline{\text{Ext}}_T^j(T_{\geq s}, N) = \underline{\text{Ext}}_T^j(T_{\geq s_0}, N).$$

Similarly to Proposition 6.2.5(2), if $j \geq 1$ then

$$\underline{H}_{\text{Qgr-}T}^j(\mathcal{N}) \cong \underline{\text{Ext}}_T^{j+1}(T/T_{\geq s_0}, N),$$

and there is an exact sequence

$$0 \rightarrow \underline{\text{Hom}}_T(T/T_{\geq s_0}, N) \rightarrow N \rightarrow \underline{H}_{\text{Qgr-}T}^0(\mathcal{N}) \rightarrow \underline{\text{Ext}}_T^1(T/T_{\geq s_0}, N) \rightarrow 0.$$

Since T satisfies χ , therefore, $\underline{H}_{\text{Qgr-}T}^i(\mathcal{N})_{\geq 0}$ is a finitely generated T -module, and thus $\underline{H}_{\text{Qgr-}T}^i(\mathcal{N})$ is finitely generated over $T_0 = B$. Further, if $i \geq 1$ then $\underline{H}_{\text{Qgr-}T}^i(\mathcal{N})_{\geq 0} \cong H_{T_+}^{i+1}(N)_{\geq 0}$ is torsion, and as it is finitely generated it must be right bounded. \square

6.4. The local cohomology of T . It will clearly be useful to be able to compute some (local) cohomology groups. The goal of this section is to show that the Čech complex $\check{C}(t_0, \dots, t_n; T)$ of T is exact in degree 0. Recall from (5.3.5) that we can embed T into the quantum polynomial ring $\mathbb{k}_{q^\bullet}[\mathbf{a}, \mathbf{b}, \mathbf{s}] \subseteq D$, where D is defined in (5.3.1), by setting

$$x = \mathbf{a}, \quad y = q^\bullet \mathbf{a}^n \mathbf{b}^{n+1}, \quad z = q^\bullet \mathbf{a} \mathbf{b}, \quad t_i = q^\bullet \mathbf{a}^{\binom{i}{2}} \mathbf{b}^{\binom{i+1}{2}} \mathbf{s},$$

for suitable scalars q^\bullet , and that this induces the toric grading on T , where $\deg \mathbf{a}^i \mathbf{b}^j \mathbf{s}^k = (i, j, k)$. In particular, the third component records the projective grading of T . Thus, any graded localisation of T can be viewed as living inside the quantum torus D , and the degree 0 part (with respect to the third component of the grading, i.e. the projective grading) of any such ring can be viewed as living inside the quantum division ring $Q(\mathbb{k}_{q^{n+1}}[\mathbf{a}^{\pm 1}, \mathbf{b}^{\pm 1}])$. (Here we recall from (5.3.1) that \mathbf{a} and \mathbf{b} q^{n+1} -commute.) Therefore we have the following:

Lemma 6.4.1. (1) For $0 \leq k \leq n$, we have

$$T[t_k^{-1}]_0 = \begin{cases} \mathbb{k}_{q^{n+1}}[x, t_1 t_0^{-1}] & \text{if } k = 0, \\ \mathbb{k}_{q^{n+1}}[t_{k-1} t_k^{-1}, t_{k+1} t_k^{-1}] & \text{if } 1 \leq k \leq n-1, \\ \mathbb{k}_{q^{n+1}}[t_{n-1} t_n^{-1}, y] & \text{if } k = n. \end{cases} \quad (6.4.2)$$

If we identify T with T_{toric} , then

$$T[t_k^{-1}]_0 = \mathbb{k}_{q^{n+1}}[\mathbf{a}^{-(k-1)} \mathbf{b}^{-k}, \mathbf{a}^k \mathbf{b}^{k+1}].$$

In particular, the given generators q^{n+1} -commute.

(2) For $0 \leq k \leq n-1$, we have

$$T[t_k^{-1}, t_{k+1}^{-1}]_0 = \begin{cases} \mathbb{k}_{q^{n+1}}[x, (t_0 t_1^{-1})^{\pm 1}] & \text{if } k = 0, \\ \mathbb{k}_{q^{n+1}}[t_{k-1} t_k^{-1}, (t_k t_{k+1}^{-1})^{\pm 1}] & \text{if } 1 \leq k \leq n. \end{cases}$$

If we identify T with T_{toric} , then

$$T[t_k^{-1}, t_{k+1}^{-1}]_0 = \mathbb{k}_{q^{n+1}}[\mathbf{a}^{-(k-1)} \mathbf{b}^{-k}, (\mathbf{a}^k \mathbf{b}^{k+1})^{\pm 1}]. \quad (6.4.3)$$

(3) For $0 \leq j < k \leq n$ with $k - j \geq 2$ we have

$$T[t_j^{-1}, t_k^{-1}]_0 = \mathbb{k}_{q^{n+1}}[\mathbf{a}^{\pm 1}, \mathbf{b}^{\pm 1}]. \quad (6.4.4)$$

(4) If $J \subseteq \{t_0, \dots, t_n\}$ with $\#J \geq 3$, then

$$T[t_J^{-1}] = \mathbb{K}_{q^{n+1}}[\mathbf{a}^{\pm 1}, \mathbf{b}^{\pm 1}]. \quad (6.4.5)$$

Proof. (1) It is clear that any element of $T[t_k^{-1}]_0$ is a linear combination of products of x, y, z and elements of the form $t_j t_k^{-1}$, where $0 \leq j \leq n$. In each case, the claimed generators of $T[t_k^{-1}]_0$ are monomials in \mathbf{a}, \mathbf{b} , and \mathbf{s} , and hence generate a subalgebra of D which is isomorphic to a quantum plane. We write this as $\mathbb{K}_{q^\bullet}[x, t_1 t_0^{-1}] \subseteq T[t_0^{-1}]_0$, for example, and have similar containments in the other cases.

For the reverse inclusion, we consider the three cases separately, beginning with $k = 0$; the case $k = n$ is similar and hence omitted. The relations in T allow us to write

$$z = q^{-\alpha_1} x t_1 t_0^{-1}, \quad y = q^{\beta_0} z^n t_1 t_0^{-1},$$

the first of which clearly lies in $\mathbb{K}_{q^\bullet}[x, t_1 t_0^{-1}]$, from which it follows that the same is true of y . Moreover, for $2 \leq j \leq n$, the relation $t_0 t_j = q^\bullet z^{j-1} t_1 t_{j-1}$ implies that

$$t_j t_0^{-1} = q^\bullet z^{j-1} (t_1 t_0^{-1}) (t_{j-1} t_0^{-1}),$$

and induction shows that $t_j t_0^{-1} \in \mathbb{K}_{q^\bullet}[x, t_1 t_0^{-1}]$.

Now suppose that $1 \leq k \leq n$. The relation $t_{k-1} t_{k+1} = q^\bullet z t_k^2$ in T gives

$$z = q^\bullet (t_{i-1} t_i^{-1}) (t_{i+1} t_i^{-1})$$

in $T[t_k^{-1}]$, which lies in $\mathbb{K}_{q^\bullet}[t_{k-1} t_k^{-1}, t_{k+1} t_k^{-1}]$. We also have

$$x = q^\bullet z^k t_{k-1} t_k^{-1}, \quad y = q^\bullet z^{n-k} t_{k+1} t_k^{-1}$$

from which it follows that $x, y \in \mathbb{K}_{q^\bullet}[t_{k-1} t_k^{-1}, t_{k+1} t_k^{-1}]$. Moreover, if $j \geq k+2$, then $t_k t_j = q^\bullet z^{j-k-1} t_{k+1} t_{j-1}$, and hence

$$t_j t_k^{-1} = q^\bullet z^{j-k-1} (t_{k+1} t_k^{-1}) (t_{j-1} t_k^{-1}),$$

and induction shows that $t_j t_k^{-1} \in \mathbb{K}_{q^\bullet}[t_{k-1} t_k^{-1}, t_{k+1} t_k^{-1}]$. Similarly, one can show that $t_j t_k^{-1} \in \mathbb{K}_{q^\bullet}[t_{k-1} t_k^{-1}, t_{k+1} t_k^{-1}]$ when $j \leq k-2$, establishing the claimed equality.

Finally, it is straightforward to check that the given generators are identified with $\mathbf{a}^{-(k-1)} \mathbf{b}^{-k}$ and $\mathbf{a}^k \mathbf{b}^{k+1}$ under the toric embedding, which gives the final equality. Direct calculation shows that these elements q^{n+1} -commute:

$$\mathbf{a}^k \mathbf{b}^{k+1} \cdot \mathbf{a}^{-(k-1)} \mathbf{b}^{-k} = q^{-(n+1)((k+1)(k-1)-k^2)} \mathbf{a}^{-(k-1)} \mathbf{b}^{-k} \cdot \mathbf{a}^k \mathbf{b}^{k+1} = q^{n+1} \mathbf{a}^{-(k-1)} \mathbf{b}^{-k} \cdot \mathbf{a}^k \mathbf{b}^{k+1}.$$

(2) It is clear that, for all values of k , the right hand side is contained in the left hand side; we now consider the reverse inclusion. Suppose that $1 \leq k \leq n-1$. By part (1), we have

$$T[t_k^{-1}]_0 = \mathbb{K}_{q^{n+1}}[t_{k-1} t_k^{-1}, t_{k+1} t_k^{-1}] \quad \text{and} \quad T[t_{k+1}^{-1}]_0 = \mathbb{K}_{q^{n+1}}[t_k t_{k+1}^{-1}, t_{k+2} t_{k+1}^{-1}],$$

so that $T[t_k^{-1}, t_{k+1}^{-1}]_0$ is the subring generated by these elements:

$$T[t_k^{-1}, t_{k+1}^{-1}]_0 = \mathbb{K}\langle t_{k-1} t_k^{-1}, (t_k t_{k+1}^{-1})^{\pm 1}, t_{k+2} t_{k+1}^{-1} \rangle,$$

It therefore suffices to show that $t_{k+2} t_{k+1}^{-1}$ lies in $\mathbb{K}_{q^{n+1}}[t_{k-1} t_k^{-1}, (t_k t_{k+1}^{-1})^{\pm 1}]$. From the relations in T , we have

$$t_{k-1} t_{k+1}^3 = q^\bullet z t_k^2 t_{k+1}^2 = q^\bullet t_k^3 t_{k+2},$$

and hence

$$t_{k+2} t_{k+1}^{-1} = q^\bullet (t_{k-1} t_k^{-1}) (t_{k+1} t_k^{-1})^2 \in \mathbb{K}_{q^{n+1}}[t_{k-1} t_k^{-1}, (t_k t_{k+1}^{-1})^{\pm 1}],$$

as desired.

The cases $k = 0$ and $k = n$ are similar; we prove $k = 0$. In this case, it suffices to show that $t_2 t_1^{-1}$ lies in $\mathbb{K}_{q^{n+1}}[x, (t_0 t_1^{-1})^{\pm 1}]$. To see this, note that we have

$$x t_1^3 = q^\bullet z t_0 t_1^2 = q^\bullet t_0^2 t_2,$$

and hence

$$t_2 t_1^{-1} = q^\bullet x (t_1 t_0^{-1})^2 \in \mathbb{k}_{q^{n+1}}[x, (t_0 t_1^{-1})^{\pm 1}].$$

(3) It suffices to show that $\mathbf{a}^{\pm 1}, \mathbf{b}^{\pm 1} \in T[t_j^{-1}, t_k^{-1}]_0$ when $0 \leq j < k \leq n$ with $k - j \geq 2$. Since $x \in T$, we have $\mathbf{a} \in T[t_j^{-1}, t_k^{-1}]_0$. We also have

$$T[t_j^{-1}, t_k^{-1}]_0 \ni z^{k-1} (t_{k-1} t_k^{-1}) = q^\bullet \mathbf{a}^{k-1} \mathbf{b}^{k-1} \mathbf{a}^{-(k-1)} \mathbf{b}^{-k} = q^\bullet \mathbf{b}^{-1},$$

so that $\mathbf{b}^{-1} \in T[t_j^{-1}, t_k^{-1}]_0$. Additionally, we have

$$T[t_j^{-1}, t_k^{-1}]_0 \ni (t_{j+1} t_j^{-1})^{k-1} (t_{k-1} t_k^{-1})^j = q^\bullet (\mathbf{a}^j \mathbf{b}^{j+1})^{k-1} (\mathbf{a}^{-(k-1)} \mathbf{b}^{-k})^j = q^\bullet \mathbf{b}^{k-j-1},$$

where $k - j - 1 \geq 1$. Multiplying by a suitable power of \mathbf{b}^{-1} shows that $\mathbf{b} \in T[t_j^{-1}, t_k^{-1}]_0$. Finally,

$$T[t_j^{-1}, t_k^{-1}]_0 \ni \mathbf{a}^{k-2} \mathbf{b}^k (t_{k-1} t_k^{-1}) = q^\bullet \mathbf{a}^{k-2} \mathbf{b}^k \mathbf{a}^{-(k-1)} \mathbf{b}^{-k} = q^\bullet \mathbf{a}^{-1},$$

so that $\mathbf{a}^{-1} \in T[t_j^{-1}, t_k^{-1}]_0$. Thus $T[t_j^{-1}, t_k^{-1}]_0 = \mathbb{k}_{q^{n+1}}[\mathbf{a}^{\pm 1}, \mathbf{b}^{\pm 1}]$.

(4) This follows from (3). \square

The computations in Lemma 6.4.1 show that $\text{qgr-}T$ is “locally nonsingular” and so it is natural to expect that it has finite homological dimension. We will prove that this is the case in Section 7.

Let $\check{C}^\bullet = \check{C}(t_0, \dots, t_n; T)_0$. Note that for all $(i, j) \in \mathbb{Z}^2$ and $s \in \mathbb{N}$ we have

$$\dim \check{C}_{(i,j)}^s \leq \binom{n+1}{s} < \infty$$

from Lemma 6.4.1(4).

Lemma 6.4.6. *For all $(i, j) \in \mathbb{Z}^2$, the Euler characteristic of $\check{C}_{(i,j)}^\bullet$ is 0: that is,*

$$\sum_{k \geq 0} (-1)^k \dim \check{C}_{(i,j)}^k = 0.$$

Proof. If $X \subseteq \mathbb{k}_{q^{n+1}}[\mathbf{a}^\pm, \mathbf{b}^\pm]$ is bigraded, define $\text{supp}(X) = \{(i, j) \mid \mathbf{a}^i \mathbf{b}^j \in X\}$.

We now describe a partition of \mathbb{Z}^2 . Let $Q_{\text{I}} = \mathbb{N} \times \mathbb{N} \setminus \{(0, 0)\}$ be the “first quadrant” and let $Q_{\text{III}} = \{(i, j) \in \mathbb{Z} \times \mathbb{Z} \mid i, j < 0\}$ be the “third quadrant”. For $1 \leq k \leq n$ let

$$U_k = \left\{ (i, j) \in Q_{\text{I}} \mid \frac{k+1}{k} < \frac{j}{i} \leq \frac{k}{k-1} \right\}$$

and also define

$$W = \left\{ (i, j) \in Q_{\text{I}} \mid \frac{j}{i} \leq \frac{n+1}{n} \right\} \cup \{0\}.$$

Moreover, let

$$L_0 = \{(i, j) \in \mathbb{Z}^2 \mid i \geq 0, j < 0\}$$

and, for $1 \leq k \leq n-1$, let

$$L_k = \left\{ (i, j) \in Q_{\text{III}} \mid \frac{k+1}{k} \leq \frac{j}{i} < \frac{k}{k-1} \right\}.$$

Finally, set

$$Z = \left\{ (i, j) \in \mathbb{Z}^2 \mid i < 0, \frac{j}{i} < \frac{n}{n-1} \right\}.$$

We observe that as $T_0 = \mathbb{k}\langle \mathbf{a}, \mathbf{a}^n \mathbf{b}^{n+1}, \mathbf{ab} \rangle \subseteq \mathbb{k}_{q^{n+1}}[\mathbf{a}, \mathbf{b}]$ we have

$$\text{supp } T_0 = \mathbb{N} \cdot (1, 0) + \mathbb{N} \cdot (n, n+1) + \mathbb{N} \cdot (1, 1) = W.$$

Let $0 \leq k \leq n$. From (6.4.2) we calculate:

$$\text{supp } T[t_k^{-1}]_0 = \mathbb{N} \cdot (-(k-1), -k) + \mathbb{N} \cdot (k, k+1) = L_{k-1} \sqcup \dots \sqcup L_0 \sqcup W \sqcup U_n \sqcup \dots \sqcup U_{k+1}.$$

Likewise, (6.4.3) tells us that, if $0 \leq k \leq n-1$, then

$$\text{supp } T[t_k^{-1}, t_{k+1}^{-1}]_0 = \mathbb{N} \cdot (-(k-1), -k) + \mathbb{Z} \cdot (k, k+1) = L_k \sqcup \dots \sqcup L_0 \sqcup W \sqcup U_n \sqcup \dots \sqcup U_{k+1}.$$

Combining these computations with (6.4.4) and (6.4.5) we obtain that

$$\dim \check{C}_{(i,j)}^s = \begin{cases} \binom{n+1}{s} & \text{if } s \geq 3 \text{ or } (i,j) \in W, \\ \binom{n}{2} & \text{if } s = 2 \text{ and } (i,j) \in Z, \\ \binom{n}{2} + k & \text{if } s = 2 \text{ and } (i,j) \in U_k, \\ \binom{n}{2} + n - k & \text{if } s = 2 \text{ and } (i,j) \in L_k, \\ k & \text{if } s = 1 \text{ and } (i,j) \in U_k, \\ n - k & \text{if } s = 1 \text{ and } (i,j) \in L_k, \\ 0 & \text{otherwise.} \end{cases}$$

The result follows. \square

We can now compute the degree 0 local cohomology of T .

Proposition 6.4.7. *The Čech complex \check{C}^\bullet is exact, and $H_{T_+}^s(T)_0 = 0$ for all s .*

Proof. We first observe that $\dim H_{T_+}^s(T)_0$ does not depend on q . To see this, note that for any $J \subseteq [n]$ and any $(i,j) \in \mathbb{Z}^2$, $T[t_J^{-1}]_{(i,j)}$ is either 0 or $\mathbb{k}\mathbf{a}^i\mathbf{b}^j$, and which occurs is independent of q . Thus if we define

$$m_J^{ij} = \begin{cases} \mathbf{a}^i\mathbf{b}^j & \text{if } T[x_J^{-1}]_{(i,j)} \neq 0, \\ 0 & \text{otherwise,} \end{cases}$$

then $d_s : \check{C}^s \rightarrow \check{C}^{s+1}$ is given by $d_s(m_J^{(i,j)}) = \sum_{k \notin J} (-1)^{o_J(k)} m_{J \cup \{k\}}^{(i,j)}$, which again does not depend on q . It thus suffices to compute $H_{T_+}^s(T)_0$ if $q = 1$.

If $q = 1$ then $\tilde{X} = \text{Proj}(T)$ is the minimal resolution of $X = \mathbb{k}^2/C_{n+1}$, constructed via quiver GIT as explained in [Wem16]. The singularity of X is rational by [Art66]; that is, if $\pi : \tilde{X} \rightarrow X$ is the morphism associated to the resolution then

$$H^s(\tilde{X}, \mathcal{O}_{\tilde{X}}) = \mathbb{R}^s \pi_* \mathcal{O}_{\tilde{X}} = \begin{cases} 0 & \text{if } s \geq 1, \\ \mathcal{O}_X & \text{if } s = 0. \end{cases}$$

But $\text{coh } \tilde{X} \simeq \text{qgr-}T$ by [Sta25, Tag 0BXD] so by Proposition 6.2.5(2) for $s \geq 2$ we have $H_{T_+}^s(T)_0 \cong H^{s-1}(\tilde{X}, \mathcal{O}_{\tilde{X}})$, giving the result for $s \geq 2$.

As T is a domain, $H_{T_+}^0(T) = 0$. The complex \check{C}^\bullet is thus exact except possibly at $s = 1$. Applying Lemma 6.4.6 we see that \check{C}^\bullet is exact, as claimed. \square

Corollary 6.4.8. *Let \mathcal{N} be a finitely generated \mathbb{Z}^3 -graded right T -module that is locally free in the sense that $\mathcal{N}[t_i^{-1}]_0$ is a free $T[t_i^{-1}]_0$ -module for all $0 \leq i \leq n$. Then $H_{T_+}^s(\mathcal{N})_0 = 0$ for all $s \geq 3$.*

Proof. Since \mathcal{N} is locally free, the Čech complex for \mathcal{N} is isomorphic to that of $T^{\oplus d}$ in cohomological degrees ≥ 2 , where d is the uniform rank of \mathcal{N} . Thus if $s \geq 3$ then $H_{T_+}^s(\mathcal{N})_0 \cong H_{T_+}^s(T)_0^d$. The result follows from Proposition 6.4.7. \square

7. A SYSTEM FOR LOCALISING T -MODULES

Let $q \in \mathbb{k}^\times$ be arbitrary and let $T = T_q$. In this section we prove two main results: first, that many computations in $\text{qgr-}T$ can be done locally, i.e. on “charts”; and, second, that $\text{qgr-}T$ has finite homological dimension. This is done by establishing an equivalence of categories between $\text{qgr-}T$ and the category of coherent sheaves over a certain noncommutative ringed space \mathfrak{X} .

7.1. A noncommutative toric variety equivalent to $\text{qgr-}T$. By (5.3.5), we can view $T \subseteq D$ via the toric embedding with, up to scalars,

$$x = \mathbf{a}, \quad y = \mathbf{a}^n \mathbf{b}^{n+1}, \quad z = \mathbf{a}\mathbf{b}, \quad t_i = \mathbf{a}^{\binom{i}{2}} \mathbf{b}^{\binom{i+1}{2}} \mathbf{s}.$$

Any two of the generators $\{x, y, z, t_0, \dots, t_n\}$ q^\bullet -commute, and as a result the generators of T are all normal.

Define a topological space \mathfrak{X} to be the set of \mathbb{Z}^3 -graded prime ideals of T which do not contain the irrelevant ideal $T_+ = (t_0, \dots, t_n)$, in the Zariski topology.

Lemma 7.1.1. *The elements of \mathfrak{X} are:*

$$\begin{aligned} 0 \\ \mathfrak{l}_0 &= \langle y, z, t_1, \dots, t_n \rangle \\ \mathfrak{l}_i &= \langle x, y, z, t_0, \dots, t_{i-2}, t_{i+1}, \dots, t_n \rangle \quad \text{for } 1 \leq i \leq n \\ \mathfrak{l}_{n+1} &= \langle x, z, t_0, \dots, t_{n-1} \rangle \\ \mathfrak{p}_i &= \langle x, y, z, t_0, \dots, t_{i-1}, t_{i+1}, \dots, t_n \rangle \quad \text{for } 0 \leq i \leq n. \end{aligned}$$

In particular, \mathfrak{X} is a two-dimensional topological space.

Remark 7.1.2. The reason for the notation here is that we may define cyclic T -modules $L_i = T/\mathfrak{l}_i$ and $P_i = T/\mathfrak{p}_i$. We will see in Section 9 that these are, respectively, line and point modules, and that they have the (noncommutative) intersection theory of an A_n Dynkin diagram.

Proof of Lemma 7.1.1. We view T as a subring of D via the toric embedding of Section 5.3. For any $\vec{n} \in \mathbb{N}^3$ we have $\dim T_{\vec{n}} \leq 1$ from the inclusion $T \subseteq D$. Thus an ideal of T is \mathbb{Z}^3 -graded exactly when it is generated by monomials in $\mathbf{a}, \mathbf{b}, \mathbf{s}$; but an element of T is a monomial in $\mathbf{a}, \mathbf{b}, \mathbf{s}$ exactly when it is a monomial in x, y, z, t_0, \dots, t_n . As these elements are all normal, an ideal \mathfrak{n} of T is \mathbb{Z}^3 -graded and prime if and only if \mathfrak{n} is generated by some subset of $\{x, y, z, t_0, \dots, t_n\}$.

Let $0 \neq \mathfrak{p} \in \mathfrak{X}$. If $z \notin \mathfrak{p}$ then from the relation $xy = q^\bullet z^{n+1}$ we see that $x, y \notin \mathfrak{p}$. But now some $t_i \in \mathfrak{p}$ as $\mathfrak{p} \neq 0$, and then from the relations $t_{i-1}y = q^\bullet t_i z^{n-i+1}$ and $t_{i+1}x = q^\bullet t_i z^{i+1}$ and the assumption that $z \notin \mathfrak{p}$ we have $t_0, \dots, t_n \in \mathfrak{p}$. This contradicts the assumption that $\mathfrak{p} \not\supseteq T_+$. It follows that $z \in \mathfrak{p}$. Now for all $1 \leq i < j \leq n-1$ we have $t_{i-1}t_{j+1} = q^\bullet z^{j-i+1}t_it_j \in \mathfrak{p}$, so if $0 \leq i \leq j \leq n$ with $j \geq i+2$ then $t_i \in \mathfrak{p}$ or $t_j \in \mathfrak{p}$.

By hypothesis there is some $t_i \notin \mathfrak{p}$; let i be the minimal such index. We therefore have $\mathfrak{p} \supseteq \langle t_0, \dots, t_{i-1}, t_{i+2}, \dots, t_n \rangle$ by the observation in the previous paragraph. If $i \leq n-1$ then $t_i y = q^\bullet t_{i+1} z^{n-i} \in \mathfrak{p}$ so $y \in \mathfrak{p}$ as $t_i \notin \mathfrak{p}$. Likewise if $i \geq 1$ then $x \in \mathfrak{p}$. Thus if $1 \leq i \leq n-1$ then $\mathfrak{p} \supseteq \mathfrak{l}_{i+1}$ and

$$\mathfrak{p} = \begin{cases} \mathfrak{l}_{i+1} & \text{if } t_{i+1} \notin \mathfrak{p}, \\ \mathfrak{p}_i & \text{if } t_{i+1} \in \mathfrak{p}. \end{cases}$$

If $i = 0$ then $\mathfrak{p} \supseteq \langle z, y, t_2, \dots, t_n \rangle$. Since $xt_1 = q^\bullet zt_0 \in \mathfrak{p}$, either $x \in \mathfrak{p}$ or $t_1 \in \mathfrak{p}$. If $x \notin \mathfrak{p}$ then $\mathfrak{p} = \mathfrak{l}_0$. If $t_1 \notin \mathfrak{p}$ then $\mathfrak{p} = \mathfrak{l}_1$. If $x, t_1 \in \mathfrak{p}$ then $\mathfrak{p} = \mathfrak{p}_0$. Finally, if $i = n$ then $\mathfrak{p} \supseteq \mathfrak{l}_{n+1}$ and

$$\mathfrak{p} = \begin{cases} \mathfrak{l}_{n+1} & \text{if } y \notin \mathfrak{p}, \\ \mathfrak{p}_n & \text{if } y \in \mathfrak{p}. \end{cases} \quad \square$$

We now make \mathfrak{X} into a ringed space by defining a sheaf $\mathcal{O}_{\mathfrak{X}}$ of noncommutative rings on \mathfrak{X} . Our conventions for ringed spaces will in general follow [Har77], except, of course, that our rings are noncommutative.

For notational purposes let $t_{n+1} = y$ and $t_{-1} = x$. Let $I \subseteq \{-1, \dots, n+1\}$, and let

$$V(I) = V\left(\prod_{i \in I} t_i\right) = \left\{ \mathfrak{p} \in \mathfrak{X} \mid \prod_{i \in I} t_i \in \mathfrak{p} \right\}.$$

The subsets $V(I)$ generate a topology on \mathfrak{X} , where the closed subsets of \mathfrak{X} are the (necessarily finite) intersections of the $V(I)$. If $I \subseteq \{-1, \dots, n+1\}$, define U_I to be $\mathfrak{X} \setminus V(I)$. We refer to the U_I with $I \neq \emptyset$ as *principal open subsets of \mathfrak{X}* or as *principal open sets*. We will omit curly braces from our notation and write, for example, U_i for $U_{\{i\}}$ and U_{ij} for $U_{\{i,j\}}$.

Let U_I be a principal open set, and define the *coordinate ring of U_I* to be

$$\mathcal{O}_{\mathfrak{X}}(U_I) = T[t_i^{-1} \mid i \in I]_0.$$

Recall that the $\mathcal{O}_{\mathfrak{X}}(U_I)$ were computed in Lemma 6.4.1. In particular, $\mathcal{O}_{\mathfrak{X}}(U_i) \cong \mathbb{k}_{q^{n+1}}[u, v]$ for all i .

As an intersection of principal open sets is again principal, any open subset of \mathfrak{X} has a cover by principal open sets. If V is any non-principal open subset of \mathfrak{X} write $V = \bigcup U_J$ where the U_J are principal, and put

$$\mathcal{O}_{\mathfrak{X}}(V) = \bigcap_J \mathcal{O}_{\mathfrak{X}}(U_J).$$

As all the $\mathcal{O}_{\mathfrak{X}}(U_J)$ are subalgebras of $D_0 = \mathbb{k}_{q^{n+1}}[\mathbf{a}^{\pm 1}, \mathbf{b}^{\pm 1}]$, this intersection makes sense.

Now if $V \subseteq U$ are open subsets of \mathfrak{X} then there is an inclusion $\mathcal{O}_{\mathfrak{X}}(U) \subseteq \mathcal{O}_{\mathfrak{X}}(V) \subseteq D_0$ so that $\mathcal{O}_{\mathfrak{X}}(V)$ is an $\mathcal{O}_{\mathfrak{X}}(U)$ -module. These inclusions make $\mathcal{O}_{\mathfrak{X}}$ a sheaf of noncommutative algebras on \mathfrak{X} , and we will refer to \mathfrak{X} as a *noncommutative ringed space*. If we need to keep track of q , we will refer to \mathfrak{X} as \mathfrak{X}_q .

Remark 7.1.3. In [Kaw24, Kaw25], Kawamata gives a definition which generalises our construction of \mathfrak{X} : he shows how to construct a *noncommutative scheme* or *NC scheme* from a collection of associative algebras parameterised by a poset. The main result of [Kaw25] is that a tilting bundle F° on a commutative scheme X deforms over an Artin local base to a tilting bundle F on any NC scheme \mathfrak{X} that deforms X , and that $\text{End}_{\mathfrak{X}}(F)$ is derived equivalent to \mathfrak{X} .

If we vary q , then \mathfrak{X}_q is a NC scheme over $\text{Spec } \mathbb{k}[q, q^{-1}]$, in Kawamata's sense. Further, at $q = 1$ we recover the minimal resolution of the Kleinian singularity X_1 , regarded as a toric variety. Thus it is not surprising that $\mathfrak{X}_q \simeq \text{qgr-}T_q$ has a tilting bundle, although note that [Kaw25] only considers formal deformations over an Artin local base.

We make the expected definition of an $\mathcal{O}_{\mathfrak{X}}$ -module: a (left or right) $\mathcal{O}_{\mathfrak{X}}$ -module is a sheaf \mathcal{M} of abelian groups on \mathfrak{X} so that each $\mathcal{M}(U)$ is a (left or right) $\mathcal{O}_{\mathfrak{X}}(U)$ -module and if $V \subseteq U$ are open subsets of \mathfrak{X} then the restriction maps $\mathcal{M}(U) \rightarrow \mathcal{M}(V)$ are compatible with the restriction $\mathcal{O}_{\mathfrak{X}}(U) \subseteq \mathcal{O}_{\mathfrak{X}}(V)$. Let \mathcal{M}, \mathcal{N} be $\mathcal{O}_{\mathfrak{X}}$ -modules. The notion of a morphism $\mathcal{M} \rightarrow \mathcal{N}$ of $\mathcal{O}_{\mathfrak{X}}$ -modules and thus of $\text{Hom}_{\mathfrak{X}}(\mathcal{M}, \mathcal{N})$ is the obvious generalisation of the standard commutative definition; cf. [Har77, p. 104]. The category of left, respectively right, $\mathcal{O}_{\mathfrak{X}}$ -modules is called $\mathcal{O}_{\mathfrak{X}}\text{-Mod}$, respectively $\text{Mod-}\mathcal{O}_{\mathfrak{X}}$. By restriction, each open subset U of \mathfrak{X} is also a noncommutative ringed space and we thus have categories $\mathcal{O}_U\text{-Mod}$ and $\text{Mod-}\mathcal{O}_U$ of \mathcal{O}_U -modules, where \mathcal{O}_U is our notation for $\mathcal{O}_{\mathfrak{X}}|_U$.

Note that $\mathfrak{X} = \bigcup_{i=0}^n U_i$. Concretely, we have

$$U_i = \{0, \mathfrak{l}_i, \mathfrak{l}_{i+1}, \mathfrak{p}_i\} \quad \text{for } 0 \leq i \leq n, \quad U_{-1} = \{0, \mathfrak{l}_0\}, \quad U_{n+1} = \{0, \mathfrak{l}_{n+1}\}.$$

We will see that principal open sets are affine in the appropriate sense, and use them to define (quasi)coherent left and right $\mathcal{O}_{\mathfrak{X}}$ -modules. We will only give definitions on the right as those for the left are the same, *mutatis mutandis*.

Let us consider U_i for $0 \leq i \leq n$. Define a functor $\sim: \text{Mod-}\mathcal{O}_X(U_i) \rightarrow \text{Mod-}\mathcal{O}_U$ as follows: if V is an open subset of U_i and $M \in \text{Mod-}\mathcal{O}_X(U_i)$ then set $\widetilde{M}(V) := M \otimes_{\mathcal{O}_X(U_i)} \mathcal{O}_{U_i}(V)$. Now, \widetilde{M} is certainly a presheaf, but note that U_i has no proper covers: if $\{V_j\}$ is a set of open subsets of U_i with $\bigcup_j V_j = U_i$ then some $V_j = U_i$. This means the glueing and vanishing sheaf conditions are trivial and any presheaf on U_i is automatically a sheaf. In particular, \widetilde{M} is a sheaf.

Fix I and let $U = U_I$ and $C = \mathcal{O}_X(U_I)$. Given C -modules M, N and a C -module homomorphism $\rho: M \rightarrow N$, we define $\widetilde{\rho}: \widetilde{M} \rightarrow \widetilde{N}$ as follows. Let V be an open subset of U . Then $\widetilde{\rho}(V): \widetilde{M}(V) \rightarrow \widetilde{N}(V)$ is given by $1 \otimes \rho$ for left modules and $\rho \otimes 1$ for right modules. Note that the functor \sim is exact and fully faithful, with left inverse given by $\mathcal{M} \mapsto \mathcal{M}(U)$.

Summarising the construction and discussion of \sim we have:

Proposition 7.1.4. *Let U_I be a principal open set. For every right $\mathcal{O}_{\mathfrak{X}}(U_I)$ -module M , the assignment $V \mapsto M \otimes_{\mathcal{O}_{\mathfrak{X}}(U_I)} \mathcal{O}_{\mathfrak{X}}(V)$ defines a sheaf \widetilde{M} of right \mathcal{O}_{U_I} -modules. This assignment induces an equivalence of categories $\text{Mod-}\mathcal{O}_{\mathfrak{X}}(U_I) \rightarrow \text{Mod-}\mathcal{O}_{U_I}$, with quasi-inverse given by $\mathcal{M} \mapsto \mathcal{M}(U_I)$. Analogous results hold on the left. \square*

We say that a (left or right) $\mathcal{O}_{\mathfrak{X}}$ -module \mathcal{M} is *quasicohherent* if for all i there is an $\mathcal{O}_X(U_i)$ -module M_{U_i} so that $\mathcal{M}|_{U_i} \cong \widetilde{M_{U_i}}$, and *coherent* if the M_{U_i} are finitely generated. The full subset

of $\mathcal{O}_{\mathfrak{X}}\text{-Mod}$ of quasicoherent (respectively, coherent) sheaves is denoted $\mathcal{O}_{\mathfrak{X}}\text{-Mod}$ (respectively $\mathcal{O}_{\mathfrak{X}}\text{-mod}$). Likewise on the right we define $\text{Mod-}\mathcal{O}_{\mathfrak{X}}$ and $\text{mod-}\mathcal{O}_{\mathfrak{X}}$. These categories are abelian given the standard notions of sheaf kernel (which is the same as presheaf kernel) and sheaf cokernel of a morphism of sheaves on \mathfrak{X} . Likewise, the image of a morphism of (quasi)coherent $\mathcal{O}_{\mathfrak{X}}$ -modules is (quasi)coherent.

We now define a functor $\pi : \text{Gr-}T \rightarrow \text{Mod-}\mathcal{O}_{\mathfrak{X}}$ as follows. Given a graded T -module M , the sheaf πM is defined by $\pi M(U_i) = M[t_i^{-1}]_0$ for $0 \leq i \leq n$. For any open $V \subseteq U_i$, define $\pi M(V) = \widetilde{M(U_i)}(V)$ as in Proposition 7.1.4. For any other open set V , use the glueing axiom of sheaves to define $\pi M(V)$. By construction, πM is quasicoherent.

A homomorphism $f : M \rightarrow N$ of graded T -modules induces a morphism of $\mathcal{O}_{\mathfrak{X}}$ -modules $\pi M \rightarrow \pi N$ in the obvious way.

Lemma 7.1.5. $\pi M = 0$ if and only if M is T_+ -torsion.

Proof. By Proposition 7.1.4, $\pi M = 0$ if and only if $\pi M(U_i) = 0$ for all $0 \leq i \leq n$. By definition, this is if and only if all $M[t_i^{-1}]_0 = 0$, or if and only if all $M[t_i^{-1}] = 0$ as the rings $T[t_i^{-1}]$ are strongly graded. This is if and only if for all homogeneous $m \in M$ there is $r \in \mathbb{N}$ so that $mt_i^r = 0$ for all $0 \leq i \leq n$. But the chain of ideals $\{(t_0^r, \dots, t_n^r)\}_{r \geq 0}$ is cofinal with $\{T_{\geq r}\}$, so this last condition occurs if and only if M is torsion. \square

Corollary 7.1.6. *There is an induced functor $\bar{\pi} : \text{Qgr-}T \rightarrow \text{Mod-}\mathcal{O}_{\mathfrak{X}}$.* \square

We will show that $\bar{\pi}$ is an equivalence of categories. To construct the inverse functor to $\bar{\pi}$, we need a bit more notation.

First, if \mathcal{M} is a right $\mathcal{O}_{\mathfrak{X}}$ -module and \mathcal{N} is an $\mathcal{O}_{\mathfrak{X}}$ -bimodule, we can define the right $\mathcal{O}_{\mathfrak{X}}$ -module $\mathcal{M} \otimes_{\mathfrak{X}} \mathcal{N}$ in the usual way, as the sheafification of the presheaf given by $\mathcal{M} \otimes_{\mathfrak{X}} \mathcal{N}(U_I) = \mathcal{M}(U_I) \otimes_{\mathcal{O}_{\mathfrak{X}}(U_I)} \mathcal{N}(U_I)$. For any $d \in \mathbb{Z}$, let $\mathcal{O}(d) := \pi T(d)$. This is an invertible $\mathcal{O}_{\mathfrak{X}}$ -bimodule under the obvious convention, as $T(d)$ is an invertible T -bimodule. The inverse of $\mathcal{O}(d)$ is $\mathcal{O}(-d)$. Note that $\{t_0, \dots, t_n\} \subseteq H^0(\mathfrak{X}, \mathcal{O}(1))$. Given a right $\mathcal{O}_{\mathfrak{X}}$ -module \mathcal{M} , let $\mathcal{M}(d) := \mathcal{M} \otimes_{\mathfrak{X}} \mathcal{O}(d)$.

Now, given $\mathcal{M} \in \text{Mod-}\mathcal{O}_{\mathfrak{X}}$, define

$$\Gamma_*(\mathcal{M}) := \bigoplus_{d \in \mathbb{Z}} H^0(\mathfrak{X}, \mathcal{M}(d)),$$

where $H^0(\mathfrak{X}, -)$ is the global section functor. We claim that $\Gamma_*(\mathcal{M})$ is a (graded) right T -module. To see this, let $m \in \Gamma_*(\mathcal{M})_d$ and let $t \in T_e$. Now, m is determined by a compatible choice of

$$m_i \in \mathcal{M}(U_i) \otimes_{\mathcal{O}_{\mathfrak{X}}(U_i)} \mathcal{O}(d)(U_i) = \mathcal{M}(U_i) \otimes_{\mathcal{O}(U_i)} T[t_i^{-1}]_d$$

for $0 \leq i \leq n$. Thus we may write $m_i = b_i \otimes r_i$, where $r_i \in T[t_i^{-1}]_d$. Then $mt = (m_i t)_i$, where $m_i t = b_i \otimes r_i t \in \mathcal{M}(U_i) \otimes_{\mathcal{O}(U_i)} T[t_i^{-1}]_{d+e}$. The compatibility relations on the m_i ensure that the $m_i t$ are also compatible, that is that $mt \in H^0(\mathfrak{X}, \mathcal{M}(d+e))$, as needed.

Before proving that Γ_* induces a quasi-inverse equivalence to $\bar{\pi}$, we need a lemma.

Lemma 7.1.7. (cf. [Har77, Lemma II.5.14]) *Let $\mathcal{M} \in \text{Mod-}\mathcal{O}_{\mathfrak{X}}$, and let $m_i \in \mathcal{M}(U_i)$ for some $0 \leq i \leq n$. Then for some d , $m_i t_i^d \in \mathcal{M}(d)(U_i)$ extends to a global section of $\mathcal{M}(d)$.*

Proof. We will be careful with the restriction maps: for $0 \leq i, j \leq n$, let $\varphi_{ij} : \mathcal{M}(U_i) \rightarrow \mathcal{M}(U_{ij})$ be the restriction. For $j \neq i$, define $m_j = \varphi_{ij}(m_i)$. As $\mathcal{M}(U_{ij}) = \mathcal{M}(U_j)[(t_i t_j^{-1})^{-1}]$, for some $d_j \in \mathbb{N}$ and $n_j \in \mathcal{M}(U_j)$ we have $m_j = n_j (t_i t_j^{-1})^{-d_j}$, so $m_j t_i^{d_j} = n_j t_j^{d_j} \in \mathcal{M}(d_j)(U_j)$. Taking $d = \max_j d_j$ we see that $m_j t_i^d \in \mathcal{M}(d)(U_j)$ for all j . These choices are compatible, so give a global section of $\mathcal{M}(d)$ extending $m_i t_i^d$. \square

Proposition 7.1.8. *Let $\mathcal{M} \in \text{Mod-}\mathcal{O}_{\mathfrak{X}}$. Then $\pi \Gamma_* \mathcal{M} \cong \mathcal{M}$.*

Proof. We first define a map $\psi : \pi \Gamma_*(\mathcal{M}) \rightarrow \mathcal{M}$. By the local nature of a sheaf and by Proposition 7.1.4, it is enough to define ψ on the open sets U_i for $0 \leq i \leq n$: that is, to define

$$\psi_i : \pi \Gamma_*(\mathcal{M})(U_i) \rightarrow \mathcal{M}(U_i)$$

for all $0 \leq i \leq n$. Now, an element of $\pi\Gamma_*(\mathcal{M})(U_i)$ has the form mt_i^{-d} where $d \in \mathbb{Z}$ and $m \in \mathcal{M}(d)(U_i)$. But $t_i^{-d} \in \mathcal{O}(-d)(U_i) = T[t_i^{-1}]_{-d}$, so we may regard mt_i^{-d} as an element of $\mathcal{M}(d) \otimes_{\mathfrak{X}} \mathcal{O}(-d)(U_i)$. Let $\psi_i(m)$ be the image of m under the natural isomorphism

$$\mathcal{M}(d) \otimes_{\mathfrak{X}} \mathcal{O}(-d) \rightarrow \mathcal{M}.$$

Note that $\psi_i(m) \neq 0$ if $m \neq 0$. Thus ψ is injective. That ψ is surjective follows from Lemma 7.1.7. \square

We now use local cohomology to show that $\bar{\pi}$ and Γ_* are quasi-inverse equivalences. Recall that $\pi : \text{Gr-}T \rightarrow \text{Qgr-}T$ is the natural (quotient) functor.

Proposition 7.1.9. *Let $M \in \text{Gr-}T$. Then in $\text{Gr-}T$ we have $\Gamma_*\pi M \cong \underline{H}_{\text{Qgr-}T}^0(\pi M)$, which is isomorphic to M in $\text{Qgr-}T$.*

Proof. That M and $\underline{H}_{\text{Qgr-}T}^0(\pi M)$ are isomorphic in $\text{Qgr-}T$ is Corollary 6.2.6.

To complete the proof, let $\mathcal{M} = \pi M$. It suffices to show that we have an isomorphism $H^0(\mathfrak{X}, \mathcal{M}) \cong \varinjlim \text{Hom}_{\text{Gr-}T}(T_{\geq d}, M)$. Now $H^0(\mathfrak{X}, \mathcal{M})$ may be identified with

$$\left\{ m = (m_i) \in \bigoplus_{i=0}^n M[t_i^{-1}]_0 \mid \varphi_{ij}(m_i) = \varphi_{ji}(m_j) \text{ for all } i, j \right\}, \quad (7.1.10)$$

where $\varphi_{ij} : \mathcal{M}(U_i) \rightarrow \mathcal{M}(U_{ij})$ is restriction.

Let $f \in \text{Hom}_{\text{Gr-}T}(T_{\geq d}, M)$. Then $n_i := f(t_i^d) \in M$ for all i , and $n_i t_i^{-d} = n_j t_j^{-d}$ as elements of $M[t_i^{-1}, t_j^{-1}]_0 = \mathcal{M}(U_{ij})$. Thus the restrictions of $n_i t_i^{-d}$ and $n_j t_j^{-d}$ give the same element of $\mathcal{M}(U_{ij})$, and so the $n_i t_i^{-d}$ glue to give an element of $H^0(\mathfrak{X}, \mathcal{M})$. This element is clearly nonzero if $f \neq 0$. In other words, we have constructed an injective homomorphism $\varinjlim \text{Hom}_{\text{Gr-}T}(T_{\geq d}, M) \rightarrow H^0(\mathfrak{X}, \mathcal{M})$.

To see this map is surjective, let $m = (m_i) \in H^0(\mathfrak{X}, \mathcal{M})$ as in (7.1.10). There are $d \in \mathbb{N}$ and $n_i \in M_d$ so that we can write $m_i = n_i t_i^{-d}$ for all i . This induces a degree 0 homomorphism $\lambda_i : t_i^d T \rightarrow M$ defined by $\lambda_i(t_i^d r) = n_i r$.

The compatibility (7.1.10) between the m_i means that $n_i t_i^{-d} = n_j t_j^{-d} \in M[t_i^{-1}, t_j^{-1}]_0$, or that $n_i t_j^d = q^\bullet n_j t_i^d$, where $t_i^d t_j^d = q^\bullet t_j^d t_i^d$. Thus on $t_i^d T \cap t_j^d T = t_i^d t_j^d T$ we have

$$\lambda_i(t_i^d t_j^d r) = n_i t_j^d r = q^\bullet n_j t_i^d r = \lambda_j(q^\bullet t_j^d t_i^d r) = \lambda_j(t_i^d t_j^d r),$$

and λ_i, λ_j agree. They thus glue to give a well-defined map $\lambda \in \text{Hom}_{\text{Gr-}T}(\sum_j t_j^d T, M)$ defined by $\lambda(\sum_j t_j^d r_j) = \sum_j n_j r_j$. To finish, we just note that $\sum_j t_j^d T \supseteq T_{\geq r}$ for appropriate r . \square

Combining Propositions 7.1.8 and 7.1.9. we have proved:

Theorem 7.1.11. *The functors Γ_* and $\bar{\pi}$ induce quasi-inverse equivalences between $\text{Mod-}\mathcal{O}_{\mathfrak{X}}$ and $\text{Qgr-}T$, which restrict to equivalences between $\text{mod-}\mathcal{O}_{\mathfrak{X}}$ and $\text{qgr-}T$.*

Proof. All that is left is to prove that if $M \in \text{gr-}T$, then the image of $\Gamma_*\pi M$ is in $\text{qgr-}T$. This follows from Proposition 6.3.9, as $M_{\geq \ell}$ is finitely generated for all ℓ . \square

7.2. Finite homological dimension. We now turn to proving that $\text{qgr-}T$ (or equivalently, $\text{mod-}\mathcal{O}_{\mathfrak{X}}$) has finite homological dimension. We will use a version of the standard Grothendieck spectral sequence relating Ext and $\mathcal{E}xt$; parts of the commutative proof go through with little change, but there are some important differences.

As $\text{Qgr-}T \simeq \text{Mod-}\mathcal{O}_{\mathfrak{X}}$, by Lemma 6.2.4 $\text{Mod-}\mathcal{O}_{\mathfrak{X}}$ has enough injectives. We wish to understand injective objects in $\text{Mod-}\mathcal{O}_{\mathfrak{X}}$ in more detail. Here we have:

Lemma 7.2.1. *Let \mathcal{I} be a (right or left) $\mathcal{O}_{\mathfrak{X}}$ -module. Then \mathcal{I} is injective if and only if the stalk $\mathcal{I}_{\mathfrak{p}}$ is an injective $\mathcal{O}_{\mathfrak{X}, \mathfrak{p}}$ -module for all $\mathfrak{p} \in \mathfrak{X}$.*

Before proving Lemma 7.2.1, we need an auxiliary lemma. The stalk $\mathcal{O}_{\mathfrak{X}, \mathfrak{p}}$ is not a local ring, except in the weak sense of being \mathbb{Z}^3 -graded local. Instead we have:

Lemma 7.2.2. *Let $\mathfrak{p} \in \mathfrak{X}$. Then*

$$\mathcal{O}_{\mathfrak{X}, \mathfrak{p}} \cong \begin{cases} \mathbb{k}_{q^\bullet}[u^\pm, v^\pm] & \text{if } \mathfrak{p} = 0, \\ \mathbb{k}_{q^\bullet}[u^\pm, v] & \text{if } \mathfrak{p} = \mathfrak{l}_i, \\ \mathbb{k}_{q^\bullet}[u, v] & \text{if } \mathfrak{p} = \mathfrak{p}_i. \end{cases}$$

Further, there is an open subset $V_{\mathfrak{p}}$ of \mathfrak{X} so that $\mathcal{O}_{\mathfrak{X}, \mathfrak{p}} = \mathcal{O}_{\mathfrak{X}}(V_{\mathfrak{p}})$.

Proof. By definition, the stalk $\mathcal{O}_{\mathfrak{X}, \mathfrak{p}}$ is

$$\varinjlim_{\{V | \mathfrak{p} \in V, V \text{ open}\}} \mathcal{O}_{\mathfrak{X}}(V) = \sum_V \mathcal{O}_{\mathfrak{X}}(V).$$

(Here recall that each $\mathcal{O}_{\mathfrak{X}}(V)$ is a subalgebra of $\mathbb{k}_{q^{n+1}}[\mathbf{a}^\pm, \mathbf{b}^\pm]$ with the restriction maps all being inclusions.) But as \mathfrak{X} is a finite space, the intersection

$$V_{\mathfrak{p}} := \bigcap \{V \mid \mathfrak{p} \in V, V \text{ open}\}$$

is itself open, so $\mathcal{O}_{\mathfrak{X}, \mathfrak{p}} = \mathcal{O}_{\mathfrak{X}}(V_{\mathfrak{p}})$. Further, we have

$$\begin{aligned} V_0 &= \{0\}, \\ V_{\mathfrak{l}_i} &= \{\mathfrak{l}_i, 0\}, \\ V_{\mathfrak{p}_i} &= \{\mathfrak{p}_i, \mathfrak{l}_i, \mathfrak{l}_{i+1}, 0\}, \end{aligned}$$

and the result follows, using the computations in Lemma 6.4.1. \square

Proof of Lemma 7.2.1. (\Leftarrow) Let \mathcal{M}, \mathcal{N} be $\mathcal{O}_{\mathfrak{X}}$ -modules and let $\gamma : \mathcal{M} \rightarrow \mathcal{N}$ and $\iota : \mathcal{M} \rightarrow \mathcal{I}$ be morphisms, where γ is monic. Now, γ being monic means by definition that its sheaf kernel is trivial; that is, that γ is injective on stalks. So for all $\mathfrak{p} \in \mathfrak{X}$ we have $\gamma_{\mathfrak{p}} : \mathcal{M}_{\mathfrak{p}} \hookrightarrow \mathcal{N}_{\mathfrak{p}}$ and $\iota_{\mathfrak{p}} : \mathcal{M}_{\mathfrak{p}} \rightarrow \mathcal{I}_{\mathfrak{p}}$. As \mathcal{I} has injective stalks, $\iota_{\mathfrak{p}}$ extends to create a commutative diagram

$$\begin{array}{ccc} \mathcal{M}_{\mathfrak{p}} & \xrightarrow{\gamma_{\mathfrak{p}}} & \mathcal{N}_{\mathfrak{p}} \\ \iota_{\mathfrak{p}} \downarrow & \swarrow j_{\mathfrak{p}} & \\ \mathcal{I}_{\mathfrak{p}} & & \end{array}$$

The $j_{\mathfrak{p}}$ glue to give a morphism $j : \mathcal{N} \rightarrow \mathcal{I}$ of $\mathcal{O}_{\mathfrak{X}}$ -modules so that

$$\begin{array}{ccc} \mathcal{M} & \xrightarrow{\gamma} & \mathcal{N} \\ \iota \downarrow & \swarrow j & \\ \mathcal{I} & & \end{array}$$

commutes, as needed.

(\Rightarrow) The restriction of an injective sheaf to an open subset is injective, so $\mathcal{I}|_{V_{\mathfrak{p}}}$ is injective for all $\mathfrak{p} \in \mathfrak{X}$. The result follows from Lemma 7.2.2. \square

Remark 7.2.3. Note that by the stalk criterion of Lemma 7.2.1, if $\mathcal{I} \in \text{Mod-}\mathcal{O}_{\mathfrak{X}}$ is injective, then \mathcal{I} is also injective considered as an object of $\mathfrak{Mod-}\mathcal{O}_{\mathfrak{X}}$.

As in the commutative setting, if \mathcal{M}, \mathcal{N} are quasicoherent sheaves on \mathfrak{X} then the presheaf $U \mapsto \text{Hom}_{\mathcal{O}_U}(\mathcal{M}|_U, \mathcal{N}|_U)$ is a sheaf (of abelian groups), which we call $\mathcal{H}om_{\mathfrak{X}}(\mathcal{M}, \mathcal{N})$. As usual, we have $\text{Hom}_{\mathfrak{X}}(\mathcal{M}, \mathcal{N}) = H^0(\mathfrak{X}, \mathcal{H}om_{\mathfrak{X}}(\mathcal{M}, \mathcal{N}))$.

For our Grothendieck spectral sequence, we will need that $\mathcal{H}om_{\mathfrak{X}}(\mathcal{M}, -)$ takes injective objects to Γ -acyclic objects.

Proposition 7.2.4. *Let I be a graded injective T -module and let \mathcal{M} be a quasicoherent $\mathcal{O}_{\mathfrak{X}}$ -module. Then $\mathcal{H}om_{\mathfrak{X}}(\mathcal{M}, \pi I)$ is a flasque sheaf on \mathfrak{X} .*

Proof. Without loss of generality I is graded uniform and T_+ -torsionfree. There is thus a finitely generated graded uniform submodule H of I and a non-irrelevant graded prime ideal \mathfrak{q} of T that is the assassinator of H , by [GW04, Lemma 5.26]. (Note that here we are using the projective \mathbb{Z} -grading of T where the t_i have degree 1 and x, y, z have degree 0.) Let $\mathcal{I} = \pi I$ and $\mathcal{H} = \pi H$.

Let $V \subseteq U$ be open subsets of \mathfrak{X} . We need to show:

$$\mathrm{Hom}_U(\mathcal{M}|_U, \mathcal{I}|_U) \rightarrow \mathrm{Hom}_V(\mathcal{M}|_V, \mathcal{I}|_V) \text{ is surjective,} \quad (7.2.5)$$

where the map is induced by restriction. Writing $Z = \mathfrak{X} \setminus V$, we have $Z = V(\mathfrak{n})$ where \mathfrak{n} is a non-irrelevant \mathbb{Z}^3 -graded ideal of T . We make and prove two claims.

Claim 1: If $\mathfrak{q} \supseteq \mathfrak{n}$ then $\mathcal{I}|_V = 0$.

To see this, it suffices to show that $\mathcal{I}|_{V'} = 0$ where V' is a principal open subset of V . There is a \mathbb{Z}^3 -homogeneous $w \in \mathfrak{n}$ so that w is invertible in $\mathcal{O}_{\mathfrak{X}}(V')$ and, because $w \in \mathfrak{q}$ and $H\mathfrak{q} = 0$, we have $\mathcal{H}(V') = 0$ by the equivalence from Proposition 7.1.4 between $\mathrm{Mod}\text{-}\mathcal{O}_{V'}$ and $\mathrm{Mod}\text{-}\mathcal{O}(V')$. Thus $\mathcal{I}(V') = 0$ as \mathcal{H} is essential in \mathcal{I} , establishing the claim.

Claim 2: If $\mathfrak{q} \not\supseteq \mathfrak{n}$ and if $\mathcal{F} \in \mathrm{Mod}\text{-}\mathcal{O}_{\mathfrak{X}}$ is supported on Z in the sense that

$$Z \supseteq \{\mathfrak{p} \in \mathfrak{X} \mid \mathcal{F}_{\mathfrak{p}} \neq 0\},$$

then $\mathrm{Hom}_U(\mathcal{F}|_U, \mathcal{I}|_U) = 0$.

To establish this, we may assume without loss of generality that U is principal, by the local nature of a sheaf, so we seek to show that $\mathrm{Hom}_U(\mathcal{F}(U), \mathcal{I}(U)) = 0$, using Proposition 7.1.4. Now, \mathfrak{q} corresponds to a prime ideal P of $C = \mathcal{O}_{\mathfrak{X}}(U)$, which is the assassinator of the essential submodule $\mathcal{H}(U)$ of $\mathcal{I}(U)$, and \mathfrak{n} corresponds to a \mathbb{Z}^2 -graded ideal N of C with $P \not\supseteq N$. Also, $\mathcal{F}(U)$ is w -torsion for any \mathbb{Z}^2 -homogeneous $w \in N$. Choose some such $w \in N \setminus P$ and let $\gamma \in \mathrm{Hom}_C(\mathcal{F}(U), \mathcal{I}(U)) = \mathrm{Hom}_U(\mathcal{F}(U), \mathcal{I}(U))$. If $\gamma \neq 0$ then P is the annihilator of the nonzero module $\mathrm{im} \gamma \cap \mathcal{H}(U)$. This contradicts the fact that $\mathrm{im} \gamma$ is also w -torsion and $w \notin P$, and thus $\mathrm{Hom}_U(\mathcal{F}|_U, \mathcal{I}|_U) = 0$, as claimed.

We return to the proof of the proposition. We must establish (7.2.5). By Claim 1, we may assume without loss of generality that $\mathfrak{n} \not\supseteq \mathfrak{q}$. We will show that (7.2.5) is an equality.

Let $j : V \rightarrow U$ be the inclusion and let \mathcal{K} be the kernel of the natural map $\mathcal{I}|_U \rightarrow j_*(\mathcal{I}|_V)$. Now \mathcal{K} is supported on Z by definition and by Claim 2 the inclusion $\mathcal{K} \subseteq \mathcal{I}|_U$ must be the zero map; that is, $\mathcal{K} = 0$. So $\mathcal{I}|_U \subseteq j_*(\mathcal{I}|_V)$ and as $\mathcal{I}|_U$ is injective the two must be equal.

Now consider the equivalent restriction map $\varphi : \mathcal{M}|_U \rightarrow j_*(\mathcal{M}|_V)$ for \mathcal{M} . Let $\mathcal{A} = \ker \varphi$ and $\mathcal{B} = \mathrm{coker} \varphi$. Both are supported on Z and so, by Claim 2, $\mathrm{Hom}_U(\mathcal{A}, \mathcal{I}|_U) = \mathrm{Hom}_U(\mathcal{B}, \mathcal{I}|_U) = 0$. Using injectivity of $\mathcal{I}|_U$ we see that

$$\mathrm{Hom}_U(\mathcal{M}|_U, \mathcal{I}|_U) = \mathrm{Hom}_U(j_*(\mathcal{M}|_V), \mathcal{I}|_U) = \mathrm{Hom}_U(j_*(\mathcal{M}|_V), j_*(\mathcal{I}|_V)).$$

This last is easily seen to be equal to $\mathrm{Hom}_V(\mathcal{M}|_V, \mathcal{I}|_V)$. \square

As $\mathrm{Mod}\text{-}\mathfrak{X}$ has enough injectives, we may construct the right derived functors of $\mathrm{Hom}_{\mathfrak{X}}(\mathcal{M}, -)$ and $\mathrm{Hom}_{\mathfrak{X}}(\mathcal{M}, -)$, which we call $\mathrm{Ext}_{\mathfrak{X}}^i(\mathcal{M}, -)$ and $\mathcal{E}xt_{\mathfrak{X}}^i(\mathcal{M}, -)$ respectively. We thus define the groups $\mathrm{Ext}_{\mathfrak{X}}^i(\mathcal{M}, \mathcal{N})$ and the sheaves $\mathcal{E}xt_{\mathfrak{X}}^i(\mathcal{M}, \mathcal{N})$. Note that we can do this entirely with resolutions using injectives of the form πI for I a graded-injective T -module, by Lemma 6.2.4 and the equivalence of categories Theorem 7.1.11.

Remark 7.2.6. Note that $\mathfrak{Mod}\text{-}\mathfrak{X}$ also has enough injectives, using the construction of [Har77, Proposition III.2.2]. Thus we can derive $\mathrm{Hom}_{\mathfrak{X}}(\mathcal{M}, -)$ either as a functor from $\mathfrak{Mod}\text{-}\mathfrak{X} \rightarrow \mathrm{Ab}$ or as a functor from $\mathrm{Mod}\text{-}\mathfrak{X} \rightarrow \mathrm{Ab}$. By Remark 7.2.3 the two agree on $\mathrm{Mod}\text{-}\mathfrak{X}$, and similarly for $\mathcal{H}om$.

If we regard \mathcal{M} simply as a sheaf of abelian groups on the topological space \mathfrak{X} then we have the usual sheaf cohomology $H^i(\mathfrak{X}, \mathcal{M})$ by deriving the global section functor $\Gamma = H^0(\mathfrak{X}, -)$. Of course, $\Gamma(\mathcal{M}) = H^0(\mathfrak{X}, \mathcal{M}) = \mathrm{Hom}_{\mathfrak{X}}(\mathcal{O}_{\mathfrak{X}}, \mathcal{M})$. As injective objects in $\mathrm{Mod}\text{-}\mathfrak{X}$ are flasque, by Proposition 7.2.4 with $\mathcal{M} = \mathcal{O}_{\mathfrak{X}}$, they are Γ -acyclic by [Har77, Proposition III.2.5]. We can thus compute $H^i(\mathfrak{X}, \mathcal{M})$ by taking an injective resolution of \mathcal{M} in $\mathrm{Mod}\text{-}\mathfrak{X}$: in other words,

$$H^i(\mathfrak{X}, \mathcal{M}) = \mathrm{Ext}_{\mathfrak{X}}^i(\mathcal{O}_{\mathfrak{X}}, \mathcal{M}).$$

We can now prove our desired nonsingularity result for \mathfrak{X} .

Theorem 7.2.7. *The categories $\text{Qgr-}T$ and $\text{Mod-}\mathfrak{X}$ have finite homological dimension. In particular, if $s > 4$ then $\text{Ext}_{\mathfrak{X}}^s(\mathcal{F}, \mathcal{G}) = 0$ for all quasicoherent sheaves \mathcal{F}, \mathcal{G} on \mathfrak{X} .*

Proof. This is a standard Grothendieck spectral sequence. We have

$$\text{Hom}_{\mathfrak{X}}(\mathcal{F}, \mathcal{G}) = \Gamma(\mathcal{H}om_{\mathfrak{X}}(\mathcal{F}, \mathcal{G})).$$

By Proposition 7.2.4, $\mathcal{H}om_{\mathfrak{X}}(\mathcal{F}, -)$ sends injective objects in $\text{Mod-}\mathfrak{X}$ to flasque, and thus Γ -acyclic, sheaves. By [Wei94, Theorem 5.8.3], there is a convergent spectral sequence

$$H^p(\mathfrak{X}, \text{Ext}_{\mathfrak{X}}^r(\mathcal{F}, \mathcal{G})) \Rightarrow \text{Ext}_{\mathfrak{X}}^{p+r}(\mathcal{F}, \mathcal{G}). \quad (7.2.8)$$

Now, by Grothendieck vanishing [Har77, Theorem III.2.7] and our observation that $\dim \mathfrak{X} = 2$, we have $H^p(\mathfrak{X}, \mathcal{E}) = 0$ for any $p \geq 3$ and any sheaf \mathcal{E} of abelian groups on \mathfrak{X} , and so in particular for $\mathcal{E} = \text{Ext}_{\mathfrak{X}}^r(\mathcal{F}, \mathcal{G})$. And since by Lemma 6.4.1 $\mathcal{O}_{\mathfrak{X}}(U_j)$ has global dimension 2 for all $0 \leq j \leq n$, for $r \geq 3$ we have $\text{Ext}_{\mathfrak{X}}^r(\mathcal{F}, \mathcal{G})(U_j) = 0$ for all $0 \leq j \leq n$ and thus $\text{Ext}_{\mathfrak{X}}^r(\mathcal{F}, \mathcal{G}) = 0$. The result follows. \square

Theorem 7.2.7 and Proposition 6.2.2 establish that the morphism $\phi = (\phi^*, \phi_*) : \text{Qgr-}T \rightarrow \text{Mod-}B$ can legitimately be considered a resolution of singularities of B . It turns out that ϕ is a weakly crepant categorical resolution of B in the sense discussed in Section 2.3. This can be proved directly, but it is easier to first prove that $\text{qgr-}T$ and $\text{mod-}\Lambda$ are derived equivalent and then use the corresponding facts about $\psi : \text{Mod-}\Lambda \rightarrow \text{mod-}B$, proven in Proposition 3.3.3.

8. A DERIVED EQUIVALENCE OF RESOLUTIONS

In this section, we give our noncommutative version of the derived geometric McKay correspondence: that is, we show that $\text{qgr-}T$ is derived equivalent to $\text{mod-}\Lambda$. We will do this by constructing a tilting object in $\text{qgr-}T$. We then use this derived equivalence to show that $\text{qgr-}T$ is also a weakly crepant categorical resolution of $\text{mod-}B$.

8.1. The modules of interest and their first properties. We begin by defining some modules that will play an important role for us.

Definition 8.1.1. For $1 \leq i \leq n$, define

$$\begin{aligned} \mathcal{M}^{(i)} &:= xT + z^i T, \\ \mathcal{N}^{(i)} &:= \sum_{j=1}^n z^{-m_{ij}} t_j T, \quad \text{where } m_{ij} := \min\{i, j\}, \end{aligned}$$

viewed as objects in $\text{gr-}T$ or $\text{qgr-}T$. Note that, since $\mathcal{M}^{(i)}$ and $\mathcal{N}^{(i)}$ are generated by normal elements, they are both graded T -bimodules. It will also be convenient to write $\mathcal{M}^{(0)} = T = \mathcal{N}^{(0)}$ and to set

$$\mathcal{M} := \bigoplus_{i=0}^n \mathcal{M}^{(i)}.$$

Our first goal is to show that the modules $\mathcal{M}^{(i)}$ and $\mathcal{N}^{(i)}$ are mutually inverse in $\text{qgr-}T$, in an appropriate sense. We begin by proving some local behaviour of these modules.

Lemma 8.1.2. *We have*

$$\mathcal{M}^{(i)}[t_k^{-1}]_0 = \begin{cases} x \mathcal{O}(U_k) & \text{if } 0 \leq k \leq i-1, \\ z^i \mathcal{O}(U_k) & \text{if } i \leq k \leq n. \end{cases}$$

Proof. It is clear that $\mathcal{M}^{(i)}[t_k^{-1}]_0 = x \mathcal{O}(U_k) + z^i \mathcal{O}(U_k)$. If $0 \leq k \leq i-1$, then

$$z^i = z^{k+1} t_k t_k^{-1} z^{i-k-1} = q^\bullet x t_{k+1} t_k^{-1} z^{i-k-1} \in x \mathcal{O}(U_k).$$

If instead $i \leq k \leq n$, we have

$$x = xt_k t_k^{-1} = q^\bullet z^k t_{k-1} t_k^{-1} = q^\bullet z^i z^{k-i} t_{k-1} t_k^{-1} \in z^i \mathcal{O}(U_k).$$

In either case, the result now follows. \square

Lemma 8.1.3. *We have*

$$\mathcal{N}^{(i)}[t_k^{-1}]_0 = \begin{cases} x^{-1} \mathcal{O}(U_k) & \text{if } 0 \leq k \leq i-1, \\ z^{-i} \mathcal{O}(U_k) & \text{if } i \leq k \leq n. \end{cases}$$

Proof. As in the previous lemma, it is clear that $\mathcal{N}^{(i)}[t_k^{-1}]_0 = \sum_{j=1}^n z^{-m_{ij}} t_j t_k^{-1} \mathcal{O}(U_k)$. First let $0 \leq k \leq i-1$, so that $m_{i,k+1} = k+1$. Then

$$x^{-1} = q^\bullet z^{-(k+1)} t_{k+1} t_k^{-1} = q^\bullet z^{-m_{i,k+1}} t_{k+1} t_k^{-1} \in \mathcal{N}^{(i)}[t_k^{-1}]_0.$$

Conversely, for $1 \leq j \leq n$ we have

$$z^{-m_{ij}} t_j t_k^{-1} = q^\bullet x^{-1} z^{-m_{ij}} x t_j t_k^{-1} = q^\bullet x^{-1} z^{j-m_{ij}} t_{j-1} t_k^{-1} \in x^{-1} \mathcal{O}(U_k),$$

so that $\mathcal{N}^{(i)}[t_k^{-1}]_0 = x^{-1} \mathcal{O}(U_k)$.

If instead $i \leq k \leq n$, then $m_{ik} = i$, so that

$$z^{-i} = z^{-m_{ik}} t_k t_k^{-1} \in \mathcal{N}^{(i)}[t_k^{-1}]_0.$$

On the other hand, for $1 \leq j \leq n$ we have

$$z^{-m_{ij}} t_j t_k^{-1} = q^\bullet z^{-i} z^{i-m_{ij}} t_j t_k^{-1} \in z^{-i} \mathcal{O}(U_k),$$

from which it follows that $\mathcal{N}^{(i)}[t_k^{-1}]_0 = z^{-i} \mathcal{O}(U_k)$. \square

Corollary 8.1.4. *For $1 \leq i \leq n$, $\mathcal{M}^{(i)}$ and $\mathcal{N}^{(i)}$ are mutually inverse in $\text{qgr-}T$ in the sense that $\mathcal{M}^{(i)} \otimes_T \mathcal{N}^{(i)} \cong \mathcal{N}^{(i)} \otimes_T \mathcal{M}^{(i)} \cong T$ in $\text{qgr-}T$. As a result, $-\otimes_T \mathcal{M}^{(i)}$ and $-\otimes_T \mathcal{N}^{(i)}$ are (quasi-inverse) autoequivalences of $\text{qgr-}T$, and similarly on the left.*

Proof. Since there is an equivalence $\text{qgr-}T \simeq \text{mod-}\mathcal{O}_{\mathfrak{X}}$, it suffices to prove these isomorphisms locally; this follows quickly from Lemmata 8.1.2 and 8.1.3. \square

We next aim to show that the tensor product

$$\mathcal{M}_1 \otimes_T \cdots \otimes_T \mathcal{M}_n$$

is isomorphic to $T(1)$ in $\text{qgr-}T$; geometrically, this means that this object is “ample”. We first require an easy lemma:

Lemma 8.1.5. *For $0 \leq k \leq n$, we have $T[t_k^{-1}]_1 = t_k \mathcal{O}(U_k)$.*

Proof. It is clear that $t_k \mathcal{O}(U_k) \subseteq T[t_k^{-1}]_1$. For the reverse inclusion, it suffices to show that $t_0, \dots, t_n \in t_k \mathcal{O}(U_k)$, but this is clear, since $t_i = t_k(t_k^{-1} t_i) \in t_k \mathcal{O}(U_k)$ for $0 \leq i \leq n$. \square

Recall that the graded quotient ring of T (in the projective grading) is

$$Q_{\text{gr}}(T) = Q_{\text{gr}}^{\mathbb{Z}}(T) = T[h^{-1} \mid 0 \neq h \in T_d \text{ for some } d \in \mathbb{N}].$$

Lemma 8.1.6. *The graded module homomorphism*

$$\beta : \mathcal{M}^{(1)} \otimes_T \mathcal{M}^{(2)} \otimes_T \cdots \otimes_T \mathcal{M}^{(n)} \rightarrow Q_{\text{gr}}(T)(1), \quad \beta(m_1 \otimes \cdots \otimes m_n) = m_1 m_2 \cdots m_n t_0 x^{-n}$$

has image in $T(1)$.

Proof. It suffices to show that the generators of the domain of β get sent to an element of T . Suppose that $m_1 \otimes \cdots \otimes m_n$ is such that each m_j is either x or z^j , and suppose that exactly $n-i$ of m_1, \dots, m_n are equal to x . Then the product of the remaining m_j is equal to z^r for some r , where necessarily $r \geq 1 + 2 + \cdots + (i-1) + i = \frac{1}{2}i(i+1)$. Then

$$\begin{aligned} \beta(m_1 \otimes \cdots \otimes m_n) &= q^\bullet z^r x^{n-i} t_0 x^{-n} \\ &= q^\bullet z^{r - \frac{1}{2}i(i+1)} (z^i x^{-1}) (z^{i-1} x^{-1}) \cdots (z^2 x^{-1}) (z^1 x^{-1}) t_0 \end{aligned}$$

$$\begin{aligned}
&= q^\bullet z^{r-\frac{1}{2}i(i+1)} (z^i x^{-1}) (z^{i-1} x^{-1}) \dots (z^2 x^{-1}) t_1 \\
&= q^\bullet z^{r-\frac{1}{2}i(i+1)} (z^i x^{-1}) (z^{i-1} x^{-1}) \dots (z^3 x^{-1}) t_2 \\
&\quad \vdots \\
&= q^\bullet z^{r-\frac{1}{2}i(i+1)} z^i x^{-1} t_{i-1} \\
&= q^\bullet z^{r-\frac{1}{2}i(i+1)} t_i,
\end{aligned}$$

which lies in T . □

Proposition 8.1.7. *In $\text{qgr-}T$, we have*

$$\mathcal{M}^{(1)} \otimes_T \mathcal{M}^{(2)} \otimes_T \dots \otimes_T \mathcal{M}^{(n)} \cong T(1).$$

Proof. Let β be as in Lemma 8.1.6 and fix $0 \leq k \leq n$. From the equivalence $\text{qgr-}T \simeq \text{mod-}\mathcal{O}_{\mathfrak{X}}$ it suffices to show that β is locally an isomorphism onto $T(1)$. Localising β at t_k , by Lemma 8.1.2 in degree 0 we obtain

$$\tilde{\beta} : \underbrace{\left(z \mathcal{O}(U_k) \otimes z^2 \mathcal{O}(U_k) \otimes \dots \otimes z^k \mathcal{O}(U_k) \right)}_{k \text{ terms}} \otimes \underbrace{\left(x \mathcal{O}(U_k) \otimes \dots \otimes x \mathcal{O}(U_k) \right)}_{n-k \text{ terms}} \rightarrow t_k \mathcal{O}(U_k),$$

where the tensor products are over $\mathcal{O}(U_k)$. The domain of this map is free of rank 1, with generator $z^{\frac{1}{2}k(k+1)} x^{n-k}$, and we can identify it with the map

$$\tilde{\beta} : z^{\frac{1}{2}k(k+1)} x^{n-k} \mathcal{O}(U_k) \rightarrow t_k \mathcal{O}(U_k), \quad m \mapsto t_0 x^{-n} m.$$

By the proof of Lemma 8.1.6, we have

$$\tilde{\beta} \left(z^{\frac{1}{2}k(k+1)} x^{n-k} \right) = t_0 x^{-n} z^{\frac{1}{2}k(k+1)} x^{n-k} = q^\bullet (z^k x^{-1}) (z^{k-1} x^{-1}) \dots (z x^{-1}) t_0 = q^\bullet t_k,$$

which shows that $\tilde{\beta}$ is an isomorphism. Thus β is locally an isomorphism, and hence an isomorphism in $\text{qgr-}T$. □

We end this section by identifying some short exact sequences of $\text{qgr-}T$ -modules in which the modules $\mathcal{M}^{(i)}$ and $\mathcal{N}^{(i)}$ sit.

Proposition 8.1.8. *Write*

$$\gamma : T \rightarrow T, \quad \gamma(t) = x t x^{-1},$$

which is an automorphism of T , and which restricts to an automorphism of any right (or left) ideal of T . Then, for any $1 \leq i, j \leq n$, the sequence

$$0 \rightarrow T \xrightarrow{t \mapsto \begin{pmatrix} \gamma^{-1}(z^i) t \\ -x t \end{pmatrix}} \mathcal{M}^{(i)} \oplus \mathcal{M}^{(j)} \xrightarrow{\begin{pmatrix} m \\ m' \end{pmatrix} \mapsto \gamma(m) \otimes x + z^i \otimes m'} \mathcal{M}^{(i)} \otimes_T \mathcal{M}^{(j)} \rightarrow 0 \quad (8.1.9)$$

is exact in $\text{qgr-}T$.

Proof. It is not immediately obvious, but nevertheless straightforward to check, that the second map in (8.1.9) is T -linear and that this sequence forms a complex. It remains to verify exactness, and it suffices to check this locally by the equivalence $\text{qgr-}T \simeq \text{mod-}\mathcal{O}_{\mathfrak{X}}$. Assume first that $i \leq j$. Localising at t_k , we obtain the following sequences:

$$\begin{aligned}
0 &\rightarrow \mathcal{O}(U_k) \rightarrow x \mathcal{O}(U_k) \oplus x \mathcal{O}(U_k) \rightarrow x^2 \mathcal{O}(U_k) \rightarrow 0 && \text{if } 0 \leq k \leq i-1 \\
0 &\rightarrow \mathcal{O}(U_k) \rightarrow z^i \mathcal{O}(U_k) \oplus x \mathcal{O}(U_k) \rightarrow x z^i \mathcal{O}(U_k) \rightarrow 0 && \text{if } i \leq k \leq j-1 \\
0 &\rightarrow \mathcal{O}(U_k) \rightarrow z^i \mathcal{O}(U_k) \oplus z^j \mathcal{O}(U_k) \rightarrow z^{i+j} \mathcal{O}(U_k) \rightarrow 0 && \text{if } j \leq k \leq n,
\end{aligned}$$

each of which are exact. We obtain similar exact sequences when $i > j$. The result now follows. □

Corollary 8.1.10. *For any $1 \leq i, j \leq n$, the sequence*

$$0 \rightarrow \mathcal{N}^{(j)} \xrightarrow{n \mapsto \begin{pmatrix} \gamma^{-1}(z^i) \otimes n \\ -xn \end{pmatrix}} (\mathcal{M}^{(i)} \otimes_T \mathcal{N}^{(j)}) \oplus T \xrightarrow{\begin{pmatrix} m \otimes n \\ r \end{pmatrix} \mapsto \gamma(m)xn + z^i r} \mathcal{M}^{(i)} \rightarrow 0 \quad (8.1.11)$$

is exact in $\text{qgr-}T$. In particular, for any $1 \leq i \leq n$, the sequence

$$0 \rightarrow \mathcal{N}^{(i)} \xrightarrow{\begin{pmatrix} \gamma^{-1}(z^i) \\ -x \end{pmatrix}} T^{\oplus 2} \xrightarrow{\begin{pmatrix} x & z^i \end{pmatrix}} \mathcal{M}^{(i)} \rightarrow 0 \quad (8.1.12)$$

is exact in $\text{qgr-}T$, where the maps are given by multiplication from the left by the indicated matrices.

Proof. There is a commutative diagram

$$\begin{array}{ccccccc} 0 \rightarrow T \otimes \mathcal{N}^{(j)} & \xrightarrow{t \otimes n \mapsto \begin{pmatrix} \gamma^{-1}(x^i) t \otimes n \\ -xt \otimes n \end{pmatrix}} & (\mathcal{M}^{(i)} \otimes \mathcal{N}^{(j)}) \oplus (\mathcal{M}^{(j)} \otimes \mathcal{N}^{(j)}) & \xrightarrow{\begin{pmatrix} m \otimes n \\ m' \otimes n' \end{pmatrix} \mapsto \gamma(m) \otimes x \otimes n + z^i \otimes m' \otimes n'} & \mathcal{M}^{(i)} \otimes \mathcal{M}^{(j)} \otimes \mathcal{N}^{(j)} & \rightarrow 0 \\ \downarrow t \otimes n \mapsto tn & & \downarrow \begin{pmatrix} m \otimes n \\ m' \otimes n' \end{pmatrix} \mapsto \begin{pmatrix} m \otimes n \\ m' \otimes n' \end{pmatrix} & & \downarrow m \otimes m' \otimes n \mapsto m(m' n') & \\ 0 \rightarrow \mathcal{N}^{(j)} & \xrightarrow{n \mapsto \begin{pmatrix} \gamma^{-1}(z^i) \otimes n \\ -xn \end{pmatrix}} & (\mathcal{M}^{(i)} \otimes \mathcal{N}^{(j)}) \oplus T & \xrightarrow{\begin{pmatrix} m \otimes n \\ t \end{pmatrix} \mapsto \gamma(m)xn + z^i r} & \mathcal{M}^{(i)} & \rightarrow 0 \end{array}$$

where the top row is obtained by applying the exact functor $- \otimes_T \mathcal{N}^{(j)}$ to (8.1.9), and hence is exact, and whose vertical arrows are isomorphisms. This implies exactness of the bottom row, which is precisely (8.1.11). Finally, exactness of (8.1.12) follows by specialising to the case $j = i$. \square

8.2. \mathcal{M} is a tilting object. Recall our notation

$$M := \bigoplus_{i=0}^n M^{(i)} \quad \text{and} \quad \mathcal{M} := \bigoplus_{i=0}^n \mathcal{M}^{(i)}$$

from Sections 3.1 and 8.1. The goal of this subsection is to determine $\text{Ext}_{\text{Qgr-}T}^i(\mathcal{M}, \mathcal{M})$ for all $i \geq 1$. We first consider the case $i = 0$, i.e. we determine the endomorphism ring of \mathcal{M} . We begin with some preparatory lemmata.

Lemma 8.2.1. *As right T -modules, $\mathcal{M}^{(i)} \cong (M^{(i)} \otimes_B T) / \text{tors}(M^{(i)} \otimes_B T)$. In particular, there is a short exact sequence of T -modules*

$$0 \rightarrow \text{tors}(M^{(i)} \otimes_B T) \rightarrow M^{(i)} \otimes_B T \rightarrow \mathcal{M}^{(i)} \rightarrow 0. \quad (8.2.2)$$

Proof. We show that

$$M^{(i)} \otimes_B T[t_j^{-1}]_0 \cong \mathcal{M}^{(i)}[t_j^{-1}]_0, \quad (8.2.3)$$

and the statement will then follow from the category equivalence of Theorem 7.1.11. To see this, first note from Lemma 3.1.6 that $M^{(i)}$ is freely presented via the exact sequence

$$B^2 \xrightarrow{\begin{pmatrix} q^{i(n+1)} z^i & -y \\ -x & q^{-\binom{n+1}{2}} z^{n+1-i} \end{pmatrix}} B^2 \xrightarrow{\begin{pmatrix} x & z^i \end{pmatrix}} M^{(i)} \rightarrow 0.$$

Applying the right exact functor $- \otimes_B T[t_j^{-1}]_0$ gives a commutative diagram:

$$\begin{array}{ccccccc} B^2 \otimes_B T[t_j^{-1}]_0 & \longrightarrow & B^2 \otimes_B T[t_j^{-1}]_0 & \longrightarrow & M^{(i)} \otimes T[t_j^{-1}]_0 & \longrightarrow & 0 \\ \cong \downarrow & & \cong \downarrow & & \downarrow & & \\ (T[t_j^{-1}]_0)^2 & \longrightarrow & (T[t_j^{-1}]_0)^2 & \longrightarrow & X & \longrightarrow & 0 \end{array}$$

where the top row is exact and the module X is a suitable cokernel. The bottom row is simply a free presentation of $\mathcal{M}^{(i)}[t_j^{-1}]_0$, and the dashed arrow exists and is an isomorphism by the Five Lemma. Thus $M^{(i)} \otimes T[t_j^{-1}]_0 \cong \mathcal{M}^{(i)}[t_j^{-1}]_0$, as desired. \square

Lemma 8.2.4. *We have $Q_{\text{gr}}^{\mathbb{Z}}(T)_0 = Q(B)$.*

Proof. Since $B = T_0$, the inclusion $Q(B) \subseteq Q_{\text{gr}}(T)_0$ is clear, so we consider only the reverse inclusion. It suffices to show that any $t_i t_j^{-1}$ lies in $Q(B)$. We prove this by induction on $|j - i|$, with the case $|j - i| = 0$ being clear. If $|j - i| = 1$, then either $j = i - 1$ or $j = i + 1$. In the former case, the relation $xt_i = q^\bullet z^i t_{i-1}$ implies $t_i t_{i-1}^{-1} = q^\bullet x^{-1} z^i \in Q(B)$. Similarly, in the latter case we have $t_i t_{i+1}^{-1} = q^\bullet y^{-1} z^{n-i} \in Q(B)$.

Now assume that $|j - i| \geq 2$; again, we consider two cases. If $j > i$ then

$$t_i t_j^{-1} = t_i t_{j-2} t_{j-2}^{-1} t_j^{-1} = q^\bullet t_i t_{j-2} z^{-1} t_{j-1}^{-2} = q^\bullet z^{-1} t_i t_{j-1}^{-1} t_{j-2} t_{j-1}^{-1},$$

which lies in $Q(B)$ by the induction hypothesis. Similarly, if $i > j$ then

$$t_i t_j^{-1} = t_i t_{i-2} t_{i-2}^{-1} t_j^{-1} = q^\bullet z t_{i-1}^2 t_{i-2}^{-1} t_j^{-1} = q^\bullet z t_{i-1} t_{i-2}^{-1} t_{i-1} t_j^{-1},$$

which again lies in $Q(B)$ by the induction hypothesis. \square

Lemma 8.2.5. *For $0 \leq i \leq n$ we have:*

- (1) $\text{Hom}_{Q_{\text{gr}}-T}(T, \mathcal{M}^{(i)}) = M^{(i)}$; and
- (2) $H_{T_+}^1(\mathcal{M})_0 = 0$.

Proof. By Proposition 6.2.5(2) it suffices to prove that $H_{T_+}^1(\mathcal{M}^{(i)})_0 = 0$ for $0 \leq i \leq n$. We begin by showing that $H_{T_+}^1(\mathcal{M}^{(i)})_0$ is an $xB + yB$ -torsion B -module. Since local cohomology commutes with localisation (see, for example, [Rot09, Theorem 10.67], the proof of which holds in our situation with minor modifications) we obtain

$$H_{T_+}^1(\mathcal{M}^{(i)})_0[x^{-1}] \cong H_{T_+}^1(T)_0[x^{-1}],$$

since it is clear from the definition of $\mathcal{M}^{(i)}$ that $\mathcal{M}^{(i)}[x^{-1}] = T[x^{-1}]$. The right hand side is 0 by Proposition 6.4.7, so $H_{T_+}^1(\mathcal{M}^{(i)})_0$ has x -torsion and, similarly, it has y -torsion. In particular, $\text{GKdim } H_{T_+}^1(\mathcal{M}^{(i)})_0 = 0$.

Now, since $M_B^{(i)}$ is maximal Cohen-Macaulay and B is Auslander–Gorenstein of dimension 2, $M_B^{(i)}$ is reflexive. By [RSS15, Lemma 4.11(1)], if M' is a B -module satisfying $M^{(i)} \subseteq M' \subseteq Q(B)$ with $\text{GKdim } M'/M^{(i)} = 0$, then $M' = M^{(i)}$. We seek to apply this result when $M' := \text{Hom}_{Q_{\text{gr}}-T}(T, \mathcal{M}^{(i)})$. Note that we have

$$M' \subseteq \text{Hom}_{Q_{\text{gr}}-T}(T, Q_{\text{gr}}^{\mathbb{Z}}(T)) = \text{End}_{Q_{\text{gr}}-T}(Q_{\text{gr}}^{\mathbb{Z}}(T)) = Q_{\text{gr}}^{\mathbb{Z}}(T)_0 = Q(B),$$

where the last equality is Lemma 8.2.4.

We now show that $\text{GKdim } M'/M^{(i)} = 0$. Consider the exact sequence in Proposition 6.2.5(2) with $N = \mathcal{M}^{(i)}$ and take degree 0. Since $\mathcal{M}^{(i)}$ is torsion-free and $\mathcal{M}_0^{(i)} = M^{(i)}$ we obtain an exact sequence

$$0 \rightarrow M^{(i)} \rightarrow M' \rightarrow H_{T_+}^1(\mathcal{M}^{(i)})_0 \rightarrow 0. \quad (8.2.6)$$

Thus

$$\text{GKdim } M'/M^{(i)} \leq 0,$$

by the first paragraph of the proof. The second paragraph implies $M' = M^{(i)}$, establishing the result. \square

Corollary 8.2.7. $\text{Hom}_{\text{Qgr-}T}(T, \mathcal{M}) \cong M$.

Proof. This follows from Lemma 8.2.5 (1). \square

Lemma 8.2.8. *We have $H_{T_+}^i(\mathcal{M}^{(j)}) = 0 = H_{T_+}^i(\mathcal{N}^{(j)})$ for all $i \geq 3$ and all $0 \leq j \leq n$.*

Proof. This follows from Corollary 6.4.8 and Corollary 8.1.4. \square

With these in hand, we are able to determine the desired endomorphism ring.

Proposition 8.2.9. $\text{End}_{\text{Qgr-}T}(\mathcal{M}) \cong \text{End}_B(M) = \Lambda$.

Proof. Since \mathcal{M} is torsion-free, by applying $\text{Hom}_{\text{Gr-}T}(-, \mathcal{M})$ to the sum of the short exact sequences (8.2.2), we obtain

$$\text{End}_{\text{Gr-}T}(\mathcal{M}) \cong \text{Hom}_{\text{Gr-}T}(M \otimes_B T, \mathcal{M}).$$

We then have a chain of isomorphisms

$$\begin{aligned} \text{End}_{\text{Qgr-}T}(\mathcal{M}) &\cong \lim_{n \rightarrow \infty} \text{Hom}_{\text{Gr-}T}(\mathcal{M}_{\geq n}, \mathcal{M}) \\ &\cong \lim_{n \rightarrow \infty} \text{Hom}_{\text{Gr-}T}(M \otimes_B T_{\geq n}, \mathcal{M}) \\ &\cong \lim_{n \rightarrow \infty} \text{Hom}_B(M, \text{Hom}_{\text{Gr-}T}(T_{\geq n}, \mathcal{M})) \\ &\cong \text{Hom}_B(M, \lim_{n \rightarrow \infty} \text{Hom}_{\text{Gr-}T}(T_{\geq n}, \mathcal{M})) \\ &\cong \text{Hom}_B(M, \text{Hom}_{\text{Qgr-}T}(T, \mathcal{M})) \\ &\cong \text{End}_B(M), \end{aligned}$$

where we have used Corollary 8.2.7 and the fact that $\text{Hom}_B(M, -)$ commutes with direct limits since M is finitely presented. \square

Finally, we are able to show that the higher Ext groups of \mathcal{M} in $\text{Qgr-}T$ vanish.

Proposition 8.2.10. *For $s \geq 1$, we have*

$$\text{Ext}_{\text{Qgr-}T}^s(\mathcal{M}, \mathcal{M}) = 0.$$

Proof. We have

$$\begin{aligned} \text{Ext}_{\text{Qgr-}T}^s(\mathcal{M}, \mathcal{M}) &= \bigoplus_{j,k=0}^n \text{Ext}_{\text{Qgr-}T}^s(\mathcal{M}^{(j)}, \mathcal{M}^{(k)}) \\ &= \bigoplus_{j,k=0}^n \text{Ext}_{\text{Qgr-}T}^s(T, \mathcal{M}^{(k)} \otimes_T \mathcal{N}^{(j)}) \\ &= \bigoplus_{j,k=0}^n H_{\text{Qgr-}T}^s(\mathcal{M}^{(k)} \otimes_T \mathcal{N}^{(j)}), \end{aligned}$$

and from (8.1.11) and Proposition 6.4.7 it suffices to prove that $H_{\text{Qgr-}T}^s(\mathcal{M}^{(i)}) = H_{\text{Qgr-}T}^s(\mathcal{N}^{(i)}) = 0$ for $i, s \geq 1$. Taking cohomology of (8.1.12) and using Proposition 6.4.7 we obtain an exact sequence

$$0 \rightarrow H_{\text{Qgr-}T}^0(\mathcal{N}^{(i)}) \rightarrow B^{\oplus 2} \rightarrow H_{\text{Qgr-}T}^0(\mathcal{M}^{(i)}) \rightarrow H_{\text{Qgr-}T}^1(\mathcal{N}^{(i)}) \rightarrow 0 \quad (8.2.11)$$

and an isomorphism

$$H_{\text{Qgr-}T}^1(\mathcal{M}^{(i)}) \cong H_{\text{Qgr-}T}^2(\mathcal{N}^{(i)}).$$

This last is 0 by Corollary 8.1.4 and Lemma 8.2.8, so we consider (8.2.11). Now, $H_{\text{Qgr-}T}^0(\mathcal{M}^{(i)}) = M^{(i)}$, and applying the proof of Proposition 8.2.9 to $\text{Hom}_{\text{Qgr-}T}(\mathcal{M}^{(i)}, T)$ shows that

$$H_{\text{Qgr-}T}^0(\mathcal{N}^{(i)}) \cong \text{Hom}_{\text{Qgr-}T}(\mathcal{M}^{(i)}, T) \cong \text{Hom}_B(M^{(i)}, T).$$

By Lemma 3.1.7, this is $M^{(n+1-i)}$ and so (8.2.11) becomes

$$0 \rightarrow M^{(n+1-i)} \rightarrow B^2 \rightarrow M^{(i)} \rightarrow H_{\text{qgr-}T}^1(\mathcal{N}^{(i)}) \rightarrow 0.$$

The first three terms of this are from the short exact sequence of Lemma 3.1.6 and one can check that the maps between them are also those of Lemma 3.1.6, so $H_{\text{qgr-}T}^1(\mathcal{N}^{(i)}) = 0$. \square

8.3. The proof of the derived equivalence. In this subsection we prove that the two resolutions $\text{qgr-}T$ and $\text{mod-}\Lambda$ of the singularity B are derived equivalent.

Let $\langle \mathcal{M} \rangle$ denote the subcategory of $\text{qgr-}T$ given by the closure of \mathcal{M} under taking finite direct sums, direct summands, and kernels of epimorphisms. (We caution that this is nonstandard notation.)

Proposition 8.3.1.

(1) Let $k \in \mathbb{N}$ and let $i_1, \dots, i_k \in \{0, \dots, n\}$. Then

$$\mathcal{N}^{(i_1)} \otimes_T \dots \otimes_T \mathcal{N}^{(i_k)} \in \langle \mathcal{M} \rangle.$$

(2) If $s > 0$ then $T(-s) \in \langle \mathcal{M} \rangle$.

Proof.

(1) We prove this by induction on k . By (8.1.12), $\mathcal{N}^{(i)} \in \langle \mathcal{M} \rangle$ for $0 \leq i \leq n$. Now suppose that $k \geq 2$, and consider the short exact sequence in $\text{qgr-}T$

$$0 \rightarrow T \rightarrow \mathcal{M}^{(i_{k-1})} \oplus \mathcal{M}^{(i_k)} \rightarrow \mathcal{M}^{(i_{k-1})} \otimes_T \mathcal{M}^{(i_k)} \rightarrow 0$$

from (8.1.9). Applying the functors $\mathcal{N}^{(i_1)} \otimes_T \dots \otimes_T \mathcal{N}^{(i_{k-1})} \otimes_T -$ and $- \otimes_T \mathcal{N}^{(i_k)}$, both of which are exact since the $\mathcal{N}^{(i)}$ are invertible objects in $\text{qgr-}T$, yields an exact sequence

$$\begin{aligned} 0 \rightarrow \mathcal{N}^{(i_1)} \otimes_T \dots \otimes_T \mathcal{N}^{(i_k)} &\rightarrow (\mathcal{N}^{(i_1)} \otimes_T \dots \otimes_T \mathcal{N}^{(i_{k-2})} \otimes_T \mathcal{N}^{(i_k)}) \oplus (\mathcal{N}^{(i_1)} \otimes_T \dots \otimes_T \mathcal{N}^{(i_{k-1})}) \\ &\rightarrow \mathcal{N}^{(i_1)} \otimes_T \dots \otimes_T \mathcal{N}^{(i_{k-2})} \rightarrow 0. \end{aligned}$$

The middle and right-hand terms both lie in $\langle \mathcal{M} \rangle$ by the induction hypothesis, and so the same is true of $\mathcal{N}^{(i_1)} \otimes_T \dots \otimes_T \mathcal{N}^{(i_k)}$.

(2) By Proposition 8.1.7, we have $\mathcal{M}^{(1)} \otimes_T \dots \otimes_T \mathcal{M}^{(n)} \cong T(1)$ in $\text{qgr-}T$, and so $\mathcal{N}^{(n)} \otimes_T \dots \otimes_T \mathcal{N}^{(1)} \cong T(-1)$. Hence in $\text{qgr-}T$ we have

$$(\mathcal{N}^{(n)} \otimes_T \dots \otimes_T \mathcal{N}^{(1)})^{\otimes Ts} \cong T(-s),$$

and this object lies in $\langle \mathcal{M} \rangle$ by part (1). \square

A set of objects S in an abelian category \mathcal{A} is said to *generate* \mathcal{A} if, for every object $A \in \mathcal{A}$, there exist objects $B_1, \dots, B_k \in S$ and an epimorphism

$$B_1 \oplus \dots \oplus B_k \twoheadrightarrow A.$$

Lemma 8.3.2. (cf. [AZ94, Proposition 4.4]) $\text{qgr-}T$ is generated by $\{T(-s) \mid s > 0\}$.

Proof. The proof of [AZ94, Proposition 4.4(1)] works in our situation but, as it is short, we give it here. Let $\mathcal{F} = \pi(F)$ be an object of $\text{qgr-}T$, where $F \in \text{gr-}T$. Then $\mathcal{F} = \pi(F_{\geq 1})$ as $F/F_{\geq 1}$ is torsion. Taking a set of homogeneous generators for $F_{\geq 1}$, we obtain a surjection

$$T(-s_1) \oplus \dots \oplus T(-s_k) \rightarrow F_{\geq 1}$$

of graded right T -modules for some positive integers s_1, \dots, s_k . Since the quotient functor $\pi : \text{gr-}T \rightarrow \text{qgr-}T$ is exact this gives a surjection

$$T(-s_1) \oplus \dots \oplus T(-s_k) \rightarrow \mathcal{F}$$

in $\text{qgr-}T$, as desired. \square

We are now in a position to prove our noncommutative McKay correspondence.

Theorem 8.3.3. \mathcal{M} is a tilting object for $D^b(\text{qgr-}T)$ and the functor $\mathbb{R}\text{Hom}_{\text{Qgr-}T}(\mathcal{M}, -)$ induces a triangle equivalence

$$D^b(\text{qgr-}T) \rightarrow D^b(\text{mod-}\Lambda) = \mathcal{D}(\Lambda).$$

Proof. This is a direct consequence of [Kel07, Proposition 4.6], once we verify the hypotheses of that result. We need to show the following:

- (1) $\text{Qgr-}T$ is locally noetherian and \mathcal{M} is a noetherian object in $\text{Qgr-}T$;
- (2) $\text{Qgr-}T$ has finite homological dimension;
- (3) $\text{Ext}_{\text{Qgr-}T}^i(\mathcal{M}, \mathcal{M}) = 0$ for all $i > 0$;
- (4) $\langle \mathcal{M} \rangle$ contains a set of generators of $\text{Qgr-}T$;
- (5) $\text{End}_{\text{Qgr-}T}(\mathcal{M}) \cong \Lambda$; and
- (6) Λ is noetherian.

As T is noetherian, $\text{Qgr-}T$ is a locally noetherian category, and \mathcal{M} is noetherian since it is a finitely generated T -module, establishing (1). Properties (2) and (3) are proved in Theorem 7.2.7 and Proposition 8.2.10, respectively. Combining Proposition 8.3.1 and Lemma 8.3.2 shows that (4) is satisfied. Finally, (5) is established in Proposition 8.2.9. The result now follows since $\Lambda \cong A \# G$ is noetherian. \square

Remark 8.3.4. It is possible to define a triangulated functor

$$- \otimes_{\Lambda}^{\mathbb{L}} \mathcal{M} : \mathcal{D}^b(\Lambda) \rightarrow D^b(\text{qgr-}T)$$

using \otimes_{Λ} on charts of the noncommutative toric variety \mathfrak{X} and the equivalence of Theorem 7.1.11 between $\text{mod-}\mathcal{O}_{\mathfrak{X}}$ and $\text{qgr-}T$. One can then show that $- \otimes_{\Lambda}^{\mathbb{L}} \mathcal{M}$ is quasi-inverse to the functor $\mathbb{R}\text{Hom}_{\text{Qgr-}T}(\mathcal{M}, -)$ from Theorem 8.3.3. Since we will not need the quasi-inverse explicitly, we omit the details.

Let us now give the promised proof that the map $\phi : \text{qgr-}T \rightarrow \text{mod-}B$ is a weakly crepant categorical resolution.

Recall that $\phi_* = H_{\text{Qgr-}T}^0(-) : \text{qgr-}T \rightarrow \text{mod-}B$ and $\phi^* = \pi(- \otimes_B T) : \text{mod-}B \rightarrow \text{qgr-}T$, and that (ϕ^*, ϕ_*) are an adjoint pair. We will denote the associated right derived functor $D^b(\text{qgr-}T) \rightarrow D(B)$ also by ϕ_* . Likewise, we will use $\phi^* : D(B) \rightarrow D(\text{qgr-}T)$ to denote the triangulated functor associated to the abelian functor ϕ^* , which we note is right exact as a consequence of Proposition 6.2.1. It will be helpful to have more compact notation for the triangle equivalence between $D^b(\text{qgr-}T)$ and $\mathcal{D}(\Lambda)$, so temporarily let Θ be the functor $\mathbb{R}\text{Hom}_{\text{Qgr-}T}(\mathcal{M}, -)$ from Theorem 8.3.3.

The following diagram records the action of these functors:

$$\begin{array}{ccc}
 D(\text{Mod-}\Lambda) & \xleftarrow[\cong]{\Theta} & D(\text{Qgr-}T) \\
 \psi^* \swarrow & & \nearrow \phi^* \\
 & D(\text{Mod-}B) & \\
 \searrow \psi_* & & \swarrow \phi_*
 \end{array} \tag{8.3.5}$$

Note that (8.3.5) is not commutative: for example, $\Theta(\mathcal{M}) = \Lambda \neq \psi^* \phi_*(\mathcal{M}) = \Lambda e_0 \Lambda$. The quotient $\Lambda / \Lambda e_0 \Lambda$ is the type \mathbb{A}_n preprojective algebra, by [Cra18, Theorem 8.2], which has dimension $\binom{n+2}{3}$ and in particular is nonzero.

However, there are some nice relations between these functors, which we give in the next two results.

Lemma 8.3.6. The functors $\Theta \phi^*$ and ψ^* from $D(B) \rightarrow \mathcal{D}(\Lambda)$ agree on $\text{Perf}(B)$.

Proof. We use the criteria of Lemma 3.3.1. First,

$$\Theta \phi^* B = \Theta(\pi T) = \text{Hom}_{\text{qgr-}T}(\mathcal{M}, \pi T) = M^* = \psi^* B,$$

where we have used Proposition 8.2.9 for the isomorphism between M^* and $\text{Hom}_{\text{Qgr-}T}(\mathcal{M}, \pi T)$. If $b \in B$, let $\lambda_b = (b \cdot -)$ denote the corresponding endomorphism of B_B . Note that λ_b also gives an element of $\Lambda = \text{End}_\Lambda(\Lambda)$, from the inclusion $B = e_0 \Lambda e_0 \subseteq \Lambda$. Then

$$\Theta \phi^*(\lambda_b) : \text{Hom}_{\text{Qgr-}T}(\mathcal{M}, \pi T) \rightarrow \text{Hom}_{\text{Qgr-}T}(\mathcal{M}, \pi T)$$

is given by $f \mapsto \lambda_b \circ f$. On the other hand, $\psi^*(\lambda_b) = \lambda_b \otimes 1 : B \otimes_B M^* \rightarrow B \otimes_B M^*$ is also given by left multiplication by b . The result follows by Lemma 3.3.1. \square

Lemma 8.3.7. *The functors ϕ_* and $\psi_* \Theta$ from $D(\text{Qgr-}T) \rightarrow D(R)$ agree on $D^b(\text{qgr-}T)$.*

Proof. Let us first check that these functors agree on \mathcal{M} and on endomorphisms of \mathcal{M} . We have $\phi_* \mathcal{M} = H_{\text{Qgr-}T}^0(\mathcal{M}) = M$, and $\psi_* \Theta \mathcal{M} = \psi_* \Lambda = \Lambda \otimes_\Lambda M = M$. Since $\psi_* \Lambda = M = \phi_* \mathcal{M}$ and $\Lambda = \Theta(\mathcal{M})$, there are homomorphisms

$$\begin{array}{ccc} \Lambda = \text{End}_\Lambda(\Lambda) & \xleftarrow{\Theta} & \text{End}_{\text{Qgr-}T}(\mathcal{M}) \\ & \searrow \psi_* & \swarrow \phi_* \\ & \text{End}_B(M) & \end{array}$$

These maps induce the canonical identifications, where we have used Proposition 8.2.9 to identify $\text{End}_B(M)$ with Λ . It follows that the functors agree on endomorphisms of \mathcal{M} .

Since Λ has finite global dimension, we obtain $\mathcal{D}(\Lambda) = \text{Perf}(\Lambda)$, which is the image under Θ of $\text{Perf}(D(\text{Qgr-}T)) = D^b(\text{qgr-}T)$. Further, $\Lambda = \Theta(\mathcal{M})$. The result follows from applying Lemma 3.3.1 to $\text{Perf}(D(\text{Qgr-}T))$, using that Θ is a triangle equivalence. \square

We can now prove that ϕ is a weakly crepant categorical resolution.

Theorem 8.3.8. *The map $\phi = (\phi^*, \phi_*)$ of noncommutative spaces is a weakly crepant categorical resolution of $D(B)$.*

Proof. We have seen in Proposition 6.2.1 that (ϕ^*, ϕ_*) are an adjoint pair and so do give a map of noncommutative spaces. It follows from Theorem 8.3.3 and Remark 3.2.7 that $D^b(\text{qgr-}T)$ is smooth. To show that ϕ is a categorical resolution, it remains to verify that $\phi_* \phi^* \cong \text{id}_{\text{Perf}(B)}$. This can be checked directly, or using Lemmata 8.3.6 and 8.3.7: we have

$$\phi_* \phi^* \cong \psi_* \Theta \phi^* \cong \psi_* \psi^*$$

as functors from $\text{Perf}(B)$ to $D(B)$. This last is naturally isomorphic to $\text{id}_{\text{Perf}(B)}$ by Proposition 3.3.3.

To show that ϕ is weakly crepant, we need that $\phi^* : \text{Perf}(B) \rightarrow D(\text{Qgr-}T)$ is also a right adjoint to ϕ_* , that is, that there are natural isomorphisms

$$\mathbb{R}\text{Hom}_B(\phi_* \mathcal{F}, N) \cong \mathbb{R}\text{Hom}_{\text{Qgr-}T}(\mathcal{F}, \phi^* N) \quad (8.3.9)$$

for all $N \in \text{Perf}(B)$ and $\mathcal{F} \in D(\text{Qgr-}T)$. But by Proposition 3.3.3 ψ is weakly crepant, and Lemmata 8.3.6 and 8.3.7 show that Θ and quasi-inverse transport the corresponding adjunction to (8.3.9). \square

We believe that $D(\text{Qgr-}T)$ is also a strongly crepant categorical resolution of $\mathcal{D}(B)$, as with $D(\Lambda)$, but again we have not been able to prove this.

9. INTERSECTION THEORY

Recall that, given suitable finiteness on Hom and Ext groups, one may define an “intersection product” on an abelian category \mathbf{C} with enough injectives by

$$\mathcal{A} \cdot \mathcal{B} = \sum_{i \geq 0} (-1)^{i+1} \dim_{\mathbb{k}} \text{Ext}_{\mathbf{C}}^i(\mathcal{A}, \mathcal{B}).$$

(This is, for example, the intersection product defined in [MS01].) We say that $\mathcal{A} \cdot \mathcal{B}$ is *well-defined* if $\dim_{\mathbb{k}} \text{Ext}_{\mathbb{C}}^i(\mathcal{A}, \mathcal{B}) < \infty$ for all i and all but finitely many $\text{Ext}_{\mathbb{C}}^i(\mathcal{A}, \mathcal{B})$ are zero. In this section, we consider the intersection theory of the “exceptional locus” of the map $\phi : \text{qgr-}T \rightarrow \text{mod-}B$ defined in Section 6.2 and show that it consists of a chain of n lines, each with self-intersection -2 : in other words, the classic resolution graph of an \mathbb{A}_n singularity.

In this section, for $1 \leq i \leq n$ let $L_i = T/\mathfrak{l}_i$, where \mathfrak{l}_i is defined in Lemma 7.1.1. We comment briefly on the term “exceptional locus of ϕ ”. Each of the L_i is annihilated by the ideal $\langle x, y, z \rangle$ of B and so is supported on the singularity of B . By Proposition 6.2.2 this is, morally, the locus where $\phi : \text{qgr-}T \rightarrow B$ is not an isomorphism.

Our analysis of the L_i begins with the observation that the L_i are line modules.

Lemma 9.1. *The composition $\mathbb{k}_q \bullet [t_{i-1}, t_i] \rightarrow T \rightarrow L_i$ is an isomorphism of graded rings, and thus L_i has Hilbert series*

$$h_{L_i}(t) = (1 - t)^{-2}. \quad \square$$

Remark 9.2. The homomorphisms $T \rightarrow L_i$ give an inclusion of $\text{qgr-}L_i$ in $\text{qgr-}T$. Since, as is well-known, $\text{qgr-}\mathbb{k}_q \bullet [x, y] \simeq \text{coh } \mathbb{P}^1$ [AVdB90], it is reasonable to think of the L_i as being projective lines inside $\text{qgr-}T$.

The main goal of this section is to prove that the line modules L_i have the intersection theory of an \mathbb{A}_n singularity. More specifically:

Theorem 9.3. *Let $1 \leq i, j \leq n$.*

- (1) *The intersection product $L_i \cdot L_j$ is well-defined.*
- (2) *We have*

$$L_i \cdot L_j = \begin{cases} -2 & \text{if } i = j, \\ 1 & \text{if } |i - j| = 1, \\ 0 & \text{otherwise.} \end{cases}$$

The proof of this result will take up most of the section. Our first step will be to prove that the line ideals \mathfrak{l}_j are invertible in $\text{qgr-}T$, in the sense of Corollary 8.1.4. Our candidate for the inverse of \mathfrak{l}_j will be

$$\mathcal{J}_j = T + z^{-1}t_{j-1}t_jT, \quad 1 \leq j \leq n.$$

(Note that $\mathcal{J}_j \subseteq Q_{\text{gr}}(T)$.) We first require some preparatory lemmata.

Lemma 9.4. *The multiplication map $\mathcal{J}_j \otimes_T \mathfrak{l}_j \rightarrow Q_{\text{gr}}(T)$ has image in T .*

Proof. It suffices to show that multiplication by $z^{-1}t_{j-1}t_j$ maps the generators of \mathfrak{l}_j into T . This is clear for z . Using the relations in T , we have

$$\begin{aligned} z^{-1}t_{j-1}t_j \cdot x &= q^\bullet z^{-1}t_{j-1}z^j t_{j-1} = q^\bullet z^{j-1}t_{j-1}^2, \\ z^{-1}t_{j-1}t_j \cdot y &= q^\bullet z^{-1}t_j z^{n-j+1}t_j = q^\bullet z^{n-j}t_j^2, \end{aligned}$$

both of which lie in T since $1 \leq j \leq n$. Moreover, for $k \neq j, j-1$, we have

$$z^{-1}t_{j-1}t_j \cdot t_k = \begin{cases} q^\bullet z^{-1}t_j z^{k-j}t_j t_{k-1} = q^\bullet z^{k-j-1}t_j^2 t_{k-1} & \text{if } k \geq j+1, \\ q^\bullet z^{-1}t_{j-1}z^{j-k-1}t_{k+1}t_{j-1} = q^\bullet z^{j-k-2}t_{j-1}^2 t_{k+1} & \text{if } k \leq j-2, \end{cases}$$

and this lies in T in both cases. \square

Recall our notation $\mathcal{O}(U_k) = T[t_k^{-1}]_0$.

Lemma 9.5. *We have*

$$\mathfrak{l}_j[t_k^{-1}]_0 = \begin{cases} \mathcal{O}(U_k) & \text{if } k \neq j, j-1, \\ \mathcal{O}(U_k)t_{k+1}t_k^{-1} & \text{if } k = j, \\ \mathcal{O}(U_k)t_{k-1}t_k^{-1} & \text{if } k = j-1, \end{cases}$$

with the understanding that $t_{-1}t_0^{-1} := x$ and $t_nt_{n+1}^{-1} := y$.

We warn the reader that the convention on t_{-1} and t_n in the lemma is slightly different from the convention in Section 7.1.

Proof. We necessarily have $\mathfrak{l}_j[t_k^{-1}]_0 \subseteq \mathcal{O}(U_k)$. If $k \neq j, j-1$, then $1 = t_k t_k^{-1} \in \mathfrak{l}_j[t_k^{-1}]_0$, so we have equality.

Now suppose that $k = j$; in particular, we have $k \geq 1$. It suffices to show that

$$x, y, z, t_\ell t_k^{-1} \in \mathcal{O}(U_k) t_{k+1} t_k^{-1} \quad \text{for } \ell \in \{0, 1, \dots, k-2, k+1, \dots, n\}.$$

We need to consider the cases $k \neq n$ and $k = n$ separately.

First suppose that $k \neq n$. Then, using the relations in T and the fact that $1 \leq k \leq n-1$, we have

$$z = q^\bullet t_{k-1} t_k^{-1} t_{k+1} t_k^{-1} \in \mathcal{O}(U_k) t_{k+1} t_k^{-1}.$$

This also implies that $x, y \in \mathcal{O}(U_k) t_{k+1} t_k^{-1}$, since

$$x = q^\bullet z^k t_{k-1} t_k^{-1} \quad \text{and} \quad y = q^\bullet z^{n-k} t_{k+1} t_k^{-1}$$

again using that $1 \leq k \leq n-1$. To show that $t_\ell t_k^{-1} \in \mathcal{O}(U_k) t_{k+1} t_k^{-1}$, we consider two cases:

$$\begin{aligned} \ell \geq k+1: \quad t_k t_\ell &= q^\bullet z^{\ell-k-1} t_{k+1} t_{\ell-1} \Rightarrow t_\ell t_k^{-1} = q^\bullet z^{\ell-k-1} t_{k+1} t_k^{-1} t_{\ell-1} t_k^{-1}, \\ \ell \leq k-2: \quad t_\ell t_k &= q^\bullet z^{k-\ell-1} t_{\ell+1} t_{k-1} \Rightarrow t_\ell t_k^{-1} = q^\bullet z^{k-\ell-1} t_{\ell+1} t_k^{-1} t_{k-1} t_k^{-1}; \end{aligned}$$

in the first case, this element plainly lies in $\mathcal{O}(U_k) t_{k+1} t_k^{-1}$, while the same is true in the latter case since $z \in \mathcal{O}(U_k) t_{k+1} t_k^{-1}$ and $k-\ell-1 \geq 1$.

Now suppose that $k = n$. We need to show that $\mathfrak{l}_n[t_n^{-1}]_0 = \mathcal{O}(U_n)y$. We have $z = q^\bullet y t_{n-1} t_n^{-1} \in \mathcal{O}(U_n)y$ and

$$x = q^\bullet y^{-1} z^{n+1} = q^\bullet y^n (t_{n-1} t_n^{-1})^{n+1} \in \mathcal{O}(U_n)y.$$

Finally, for $\ell \leq n-2$:

$$t_\ell t_n = q^\bullet z^{n-\ell-1} t_{\ell+1} t_{n-1} \Rightarrow t_\ell t_n^{-1} = q^\bullet z^{n-\ell-1} t_{\ell+1} t_n^{-1} t_{n-1} t_n^{-1} \in \mathcal{O}(U_n),$$

since $z^{n-\ell-1} \in \mathcal{O}(U_n)$.

The case $k = j-1$ is similar, except one must consider the subcases $k = 0$ and $k \neq 0$ separately; we omit the proof. \square

Lemma 9.6. *We have*

$$\mathcal{J}_j[t_i^{-1}]_0 = \begin{cases} \mathcal{O}(U_i) & \text{if } i \neq j, j-1, \\ \mathcal{O}(U_i) t_i t_{i+1}^{-1} & \text{if } i = j, \\ \mathcal{O}(U_i) t_i t_{i-1}^{-1} & \text{if } i = j-1, \end{cases}$$

again with the understanding that $t_{-1} t_0^{-1} := x$ and $t_n t_{n+1}^{-1} := y$.

Proof. First suppose that $i \neq j, j-1$; we need to show that $\mathcal{J}_j[t_i^{-1}]_0 = \mathcal{O}(U_i)$. Since $T \subseteq \mathcal{J}$, the inclusion $\mathcal{O}(U_i) \subseteq \mathcal{J}_j[t_i^{-1}]_0$ is clear. For the reverse inclusion, it suffices to show that $z^{-1} t_{j-1} t_j t_i^{-2} \in \mathcal{O}(U_i)$. Since $i \neq j, j-1$, we have $j-i \geq 2$ or $i-j \geq 1$. If $j-i \geq 2$, then the relations in T give

$$\begin{aligned} t_i t_j &= q^\bullet z^{j-i-1} t_{i+1} t_{j-1} \Rightarrow t_j t_i^{-1} = q^\bullet z^{j-i-1} t_{i+1} t_{j-1} t_i^{-2} \\ &\Rightarrow z^{-1} t_{j-1} t_j t_i^{-2} = q^\bullet z^{j-i-2} t_{j-1}^2 t_{i+1} t_i^{-3} \in \mathcal{O}(U_i). \end{aligned}$$

If instead $i-j \geq 1$, then

$$\begin{aligned} t_{j-1} t_i &= q^\bullet z^{i-j} t_j t_{i-1} \Rightarrow t_{j-1} t_i^{-1} = q^\bullet z^{i-j} t_j t_{i-1} t_i^{-2} \\ &\Rightarrow z^{-1} t_{j-1} t_j t_i^{-2} = q^\bullet z^{i-j-1} t_j^2 t_{i-1} t_i^{-3} \in \mathcal{O}(U_i). \end{aligned}$$

Now suppose $i = j$, so that we need to show $\mathcal{J}_i[t_i^{-1}]_0 = \mathcal{O}(U_i) t_i t_{i+1}^{-1}$. We consider the cases $1 \leq i \leq n-1$ and $i = n$ separately.

First suppose that $1 \leq i \leq n-1$. In this case, the claim essentially follows after observing that

$$1 = t_{i+1}t_i^{-1} \cdot t_it_{i+1}^{-1} \quad \text{and} \quad q^\bullet z^{-1}t_{i-1}t_it_i^{-2} = t_it_{i+1}^{-1}.$$

If instead $i = n$, then we need to show that $\mathcal{J}_i[t_i^{-1}]_0 = \mathcal{O}(U_i)y^{-1}$. This follows since

$$1 = y \cdot y^{-1} \quad \text{and} \quad z^{-1}t_{n-1}t_nt_n^{-2} = z^{-1}t_{n-1}t_n^{-1} = q^\bullet y^{-1}.$$

The case $i = j-1$ is similar and therefore omitted; we remark that the proof in this case requires analysing the subcases $j = 1$ and $2 \leq j \leq n$ separately. \square

We can now establish invertibility of the line ideals \mathfrak{l}_j :

Proposition 9.7. *The \mathfrak{l}_j are invertible T -bimodules in $\text{qgr-}T$, in the sense of Corollary 8.1.4.*

Proof. As before, Theorem 7.1.11 allows us to prove this locally, and this is precisely the content of Lemma 9.6. \square

Our basic strategy to prove Theorem 9.3 is to use the local-to-global spectral sequence (7.2.8). We thus need to compute $\text{Ext}_{\mathfrak{X}}^s(\pi L_i, \pi L_j)$. Since L_i, L_j are T -bimodules, this will be an $\mathcal{O}_{\mathfrak{X}}$ -bimodule. We first compute:

Lemma 9.8. *Identify $\mathcal{O}(U_k)$ with $\mathbb{k}_q^\bullet[t_{k-1}t_k^{-1}, t_{k+1}t_k^{-1}]$, using Lemma 6.4.1(1). Then*

$$(\pi L_i)(U_k) = \begin{cases} \mathbb{k}_q^\bullet[t_{i-2}t_{i-1}^{-1}, t_it_{i-1}^{-1}]/(t_{i-2}t_{i-1}^{-1}) & \text{if } k = i-1, \\ \mathbb{k}_q^\bullet[t_{i-1}t_i^{-1}, t_{i+1}t_i^{-1}]/(t_{i+1}t_i^{-1}) & \text{if } k = i, \\ 0 & \text{otherwise.} \end{cases}$$

Further, $(\pi L_i)(U_k)$ is a GK 1-critical $\mathcal{O}(U_k)$ -module provided that it is nonzero, i.e. it has GK-dimension 1 and any proper quotient is finite-dimensional.

Proof. This follows directly from Lemma 9.5. \square

Corollary 9.9. *If $|j-i| \geq 2$ then $\text{Ext}_{\mathfrak{X}}^s(\pi L_i, \pi L_j) = 0$ for all s and so $L_i \cdot L_j = 0$ is well-defined.*

Proof. This follows from (7.2.8) and the observation that on all charts U_k either $\pi L_i(U_k) = 0$ or $\pi L_j(U_k) = 0$. \square

We further have the following computation, which we give without proof.

Lemma 9.10. *Let $C = \mathbb{k}_q[u, v]$ be a quantum polynomial ring, let L, L' be $C/(u)$ and $C/(v)$, respectively, and let $P = \mathbb{k}_q[u, v]/\langle u, v \rangle$. Then*

$$\text{Hom}_C(L, L) \cong L \cong \text{Ext}_C^1(L, L), \quad \text{Ext}_C^1(L, L') \cong P,$$

and all other $\text{Ext}_C^s(L, L)$ and $\text{Ext}_C^s(L, L')$ vanish. Thus

$$\text{Ext}_{\mathfrak{X}}^s(\pi L_i, \pi L_{i+1}) = \begin{cases} \pi P_i & \text{if } s = 1 \\ 0 & \text{otherwise,} \end{cases}$$

where $P_i = T/(\mathfrak{l}_i + \mathfrak{l}_{i+1})$. \square

Lemma 9.11. *We have*

$$\text{Ext}_{\mathfrak{X}}^s(\pi L_i, \pi L_{i+1}) = \begin{cases} \mathbb{k} & \text{if } s = 1, \\ 0 & \text{otherwise,} \end{cases}$$

so $L_i \cdot L_{i+1} = 1$ is well-defined.

Proof. By Lemma 9.10, the local-to-global spectral sequence (7.2.8) collapses to give

$$\text{Ext}_{\mathfrak{X}}^s(\pi L_i, \pi L_{i+1}) = H^{s-1}(\mathfrak{X}, \text{Ext}_{\mathfrak{X}}^1(\pi L_i, \pi L_{i+1})) \cong H_{\text{qgr-}T}^{s-1}(P_i).$$

A local cohomology computation gives the result. \square

We now complete the proof of Theorem 9.3.

Proposition 9.12.

$$\mathcal{E}xt_{\mathfrak{X}}^s(\pi L_i, \pi L_i) \cong \begin{cases} \pi L_i & \text{if } s = 0, \\ \pi L_i(-2) & \text{if } s = 1, \\ 0 & \text{otherwise,} \end{cases}$$

and $L_i \cdot L_i = -2$ is well-defined.

Proof. Consider the exact sequence $0 \rightarrow \mathfrak{l}_i \rightarrow T \rightarrow L_i \rightarrow 0$ and the associated long exact sequence

$$0 \rightarrow \mathcal{H}om_{\mathfrak{X}}(\pi L_i, \pi L_i) \xrightarrow{f} \mathcal{H}om_{\mathfrak{X}}(\pi T, \pi L_i) \xrightarrow{g} \mathcal{H}om_{\mathfrak{X}}(\pi \mathfrak{l}_i, \pi L_i) \xrightarrow{h} \mathcal{E}xt_{\mathfrak{X}}^1(\pi L_i, \pi L_i) \rightarrow 0, \quad (9.13)$$

where all other terms in the long exact sequence vanish. On each open set U_k , by Lemma 9.10 we see, first that f, h are isomorphisms and $g = 0$, and second that

$$\mathcal{E}xt_{\mathfrak{X}}^1(\pi L_i, \pi L_i)(U_k) \cong (\pi L_i)(U_k) \quad \text{for all } k. \quad (9.14)$$

Thus f is a global isomorphism and

$$\mathcal{H}om_{\mathfrak{X}}(\pi L_i, \pi L_i) \cong \mathcal{H}om_{\mathfrak{X}}(\pi T, \pi L_i) \cong \pi L_i.$$

We can therefore compute $\mathcal{E}xt_{\mathfrak{X}}^1(\pi L_i, \pi L_i)$ as $\mathcal{H}om_{\mathfrak{X}}(\pi \mathfrak{l}_i, \pi L_i)$. This object is isomorphic to $\pi(\mathcal{J}_i \otimes_T L_i)$ by Proposition 9.7 and the equivalence $\text{qgr-}\mathcal{T} \simeq \text{mod-}\mathcal{O}_{\mathfrak{X}}$ in Theorem 7.1.11. Again using Proposition 9.7 $\mathcal{J}_i \otimes_T L_i \cong \mathcal{J}_i / \mathcal{J}_i \mathfrak{l}_i$ is isomorphic in $\text{qgr-}T$ to

$$\mathcal{J}_i / T \cong \frac{z^{-1}t_{i-1}t_i T}{z^{-1}t_{i-1}t_i T \cap T}.$$

Since $z^{-1}t_{i-1}t_i \mathfrak{l}_i \subseteq T$, there is a surjection

$$\frac{z^{-1}t_{i-1}t_i T}{z^{-1}t_{i-1}t_i \mathfrak{l}_i} \twoheadrightarrow \frac{z^{-1}t_{i-1}t_i T}{z^{-1}t_{i-1}t_i T \cap T}. \quad (9.15)$$

Applying $\pi(-)(U_k)$, both terms in (9.15) become isomorphic to $(\pi L_i)(U_k)$. By the GK-criticality in Lemma 9.8, the surjection (9.15) must be an isomorphism on all charts and thus an isomorphism in $\text{qgr-}T$. Thus

$$\mathcal{E}xt_{\mathfrak{X}}^1(\pi L_i, \pi L_i) \cong \pi\left(\frac{z^{-1}t_{i-1}t_i T}{z^{-1}t_{i-1}t_i \mathfrak{l}_i}\right) \cong \pi(T(-2)/\mathfrak{l}_i(-2)) \cong \pi(L_i(-2)).$$

By the above computations, the E_2 -page of (7.2.8) for $\text{Ext}_{\mathfrak{X}}^s(\pi L_i, \pi L_i)$ is

$$H^0(\mathfrak{X}, \pi L_i(-2)) \quad H^1(\mathfrak{X}, \pi L_i(-2))$$

$$H^0(\mathfrak{X}, \pi L_i) \quad H^1(\mathfrak{X}, \pi L_i)$$

where all the maps are zero as we are on the E_2 -page. Thus

$$\begin{aligned} \mathcal{H}om_{\mathfrak{X}}(\pi L_i, \pi L_i) &\cong H^0(\mathfrak{X}, \pi L_i), \\ \mathcal{E}xt_{\mathfrak{X}}^1(\pi L_i, \pi L_i) &\cong H^0(\mathfrak{X}, \pi L_i(-2)) \oplus H^1(\mathfrak{X}, \pi L_i), \\ \mathcal{E}xt_{\mathfrak{X}}^2(\pi L_i, \pi L_i) &\cong H^1(\mathfrak{X}, \pi L_i(-2)). \end{aligned}$$

We see that

$$L_i \cdot L_i = -\chi(L_i) + \chi(L_i(-2)),$$

where for a graded T -module M we have

$$\chi(M) = \sum_{i \geq 0} (-1)^i \dim_{\mathbb{k}} H^i(\mathfrak{X}, \pi M) = \sum_{i \geq 0} (-1)^i \dim_{\mathbb{k}} H_{\text{Qgr-}T}^i(\pi M),$$

where the second equality follows from the equivalence between $\text{mod-}\mathcal{O}_{\mathfrak{X}}$ and $\text{qgr-}T$.

We will now show that $\chi(L_i(n)) = n+1$ for all $n \in \mathbb{Z}$, from which the fact that $L_i \cdot L_i = -2$ will follow. We calculate this via a local cohomology computation. If $k \notin \{i-1, i\}$ then $L_i[t_k^{-1}] = 0$. The Čech complex for L_i is thus

$$0 \rightarrow L_i \rightarrow L_i[t_{i-1}^{-1}] \oplus L_i[t_i^{-1}] \rightarrow L_i[t_{i-1}^{-1}, t_i^{-1}] \rightarrow 0.$$

From Lemma 9.1, this is the complex

$$0 \rightarrow \mathbb{k}[t_{i-1}, t_i] \rightarrow \mathbb{k}[t_{i-1}^{\pm 1}, t_i] \oplus \mathbb{k}[t_{i-1}, t_i^{\pm 1}] \rightarrow \mathbb{k}[t_{i-1}^{\pm 1}, t_i^{\pm 1}] \rightarrow 0$$

of \mathbb{k} -vector spaces. Thus $H_{T_+}^s(L_i) = 0$ for $s \neq 2$, and

$$\dim_{\mathbb{k}} H_{T_+}^2(L_i)_n = \begin{cases} 0 & \text{if } n \geq 0, \\ -(n+1) & \text{if } n \leq -1. \end{cases} \quad (9.16)$$

That $\chi(L_i(n)) = n+1$ follows from (9.16) and Lemma 9.1. \square

Remark 9.17. The computations in Proposition 9.12 show that

$$H_{\text{Qgr-}T}^{\bullet}(L_i(-1)) = \mathbb{R}\phi_*(L_i(-1)) = 0$$

for $1 \leq i \leq n$. Thus the $L_i(-1)$ are contained in the kernel of ϕ_* . We conjecture that the modules $L_1(-1), \dots, L_n(-1)$ generate the kernel of ϕ_* . This is the subject of ongoing research.

REFERENCES

- [AF92] F. W. Anderson and K. R. Fuller, *Rings and categories of modules*, second ed., Graduate Texts in Mathematics, vol. 13, Springer-Verlag, New York, 1992.
- [Art66] M. Artin, *On isolated rational singularities of surfaces*, Amer. J. Math. **88** (1966), 129–136.
- [AS87] M. Artin and W. F. Schelter, *Graded algebras of global dimension 3*, Adv. in Math. **66** (1987), no. 2, 171–216.
- [Aus62] M. Auslander, *On the purity of the branch locus*, American Journal of Mathematics **84** (1962), no. 1, 116–125.
- [AVdB90] M. Artin and M. Van den Bergh, *Twisted homogeneous coordinate rings*, J. Algebra **133** (1990), no. 2, 249–271.
- [AZ94] M. Artin and J. J. Zhang, *Noncommutative projective schemes*, Advances in Mathematics **109** (1994), no. 2, 228–287.
- [BH93] W. Bruns and J. Herzog, *Cohen-Macaulay rings*, Cambridge Studies in Advanced Mathematics, vol. 39, Cambridge University Press, Cambridge, 1993.
- [BKR01] T. Bridgeland, A. King, and M. Reid, *The McKay correspondence as an equivalence of derived categories*, J. Amer. Math. Soc. **14** (2001), no. 3, 535–554.
- [BM17] K. A. Brown and M. J. Macleod, *The Cohen–Macaulay property for noncommutative rings*, Algebras and Representation Theory **20** (2017), 1433–1465.
- [BS13] M. P. Brodmann and R. Y. Sharp, *Local cohomology*, second ed., Cambridge Studies in Advanced Mathematics, vol. 136, Cambridge University Press, Cambridge, 2013.
- [CBH98] W. Crawley-Boevey and M. P. Holland, *Noncommutative deformations of Kleinian singularities*, Duke Mathematical Journal **92** (1998), no. 3, 605–636.
- [CGW25] S. Crawford, J. Gaddis, and R. Won, *Actions of Taft algebras on noetherian down-up algebras*, Transformation Groups (2025), 1–46.
- [CKWZ16] K. Chan, E. Kirkman, C. Walton, and J. J. Zhang, *Quantum binary polyhedral groups and their actions on quantum planes*, J. Reine Angew. Math. **719** (2016), 211–252.
- [CKWZ18] ———, *McKay correspondence for semisimple Hopf actions on regular graded algebras, I*, Journal of Algebra **508** (2018), 512–538.
- [CKWZ19] ———, *McKay correspondence for semisimple Hopf actions on regular graded algebras. II*, Journal of Noncommutative Geometry **13** (2019), no. 1, 87–114.
- [Cra18] S. Crawford, *Singularities of noncommutative surfaces*, Ph.D. thesis, The University of Edinburgh, 2018.
- [Cra22] ———, *Superpotentials and quiver algebras for semisimple Hopf actions*, Algebras and Representation Theory (2022), 1–44.
- [DV34] P. Du Val, *On isolated singularities of surfaces which do not affect the conditions of adjunction (Part I.)*, Mathematical Proceedings of the Cambridge Philosophical Society **30** (1934), no. 4, 453–459.
- [Gro61] A. Grothendieck, *Éléments de géométrie algébrique, III. Étude cohomologique des faisceaux cohérents, I*, vol. 11, Inst. Hautes Études Sci. Publ. Math., 1961.
- [GS00] K. R. Goodearl and J. T. Stafford, *The graded version of Goldie’s theorem*, Algebra and Its Applications, vol. 259, American Mathematical Soc., 2000, pp. 237–240.
- [GW04] K. R. Goodearl and R. B. Warfield, Jr., *An introduction to noncommutative Noetherian rings*, second ed., London Mathematical Society Student Texts, vol. 61, Cambridge University Press, Cambridge, 2004.
- [Har77] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, No. 52, Springer-Verlag, New York-Heidelberg, 1977.

- [Ing16] C. Ingalls, *Quantum toric varieties*, unpublished personal communication, 2016.
- [IW13] O. Iyama and M. Wemyss, *On the noncommutative Bondal–Orlov conjecture*, Journal für die reine und angewandte Mathematik (Crelles Journal) **2013** (2013), no. 683, 119–128.
- [Kaw24] Y. Kawamata, *Non-commutative deformations of derived McKay correspondence for $A(n)$ singularities*, arXiv preprint arXiv:2410.15340 (2024).
- [Kaw25] ———, *Non-commutative deformations of tilting bundles and the derived McKay correspondence*, arXiv preprint arXiv:2505.07174 (2025).
- [Kel07] B. Keller, *Derived categories and tilting*, London Mathematical Society Lecture Note Series **332** (2007), 49.
- [KL00] G. R. Krause and T. H. Lenagan, *Growth of algebras and Gelfand–Kirillov dimension*, revised ed., Graduate Studies in Mathematics, vol. 22, American Mathematical Society, Providence, RI, 2000.
- [Kuz08] A. Kuznetsov, *Lefschetz decompositions and categorical resolutions of singularities*, Selecta Mathematica **13** (2008), no. 4, 661.
- [KV00] M. Kapranov and E. Vasserot, *Kleinian singularities, derived categories and Hall algebras*, Math. Ann. **316** (2000), no. 3, 565–576.
- [LW12] G. J. Leuschke and R. Wiegand, *Cohen–Macaulay representations*, Mathematical Surveys and Monographs, vol. 181, American Mathematical Society, Providence, RI, 2012.
- [Mat89] H. Matsumura, *Commutative ring theory*, second ed., Cambridge Studies in Advanced Mathematics, vol. 8, Cambridge University Press, Cambridge, 1989, Translated from the Japanese by M. Reid.
- [McK80] J. McKay, *Graphs, singularities, and finite groups*, Proc. Symp. Pure Math, vol. 37, 1980, pp. 183–186.
- [MS01] I. Mori and S. P. Smith, *Bézout’s theorem for non-commutative projective spaces*, J. Pure Appl. Algebra **157** (2001), no. 2-3, 279–299.
- [QWZ19] X.-S. Qin, Y.-H. Wang, and J. J. Zhang, *Noncommutative quasi-resolutions*, Journal of Algebra **536** (2019), 102–148.
- [Rot09] J. J. Rotman, *An introduction to homological algebra*, second ed., Universitext, Springer, New York, 2009.
- [RRZ14] M. Reyes, D. Rogalski, and J. J. Zhang, *Skew Calabi–Yau algebras and homological identities*, Advances in Mathematics **264** (2014), 308–354.
- [RSS15] D. Rogalski, S. J. Sierra, and J. T. Stafford, *Classifying orders in the Sklyanin algebra*, Algebra Number Theory **9** (2015), no. 9, 2055–2119.
- [RVdB89] I. Reiten and M. Van den Bergh, *Two-dimensional tame and maximal orders of finite representation type*, Mem. Amer. Math. Soc. **80** (1989), no. 408, viii+72.
- [Smi04] S. P. Smith, *Maps between non-commutative spaces*, Trans. Amer. Math. Soc. **356** (2004), no. 7, 2927–2944.
- [Sta25] The Stacks project authors, *The Stacks project*, <https://stacks.math.columbia.edu>, 2025.
- [VdB04a] M. Van den Bergh, *Non-commutative crepant resolutions*, The legacy of Niels Henrik Abel, Springer, Berlin, 2004, pp. 749–770.
- [VdB04b] ———, *Three-dimensional flops and non-commutative rings*, Duke Mathematical Journal **122** (2004), 423–455.
- [VdB23] ———, *Noncommutative crepant resolutions, an overview*, ICM—International Congress of Mathematicians. Vol. 2. Plenary lectures, EMS Press, Berlin, 2023, pp. 1354–1391.
- [Wei94] C. A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994.
- [Wem16] M. Wemyss, *Noncommutative resolutions*, Noncommutative algebraic geometry, Math. Sci. Res. Inst. Publ., vol. 64, Cambridge Univ. Press, New York, 2016, pp. 239–306.
- [YZ06] A. Yekutieli and J. J. Zhang, *Homological transcendence degree*, Proceedings of the London Mathematical Society **93** (2006), no. 1, 105–137.

THE UNIVERSITY OF MANCHESTER, ALAN TURING BUILDING, OXFORD ROAD, MANCHESTER, M13 9PL, UNITED KINGDOM

HEILBRONN INSTITUTE FOR MATHEMATICAL RESEARCH, BRISTOL, UNITED KINGDOM
 Email address: simon.crawford@manchester.ac.uk

SCHOOL OF MATHEMATICS, UNIVERSITY OF EDINBURGH, EDINBURGH EH9 3FD, UNITED KINGDOM
 Email address: s.sierra@ed.ac.uk