

# MONOTONICITY OF SOLUTIONS TO SECOND ORDER LINEAR DIFFERENCE EQUATIONS WITH CONSTANT COEFFICIENTS

YOSHIKI GOTO AND GENKI SHIBUKAWA

ABSTRACT. We describe some monotone properties of solutions to second order linear difference equations with real constant coefficients. As an application, we give a characterization of the Fibonacci numbers.

## 1. INTRODUCTION

The Fibonacci numbers

$$F_0 := 0, \quad F_1 := 1, \quad F_{n+2} - F_{n+1} - F_n = 0$$

can be regarded as the special values of the complete homogeneous symmetric polynomials

$$h_n(x, y) := \frac{x^{n+1} - y^{n+1}}{x - y} = \sum_{j=0}^n x^j y^{n-j}, \quad h_{-1}(x, y) := 0, \quad h_0(x, y) := 1$$

in two variables with

$$x = -2 \cos \left( \frac{4\pi}{5} \right) = \frac{1 + \sqrt{5}}{2}, \quad y = -2 \cos \left( \frac{2\pi}{5} \right) = \frac{1 - \sqrt{5}}{2},$$

that is,

$$F_{n+1} = h_n \left( -2 \cos \left( \frac{4\pi}{5} \right), -2 \cos \left( \frac{2\pi}{5} \right) \right) = \frac{1}{\sqrt{5}} \left[ \left( \frac{1 + \sqrt{5}}{2} \right)^{n+1} - \left( \frac{1 - \sqrt{5}}{2} \right)^{n+1} \right].$$

For the Fibonacci numbers, we easily check that the following inequalities which we call monotone properties hold:

- **monotone property 1** (positive monotone non-decreasing property):

$$F_0 = 0 < F_{n+1} \leq F_{n+2}; \tag{1.1}$$

- **monotone property 2:**

$$\left| \frac{1 + \sqrt{5}}{2} - \frac{F_{n+1}}{F_n} \right| \geq \left| \frac{1 + \sqrt{5}}{2} - \frac{F_{n+2}}{F_{n+1}} \right|; \tag{1.2}$$

- **monotone property 3:**

$$\left| F_n \frac{1 + \sqrt{5}}{2} - F_{n+1} \right| \geq \left| F_{n+1} \frac{1 + \sqrt{5}}{2} - F_{n+2} \right|. \tag{1.3}$$

The Lucas numbers

$$L_0 := 2, \quad L_1 := 1, \quad L_{n+2} - L_{n+1} - L_n = 0$$

can be regarded as the special values of the power sum symmetric polynomials

$$p_n(x, y) := x^n + y^n, \quad p_0(x, y) = 2$$

---

*Date:* November 18, 2025.

*2020 Mathematics Subject Classification.* 05E05, 11B39, 39A06.

*Key words and phrases.* Fibonacci numbers; symmetric polynomials; inequalities.

in two variables:

$$L_n = p_n \left( -2 \cos \left( \frac{4\pi}{5} \right), -2 \cos \left( \frac{2\pi}{5} \right) \right) = \left( \frac{1 + \sqrt{5}}{2} \right)^n + \left( \frac{1 - \sqrt{5}}{2} \right)^n.$$

For the Lucas numbers, since we have

$$L_0 = 2 > L_1 = 1 < L_2 = 3 < L_3 = 4 < \dots, \\ \left| \frac{1 + \sqrt{5}}{2} - \frac{1}{2} \right| = 1.11 \dots \leq \left| \frac{1 + \sqrt{5}}{2} - 3 \right| = 1.38 \dots \geq \left| \frac{1 + \sqrt{5}}{2} - \frac{4}{3} \right| = 0.28 \dots \geq \dots,$$

the monotone properties 1 and 2 do not hold. However, because of

$$\left| L_n \frac{1 + \sqrt{5}}{2} - L_{n+1} \right| = \left| \frac{1 - \sqrt{5}}{2} \right|^n \left| \frac{1 + \sqrt{5}}{2} - \frac{1 - \sqrt{5}}{2} \right| = \sqrt{5} \left| \frac{1 - \sqrt{5}}{2} \right|^n = \sqrt{5} (0.61 \dots)^n,$$

the monotone property 3 holds.

In this paper, we study the monotone properties 1–3 of a solution to a second order linear difference equation with real constant coefficients. For a first order linear difference equation, the problem is easy. Let us consider

$$a_0 := c_0 \in \mathbb{R} \setminus \{0\}, \quad a_{n+1} - aa_n = 0, \quad a \in \mathbb{R} \setminus \{0\}.$$

For the solution

$$a_n = c_0 a^n,$$

the monotone property 1 is equivalent to the condition  $c_0 > 0$  and  $a \geq 1$ . The monotone properties 2 and 3 always hold, since

$$\left| a - \frac{a_{n+1}}{a_n} \right| = 0, \quad |a_n a - a_{n+1}| = 0.$$

Now we consider a second order linear difference equation with real constant coefficients, with initial values  $(0, 0) \neq (c_0, c_1) \in \mathbb{R}^2$  or  $(0, 0) \neq (c_{-1}, c_0) \in \mathbb{R}^2$ :

$$a_0 := c_0, \quad a_1 := c_1, \quad a_{n+2} - aa_{n+1} + ba_n = 0, \quad a, b \in \mathbb{R}, \quad ab \neq 0. \quad (1.4)$$

In general, conditions for the monotone properties 1–3 of its solution are not so simple. As an example, we consider the following two difference equations with initial values  $c_{-1} := 0, c_0 := 1$ , which are given as the special values of the complete homogeneous symmetric polynomials  $h_n(x, y)$ .

**Example 1.1.** The solution to (1.4) for  $a = 1, b = 1/4, a_{-1} := 0, a_0 := 1$  is

$$a_n = (n + 1)2^{-n}.$$

Because of

$$a_n \geq a_{n+1},$$

the monotone property 1 does not hold. On the other hand, by

$$\left| \frac{1}{2} - \frac{a_{n+1}}{a_n} \right| = \frac{1}{2} \frac{1}{n + 1}, \quad \left| a_n \frac{1}{2} - a_{n+1} \right| = \frac{1}{2^{n+1}},$$

the monotone properties 2 and 3 hold.

**Example 1.2.** The solution to (1.4) for  $a = 1, b = -3, a_{-1} := 0, a_0 := 1$  is

$$a_n = \frac{1}{\sqrt{13}} \left[ \left( \frac{1 + \sqrt{13}}{2} \right)^{n+1} - \left( \frac{1 - \sqrt{13}}{2} \right)^{n+1} \right].$$

By the definition (1.4), it satisfies the monotone property 1. However, the monotone properties 2 and 3 do not hold, since we have

$$\left| \frac{1 + \sqrt{13}}{2} - 1 \right| = 1.30 \dots \leq \left| \frac{1 + \sqrt{13}}{2} - 4 \right| = 1.69 \dots \geq \left| \frac{1 + \sqrt{13}}{2} - \frac{7}{4} \right| = 0.55 \dots \geq \dots,$$

$$\left| \frac{1 + \sqrt{13}}{2} - 1 \right| = 1.30 \dots \leq \left| \frac{1 + \sqrt{13}}{2} - 4 \right| = 1.69 \dots \leq \left| 4 \frac{1 + \sqrt{13}}{2} - 7 \right| = 2.21 \dots \leq \dots.$$

Further, as in the case of the Lucas numbers, even if the monotone properties 1 and 2 do not hold, sometimes the properties hold for sufficiently large  $n$ .

Let  $a_n$  be the solution to the difference equation (1.4), and let

$$\alpha_{\pm} := \frac{a \pm \sqrt{a^2 - 4b}}{2}$$

be the roots of the characteristic polynomial  $x^2 - ax + b$ . Here, when  $a^2 - 4b < 0$ , we set  $\sqrt{a^2 - 4b} = \sqrt{-1}\sqrt{4b - a^2}$ . Note that the assumption  $ab \neq 0$  implies

$$\alpha_{\pm} \neq 0, \tag{1.5}$$

$$|\alpha_+| = |\alpha_-| \Leftrightarrow \alpha_+ = \alpha_-. \tag{1.6}$$

We also use notations  $\alpha, \beta$  for the characteristic roots (i.e.,  $\{\alpha, \beta\} = \{\alpha_+, \alpha_-\}$ ) so that

$$|\alpha| \geq |\beta|.$$

We first give conditions under which the following inequalities (1.7) and (1.10) hold for sufficiently large  $n$ .

**Theorem 1.3.** *There exists a non-negative integer  $n_0$  such that*

$$a_n \leq a_{n+1} \quad (n_0 \leq n), \tag{1.7}$$

*if and only if  $a^2 - 4b \geq 0$  and one of the following two conditions hold:*

$$1 \neq \alpha_+ > 0, \quad a > 0, \quad (\alpha_+ - 1)(c_1 - c_0\alpha_-) > 0; \tag{1.8}$$

*or*

$$\alpha_+ = 1, \quad a_0 \leq a_1 \leq a_2. \tag{1.9}$$

**Theorem 1.4.** *There exists a non-negative integer  $n_0$  such that*

$$\left| \alpha - \frac{a_{n+1}}{a_n} \right| \geq \left| \alpha - \frac{a_{n+2}}{a_{n+1}} \right| \quad (n_0 \leq n), \tag{1.10}$$

*if and only if  $a^2 - 4b \geq 0$  holds.*

We next discuss three monotone properties 1, 2 and 3 (respectively, Theorem 1.5, Theorem 1.7 and Theorem 1.8).

**Theorem 1.5.** *For a non-negative integer  $k$ , the inequality*

$$a_n \leq a_{n+1} \quad (n \geq k - 1) \tag{1.11}$$

*holds if and only if  $a^2 - 4b \geq 0$  and one of the following two conditions hold:*

$$a_{k-1} \leq a_k \leq a_{k+1}, \quad 1 \neq \alpha_+ > 0, \quad a > 0, \quad (\alpha_+ - 1)(c_1 - c_0\alpha_-) > 0, \tag{1.12}$$

*or*

$$a_{k-1} \leq a_k \leq a_{k+1}, \quad \alpha_+ = 1. \tag{1.13}$$

*In particular, the inequality  $a_n \leq a_{n+1}$  holds for any non-negative integer  $n$  if and only if  $a^2 - 4b \geq 0$  and one of the following two conditions hold:*

$$a_{-1} \leq a_0 \leq a_1, \quad 1 \neq \alpha_+ > 0, \quad a > 0, \quad (\alpha_+ - 1)(c_1 - c_0\alpha_-) > 0, \tag{1.14}$$

or

$$a_{-1} \leq a_0 \leq a_1, \quad \alpha_+ = 1. \quad (1.15)$$

As a corollary of Theorem 1.5, we obtain a condition equivalent to the monotone property 1 for the special values of the complete homogeneous symmetric polynomials.

**Corollary 1.6.** *Let the initial values be  $a_{-1} := 0$ ,  $a_0 \neq 0$ . The inequality*

$$a_{-1} = 0 < a_0 \leq a_n \leq a_{n+1} \quad (1.16)$$

*holds for any non-negative integer  $n$  if and only if we have  $a^2 - 4b \geq 0$  and*

$$a_0 > 0, \quad a \geq 1, \quad \alpha_+ \geq 1. \quad (1.17)$$

**Theorem 1.7.** *Let the initial values be  $a_{-1} := 0$ ,  $a_0 \neq 0$ . The inequality*

$$\left| \alpha - \frac{a_{n+1}}{a_n} \right| \geq \left| \alpha - \frac{a_{n+2}}{a_{n+1}} \right| \quad (1.18)$$

*holds for any non-negative integer  $n$  if and only if we have  $a^2 - 4b \geq 0$  and*

$$|\alpha + \beta| = |a| \geq |\beta|. \quad (1.19)$$

**Theorem 1.8.** *The inequality*

$$|a_n \alpha - a_{n+1}| \geq |a_{n+1} \alpha - a_{n+2}| \quad (1.20)$$

*holds for any non-negative integer  $n$  if and only if*

$$|\beta| \leq 1. \quad (1.21)$$

This paper is arranged as follows. In Section 2, we give some lemmas which are used in the proof of main results. In Section 3, we prove our main results. In Section 4, we determine a class of special values of  $h_n(x, y)$  which satisfy the monotone properties 1–3, and give a characterization of the Fibonacci numbers.

## 2. PRELIMINARIES

Throughout this paper,  $a_n$  is the solution to the difference equation (1.4).

**Lemma 2.1.** *We assume  $a^2 - 4b \geq 0$ . We have*

$$|\alpha_+| - |\alpha_-| = \begin{cases} \sqrt{a^2 - 4b} & (a > 0, b > 0) \\ -\sqrt{a^2 - 4b} & (a < 0, b > 0) \\ a & (b < 0) \end{cases}.$$

*Especially, we obtain*

$$a > 0, a^2 - 4b > 0 \quad \Leftrightarrow \quad |\alpha_+| = \alpha_+ > |\alpha_-|, \quad (2.1)$$

$$a < 0, a^2 - 4b > 0 \quad \Leftrightarrow \quad |\alpha_+| < |\alpha_-|. \quad (2.2)$$

*Proof.* We first assume  $a > 0$  and  $b > 0$ . Then we have  $a > \sqrt{a^2 - 4b} \geq 0$ , and hence, we obtain

$$|\alpha_+| - |\alpha_-| = \alpha_+ - \alpha_- = \sqrt{a^2 - 4b}.$$

Next, we assume  $a < 0$  and  $b > 0$ . Because of  $a + \sqrt{a^2 - 4b} < 0$  and  $a - \sqrt{a^2 - 4b} < 0$ , we obtain

$$|\alpha_+| - |\alpha_-| = -\alpha_+ + \alpha_- = -\sqrt{a^2 - 4b}.$$

Finally, we assume  $b < 0$ . Then we have  $\sqrt{a^2 - 4b} > a$ , which implies

$$|\alpha_+| - |\alpha_-| = \alpha_+ + \alpha_- = a.$$

□

**Lemma 2.2.** *For any non-negative integer  $n$ , we have*

$$a_n = c_0 h_n(\alpha_+, \alpha_-) + (c_1 - ac_0) h_{n-1}(\alpha_+, \alpha_-) \quad (2.3)$$

$$\begin{aligned} &= c_0 h_n(\alpha_+, \alpha_-) - bc_{-1} h_{n-1}(\alpha_+, \alpha_-) \\ &= \frac{(c_0 \alpha_+ + c_1 - ac_0) \alpha_+^n - (c_0 \alpha_- + c_1 - ac_0) \alpha_-^n}{\sqrt{a^2 - 4b}} \\ &= \frac{(c_1 - c_0 \alpha_-) \alpha_+^n - (c_1 - c_0 \alpha_+) \alpha_-^n}{\sqrt{a^2 - 4b}}. \end{aligned} \quad (2.4)$$

In particular, in the case of  $\alpha = \alpha_+ = \alpha_-$ , we have

$$\begin{aligned} a_n &= c_0(n+1)\alpha^n + (c_1 - ac_0)n\alpha^{n-1} \\ &= c_0(1-n)\alpha^n + c_1 n\alpha^{n-1}. \end{aligned} \quad (2.5)$$

*Proof.* We prove only (2.3); the others easily follow from this. By the definition, we have  $h_{-1}(\alpha_+, \alpha_-) = 0$ ,  $h_0(\alpha_+, \alpha_-) = 1$  and  $h_1(\alpha_+, \alpha_-) = a$ . Thus, (2.3) holds for  $n = 0, 1$ . Further, we have

$$\begin{aligned} &h_{n+2}(\alpha_+, \alpha_-) - ah_{n+1}(\alpha_+, \alpha_-) + bh_n(\alpha_+, \alpha_-) \\ &= \frac{(\alpha_+^{n+3} - a\alpha_+^{n+2} + b\alpha_+^{n+1}) - (\alpha_-^{n+3} - a\alpha_-^{n+2} + b\alpha_-^{n+1})}{\alpha_+ - \alpha_-} = 0, \end{aligned}$$

which implies that (2.3) holds for any non-negative integer  $n$ .  $\square$

**Lemma 2.3.** *If we assume  $c_0 c_1 \neq 0$  and  $a^2 - 4b \geq 0$ , then the solution  $a_n$  of the difference equation (1.4) is not zero, except for at most one  $n$ . In particular, we have  $a_n \neq 0$  for sufficiently large  $n$ .*

*Proof.* First, we assume  $a^2 - 4b = 0$ . In this case, we have  $\alpha_+ = \alpha_- = \alpha$ , and hence,

$$a_n = c_0(1-n)\alpha^n + c_1 n\alpha^{n-1} = \alpha^{n-1}(c_0\alpha - (c_0\alpha - c_1)n) = 0$$

implies  $c_0\alpha = (c_0\alpha - c_1)n$ . We may assume  $c_0\alpha - c_1 \neq 0$ , since  $c_0\alpha - c_1 = 0$  implies  $c_0 = c_1 = 0$ . We obtain

$$n = \frac{c_0\alpha}{c_0\alpha - c_1},$$

and the number of such integers  $n$  is at most one. Next, we assume  $a^2 - 4b > 0$ . Then,

$$a_n = \frac{(c_1 - c_0\alpha_-)\alpha_+^n - (c_1 - c_0\alpha_+)\alpha_-^n}{\sqrt{a^2 - 4b}} = 0$$

implies

$$(c_1 - c_0\alpha_-)\alpha_+^n = (c_1 - c_0\alpha_+)\alpha_-^n. \quad (2.6)$$

By  $\alpha_+ \neq \alpha_-$  and (1.5), at least one of  $c_1 - c_0\alpha_+$  and  $c_1 - c_0\alpha_-$  are not zero. The equality (2.6) is equivalent to

$$c_1 - c_0\alpha_- = (c_1 - c_0\alpha_+) \left( \frac{\alpha_-}{\alpha_+} \right)^n,$$

and the number of such integers  $n$  is at most one.  $\square$

**Lemma 2.4.** *We assume  $c_0 c_1 \neq 0$ . The ratio  $a_{n+1}/a_n$  of consecutive solutions converges if and only if  $a^2 - 4b \geq 0$  holds. Moreover, in this case, the limit is give by*

$$\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = \begin{cases} \alpha & (c_1 - c_0\beta \neq 0) \\ \beta & (c_1 - c_0\beta = 0) \end{cases}. \quad (2.7)$$

*Proof.* First, we prove the “if” part. We assume  $a^2 - 4b < 0$ , that is,  $\alpha_{\pm} \notin \mathbb{R}$ . Since there exist  $\theta \in \mathbb{R} \setminus \mathbb{Z}$  such that

$$\alpha_{\pm} = |b|^{\frac{1}{2}} e^{\pm \sqrt{-1}\pi\theta},$$

we have

$$\frac{a_{n+1}}{a_n} = |b|^{\frac{1}{2}} \frac{c_0 |b|^{\frac{1}{2}} \sin(\pi(n+2)\theta) + (c_1 - ac_0) \sin(\pi(n+1)\theta)}{c_0 |b|^{\frac{1}{2}} \sin(\pi(n+1)\theta) + (c_1 - ac_0) \sin(\pi n\theta)}.$$

If  $\theta = \frac{N}{M} \in \mathbb{Q}$ , then  $a \neq 0$  implies  $M \geq 3$ , and hence, the ratio  $a_{n+1}/a_n$  does not converge because of  $a_{n+M} = (-1)^N a_n$ . On the other hand, if we assume  $\theta \notin \mathbb{Q}$ , then the ratio  $a_{n+1}/a_n$  does not converge by Kronecker's approximation theorem (see [2, Theorem 439]).

Next, we prove the “only if” part. In the case of  $|\alpha| > |\beta|$ , we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} &= \lim_{n \rightarrow \infty} \frac{(c_1 - c_0\beta)\alpha^{n+1} - (c_1 - c_0\alpha)\beta^{n+1}}{(c_1 - c_0\beta)\alpha^n - (c_1 - c_0\alpha)\beta^n} \\ &= \begin{cases} \alpha & (c_1 - c_0\beta \neq 0) \\ \beta & (c_1 - c_0\beta = 0) \end{cases}. \end{aligned}$$

We assume  $|\alpha| = |\beta|$ . By (1.6), we have  $\alpha = \beta = \frac{a}{2}$ , and hence, we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} &= \lim_{n \rightarrow \infty} \frac{-c_0 n \alpha^{n+1} + c_1(n+1)\alpha^n}{c_0(1-n)\alpha^n + c_1 n \alpha^{n-1}} \\ &= \alpha \lim_{n \rightarrow \infty} \frac{n(c_1 - c_0\alpha) + c_1}{n(c_1 - c_0\alpha) + c_0\alpha} \\ &= \begin{cases} \alpha & (c_1 - c_0\alpha \neq 0) \\ \frac{c_1}{c_0} = \alpha & (c_1 - c_0\alpha = 0) \end{cases}. \end{aligned}$$

□

### 3. PROOFS OF MAIN RESULTS

*Proof of Theorem 1.3.* We prove the theorem by considering the following eight cases:

- (a1)  $\alpha_+ > 1$ ,  $a^2 - 4b > 0$ ,  $a > 0$ ; (a2)  $\alpha_+ > 1$ ,  $a^2 - 4b > 0$ ,  $a < 0$ ; (a3)  $\alpha_+ > 1$ ,  $a^2 - 4b = 0$ ;  
 (b1)  $\alpha_+ < 1$ ,  $a^2 - 4b > 0$ ,  $a > 0$ ; (b2)  $\alpha_+ < 1$ ,  $a^2 - 4b > 0$ ,  $a < 0$ ; (b3)  $\alpha_+ < 1$ ,  $a^2 - 4b = 0$ ;  
 (c1)  $\alpha_+ = 1$ ,  $a^2 - 4b > 0$ ; (c2)  $\alpha_+ = 1$ ,  $a^2 - 4b = 0$ .

(a1) By (2.4), we have

$$a_{n+1} - a_n = \frac{\alpha_+^n(\alpha_+ - 1)(c_1 - c_0\alpha_-) - \alpha_-^n(\alpha_- - 1)(c_1 - c_0\alpha_+)}{\sqrt{a^2 - 4b}} \quad (3.1)$$

$$= \frac{\alpha_+^n}{\sqrt{a^2 - 4b}} \left[ (\alpha_+ - 1)(c_1 - c_0\alpha_-) - \left( \frac{\alpha_-}{\alpha_+} \right)^n (\alpha_- - 1)(c_1 - c_0\alpha_+) \right] \quad (3.2)$$

Since the inequality (2.1)  $|\alpha_+| > |\alpha_-|$  holds by the assumption  $a > 0$ , the absolute value of the second term of the right-hand side converges to zero. Therefore,  $a_n \leq a_{n+1}$  holds for sufficiently large  $n$  if and only if  $(\alpha_+ - 1)(c_1 - c_0\alpha_-) > 0$  holds.

(a2) We rewrite (3.1) as

$$a_{n+1} - a_n = \frac{\alpha_-^n}{\sqrt{a^2 - 4b}} \left[ \left( \frac{\alpha_+}{\alpha_-} \right)^n (\alpha_+ - 1)(c_1 - c_0\alpha_-) - (\alpha_- - 1)(c_1 - c_0\alpha_+) \right]. \quad (3.3)$$

The inequality (2.2)  $|\alpha_+| < |\alpha_-|$  holds by the assumption  $a < 0$ . Since  $\alpha_+ > 1$  and  $a = \alpha_+ + \alpha_- < 0$  imply  $\alpha_- < -1$ , the value of (3.3) takes both positive and negative numbers. Thus, (1.7) does not hold.

(a3) By (2.5), we have

$$\begin{aligned} a_{n+1} - a_n &= c_0\alpha_+^n((n+2)\alpha_+ - (n+1)) + (c_1 - ac_0)\alpha_+^{n-1}((n+1)\alpha_+ - n) \\ &= ((n+1)\alpha_+ - n)(c_1 - c_0\alpha_-)\alpha_+^{n-1} + c_0\alpha_+^n(\alpha_+ - 1) \end{aligned}$$

$$= \alpha_+^{n-1} \left[ (n+1) \left( \alpha_+ - 1 + \frac{1}{n+1} \right) (c_1 - c_0 \alpha_-) + c_0 \alpha_+ (\alpha_+ - 1) \right]. \quad (3.4)$$

Since the term  $(n+1) \left( \alpha_+ - 1 + \frac{1}{n+1} \right)$  can take an arbitrarily large number,  $a_n \leq a_{n+1}$  holds for sufficiently large  $n$  if and only if  $c_1 - c_0 \alpha_- > 0$  holds.

(b1) In the case of  $\alpha_+ < 0$ , the value of (3.2) takes both positive and negative numbers, and hence, (1.7) does not hold. In the case of  $0 < \alpha_+ < 1$ , the inequality  $a_n \leq a_{n+1}$  holds for sufficiently large  $n$  if and only if  $(\alpha_+ - 1)(c_1 - c_0 \alpha_-) \geq 0$  holds.

(b2) Similarly to (a2), by using  $\alpha_- < 0$  and (3.3), we can show that (1.7) does not hold.

(b3) For sufficiently large  $n$ , the value of  $(n+1) \left( \alpha_+ - 1 + \frac{1}{n+1} \right)$  in (3.4) takes a negative number with arbitrarily large absolute value. In the case of  $\alpha_+ < 0$ , the inequality (1.7) does not hold. In the case of  $0 < \alpha_+ < 1$ , the inequality  $a_n \leq a_{n+1}$  holds for sufficiently large  $n$  if and only if  $c_1 - c_0 \alpha_- \leq 0$  holds.

(c1) Because of

$$b = \alpha_- = a - \alpha_+ = a - 1,$$

we have

$$a_{n+1} - a_n = (a - 1)(a_n - a_{n-1}) = (a - 1)^n (a_1 - a_0). \quad (3.5)$$

This implies

$$a_0 \leq a_1, a \geq 1 \quad \Leftrightarrow \quad a_{n+1} - a_n \geq 0 \quad (n \in \mathbb{Z}_{\geq 0}).$$

(c2) In this case, we have  $\alpha_+ = \alpha_- = 1$  and  $a = 2$ . Thus, (3.5) implies

$$a_0 \leq a_1 \quad \Leftrightarrow \quad a_{n+1} - a_n \geq 0 \quad (n \in \mathbb{Z}_{\geq 0}).$$

□

*Proof of Theorem 1.4.* We first assume  $c_1 \neq c_0 \beta$ . By (2.7), the ratio  $a_{n+1}/a_n$  converges to  $\alpha$ . Since we have

$$\left| \alpha - \frac{a_{n+2}}{a_{n+1}} \right| = \left| \alpha - a + b \frac{a_n}{a_{n+1}} \right| = \left| \frac{-b}{\alpha} + b \frac{a_n}{a_{n+1}} \right| = \left| \frac{b}{\alpha} \frac{a_n}{a_{n+1}} \right| \left| \alpha - \frac{a_{n+1}}{a_n} \right|,$$

the inequality (1.10) is equivalent to

$$\left| \frac{b}{\alpha} \frac{a_n}{a_{n+1}} \right| = \left| \beta \frac{a_n}{a_{n+1}} \right| \leq 1. \quad (3.6)$$

If  $a^2 - 4b > 0$  holds, it easily follows from  $|\alpha| > |\beta|$  that (3.6) holds for sufficiently large  $n$ . We assume  $a^2 - 4b = 0$ . By  $\alpha = \beta$  and (2.5), we have

$$\begin{aligned} \left| \beta \frac{a_n}{a_{n+1}} \right| &= \left| \alpha \frac{c_0(n+1)\alpha^n + (c_1 - ac_0)n\alpha^{n-1}}{c_0(n+2)\alpha^{n+1} + (c_1 - ac_0)(n+1)\alpha^n} \right| \\ &= \left| \frac{c_0(n+1)\alpha + (c_1 - ac_0)n}{c_0(n+2)\alpha + (c_1 - ac_0)(n+1)} \right| \\ &= \left| \frac{c_0\alpha + (c_1 - c_0\alpha)n}{c_1 + (c_1 - c_0\alpha)n} \right|. \end{aligned}$$

By the assumption  $c_1 \neq c_0\alpha$ , we have

$$\left| \beta \frac{a_n}{a_{n+1}} \right| = 1 - \frac{1}{1 + \frac{c_1 - c_0\alpha}{c_1 - c_0\alpha} \frac{1}{n}} \frac{1}{n} \leq 1$$

for sufficiently large  $n$ , and hence, the inequality (1.10) holds.

Next, we consider the case of  $c_1 = c_0\beta$ . In this case, we have

$$a_n = c_0\beta^n$$

for any non-negative integer  $n$ . Indeed, the claim obviously holds for  $n = 0, 1$ , and if we assume it holds for  $n$  and  $n + 1$  then we have

$$a_{n+2} = aa_{n+1} - ba_n = c_0\beta^n((\alpha + \beta)\beta - \alpha\beta) = c_0\beta^{n+2}.$$

Therefore, for any non-negative integer  $n$ , we have

$$\left| \alpha - \frac{a_{n+1}}{a_n} \right| = |\alpha - \beta| \geq 0,$$

and hence, the inequality (1.10) holds.  $\square$

*Proof of Theorem 1.5.* We first assume  $\alpha_+ = 1$ . In this case, the “if” part of the proof is obvious. We show the “only if” part. By

$$a_{k+1} - a_k = (a - 1)a_k - ba_{k-1} = (a - 1)(a_k - a_{k-1}) \geq 0,$$

we obtain  $a - 1 \geq 0$ . Thus, the inequality  $0 < a_n \leq a_{n+1}$  holds for any  $n \geq k - 1$ .

Next, we assume  $\alpha_+ \neq 1$ . The “if” part clearly follows from Theorem 1.3. We show the “only if” part. Namely, we assume the condition (1.12). Note that the inequality (2.1)  $|\alpha_+| = \alpha_+ \geq |\alpha_-|$  holds by  $a > 0$ . The condition  $a_{k-1} \leq a_k \leq a_{k+1}$  implies

$$\begin{aligned} a_k - a_{k-1} &= \frac{\alpha_+^{k-1}}{\sqrt{a^2 - 4b}} \left[ (\alpha_+ - 1)(c_1 - c_0\alpha_-) - \left( \frac{\alpha_-}{\alpha_+} \right)^{k-1} (\alpha_- - 1)(c_1 - c_0\alpha_+) \right] \geq 0, \\ a_{k+1} - a_k &= \frac{\alpha_+^k}{\sqrt{a^2 - 4b}} \left[ (\alpha_+ - 1)(c_1 - c_0\alpha_-) - \left( \frac{\alpha_-}{\alpha_+} \right)^k (\alpha_- - 1)(c_1 - c_0\alpha_+) \right] \geq 0. \end{aligned}$$

In the following, let  $m$  be an arbitrary positive integer.

(A1) In the case of  $(\alpha_- - 1)(c_1 - c_0\alpha_+) \geq 0$ ,  $\frac{\alpha_-}{\alpha_+} > 0$ , we have

$$\begin{aligned} a_{k+m+1} - a_{k+m} &= \frac{\alpha_+^{k+m}}{\sqrt{a^2 - 4b}} \left[ (\alpha_+ - 1)(c_1 - c_0\alpha_-) - \left( \frac{\alpha_-}{\alpha_+} \right)^{k+m} (\alpha_- - 1)(c_1 - c_0\alpha_+) \right] \\ &\geq \frac{\alpha_+^{k+m}}{\sqrt{a^2 - 4b}} \left[ (\alpha_+ - 1)(c_1 - c_0\alpha_-) - \left( \frac{\alpha_-}{\alpha_+} \right)^k (\alpha_- - 1)(c_1 - c_0\alpha_+) \right] \geq 0. \end{aligned}$$

(A2) We consider the case of  $(\alpha_- - 1)(c_1 - c_0\alpha_+) \geq 0$ ,  $\frac{\alpha_-}{\alpha_+} < 0$ . If  $k$  is odd, then  $a_{k+2m+1} - a_{k+2m} \geq 0$  is obvious, and we have

$$\begin{aligned} a_{k+2m} - a_{k+2m-1} &= \frac{\alpha_+^{k+2m-1}}{\sqrt{a^2 - 4b}} \left[ (\alpha_+ - 1)(c_1 - c_0\alpha_-) - \left( \frac{\alpha_-}{\alpha_+} \right)^{k+2m-1} (\alpha_- - 1)(c_1 - c_0\alpha_+) \right] \\ &= \frac{\alpha_+^{k+2m-1}}{\sqrt{a^2 - 4b}} \left[ (\alpha_+ - 1)(c_1 - c_0\alpha_-) - \left( \frac{\alpha_-}{\alpha_+} \right)^{k-1} (\alpha_- - 1)(c_1 - c_0\alpha_+) \right] \\ &\quad + \frac{\alpha_+^{k+2m-1}}{\sqrt{a^2 - 4b}} \left( \frac{\alpha_-}{\alpha_+} \right)^{k-1} \left[ 1 - \left( \frac{\alpha_-}{\alpha_+} \right)^{2m} \right] (\alpha_- - 1)(c_1 - c_0\alpha_+) \geq 0. \end{aligned}$$

If  $k$  is even, then  $a_{k+2m} - a_{k+2m-1} \geq 0$  is obvious, and we have

$$\begin{aligned} a_{k+2m+1} - a_{k+2m} &= \frac{\alpha_+^{k+2m}}{\sqrt{a^2 - 4b}} \left[ (\alpha_+ - 1)(c_1 - c_0\alpha_-) - \left( \frac{\alpha_-}{\alpha_+} \right)^{k+2m} (\alpha_- - 1)(c_1 - c_0\alpha_+) \right] \\ &= \frac{\alpha_+^{k+2m}}{\sqrt{a^2 - 4b}} \left[ (\alpha_+ - 1)(c_1 - c_0\alpha_-) - \left( \frac{\alpha_-}{\alpha_+} \right)^k (\alpha_- - 1)(c_1 - c_0\alpha_+) \right] \\ &\quad + \frac{\alpha_+^{k+2m}}{\sqrt{a^2 - 4b}} \left( \frac{\alpha_-}{\alpha_+} \right)^k \left[ 1 - \left( \frac{\alpha_-}{\alpha_+} \right)^{2m} \right] (\alpha_- - 1)(c_1 - c_0\alpha_+) \geq 0. \end{aligned}$$



(B1) In the case of  $(\alpha_- - 1)(c_1 - c_0\alpha_+) \leq 0$ ,  $\frac{\alpha_-}{\alpha_+} > 0$ , the inequality  $a_{k+m+1} - a_{k+m} \geq 0$  holds obviously.  
 (B2) We consider the case of  $(\alpha_- - 1)(c_1 - c_0\alpha_+) \leq 0$ ,  $\frac{\alpha_-}{\alpha_+} < 0$ . If  $k$  is odd, then  $a_{k+2m} - a_{k+2m-1} \geq 0$  is obvious, and we have

$$\begin{aligned} a_{k+2m+1} - a_{k+2m} &= \frac{\alpha_+^{k+2m}}{\sqrt{a^2 - 4b}} \left[ (\alpha_+ - 1)(c_1 - c_0\alpha_-) - \left(\frac{\alpha_-}{\alpha_+}\right)^{k+2m} (\alpha_- - 1)(c_1 - c_0\alpha_+) \right] \\ &= \frac{\alpha_+^{k+2m}}{\sqrt{a^2 - 4b}} \left[ (\alpha_+ - 1)(c_1 - c_0\alpha_-) - \left(\frac{\alpha_-}{\alpha_+}\right)^k (\alpha_- - 1)(c_1 - c_0\alpha_+) \right] \\ &\quad + \frac{\alpha_+^{k+2m}}{\sqrt{a^2 - 4b}} \left(\frac{\alpha_-}{\alpha_+}\right)^k \left[ 1 - \left(\frac{\alpha_-}{\alpha_+}\right)^{2m} \right] (\alpha_- - 1)(c_1 - c_0\alpha_+) \geq 0. \end{aligned}$$

If  $k$  is even, then  $a_{k+2m+1} - a_{k+2m} \geq 0$  is obvious, and we have

$$\begin{aligned} a_{k+2m} - a_{k+2m-1} &= \frac{\alpha_+^{k+2m-1}}{\sqrt{a^2 - 4b}} \left[ (\alpha_+ - 1)(c_1 - c_0\alpha_-) - \left(\frac{\alpha_-}{\alpha_+}\right)^{k+2m-1} (\alpha_- - 1)(c_1 - c_0\alpha_+) \right] \\ &= \frac{\alpha_+^{k+2m-1}}{\sqrt{a^2 - 4b}} \left[ (\alpha_+ - 1)(c_1 - c_0\alpha_-) - \left(\frac{\alpha_-}{\alpha_+}\right)^{k-1} (\alpha_- - 1)(c_1 - c_0\alpha_+) \right] \\ &\quad + \frac{\alpha_+^{k+2m-1}}{\sqrt{a^2 - 4b}} \left(\frac{\alpha_-}{\alpha_+}\right)^{k-1} \left[ 1 - \left(\frac{\alpha_-}{\alpha_+}\right)^{2m} \right] (\alpha_- - 1)(c_1 - c_0\alpha_+) \geq 0. \end{aligned}$$

□

By the proof of Theorem 1.5, we conclude that if the inequality (1.11) holds for  $n = k - 1$ ,  $n = k$  and for sufficiently large  $n$ , then it holds for any  $n \geq k - 1$ .

**Remark 3.1.** In Hartman-Aurel [3], [4], some sufficient conditions for the monotone property  $y_n < y_{n+1}$  of solutions to the linear ordinary difference equations are given. If we apply the result for our difference equation (1.4), we obtain a sufficient condition for the property:

$$a - 1 - b > 0, \quad b > 0.$$

Indeed, it is obvious since we have

$$a_{n+2} - a_{n+1} = (a - 1 - b)a_{n+1} + b(a_{n+1} - a_n).$$

*Proof of Corollary 1.6.* Note that  $a_1 = aa_0$ . We apply Theorem 1.5 for  $k = 0$ . The condition (1.12) is reduced into

$$0 < a_{-1} = 0 < a_0 \leq a_1 = aa_0, \quad 1 \neq \alpha_+ > 0, \quad a > 0, \quad (\alpha_+ - 1)(c_1 - c_0\alpha_-) = a_0\alpha_+(\alpha_+ - 1) > 0,$$

which is equivalent to  $a_0 > 0$ ,  $a \geq 1$ ,  $\alpha_+ > 1$ . The condition (1.13) is equivalent to  $a_0 > 0$ ,  $a \geq 1$ ,  $\alpha_+ = 1$ . Summarizing them, we obtain the condition (1.17). □

**Remark 3.2.** Note that under the condition (1.13) in Corollary 1.6, we have



$$\alpha_+ = \alpha, \quad \alpha_- = \beta.$$

In this situation, two domains

$$D_1 := \{(\alpha, \beta) \in \mathbb{R}^2 \mid \alpha + \beta \geq 1, \alpha \geq 1, |\alpha| \geq |\beta|\}, \quad (3.7)$$

$$D'_1 := \{(a, b) \in \mathbb{R}^2 \mid a \geq 1, \alpha \geq 1, |\alpha| \geq |\beta|\} \quad (3.8)$$

are drawn as in Figures 1 and 2, respectively.

FIGURE 1.  $D_1$ FIGURE 2.  $D'_1$ 

*Proof of Theorem 1.7.* First, we show the “only if” part of the proof. Namely, we assume  $a^2 - 4b \geq 0$  and  $|\alpha + \beta| = |a| \geq |\beta|$ . Similarly to the proof of Theorem 1.4, for any  $n$ , the inequality (1.10) is equivalent to (3.6), that is,

$$\left| \frac{b}{\alpha} \frac{a_n}{a_{n+1}} \right| = \left| \beta \frac{\alpha^{n+1} - \beta^{n+1}}{\alpha^{n+2} - \beta^{n+2}} \right| \leq 1.$$

We consider the following five cases:

- (i)  $|\alpha| = |\beta|$ ;      (ii-1)  $|\alpha| > |\beta|$ ,  $\alpha > 0$ ,  $\beta > 0$ ;      (ii-2)  $|\alpha| > |\beta|$ ,  $\alpha < 0$ ,  $\beta < 0$ ;
- (ii-3)  $|\alpha| > |\beta|$ ,  $\alpha > 0$ ,  $\beta < 0$ ;      (ii-4)  $|\alpha| > |\beta|$ ,  $\alpha < 0$ ,  $\beta > 0$ .

(i) Note that  $\alpha = \beta \neq 0$  by (1.6). Since we have

$$a_n = a_0(n+1)\alpha^n, \quad \frac{a_{n+1}}{a_n} = \frac{n+2}{n+1}\alpha,$$

we obtain

$$\left| \alpha - \frac{a_{n+2}}{a_{n+1}} \right| = \frac{|\alpha|}{n+2} \leq \frac{|\alpha|}{n+1} = \left| \alpha - \frac{a_{n+1}}{a_n} \right|$$

for any non-negative integer  $n$ .

(ii-1) In this case, we have  $\alpha > \beta > 0$ . The condition (1.19) is reduced into  $\alpha + \beta \geq \beta$ , and hence, we have

$$|\alpha^{n+2} - \beta^{n+2}| - |\beta(\alpha^{n+1} - \beta^{n+1})| = \alpha^{n+1}(\alpha - \beta) \geq 0.$$

(ii-2) The inequality (3.6) can be shown similarly to (ii-1).

(ii-3) We have  $|\alpha| = \alpha > -\beta = |\beta|$  and  $|\alpha + \beta| = \alpha + \beta \geq -\beta = |\beta|$ . Thus, we obtain

$$\begin{aligned} |\alpha^{n+2} - \beta^{n+2}| - |\beta(\alpha^{n+1} - \beta^{n+1})| &= \alpha^{n+1}(\alpha + \beta) - 2\beta^{n+2} \\ &= \alpha^{n+1}(\alpha + 2\beta) - \beta(\alpha^{n+1} + 2\beta^{n+1}). \end{aligned} \quad (3.9)$$

Since the inequality  $\alpha^{n+1} + 2\beta^{n+1} \geq 0$  holds for any non-negative integer  $n$ , the right-hand side of (3.9) is non-negative.

(ii-4) By setting  $\alpha' := -\alpha$  and  $\beta' := -\beta$ , we can prove (3.6) similarly to (ii-3).

Next, we show the “if” part. The inequality (3.6) for  $n = 0$  is reduced into

$$\left| \frac{\beta}{\alpha + \beta} \right| \leq 1,$$

which implies  $\alpha + \beta \neq 0$  and  $|\beta| \leq |\alpha + \beta|$ . Further, by Theorem 1.4, the inequality (1.10) holds for sufficiently large  $n$  if and only if  $a^2 - 4b \geq 0$  holds.  $\square$

**Remark 3.3.** As an analogy of Theorem 1.7, we consider a necessary condition for arbitrary initial values:  $a^2 - 4b \geq 0$  and

$$\left| \frac{\beta c_0}{c_1} \right| \leq 1.$$

However, it is not a sufficient condition. Indeed, if we set

$$c_0 = 1, \quad c_1 = 3, \quad \alpha = 2.1, \quad \beta = -2,$$

then we have  $a_2 = (2.1 - 2)3 + 4.2 = 4.5$  and

$$\left| \frac{\beta a_0}{a_1} \right| = \left| \frac{\beta c_0}{c_1} \right| = \frac{2}{3} \leq 1, \quad \left| \frac{\beta a_1}{a_2} \right| = \frac{4}{3} \geq 1.$$

**Remark 3.4.** Two domains

$$D_2 := \{(\alpha, \beta) \in \mathbb{R}^2 \mid |\alpha + \beta| \geq |\beta|, |\alpha| \geq |\beta|\}, \quad (3.10)$$

$$D'_2 := \{(a, b) \in \mathbb{R}^2 \mid |a| \geq |\beta|, |\alpha| \geq |\beta|\} \quad (3.11)$$

corresponding to the condition (1.19) of Theorem 1.7 are drawn as in Figures 3 and 4, respectively.




FIGURE 3.  $D_2$




FIGURE 4.  $D'_2$

**Remark 3.5.** Theorem 1.7 can be translated into the property of a solution to a difference Riccati equation with real constant coefficients:

$$b_{n+1} = \frac{xb_n + y}{zb_n + w}. \quad (3.12)$$

Indeed, by setting

$$b_n := \frac{a_{n+1}}{a_n}$$

and dividing the difference equation (1.4) by  $a_{n+1}$ , we obtain

$$b_{n+1} = a - b \frac{1}{b_n} = \frac{ab_n - b}{b_n}. \quad (3.13)$$

Theorem 1.7 gives a condition under which the solution of the difference Riccati equation (3.13) with initial value

$$b_0 = \frac{a_1}{a_0} = a$$

approaches to the fixed point monotonically.

*Proof of Theorem 1.8.* For the solution  $a_n$  to the difference equation (1.4) with arbitrary initial values, we have

$$\begin{aligned}
|a_n\alpha - a_{n+1}| &= |c_0(h_n(\alpha, \beta)\alpha - h_{n+1}(\alpha, \beta)) + (c_1 - ac_0)(h_{n-1}(\alpha, \beta)\alpha - h_n(\alpha, \beta))| \\
&= \left| c_0 \frac{\beta^{n+2} - \alpha\beta^{n+1}}{\alpha - \beta} + (c_1 - ac_0) \frac{\beta^{n+1} - \alpha\beta^n}{\alpha - \beta} \right| \\
&= |-c_0\beta^{n+1} - (c_1 - ac_0)\beta^n| \\
&= |\beta|^n |c_0\beta + (c_1 - ac_0)| \\
&= |c_1 - c_0\alpha| |\beta|^n.
\end{aligned}$$

If we assume

$$\begin{aligned}
|a_n\alpha - a_{n+1}| - |a_{n+1}\alpha - a_{n+2}| &= |c_1 - c_0\alpha| |\beta|^n - |c_1 - c_0\alpha| |\beta|^{n+1} \\
&= |c_1 - c_0\alpha| |\beta|^n (1 - |\beta|) \geq 0,
\end{aligned} \tag{3.14}$$

then we obtain  $|\beta| \leq 1$ . Conversely, if we assume  $|\beta| \leq 1$ , then the inequality (3.14) holds.  $\square$

**Remark 3.6.** If we assume that the monotone property 1 (1.16) and the monotone property 3 (1.20) hold for any non-negative integer  $n$ , then the monotone property 2 (1.10) also holds, since we have

$$\left| \alpha - \frac{a_{n+2}}{a_{n+1}} \right| \leq \left| \frac{a_n}{a_{n+1}} \right| \left| \alpha - \frac{a_{n+1}}{a_n} \right| \leq \left| \alpha - \frac{a_{n+1}}{a_n} \right|$$

(see Proposition 4.1). However, the converse is not true. For example, although the solution in Example 1.1 satisfies (1.10) and (1.20), it does not satisfy (1.16).

**Remark 3.7.** We assume  $a, b \in \mathbb{Z}$ . Suppose that the characteristic polynomial  $x^2 - ax + b$  is irreducible over  $\mathbb{Z}$ . In this case, the condition (1.21) is  $\beta \neq 0$  and  $|\beta| < 1$ , which implies that  $\alpha$  is a quadratic integer greater than 1 and is a Pisot number [1]. Namely, the inequality (1.20) holds for any initial values  $c_0, c_1$  and for any  $n$  if and only if  $\alpha$  is a Pisot number.

**Remark 3.8.** Two domains

$$D_3 := \{(\alpha, \beta) \in \mathbb{R}^2 \mid |\beta| \leq 1, |\alpha| \geq |\beta|\}, \tag{3.15}$$

$$D'_3 := \{(a, b) \in \mathbb{R}^2 \mid |\beta| \leq 1, |\alpha| \geq |\beta|\} \tag{3.16}$$

corresponding the condition (1.21) of Theorem 1.8 are drawn as in Figures 5 and 6, respectively.




FIGURE 5.  $D_3$




FIGURE 6.  $D'_3$

## 4. CONCLUDING REMARKS



One of the motivations of our study is to characterize the Fibonacci numbers by monotone properties. As a generalization of the Fibonacci numbers, the solution  $a_n$  to the difference equation (1.4), especially that with initial values  $a_{-1} := 0$ ,  $a_0 := 1$  (i.e., the special values of the complete homogeneous symmetric polynomials), has been studied well (e.g., [5]). On the other hand, in previous studies, it seems that the monotone properties 1–3 were not paid attention to. By our main results, we determine integer solutions  $a_n$  satisfying the monotone properties (1.16), (1.18) and (1.20) which are analogies of the inequalities (1.1), (1.2) and (1.3) for the Fibonacci numbers.

Indeed, it suffices to restrict the characteristic roots (resp. the coefficients) of (1.4) to the intersection  $D := D_1 \cap D_2 \cap D_3$  (resp.  $D' := D'_1 \cap D'_2 \cap D'_3$ ) of the domains  $D_1$ ,  $D_2$  and  $D_3$  (resp.  $D'_1$ ,  $D'_2$  and  $D'_3$ ) in Remarks 3.2, 3.4 and 3.8. By easy calculations, we obtain the following proposition.

**Proposition 4.1.** *We have*

$$D := D_1 \cap D_2 \cap D_3 = \{(\alpha, \beta) \in \mathbb{R}^2 \mid \alpha + \beta \geq 1, \alpha \geq 1, |\beta| \leq 1\}, \quad (4.1)$$

$$D' := D'_1 \cap D'_2 \cap D'_3 = \{(a, b) \in \mathbb{R}^2 \mid a \geq 1, a - 1 \geq b \geq -a - 1\}. \quad (4.2)$$

FIGURE 7.  $D$ FIGURE 8.  $D'$ 

Further, we note the following lemma.

**Lemma 4.2.** *Let  $(a, b) \in \mathbb{Z}^2 \cap D'$ . Then, we have the following equivalence:*

$$x^2 - ax + b \text{ is irreducible over } \mathbb{Z} \Leftrightarrow b \neq -a - 1, 0, a - 1. \quad (4.3)$$

*Proof.* It is obvious that if  $b = -a - 1, 0, a - 1$ , the polynomial  $x^2 - ax + b$  is reducible over  $\mathbb{Z}$ . We prove the converse. Assume that  $x^2 - ax + b$  is reducible over  $\mathbb{Z}$ . Then, there exists an integer  $k$  such that

$$x^2 - ax + b = (x - k)(x - a + k).$$

By  $(k, a - k) \in \mathbb{Z}^2 \cap D'$  or  $(a - k, k) \in \mathbb{Z}^2 \cap D'$ , Proposition 4.1 implies that  $|a - k| \leq 1$  or  $|k| \leq 1$  holds. Therefore, we obtain  $k = a - 1, a, a + 1$  or  $k = -1, 0, 1$ , and hence, the value of  $b = k(a - k)$  takes only  $b = -a - 1, 0, a - 1$ .  $\square$

We list difference equations (1.4) such that the coefficients  $(a, b) \in \mathbb{Z}^2 \cap D'$  satisfy the condition (4.3):

$$a = 1 : a_{n+2} = a_{n+1} + a_n,$$

$$a = 2 : a_{n+2} = 2a_{n+1} + 2a_n, \quad a_{n+2} = 2a_{n+1} + a_n,$$

$$a = 3 : a_{n+2} = 3a_{n+1} + 3a_n, \quad a_{n+2} = 3a_{n+1} + 2a_n, \quad a_{n+2} = 3a_{n+1} + a_n, \quad a_{n+2} = 3a_{n+1} - a_n,$$

⋮

The first equation is the recursion of the Fibonacci numbers. The second equation in  $a = 2$  is the recursion of the Pell numbers. The recursion of the Fibonacci numbers is characterized by

$$\{(a, b) \in \mathbb{Z}^2 \cap \partial D' \mid (a, b) \text{ satisfies (4.3)}\} = \{(1, -1)\},$$

where  $\partial D'$  means the boundary of the domain  $D'$ . From the viewpoint of the monotone properties and irreducibility over  $\mathbb{Z}$ , the Fibonacci numbers are very special. Thus, the above recursions with initial values  $a_{-1} := 0$ ,  $a_0 := 1$  give appropriate generalizations of the Fibonacci numbers.

*Acknowledgement.* The authors are grateful to Professor Shigeki Akiyama who gave an opportunity to consider Theorem 1.8. We thank Professor Kentaro Mitsui for pointing out the interpretation in Remark 3.5. This work was supported by JST CREST Grant Number JP19209317 and JSPS KAKENHI Grant Number 21K13808.

## REFERENCES

- [1] M.J. Bertin, A. Decomps-Guilloux, M. Grandet-Hugot, M. Pathiaux-Delefosse, J.P. Schreiber: *Pisot and Salem Numbers*, Birkhauser, Basel, (1992).
- [2] G.H. Hardy and E.M. Wright: *An introduction to the theory of numbers. Fifth edition*, The Clarendon Press, Oxford University Press, New York, 1979.
- [3] P. Hartman and W. Aurel: *On linear difference equations of second order*, Amer. J. Math. **72**-1, (1950) pp124–128.
- [4] P. Hartman and W. Aurel: *Linear differential and difference equations with monotone solutions*, Amer. J. Math. **75**, (1953) pp731–743.
- [5] T. Koshy: *Fibonacci and Lucas Numbers with Applications Volume 1 Second Edition*, Wiley, (2017).

(Goto) OTARU UNIVERSITY OF COMMERCE, 3-5-21, MIDORI, OTARU, HOKKAIDO, 047-8501, JAPAN  
*Email address:* goto@res.otaru-uc.ac.jp

(Shibukawa) KITAMI INSTITUTE OF TECHNOLOGY, 165, KOEN-CHO, KITAMI, HOKKAIDO, 090-8507, JAPAN  
*Email address:* g-shibukawa@mail.kitami-it.ac.jp