

Characterization of t-norms on normal convex functions

Jie Sun*

School of Mathematics, Southwest Minzu University, Chengdu 610041, China

Abstract

Type-2 fuzzy set (T2 FS) were introduced by Zadeh in 1965, and the membership degrees of T2 FSs are type-1 fuzzy sets (T1 FSs). Owing to the fuzziness of membership degrees, T2 FSs can better model the uncertainty of real life, and thus, type-2 rule-based fuzzy systems (T2 RFSs) become hot research topics in recent decades. In T2 RFS, the compositional rule of inference is based on triangular norms (t-norms) defined on complete lattice $(\mathbf{L}, \sqsubseteq)$ (\mathbf{L} is the set of all convex normal functions from $[0, 1]$ to $[0, 1]$, and \sqsubseteq is the so-called convolution order). Hence, the choice of t-norm on $(\mathbf{L}, \sqsubseteq)$ may influence the performance of T2 RFS. Therefore, it is significant to broad the set of t-norms among which domain experts can choose most suitable one. To construct t-norms on $(\mathbf{L}, \sqsubseteq)$, the mainstream method is convolution $*_{\Delta}$ which is induced by two operators $*$ and Δ on the unit interval $[0, 1]$. A key problem appears naturally, when convolution $*_{\Delta}$ is a t-norm on $(\mathbf{L}, \sqsubseteq)$. This paper has solved this problem completely. Moreover, note that the computational complexity of operators prevent the application of T2 RFSs. This paper also provides one kind of convolutions which are t-norms on $(\mathbf{L}, \sqsubseteq)$ and extremely easy to calculate.

Keywords: Type-2 fuzzy sets, type-2 rule-based fuzzy systems, triangular norms, convolutions, convex normal functions

1. Introduction

Fuzzy set theory were introduced by Zadeh in 1965 [38]. Later on, one rule-based fuzzy system (RFS) were given by Mamdani and Assilian [22, 23] to control a steam engine, that is the earliest application of fuzzy sets in real world. Until now RFSs have been successfully applied to a quantity of industrial processes, see e.g. [25, 26, 28, 32].

A RFS contains four components—rules, fuzzifier, inference, and output processor as shown in Figure 1. Once the rules have been established, the fuzzy system can be viewed as a mapping from inputs to outputs. When T1 (T2, resp.) FSs are used to model the linguistic variable that appears in rule's antecedent and consequent, the RFS is called a T1 (T2, resp.) RFS. In recent decades, many authors have studied T2 RFSs [2, 18, 24, 29], especially interval-valued (IV) RFSs [3, 6] (since each IV FS can be expressed by a T2 FS, IV RFSs can be viewed as one kind of T2 RFSs). Note that in the research of T2 RFSs, the membership degrees of T2 FSs are usually assumed to be convex normal functions from $[0, 1]$ to $[0, 1]$.

Triangular norms on the unit interval $[0, 1]$ were introduced by Schweizer and Sklar [27]. Due to the close connection between order theory and fuzzy set theory, see e.g. [8], t-norms on generally partially ordered sets (posets) have been studied by many authors, see e.g. [19, 4, 41].

Triangular norms on complete lattices play an important role in the inference block of RFS. For example, suppose that there is only one rule in Figure 1, that is

IF input is A , THEN output is B .

According to Mamdani implication (see [24, 23])

$$\mu_{A \rightarrow B}(x, y) = A(x) * B(y)$$

*Corresponding author.

Email address: jiesun1027@163.com (Jie Sun)




Figure 1: Rule-based fuzzy system

and Zadeh's compositional rule of inference (see [7, 17, 39]), the inference block assigns a fuzzy input set A' to a fuzzy output set B' given by

$$B'(y) = \sup\{A'(x) * A(x) * B(y) \mid x \in X\},$$

where $*$ is a t-norm on certain complete lattice. Concretely,

- In T1 RFS, all fuzzy sets A, B, A', B' are type-1 fuzzy sets, and $*$ is a t-norm on complete lattice $([0, 1], \leq)$ such as minimum or product.
- In IV RFS, all fuzzy sets A, B, A', B' are interval-valued fuzzy sets, and $*$ is a t-norm on complete lattice $(\mathbf{I}^{[2]}, \leq_I)$ ($\mathbf{I}^{[2]}$ is the set of all closed subintervals of $[0, 1]$, and \leq_I is the interval-valued order [5]).
- In T2 RFS, all fuzzy sets A, B, A', B' are type-2 fuzzy sets whose membership degrees are elements of \mathbf{L} , and $*$ is a t-norm on complete lattice $(\mathbf{L}, \sqsubseteq)$.

Moreover, there exists order-preserving embedding from lattice $([0, 1], \leq)$ to lattice $(\mathbf{I}^{[2]}, \leq_I)$, from lattice $(\mathbf{I}^{[2]}, \leq_I)$ to lattice $(\mathbf{L}, \sqsubseteq)$, respectively, which means that IV RFS is the generalization of T1 RFS, and T2 RFS is the generalization of IV RFS to some extent.

As mentioned above, t-norms play an important role in RFS and may influence performance of RFS. To provide more choice for design of RFS, it is necessary to study construction of t-norms. However, though the construction of t-norms on lattice $([0, 1], \leq)$ and $(\mathbf{I}^{[2]}, \leq_I)$ have been deeply studied for T1 RFSs and IV RFSs, respectively, the same has not happened with t-norms on $(\mathbf{L}, \sqsubseteq)$ for T2 RFSs.

To construct operators on \mathbf{L} or $\mathbf{M} = [0, 1]^{[0,1]}$ (all maps from $[0, 1]$ to $[0, 1]$), the mainstream method is based on convolution (see Definition 2.6), which has a close relation with Zadeh's extension principle [40]. Convolution were widely studied as (generalized) extended t-norm [13, 36, 42], uninorm [21, 37], implication [34, 35], aggregation [30, 31], overlap function [16, 20], etc. In particular, through the research on convolutions \wedge_Δ (extended minimum) and \vee_Δ (extended maximum), a series of significant properties of poset $(\mathbf{M}, \sqsubseteq)$ and its subposets, such as $(\mathbf{L}, \sqsubseteq)$, have been obtained by Harding, Walker and Walker (see [10, 11, 33]). Besides, Zhang and Hu [43] showed that one kind of convolution $*_\Delta$ can be calculated through α -cut. Hu and Wang [15] provided an equivalent characterization of convolution order \sqsubseteq on \mathbf{L} through the strong α -cut, that makes the convolution order \sqsubseteq more intuitive.

To construct t-norms on $(\mathbf{L}, \sqsubseteq)$ for T2 RFS, a key problem appears naturally, when convolution is a t-norm on $(\mathbf{L}, \sqsubseteq)$. In 2014, Hernández, Cubillo and Torres-Blanc [12] provided sufficient but not necessary conditions under which convolution is a t_r -norm (t-norm satisfying more restrictive axioms) on $(\mathbf{L}, \sqsubseteq)$. In this paper, we completely solve the above problem. The main results are listed as follows.

- 1) The necessary and sufficient conditions, under which convolution is a t-norm on $(\mathbf{L}, \sqsubseteq)$, is obtained, see Theorem 3.5 and Theorem 4.19. A large number of t-norms on $(\mathbf{L}, \sqsubseteq)$ can be constructed by Theorem 4.19. As an application, experts can choice t-norms more flexibly to design T2 RFS.

- 2) One kind of convolutions are obtained which are t-norms on $(\mathbf{L}, \sqsubseteq)$ and extremely easy to calculate, see Definition 4.15 and Proposition 4.21. As a result, T2 RFSs, employing this kind of convolutions, are relatively easy to compute.
- 3) New equivalent characterization of convolution order \sqsubseteq are given, see Proposition 4.11. This result may promote the research of aggregation operators (one kind of monotone functions) on $(\mathbf{L}, \sqsubseteq)$. Aggregation refers to the process of combining several values into a single one, and are used for the fusion of information [9]. In particular, arithmetic mean is an aggregation operator.

The paper are organized as follows. Section II reviews definitions and properties of t-norm, t_r -norm, type-2 fuzzy set and convolution. Section III shows that $*_{\Delta}$ is a t-norm if and only if it is a t_r -norm, and provides the necessary conditions under which convolution $*_{\Delta}$ is a t-norm. Section IV proves that the necessary conditions given in Section III is also sufficient. Section V provides the necessary and sufficient conditions under which $*_{\Delta}$ is a t-conorm or t_r -conorm.

2. Preliminaries

In this section, we recall some definitions and properties which are used in the sequel.

2.1. T-norm and L-fuzzy set

Let $*$ be a binary operation on X . The algebra $(X, *)$ is called a *semigroup*, if for all $x, y, z \in X$, $(x * y) * z = x * (y * z)$. The algebra $(X, *)$ is called *commutative*, if for all $x, y \in X$, $x * y = y * x$. An element $e \in X$ is called the *unit* of the algebra $(X, *)$, if $e * x = x * e = x$ for all $x \in X$.

Let \leq be a partial order on X . The triple $(X, *, \leq)$ is called a *partially ordered semigroup* or *posemigroup* for brief, if $(P, *)$ is a semigroup such that $x * y \leq x * z$, $y * x \leq z * x$ whenever $y \leq z$.

Definition 2.1. Let $(P, \leq, 0_P, 1_P)$ be a bounded poset. A binary operation $* : P^2 \longrightarrow P$ is called a t-norm (t-conorm resp.), if $(P, *, \leq)$ is a commutative posemigroup with 1_P (0_P resp.) being the unit.

Two posemigroups $(P_1, *_1, \leq_1)$ and $(P_2, *_2, \leq_2)$ are called to be *isomorphic*, if there is an order isomorphism $\tau : P_1 \longrightarrow P_2$ such that for all $x, y \in P_1$,

$$\tau(x *_1 y) = \tau(x) *_2 \tau(y).$$

In particular, two t-norms $*_1 : P_1^2 \longrightarrow P_1$ and $*_2 : P_2^2 \longrightarrow P_2$ are called to be isomorphic, if the two posemigroups $(P_1, *_1, \leq_1)$ and $(P_2, *_2, \leq_2)$ are isomorphic.

Proposition 2.2. Suppose that posemigroups $(P_1, *_1, \leq_1)$ and $(P_2, *_2, \leq_2)$ are isomorphic. Then $*_1$ is a t-norm on (P_1, \leq_1) if and only if $*_2$ is a t-norm on (P_2, \leq_2) .

Proposition 2.3. Let $*_1$ and $*_2$ be two binary functions on bounded posets (P_1, \leq_1) and (P_2, \leq_2) , respectively. If there exists an order-reversing isomorphism $\tau : P_1 \longrightarrow P_2$ satisfying for all $x, y \in P_1$,

$$\tau(x *_1 y) = \tau(x) *_2 \tau(y),$$

then $*_1$ is a t-norm on (P_1, \leq_1) if and only if $*_2$ is a t-conorm on (P_2, \leq_2) .

Example 2.4. The product t-norm $T_p : [0, 1]^2 \longrightarrow [0, 1]$ and the Łukasiewicz t-norm $T_L : [0, 1]^2 \longrightarrow [0, 1]$, are respectively given by

$$T_p(x, y) = x \cdot y, \quad T_L(x, y) = \max\{x + y - 1, 0\}.$$

Suppose that $0 \leq a < b \leq 1$. A t-norm $* : [a, b]^2 \longrightarrow [a, b]$ is called

- *continuous*, if it is continuous as a two binary function.
- *border continuous*, if it is continuous on the boundary of the unit square $[a, b]^2$, i.e. on the set of $[a, b]^2 \setminus (a, b)^2$.

- *left-continuous*, if it is left continuous in each component.
- *conditionally cancellative*, if $x_1 * y = x_2 * y > a$ implies $x_1 = x_2$.
- *Archimedean*, if for any $x, y \in (a, b)$, there exists $n \in \mathbb{N}$ such that $x_*^{(n)} \leq y$.

Let $*$: $[0, 1]^2 \longrightarrow [0, 1]$ be a binary map. For $a, b \in [0, 1]$, define

$$a * b^- = \sup\{a * t \mid t < b\}$$

In particular, $a * 0^- = \sup \emptyset = 0$.

The following proposition is obtained from the monograph [1].

Proposition 2.5. *Let $*$: $[0, 1]^2 \longrightarrow [0, 1]$ be a t-norm. Then*

- (i) *$*$ is continuous if and only if it is continuous in each component.*
- (ii) *$*$ is border continuous if and only if for all $a \in [0, 1]$, $a * 1^- = a * 1$.*
- (iii) *$*$ is left-continuous implies that $*$ is border continuous.*

Let L be a complete lattice. An L -fuzzy set A is determined by a membership function

$$\mu_A : X \longrightarrow L$$

where $\mu_A(x)$ denote the membership degree of $x \in X$ to A (see [8]).

In particular, type-1 fuzzy sets are exactly $[0, 1]$ -fuzzy sets. Interval-valued fuzzy sets are exactly $\mathbf{I}^{[2]}$ -fuzzy sets, where the complete lattice $\mathbf{I}^{[2]}$ denotes the set of all closed subintervals of $[0, 1]$ equipped with the order \leq_I defined by $[a_1, b_1] \leq_I [a_2, b_2]$, if $a_1 \leq b_1$ and $a_2 \leq b_2$.

2.2. Type-2 fuzzy set and convolution

A type-2 fuzzy set A is determined by a membership function

$$\mu_A : X \longrightarrow \mathbf{M} = [0, 1]^{[0,1]} = \text{Map}([0, 1], [0, 1]).$$

A standard method to construct operations on \mathbf{M} is the convolution.

Definition 2.6. Let $*$: $[0, 1]^n \longrightarrow [0, 1]$ and Δ : $[0, 1]^n \longrightarrow [0, 1]$ be two n -ary operations on $[0, 1]$. The convolution induced by $*$ and Δ is an n -ary operation $*_{\Delta} : \mathbf{M}^n \longrightarrow \mathbf{M}$ given by

$$*_{\Delta}(f_1, \dots, f_n)(x) = \bigvee_{*(y_1, \dots, y_n)=x} \Delta(f(y_1), \dots, f(y_n)).$$

In particular, if there not exist $(y_1, y_2, \dots, y_n) \in [0, 1]^n$ such that $*(y_1, \dots, y_n) = x$, let

$$*_{\Delta}(f_1, \dots, f_n)(x) = \bigvee \emptyset = 0.$$

Let $*$ be an n -ary operation on $[0, 1]$. The Zadeh extension of $*$ is exactly the convolution $*_{\wedge} : \mathbf{M}^n \longrightarrow \mathbf{M}$. The Δ -extension of $*$ is exactly the convolution $*_{\Delta} : \mathbf{M}^n \longrightarrow \mathbf{M}$, where Δ is usually a t-norm on $[0, 1]$. The Δ -extension is also called generalized extension.

For each $\alpha \in [0, 1]$, the α -cut and strong α -cut of $f \in \mathbf{M}$ are respectively given by

$$f^{\alpha} = \{x \in [0, 1] \mid f(x) \geq \alpha\},$$

$$f^{\hat{\alpha}} = \{x \in [0, 1] \mid f(x) > \alpha\}.$$

If $*$ be a binary function on $[0, 1]$, for any $A, B \subseteq [0, 1]$, let $A * B$ denote the set

$$\{x * y \mid x \in A, y \in B\}.$$

Under the case that $*$: $[0, 1]^2 \longrightarrow [0, 1]$ is surjective and Δ : $[0, 1]^2 \longrightarrow [0, 1]$ is a t-norm, the convolution $*_{\Delta}$ can be represented through α -cut, see [43, Theorem 3.1 (2)]. In the following, we give a more general result.

Proposition 2.7. *If $*$ and Δ are two binary functions on $[0, 1]$ and Δ is monotone increasing in each place, then for any $f, g \in \mathbf{M}$ and $\alpha \in [0, 1]$,*

$$(f *_\Delta g)^{\hat{\alpha}} = \bigcup_{\alpha_1 \Delta \alpha_2 > \alpha} f^{\alpha_1} * g^{\alpha_2}.$$

Proof. Firstly, we show that $(f *_\Delta g)^{\hat{\alpha}} \subseteq \bigcup_{\alpha_1 \Delta \alpha_2 > \alpha} f^{\alpha_1} * g^{\alpha_2}$. For any $x_0 \in (f *_\Delta g)^{\hat{\alpha}}$, since $(f *_\Delta g)(x_0) > \alpha$, i.e.

$$(f *_\Delta g)(x_0) = \bigvee_{y*z=x_0} f(y)\Delta g(z) > \alpha$$

then there exist $y_0, z_0 \in [0, 1]$ such that $y_0 * z_0 = x$ and $f(y_0)\Delta g(z_0) > \alpha$. Put $\alpha_1 = f(y_0)$ and $\alpha_2 = g(z_0)$, then

$$x = y_0 * z_0 \in f^{\alpha_1} * g^{\alpha_2} \subseteq \bigcup_{\alpha_1 \Delta \alpha_2 > \alpha} f^{\alpha_1} * g^{\alpha_2}.$$

Conversely, for any $x \in \bigcup_{\alpha_1 \Delta \alpha_2 > \alpha} f^{\alpha_1} * g^{\alpha_2}$, there exist $\alpha_1, \alpha_2 \in [0, 1]$ such that $\alpha_1 \Delta \alpha_2 > \alpha$ and $x \in f^{\alpha_1} * g^{\alpha_2}$. Due to $x \in f^{\alpha_1} * g^{\alpha_2}$, there exist $y_0, z_0 \in [0, 1]$ such that $x = y_0 * z_0$, $f(y_0) \geq \alpha_1$ and $g(z_0) \geq \alpha_2$. Therefore,

$$(f *_\Delta g)(x) = \bigvee_{y*z=x} f(y)\Delta g(z) \geq f(y_0)\Delta g(z_0) \geq \alpha_1 \Delta \alpha_2 > \alpha,$$

thus, $x \in (f *_\Delta g)^{\hat{\alpha}}$. □

Proposition 2.8. *For $f, g \in \mathbf{M}$, the following statements are equivalent.*

- (i) $f = g$.
- (ii) $f^\alpha = g^\alpha$ for all $\alpha \in (0, 1]$.
- (iii) $f^{\hat{\alpha}} = g^{\hat{\alpha}}$ for all $\alpha \in [0, 1)$.

There exist two important partial orders \sqsubseteq and \leq on \mathbf{M} induced by the convolutions \wedge_Δ (extended minimum) and \vee_Δ (extended maximum), respectively.

Definition 2.9. The partial orders \sqsubseteq and \leq on \mathbf{M} defined by

$$\begin{aligned} f &\sqsubseteq g, \text{ if } f \wedge_\Delta g = f, \\ f &\leq g, \text{ if } f \vee_\Delta g = g, \end{aligned}$$

are called to be *convolution orders*.

It is known that $(\mathbf{M}, \sqsubseteq)$ is an inf semilattice, (\mathbf{M}, \leq) is a sup semilattice, and the convolution orders \sqsubseteq and \leq are distinct on \mathbf{M} (see [33]).

For $A \subseteq [0, 1]$, let $\bar{A} : [0, 1] \rightarrow [0, 1]$ denote the *characteristic function* of A , that is

$$\bar{A}(x) = \begin{cases} 1, & \text{if } x \in A, \\ 0, & \text{otherwise.} \end{cases}$$

In particular, for $a \in [0, 1]$, let \bar{a} denote the characteristic function of the singleton $\{a\}$.

Proposition 2.10. [33] *For any $f \in \mathbf{M}$, $\bar{0} \leq f$ and $f \sqsubseteq \bar{1}$.*

Unfortunately, $(\mathbf{M}, \sqsubseteq, \bar{0}, \bar{1})$ is not a bounded poset since $\bar{0}$ is not the minimum, and $(\mathbf{M}, \leq, \bar{0}, \bar{1})$ is not a bounded poset since $\bar{1}$ is not the maximum.

Note that some significant results of the subposets of $(\mathbf{M}, \sqsubseteq)$ (or (\mathbf{M}, \leq)) are obtained by Harding, Walker and Walker (see [10, 11, 33]). In the following, we recall these subposets.

An element f of \mathbf{M} is called *normal*, if

$$\sup\{f(x) \mid x \in [0, 1]\} = 1.$$

Let \mathbf{N} denote the set of all normal elements of \mathbf{M} .

Proposition 2.11. [12] For each $f \in \mathbf{M}$, $\bar{0} \sqsubseteq f \Leftrightarrow f \leq \bar{1} \Leftrightarrow f \in \mathbf{N}$.

The conditions, under which the subset of \mathbf{M} is a bounded poset, can be obtained from Proposition 2.10 and Proposition 2.11 immediately.

Proposition 2.12. Let $\{\bar{0}, \bar{1}\} \subseteq P \subseteq \mathbf{M}$. Then $(P, \sqsubseteq, \bar{0}, \bar{1})$ (or $(P, \leq, \bar{0}, \bar{1})$) is a bounded poset if and only if $P \subseteq \mathbf{N}$.

An element f of \mathbf{M} is called *convex*, if

$$x \leq y \leq z \Rightarrow f(x) \wedge f(z) \leq f(y).$$

Let \mathbf{C} denote the set of all convex elements of \mathbf{M} .

For $f \in \mathbf{M}$, let $f^L, f^R \in \mathbf{M}$ be given by

$$f^L(x) = \sup\{f(t) \mid t \leq x\}, \quad f^R(x) = \sup\{f(t) \mid t \geq x\}.$$

There exist some equivalent statements of the concept *convex*. (see e.g. [33]).

Proposition 2.13. For $f \in \mathbf{M}$, the following statements are equivalent.

- (i) f is convex.
- (ii) $f = f^L \wedge f^R$.
- (iii) f^α is a convex set for each $\alpha \in (0, 1]$.
- (iv) $f^{\hat{\alpha}}$ is a convex set for each $\alpha \in [0, 1)$.
- (v) f is the minimum of a monotone increasing function and a monotone decreasing one.

Let \mathbf{L} denote the set of all normal convex elements of \mathbf{M} . Then the convolution orders \sqsubseteq and \leq coincide on \mathbf{L} (see [33, Proposition 19, Proposition 37]), and the bounded poset $(\mathbf{L}, \sqsubseteq, \bar{0}, \bar{1})$ is a completely distributive lattice (see [10]).

Proposition 2.14. [10] For any $f, g \in \mathbf{L}$, $f \sqsubseteq g$ if and only if $g^L \leq f^L$ and $f^R \leq g^R$.

An element f of \mathbf{M} is called *upper semicontinuous*, if f^α is closed for any $\alpha \in [0, 1]$. Let \mathbf{L}_u denote the set of all normal, convex and upper semicontinuous elements of \mathbf{M} . The bounded poset $(\mathbf{L}_u, \sqsubseteq, \bar{0}, \bar{1})$ is a completely distributive lattice (see [11]).

Let \mathbf{J} denote the set of all characteristic functions of singletons in $[0, 1]$, i.e.

$$\mathbf{J} = \{\bar{x} \mid x \in [0, 1]\}.$$

There exists an order isomorphism between $([0, 1], \leq)$ and $(\mathbf{J}, \sqsubseteq)$, that is

$$\tau : [0, 1] \longrightarrow \mathbf{J}, \quad x \mapsto \bar{x}.$$

Let $\mathbf{J}^{[2]}$ denote the set of all characteristic functions of all closed subintervals of $[0, 1]$, i.e.

$$\mathbf{J}^{[2]} = \{\overline{[a, b]} \mid 0 \leq a \leq b \leq 1\}.$$

There also exists an order isomorphism between $(\mathbf{I}^{[2]}, \leq_I)$ and $(\mathbf{J}^{[2]}, \sqsubseteq)$, that is

$$\tau : \mathbf{I}^{[2]} \longrightarrow \mathbf{J}^{[2]}, \quad [a, b] \mapsto \overline{[a, b]}.$$

Obviously,

$$\mathbf{J} \subseteq \mathbf{J}^{[2]} \subseteq \mathbf{L}_u \subseteq \mathbf{L} = \mathbf{C} \cap \mathbf{N} \subseteq \mathbf{M}.$$

Definition 2.15. [12] A binary operation $T : \mathbf{L}^2 \longrightarrow \mathbf{L}$ is called a t_r -norm on $(\mathbf{L}, \sqsubseteq)$, if T is a t-norm on $(\mathbf{L}, \sqsubseteq)$, T is closed on \mathbf{J} and $\mathbf{J}^{[2]}$, respectively, and $T(\overline{[0, 1]}, \overline{[a, b]}) = \overline{[0, b]}$.

A binary operation $S : \mathbf{L}^2 \longrightarrow \mathbf{L}$ is called a t_r -conorm on $(\mathbf{L}, \sqsubseteq)$, if S is a t-conorm on $(\mathbf{L}, \sqsubseteq)$, S is closed on \mathbf{J} and $\mathbf{J}^{[2]}$, respectively, and $S(\overline{[0, 1]}, \overline{[a, b]}) = \overline{[a, 1]}$.

There exist two reasons to concern about the poset $(\mathbf{L}, \sqsubseteq)$. On the one hand, the convolution orders \sqsubseteq and \leq coincide on \mathbf{L} , and the bounded poset $(\mathbf{L}, \sqsubseteq, \bar{0}, \bar{1})$ is a complete lattice. On the other hand, $\mathbf{J} \subseteq \mathbf{J}^{[2]} \subseteq \mathbf{L}$ means that \mathbf{L} -fuzzy sets are generalization of type-1 fuzzy sets and interval-valued fuzzy sets.

In the following context, symbols $*$ and Δ always denote any two binary operations on $[0, 1]$, and $*_{\Delta} : \mathbf{M}^2 \longrightarrow \mathbf{M}$ denotes the convolution induced by $*$ and Δ , i.e.

$$(f *_{\Delta} g)(x) = \bigvee_{y * z = x} f(y) \Delta g(z). \quad (2.1)$$

3. Necessity

In this section, we present the necessary conditions such that $*_{\Delta}$ is a t-norm on $(\mathbf{L}, \sqsubseteq)$. Besides, we prove that convolution $*_{\Delta}$ is a t-norm on $(\mathbf{L}, \sqsubseteq)$ if and only if it is a t_r -norm on $(\mathbf{L}, \sqsubseteq)$.

In order to prove Proposition 3.4, we prove some relevant lemmas as follows.

Lemma 3.1. *If for all $x \in [0, 1]$, $\bar{1} *_{\Delta} \bar{x} = \bar{x}$, then for any $A, B \subseteq [0, 1]$,*

$$\bar{A} *_{\Delta} \bar{B} = \overline{A * B}.$$

Proof. Firstly, we show that for any $a, b \in \{0, 1\}$, $a \Delta b = a \wedge b$. If $1 \Delta 0 > 0$, then for any two distinct $x, y \in [0, 1]$, by (2.1)

$$(\bar{1} *_{\Delta} \bar{x})(1 * y) \geq \bar{1}(1) \Delta \bar{x}(y) = 1 \Delta 0 > 0,$$

thus,

$$1 * y \in (\bar{1} *_{\Delta} \bar{x})^{\hat{0}} = \bar{x}^{\hat{0}} = \{x\},$$

so $1 * y = x$. However, it is impossible that for any two distinct $x, y \in [0, 1]$, $1 * y = x$. Therefore, $1 \Delta 0 = 0$. Similarly, we can get

$$0 \Delta 1 = 0 \Delta 0 = 0.$$

Due to

$$\begin{aligned} 1 &= (\bar{1} *_{\Delta} \bar{x})(x) \\ &= \bigvee_{y * z = x} \bar{1}(y) \Delta \bar{x}(z) \\ &\leq \sup(\text{Ran}(\bar{1}) \Delta \text{Ran}(\bar{x})) \\ &= \max(\{0, 1\} \Delta \{0, 1\}) \\ &= \max\{0 \Delta 0, 0 \Delta 1, 1 \Delta 0, 1 \Delta 1\} \\ &= 1 \Delta 1, \end{aligned}$$

we have $1 \Delta 1 = 1$.

Since for any $a, b \in \{0, 1\}$, $a \Delta b = a \wedge b$, it is easy to check that for any $A, B \subseteq [0, 1]$, $\bar{A} *_{\Delta} \bar{B} = \overline{A * B}$ by (2.1). \square

Lemma 3.2. *Let $\mathbf{J}^{[2]} \subseteq P \subseteq \mathbf{L}$. If $*_{\Delta}$ is a t-norm on (P, \sqsubseteq) , then $*_{\Delta}$ is closed on \mathbf{J} and $\mathbf{J}^{[2]}$, respectively, and $*$ is a continuous t-norm.*

Proof. Firstly, $*_{\Delta}$ is closed on \mathbf{J} . In fact, since $*_{\Delta}$ is a t-norm, $\bar{1} *_{\Delta} \bar{x} = \bar{x}$ holds for all $x \in [0, 1]$. By Lemma 3.1, for all $x, y \in [0, 1]$,

$$\bar{x} *_{\Delta} \bar{y} = \overline{x * y}. \quad (3.1)$$

thus $*_{\Delta}$ is closed on \mathbf{J} . Trivially, $*_{\Delta}$ is a t-norm on $(\mathbf{J}, \sqsubseteq)$.

Next, $*_{\Delta}$ is closed on $\mathbf{J}^{[2]}$. Given any $A, B \in \mathbf{I}^{[2]}$, by Lemma 3.1,

$$\overline{A * B} = \bar{A} *_{\Delta} \bar{B} \in P \subseteq \mathbf{L}$$

which means

$$A * B \in \mathbf{I}^{[2]},$$

i.e.

$$\overline{A * B} \in \mathbf{J}^{[2]}.$$

Hence, $*_{\Delta}$ is closed on $\mathbf{J}^{[2]}$.

Finally, $*$ is continuous t-norm. It follows from (3.1) that

$$\tau : [0, 1] \longrightarrow \mathbf{J}, \quad x \mapsto \bar{x}$$

is an isomorphism between posemigroups $([0, 1], *, \leq)$ and $(\mathbf{J}, *_{\Delta}, \sqsubseteq)$. This, together with that $*_{\Delta}$ is a t-norm on $(\mathbf{J}, \sqsubseteq)$, implies that $*$ is a t-norm on $([0, 1], \leq)$ by Proposition 2.2. In the above, we have shown that for any $A, B \in \mathbf{I}^{[2]}$,

$$A * B \in \mathbf{I}^{[2]}.$$

Hence, for each $a \in [0, 1]$,

$$\{a\} * [0, 1] \in \mathbf{I}^{[2]},$$

which means that the section

$$a * (-) : [0, 1] \longrightarrow [0, 1]$$

is continuous. Therefore, t-norm $*$ is continuous. \square

Lemma 3.3. *Let $\mathbf{L}_{\mathbf{u}} \subseteq P \subseteq \mathbf{L}$. If $*_{\Delta}$ is a t-norm on (P, \sqsubseteq) , then Δ is a t-norm.*

Proof. For each $a \in [0, 1]$, define $p_a \in \mathbf{L}_{\mathbf{u}}$ as follows,

$$p_a(x) = \begin{cases} 1, & x = 0, \\ a, & \text{otherwise.} \end{cases}$$

Define the set $\mathcal{P} = \{p_a \mid a \in [0, 1]\}$ and the order-preserving isomorphism $p : ([0, 1], \leq) \longrightarrow (\mathcal{P}, \sqsubseteq)$,

$$p(a) = p_a.$$

By Lemma 3.2, $*$ is a t-norm. Then by (2.1), it is not difficult to check that for all $a, b \in [0, 1]$,

$$p(a) *_{\Delta} p(b) = p(a \Delta b),$$

thus, t-norm $*_{\Delta}$ is closed on \mathcal{P} , and $(\mathcal{P}, *_{\Delta}, \sqsubseteq)$ is a commutative posemigroup. Since the map p is an isomorphism between commutative posemigroups $([0, 1], \Delta, \leq)$ and $(\mathcal{P}, *_{\Delta}, \sqsubseteq)$, and for all $a \in [0, 1]$,

$$a = p_a(1) = (p_a *_{\Delta} \bar{1})(1) = p_a(1) \Delta \bar{1}(1) = a \Delta 1,$$

then Δ is a t-norm. \square

Proposition 3.4. *If $*_{\Delta}$ is a t-norm on $(\mathbf{L}_{\mathbf{u}}, \sqsubseteq)$ or $(\mathbf{L}, \sqsubseteq)$, then*

(i) *$*$ is a continuous t-norm and Δ is a t-norm.*

(ii) *$*_{\Delta}$ is closed on \mathbf{J} and $\mathbf{J}^{[2]}$, respectively.*

(iii) *$\overline{[0, 1]} *_{\Delta} \overline{[a, b]} = \overline{[0, b]}$.*

Proof. (i) and (ii) are straightforward from Lemma 3.2 and Lemma 3.3. By the continuity of $*$ and Lemma 3.1, (iii) holds. \square

Theorem 3.5. *The convolution $*_{\Delta}$ is a t_r -norm on $(\mathbf{L}, \sqsubseteq)$ if and only if it is a t-norm on $(\mathbf{L}, \sqsubseteq)$.*

Proof. Obviously, each t_r -norm is a t-norm. Another direction holds from Proposition 3.4. \square

Proposition 3.6. *If $*_{\Delta}$ is a t-norm on $(\mathbf{L}, \sqsubseteq)$, then Δ is a border continuous t-norm.*

Proof. From Proposition 3.4, $*$ is a continuous t-norm and Δ is a t-norm. For any $a \in [0, 1]$, let $f, g, h \in \mathbf{L}$ be given by $f \equiv 1$,

$$g(x) = \begin{cases} x, & x \in [0, 1), \\ 0, & x = 1, \end{cases}$$

$$h(x) = \begin{cases} 1, & x = 0, \\ a, & x \in (0, 1]. \end{cases}$$

Claim 1, for each $x_0 \in (0, 1)$, $(f *_\Delta g) *_\Delta h(x_0) \geq a$. In fact, for each $y \in [x_0, 1)$, since $*$ is continuous, there exists $z \in [0, 1]$ such that

$$y * z = x_0,$$

and thus,

$$(f *_\Delta g)(x_0) = \bigvee_{s * t = x_0} f(s) * g(t) \geq f(z) \Delta g(y) = 1 \Delta g(y) = y.$$

Hence,

$$(f *_\Delta g)(x_0) = 1.$$

Therefore,

$$((f *_\Delta g) *_\Delta h)(x_0) \geq (f *_\Delta g)(x_0) * h(1) = 1 * a = a.$$

Claim 2, for each $x_0 \in (0, 1)$, $(f *_\Delta (g *_\Delta h))(x_0) \leq a \Delta 1^-$. For this purpose, we only need to check that for any $y_0, z_0 \in [0, 1]$ satisfying $y_0 * z_0 = x_0$,

$$f(y_0) * (g *_\Delta h)(z_0) \leq a \Delta 1^-. \quad (3.2)$$

In fact, for all $z_1, z_2 \in [0, 1]$ satisfying

$$z_1 * z_2 = z_0,$$

since

$$z_1 * z_2 * y_0 = z_0 * y_0 = x_0 > 0,$$

then $z_2 > 0$, thus,

$$g(z_1) \Delta h(z_2) = g(z_1) \Delta a \leq a \Delta 1^-.$$

In the above, we have shown that for all $z_1, z_2 \in [0, 1]$ satisfying $z_1 * z_2 = z_0$,

$$g(z_1) \Delta h(z_2) \leq a \Delta 1^-.$$

Hence,

$$(g *_\Delta h)(z_0) \leq a \Delta 1^-.$$

Therefore, (3.2) holds.

By claim 1 and claim 2, for any $x_0 \in (0, 1)$,

$$a \leq ((f *_\Delta g) *_\Delta h)(x_0) = (f *_\Delta (g *_\Delta h))(x_0) \leq a \Delta 1^-.$$

On the other hand, since Δ is a t-norm, $a \geq a \Delta 1^-$. Therefore, $a = a \Delta 1^-$, i.e. t-norm Δ is border continuous. \square

Proposition 3.7. [1] Suppose that $0 \leq a < b \leq 1$. A t-norm $*$: $[a, b]^2 \longrightarrow [a, b]$ is Archimedean continuous if and only if it is isomorphic to product t-norm T_P or Łukasiewicz t-norm T_L .

Proposition 3.8. [1] A binary function $T : [0, 1]^2 \longrightarrow [0, 1]$ is a continuous t-norm if and only if there exists a family of pairwise disjoint open subintervals $(a_\alpha, b_\alpha)_{\alpha \in A}$ of $[0, 1]$ and a family of continuous Archimedean t-norms $T_\alpha : [a_\alpha, b_\alpha]^2 \longrightarrow [a_\alpha, b_\alpha]$ such that T can be represented by

$$T(x, y) = \begin{cases} T_\alpha(x, y), & (x, y) \in [a_\alpha, b_\alpha]^2, \\ x \wedge y, & \text{otherwise.} \end{cases}$$

Obviously, if the index set $A = \emptyset$, then $T = \wedge$ in the above proposition.

Lemma 3.9. *Let $*$ be a continuous t-norm not equal to \wedge . Then there exist $0 \leq \alpha < \beta \leq 1$ such that the restriction of $*$ on $[\alpha, \beta]$ is a continuous Archimedean t-norm. Besides, the following statements hold.*

- (i) *If $a * b \in (\alpha, \beta)$, then $a, b \in [\alpha, \beta]$.*
- (ii) *if $a_1 * b_1 = a_2 * b_2 \in (\alpha, \beta)$, then $a_1 > a_2$ implies $b_1 < b_2$.*

Proof. By Proposition 3.8, (i) holds immediately. Since the restriction of $*$ on $[\alpha, \beta]$ is isomorphic to T_p or T_L , it is conditionally cancellative. Hence, (ii) holds. \square

Proposition 3.10. *If $*_{\Delta}$ is a t-norm on $(\mathbf{L}, \sqsubseteq)$ and $*$ $\neq \wedge$, then Δ is a left-continuous t-norm.*

Proof. By Proposition 3.6, Δ is a border continuous t-norm. Given any $u, v \in (0, 1)$, we shall show $u \Delta v = u \Delta v^-$. Hence, Δ is a left-continuous t-norm.

By Proposition 3.4, $*$ is a continuous t-norm. Since $*$ $\neq \wedge$, there exist $0 \leq \alpha < \beta \leq 1$ such that the restriction of $*$ on $[\alpha, \beta]$ is a continuous Archimedean t-norm. Choose arbitrary $\lambda \in (\alpha, \beta)$ satisfying

$$\lambda_*^{(3)} \in (\alpha, \beta), \quad (3.3)$$

where $\lambda_*^{(3)}$ denotes $\lambda * \lambda * \lambda$. Let the functions $f, g, h \in \mathbf{L}$ be given by

$$f(x) = \begin{cases} \max\{0, x - \lambda + v\}, & x \in (\lambda * \lambda, \lambda), \\ 1, & x = \lambda, \\ 0, & \text{otherwise,} \end{cases}$$

$$g(x) = \begin{cases} -x + \lambda + 1, & x \in (\lambda, 1], \\ 0, & \text{otherwise,} \end{cases}$$

$$h(x) = \begin{cases} u, & x \in [\lambda, 1), \\ 1, & x = 1, \\ 0, & \text{otherwise.} \end{cases}$$

Claim 1, $((f *_{\Delta} g) *_{\Delta} h)(\lambda_*^{(3)}) \geq u \Delta v$. Since the restriction of $*$ on $[\alpha, \beta]$ is a t-norm isomorphic to T_p or T_L , and

$$\lambda * \lambda \in (\alpha, \beta),$$

then there exists a strictly increasing sequence $\{x_n\}_{n \in \mathbb{N}}$ and a strictly decreasing sequence $\{y_n\}_{n \in \mathbb{N}}$ in (α, β) , both of them converge to λ , and for all $n \in \mathbb{N}$

$$x_n * y_n = \lambda * \lambda. \quad (3.4)$$

It is easy to see that both $\{f(x_n)\}_{n \in \mathbb{N}}$ and $\{g(y_n)\}_{n \in \mathbb{N}}$ are monotone increasing, and

$$\lim_{n \rightarrow \infty} f(x_n) = f(\lambda^-) = v, \quad \lim_{n \rightarrow \infty} g(y_n) = g(\lambda^+) = 1.$$

Since Δ is a border continuous t-norm,

$$\lim_{n \rightarrow \infty} f(x_n) \Delta g(y_n) = \lim_{n \rightarrow \infty} f(x_n) \Delta \lim_{n \rightarrow \infty} g(y_n) = v \Delta 1 = v.$$

This, together with (3.4), implies that

$$(f *_{\Delta} g)(\lambda * \lambda) = \bigvee_{y * z = \lambda * \lambda} f(y) \Delta g(z) \geq \lim_{n \rightarrow \infty} f(x_n) \Delta g(y_n) = v.$$

Hence,

$$((f *_{\Delta} g) *_{\Delta} h)(\lambda_*^{(3)}) \geq (f *_{\Delta} g)(\lambda * \lambda) \Delta h(\lambda) \geq v \Delta u.$$

Claim 2, $(f *_\Delta (g *_\Delta h))(\lambda_*^{(3)}) \leq u \Delta v^-$. For this purpose, we only need to show that for all $s, t \in [0, 1]$ satisfying

$$s * t = \lambda_*^{(3)}, \quad (3.5)$$

the following inequality holds

$$f(s) \Delta (g *_\Delta h)(t) \leq u \Delta v^- \quad (3.6)$$

case by case.

Case 1, $s \notin (\lambda * \lambda, \lambda]$. Since $f(s) = 0$, then (3.6) holds.

Case 2, $s = \lambda$. Since the restriction of $*$ on $[\alpha, \beta]$ is a conditionally cancellative t-norm, $s * t = \lambda_*^{(3)}$, and $\lambda_*^{(3)} \in (\alpha, \beta)$, then

$$t = \lambda * \lambda.$$

Hence, to prove (3.6), we only need to show

$$(g *_\Delta h)(\lambda * \lambda) = 0. \quad (3.7)$$

For all $b, c \in [0, 1]$ satisfying

$$b * c = \lambda * \lambda,$$

if $b \leq \lambda$, then

$$g(b) \Delta h(c) = 0 \Delta h(c) = 0,$$

if $b > \lambda$, since $b * c = \lambda * \lambda$, we have $c < \lambda$ from Lemma 3.9, and thus,

$$g(b) \Delta h(c) = g(b) \Delta 0 = 0.$$

In the above, we have shown that for all $b, c \in [0, 1]$ satisfying $b * c = \lambda * \lambda$,

$$g(b) \Delta h(c) = 0.$$

Hence, (3.7) holds.

Case 3, $s \in (\lambda * \lambda, \lambda)$. Since

$$f(s) < f(\lambda^-) = v,$$

to prove (3.6), we only need to show

$$(g *_\Delta h)(t) \leq u. \quad (3.8)$$

Suppose that $b, c \in [0, 1]$ and

$$b * c = t.$$

If $c < 1$, then $h(c) \leq u$, hence

$$g(b) \Delta h(c) \leq u.$$

If $c = 1$, then $b = t$. In this case, on the one hand,

$$g(b) \Delta h(c) = g(t) \Delta h(1) = g(t) \Delta 1 = g(t). \quad (3.9)$$

On the other hand, by (3.3) and (3.5),

$$s * t = \lambda_*^{(3)} \in (\alpha, \beta).$$

This, together with $s > \lambda * \lambda$, implies $t < \lambda$ from Lemma 3.9. Due to $t < \lambda$,

$$g(t) = 0.$$

Therefore, by (3.9),

$$g(b) \Delta h(c) = g(t) = 0 \leq u.$$

In the above, we have shown that for all $b, c \in [0, 1]$ with $b * c = t$,

$$g(b) \Delta h(c) \leq u.$$

Hence, (3.8) holds.

By claim 1 and claim 2,

$$u \Delta v = ((f *_\Delta g) *_\Delta h)(\lambda_*^{(3)}) = (f *_\Delta (g *_\Delta h))(\lambda_*^{(3)}) \leq u \Delta v^-.$$

Since Δ is a t-norm, $u \Delta v \geq u \Delta v^-$. Hence, $u \Delta v = u \Delta v^-$. Therefore, Δ is left-continuous. \square

As a conclusion of this section, we give the necessary conditions such that $*_{\Delta}$ is a t-norm on $(\mathbf{L}, \sqsubseteq)$.

Proposition 3.11. *Let $*_{\Delta}$ be a t-norm on $(\mathbf{L}, \sqsubseteq)$. Then $*$ is a continuous t-norm, and Δ is a border continuous t-norm. Besides, Δ is left-continuous whenever $*$ $\neq \wedge$.*

Proof. It follows from Proposition 3.4, Proposition 3.6 and Proposition 3.10. \square

4. Sufficiency

In this section, we show that the necessary conditions given in Proposition 3.11 is also sufficient such that $*_{\Delta}$ is a t-norm on $(\mathbf{L}, \sqsubseteq)$. Hence, we obtain the necessary and sufficient conditions as desired.

4.1. Closure

In this subsection, we give the sufficient conditions such that $*_{\Delta}$ is closed on \mathbf{L} .

Lemma 4.1. *Suppose that $\{A_i\}_{i \in \Lambda}$ is a family of convex subsets of $[0, 1]$. If $A_i \cap A_j \neq \emptyset$ whenever $A_i \neq \emptyset$ and $A_j \neq \emptyset$, then $\bigcup_{i \in \Lambda} A_i$ is convex.*

Proof. For any $i, j \in \Lambda$, $x \in A_i, y \in A_j$, since $A_i \cap A_j \neq \emptyset$, and both A_i and A_j are convex, then $A_i \cup A_j$ is convex, hence

$$[x \wedge y, x \vee y] \subseteq A_i \cup A_j \subseteq \bigcup_{i \in \Lambda} A_i.$$

Therefore, $\bigcup_{i \in \Lambda} A_i$ is convex. \square

Proposition 4.2. *If t-norm $*$ is continuous and t-norm Δ is continuous at $(1, 1)$, then $*_{\Delta}$ is closed on \mathbf{L} .*

Proof. For any $f, g \in \mathbf{L}$, $f *_{\Delta} g$ is normal. In fact, for any $a \in [0, 1]$, since Δ is continuous at $(1, 1)$, then there exist $b_1, c_1 \in [0, 1]$ satisfying $b_1 \Delta c_1 > a$. By Proposition 2.7,

$$(f *_{\Delta} g)^{\hat{a}} = \bigcup_{b \Delta c > a} f^b * g^c \supseteq f^{b_1} * g^{c_1} \neq \emptyset.$$

Therefore, $f *_{\Delta} g$ is normal.

Next, $f *_{\Delta} g$ is convex. Given any $a \in [0, 1]$, we claim that for all $b_1, b_2, c_1, c_2 \in [0, 1]$, if $b_1 \Delta c_1 > a$, $b_2 \Delta c_2 > a$, and both $f^{b_1} * g^{c_1}$ and $f^{b_2} * g^{c_2}$ are non-empty, then $(f^{b_1} * g^{c_1}) \cap (f^{b_2} * g^{c_2})$ is non-empty. In fact, on the one hand,

$$f^{b_1} * g^{c_1} \supseteq (f^{b_1} \cap f^{b_2}) * (g^{c_1} \cap g^{c_2})$$

and

$$f^{b_2} * g^{c_2} \supseteq (f^{b_1} \cap f^{b_2}) * (g^{c_1} \cap g^{c_2})$$

imply that

$$(f^{b_1} * g^{c_1}) \cap (f^{b_2} * g^{c_2}) \supseteq (f^{b_1} \cap f^{b_2}) * (g^{c_1} \cap g^{c_2}) = f^{b_1 \vee b_2} * g^{c_1 \vee c_2}.$$

On the other hand, since $f^{b_1} * g^{c_1}$ and $f^{b_2} * g^{c_2}$ are non-empty, all $f^{b_i}, g^{c_i}, i = 1, 2$ are non-empty. Therefore, $(f^{b_1} * g^{c_1}) \cap (f^{b_2} * g^{c_2})$ is non-empty as desired.

By Lemma 4.1, the above claim implies that

$$\bigcup_{b \Delta c > a} f^b * g^c = (f *_{\Delta} g)^{\hat{a}}$$

is convex. By Proposition 2.13, $f *_{\Delta} g$ is convex. Hence, $f *_{\Delta} g \in \mathbf{L}$. \square

4.2. Monotone

In this subsection, we show the sufficient conditions such that $*_{\Delta} : \mathbf{L}^2 \longrightarrow \mathbf{L}$ is monotone increasing in each component.

A non-empty convex subset of $[0, 1]$ is called an *subinterval* of $[0, 1]$. Let \mathcal{I} denote the set of all subinterval of $[0, 1]$. Recall that $\mathbf{I}^{[2]}$ is the set of all closed subintervals of $[0, 1]$, hence $\mathbf{I}^{[2]} \subsetneq \mathcal{I}$.

Define the relation \preceq on \mathcal{I} as follows.

$$A \preceq B, \text{ if } A \wedge B = A.$$

Recall that $A \wedge B = \{x \wedge y \mid x \in A, y \in B\}$.

Example 4.3. Let $0 \leq a_1 < b_1 \leq 1$ and $0 \leq a_2 < b_2 \leq 1$. If $a_1 \leq a_2, b_1 \leq b_2$, then $[a_1, b_1] \preceq [a_2, b_2]$ and $(a_1, b_1) \preceq (a_2, b_2)$.

The following property were provided by Hu and Wang [15]. Here, we give a new proof.

Proposition 4.4. For all $f, g \in \mathbf{L}$, $f \sqsubseteq g$ if and only if $f^{\hat{a}} \preceq g^{\hat{a}}$ for all $a \in [0, 1]$.

Proof. For any $a \in [0, 1]$,

$$(f \wedge g)^{\hat{a}} = \bigcup_{b \wedge c > a} f^b \wedge g^c = \left(\bigcup_{b > a} f^b \right) \wedge \left(\bigcup_{c > a} g^c \right) = f^{\hat{a}} \wedge g^{\hat{a}},$$

thus $f \sqsubseteq g$

$$\Leftrightarrow f \wedge g = f$$

$$\Leftrightarrow (f \wedge g)^{\hat{a}} = f^{\hat{a}} \text{ for all } a \in [0, 1]$$

$$\Leftrightarrow f^{\hat{a}} \wedge g^{\hat{a}} = f^{\hat{a}} \text{ for all } a \in [0, 1]$$

$$\Leftrightarrow f^{\hat{a}} \preceq g^{\hat{a}} \text{ for all } a \in [0, 1].$$

□

Definition 4.5. For any non-empty subset A of $[0, 1]$, define

$$\uparrow A = \bigcup_{x \in A} [x, 1]$$

and

$$\downarrow A = \bigcup_{x \in A} [0, x].$$

Obviously, for each $A \in \mathcal{I}$, $A = \uparrow A \cap \downarrow A$. Briefly, we write $\downarrow a$ and $\uparrow a$ instead of $\downarrow \{a\}$ and $\uparrow \{a\}$, respectively.

Lemma 4.6. For any $A, B \in \mathcal{I}$, $A \wedge B \in \mathcal{I}$. Besides, $A = B$ if and only if $\uparrow A = \uparrow B$ and $\downarrow A = \downarrow B$.

Lemma 4.7. For any $A, B \in \mathcal{I}$,

$$\uparrow(A \wedge B) = \uparrow A \cup \uparrow B, \quad \downarrow(A \wedge B) = \downarrow A \cap \downarrow B.$$

Proof. $x \in \uparrow(A \wedge B)$.

$$\Leftrightarrow \text{there exist } a \in A, b \in B \text{ such that } x \geq a \wedge b.$$

$$\Leftrightarrow \text{there exist } a \in A, b \in B \text{ such that } x \geq a \text{ or } x \geq b.$$

$$\Leftrightarrow x \in \uparrow A \text{ or } x \in \uparrow B.$$

$$\Leftrightarrow x \in \uparrow A \cup \uparrow B.$$

Similarly, $\downarrow(A \wedge B) = \downarrow A \cap \downarrow B$.

□

Lemma 4.8. For $A, B \in \mathcal{I}$, $A \preceq B$ if and only if $\uparrow B \subseteq \uparrow A$ and $\downarrow A \subseteq \downarrow B$.

Proof. $A \preceq B$.

$$\Leftrightarrow A \wedge B = A.$$

$$\Leftrightarrow \uparrow(A \wedge B) = \uparrow A \text{ and } \downarrow(A \wedge B) = \downarrow A \text{ (by Lemma 4.6).}$$

$$\Leftrightarrow \uparrow A \cup \uparrow B = \uparrow A \text{ and } \downarrow A \cap \downarrow B = \downarrow A \text{ (by Lemma 4.7).}$$

$$\Leftrightarrow \uparrow B \subseteq \uparrow A \text{ and } \downarrow A \subseteq \downarrow B.$$

□

Lemma 4.9. Suppose that $\{A_i\}_{i \in \Lambda}$ is a family of subinterval of $[0, 1]$. Then

$$\uparrow(\bigcup_{i \in \Lambda} A_i) = \bigcup_{i \in \Lambda} \uparrow A_i, \quad \downarrow(\bigcup_{i \in \Lambda} A_i) = \bigcup_{i \in \Lambda} \downarrow A_i.$$

Furthermore, if $\bigcap_{i \in \Lambda} A_i \neq \emptyset$, then

$$\uparrow(\bigcap_{i \in \Lambda} A_i) = \bigcap_{i \in \Lambda} \uparrow A_i, \quad \downarrow(\bigcap_{i \in \Lambda} A_i) = \bigcap_{i \in \Lambda} \downarrow A_i$$

Proof. It is trivial that

$$\uparrow(\bigcup_{i \in \Lambda} A_i) = \bigcup_{i \in \Lambda} \uparrow A_i, \quad \downarrow(\bigcup_{i \in \Lambda} A_i) = \bigcup_{i \in \Lambda} \downarrow A_i.$$

In the following, we shall show that if $\bigcap_{i \in \Lambda} A_i \neq \emptyset$, then

$$\uparrow(\bigcap_{i \in \Lambda} A_i) = \bigcap_{i \in \Lambda} \uparrow A_i.$$

Obviously,

$$\uparrow(\bigcap_{i \in \Lambda} A_i) \subseteq \bigcap_{i \in \Lambda} \uparrow A_i.$$

Conversely, suppose

$$\bigcap_{i \in \Lambda} \uparrow A_i \not\subseteq \uparrow(\bigcap_{i \in \Lambda} A_i),$$

i.e. there exists $x_0 \in \bigcap_{i \in \Lambda} \uparrow A_i$ satisfying $x_0 \notin \uparrow(\bigcap_{i \in \Lambda} A_i)$. In this case, given any $a_0 \in \bigcap_{i \in \Lambda} A_i$, since $x_0 \notin \uparrow(\bigcap_{i \in \Lambda} A_i)$, then $x_0 < a_0$. For each $i \in \Lambda$, since $x_0 \in \bigcap_{j \in \Lambda} \uparrow A_j \subseteq \uparrow A_i$, there exists $a_i \in A_i$ such that $x_0 \geq a_i$. Since $a_i \leq x_0 < a_0$, where both a_i and a_0 belong to the convex set A_i , then $x_0 \in A_i$. Hence, $x_0 \in \bigcap_{i \in \Lambda} A_i$, which is contradict to the assumption $x_0 \notin \uparrow(\bigcap_{i \in \Lambda} A_i)$. Therefore,

$$\bigcap_{i \in \Lambda} \uparrow A_i \subseteq \uparrow(\bigcap_{i \in \Lambda} A_i),$$

thus

$$\bigcap_{i \in \Lambda} \uparrow A_i = \uparrow(\bigcap_{i \in \Lambda} A_i).$$

Dually, if $\bigcap_{i \in \Lambda} A_i \neq \emptyset$, then

$$\downarrow(\bigcap_{i \in \Lambda} A_i) = \bigcap_{i \in \Lambda} \downarrow A_i.$$

□

Proposition 4.10. Suppose that $\{A_i\}_{i \in \Lambda}$ and $\{B_i\}_{i \in \Lambda}$ are two families of subintervals of $[0, 1]$, and $A_i \preceq B_i$ for each $i \in \Lambda$. If both $\bigcup_{i \in \Lambda} A_i$ and $\bigcup_{i \in \Lambda} B_i$ are convex, then

$$\bigcup_{i \in \Lambda} A_i \preceq \bigcup_{i \in \Lambda} B_i$$

If both $\bigcap_{i \in \Lambda} A_i$ and $\bigcap_{i \in \Lambda} B_i$ are non-empty, then

$$\bigcap_{i \in \Lambda} A_i \preceq \bigcap_{i \in \Lambda} B_i.$$

Proof. By Lemma 4.8, for any $i \in \Lambda$,

$$\uparrow B_i \subseteq \uparrow A_i, \quad \downarrow A_i \subseteq \downarrow B_i.$$

Hence,

$$\bigcup_{i \in \Lambda} \uparrow B_i \subseteq \bigcup_{i \in \Lambda} \uparrow A_i, \quad \bigcup_{i \in \Lambda} \downarrow A_i \subseteq \bigcup_{i \in \Lambda} \downarrow B_i. \quad (4.1)$$

If both $\bigcup_{i \in \Lambda} A_i$ and $\bigcup_{i \in \Lambda} B_i$ are convex, by (4.1) and Lemma 4.9,

$$\uparrow(\bigcup_{i \in \Lambda} B_i) \subseteq \uparrow(\bigcup_{i \in \Lambda} A_i), \quad \downarrow(\bigcup_{i \in \Lambda} A_i) \subseteq \downarrow(\bigcup_{i \in \Lambda} B_i).$$

This, together with Lemma 4.8, implies

$$\bigcup_{i \in \Lambda} A_i \preceq \bigcup_{i \in \Lambda} B_i.$$

Similarly, if both $\bigcap_{i \in \Lambda} A_i$ and $\bigcap_{i \in \Lambda} B_i$ are non-empty, then

$$\bigcap_{i \in \Lambda} A_i \preceq \bigcap_{i \in \Lambda} B_i.$$

□

Proposition 4.11. *For all $f, g \in \mathbf{L}$, the following statements are equivalent.*

- (i) $f \sqsubseteq g$.
- (ii) $f^{\hat{a}} \preceq g^{\hat{a}}$ for all $a \in (0, 1)$.
- (iii) $f^a \preceq g^a$ for all $a \in (0, 1)$.

Proof. (i) \Rightarrow (ii). It follows from Proposition 4.4 immediately.

(ii) \Rightarrow (iii). For any $a \in (0, 1)$, by Proposition 4.10,

$$f^a = \bigcap_{b \in (0, a)} f^{\hat{b}} \preceq \bigcap_{b \in (0, a)} g^{\hat{b}} = g^a.$$

(iii) \Rightarrow (i). For any $a \in [0, 1)$, by Proposition 4.10,

$$f^{\hat{a}} = \bigcup_{b \in (a, 1)} f^b \preceq \bigcup_{b \in (a, 1)} g^b = g^{\hat{a}}.$$

By Proposition 4.4, $f \sqsubseteq g$.

□

Lemma 4.12. *If $*$ is a continuous t-norm, then for $A, B, C \in \mathcal{I}$, $A \preceq B$ implies $A * C \preceq B * C$.*

Proof. If $A \preceq B$, then $\uparrow A \subseteq \uparrow B$ and $\downarrow B \subseteq \downarrow A$, hence,

$$\uparrow(A * C) = \uparrow A * \uparrow C \subseteq \uparrow B * \uparrow C = \uparrow(B * C),$$

$$\downarrow(B * C) = \downarrow B * \downarrow C \subseteq \downarrow A * \downarrow C = \downarrow(A * C).$$

Therefore, $A * C \preceq B * C$.

□

Proposition 4.13. *Suppose that $*$ is a continuous t-norm and Δ is a border continuous t-norm. Then for all $f_1, f_2, g \in \mathbf{L}$, $f_1 \sqsubseteq f_2$ implies $f_1 *_{\Delta} g \sqsubseteq f_2 *_{\Delta} g$.*

Proof. On the one hand, for all $a \in (0, 1)$ and $i = 1, 2$,

$$(f_i *_{\Delta} g)^{\hat{a}} = \bigcup_{b \Delta c > a} f_i^b * g^c = \bigcup_{b \Delta c > a, (b, c) \in (0, 1)^2} f_i^b * g^c,$$

where the second equality follows from the fact that $a > 0$, t-norm Δ is continuous at $(1, 1)$, and for all $\alpha, \beta \in [0, 1]$, $\alpha \leq \beta$ implies $f_i^{\beta} \subseteq f_i^{\alpha}$. On the other hand, by Proposition 4.11, $f_1^b \preceq f_2^b$ for all $b \in (0, 1)$, then by Lemma 4.12, for all $b, c \in (0, 1)$,

$$f_1^b * g^c \preceq f_2^b * g^c.$$

From Proposition 4.10, for all $a \in (0, 1)$,

$$(f_1 *_{\Delta} g)^{\hat{a}} = \bigcup_{b \Delta c > a, (b, c) \in (0, 1)^2} f_1^b * g^c \preceq \bigcup_{b \Delta c > a, (b, c) \in (0, 1)^2} f_2^b * g^c = (f_2 *_{\Delta} g)^{\hat{a}}.$$

Hence, $f_1 *_{\Delta} g \sqsubseteq f_2 *_{\Delta} g$.

□

4.3. Associativity

In this subsection, we give the sufficient conditions such that $*_{\Delta}$ is associative on \mathbf{L} .

Proposition 4.14. *Let $*$ and Δ be two t -norms. If Δ is left-continuous, then $*_{\Delta}$ is associative on \mathbf{M} .*

Proof. Let $f, g, h \in \mathbf{M}$. For any $a \in [0, 1]$,

$$\begin{aligned} ((f *_{\Delta} g) *_{\Delta} h)(a) &= \sup\{(f *_{\Delta} g)(u) \Delta h(z) \mid u * z = a\} \\ &= \sup\{\sup\{f(x) \Delta g(y) \mid x * y = u\} \Delta h(z) \mid u * z = a\} \\ &= \sup\{\sup\{f(x) \Delta g(y) \Delta h(z) \mid x * y = u\} \mid u * z = a\} \\ &= \sup\{f(x) \Delta g(y) \Delta h(z) \mid x * y * z = u\}. \end{aligned}$$

Similarly,

$$(f *_{\Delta} (g *_{\Delta} h))(a) = \sup\{f(x) \Delta g(y) \Delta h(z) \mid x * y * z = u\}.$$

Hence, $*_{\Delta}$ is associative on \mathbf{M} . □

Recall that for each $f \in \mathbf{L}$, $f = f^L \wedge f^R$, where f^L is monotone increasing and f^R is monotone decreasing.

Definition 4.15. For each $f \in \mathbf{L}$, the subinterval f^+ of $[0, 1]$ is defined by

$$f^+ = \cup\{[0, a] \mid f \text{ is monotone increasing on } [0, a]\}$$

Lemma 4.16. *For each $f \in \mathbf{L}$, $x \in f^+$ if and only if $f^R(x) = 1$.*

Proof. If $x \in f^+$, then f is monotone increasing on $[0, x]$, thus

$$f^R(x) = \bigvee f(\uparrow x) = \bigvee f(\uparrow 0) = 1.$$

Conversely, suppose that $f^R(x) = 1$. For any $a \leq x$, since f^R is monotone decreasing,

$$f^R(a) \geq f^R(x) = 1,$$

then

$$f(a) = f^L(a) \wedge f^R(a) = f^L(a) \wedge 1 = f^L(a),$$

thus

$$f|_{[0, x]} = f^L|_{[0, x]}.$$

Therefore, f is monotone increasing on $[0, x]$, i.e. $x \in f^+$. □

Proposition 4.17. *If Δ is a border-continuous t -norm, then \wedge_{Δ} is associative on \mathbf{L} .*

Proof. Claim 1, for all $f, g \in \mathbf{L}$,

$$(f \wedge_{\Delta} g)(x) = \begin{cases} f(x) \vee g(x), & x \in f^+ \cap g^+, \\ f(x), & x \in g^+ \setminus f^+, \\ g(x), & x \in f^+ \setminus g^+, \\ f(x) \Delta g(x), & x \notin f^+ \cup g^+. \end{cases}$$

It is easy to see

$$(f \wedge_{\Delta} g)(x) = \bigvee_{a \wedge b = x} f(a) \Delta g(b) = \max\{\bigvee (f(x) \Delta g(\uparrow x)), \bigvee (f(\uparrow x) \Delta g(x))\}.$$

For all $x \in [0, 1]$, let

$$F_1(x) = \bigvee (f(x) \Delta g(\uparrow x)),$$

$$F_2(x) = \bigvee (f(\uparrow x) \Delta g(x)),$$

then

$$(f \wedge_{\Delta} g)(x) = F_1(x) \vee F_2(x). \quad (4.2)$$

In the following, we shall show that

$$F_1(x) = \begin{cases} f(x), & x \in g^+, \\ f(x) \Delta g(x), & x \notin g^+, \end{cases} \quad (4.3)$$

$$F_2(x) = \begin{cases} g(x), & x \in f^+, \\ f(x) \Delta g(x), & x \notin f^+. \end{cases}$$

These, together with (4.2), imply that claim 1 holds.

If $x \in g^+$, by Lemma 4.16,

$$g^R(x) = \bigvee g(\uparrow x) = 1.$$

Since Δ is border continuous,

$$F_1(x) = \bigvee (f(x) \Delta g(\uparrow x)) = f(x) \Delta \bigvee g(\uparrow x) = f(x).$$

If $x \notin g^+$, since g is monotone decreasing on $[x, 1]$,

$$\max g(\uparrow x) = g(x),$$

thus

$$F_1(x) = \bigvee (f(x) \Delta g(\uparrow x)) = f(x) \Delta g(x).$$

Hence, (4.3) holds. The proof of F_2 is similar.

Claim 2, for $f, g \in \mathbf{L}$, $(f \wedge_{\Delta} g)^+ = f^+ \cap g^+$. Since both intervals f^+ and g^+ contain the point 0, suppose

$$f^+ \subseteq g^+$$

without loss of the generality. By claim 1,

$$(f \wedge_{\Delta} g)(x) = f(x) \vee g(x), \quad \text{for } x \in f^+ \cap g^+.$$

This, together with that both f and g are monotone increasing on $f^+ \cap g^+$, implies that $f \wedge_{\Delta} g$ is monotone increasing on $f^+ \cap g^+$. Hence,

$$f^+ \cap g^+ \subseteq (f \wedge_{\Delta} g)^+.$$

Conversely, in order to show

$$(f \wedge_{\Delta} g)^+ \subseteq f^+ \cap g^+ = f^+,$$

given any $x_0 \in [0, 1] \setminus f^+$, we shall show that $x_0 \notin (f \wedge_{\Delta} g)^+$. Hence, claim 2 holds.

In fact, for each $x \in [0, 1] \setminus f^+$, since $f^+ \subseteq g^+$, then either $x \in g^+ \setminus f^+$ or $x \notin g^+ = f^+ \cup g^+$. In each case, we always have

$$(f \wedge_{\Delta} g)(x) \leq f(x)$$

from claim 1. This, together with the fact $\uparrow x_0 \subseteq [0, 1] \setminus f^+$, implies

$$\bigvee (f \wedge_{\Delta} g)(\uparrow x_0) \leq \bigvee f(\uparrow x_0).$$

Due to $x_0 \notin f^+$, then $f^R(x_0) < 1$, hence

$$(f \wedge_{\Delta} g)^R(x_0) = \bigvee (f \wedge_{\Delta} g)(\uparrow x_0) \leq \bigvee f(\uparrow x_0) = f^R(x_0) < 1,$$

which implies

$$x_0 \notin (f \wedge_{\Delta} g)^+$$

as desired.

For all $f, g, h \in \mathbf{L}$, by claim 1 and claim 2, one easily obtains the following results.

If $x \in f^+ \cap g^+ \cap h^+$, then

$$((f \wedge_{\Delta} g) \wedge_{\Delta} h)(x) = (f \wedge_{\Delta} g)(x) \vee h(x) = f(x) \vee g(x) \vee h(x),$$

$$(f \wedge_{\Delta} (g \wedge_{\Delta} h))(x) = f(x) \vee (g \wedge_{\Delta} h)(x) = f(x) \vee g(x) \vee h(x).$$

If $x \in (g^+ \cap h^+) \setminus f^+$, then

$$((f \wedge_{\Delta} g) \wedge_{\Delta} h)(x) = (f \wedge_{\Delta} g)(x) = f(x),$$

$$(f \wedge_{\Delta} (g \wedge_{\Delta} h))(x) = f(x).$$

If $x \in f^+ \setminus (g^+ \cup h^+)$, then

$$((f \wedge_{\Delta} g) \wedge_{\Delta} h)(x) = (f \wedge_{\Delta} g)(x) \Delta h(x) = g(x) \Delta h(x),$$

$$(f \wedge_{\Delta} (g \wedge_{\Delta} h))(x) = (g \wedge_{\Delta} h)(x) = g(x) \Delta h(x).$$

If $x \notin f^+ \cup g^+ \cup h^+$, then

$$((f \wedge_{\Delta} g) \wedge_{\Delta} h)(x) = (f \wedge_{\Delta} g)(x) \Delta h(x) = f(x) \Delta g(x) \Delta h(x),$$

$$(f \wedge_{\Delta} (g \wedge_{\Delta} h))(x) = f(x) \Delta (g \wedge_{\Delta} h)(x) = f(x) \Delta g(x) \Delta h(x).$$

Similarly, in other cases, the following equality still holds,

$$((f \wedge_{\Delta} g) \wedge_{\Delta} h)(x) = (f \wedge_{\Delta} (g \wedge_{\Delta} h))(x).$$

Therefore, \wedge_{Δ} is associative on \mathbf{L} . □

4.4. Necessary and sufficient conditions

In this subsection, we give the necessary and sufficient conditions under which $*_{\Delta}$ is a t-norm on $(\mathbf{L}, \sqsubseteq)$.

Proposition 4.18. [14] *Let $*$ and Δ be two t-norms. Then convolution $*_{\Delta} : \mathbf{M}^2 \longrightarrow \mathbf{M}$ is commutative and $\bar{1}$ is the unit.*

Theorem 4.19. *The convolution $*_{\Delta}$ is a t-norm on $(\mathbf{L}, \sqsubseteq)$ if and only if all the following conditions hold.*

- (i) $*$ is a continuous t-norm.
- (ii) If $* \neq \wedge$, then Δ is a left-continuous t-norm.
- (iii) If $* = \wedge$, then Δ is a border continuous t-norm.

Proof. (\Rightarrow) It is clear from Proposition 3.11.

(\Leftarrow) By Proposition 4.2, $*_{\Delta}$ is closed on \mathbf{L} . By Proposition 4.18, $*_{\Delta}$ is commutative and has the unit $\bar{1}$. By Proposition 4.13, $*_{\Delta}$ is monotone increasing in each component. If $* \neq \wedge$, then Δ is left-continuous, and thus $*_{\Delta}$ is associative by Proposition 4.14. If $* = \wedge$, then $*_{\Delta} = \wedge_{\Delta}$ is associative by Proposition 4.17. Therefore, $*_{\Delta}$ is a t-norm on $(\mathbf{L}, \sqsubseteq)$. □

Corollary 4.20. *If $*$ is a continuous t-norm and Δ is a left-continuous t-norm, then $*_{\Delta}$ is a t-norm on $(\mathbf{L}, \sqsubseteq)$.*

In general, the computation of convolution is very complex. Here, we provide one kind of convolutions which are t-norms on $(\mathbf{L}, \sqsubseteq)$ and are easily to calculate.

Proposition 4.21. *Let Δ be a border-continuous t-norm. Then \wedge_{Δ} is a t-norm on $(\mathbf{L}, \sqsubseteq)$, and for all $f, g \in \mathbf{L}$,*

$$(f \wedge_{\Delta} g)(x) = \begin{cases} f(x) \vee g(x), & x \in f^+ \cap g^+, \\ f(x), & x \in g^+ \setminus f^+, \\ g(x), & x \in f^+ \setminus g^+, \\ f(x) \Delta g(x), & x \notin f^+ \cup g^+. \end{cases}$$

Proof. It is straightforward from Theorem 4.19 and the proof of Proposition 4.17. □

5. Triangular conorms

In this section, we give the necessary and sufficient conditions under which $*_{\Delta}$ is a t-conorm or t_r -conorm on $(\mathbf{L}, \sqsubseteq)$.

Define $N : [0, 1] \longrightarrow [0, 1]$,

$$N(x) = 1 - x.$$

$$\neg : \mathbf{M} \longrightarrow \mathbf{M}$$

$$\neg f = f \circ N.$$

$$\bar{*} : [0, 1]^2 \longrightarrow [0, 1],$$

$$\bar{*}(x, y) = N(N(x) * N(y)).$$

Obviously, $\neg(\neg f) = f$, $\bar{*} = *$, and $*$ is a t-norm if and only if $\bar{*}$ is a t-conorm.

Remark 5.1. Since N is an isomorphism between the algebras $([0, 1], *)$ and $([0, 1], \bar{*})$, i.e. for all $x, y \in [0, 1]$,

$$N(x * y) = N(x) \bar{*} N(y),$$

then for any $A, B \subseteq [0, 1]$,

$$N(A * B) = N(A) \bar{*} N(B).$$

Lemma 5.2. For any $f \in \mathbf{M}$ and $a \in [0, 1]$,

$$(\neg f)^a = N(f^a), \quad (\neg f)^{\hat{a}} = N(f^{\hat{a}}).$$

Proof. $x \in (\neg f)^a \Leftrightarrow (\neg f)(x) \geq a \Leftrightarrow f(N(x)) \geq a \Leftrightarrow N(x) \in f^a \Leftrightarrow N(N(x)) \in N(f^a) \Leftrightarrow x \in N(f^a)$. Similarly, $(\neg f)^{\hat{a}} = N(f^{\hat{a}})$. \square

Proposition 5.3. If Δ is monotone increasing in each place, then $\neg : (\mathbf{L}, \sqsubseteq) \longrightarrow (\mathbf{L}, \sqsubseteq)$ is an order-reversing isomorphism such that for all $f, g \in \mathbf{L}$,

$$\neg(f *_{\Delta} g) = (\neg f) \bar{*}_{\Delta} (\neg g).$$

Proof. Firstly, for any binary function $*$ on $[0, 1]$ and $a \in [0, 1]$,

$$\begin{aligned} (\neg(f *_{\Delta} g))^{\hat{a}} &= N((f *_{\Delta} g)^{\hat{a}}) \text{ (by Lemma 5.2)} \\ &= N\left(\bigcup_{b \Delta c > a} f^b * g^c\right) \text{ (by Proposition 2.7)} \\ &= \bigcup_{b \Delta c > a} N(f^b * g^c) \\ &= \bigcup_{b \Delta c > a} N(f^b) \bar{*} N(g^c) \text{ (by Remark 5.1)} \\ &= \bigcup_{b \Delta c > a} (\neg f)^b \bar{*} (\neg g)^c \\ &= ((\neg f) \bar{*}_{\Delta} (\neg g))^{\hat{a}}, \end{aligned}$$

hence,

$$\neg(f *_{\Delta} g) = (\neg f) \bar{*}_{\Delta} (\neg g). \quad (5.1)$$

Next, we show that \neg is an order-reversing isomorphism. Obviously, $\neg : \mathbf{L} \longrightarrow \mathbf{L}$ is a bijection. For any $f, g \in \mathbf{L}$, if $f \sqsubseteq g$, then $f \vee_{\Delta} g = g$. By (5.1),

$$\neg g = \neg(f \vee_{\Delta} g) = (\neg f) \bar{\vee}_{\Delta} (\neg g) = (\neg f) \wedge_{\Delta} (\neg g),$$

which means $\neg g \sqsubseteq \neg f$. Hence, \neg is an order-reversing isomorphism. \square

Proposition 5.4. If $*_{\Delta}$ is a t-conorm on $(\mathbf{L}_{\mathbf{u}}, \sqsubseteq)$ or $(\mathbf{L}, \sqsubseteq)$, then

- (i) $*$ is a continuous t -conorm and Δ is a t -norm.
- (ii) $*_{\Delta}$ is closed on \mathbf{J} and $\mathbf{J}^{[2]}$, respectively.
- (iii) $\overline{[0, 1]} *_{\Delta} \overline{[a, b]} = \overline{[a, b]}$.

Proof. Similar to the proof of Proposition 3.4. □

Proposition 5.5. $*_{\Delta}$ is a t -norm on $(\mathbf{L}, \sqsubseteq)$ if and only if $\bar{*}_{\Delta}$ is a t -conorm on $(\mathbf{L}, \sqsubseteq)$.

Proof. If $*_{\Delta}$ is a t -norm, then Δ is a t -norm from Proposition 3.4. By Proposition 2.3 and Proposition 5.3, $\bar{*}_{\Delta}$ is a t -conorm. Another direction is similar. □

Theorem 5.6. The convolution $*_{\Delta}$ is a t -conorm on $(\mathbf{L}, \sqsubseteq)$ if and only if all the following conditions hold.

- (i) $*$ is a continuous t -conorm.
- (ii) If $*$ $\neq \vee$, then Δ is a left-continuous t -norm.
- (iii) If $*$ $= \vee$, then Δ is a border continuous t -norm.

Proof. It follows from Theorem 4.19 and Proposition 5.5 immediately. □

Theorem 5.7. The convolution $*_{\Delta}$ is a t_r -conorm on $(\mathbf{L}, \sqsubseteq)$ if and only if $*_{\Delta}$ is a t -conorm on $(\mathbf{L}, \sqsubseteq)$.

Proof. It is straightforward from Proposition 5.4. □

6. Convolution

In this study, we have investigated the construction of t -norms on $(\mathbf{L}, \sqsubseteq)$ for T2 RFSs. The main results are as follows:

- 1) Under the condition that both $*$ and Δ are two binary operation on $[0, 1]$, and Δ is surjective, we have provided the necessary and sufficient conditions under which $*_{\Delta}$ is a t -norm (t_r -norm) on $(\mathbf{L}, \sqsubseteq)$, see Theorem 3.5 and Theorem 4.19. Dual results of t -conorms and t_r -conorms have been obtained.
- 2) When Δ is a border-continuous t -norm, the calculation of t -norm \wedge_{Δ} can be expressed simply, see Proposition 4.21.
- 3) We have provided the equivalent characterization of convolution order \sqsubseteq on \mathbf{L} through the α -cut and the strong α -cut, respectively, see Proposition 4.11.

For the further work, we will concentrate on characterizations of t -norms on $(\mathbf{L}_{\mathbf{u}}, \sqsubseteq)$.

Acknowledgement

The author acknowledges the support of the Southwest Minzu University Research Startup Funds (No:RQD2024011).

References

- [1] C. Alsina, M. Frank, and B. Schweizer, *Associative functions: triangular norms and copulas*, World Scientific, 2006.
- [2] S. Cherif, N. Baklouti, H. Hagrass, and A. Alimi, “Novel intuitionistic-based interval type-2 fuzzy similarity measures with application to clustering,” *IEEE Transactions on Fuzzy Systems*, vol. 30, no. 5, pp. 1260–1271, 2021.
- [3] C. Cornelis, G. Deschrijver, E. Kerre, “Advances and challenges in interval-valued fuzzy logic,” *Fuzzy sets and systems*, vol. 157, no. 5, pp. 622–627, 2006.
- [4] G. Deschrijver, C. Cornelis, and E. Kerre, “On the representation of intuitionistic fuzzy t-norms and t-conorms,” *IEEE transactions on fuzzy systems*, vol. 12, no. 1, pp. 45–61, 2004.
- [5] G. Deschrijver and E. Kerre, “On the relationship between some extensions of fuzzy set theory,” *Fuzzy sets and systems*, vol. 133, no. 2, pp. 227–35, 2003.
- [6] Z. Feng and P. Shi, “Admissibilization of singular interval-valued fuzzy systems,” *IEEE Transactions on Fuzzy Systems*, vol. 25, no. 6, pp. 1765–1776, 2016.
- [7] R. Fullér and H. Zimmermann, “On Zadeh’s compositional rule of inference,” *In Fuzzy Logic: State of the Art*, Dordrecht: Springer Netherlands, 1993.
- [8] J. Goguen, “L-fuzzy sets,” *Journal of mathematical analysis and applications*, vol. 18, no. 1, pp. 145–174, 1967.
- [9] M. Grabisch, J. Marichal, R. Mesiar, and E. Pap, *Aggregation functions*, Cambridge University Press, 2009.
- [10] J. Harding, C. Walker, and E. Walker, “Lattices of convex normal functions,” *Fuzzy Sets and Systems*, vol. 159, no. 9, pp. 1061–1071, 2008.
- [11] J. Harding, C. Walker, and E. Walker, “Convex normal functions revisited,” *Fuzzy Sets and Systems*, vol. 161, no. 9, pp. 1343–1349, 2010.
- [12] P. Hernández, S. Cubillo, and C. Torres-Blanc, “On t-norms for type-2 fuzzy sets,” *IEEE Transactions on Fuzzy Systems*, vol. 23, no. 4, pp. 1155–1163, 2014.
- [13] P. Hernández, S. Cubillo, and C. Torres-Blanc, “A complementary study on general interval type-2 fuzzy sets,” *IEEE Transactions on Fuzzy Systems*, vol. 30, no. 11, pp. 5034–5043, 2022.
- [14] B. Hu and C. Kwong, “On type-2 fuzzy sets and their t-norm operations,” *Information sciences*, vol. 255, pp. 58–81, 2014.
- [15] B. Hu and C. Wang, “On type-2 fuzzy relations and interval-valued type-2 fuzzy sets,” *Fuzzy Sets and Systems*, vol. 236, pp. 1–32, 2014.
- [16] Z. Jia and J. Qiao, “Extension operators for type-2 fuzzy sets derived from overlap functions,” *Fuzzy Sets and Systems*, vol. 451, pp. 130–156, 2022.
- [17] E. Klement, R. Mesiar, and E. Pap, *Triangular Norms*, Springer Science & Business Media, 2013.
- [18] T. Kumbasar and H. Hagrass, “A self-tuning zSlices-based general type-2 fuzzy PI controller,” *IEEE Transactions on Fuzzy Systems*, vol. 23, no. 4, pp. 991–1013, 2014.
- [19] H. Lai and J. Sun, “The construction of t-norms on product posets by distributive t-subnorms,” *Fuzzy Sets and Systems*, vol. 467, p. 108567, 2023.
- [20] Z. Liu and X. Wang, “On the extensions of overlap functions and grouping functions to fuzzy truth values,” *IEEE Transactions on Fuzzy Systems*, vol. 29, no. 6, pp. 1423–1430, 2020.

- [21] Z. Liu and X. Wang, "The distributivity of extended uninorms over extended overlap functions on the membership functions of type-2 fuzzy sets," *Fuzzy Sets and Systems*, vol. 448, pp. 94–106, 2022.
- [22] E. Mamdani, "Applications of fuzzy algorithms for simple dynamic plant," *Proceedings of the Institution of Electrical Engineers*, vol. 121, no. 12, pp. 1585–1588, 1974.
- [23] E. Mamdani and S. Assilian, "An experiment in linguistic synthesis with a fuzzy logic controller," *International Journal of Machine Studies*, vol. 7, pp. 1–13, 1975.
- [24] J. Mendel, *Explainable Uncertain Rule-Based Fuzzy Systems*, Springer, 2024.
- [25] R. Precup and H. Hellendoorn, "A survey on industrial applications of fuzzy control," *Computers in industry*, vol. 62, no. 3, pp. 213–226, 2011.
- [26] R. Precup, A. Nguyen, and S. Blažič, "A survey on fuzzy control for mechatronics applications," *International Journal of Systems Science*, vol. 55, no. 4, pp. 771–813, 2024.
- [27] B. Schweizer, A. Sklar, *Probabilistic Metric Spaces*, Courier Corporation, 2011.
- [28] M. Sugeno, "An introductory survey of fuzzy control," *Information Sciences*, vol. 36, no. 1-2, pp. 59–83, 1985.
- [29] P. Sussner, "Complete Lattices as Frameworks for Interval and General Type-2 Fuzzy Inference Systems," *IEEE Transactions on Fuzzy Systems*, doi: 10.1109/TFUZZ.2025.3545295.
- [30] Z. Takáč, "Aggregation of fuzzy truth values," *Information Sciences*, vol. 271, pp. 1–13, 2014.
- [31] C. Torres-Blanc, S. Cubillo, and P. Hernández, "Aggregation operators on type-2 fuzzy sets," *Fuzzy Sets and Systems*, vol. 324, pp. 74–90, 2017.
- [32] A. Van der Wal, "Application of fuzzy logic control in industry," *Fuzzy Sets and Systems*, vol. 74, no. 1, pp. 33–41, 1995.
- [33] C. Walker and E. Walker, "The algebra of fuzzy truth values," *Fuzzy sets and systems*, vol. 149, no. 2, pp. 309–348, 2005.
- [34] C. Wang and B. Hu, "Generalized extended fuzzy implications," *Fuzzy Sets and Systems*, vol. 268, pp. 93–109, 2015.
- [35] C. Wang and L. Wan, "Type-2 fuzzy implications and fuzzy-valued approximation reasoning," *International Journal of Approximate Reasoning*, vol. 102, pp. 108–122, 2018.
- [36] X. Wu and G. Chen, "Answering an open problem on t-norms for type-2 fuzzy sets," *Information Sciences*, vol. 522, pp. 124–133, 2020.
- [37] A. Xie, "On the extension of nullnorms and uninorms to fuzzy truth values," *Fuzzy Sets and Systems*, vol. 352, pp. 92–118, 2018.
- [38] L. Zadeh, "Fuzzy sets," *Information and Control*, vol. 8, no. 3, pp. 338–353, 1965.
- [39] L. Zadeh, "Outline of a new approach to the analysis of complex systems and decision processes," *IEEE Transactions on Fuzzy Systems*, vol. 31 no. 1, pp. 28–44, 1973.
- [40] L. Zadeh, "The concept of a linguistic variable and its application to approximate reasoning—I," *Information sciences*, vol. 8, no. 3, pp. 199–249, 1975.
- [41] D. Zhang, "Triangular norms on partially ordered sets," *Fuzzy Sets and Systems*, vol. 153, no. 2, pp. 195–209, 2005.
- [42] W. Zhang and B. Hu, "The distributive laws of convolution operations over meet-convolution and join-convolution on fuzzy truth values," *IEEE Transactions on Fuzzy Systems*, vol. 29, no. 2, pp. 415–426, 2019.

- [43] W. Zhang and B. Hu, “New results on α -cuts of type-2 fuzzy sets,” Fuzzy Sets and Systems, vol. 498, p. 109152, 2025.