

Generalizing the Shell Theorem to Constant Curvature Spaces in All Dimensions and Topologies

Ava K. Tse,^{1,*} Olivia M. Markowich,^{1,*} and Trung V. Phan^{2,†}

¹*Scripps College, Claremont, CA 91711, USA.*

²*Department of Natural Sciences, Scripps and Pitzer Colleges,
Claremont Colleges Consortium, Claremont, CA 91711, USA*

(Dated: November 27, 2025)

A gravitational potential has the *spherical property* when the field outside any uniform spherical shell is indistinguishable from that of a point mass at the center. We present the general potentials that possess this property on constant curvature spaces, using the Euler-Poisson-Darboux identity for spherical means. Our results are consistent with known findings in flat three-dimensional space and reduce to Gurzadyan’s cosmological theorem when the *rescaling factor* is exactly 1. Our approach naturally extends to *nontrivial spatial topologies*.

In cosmology, Gurzadyan’s theorem characterizes the full class of potentials for which a uniform spherical mass, seen from outside, is gravitationally indistinguishable from a point mass at its center [1–4]. This extends the first *shell theorem* in Newtonian classical gravity (point-mass equivalence for the inverse-square law) [5], first established in [6, 7] and later generalized under the name *spherical property*, where the exterior field of a uniform shell coincides with that of a central point mass up to a shell-radius-dependent *rescaling factor* [8, 9] (see Fig. 1A).¹ These results were obtained in three-dimensional Euclidean space \mathbb{R}^3 ; here we further extend them to arbitrary n -dimensional constant curvature spaces, also known as spherical forms, i.e. spherical \mathbb{S}^n , Euclidean \mathbb{R}^n , and hyperbolic \mathbb{H}^n . This extension is made possible by linking the *spherical property* to the Euler–Poisson–Darboux identity – a connection that, to the best of our knowledge, is absent from standard references.

We start by considering the situation in flat space \mathbb{R}^n . Let $\phi(r)$ denote the *translationally invariant and pairwise* potential as a function of the Euclidean separation r ; then, outside a uniform spherical shell of radius b , the potential at a *test point* located at position \vec{a} away

* These authors contributed equally.

† tphan@natsci.claremont.edu

¹ equivalently, the field outside a uniform solid sphere is indistinguishable from that of a central point mass.




FIG. 1. **Generalizing shell theorem to sphere forms in all dimensions and topologies.**

Here, we focus on translationally invariant pairwise interaction potentials. **(A)** The generalized shell theorem: outside a uniform spherical shell, the field is indistinguishable from that of a central point mass, scaled by a radius-dependent factor. **(B)** The field at a *test point* can be calculated by summing the contributions from all surface elements of the spherical shell. **(C)** A list of geometries we are interested in: the Euclidean flat space \mathbb{R}^n , the spherical space \mathbb{S}^n , the hyperbolic space \mathbb{H}^n , and the hypercubic toroidal space \mathbb{T}^n .

from the shell center is given by the spherical mean over all shell elements (see Fig. 1B):

$$\Phi(\vec{a}, b) = \frac{\oint_{S^{n-1}(b)} d\sigma(\hat{\omega}) \phi(|\vec{a} - b\hat{\omega}|)}{\text{Vol}[S^{n-1}(b)]}, \quad (1)$$

where $S^{n-1}(b)$ is the $(n-1)$ -dimensional spherical surface of radius b , $\text{Vol}[S^{n-1}(b)]$ is its total

surface volume, $\hat{\omega}$ is a unit direction vector on the sphere, $d\sigma(\hat{\omega})$ is the surface-area element, and $|\circ|$ denotes the Euclidean norm of a vector quantity \circ (so $a = |\vec{a}| > b$ and $|\hat{\omega}| = 1$). The Euler-Poisson-Darboux (EPD) identity for this spherical mean – at least, away from singularities – is given by [10]:

$$\partial_b^2 \Phi(\vec{a}, b) + \frac{(n-1)}{b} \partial_b \Phi(\vec{a}, b) = \Delta_{\vec{a}} \Phi(\vec{a}, b) . \quad (2)$$

which is partial differential equation (PDE) that any spherical mean function $\Phi(\vec{a}, b)$ should obey. The operator $\Delta_{\vec{a}}$ is the Laplacian with respect to the variable \vec{a} . As a demonstration, we derive this equation for $n = 3$ in Appendix A.

To satisfy the *spherical property*, which specifies how the induced force behaves, we need the potential to take the form:²

$$\Phi(\vec{a}, b) = \phi(a)M(b) + C(b) ,$$

where $M(b)$ and $C(b)$ are functions to be determined;³ $M(b)$ is the *rescaling factor* of the mass (as represented in Fig. 1A2). Place this expression in Eq. (2) gives:

$$-\lambda(b)\phi(a) - \rho(b) = \Delta_{\vec{a}}\phi(a) = \partial_a^2 \phi(a) + \frac{(n-1)}{a} \partial_a \phi(a) \quad (3)$$

where

$$\lambda(b) = -\frac{\partial_b^2 M(b) + \frac{(n-1)}{b} \partial_b M(b)}{M(b)} , \quad \rho(b) = -\frac{\partial_b^2 C(b) + \frac{(n-1)}{b} \partial_b C(b)}{M(b)} . \quad (4)$$

Since the right hand side (RHS) of Eq. (3) does not depend of b , the left hand side (LHS) should also not, therefore $\lambda(b)$ and $\rho(b)$ are constants, i.e. $\lambda(b) = \lambda$ and $\rho(b) = \rho$. We can then solve the PDE in Eq. (3) to obtain the potential function $\phi(a)$ that possesses the *spherical property*, which can be divided into two different cases:

- $\lambda \neq 0$: After a constant shift

$$\phi(a) \rightarrow \phi(a) - \rho/\lambda ,$$

Eq. (3) becomes the Helmholtz equation [11]. For $\lambda > 0$, the general solution is given by a linear combination of the Bessel functions:

$$\phi(a) = A_1 a^{-(n/2-1)} J_{n/2-1}(\lambda^{1/2} a) + A_2 a^{-(n/2-1)} Y_{n/2-1}(\lambda^{1/2} a) ,$$

² so that the force, given by $\vec{F}(\vec{a}, b) = \vec{\nabla}_{\vec{a}} \Phi(\vec{a}, b)$, satisfies $\vec{F}(\vec{a}, b) = \vec{f}(\vec{a})M(b)$, where $\vec{f}(\vec{a}) = \vec{\nabla}_{\vec{a}} \phi(a)$.

³ in the limit $b \rightarrow 0$, Eq. (1) leads to $\Phi(\vec{a}, 0) = \phi(|\vec{a}|)$, and thus $M(0) = 1$ and $C(0) = 0$.

where the A -values are arbitrary constants; and for $\lambda < 0$, it is obtained by analytic continuation (equivalently, in terms of the modified Bessel functions) [12]. When $n = 3$, this expression – after some algebraic manipulation – can be written as

$$\phi(a) = A_3 a^{-1} \sinh(qa) + A_4 a^{-1} \cosh(qa) \quad \text{for } \lambda = q^2 > 0 \quad (5)$$

or

$$\phi(a) = A_5 a^{-1} \sin(pa) + A_6 a^{-1} \cos(pa) \quad \text{for } \lambda = -p^2 < 0 . \quad (6)$$

The Yukawa potential, which appears in nuclear interactions (e.g. one-pion exchange) [13], is a special case of Eq. (5).

- $\lambda = 0$: This is the case when the *rescale factor* is exactly 1. Eq. (3) becomes the Poisson equation with constant source term ρ , therefore the general solution for $\phi(a)$ is a linear combination between the particular solution (a quadratic function) and the fundamental solution (a Green's function, which is harmonic off the singularity):

$$\phi(a) = -\frac{\rho}{2n} a^2 + A_7 a^{2-n} + A_8 , \quad (7)$$

where the A -values are arbitrary constants. When $n = 3$, this expression is the sum of Hookean and Coulombic potentials:

$$\phi(a) = -\frac{\rho}{2n} a^2 + A_7 a^{-1} + A_8 . \quad (8)$$

For a sanity check, we note that Eq. (5), Eq. (6), and Eq. (8) agree with known results in \mathbb{R}^3 space [8, 9], and Eq. (8) is the same with Gurzadyan's cosmological theorem [1–4, 6, 7]. To find the corresponding function $M(b)$, we solve Eq. (4) with boundary conditions $M(0) = 1$ and $\partial_b M(0) = 0$, which are direct consequences of Eq. (1); the solutions are provided in the Appendix B.

On curved spaces (see Fig. 1C), the analysis mirrors the flat case: replace the Euclidean distance with the geodesic distance and $\Delta_{\vec{a}}$ with the Laplace–Beltrami operator. The Euclidean Euler–Poisson–Darboux identity, i.e. Eq. (2), changes into

$$\partial_b^2 \Phi(\vec{a}, b) + \frac{(n-1)}{b} \cot(b) \partial_b \Phi(\vec{a}, b) = \Delta_{\vec{a}} \Phi(\vec{a}, b)$$

for spherical space \mathbb{S}^n (positive spatial curvature), and

$$\partial_b^2 \Phi(\vec{a}, b) + \frac{(n-1)}{b} \coth(b) \partial_b \Phi(\vec{a}, b) = \Delta_{\vec{a}} \Phi(\vec{a}, b)$$

for hyperbolic space \mathbb{H}^n (negative spatial curvature) [14]. The class of potentials $\phi(a)$ with *spherical property* in these spaces are solutions of an equation similar to Eq. (3), i.e.

$$-\lambda\phi(a) - \rho = \Delta_{\vec{a}}\phi(a) , \quad (9)$$

where λ and ρ are constants. Therefore:

- In *compact topology* \mathbb{S}^n spaces: For $\lambda \neq 0$, $\phi(a)$ is a linear combination of Gegenbauer functions [12]; and for $\lambda = 0$, since local smoothness of $\phi(a)$ in this compact manifold requires $\rho = 0$, $\phi(a)$ only corresponds to the fundamental solution.
- In *open topology* \mathbb{H}^n spaces: For $\lambda \neq 0$, $\phi(a)$ is a linear combination of associated Legendre functions [12]; and for $\lambda = 0$, $\phi(a)$ is a linear combination between the particular solution and the fundamental solution.

We show the details in Appendix C.

Observations indicate that our universe possesses spatial flatness alongside positive space-time curvature driven by cosmic expansion [15]. An interesting direction to further explore after this work is considering more *nontrivial spatial topologies* [16] (and also for *nonspherical shell*), in which we can apply the same approach in this work. In general, the EPD identity takes the vector form – i.e. Eq. (9) with the potential as a function of the full displacement \vec{a} rather than only its norm:

$$-\lambda\phi(\vec{a}) - \rho = \Delta_{\vec{a}}\phi(\vec{a}) . \quad (10)$$

Because this differential relation is local, it remains unchanged; only the global geometry differs. We work out an example (for generalizing the spherical shell theorem) in flat hypercubic torus \mathbb{T}^n spaces (see Fig. 1C) – with common compactification length L in all directions – in Appendix D.

-
- [1] V. G. Gurzadyan, The Observatory **105**, 42 (1985).
 - [2] B. C. Reed, American Journal of Physics **90**, 394 (2022).
 - [3] C. Carimalo, American Journal of Physics **91**, 1031 (2023).
 - [4] T. V. Phan, arXiv preprint arXiv:2504.02897 (2025).

- [5] I. Newton, *Philosophiae naturalis principia mathematica*, Vol. 1 (G. Brookman, 1833).
- [6] I. N. Sneddon and C. K. Thornhill, Proceedings of the Cambridge Philosophical Society **45**, 318 (1949).
- [7] C. W. Kilmister, The Mathematical Gazette **58**, 129 (1974).
- [8] C. J. Chapman, The Mathematical Gazette **67**, 14 (1983).
- [9] A. Barnes and C. K. Keogh, The Mathematical Gazette **68**, 138 (1984).
- [10] F. John, *Plane waves and spherical means applied to partial differential equations* (Courier Corporation, 2004).
- [11] A. Sommerfeld, *Partial differential equations in physics*, Vol. 1 (Academic press, 1949).
- [12] M. Abramowitz and I. A. Stegun, *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, Vol. 55 (US Government printing office, 1948).
- [13] J. Rowlinson, Physica A: Statistical Mechanics and its Applications **156**, 15 (1989).
- [14] L. V. Nguyen, arXiv preprint arXiv:1201.0314 (2011).
- [15] N. Aghanim *et al.*, Astron. Astrophys **641**, A6 (2020).
- [16] J.-P. Luminet, J. R. Weeks, A. Riazuelo, R. Lehoucq, and J.-P. Uzan, Nature **425**, 593 (2003).

Appendix A: EPD identity in \mathbb{R}^3

In \mathbb{R}^3 , the direction unit vector can be written in geographic angles (θ, φ) as

$$\hat{\omega} = (\sin \theta \cos \varphi, \sin \theta \sin \varphi, \cos \theta) ,$$

and Eq. (1) becomes:

$$\Phi(\vec{a}, b) = \frac{1}{4\pi} \int_0^\pi d\theta \sin \theta \oint d\varphi \phi(\vec{a} - b\hat{\omega}) .$$

Note that here we consider a general $(\vec{a} - b\hat{\omega})$ -dependence, not just on the norm. Our aim is to prove the Euler-Poisson-Darboux identity:

$$\partial_b^2 \Phi(\vec{a}, b) + \frac{2}{b} \partial_b \Phi(\vec{a}, b) = \Delta_{\vec{a}} \Phi(\vec{a}, b) .$$

Let us define $\vec{R} = \vec{a} - b\hat{\omega}$, $\vec{r} = \vec{a} - \vec{R}$, and $r = |\vec{r}| = b$. The Laplacian $\Delta_{\vec{r}}$ in the spherical coordinates (r, θ, φ) centered at \vec{a} is given by:

$$\Delta_{\vec{r}} = \partial_r^2 + \frac{2}{r} \partial_r + \frac{1}{r^2} \Delta_{(\theta, \varphi)} \quad \text{where} \quad \Delta_{(\theta, \varphi)} = \frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta) + \frac{1}{\sin^2 \theta} \partial_\varphi^2 .$$

Due to translational invariance, we have $\Delta_{\vec{a}}\phi(\vec{a} - b\hat{\omega}) = \Delta_{\vec{R}}\phi(\vec{R})$, therefore:

$$\begin{aligned}\Delta_{\vec{a}}\Phi(\vec{a}, b) &= \frac{1}{4\pi} \int_0^\pi d\theta \sin \theta \oint d\varphi \left(\partial_r^2 + \frac{2}{r} \partial_r + \frac{1}{r^2} \Delta_{(\theta, \varphi)} \right) \phi(\vec{a} - b\hat{\omega}) \\ &= \frac{1}{4\pi} \int_0^\pi d\theta \sin \theta \oint d\varphi \left(\partial_b^2 + \frac{2}{b} \partial_b + \frac{1}{b^2} \Delta_{(\theta, \varphi)} \right) \phi(\vec{a} - b\hat{\omega}) .\end{aligned}\tag{A1}$$

In the above equation, to go from the first- to the second-line we use $r = b$, e.g. any variation along the ray \vec{r} is equivalent to varying b , hence $\partial_r = \partial_b$ and $\partial_r^2 = \partial_b^2$.

For the angular Laplacian $\Delta_{(\theta, \varphi)}$ acting on $\phi(\vec{a} - b\hat{\omega})$, we can treat it as a function of the geographic angle variables (θ, φ) , i.e. $\phi(\vec{a} - b\hat{\omega}) = f(\theta, \varphi)$. Since Eq. (A1) can be separated into:

$$\Delta_{\vec{a}}\Phi(\vec{a}, b) = \left(\partial_b^2 + \frac{2}{b} \partial_b \right) \Phi(\vec{a}, b) + I ,\tag{A2}$$

the integration part I can be written as:

$$\begin{aligned}I &= \frac{1}{4\pi b^2} \int_0^\pi d\theta \sin \theta \oint d\varphi \Delta_{(\theta, \varphi)} f(\theta, \varphi) \\ &= \frac{1}{4\pi b^2} \left\{ \oint d\varphi \int_0^\pi d\theta \partial_\theta [\sin \theta \partial_\theta f(\theta, \varphi)] + \int_0^\pi \frac{d\theta}{\sin \theta} \oint d\varphi \partial_\varphi^2 f(\theta, \varphi) \right\}\end{aligned}\tag{A3}$$

The second term here is trivially 0, while the first term after the $\int d\theta$ integration gives

$$[\sin \theta \partial_\theta f(\theta, \varphi)] \Big|_0^\pi ,$$

which is equal to 0 since $\sin \theta = 0$ for both $\theta = 0$ and $\theta = \pi$. Together, $I = 0$, and what are left of Eq. (A2) is exactly the EPD identity.

Appendix B: The *rescaling factor* $M(b)$ in \mathbb{R}^n

We want to find the function $M(b)$ that satisfies the boundary conditions $M(0) = 1$ and $\partial_b M(0) = 0$ while solving the following differential equation:

$$\partial_b^2 M(b) + \frac{n-1}{b} \partial_b M(b) + \lambda M(b) = 0 ,\tag{B1}$$

which is identical to Eq. (4) up to simple algebraic rearrangement. Here we present the solutions for different values of λ :

- $\lambda = 0$: The *rescaling factor* is the identity, i.e. $M(b) = 1$.

- $\lambda > 0$: Define $\lambda = q^2 > 0$, then the *rescaling factor* is given by:

$$M(b) = \Gamma(n/2) (2/qb)^{n/2-1} J_{n/2-1}(qb) . \quad (\text{B2})$$

When $n = 3$, this expression becomes $M(b) = \sin(qb)/qb$.

- $\lambda < 0$: Define $\lambda = -p^2 < 0$, then the *rescaling factor* can be written using modified Bessel function:

$$M(b) = \Gamma(n/2) (2/qb)^{n/2-1} I_{n/2-1}(qb) . \quad (\text{B3})$$

When $n = 3$, this expression becomes $M(b) = \sinh(pb)/pb$.

Appendix C: Potentials possess *spherical property* in \mathbb{S}^n and \mathbb{H}^n

In this Appendix, the A -values are arbitrary constants.

Consider \mathbb{S}^n spaces with curvature $\kappa = k^2 > 0$. For $\lambda \neq 0$, $\phi(a)$ is a linear combination of Gegenbauer functions [12]:

$$\phi(a) = A_9 C_\nu^\alpha [\cos(ka)] + A_{10} Q_\nu^\alpha [\cos(ka)] - \rho/\lambda ,$$

in which

$$\alpha = (n-1)/2 , \quad \nu(\nu+n-1) = \lambda/k^2 .$$

For $\lambda = 0$, local smoothness of $\phi(a)$ in this compact manifold requires $\rho = 0$,⁴ therefore $\phi(a)$ corresponds to a fundamental solution:

$$\phi(a) = A_{11} [2 \sin(ka/2)]^{2-n} + A_{12} .$$

Next, we look at \mathbb{H}^n spaces with curvature $\kappa = -l^2 < 0$: For $\lambda \neq 0$, $\phi(a)$ is a linear combination of associated Legendre functions [12]:

$$\phi(a) = A_{13} P_{-1/2+i\tau}^{-\alpha} [\cosh(la)] + A_{14} Q_{-1/2+i\tau}^{-\alpha} [\cosh(la)] - \rho/\lambda ,$$

in which

$$\alpha = (n-1)/2 , \quad \tau^2 = \lambda/l^2 - \alpha^2 .$$

⁴ since we always have $\int_{\mathbb{S}^n} d^n \vec{a} \Delta_{\vec{a}} \phi(a) = 0$ (properties on a compact manifold), thus from $\Delta_{\vec{a}} \phi(a) = -\rho$ we obtain $\int_{\mathbb{S}^n} d^n \vec{a} (-\rho) = -\rho \text{Vol}(\mathbb{S}^n) = 0$, which leads to $\rho = 0$.

For $\underline{\lambda=0}$, $\phi(a)$ can be found as a linear combination between the particular solution (regular) and the fundamental solution (singular):

$$\phi(a) = \rho \int_0^a da' \frac{\int_0^{a'} da'' S(a'')}{S(a')} + A_{15} \int_a^\infty \frac{da'}{S(a')} + A_{16} ,$$

where we define $S(a) = [\sinh(la)]^{n-1}$. When $n = 3$, this admits a closed form expression:

$$\phi(a) = \frac{\rho}{2l} a \coth(la) + A_{15} [\coth(la) - 1] + A_{16} .$$

Appendix D: Potentials possess *spherical property* in \mathbb{T}^n

Consider hypercubic \mathbb{T}^n spaces with a common compactification length L . We want to find to general solution $\phi(\vec{a})$ of Eq. (10).

For $\underline{\lambda \neq 0}$, $\phi(\vec{a})$ can only be the constant function $-\rho/\lambda$ (so no force) unless λ is an eigenvalue of $\Delta_{\vec{a}}$, i.e. $\lambda = (2\pi|\vec{m}|/L)^2$ with $\vec{m} \in \mathbb{Z}^n \setminus \{0\}$. When this is the case, the most general potential is of the form:

$$\phi(\vec{a}) = c_{\vec{m}} \cos [2\pi(\vec{m}.\vec{a})/L] + s_{\vec{m}} \sin [2\pi(\vec{m}.\vec{a})/L] - \rho/\lambda , \quad (\text{D1})$$

where $c_{\vec{m}}$ and $s_{\vec{m}}$ are arbitrary constants.

For $\underline{\lambda=0}$, similar to the argument made in Appendix C for \mathbb{S}^n spaces, local smoothness of $\phi(\vec{a})$ in this compact manifold (or any compact manifold) requires $\rho = 0$, therefore $\phi(\vec{a})$ corresponds to a fundamental solution:

$$\phi(a) = B_1 \sum_{\vec{m} \in \mathbb{Z}^n \setminus \{0\}} \frac{\exp [i2\pi(\vec{m}.\vec{a})/L]}{(2\pi|\vec{m}|/L)^2} + B_2 ,$$

where i is the unit imaginary number and the B -values are arbitrary constants.