

(2) The simple R -module S_3 is 2-reflexive but not reflexive.

Proof. Let e_i denote the primitive idempotents corresponding to the vertices i in the quiver of R .

(1) Note that we have $\Omega^1(N) \cong S_2$ and we have the exact sequence

$$0 \rightarrow S_3 \rightarrow e_2 R \rightarrow S_2 \rightarrow 0$$

and applying the functor $\text{Hom}_R(-, R)$ to this short exact sequence we obtain the exact sequence

$$0 \rightarrow \text{Hom}_R(S_2, R) \rightarrow \text{Hom}_R(e_2 R, R) \rightarrow \text{Hom}_R(S_3, R) \rightarrow \text{Ext}_R^1(S_2, R) \rightarrow 0.$$

Now the left R -module $\text{Hom}_R(S_2, R)$ has dimension vector $[1, 0, 0, 0]$, the left R -module $\text{Hom}_R(e_2 R, R)$ has dimension vector $[1, 1, 0, 0]$ and the left R -module $\text{Hom}_R(S_3, R)$ has dimension vector $[1, 1, 0, 0]$. This implies that the left R -module $\text{Ext}_R^2(N, R) \cong \text{Ext}_R^1(S_2, R)$ has dimension vector $[1, 0, 0, 0]$ and thus is a non-zero projective left R -module.

(2) Let M be a general R -module such that M is 1-torsionfree (meaning it is a submodule of a projective module), then M is 2-reflexive if and only if the R -module $\text{Ext}_R^2(\text{Tr } M, R)$ is projective by the proposition on page 143 in [AB]. Now the R -module S_3 is 1-torsionfree since it is a submodule of the indecomposable projective R -module $e_2 R$. Now we calculate $\text{Ext}_R^2(\text{Tr } S_3, R)$ and show that this module is projective to conclude that S_3 is 2-reflexive. A minimal projective resolution of S_3 is given by

$$0 \rightarrow e_4 R \rightarrow e_3 R \rightarrow S_3 \rightarrow 0$$

and applying the functor $\text{Hom}_R(-, R)$ to this exact sequence gives

$$0 \rightarrow \text{Hom}_R(S_3, R) \rightarrow \text{Hom}_R(e_3 R, R) \rightarrow \text{Hom}_R(e_4 R, R) \rightarrow \text{Tr}(S_3) \rightarrow 0.$$

Now $\text{Hom}_R(S_3, R)$ has dimension vector $[1, 1, 0, 0]$, $\text{Hom}_R(e_3 R, R)$ has dimension vector $[1, 1, 1, 0]$ and $\text{Hom}_R(e_4 R, R)$ has dimension vector $[0, 0, 1, 1]$. Thus $\text{Tr}(S_3)$ has dimension vector $[0, 0, 0, 1]$ and $\Omega^1(\text{Tr}(S_3))$ is the left simple R -module Y with dimension vector $[0, 0, 1, 0]$. We have the exact sequence of left R -modules:

$$0 \rightarrow Y_2 \rightarrow R e_3 \rightarrow Y \rightarrow 0,$$

where Y_2 is the left R -module with dimension vector $[1, 1, 0, 0]$. Applying the functor $\text{Hom}_R(-, R)$ to this exact sequence gives the exact sequence:

$$0 \rightarrow \text{Hom}_R(Y, R) \rightarrow \text{Hom}_R(R e_3, R) \rightarrow \text{Hom}_R(Y_2, R) \rightarrow \text{Ext}_R^1(Y, R) \rightarrow 0.$$

Now $\text{Hom}_R(Y, R)$ has dimension vector $[0, 0, 0, 1]$, $\text{Hom}_R(R e_3, R)$ has dimension vector $[0, 0, 1, 1]$ and $\text{Hom}_R(Y_2, R)$ has dimension vector $[1, 1, 2, 0]$. Thus the R -module $\text{Ext}_R^1(Y, R)$ has dimension vector $[1, 1, 1, 0]$. Now the module $\text{Hom}_R(Y_2, R)$ with dimension vector $[1, 1, 2, 0]$ is isomorphic to the R -module $S_3 \oplus e_1 R$ and $\text{Ext}_R^1(Y, R)$ being a quotient of $\text{Hom}_R(Y_2, R) \cong S_3 \oplus e_1 R$ with dimension vector $[1, 1, 1, 0]$ implies that $\text{Ext}_R^1(Y, R) \cong e_1 R$ is projective. Thus $\text{Ext}_R^2(\text{Tr}(S_3), R) \cong \text{Ext}_R^1(Y, R)$ is projective and S_3 is 2-reflexive. What is left to show is that S_3 is not reflexive. But S_3 being reflexive is equivalent to $\text{Ext}_R^i(\text{Tr}(S_3), R) = 0$ for $i = 1, 2$, see for example [ARS, corollary 3.3 in chapter IV.]. But we just saw that $\text{Ext}_R^2(\text{Tr}(S_3), R)$ is non-zero and thus S_3 can not be reflexive. \square

1. ACKNOWLEDGEMENT

The algebra and the modules N and S_3 in Proposition 0.2 were found and verified using the GAP-package [QPA].

REFERENCES

- [AB] Auslander, M.; Bridger, M.: *Stable module theory*, Memoirs of the American Mathematical Society, No. 94 American Mathematical Society, Providence, R.I. 1969, 146 pp.
- [ARS] Auslander, M.; Reiten, I.; Smalø, S. O.: *Representation theory of Artin algebras*. Cambridge Studies in Advanced Mathematics, Volume 36, Cambridge University Press, 1997.
- [ASS] Assem, I.; Simson, D.; Skowronski, A.: *Elements of the representation theory of associative algebras. Vol. 1: Techniques of representation theory*. London Mathematical Society Student Texts 65. Cambridge: Cambridge University Press, 458 p. (2006).
- [CM] Cruz, T.; Marczinzik, R.: *A new formula for the classical dominant dimension using bimodules*. <https://arxiv.org/abs/2508.18398>
- [E] Esnault, H.: *Reflexive modules on quotient surface singularities*. J. Reine Angew. Math., 362:63-71, 1985.

- [EK] Esnault, H; Knörrer, H.: *Reflexive modules over rational double points*. Math. Ann., 272:545-548, 1985.
- [QPA] The QPA-team, QPA - Quivers, path algebras and representations - a GAP package, Version 1.33; 2022 <https://folk.ntnu.no/oyvinso/QPA/>.

(Marczinzik) MATHEMATICAL INSTITUTE OF THE UNIVERSITY OF BONN, ENDENICHER ALLEE 60, 53115 BONN, GERMANY
Email address: `marczire@math.uni-bonn.de`