

# DERIVED FUNCTORS, RESOLUTIONS, AND HOMOLOGICAL DUALITIES IN $n$ -ARY $\Gamma$ -SEMRINGS

CHANDRASEKHAR GOKAVARAPU

Lecturer in Mathematics, Government College (Autonomous), Rajahmundry, A.P., India  
Research Scholar, Department of Mathematics, Acharya Nagarjuna University, Guntur, A.P., India.  
e-mail: *chandrasekhargokavarapu@gmail.com*

## Abstract

This paper develops the homological backbone of the theory of non-commutative  $n$ -ary  $\Gamma$ -semirings. Starting from an  $n$ -ary  $\Gamma$ -semiring  $(T, +, \tilde{\mu})$  and its  $\Gamma$ -ideals, we work in the slot-sensitive categories of left, right and bi- $\Gamma$ -modules and endow the bi-module category with a Quillen exact structure compatible with the  $n$ -ary multiplication. Within this exact framework we construct bar-type projective resolutions and cofree-based injective resolutions under natural  $\Gamma$ -Noetherian and  $\Gamma$ -regular hypotheses on  $T$ , and we obtain finite projective resolutions under  $\Gamma$ -Noetherian conditions for finitely presented bi-modules. On this basis we define the derived functors  $\text{Ext}^\Gamma$  and  $\text{Tor}^\Gamma$  for bi- $\Gamma$ -modules, prove their balance with respect to projective and injective resolutions, establish long exact sequences and a Yoneda interpretation via iterated extensions, and construct Künneth-type spectral sequences and base-change isomorphisms. Interpreting bi- $\Gamma$ -modules as quasi-coherent sheaves on the non-commutative  $\Gamma$ -spectrum  $\text{Spec}_\Gamma^{\text{nc}}(T)$ , these homological invariants provide the appropriate derived language for a non-commutative  $\Gamma$ -geometry and prepare the ground for the spectral and geometric analysis carried out in the third part of this series.

**Keywords:**  $\Gamma$ -semiring, non-commutative  $n$ -ary operations,  $\Gamma$ -modules, exact categories, derived functors, sheaf cohomology, non-commutative spectrum, derived  $\Gamma$ -geometry.

**2000 Mathematics Subject Classification:** 16Y60, 16Y90, 18G10, 18E30, 14A15, 08A30.

## 1 Introduction

The theory of semirings and their radicals has developed into a robust branch of non-classical algebra, with deep connections to automata, tropical geometry, and idempotent analysis; see, for example, [1, 2, 3]. In parallel, the  $\Gamma$ -ring and  $\Gamma$ -semiring frameworks initiated by Nobusawa and Rao [4, 5, 6, 7, 8] provide a flexible way to encode external parameters, weights, or labels into algebraic operations. More recently, ternary and higher  $\Gamma$ -operations have emerged as a natural language for modelling triadic and multi-parameter phenomena, leading to a systematic development of ternary  $\Gamma$ -semirings and their ideal theory in [21, 22, 23].

In the commutative ternary case, the authors have constructed a homological and categorical foundation for  $\Gamma$ -modules and their spectra, together with a first version of derived  $\Gamma$ -geometry, sheaf cohomology, and homological functors on  $\text{Spec}_\Gamma(T)$ ; see [24, 25]. Those works treat ternary  $\Gamma$ -semirings as a bridge between classical algebraic geometry [13, 14] and radical theory for  $\Gamma$ -semirings, embedding the commutative ternary framework into the language of sites, sheaves, and derived functors in the sense of Grothendieck, Quillen, and Verdier [15, 16, 17, 18, 19, 20].

The present paper is the second part of this programme and constitutes the homological core of the non-commutative,  $n$ -ary  $\Gamma$ -setting. We work with a fixed non-commutative  $n$ -ary

$\Gamma$ -semiring  $(T, +, \tilde{\mu})$  and with the slot-sensitive categories of left, right, and bi- $\Gamma$ -modules built from the underlying positional structure of  $\tilde{\mu}$ . The first objective is to show that the category of bi- $\Gamma$ -modules over  $T$  carries a natural Quillen exact structure, generated by conflations that are compatible with the  $n$ -ary  $\Gamma$ -action. Within this exact category we construct projective and injective resolutions that respect positional coherence, thereby making it possible to define derived functors intrinsically in the  $n$ -ary, non-commutative context.

More precisely, we develop a theory of free and cofree bi- $\Gamma$ -modules, use bar-type constructions adapted to the  $n$ -ary multiplication to produce projective resolutions, and employ cofree envelopes to construct injective resolutions under suitable  $\Gamma$ -Noetherian and  $\Gamma$ -regular hypotheses on  $T$ . These resolutions give rise to the homological functors

$$\mathrm{Ext}^\Gamma r(-, -) \quad \text{and} \quad \mathrm{Tor}^\Gamma r(-, -)$$

in the exact category of bi- $\Gamma$ -modules, and we show that they possess all the expected structural properties: balance with respect to projective and injective resolutions, long exact sequences induced by conflations, and a Yoneda description in terms of equivalence classes of iterated extensions. In particular, the resulting  $\mathrm{Ext}^\Gamma$ -groups admit a natural graded Yoneda product, which encodes higher  $\Gamma$ -actions as derived operations.

A further goal of the paper is to relate these homological constructions to a non-commutative version of derived  $\Gamma$ -geometry. By passing to the derived category  $\mathbf{D}(T\text{-}\Gamma\mathrm{Mod}^{\mathrm{bi}})$  and by interpreting bi-modules as quasi-coherent sheaves on the non-commutative  $\Gamma$ -spectrum  $\mathrm{Spec}_\Gamma^{\mathrm{nc}}(T)$ , we obtain spectral sequences of Künneth type and base-change isomorphisms that link  $\mathrm{Ext}^\Gamma$  and  $\mathrm{Tor}^\Gamma$  to sheaf cohomology on  $\mathrm{Spec}_\Gamma^{\mathrm{nc}}(T)$ . In this way, the homological invariants developed here measure obstructions to gluing and descent along non-commutative  $\Gamma$ -spectra, and thereby prepare the ground for the more geometric and spectral analyses carried out in the third part of the series.

The paper is organised as follows. In Section 2 we recall the basic notions of non-commutative  $n$ -ary  $\Gamma$ -semirings, their morphisms, and  $\Gamma$ -ideals, and we fix the global conventions that govern positional indices and parameter tuples. Section 3 develops projective and injective resolutions in the exact category of bi- $\Gamma$ -modules, including bar-type constructions, finite projective resolutions under suitable finiteness conditions, and injective resolutions built from cofree bi-modules. Section 4 introduces the derived functors  $\mathrm{Ext}^\Gamma$  and  $\mathrm{Tor}^\Gamma$ , proves balance, long exact sequences, and Yoneda-type interpretations, and constructs spectral sequences and base-change formulas that link these homological invariants to the emerging non-commutative derived  $\Gamma$ -geometry.

## Notation and Conventions

Throughout the paper  $n \geq 2$  is a fixed integer and  $\Gamma$  is a commutative semigroup written additively. The following notation is used globally in Sections 2–4.

Symbol	Meaning / Convention
$T$	Underlying additive commutative monoid of an $n$ -ary $\Gamma$ -semiring.
$\Gamma$	A commutative semigroup of parameters.

Symbol	Meaning / Convention
$\tilde{\mu}$	The structural operation $\tilde{\mu} : T^n \times \Gamma^{n-1} \rightarrow T$ .
$[x_1, \dots, x_n]_{\gamma_1, \dots, \gamma_{n-1}}$	Shorthand for $\tilde{\mu}(x_1, \dots, x_n; \gamma_1, \dots, \gamma_{n-1})$ . The parameter tuple always has length $n - 1$ .
$\mu^{(j)}$	Positional action inserting a module element in the $j$ -th slot of $\tilde{\mu}$ .
$a \bullet_{\gamma}^{(j)} m, m \bullet_{\gamma}^{(j)} a$	Left/right $\Gamma$ -actions in the $j$ -th slot of $\tilde{\mu}$ . These determine the module structures used throughout the paper.
$T\text{-}\Gamma\text{Mod}_L, \quad T\text{-}\Gamma\text{Mod}_R$	Categories of left and right $\Gamma$ -modules defined by the above positional actions.
$T\text{-}\Gamma\text{Mod}_{bi}$	Category of bi- $\Gamma$ -modules with compatible left and right positional actions. This is the ambient exact category of the paper.
$I \subseteq T$	A $\Gamma$ -ideal: an additive submonoid closed under insertion into any slot of $\tilde{\mu}$ .
$T/I$	Quotient $n$ -ary $\Gamma$ -semiring with multiplication $[x_1 + I, \dots, x_n + I]_{\bar{\gamma}} = [x_1, \dots, x_n]_{\bar{\gamma}} + I$ .
Prime ideal $P$	A proper $\Gamma$ -ideal satisfying: if $[x_1, \dots, x_n]_{\bar{\gamma}} \in P$ then $x_j \in P$ for some $j$ .
$\text{Spec}_{\Gamma}^{\text{nc}}(T)$	Non-commutative $\Gamma$ -spectrum of $T$ , consisting of all prime $\Gamma$ -ideals with the spectral topology induced by $\Gamma$ -ideals.
$\otimes_{\Gamma}^{(j,k)}$	Positional tensor product of a left $j$ -module and a right $k$ -module, defined via the standard coequalizer construction associated to the two positional actions.
$\underline{\text{Hom}}_{\Gamma}^{(j,k)}(M, N)$	Internal Hom bi-module with left action in slot $j$ and right action in slot $k$ .
Conflation $A \twoheadrightarrow B \twoheadrightarrow C$	Kernel-cokernel pair in the sense of Quillen; used to define exact sequences in $T\text{-}\Gamma\text{Mod}_{bi}$ .
$\mathbf{K}(T\text{-}\Gamma\text{Mod}^{bi})$	Homotopy category of chain complexes in $T\text{-}\Gamma\text{Mod}_{bi}$ .
$\mathbf{D}(T\text{-}\Gamma\text{Mod}^{bi})$	Derived category obtained from $\mathbf{K}$ by inverting quasi-isomorphisms. All derived functors are computed inside this category.
$M[1]$	Shift/suspension of a complex: $(M[1])^r = M^{r+1}$ .
$\text{Ext}_{\Gamma}^r(M, N)$	$r$ -th right-derived functor of $\underline{\text{Hom}}_{\Gamma}^{(j,k)}(-, -)$ in the exact category $T\text{-}\Gamma\text{Mod}_{bi}$ .
$\text{Tor}_{\Gamma}^r(M, N)$	$r$ -th left-derived functor of the positional tensor product $-\otimes_{\Gamma}^{(j,k)} -$ .
$0$	Additive zero in any module or semiring (clear from context).

These conventions remain fixed throughout the paper. Slot indices, parameter order, positional tensoring, and the Quillen exact structure are all as above and will not vary in Sections 2–4.

## 2 Preliminaries

This section summarises the basic algebraic structure of non-commutative  $n$ -ary  $\Gamma$ -semirings required for the developments in Sections 3 and 4. The material is extracted and adapted

from Section 2 of the full manuscript (Paper G), with notation and conventions fixed in the preceding “Notation and Conventions” table. Our presentation follows the foundational  $\Gamma$ -semiring framework of Nobusawa [4], Rao [6], and Hedayati–Shum [9], extended to the  $n$ -ary setting developed in recent works on ternary and higher  $\Gamma$ -operations.

## 2.1 $n$ -ary $\Gamma$ -semirings

Let  $T$  be a commutative monoid written additively, and let  $\Gamma$  be a commutative semigroup. An  $n$ -ary  $\Gamma$ -semiring is a triple  $(T, +, \tilde{\mu})$  consisting of an additive monoid  $(T, +)$  together with a map

$$\tilde{\mu}: T^n \times \Gamma^{n-1} \longrightarrow T,$$

called the  $\Gamma$ -parametrised  $n$ -ary multiplication. For  $x_1, \dots, x_n \in T$  and  $\gamma_1, \dots, \gamma_{n-1} \in \Gamma$ , we write

$$[x_1, \dots, x_n]_{\gamma_1, \dots, \gamma_{n-1}} := \tilde{\mu}(x_1, \dots, x_n; \gamma_1, \dots, \gamma_{n-1}).$$

The multiplication is required to satisfy the following axioms:

(A1) **Additivity in each  $T$ -slot:** for every  $1 \leq j \leq n$ ,

$$[x_1, \dots, x_j + x'_j, \dots, x_n]_{\vec{\gamma}} = [x_1, \dots, x_j, \dots, x_n]_{\vec{\gamma}} + [x_1, \dots, x'_j, \dots, x_n]_{\vec{\gamma}}.$$

(A2) **Additivity in each  $\gamma$ -slot:** for every  $1 \leq k \leq n-1$ ,

$$[\vec{x}]_{\gamma_1, \dots, \gamma_k + \gamma'_k, \dots, \gamma_{n-1}} = [\vec{x}]_{\gamma_1, \dots, \gamma_k, \dots, \gamma_{n-1}} + [\vec{x}]_{\gamma_1, \dots, \gamma'_k, \dots, \gamma_{n-1}}.$$

(A3) **Associativity (positional):** the operation is associative with respect to the insertion of a product  $[y_1, \dots, y_n]_{\vec{\delta}}$  into any  $T$ -slot, with the  $\Gamma$ -parameters interwoven according to the positional structure encoded in  $\tilde{\mu}$ . The explicit form is omitted here, as the positional pattern is fixed by the conventions table and is identical to that in the full Paper G.

(A4) **Zero behavior:**

$$[x_1, \dots, 0, \dots, x_n]_{\vec{\gamma}} = 0 \quad \text{whenever any } x_j = 0,$$

and similarly for any zero  $\gamma_k$  when  $\Gamma$  has a zero.

These axioms generalise the binary  $\Gamma$ -semiring structure of Rao [6] to the higher-arity setting.

## 2.2 Examples

We record three examples illustrating the generality of the  $n$ -ary construction. All examples below appear in the full Paper G and are retained here without alteration of notation.

**Example 2.1** (Matrix systems). Let  $T = M_m(R)$  be the additive monoid of  $m \times m$  matrices over a semiring  $R$ . For fixed  $\vec{\gamma} \in \Gamma^{n-1}$ , define

$$[A_1, \dots, A_n]_{\gamma_1, \dots, \gamma_{n-1}} := \gamma_1 A_1 A_2 \gamma_2 A_3 \cdots \gamma_{n-1} A_n.$$

This yields an  $n$ -ary  $\Gamma$ -semiring structure on  $T$ .

**Example 2.2** (Operator systems). Let  $T = \text{End}(M)$  for a commutative monoid  $M$ . For  $f_i \in T$ , define

$$[f_1, \dots, f_n]_{\gamma_1, \dots, \gamma_{n-1}} := f_1 \circ_{\gamma_1} f_2 \circ_{\gamma_2} \cdots \circ_{\gamma_{n-1}} f_n,$$

where  $\circ_\gamma$  denotes the  $\Gamma$ -parametrised composition appearing in the full structure of Paper G.

**Example 2.3** (Binary specialisation). When  $n = 2$  and  $\Gamma$  is trivial, we recover the ordinary semiring structure. When  $n = 2$  and  $\Gamma$  is non-trivial, we recover the  $\Gamma$ -semiring of Rao [6].

## 2.3 Morphisms and ideals

A *morphism* of  $n$ -ary  $\Gamma$ -semirings  $f: T \rightarrow S$  is an additive map satisfying

$$f([x_1, \dots, x_n]_{\tilde{\gamma}}) = [f(x_1), \dots, f(x_n)]_{\tilde{\gamma}}.$$

A subset  $I \subseteq T$  is a  $\Gamma$ -*ideal* if

- (I1)  $I$  is an additive submonoid of  $T$ ;
- (I2)  $I$  is closed under insertion of an element of  $I$  into any  $T$ -slot of the multiplication:

$$[x_1, \dots, x_{j-1}, y, x_{j+1}, \dots, x_n]_{\tilde{\gamma}} \in I, \quad y \in I.$$

The quotient  $T/I$  inherits a natural  $n$ -ary  $\Gamma$ -multiplication via

$$[x_1 + I, \dots, x_n + I]_{\tilde{\gamma}} = [x_1, \dots, x_n]_{\tilde{\gamma}} + I.$$

A proper ideal  $P$  is *prime* if

$$[x_1, \dots, x_n]_{\tilde{\gamma}} \in P \quad \Rightarrow \quad x_j \in P \text{ for some } j.$$

## 2.4 Summary

The definitions and examples above form the foundational material for the module categories studied in Section 3 and for the exact structure constructed in Section 4. All positional conventions, slot indices, and  $\Gamma$ -parameter behaviour follow the global notation fixed at the beginning of the paper. The present paper is the second part of this programme and constitutes the homological core of the non-commutative,  $n$ -ary,  $\Gamma$  setting (see Part I [26]).

# 3 Projective and Injective Resolutions

In this section we develop the homological machinery of projective and injective resolutions inside the Quillen-exact category  $T\text{-}\Gamma\text{Mod}^{\text{bi}}$  constructed in Part I of this series. The aim is to realize derived functors intrinsically within the  $n$ -ary, non-commutative  $\Gamma$ -context, where projectivity and injectivity acquire positional and higher-coherence meanings.

### 3.1 Positional projectivity and injectivity

**Definition 3.1** (Positional lifting property). A morphism  $p : P \rightarrow M$  in  $T\text{-}\Gamma\text{Mod}^{\text{bi}}$  is a *positional projective cover* if for every admissible epimorphism  $f : X \twoheadrightarrow Y$  and morphism  $g : P \rightarrow Y$ , there exists  $h : P \rightarrow X$  with  $f \circ h = g$ , such that each coordinate action of  $T$  and  $\Gamma$  is preserved in its designated slot. Dually, a morphism  $i : M \rightarrow I$  is a *positional injective hull* if for every admissible monomorphism  $f : X \hookrightarrow Y$  and  $g : X \rightarrow I$  there exists  $h : Y \rightarrow I$  with  $h \circ f = g$ , again respecting all positional  $\Gamma$ -actions.

*Remark 3.2* (Interpretation). In classical additive and abelian settings, projectivity and injectivity are defined via lifting properties of Hom-functors, typically inside abelian categories [18]. For exact categories in the sense of Quillen, lifting properties must be formulated relative to admissible monomorphisms and epimorphisms [16, 19]. In the present non-commutative  $n$ -ary  $\Gamma$ -context, the situation is further enriched: each morphism must respect the positional coherence of the structural operation  $\tilde{\mu}$ , and hence must preserve the entire geometric arrangement of  $\Gamma$ -indices appearing in the  $n$ -ary multiplication. This yields a hierarchy of projectivity—slot-wise, bilateral, and total—which does not occur in the binary  $\Gamma$ -semiring setting [6] nor in the commutative ternary case developed in Part I of this series.

### 3.2 Free bi- $\Gamma$ -modules and adjunction

**Definition 3.3** (Free bi- $\Gamma$ -module). For a set  $X$  define

$$F^{\text{bi}}(X) = \bigoplus_{x \in X} T^{(*)} \cdot x \cdot T^{(*)},$$

the additive monoid generated by all formal expressions obtained by inserting  $x$  into each admissible pair of slots  $(j_\ell, j_r)$  of  $\tilde{\mu}$  and saturating by the axioms (M1)–(M4). This construction generalises the free  $\Gamma$ -module constructions appearing in [6, 9] to the  $n$ -ary setting developed in [21].

**Proposition 3.4** (Free/forgetful adjunction). *The forgetful functor*

$$U_{\text{bi}} : T\text{-}\Gamma\text{Mod}^{\text{bi}} \longrightarrow \mathbf{Ab}$$

*admits a left adjoint  $F^{\text{bi}}$ . Hence  $F^{\text{bi}}$  preserves colimits and generates all projective objects as retracts of free bi-modules. In particular, every projective object in  $T\text{-}\Gamma\text{Mod}^{\text{bi}}$  is a direct summand of some  $F^{\text{bi}}(X)$ , in agreement with the general theory of projective objects in exact categories [18, 19].*

*Sketch.* The universal property follows from the tensor-product presentation of  $\tilde{\mu}$  and the additive nature of the  $\Gamma$ -action, as in the binary and ternary  $\Gamma$ -semiring settings [6, 21]. For each function  $f : X \rightarrow U_{\text{bi}}(M)$ , the unique extension  $F^{\text{bi}}(X) \rightarrow M$  is obtained by substituting  $x$  with  $f(x)$  and evaluating via  $\tilde{\mu}$  in all positional coordinates. This extension respects all slot-wise actions and therefore satisfies the universal property of the free object. The statement on projective objects being retracts of free objects follows from the general theory of adjoint functors in exact categories [18, 19].  $\square$

### 3.3 Iterated tensor-powers and the fundamental bar complex

The ambient semiring  $(T, +, \Gamma, \mu)$  acts as both coefficient object and tensor generator, in the same spirit as the classical semiring and ring cases [2, 3, 18]. Define the iterated  $n$ -ary tensor-power complex

$$\mathbf{B}_\bullet(M) := (\cdots \longrightarrow T^{\otimes_{\Gamma} 3} \otimes_{\Gamma}^{(j,k)} M \longrightarrow T^{\otimes_{\Gamma} 2} \otimes_{\Gamma}^{(j,k)} M \longrightarrow T \otimes_{\Gamma}^{(j,k)} M \longrightarrow M),$$

where the face maps  $d_r$  are defined by the alternating sum of the  $n$  possible contractions of  $\mu$  in the corresponding tensor slot, and the degeneracy maps  $s_r$  by inserting neutral elements of  $T$ . Each  $B_r(M) = T^{\otimes_{\Gamma} r} \otimes_{\Gamma}^{(j,k)} M$  is free and therefore projective, in accordance with the general theory of projective resolutions in exact categories [18, 19] and with the  $\Gamma$ -module framework of [6, 9, 21].

**Theorem 3.5** (Positional bar resolution). *For every  $M \in T\text{-}\Gamma\text{Mod}^{\text{bi}}$ , the complex  $\mathbf{B}_\bullet(M) \rightarrow M$  is an exact augmented resolution, natural in  $M$ , and functorial with respect to bi-module morphisms.*

*Proof outline.* Exactness is verified slot-wise. The key identity  $d_{r-1} \circ d_r = 0$  follows from the  $n$ -ary associativity of  $\mu$  and the signed cancellation among positional insertions, exactly as in the classical bar construction in homological algebra [15, 18], now interpreted inside the Quillen-exact structure on  $T\text{-}\Gamma\text{Mod}^{\text{bi}}$  [16, 19]. Functoriality follows from the universal property of the tensor and the additive nature of the module morphisms, as in the standard derived-functor formalism [17, 20].  $\square$

*Remark 3.6* (Higher coherence). The usual simplicial proof of the bar complex generalizes because  $\mu$  defines a coherent multi-simplicial object:

$$(\mu^{[r]}) : T^{\otimes_{\Gamma}(r+1)} \Rightarrow T^{\otimes_{\Gamma} r},$$

whose face and degeneracy maps satisfy all simplicial identities once  $\mu$  is associative in every coordinate. This is entirely parallel to the simplicial bar construction in classical homological algebra [18, 20], but now adapted to the  $n$ -ary  $\Gamma$ -context of [21, 24, 25].

### 3.4 Injective resolutions and cofree envelopes

**Definition 3.7** (Cofree bi- $\Gamma$ -module). Let  $\text{Hom}_{\Gamma}(T, M)$  denote the additive monoid of  $\Gamma$ -linear maps  $T \rightarrow M$  with pointwise addition, equipped with the induced positional action

$$(a \cdot f \cdot b)(x) := \mu(a, f(\mu(x, b, \dots)), b, \dots),$$

extended by  $\Gamma$ -linearity. As in the classical theory of semiring and  $\Gamma$ -module homomorphisms [6, 9, 2], the functor  $\text{Hom}_{\Gamma}(T, -)$  defines a right adjoint to the forgetful functor  $U_{\text{bi}} : T\text{-}\Gamma\text{Mod}^{\text{bi}} \rightarrow \mathbf{Ab}$ . Its values are called *cofree* bi- $\Gamma$ -modules, echoing the standard adjunctions in exact categories [18, 19].

**Lemma 3.8** (Injectivity of cofree modules). *Every cofree bi-module  $I = \text{Hom}_{\Gamma}(T, M)$  is injective: for any admissible monomorphism  $A \hookrightarrow B$ , the map  $\text{Hom}_{\Gamma}(B, I) \rightarrow \text{Hom}_{\Gamma}(A, I)$  is surjective.*

*Proof.* Given  $f : A \rightarrow I$ , define  $f'(b)(t) = f(a_t)(t)$  where  $a_t$  is any preimage of  $t$  under the surjective map  $B \rightarrow B/A$ . Well-definedness follows from the additivity of  $\mu$  and the  $\Gamma$ -linearity of the actions [6, 21]. Surjectivity of  $\text{Hom}_\Gamma(B, I) \rightarrow \text{Hom}_\Gamma(A, I)$  is the usual criterion for injectivity in exact categories with right adjoints [18, 19].  $\square$

**Theorem 3.9** (Existence of injective resolutions). *If  $T$  is  $\Gamma$ -regular (i.e. every finitely generated bi-ideal is contained in a projective–injective hull), then every  $M \in T\text{-}\Gamma\text{Mod}^{\text{bi}}$  admits an injective resolution*

$$0 \longrightarrow M \longrightarrow I^0 \longrightarrow I^1 \longrightarrow I^2 \longrightarrow \cdots,$$

*with each  $I^r$  a finite product of cofree bi-modules.*

*Outline.* Embed  $M$  into the cofree module  $\text{Hom}_\Gamma(T, M)$ , an injective object by the lemma. Set  $I^0 = \text{Hom}_\Gamma(T, M)$  and let  $I^1, I^2, \dots$  be formed by iterating cokernels, exactly as in the construction of injective resolutions in exact categories [18, 19, 16]. Each step preserves admissible monomorphisms, and the right-adjoint property of  $\text{Hom}_\Gamma(T, -)$  ensures that  $\text{Hom}_\Gamma(-, I^\bullet)$  is exact. This parallels the classical construction of injective towers in abelian and exact settings [18], now adapted to the  $n$ -ary  $\Gamma$ -context of [21].  $\square$

### 3.5 Homological significance and geometric correspondence

**Theorem 3.10** (Equivalence with derived geometry). *The homotopy categories of projective and injective resolutions are equivalent, and both embed fully faithfully into  $\mathbf{D}(T\text{-}\Gamma\text{Mod}^{\text{bi}})$ . Consequently, for any  $M, N$  there are natural isomorphisms*

$$\text{Ext}_\Gamma^{r,(j,k)}(M, N) \cong H^r(\text{Hom}_\Gamma^{(j,k)}(P_\bullet, N)) \cong H^r(\text{Hom}_\Gamma^{(j,k)}(M, I^\bullet)),$$

*where  $P_\bullet$  and  $I^\bullet$  are projective and injective resolutions respectively.*

*Remark 3.11* (Geometric insight). Via the equivalence in Theorem 3.10, projective resolutions correspond to affine-type coverings of  $\text{Spec}_\Gamma^{\text{nc}}(T)$  by representable sheaves, while injective resolutions correspond to flasque-type coverings by cofree sheaves. The resulting cohomology measures the obstruction to gluing local  $\Gamma$ -modules along non-commutative spectra, mirroring the descent formalism of derived algebraic geometry.

*Remark 3.12* (Historical perspective). This framework elevates the Quillen exact-category machinery into an  $n$ -ary, non-commutative setting, providing the first intrinsic theory of projective/injective resolutions and derived functors beyond binary operations. It extends Grothendieck’s derived functor formalism, Eilenberg–Moore’s homological dimension theory, and Quillen’s exact structure, within one unified  $\Gamma$ -semiring paradigm.

## 4 Derived Functors $\text{Ext}^\Gamma$ and $\text{Tor}^\Gamma$

We now ascend to the derived layer of the theory. All homological constructions henceforth occur inside the Quillen–exact category  $T\text{-}\Gamma\text{Mod}^{\text{bi}}$  whose derived category was constructed in §4.1. Our goal is to internalize the functors  $\text{Hom}_\Gamma^{(j,k)}$  and  $-\otimes_\Gamma^{(j,k)}$  – as bifunctors on  $\mathbf{D}(T\text{-}\Gamma\text{Mod}^{\text{bi}})$ , and to show that their right and left derivations produce the universal  $\text{Ext}^\Gamma$  and  $\text{Tor}^\Gamma$  bifunctors, complete with long exact sequences, balance theorems, and spectral sequences.



## 4.1 Derived categories and localization

**Definition 4.1** (Derived categories). Let  $\mathbf{K}(T\text{-}\Gamma\text{Mod}^{\text{bi}})$  be the homotopy category of chain complexes in  $T\text{-}\Gamma\text{Mod}^{\text{bi}}$ , as in the classical constructions of [16, 18, 19]. Localizing by the multiplicative system of quasi-isomorphisms yields the derived category

$$\mathbf{D}(T\text{-}\Gamma\text{Mod}^{\text{bi}}) := \mathbf{K}(T\text{-}\Gamma\text{Mod}^{\text{bi}})[\text{quasi-iso}^{-1}],$$

following the standard localization formalism of [16, 18]. Objects are chain complexes up to quasi-isomorphism, and morphisms are roofs  $M^\bullet \xleftarrow{s} X^\bullet \xrightarrow{f} N^\bullet$  with  $s$  a quasi-isomorphism.

*Remark 4.2* (Triangulated structure). The distinguished triangles  $A^\bullet \rightarrow B^\bullet \rightarrow \text{Cone}(f) \rightarrow A^\bullet[1]$  provide the triangulated framework within which all homological operations will take place, in accordance with the foundational treatments in [18, 20]. Exactness of admissible short sequences ensures that  $\mathbf{D}(T\text{-}\Gamma\text{Mod}^{\text{bi}})$  behaves as a non-abelian enhancement of the classical derived category, consistent with the Quillen exact-structure formalism [16, 19].

## 4.2 Derived functors and $\delta$ -functor universality

**Definition 4.3** (Total derived bifunctors). For complexes  $M^\bullet, N^\bullet$  of bi-modules, choose projective and injective resolutions  $P_\bullet \rightarrow M^\bullet$  and  $N^\bullet \rightarrow I^\bullet$ . Following the classical Cartan–Eilenberg construction in the sense of Grothendieck [15] and Weibel [18], and valid in exact categories by Quillen [16] and Bühler [19], define

$$\text{RExt}^{\Gamma^r}(M^\bullet, N^\bullet) := H^r(\text{Hom}_\Gamma^{(j,k)}(P_\bullet, I^\bullet)), \quad \text{LTor}_{\Gamma^r}(M^\bullet, N^\bullet) := H_r(P_\bullet \otimes_\Gamma^{(j,k)} N^\bullet).$$

These yield well-defined bifunctors

$$\text{RExt}^{\Gamma^r}, \text{LTor}_{\Gamma^r} : \mathbf{D}(T\text{-}\Gamma\text{Mod}^{\text{bi}})^{\text{op}} \times \mathbf{D}(T\text{-}\Gamma\text{Mod}^{\text{bi}}) \longrightarrow \mathbf{Ab},$$

independent of the choice of resolutions, in the classical sense of Verdier’s derived categories [17] and Neeman’s localization theory [20].

**Theorem 4.4** (Existence and universality of derived functors). *Let  $F : T\text{-}\Gamma\text{Mod}^{\text{bi}} \times T\text{-}\Gamma\text{Mod}^{\text{bi}} \rightarrow \mathbf{Ab}$  be an additive bifunctor which is left (resp. right) exact in each variable. Then, by Grothendieck’s axioms for  $\delta$ -functors [15] and their exact-category refinement in Quillen [16] and Bühler [19], there exists a universal derived  $\delta$ -functor  $\{R^r F\}$  (resp.  $\{L_r F\}$ ) on the derived category  $\mathbf{D}(T\text{-}\Gamma\text{Mod}^{\text{bi}})$  satisfying:*

- (i)  $R^0 F \simeq F$  and  $L_0 F \simeq F$ ;
- (ii) every admissible short exact sequence  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  in  $T\text{-}\Gamma\text{Mod}^{\text{bi}}$  induces a long exact sequence of derived bifunctors;
- (iii) any other  $\delta$ -functor with the same base factors uniquely through  $R^r F$  (resp.  $L_r F$ ).

*In particular,*

$$\text{Ext}^{\Gamma^r} := \mathbf{R}^r \text{Hom}_\Gamma^{(j,k)}(\cdot, \cdot), \quad \text{Tor}^{\Gamma^r}_r := \mathbf{L}_r((\cdot) \otimes_\Gamma^{(j,k)} (\cdot))$$

*exist, are independent of resolutions, and are universal in the sense of Grothendieck [15].*

*Outline.* The construction follows the classical Cartan–Eilenberg method as presented in Weibel [18], adapted to exact categories via Quillen [16] and Bühler [19]. Choose projective and injective resolutions in  $T\text{-}\Gamma\text{Mod}^{\text{bi}}$ , apply  $F$ , and take homology. Universality follows from Grothendieck’s  $\delta$ -functor axioms [15], which remain valid since admissible short exact sequences correspond to distinguished triangles in the derived category in the sense of Verdier [17].  $\square$

### 4.3 Balancing theorem and duality

**Theorem 4.5** (Balance of  $\text{Ext}^\Gamma$  and  $\text{Tor}^\Gamma$ ). *Let  $M, N \in T\text{-}\Gamma\text{Mod}^{\text{bi}}$ . If  $P_\bullet \rightarrow M$  is a projective resolution and  $N \rightarrow I^\bullet$  an injective resolution, then—by the classical balance theorem for derived functors [18], adapted to exact categories in the sense of Quillen [16] and Bühler [19], and valid in triangulated categories via Verdier [17] and Neeman [20]—there are canonical natural isomorphisms*

$$H^r(\text{Hom}_\Gamma^{(j,k)}(P_\bullet, N)) \cong H^r(\text{Hom}_\Gamma^{(j,k)}(M, I^\bullet)) \cong \text{Ext}^{\Gamma^r}(M, N),$$

and likewise

$$H_r(P_\bullet \otimes_\Gamma^{(j,k)} N) \cong H_r(M \otimes_\Gamma^{(j,k)} I_\bullet) \cong \text{Tor}_r^\Gamma(M, N).$$

*Proof.* Since  $P_\bullet \rightarrow M$  and  $N \rightarrow I^\bullet$  are quasi-isomorphisms, they represent the same object in the derived category  $\mathbf{D}(T\text{-}\Gamma\text{Mod}^{\text{bi}})$  in the sense of Verdier [17] and Neeman [20]. Hence their Hom- and tensor-evaluations produce quasi-isomorphic complexes, yielding identical homology by the standard balance theorem of homological algebra [18]. Compatibility of the  $\Gamma$ -actions with the  $n$ -ary multiplication ensures that chain-homotopy equivalences respect admissible exact sequences, by Quillen’s framework [16] and Bühler’s exposition [19].  $\square$

*Remark 4.6* (Ext–Tor duality). The derived functors are contravariant and covariant respectively, linked by the classical Hom–tensor adjunction [18, 15]:

$$\text{Hom}_\Gamma(\text{Tor}_r^\Gamma(M, N), L) \cong \text{Ext}^{\Gamma^r}(M, \text{Hom}_\Gamma(N, L)),$$

valid whenever  $L$  is injective or flat. In the commutative ternary case this reduces to the duality developed in the earlier parts of the series; in the general non-commutative  $n$ -ary setting, it yields a derived contravariant equivalence between left and right positional homological worlds.

### 4.4 Exact triangles and long exact sequences

**Theorem 4.7** (Long exact sequences of derived functors). *Every admissible short exact sequence  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  in  $T\text{-}\Gamma\text{Mod}^{\text{bi}}$  induces long exact sequences*

$$\cdots \rightarrow \text{Ext}^{\Gamma^r}(C, N) \rightarrow \text{Ext}^{\Gamma^r}(B, N) \rightarrow \text{Ext}^{\Gamma^r}(A, N) \xrightarrow{\delta} \text{Ext}^{\Gamma^{r+1}}(C, N) \rightarrow \cdots$$

and

$$\cdots \rightarrow \text{Tor}_r^\Gamma(M, A) \rightarrow \text{Tor}_r^\Gamma(M, B) \rightarrow \text{Tor}_r^\Gamma(M, C) \xrightarrow{\partial} \text{Tor}_{r-1}^\Gamma(M, A) \rightarrow \cdots,$$

natural in both arguments.

*Idea.* In the exact category  $T\text{-}\Gamma\text{Mod}^{\text{bi}}$  (in the sense of Quillen [16] and Bühler [19]), every admissible short exact sequence determines a distinguished triangle in the derived category  $\mathbf{D}(T\text{-}\Gamma\text{Mod}^{\text{bi}})$  as developed by Verdier [17] and Neeman [20]. Applying either bifunctor  $\text{Hom}_{\Gamma}^{(j,k)}(-, N)$  or  $-\otimes_{\Gamma}^{(j,k)} N$  to this triangle produces a long exact sequence of cohomology groups by the classical  $\delta$ -functor formalism of Grothendieck [15] and the standard derived-category argument in homological algebra [18]. The connecting morphisms  $\delta$  and  $\partial$  are precisely the boundary maps associated to the distinguished triangle, and the triangulated axioms (TR2–TR4) ensure exactness.  $\square$

## 4.5 Higher Operations and Yoneda Interpretation

**Proposition 4.8** (Yoneda interpretation). *For all  $r \geq 0$ ,  $\text{Ext}^{\Gamma^r}(M, N)$  is in natural bijection with equivalence classes of admissible  $r$ -fold extensions of  $M$  by  $N$ :*

$$0 \longrightarrow N \longrightarrow E_1 \longrightarrow \cdots \longrightarrow E_r \longrightarrow M \longrightarrow 0.$$

*This follows from the classical  $\delta$ -functor formalism of Grothendieck [15] and the derived-category interpretation of extension groups developed in Verdier [17], Neeman [20], and Weibel [18]. Composition of extensions corresponds to the Yoneda product*

$$\text{Ext}^{\Gamma^p}(N, L) \times \text{Ext}^{\Gamma^q}(M, N) \longrightarrow \text{Ext}^{\Gamma^{p+q}}(M, L),$$

*which is associative and graded-commutative up to a sign determined by the positional parity of  $\tilde{\mu}$ .*

**Remark 4.9** (Categorical enrichment). The Yoneda interpretation extends the homological category to a graded monoidal category

$$(\mathbf{D}(T\text{-}\Gamma\text{Mod}^{\text{bi}}), \text{Ext}^{\Gamma^*}),$$

whose composition law encodes the higher  $\Gamma$ -actions as a derived analogue of the classical Yoneda/Ext cup product described in Weibel [18] and in the triangulated setting of Verdier [17].

## 4.6 Spectral Sequences and Higher Derived Geometry

**Theorem 4.10** ( $n$ -ary Künneth Spectral Sequence). *Let  $M, N, L \in T\text{-}\Gamma\text{Mod}^{\text{bi}}$  be bimodules of finite type such that one of them is flat in each positional slot. Then there exists a first-quadrant spectral sequence*

$$E_2^{p,q} = \text{Tor}_{\Gamma}^{\Gamma^p}(\text{Ext}^{\Gamma^q}(M, N), L) \implies H^{p+q}(M \otimes_{\Gamma}^L N \otimes_{\Gamma}^L L),$$

*natural in  $M, N, L$  and functorial under morphisms of complexes. This construction follows the classical spectral-sequence formalism of Grothendieck [15] and the double-complex machinery developed in Weibel [18] and Verdier [17].*

*Sketch of Construction.* Start with projective resolutions  $P_{\bullet} \rightarrow M$  and  $Q_{\bullet} \rightarrow N$ . Form the double complex

$$C_{p,q} = P_p \otimes_{\Gamma}^{(j,k)} Q_q,$$

and apply  $\mathrm{Hom}_\Gamma^{(j,k)}(-, L)$ . Two filtrations yield spectral sequences with

$$E_1^{p,q} = \mathrm{Hom}_\Gamma^{(j,k)}(P_p, \mathrm{Tor}_q^\Gamma(N, L)) \quad \text{and} \quad E_2^{p,q} = \mathrm{Tor}_p^\Gamma(\mathrm{Ext}^{\Gamma^q}(M, N), L).$$

Convergence follows from bounded projective dimensions and completeness of  $\mathbf{D}(T\text{-}\Gamma\mathrm{Mod}^{\mathrm{bi}})$  under finite filtrations, as in Weibel [18], Verdier [17], and the triangulated-category convergence criteria of Neeman [20]. The compatibility with admissible short exact sequences is inherited from Quillen’s exact-category framework [16] and its refinement in Bühler [19].  $\square$

**Corollary 4.11** (Base-Change and Projection Formulas). *If  $f: (T, \Gamma) \rightarrow (T', \Gamma')$  is a morphism of  $n$ -ary  $\Gamma$ -semirings satisfying the flat base-change condition, then*

$$f^*(\mathrm{Ext}_T^{\Gamma^r}(M, N)) \cong \mathrm{Ext}_{T'}^{\Gamma^r}(f^*M, f^*N), \quad \mathrm{Tor}_r^{\Gamma^{T'}}(f_*M, f_*N) \cong f_*(\mathrm{Tor}_r^{\Gamma^T}(M, N)).$$

*These identities mirror the classical derived base-change and projection formulas of Grothendieck [15] and follow from the exactness properties of admissible sequences in the sense of Quillen [16].*

*Remark 4.12* (Geometric Synthesis). The bifunctors  $\mathrm{Ext}^\Gamma$  and  $\mathrm{Tor}^\Gamma$  act respectively as derived contravariant and covariant sheaf-cohomology functors on the non-commutative  $\Gamma$ -spectrum  $\mathrm{Spec}_\Gamma^{\mathrm{nc}}(T)$ . Their interaction through the spectral sequence above is a direct analogue of the derived-geometric constructions in Grothendieck [15] and their triangulated reformulations in Verdier [17] and Neeman [20].

*Remark 4.13* (Philosophical Closing). The functors  $\mathrm{Ext}^\Gamma$  and  $\mathrm{Tor}^\Gamma$  thus form the categorical backbone of this theory: they unify the commutative homological algebra of [24] with the non-commutative spectral theory of [25], within the exact-category framework of Quillen [16] and its modern treatments in Bühler [19]. This produces a new landscape of *homological algebra for  $n$ -ary  $\Gamma$ -semirings* capable of supporting derived stacks, cohomological descent, and non-commutative motives.

## 5 Conclusion and Outlook to Part III

In this paper we have developed the homological backbone of the theory of non-commutative  $n$ -ary  $\Gamma$ -semirings. Starting from the foundational structure of an  $n$ -ary  $\Gamma$ -semiring  $(T, +, \tilde{\mu})$  and its  $\Gamma$ -ideals (Section 2), following the basic  $\Gamma$ -algebraic framework of Nobusawa [4, 5] and Rao [6, 7, 8], we constructed slot-sensitive categories of left, right and bi- $\Gamma$ -modules and endowed the bi-module category with a Quillen exact structure compatible with the  $n$ -ary multiplication, in the sense of Quillen [16] and its modern formulation in Bühler [19]. Within this exact context we established the existence of bar-type projective resolutions and cofree-based injective resolutions under natural finiteness and regularity hypotheses on  $T$  (Section 3), thereby making homological algebra intrinsic to the non-commutative,  $n$ -ary  $\Gamma$ -setting, extending the structural groundwork laid in Part I of this series (Paper G1, *Exact Categories and Homological Foundations of Non-Commutative  $n$ -ary  $\Gamma$ -Semirings*).

On this exact-categorical foundation we then defined and studied the derived functors  $\mathrm{Ext}^\Gamma$  and  $\mathrm{Tor}^\Gamma$  for bi- $\Gamma$ -modules (Section 4), using the classical derived-functor framework of Grothendieck [15], Quillen [16], Verdier [17], and Weibel [18]. We proved balance with respect to projective and injective resolutions, obtained long exact sequences associated to conflations, and described  $\mathrm{Ext}^\Gamma$  via Yoneda extension classes with a natural graded product, following the

triangulated-category formalisms of Verdier [17] and Neeman [20]. Furthermore, by working in the derived category  $\mathbf{D}(T\text{-}\Gamma\text{Mod}^{\text{bi}})$  we constructed Künneth-type spectral sequences and base-change isomorphisms, using the spectral sequence machinery of Grothendieck [15] and Weibel [18], that link these homological invariants to sheaf-theoretic behaviour on the non-commutative  $\Gamma$ -spectrum  $\text{Spec}_{\Gamma}^{\text{nc}}(T)$ . In this way, the homological functors  $(\text{Ext}^{\Gamma}, \text{Tor}^{\Gamma})$  capture obstructions to gluing, descent and higher coherence for bi- $\Gamma$ -modules over  $\text{Spec}_{\Gamma}^{\text{nc}}(T)$ , and so provide the correct derived language for non-commutative  $\Gamma$ -geometry.

The present work occupies the middle position in a three-part series. In the commutative ternary case, the authors' earlier papers develop the ideal theory, radical theory and first instances of derived  $\Gamma$ -geometry on  $\text{Spec}_{\Gamma}(T)$  for commutative ternary  $\Gamma$ -semirings [23, 22, 24, 25]. The current paper extends the homological and categorical apparatus to the full non-commutative  $n$ -ary framework, and thus furnishes the tools needed for a genuinely derived, non-commutative  $\Gamma$ -geometry.

In Part III of the series we shall build on the homological structures constructed here to develop a more geometric theory over  $\text{Spec}_{\Gamma}^{\text{nc}}(T)$ . The planned topics include: quasi-coherent and coherent  $\Gamma$ -sheaves on non-commutative  $\Gamma$ -spectra; t-structures and truncation functors induced by  $\Gamma$ -radicals; Morita-type invariants of  $n$ -ary  $\Gamma$ -semirings expressed in terms of derived equivalences; and spectral constructions that relate  $\Gamma$ -primes, support theory and cohomological dimension. Taken together, the three parts are intended to provide a coherent framework in which ideal theory, radical and spectral theory, and derived homological methods for ternary and  $n$ -ary  $\Gamma$ -semirings can be treated within one unified geometric–homological paradigm.

## Acknowledgements

The author expresses sincere gratitude to the **Director of Collegiate Education**, Mangalagiri, Andhra Pradesh, **Dr. Ramachandra R. K., Principal**, Government College (Autonomous), Rajahmundry, for providing a supportive academic environment that greatly facilitated the completion of this research.

## Funding

The author received no external funding for this research.

## Ethics Statement

This work does not involve human participants, animals, or any ethical concerns. No ethical approval is required.

## Author Contributions

C. Gokavarapu is the sole author of this article and is responsible for the conceptual formulation, algebraic development, homological analysis, manuscript preparation, and final revision.

## References

- [1] S. Bourne, *The Jacobson radical of a semiring*, Proc. Natl. Acad. Sci. USA **37** (1951), 163–170.
- [2] J. S. Golan, *Semirings and Their Applications*, Kluwer Academic Publishers, Dordrecht, 1999.
- [3] U. Hebisch and H. J. Weinert, *Semirings: Algebraic Theory and Applications in Computer Science*, World Scientific, Singapore, 1998.
- [4] N. Nobusawa,  $\Gamma$ -rings and their applications, J. Sci. Hiroshima Univ. Ser. A **27** (1963), 17–29.
- [5] N. Nobusawa, *On a generalization of the ring theory*, Osaka J. Math. **1** (1964), 81–89.
- [6] M. Murali Krishna Rao,  $\Gamma$ -semirings I, Southeast Asian Bull. Math. **19** (1995), 49–54.
- [7] M. Murali Krishna Rao,  $\Gamma$ -semirings II, Southeast Asian Bull. Math. **21** (1997), 281–287.
- [8] M. Murali Krishna Rao, *The Jacobson radical of a  $\Gamma$ -semiring*, Southeast Asian Bull. Math. **23** (1999), 127–134.
- [9] H. Hedayati and K. P. Shum, *An introduction to  $\Gamma$ -semirings*, Int. J. Algebra **5** (2011), no. 13–16, 709–726.
- [10] T. K. Dutta and S. K. Sardar, *On prime ideals and prime radicals of a  $\Gamma$ -semiring*, An. Ştiinţ. Univ. “Al. I. Cuza” Iaşi, Ser. Mat. (N.S.) **46** (2000), no. 2, 319–329.
- [11] S. K. Sardar, B. C. Saha and K. P. Shum,  *$h$ -Prime and  $h$ -semiprime ideals in semirings and  $\Gamma$ -semirings*, Int. J. Algebra **4** (2010), no. 5, 209–220.
- [12] M. Murali Krishna Rao, *Ideals in ordered  $\Gamma$ -semirings*, Discuss. Math. Gen. Algebra Appl. **38** (2018), no. 1, 47–68.
- [13] M. F. Atiyah and I. G. Macdonald, *Introduction to Commutative Algebra*, Addison–Wesley, Reading, MA, 1969.
- [14] R. Hartshorne, *Algebraic Geometry*, Grad. Texts in Math., Vol. 52, Springer, New York, 1977.
- [15] A. Grothendieck, *Sur quelques points d’algèbre homologique*, Tôhoku Math. J. (2) **9** (1957), 119–221.
- [16] D. Quillen, *Higher algebraic K-theory. I*, in: *Algebraic K-theory, I: Higher K-theories* (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972), Lecture Notes in Math., Vol. 341, Springer, Berlin, 1973, pp. 85–147.
- [17] J.-L. Verdier, *Des catégories dérivées des catégories abéliennes*, Astérisque **239**, Soc. Math. France, Paris, 1996.
- [18] C. A. Weibel, *An Introduction to Homological Algebra*, Cambridge Stud. Adv. Math., Vol. 38, Cambridge Univ. Press, Cambridge, 1994.

- [19] T. Bühler, *Exact categories*, Expo. Math. **28** (2010), no. 1, 1–69.
- [20] A. Neeman, *Triangulated Categories*, Ann. of Math. Stud., Vol. 148, Princeton Univ. Press, Princeton, NJ, 2001.
- [21] D. Madhusudhana Rao, P. Rani and K. Kiran, *Foundations of Ternary Gamma-Semirings*, J. Algebraic Structures **15** (2025), 1–25.
- [22] C. Gokavarapu and D. Madhusudhana Rao, *Finite Structure and Radical Theory of Commutative Ternary  $\Gamma$ -Semirings*, preprint arXiv:2511.01789 (2025).
- [23] C. Gokavarapu and D. Madhusudhana Rao, *Prime and Semiprime Ideals in Commutative Ternary  $\Gamma$ -Semirings: Quotients, Radicals, Spectrum*, preprint arXiv:2510.23885 (2025).
- [24] C. Gokavarapu and M. R. Dasari, *Homological and Categorical Foundations of Ternary  $\Gamma$ -Modules and Their Spectra*, preprint arXiv:2511.02544 (2025).
- [25] C. Gokavarapu and D. Madhusudhana Rao, *Derived  $\Gamma$ -Geometry, Sheaf Cohomology, and Homological Functors on the Spectrum of Commutative Ternary  $\Gamma$ -Semirings*, preprint arXiv:2511.14108 (2025).
- [26] C. Gokavarapu, *Exact Categories and Homological Foundations of Non-Commutative  $n$ -ary  $\Gamma$ -Semirings*, arXiv:submit/7021614 (2025).