

ON THE 2-TORSION IN CLASS GROUPS OF NUMBER FIELDS

DANTE BONOLIS

ABSTRACT. In 2020, Bhargava, Shankar, Taniguchi, Thorne, Tsimerman, and Zhao proved that for a finite extension K/\mathbb{Q} of degree $n \geq 5$, the size of the 2-torsion class group is bounded by $\#h_2(K) = O_{n,\varepsilon}(D_K^{\frac{1}{2}-\frac{1}{2n}+\varepsilon})$, where D_K is the absolute discriminant of K . In the present paper, we improve their bound by proving that $\#h_2(K) = O_{n,\varepsilon}(D_K^{\frac{1}{2}-\frac{1}{2n}-\delta_K+\varepsilon})$, for a constant $\delta_K \geq \frac{1}{28n} - \frac{3}{28n(n-1)}$.

1. INTRODUCTION

For a given number field K/\mathbb{Q} of degree n and for any ℓ prime, a classical problem in algebraic number theory is to bound the size of $h_\ell(K)$, the ℓ -torsion subgroup of the class group $h(K)$. Since $\#h_\ell(K) \leq \#h(K)$, from the Brauer-Siegel Theorem (see [Lan94, Chapter XVI]) one deduces that

$$\#h_\ell(K) = O_{n,\varepsilon}(D_K^{1/2}(\log D_K)^{n-1}),$$

where D_K denotes the absolute value of the discriminant of K over \mathbb{Q} . Recently, Lemke Oliver and Zaman ([LZ25, Corollary 1]) showed that for every ℓ , one has $\#h_\ell(K) = o(D_K^{\frac{1}{2}})$ (and where the saving is an explicit power of $\log(D_K)$). On the other hand, it is conjectured that

$$\#h_\ell(K) = O_{n,\varepsilon}(D_K^\varepsilon), \tag{1}$$

for any $\varepsilon > 0$. This conjecture remains widely open; the only case currently known is $(n, \ell) = (2, 2)$, which follows from Gauss' genus theory. In fact, even establishing the existence of a constant $\delta_{n,\ell} > 0$ such that for any number field K of degree n , we have

$$\#h_\ell(K) \ll D_K^{\frac{1}{2}-\delta_{n,\ell}+\varepsilon}$$

It is already a challenging problem. In this paper, we focus on the case $\ell = 2$. It was shown in [BSTTZ20, Theorem 1.1] that one may take

$$\delta_{n,2} = \begin{cases} 0.2874\dots & \text{if } n = 3, 4, \\ \frac{1}{2n} & \text{if } n \geq 5. \end{cases}$$

Our main result is the following:

Theorem 1.1. *Let K/\mathbb{Q} be a number field of degree $n \geq 5$. Then, there exists $\delta_K > 0$ such that for any $\varepsilon > 0$, one has*

$$\#h_2(K) = O_{n,\varepsilon}(D_K^{\frac{1}{2}-\frac{1}{2n}-\delta_K+\varepsilon}).$$

Moreover we have that $\delta_K \geq \frac{1}{28n} - \frac{3}{28n(n-1)}$.

Date: December 1, 2025.

2. OUTLINE OF THE PROOF

We start by fixing some notation: let \mathcal{O}_K be the ring of integers of K . It is a well-known fact that \mathcal{O}_K is a \mathbb{Z} -lattice of rank n . Moreover, we have $\text{Disc}(\mathcal{O}_K) = D_K^{\frac{1}{2}}$. For every $\alpha \in \mathcal{O}_K$, consider the norm:

$$\|\alpha\| = \sum_{\sigma \in \text{Emb}(K/\mathbb{Q})} |\sigma(\alpha)|,$$

where $\text{Emb}(K/\mathbb{Q})$ denotes the set of embeddings of K into \mathbb{C} fixing \mathbb{Q} . Since \mathcal{O}_K is a lattice, we can find a basis $\{\alpha_{1,K} = 1, \alpha_{2,K}, \dots, \alpha_{n,K}\}$ which is Minkowski-reduced, i.e. such that $1 \leq \|\alpha_{2,K}\| \leq \dots \leq \|\alpha_{n,K}\|$, and for each $i \leq n-1$, $\|\alpha_{i,K}\| \ll \|v\|$ for every $v \notin \text{Span}\{1, \alpha_{2,K}, \dots, \alpha_{i-1,K}\}$. Observe that this implies that

$$\|\alpha_{2,K}\| \cdots \|\alpha_{n,K}\| \asymp D_K^{\frac{1}{2}}. \quad (2)$$

Finally, for each $i = 2, \dots, n$, we denote by $\delta_{i,K}$ the real positive number such that $\|\alpha_{i,K}\| = D_K^{\delta_{i,K}}$. At this point, it is useful to record the following

Proposition 2.1. *Let K/\mathbb{Q} be a finite extension of degree n . Then*

- i) $\|\alpha_{n,K}\| = O_n(D_K^{\frac{1}{n}})$. In particular we have $0 \leq \delta_{i,K} \leq \frac{1}{n} + o(1)$, for each $2 \leq i \leq n$;
- ii) $\|\alpha_{2,K}\| = O_n(D_K^{\frac{1}{2(n-1)}})$. In particular we have $0 \leq \delta_{2,K} \leq \frac{1}{2(n-1)} + o(1)$.

Proof. We start with (i). The bound $\|\alpha_{n,K}\| = O_n(D_K^{\frac{1}{n}})$ is proved in [BSTTZ20, Theorem 1.6]. Since the basis $\{1, \alpha_{2,K}, \dots, \alpha_{n,K}\}$ is Minkowski-reduced, we have $\|\alpha_{i,K}\| \leq \|\alpha_{n,K}\| = O(D_K^{\frac{1}{n}})$ for every $i \in \{2, \dots, n\}$. Hence, $\delta_{i,K} \leq \frac{1}{n} + o(1)$ for every $i \in \{2, \dots, n\}$, as claimed. For (ii), it is enough to observe that by (2),

$$D_K^{\frac{1}{2}} \gg_n \|\alpha_{2,K}\| \cdots \|\alpha_{n,K}\| \geq \|\alpha_{2,K}\|^{n-1}.$$

□

Our starting point is the same as [BSTTZ20, Theorem 1.1]: The size of the 2-torsion of a number field extension is bounded by

$$\#h_2(K) \ll \# \left\{ (y, z_1, \dots, z_n) \in \mathbb{Z}^{n+1} : \begin{array}{l} |z_i| \ll \frac{D_K^{\frac{1}{n}}}{\|\alpha_{i,K}\|}, \quad \text{for } i = 1, \dots, n \\ y^2 = N_K(z_1 + z_2\alpha_{2,K} + \dots + z_n\alpha_{n,K}) \end{array} \right\}.$$

In [BSTTZ20], the bound for $\#h_2(K)$ (when $n \geq 5$) is achieved by fixing a choice of z_2, \dots, z_n and by using the Bombieri-Pila bound ([BP89]) to count integral solutions (y, x_1) of the curve,

$$Y^2 = N_K(X_1 + z_2\alpha_{2,K} + \dots + z_n\alpha_{n,K}),$$

in the box $[-D_K^{\frac{1}{2}}, D_K^{\frac{1}{2}}] \times [-D_K^{\frac{1}{n}}, D_K^{\frac{1}{n}}]$. Similarly to Salberger's approach ([Sal22]), we shall instead fix only z_3, \dots, z_n and count integral solutions (y, x_1, x_2) on the surface

$$Y^2 = N_K(X_1 + X_2\alpha_{2,K} + z_3\alpha_{3,K} \cdots + z_n\alpha_{n,K}),$$

in the box

$$[-D_K^{\frac{1}{2}}, D_K^{\frac{1}{2}}] \times [-D_K^{\frac{1}{n}}, D_K^{\frac{1}{n}}] \times [-D_K^{\frac{1}{n}-\delta_{2,K}}, D_K^{\frac{1}{n}-\delta_{2,K}}]. \quad (3)$$

One of the main difficulties in using this approach is that the box in (3) has edges with highly unequal sizes: for instance, when $\frac{1}{2n} \leq \delta_{2,K} \leq \frac{1}{2(n-1)} + o(1)$, the range of the variable

X_2 is smaller than the square-root threshold of the range of the variable X_1 , making the analysis significantly more challenging. As a result, we are going to use a different approach depending on the size of $\delta_{2,K}$.

2.1. The case $\delta_{2,K} \leq \frac{1}{2n}$. In this case, it will be enough to deploy the classical square sieve as introduced in [HB84], getting

Theorem 2.2. *Let $n \geq 5$ and $\varepsilon > 0$. For any number field K of degree n , we have*

$$\#h_2(K) \ll_{n,\varepsilon} D_K^{\frac{1}{2} - \frac{1}{2n} - \left(\frac{1}{6n} - \frac{\delta_{2,K}}{3}\right) + \varepsilon}$$

That is, when $0 \leq \delta_{2,K} \leq \frac{1}{2n}$, Theorem 1.1 holds for $\delta_K = \frac{1}{6n} - \frac{\delta_{2,K}}{3}$. Moreover, since $\frac{1}{6n} - \frac{\delta_{2,K}}{3} < 0$ when $\delta_{2,K} \geq \frac{1}{2n}$, it follows the bound in Theorem 2.2 is actually worst than the one in [BSTTZ20, Theorem 1.1] in the range $\frac{1}{2n} \leq \delta_{2,K} \leq \frac{1}{2(n-1)} + o(1)$; hence, Theorem 2.2 alone would not suffice to prove Theorem 1.1.

2.2. The case $\frac{1}{2n} \leq \delta_{2,K} \leq \frac{1}{2(n-1)} + o(1)$. As mentioned earlier, this case will be the most complicated due to the fact that the range of the variable X_2 is considerably smaller than the one of X_1 . To address this, we combine the square sieve with the q -analogue of van der Corput (as in [Pie06]), proving

Theorem 2.3. *Let $n \geq 5$ be an integer and $\varepsilon > 0$. For any number field K of degree n , we have*

$$\#h_2(K) \ll_{n,\varepsilon} D_K^{\frac{1}{2} - \frac{1}{2n} - \left(\frac{1}{7n} - \frac{3\delta_{2,K}}{14}\right) + \varepsilon},$$

provided that $\frac{1}{4n} \leq \delta_{2,K}$.

That is, when $\frac{1}{4n} \leq \delta_{2,K} \leq \frac{1}{2(n-1)} + o(1)$, Theorem 1.1 holds for $\delta_K = \frac{1}{7n} - \frac{3\delta_{2,K}}{14}$. Thus, by defining

$$\delta_K = \max \left\{ \frac{1}{6n} - \frac{\delta_{2,K}}{3}, \frac{1}{7n} - \frac{3\delta_{2,K}}{14} \right\} \geq \frac{1}{28n} - \frac{3}{28n(n-1)},$$

one finally proves Theorem 1.1.

The main ingredient in proving Theorem 2.2 is the following stratification result for exponential sums

Proposition 2.4. *Let $F(X, Y) \in \mathbb{F}_p[X, Y]$ a square free polynomial such that*

$$F[X, Y] = \prod_{i=1}^n (X + a_i Y + b_i) \text{ over } \overline{\mathbb{F}}_p,$$

and assume that the following conditions hold:

- i) *There exists $i \in \{2, \dots, n\}$ such that $a_1 \neq a_i$.*
- ii) *$\frac{a_i - a_1}{a_j - a_1} \cdot (b_j - b_1) \neq b_i - b_1$ for any i, j such that $a_j \neq a_1$ and $i \neq j$.*
- iii) *For every i, j with $i \neq j$, $b_i \neq b_j$.*

Then there exists $A_s \subset \{(r, t) \in \mathbb{F}_p \times \mathbb{F}_p\}$ such that

i) for any $(r, t) \in (\mathbb{F}_p \times \mathbb{F}_p) \setminus A_s$, one has

$$\left| \sum_{(x,y,z) \in \mathbb{F}_p^3} \left(\frac{F(x,y)}{p} \right) \left(\frac{F(z,y+r)}{p} \right) e_p(s(x-z) + ty) \right| \ll_n p^{3/2}(s, t, p)^{1/2}.$$

ii) $\#A_s = O_n(1)$.

When $(s, p) = 1$, Proposition 2.4, implies that the exponential sum

$$\sum_{(x,y,z) \in \mathbb{F}_p^3} \left(\frac{F(x,y)}{p} \right) \left(\frac{F(z,y+r)}{p} \right) e_p(s(x-z) + ty) \quad (4)$$

exhibits square root cancellation for all (r, t) which are not contained in a set of codimension 0 in \mathbb{F}_p^2 . Then Proposition 2.4 is the analogue of showing that the A -number associated to the exponential sum 4, in the sense of [Kat89], is not zero. We remark that, in general, determining whether the A -number associated with an exponential sum does not vanishes is a difficult problem (see [Kat94], [FK01]).

3. PREVIOUS RESULTS

The size of the $h_\ell(K)$, for a given number field K of degree n , has been the object of extensive studies in the past years. As we mentioned in the introduction, the bound in (1) has been established only in the case $n, \ell = 2$. Many efforts have been made to prove power-saving bounds of the form

$$\#h_\ell(K) \ll_{n,\varepsilon} D_K^{\frac{1}{2} - \delta_{n,\ell} + \varepsilon}.$$

We can summarize all the cases in which some power-saving bound has been achieved in the following table:

Source	$\delta_{n,\ell}$	\mathbf{n}	ℓ	Other restrictions
Gauss	$1/2$	2	2	
[Pie05, Pie06]	$1/56$	2	3	
[HV06]	$0.0582 \dots$	2	3	
[EV07]	$1/6$	2, 3	3	
[CK25]	0.1806	2	3	
[HB24]	$1/(2\ell)$	2, 3	prime	$p \mid D_K \Rightarrow p \leq D_K^{\delta_\ell}$
[HB24]	$1/(4\ell)$	3	prime	pure cubic extension
[BSTTZ20]	$0.2215 \dots$	3, 4	2	
[EV07]	$1/168$	4	3	$\text{Gal}(K/\mathbb{Q}) \cong A_4$ or S_4
[EV07]	δ	4	3	$\text{Gal}(K/\mathbb{Q}) \cong C_4, V_4$, or D_4
[BSTTZ20]	$1/(2n)$	≥ 5	2	
[Sal22]	$\frac{2 - n\delta_{2,K} + 2\sqrt{1 - 2n\delta_{2,K}}}{n}$	≥ 5	2	$\delta_{2,K} \leq 1/(2n)$
[KW22]	$1/2$	p^r	p^s	$\text{Gal}(K/\mathbb{Q}) \cong G$, any p -group G
[Wan21, Wan20]	$\delta_{G,\ell}$	powerful > 2		$\text{Gal}(K/\mathbb{Q}) \cong G$, many nilp. grps G

We remark that Theorem 2.2 improves upon [Sal22] for all number field K with $0 \leq \delta_{2,K} \leq \frac{1}{2n}$. Significant efforts have been made to achieve unconditional improvements on

the trivial bound *on average across families of number fields*: [HP17], [EPW17], [Wid18], [FW18], [PTBW20], [An20], [FW21], [TZ23], [LOTZ24], [KT24], [LOWW25], and [CK25].

3.1. Some application of Theorem 2.3. We conclude the introduction by presenting some consequences of Theorem 2.3.

Corollary 3.1. *Let $\mathbb{Q} \subset F$ be a finite extension of \mathbb{Q} . Then for every $\mathbb{Q} \subset F \subset K$ extension of degree n , we have*

$$\#h_2(K) \ll_{D_F, n, \varepsilon} D_K^{\frac{1}{2} - \frac{2}{3n} + \varepsilon}.$$

Proof. Since $F \subset K$, one has that $\|\alpha_{2,K}\| \leq \|\alpha_{2,F}\|$. Hence, Theorem 2.3 implies

$$h_2(K) \ll_{n, \varepsilon} D_K^{\frac{1}{2} - \frac{1}{2n} - \left(\frac{1}{6n} - \frac{\delta_{2,K}}{3}\right) + \varepsilon} = D_K^{\frac{1}{2} - \frac{2}{3n} + \varepsilon} \|\alpha_{2,K}\|^{\frac{1}{3}} \leq D_K^{\frac{1}{2} - \frac{2}{3n} + \varepsilon} \|\alpha_{2,F}\|^{\frac{1}{3}} \ll_{D_F} D_K^{\frac{1}{2} - \frac{2}{3n} + \varepsilon}.$$

□

Corollary 3.2. *Let $\varepsilon > 0$ and let q be a prime number. Then*

$$\#h_2(\mathbb{Q}[\sqrt[n]{q}]) \ll_{n, \varepsilon} D_{\mathbb{Q}[\sqrt[n]{q}]}^{\frac{1}{2} - \frac{2}{3n} + \frac{1}{3n(n-1)} + \varepsilon}$$

Proof. One has $\|\alpha_{2,K}\| \ll q^{\frac{1}{n}}$ and $D_{\mathbb{Q}[\sqrt[n]{q}]} = q^{n-1}$. Then,

$$\|\alpha_{2,K}\| \ll q^{\frac{1}{n}} = (q^{n-1})^{\frac{1}{n(n-1)}} = D_{\mathbb{Q}[\sqrt[n]{q}]}^{\frac{1}{n(n-1)}},$$

which implies $\delta_{\mathbb{Q}[\sqrt[n]{q}], 2} \leq \frac{1}{n(n-1)}$. Hence, an application of Theorem 2.3 complete the proof of the Corollary. □

4. PROOF OF THEOREM 1.1

4.1. Some Lemmas. As mentioned in the introduction, the idea is to deploy the square sieve to detect how often the norm

$$N_K(X_1 + X_2\alpha_{2,K} + z_3\alpha_{3,K} + \cdots + z_n\alpha_{n,K}) =: F_{\mathbf{z}}(X_1, X_2),$$

is a square, for $(x_1, x_2) \in \left[-D_K^{\frac{1}{n}}, D_K^{\frac{1}{n}}\right] \times \left[-D_K^{\frac{1}{n} - \delta_{2,K}}, D_K^{\frac{1}{n} - \delta_{2,K}}\right] \cap \mathbb{Z}^2$, and where

$$\mathbf{z} \in \left[-D_K^{\frac{1}{n} - \delta_{3,K}}, D_K^{\frac{1}{n} - \delta_{3,K}}\right] \times \cdots \times \left[-D_K^{\frac{1}{n} - \delta_{n,K}}, D_K^{\frac{1}{n} - \delta_{n,K}}\right] \cap \mathbb{Z}^{n-2}$$

is fixed. Before applying the sieve, we must exclude certain choices of \mathbf{z} for which our method does not apply. Namely:

- (i) Those $\mathbf{z} \in \mathbb{Z}^{n-2}$ such that the polynomial $N_K(X_1 + X_2\alpha_{2,K} + z_3\alpha_{3,K} + \cdots + z_n\alpha_{n,K})$ is a square, i.e., the equation

$$Y^2 = N_K(X_1 + X_2\alpha_{2,K} + z_3\alpha_{3,K} + \cdots + z_n\alpha_{n,K})$$

has a solution for every choice of X_1, X_2 . We denote the set of these points as Z_{\square} .

- (ii) Those $\mathbf{z} \in \mathbb{Z}^{n-2}$ for which the the polynomial $F_{\mathbf{z}}(X_1, X_2)$ does not satisfies the hypotheses of Proposition 2.4 over $\overline{\mathbb{Q}}$. We denote the set of these points as Z_F .

To simplify the notation, from now on we denote

$$\mathcal{B} = \left[-D_K^{\frac{1}{n}}, D_K^{\frac{1}{n}}\right] \times \left[-D_K^{\frac{1}{n} - \delta_{2,K}}, D_K^{\frac{1}{n} - \delta_{2,K}}\right],$$

and

$$\mathcal{C} = \left[-D_K^{\frac{1}{n} - \delta_{3,K}}, D_K^{\frac{1}{n} - \delta_{3,K}}\right] \times \cdots \times \left[-D_K^{\frac{1}{n} - \delta_{n,K}}, D_K^{\frac{1}{n} - \delta_{n,K}}\right].$$

4.1.1. *The contribution of Z_\square .* We start by recording the following Lemma, which is proved in [BSTTZ20, Bottom of page 6] and will be useful in bounding the contribution of the $\mathbf{z} \in Z_\square$:

Lemma 4.1. *Let K/\mathbb{Q} be a number field and assume that there exists $\mathbb{Q} \subset F \subset K$, such that $[K : F] = 2$. Then $\#h_2(K) \ll_\varepsilon D_K^{\frac{1}{4}+\varepsilon}$.*

Corollary 4.2. *Assume that K/\mathbb{Q} does not contain a subfield of index 2. Then Z_\square is empty if $n = 5, 6, 7$, and*

$$\sum_{\mathbf{z} \in \mathcal{C} \cap Z_\square} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 \ll_{n, \varepsilon} D_K^{\frac{1}{4} + \frac{1}{n} + \varepsilon},$$

for $n \geq 8$.

Proof. If the polynomial $F_{\mathbf{z}}(X_1, X_2)$ is a square, then $F_{\mathbf{z}}(X_1, x_2) = N_{K/\mathbb{Q}}(X_1 + x_2\alpha_{2,K} + z_3x_3 + \cdots + z_n\alpha_{n,K})$ is a square for every value of $x_2 \in \mathbb{Z}$. On the other hand, since we are assuming that K does not contain any subfield of index 2, one can find (x_2, \mathbf{z}) such that $F_{\mathbf{z}}(X_1, x_2) = N_{K/\mathbb{Q}}(X_1 + x_2\alpha_{2,K} + z_3x_3 + \cdots + z_n\alpha_{n,K})$ is a square only if there exists $F \subset K$ such that $[K : F] > 2$ even (see [BSTTZ20, proof of Lemma 4.1]). Hence, for $n = 5, 6, 7$ the set Z_\square is actually empty. Let us assume now $n \geq 8$. We start by writing

$$\begin{aligned} \sum_{\mathbf{z} \in \mathcal{C} \cap Z_\square} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 &= \sum_{\substack{\mathbf{z} \in \mathcal{C} \cap \mathbb{Z}^{n-2} \\ F_{\mathbf{z}}(X_1, X_2) = \square}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 \\ &\leq \sum_{\substack{(x_2, \mathbf{z}) \in \mathbb{Z}^{n-1} \\ (x_2, \mathbf{z}) \in [-D_K^{\frac{1}{n} + \delta_2, K}, D_K^{\frac{1}{n} + \delta_2, K}] \times \mathcal{C} \\ F_{\mathbf{z}}(X_1, x_2) = \square}} \sum_{\substack{|x_1| \leq D_K^{\frac{1}{n}} \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1. \end{aligned}$$

An application of [BSTTZ20, Lemma 4.1] leads to

$$\sum_{\substack{(x_2, \mathbf{z}) \in \mathbb{Z}^{n-1} \\ (x_2, \mathbf{z}) \in [-D_K^{\frac{1}{n} + \delta_2, K}, D_K^{\frac{1}{n} + \delta_2, K}] \times \mathcal{C} \\ F_{\mathbf{z}}(X_1, x_2) = \square}} 1 \ll_{n, \varepsilon} D_K^{\frac{1}{4} + \varepsilon}$$

Hence

$$\sum_{\substack{(x_2, \mathbf{z}) \in \mathbb{Z}^{n-1} \\ (x_2, \mathbf{z}) \in [-D_K^{\frac{1}{n} + \delta_2, K}, D_K^{\frac{1}{n} + \delta_2, K}] \times \mathcal{C} \\ F_{\mathbf{z}}(X_1, x_2) = \square}} \sum_{\substack{|x_1| \leq D_K^{\frac{1}{n}} \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 \ll_{n, \varepsilon} D_K^{\frac{1}{4} + \frac{1}{n} + \varepsilon},$$

as claimed. \square

4.1.2. *The contribution of Z_F .* We now bound the contribution of the vectors in $\mathcal{C} \cap Z_F$. Since we have

$$\begin{aligned} F_{\mathbf{z}}[X_1, X_2] &= \prod_{i=1}^n (X_1 + \alpha_{2,K} X_2 + z_i) \text{ over } \overline{\mathbb{Q}} \\ &= \prod_{\sigma \in \text{Emb}(K)} (X_1 + \sigma(\alpha_{2,K}) X_2 + \sigma(z)) \text{ over } \overline{\mathbb{Q}}, \end{aligned}$$

the polynomial $F_{\mathbf{z}}[X_1, X_2]$ satisfies the hypothesis of Proposition 2.4 over $\overline{\mathbb{Q}}$, if

- a) the element z does not belong in any intermediate field $\mathbb{Q} \subset F \subset K$ (this will also guarantees that $F_{\mathbf{z}}$ is square-free);
- b) for every $\sigma, \tau \in \text{Emb}(K)$, with $\sigma \neq \tau$, and $\tau(\alpha_{2,K}) \neq \alpha$, one has

$$\sigma(z) - z \neq \frac{\sigma(\alpha_{2,K}) - \alpha_{2,K}}{\tau(\alpha_{2,K}) - \alpha_{2,K}}(\tau(z) - z).$$

Notice that, since $\alpha_2 \in K \setminus \mathbb{Q}$, the polynomial F always satisfies condition (i) of Proposition 2.4. We start with the following:

Lemma 4.3. *Let $\lambda \in K \setminus \{0, 1\}$, and let $\sigma, \tau \in \text{Emb}(K/\mathbb{Q})$ such that $\sigma \neq \tau$. Let $V_{\sigma, \tau, \lambda}$ be the \mathbb{Q} -vector space defined as*

$$V_{\sigma, \tau, \lambda} = \left\{ x \in K : x = \frac{1}{1-\lambda}\sigma(x) - \frac{\lambda}{1-\lambda}\tau(x) \right\}.$$

Then $\dim_{\mathbb{Q}}(V_{\sigma, \tau, \lambda}) \leq \frac{2n}{3}$.

Proof. Let L be the Galois closure of K , observe that we can extend any embedding $\text{Emb}(K/\mathbb{Q})$ to an element of $\text{Gal}(L/\mathbb{Q})$: for every $\rho \in \text{Emb}(K/\mathbb{Q})$, we will denote $\tilde{\rho}$ one of its extension to the whole L . Consider the linear map

$$\varphi : K \rightarrow L,$$

defined as $\varphi(x) = x - \frac{1}{1-\lambda}\sigma(x) + \frac{\lambda}{1-\lambda}\tau(x)$. Then $V_{\sigma, \tau, \lambda} = \text{Ker}(\varphi)$. Moreover, denote by $W_{\sigma, \tau, \lambda} = \text{Im}(\varphi)$. Let $d = \dim V_{\sigma, \tau, \lambda}$. Then $\dim_{\mathbb{Q}} W_{\sigma, \tau, \lambda} = n - d$, and let $\{h_1, \dots, h_{n-d}\}$ be a basis of for $W_{\sigma, \tau, \lambda}$. Let $\rho_1, \dots, \rho_{n-d+1} \in \text{Emb}(K/\mathbb{Q})$ distinct morphisms and consider the system

$$\begin{cases} X_1 \tilde{\rho}_1(h_1) + \dots + X_{n-d+1} \tilde{\rho}_{n-d+1}(h_1) = 0 \\ X_1 \tilde{\rho}_1(h_2) + \dots + X_{n-d+1} \tilde{\rho}_{n-d+1}(h_2) = 0 \\ \vdots \\ X_1 \tilde{\rho}_1(h_{n-d}) + \dots + X_{n-d+1} \tilde{\rho}_{n-d+1}(h_{n-d}) = 0 \end{cases}$$

Since this system has $n - d$ equations in $n - d + 1$ variables, one of the columns has to be a linear combination of the others. Hence, without loss of generality:

$$\begin{pmatrix} \tilde{\rho}_{n-d+1}(h_1) \\ \tilde{\rho}_{n-d+1}(h_2) \\ \vdots \\ \tilde{\rho}_{n-d+1}(h_{n-d}) \end{pmatrix} = \mu_1 \begin{pmatrix} \tilde{\rho}_1(h_1) \\ \tilde{\rho}_1(h_2) \\ \vdots \\ \tilde{\rho}_1(h_{n-d}) \end{pmatrix} + \dots + \mu_{n-d} \begin{pmatrix} \tilde{\rho}_{n-d}(h_1) \\ \tilde{\rho}_{n-d}(h_2) \\ \vdots \\ \tilde{\rho}_{n-d}(h_{n-d}) \end{pmatrix}$$

for some μ_1, \dots, μ_{n-d} . Knowing that h_1, \dots, h_{n-d} is a basis for $W_{\sigma, \tau, \lambda}$ and that φ vanishes on $V_{\sigma, \tau, \lambda}$, one has

$$\tilde{\rho}_{n-d+1} \circ \varphi = \mu_1(\tilde{\rho}_1 \circ \varphi) + \dots + \mu_{n-d}(\tilde{\rho}_{n-d} \circ \varphi).$$

By contradiction, assume that $d > \frac{2n}{3}$. Then, $n - d < \frac{n}{3}$. We claim that for every $k < \frac{n}{3}$, we can find $A_k = \{\text{Id} = \rho_1, \rho_2, \dots, \rho_{k+1}\} \subset \text{Emb}(K/\mathbb{Q})$, such that:

$$\rho_i \neq \rho_j, \quad \rho_i \neq \tilde{\rho}_j \circ \sigma, \quad \rho_i \neq \tilde{\rho}_j \circ \tau \quad \text{for any } j < i. \quad (5)$$

We do this by induction, when $k = 0$, we can take $\{\text{Id}\}$. Assume the claim is true for $k - 1$, then the set

$$B_{k-1} = \bigcup_{\rho \in A_{k-1}} \{\rho, \tilde{\rho} \circ \sigma, \tilde{\rho} \circ \tau\},$$

has cardinality $\#B_{k-1} \leq 3k < n$ (we are assuming $k < \frac{n}{3}$). Since $\text{Emb}(K/\mathbb{Q}) = n$, we can find $\rho_{k+1} \in \text{Emb}(K/\mathbb{Q}) \setminus A_k$ which proves the claim. Since $n - d < \frac{n}{3}$, the claim implies that there exists $A_{n-d} = \{\text{Id} = \rho_1, \dots, \rho_{n-d+1}\} \subset \text{Emb}(K/\mathbb{Q})$ such that for any $i = 1, \dots, n - d + 1$

$$\rho_i \neq \rho_j, \quad \rho_i \neq \tilde{\rho}_j \circ \sigma, \quad \rho_i \neq \tilde{\rho}_j \circ \tau \quad \text{for any } j < i.$$

Since we have $n - d + 1$ endomorphisms, by the previous part (reordering the morphisms if necessary), we have that

$$\tilde{\rho}_{n-d+1} \circ \varphi = \mu_1(\tilde{\rho}_1 \circ \varphi) + \dots + \mu_{n-d}(\tilde{\rho}_{n-d} \circ \varphi).$$

On the other hand, using the definition of φ , one would get that

$$\begin{aligned} \rho_{n-d+1} = & -\frac{1}{1 - \rho_{n-d+1}(\lambda)}(\tilde{\rho}_{n-d+1} \circ \sigma) + \frac{\rho_{n-d+1}(\lambda)}{1 - \rho_{n-d+1}(\lambda)}(\tilde{\rho}_i \circ \tau) \\ & + \sum_{i=1}^{n-d} \mu_i \left(\rho_i - \frac{1}{1 - \rho_i(\lambda)}(\tilde{\rho}_i \circ \sigma) + \frac{\rho_i(\lambda)}{1 - \rho_i(\lambda)}(\tilde{\rho}_i \circ \tau) \right), \end{aligned} \quad (6)$$

which would contradict the Artin-Dedekind's linear independence of characters ([Rot98, Page 76, Lemma 76]) since, by construction, the character ρ_{n-d+1} does not appear in the right-hand side of (6). \square

We recall that, for every $\mathbf{z} \in \mathbb{Z}^{n-2}$, we have $z = z_3\alpha_{3,K} + \dots + z_n\alpha_{n,K} \in K$. Moreover, in what follow we denote

$$E_{K,3} = \sum_{3 \leq j \leq \frac{2n}{3}} \left(\frac{1}{n} - \delta_{j,K} \right).$$

Corollary 4.4. *Let K/\mathbb{Q} be a number field of degree $n \geq 5$ which does not contain any subfield of index 2. Let $\sigma, \tau \in \text{Emb}(K/\mathbb{Q})$ be two distinct element in $\text{Emb}(K/\mathbb{Q})$ such that $\tau(\alpha_{2,K}) \neq \alpha_{2,K}$. Then*

$$\# \left\{ \mathbf{z} \in \mathbb{C} \cap \mathbb{Z}^{n-2} : \sigma(z) - z = \frac{\sigma(\alpha_{2,K}) - \alpha_{2,K}}{\tau(\alpha_{2,K}) - \alpha_{2,K}} \cdot (\tau(z) - z) \right\} \leq D_K^{E_{K,3}}$$

Proof. Consider the vector space

$$V_{\sigma,\tau} = \left\{ x \in K : \sigma(x) - x = \frac{\sigma(\alpha_{2,K}) - \alpha_{2,K}}{\tau(\alpha_{2,K}) - \alpha_{2,K}} \cdot (\tau(x) - x) \right\}.$$

We first handle the case when $\frac{\sigma(\alpha_{2,K}) - \alpha_{2,K}}{\tau(\alpha_{2,K}) - \alpha_{2,K}} = 1$. In this case, x satisfies $(\tau^{-1} \circ \sigma)(x) = x$, i.e. $x \in K^{\tau^{-1} \circ \sigma}$ which is a \mathbb{Q} -vectorspace of dimension at most $\frac{n}{3}$ (since we are assuming that K does not contain any subfield of index 2). Since $1 \in V_{\sigma,\tau}$, we have

$$\dim(\{(z_3, \dots, z_n) : z_3\alpha_{3,K} + \dots + z_n\alpha_{n,K} \in V_{\sigma,\tau}\}) \leq \frac{n}{3} - 1.$$

Thus,

$$\#\{\mathbf{z} \in \mathbb{C} \cap \mathbb{Z}^{n-2} : z \in V_{\sigma,\tau}\} \leq D_K^{\sum_{3 \leq j \leq \frac{n}{3}+1} (\frac{1}{n} - \delta_{j,K})} \leq D_K^{E_{K,3}},$$

since when $n \geq 5$, $\frac{n}{3} + 1 \leq \frac{2n}{3}$. We now assume $\frac{\sigma(\alpha_{2,K}) - \alpha_{2,K}}{\tau(\alpha_{2,K}) - \alpha_{2,K}} \neq 1$. In this case, we can apply Lemma 4.3 with $\lambda = \frac{\sigma(\alpha_{2,K}) - \alpha_{2,K}}{\tau(\alpha_{2,K}) - \alpha_{2,K}}$, getting $\dim_{\mathbb{Q}} V \leq \frac{2n}{3}$. Moreover, one has that $1, \alpha_{2,K} \in V_{\sigma,\tau}$, thus

$$\dim(\{(x_3, \dots, x_n) : x_3\alpha_{3,K} + \dots + x_n\alpha_{n,K} \in V\}) = \frac{2n}{3} - 2.$$

Thus, it follows that

$$\#(V_{\sigma,\tau} \cap \mathcal{C} \cap \mathbb{Z}^{n-2}) \leq D_K^{E_{K,3}},$$

as claimed. \square

Lemma 4.5. *Some notation as Corollary 4.13. Then*

$$\#\{\mathbf{z} \in \mathcal{C} \cap \mathbb{Z}^{n-2} : \sigma(z) = z, \text{ for some } \sigma \in \text{Emb}(K/\mathbb{Q}) \setminus \{\text{Id}\}\} \ll D_K^{E_{K,3}}$$

Proof. One has that

$$\#\{x \in K : \sigma(x) = x, \text{ for some } \sigma \in \text{Emb}(K/\mathbb{Q}) \setminus \{\text{Id}\}\} = \bigcup_{\substack{\sigma \in \text{Emb}(K/\mathbb{Q}) \\ \sigma \neq \text{Id}}} K^\sigma,$$

where K^σ is the field of all the element in K which are fixed by σ . Since $[K : K^\sigma] \geq 3$ (we are assuming that K does not contain any intermediate field of index 2), then K^σ is a \mathbb{Q} -vector field of dimension $\leq \frac{n}{3}$, for every $\sigma \in \text{Emb}(K/\mathbb{Q}) \setminus \{\text{Id}\}$. Moreover, since $1 \in K^\sigma$

$$\dim(\{(z_3, \dots, z_n) : z_3\alpha_{3,K} + \dots + z_n\alpha_{n,K} \in K^\sigma\}) \leq \frac{n}{3} - 1.$$

Hence,

$$\begin{aligned} \#\{\mathbf{z} \in \mathcal{C} \cap \mathbb{Z}^{n-2} : \sigma(z) = z, \text{ for some } \sigma \in \text{Emb}(K/\mathbb{Q}) \setminus \{\text{Id}\}\} &\ll D_K^{\sum_{3 \leq j \leq \frac{n}{3}+1} (\frac{1}{n} - \delta_{j,K})} \\ &\ll D_K^{\sum_{3 \leq j \leq \frac{2n}{3}} (\frac{1}{n} - \delta_{j,K})}, \end{aligned}$$

since when $n \geq 5$, $\frac{n}{3} + 1 \leq \frac{2n}{3}$ \square

We are finally ready to prove

Corollary 4.6. *Assume that K/\mathbb{Q} does not contain a subfield of index 2, and $\delta_{2,K} \geq \frac{1}{4n}$. Then*

$$\sum_{\substack{\mathbf{z} \in \mathcal{C} \\ \mathbf{z} \in Z_F \setminus Z_{\square}}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 \ll_{n,\varepsilon} D_K^{\frac{1}{2} - \frac{1}{2n} - \frac{1}{7n}}$$

Proof. We have:

$$Z_F = \left(\bigcup_{\substack{\sigma, \tau \in \text{Emb}(K/\mathbb{Q}) \\ \sigma \neq \tau \\ \tau(\alpha_{2,K}) \neq \alpha_{2,K}}} V_{\sigma,\tau} \right) \cup \left(\bigcup_{\substack{\sigma \in \text{Emb}(K/\mathbb{Q}) \\ \sigma \neq \text{Id}}} K^\sigma \right) \cap \mathbb{Z}^{n-2},$$

where

$$V_{\sigma,\tau} = \left\{ \mathbf{z} \in \mathcal{C} : \sigma(z) - z = \frac{\sigma(\alpha_{2,K}) - \alpha_{2,K}}{\tau(\alpha_{2,K}) - \alpha_{2,K}} \cdot (\tau(z) - z) \right\}.$$

Moreover, we recall that

$$\begin{aligned} E_{K,3} &= \sum_{3 \leq j \leq \frac{2n}{3}} \left(\frac{1}{n} - \delta_{j,K} \right). \\ \sum_{\substack{\mathbf{z} \in \mathcal{C} \\ \mathbf{z} \in Z_F \setminus Z_\square}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 &= \sum_{\substack{\mathbf{z} \in \mathcal{C} \cap Z_F \\ F_{\mathbf{z}}(X_1, X_2) \neq \square}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 \\ &= \sum_{\substack{\mathbf{z} \in \mathcal{C} \cap Z_F \\ F_{\mathbf{z}}(X_1, X_2) \neq \square}} \sum_{\substack{|x_2| \leq D_K^{\frac{1}{n} - \delta_{2,K}} \\ |x_1| \leq D_K^{\frac{1}{n}} \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 \\ &\ll_{n,\varepsilon} D_K^{\frac{1}{2n}} \cdot D_K^{\frac{1}{n} - \delta_{2,K}} \cdot \#(\mathcal{C} \cap Z_F) \end{aligned}$$

where in the last step, we used the Bombieri-Pila bound ([BP89]). Using Corollary 4.4 and Lemma 4.5, one gets

$$\#(\mathcal{C} \cap Z_F) \ll D_K^{E_{K,3}}.$$

On the other hand, (2), implies that

$$D_K^{\frac{1}{2n} + (\frac{1}{n} - \delta_{2,K}) + E_{K,3}} = D_K^{\frac{1}{2n} + \sum_{2 \leq j \leq \frac{2n}{3}} (\frac{1}{n} - \delta_{j,K})} \ll D_K^{\frac{1}{2} - \frac{1}{2n} - \sum_{j > \frac{2n}{3}} (\frac{1}{n} - \delta_{j,K})}.$$

If

$$\sum_{j > \frac{2n}{3} - 2} \left(\frac{1}{n} - \delta_{j,K} \right) < \frac{1}{7n},$$

then

$$\sum_{j > \frac{2n}{3} - 2} \delta_{j,K} < \frac{1}{3} + \frac{1}{n} - \frac{1}{7n}.$$

Since from (2) we have

$$\sum_{j=2}^n \delta_{j,K} = \frac{1}{2} + o(1);$$

This would imply that

$$\left\lfloor \frac{2n}{3} - 2 \right\rfloor \delta_{2,K} \leq \sum_{j=2}^{\lfloor \frac{2n}{3} - 2 \rfloor} \delta_{j,K} = \frac{1}{2} - \sum_{j > \frac{2n}{3} - 2} \delta_{j,K} < \frac{1}{2} - \frac{1}{3} - \frac{1}{n} + \frac{1}{7n} \leq \frac{1}{6} - \frac{6}{7n}$$

Thus, one deduces that

$$\delta_{2,K} < \left(\frac{1}{6} - \frac{6}{7n} \right) \cdot \left(\frac{2n}{3} - 3 \right)^{-1} \leq \frac{1}{4n}, \quad \text{when } n \geq 5$$

which would be in contradiction with the fact that we are assuming $\delta_{2,K} \geq \frac{1}{4n}$. Hence, we have

$$\sum_{j > \frac{2n}{3} - 2} \left(\frac{1}{n} - \delta_{j,K} \right) \geq \frac{1}{7n},$$

which leads to

$$\sum_{\substack{\mathbf{z} \in \mathcal{C} \cap Z_F \\ F_{\mathbf{z}}(X_1, X_2) \neq \square}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 \ll_{n, \varepsilon} D_K^{\frac{1}{2} - \frac{1}{2n} - \frac{1}{7n}},$$

which conclude the proof of the Lemma. \square

4.2. Preparation for the sieve: smooth weights. As previously mentioned, our strategy is to fix $\mathbf{z} = (z_3, \dots, z_n)$ and study how often

$$F_{\mathbf{z}}(x_1, x_2) = N_K(X_1 + X_2\alpha_{2,K} + z_3\alpha_{3,K} + \dots + z_n\alpha_{n,K})$$

is a square, as (x_1, x_2) ranges over the box

$$\mathcal{B}_{\mathbf{z}} := \left[-D_K^{\frac{1}{n}}, D_K^{\frac{1}{n}}\right] \times \left[-D_K^{\frac{1}{n} - \delta_{2,K}}, D_K^{\frac{1}{n} - \delta_{2,K}}\right].$$

Having this in mind, define

$$a_{\mathbf{z}}(n) = \sum_{\substack{(x_1, x_2) \in \mathcal{B}_{\mathbf{z}} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = n}} 1.$$

Then,

$$\#\{(x_1, x_2) \in \mathcal{B}_{\mathbf{z}} : F_{\mathbf{z}}(x_1, x_2) = \square\} \leq \sum_{k \in \mathbb{N}} a_{\mathbf{z}}(k^2). \quad (7)$$

Our goal is to estimate the right-hand side of (7) using the square sieve (including its version for composite moduli). As we will see in the upcoming sections, after applying the sieve lemma, we will need to apply the Poisson summation formula. For that reason, rather than working directly with the sequence $(a_{\mathbf{z}}(n))_{n \in \mathbb{N}}$, we will instead introduce suitable smooth weights. We begin by choosing two smooth functions $g_1, g_2 : \mathbb{R} \rightarrow \mathbb{R}_{\geq 0}$ satisfying the following properties:

- i) g_1 is supported on $\left[-D_K^{\frac{1}{n}}, D_K^{\frac{1}{n}}\right]$ and g_2 is supported on $\left[-D_K^{\frac{1}{n} - \delta_{2,K}}, D_K^{\frac{1}{n} - \delta_{2,K}}\right]$
- ii) for any $k \geq 0$, we have that

$$\left| \frac{\partial^k g_1}{\partial x_1^k} \right| \ll D_K^{-\frac{k}{n}}, \quad \left| \frac{\partial^k g_2}{\partial x_2^k} \right| \ll D_K^{-k(\frac{1}{n} - \delta_{2,K})}. \quad (8)$$

From the condition above, one deduces the following bounds on the Fourier transforms of g_1 , and g_2

$$|\hat{g}_1(u)| \ll D_K^{\frac{1}{n}} \left| 1 + u D_K^{\frac{1}{n}} \right|^{-2}, \quad |\hat{g}_2(u)| \ll D_K^{\frac{1}{n} - \delta_{2,K}} \left| 1 + u D_K^{\frac{1}{n} - \delta_{2,K}} \right|^{-2}.$$

Then, for any $n \geq 0$ we define

$$\omega_{\mathbf{z}}(n) = \sum_{\substack{(x_1, x_2) \in \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = n}} g_1(x_1) g_2(x_2). \quad (9)$$

We remark that when $n = 0$,

$$\omega_{\mathbf{z}}(n) = \begin{cases} g_1(0)g_2(0) & \text{if } \mathbf{z} = \mathbf{0} \\ 0 & \text{otherwise,} \end{cases}$$

since $F_{\mathbf{z}}(x_1, x_2) = N_K(x_1 + x_2\alpha_{2,K} + z_3\alpha_{3,K} + \dots + z_n\alpha_{n,K})$ and $x \in K$ is such that $N_K(x) = 0$ if and only if $x = 0$.

4.3. The square sieve: proof of Theorem 2.3. We record the following Sieve Lemma due to Heath-Brown [HB84]:

Lemma 4.7. *Let \mathcal{P} be a finite set of P prime numbers. Let $(\omega(n))_{n \in \mathbb{N}}$ be a sequence of non negative number such that $\omega(n) = 0$ if either $n = 0$, or $n > e^P$. Then*

$$\sum_k \omega(k^2) \ll \frac{1}{P} \sum_n \omega(n) + \frac{1}{P^2} \sum_{\substack{p, q \in \mathcal{P} \\ p \neq q}} \left| \sum_n \omega(n) \left(\frac{n}{pq} \right) \right|.$$

We will use the sieve inequality in Lemma 4.7 to show:

Lemma 4.8. *Let $\mathbf{z} \in \mathbb{Z}^{n-2} \setminus \{\mathbf{0}\}$ such that $F_{\mathbf{z}}(X_1, X_2)$ is not a square in $\mathbb{Z}[X_1, X_2]$, and consider the sequence $(\omega_{\mathbf{z}}(n))_{n \in \mathbb{N}}$ defined in (9). Then,*

$$\sum_k \omega_{\mathbf{z}}(k^2) \ll_{n, \varepsilon} D_K^{\frac{4}{3n} - \frac{2\delta_{2,K}}{3} + \varepsilon}.$$

To prove Lemma 4.8, we start by applying Lemma 4.7 to the sequence $(\omega_{\mathbf{z}}(n))_n$:

$$\sum_k \omega_{\mathbf{z}}(k^2) \ll \frac{1}{P} \sum_n \omega_{\mathbf{z}}(n) + \frac{1}{P^2} \sum_{\substack{p, q \in \mathcal{P} \\ p \neq q}} \left| \sum_n \omega_{\mathbf{z}}(n) \left(\frac{n}{pq} \right) \right|, \quad (10)$$

provided that $P > (\log D_K)^2$. By the definition of the sequence $\omega_{\mathbf{z}}(n)$, it follows that

$$\frac{1}{P} \sum_n \omega_{\mathbf{z}}(n) \ll \frac{D_K^{\frac{2}{n} - \delta_{2,K}}}{P}. \quad (11)$$

Hence, it remains to bound the second term on the left-hand side of (10). We start by observing that

$$\begin{aligned} S(\mathbf{z}, p, q) &= \sum_{n \in \mathbb{N}} \omega_{\mathbf{z}}(n) \left(\frac{n}{pq} \right) \\ &= \sum_{(x_1, x_2) \in \mathbb{Z}^2} g_1(x_1) g_2(x_2) \left(\frac{F_{\mathbf{z}}(x_1, x_2)}{pq} \right) \\ &= \sum_{(a_1, a_2) \mid (pq)} \left(\frac{F_{\mathbf{z}}(a_1, a_2)}{pq} \right) \sum_{\substack{x_1 \in \mathbb{Z} \\ x_1 \equiv a_1 \pmod{pq}}} g_1(x_1) \sum_{\substack{x_2 \in \mathbb{Z} \\ x_2 \equiv a_2 \pmod{pq}}} g_2(x_2) \\ &= \frac{1}{(pq)^2} \sum_{(u_1, u_2) \in \mathbb{Z}^2} \hat{g}_1 \left(\frac{u_1}{pq} \right) \hat{g}_2 \left(\frac{u_2}{pq} \right) T(\mathbf{u}, \mathbf{z}; pq), \end{aligned} \quad (12)$$

where for every $r \geq 2$ square-free and $\mathbf{v} \in \mathbb{Z}^2$,

$$T(\mathbf{v}, \mathbf{z}; r) = \sum_{(a_1, a_2) \mid (r)} \left(\frac{F_{\mathbf{z}}(a_1, a_2)}{r} \right) e_r(a_1 v_1 + a_2 v_2).$$

An application of the Chinese Remainder Theorem shows that

$$S(\mathbf{z}, p, q) = \frac{1}{(pq)^2} \sum_{(u_1, u_2) \in \mathbb{Z}^2} \hat{g}_1\left(\frac{u_1}{pq}\right) \hat{g}_2\left(\frac{u_2}{pq}\right) T(\bar{q}\mathbf{u}, \mathbf{z}; p) T(\bar{p}\mathbf{u}, \mathbf{z}; q).$$

4.3.1. *The exponential sum $T(\mathbf{v}, \mathbf{z}; r)$.* In the next session (see Lemma 5.6), we shall prove the following

Lemma 4.9. *Let $\mathbf{z} = (z_3, \dots, z_n) \in \mathbb{Z}^{n-2}$, such that $F_{\mathbf{z}}(X_1, X_2)$ is not a square. Then for every prime number p such that*

- i) the polynomial $F_{\mathbf{z}}(X_1, X_2) \pmod{p}$ is not a square;*
- ii) $p \nmid N_K(\sigma(\alpha_2) - \alpha_2)$, for every $\sigma \in \text{Emb}(K/\mathbb{Q})$ such that $\sigma(\alpha_2) \neq \alpha_2$.*

$$|T(\mathbf{v}, \mathbf{z}; p)| \ll_n p(p, v_1, v_2)^{\frac{1}{2}}.$$

As a consequence of Lemma 4.9, we obtain

Corollary 4.10. *Let $Q > 0$ and let p, q be primes such that $Q < p, q < 2Q$ and assume*

- i) the polynomials $F_{\mathbf{z}}(X_1, X_2) \pmod{p}$, $F_{\mathbf{z}}(X_1, X_2) \pmod{q}$ are not squares in \mathbb{F}_p and \mathbb{F}_q respectively;*
- ii) $p, q \nmid N_K(\sigma(\alpha_2) - \alpha_2)$, for every $\sigma \in \text{Emb}(K/\mathbb{Q})$ such that $\sigma(\alpha_2) \neq \alpha_2$.*

Then

$$S(\mathbf{z}, pq) \ll_n Q^2 + \frac{D_K^{\frac{2}{n}-\delta_{2,K}}}{Q},$$

provided that $D_K^{\frac{1}{2n}} \leq Q \leq D_K^{\frac{1}{n}-\delta_{2,K}}$.

Proof. Since the bound for $T(\mathbf{u}, \mathbf{z}, pq)$ depends on the G.C.D. between u_1, u_2 and pq we start by rewriting $S(\mathbf{z}, pq)$ as

$$S(\mathbf{z}, pq) = \frac{1}{(pq)^2} \sum_{\substack{\mathbf{u} \in \mathbb{Z}^2 \\ (\mathbf{u}, pq)=1}} \hat{g}_1\left(\frac{u_1}{pq}\right) \hat{g}_2\left(\frac{u_2}{pq}\right) T(\mathbf{u}, \mathbf{z}; pq) + \frac{1}{(pq)^2} \sum_{\substack{\mathbf{u} \in \mathbb{Z}^2 \\ (\mathbf{u}, pq)>1}} \hat{g}_1\left(\frac{u_1}{pq}\right) \hat{g}_2\left(\frac{u_2}{pq}\right) T(\mathbf{u}, \mathbf{z}; pq)$$

From (8), it follows that

$$\left| \hat{g}_1\left(\frac{u_1}{pq}\right) \hat{g}_2\left(\frac{u_2}{pq}\right) \right| \ll D_K^{\frac{2}{n}-\delta_{2,K}} \left(1 + \frac{D_K^{\frac{1}{n}} u_1}{pq}\right)^{-2} \cdot \left(1 + \frac{D_K^{\frac{1}{n}-\delta_{2,K}} u_2}{pq}\right)^{-2}.$$

On the other hand, by applying Lemma 4.9, one gets

$$\begin{aligned} \frac{1}{(pq)^2} \sum_{\substack{\mathbf{u} \in \mathbb{Z}^2 \\ (\mathbf{u}, pq)=1}} |T(\mathbf{u}, \mathbf{z}; pq)| \cdot \left| \hat{g}_1\left(\frac{u_1}{pq}\right) \hat{g}_2\left(\frac{u_2}{pq}\right) \right| &\ll_n \\ &\ll_n pq \cdot \frac{D_K^{\frac{2}{n}-\delta_{2,K}}}{(pq)^2} \sum_{\mathbf{u} \in \mathbb{Z}^2} \left(1 + \frac{D_K^{\frac{1}{n}} u_1}{pq}\right)^{-2} \cdot \left(1 + \frac{D_K^{\frac{1}{n}-\delta_{2,K}} u_2}{pq}\right)^{-2} \\ &\ll Q^2, \end{aligned}$$

provided that $D_K^{\frac{1}{n}} \leq pq \leq 4Q^2$. We now deal with the contributions of those \mathbf{u} for which $(\mathbf{u}, pq) > 1$:

$$\begin{aligned} \frac{1}{(pq)^2} \sum_{\substack{\mathbf{u} \in \mathbb{Z}^2 \\ (\mathbf{u}, pq) > 1}} \hat{g}_1\left(\frac{u_1}{pq}\right) \hat{g}_2\left(\frac{u_2}{pq}\right) T(\mathbf{u}, \mathbf{z}; pq) &\ll_n (pq)^{\frac{3}{2}} \cdot \frac{1}{(pq)^2} \sum_{\substack{\mathbf{u} \in \mathbb{Z}^2 \\ p|\mathbf{u}}} \hat{g}_1\left(\frac{u_1}{pq}\right) \hat{g}_2\left(\frac{u_2}{pq}\right) \\ &+ (pq)^{\frac{3}{2}} \cdot \frac{1}{(pq)^2} \sum_{\substack{\mathbf{u} \in \mathbb{Z}^2 \\ q|\mathbf{u}}} \hat{g}_1\left(\frac{u_1}{pq}\right) \hat{g}_2\left(\frac{u_2}{pq}\right). \end{aligned}$$

Provided that $q \leq D_n^{\frac{1}{n}-\delta_{2,K}}$, one has

$$\begin{aligned} \sum_{\substack{\mathbf{u} \in \mathbb{Z}^2 \\ q|\mathbf{u}}} \hat{g}_1\left(\frac{u_1}{pq}\right) \hat{g}_2\left(\frac{u_2}{pq}\right) &\ll D_K^{\frac{2}{n}-\delta_{2,K}} \sum_{\substack{\mathbf{u} \in \mathbb{Z}^2 \\ q|\mathbf{u}}} \left(1 + \frac{D_K^{\frac{1}{n}} u_1}{pq}\right)^{-2} \cdot \left(1 + \frac{D_K^{\frac{1}{n}-\delta_{2,K}} u_2}{pq}\right)^{-2} \\ &\ll D_K^{\frac{2}{n}-\delta_{2,K}}, \end{aligned}$$

which is enough to complete the proof of the Lemma. \square

We are finally ready to complete the proof of Lemma 4.8: let Q be such that $D_K^{\frac{1}{2n}} \leq Q \leq D_K^{\frac{1}{n}-\delta_{2,K}}$ and take

$$\mathcal{P} = \left\{ p : \begin{array}{l} Q \leq p \leq 2Q \\ F_{\mathbf{z}}(X_1, X_2) \pmod{p} \text{ is not a square in } \mathbb{F}_p \\ p \nmid N_K(\sigma(\alpha_2) - \alpha_2) \text{ for all } \sigma \in \text{Emb}(K/\mathbb{Q}) \text{ such that } \sigma(\alpha_2) \neq \alpha_2 \end{array} \right\}.$$

Then combining (10), (11), and Corollary 4.10, one gets

$$\sum_{n \in \mathbb{Z}} \omega_{\mathbf{z}}(k^2) \ll_n \frac{D_K^{\frac{2}{n}-\delta_{2,K}} \log Q}{Q} + Q^2.$$

The optimal choice here is when $Q = D_K^{\frac{2}{3n}-\frac{\delta_{2,K}}{3}}$. So we get

$$\sum_{n \in \mathbb{Z}} \omega_{\mathbf{z}}(k^2) \ll_{n,\varepsilon} D_K^{\frac{4}{3n}-\frac{2\delta_{2,K}}{3}+\varepsilon},$$

provided that $D_K^{\frac{1}{n}} \leq D_K^{\frac{4}{3n}-\frac{2\delta_{2,K}}{3}}$ and $D_K^{\frac{2}{3n}-\frac{\delta_{2,K}}{3}} \leq D_n^{\frac{1}{n}-\delta_{2,K}}$, which is satisfied as soon as $\delta_{2,K} \leq \frac{1}{2n}$.

4.3.2. Proof of Theorem 2.2. We can now prove Theorem 2.2. Thanks to Lemma 4.1, we may assume that K does not contain a subfield of index 2. We recall that

$$\mathcal{C} = \left[-D_K^{\frac{1}{n}-\delta_{3,K}}, D_K^{\frac{1}{n}-\delta_{3,K}}\right] \times \cdots \times \left[-D_K^{\frac{1}{n}-\delta_{n,K}}, D_K^{\frac{1}{n}-\delta_{n,K}}\right],$$

and that

$$Z_{\square} = \{\mathbf{z} \in \mathbb{Z}^{n-2} : F_{\mathbf{z}}(X_1, X_2) \text{ is a square}\}.$$

Then

$$\begin{aligned}
\#h_2(K) &\ll \# \left\{ (y, z_1, \dots, z_n) \in \mathbb{Z}^{n+1} : \begin{array}{l} |z_i| \ll D_K^{\frac{1}{n}-\delta_{i,K}}, \quad \text{for } i = 1, \dots, n \\ y^2 = N_K(z_1 + z_2\alpha_{2,K} + \dots + z_n\alpha_{n,K}) \end{array} \right\} \\
&= \sum_{\mathbf{z} \in \mathcal{C} \cap \mathbb{Z}^{n-2}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 \\
&= \sum_{\mathbf{z} \in \mathcal{C} \cap Z_{\square}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 + \sum_{\substack{\mathbf{z} \in \mathcal{C} \cap \mathbb{Z}^{n-2} \\ \mathbf{z} \notin Z_{\square}}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1.
\end{aligned} \tag{13}$$

For the second term in (13), we can use Lemma 4.8, getting

$$\begin{aligned}
\sum_{\substack{\mathbf{z} \in \mathcal{C} \cap \mathbb{Z}^{n-2} \\ F_{\mathbf{z}}(X_1, X_2) \neq \square}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 &\ll_{n, \varepsilon} \#(\mathcal{C} \cap \mathbb{Z}) \cdot D_K^{\frac{4}{3n} - \frac{2\delta_{2,K}}{3} + \varepsilon} \\
&= D_K^{\sum_{j=3}^n (\frac{1}{n} - \delta_{j,K}) + \frac{4}{3n} - \frac{2\delta_{2,K}}{3} + \varepsilon} \\
&= D_K^{\frac{1}{2} - \frac{2}{3n} + \frac{\delta_{2,K}}{3} + \varepsilon}.
\end{aligned}$$

On the other, thanks to Lemma 4.2 we know that

$$\sum_{\mathbf{z} \in \mathcal{C} \cap Z_{\square}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 \ll_{n, \varepsilon} D_K^{\frac{1}{4} + \frac{1}{n} + \varepsilon},$$

when $n \geq 8$ and zero otherwise. Since $\frac{1}{4} + \frac{1}{n} \leq \frac{1}{2} - \frac{2}{3n}$, for $n > 6$, we complete the proof of Theorem 2.2.

4.4. The square sieve for composite moduli: bounding $h_2(K)$ when $\delta_{2,K}$ is large.
In this section, we are going to prove Theorem 2.3. As mentioned in the introduction, we do this by deploying the square sieve for composite moduli. We record here the sieve Lemma as stated in [HP12, Lemma 2]

Lemma 4.11. *Let $A = |\mathcal{A}|$, $U = |\mathcal{U}|$, and $V = |\mathcal{V}|$, so that $A = UV$. Furthermore, assume that $V^3 \ll A$. Let ω be a non-negative weight such that $\omega(n) = 0$ for $n > \exp \min(U, V)$, then*

$$\begin{aligned}
\sum_{n \neq 0} \omega(n) &\ll \frac{1}{A} \sum_{n \neq 0} \omega(n) + \frac{1}{A^2} \sum_{v, v' \in \mathcal{V}} \sum_{\substack{u, u' \in \mathcal{U} \\ u \neq u'}} \left| \sum_{n \in \mathbb{Z}} \left(\frac{n}{uu'} \right) \left(\frac{n}{vv'} \right) \omega(n) \right| \\
&\quad + \frac{U}{A^2} \sum_{\substack{v, v' \in \mathcal{V} \\ v \neq v'}} \left| \sum_{n \in \mathbb{Z}} \left(\frac{n}{vv'} \right) \omega(n) \right|.
\end{aligned}$$

In the remainder of this section we will use Lemma 4.11, to prove the following

Lemma 4.12. *Let $\mathbf{z} \in \mathbb{Z}^{n-2} \setminus \{\mathbf{0}\}$ satisfying the following condition:*

- i) *The polynomial $F_{\mathbf{z}}(X_1, X_2)$ is not a square in $\mathbb{Z}[X_1, X_2]$.*

ii) For every $\sigma, \tau \in \text{Emb}(K/\mathbb{Q})$, with $\sigma \neq \tau$ and $\tau(\alpha_{2,K}) \neq \alpha_{2,K}$,

$$\sigma(z) - z \neq \frac{\sigma(\alpha_{2,K}) - \alpha_{2,K}}{\tau(\alpha_{2,K}) - \alpha_{2,K}}(\tau(z) - z),$$

iii) For every $\sigma \in \text{Emb}(K/\mathbb{Q}) \setminus \{\text{Id}\}$, $z \notin K^\sigma$.

Let $(\omega_{\mathbf{z}}(n))_n$ be the sequence defined in (7). Then,

$$\sum_k \omega_{\mathbf{z}}(k^2) \ll_{n,\varepsilon} D_K^{\frac{45}{28n} - \frac{11}{14}\delta_{2,K} + \varepsilon}$$

and ultimately, use Lemma 4.12 to prove Theorem 2.3.

To prove Lemma 4.12, we start by applying Lemma 4.11, getting

$$\begin{aligned} \sum_{n \in \mathbb{Z}} \omega_{\mathbf{z}}(n^2) &= \frac{1}{A} \sum_n \omega_{\mathbf{z}}(n) + \frac{1}{A^2} \sum_{v,v' \in \mathcal{V}} \sum_{\substack{u,u' \in \mathcal{U} \\ u \neq u'}} \left| \sum_{n \in \mathbb{Z}} \left(\frac{n}{uu'} \right) \left(\frac{n}{vv'} \right) \omega_{\mathbf{z}}(n) \right| \\ &\quad + \frac{U}{A^2} \sum_{\substack{v,v' \in \mathcal{V} \\ v \neq v'}} \left| \sum_{n \in \mathbb{Z}} \left(\frac{n}{vv'} \right) \omega_{\mathbf{z}}(n) \right|. \end{aligned} \tag{14}$$

where \mathcal{U}, \mathcal{V} are finite families of primes that will be chosen later. Again, the first term on the right-hand side of (14) can be easily bounded by

$$\frac{1}{A} \sum_n \omega_{\mathbf{z}}(n) \ll \frac{D_K^{\frac{2}{n} - \delta_{2,K}}}{A}. \tag{15}$$

Hence, we need to focus on the two other terms. We will refer to the term

$$M = \frac{1}{A^2} \sum_{v,v' \in \mathcal{V}} \sum_{\substack{u,u' \in \mathcal{U} \\ u \neq u'}} \left| \sum_{n \in \mathbb{Z}} \left(\frac{n}{uu'} \right) \left(\frac{n}{vv'} \right) \omega_{\mathbf{z}}(n) \right|, \tag{16}$$

as the main sieve and to the term

$$N = \frac{U}{A^2} \sum_{\substack{v,v' \in \mathcal{V} \\ v \neq v'}} \left| \sum_{n \in \mathbb{Z}} \left(\frac{n}{vv'} \right) \omega_{\mathbf{z}}(n) \right|,$$

as the prime sieve.

Proof of Lemma 4.12: the main sieve. In what follows, we denote $q_1 = uu'$, $q_2 = vv'$. We start again from

$$S(\mathbf{z}; q_1 q_2) = \sum_{(x_1, x_2) \in \mathbb{Z}^2} g_1(x_1) g_2(x_2) \left(\frac{F_{\mathbf{z}}(x_1, x_2)}{q_1 q_2} \right)$$

but this time, the first step will be to extend only the summation in x_1 over all the classes modulo q_1q_2 :

$$\begin{aligned} S(\mathbf{z}; q_1q_2) &= \sum_{x_2 \in \mathbb{Z}} g_2(x_2) \sum_{y_1 \pmod{q_1q_2}} \left(\frac{F_{\mathbf{z}}(y_1, x_2)}{q_1q_2} \right) \sum_{\substack{x_2 \in \mathbb{Z} \\ x_1 \equiv y_1 \pmod{q_1q_2}}} g_1(x_1) \\ &= \frac{1}{q_1q_2} \sum_{u_1} \hat{g}_1 \left(\frac{u_1}{q_1q_2} \right) S(\mathbf{z}, u_1; q_1q_2), \end{aligned}$$

where

$$S(\mathbf{z}, u_1; q_1q_2) = \sum_{x_2 \in \mathbb{Z}} g_2(x_2) Z(u_1, x_2; q_1q_2),$$

and where for every r square-free number,

$$Z(u_1, x_2; r) = \sum_{y_1 \pmod{r}} \left(\frac{F_{\mathbf{z}}(y_1, x_2)}{r} \right) e_r(u_1 y_1).$$

Again, applying the Chinese Remainder Theorem, one has that

$$S(\mathbf{z}, u_1; q_1q_2) = \sum_{x_2 \in \mathbb{Z}} g_2(x_2) Z(u_1 \overline{q_2}, x_2; q_1) Z(u_1 \overline{q_1}, x_2; q_2)$$

We now deploy the q -analogue of the Van der Corput method: we denote $H = \lfloor \frac{D_K^{\frac{1}{n}-\mu_2}}{q_2} \rfloor$. Then

$$\begin{aligned} HS(\mathbf{z}, u_1; q_1q_2) &= \sum_{h \in [0, H]} \sum_{x_2 \in \mathbb{Z}} g_2(x_2 + q_2 h) Z(u_1 \overline{q_2}, x_2 + q_2 h; q_1) Z(u_1 \overline{q_1}, x_2 + q_2 h; q_2) \\ &= \sum_{x_2 \in I} Z(u_1 \overline{q_1}, x_2; q_2) \sum_{h \in [0, H]} g_2(x_2 + q_2 h) Z(u_1 \overline{q_2}, x_2 + q_2 h; q_1), \end{aligned}$$

where $I = [D_K^{\frac{1}{n}-\delta_2, \kappa} - q_2 H, D_K^{\frac{1}{n}-\delta_2, \kappa} - q_2]$. Applying Cauchy-Schwarz,

$$H^2 |S(\mathbf{z}, u_1; q_1q_2)|^2 \ll \Sigma_1 \Sigma_2,$$

where

$$\begin{aligned} \Sigma_1 &= \sum_{x_2 \in I} |Z(u_1 \overline{q_1}, x_2; q_2)|^2 \\ \Sigma_2 &= \sum_{x_2 \in I} \left| \sum_{h \in [0, H]} g_2(x_2 + q_2 h) Z(u_1 \overline{q_2}, x_2 + q_2 h; q_1) \right|^2. \end{aligned}$$

Expanding the square in Σ_2 , we get

$$\Sigma_2 = \sum_{h_1, h_2 \in [0, H]} R(h_1, h_2),$$

where

$$R(h_1, h_2) = \sum_{x_2 \in \mathbb{Z}} g_2(x_2 + q_2 h_1) g_2(x_2 + q_2 h_2) Z(u_1 \overline{q_2}, x_2 + q_2 h_1; q_1) \overline{Z(u_1 \overline{q_2}, x_2 + q_2 h_2; q_1)}.$$

Since $R(h_1, h_2) = R(h_1 - h_2, 0)$, we get that

$$\begin{aligned}\Sigma_2 &= \sum_{h_1, h_2 \in [0, H]} R(h_1 - h_2, 0) \\ &= \sum_{h \in [-H, H]} (H - |h|) R(h, 0) \\ &\ll H \sum_{h \in [-H, H]} \left| \sum_{x_2 \in \mathbb{Z}} g_2(x_2) g_2(x_2 + q_2 h) Z(u_1 \overline{q_2}, x_2; q_1) \overline{Z(u_1 \overline{q_2}, x_2 + q_2 h; q_1)} \right|,\end{aligned}$$

Then

$$\Sigma_2 \leq H(\Sigma_{2A} + \Sigma_{2B}),$$

where,

$$\begin{aligned}\Sigma_{2A} &= \sum_{x_2 \in \mathbb{Z}} (g(x_2))^2 |Z(u_1 \overline{q_2}, x_2; q_1)|^2 \\ \Sigma_{2B} &= \sum_{\substack{h \in [-H, H] \\ h \neq 0}} \left| \sum_{x_2 \in \mathbb{Z}} g_2(x_2) g_2(x_2 + q_2 h) Z(u_1 \overline{q_2}, x_2; q_1) \overline{Z(u_1 \overline{q_2}, x_2 + q_2 h; q_1)} \right|.\end{aligned}$$

An application of Lemma 5.4 gives

$$\Sigma_1 \ll_n D_K^{\frac{1}{n} - \delta_2, K} q_2, \quad \Sigma_{2A} \ll_n D_K^{\frac{1}{n} - \delta_2, K} q_1,$$

provided that $\mathcal{U}, \mathcal{V} \subset \{p : F(X, Y) \pmod p \text{ is not a square}\}$. Thus, it remains to estimate Σ_{2B} . To do this, we first denote $g_h(x_2) = g_2(x_2) g_2(x_2 + q_2 h)$. Applying the Poisson summation formula, one gets

$$\Sigma_{2B} \ll \frac{1}{q_1} \sum_{h \in [-H, H]} \left| \sum_{u_2 \in \mathbb{Z}} \hat{g}_h\left(\frac{u_2}{q_1}\right) W(u_1 \overline{q_2}, u_2, q_2 h; q_1) \right|$$

where

$$W(s, t, r; q_1) = \sum_{x, y, w \pmod{q_1}} \left(\frac{F_{\mathbf{z}}(x, y)}{q_1} \right) \left(\frac{F_{\mathbf{z}}(w, y + r)}{q_1} \right) e_p(s(x - w) + ty)$$

Applying Proposition 2.4, we have

Corollary 4.13. *Let $\mathbf{z} = (z_3, \dots, z_n)$, and $q_1 = uu'$ with u, u' primes such that for every $\sigma \in \text{Emb}(K/\mathbb{Q})$ with $\sigma \neq \tau$:*

- i) *if $\sigma(\alpha_2) \neq \alpha_2$, then $u, u' \nmid N_K(\sigma(\alpha_2) - \alpha_2)$,*
- ii) *if $\sigma \neq \text{Id}$, then $u, u' \nmid \sigma(z) - z$,*
- iii) *if $\tau(\alpha_{2, K}) \neq \alpha_2$, then*

$$u, u' \nmid N_K \left(\left(\sigma(z) - z - \frac{\sigma(\alpha_{2, K}) - \alpha_{2, K}}{\tau(\alpha_{2, K}) - \alpha_{2, K}} \cdot (\tau(z) - z) \right) \right)$$

Then, for any $s \in \mathbb{Z}/q_1\mathbb{Z}$, there exists $A_{s, u} \subset \{(r, t) \in (\mathbb{Z}/u\mathbb{Z}) \times (\mathbb{Z}/u\mathbb{Z})\}$, and $A_{s, u'} \subset \{(r, t) \in (\mathbb{Z}/u'\mathbb{Z}) \times (\mathbb{Z}/u'\mathbb{Z})\}$, such that

a) for any (r, t) such that $(r, t) \bmod u \notin A_{s,u}$, and $(r, t) \bmod u \notin A_{s,u'}$, one has

$$\left| \sum_{x,y,w \pmod{q_1}} \left(\frac{F_{\mathbf{z}}(X_1, X_2)}{q_1} \right) \left(\frac{F_{\mathbf{z}}(w, y+r)}{q_1} \right) e_{q_1}(s(x-w) + ty) \right| \ll_n q_1^{3/2}(s, t, q_1)^{\frac{1}{2}}.$$

b) $\#A_{s,u} = O_n(1)$, and $\#A_{s,u'} = O_n(1)$.

We are ready to bound the contribution of Σ_{2B} . First we recall that $q_1 = uu'$ with $u, u' \in \mathcal{U}$. From now on we set

$$\mathcal{U} = \left\{ u : \begin{array}{l} U \leq u \leq 2U \\ u \nmid N_K(\sigma(\alpha_2) - \alpha_2), \forall \sigma : \sigma(\alpha_2) \neq \alpha_2 \\ u \nmid N_K(\sigma(z) - z), \forall \sigma \in \text{Emb}(K/\mathbb{Q}) \setminus \{\text{Id}\} \\ u \nmid N_K \left(\left(\sigma(z) - z - \frac{\sigma(\alpha_{2,K}) - \alpha_{2,K}}{\tau(\alpha_{2,K}) - \alpha_{2,K}} \cdot (\tau(z) - z) \right) \right), \forall \sigma, \tau \in \text{Emb}(K/\mathbb{Q}) : \tau(\alpha_{2,K}) \neq \alpha_2 \end{array} \right\},$$

Moreover, we assume that $H \leq U$: in this way, it is guaranteed that $h \neq 0$, implies that $h \neq 0 \pmod{u}$ for every $U \leq u$. Since the bound for $W(u_1 \overline{q_2}, u_2, q_2 h; q_1)$ in Corollary 4.13 depends on the G.C.D. of $q_1, u_1 \overline{q_2}$, and $u_2 \overline{q_2}$. We have to handle two cases

i) the case when $(u_1, q_1) = 1$. We start by applying Corollary 4.13, getting

$$\begin{aligned} \Sigma_{2B} &\ll \frac{1}{q_1} \sum_{\substack{h \in [-H, H] \\ h \neq 0}} \left| \sum_{\substack{u_2 \in \mathbb{Z} \\ (q_2 h, u_2) \notin A_{u_1 \overline{q_2}, u} \\ (q_2 h, u_2) \notin A_{u_1 \overline{q_2}, u'}}} \hat{g}_h \left(\frac{u_2}{q_1} \right) W(u_1 \overline{q_2}, u_2, q_2 h; q_1) \right| \\ &+ \frac{1}{q_1} \sum_{\substack{h \in [-H, H] \\ h \neq 0}} \left| \sum_{\substack{u_2 \in \mathbb{Z} \\ (q_2 h, u_2) \in A_{u_1 \overline{q_2}, u}}} \hat{g}_h \left(\frac{u_2}{q_1} \right) W(u_1 \overline{q_2}, u_2, q_2 h; q_1) \right| \\ &+ \frac{1}{q_1} \sum_{\substack{h \in [-H, H] \\ h \neq 0}} \left| \sum_{\substack{u_2 \in \mathbb{Z} \\ (q_2 h, u_2) \in A_{u_1 \overline{q_2}, u'}}} \hat{g}_h \left(\frac{u_2}{q_1} \right) W(u_1 \overline{q_2}, u_2, q_2 h; q_1) \right| \\ &\ll_n H q_1^{\frac{3}{2}} + q_1^2 \cdot \frac{1}{q_1} \sum_{\substack{h \in [-H, H] \\ h \neq 0}} \sum_{\substack{u_2 \in \mathbb{Z} \\ (q_2 h, u_2) \in A_{u_1 \overline{q_2}, u}}} \left| \hat{g}_h \left(\frac{u_2}{q_1} \right) \right| \\ &+ q_1^2 \cdot \frac{1}{q_1} \sum_{\substack{h \in [-H, H] \\ h \neq 0}} \sum_{\substack{u_2 \in \mathbb{Z} \\ (q_2 h, u_2) \in A_{u_1 \overline{q_2}, u'}}} \left| \hat{g}_h \left(\frac{u_2}{q_1} \right) \right|, \end{aligned}$$

Provided that we have $q_1 \geq D_K^{\frac{1}{n}-\delta_{2,K}}$; now we have

$$\begin{aligned} \sum_{\substack{h \in [-H, H] \\ h \neq 0}} \sum_{\substack{u_2 \in \mathbb{Z} \\ (q_2 h, u_2) \in A_{u_1 \overline{q_2}, u}}} \left| \hat{g}_h \left(\frac{u_2}{q_1} \right) \right| &\leq \sum_{(r, t) \in A_{u_1 \overline{q_2}, u}} \sum_{\substack{h \in [-H, H] \\ h \neq 0 \\ q_2 h = r(u)}} \sum_{\substack{u_2 \in \mathbb{Z} \\ u_2 = t(u)}} \left| \hat{g}_h \left(\frac{u_2}{q_1} \right) \right| \\ &\ll D_K^{\frac{1}{n}-\delta_{2,K}} \sum_{(r, t) \in A_{u_1 \overline{q_2}, u}} \sum_{\substack{h \in [-H, H] \\ q_2 h = r(u)}} \sum_{\substack{u_2 \in \mathbb{Z} \\ u_2 = t(u)}} \left| 1 + \frac{u_2 D_K^{\frac{1}{n}-\delta_{2,K}}}{q_1} \right|^{-2} \\ &\ll_n D_K^{\frac{1}{n}-\delta_{2,K}}, \end{aligned}$$

provided that $\frac{q_1}{u} \leq D_K^{\frac{1}{n}-\delta_{2,K}}$, i.e. $q_1^{\frac{1}{2}} \ll D_K^{\frac{1}{n}-\delta_{2,K}}$. Moreover, recalling that $H = \lfloor \frac{D_K^{\frac{1}{n}-\delta_{2,K}}}{q_2} \rfloor$, we conclude that

$$\begin{aligned} q_1^2 \cdot \frac{1}{q_1} \sum_{h \in [-H, H]} \sum_{\substack{u_2 \in \mathbb{Z} \\ (q_2 h, u_2) \in A_{u_1 \overline{q_2}, u}}} \left| \hat{g}_h \left(\frac{u_2}{q_1} \right) \right| &\ll_n D_K^{\frac{1}{n}-\delta_{2,K}} q_1 \\ &\leq H q_2 q_1 \\ &\leq H q_1^{\frac{3}{2}}, \end{aligned}$$

provided that $\max\{q_2^2, D_K^{\frac{1}{n}-\delta_{2,K}}\} \leq q_1 \leq D_K^{\frac{2}{n}-2\delta_{2,K}}$ (notice that $\max\{q_2^2, D_K^{\frac{1}{n}-\delta_{2,K}}\} \leq q_1$ guarantees that $H \leq q_1$). Hence, in this case, we have

$$\Sigma_{2B} \ll_n H q_1^{\frac{3}{2}},$$

provided that $\max\{q_2^2, D_K^{\frac{1}{n}-\delta_{2,K}}\} \leq q_1 \leq D_K^{\frac{2}{n}-2\delta_{2,K}}$.

ii) The case $(u_1, q_1) > 1$. We start by writing

$$\begin{aligned} \Sigma_{2B} &\ll \frac{1}{q_1} \sum_{\substack{h \in [-H, H] \\ h \neq 0}} \left| \sum_{\substack{u_2 \in \mathbb{Z} \\ (u_2, q_1) = 1}} \hat{g}_h \left(\frac{u_2}{q_1} \right) W(u_1 \overline{q_2}, u_2, q_2 h; q_1) \right| \\ &\quad + \frac{1}{q_1} \sum_{\substack{h \in [-H, H] \\ h \neq 0}} \left| \sum_{\substack{u_2 \in \mathbb{Z} \\ (u_2, q_1) > 1}} \hat{g}_h \left(\frac{u_2}{q_1} \right) W(u_1 \overline{q_2}, u_2, q_2 h; q_1) \right| \end{aligned} \tag{17}$$

observe that the contribution of the u_2 coprime q_1 can be bounded as in (i), getting

$$\frac{1}{q_1} \sum_{\substack{h \in [-H, H] \\ h \neq 0}} \left| \sum_{\substack{u_2 \in \mathbb{Z} \\ (u_2, q_1) = 1}} \hat{g}_h \left(\frac{u_2}{q_1} \right) W(u_1 \overline{q_2}, u_2, q_2 h; q_1) \right| \ll_n H q_1^{\frac{3}{2}},$$

provided that $\max\{q_2^2, D_K^{\frac{1}{n}-\delta_{2,K}}\} \leq q_1 \leq D_K^{\frac{2}{n}-2\delta_{2,K}}$. For the contribution of the second sum appearing on the right-hand side of (17), we are going to use the trivial bound

$\ll q_1^2$ for the exponential sum (see Lemma 5.7), getting

$$\begin{aligned} & \frac{1}{q_1} \sum_{\substack{h \in [-H, H] \\ h \neq 0}} \left| \sum_{\substack{u_2 \in \mathbb{Z} \\ (u_2, q_1) > 1}} \hat{g}_h \left(\frac{u_2}{q_1} \right) W(u_1 \bar{q}_2, u_2, q_2 h; q_1) \right| \\ & \ll_n q_1 \sum_{\substack{h \in [-H, H] \\ h \neq 0}} \sum_{\substack{u_2 \in \mathbb{Z} \\ (u_2, q_1) > 1}} \left| \hat{g}_h \left(\frac{u_2}{q_1} \right) \right| \\ & \ll H \cdot q_1 \cdot D_K^{\frac{1}{n} - \delta_{2,K}}, \end{aligned}$$

provided that $q_1^{\frac{1}{2}} \leq D_K^{\frac{1}{n} - \delta_{2,K}}$. Thus

$$\frac{1}{q_1} \sum_{h \in [-H, H]} \left| \sum_{\substack{u_2 \in \mathbb{Z} \\ (u_2, q_1) > 1}} \hat{g}_h \left(\frac{u_2}{q_1} \right) W(u_1 \bar{q}_2, u_2, q_2 h; q_1) \right| \ll_n H D_K^{\frac{1}{n} - \delta_{2,K}} q_1,$$

provided that $\max\{q_2^2, D_K^{\frac{1}{n} - \delta_{2,K}}\} \leq q_1 \leq D_K^{\frac{2}{n} - 2\delta_{2,K}}$.

Combining the estimates for Σ_1, Σ_{2A} , and Σ_{2B} , we conclude

Lemma 4.14. *Let $\mathbf{z} \in \mathbb{Z}^{n-2} \setminus \{\mathbf{0}\}$ satisfying the following condition:*

i) For every $\sigma, \tau \in \text{Emb}(K/\mathbb{Q})$, with $\sigma \neq \tau$ and $\tau(\alpha_{2,K}) \neq \alpha_{2,K}$,

$$\sigma(z) - z \neq \frac{\sigma(\alpha_{2,K}) - \alpha_{2,K}}{\tau(\alpha_{2,K}) - \alpha_{2,K}} (\tau(z) - z),$$

ii) For every $\sigma \in \text{Emb}(K/\mathbb{Q}) \setminus \{\text{Id}\}$, $z \notin K^\sigma$.

Then, provided that $H < q_1$ and $q_2^2 = q_1$, one has

$$S(\mathbf{z}, u_1; q_1 q_2) \ll_n \begin{cases} D_K^{\frac{1}{2n} - \frac{\delta_{2,K}}{2}} q_1 & \text{if } (u_1, q_1) = 1 \\ \frac{D_K^{\frac{1}{n} - \delta_{2,K}} q_1}{q_2^{\frac{1}{2}}} & \text{otherwise.} \end{cases}$$

provided that $\max\{q_2^2, D_K^{\frac{1}{n} - \delta_{2,K}}\} \leq q_1 \leq D_K^{\frac{2}{n} - 2\delta_{2,K}}$, and $D_K^{\frac{1}{2n} - \delta_{2,K}} \leq q_2 q_1^{\frac{1}{2}}$

Then we can finally bound the main sieve. From now on we set

$$\mathcal{V} = \left\{ v : \begin{array}{l} V \leq v \leq 2V \\ F_{\mathbf{z}}(X_1, X_2) \pmod{v} \text{ is not a square in } \mathbb{F}_v \\ v \nmid N_K(\sigma(\alpha_2) - \alpha_2) \text{ for all } \sigma \text{ such that } \sigma(\alpha_2) \neq \alpha_2 \end{array} \right\},$$

Moreover, we require $D_K^{\frac{1}{2n} - \frac{\delta_{2,K}}{2}} \leq U \leq D_K^{\frac{1}{n} - \delta_{2,K}}$ and $U = V^2$, and $UV^2 \leq D_K^{\frac{1}{n}} \leq (UV)^2$. We have

$$S(\mathbf{z}; q_1 q_2) = \frac{1}{q_1 q_2} \sum_{\substack{u_1 \in \mathbb{Z} \\ (u_1, q_1) = 1}} \hat{g}_1 \left(\frac{u_1}{q_1 q_2} \right) S(\mathbf{z}, u_1; q_1 q_2) + \frac{1}{q_1 q_2} \sum_{\substack{u_1 \in \mathbb{Z} \\ (u_1, q_1) > 1}} \hat{g}_1 \left(\frac{u_1}{q_1 q_2} \right) S(\mathbf{z}, u_1; q_1 q_2)$$

We start by dealing with the contribution of those u_1 such that $(u_1, q_1) > 1$. An application of Lemma 4.14 combined with our assumption that $UV^2 \leq D_K^{\frac{1}{n}}$, gives

$$\begin{aligned} \frac{1}{q_1 q_2} \sum_{\substack{u_1 \in \mathbb{Z} \\ (u_1, q_1) > 1}} \hat{g}_1 \left(\frac{u_1}{q_1 q_2} \right) S(\mathbf{z}, u_1; q_1 q_2) &\ll_n \frac{D_K^{\frac{2}{n} - \delta_{2,K}}}{q_2^{\frac{3}{2}}} \sum_{\substack{u_1 \in \mathbb{Z} \\ (u_1, q_1) > 1}} \left| \hat{g}_1 \left(\frac{u_1}{q_1 q_2} \right) \right| \\ &\ll \frac{D_K^{\frac{2}{n} - \delta_{2,K}}}{UV} \end{aligned}$$

Let us turn our attention those u_1 such that $(u_1, q_1) = 1$. Applying Lemma 4.14, combined with our assumption that $(UV)^2 \geq D_K^{\frac{1}{n}}$, gives

$$\frac{1}{q_1 q_2} \sum_{\substack{u_1 \in \mathbb{Z} \\ (u_1, q_1) = 1}} \hat{g}_1 \left(\frac{u_1}{q_1 q_2} \right) S(\mathbf{z}, u_1; q_1 q_2) \ll_n D_K^{\frac{1}{2n} - \frac{\delta_{2,K}}{2}} U^2$$

Combining the two bounds, we get

$$S(\mathbf{z}; q_1 q_2) \ll_n D_K^{\frac{1}{2n} - \frac{\delta_{2,K}}{2}} U^2 + \frac{D_K^{\frac{2}{n} - \delta_{2,K}}}{UV}$$

Inserting this bound to the main sieve (16), we get

$$\frac{1}{A^2} \sum_{v, v' \in \mathcal{V}} \sum_{\substack{u, u' \in \mathcal{U} \\ u \neq u'}} \left| \sum_{n \in \mathbb{Z}} \left(\frac{n}{uu'} \right) \left(\frac{n}{vv'} \right) \omega_{\mathbf{z}}(n) \right| \ll_n D_K^{\frac{1}{2n} - \frac{\delta_{2,K}}{2}} U^2 + \frac{D_K^{\frac{2}{n} - \delta_{2,K}}}{UV}. \quad (18)$$

4.4.1. *Proof of Lemma 4.12: the prime sieve.* We briefly discuss now how to handle the Prime sieve

$$N = \frac{U}{A^2} \sum_{\substack{v, v' \in \mathcal{V} \\ v \neq v'}} \left| \sum_{n \in \mathbb{Z}} \left(\frac{n}{vv'} \right) \omega_{\mathbf{z}}(n) \right|.$$

Note that the inner sum in the right-hand side is the same as the one appearing in the second term of the right-hand side of (10). Since we have

$$\mathcal{V} = \left\{ v : \begin{array}{l} V \leq v \leq 2V \\ F_{\mathbf{z}}(X_1, X_2) \pmod{v} \text{ is not a square in } \mathbb{F}_v \\ v \nmid N_K(\sigma(\alpha_2) - \alpha_2) \text{ for all } \sigma \text{ such that } \sigma(\alpha_2) \neq \alpha_2 \end{array} \right\},$$

arguing as in Lemma 4.10, one gets

$$\left| \sum_{n \in \mathbb{Z}} \left(\frac{n}{vv'} \right) \omega_{\mathbf{z}}(n) \right| \ll_n \frac{D_K^{\frac{2}{n} - \delta_{2,K}}}{vv'} \sum_{\mathbf{u} \in \mathbb{Z}^2} \left(1 + \frac{D_K^{\frac{1}{n}} u_1}{vv'} \right)^{-2} \cdot \left(1 + \frac{D_K^{\frac{1}{n} - \delta_{2,K}} u_2}{vv'} \right)^{-2}. \quad (19)$$

Since $v'v \leq 4V^2 \ll D_K^{\frac{1}{n} - \delta_{2,K}}$, the inner sum in (19) is a $O(1)$. Thus

$$\left| \sum_{n \in \mathbb{Z}} \left(\frac{n}{vv'} \right) \omega_{\mathbf{z}}(n) \right| \ll_n \frac{D_K^{\frac{2}{n} - \delta_{2,K}}}{V^2}$$

Hence,

$$\frac{U}{A^2} \sum_{\substack{v, v' \in \mathcal{V} \\ v \neq v'}} \left| \sum_{n \in \mathbb{Z}} \left(\frac{n}{vv'} \right) \omega_{\mathbf{z}}(n) \right| \ll_n \frac{D_K^{\frac{2}{n} - \delta_{2,K}}}{UV^2}, \quad (20)$$

which is acceptable.

4.4.2. *Proof of Lemma 4.12: conclusion.* Combining (14) with (18), and (15), we obtain

$$\sum_k \omega_{\mathbf{z}}(k^2) \ll_n \frac{D_K^{\frac{2}{n} - \delta_{2,K}}}{UV} + D_K^{\frac{1}{2n} - \frac{\delta_{2,K}}{2}} U^2.$$

Imposing $U = V^2$, the bound above is optimal as soon as

$$\frac{D_K^{\frac{2}{n} - \delta_{2,K}}}{V^3} = D_K^{\frac{1}{2n} - \frac{\delta_{2,K}}{2}} V^4$$

which implies

$$V = D_K^{\frac{1}{7n} + \frac{1}{14}(\frac{1}{n} - \delta_{2,K})}$$

and concludes the proof of Lemma 4.12.

4.4.3. *Proof of Theorem 2.3.* We can now prove Theorem 2.3. Again, thanks to Lemma 4.1, we may assume that K does not contain a subfield of index 2. We recall that

$$\mathcal{C} = \left[-D_K^{\frac{1}{n} - \delta_{3,K}}, D_K^{\frac{1}{n} - \delta_{3,K}} \right] \times \cdots \times \left[-D_K^{\frac{1}{n} - \delta_{n,K}}, D_K^{\frac{1}{n} - \delta_{n,K}} \right],$$

and that

$$Z_{\square} = \{\mathbf{z} \in \mathbb{Z}^{n-2} : F_{\mathbf{z}}(X_1, X_2) \text{ is a square}\};$$

$$Z_F = \{\mathbf{z} \in \mathbb{Z}^{n-2} : F_{\mathbf{z}}(X_1, X_2) \text{ does not satisfy the hypothesis of Proposition 2.4}\}$$

We start by writing

$$\begin{aligned} h_2(K) &\ll \# \left\{ (y, z_1, \dots, z_n) \in \mathbb{Z}^{n+1} : \begin{array}{l} |z_i| \ll D_K^{\frac{1}{n} - \delta_{i,K}}, \text{ for } i = 1, \dots, n \\ y^2 = N_K(z_1 + z_2 \alpha_{2,K} + \cdots + z_n \alpha_{n,K}) \end{array} \right\} \\ &= \sum_{\mathbf{z} \in \mathcal{C} \cap \mathbb{Z}^{n-2}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 \\ &= \sum_{\mathbf{z} \in \mathcal{C} \cap Z_{\square}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 + \sum_{\substack{\mathbf{z} \in \mathcal{C} \\ \mathbf{z} \in Z_F \setminus Z_{\square}}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 + \sum_{\substack{\mathbf{z} \in \mathcal{C} \cap \mathbb{Z}^{n-2} \\ \mathbf{z} \notin Z_{\square} \cup Z_F}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1. \end{aligned} \quad (21)$$

For the first two terms we have the bounds

$$\sum_{\mathbf{z} \in \mathcal{C} \cap Z_{\square}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 = \begin{cases} 0 & \text{if } n = 5, 6 \\ O_{n,\varepsilon} \left(D_K^{\frac{1}{4} + \frac{1}{n}} \right) & \text{if } n > 6 \end{cases}, \quad \sum_{\substack{\mathbf{z} \in \mathcal{C} \\ \mathbf{z} \in Z_F \setminus Z_{\square}}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 \ll_{n,\varepsilon} D_K^{\frac{1}{2} - \frac{1}{2n} - \frac{1}{7n}},$$

thanks to Corollary 4.2 and Corollary 4.6. For the third term in (21), we can use Lemma 4.12, getting

$$\begin{aligned}
\sum_{\substack{\mathbf{z} \in \mathcal{C} \cap \mathbb{Z}^{n-2} \\ \mathbf{z} \notin Z_{\square} \cup Z_F}} \sum_{\substack{(x_1, x_2) \in \mathcal{B} \cap \mathbb{Z}^2 \\ F_{\mathbf{z}}(x_1, x_2) = \square}} 1 &\ll_{n, \varepsilon} \#(\mathcal{C} \cap \mathbb{Z}^{n-2}) \cdot D_K^{\frac{45}{28n} - \frac{11}{14} \delta_{2, K}} \\
&\ll D_K^{\sum_{j=3}^n \left(\frac{1}{n} - \delta_{j, K} \right) + \frac{1}{2} - \frac{1}{2n} - \left(\frac{1}{7n} - \frac{3\delta_{2, K}}{14} \right) + \varepsilon} \\
&= D_K^{\frac{1}{2} - \frac{1}{2n} - \left(\frac{1}{7n} - \frac{3\delta_{2, K}}{14} \right) + \varepsilon}.
\end{aligned}$$

which completes the proof of Theorem 2.3.

5. THE RELEVANT EXPONENTIAL SUMS.

This section is dedicated to prove the key exponential sums used in proof of Lemma 4.9 and Proposition 2.4.

5.1. Notions of the formalism of ℓ -adic trace functions. We first recall some notions of the formalism of ℓ -adic trace functions. We are assuming the definition of a constructible ℓ -adic sheaf, for which, as well as for a general introduction on the subject, we refer the reader to [Del80, Section 1.1] and [Kat80, Section 3.4.2]. We start by introducing the notion of a trace function attached to a constructible ℓ -adic sheaf as in [Kat90, 7.3.7].

In the following $p, \ell > 2$ and $q = p^k$ for some $k \geq 1$ are distinct prime numbers and $\iota : \overline{\mathbb{Q}}_{\ell} \hookrightarrow \mathbb{C}$ is a fixed embedding. Let \mathcal{F} be a constructible ℓ -adic sheaf on $\overline{\mathbb{A}}_{\mathbb{F}_q}^1$. For any $x \in \overline{\mathbb{A}}_{\mathbb{F}_q}^1(\mathbb{F}_{q^n})$ one defines

$$t_{\mathcal{F}, n}(x) := \iota(\text{Tr}(\text{Fr}_{q^n} | \mathcal{F}_{\bar{x}})),$$

where Fr_{q^n} is the geometric Frobenius automorphism of \mathbb{F}_{q^n} and $\mathcal{F}_{\bar{x}}$ is the stalk of \mathcal{F} at a geometric point \bar{x} over x . The function $t_{\mathcal{F}, n}$ is called *the trace function attached to \mathcal{F} over \mathbb{F}_{q^n}* . If there is not ambiguity, we denote by $t_{\mathcal{F}}$ the trace function $t_{\mathcal{F}, 1}$.

Definition 5.1. Let \mathcal{F} be a constructible ℓ -adic sheaf on $\overline{\mathbb{A}}_{\mathbb{F}_q}^1$ and $j : U \hookrightarrow \overline{\mathbb{A}}_{\mathbb{F}_q}^1$ the largest dense open subset of $\overline{\mathbb{A}}_{\mathbb{F}_q}^1$ where \mathcal{F} is lisse.

- i) The sheaf \mathcal{F} is said to be a *middle-extension ℓ -adic sheaf* if $\mathcal{F} = j_* j^* \mathcal{F}$ (see [Kat80, 4.4, 4.5] for the definition of $j_* j^* \mathcal{F}$ and its basic properties).
- ii) The sheaf \mathcal{F} is said to be a *middle-extension ℓ -adic sheaf, punctually pure of weight 0* if it is a middle-extension sheaf and if for every $n \geq 1$ and every $x \in U(\mathbb{F}_{q^n})$, the images of the eigenvalues of $(\text{Fr}_{q^n} | \mathcal{F}_{\bar{x}})$ via the fixed embedding $\iota : \overline{\mathbb{Q}}_{\ell} \hookrightarrow \mathbb{C}$, are complex numbers of modulus 1.

Remark 1. By [FKM15, Definition 1.12], we will refer to trace functions attached to punctually pure of weight 0 middle-extension ℓ -adic sheaves on $\overline{\mathbb{A}}_{\mathbb{F}_q}^1$ as trace functions.

Definition 5.2 ([FKM19], pp. 4 – 6). Let \mathcal{F} be a constructible ℓ -adic sheaf on $\overline{\mathbb{A}}_{\mathbb{F}_q}^1$ and $j : U \hookrightarrow \overline{\mathbb{A}}_{\mathbb{F}_q}^1$ the largest dense open subset of $\overline{\mathbb{A}}_{\mathbb{F}_q}^1$ where \mathcal{F} is lisse. The *conductor of \mathcal{F}* is defined as

$$c(\mathcal{F}) := \text{rank}(\mathcal{F}) + |\text{sing}(\mathcal{F})| + \sum_{x \in \overline{\mathbb{P}}_{\mathbb{F}_q}^1(\mathbb{F}_q)} \text{Swan}_x(j_* j^* \mathcal{F}) + \dim H_c^0(\overline{\mathbb{A}}_{\mathbb{F}_q}^1, \mathcal{F}),$$

where

- i) $\text{rank}(\mathcal{F}) := \dim \mathcal{F}_x$, for any x where \mathcal{F} is lisse.
- ii) $\text{sing}(\mathcal{F}) := \{x \in \overline{\mathbb{P}}_{\mathbb{F}_q}^1(\overline{\mathbb{F}}_q) : \mathcal{F} \text{ is not lisse at } x\}$.
- iii) For any $x \in \overline{\mathbb{P}}_{\mathbb{F}_q}^1(\overline{\mathbb{F}}_q)$, $\text{Swan}_x(j_*j^*\mathcal{F})$ is the *Swan conductor of \mathcal{F} at x* (see [Kat88, Chapter 1] for the definition of the Swan conductor).

Remark 2. Recall that if \mathcal{F} is a middle-extension sheaf then

$$c(\mathcal{F}) = \text{rank}(\mathcal{F}) + |\text{sing}(\mathcal{F})| + \sum_{x \in \overline{\mathbb{P}}_{\mathbb{F}_q}^1(\overline{\mathbb{F}}_q)} \text{Swan}_x(\mathcal{F}),$$

since in this case $\mathcal{F} \cong j_*j^*\mathcal{F}$ and $\dim H_c^0(\overline{\mathbb{A}}_{\mathbb{F}_q}^1, \mathcal{F}) = 0$.

Finally, we recall the following construction:

Example 5.3. let $F[X, Y] \in \mathbb{F}_q[X, Y]$ be a square-free polynomial of degree n . Then for any non trivial multiplicative character over \mathbb{F}_q^\times , χ , one can consider the following $t : \mathbb{F}_q \rightarrow \mathbb{C}$

$$x \mapsto \sum_{y \in \mathbb{F}_q} \chi(F(x, y)).$$

It turns out that we can see this function as a trace function: combining [FKM19, Definition 5.9, Theorem 5.10] and arguing like in [FKM19, Corollary 5.10] one shows that there exists a middle extension ℓ -adic sheaf $\mathcal{H}_{F, \chi}$ pure of weight 1 such that

$$t_{\mathcal{H}_{F, \chi}, k}(x) = \sum_{y \in \mathbb{F}_{q^k}} \chi_k(F(x, y)) + O(1) \quad \text{for all } x \in \mathbb{F}_{q^k} \setminus (\text{sing}(\mathcal{H}_{F, \chi}) \cap \mathbb{F}_{q^k}),$$

where $\chi_k = \chi \circ \text{Norm}_{\mathbb{F}_{q^k}/\mathbb{F}_q}$. Moreover, one has that

- i) $\text{sing}(\mathcal{H}_{F, \chi}) \cap \overline{\mathbb{A}}_{\mathbb{F}_q}^1(\overline{\mathbb{F}}_q) = \{y \in \overline{\mathbb{A}}_{\mathbb{F}_q}^1(\overline{\mathbb{F}}_q) : F(X, y) \text{ is not square-free}\}$
- ii) $\text{rank}(\mathcal{H}_{F, \chi}) = \dim(H^1(\overline{\mathbb{A}}_{\mathbb{F}_q}^1, \mathcal{L}_{\chi(F(X, Y))}))$ for any x where \mathcal{H} is lisse.
- iii) $C(\mathcal{H}_{F, \chi}) \ll_n 1$

5.1.1. *A 1-dimensional exponential sum.* We conclude this section by recording the following Lemma about one dimensional exponential sums

Lemma 5.4. Let $F[X, Y] \in \mathbb{F}_q[X, Y]$ be a square free polynomial. Then

$$\left| \sum_{x \in \mathbb{F}_p} \left(\frac{F(x, y)}{p} \right) e_p(sx) \right| \ll_{\deg F} p^{\frac{1}{2}},$$

for all but $O_{\deg F}(1)$ values of y .

Proof. If $s \neq 0$, then the result follows from [We48]. If $s = 0$, then from [IK04, Theorem 11.23], we have that

$$\left| \sum_{x \in \mathbb{F}_p} \left(\frac{F(x, y)}{p} \right) \right| \ll_{\deg F} p^{\frac{1}{2}},$$

for every y such that $F(X, y) \in \mathbb{F}_p[X]$ is not a square. it rests to show that $F(X, y)$ can be a square only for $O_{\deg F}(1)$ values of y : since we are assuming $F(X, Y) \in \mathbb{F}_q(Y)[X]$ is square free it follows that $\text{disc}(F) \in \mathbb{F}[Y] \neq 0$. Moreover,

$$F(X, y) \text{ is square-free} \iff \text{disc}(F)(y) \neq 0.$$

Thus $F(X, y)$ is square-free for but $O_{\deg F}(1)$ values of y . \square

5.2. Hooley's Theorem for exponential sums. Both the proofs of Proposition 2.4 and Lemma 4.9 rely on a general procedure due to Hooley [Ho82] (see also ([BB23, Theorem A.1])), which allows one to estimate a very general family of exponential sums over a finite field, provided that one can control the seconds moment of an appropriate counting function.

Theorem 5.5 (Hooley). *Let $F, G_1, \dots, G_k \in \mathbb{Z}[X_1, \dots, X_m]$ be polynomials of degree at most d and let*

$$S = \sum_{\substack{x \in \mathbb{F}_p^m \\ G_1(\mathbf{x}) = \dots = G_k(\mathbf{x}) = 0}} e_p(F(\mathbf{x})).$$

For each $r \geq 1$ and $\tau \in \mathbb{F}_{p^r}$, write

$$N_r(\tau) = \# \{ \mathbf{x} \in \mathbb{F}_{p^r}^m : G_1(\mathbf{x}) = \dots = G_k(\mathbf{x}) = 0, F(\mathbf{x}) = \tau \}. \quad (22)$$

If there exist $N_r \in \mathbb{R}$ such that

$$\sum_{\tau \in \mathbb{F}_{p^r}} |N_r(\tau) - N_r|^2 \ll_{d,k,m} p^{\kappa r},$$

where $\kappa \in \mathbb{Z}$ is independent of r , then $S \ll_{d,k,m} p^{\kappa/2}$.

5.3. Proof of Lemma 4.9. We are now ready to prove Lemma 4.9 which is a direct consequence of:

Lemma 5.6. *Let $F[X, Y] \in \mathbb{F}_q[X, Y]$ be a square-free polynomial such that*

$$F[X, Y] = \prod_{i=1}^n (X + a_i Y + b_i) \text{ over } \overline{\mathbb{F}}_q.$$

Moreover assume that $a_i \neq a_1$ for some $i \in \{2, \dots, n\}$. Then for any $s, t \in \mathbb{F}_p$, we have

$$\left| \sum_{x,y \in \mathbb{F}_p} \left(\frac{F(x, y)}{p} \right) e_p(sx + ty) \right| \ll_n p(p, r, s)^{\frac{1}{2}}.$$

Proof. We start by dealing with the case when $r = s = 0$: we have

$$\sum_{x,y \in \mathbb{F}_p} \left(\frac{F(x, y)}{p} \right) = \sum_{y \in \mathbb{F}_p} \sum_{x \in \mathbb{F}_p} \left(\frac{F(x, y)}{p} \right).$$

Thanks to Lemma 5.4, we get

$$\left| \sum_{x \in \mathbb{F}_p} \left(\frac{F(x, y)}{p} \right) \right| \ll_n \sqrt{p}, \quad (23)$$

for all but $O_n(1)$ values of y . Let us denote by S the set of those y for which the bound (23) does not hold. Hence,

$$\sum_{y \in \mathbb{F}_p} \sum_{x \in \mathbb{F}_p} \left(\frac{F(x, y)}{p} \right) = \sum_{y \in \mathbb{F}_p \setminus S} \sum_{x \in \mathbb{F}_p} \left(\frac{F(x, y)}{p} \right) + \sum_{y \in S} \sum_{x \in \mathbb{F}_p} \left(\frac{F(x, y)}{p} \right) \\ \ll_n p^{\frac{3}{2}}.$$

Now we deal with the case $(s, t) \neq (0, 0)$. We may assume $s \neq 0$ (the case $t \neq 0$ is similar). We start by observing that

$$\sum_{x, y \in \mathbb{F}_p} \left(\frac{F(x, y)}{p} \right) e_p(sx + ty) = \sum_{\substack{w, x, y \in \mathbb{F}_p \\ w^2 = F(x, y)}} e_p(sx + ty).$$

With the goal of using Theorem 5.5, we are going to compute

$$N_m(k) = \#\{(w, x, y) \in \mathbb{F}_{p^m}^3 : w^2 = F(x, y), sx + ty = k\} \\ = \#\{(w, x) \in \mathbb{F}_{p^m}^3 : w^2 = F(x, s'x + k')\},$$

for every $m \geq 1$, where $s' = -\bar{t}s$, and $k' = \bar{t}k$. We claim that $F(X, s'X + k')$ is square-free for all but $O_n(1)$ values of k' . With this in hand, let us see how we can finish the Lemma: if the polynomial $F(X, s'X + k')$ is square-free, then the polynomial $W^2 = F(X, s'X + k')$ is irreducible. Provided that $F(X, s'X + k')$ is not a square, it follows from the Lang-Weil bound that

$$\#\{(w, x) \in \mathbb{F}_{p^m}^3 : w^2 = F(x, s'x + k')\} = p^m + O_n(p^{\frac{m}{2}}).$$

Thus,

$$\sum_{k \in \mathbb{F}_{p^m}} |N_m(k) - p^m|^2 \ll_n p^{2m},$$

which leads to

$$\left| \sum_{x, y \in \mathbb{F}_p} \left(\frac{F(x, y)}{p} \right) e_p(sx + ty) \right| \ll_n p,$$

thanks to Theorem 5.5. It rests to prove that $F(X, s'X + k')$ is square-free for all but $O_n(1)$ values of k' . First observe that

$$F(X, s'X + k') = A_{s', k'} G_{s', k'}(X),$$

where

$$A_{s', k'} = \prod_{i: a_i s' = -1} (b_i + a_i k'), \quad G_{s', k'}(X) = \prod_{\substack{i=1 \\ a_i s' \neq -1}}^n ((1 + a_i s')X + a_i k' + b_i).$$

Since we are assuming that there exists i such that $a_i \neq a_1$, then $G_{s', k'}(X)$ is a non-constant polynomial. Moreover, $F(X, s'X + k')$ is square-free if and only if $G_{s', k'}(X)$ is. On the other hand, $G_{s', k'}(X)$ is not square-free if and only if two distinct factors

$$(1 + a_i s')X + a_i k' + b_i, \quad (1 + a_j s')X + a_j k' + b_j$$

are proportional. These two linear equations are proportional if and only if

$$\frac{a_i k' + b_i}{(1 + a_i s')} = \frac{a_j k' + b_j}{(1 + a_j s')}$$

which is true if and only if

$$k'(a_i - a_j) = b_j(1 + a_i s') - b_i(1 + a_j s').$$

If $a_i - a_j \neq 0$, the equation above is satisfied for a unique value of k' . If $a_i = a_j$, the equation above become $(b_j - b_i)(1 + a_i s') = 0$ which would implies $b_j - b_i = 0$ since every factor appearing in the decomposition of $G_{s',k'}(X)$ satisfies $1 + a_i s' \neq 0$. On the other hand, since we are assuming that $F(X, Y)$ is square-free, if two factors

$$X + a_i Y + b_i, \quad X + a_j Y + b_j,$$

are such that $a_i = a_j$, then necessarily one has $b_i - b_j \neq 0$. Hence, $G_{s',k'}(X)$ is square free for all but $O_n(1)$ values of k' as claimed. \square

Proof of Lemma 4.9. Since $F_{\mathbf{z}} = N_K(X_1 + X_2 \alpha_2 + z)$, with $z = z_3 \alpha_3 + \cdots + z_n \alpha_n$, it follows that

$$F_{\mathbf{z}}(X_1, X_2) = \prod_{\sigma \in \text{Emb}(K/\mathbb{Q})} (X_1 + X_2 \sigma(\alpha_2) + \sigma(z)).$$

Now let $F_{\mathbf{z}} = H_{\mathbf{z}}^2 G_{\mathbf{z}}$ with $H_{\mathbf{z}}, G_{\mathbf{z}} \in \mathbb{Q}[X, Y]$, monic and $G_{\mathbf{z}}$ square-free. Since

$$\sum_{x_1, x_2 \in \mathbb{F}_p} \left(\frac{F_{\mathbf{z}}(x_1, x_2)}{p} \right) e_p(sx_1 + tx_2) = \sum_{x_1, x_2 \in \mathbb{F}_p} \left(\frac{G_{\mathbf{z}}(x_1, x_2)}{p} \right) e_p(sx_1 + tx_2) + O_n(p),$$

it is enough to show that $G_{\mathbf{z}}(X_1, X_2)$ satisfies the hypothesis of Lemma 5.6. Since we are assuming $F_{\mathbf{z}} \bmod p$ is not a square, it follows that

$$G_{\mathbf{z}}(X_1, X_2) = \prod_{\sigma \in \mathcal{E}} (X_1 + X_2 \sigma(\alpha_2) + \sigma(z)),$$

for some $\emptyset \neq \mathcal{E} \subset \text{Emb}(K/\mathbb{Q})$. Let $\sigma \in \mathcal{E}$. Since $\sigma(\alpha_2) \notin \mathbb{Q}$, we can find $\tau \in \text{Emb}(K/\mathbb{Q})$ such that $\tau(\sigma(\alpha_2)) \neq \sigma(\alpha_2)$. On the other hand,

$$\tau((X_1 + X_2 \sigma(\alpha_2) + \sigma(z)) | \tau(G_{\mathbf{z}}) = G_{\mathbf{z}}.$$

Hence, $G_{\mathbf{z}}$ satisfies the hypothesis of lemma 5.6, since it has two linear factors

$$X_1 + X_2 \sigma(\alpha_2) + \sigma(z), \quad X_1 + X_2 \tau \circ \sigma(\alpha_2) + \tau \circ \sigma(z),$$

such that $\sigma(\alpha_2) \neq \tau \circ \sigma(\alpha_2)$. \square

5.4. Proof of Proposition 2.4. This section is devoted to bounding the exponential sum

$$\sum_{(x, y, z) \in \mathbb{F}_p^3} \left(\frac{F(x, y)}{p} \right) \left(\frac{F(z, y + r)}{p} \right) e_p(s(x - z) + ty)$$

for a given square-free polynomial $F(X, Y)$. We begin by establishing the following general estimate:

Lemma 5.7. *Let $F \in \mathbb{F}_p[X, Y]$ be a square-free polynomial with $\deg_X F \geq 1$. Then*

$$\left| \sum_{(x, y, z) \in \mathbb{F}_p^3} \left(\frac{F(x, y)}{p} \right) \left(\frac{F(z, y + r)}{p} \right) e_p(s(x - z) + ty) \right| \ll_{\deg F} p^2$$

Proof. We start by rewriting

$$\begin{aligned} & \sum_{(x,y,z) \in \mathbb{F}_p^3} \left(\frac{F(x,y)}{p} \right) \left(\frac{F(z,y+r)}{p} \right) e_p(s(x-z) + ty) \\ &= \sum_{y \in \mathbb{F}_p} e_p(ty) \left(\sum_{x \in \mathbb{F}_p} \left(\frac{F(x,y)}{p} \right) e_p(sx) \right) \left(\sum_{z \in \mathbb{F}_p} \left(\frac{F(z,y+r)}{p} \right) e_p(-sz) \right). \end{aligned}$$

Thanks to Lemma 5.4, there exists $S \subset \mathbb{F}_p$, of cardinality $\#S = O_{\deg F}(1)$ and such that

$$\left| \sum_{x \in \mathbb{F}_p} \left(\frac{F(x,y)}{p} \right) e_p(sx) \right| \ll_{\deg F} \sqrt{p}, \quad \left| \sum_{z \in \mathbb{F}_p} \left(\frac{F(z,y+r)}{p} \right) e_p(-sz) \right| \ll_{\deg F} \sqrt{p},$$

for every $y \notin S$. Then

$$\begin{aligned} & \sum_{(x,y,z) \in \mathbb{F}_p^3} \left(\frac{F(x,y)}{p} \right) \left(\frac{F(z,y+r)}{p} \right) e_p(s(x-z) + ty) \\ &= \sum_{y \in \mathbb{F}_p \setminus S} e_p(ty) \left(\sum_{x \in \mathbb{F}_p} \left(\frac{F(x,y)}{p} \right) e_p(sx) \right) \left(\sum_{z \in \mathbb{F}_p} \left(\frac{F(z,y+r)}{p} \right) e_p(-sz) \right) \\ & \quad + \sum_{y \in S} e_p(ty) \left(\sum_{x \in \mathbb{F}_p} \left(\frac{F(x,y)}{p} \right) e_p(sx) \right) \left(\sum_{z \in \mathbb{F}_p} \left(\frac{F(z,y+r)}{p} \right) e_p(-sz) \right) \\ & \ll_{\deg F} p^2. \end{aligned}$$

□

Lemma 5.8. *Let $F[X, Y] \in \mathbb{F}_q[X, Y]$ be a square-free polynomial such that*

$$F[X, Y] = \prod_{i=1}^n (X + a_i Y + b_i) \text{ over } \overline{\mathbb{F}}_q.$$

Moreover, assume that the following conditions hold:

- i) There exists $i \in \{2, \dots, n\}$ such that $a_1 \neq a_i$.*
- ii) $\frac{a_i - a_1}{a_j - a_1} \cdot (b_j - b_1) \neq b_i - b_1$ for any i, j such that $a_j \neq a_1$ and $i \neq j$.*

Then the sheaf $\mathcal{H}_{F,\chi}$ does not have any irreducible component that is geometrically trivial for every multiplicative character χ of \mathbb{F}_p^\times , i.e.

$$\left| \sum_{(x,y) \in \mathbb{F}_p^2} \chi(F(x,y)) \right| \ll_n p,$$

for every multiplicative character χ of \mathbb{F}_p^\times .

Proof. Let $\mathcal{H}_{F,\chi} = \bigoplus_j \mathcal{F}_j$ be the decomposition of $\mathcal{H}_{F,\chi}$ into irreducible components. Define $J = \{j : \mathcal{F}_j \cong_{\text{geom}} \mathbb{Q}_z\}$. For any $r \geq 1$, we have

$$\sum_{x \in \mathbb{F}_{q^r}} t_{\mathcal{H}_{F,\chi},r}(x) = q^{\frac{3r}{2}} \sum_{j \in J} \omega_j^r + O(q^r),$$

where $|\omega_j| = 1$ for every $j \in J$, thanks to Deligne's resolution of the Riemann hypothesis over finite fields ([FKMS19, Theorem 5.2]). On the other hand, for every $m > 0$, we have

$$\frac{1}{R} \sum_{r \leq R} \left| \frac{1}{q^{\frac{3mr}{2}}} \sum_{x \in \mathbb{F}_{q^{mr}}} t_{\mathcal{H}_{F,\chi},mr}(x) \right|^2 = \frac{1}{R} \sum_{r \leq R} \left| \sum_{j \in J} w_j^{mr} \right|^2 + \frac{1}{q^{\frac{mr}{2}} R}$$

Since w_j are all root of unity we get

$$\lim_{R \rightarrow \infty} \frac{1}{R} \sum_{r \leq R} \left| \sum_{j \in J} w_j^{mr} \right|^2 \geq \#J.$$

which leads to

$$\lim_{R \rightarrow \infty} \frac{1}{R} \sum_{r \leq R} \left| \frac{1}{q^{\frac{3mr}{2}}} \sum_{x \in \mathbb{F}_{q^{mr}}} t_{\mathcal{H}_{F,\chi},mr}(x) \right|^2 \geq \#J. \quad (24)$$

We claim that we can find m such that for every $r \geq 1$,

$$\sum_{x \in \mathbb{F}_{q^{mr}}} t_{\mathcal{H}_{F,\chi},mr}(x) = O_n(q^{mr}).$$

Assuming the claim, (24) implies that $\#J = 0$, as stated in the Lemma. Hence, it rests to prove the claim. First, observe that, for every $m \geq 1$

$$t_{\mathcal{H}_{F,\chi},m}(x) = \sum_{y \in \mathbb{F}_{q^m}} \chi_m(F(x, y)) + O(1) \quad \text{for all } x \in \mathbb{F}_{q^m} \setminus (\text{Sing}(\mathcal{H}_{F,\chi}) \cap \mathbb{F}_{q^m}),$$

So it is enough to prove that for a suitable $m \geq 1$ and for every $r \geq 1$,

$$\sum_{x, y \in \mathbb{F}_{q^{mr}}} \chi_{mr}(F(x, y)) = O_n(q^{mr}).$$

Let m be an integer such that $F[X, Y]$ totally splits over \mathbb{F}_{q^m} . Then for every $r \geq 1$ we have

$$F(X, Y) = \prod_{i=1}^n (X + a_i Y + b_i), \text{ over } \mathbb{F}_{q^{mr}}.$$

Hence

$$\sum_{x, y \in \mathbb{F}_{q^{mr}}} \chi_{mr}(F(x, y)) = \sum_{x, y \in \mathbb{F}_{q^{mr}}} \prod_{i=1}^n \chi_{mr}(x + a_i y + b_i).$$

Using the change of variables $z = x + a_1 y$, we can rewrite

$$F(x, y) = g(z)G(z, y),$$

where

$$g(z) = \prod_{i: a_i = a_1} (z + b_i), \quad G(z, y) = \prod_{i: a_i \neq a_1} (z + (a_i - a_1)y + b_i).$$

Since we are assuming that

- the polynomial

$$F[X, Y] = \prod_{i=1}^n (X + a_i Y + b_i) \text{ over } \overline{\mathbb{F}_q},$$

is square free, and hence whenever $a_i = a_j$ for $i \neq j$, one has $b_i \neq b_j$,

- $a_i \neq a_1$ for some $i \in \{2, \dots, n\}$,

it follows that $G(Z, Y)$ is a non-constant square-free polynomial. Thus, we can rewrite the sum above as

$$\sum_{z, y \in \mathbb{F}_{q^{mr}}} \chi_{mr}(g(z)) \chi_{mr}(G(z, y)) = \sum_{z \in \mathbb{F}_{q^{mr}}} t_{\mathcal{L}_{\chi_{mr}(g(Z))}}(z) t_{\mathcal{H}_{G, \chi}}(z),$$

Where:

- $\mathcal{L}_{\chi_{mr}(g(Z))}$ is the Kummer sheaf over $\overline{\mathbb{A}}_{\mathbb{F}_q^{mr}}^1$ associated with the function $\chi_{mr}(g(z))$, with singularities $\text{Sing}(\mathcal{L}_{\chi_{mr}(g(Z))}) = \{\infty\} \cup \{-b_i : i \text{ such that } a_i = a_1\}$.
- the sheaf $\mathcal{H}_{G, \chi_{mr}}$ is the one in Example 5.3 associated to the polynomial $G(Y, Z)$. In particular, $\text{sing}(\mathcal{H}_{G, \chi_{mr}}) \cap \overline{\mathbb{A}}_{\mathbb{F}_q}^1(\overline{\mathbb{F}}_q) = \{z \in \overline{\mathbb{A}}_{\mathbb{F}_q}^1(\overline{\mathbb{F}}_q) : G(z, Y) \text{ is not square-free}\}$.

We claim that the sheaf $\mathcal{H}_{G, \chi_{mr}}$ is lisse at $-b_1$. To do this, it is enough to show that the polynomial $G(-b_1, Y)$ is square-free: for distinct i, j such that $a_j \neq a_1$, the factors

$$(a_i - a_1)Y + (b_i - b_1), \quad (a_j - a_1)Y + (b_j - b_1)$$

are proportional only if

$$\frac{a_i - a_1}{a_j - a_1} \cdot (b_j - b_1) \neq b_i - b_1,$$

but this is not possible thanks to hypothesis (ii). Since $-b_1 \in \text{Sing}(\mathcal{L}_{\chi_{mr}(g(Z))})$ but $-b_1 \notin \text{sing}(\mathcal{H}_{G, \chi_{mr}})$ and $\mathcal{L}_{\chi_{mr}(g(Z))}$ is irreducible, the sheaf $\mathcal{L}_{\chi_{mr}(g(Z))}$ does not appear in the decomposition $\mathcal{H}_{F, \chi} = \bigoplus_j \mathcal{F}_j$. Then, the semi-orthogonality of trace functions ([FKMS19, Theorem 5.2]) implies that

$$\sum_{x, y \in \mathbb{F}_{q^{mr}}} \chi_{mr}(F(x, y)) = \sum_{z \in \mathbb{F}_{q^{mr}}} t_{\mathcal{L}_{\chi_{mr}(g(Z))}}(z) t_{\mathcal{H}_{G, \chi_{mr}}}(z) \ll_n q^{mr},$$

as claimed. □

We are finally ready to prove:

Proposition 2.4. *Let $F(X, Y) \in \mathbb{F}_p[X, Y]$ a square free polynomial such that*

$$F[X, Y] = \prod_{i=1}^n (X + a_i Y + b_i) \text{ over } \overline{\mathbb{F}}_p,$$

and assume that the following conditions hold:

- i) *There exists $i \in \{2, \dots, n\}$ such that $a_1 \neq a_i$.*
- ii) *$\frac{a_i - a_1}{a_j - a_1} \cdot (b_j - b_1) \neq b_i - b_1$ for any i, j such that $a_j \neq a_1$ and $i \neq j$.*
- iii) *For every i, j with $i \neq j$, $b_i \neq b_j$.*

Moreover, assume that $a_i \neq 0$ for every $i = 1, \dots, n$. Let $s \in \mathbb{F}_p$, then there exists $A_s \subset \{(r, t) \in \mathbb{F}_p \times \mathbb{F}_p\}$ such that

- i) *for any $(r, t) \in (\mathbb{F}_p \times \mathbb{F}_p) \setminus A_s$ and , one has*

$$\left| \sum_{(x, y, z) \in \mathbb{F}_p^3} \left(\frac{F(x, y)}{p} \right) \left(\frac{F(z, y + r)}{p} \right) e_p(s(x - z) + ty) \right| \ll_n p^{3/2} (s, t, p)^{1/2}.$$

- ii) *$\#A_s = O_n(1)$.*

Proof. The case $s = t = 0$ is a special case of Lemma 5.7. Hence, we may assume $(s, t) \neq (0, 0)$. We start by proving the case when $s \neq 0$. We begin by noticing that:

$$\sum_{x,y,z \in \mathbb{F}_p^3} \left(\frac{F(x,y)}{p} \right) \left(\frac{F(z,y+r)}{p} \right) e_p(s(x-z) + ty) = \sum_{\substack{w,x,y,z \in \mathbb{F}_p^4 \\ w^2 = F(x,y)F(z,y+r)}} e_p(s(x-z) + ty).$$

Then, thanks to Theorem 5.5, it is enough to estimate

$$\begin{aligned} N_m(k) &= |\{(w, x, y, z) \in \mathbb{F}_{p^m} : w^2 = F(x, y)F(z, y+r), s(x-z) + ty = k\}| \\ &= |\{(w, x, y) \in \mathbb{F}_{p^m} : w^2 = F(x, z)F(x + \bar{s}(ty - k), y+r)\}|, \end{aligned}$$

for any $m \geq 1$ and any $k \in \mathbb{F}_{p^m}$. First observe that over $\bar{\mathbb{F}}_p$ we can decompose $F(X, Y)F(X + \bar{s}(tY - k), Y + r)$ as

$$F(X, Y)F(X + \bar{s}(tY - k), Y + r) = \prod_{i=1}^n (X + a_i Y + b_i) \prod_{i=1}^n (X + (a_i + \bar{t}s)Y + b_i + a_i r - \bar{t}k). \quad (25)$$

We start by showing that there exists $A_s \subset \{(r, t) \in \mathbb{F}_p^2\}$ such that $\#A_s = O_n(1)$ and such that for any $(r, t) \notin A_s$ the polynomial $F(X, Y)F(X + \bar{s}(tY - k), Y + r)$ is not a square for any k . The polynomial $F(X, Y)$ is square-free by hypothesis. The polynomial $F(X + \bar{s}(tY - k), Y + r)$ is also square-free since it is obtained by composing $F(X, Y)$ with a linear function. From the decomposition in (25) it follows that if $F(X, Y)F(X + \bar{s}(tY - k), Y + r)$ is a square, then there exists a permutation $\alpha \in S_n \setminus \{\text{Id}\}$ such that for every $i = 1, \dots, n$ the factors

$$X + a_i Y + b_i, \quad X + (a_{\alpha(i)} + \bar{s}t)Y + b_{\alpha(i)} + a_{\alpha(i)}r - \bar{s}k,$$

are proportional. From this we deduce that for every $i = 1, \dots, n$ one has

$$\begin{cases} s(a_i - a_{\alpha(i)}) - t = 0 \\ s(b_i - b_{\alpha(i)} - a_{\alpha(i)}r) = k. \end{cases}$$

Now let $i = \alpha_1$ and j such that $\alpha(j) = 1$. Then s and t are the solutions of the system

$$\begin{cases} s(a_1 - a_i) - t = 0 \\ s(b_1 - b_i - a_i r) = k \\ s(a_j - a_1) - t = 0 \\ s(b_j - b_1 - a_1 r) = k. \end{cases}$$

Hence, from the system above, we deduce that

$$\begin{cases} t = s(a_1 - a_i), \\ r = \frac{2b_1 - b_i - b_j}{a_1 - a_j} \end{cases}$$

Now, if we define

$$A_s = \left\{ (t, r) \in (\mathbb{F}_p^\times)^2 : t = s(a_1 - a_i), \quad r = \frac{2b_1 - b_i - b_j}{a_1 - a_j} \text{ for any } i, j = 1, \dots, n \text{ s.t. } a_i \neq a_1 \right\},$$

it follows that for any $(t, r) \notin A_s$ and any $k \in \mathbb{F}_{p^m}$ the polynomial $W^2 = F(X, Y)F(X + \bar{s}(tY - k), Y + r)$ is irreducible for any k . Hence, using the Lang-Weil bound, we get that

$$N_m(k) = p^{2m} + O_n(p^{3m/2}),$$

for any $k \in \mathbb{F}_{p^m}$, provided that $(t, r) \notin A_s$. From now on, we always assume that $(t, r) \notin A_s$. Moreover, to lighten the notation, we denote by $s' = \bar{s}t$, $k' = \bar{s}k$. The next step is to show that the polynomial $F(X, Y)F(X + s'Y - k', Y + r)$ satisfies the hypothesis of Lemma 5.8 for all but $O_n(1)$ choices of k . Notice that the polynomials $F(X, Y)$ and $F(X + s'Y - k', Y + r)$ both satisfy the hypothesis of Lemma 5.8: the polynomial $F(X, Y)$ by hypothesis, and $F(X + s'Y - k', Y + r)$ because it is obtained by $F(X, Y)$ using a change of variables.

We start by showing that the polynomial in (25) is square-free for all but $O_n(1)$ many k' : since both $F(X, Y)$ and $F(X + s'Y - k', Y + r)$ are square-free, we need only check that for all but $O_n(1)$ values of k' and for every i, j , the polynomials

$$X + a_iY + b_i, \quad X + (a_j + s')Y + b_j + a_jr - k', \quad (26)$$

are not proportional, which is guaranteed if $k' \neq b_i - b_j + a_jr$.

Next, we verify that the polynomial $F(X, Y)F(X + s'Y - k', Y + r)$ also satisfies the second condition in Lemma 5.8 for all but $O_n(1)$ many k' . Since $F(X, Y)$ and $F(X + s'Y - k', Y + r)$ already satisfy this condition individually by hypothesis, it remains to ensure that any pair of linear factors coming from different components of the product still satisfy condition (ii) of Lemma 5.8. That is, we must check that given two such factors,

$$X + a_iY + b_i, \quad X + (a_j + s')Y + b_j + a_jr - k',$$

with $a_i \neq a_1$ and $a_j + s' \neq a_1$, they satisfy

$$\frac{b_j + a_jr - k' - b_1}{a_j + s' - a_1} = \frac{b_i - b_1}{a_i - a_1}, \quad (27)$$

for all but $O_n(1)$ values of k' : since we are assuming $a_1 \neq a_i$, equation (27) can be satisfied by only one value of k' . Therefore, for all but $O_n(1)$ many k' , the polynomial $F(X, Y)F(X + s'Y - k', Y + r)$ satisfies both conditions of Lemma 5.8.

Hence, if we denote by χ_m the unique character of order 2 over \mathbb{F}_{p^m} , it follows from Lemma 5.8 that

$$N_m(k) = p^{2m} + \sum_{x, y} \chi_m(F(x, y)F(x + s'y - k', y + r)) = p^{2m} + O_n(p^m),$$

for all but $O_n(1)$ values of k . Thus, we get the result by applying Theorem 5.5.

We now deal with the case $s = 0$, $t \neq 0$. In this case, the exponential sum becomes

$$\sum_{x, y, z \in \mathbb{F}_p^3} \left(\frac{F(x, y)}{p} \right) \left(\frac{F(z, y + r)}{p} \right) e_p(ty) = \sum_{\substack{w, x, y, z \in \mathbb{F}_p^4 \\ w^2 = F(x, y)F(z, y + r)}} e_p(ty).$$

With the goal of using Hooley's Theorem, for any $m \geq 1$ and any $k \in \mathbb{F}_{p^m}$ we have to count

$$\begin{aligned}
N_m(k) &= |\{(w, x, y, z) \in \mathbb{F}_{p^m} : w^2 = F(x, y)F(z, y + r), sy = k\}| \\
&= |\{(w, x, y) \in \mathbb{F}_{p^m} : w^2 = F(x, \bar{t}k)F(z, \bar{t}k + r)\}| \\
&= p^{2m} + \sum_{(x, z) \in \mathbb{F}_{p^m}^2} \chi_m(F(x, \bar{t}k)F(z, \bar{t}k + r)) \\
&= p^{2m} + \left(\sum_{x \in \mathbb{F}_{p^m}} \chi(F(X, \bar{t}k)) \right) \left(\sum_{z \in \mathbb{F}_{p^m}} \chi_m(F(z, \bar{t}k + r)) \right).
\end{aligned}$$

We claim that for every value of $k \neq 0$ the polynomials $F(X, \bar{t}k)$ and $F(Z, \bar{t}k + r)$ are not squares. Let us first show how from the claim we can conclude the proof of the proposition: Since $F(X, \bar{t}k)$ and $F(Z, \bar{t}k + r)$ are not square for every k , we have ([IK04, Theorem 11.23])

$$\left| \sum_{x \in \mathbb{F}_{p^m}} \chi(F(X, \bar{t}k)) \right| \ll_n p^{\frac{m}{2}}, \quad \left| \sum_{z \in \mathbb{F}_{p^m}} \chi_m(F(z, \bar{t}k + r)) \right| \ll_n p^{\frac{m}{2}}$$

Hence,

$$N_m(k) = p^{2m} + O_n(p^m),$$

for every k . Hence, applying Theorem 5.5, we get

$$\left| \sum_{x, y, z \in \mathbb{F}_p^3} \left(\frac{F(x, y)}{p} \right) \left(\frac{F(z, y + r)}{p} \right) e_p(ty) \right| \ll_n p.$$

It rests to prove the claim. To do this it is enough to show that $F(X, \bar{t}k)$ is not a square for every value of k . Since

$$F(X, \bar{t}k) = \prod_{i=1}^n (X + \bar{t}ka_i + b_i) \text{ over } \overline{\mathbb{F}_p}$$

the polynomial $F(X, \bar{t}k)$ is a square if and only if there exists $\alpha \in S_n \setminus \{\text{Id}\}$ such that for every i, \dots, n the factors

$$X + \bar{t}ka_i + b_i, \quad X + \bar{t}ka_{\alpha(i)} + b_{\alpha(i)} \text{ are equal for every } i \in \{1, \dots, n\},$$

that is if and only if

$$k(a_i - a_{\alpha(i)}) = t(b_{\alpha(i)} - b_i) \text{ for every } i \in \{1, \dots, n\}, \quad (28)$$

Now we distinguish two cases:

- a) the case $k \neq 0$. By contradiction, assume that (28) holds. First observe that if such α exists, then $a_{\alpha(i)} \neq a_i$ for all i : indeed, if $a_{\alpha(i)} = a_i$ for some $i \in \{1, \dots, n\}$, then (28) is satisfied only if $b_i = b_{\alpha(i)}$ which is not possible in virtue of assumption (iii). Now let i, j such that $\alpha(i) = 1$ and $\alpha(1) = j$. Then since we are assuming that (28) holds, we would have

$$k = t \frac{b_i - b_1}{a_i - a_1} = t \frac{b_j - b_1}{a_j - a_1},$$

which would be in contradiction with assumption (ii).

b) the case $k = 0$: (28) is true if and only if

$$b_i = b_{\alpha(i)} \text{ for all } i ,$$

but this is not possible since $\alpha \neq \text{Id}$ and our assumption (iii). □

ACKNOWLEDGMENTS

This project began while the author was affiliated with University of Basel, and it originated as a joint project with Pierre Le Boudec. Although P. Le Boudec has decided to moved away from academia and preferred not to be listed as a co-author, the author is deeply grateful for his contribution to the paper. The author would also like to thank Julian Demeio for helpful discussions, which led to a simplification of Lemma 4.3. Thanks are due as well to Tim Browning and Lillian Pierce for their valuable comments on an earlier version of the paper, which helped improve the exposition of the material.

REFERENCES

- [An20] C. An, *ℓ -torsion in class group of certain D_4 -quartic fields*. J. Théor. Nombres Bordeaux, Vol. 32 No. 1, (2020), 1–23.
- [BSTTZ20] M. Bhargava, A. Shankar, T. Taniguchi, F. Thorne, J. Tsimerman and Y. Zhao, *Bounds on 2-torsion in class groups of number fields and integral points on elliptic curves*. J. Amer. Math. Soc., Vol. 33 No. 4, (2020), 1087–1099.
- [BP89] E. Bombieri, and J. Pila. *The number of integral points on arcs and ovals*. Duke Math. J., Vol. 59 No. 2, (1989), 337–357.
- [BB23] D. Bonolis, and T. Browning. *Uniform bounds for rational points on hyperelliptic fibrations*. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), Vol. 24 No. 1, (2023), 173–204.
- [CK25] S. Chan and P. Koymans, *A new pointwise bound for 3-torsion of class groups*. preprint (2025)
- [Del74] P. Deligne, *La conjecture de Weil. I*. Inst. Hautes Études Sci. Publ. Math., No. 43, (1974), 273–307.
- [Del80] P. Deligne, *La conjecture de Weil. II*. Inst. Hautes Études Sci. Publ. Math., No. 52, (1980), 137–252.
- [Dw60] B. Dwork, *On the rationality of the zeta function of an algebraic variety*. American J. Math., No. 82, (1960), 631–648.
- [EPW17] J.S. Ellenberg, L.B. Pierce, and M.M. Wood, *On ℓ -torsion in class groups of number fields*. Algebra Number Theory, Vol. 11 No. 8, (2017), 1739–1778.
- [EV07] J.S. Ellenberg, A. Venkatesh, *Reflection principles and bounds for class group torsion*. Int. Math. Res. Not. IMRN, No. 1, (2007), 18 pp.
- [FK01] É. Fouvry and N. Katz, *A general stratification theorem for exponential sums, and applications*. J. Reine Angew. Math., Vol. 540, (2001), 115–166.
- [FKM15] É. Fouvry, E. Kowalski, and P. Michel, *Algebraic twists of modular forms and Hecke orbits*. Geom. Funct. Anal., Vol. 25 No. 2, (2015), 580–657.
- [FKM19] É. Fouvry, E. Kowalski, and P. Michel, *On the conductor of cohomological transforms*. Ann. Fac. Sci. Toulouse Math. Vol. 30, No. 1, (2021), 203–254.
- [FKMS19] É. Fouvry, E. Kowalski, P. Michel, and W. Sawin *Lectures on applied ℓ -adic cohomology*. Contemp. Math., Vol. 740, (2019), 113–195.
- [FW18] C. Frei and M. Widmer, *Average bounds for the ℓ -torsion in class groups of cyclic extensions*. Res. Number Theory, Vol. 4 No. 3, (2018), 25 pp.
- [FW21] C. Frei and M. Widmer, *Averages and higher moments for the ℓ -torsion in class groups*. Math. Ann., Vol. 379 No. 3-4, (2021), 1205–1229.
- [HB84] D.R. Heath-Brown, *The Square Sieve and Consecutive square-free Numbers*. Math. Ann., Vol. 266 No. 3, (1984), 251–259.
- [HB24] D.R. Heath-Brown, *ℓ -Torsion in Class Groups via Dirichlet L -functions*. preprint, 2024.
- [HP12] D.R. Heath-Brown, L.B. Pierce *Counting rational points on smooth cyclic covers*. J. Number Theory, Vol. 132 No. 8, (2012), 1741–1757

- [HP17] D.R. Heath-Brown and L.B. Pierce, *Averages and moments associated to class numbers of imaginary quadratic fields*. Comp. Math., Vol. 153 No. 11, (2017), 2287–2309.
- [HV06] H.A. Helfgott, A. Venkatesh, *Integral points on elliptic curves and 3-torsion in class groups*. J. Amer. Math. Soc., Vol. 19 No. 3, (2006), 527–550.
- [Ho82] C. Hooley, *On exponential sums and certain of their applications*. Journées Arithmétiques (1980) LMS Lecture Notes Vol. 56 (1982), Cambridge Univ. Press, 92–122.
- [IK04] H. Iwaniec and Emmanuel Kowalski *Analytic number theory*. American Mathematical Society Colloquium Publications Vol. 53, American Mathematical Society Colloquium Publications (2004), xii+615 pp.
- [Kat80] N.M. Katz, *Sommes exponentielles*. Astérisque, Vol. 79, (1980), 1–209.
- [Kat88] N.M. Katz, *Gauss sums, Kloosterman sums, and monodromy groups*. Annals of Mathematics Studies Vol. 116, (1988), x+246 pp.
- [Kat89] N.M. Katz, *Perversity and exponential sums*. Algebraic number theory-in honor of K. Iwasawa, Adv. Stud. Pure Math., Vol. 17 (1989), 209–259
- [Kat90] N.M. Katz, *Exponential sums and differential equations*. Annals of Mathematics Studies Vol. 124, (1990), xii+430 pp.
- [Kat94] N.M. Katz, *Perversity and exponential sums. II. Estimates for and inequalities among A-numbers*. Barsotti Symposium in Algebraic Geometry (Abano Terme, Academic Press (1994), 205–252
- [KW22] J. Klüners, J. Wang, *ℓ -torsion bounds for the class group of number fields with an ℓ -group as Galois group*. Proc. Amer. Math. Soc., Vol. 150 No. 7 (2022), 2793–2805.
- [KT24] P. Koymans and J. Thorner, *Bounds for moments of ℓ -torsion in class groups*. Math. Ann., Vol. 390 No. 2 (2024), 3221–3237.
- [Lan94] S. Lang, *Algebraic number theory*. Graduate Texts in Mathematics, Vol. 110, Springer-Verlag, New York, (1994), xiv+357 pp.
- [LOTZ24] R.J. Lemke Oliver, J. Thorner, and A. Zaman, *An approximate form of Artin’s holomorphy conjecture and non-vanishing of Artin L-functions*. Invent. Math., Vol. 235 No. 3 (2024), 893–971.
- [LOWW25] R.J. Lemke Oliver, J. Wang, M. M. Wood, *The average size of 3-torsion in class groups of 2-extensions*. Forum Math. Pi, Vol. 13, (2025), 1–43.
- [LZ25] R.J. Lemke Oliver and A. Zaman, *Improving the trivial bound for ℓ -torsion in class groups*. Preprint, (2025).
- [Pie05] L.B. Pierce, *The 3-part of class numbers of quadratic fields*. J. London Math. Soc. (2), Vol. 71 No. 3, (2005), 579–598.
- [Pie06] L.B. Pierce, *A bound for the 3-part of class numbers of quadratic fields by means of the square sieve*. Forum Math., Vol. 18 No. 4, (2006), 677–698.
- [PTBW20] L.B. Pierce, R. Turnage-Butterbaugh, and M.M. Wood, *An effective Chebotarev density theorem for families of number fields, with an application to ℓ -torsion in class groups*. Invent. Math., Vol. 219 No. 2, (2020), 701–778.
- [Rot98] J. Rotman, *Galois theory*. Universitext Springer-Verlag, second edition, (1998), xiv+157 pp.
- [Sal22] P. Salberger *Bounds on 2-torsion in class groups over number fields*. Oberwolfach report, Vol.50 (2022), 1252–1252.
- [TZ23] J. Thorner and A. Zaman, *A zero density estimate for Dedekind zeta functions*. Int. Math. Res. Not. IMRN, No. 8, (2023), 6739–6761.
- [Wan20] J. Wang, *Pointwise Bound for ℓ -torsion in Class Groups II: Nilpotent Extensions*. Preprint (2020).
- [Wan21] J. Wang, *Pointwise Bound for ℓ -torsion in Class Groups: Elementary abelian extensions*. J. Reine Angew. Math., Vol. 2021 No. 773, (2021), 129–151.
- [We48] A. Weil, *On some exponential sums*. Proc. Nat. Acad. Sci. U.S.A., Vol. 34 No. 5, (1948), 204–207.
- [Wid18] M. Widmer, *Bounds for the ℓ -torsion in class groups*. Bull. Lond. Math. Soc., Vol. 50 No. 1, (2018), 124–131.

TU GRAZ, STYREGASSE 30, GRAZ, AUSTRIA 8010
 Email address: dante.bonolis@tugraz.at