

HYPERBOLIC $O(N)$ LINEAR SIGMA MODEL AND ITS MEAN-FIELD LIMIT

RUOYUAN LIU, SHAO LIU, AND TADAHIRO OH

ABSTRACT. We study large N limits of the hyperbolic $O(N)$ linear sigma model (HLSM $_N$) on the two-dimensional torus \mathbb{T}^2 , namely, a system of N interacting stochastic damped nonlinear wave equations (SdNLW) with coupled cubic nonlinearities. After establishing (pathwise) global well-posedness of HLSM $_N$ and the limiting equation, called the mean-field SdNLW, we first establish convergence of HLSM $_N$ to the mean-field SdNLW with general initial data (under a suitable assumption). In particular, for the local-in-time convergence, we obtain a convergence rate of order $N^{-\frac{1}{2}+\varepsilon}$ for any $\varepsilon > 0$. We then show that the invariant Gibbs dynamics for HLSM $_N$ converges to that for the mean-field SdNLW with a convergence rate of order $N^{-\frac{1}{2}+\varepsilon}$ on any large time intervals.

CONTENTS

1. Introduction	2
1.1. Hyperbolic $O(N)$ linear sigma model	2
1.2. Large N limit of the hyperbolic $O(N)$ linear sigma model	3
1.3. Large N limit of the invariant Gibbs dynamics	9
2. Notations and preliminary lemmas	13
2.1. Notations	13
2.2. Product estimates	15
2.3. I -operator and basic commutator estimates	15
2.4. Stochastic convolutions and their Wick powers	16
3. Well-posedness of the hyperbolic $O(N)$ linear sigma model	21
3.1. Local well-posedness	21
3.2. Preliminary estimates	25
3.3. Global well-posedness	28
4. Well-posedness of the mean-field SdNLW	32
4.1. Local well-posedness	33
4.2. Global well-posedness	35
5. Mean-field limit of the hyperbolic $O(N)$ linear sigma model	38
5.1. Law of large numbers	38
5.2. Proof of Theorem 1.4 (i)	43
5.3. On the convergence rate under a higher moment assumption	52
6. Mean-field limit of invariant Gibbs dynamics	55
6.1. On the Gibbsian initial data	55
6.2. Convergence of invariant Gibbs dynamics	58

2020 *Mathematics Subject Classification.* 35L71, 60H15, 35R60.

Key words and phrases. $O(N)$ linear sigma model; stochastic nonlinear wave equation; wave equation; Gibbs measure; mean-field limit; stochastic quantization; canonical stochastic quantization.

1. INTRODUCTION

1.1. Hyperbolic $O(N)$ linear sigma model. In this paper, we consider the following system of N interacting stochastic damped nonlinear wave equations (SdNLW) with coupled cubic nonlinearities, posed on the two-dimensional torus $\mathbb{T}^2 = (\mathbb{R}/2\pi\mathbb{Z})^{2,1}$

$$(\partial_t^2 + \partial_t + m - \Delta)u_{N,j} = -\frac{1}{N} \sum_{k=1}^N u_{N,k}^2 u_{N,j} + \sqrt{2}\xi_j, \quad j = 1, \dots, N, \quad (1.1)$$

where $m > 0$ and $\{\xi_j\}_{j \in \mathbb{N}}$ is a family of independent space-time white noises on $\mathbb{R}_+ \times \mathbb{T}^2$. Our main goal is to study the large N limit of solutions to (1.1) and show that (under suitable assumptions on initial data), for each fixed $j \in \mathbb{N}$, a solution $u_{N,j}$ to (1.1) converges in probability to a solution u_j to the following mean-field SdNLW on \mathbb{T}^2 as $N \rightarrow \infty$:

$$(\partial_t^2 + \partial_t + m - \Delta)u_j = -\mathbb{E}[u_j^2] u_j + \sqrt{2}\xi_j. \quad (1.2)$$

See Theorems 1.4 and 1.6 below for the main statements.

In quantum field theory, there are natural models where the field $\vec{u}_N = (u_{N,1}, \dots, u_{N,N})$ takes values in an N -dimensional space, and it is of importance to investigate their asymptotic behavior as $N \rightarrow \infty$. Such a large N limit often leads to a substantial simplification and has been studied for many models. In [60], Wilson considered the following N -component generalization of the Φ_2^4 -measure:

$$\begin{aligned} d\rho_N = Z_N^{-1} \exp \left(- \int_{\mathbb{T}^2} \frac{1}{2} \sum_{j=1}^N |\nabla u_{N,j}|^2 + \frac{m}{2} \sum_{j=1}^N u_{N,j}^2 dx \right. \\ \left. - \frac{1}{4N} \int_{\mathbb{T}^2} \left(\sum_{j=1}^N u_{N,j}^2 \right)^2 dx \right) d\vec{u}_N, \end{aligned} \quad (1.3)$$

known as the $O(N)$ linear sigma model. Here, the word “linear” refers to the fact that the target space \mathbb{R}^N is a linear space. In [47], Shen, Smith, Zhu, and Zhu studied the large N limit of the $O(N)$ linear sigma model ρ_N via the method of stochastic quantization. Namely, they studied the following coupled system of stochastic nonlinear heat equations (SNLH) on \mathbb{T}^2 :

$$(\partial_t + m - \Delta)u_{N,j} = -\frac{1}{N} \sum_{k=1}^N u_{N,k}^2 u_{N,j} + \sqrt{2}\xi_j. \quad (1.4)$$

It is easy to see that the measure ρ_N in (1.3) is formally invariant under the dynamics of (1.4) and, for this reason, we refer to (1.4) as the parabolic $O(N)$ linear sigma model. In [47], the authors showed that a solution $u_{N,j}$ to (1.4) converges to the following mean-field SNLH:

$$(\partial_t + m - \Delta)u_j = -\mathbb{E}[u_j^2] u_j + \sqrt{2}\xi_j$$

(under suitable assumptions on initial data). This dynamical result in turn implied convergence of the invariant measure ρ_N in (1.3) to the limiting Gaussian measure $\mu_1^{\otimes \mathbb{N}}$ (in a suitable sense),

¹In this formal discussion, we ignore the issue of renormalization.

provided that $m > 0$ is sufficiently large. Here, μ_1 denotes the massive Gaussian free field measure with mass $m > 0$:

$$d\mu_1 = Z^{-1} \exp \left(-\frac{1}{2} \|(m - \Delta)^{\frac{1}{2}} u\|_{L^2}^2 \right) du. \quad (1.5)$$

See [47, Theorem 1.2] for a precise statement. In a recent work [14], Delgadino and Smith extended the latter result on the measure convergence to arbitrarily small masses $m > 0$ by making use of Talagrand's inequality [54, 44, 17]. See [48, 49] for further results on the (parabolic) $O(N)$ linear sigma model. See also a nice survey paper [46] on the subject, including physical and mathematical backgrounds on the large N methods and $\frac{1}{N}$ expansions, and the references therein.

As in the scalar case ($N = 1$) studied in [23], the SdNLW system (1.1) corresponds to the hyperbolic counterpart of the parabolic $O(N)$ linear sigma model (1.4). In particular, the SdNLW system (1.1) is the so-called canonical stochastic quantization of the $O(N)$ linear sigma model ρ_N in (1.3); see [45] and also Subsection 1.3 below. For this reason, we refer to the SdNLW system (1.1) as the *hyperbolic $O(N)$ linear sigma model* (HLSM $_N$) in the following

Starting from the work [41, 21], there has been a significant progress on the study of singular stochastic (damped) nonlinear wave equations, broadly interpreted with stochastic forcing and/or random initial data in recent years; see [15, 22, 36, 23, 56, 32, 31, 33, 6, 34, 37, 35, 43, 53, 9, 42, 7, 61]. Nevertheless, to the best of our knowledge, there is no well-posedness result on a wave equation with a mean-field nonlinearity, in particular in the singular setting (such as the mean-field SdNLW (1.2)), not to mention the large N limit (= the mean-field limit) of a system of nonlinear wave equations. Thus, the current paper presents the first such results on the hyperbolic $O(N)$ linear sigma model (1.1) and its mean-field limit (1.2).

1.2. Large N limit of the hyperbolic $O(N)$ linear sigma model. We now take a closer look at HLSM $_N$ (1.1) and its (formal) mean-field limit (1.2). Given $j \in \mathbb{N}$, let Ψ_j denote the stochastic convolution, satisfying the following linear stochastic damped wave equation with zero initial data:

$$\begin{cases} (\partial_t^2 + \partial_t + m - \Delta)\Psi_j = \sqrt{2}\xi_j \\ (\Psi_j, \partial_t \Psi_j)|_{t=0} = (0, 0), \end{cases} \quad (1.6)$$

where $\{\xi_j\}_{j \in \mathbb{N}}$ is the family of independent space-time white noises as in (1.1). As seen in [21, 23], $\Psi_j(t)$ is almost surely a distribution of regularity² $-\varepsilon$ and thus we need to introduce renormalization in considering its powers.

Let $\mathbf{u}_N = \{u_{N,j}\}_{j=1}^N$ be a solution to HLSM $_N$ (1.1). Proceeding with the first order expansion [28, 5, 12]:

$$u_{N,j} = \Psi_j + v_{N,j}, \quad j = 1, \dots, N, \quad (1.7)$$

we see that the residual term $\mathbf{v}_N = \{v_{N,j}\}_{j=1}^N$ satisfies the following SdNLW system:

$$\begin{aligned} (\partial_t^2 + \partial_t + m - \Delta)v_{N,j} = & -\frac{1}{N} \sum_{k=1}^N \left(v_{N,k}^2 v_{N,j} + 2\Psi_k v_{N,k} v_{N,j} + v_{N,k}^2 \Psi_j \right. \\ & \left. + : \Psi_k^2 : v_{N,j} + 2v_{N,k} : \Psi_k \Psi_j : + : \Psi_k^2 \Psi_j : \right), \end{aligned} \quad (1.8)$$

²In describing regularities of functions and distributions, we use $\varepsilon > 0$ to denote an arbitrarily small constant.

where we have already applied Wick renormalization to the powers of the stochastic convolution Ψ_j :

$$\begin{aligned} :\Psi_k\Psi_j: &= \begin{cases} :\Psi_k^2:, & \text{if } k = j, \\ \Psi_k\Psi_j, & \text{if } k \neq j, \end{cases} \\ :\Psi_k^2\Psi_j: &= \begin{cases} :\Psi_k^3:, & \text{if } k = j, \\ :\Psi_k^2:\Psi_j, & \text{if } k \neq j, \end{cases} \end{aligned} \quad (1.9)$$

where $:\Psi_k^\ell:$ on the right-hand side denotes the standard Wick power. See (2.12) and (2.13) for the precise definition via a regularization and limiting procedure. See Subsection 2.4 for a further discussion. Then, by setting

$$\begin{aligned} :u_{N,k}^2 u_{N,j}: &\stackrel{\text{def}}{=} v_{N,k}^2 v_{N,j} + 2\Psi_k v_{N,k} v_{N,j} + v_{N,k}^2 \Psi_j \\ &+ :\Psi_k^2: v_{N,j} + 2v_{N,k} :\Psi_k\Psi_j: + :\Psi_k^2\Psi_j:, \end{aligned}$$

we obtain the renormalized version of HLSM_N (1.1):

$$(\partial_t^2 + \partial_t + m - \Delta)u_{N,j} = -\frac{1}{N} \sum_{k=1}^N :u_{N,k}^2 u_{N,j}: + \sqrt{2}\xi_j. \quad (1.10)$$

As usual in practice [21, 23], we say that $\mathbf{u}_N = \{u_{N,j}\}_{j=1}^N$ is a solution to (1.10) if $u_{N,j}$, $j = 1, \dots, N$, has the structure (1.7), where $\mathbf{v}_N = \{v_{N,j}\}_{j=1}^N$ is a solution to (1.8).

Before proceeding further, we introduce some notations; see also Subsection 2.1. Given $s \in \mathbb{R}$, we set

$$\mathcal{H}^s(\mathbb{T}^2) = H^s(\mathbb{T}^2) \times H^{s-1}(\mathbb{T}^2).$$

For a function space B , we denote by $B^{\otimes N}$ as the N -fold product of the space B with itself: $B^{\otimes N} = \underbrace{B \times \dots \times B}_{N \text{ times}}$. Given a Banach space B and $N \in \mathbb{N}$, we define the $\mathcal{A}_N B$ -norm, denoting

the ℓ^2 -average, by setting

$$\|a\|_{\mathcal{A}_N B}^2 = \frac{1}{N} \sum_{j=1}^N \|a_j\|_B^2 \quad (1.11)$$

for $a = \{a_j\}_{j=1}^N \in B^{\otimes N}$. We may write $\mathcal{A}_{N,j}$ to emphasize that the ℓ^2 -average is taken over the index j . When $B = \mathbb{R}$, we simply set $\mathcal{A}_N = \mathcal{A}_N \mathbb{R}$.

We state pathwise local and global well-posedness of HLSM_N (1.10).

Theorem 1.1. (i) *Let $m > 0$ and $\frac{1}{2} \leq s < 1$. Then, given $N \in \mathbb{N}$, HLSM_N (1.10) is locally well-posed in $(\mathcal{H}^s(\mathbb{T}^2))^{\otimes N}$. More precisely, given $(\mathbf{v}_N^{(0)}, \mathbf{v}_N^{(1)}) = \{(v_{N,j}^{(0)}, v_{N,j}^{(1)})\}_{j=1}^N \in (\mathcal{H}^s(\mathbb{T}^2))^{\otimes N}$, there exists a unique solution*

$$(\mathbf{v}_N, \partial_t \mathbf{v}_N) = \{(v_{N,j}, \partial_t v_{N,j})\}_{j=1}^N \in \mathcal{A}_N C([0, \tau]; \mathcal{H}^s(\mathbb{T}^2)) \quad (1.12)$$

to (1.8) with $(\mathbf{v}_N, \partial_t \mathbf{v}_N)|_{t=0} = (\mathbf{v}_N^{(0)}, \mathbf{v}_N^{(1)})$ on the time interval $[0, \tau]$ for some almost surely positive time $\tau > 0$.

(ii) *Let $m > 0$ and $\frac{4}{5} < s < 1$. Then, given $N \in \mathbb{N}$, HLSM_N (1.10) is globally well-posed in $(\mathcal{H}^s(\mathbb{T}^2))^{\otimes N}$. More precisely, given $(\mathbf{v}_N^{(0)}, \mathbf{v}_N^{(1)}) = \{(v_{N,j}^{(0)}, v_{N,j}^{(1)})\}_{j=1}^N \in (\mathcal{H}^s(\mathbb{T}^2))^{\otimes N}$, there exists*

a unique global-in-time solution $(\mathbf{v}_N, \partial_t \mathbf{v}_N) \in (C(\mathbb{R}_+; \mathcal{H}^s(\mathbb{T}^2)))^{\otimes N}$ to (1.8) with initial data $(\mathbf{v}_N^{(0)}, \mathbf{v}_N^{(1)})$, almost surely.

Theorem 1.1 follows from a straightforward modification of the corresponding results in the scalar case (i.e. $N = 1$) in [23]. We present its proof in Section 3. The local well-posedness result (Theorem 1.1 (i)) follows from Proposition 3.1 on local well-posedness of the perturbed SdNLW system (3.2) with Lemma 2.5 on the regularity properties of the Wick powers of the stochastic convolution Ψ_j defined in (1.6). Our proof does not make use of any auxiliary function space and thus the uniqueness of a solution $(\mathbf{v}_N, \partial_t \mathbf{v}_N)$ holds in the entire class (1.12). See Remark 3.2 (i). We also point out that our argument yields a uniform (in N) local existence time (in the sense of Proposition 3.1).

Let us now turn our attention to the pathwise global well-posedness result (Theorem 1.1 (ii)). Given $(\mathbf{u}_N, \partial_t \mathbf{u}_N) = \{(u_{N,j}, \partial_t u_{N,j})\}_{j=1}^N$, define the energy functional $E_N(\mathbf{u}_N, \partial_t \mathbf{u}_N)$ by setting

$$\begin{aligned} E_N(\mathbf{u}_N, \partial_t \mathbf{u}_N) &= \frac{1}{2N} \sum_{j=1}^N \int_{\mathbb{T}^2} |\nabla u_{N,j}|^2 + m u_{N,j}^2 + (\partial_t u_{N,j})^2 dx \\ &\quad + \frac{1}{4N^2} \int_{\mathbb{T}^2} \left(\sum_{j=1}^N u_{N,j}^2 \right)^2 dx \\ &= \frac{1}{2} \int_{\mathbb{T}^2} \|\nabla u_{N,j}\|_{\mathcal{A}_N}^2 + m \|u_{N,j}\|_{\mathcal{A}_N}^2 + \|\partial_t u_{N,j}\|_{\mathcal{A}_N}^2 dx \\ &\quad + \frac{1}{4} \int_{\mathbb{T}^2} \|u_{N,j}\|_{\mathcal{A}_N}^4 dx, \end{aligned} \tag{1.13}$$

where $\mathcal{A}_N = \mathcal{A}_N \mathbb{R}$ is as in (1.11). The energy functional $E_N(\mathbf{u}_N, \partial_t \mathbf{u}_N)$ is obtained by simply replacing the absolute value (= the norm on \mathbb{R}) in the energy:

$$E(u, \partial_t u) = \frac{1}{2} \int_{\mathbb{T}^2} |\nabla u|^2 + m u^2 + (\partial_t u)^2 dx + \frac{1}{4} \int_{\mathbb{T}^2} u^4 dx \tag{1.14}$$

for the scalar (deterministic) NLW:

$$(\partial_t^2 + m - \Delta)u = -u^3 \tag{1.15}$$

by the $\mathcal{A}_N \mathbb{R}$ -norm. As such, it is easy to check that $E_N(\mathbf{u}_N, \partial_t \mathbf{u}_N)$ is conserved under the following (deterministic) NLW system:

$$(\partial_t^2 + m - \Delta)u_{N,j} = -\frac{1}{N} \sum_{k=1}^N u_{N,k}^2 u_{N,j}, \quad j = 1, \dots, N. \tag{1.16}$$

With $\mathbf{v}_N = \partial_t \mathbf{u}_N$, we can write (1.16) as the following Hamiltonian formulation:

$$\partial_t \begin{pmatrix} \mathbf{u}_N \\ \mathbf{v}_N \end{pmatrix} = \begin{pmatrix} 0 & \mathbf{Id}_N \\ -\mathbf{Id}_N & 0 \end{pmatrix} \nabla_{(\mathcal{A}_N L_x^2)^{\otimes 2}} E_N(\mathbf{u}_N, \mathbf{v}_N), \tag{1.17}$$

where \mathbf{Id}_N is the $N \times N$ identity matrix and $\nabla_{(\mathcal{A}_N L_x^2)^{\otimes 2}} E_N(\mathbf{u}_N, \mathbf{v}_N)$ is the L^2 -gradient (= the Riesz representation of the Fréchet derivative of $E_N(\mathbf{u}_N, \mathbf{v}_N)$) with respect to the $(\mathcal{A}_N L^2(\mathbb{T}^2))^{\otimes 2}$ -inner product.

As observed in [23], there are two difficulties in proving global well-posedness of (1.8) (see Proposition 3.7), using the energy $E_N(\mathbf{u}_N, \partial_t \mathbf{u}_N)$ in (1.13). First of all, due to the roughness of the space-time white noises $\{\xi_j\}_{j=1}^N$, the local-in-time solution $(\mathbf{v}_N, \partial_t \mathbf{v}_N)(t)$ constructed in

Theorem 1.1 (i) does *not* belong to $(\mathcal{H}^1(\mathbb{T}^2))^{\otimes N}$, namely $E_N(\mathbf{v}_N, \partial_t \mathbf{v}_N)(t) = \infty$ for $t > 0$, almost surely. The second issue comes from the fact that $(\mathbf{v}_N, \partial_t \mathbf{v}_N)$ does not solve the NLW system (1.16) but it satisfies the perturbed NLW system (1.8), where we view the terms involving (the Wick powers of) Ψ_j as a perturbation. If each of these difficulties appears alone, then we could overcome the issue by applying the I -method [11] in the former case, and by applying a Gronwall-type argument after Burq and Tzvetkov [10] in the latter case. In our current setting, however, we need to handle both of the difficulties at the same time. Our strategy is to adapt the hybrid approach combining the I -method with a Gronwall-type argument, introduced in [23] by Gubinelli, Koch, Tolomeo, and the third author, to the current vector-valued setting. In Subsections 3.2 and 3.3, we present details of a proof of Theorem 1.1 (ii) by establishing uniform (in N) estimates. Our argument also yields a uniform (in N) double exponential growth bound; see Remark 3.10.

Next, we turn our attention to the well-posedness issue of the mean-field SdNLW (1.2). Take $N \rightarrow \infty$ in (1.8). Heuristically, in view of the law of large numbers (see Subsection 5.1), the empirical average $\frac{1}{N} \sum_{1 \leq k \leq N}$ in (1.8) gets replaced by an expectation \mathbb{E} . Thus, in the limit as $N \rightarrow \infty$, we formally obtain the following limiting mean-field equation:

$$(\partial_t^2 + \partial_t + m - \Delta)v_j = -\mathbb{E}[v_j^2]v_j - 2\mathbb{E}[\Psi_j v_j]v_j - \mathbb{E}[v_j^2]\Psi_j - 2\mathbb{E}[v_j \Psi_j]\Psi_j, \quad (1.18)$$

We note that the fourth and sixth terms on the right-hand side of (1.8) vanish in the limit, since they (formally) converge to $\mathbb{E}[\Psi_j^2:]v_j$ and $\mathbb{E}[\Psi_j^2:] \Psi_j$, respectively, which are both 0 in view of $\mathbb{E}[\Psi_j^2:] = 0$. By setting

$$u_j = \Psi_j + v_j, \quad (1.19)$$

we obtain the following renormalized version of the mean-field SdNLW (1.2):

$$(\partial_t^2 + \partial_t + m - \Delta)u_j = -\mathbb{E}[u_j^2 - \Psi_j^2]u_j + \sqrt{2}\xi_j. \quad (1.20)$$

As in the HLSM $_N$ case, we say that u_j is a solution to (1.20) if u_j has the structure (1.19), where v_j is a solution to (1.18).

Next, we state local and global well-posedness of the mean-field SdNLW (1.20).

Theorem 1.2. (i) *Let $m > 0$, $\frac{1}{2} \leq s < 1$, and $2 \leq p < \infty$. Then, the mean-field SdNLW (1.20) is locally well-posed in $L^p(\Omega; \mathcal{H}^s(\mathbb{T}^2))$. More precisely, given $(v^{(0)}, v^{(1)}) \in L^p(\Omega; \mathcal{H}^s(\mathbb{T}^2))$, there exists a unique solution*

$$(v, \partial_t v) \in L^p(\Omega; C([0, \tau]; \mathcal{H}^s(\mathbb{T}^2))) \quad (1.21)$$

to (1.18) with $(v, \partial_t v)|_{t=0} = (v^{(0)}, v^{(1)})$ on the time interval $[0, \tau]$ for some $\tau > 0$.

(ii) *Let $m > 0$ and $\frac{4}{5} < s < 1$. Then, the mean-field SdNLW (1.20) is globally well-posed in $L^2(\Omega; \mathcal{H}^s(\mathbb{T}^2))$. More precisely, given any $(v^{(0)}, v^{(1)}) \in L^2(\Omega; \mathcal{H}^s(\mathbb{T}^2))$ and $T > 0$, there exists a unique global-in-time solution $(v, \partial_t v) \in L^2(\Omega; C([0, T]; \mathcal{H}^s(\mathbb{T}^2)))$ to (1.18) with $(v, \partial_t v)|_{t=0} = (v^{(0)}, v^{(1)})$.*

We note that the uniqueness of a solution $(v, \partial_t v)$ to (1.18) holds in the entire class (1.21). See Remark 4.2 (i). As in the case of HLSM $_N$ (1.10) (Theorem 1.1), Theorem 1.2 follows from adapting the argument for the scalar case [23] to the current mean-field setting. There is, however, one seemingly problematic term $\mathbb{E}[v_j \Psi_j]\Psi_j$ in (1.18) which does not come with renormalization. Such a term does not appear in the scalar case [23] or in (1.8), representing a new aspect of the current mean-field model, as compared to the hyperbolic $O(N)$ linear sigma

model. See a proof of Proposition 4.1. See also Subsection 4.2 for a globalization argument where we encounter similar issues, specific to the current mean-field setting.

As for the global well-posedness (Theorem 1.2 (ii)), the main strategy is to adapt the hybrid I -method approach [23] to the current mean-field setting, starting with the following energy functional:

$$\begin{aligned}\mathcal{E}(u, \partial_t u) &= \frac{1}{2} \mathbb{E} \left[\int_{\mathbb{T}^2} |\nabla u|^2 + mu^2 + (\partial_t u)^2 dx \right] + \frac{1}{4} \int_{\mathbb{T}^2} (\mathbb{E}[u^2])^2 dx \\ &= \frac{1}{2} \int_{\mathbb{T}^2} \|\nabla u\|_{L_\omega^2}^2 + m\|u\|_{L_\omega^2}^2 + \|\partial_t u\|_{L_\omega^2}^2 dx + \frac{1}{4} \int_{\mathbb{T}^2} \|u\|_{L_\omega^2}^4 dx.\end{aligned}\tag{1.22}$$

Note that the energy $\mathcal{E}(u, \partial_t u)$ is obtained by simply replacing the absolute value (= the norm on \mathbb{R}) in the definition (1.14) of the energy $E(u, \partial_t u)$ for the scalar (deterministic) NLW (1.15) by the $L^2(\Omega)$ -norm. It is easy to check that $\mathcal{E}(u, \partial_t u)$ is conserved under the mean-field NLW:

$$(\partial_t^2 + m - \Delta)u = -\mathbb{E}[u^2]u.\tag{1.23}$$

Remark 1.3. With $v = \partial_t u$, (1.23) is equivalent to the following Hamiltonian formulation:

$$\partial_t \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \nabla_{(L_{\omega,x}^2)^{\otimes 2}} \mathcal{E}(u, v),$$

where $\nabla_{(L_{\omega,x}^2)^{\otimes 2}} \mathcal{E}(u, v)$ is the L^2 -gradient with respect to the $(L^2(\Omega; L^2(\mathbb{T}^2)))^{\otimes 2}$ -inner product.

Finally, we are ready to state a convergence result for HLSM $_N$ (1.10) to the mean-field SdNLW (1.20) with general initial data (under some assumptions).

Theorem 1.4. (i) Let $m > 0$ and $\frac{4}{5} < s < 1$. Let $\{(u_{N,j}^{(0)}, u_{N,j}^{(1)})\}_{j=1}^N \in (\mathcal{H}^s(\mathbb{T}^2))^{\otimes N}$, $N \in \mathbb{N}$, and $\{(u_j^{(0)}, u_j^{(1)})\}_{j \in \mathbb{N}} \in (\mathcal{H}^s(\mathbb{T}^2))^{\otimes \mathbb{N}}$ be sequences of random functions such that

- $(u_j^{(0)}, u_j^{(1)}) \in L^2(\Omega; \mathcal{H}^s(\mathbb{T}^2))$, $j \in \mathbb{N}$, are independent and identically distributed,
- for each fixed $j \in \mathbb{N}$, we have

$$\|(u_{N,j}^{(0)} - u_j^{(0)}, u_{N,j}^{(1)} - u_j^{(1)})\|_{\mathcal{H}^s(\mathbb{T}^2)} \longrightarrow 0\tag{1.24}$$

in probability as $N \rightarrow \infty$, and

$$\|(u_{N,j}^{(0)} - u_j^{(0)}, u_{N,j}^{(1)} - u_j^{(1)})\|_{\mathcal{A}_N \mathcal{H}^s(\mathbb{T}^2)} \longrightarrow 0\tag{1.25}$$

in probability as $N \rightarrow \infty$.

Let $\{(u_{N,j}, \partial_t u_{N,j})\}_{j=1}^N$ and $\{(u_j, \partial_t u_j)\}_{j \in \mathbb{N}}$ be the global-in-time solutions to (1.10) and (1.20) with initial data $\{(u_{N,j}^{(0)}, u_{N,j}^{(1)})\}_{j=1}^N$ and $\{(u_j^{(0)}, u_j^{(1)})\}_{j \in \mathbb{N}}$ constructed in Theorems 1.1 and 1.2, respectively. Then, given any $T > 0$, we have, for each fixed $j \in \mathbb{N}$,

$$\|(u_{N,j}, \partial_t u_{N,j}) - (u_j, \partial_t u_j)\|_{C_T \mathcal{H}_x^s} \longrightarrow 0\tag{1.26}$$

in probability as $N \rightarrow \infty$, and

$$\|(u_{N,j}, \partial_t u_{N,j}) - (u_j, \partial_t u_j)\|_{\mathcal{A}_{N,j} C_T \mathcal{H}_x^s} \longrightarrow 0\tag{1.27}$$

in probability as $N \rightarrow \infty$.

(ii) Let $m > 0$ and $\frac{1}{2} \leq s < 1$. Let $\{(u_{N,j}^{(0)}, u_{N,j}^{(1)})\}_{j=1}^N \in (\mathcal{H}^s(\mathbb{T}^2))^{\otimes N}$, $N \in \mathbb{N}$, and $\{(u_j^{(0)}, u_j^{(1)})\}_{j \in \mathbb{N}} \in (\mathcal{H}^s(\mathbb{T}^2))^{\otimes \mathbb{N}}$ be sequences of random functions such that

- $(u_j^{(0)}, u_j^{(1)}) \in L^6(\Omega; \mathcal{H}^s(\mathbb{T}^2))$, $j \in \mathbb{N}$, are independent and identically distributed,
- there exists small $\varepsilon_0 > 0$ such that, for each fixed $j \in \mathbb{N}$, we have

$$N^{\frac{1}{2}-\varepsilon_0} \|(u_{N,j}^{(0)} - u_j^{(0)}, u_{N,j}^{(1)} - u_j^{(1)})\|_{\mathcal{H}^s(\mathbb{T}^2)} \longrightarrow 0 \quad (1.28)$$

in probability as $N \rightarrow \infty$, and

$$N^{\frac{1}{2}-\varepsilon_0} \|(u_{N,j}^{(0)} - u_j^{(0)}, u_{N,j}^{(1)} - u_j^{(1)})\|_{\mathcal{A}_N \mathcal{H}^s(\mathbb{T}^2)} \longrightarrow 0 \quad (1.29)$$

in probability as $N \rightarrow \infty$.

Let $\{(u_{N,j}, \partial_t u_{N,j})\}_{j=1}^N$ and $\{(u_j, \partial_t u_j)\}_{j \in \mathbb{N}}$ be the local-in-time solutions to (1.10) and (1.20) with initial data $\{(u_{N,j}^{(0)}, u_{N,j}^{(1)})\}_{j=1}^N$ and $\{(u_j^{(0)}, u_j^{(1)})\}_{j \in \mathbb{N}}$ constructed in Theorems 1.1 and 1.2, respectively. Suppose that u_j , $j \in \mathbb{N}$, exists on the time interval $[0, T]$ for some $T > 0$ and that

$$v_j = u_j - \Psi_j \in L^6(\Omega; C([0, T]; H^s(\mathbb{T}^2))), \quad j \in \mathbb{N}. \quad (1.30)$$

Then, we have for each fixed $j \in \mathbb{N}$,

$$N^{\frac{1}{2}-\varepsilon_0} \|(u_{N,j}, \partial_t u_{N,j}) - (u_j, \partial_t u_j)\|_{C_T \mathcal{H}_x^s} \longrightarrow 0 \quad (1.31)$$

in probability as $N \rightarrow \infty$, and

$$N^{\frac{1}{2}-\varepsilon_0} \|(u_{N,j}, \partial_t u_{N,j}) - (u_j, \partial_t u_j)\|_{\mathcal{A}_{N,j} C_T \mathcal{H}_x^s} \longrightarrow 0 \quad (1.32)$$

in probability as $N \rightarrow \infty$.

Theorem 1.4 (i) establishes global-in-time convergence of HLSM $_N$ (1.10) to the mean-field SdNLW (1.20) but it does not provide a rate of convergence. On the other hand, Theorem 1.4 (ii) only establishes local-in-time convergence but it provides a convergence rate of $N^{-\frac{1}{2}+\varepsilon_0}$; see (1.31) and (1.32). In view of Theorem 1.2 (i), we note that the higher moment assumption (1.30) indeed holds true, at least for small $T > 0$ (and hence is not an assumption for short times).

We present a proof of Theorem 1.4 in Section 5. The main difficulty for proving the global-in-time convergence result (Theorem 1.4 (i)) comes from the lack of a higher moment control on a global solution v_j to the mean-field SdNLW (1.18). Namely, due to the use of the energy $\mathcal{E}(v_j, \partial_t v_j)$ in (1.22) which comes with the $L^2(\Omega)$ -norm, Theorem 1.2 (ii) only provides the second moment bound for the residual part $v_j = u_j - \Psi_j$. This is in sharp contrast to the parabolic setting studied in [47], where a higher moment bound is available; see [47, Lemma 3.4]. We overcome this difficulty in the current hyperbolic/dispersive problem by directly establishing law-of-large-numbers type lemmas (Lemmas 5.1, 5.2, and 5.4) in Subsection 5.1, assuming only the second moment bound on v_j . We note that, due to the lack of a higher moment control, we can not rely on a simple orthogonality argument (as in Lemmas 5.5 and 5.6; see a discussion below) and thus we need to proceed with more care, invoking the (strong) law of large numbers in the Banach space setting; see [26, Chapter 7]. Once we prove the key law-of-large-numbers type lemmas (Lemmas 5.1, 5.2, and 5.4), Theorem 1.4 (i) follows from a modification of the pathwise local well-posedness argument with Lemmas 5.1, 5.2, and 5.4; see Subsection 5.2 for details. Note that, in the parabolic case [47], the convergence was shown by an energy estimate which is not available in the current hyperbolic/dispersive setting (recall that the energy is almost surely infinite since $(v_{N,j}(t), \partial_t v_{N,j}(t)) \notin \mathcal{H}^1(\mathbb{T}^2)$, almost surely).

We point out that, in handling the term I_3^N in [47, (4.5)], the authors also established a law-of-large-numbers type result; see [47, Step 4 on p. 163]. Their argument, however, relies on

a square root gain via orthogonality (see [47, (4.1)]) which requires a higher moment bound. In proving Theorem 1.4(ii) under the higher moment assumption, we also establish law-of-large-numbers type lemmas (Lemmas 5.5 and 5.6) but their proofs are much simpler than those in Subsection 5.1 thanks to the higher moment control, which allows us to exploit orthogonality and obtain a square root gain just as in [47]. See Subsection 5.3.

Remark 1.5. The results stated in this subsection (Theorems 1.1, 1.2, and 1.4) hold for general initial data (under suitable assumptions). In particular, we do not make use of invariant Gibbs measures. As such, the damping term $\partial_t u_{N,j}$ and $\sqrt{2}$ on the noises in (1.10) (and also in (1.8), (1.18), and (1.20)) do not play any role, and the corresponding results hold for the undamped case. We also note that these results also hold for the case $m = 0$ (without the damping term), where a separate (but not difficult) analysis at the frequency 0 is required; see, for example, [21]. In the next subsection, we consider invariant Gibbs dynamics, for which the damping term, $\sqrt{2}$ on the noises, and the positivity of m are all necessary. We chose to treat the damped case in this subsection such that we have a uniform presentation throughout the paper.

1.3. Large N limit of the invariant Gibbs dynamics. In this subsection, we consider the hyperbolic $O(N)$ linear sigma model (1.10) and its mean-field limit (1.20) at Gibbs equilibria.

The Gibbs measure $\vec{\rho}_N$ for the coupled NLW system (1.16) is formally given by

$$d\vec{\rho}_N(\mathbf{u}_N, \partial_t \mathbf{u}_N) = Z_N^{-1} e^{-N \cdot E_N(\mathbf{u}_N, \partial_t \mathbf{u}_N)} d\mathbf{u}_N d(\partial_t \mathbf{u}_N),$$

where $(\mathbf{u}_N, \partial_t \mathbf{u}_N) = \{(u_{N,j}, \partial_t u_{N,j})\}_{j=1}^N$ and $E_N(\mathbf{u}_N, \partial_t \mathbf{u}_N)$ is as in (1.13). Let us rewrite $\vec{\rho}_N$ as a weighted Gaussian measure in the following. Given $m > 0$ and $s \in \mathbb{R}$, define the fractional Gaussian field μ_s with mass $m > 0$:

$$d\mu_s = Z^{-1} \exp \left(-\frac{1}{2} \|(m - \Delta)^{\frac{s}{2}} u\|_{L^2}^2 \right) du. \quad (1.33)$$

Recall that μ_s is a Gaussian probability measure on $H^{s-1-\varepsilon}(\mathbb{T}^2) \setminus H^{s-1}(\mathbb{T}^2)$. When $s = 1$, it corresponds to the massive Gaussian free field μ_1 defined in (1.5), while it corresponds to the white noise measure when $s = 0$. With this notation, we can write $\vec{\rho}_N$ as

$$\begin{aligned} d\vec{\rho}_N(\mathbf{u}_N, \partial_t \mathbf{u}_N) &= Z_N^{-1} \exp \left(-\frac{1}{4N} \int_{\mathbb{T}^2} \left(\sum_{j=1}^N u_{N,j}^2 \right)^2 dx \right) d\mu_1^{\otimes N}(\mathbf{u}_N) \otimes d\mu_0^{\otimes N}(\partial_t \mathbf{u}_N) \\ &= d\rho_N(\mathbf{u}_N) \otimes d\mu_0^{\otimes N}(\partial_t \mathbf{u}_N), \end{aligned} \quad (1.34)$$

where ρ_N is the $O(N)$ linear sigma model in (1.3). Using the notation in (1.17) (with $\mathbf{v}_N = \partial_t \mathbf{u}_N$), let us now rewrite the (formal)³ HLSM _{N} (1.1) as

$$\partial_t \begin{pmatrix} \mathbf{u}_N \\ \mathbf{v}_N \end{pmatrix} = \begin{pmatrix} 0 & \mathbf{Id}_N \\ -\mathbf{Id}_N & 0 \end{pmatrix} \nabla_{(\mathcal{A}_N L_x^2)^{\otimes 2}} E_N(\mathbf{u}_N, \mathbf{v}_N) + \begin{pmatrix} 0 \\ -\mathbf{v}_N + \sqrt{2} \boldsymbol{\xi}_N \end{pmatrix}$$

where $\boldsymbol{\xi}_N = \{\xi_j\}_{j=1}^N$. Namely, HLSM _{N} (1.1) is written as a superposition of the coupled NLW system (1.16) and an Ornstein-Uhlenbeck process (for the \mathbf{v}_N -component), both of which (formally) preserves the Gibbs measure $\vec{\rho}_N$. As a result, we see that the Gibbs measure $\vec{\rho}_N$ in (1.34) is (formally) invariant under the dynamics of HLSM _{N} (1.1) (which can be made rigorous with a renormalization and a frequency-truncation; see [23, Section 4]).

³Namely, without renormalization.

Note that $\mu_1^{\otimes N}$ in (1.34) is supported on (vectors of) distributions of negative regularity. Namely, the interaction potential in (1.34) requires a renormalization. At the level of the $O(N)$ linear sigma model, this leads to

$$d\rho_N(\mathbf{u}_N) = Z_N^{-1} \exp \left(-\frac{1}{4N} \int_{\mathbb{T}^2} : \left(\sum_{j=1}^N u_{N,j}^2 \right)^2 : dx \right) d\mu_1^{\otimes N}(\mathbf{u}_N), \quad (1.35)$$

where

$$: \left(\sum_{j=1}^N u_{N,j}^2 \right)^2 : = \sum_{k,j=1}^N : u_{N,k}^2 u_{N,j}^2 :$$

with

$$: u_{N,k}^2 u_{N,j}^2 : = \begin{cases} : u_{N,k}^4 :, & \text{if } k = j, \\ : u_{N,k}^2 : : u_{N,j}^2 :, & \text{if } k \neq j. \end{cases}$$

Here, $:u_k^\ell:$ on the right-hand side denotes the standard Wick power. By adapting the argument from the scalar case ([29, 13, 40]) to the current vector-valued setting, we can realize ρ_N in (1.35) as a probability measure on $(H^{-\varepsilon}(\mathbb{T}^2))^{\otimes N}$; see also [47, Remark 5.8]. The renormalized Gibbs measure $\vec{\rho}_N = \rho_N \otimes \mu_0^{\otimes N}$ (with ρ_N in (1.35)) is then given by

$$\begin{aligned} d\vec{\rho}_N(\mathbf{u}_N, \partial_t \mathbf{u}_N) \\ = Z_N^{-1} \exp \left(-\frac{1}{4N} \int_{\mathbb{T}^2} : \left(\sum_{j=1}^N u_{N,j}^2 \right)^2 : dx \right) d\mu_1^{\otimes N}(\mathbf{u}_N) \otimes d\mu_0^{\otimes N}(\partial_t \mathbf{u}_N). \end{aligned} \quad (1.36)$$

Let us now turn our attention to the dynamical problem. Let $\{(\phi_j^{(0)}, \phi_j^{(1)})\}_{j \in \mathbb{N}}$ be a family of independent $\mathcal{D}'(\mathbb{T}^2)$ -valued random variables, independent of the family $\{\xi_j\}_{j \in \mathbb{N}}$ of the space-time white noises, satisfying

$$\text{Law}(\phi_j^{(0)}) = \mu_1 \quad \text{and} \quad \text{Law}(\phi_j^{(1)}) = \mu_0, \quad j \in \mathbb{N}, \quad (1.37)$$

where μ_s is as in (1.33). Here, $\text{Law}(X)$ denotes the law of a random variable X . Given $j \in \mathbb{N}$, let Φ_j be the solution to the following linear stochastic damped wave equation:

$$\begin{cases} (\partial_t^2 + \partial_t + m - \Delta)\Phi_j = \sqrt{2}\xi_j \\ (\Phi_j, \partial_t \Phi_j)|_{t=0} = (\phi_j^{(0)}, \phi_j^{(1)}). \end{cases} \quad (1.38)$$

A straightforward computation shows that

$$\text{Law}(\Phi_j(t), \partial_t \Phi_j(t)) = \text{Law}(\phi_j^{(0)}, \phi_j^{(1)}) = \mu_1 \otimes \mu_0$$

for any $t \in \mathbb{R}_+$.

Our goal is to study the convergence issue of (the renormalized version of) HL SM_N (1.1) with the Gibbsian initial data:

$$\text{Law} \left(\{(u_{N,j}(0), \partial_t u_{N,j}(0))\}_{j=1}^N \right) = \vec{\rho}_N = \rho_N \otimes \mu_0^{\otimes N}, \quad (1.39)$$

which we assume to be independent of the family $\{\xi_j\}_{j \in \mathbb{N}}$ of the space-time white noises. Moreover, we impose that

$$\partial_t u_{N,j}(0) = \phi_j^{(1)}, \quad j = 1, \dots, N. \quad (1.40)$$

By the first order expansion:

$$u_{N,j} = \Phi_j + v_{N,j},$$

the residual term $v_{N,j}$ satisfies

$$\begin{aligned} (\partial_t^2 + \partial_t + m - \Delta)v_{N,j} = & -\frac{1}{N} \sum_{k=1}^N \left(v_{N,k}^2 v_{N,j} + 2\Phi_k v_{N,k} v_{N,j} + v_{N,k}^2 \Phi_j \right. \\ & \left. + : \Phi_k^2 : v_{N,j} + 2v_{N,k} : \Phi_k \Phi_j : + : \Phi_k^2 \Phi_j : \right) \end{aligned} \quad (1.41)$$

with initial data (recall (1.40))

$$(v_{N,j}, \partial_t v_{N,j})|_{t=0} = (v_{N,j}^{(0)}, 0) = (u_{N,j}(0) - \phi_j^{(0)}, 0),$$

where we have already applied Wick renormalization to the powers of the stochastic convolution Φ_j ; see (2.15) for the more precise definition. Then, by setting

$$\begin{aligned} :u_{N,k}^2 u_{N,j}: & \stackrel{\text{def}}{=} v_{N,k}^2 v_{N,j} + 2\Phi_k v_{N,k} v_{N,j} + v_{N,k}^2 \Phi_j \\ & + : \Phi_k^2 : v_{N,j} + 2v_{N,k} : \Phi_k \Phi_j : + : \Phi_k^2 \Phi_j :, \end{aligned}$$

we obtain the renormalized version of HLSM $_N$ (1.1):

$$(\partial_t^2 + \partial_t + m - \Delta)u_{N,j} = -\frac{1}{N} \sum_{k=1}^N :u_{N,k}^2 u_{N,j}: + \sqrt{2}\xi_j \quad (1.42)$$

with the Gibbsian initial data (1.39). In the current setting, the limiting mean-field equation is given by

$$(\partial_t^2 + \partial_t + m - \Delta)u_j = -\mathbb{E}[u_j^2 - \Phi_j^2] u_j + \sqrt{2}\xi_j. \quad (1.43)$$

Compare (1.43) with (1.20). Consider (1.43) with initial data $(\phi_j^{(0)}, \phi_j^{(1)})$. We first note that $u_j \equiv \Phi_j$ satisfies (1.43) with $(u_j, \partial_t u_j) = (\phi_j^{(0)}, \phi_j^{(1)})$. Under the first order expansion $u_j = \Phi_j + v_j$, this implies that $(v_j, \partial_t v_j) \equiv (0, 0)$ is a global-in-time solution to

$$(\partial_t^2 + \partial_t + m - \Delta)v_j = -\mathbb{E}[v_j^2]v_j - 2\mathbb{E}[\Phi_j v_j]v_j - \mathbb{E}[v_j^2]\Phi_j - 2\mathbb{E}[v_j \Phi_j]\Phi_j \quad (1.44)$$

with zero initial data, which trivially implies

$$(v_j, \partial_t v_j) \in L^p(\Omega; C([0, T]; \mathcal{H}^s(\mathbb{T}^2))) \quad (1.45)$$

for any $s \in \mathbb{R}$ and $T > 0$. By noting that Theorem 1.2 also applies to (1.43) and (1.44) and iteratively applying Theorem 1.2 (i), we then conclude that $(u_j, \partial_t u_j) \equiv (\Phi_j, \partial_t \Phi_j)$ is the unique solution in the class:

$$(\Phi_j, \partial_t \Phi_j) \in L^p(\Omega; C([0, T]; \mathcal{H}^s(\mathbb{T}^2))) \quad (1.46)$$

for any $\frac{1}{2} \leq s < 1$ and $T > 0$.

We now state a convergence result of the invariant Gibbs dynamics of HLSM $_N$ (1.42).

Theorem 1.6. (i) *Let $m > 0$ and $\frac{4}{5} < s < 1$. Then, given $N \in \mathbb{N}$, HLSM $_N$ (1.42) is globally well-posed in $(\mathcal{H}^s(\mathbb{T}^2))^{\otimes N}$. More precisely, given $(\mathbf{v}_N^{(0)}, \mathbf{v}_N^{(1)}) = \{(v_{N,j}^{(0)}, v_{N,j}^{(1)})\}_{j=1}^N \in (\mathcal{H}^s(\mathbb{T}^2))^{\otimes N}$, there exists a unique global-in-time solution*

$$(\mathbf{v}_N, \partial_t \mathbf{v}_N) = \{(v_{N,j}, \partial_t v_{N,j})\}_{j=1}^N \in (C(\mathbb{R}_+; \mathcal{H}^s(\mathbb{T}^2)))^{\otimes N}$$

to (1.41) with initial data $(\mathbf{v}_N^{(0)}, \mathbf{v}_N^{(1)})$, almost surely.

(ii) Let $m > 0$. Then, for each $N \in \mathbb{N}$, the Gibbs measure $\vec{\rho}_N = \rho_N \otimes \mu_0^{\otimes N}$ defined in (1.36) is invariant under the dynamics of (1.42). More precisely, for each $t \in \mathbb{R}_+$, we have

$$\text{Law} \left(\{ (u_{N,j}(t), \partial_t u_{N,j}(t)) \}_{j=1}^N \right) = \vec{\rho}_N.$$

(iii) Let $m > 0$. There exists a probability space (Ω', \mathbb{P}') such that the following statements hold.

(iii.a) There exists a family $\{(\phi_j^{(0)}, \phi_j^{(1)})\}_{j \in \mathbb{N}}$ of independent $\mathcal{D}'(\mathbb{T}^2)$ -valued random variables on Ω' , independent of the family $\{\xi_j\}_{j \in \mathbb{N}}$ of the space-time white noises, satisfying (1.37).

(iii.b) For each $N \in \mathbb{N}$, there exists a family $\{v_{N,j}^{(0)}\}_{j \in \mathbb{N}}$ of H^1 -valued random variables on Ω' , independent of the family $\{\xi_j\}_{j \in \mathbb{N}}$ of the space-time white noises, satisfying

$$\sup_{j=1, \dots, N} \left\| \|v_{N,j}^{(0)}\|_{H_x^1} \right\|_{L^2(\Omega')} \lesssim N^{-\frac{1}{2}}, \quad (1.47)$$

such that

$$\text{Law} \left(\{ (v_{N,j}^{(0)} + \phi_j^{(0)}, \phi_j^{(1)}) \}_{j=1}^N \right) = \vec{\rho}_N.$$

(iii.c) Let $\{(u_{N,j}, \partial_t u_{N,j})\}_{j=1}^N$ be the global-in-time solution to (1.42) with the Gibbsian initial data $\{(v_{N,j}^{(0)} + \phi_j^{(0)}, \phi_j^{(1)})\}_{j=1}^N$ constructed in (iii.b). Then, given any $\varepsilon_0 > 0$, we have, for each fixed $j \in \mathbb{N}$,

$$N^{\frac{1}{2}-\varepsilon_0} \|(u_{N,j}, \partial_t u_{N,j}) - (\Phi_j, \partial_t \Phi_j)\|_{C_T \mathcal{H}_x^s} \longrightarrow 0 \quad (1.48)$$

in probability as $N \rightarrow \infty$, and

$$N^{\frac{1}{2}-\varepsilon_0} \|(u_{N,j}, \partial_t u_{N,j}) - (\Phi_j, \partial_t \Phi_j)\|_{A_{N,j} C_T \mathcal{H}_x^s} \longrightarrow 0 \quad (1.49)$$

in probability as $N \rightarrow \infty$.

Theorem 1.6 (iii) implies propagation of chaos for HLSM_N (1.42). In view of (1.47) and the independence of $\{(\phi_j^{(0)}, \phi_j^{(1)})\}_{j \in \mathbb{N}}$, the Gibbsian initial data $\{(v_{N,j}^{(0)} + \phi_j^{(0)}, \phi_j^{(1)})\}_{j=1}^N$ (which is coupled) is asymptotically chaotic (namely, components becomes independent as $N \rightarrow \infty$). The convergence claims (1.48) and (1.49) show that the solution $\{(u_{N,j}(t), \partial_t u_{N,j}(t))\}_{j=1}^N$ is also asymptotically chaotic for each $t > 0$. See [51, Definition 2.1 on p.177 and Theorem 3.1 on p.203] for a further discussion on propagation of chaos.

Noting that the regularity properties of Φ_j is essentially the same as Ψ_j (see Lemmas 2.5 and 2.7), Theorem 1.6 (i) follows from a slight modification of the proof of Theorem 1.1 presented in Section 3 and thus we omit details. As for Theorem 1.6 (ii), we first note that Bourgain's invariant measure argument [4, 5] is not needed here in view of the pathwise global well-posedness (Theorem 1.6 (i)). The claimed invariance of the Gibbs measure $\vec{\rho}_N$ under (1.42) follows from the invariance of the frequency-truncated version of the Gibbs measure $\vec{\rho}_N$ and a PDE approximation argument; see, for example, [23, 37]. As this is standard in the field by now, we omit details.

The construction of the initial data sets claimed in Theorem 1.6 (iii.a) and (iii.b) is based on the results in [47, 14] (see Lemma 6.2) and follows from a key proposition (Proposition 6.1). We present details in Subsection 6.1. In view of (1.45) and (1.47), the convergence result (Theorem 1.6 (iii.c)) follows from a straightforward modification of the proof of Theorem 1.4 (ii). See Subsection 6.2 for details.

Remark 1.7. In [57], Tolomeo proved ergodicity of the invariant Gibbs measures for the defocusing SdNLW on \mathbb{T}^2 . A slight variation of the argument in [57] yields ergodicity of the Gibbs measure $\bar{\rho}_N$ for HLSM $_N$ (1.42). It would be of interest to study ergodic properties of the invariant Gaussian measure $\mu_1 \otimes \mu_0$ for the mean-field SdNLW (1.43) on \mathbb{T}^2 . Note that such a result is known in the parabolic case; see [47].

Remark 1.8. With a slight modification of the proof, Theorem 1.6 extends to the NLW system (1.16) (with a proper renormalization) with the Gibbsian initial data in (1.39), converging to its mean-field limit (1.23), which is nothing but the linear dynamics:

$$\begin{cases} (\partial_t^2 + m - \Delta)Z_j = 0 \\ (Z_j, \partial_t Z_j)|_{t=0} = (\phi_j^{(0)}, \phi_j^{(1)}). \end{cases}$$

with $\{(\phi_j^{(0)}, \phi_j^{(1)})\}_{j \in \mathbb{N}}$ as in (1.37).

Remark 1.9. (i) In this paper, we considered the defocusing case, namely, with the $-$ sign in front of the nonlinearity $\frac{1}{N} \sum_{k=1}^N u_{N,k}^2 u_{N,j}$ in (1.1). We note that the local-in-time results (Theorems 1.1 (i), 1.2 (i), and 1.4 (ii)) hold true even in the focusing case (i.e. with the $+$ sign in front of the nonlinearity $\frac{1}{N} \sum_{k=1}^N u_{N,k}^2 u_{N,j}$ in (1.1)).

As for Theorem 1.6 (iii) on the convergence of invariant Gibbs dynamics, it does not extend to the focusing case, since, even in the scalar case ($N = 1$), the focusing Φ_2^4 -measure does not exist as a probability measure (even with a (Wick renormalized) L^2 -cutoff (as in [25, 5]) or a taming by a power of the Wick renormalized L^2 -norm); see [8, 38, 20]. See also [39, 33, 34] for an overview of the focusing Gibbs measure (non-)construction and the references therein. On the other hand, in the case of the quadratic nonlinearity, the Gibbs measure exists in the scalar case ($N = 1$; see [38]), which seems to extend to the $O(N)$ linear sigma model (with a taming by a power of the Wick renormalized L^2 -norm, which is suitable for studying wave dynamics). Thus, it would be of interest to investigate if an analogue of Theorem 1.6 (iii) holds in this case.

(ii) In the survey paper [46, Section 9], Shen proposed to study the case of infinite volume, where there is no known result at this point. In the wave case, thanks to the finite speed of propagation, we may combine the ideas from [56] and the current paper to extend Theorem 1.4 for general initial data to HLSM $_N$ (1.10) posed on the plane \mathbb{R}^2 . We will address this issue in a forthcoming work.

2. NOTATIONS AND PRELIMINARY LEMMAS

In Subsections 2.1 and 2.2, we introduce basic notations and recall the product estimates. In Subsection 2.3, we then recall the definition of the I -operator along with its basic properties and commutator estimates. In Subsection 2.4, we go over the definitions of the stochastic convolutions Ψ_j and Φ_j in (1.6) and (1.38), respectively, and their Wick powers along with their regularity properties.

2.1. Notations. Throughout this paper, the value of mass $m > 0$ is fixed and thus we often suppress m -dependence in the remaining part of the paper. In describing regularities of functions and distributions, we use $\varepsilon > 0$ to denote an arbitrarily small constant. Constants in various estimates depend on this ε but, for simplicity of notation, we suppress such ε -dependence. When it is clear from the context, we often suppress dependence on other parameters such as s and t .

For $A, B > 0$, we use $A \lesssim B$ to mean that there exists $C > 0$ such that $A \leq CB$. By $A \sim B$, we mean that $A \lesssim B$ and $B \lesssim A$. We may write \lesssim_α and \sim_α to emphasize the dependence on an external parameter α . We use $A \ll B$ to mean $A \leq c_0 B$ for some small $c_0 > 0$. We use $C > 0$ to denote various constants, which may vary line by line, and we may write C_α or $C(\alpha)$ to signify dependence on an external parameter α .

In dealing with a solution u to a wave equation, we use \vec{u} to denote a pair $(u, \partial_t u)$. With a slight abuse of notation, given a vector (f_1, \dots, f_N) , we often write it as $\{f_j\}_{j=1}^N$ or \vec{f} . We apply a similar convention for a vector (f_1, f_2, \dots) of infinite length and write it as $\{f_j\}_{j=1}^\infty$ or \vec{f} . Namely, we use the vectorial notation for two different meanings: $\vec{u} = (u, \partial_t u)$ and $\vec{f} = (f_1, \dots, f_N) = \{f_j\}_{j=1}^N$, but its meaning is clear from the context.

Given a Banach space B and $N \in \mathbb{N}$, let the $\mathcal{A}_N B$ -norm of $a = \{a_j\}_{j=1}^N \in B^{\otimes N}$ be the ℓ^2 -average of the B -norms of a_j , defined in (1.11). When there is no confusion, we also write $\|a_j\|_{\mathcal{A}_N B}$ for $\|a\|_{\mathcal{A}_N B} = \|a_j\|_{\mathcal{A}_{N,j} B}$. By Jensen's inequality (or Cauchy-Schwarz's inequality), we have

$$\frac{1}{N} \sum_{j=1}^N \|a_j\|_B \leq \|a\|_{\mathcal{A}_N B} = \|a_j\|_{\mathcal{A}_N B}. \quad (2.1)$$

Similarly, we define the $\mathcal{A}_N^{(2)} B$ -norm by setting

$$\|a\|_{\mathcal{A}_N^{(2)} B}^2 = \|a\|_{\mathcal{A}_{N,j} \mathcal{A}_{N,k} B}^2 = \frac{1}{N^2} \sum_{j,k=1}^N \|a_{j,k}\|_B^2 \quad (2.2)$$

for $a = \{a_{j,k}\}_{j,k=1}^N \in B^{\otimes N^2}$. When $B = \mathbb{R}$, we simply set $\mathcal{A}_N = \mathcal{A}_N \mathbb{R}$ and $\mathcal{A}_N^{(2)} = \mathcal{A}_N^{(2)} \mathbb{R}$.

By convention, we endow \mathbb{T}^2 with the normalized Lebesgue measure $dx_{\mathbb{T}^2} = (2\pi)^{-2} dx$ such that we do not need to carry factors involving 2π . For simplicity of notation, we use dx to denote the normalized Lebesgue measure $dx_{\mathbb{T}^2}$ on \mathbb{T}^2 in the following. We write \hat{f} or $\mathcal{F}(f) = \mathcal{F}_x(f)$ to denote the Fourier transform of f on \mathbb{T}^2 . Given $M \in \mathbb{N}$, we denote by \mathbf{P}_M the (spatial) frequency projector onto frequencies $\{n \in \mathbb{Z}^2 : |n| \leq M\}$ and we set $\mathbf{P}_M^\perp = \mathbf{Id} - \mathbf{P}_M$.

We use $\mathcal{D}(t)$ to denote the linear damped wave propagator defined by

$$\mathcal{D}(t) = e^{-\frac{t}{2}} \frac{\sin(t\llbracket \nabla \rrbracket)}{\llbracket \nabla \rrbracket}, \quad (2.3)$$

where

$$\llbracket \nabla \rrbracket = \llbracket \nabla \rrbracket_m = \sqrt{m - \frac{1}{4} - \Delta}.$$

For $n \in \mathbb{Z}^2$, we also set

$$\llbracket n \rrbracket = \llbracket n \rrbracket_m = \sqrt{m - \frac{1}{4} + |n|^2}.$$

(When $0 < m \leq \frac{1}{4}$, the operator $\mathcal{D}(t)$ in (2.3) behaves slightly differently at the frequency $n = 0$. However, it does not affect our analysis.) Given $t_0 \in \mathbb{R}_+$, consider the following linear damped wave equation (with a nice function F on $\mathbb{R}_+ \times \mathbb{T}^2$):

$$\begin{cases} (\partial_t^2 + \partial_t + m - \Delta)u = F \\ (u, \partial_t u)|_{t=t_0} = (f, g). \end{cases} \quad (2.4)$$

Then, a solution u to (2.4) is given by

$$u(t) = S(t - t_0)(f, g) + \mathcal{I}_{t_0}(F)(t),$$

where $S(t)$ is the linear propagator given by

$$S(t)(f, g) = \partial_t \mathcal{D}(t)f + \mathcal{D}(t)(f + g) \quad (2.5)$$

and \mathcal{I}_{t_0} denotes the damped wave Duhamel integral operator given by

$$\mathcal{I}_{t_0}(f)(t) = \int_{t_0}^t \mathcal{D}(t - t')f(t')dt' = \int_{t_0}^t e^{-\frac{t-t'}{2}} \frac{\sin((t-t')\llbracket \nabla \rrbracket)}{\llbracket \nabla \rrbracket} f(t')dt'. \quad (2.6)$$

When $t_0 = 0$, we set $\mathcal{I} = \mathcal{I}_0$ for simplicity.

Given $s \in \mathbb{R}$ and $1 \leq r \leq \infty$, we define the L^r -based Sobolev space $W^{s,r}(\mathbb{T}^2)$ as the closure of $C^\infty(\mathbb{T}^2)$ under the norm:

$$\|f\|_{W^{s,r}} = \|\langle \nabla \rangle^s f\|_{L^r} = \|\mathcal{F}^{-1}(\langle \cdot \rangle^s \widehat{f})\|_{L^r}$$

such that $W^{s,r}(\mathbb{T}^2)$ is separable even when $r = \infty$. Here, $\langle \cdot \rangle = (1 + |\cdot|^2)^{\frac{1}{2}}$. When $r = 2$, we set $H^s(\mathbb{T}^2) = W^{s,2}(\mathbb{T}^2)$. We also define $\mathcal{H}^s(\mathbb{T}^2) = H^s(\mathbb{T}^2) \times H^{s-1}(\mathbb{T}^2)$ via the norm:

$$\|(f, g)\|_{\mathcal{H}^s} = \|f\|_{H^s} + \|g\|_{H^{s-1}}.$$

We will often use the short-hand notations such as $C_T H_x^s$ and $L_\omega^p H_x^s$ for $C([0, T]; H^s(\mathbb{T}^2))$ and $L^p(\Omega; H^s(\mathbb{T}^2))$, respectively. Given an interval $I \subset \mathbb{R}$, we also write $C_I H_x^s$ for $C(I; H^s(\mathbb{T}^2))$, etc. Lastly, given $1 \leq p \leq \infty$, we denote by p' its Hölder conjugate (i.e. $\frac{1}{p} + \frac{1}{p'} = 1$).

2.2. Product estimates. We recall the following product estimates for Sobolev spaces; see [3] for a proof. See also [21].

Lemma 2.1. *Let $s > 0$.*

(i) (fractional Leibniz rule). *Let $1 < p_1, p_2, q_1, q_2, r \leq \infty$ be such that $\frac{1}{r} = \frac{1}{p_1} + \frac{1}{q_1} = \frac{1}{p_2} + \frac{1}{q_2}$. Then, we have*

$$\|fg\|_{W^{s,r}} \lesssim \|f\|_{W^{s,p_1}} \|g\|_{L^{q_1}} + \|f\|_{L^{p_2}} \|g\|_{W^{s,q_2}}.$$

(ii) *Let $1 < p \leq \infty$ and $1 < q, r < \infty$ be such that $\frac{1}{p} + \frac{1}{q} \leq \frac{s}{2} + \frac{1}{r}$ and $q, r' \geq p'$. Then, we have*

$$\|fg\|_{W^{-s,r}} \lesssim \|f\|_{W^{-s,p}} \|g\|_{W^{s,q}}.$$

2.3. I -operator and basic commutator estimates. Fix $0 < s < 1$. Given $M \in \mathbb{N}$, we define a smooth, radially symmetric, non-increasing (in $|\xi|$) multiplier $\mathbf{m} = \mathbf{m}_{s,M} \in C^\infty(\mathbb{R}^2; [0, 1])$, satisfying

$$\mathbf{m}_{s,M}(\xi) = \begin{cases} 1, & \text{if } |\xi| \leq M, \\ \left(\frac{M}{|\xi|}\right)^{1-s}, & \text{if } |\xi| \geq 2M. \end{cases}$$

We then define the I -operator $I = I_{s,M}$ to be the Fourier multiplier operator with the multiplier $\mathbf{m}_{s,M}$:

$$\widehat{I_{s,M} f}(n) = \mathbf{m}_{s,M}(n) \widehat{f}(n), \quad n \in \mathbb{Z}^2. \quad (2.7)$$

As it is customary in the field, we suppress the s -dependence and usually use I (or I_M to denote the M -dependence in an explicit manner) in the following. Note that $If \in H^1(\mathbb{T}^2)$ if and only if $f \in H^s(\mathbb{T}^2)$ with the bound:

$$\|f\|_{H^s} \lesssim \|If\|_{H^1} \lesssim M^{1-s} \|f\|_{H^s}. \quad (2.8)$$

Moreover, from the Littlewood-Paley theorem, we have

$$\|If\|_{W^{s_0+s_1,p}} \lesssim M^{s_1} \|f\|_{W^{s_0,p}} \quad (2.9)$$

for any $s_0 \in \mathbb{R}$, $0 \leq s_1 \leq 1 - s$, and $1 < p < \infty$.

We now recall the commutator estimates. See [23, Lemmas 3.1 and 3.3] for the proofs.

Lemma 2.2. *Let $\frac{2}{3} \leq s < 1$. Then, we have*

$$\|I(f^2g) - (If)^2Ig\|_{L^2} \lesssim M^{2-3s} \|If\|_{H^1}^2 \|Ig\|_{H^1},$$

where the implicit constant is independent of $M \in \mathbb{N}$.

Lemma 2.3. *Let $\frac{2}{3} \leq s < 1$. Then, given small $\delta > 0$, there exist small $\sigma_0 = \sigma_0(\delta) > 0$ and $p_0 = p_0(\delta) \gg 1$ such that*

$$\begin{aligned} \|I(fh) - (If)(Ih)\|_{L^2} &\lesssim M^{-\frac{s}{2}+\delta} \|If\|_{H^1} \|h\|_{W^{-\sigma_0,p_0}}, \\ \|I(fgh) - (If)(Ig)(Ih)\|_{L^2} &\lesssim M^{-s+\frac{1}{2}+\delta} \|If\|_{H^1} \|Ig\|_{H^1} \|h\|_{W^{-\sigma_0,p_0}} \end{aligned}$$

for any sufficiently large $M \gg 1$, where the implicit constant is independent of $M \gg 1$.

2.4. Stochastic convolutions and their Wick powers. In this subsection, by first recalling some basic tools from stochastic analysis (see, for example, [50, 24, 30]), we go over the definitions and basic regularity properties of the stochastic convolutions Ψ_j and Φ_j defined in (1.6) and (1.38), respectively, and their Wick powers, formally defined in (1.9).

First, recall the Hermite polynomials $H_k(x; \sigma)$ of degree k with variance parameter $\sigma > 0$, defined through the generating function:

$$e^{tx - \frac{1}{2}\sigma t^2} = \sum_{k=0}^{\infty} \frac{t^k}{k!} H_k(x; \sigma).$$

For readers' convenience, we write out the first few Hermite polynomials:

$$H_0(x; \sigma) = 1, \quad H_1(x; \sigma) = x, \quad H_2(x; \sigma) = x^2 - \sigma, \quad H_3(x; \sigma) = x^3 - 3\sigma x.$$

Next, we recall the Wiener chaos estimate. Let (H, B, μ) be an abstract Wiener space. Namely, μ is a Gaussian measure on a separable Banach space B with $H \subset B$ as its Cameron-Martin space. Given a complete orthonormal system $\{e_j\}_{j \in \mathbb{N}} \subset B^*$ of $H^* = H$, we define a polynomial chaos of order $k \in \mathbb{N} \cup \{0\}$ to be an element of the form $\prod_{j=1}^{\infty} H_{k_j}(\langle x, e_j \rangle)$, where $x \in B$, $k_j \neq 0$ for only finitely many j 's, $k = \sum_{j=1}^{\infty} k_j$, H_{k_j} is the Hermite polynomial of degree k_j , and $\langle \cdot, \cdot \rangle = {}_B \langle \cdot, \cdot \rangle_{B^*}$ denotes the B - B^* duality pairing. We then denote the closure of the span of polynomial chaoses of order k under $L^2(B, \mu)$ by \mathcal{H}_k . The elements in \mathcal{H}_k are called homogeneous Wiener chaoses of order k . We also set

$$\mathcal{H}_{\leq k} = \bigoplus_{j=0}^k \mathcal{H}_j$$

for $k \in \mathbb{N}$.

As a consequence of the hypercontractivity of the Ornstein-Uhlenbeck semigroup due to Nelson [29], we have the following Wiener chaos estimate ([50, Theorem I.22]). See also [55, Proposition 2.4].

Lemma 2.4. *Let $k \in \mathbb{N}$. Then, we have*

$$\|X\|_{L^p(\Omega)} \leq (p-1)^{\frac{k}{2}} \|X\|_{L^2(\Omega)}$$

for any $p \geq 2$ and $X \in \mathcal{H}_{\leq k}$.

Next, we provide a precise meaning of the stochastic convolutions Ψ_j and Φ_j and define their Wick powers. Let $\{\xi_j\}_{j \in \mathbb{N}}$ be a family of independent space-time white noises on $\mathbb{R}_+ \times \mathbb{T}^2$ as in (1.1). Given $j \in \mathbb{N}$ and $n \in \mathbb{Z}^2$, define B_n^j by setting $B_n^j(t) = \langle \xi_j, \mathbf{1}_{[0,t]} \cdot e^{in \cdot x} \rangle_{t,x}$, where $\langle \cdot, \cdot \rangle_{t,x}$ denotes the duality pairing on $\mathbb{R}_+ \times \mathbb{T}^2$. It is easy to see that $\{B_n^j\}_{j \in \mathbb{N}, n \in \mathbb{Z}^2}$ is a family of independent complex-valued Brownian motions conditioned that $B_{-n}^j = \overline{B_n^j}$ for each $j \in \mathbb{N}$ and $n \in \mathbb{Z}^2$. Note that we have

$$\text{Var}(B_n^j(t)) = \mathbb{E} \left[\langle \xi_j, \mathbf{1}_{[0,t]} \cdot e^{in \cdot x} \rangle_{t,x} \overline{\langle \xi_j, \mathbf{1}_{[0,t]} \cdot e^{in \cdot x} \rangle_{t,x}} \right] = \|\mathbf{1}_{[0,t]} \cdot e^{in \cdot x}\|_{L^2_{t,x}}^2 = t$$

for any $n \in \mathbb{Z}^2$. We then define a family $\{W^j\}_{j \in \mathbb{N}}$ of independent cylindrical Wiener processes on $L^2(\mathbb{T}^2)$ by setting

$$W^j(t) = \sum_{n \in \mathbb{Z}^2} B_n^j(t) e^{in \cdot x}.$$

With these notations, we can write Ψ_j in (1.6) as

$$\begin{aligned} \Psi_j(t) &= \sqrt{2} \int_0^t \mathcal{D}(t-t') \xi_j(dt') \\ &= \sqrt{2} \int_0^t \mathcal{D}(t-t') dW^j(t'), \end{aligned} \tag{2.10}$$

where $\mathcal{D}(t)$ is the linear damped wave propagator defined in (2.3). Given $j, M \in \mathbb{N}$, set $\Psi_{j,M} = \mathbf{P}_M \Psi_j$. Then, given $(t, x) \in \mathbb{R}_+ \times \mathbb{T}^2$, we see that $\Psi_{j,M}(t, x)$ is a mean-zero Gaussian random variable with variance:

$$\begin{aligned} \sigma_M(t) &\stackrel{\text{def}}{=} \mathbb{E}[\Psi_{j,M}^2(t, x)] \\ &= \sum_{|n| \leq M} \left(\frac{1 - e^{-t}}{\llbracket n \rrbracket^2} - \frac{e^{-t} 2 \sin(2t \llbracket n \rrbracket)}{\llbracket n \rrbracket (1 + 4 \llbracket n \rrbracket^2)} - \frac{1 - e^{-t} \cos(2t \llbracket n \rrbracket)}{\llbracket n \rrbracket^2 (1 + 4 \llbracket n \rrbracket^2)} \right) \\ &\sim_{m,t} \log M \longrightarrow \infty, \end{aligned} \tag{2.11}$$

as $M \rightarrow \infty$. Note that, while σ_M depends on $t \in \mathbb{R}_+$, it is independent of $x \in \mathbb{T}^2$.

Given $k, j \in \mathbb{N}$, we define the Wick powers by setting

$$\begin{aligned} :\Psi_{k,M} \Psi_{j,M}: &= \begin{cases} H_2(\Psi_{j,M}; \sigma_M), & \text{if } k = j, \\ \Psi_{k,M} \Psi_{j,M}, & \text{if } k \neq j, \end{cases} \\ :\Psi_{k,M}^2 \Psi_{j,M}: &= \begin{cases} H_3(\Psi_{j,M}; \sigma_M), & \text{if } k = j, \\ H_2(\Psi_{k,M}; \sigma_M) \Psi_{j,M}, & \text{if } k \neq j, \end{cases} \end{aligned} \tag{2.12}$$

where $H_\ell(x; \sigma)$ is the Hermite polynomial of degree ℓ with variance parameter σ . Then, the Wick powers, formally given in (1.9), are defined by

$$\begin{aligned} :\Psi_k \Psi_j: &= \lim_{M \rightarrow \infty} :\Psi_{k,M} \Psi_{j,M}:, \\ :\Psi_k^2 \Psi_j: &= \lim_{M \rightarrow \infty} :\Psi_{k,M}^2 \Psi_{j,M}:. \end{aligned} \tag{2.13}$$

Proceeding as in [21, 22], it is easy to see that, given any $\varepsilon > 0$, the limits in (2.13) exist in $C(\mathbb{R}_+; W^{-\varepsilon, \infty}(\mathbb{T}^2))$ almost surely. See Lemma 2.5.

Next, we turn our attention to Φ_j in (1.38). With the notations above, we have

$$\Phi_j(t) = S(t)(\phi_j^{(0)}, \phi_j^{(1)}) + \sqrt{2} \int_0^t \mathcal{D}(t-t') dW^j(t'), \quad (2.14)$$

where $S(t)$ is as in (2.5) and $\{(\phi_j^{(0)}, \phi_j^{(1)})\}_{j \in \mathbb{N}}$ is a family of independent $\mathcal{D}'(\mathbb{T}^2)$ -valued random variables, satisfying (1.37), which is assumed to be independent of $\{\xi_j\}_{j \in \mathbb{N}}$. Given $M \in \mathbb{N}$ and $j \in \mathbb{N}$, we set $\Phi_{j,M} = \mathbf{P}_M \Phi_j$. Then, using (2.11), it is easy to see that, given $(t, x) \in \mathbb{R}_+ \times \mathbb{T}^2$, $\Phi_{j,M}(t, x)$ is a mean-zero Gaussian random variable with variance:

$$\alpha_M \stackrel{\text{def}}{=} \mathbb{E}[\Phi_{j,M}^2(t, x)] = \sum_{\substack{n \in \mathbb{Z}^2 \\ |n| \leq M}} \frac{1}{m + |n|^2} \sim_m \log M \longrightarrow \infty,$$

as $M \rightarrow \infty$. Note that α_M is independent of $(t, x) \in \mathbb{R}_+ \times \mathbb{T}^2$. Then, as in (2.13), we define the Wick powers by setting

$$\begin{aligned} :\Phi_k \Phi_j: &= \lim_{M \rightarrow \infty} :\Phi_{k,M} \Phi_{j,M}:, \\ :\Phi_k^2 \Phi_j: &= \lim_{M \rightarrow \infty} :\Phi_{k,M}^2 \Phi_{j,M}:, \end{aligned} \quad (2.15)$$

where the limits exist almost surely in $C(\mathbb{R}_+; W^{-\varepsilon, \infty}(\mathbb{T}^2))$ for any $\varepsilon > 0$. Here, $:\Phi_{k,M} \Phi_{j,M}:$ and $:(\Phi_{k,M})^2 \Phi_{j,M}:$ are as in (2.12), but with the variance parameter α_M instead of σ_M .

Before we state lemmas on regularity properties of the stochastic convolutions and their Wick powers, we need to introduce some notations. Fix $N \in \mathbb{N}$. Given $\varepsilon > 0$ and an interval $J \subset \mathbb{R}_+$, we define the space $\mathcal{Z}_N^\varepsilon(J)$ of enhanced data sets \mathbf{Z}_N of the form

$$\mathbf{Z}_N = \{Z_j^{(1)}, Z_{k,j}^{(2)}, Z_{k,j}^{(3)}\}_{j,k=1}^N \quad (2.16)$$

by setting

$$\begin{aligned} \|\mathbf{Z}_N\|_{\mathcal{Z}_N^\varepsilon(J)} &= \|Z_j^{(1)}\|_{\mathcal{A}_N C_J W_x^{-\varepsilon, \infty}} + \|Z_{k,k}^{(2)}\|_{\mathcal{A}_N C_J W_x^{-\varepsilon, \infty}} \\ &\quad + \|Z_{k,j}^{(2)}\|_{\mathcal{A}_N^{(2)} C_J W_x^{-\varepsilon, \infty}} + \|Z_{k,j}^{(3)}\|_{\mathcal{A}_N^{(2)} C_J W_x^{-\varepsilon, \infty}}, \end{aligned} \quad (2.17)$$

where $\mathcal{A}_N B$ and $\mathcal{A}_N^{(2)} B$ are the ℓ^2 -averages defined in (1.11) and (2.2). When $J = [0, T]$, with a slight abuse of notation, we simply set

$$\mathcal{Z}_N^\varepsilon(T) = \mathcal{Z}_N^\varepsilon([0, T]). \quad (2.18)$$

For our application, we take

$$Z_j^{(1)} = \Psi_j, \quad Z_{k,j}^{(2)} = :\Psi_k \Psi_j:, \quad \text{and} \quad Z_{k,j}^{(3)} = :\Psi_k^2 \Psi_j: \quad (2.19)$$

in studying (1.8), while we set

$$Z_j^{(1)} = \Phi_j, \quad Z_{k,j}^{(2)} = :\Phi_k \Phi_j:, \quad \text{and} \quad Z_{k,j}^{(3)} = :\Phi_k^2 \Phi_j: \quad (2.20)$$

in studying (1.41).

The first lemma states the basic regularity properties of these stochastic terms.

Lemma 2.5. (i) *Given $j, k, M \in \mathbb{N}$, set*

$$Z_M^{(1)} = \Psi_{j,M}, \quad Z_M^{(2)} = :\Psi_{k,M} \Psi_{j,M}:, \quad \text{and} \quad Z_M^{(3)} = :\Psi_{k,M}^2 \Psi_{j,M}:,$$

where $\Psi_{j,M} = \mathbf{P}_M \Psi_j$, and $:\Psi_{k,M} \Psi_{j,M}:$ and $:\Psi_{k,M}^2 \Psi_{j,M}:$ are as in (2.12). Let $\ell = 1, 2, 3$. Then, given any $T, \varepsilon > 0$ and finite $p \geq 1$, $Z_M^{(\ell)}$ converges to some limit $Z^{(\ell)}$ in

$$L^p(\Omega; C([0, T]; W^{-\varepsilon, \infty}(\mathbb{T}^2)))$$

and also almost surely in $C(\mathbb{R}_+; W^{-\varepsilon, \infty}(\mathbb{T}^2))$, endowed with the compact-open topology in time, as $M \rightarrow \infty$, where $Z^{(\ell)}$ is given by

$$Z^{(1)} = \Psi_j, \quad Z^{(2)} = :\Psi_k \Psi_j:, \quad \text{and} \quad Z^{(3)} = :\Psi_k^2 \Psi_j:.$$

Moreover, there exist $C, c > 0$ such that

$$\mathbb{P}\left(\|Z^{(\ell)}\|_{C_J W_x^{-\varepsilon, \infty}} > \lambda\right) \leq C \exp\left(-c\lambda^{\frac{2}{\ell}}\right) \quad (2.21)$$

for any $\lambda > 0$ and any interval $J \subset \mathbb{R}_+$ with $|J| = 1$, uniformly in J and $j, k \in \mathbb{N}$.

(ii) Given $N \in \mathbb{N}$, let \mathbf{Z}_N be as in (2.16), where $Z_j^{(1)}$, $Z_{k,j}^{(2)}$, and $Z_{k,j}^{(3)}$ are as in (2.19). Then, given $\varepsilon > 0$, there exist $C, c > 0$ such that

$$\mathbb{P}\left(\|\mathbf{Z}_N\|_{\mathcal{Z}_N^\varepsilon(J)} > \lambda\right) \leq C \exp\left(-c \frac{\lambda^{\frac{2}{3}}}{(1 + |J|)^{\frac{2}{3}}}\right) \quad (2.22)$$

for any interval $J \subset \mathbb{R}_+$ and $\lambda > 0$, uniformly in $N \in \mathbb{N}$, where the $\mathcal{Z}_N^\varepsilon(J)$ -norm is as in (2.17).

(iii) The claims in Parts (i) and (ii) hold even if we replace Ψ_j and its Wick powers by Φ_j defined in (2.14) with the corresponding Wick powers.

Proof. (i) By noting that $Z_M^{(\ell)}, Z^{(\ell)} \in \mathcal{H}_\ell$, the claims in Part (i) follows from a slight modification of the proof of [23, Lemma 2.3], based on the Wiener chaos estimate (Lemma 2.4) and the Garsia-Rodemich-Rumsey inequality ([23, Lemma 2.2]). Since we work with the damped wave equation, there is no growth in time in computing the p th moments of the stochastic terms, which is the reason that the right-hand side of (2.21) is independent of time; compare this with [23, (2.7)]. See also [21, Proposition 2.1] and [22, Lemma 3.1], where the latter utilizes [35, Proposition 2.7 and Lemma A.1].

(ii) For simplicity of the presentation, we only consider the case $J = [0, T]$. We first consider the contribution from the last term on the right-hand side of (2.17). Write $[0, T] = \bigcup_{\ell=0}^{[T]} J_\ell$, where $J_\ell = [\ell, \ell + 1]$ for $\ell = 0, \dots, [T] - 1$ and $J_{[T]} = [T, T]$. Here, $[T]$ denotes the integer part of T .

From (2.21), we have

$$\sup_{\ell=0, \dots, [T]} \sup_{j, k \in \mathbb{N}} \|Z_{k,j}^{(3)}\|_{L_\omega^p C_{J_\ell} W_x^{-\varepsilon, \infty}} \lesssim p^{\frac{3}{2}}.$$

Then, by Minkowski's integral inequality, we obtain

$$\begin{aligned} \left\| \|Z_{k,j}^{(3)}\|_{\mathcal{A}_N^{(2)} C_T W_x^{-\varepsilon, \infty}} \right\|_{L^p(\Omega)} &\leq \sum_{\ell=0}^{[T]} \left\| \|Z_{k,j}^{(3)}\|_{L_\omega^p C_{J_\ell} W_x^{-\varepsilon, \infty}} \right\|_{\mathcal{A}_N^{(2)}} \\ &\lesssim p^{\frac{3}{2}} (1 + T) \end{aligned} \quad (2.23)$$

for any finite $p \geq 2$, where the implicit constant is independent of $N \in \mathbb{N}$. By a crude estimate, we see that the other terms on the right-hand side of (2.17) also satisfy the same bound. Hence, the tail estimate (2.22) follows from (2.23) and Chebyshev's inequality (see the proof of [2, Lemma 3]⁴); see also [58, Lemma 4.5].

⁴Lemma 2.1 in the arXiv version.

(iii) A straightforward modification yields the corresponding results for Φ_j , and thus we omit details. \square

Remark 2.6. (i) As a consequence of (2.21), we have the following moment bound:

$$\mathbb{E} \left[\|Z^{(\ell)}\|_{C_J W_x^{-\varepsilon, \infty}}^p \right] \lesssim p^{\frac{\ell}{2}}$$

for any finite $p \geq 1$ and any interval $J \subset \mathbb{R}_+$ with $|J| = 1$, uniformly in J and $j, k \in \mathbb{N}$.

(ii) In Section 5, we need to deal with the product $\Psi_k \mathbf{P}_M^\perp \Psi_j$ with $k \neq j$, where $\mathbf{P}_M^\perp = \mathbf{Id} - \mathbf{P}_M$. It follows from a slight modification of the proof of Lemma 2.5 that there exists small $\theta > 0$ such that

$$\mathbb{E} \left[\|\Psi_k \mathbf{P}_M^\perp \Psi_j\|_{C_T W_x^{-\varepsilon, \infty}}^p \right] \lesssim_T p M^{-\theta p} \quad (2.24)$$

for any finite $p \geq 1$, uniformly in $j, k, M \in \mathbb{N}$.

Lastly, we state a variant of [23, Lemma 2.4] and its consequence (see [23, Subsection 3.2]) on the logarithmically divergent nature of the stochastic convolution Ψ_j in (2.10).

Lemma 2.7. *Given $0 < s < 1$ and $M \in \mathbb{N}$, let $I = I_{s, M}$ be the I -operator defined in (2.7). Given $j \in \mathbb{N}$ and $(t, x) \in \mathbb{R}_+ \times \mathbb{T}^2$, $I\Psi_j(t, x)$ is a mean-zero Gaussian random variable, satisfying*

$$\mathbb{E}[(I\Psi_j(t, x))^2] \lesssim_s \log\langle M \rangle, \quad (2.25)$$

uniformly in $j \in \mathbb{N}$ and $(t, x) \in \mathbb{R}_+ \times \mathbb{T}^2$. Furthermore, given $N \in \mathbb{N}$, there exists a random variable R_N and $\theta = \theta(s) \gg 1$, independent of $N \in \mathbb{N}$, such that

$$\sup_{N \in \mathbb{N}} \mathbb{E}[R_N] < \infty \quad (2.26)$$

and

$$\|I\Psi_j\|_{L_{[t_1, t_2], x}^r \mathcal{A}_N} \leq r e^{\frac{1}{r} \theta T \log\langle M \rangle} R_N^{\frac{1}{r}} \quad (2.27)$$

for any $1 \leq r < \infty$, $0 \leq t_1 \leq t_2 \leq T$, and $M \in \mathbb{N}$, uniformly in $N \in \mathbb{N}$, where $L_{[t_1, t_2], x}^r \mathcal{A}_N = L_{[t_1, t_2]}^r L_x^r \mathcal{A}_N$.

Proof. The bound (2.25) follows from a straightforward modification of the proof of [23, Lemma 2.4] and thus we omit details.

Given $N \in \mathbb{N}$, define $R_N = R_N(\theta)$ by setting

$$R_N = \sum_{M=1}^{\infty} \sum_{\ell=1}^{\infty} e^{-\theta \ell \log\langle M \rangle} \int_0^\ell \int_{\mathbb{T}^2} e^{\|I\Psi_j(t, x)\|_{\mathcal{A}_N}} dx dt, \quad (2.28)$$

where $\theta = \theta(s) \gg 1$ (to be chosen later) and $\mathcal{A}_N = \mathcal{A}_N \mathbb{R}$ is as in (1.11). Then, from (2.28), we have

$$\begin{aligned} \|I\Psi_j\|_{L_{[t_1, t_2], x}^r \mathcal{A}_N}^r &= \int_{t_1}^{t_2} \int_{\mathbb{T}^2} \|I\Psi_j(t, x)\|_{\mathcal{A}_N}^r dx dt \leq r! \int_0^T \int_{\mathbb{T}^2} e^{\|I\Psi_j(t, x)\|_{\mathcal{A}_N}} dx dt \\ &\leq r! e^{\theta T \log\langle M \rangle} R_N \end{aligned} \quad (2.29)$$

for any $1 \leq r < \infty$, $0 \leq t_1 \leq t_2 \leq T$, and $M \in \mathbb{N}$, uniformly in $N \in \mathbb{N}$. By noting $r! \leq r^r$, the bound (2.27) follows from (2.29).

It remains to prove (2.26). Note that the function $e^{(1+|x|)^{\frac{1}{2}}}$ is convex. From (1.11) and Jensen's inequality, we have

$$\begin{aligned} e^{\|I\Psi_j(t,x)\|_{\mathcal{A}_N}} &\leq \exp\left(\left\{1 + \frac{1}{N} \sum_{j=1}^N (I\Psi_j(t,x))^2\right\}^{\frac{1}{2}}\right) \\ &\leq \frac{1}{N} \sum_{j=1}^N e^{1+|I\Psi_j(t,x)|}. \end{aligned}$$

Then, from (2.28), (2.25), and the fact that $I\Psi_j(t,x)$ is a mean-zero Gaussian random variable, satisfying (2.25), we obtain

$$\begin{aligned} \mathbb{E}[R_N] &\lesssim \frac{1}{N} \sum_{j=1}^N \sum_{M=1}^{\infty} \sum_{\ell=1}^{\infty} e^{-\theta\ell \log\langle M \rangle} \int_0^\ell \int_{\mathbb{T}^2} \mathbb{E}\left[e^{|I\Psi_j(t,x)|}\right] dx dt \\ &\lesssim \frac{1}{N} \sum_{j=1}^N \sum_{M=1}^{\infty} \sum_{\ell=1}^{\infty} e^{-\theta\ell \log\langle M \rangle} \cdot \ell e^{C_s \log\langle M \rangle} \\ &\lesssim 1, \end{aligned}$$

uniformly in $N \in \mathbb{N}$, where the last step follows from taking sufficiently large $\theta = \theta(s) \gg 1$. This proves (2.26). \square

3. WELL-POSEDNESS OF THE HYPERBOLIC $O(N)$ LINEAR SIGMA MODEL

In this section, we prove pathwise global well-posedness of HLSM_N (1.10) (Theorem 1.1). In fact, by the first order expansion (1.7), we show global well-posedness of the SdNLW system (1.8) for $\mathbf{v}_N = \{v_{N,j}\}_{j=1}^N$, which we write out below for readers' convenience:

$$\begin{aligned} (\partial_t^2 + \partial_t + m - \Delta)v_{N,j} &= -\frac{1}{N} \sum_{k=1}^N \left(v_{N,k}^2 v_{N,j} + 2\Psi_k v_{N,k} v_{N,j} + v_{N,k}^2 \Psi_j \right. \\ &\quad \left. + :\Psi_k^2: v_{N,j} + 2v_{N,k} : \Psi_k \Psi_j : + : \Psi_k^2 \Psi_j : \right) \end{aligned} \quad (3.1)$$

for $j = 1, \dots, N$.

In Subsection 3.1, we prove local well-posedness of (3.1) via a standard contraction argument. In Subsection 3.2, we then establish some preliminary uniform (in N) estimates for the modified energy $E_N(t)$ defined in (3.15). Finally, in Subsection 3.3, we prove global well-posedness of (3.1), with uniform (in N) bounds, by adapting the hybrid I -method introduced in [23] to the current vector-valued setting. We note that a slight modification of the argument presented in this section also yields pathwise global well-posedness for (1.42) (Theorem 1.6 (i)). We omit details.

3.1. Local well-posedness. In this subsection, we prove local well-posedness of (3.1). In order to separate the deterministic and probabilistic components of the argument, we consider the following system of SdNLW with a given *deterministic* enhanced data set $\mathbf{Z}_N =$

$\{Z_j^{(1)}, Z_{k,j}^{(2)}, Z_{k,j}^{(3)}\}_{j,k=1}^N$ as in (2.16):

$$\begin{aligned} (\partial_t^2 + \partial_t + m - \Delta)v_{N,j} = & -\frac{1}{N} \sum_{k=1}^N \left(v_{N,k}^2 v_{N,j} + 2Z_k^{(1)} v_{N,k} v_{N,j} + v_{N,k}^2 Z_j^{(1)} \right. \\ & \left. + Z_{k,k}^{(2)} v_{N,j} + 2v_{N,k} Z_{k,j}^{(2)} + Z_{k,j}^{(3)} \right) \end{aligned} \quad (3.2)$$

for $j = 1, \dots, N$. Under (2.19), the system (3.2) reduces to (3.1), while it reduces to (1.41) under (2.20).

Given an interval $J \subset \mathbb{R}_+$, define the $X_N^s(J)$ -norm by setting

$$\begin{aligned} \|\vec{\mathbf{v}}_N\|_{X_N^s(J)} &= \|(\mathbf{v}_N, \partial_t \mathbf{v}_N)\|_{\mathcal{A}_N C_J \mathcal{H}_x^s} \\ &= \|(v_{N,j}, \partial_t v_{N,j})\|_{\mathcal{A}_{N,j} C_J \mathcal{H}_x^s}. \end{aligned} \quad (3.3)$$

When $J = [0, \tau]$, with a slight abuse of notation, we set $X_N^s(\tau) = X_N^s([0, \tau])$.

Proposition 3.1. *Let $m > 0$ and $\frac{1}{2} \leq s < 1$ and $T \geq 1$. Given $N \in \mathbb{N}$, suppose that*

$$\mathbf{Z}_N = \{Z_j^{(1)}, Z_{k,j}^{(2)}, Z_{k,j}^{(3)}\}_{j,k=1}^N \in \mathcal{Z}_N^\varepsilon(T),$$

and $(\mathbf{v}_N^{(0)}, \mathbf{v}_N^{(1)}) = \{(v_{N,j}^{(0)}, v_{N,j}^{(1)})\}_{j=1}^N \in \mathcal{A}_N \mathcal{H}^s(\mathbb{T}^2)$, where $\mathcal{Z}_N^\varepsilon(T)$ and $\mathcal{A}_N B$ are as in (2.17) and (1.11), respectively. Then, given any $t_0 \in [0, T]$, there exists a unique solution $\vec{\mathbf{v}}_N = (\mathbf{v}_N, \partial_t \mathbf{v}_N) = \{(v_{N,j}, \partial_t v_{N,j})\}_{j=1}^N$ to (3.2) with $\vec{\mathbf{v}}_N|_{t=t_0} = (\mathbf{v}_N^{(0)}, \mathbf{v}_N^{(1)})$ on the time interval $[t_0, t_0 + \tau] \cap [0, T]$, where the local existence time

$$\tau = \tau \left(\|(v_{N,j}^{(0)}, v_{N,j}^{(1)})\|_{\mathcal{A}_N \mathcal{H}_x^s}, \|\mathbf{Z}_N\|_{\mathcal{Z}_N^\varepsilon(T)} \right) > 0$$

does not depend on $N \in \mathbb{N}$ in an explicit manner. Moreover, the solution $\vec{\mathbf{v}}_N$ is unique in the entire class $X_N^s([t_0, (t_0 + \tau) \wedge T])$, where $a \wedge b = \min(a, b)$.

Theorem 1.1 (i) follows from Proposition 3.1, (2.19), and Lemma 2.5. We omit details.

Remark 3.2. (i) As pointed out in Section 1, by using the Strichartz estimates as in [21], it is possible to prove local well-posedness of (3.2) for $s > \frac{1}{4}$. However, the uniqueness guaranteed by such an argument holds only in the intersection of $X_N^s(\tau \wedge (T - t_0))$ with a suitable ℓ^2 -averaged version of Strichartz spaces. A similar remark holds for the local well-posedness argument of the mean-field SdNLW (1.20) (see Proposition 4.1) for which unconditional uniqueness of solutions plays a key role in Section 6. Since we prefer to have a uniform treatment for both HLSM_N (1.10) and the mean-field SdNLW (1.20), we restrict our attention to $\frac{1}{2} \leq s < 1$ in Proposition 3.1 (and also in Proposition 4.1), guaranteeing unconditional uniqueness of solutions.

(ii) Let $T \geq 1$. Given $N \in \mathbb{N}$, let \mathbf{v}_N be the local-in-time solution to (3.2) constructed in Proposition 3.1. Then, by denoting by T_* its maximal existence time, the following blowup alternative holds:

$$T_* = T \quad \text{or} \quad \lim_{t \rightarrow T_*^-} \|(\mathbf{v}_N, \partial_t \mathbf{v}_N)(t)\|_{\mathcal{A}_N \mathcal{H}_x^s} = \infty. \quad (3.4)$$

We now present a proof of Proposition 3.1.

Proof of Proposition 3.1. The claimed local well-posedness of (3.2) follows from a straightforward adaptation of the proof of [23, Proposition 4.1]. For readers' convenience, however, we present some details, in particular to show that the local existence time $\tau > 0$ does not depend on $N \in \mathbb{N}$ in an explicit manner.

Without loss of generality, we assume that $t_0 = 0$. Fix $N \in \mathbb{N}$. By writing (3.1) in the Duhamel formulation, we have

$$v_{N,j}(t) = S(t)(v_{N,j}^{(0)}, v_{N,j}^{(1)}) - \sum_{\ell=1}^6 \mathcal{I}(A_{j,\ell})(t), \quad j = 1, \dots, N, \quad (3.5)$$

where $S(t)$ and $\mathcal{I} = \mathcal{I}_0$ are as in (2.5) and (2.6), respectively, and $\{A_{j,\ell} = A_{j,\ell}(N)\}_{j=1}^N$, $\ell = 1, \dots, 6$, are given by

$$\begin{aligned} A_{j,1} &= \frac{1}{N} \sum_{k=1}^N v_{N,k}^2 v_{N,j}, & A_{j,2} &= \frac{1}{N} \sum_{k=1}^N 2Z_k^{(1)} v_{N,k} v_{N,j}, \\ A_{j,3} &= \frac{1}{N} \sum_{k=1}^N v_{N,k}^2 Z_j^{(1)}, & A_{j,4} &= \frac{1}{N} \sum_{k=1}^N Z_{k,k}^{(2)} v_{N,j}, \\ A_{j,5} &= \frac{1}{N} \sum_{k=1}^N 2v_{N,k} Z_{k,j}^{(2)}, & A_{j,6} &= \frac{1}{N} \sum_{k=1}^N Z_{k,j}^{(3)}. \end{aligned}$$

Fix $0 < \tau \leq 1$ and small $\varepsilon > 0$ such that $s \leq 1 - \varepsilon$. By Sobolev's and Hölder's inequalities with (3.3), we have

$$\begin{aligned} \|\vec{\mathcal{I}}(A_{j,1})\|_{X_{N,j}^s(\tau)} &\lesssim \tau \frac{1}{N} \sum_{k=1}^N \|v_{N,k}^2 v_{N,j}\|_{\mathcal{A}_{N,j} C_\tau H_x^{s-1}} \\ &\lesssim \tau \frac{1}{N} \sum_{k=1}^N \|v_{N,k}\|_{C_\tau L_x^{\frac{6}{2-s}}}^2 \|v_{N,j}\|_{\mathcal{A}_{N,j} C_\tau L_x^{\frac{6}{2-s}}} \\ &\leq \tau \|\vec{\mathbf{v}}_N\|_{X_N^s(\tau)}^3, \end{aligned} \quad (3.6)$$

provided that $\frac{1}{2} \leq s (< 1)$. Here, $\vec{\mathcal{I}}(A_{j,1})$ is given by

$$\vec{\mathcal{I}}(A_{j,1}) = (\mathcal{I}(A_{j,1}), \partial_t \mathcal{I}(A_{j,1})) \quad (3.7)$$

and we use analogous notations in the following. By (3.3), Sobolev's inequality, Lemma 2.1 (ii), the fractional Leibniz rule (Lemma 2.1 (i)), Sobolev's inequality, and Cauchy-Schwarz's inequality (in k) with (2.17), we have

$$\begin{aligned} \|\vec{\mathcal{I}}(A_{j,2})\|_{X_{N,j}^s(\tau)} &\lesssim \tau \frac{1}{N} \sum_{k=1}^N \|Z_k^{(1)} v_{N,k} v_{N,j}\|_{\mathcal{A}_{N,j} C_\tau W_x^{-\varepsilon, \frac{2}{2-s-\varepsilon}}} \\ &\lesssim \tau \frac{1}{N} \sum_{k=1}^N \|Z_k^{(1)}\|_{C_T W_x^{-\varepsilon, \infty}} \|v_{N,k} v_{N,j}\|_{\mathcal{A}_{N,j} C_\tau W_x^{\varepsilon, \frac{2}{2-s-\varepsilon}}} \\ &\lesssim \tau \frac{1}{N} \sum_{k=1}^N \|Z_k^{(1)}\|_{C_T W_x^{-\varepsilon, \infty}} \|v_{N,k}\|_{C_\tau H_x^{\frac{s}{2} + \frac{3}{2}\varepsilon}} \|v_{N,j}\|_{\mathcal{A}_{N,j} C_\tau H_x^{\frac{s}{2} + \frac{3}{2}\varepsilon}} \\ &\leq \tau \|\mathbf{Z}_N\|_{\mathcal{Z}_N^s(T)} \|\vec{\mathbf{v}}_N\|_{X_N^s(\tau)}^2, \end{aligned} \quad (3.8)$$

provided that $3\varepsilon \leq s \leq 1 - \varepsilon$. A similar computation yields

$$\|\vec{\mathcal{I}}(A_{j,3})\|_{X_{N,j}^s(\tau)} \lesssim \tau \|\mathbf{Z}\|_{\mathcal{Z}_N^s(T)} \|\vec{\mathbf{v}}_N\|_{X_N^s(\tau)}^2. \quad (3.9)$$

By Sobolev's inequality, Lemma 2.1 (ii), and (2.1) with (2.17), we have

$$\begin{aligned}
\|\vec{\mathcal{I}}(A_{j,4})\|_{X_{N,j}^s(\tau)} &\lesssim \tau \frac{1}{N} \sum_{k=1}^N \|Z_{k,k}^{(2)} v_{N,j}\|_{\mathcal{A}_{N,j} C_\tau W_x^{-\varepsilon, \frac{2}{2-s-\varepsilon}}} \\
&\lesssim \tau \frac{1}{N} \sum_{k=1}^N \|Z_{k,k}^{(2)}\|_{C_T W_x^{-\varepsilon, \infty}} \|v_{N,j}\|_{\mathcal{A}_{N,j} C_\tau W_x^{\varepsilon, \frac{2}{2-s-\varepsilon}}} \\
&\lesssim \tau \|\mathbf{Z}_N\|_{\mathcal{Z}_N^\varepsilon(T)} \|\vec{\mathbf{v}}_N\|_{X_N^s(\tau)},
\end{aligned} \tag{3.10}$$

provided that $\varepsilon \leq s \leq 1 - \varepsilon$. Similarly, by Cauchy-Schwarz's inequality (in k), we have

$$\begin{aligned}
\|\vec{\mathcal{I}}(A_{j,5})\|_{X_{N,j}^s(\tau)} &\lesssim \tau \frac{1}{N} \sum_{k=1}^N \|v_{N,k} Z_{k,j}^{(2)}\|_{\mathcal{A}_{N,j} C_\tau H_x^{s-1}} \\
&\lesssim \tau \|\mathbf{Z}_N\|_{\mathcal{Z}_N^\varepsilon(T)} \|\vec{\mathbf{v}}_N\|_{X_N^s(\tau)}.
\end{aligned} \tag{3.11}$$

Lastly, it follows from (3.3), (2.1), and (2.2) with (2.17) that

$$\begin{aligned}
\|\vec{\mathcal{I}}(A_{j,6})\|_{X_{N,j}^s(\tau)} &\lesssim \tau \left\| \frac{1}{N} \sum_{k=1}^N Z_{k,j}^{(3)} \right\|_{\mathcal{A}_{N,j} C_T W_x^{s-1, \infty}} \leq \tau \|Z_{k,j}^{(3)}\|_{\mathcal{A}_{N,j}^{(2)} C_T W_x^{-\varepsilon, \infty}} \\
&\leq \tau \|\mathbf{Z}_N\|_{\mathcal{Z}_N^\varepsilon(T)},
\end{aligned} \tag{3.12}$$

provided that $s - 1 \leq -\varepsilon$.

Denote the right-hand side of (3.5) by $\Gamma_{N,j}(\vec{\mathbf{v}}_N)$, and set

$$\vec{\mathbf{\Gamma}}_N(\vec{\mathbf{v}}_N) = \{(\Gamma_{N,j}(\vec{\mathbf{v}}_N), \partial_t \Gamma_{N,j}(\vec{\mathbf{v}}_N))\}_{j=1}^N.$$

Then, putting (3.5)-(3.12) together, we obtain

$$\begin{aligned}
\|\vec{\mathbf{\Gamma}}_N(\vec{\mathbf{v}}_N)\|_{X_N^s(\tau)} &\leq C_1 \|(v_{N,j}^{(0)}, v_{N,j}^{(1)})\|_{\mathcal{A}_{N,j} \mathcal{H}_x^s} + C_2 \tau \|\vec{\mathbf{v}}_N\|_{X_N^s(\tau)}^3 \\
&\quad + C_3 \tau \|\mathbf{Z}_N\|_{\mathcal{Z}_N^\varepsilon(T)} \left(1 + \|\vec{\mathbf{v}}_N\|_{X_N^s(\tau)}\right)^2
\end{aligned}$$

for any $\vec{\mathbf{v}}_N \in X_N^s(\tau)$. A similar computation yields the following difference estimate:

$$\begin{aligned}
&\|\vec{\mathbf{\Gamma}}_N(\vec{\mathbf{v}}_N) - \vec{\mathbf{\Gamma}}_N(\vec{\mathbf{w}}_N)\|_{X_N^s(\tau)} \\
&\lesssim \tau \left(1 + \|\mathbf{Z}_N\|_{\mathcal{Z}_N^\varepsilon(T)}\right) \\
&\quad \times \left(1 + \|\vec{\mathbf{v}}_N\|_{X_N^s(\tau)} + \|\vec{\mathbf{w}}_N\|_{X_N^s(\tau)}\right)^2 \|\vec{\mathbf{v}}_N - \vec{\mathbf{w}}_N\|_{X_N^s(\tau)}
\end{aligned}$$

for any $\vec{\mathbf{v}}_N, \vec{\mathbf{w}}_N \in X_N^s(\tau)$. Therefore, by choosing

$$\tau = \tau \left(\|(v_{N,j}^{(0)}, v_{N,j}^{(1)})\|_{\mathcal{A}_N \mathcal{H}^s}, \|\mathbf{Z}_N\|_{\mathcal{Z}_N^\varepsilon(T)} \right) > 0$$

sufficiently small, we conclude that $\vec{\mathbf{\Gamma}}_N$ is a contraction in the (closed) ball $B_R \subset X_N^s(\tau)$ of radius $R = 2C_1 \|(v_{N,j}^{(0)}, v_{N,j}^{(1)})\|_{\mathcal{A}_N \mathcal{H}^s} + 1$. At this point, the uniqueness holds only in the ball B_R but by a standard continuity argument, we can extend the uniqueness to hold in the entire class $X_N^s(\tau)$. We omit details. This concludes the proof of Proposition 3.1. \square

3.2. Preliminary estimates. In this subsection, we establish preliminary uniform (in N) estimates for proving global well-posedness of HLSM_N (1.10) by pathwise analysis.

Fix a target time $T \gg 1$. In view of the blowup alternative (3.4) and (2.8), almost sure global well-posedness of (3.1) on $[0, T]$ follows once we show the following bound:

$$\sup_{0 \leq t \leq T} \|(Iv_{N,j}, \partial_t Iv_{N,j})(t; \omega)\|_{\mathcal{A}_N \mathcal{H}_x^1} \leq C(\omega, T) \quad (3.13)$$

for some almost surely finite constant $C(\omega, T) > 0$, where $\{Iv_{N,j}\}_{j=1}^N$ is a solution to the following I -SdNLW system:

$$\begin{aligned} & (\partial_t^2 + \partial_t + m - \Delta)Iv_{N,j} \\ &= -\frac{1}{N} \sum_{k=1}^N \left(I(v_{N,k}^2 v_{N,j}) + 2I(\Psi_k v_{N,k} v_{N,j}) + I(v_{N,k}^2 \Psi_j) \right. \\ & \quad \left. + I(\Psi_k^2 : v_{N,j}) + 2I(v_{N,k} : \Psi_k \Psi_j) + I(\Psi_k^2 \Psi_j) \right). \end{aligned} \quad (3.14)$$

Here, I denotes the I -operator defined in (2.7). Define the modified energy $E_N(t)$ by setting

$$E_N(t) = E_N(I\mathbf{v}_N, \partial_t I\mathbf{v}_N)(t), \quad (3.15)$$

where E_N is the energy defined in (1.13). Then, (3.13) follows once we prove

$$\sup_{0 \leq t \leq T} E_N(t; \omega) \leq C'(\omega, T)$$

for some almost surely finite constant $C'(\omega, T) > 0$.

Let $t_2 \geq t_1 \geq 0$. From (3.15) and (3.14), we have

$$\begin{aligned} E_N(t_2) - E_N(t_1) &= \sum_{\ell=1}^4 B_\ell - \int_{t_1}^{t_2} \int_{\mathbb{T}^2} \frac{1}{N} \sum_{j=1}^N (\partial_t Iv_{N,j})^2 dx dt \\ &\leq \sum_{\ell=1}^4 B_\ell, \end{aligned} \quad (3.16)$$

where $B_\ell = B_\ell(N)$, $\ell = 1, \dots, 4$, are given by

$$\begin{aligned} B_1 &= \frac{1}{N^2} \sum_{j,k=1}^N \int_{t_1}^{t_2} \int_{\mathbb{T}^2} \partial_t Iv_{N,j} \left((Iv_{N,k})^2 Iv_{N,j} - I(v_{N,k}^2 v_{N,j}) \right) dx dt \\ B_2 &= -\frac{1}{N^2} \sum_{j,k=1}^N \int_{t_1}^{t_2} \int_{\mathbb{T}^2} \partial_t Iv_{N,j} \left(2I(\Psi_k v_{N,k} v_{N,j}) + I(v_{N,k}^2 \Psi_j) \right) dx dt, \\ B_3 &= -\frac{1}{N^2} \sum_{j,k=1}^N \int_{t_1}^{t_2} \int_{\mathbb{T}^2} \partial_t Iv_{N,j} \left(I(\Psi_k^2 : v_{N,j}) + 2I(v_{N,k} : \Psi_k \Psi_j) \right) dx dt, \\ B_4 &= -\frac{1}{N^2} \sum_{j,k=1}^N \int_{t_1}^{t_2} \int_{\mathbb{T}^2} \partial_t Iv_{N,j} \sum_{k=1}^N I(\Psi_k^2 \Psi_j) dx dt. \end{aligned} \quad (3.17)$$

In (3.16) and (3.17), we suppressed t -dependence for notational simplicity; we apply the same convention in the following. The first term B_1 is the main commutator term, appearing in the standard application of the I -method, while the terms B_2 , B_3 , and B_4 represent the contributions from the perturbative terms in (3.14) which are to be controlled by a Gronwall-type argument.

As seen in [23], the key contribution from the perturbative terms comes from B_2 (or rather B'_2 defined in (3.19) below).

In the following, we establish uniform (in $N \in \mathbb{N}$) bounds on B_ℓ , $\ell = 1, \dots, 4$.

Lemma 3.3. *Let $\frac{2}{3} \leq s < 1$. Then, we have*

$$|B_1| \lesssim M^{2-3s} \int_{t_1}^{t_2} E_N^2(t) dt \quad (3.18)$$

for any $t_2 \geq t_1 \geq 0$, where the implicit constant is independent of $N, M \in \mathbb{N}$.

Proof. The bound (3.18) follows from Cauchy-Schwarz's inequality in x , Lemma 2.2, and Cauchy-Schwarz's inequality in j and k with (3.15). \square

By formally distributing the I -operator to each factor of B_2 and B_3 in (3.17), we obtain the following terms:

$$\begin{aligned} B'_2 &= -\frac{1}{N^2} \sum_{j,k=1}^N \int_{t_1}^{t_2} \int_{\mathbb{T}^2} \partial_t I v_{N,j} \\ &\quad \times \left(2(I\Psi_k)(Iv_{N,k})(Iv_{N,j}) + (Iv_{N,k})^2 I\Psi_j \right) dx dt, \\ B'_3 &= -\frac{1}{N^2} \sum_{j,k=1}^N \int_{t_1}^{t_2} \int_{\mathbb{T}^2} \partial_t I v_{N,j} \\ &\quad \times \left((I(\Psi_k^2)) I v_{N,j} + 2Iv_{N,k}(I(\Psi_k \Psi_j)) \right) dx dt. \end{aligned} \quad (3.19)$$

We first establish the following commutator estimates.

Lemma 3.4. *For $\ell = 2, 3$, let B_ℓ and B'_ℓ be as in (3.17) and (3.19), respectively. Let $\frac{2}{3} \leq s < 1$. Then, given $T \gg 1$ and small $\delta > 0$, we have*

$$|B_2 - B'_2| \lesssim M^{-s+\frac{1}{2}+\delta} V_N^{(1)}(T, \delta) \int_{t_1}^{t_2} E_N^{\frac{3}{2}}(t) dt, \quad (3.20)$$

$$|B_3 - B'_3| \lesssim M^{-\frac{s}{2}+\delta} V_N^{(1)}(T, \delta) \int_{t_1}^{t_2} E_N(t) dt \quad (3.21)$$

for any $0 \leq t_1 \leq t_2 \leq T$ and any sufficiently large $M \gg 1$, where the implicit constant is independent of $N \in \mathbb{N}$ and $M, T \gg 1$. Here, $V_N^{(1)}(T, \delta)$ is defined by

$$\begin{aligned} V_N^{(1)}(T, \delta) &= \|\Psi_j\|_{\mathcal{A}_N C_T W_x^{-\sigma_0, p_0}} + \|\Psi_j^2\|_{\mathcal{A}_N C_T W_x^{-\sigma_0, p_0}} \\ &\quad + \|\Psi_k \Psi_j\|_{\mathcal{A}_N^{(2)} C_T W_x^{-\sigma_0, p_0}}, \end{aligned} \quad (3.22)$$

where $\mathcal{A}_N B$ and $\mathcal{A}_N^{(2)} B$ are the ℓ^2 -averages defined in (1.11) and (2.2), while $\sigma_0 = \sigma_0(\delta) > 0$ and $p_0 = p_0(\delta) \gg 1$ are as in Lemma 2.3.

Proof. The first bound (3.20) follows from Lemma 2.3 and Cauchy-Schwarz's inequality, while the second bound (3.21) follows from Lemma 2.3, Cauchy-Schwarz's inequality, and (2.1). \square

Next, we estimate B_4 and B'_3 in (3.17) and (3.19), respectively.

Lemma 3.5. *Let B_4 and B'_3 be as in (3.17) and (3.19), respectively. Let $\frac{2}{3} \leq s < 1$. Then, given small $\gamma > 0$, we have*

$$|B_4| \lesssim M^\gamma V_N^{(2)}(T, \gamma) \int_{t_1}^{t_2} E_N^{\frac{1}{2}}(t) dt, \quad (3.23)$$

$$|B'_3| \lesssim M^\gamma V_N^{(2)}(T, \gamma) \int_{t_1}^{t_2} E_N^{\frac{3}{4}}(t) dt \quad (3.24)$$

for any $0 \leq t_0 \leq t \leq T$ and any sufficiently large $M \gg 1$, where the implicit constant is independent of $N \in \mathbb{N}$ and $M, T \gg 1$. Here, $V_N^{(2)}(T, \gamma)$ is defined by

$$\begin{aligned} V_N^{(2)}(T, \gamma) = & \|:\Psi_j^2:\|_{\mathcal{A}_N C_T W_x^{-\gamma, 4}} + \|:\Psi_k \Psi_j:\|_{\mathcal{A}_N^{(2)} C_T W_x^{-\gamma, 4}} \\ & + \|:\Psi_k^2 \Psi_j:\|_{\mathcal{A}_N^{(2)} C_T W_x^{-\gamma, 4}}, \end{aligned} \quad (3.25)$$

where $\mathcal{A}_N B$ and $\mathcal{A}_N^{(2)} B$ are the ℓ^2 -averages defined in (1.11) and (2.2).

Proof. The first bound (3.23) on B_4 follows from Cauchy-Schwarz's inequality in x and then in j and (2.1).

As for the second bound (3.24) on B'_3 , we first write

$$B'_3 = B'_{31} + B'_{32}, \quad (3.26)$$

where B'_{31} and B'_{32} are given by

$$\begin{aligned} B'_{31} &= -\frac{1}{N^2} \sum_{j,k=1}^N \int_{t_1}^{t_2} \int_{\mathbb{T}^2} \partial_t I v_{N,j} \cdot I(:\Psi_k^2:) \cdot I v_{N,j} dx dt, \\ B'_{32} &= -\frac{2}{N^2} \sum_{j,k=1}^N \int_{t_1}^{t_2} \int_{\mathbb{T}^2} \partial_t I v_{N,j} \cdot I v_{N,k} \cdot I(:\Psi_k \Psi_j:) dx dt. \end{aligned}$$

By Cauchy-Schwarz's inequality in j , Hölder's inequality in x , (3.15), (2.9), and (2.1), we have

$$\begin{aligned} |B'_{31}| &\leq \frac{1}{N^2} \sum_{k=1}^N \int_{t_1}^{t_2} \|\partial_t I v_{N,j}\|_{\ell_j^2 L_x^2} \|I v_{N,j}\|_{L_x^4 \ell_j^2} \|I(:\Psi_k^2:)\|_{L_x^4} dt' \\ &\lesssim M^\gamma \|:\Psi_k^2:\|_{\mathcal{A}_N C_T W_x^{-\gamma, 4}} \int_{t_1}^{t_2} E_N^{\frac{3}{4}}(t) dt. \end{aligned} \quad (3.27)$$

Similarly, we have

$$\begin{aligned} |B'_{32}| &\lesssim \frac{1}{N^2} \int_{t_1}^{t_2} \|\partial_t I v_{N,j}\|_{\ell_j^2 L_x^2} \|I v_{N,k}\|_{L_x^4 \ell_k^2} \|I(:\Psi_k \Psi_j:)\|_{\ell_{j,k}^2 L_x^4} dt' \\ &\lesssim M^\gamma \|:\Psi_k \Psi_j:\|_{\mathcal{A}_N^{(2)} C_T W_x^{-\gamma, 4}} \int_{t_1}^{t_2} E_N^{\frac{3}{4}}(t) dt. \end{aligned} \quad (3.28)$$

Hence, the second bound (3.24) follows from (3.26), (3.27), and (3.28) with (3.25). \square

Lastly, we estimate B'_2 in (3.19), where the logarithmically divergent nature of $I\Psi_j$ plays an important role.

Lemma 3.6. *Let B'_2 be as in (3.19). Then, given $s < 1$, there exists $c > 0$ such that*

$$|B'_2| \lesssim \left\{ \int_{t_1}^{t_2} \left(1 + E_N^{1+c\eta}(t) + \frac{\eta}{(t-t_1)^{\frac{1}{2}}} \right) dt \right\} \|I\Psi_j\|_{L_{[t_1, t_2], x}^{\eta-1} \mathcal{A}_N} \quad (3.29)$$

for any $t_2 \geq t_1 \geq 0$ and small $\eta > 0$, uniformly in $M, N \in \mathbb{N}$, where $L_{[t_1, t_2], x}^r \mathcal{A}_N = L_{[t_1, t_2]}^r L_x^r \mathcal{A}_N$.

Proof. The bound (3.29) follows from a straightforward modification of the proof of [23, Lemma 3.4] adapted to the current vector-valued setting. We first note that by interpolation, Minkowski's integral inequality, and Sobolev's inequality with (3.15), we have

$$\begin{aligned} \|Iv_{N,j}\|_{L_x^{\frac{4}{1-2\eta}} \mathcal{A}_N} &\leq \|Iv_{N,j}\|_{L_x^4 \mathcal{A}_N}^{1-4\eta} \|Iv_{N,j}\|_{L_x^8 \mathcal{A}_N}^{4\eta} \leq \|Iv_{N,j}\|_{L_x^4 \mathcal{A}_N}^{1-4\eta} \|Iv_{N,j}\|_{\mathcal{A}_N L_x^8}^{4\eta} \\ &\lesssim \|Iv_{N,j}\|_{L_x^4 \mathcal{A}_N}^{1-4\eta} \|Iv_{N,j}\|_{\mathcal{A}_N H_x^1}^{4\eta} \\ &\lesssim E_N^{\frac{1-4\eta}{4}} E_N^{2\eta} = E_N^{\frac{1+4\eta}{4}}, \end{aligned} \quad (3.30)$$

uniformly in small $\eta > 0$.

Write B'_2 in (3.19) as

$$B'_2 = B'_{21} + B'_{22},$$

where B'_{21} and B'_{22} are given by

$$\begin{aligned} B'_{21} &= -\frac{2}{N^2} \sum_{j,k=1}^N \int_{t_1}^{t_2} \int_{\mathbb{T}^2} \partial_t Iv_{N,j} \cdot (I\Psi_k)(Iv_{N,k})(Iv_{N,j}) dx dt, \\ B'_{22} &= -\frac{1}{N^2} \sum_{j,k=1}^N \int_{t_1}^{t_2} \int_{\mathbb{T}^2} \partial_t Iv_{N,j} \cdot (Iv_{N,k})^2 \cdot I\Psi_j dx dt. \end{aligned} \quad (3.31)$$

By Cauchy-Schwarz's inequality (in j and k), Hölder's inequality (in x), and (3.30), we have

$$\begin{aligned} |B'_{22}| &\leq \int_{t_1}^{t_2} \|\partial_t Iv_{N,j}\|_{L_x^2 \mathcal{A}_N} \|Iv_{N,k}\|_{L_x^{\frac{4}{1-2\eta}} \mathcal{A}_N}^2 \|I\Psi_j\|_{L_x^{\eta-1} \mathcal{A}_N} dt \\ &\lesssim \int_{t_1}^{t_2} E_N^{1+2\eta}(t) \|I\Psi_j(t)\|_{L_x^{\eta-1} \mathcal{A}_N} dt \\ &\lesssim \left(\int_{t_1}^{t_2} E_N^{\frac{1+2\eta}{1-\eta}}(t) dt \right)^{1-\eta} \|I\Psi_j\|_{L_{[t_1, t_2], x}^{\eta-1} \mathcal{A}_N} \\ &\lesssim \left\{ \int_{t_1}^{t_2} \left(E_N^{\frac{1+2\eta}{1-2\eta}}(t) + \frac{\eta}{(t-t_1)^{\frac{1}{2}}} \right) dt \right\} \|I\Psi_j\|_{L_{[t_1, t_2], x}^{\eta-1} \mathcal{A}_N} \\ &\lesssim \left\{ \int_{t_1}^{t_2} \left(1 + E_N^{1+c\eta}(t) + \frac{\eta}{(t-t_1)^{\frac{1}{2}}} \right) dt \right\} \|I\Psi_j\|_{L_{[t_1, t_2], x}^{\eta-1} \mathcal{A}_N}, \end{aligned}$$

uniformly in small $\eta > 0$, where the penultimate step follows from [23, the math display before (3.21)] as a consequence of Hölder's and Young's inequalities. The contribution from B'_{21} in (3.31) can be treated in an analogous manner, thus yielding (3.29). \square

3.3. Global well-posedness. In this subsection, we prove pathwise global well-posedness of HLSM_N (1.10) (Theorem 1.1 (ii)). In view of the first order expansion (1.7), Theorem 1.1 (ii) follows once we prove the following proposition.

Proposition 3.7. *Let $m > 0$ and $\frac{4}{5} < s < 1$. Given $N \in \mathbb{N}$, let $(\mathbf{v}_N^{(0)}, \mathbf{v}_N^{(1)}) = \{(v_{N,j}^{(0)}, v_{N,j}^{(1)})\}_{j=1}^N \in (\mathcal{H}^s(\mathbb{T}^2))^{\otimes N}$. Then, there exists a unique solution $(\mathbf{v}_N, \partial_t \mathbf{v}_N) = \{(v_{N,j}, \partial_t v_{N,j})\}_{j=1}^N$ in the class $(C(\mathbb{R}_+; \mathcal{H}^s(\mathbb{T}^2)))^{\otimes N}$ to (3.1) with initial data $(\mathbf{v}_N^{(0)}, \mathbf{v}_N^{(1)})$, almost surely.*

The rest of this subsection is devoted to a proof of Proposition 3.7, with a particular emphasis on uniformity in $N \in \mathbb{N}$; see, for example, Remark 3.9. Fix $\frac{4}{5} < s < 1$. Given small $\delta, \gamma > 0$ and a target time $T \gg 1$, it follows from (3.16), (3.17), (3.19), Lemmas 3.3, 3.4, 3.5, and 3.6 that

$$\begin{aligned}
& E_N(t_2) - E_N(t_1) \\
& \lesssim M^{2-3s} \int_{t_1}^{t_2} E_N^2(t) dt \\
& + \sum_{\ell=1}^2 M^{-\frac{1-\ell(1-s)}{2} + \delta} V_N^{(1)}(T, \delta) \int_{t_1}^{t_2} E_N^{\frac{\ell+1}{2}}(t) dt \\
& + M^\gamma V_N^{(2)}(T, \gamma) \int_{t_1}^{t_2} (1 + E_N^{\frac{3}{4}}(t)) dt \\
& + \left\{ \int_{t_1}^{t_2} \left(1 + E_N^{1+c\eta}(t) + \frac{\eta}{(t-t_1)^{\frac{1}{2}}} \right) dt \right\} \|I\Psi_j\|_{L_{[t_1, t_2], x}^{\eta-1} \mathcal{A}_N}
\end{aligned} \tag{3.32}$$

for any $0 \leq t_1 \leq t_2 \leq T$ and small $\eta > 0$, uniformly in $N \in \mathbb{N}$ and $M \gg 1$, where $V_N^{(1)}(T, \delta)$ and $V_N^{(2)}(T, \gamma)$ are as in (3.22) and (3.25), respectively.

Before we state a crucial lemma (Lemma 3.8) that allows us to proceed with an iterative argument, we introduce two auxiliary random variables. Given $N \in \mathbb{N}$, let R_N be as in Lemma 2.7 which controls $I\Psi_j$ on the last term in (3.32). The next random V_N controls the other contributions from the stochastic terms in (3.32). By choosing $\varepsilon = \min(\sigma_0, \gamma)$, it follows from (3.22) and (3.25) that

$$V_N^{(1)}(T, \delta) + V_N^{(2)}(T, \gamma) \lesssim \|\Psi_N\|_{\mathcal{Z}_N^\varepsilon(T)},$$

where $\mathcal{Z}_N^\varepsilon(T)$ is as in (2.17) and Ψ_N is given by

$$\Psi_N = \left\{ \Psi_j, : \Psi_k \Psi_j : , : \Psi_k^2 \Psi_j : \right\}_{j,k=1}^N.$$

Define a random variable V_N by

$$e^{V_N^{\frac{1}{3}}} = \sum_{\ell=0}^{\infty} e^{-\theta \ell} e^{\|\Psi_N\|_{\mathcal{Z}^\varepsilon(\ell)}^{\frac{1}{3}}} \tag{3.33}$$

for some $\theta > 0$. Then, by applying Lemma 2.5 (ii) and choosing $\theta > 0$ sufficiently large, we have

$$\sup_{N \in \mathbb{N}} \mathbb{E} \left[e^{V_N^{\frac{1}{3}}} \right] = \sup_{N \in \mathbb{N}} \sum_{\ell=1}^{\infty} e^{-\theta \ell} \mathbb{E} \left[e^{\|\Psi_N\|_{\mathcal{Z}^\varepsilon(\ell)}^{\frac{1}{3}}} \right] \lesssim \sum_{\ell=1}^{\infty} e^{-\theta \ell} e^{c\ell} < \infty. \tag{3.34}$$

From (3.33), we also have

$$\|\Psi_N\|_{\mathcal{Z}_N^\varepsilon(T)} \leq \left(V_N^{\frac{1}{3}} + \theta(1+T) \right)^3 \lesssim V_N + T^3. \tag{3.35}$$

The left-hand side of (3.35) plays a role of M_T in [23, (3.24)].

The following lemma is an analogue of [23, Proposition 3.5] in the current vector-valued setting.

Lemma 3.8. *Let $\frac{2}{3} < s < 1$ and $T \gg 1$. Given $N \in \mathbb{N}$, let R_N and V_N be as in Lemma 2.7 and (3.33), respectively, which are almost surely finite. Fix $\omega \in \Omega$ such that $R_N(\omega), V_N(\omega) < \infty$. Then, there exist $\alpha = \alpha(s)$, $\beta = \beta(s) > 0$, and $M_* = M_*(s) \in \mathbb{N}$, satisfying*

$$\beta < \alpha \leq 3s - 2, \tag{3.36}$$

such that if

$$E_N(t_0) \leq M^\beta \quad (3.37)$$

for some $0 \leq t_0 < T$ and $M \geq M_*$, then there exists small

$$\tau_N = \tau_N(s, M, T, R_N(\omega), V_N(\omega)) = \tau_N(s, M, T, \omega) > 0$$

such that

$$E_N(t) \leq M^\alpha$$

for any t satisfying $t_0 \leq t \leq \min(T, t_0 + \tau_N)$.

Remark 3.9. In Lemma 3.8, the dependence of τ_N on N only appears through R_N and V_N . In view of the uniform (in N) bounds (2.26) and (3.34) with Chebyshev's inequality, we see that Lemma 3.8 holds “uniformly in $N \in \mathbb{N}$ ” in the sense that the statistical properties of τ_N can be estimated uniformly in $N \in \mathbb{N}$.

Proof of Lemma 3.8. Once we have the bound (3.32), this proposition follows from repeating the proof of [23, Proposition 3.5] based on a double Gronwall-type inequality and we only indicate the basic idea. As in [23, (3.32) and (3.38)], define F_N by

$$\begin{aligned} F_N(t) &= \max_{t_0 \leq \tau \leq t} E_N(\tau) - E_N(t_0) + \max(E_N(t_0), M^\beta), \\ G_N(t) &= F_N(t) - 2C_0 R_N(t - t_0)^{\frac{1}{2}} \end{aligned} \quad (3.38)$$

for some suitable $C_0 > 0$. Then, after some manipulation on (3.32) with (3.38), we arrive at

$$G_N(t) - G_N(t_0) \lesssim (1 + V_N + R_N + T) \int_{t_0}^t G_N(t') (\log G(t') + T^2) dt' \quad (3.39)$$

for any $t_0 \leq t \leq \min(t_1, t_0 + t_*(C_0, R_N))$ such that $2C_0 R(t - t_0)^{\frac{1}{2}} \sim 1$, guaranteeing $G_N(t) \sim F_N(t)$; see [23, (3.39)]. From the differential inequality (3.39), we obtain a double exponential bound on $G_N(t)$, which, together with (3.37), implies the conclusion of this proposition. Lastly, we note that the condition (3.36) follows from [23, (3.36)] and that the condition $M \geq M_* = M_*(s)$ comes from [23, (3.45) and (3.46)]. \square

We conclude this section by presenting a proof of Proposition 3.7.

Proof of Proposition 3.7. This proposition follows from a straightforward modification of the proof of [23, Theorem 1.2]. We, however, present some details in order to demonstrate that the following iterative argument provides a uniform (in N) bound on the $\mathcal{A}_N \mathcal{H}_x^s$ -norm of a solution $\vec{\mathbf{v}}_N(t) = (\mathbf{v}_N, \partial_t \mathbf{v}_N)(t)$. See also Remark 3.10.

Let $\frac{4}{5} < s < 1$ and fix a target time $T \gg 1$. Moreover, given $N \in \mathbb{N}$, fix $\omega \in \Omega$ such that

$$R_N(\omega), V_N(\omega) < \infty. \quad (3.40)$$

Then, let the parameters α, β, M_* and τ_N be as in Lemma 3.8.

Fix $M_0 \gg 1$ (to be determined later) and define an increasing sequence $\{M_k\}_{k \in \mathbb{N}}$ by setting

$$M_k = M_0^{\sigma^k}$$

for some $\sigma \gg 1$. Under the assumption

$$2(1 - s) < \beta, \quad (3.41)$$

we can take $\sigma = \sigma(s, \alpha, \beta) \gg 1$ sufficiently large such that

$$M_{k+1}^{2(1-s)} M_k^\alpha + M_k^{2\alpha} \ll M_{k+1}^\beta, \quad (3.42)$$

We point out that the constraint $s > \frac{4}{5}$ appears from (3.36) and (3.41).

Given $N, M \in \mathbb{N}$ and $t \geq 0$, set

$$E_{N,M}(t) = E_N(I_M \mathbf{v}_N, \partial_t I_M \mathbf{v}_N)(t) \quad (3.43)$$

where the right-hand side is as in (1.13). Suppose that

$$E_{N,M_k}(t) \leq M_k^\alpha \quad (3.44)$$

for some $k \in \mathbb{N} \cup \{0\}$ and $t \geq 0$. Then, by (2.8), Minkowski's integral inequality, (2.9), Sobolev's inequality (with $s \geq \frac{1}{2}$), (3.44), and (3.42), we have

$$\begin{aligned} E_{N,M_{k+1}}(t) &\lesssim \|(I_{M_{k+1}} \mathbf{v}_N, \partial_t I_{M_{k+1}} \mathbf{v}_N)\|_{\mathcal{A}_N \mathcal{H}_x^1}^2 + \|I_{M_{k+1}} \mathbf{v}_N\|_{L_x^4 \mathcal{A}_N}^4 \\ &\lesssim M_{k+1}^{2(1-s)} \|(\mathbf{v}_N, \partial_t \mathbf{v}_N)\|_{\mathcal{A}_N \mathcal{H}_x^s}^2 + \|I_{M_{k+1}} \mathbf{v}_N\|_{\mathcal{A}_N L_x^4}^4 \\ &\lesssim M_{k+1}^{2(1-s)} \|(I_{M_k} \mathbf{v}_N, \partial_t I_{M_k} \mathbf{v}_N)\|_{\mathcal{A}_N \mathcal{H}_x^1}^2 + \|\mathbf{v}_N\|_{\mathcal{A}_N H_x^{\frac{1}{2}}}^4 \\ &\lesssim M_{k+1}^{2(1-s)} M_k^\alpha + \|I_{M_k} \mathbf{v}_N\|_{\mathcal{A}_N H_x^1}^4 \\ &\lesssim M_{k+1}^{2(1-s)} M_k^\alpha + M_k^{2\alpha} \\ &\ll M_{k+1}^\beta, \end{aligned} \quad (3.45)$$

where, at the fourth inequality, we use the fact that $s \geq \frac{1}{2}$ (in applying (2.8)).

Given $(\mathbf{v}_N^{(0)}, \mathbf{v}_N^{(1)}) \in \mathcal{A}_N \mathcal{H}_x^s(\mathbb{T}^2)$ and $\omega \in \Omega$, satisfying (3.40), let $\vec{\mathbf{v}}_N(\omega) = (\mathbf{v}_N(\omega), \partial_t \mathbf{v}_N(\omega))$ be the local-in-time solution to (3.1) with $\vec{\mathbf{v}}_N(\omega)|_{t=0} = (\mathbf{v}_N^{(0)}, \mathbf{v}_N^{(1)})$. In view of the blowup alternative (3.4), the desired global well-posedness of (3.1) on the time interval $[0, T]$ (for the fixed $\omega \in \Omega$) follows once we prove

$$\sup_{0 \leq t \leq T} \|\vec{\mathbf{v}}_N(t; \omega)\|_{\mathcal{A}_N \mathcal{H}_x^s} \leq C(\omega, T) < \infty. \quad (3.46)$$

In the following, we prove (3.46) by implementing an iterative argument based on Lemma 3.8.

Choose $M_0 = M_0(\mathbf{v}_N^{(0)}, \mathbf{v}_N^{(1)}, s, T) \gg 1$ such that

$$E_{N,M_0}(0) \leq M_0^\beta.$$

Then, it follows from Lemma 3.8 that

$$\sup_{0 \leq t \leq \tau_N} E_{N,M_0}(t) \leq M_0^\alpha,$$

satisfying (3.44) at time τ_N . From (3.43), (1.13), and (2.8), this in particular implies

$$\sup_{0 \leq t \leq \tau_N} \|\vec{\mathbf{v}}_N(t; \omega)\|_{\mathcal{A}_N \mathcal{H}_x^s}^2 \lesssim M_0^\alpha.$$

Moreover, from (3.45), we obtain

$$E_{N,M_1}(\tau_N) \leq M_1^\beta.$$

By applying Lemma 3.8 once again, we have

$$\sup_{\tau_N \leq t \leq 2\tau_N} E_{N,M_1}(t) \leq M_1^\alpha,$$

satisfying (3.44) at time $2\tau_N$, from which we obtain

$$\sup_{\tau_N \leq t \leq 2\tau_N} \|\vec{\mathbf{v}}_N(t; \omega)\|_{\mathcal{A}_N \mathcal{H}_x^s}^2 \lesssim M_1^\alpha$$

and

$$E_{N, M_2}(2\tau_N) \leq M_2^\beta.$$

By iterating this argument $\lceil \frac{T}{\tau_N} \rceil + 1$ times, we obtain the following bound:

$$\sup_{\ell\tau_N \leq t \leq (\ell+1)\tau_N} \|\vec{\mathbf{v}}_N(t; \omega)\|_{\mathcal{A}_N \mathcal{H}_x^s}^2 \lesssim M_\ell^\alpha$$

for $\ell = 0, \dots, \lceil \frac{T}{\tau_N} \rceil + 1$, yielding (3.46). This proves existence of the unique solution \mathbf{v}_N to (3.1) on the time interval $[0, T]$ (for the fixed $\omega \in \Omega$, satisfying (3.40)). Since the choice of $T \gg 1$ was arbitrary, we conclude almost sure global well-posedness of (3.1). \square

Remark 3.10. As pointed out in [23, Remark 3.7], the iterative argument presented above yields the following uniform (in N) double exponential bound:

$$\|\vec{\mathbf{v}}_N(t)\|_{\mathcal{A}_N \mathcal{H}_x^s} \leq C \exp \left(c \log (2 + \|\vec{\mathbf{v}}_N(0)\|_{\mathcal{A}_N \mathcal{H}_x^s}) \cdot e^{C(\omega)t^2} \right)$$

for any $t \geq 0$. See [23, Remark 3.7] for details.

4. WELL-POSEDNESS OF THE MEAN-FIELD SdNLW

In this section, we present a proof of Theorem 1.2 on local and global well-posedness of the mean-field SdNLW (1.20). By the first order expansion (1.19), we study the following equation (see (1.18)) satisfied by the residual part $v = u - \Phi$, where we drop the superscript j for notational simplicity:

$$\begin{cases} (\partial_t^2 + \partial_t + m - \Delta)v = -\mathbb{E}[v^2]v - 2\mathbb{E}[\Psi v]v - \mathbb{E}[v^2]\Psi - 2\mathbb{E}[v\Psi]\Psi \\ (v, \partial_t v)|_{t=0} = (v^{(0)}, v^{(1)}). \end{cases} \quad (4.1)$$

As pointed out in Section 1, there are no terms involving $\mathbb{E}[:\Psi^2:]$ since $\mathbb{E}[:\Psi^2:] = 0$. We note that there are certain similarities between (4.1) and (3.1) after replacing the empirical average $\frac{1}{N} \sum_{1 \leq k \leq N}$ in (3.1) with the expectation \mathbb{E} , leading to computations similar to those presented in Section 3. There are, however, fundamental differences between the mean-field SdNLW (4.1) and the SdNLW system (3.1), in particular on the treatment of the following terms:

$$\mathbb{E}[v\Psi]\Psi \quad \text{and} \quad \frac{1}{N} \sum_{k=1}^N v_{N,k} : \Psi_k \Psi_j :,$$

where the former term does *not* have an explicit renormalization; see (4.6) below. See also the proofs of Lemmas 4.4, 4.5, 4.6, and 4.7, where we apply a similar idea.

In Subsection 4.1, we prove local well-posedness of (4.1). In Subsection 4.2, we briefly discuss global well-posedness of (4.1).

4.1. Local well-posedness. In this subsection, we prove the following proposition on local well-posedness of the mean-field SdNLW (4.1) (Theorem 1.2 (i)) whose proof follows closely that of Proposition 3.1 but with one twist due to the term $\mathbb{E}[v\Psi]\Psi$; see (4.4) below.

Proposition 4.1. *Let $m > 0$ and $\frac{1}{2} \leq s < 1$. Given $2 \leq p < \infty$, let $(v^{(0)}, v^{(1)}) \in L^p(\Omega; \mathcal{H}^s(\mathbb{T}^2))$. Then, given any $t_0 \in \mathbb{R}_+$, there exists a unique solution $(v, \partial_t v) \in L^p(\Omega; C([t_0, t_0 + \tau]; \mathcal{H}^s(\mathbb{T}^2)))$ to (4.1) with $(v, \partial_t v)|_{t=t_0} = (v^{(0)}, v^{(1)})$, where the local existence time $\tau > 0$ depends on $\|(v^{(0)}, v^{(1)})\|_{L_\omega^p \mathcal{H}_x^s}$ and Ψ .*

Remark 4.2. (i) As pointed out in Remark 3.2 (i), it is possible to lower the regularity threshold by using the Strichartz estimates. We, however, restrict our attention to the range $\frac{1}{2} \leq s < 1$, which guarantees unconditional uniqueness of solutions (namely, uniqueness in the entire class $L^p(\Omega; C(\tau; \mathcal{H}^s(\mathbb{T}^2)))$), since it plays a key role in Section 6.

(ii) Let $T \geq 1$. Let $(v, \partial_t v)$ be the local-in-time solution to (4.1) constructed in Proposition 4.1. Then, by denoting by T_* its maximal existence time, the following blowup alternative holds:

$$T_* = T \quad \text{or} \quad \lim_{t \rightarrow T_*^-} \|(v, \partial_t v)(t)\|_{L_\omega^p \mathcal{H}_x^s} = \infty. \quad (4.2)$$

Proof of Proposition 4.1. Without loss of generality, we assume that $t_0 = 0$. By writing (4.1) in the Duhamel formulation, we have

$$\begin{aligned} v(t) = & S(t)(v^{(0)}, v^{(1)}) - \mathcal{I}(\mathbb{E}[v^2]v)(t) - 2\mathcal{I}(\mathbb{E}[\Psi v]v)(t) \\ & - \mathcal{I}(\mathbb{E}[v^2]\Psi)(t) - 2\mathcal{I}(\mathbb{E}[v\Psi]\Psi)(t), \end{aligned} \quad (4.3)$$

where $S(t)$ and \mathcal{I} are as in (2.5) and (2.6), respectively.

Let $2 \leq p < \infty$. Fix $0 < \tau \leq 1$ and small $\varepsilon > 0$ such that $s \leq 1 - \varepsilon$. Let us first consider the last term on the right-hand side of (4.3), involving $\mathbb{E}[v\Psi]\Psi$, which shows a new aspect as compared to the hyperbolic $O(N)$ linear sigma model (3.1) studied in Subsection 3.1. In order to handle this term, we write

$$\mathbb{E}[v\Psi]\Psi(\omega) = \mathbb{E}[v'\Psi']\Psi(\omega) = \int_{\Omega} v'(\omega')\Psi'(\omega')\Psi(\omega)\mathbb{P}(d\omega'), \quad (4.4)$$

where $(v'\Psi')$ is an independent copy of (v, Ψ) . In particular, the product $\Psi'\Psi$ makes sense *without* renormalization. As a consequence, it follows from Lemma 2.5 that

$$K_p = \|\Psi\|_{L_\omega^p C_1 W_x^{-\varepsilon, \infty}} + \|\Psi'\Psi\|_{L_\omega^p C_1 W_x^{-\varepsilon, \infty}} < \infty. \quad (4.5)$$

Recall the notation (3.7). We apply (4.4) and proceed as in (3.10) with Sobolev's inequality and Lemma 2.1 (ii). Then, from the independence of Ψ' and Ψ with $\text{Law}(v') = \text{Law}(v)$ and (4.5), we have

$$\begin{aligned} \|\vec{\mathcal{I}}(\mathbb{E}[v\Psi]\Psi)\|_{L_\omega^p C_\tau \mathcal{H}_x^s} & \lesssim \tau \left\| \|v'(\omega')\Psi'(\omega')\Psi(\omega)\|_{C_\tau W_x^{-\varepsilon, \frac{2}{2-s-\varepsilon}}} \right\|_{L_\omega^p L_{\omega'}^1} \\ & \lesssim \tau \left\| \|\Psi'(\omega')\Psi(\omega)\|_{C_1 W_x^{-\varepsilon, \infty}} \|v'(\omega')\|_{C_\tau W_x^{\varepsilon, \frac{2}{2-s-\varepsilon}}} \right\|_{L_\omega^p L_{\omega'}^1} \\ & \leq \tau \|\Psi'(\omega')\Psi(\omega)\|_{L_{\omega, \omega'}^p C_1 W_x^{-\varepsilon, \infty}} \|v'\|_{L_{\omega'}^2 C_\tau W_x^{\varepsilon, \frac{2}{2-s-\varepsilon}}} \\ & = \tau \|\Psi'(\omega)\Psi(\omega)\|_{L_\omega^p C_1 W_x^{-\varepsilon, \infty}} \|v\|_{L_\omega^2 C_\tau W_x^{\varepsilon, \frac{2}{2-s-\varepsilon}}} \\ & \lesssim \tau K_p \|v\|_{L_\omega^p C_\tau \mathcal{H}_x^s}, \end{aligned} \quad (4.6)$$

provided that $\varepsilon \leq s \leq 1 - \varepsilon$, where, at the third inequality, we used the fact that $p \geq 2$.

Let us now handle the remaining terms. Proceeding as in (3.6), we have

$$\begin{aligned}
\|\vec{\mathcal{I}}(\mathbb{E}[v^2]v)\|_{L_\omega^p C_\tau \mathcal{H}_x^s} &\lesssim \tau \|\mathbb{E}[v^2]v\|_{L_\omega^p C_\tau H_x^{s-1}} \\
&\lesssim \tau \left\| \mathbb{E} \left[\|v\|_{C_\tau L_x^{\frac{6}{2-s}}}^2 \right] \|v\|_{C_\tau L_x^{\frac{6}{2-s}}} \right\|_{L_\omega^p} \\
&\lesssim \tau \|v\|_{L_\omega^p C_\tau H_x^s}^3,
\end{aligned} \tag{4.7}$$

provided that $\frac{1}{2} \leq s (< 1)$. Proceeding as in (3.8) with Lemma 2.1 (ii) (but in a slightly different order due to the presence of the expectation), we have

$$\begin{aligned}
\|\vec{\mathcal{I}}(\mathbb{E}[\Psi v]v)\|_{L_\omega^p C_\tau \mathcal{H}_x^s} &\lesssim \tau \|\mathbb{E}[\Psi v]v\|_{L_\omega^p C_\tau W_x^{-\varepsilon, \frac{2}{2-s-\varepsilon}}} \\
&\lesssim \tau \left\| \mathbb{E} \left[\|\Psi v\|_{C_\tau W_x^{-\varepsilon, \frac{4}{2-s-\varepsilon}}} \right] \|v\|_{C_1 W_x^{\varepsilon, \frac{4}{2-s-\varepsilon}}} \right\|_{L_\omega^p} \\
&\lesssim \tau \|\Psi\|_{L_\omega^p C_1 W_x^{-\varepsilon, \infty}} \|v\|_{L_\omega^p C_\tau H_x^{\frac{s}{2} + \frac{3}{2}\varepsilon}}^2 \\
&\lesssim \tau K_p \|v\|_{L_\omega^p C_\tau H_x^s}^2,
\end{aligned} \tag{4.8}$$

provided that $3\varepsilon \leq s (< 1)$. Similarly, we have

$$\begin{aligned}
\|\vec{\mathcal{I}}(\mathbb{E}[v^2]\Psi)\|_{L_\omega^p C_\tau \mathcal{H}_x^s} &\lesssim \tau \|\mathbb{E}[v^2]\Psi\|_{L_\omega^p C_\tau W_x^{-\varepsilon, \frac{2}{2-s-\varepsilon}}} \\
&\lesssim \tau \left\| \mathbb{E} \left[\|v^2\|_{C_\tau W_x^{\varepsilon, \frac{2}{2-s-\varepsilon}}} \right] \|\Psi\|_{C_1 W_x^{-\varepsilon, \infty}} \right\|_{L_\omega^p} \\
&\lesssim \tau \|\Psi\|_{L_\omega^p C_1 W_x^{-\varepsilon, \infty}} \|v\|_{L_\omega^p C_\tau H_x^{\frac{s}{2} + \frac{3}{2}\varepsilon}}^2 \\
&\lesssim \tau K_p \|v\|_{L_\omega^p C_\tau H_x^s}^2,
\end{aligned} \tag{4.9}$$

provided that $3\varepsilon \leq s (< 1)$.

Denote the right-hand side of (4.3) by $\Gamma(\vec{v}) = \Gamma(v, \partial_t v)$, and set $\vec{\Gamma}(\vec{v}) = (\Gamma(\vec{v}), \partial_t \Gamma(\vec{v}))$. Then, putting (4.3), (4.6), (4.7), (4.8), and (4.9) together, we obtain

$$\begin{aligned}
\|\vec{\Gamma}(\vec{v})\|_{L_\omega^p C_\tau \mathcal{H}_x^s} &\leq C_1 \|(v^{(0)}, v^{(1)})\|_{L_\omega^p \mathcal{H}_x^s} + C_2 \tau \|\vec{v}\|_{L_\omega^p C_\tau \mathcal{H}_x^s}^3 \\
&\quad + C_3 \tau K_p \left(1 + \|\vec{v}\|_{L_\omega^p C_\tau \mathcal{H}_x^s} \right) \|\vec{v}\|_{L_\omega^p C_\tau \mathcal{H}_x^s}
\end{aligned}$$

for any $\vec{v} \in L^p(\Omega; C([0, \tau]; \mathcal{H}^s(\mathbb{T}^2)))$, where K_p is as in (4.5). A similar computation yields the following difference estimate:

$$\begin{aligned}
&\|\vec{\Gamma}(\vec{v}) - \vec{\Gamma}(\vec{w})\|_{L_\omega^p C_\tau \mathcal{H}_x^s} \\
&\lesssim \tau \left(\|\vec{v}\|_{L_\omega^p C_\tau \mathcal{H}_x^s}^2 + \|\vec{w}\|_{L_\omega^p C_\tau \mathcal{H}_x^s}^2 \right) \|\vec{v} - \vec{w}\|_{L_\omega^p C_\tau \mathcal{H}_x^s} \\
&\quad + \tau K_p \left(1 + \|\vec{v}\|_{L_\omega^p C_\tau \mathcal{H}_x^s} + \|\vec{w}\|_{L_\omega^p C_\tau \mathcal{H}_x^s} \right) \|\vec{v} - \vec{w}\|_{L_\omega^p C_\tau \mathcal{H}_x^s}
\end{aligned}$$

for any $\vec{v}, \vec{w} \in L^p(\Omega; C([0, \tau]; \mathcal{H}^s(\mathbb{T}^2)))$. Therefore, by choosing

$$\tau = \tau \left(\|(v^{(0)}, v^{(1)})\|_{L_\omega^p \mathcal{H}_x^s}, K_p \right) > 0$$

sufficiently small, we conclude that $\vec{\Gamma}$ is a contraction in the (closed) ball

$$B_R \subset L^p(\Omega; C([0, \tau]; \mathcal{H}^s(\mathbb{T}^2)))$$

of radius $R = 2C_1 \|(v^{(0)}, v^{(1)})\|_{L_\omega^p \mathcal{H}_x^s} + 1$. At this point, the uniqueness holds only in the ball B_R but by a standard continuity argument, we can extend the uniqueness to hold in the entire class $L^p(\Omega; C([0, \tau]; \mathcal{H}^s(\mathbb{T}^2)))$. We omit details. This concludes the proof of Proposition 4.1. \square

4.2. Global well-posedness. In this subsection, we prove global well-posedness of the mean-field SdNLW (1.20) (Theorem 1.2 (ii)). As in Subsection 3.3, our main strategy is to adapt the hybrid I -method introduced in [23] to the current mean-field setting. We note that, while the analysis in [23] and Section 3 is pathwise, our analysis presented below is carried out under the $L^2(\Omega)$ -norm (i.e. not pathwise).

Fix a target time $T \gg 1$. In view of the blowup alternative (4.2) (with $p = 2$) and (2.8), existence of the unique solution $(v, \partial_t v)$ to (4.1) on the time interval $[0, T]$ follows once we show the following bound:

$$\sup_{0 \leq t \leq T} \|(Iv, \partial_t Iv)(t)\|_{L_\omega^2 \mathcal{H}_x^1} \leq C(T)$$

for some finite constant $C(T) > 0$, where $(Iv, \partial_t Iv)$ is a solution to the following I -mean-field SdNLW:

$$\begin{aligned} & (\partial_t^2 + \partial_t + m - \Delta)Iv \\ &= -I(\mathbb{E}[v^2]v) - 2I(\mathbb{E}[\Psi v]v) - I(\mathbb{E}[v^2]\Psi) - 2I(\mathbb{E}[v\Psi]\Psi). \end{aligned} \quad (4.10)$$

For this purpose, define the modified energy $\mathcal{E}(Iv, \partial_t Iv)$:

$$\begin{aligned} \mathcal{E}(Iv, \partial_t Iv) &= \frac{1}{2} \mathbb{E} \left[\int_{\mathbb{T}^2} |\nabla Iv|^2 + m(Iv)^2 + (\partial_t Iv)^2 dx \right] + \frac{1}{4} \int_{\mathbb{T}^2} (\mathbb{E}[(Iv)^2])^2 dx \\ &= \frac{1}{2} \int_{\mathbb{T}^2} \|\nabla Iv\|_{L_\omega^2}^2 + m\|Iv\|_{L_\omega^2}^2 + \|\partial_t Iv\|_{L_\omega^2}^2 dx + \frac{1}{4} \int_{\mathbb{T}^2} \|Iv\|_{L_\omega^2}^4 dx. \end{aligned} \quad (4.11)$$

where $\mathcal{E}(v, \partial_t v)$ is as in (1.22). For notational simplicity, we set

$$\mathcal{E}(t) = \mathcal{E}(Iv, \partial_t Iv)(t).$$

As in Subsection 3.3, Theorem 1.2 (ii) on global well-posedness of the mean-field SdNLW (1.20) follows from the following proposition; compare (4.12) with (3.32).

Proposition 4.3. (i) *Let $\frac{2}{3} \leq s < 1$. Then, given small $\delta, \gamma > 0$ and $T \gg 1$, we have*

$$\begin{aligned} \mathcal{E}(t_2) - \mathcal{E}(t_1) &\lesssim M^{2-3s} \int_{t_1}^{t_2} \mathcal{E}^2(t) dt \\ &\quad + M^{-s+\frac{1}{2}+\delta} \int_{t_1}^{t_2} \mathcal{E}^{\frac{3}{2}}(t) dt \cdot \|\Psi\|_{L_\omega^2 C_T W_x^{-\sigma_0, p_0}} \\ &\quad + M^{-\frac{s}{2}+\delta} \int_{t_1}^{t_2} \mathcal{E}(t) dt \cdot \|\Psi' \Psi\|_{L_\omega^2 C_T W_x^{-\sigma_0, p_0}} \\ &\quad + M^\gamma \int_{t_1}^{t_2} \mathcal{E}^{\frac{3}{4}}(t) dt \cdot \|\Psi' \Psi\|_{L_\omega^2 C_T W_x^{-\gamma, \infty}} \\ &\quad + \left\{ \int_{t_1}^{t_2} \left(1 + \mathcal{E}^{1+c\eta}(t) + \frac{\eta}{(t-t_1)^{\frac{1}{2}}} \right) dt \right\} \|I\Psi\|_{L_{[t_1, t_2], x}^{\eta-1} L_\omega^2} \end{aligned} \quad (4.12)$$

for any $0 \leq t_1 \leq t_2 \leq T$ and small $\eta > 0$, uniformly in $N \in \mathbb{N}$ and $M, T \gg 1$, where $\sigma_0 = \sigma_0(\delta) > 0$ and $p_0 = p_0(\delta) \gg 1$ are as in Lemma 2.3. Here, Ψ' denotes an independent copy of Ψ .

(ii) *Let $\frac{4}{5} < s < 1$. Then, given $(v^{(0)}, v^{(1)}) \in \mathcal{H}^s(\mathbb{T}^2)$, there exists a unique global solution $(v, \partial_t v) \in C(\mathbb{R}_+; \mathcal{H}^s(\mathbb{T}^2))$ to (4.1) with $(v, \partial_t v)|_{t=0} = (v^{(0)}, v^{(1)})$.*

Proof. (i) Let $t_2 \geq t_1 \geq 0$. Then, from (4.11) and (4.10), we have

$$\begin{aligned}\mathcal{E}(t_2) - \mathcal{E}(t_1) &= \sum_{\ell=1}^3 \mathcal{B}_\ell - \mathbb{E} \left[\int_{t_1}^{t_2} \int_{\mathbb{T}^2} (\partial_t I v)^2 dx dt \right] \\ &\leq \sum_{\ell=1}^3 \mathcal{B}_\ell,\end{aligned}$$

where $\mathcal{B}_\ell = \mathcal{B}_\ell(N)$, $\ell = 1, \dots, 3$, are given by

$$\begin{aligned}\mathcal{B}_1 &= \mathbb{E} \left[\int_{t_1}^{t_2} \int_{\mathbb{T}^2} (\partial_t I v) \left(\mathbb{E}[(I v)^2] I v - I(\mathbb{E}[v^2] v) \right) dx dt \right], \\ \mathcal{B}_2 &= -\mathbb{E} \left[\int_{t_1}^{t_2} \int_{\mathbb{T}^2} (\partial_t I v) \left(2I(\mathbb{E}[\Psi v] v) + I(\mathbb{E}[v^2] \Psi) \right) dx dt \right], \\ \mathcal{B}_3 &= -2\mathbb{E} \left[\int_{t_1}^{t_2} \int_{\mathbb{T}^2} (\partial_t I v) I(\mathbb{E}[v \Psi] \Psi) dx dt' \right].\end{aligned}\tag{4.13}$$

Then, the energy growth bound (4.12) follows from Lemmas 4.4, 4.5, 4.6, and 4.7 presented below.

(ii) The energy growth bound (4.12) yields an analogue of Lemma 3.8, which in turn allows us to proceed with an iterative argument as in the proof of Proposition 3.7 presented in Subsection 3.3. We omit details. \square

The remaining part of this section is devoted to estimating \mathcal{B}_ℓ , $\ell = 1, \dots, 3$, which closely follows the argument presented in Subsection 3.2 but with modification coming from the viewpoint (4.4); more precisely, as in (4.4), we regard the outer expectation as an integration over $\omega \in \Omega$, while the inner expectation is over $\omega' \in \Omega$. Moreover, as in Subsection 3.2, we often suppress t -dependence for notational simplicity in the following.

The following lemma establishes the commutator estimate on the cubic term in v .

Lemma 4.4. *Let $\frac{2}{3} \leq s < 1$. Then, we have*

$$|\mathcal{B}_1| \lesssim M^{2-3s} \int_{t_1}^{t_2} \mathcal{E}^2(t) dt\tag{4.14}$$

for any $t_2 \geq t_1 \geq 0$, where the implicit constant is independent of $M \in \mathbb{N}$.

Proof. By Cauchy-Schwarz's and Minkowski's integral inequalities, we have

$$\begin{aligned}&\left\| (\partial_t I v) \left(\mathbb{E}[(I v)^2] I v - I(\mathbb{E}[v^2] v) \right) \right\|_{L_{\omega, x}^1} \\ &\leq \|\partial_t I v\|_{L_{\omega, x}^2} \left\| \mathbb{E}[(I v)^2] I v - I(\mathbb{E}[v^2] v) \right\|_{L_{\omega, x}^2} \\ &\lesssim \mathcal{E}^{\frac{1}{2}} \left\| \mathbb{E}[(I v(\omega'))^2] I v(\omega) - I(v^2(\omega') v(\omega)) \right\|_{L_x^2} \Big\|_{L_{\omega}^2 L_{\omega'}^1}.\end{aligned}\tag{4.15}$$

Then, the bound (4.14) follows from (4.15), Lemma 2.2, and (4.11). \square

By formally distributing the I -operator to each factor of \mathcal{B}_2 and \mathcal{B}_3 in (4.13), we obtain the following term:

$$\begin{aligned}\mathcal{B}'_2 &= -\mathbb{E} \left[\int_{t_1}^{t_2} \int_{\mathbb{T}^2} (\partial_t I v) \left(2\mathbb{E}[(I\Psi)(Iv)]Iv + \mathbb{E}[(Iv)^2]I\Psi \right) dx dt \right], \\ \mathcal{B}'_3 &= -2\mathbb{E}_\omega \left[\int_{t_1}^{t_2} \int_{\mathbb{T}^2} (\partial_t I v(\omega)) \mathbb{E}_{\omega'} \left[(Iv'(\omega'))(I(\Psi'(\omega')\Psi(\omega))) \right] dx dt \right],\end{aligned}\tag{4.16}$$

where (Ψ', v') is an independent copy of (Ψ, v) as in (4.4). We first establish the following commutator estimates.

Lemma 4.5. *For $\ell = 2, 3$, let \mathcal{B}_ℓ and \mathcal{B}'_ℓ be as in (4.13) and (4.16), respectively. Let $\frac{2}{3} \leq s < 1$. Then, given $T \gg 1$ and small $\delta > 0$, we have*

$$|\mathcal{B}_2 - \mathcal{B}'_2| \lesssim M^{-s+\frac{1}{2}+\delta} \int_{t_1}^{t_2} \mathcal{E}^{\frac{3}{2}}(t) dt \cdot \|\Psi\|_{L_\omega^2 C_T W_x^{-\sigma_0, p_0}},\tag{4.17}$$

$$|\mathcal{B}_3 - \mathcal{B}'_3| \lesssim M^{-\frac{s}{2}+\delta} \int_{t_1}^{t_2} \mathcal{E}(t) dt \cdot \|\Psi'\Psi\|_{L_\omega^2 C_T W_x^{-\sigma_0, p_0}}\tag{4.18}$$

for any $0 \leq t_1 \leq t_2 \leq T$, uniformly in $N \in \mathbb{N}$ and $M, T \gg 1$, where $\sigma_0 = \sigma_0(\delta) > 0$ and $p_0 = p_0(\delta) \gg 1$ are as in Lemma 2.3. Here, Ψ' denotes an independent copy of Ψ .

Proof. Proceeding as in (4.16), we have

$$\begin{aligned}& \left\| \mathbb{E}[(I\Psi)(Iv)]Iv - I(\mathbb{E}[\Psi v])v \right\|_{L_{\omega, x}^2} \\ & \leq \left\| \left\| (I\Psi(\omega'))(Iv(\omega'))(Iv(\omega)) - I(\Psi(\omega')v(\omega')v(\omega)) \right\|_{L_x^2} \right\|_{L_\omega^2 L_{\omega'}^1}.\end{aligned}\tag{4.19}$$

Then, the bound (4.17) follows from Cauchy-Schwarz's inequality (in ω and x), (4.19), and Lemma 2.3 with (4.11). The second bound (4.18) follows in a similar manner. \square

We conclude this section by estimating \mathcal{B}'_3 and \mathcal{B}'_2 in (4.16).

Lemma 4.6. *Let \mathcal{B}'_3 be as in (4.16). Let $\frac{2}{3} \leq s < 1$. Then, given small $\gamma > 0$, we have*

$$|\mathcal{B}'_3| \lesssim M^\gamma \int_{t_1}^{t_2} \mathcal{E}^{\frac{3}{4}}(t) dt \cdot \|\Psi'\Psi\|_{L_\omega^2 C_T W_x^{-\gamma, \infty}}\tag{4.20}$$

for any $0 \leq t_0 \leq t \leq T$, uniformly in $N \in \mathbb{N}$ and $M, T \gg 1$. Here, Ψ' denotes an independent copy of Ψ .

Proof. From the independence of Ψ and Ψ' , we have

$$\begin{aligned}& \left\| (\partial_t I v(\omega))(Iv'(\omega'))(I(\Psi'(\omega')\Psi(\omega))) \right\|_{L_{\omega, \omega'}^1} \\ & \leq \|\partial_t I v\|_{L_\omega^2} \|Iv\|_{L_\omega^2} \|I(\Psi'(\omega')\Psi(\omega))\|_{L_{\omega, \omega'}^2} \\ & = \|\partial_t I v\|_{L_\omega^2} \|Iv\|_{L_\omega^2} \|I(\Psi'\Psi)\|_{L_\omega^2}.\end{aligned}\tag{4.21}$$

Then, the bound (4.20) follows from proceeding as in the proof of (3.24) in Lemma 3.5 with (4.21). \square

Lemma 4.7. *Let \mathcal{B}'_2 be as in (4.16). Then, given $s < 1$, there exists $c > 0$ such that*

$$|\mathcal{B}'_2| \lesssim \left\{ \int_{t_1}^{t_2} \left(1 + \mathcal{E}^{1+c\eta}(t) + \frac{\eta}{(t-t_1)^{\frac{1}{2}}} \right) dt \right\} \|I\Psi\|_{L_{[t_1, t_2], x}^{\eta-1} L_\omega^2}\tag{4.22}$$

for any $t_2 \geq t_1 \geq 0$ and small $\eta > 0$, uniformly in $M, N \in \mathbb{N}$, where $L_{[t_1, t_2], x}^r = L_{[t_1, t_2]}^r L_x^r$.

Proof. The bound (4.22) follows from a slight modification straightforward modification of the proofs of [23, Lemma 3.4] and Lemma 3.6, and thus we only indicate required modifications.

By replacing the \mathcal{A}_N -norm in (3.30) with the $L^2(\Omega)$ -norm, we have

$$\begin{aligned} \|Iv\|_{L_x^{\frac{4}{1-2\eta}} L_\omega^2} &\lesssim \|Iv\|_{L_x^4 L_\omega^2}^{1-4\eta} \|Iv\|_{L_\omega^2 H_x^1}^{4\eta} \\ &\lesssim \mathcal{E}^{\frac{1+4\eta}{4}}, \end{aligned} \quad (4.23)$$

uniformly in small $\eta > 0$. Moreover, by Hölder's inequality, we have

$$\begin{aligned} &\left\| (\partial_t Iv(\omega)) \left(2(I\Psi(\omega'))(Iv(\omega'))(Iv(\omega)) + (Iv(\omega'))^2(I\Psi(\omega)) \right) \right\|_{L_{\omega, \omega'}^1 L_x^1} \\ &\lesssim \|\partial_t Iv\|_{L_x^2 L_\omega^2} \|Iv\|_{L_x^{\frac{4}{1-2\eta}} L_\omega^2}^2 \|I\Psi\|_{L_x^{\eta^{-1}} L_\omega^2}. \end{aligned} \quad (4.24)$$

Then, the bound (4.22) follows from repeating the rest of the argument in the proof of Lemma 3.6 with (4.23) and (4.24). \square

5. MEAN-FIELD LIMIT OF THE HYPERBOLIC $O(N)$ LINEAR SIGMA MODEL

In this section, we present a proof of Theorem 1.4 on convergence of HL SM_N (1.10) to the mean-field SdNLW (1.20). In Subsection 5.1, we present several law-of-large-numbers type lemmas. In Subsection 5.2, we then prove Theorem 1.4 (i). Lastly, under a higher moment assumption, we establish a convergence rate of (1.10) to the mean-field limit (Theorem 1.4 (ii)) in Subsection 5.3.

5.1. Law of large numbers. In this subsection, we present three law-of-large-numbers type lemmas. While the first lemma (Lemma 5.1) provides a convergence rate of the purely stochastic terms, the second and third lemmas (Lemmas 5.2 and 5.4) do not provide any rate under the second moment assumption. Under a higher moment assumption, they can be upgraded to versions with a decay rate; see Lemmas 5.5 and 5.6.

Lemma 5.1. *Given $\ell = 2, 3$ and $j, k \in \mathbb{N}$, let $Z_{k,j}^{(\ell)}$ be as in (2.19) or (2.20). Namely,*

$$Z_{k,j}^{(2)} = : \Psi_k \Psi_j : \quad \text{and} \quad Z_{k,j}^{(3)} = : \Psi_k^2 \Psi_j : \quad (5.1)$$

or

$$Z_{k,j}^{(2)} = : \Phi_k \Phi_j : \quad \text{and} \quad Z_{k,j}^{(3)} = : \Phi_k^2 \Phi_j : . \quad (5.2)$$

Then, given $\varepsilon > 0$, we have

$$\left\| \frac{1}{N} \sum_{k=1}^N Z_{k,k}^{(2)} \right\|_{L_\omega^p L_T^2 W_x^{-\varepsilon, \infty}} \lesssim_T \frac{p}{N^{\frac{1}{2}}}, \quad (5.3)$$

$$\sup_{j \in \mathbb{N}} \left\| \frac{1}{N} \sum_{k=1}^N Z_{k,j}^{(3)} \right\|_{L_\omega^p L_T^2 W_x^{-\varepsilon, \infty}} \lesssim_T \frac{p^{\frac{3}{2}}}{N^{\frac{1}{2}}}, \quad (5.4)$$

$$\left\| \frac{1}{N} \sum_{k=1}^N Z_{k,j}^{(3)} \right\|_{L_\omega^p \mathcal{A}_{N,j} L_T^2 W_x^{-\varepsilon, \infty}} \lesssim_T \frac{p^{\frac{3}{2}}}{N^{\frac{1}{2}}}, \quad (5.5)$$

uniformly in finite $p \geq 1$ and $N \in \mathbb{N}$, where $\mathcal{A}_N B$ is as in (1.11). As a consequence, given any $\varepsilon_0 > 0$, the following almost sure convergence holds; for each fixed $T > 0$, we have

$$\begin{aligned} N^{\frac{1}{2}-\varepsilon_0} \left\| \frac{1}{N} \sum_{k=1}^N Z_{k,k}^{(2)} \right\|_{L_T^2 W_x^{-\varepsilon, \infty}} &\longrightarrow 0, \\ \text{for each fixed } j \in \mathbb{N}: \quad N^{\frac{1}{2}-\varepsilon_0} \left\| \frac{1}{N} \sum_{k=1}^N Z_{k,j}^{(3)} \right\|_{L_T^2 W_x^{-\varepsilon, \infty}} &\longrightarrow 0, \\ N^{\frac{1}{2}-\varepsilon_0} \left\| \frac{1}{N} \sum_{k=1}^N Z_{k,j}^{(3)} \right\|_{\mathcal{A}_{N,j} L_T^2 W_x^{-\varepsilon, \infty}} &\longrightarrow 0, \end{aligned} \quad (5.6)$$

as $N \rightarrow \infty$.

Proof. By applying Chebyshev's inequality with (5.3), (5.4), and (5.5), we see that the almost sure convergence (5.6) follows from the Borel-Cantelli lemma. Hence, we focus on proving (5.3), (5.4), and (5.5) in the following. Moreover, we assume (5.1) since the same proof holds under (5.2).

We first look at (5.3). By Sobolev's embedding (with finite $r \gg 1$ such that $\varepsilon r > 4$), Minkowski's integral inequality (for finite $p \geq r \geq 2$), the Wiener chaos estimate (Lemma 2.4), the independence of $\{Z_{k,k}^{(2)}\}_{k=1}^N$, and Lemma 2.5 (i), we have

$$\begin{aligned} \left\| \frac{1}{N} \sum_{k=1}^N Z_{k,k}^{(2)} \right\|_{L_\omega^p L_T^2 W_x^{-\varepsilon, \infty}} &\lesssim \left\| \frac{1}{N} \sum_{k=1}^N \langle \nabla \rangle^{-\frac{\varepsilon}{2}} Z_{k,k}^{(2)} \right\|_{L_\omega^p L_T^2 L_x^r} \\ &\leq \left\| \frac{1}{N} \sum_{k=1}^N \langle \nabla \rangle^{-\frac{\varepsilon}{2}} Z_{k,k}^{(2)}(t, x) \right\|_{L_T^2 L_x^r L_\omega^p} \leq p \left\| \frac{1}{N} \sum_{k=1}^N \langle \nabla \rangle^{-\frac{\varepsilon}{2}} Z_{k,k}^{(2)}(t, x) \right\|_{L_T^2 L_x^r L_\omega^2} \\ &= \frac{p}{N^{\frac{1}{2}}} \|\langle \nabla \rangle^{-\frac{\varepsilon}{2}} Z_{k,k}^{(2)}(t, x)\|_{L_T^2 L_x^r \mathcal{A}_N L_\omega^2} \lesssim_T \frac{p}{N^{\frac{1}{2}}} \|\langle \nabla \rangle^{-\frac{\varepsilon}{2}} Z_{k,k}^{(2)}(t, x)\|_{\mathcal{A}_N L_\omega^2 C_T L_x^r} \\ &\lesssim_T \frac{p}{N^{\frac{1}{2}}}, \end{aligned} \quad (5.7)$$

yielding (5.3).

We note that (5.5) follows from (5.4) and the definition (1.11) of the $\mathcal{A}_N B$ -norm. Thus, it suffices to prove (5.4). Note from (2.12) that

$$\mathbb{E}[\Psi_k^2 \Psi_j : \Psi_{k'}^2 \Psi_j] = 0$$

unless $k = k'$. Then, the bound (5.4) follows from proceeding as in (5.7). \square

In the following two lemmas, we assume that $v_k \in L^2(\Omega; C([0, T]; H^s(\mathbb{T}^2)))$, $k \in \mathbb{N}$, which is the only integrability guaranteed by Theorem 1.2 for global-in-time solutions to the mean-field SdNLW (1.20). See Subsection 5.3 for a discussion under a higher moment assumption.

Lemma 5.2. *Let $\frac{1}{2} \leq s < 1$. Given $k \in \mathbb{N}$, let Ψ_k be the stochastic convolution defined in (2.10) and let $v_k \in L^2(\Omega; C([0, T]; H^s(\mathbb{T}^2)))$ be a solution of the mean-field SdNLW (1.18). Moreover, we assume that $(v_k(0), \partial_t v_k(0)) \in L^2(\Omega; \mathcal{H}^s(\mathbb{T}^2))$, $k \in \mathbb{N}$, are independent and identically distributed over $k \in \mathbb{N}$. Then, given $T > 0$ and small $\varepsilon > 0$, we have*

$$\left\| \frac{1}{N} \sum_{k=1}^N \left(\Psi_k v_k - \mathbb{E}[\Psi_k v_k] \right) \right\|_{L_T^2 W_x^{-\varepsilon, \frac{4}{2-s-\varepsilon}}} \longrightarrow 0, \quad (5.8)$$

$$\left\| \frac{1}{N} \sum_{k=1}^N \left(v_k^2 - \mathbb{E}[v_k^2] \right) \right\|_{L_T^2(L_x^{\frac{3}{2-s}} \cap W_x^{\varepsilon, \frac{2}{2-s-\varepsilon}})} \longrightarrow 0, \quad (5.9)$$

almost surely, as $N \rightarrow \infty$.

Proof. We first note that (Ψ_k, v_k) , $k \in \mathbb{N}$, are independent and identically distributed since Ψ_k are independent and identically distributed and so are the initial data $(v_k(0), \partial_t v_k(0))$. In particular, we have

$$R \stackrel{\text{def}}{=} \sup_{k \in \mathbb{N}} \|v_k\|_{L_\omega^2 C_T H_x^s} = \|v_1\|_{L_\omega^2 C_T H_x^s} < \infty. \quad (5.10)$$

Proceeding as in (4.8) with Lemma 2.1 (ii), Cauchy-Schwarz's inequality, Lemma 2.5 (i), Sobolev's inequality, and (5.10), we have

$$\begin{aligned} \mathbb{E} \left[\|\Psi_k v_k\|_{L_T^2 W_x^{-\varepsilon, \frac{4}{2-s-\varepsilon}}} \right] &\lesssim T^{\frac{1}{2}} \|\Psi_k\|_{L_\omega^2 C_T W_x^{-\varepsilon, \infty}} \|v_k\|_{L_\omega^2 C_T W_x^{\varepsilon, \frac{4}{2-s-\varepsilon}}} \\ &\lesssim T^{\frac{1}{2}} \|v_k\|_{L_\omega^2 C_T H_x^s} \\ &< \infty, \end{aligned}$$

uniformly in $k \in \mathbb{N}$. Hence, in view of the independence of $\{\Psi_k v_k\}_{k \in \mathbb{N}}$, the almost sure convergence (5.8) follows from the strong law of large numbers in the Banach space setting ([26, Corollary 7.10 on p. 189]).

Next, we consider (5.9). Proceeding as in (4.7) and (4.9) with the fractional Leibniz rule (Lemma 2.1 (i)), Sobolev's inequality, Cauchy-Schwarz's inequality, and (5.10), we have

$$\mathbb{E} \left[\|v_k^2\|_{L_T^2(L_x^{\frac{3}{2-s}} \cap W_x^{\varepsilon, \frac{2}{2-s-\varepsilon}})} \right] \lesssim T^{\frac{1}{2}} \|v_k\|_{L_\omega^2 C_T H_x^s}^2 < \infty, \quad (5.11)$$

uniformly in $k \in \mathbb{N}$. Hence, in view of the independence of $\{v_k^2\}_{k \in \mathbb{N}}$, the almost sure convergence (5.8) follows from the strong law of large numbers ([26, Corollary 7.10 on p. 189]). \square

Remark 5.3. From (5.11), we see that the $L^2(\Omega)$ -integrability assumption for v_k is indeed necessary for the (almost sure) convergence (5.9); see [26, Corollary 7.10 on p. 189].

Lemma 5.4. *Let $\frac{1}{2} \leq s < 1$ and $\{(\Psi_j, v_j)\}_{j \in \mathbb{N}}$ be as in Lemma 5.2. Then, given $T > 0$, we have, for each $j \in \mathbb{N}$,*

$$\left\| \frac{1}{N} \sum_{k=1}^N \left(v_k : \Psi_k \Psi_j : - \mathbb{E}[v_k \Psi_k] \Psi_j \right) \right\|_{L_T^2 H_x^{s-1}} \longrightarrow 0 \quad (5.12)$$

in probability as $N \rightarrow \infty$, and

$$\left\| \frac{1}{N} \sum_{k=1}^N \left(v_k : \Psi_k \Psi_j : - \mathbb{E}[v_k \Psi_k] \Psi_j \right) \right\|_{\mathcal{A}_{N,j} L_T^2 H_x^{s-1}} \longrightarrow 0 \quad (5.13)$$

in probability as $N \rightarrow \infty$.

Proof. We only prove (5.13) in the following, since (5.12) follows in a similar manner.

We first study the contribution to (5.13) from the case $k = j$. Let $1 < p < 2$. Then, proceeding as in (3.10) with Hölder's inequality, Lemma 2.5 (i), and (5.10), we have

$$\begin{aligned} \left\| \|v_j : \Psi_j^2 : \|_{L_T^2 H_x^{s-1}} \right\|_{L_\omega^p} &\lesssim T^{\frac{1}{2}} \|v_j\|_{L_\omega^2 C_T H_x^s} \|\Psi_j^2 : \|_{L_\omega^{\frac{2p}{2-p}} C_T W_x^{-\varepsilon, \infty}} \\ &\lesssim_p T^{\frac{1}{2}}, \end{aligned} \quad (5.14)$$

uniformly in $j \in \mathbb{N}$, provided that $0 < \varepsilon \leq \min(1-s, s)$. Let (Ψ', v') be an independent copy of (Ψ_j, v_j) . Then, given $1 < p < 2$, it follows from Minkowski's integral inequality, the independence of (Ψ', v') from Ψ_j , and proceeding as in (5.14) with (4.5) that

$$\begin{aligned} \left\| \mathbb{E}[v_j \Psi_j] \Psi_j \right\|_{L_T^2 H_x^{s-1}} \Big|_{L_\omega^p} &= \left\| \mathbb{E}_{\omega'}[v'(\omega') \Psi'(\omega')] \Psi_j(\omega) \right\|_{L_T^2 H_x^{s-1}} \Big|_{L_\omega^p} \\ &\leq \|v' \Psi' \Psi_j\|_{L_\omega^p L_T^2 H_x^{s-1}} \\ &\lesssim T^{\frac{1}{2}} \|v_j\|_{L_\omega^2 C_T H_x^s} \|\Psi' \Psi_j\|_{L_\omega^{\frac{2p}{2-p}} C_T W_x^{-\varepsilon, \infty}} \\ &\lesssim T^{\frac{1}{2}}, \end{aligned} \quad (5.15)$$

uniformly in $j \in \mathbb{N}$. Hence, given $L \gg 1$, it follows from Chebyshev's inequality with (5.14) and (5.15) that

$$\sum_{N=1}^{\infty} \mathbb{P} \left(\left\| \frac{1}{N} (v_j : \Psi_j^2 : -\mathbb{E}[v_j \Psi_j] \Psi_j) \right\|_{\mathcal{A}_{N,j} L_T^2 H_x^{s-1}} > \frac{1}{L} \right) \lesssim_T \sum_{N=1}^{\infty} \frac{L^p}{N^p} < \infty, \quad (5.16)$$

since $p > 1$. Therefore, by the Borel-Cantelli lemma, we conclude that

$$\left\| \frac{1}{N} (v_j : \Psi_j^2 : -\mathbb{E}[v_j \Psi_j] \Psi_j) \right\|_{\mathcal{A}_{N,j} L_T^2 H_x^{s-1}} \longrightarrow 0,$$

almost surely (and thus in probability) as $N \rightarrow \infty$, which shows (5.13) for the case $k = j$.

Next, we study the contribution to (5.13) from the case $k \neq j$. By taking the Fourier transform in x (suppressing the t -dependence), we have

$$\begin{aligned} \mathcal{F}_x \left((v_k \Psi_k - \mathbb{E}[v_k \Psi_k]) \Psi_j \right) (n) \\ = \sum_{n=n_1+n_2+n_3} \left(\widehat{v}_k(n_1) \widehat{\Psi}_k(n_2) - \mathbb{E}[\widehat{v}_k(n_1) \widehat{\Psi}_k(n_2)] \right) \widehat{\Psi}_j(n_3). \end{aligned}$$

Then, given $M \gg 1$ (to be chosen later), we write

$$\frac{1}{N} \sum_{\substack{k=1 \\ k \neq j}}^N (v_k \Psi_k - \mathbb{E}[v_k \Psi_k]) \Psi_j = \mathbf{I}_j + \mathbf{II}_j \quad (5.17)$$

where \mathbf{I}_j denotes the contribution from the case

$$\max(|n|, |n_1|, |n_2|, |n_3|) > M. \quad (5.18)$$

Namely, \mathbf{II}_j denotes the contribution from the case: $\max(|n|, |n_1|, |n_2|, |n_3|) \leq M$, given by

$$\mathbf{II}_j = \frac{1}{N} \sum_{\substack{k=1 \\ k \neq j}}^N \mathbf{P}_M \left((\mathbf{P}_M v_k \Psi_{k,M} - \mathbb{E}[\mathbf{P}_M v_k \Psi_{k,M}]) \Psi_{j,M} \right), \quad (5.19)$$

where $\Psi_{j,M} = \mathbf{P}_M \Psi_j$.

We first treat the first term \mathbf{I}_j . From a slight modification of (3.10), we have

$$\begin{aligned} \|v_k \Psi_k \Psi_j\|_{L_T^2 H_x^{s-1+\varepsilon}} &\lesssim T^{\frac{1}{2}} \|v_k \Psi_k \Psi_j\|_{C_T W_x^{-2\varepsilon, \frac{2}{2-s-3\varepsilon}}} \\ &\lesssim T^{\frac{1}{2}} \|v_k\|_{C_T W_x^{2\varepsilon, \frac{2}{2-s-3\varepsilon}}} \|\Psi_k \Psi_j\|_{C_T W_x^{-2\varepsilon, \infty}} \\ &\lesssim T^{\frac{1}{2}} \|v_k\|_{C_T H_x^{s-\varepsilon}} \|\Psi_k \Psi_j\|_{C_T W_x^{-2\varepsilon, \infty}}. \end{aligned} \quad (5.20)$$

Note that, thanks to (5.18), we can gain $M^{-\varepsilon}$ by losing ε -regularity. Thus, from (5.20), (5.18), Lemma 2.5(ii) (see also (2.24) in Remark 2.6(ii)), (5.20), Cauchy-Schwarz's inequality, and (5.10), we have

$$\mathbb{E} \left[\|\mathbf{I}_j\|_{\mathcal{A}_{N,j} L_T^2 H_x^{s-1}} \right] \lesssim_T M^{-\varepsilon} R,$$

uniformly in $N \in \mathbb{N}$. Then, given small $\delta > 0$, it follows from Markov's inequality that

$$\mathbb{P} \left(\|\mathbf{I}_j\|_{\mathcal{A}_{N,j} L_T^2 H_x^{s-1}} > \delta \right) \leq C_T \frac{R}{\delta M^\varepsilon} < \frac{\delta}{3}, \quad (5.21)$$

uniformly in $N \in \mathbb{N}$, by taking $M = M(\delta, T, R) \gg 1$ sufficiently large.

Next, we consider the term Π_j in (5.19), where all the frequencies are bounded by M . We first write Π_j as

$$\begin{aligned} \Pi_j &= \Pi_j^{(1)} + \Pi_j^{(2)} \\ &\stackrel{\text{def}}{=} \frac{1}{N} \sum_{k=1}^N \mathbf{P}_M \left((\mathbf{P}_M v_k \Psi_{k,M} - \mathbb{E}[\mathbf{P}_M v_k \Psi_{k,M}]) \Psi_{j,M} \right) \\ &\quad - \frac{1}{N} \mathbf{P}_M \left((\mathbf{P}_M v_j \Psi_{j,M} - \mathbb{E}[\mathbf{P}_M v_j \Psi_{j,M}]) \Psi_{j,M} \right). \end{aligned} \quad (5.22)$$

Let us first consider $\Pi_j^{(2)}$. By a slight modification of (5.14), for $1 < p < 2$, we have

$$\begin{aligned} \left\| \|\mathbf{P}_M v_j \Psi_{j,M}^2\|_{L_T^2 H_x^{s-1}} \right\|_{L_\omega^p} &\lesssim T^{\frac{1}{2}} M^{2\varepsilon} \|v_j\|_{L_\omega^2 C_T H_x^s} \|\Psi_j\|_{L_\omega^{\frac{4p}{2-p}} C_T W_x^{-\varepsilon, \infty}}^2 \\ &\lesssim_p T^{\frac{1}{2}} M^{2\varepsilon}, \end{aligned} \quad (5.23)$$

uniformly in $j \in \mathbb{N}$. Then, from (5.23) and (5.15) (which implies (5.16) with an extra factor $M^{2\varepsilon p}$ on the right-hand side) with the Borel-Cantelli lemma, we conclude that

$$\|\Pi_j^{(2)}\|_{\mathcal{A}_{N,j} L_T^2 H_x^{s-1}} \longrightarrow 0,$$

almost surely (and thus in probability) as $N \rightarrow \infty$. Namely, given small $\delta > 0$, there exists $N_1 = N_1(\delta) \gg 1$ such that

$$\mathbb{P} \left(\|\Pi_j^{(2)}\|_{\mathcal{A}_{N,j} L_T^2 H_x^{s-1}} > \delta \right) < \frac{\delta}{3} \quad (5.24)$$

for any $N \geq N_1$.

Next, we treat $\Pi_j^{(1)}$ in (5.22). From (5.10) and Lemma 2.5(i), we have

$$\begin{aligned} \mathbb{E} \left[\|\mathbf{P}_M v_k \Psi_{k,M}\|_{L_T^2 L_x^2} \right] &\lesssim M^\varepsilon T^{\frac{1}{2}} \|v_1\|_{L_\omega^2 C_T L_x^2} \|\Psi_1\|_{L_\omega^2 C_T W_x^{-\varepsilon, \infty}} \\ &< \infty, \end{aligned}$$

uniformly in $k \in \mathbb{N}$. Then, noting that $\mathbf{P}_M v_k \Psi_{k,M} - \mathbb{E}[\mathbf{P}_M v_k \Psi_{k,M}]$, $k \in \mathbb{N}$, are independent and identically distributed, it follows from [26, Corollary 7.10 on p. 189] that

$$Y_{N,M} \stackrel{\text{def}}{=} \left\| \frac{1}{N} \sum_{k=1}^N (\mathbf{P}_M v_k \Psi_{k,M} - \mathbb{E}[\mathbf{P}_M v_k \Psi_{k,M}]) \right\|_{L_T^2 L_x^2} \longrightarrow 0, \quad (5.25)$$

almost surely, as $N \rightarrow \infty$, which in particular implies that the convergence (5.25) also holds in probability. Thus, given small $\delta > 0$ and $K \gg 1$, there exists $N_2 = N_2(\delta, M, K) \gg 1$ such that

$$\mathbb{P} \left(Y_{N,M} > \delta K^{-1} \right) < \frac{\delta}{6} \quad (5.26)$$

for any $N \geq N_2$. By a crude estimate, we have

$$\|\Pi_j^{(1)}\|_{\mathcal{A}_{N,j}L_T^2H_x^{s-1}} \leq C_0M^\varepsilon Y_{N,M}\|\Psi_j\|_{\mathcal{A}_{N,j}C_TW_x^{-\varepsilon,\infty}}. \quad (5.27)$$

Given small $\delta > 0$, $T > 0$, and $M \gg 1$, it follows from Lemma 2.5 (ii) that there exists $K = K(\delta, M, T) \gg 1$ such that

$$\mathbb{P}\left(C_0M^\varepsilon\|\Psi_j\|_{\mathcal{A}_{N,j}C_TW_x^{-\varepsilon,\infty}} > K\right) < \frac{\delta}{6}, \quad (5.28)$$

uniformly in $N \in \mathbb{N}$. Hence, given small $\delta > 0$, it follows from (5.27) with (5.26) and (5.28) that we have

$$\begin{aligned} & \mathbb{P}\left(\|\Pi_j^{(1)}\|_{\mathcal{A}_{N,j}L_T^2H_x^{s-1}} > \delta\right) \\ & \leq \mathbb{P}\left(Y_{N,M} > \delta K^{-1}\right) \\ & \quad + \mathbb{P}\left(\{Y_{N,M} \leq \delta K^{-1}\} \cap \{C_0M^\varepsilon\|\Psi_j\|_{\mathcal{A}_{N,j}C_TW_x^{-\varepsilon,\infty}} > K\}\right) \\ & < \frac{\delta}{6} + \mathbb{P}\left(C_0M^\varepsilon\|\Psi_j\|_{\mathcal{A}_{N,j}C_TW_x^{-\varepsilon,\infty}} > K\right) \\ & < \frac{\delta}{3} \end{aligned} \quad (5.29)$$

for any $N \geq N_2$.

Therefore, from (5.17), (5.21), (5.22), (5.24), and (5.29), we conclude (5.13) for the case $k \neq j$. \square

5.2. Proof of Theorem 1.4 (i). In this subsection, we present a proof of Theorem 1.4 (i).

Let $\frac{4}{5} < s < 1$. Let $\{(u_{N,j}^{(0)}, u_{N,j}^{(1)})\}_{j=1}^N \in (\mathcal{H}^s(\mathbb{T}^2))^{\otimes N}$, $N \in \mathbb{N}$, and $\{(u_j^{(0)}, u_j^{(1)})\}_{j \in \mathbb{N}} \in (\mathcal{H}^s(\mathbb{T}^2))^{\otimes \mathbb{N}}$ be sequences of random functions as in Theorem 1.4 (i), where the former converges in probability to the latter in the sense of (1.24) and (1.25). In particular, we assume that $(u_j^{(0)}, u_j^{(1)}) \in L^2(\Omega; \mathcal{H}^s(\mathbb{T}^2))$, $j \in \mathbb{N}$, are independent and identically distributed.

Let $\{(u_{N,j}, \partial_t u_{N,j})\}_{j=1}^N$ and $\{(u_j, \partial_t u_j)\}_{j \in \mathbb{N}}$ be the global-in-time solutions to (1.10) and (1.20) with initial data $\{(u_{N,j}^{(0)}, u_{N,j}^{(1)})\}_{j=1}^N$ and $\{(u_j^{(0)}, u_j^{(1)})\}_{j \in \mathbb{N}}$ constructed in Theorems 1.1 and 1.2, respectively. Then, set $v_{N,j} = u_{N,j} - \Psi_j$ and $v_j = u_j - \Psi_j$ as in (1.7) and (1.19) such that $\{(v_{N,j}, \partial_t v_{N,j})\}_{j=1}^N \in (C(\mathbb{R}_+; \mathcal{H}^s(\mathbb{T}^2)))^{\otimes N}$ almost surely and $(v_j, \partial_t v_j) \in L^2(\Omega; C([0, T]; \mathcal{H}^s(\mathbb{T}^2)))$ for any $T > 0$. Then, by writing (1.8) and (1.18) in the Duhamel formulation, we have

$$\begin{aligned} v_{N,j} = S(t)(u_{N,j}^{(0)}, u_{N,j}^{(1)}) - \frac{1}{N} \sum_{k=1}^N \mathcal{I} \Big(& v_{N,k}^2 v_{N,j} + 2\Psi_k v_{N,k} v_{N,j} + v_{N,k}^2 \Psi_j \\ & + : \Psi_k^2 : v_{N,j} + 2v_{N,k} : \Psi_k \Psi_j : + : \Psi_k^2 \Psi_j : \Big) \end{aligned} \quad (5.30)$$

and

$$\begin{aligned} v_j = S(t)(u_j^{(0)}, u_j^{(1)}) \\ - \mathcal{I} \Big(\mathbb{E}[v_j^2] v_j + 2\mathbb{E}[\Psi_j v_j] v_j + \mathbb{E}[v_j^2] \Psi_j + 2\mathbb{E}[v_j \Psi_j] \Psi_j \Big), \end{aligned} \quad (5.31)$$

where $S(t)$ and \mathcal{I} are as in (2.5) and (2.6), respectively. We set

$$V_{N,j} = v_{N,j} - v_j \quad (5.32)$$

and $\vec{V}_{N,j} = (V_{N,j}, \partial_t V_{N,j})$. By taking the difference of (5.30) and (5.31), we have

$$\begin{aligned} V_{N,j}(t) &= S(t)(\vec{V}_{N,j}(0)) - \sum_{\ell=1}^6 \mathcal{I}(\mathcal{A}_{j,\ell})(t) \\ &= S(t)(u_{N,j}^{(0)} - u_j^{(0)}, u_{N,j}^{(1)} - u_j^{(1)}) - \sum_{\ell=1}^6 \mathcal{I}(\mathcal{A}_{j,\ell})(t), \end{aligned} \quad (5.33)$$

where $\{\mathcal{A}_{j,\ell} = A_{j,\ell}(N)\}_{j=1}^N$, $\ell = 1, \dots, 6$, are given by

$$\begin{aligned} \mathcal{A}_{j,1} &= \frac{1}{N} \sum_{k=1}^N v_{N,k}^2 v_{N,j} - \mathbb{E}[v_j^2] v_j, \\ \mathcal{A}_{j,2} &= \frac{1}{N} \sum_{k=1}^N 2\Psi_k v_{N,k} v_{N,j} - 2\mathbb{E}[\Psi_j v_j] v_j, \\ \mathcal{A}_{j,3} &= \frac{1}{N} \sum_{k=1}^N v_{N,k}^2 \Psi_j - \mathbb{E}[v_j^2] \Psi_j, \\ \mathcal{A}_{j,4} &= \frac{1}{N} \sum_{k=1}^N : \Psi_k^2 : v_{N,j}, \\ \mathcal{A}_{j,5} &= \frac{1}{N} \sum_{k=1}^N 2v_{N,k} : \Psi_k \Psi_j : - 2\mathbb{E}[v_j \Psi_j] \Psi_j, \\ \mathcal{A}_{j,6} &= \frac{1}{N} \sum_{k=1}^N : \Psi_k^2 \Psi_j : . \end{aligned} \quad (5.34)$$

Fix a target time $T \gg 1$ and we work on the interval $[0, T]$. We point out the following; while we take $\frac{4}{5} < s < 1$ to prove global-in-time convergence (namely, on $[0, T]$ for any given $T \gg 1$), the convergence argument presented below works for $\frac{1}{2} \leq s < 1$, *assuming* that v_j exists on $[0, T]$ (more precisely, $(v_j, \partial_t v_j) \in L^2(\Omega; C([0, T]; \mathcal{H}^s(\mathbb{T}^2)))$).

Since the parameters s, ε, T are fixed, we often drop the dependence on these parameters. In the remaining part of this subsection, we fix small $\delta > 0$.

• **Step 1:** In this first step, we establish the convergence (1.27) for each $T \gg 1$:

$$\|\vec{V}_{N,j}\|_{X_N^s(T)} = \|\vec{v}_{N,j} - \vec{v}_j\|_{X_N^s(T)} \longrightarrow 0 \quad (5.35)$$

in probability as $N \rightarrow \infty$, where $X_N^s(T)$ is as in (3.3).

Before we start estimating the terms on the right-hand side of (5.33), let us first prepare data sets indexed by $N \gg 1$. As we noted in the proof of Lemma 5.2, the solutions $\vec{v}_j = (v_j, \partial_t v_j) \in C([0, T]; \mathcal{H}^s(\mathbb{T}^2))$ to (1.18), $j \in \mathbb{N}$, are independent and identically distributed. In particular, we have

$$\sup_{j \in \mathbb{N}} \|\vec{v}_j\|_{L_\omega^2 C_T \mathcal{H}_x^s} = \|\vec{v}_1\|_{L_\omega^2 C_T \mathcal{H}_x^s} < \infty. \quad (5.36)$$

Given $N \in \mathbb{N}$ and $R \geq 1$, define the set $\Omega_{1,N}(R) \subset \Omega$ by setting

$$\Omega_{1,N}(R) = \left\{ \omega \in \Omega : \|\vec{v}_j\|_{X_N^s(T)} \leq R \right\}. \quad (5.37)$$

Then, it follows from Chebyshev's inequality, (3.3), and (5.36) that there exists $R = R(\delta) \gg 1$, independent of $N \in \mathbb{N}$, such that

$$\mathbb{P}((\Omega_{1,N}(R))^c) < \frac{\delta}{8}. \quad (5.38)$$

Given small $\eta_0 > 0$, define $\Omega_{2,N}(\eta_0) \subset \Omega$ by setting

$$\Omega_{2,N}(\eta_0) = \left\{ \omega \in \Omega : \|\vec{V}_{N,j}(0)\|_{\mathcal{A}_N \mathcal{H}_x^s} = \|(u_{N,j}^{(0)} - u_j^{(0)}, u_{N,j}^{(1)} - u_j^{(1)})\|_{\mathcal{A}_N \mathcal{H}_x^s} \leq \eta_0 \right\}. \quad (5.39)$$

Then, from the assumption (1.25) on the initial data, there exists $N_1 = N_1(\eta_0, \delta) \in \mathbb{N}$ such that

$$\mathbb{P}((\Omega_{2,N}(\eta_0))^c) < \frac{\delta}{8}. \quad (5.40)$$

for any $N \geq N_1$.

Next, we discuss controls on the stochastic terms. Given $K \geq 1$, define the set $\Omega_{3,N}(K) \subset \Omega$ by setting

$$\Omega_{3,N}(K) = \left\{ \omega \in \Omega : \|\Psi_j\|_{\mathcal{A}_N C_T W_x^{-\varepsilon, \infty}} + \|\Psi_k \Psi_j\|_{\mathcal{A}_N^{(2)} C_T W_x^{-\varepsilon, \infty}} \leq K \right\}. \quad (5.41)$$

Then, it follows from Lemma 2.5 (ii) and Chebyshev's inequality that there exists $K = K(\delta) \gg 1$, independent of $N \in \mathbb{N}$, such that

$$\mathbb{P}((\Omega_{3,N}(K))^c) < \frac{\delta}{8}. \quad (5.42)$$

Given small $\eta > 0$, define the set $\Omega_{4,N}(\eta) \subset \Omega$ by setting

$$\Omega_{4,N}(\eta) = \left\{ \omega \in \Omega : \|(\mathbf{v}, \Psi)\|_{\mathcal{Y}_N} \leq \eta \right\}, \quad (5.43)$$

where the \mathcal{Y}_N -norm is defined by

$$\begin{aligned} \|(\mathbf{v}, \Psi)\|_{\mathcal{Y}_N} &= \left\| \frac{1}{N} \sum_{k=1}^N \Psi_k^2 \right\|_{L_T^2 W_x^{-\varepsilon, \infty}} + \left\| \frac{1}{N} \sum_{k=1}^N \Psi_k^2 \Psi_j \right\|_{\mathcal{A}_{N,j} L_T^2 W_x^{-\varepsilon, \infty}} \\ &\quad + \left\| \frac{1}{N} \sum_{k=1}^N (\Psi_k v_k - \mathbb{E}[\Psi_k v_k]) \right\|_{L_T^2 W_x^{-\varepsilon, \frac{4}{2-s-\varepsilon}}} \\ &\quad + \left\| \frac{1}{N} \sum_{k=1}^N (v_k^2 - \mathbb{E}[v_k^2]) \right\|_{L_T^2 (L_x^{\frac{3}{2-s}} \cap W_x^{\varepsilon, \frac{2}{2-s-\varepsilon}})} \\ &\quad + \left\| \frac{1}{N} \sum_{k=1}^N (v_k \Psi_k \Psi_j - \mathbb{E}[v_k \Psi_k] \Psi_j) \right\|_{\mathcal{A}_{N,j} L_T^2 H_x^{s-1}}. \end{aligned} \quad (5.44)$$

Then, it follows from Lemmas 5.1, 5.2, and 5.4 that there exists $N_2 = N_2(\eta, \delta) \in \mathbb{N}$ such that

$$\mathbb{P}((\Omega_{4,N}(\eta))^c) < \frac{\delta}{8} \quad (5.45)$$

for $N \geq N_2$.

We are now ready to estimate the terms on the right-hand side of (5.33). Let $J \subset [0, T]$ be an interval with $|J| \leq 1$. With a slight abuse of notation, we define $X_N^s(J)$ by setting

$$\begin{aligned} \|\vec{v}_j\|_{X_N^s(J)} &= \|(v_j, \partial_t v_j)\|_{\mathcal{A}_N C_J \mathcal{H}_x^s} \\ &= \|(v_j, \partial_t v_j)\|_{\mathcal{A}_{N,j} C_J \mathcal{H}_x^s}, \end{aligned} \quad (5.46)$$

where the $\mathcal{A}_N B$ -norm is as in (1.11). In the following, we work on $\Omega_N(R, K, \eta_0, \eta)$ defined by

$$\Omega_N(R, K, \eta_0, \eta) = \Omega_{1,N}(R) \cap \Omega_{2,N}(\eta_0) \cap \Omega_{3,N}(K) \cap \Omega_{4,N}(\eta), \quad (5.47)$$

where $\Omega_{1,N}(R)$, $\Omega_{2,N}(\eta_0)$, $\Omega_{3,N}(K)$, and $\Omega_{4,N}(\eta)$ are as in (5.37), (5.39), (5.41), and (5.43), respectively. Before proceeding further, we note that, given small $\delta > 0$, by choosing $R = R(\delta)$ and $K = K(\delta)$ sufficiently large, it follows from (5.47) with (5.38), (5.40), (5.42), and (5.45) that

$$\mathbb{P}((\Omega_N(R, K, \eta_0, \eta))^c) < \frac{\delta}{2} \quad (5.48)$$

for any $N \geq \max(N_1, N_2)$, where $N_1(\eta_0, \delta)$ and $N_2(\eta, \delta)$ are as in (5.40) and (5.45), respectively.

From (5.34) with (5.32), we have

$$\begin{aligned} \mathcal{A}_{j,1} &= \frac{1}{N} \sum_{k=1}^N V_{N,k} v_{N,k} v_{N,j} + \frac{1}{N} \sum_{k=1}^N v_k V_{N,k} v_{N,j} \\ &\quad + \frac{1}{N} \sum_{k=1}^N v_k v_k V_{N,j} + \frac{1}{N} \sum_{k=1}^N (v_k^2 - \mathbb{E}[v_k^2]) v_j. \end{aligned} \quad (5.49)$$

Using (5.32), we replace each occurrence of $v_{N,\ell}$ in (5.49) by $V_{N,\ell} + v_\ell$, $\ell = j, k$. Then, by proceeding as in (3.6) (while keeping the k -summation inside for the last term in (5.49)) with (5.37) and (5.43), we have

$$\begin{aligned} \|\vec{\mathcal{I}}(\mathcal{A}_{j,1})\|_{X_N^s(J)} &\lesssim |J| \left(\|\vec{V}_{N,k}\|_{X_N^s(J)}^2 + \|\vec{v}_k\|_{X_N^s(J)}^2 \right) \|\vec{V}_{N,j}\|_{X_N^s(J)} \\ &\quad + |J|^{\frac{1}{2}} \left\| \frac{1}{N} \sum_{k=1}^N (v_k^2 - \mathbb{E}[v_k^2]) \right\|_{L_J^2 L_x^{\frac{3}{2-s}}} \|\vec{v}_j\|_{X_N^s(J)} \\ &\leq |J| \left(\|\vec{V}_{N,k}\|_{X_N^s(J)}^2 + R^2 \right) \|\vec{V}_{N,j}\|_{X_N^s(J)} + |J|^{\frac{1}{2}} \eta R, \end{aligned} \quad (5.50)$$

provided that $\frac{1}{2} \leq s (< 1)$.

From (5.34) with (5.32), we have

$$\begin{aligned} \mathcal{A}_{j,2} &= \frac{1}{N} \sum_{k=1}^N 2\Psi_k V_{N,k} v_{N,j} + \frac{1}{N} \sum_{k=1}^N 2\Psi_k v_k V_{N,j} \\ &\quad + \frac{2}{N} \sum_{k=1}^N (\Psi_k v_k - \mathbb{E}[\Psi_k v_k]) v_j. \end{aligned} \quad (5.51)$$

Then, by proceeding as in (3.8) (see also (4.8)) with Lemma 2.1 (ii), Sobolev's inequality, (5.37), (5.41), and (5.43), we have

$$\begin{aligned} \|\vec{\mathcal{I}}(\mathcal{A}_{j,2})\|_{X_N^s(J)} &\lesssim |J| \|\Psi_k\|_{\mathcal{A}_N C_J W_x^{-\varepsilon, \infty}} \left(\|\vec{V}_{N,k}\|_{X_N^s(J)} + \|\vec{v}_k\|_{X_N^s(J)} \right) \|\vec{V}_{N,j}\|_{X_N^s(J)} \\ &\quad + |J|^{\frac{1}{2}} \left\| \frac{1}{N} \sum_{k=1}^N (\Psi_k v_k - \mathbb{E}[\Psi_k v_k]) \right\|_{L_J^2 W_x^{-\varepsilon, \frac{4}{2-s-\varepsilon}}} \|v_j\|_{\mathcal{A}_{N,j} C_J W_x^{\varepsilon, \frac{4}{2-s-\varepsilon}}} \\ &\leq |J| K \left(\|\vec{V}_{N,k}\|_{X_N^s(J)} + R \right) \|\vec{V}_{N,j}\|_{X_N^s(J)} + |J|^{\frac{1}{2}} \eta R, \end{aligned} \quad (5.52)$$

provided that $3\varepsilon \leq s$ (< 1). Similarly, by writing

$$\mathcal{A}_{j,3} = \frac{1}{N} \sum_{k=1}^N V_{N,k} v_{N,k} \Psi_j + \frac{1}{N} \sum_{k=1}^N v_k V_{N,k} \Psi_j + \frac{1}{N} \sum_{k=1}^N \left(v_k^2 - \mathbb{E}[v_k^2] \right) \Psi_j \quad (5.53)$$

and proceeding as in (5.52) (see also (4.9)) with (5.37), (5.41), and (5.43), we have

$$\begin{aligned} & \|\vec{\mathcal{I}}(\mathcal{A}_{j,3})\|_{X_N^s(J)} \\ & \lesssim |J| \left(\|\vec{V}_{N,k}\|_{X_N^s(J)} + \|\vec{v}_k\|_{X_N^s(J)} \right) \|\vec{V}_{N,k}\|_{X_N^s(J)} \|\Psi_j\|_{\mathcal{A}_N C_J W_x^{-\varepsilon, \infty}} \\ & \quad + |J|^{\frac{1}{2}} \left\| \frac{1}{N} \sum_{k=1}^N \left(v_k^2 - \mathbb{E}[v_k^2] \right) \right\|_{L_T^2 W_x^{\varepsilon, \frac{2}{2-s-\varepsilon}}} \|\Psi_j\|_{\mathcal{A}_N C_J W_x^{-\varepsilon, \infty}} \\ & \leq |J| K \left(\|\vec{V}_{N,k}\|_{X_N^s(J)} + R \right) \|\vec{V}_{N,j}\|_{X_N^s(J)} + |J|^{\frac{1}{2}} \eta K. \end{aligned} \quad (5.54)$$

Proceeding as in (3.10) with (5.37) and (5.43), we have

$$\begin{aligned} \|\vec{\mathcal{I}}(\mathcal{A}_{j,4})\|_{X_N^s(J)} & \lesssim |J|^{\frac{1}{2}} \left\| \frac{1}{N} \sum_{k=1}^N : \Psi_k^2 : \right\|_{L_T^2 W_x^{-\varepsilon, \infty}} \left(\|\vec{V}_{N,j}\|_{X_N^s(J)} + \|\vec{v}_j\|_{X_N^s(J)} \right) \\ & \lesssim |J|^{\frac{1}{2}} \|\vec{V}_{N,j}\|_{X_N^s(J)} + |J|^{\frac{1}{2}} \eta R, \end{aligned} \quad (5.55)$$

provided that $\varepsilon \leq s \leq 1 - \varepsilon$. By writing

$$\mathcal{A}_{j,5} = \frac{2}{N} \sum_{k=1}^N V_{N,k} : \Psi_k \Psi_j : + \frac{2}{N} \sum_{k=1}^N \left(v_k : \Psi_k \Psi_j : - \mathbb{E}[v_k \Psi_k] \Psi_j \right) \quad (5.56)$$

and proceeding as in (3.10) with (5.41) and (5.43), we have

$$\begin{aligned} \|\vec{\mathcal{I}}(\mathcal{A}_{j,5})\|_{X_N^s(J)} & \lesssim |J| \| : \Psi_k \Psi_j : \|_{\mathcal{A}_N^{(2)} C_J W_x^{-\varepsilon, \infty}} \|\vec{V}_{N,k}\|_{X_N^s(J)} + |J|^{\frac{1}{2}} \eta \\ & \leq |J| K \|\vec{V}_{N,k}\|_{X_N^s(J)} + |J|^{\frac{1}{2}} \eta, \end{aligned} \quad (5.57)$$

where $A_N^{(2)} B$ is as in (2.2). Lastly, from (5.34) and (5.43), we have

$$\begin{aligned} \|\vec{\mathcal{I}}(\mathcal{A}_{j,6})\|_{X_N^s(J)} & \lesssim |J|^{\frac{1}{2}} \left\| \frac{1}{N} \sum_{k=1}^N : \Psi_k^2 \Psi_j : \right\|_{\mathcal{A}_{N,j} L_J^2 H_x^{s-1}} \\ & \leq |J|^{\frac{1}{2}} \eta. \end{aligned} \quad (5.58)$$

Therefore, putting (5.33) (but starting at time t_0), (5.50), (5.52), (5.54), (5.55), (5.57), and (5.58) together with $J = [t_0, t_1] \subset [0, T]$ such that $|J| \leq 1$, we obtain

$$\begin{aligned} \|\vec{V}_{N,j}\|_{X_N^s(J)} & \leq C_1 \|\vec{V}_{N,j}(t_0)\|_{\mathcal{A}_N \mathcal{H}_x^s} \\ & \quad + C_2 |J|^{\frac{1}{2}} K \left(\|\vec{V}_{N,k}\|_{X_N^s(J)}^2 + R^2 \right) \|\vec{V}_{N,j}\|_{X_N^s(J)} \\ & \quad + C_3 |J|^{\frac{1}{2}} \eta (R + K). \end{aligned} \quad (5.59)$$

Suppose that

$$\|\vec{V}_{N,j}\|_{X_N^s(J)} \leq 1 \quad (5.60)$$

for some interval $J \subset [0, T]$. Then, it follows from (5.59) that there exists $\tau = \tau(R, K) \in (0, 1]$ such that

$$\|\vec{V}_{N,j}\|_{X_N^s(J)} \leq 2C_1 \|\vec{V}_{N,j}(t_0)\|_{\mathcal{A}_N \mathcal{H}_x^s} + 2C_3 \eta, \quad (5.61)$$

provided that $|J| \leq \tau$.

While the claimed convergence-in-probability (5.35) follows from (5.48), the discussion above, (5.39) (with sufficiently small $\eta_0 > 0$), and a standard continuity argument, we present some details for readers' convenience, especially similar iterative arguments are needed in Step 2 and in Subsection 6.2.

Fix small $\delta > 0$. With $R = R(\delta) \gg 1$ and $K = K(\delta) \gg 1$ as in (5.38) and (5.42), let $\tau = \tau(R(\delta), K(\delta)) = \tau(\delta) > 0$ as in (5.61). Then, write $[0, T]$ as

$$[0, T] = \bigcup_{\ell=0}^{\lfloor T/\tau \rfloor} J_\ell,$$

where $J_\ell = [\ell\tau, (\ell+1)\tau] \cap [0, T]$. Let us work on the first interval $J_0 = [0, \tau]$. Note from (5.39) that

$$\|\vec{V}_{N,j}(0)\|_{\mathcal{A}_N \mathcal{H}_x^s} \leq \eta_0 \ll 1. \quad (5.62)$$

Then, in view of the continuity of $\|\vec{V}_{N,j}\|_{X_N^s([0,t])}$ in t , it follows from a standard continuity argument with (5.60), (5.61), and (5.62) that

$$\|\vec{V}_{N,j}\|_{X_N^s(J_0)} \leq 2C_1\eta_0 + 2C_3\eta,$$

which in turn implies

$$\|\vec{V}_{N,j}(\tau)\|_{\mathcal{A}_N \mathcal{H}_x^s} \leq 2C_1\eta_0 + 2C_3\eta \ll 1. \quad (5.63)$$

As a consequence, it follows from (5.63), (5.61), and a continuity argument that

$$\|\vec{V}_{N,j}(2\tau)\|_{\mathcal{A}_N \mathcal{H}_x^s} \leq \|\vec{V}_{N,j}\|_{X_N^s(J_1)} \leq (2C_1)^2\eta_0 + 2C_3(2C_1+1)\eta \ll 1,$$

which allows us to iterate the argument on the third interval J_2 .

After $\ell+1$ steps, we obtain

$$\begin{aligned} \|\vec{V}_{N,j}((\ell+1)\tau)\|_{\mathcal{A}_N \mathcal{H}_x^s} &\leq \|\vec{V}_{N,j}\|_{X_N^s(J_\ell)} \leq (2C_1)^\ell \eta_0 + 2C_3 \sum_{\ell'=0}^{\ell-1} (2C_1)^{\ell'} \cdot \eta \\ &\leq (2C_1)^\ell \eta_0 + 2C_3 \frac{(2C_1)^\ell}{2C_1-1} \eta. \end{aligned} \quad (5.64)$$

In order to cover the entire interval $[0, T]$, recalling that τ depends only on δ , it suffices to choose $\eta_0 = \eta_0(\delta)$ and $\eta = \eta(\delta)$ sufficiently small such that

$$\left(\left\lceil \frac{T}{\tau} \right\rceil + 1 \right) \left((2C_1)^{\lfloor \frac{T}{\tau} \rfloor} \eta_0 + 2C_3 \frac{(2C_1)^{\lfloor \frac{T}{\tau} \rfloor}}{2C_1-1} \eta \right) \leq \delta \ll 1. \quad (5.65)$$

Then, it follows from (5.64) and (5.65) that

$$\|\vec{V}_{N,j}\|_{\mathcal{A}_N C_T \mathcal{H}_x^s} \leq \sum_{\ell=0}^{\lfloor T/\tau \rfloor} \|\vec{V}_{N,j}\|_{X_N^s(J_\ell)} \leq \delta,$$

on $\Omega_N(R, K, \eta_0, \eta)$ defined in (5.47), for any $N \geq \max(N_1(\eta_0(\delta), \delta), N_2(\eta(\delta), \delta))$, where N_1 and N_2 are as in (5.40) and (5.45), respectively. In view of (5.48), this proves the convergence-in-probability (5.35).

• **Step 2:** In this part, we establish the convergence (1.26) for each *fixed* $j \in \mathbb{N}$:

$$\|\vec{V}_{N,j}\|_{C_T \mathcal{H}_x^s} = \|\vec{v}_{N,j} - \vec{v}_j\|_{C_T \mathcal{H}_x^s} \longrightarrow 0 \quad (5.66)$$

in probability as $N \rightarrow \infty$.

We first go over the construction of data sets indexed by $N \gg 1$. Fix $j \in \mathbb{N}$. Given $R \geq 1$, define the set $\Omega_{5,j}(R) \subset \Omega$ by setting

$$\Omega_{5,j}(R) = \left\{ \omega \in \Omega : \|\vec{v}_j\|_{C_T \mathcal{H}_x^s} \leq R \right\}. \quad (5.67)$$

Then, it follows from Chebyshev's inequality and (5.36) that there exists $R = R(\delta) \gg 1$, independent of $j \in \mathbb{N}$, such that

$$\mathbb{P}((\Omega_{5,j}(R))^c) < \frac{\delta}{10}. \quad (5.68)$$

Given small $\eta_0 > 0$, define $\Omega_{6,j,N}(\eta_0) \subset \Omega$ by setting

$$\Omega_{6,j,N}(\eta_0) = \left\{ \omega \in \Omega : \|\vec{V}_{N,j}(0)\|_{\mathcal{H}_x^s} \leq \eta_0 \right\}. \quad (5.69)$$

Then, from the assumption (1.24) on the initial data, there exists $N_3 = N_3(j, \eta_0, \delta) \in \mathbb{N}$ such that

$$\mathbb{P}((\Omega_{6,j,N}(\eta_0))^c) < \frac{\delta}{10}. \quad (5.70)$$

for any $N \geq N_3$. Given small $\eta > 0$, define the $\Omega_{7,N}(\eta) \subset \Omega$ by setting

$$\Omega_{7,N}(\eta) = \left\{ \omega \in \Omega : \|\vec{V}_{N,k}\|_{X_N^s(J)} \leq \eta \right\} \quad (5.71)$$

Then, it follows from the convergence-in-probability (5.35) established in Step 1 that, given $\delta > 0$, there exists $N_4 = N_4(\eta, \delta) \in \mathbb{N}$ such that

$$\mathbb{P}((\Omega_{7,N}(\eta))^c) < \frac{\delta}{10} \quad (5.72)$$

for any $N \geq N_4$. Given $j \in \mathbb{N}$ and $K \geq 1$, define the set $\Omega_{8,j,N}(K) \subset \Omega$ by setting

$$\Omega_{8,j,N}(K) = \left\{ \omega \in \Omega : \|\Psi_k \Psi_j\|_{\mathcal{A}_{N,k} C_T W_x^{-\varepsilon, \infty}} \leq K \right\}. \quad (5.73)$$

Then, it follows from Lemma 2.5 (i) (for the uniform (in j, k) moment bound on the $C_T W_x^{-\varepsilon, \infty}$ -norm of $\Psi_k \Psi_j$) and Chebyshev's inequality that there exists $K = K(\delta) \gg 1$, independent of $j, N \in \mathbb{N}$, such that

$$\mathbb{P}((\Omega_{8,j,N}(K))^c) < \frac{\delta}{10}. \quad (5.74)$$

Given small $\eta > 0$, define the set $\Omega_{9,j,N}(\eta) \subset \Omega$ by setting

$$\Omega_{9,j,N}(\eta) = \left\{ \omega \in \Omega : \|(\mathbf{v}, \Psi)\|_{\tilde{\mathcal{Y}}_{j,N}} \leq \eta \right\}, \quad (5.75)$$

where the $\tilde{\mathcal{Y}}_{j,N}$ -norm is defined by

$$\begin{aligned} \|(\mathbf{v}, \Psi)\|_{\tilde{\mathcal{Y}}_{j,N}} &= \left\| \frac{1}{N} \sum_{k=1}^N \Psi_k^2 \Psi_j \right\|_{L_T^2 W_x^{-\varepsilon, \infty}} \\ &\quad + \left\| \frac{1}{N} \sum_{k=1}^N \left(v_k \Psi_k \Psi_j - \mathbb{E}[v_k \Psi_k] \Psi_j \right) \right\|_{L_T^2 H_x^{s-1}}. \end{aligned} \quad (5.76)$$

Then, it follows from Lemmas 5.1 and 5.4 that there exists $N_5 = N_5(j, \eta, \delta) \in \mathbb{N}$ such that

$$\mathbb{P}((\Omega_{9,j,N}(\eta))^c) < \frac{\delta}{10} \quad (5.77)$$

for $N \geq N_5$.

We now turn to estimating the terms on the right-hand side of (5.33) for *fixed* $j \in \mathbb{N}$, which follows from a straightforward modification of the estimates in Step 1. In the following, we work on $\Omega_{j,N}(R, K, \eta_0, \eta)$ defined by

$$\begin{aligned} \Omega_{j,N}(R, K, \eta_0, \eta) &= \Omega_N(R, K, \eta_0, \eta) \cap \Omega_{5,j}(R) \cap \Omega_{6,j,N}(\eta_0) \\ &\quad \cap \Omega_{7,N}(\eta) \cap \Omega_{8,j,N}(K) \cap \Omega_{9,j,N}(\eta). \end{aligned} \quad (5.78)$$

where $\Omega_N(R, K, \eta_0, \eta)$, $\Omega_{5,j}(R)$, $\Omega_{6,j,N}(\eta_0)$, $\Omega_{7,N}(\eta)$, $\Omega_{8,j,N}(K)$, and $\Omega_{9,j,N}(\eta)$ are as in (5.47), (5.67), (5.69), (5.71), (5.73), and (5.75), respectively. Then, given small $\delta > 0$, by choosing $R = R(\delta)$ and $K = K(\delta)$ sufficiently large, it follows from (5.78) with (5.48), (5.68), (5.70), (5.72), (5.74), and (5.77) that

$$\mathbb{P}((\Omega_{j,N}(R, K, \eta))^c) < \delta \quad (5.79)$$

for any $N \geq N_*$, where

$$N_* = \max \left(N_1(\eta_0, \delta), N_2(\eta, \delta), N_3(j, \eta_0, \delta), N_4(\eta, \delta), N_5(j, \eta, \delta) \right). \quad (5.80)$$

Here, N_ℓ , $\ell = 1, \dots, 5$, are as in (5.40) (5.45), (5.70), (5.72), and (5.77), respectively.

Let $J \subset [0, T]$ be an interval with $|J| \leq 1$. Proceeding as in (5.50) with (5.49), (5.37), (5.43), (5.44), (5.67), and (5.71), we have

$$\begin{aligned} \|\vec{\mathcal{I}}(\mathcal{A}_{j,1})\|_{C_J \mathcal{H}_x^s} &\lesssim |J| \left(\|\vec{V}_{N,k}\|_{X_N^s(J)}^2 + \|\vec{v}_k\|_{X_N^s(J)}^2 \right) \|\vec{V}_{N,j}\|_{C_J \mathcal{H}_x^s} \\ &\quad + |J| \|\vec{V}_{N,k}\|_{X_N^s(J)} \left(\|\vec{V}_{N,k}\|_{X_N^s(J)} + \|\vec{v}_k\|_{X_N^s(J)} \right) \|\vec{v}_j\|_{C_J \mathcal{H}_x^s} \\ &\quad + |J|^{\frac{1}{2}} \left\| \frac{1}{N} \sum_{k=1}^N \left(v_k^2 - \mathbb{E}[v_k^2] \right) \right\|_{L_J^2 L_x^{\frac{3}{2-s}}} \|\vec{v}_j\|_{C_J \mathcal{H}_x^s} \\ &\lesssim |J| R^2 \|\vec{V}_{N,j}\|_{C_J \mathcal{H}_x^s} + |J|^{\frac{1}{2}} \eta R^2, \end{aligned} \quad (5.81)$$

provided that $\frac{1}{2} \leq s (< 1)$. Proceeding as in (5.52) with (5.51), (5.37), (5.41), (5.43), (5.44), (5.67), and (5.71), we have

$$\begin{aligned} \|\vec{\mathcal{I}}(\mathcal{A}_{j,2})\|_{C_J \mathcal{H}_x^s} &\lesssim |J| \|\Psi_k\|_{\mathcal{A}_N C_J W_x^{-\varepsilon, \infty}} \left(\|\vec{V}_{N,k}\|_{X_N^s(J)} + \|\vec{v}_k\|_{X_N^s(J)} \right) \|\vec{V}_{N,j}\|_{C_J \mathcal{H}_x^s} \\ &\quad + |J| \|\Psi_k\|_{\mathcal{A}_N C_J W_x^{-\varepsilon, \infty}} \|\vec{V}_{N,k}\|_{X_N^s(J)} \|\vec{v}_j\|_{C_J \mathcal{H}_x^s} \\ &\quad + |J|^{\frac{1}{2}} \left\| \frac{1}{N} \sum_{k=1}^N \left(\Psi_k v_k - \mathbb{E}[\Psi_k v_k] \right) \right\|_{L_J^2 W_x^{-\varepsilon, \frac{4}{2-s-\varepsilon}}} \|\vec{v}_j\|_{C_J \mathcal{H}_x^s} \\ &\lesssim |J| K R \|\vec{V}_{N,j}\|_{C_J \mathcal{H}_x^s} + |J|^{\frac{1}{2}} \eta K R, \end{aligned} \quad (5.82)$$

provided that $3\varepsilon \leq s$ (< 1). Similarly, proceeding as in (5.54) with (5.53), we have

$$\begin{aligned} & \|\vec{\mathcal{I}}(\mathcal{A}_{j,3})\|_{C_J \mathcal{H}_x^s} \\ & \lesssim |J| \left(\|\vec{V}_{N,k}\|_{X_N^s(J)} + \|\vec{v}_k\|_{X_N^s(J)} \right) \|\vec{V}_{N,k}\|_{X_N^s(J)} \|\Psi_j\|_{C_J W_x^{-\varepsilon, \infty}} \\ & \quad + |J|^{\frac{1}{2}} \left\| \frac{1}{N} \sum_{k=1}^N \left(v_k^2 - \mathbb{E}[v_k^2] \right) \right\|_{L_J^2 W_x^{\varepsilon, \frac{2}{2-s-\varepsilon}}} \|\Psi_j\|_{C_J W_x^{-\varepsilon, \infty}} \\ & \lesssim |J|^{\frac{1}{2}} \eta K R. \end{aligned} \tag{5.83}$$

Proceeding as in (5.55) with (5.34), we have

$$\begin{aligned} \|\vec{\mathcal{I}}(\mathcal{A}_{j,4})\|_{C_J \mathcal{H}_x^s} & \lesssim |J|^{\frac{1}{2}} \left\| \frac{1}{N} \sum_{k=1}^N : \Psi_k^2 : \right\|_{L_T^2 W_x^{-\varepsilon, \infty}} \left(\|\vec{V}_{N,j}\|_{C_J \mathcal{H}_x^s} + \|\vec{v}_j\|_{C_J \mathcal{H}_x^s} \right) \\ & \lesssim |J|^{\frac{1}{2}} \|\vec{V}_{N,j}\|_{C_J \mathcal{H}_x^s} + |J|^{\frac{1}{2}} \eta R, \end{aligned} \tag{5.84}$$

provided that $\varepsilon \leq s \leq 1 - \varepsilon$. Proceeding as in (5.57) with (5.56) and (5.73), we have

$$\begin{aligned} \|\vec{\mathcal{I}}(\mathcal{A}_{j,5})\|_{C_J \mathcal{H}_x^s} & \lesssim |J| \| : \Psi_k \Psi_j : \|_{\mathcal{A}_{N,k} C_J W_x^{-\varepsilon, \infty}} \|\vec{V}_{N,k}\|_{X_N^s(J)} + |J|^{\frac{1}{2}} \eta \\ & \lesssim |J|^{\frac{1}{2}} \eta K. \end{aligned} \tag{5.85}$$

Lastly, from (5.34) and (5.75), we have

$$\begin{aligned} \|\vec{\mathcal{I}}(\mathcal{A}_{j,6})\|_{C_J \mathcal{H}_x^s} & \lesssim |J|^{\frac{1}{2}} \left\| \frac{1}{N} \sum_{k=1}^N : \Psi_k^2 \Psi_j : \right\|_{L_J^2 H_x^{s-1}} \\ & \leq |J|^{\frac{1}{2}} \eta. \end{aligned} \tag{5.86}$$

Therefore, putting (5.33) (but starting at time t_0), (5.81), (5.82), (5.83), (5.84), (5.85), and (5.86) together with $J = [t_0, t_1] \subset [0, T]$ such that $|J| \leq 1$, we obtain

$$\begin{aligned} \|\vec{V}_{N,j}\|_{C_J \mathcal{H}_x^s} & \leq C_4 \|\vec{V}_{N,j}(t_0)\|_{\mathcal{H}_x^s} + C_5 |J|^{\frac{1}{2}} R (K + R) \|\vec{V}_{N,j}\|_{C_J \mathcal{H}_x^s} \\ & \quad + C_6 |J|^{\frac{1}{2}} \eta R (K + R). \end{aligned} \tag{5.87}$$

Suppose that

$$\|\vec{V}_{N,j}\|_{C_J \mathcal{H}_x^s} \leq 1$$

for some interval $J \subset [0, T]$. Then, it follows from (5.87) that there exists $\tau = \tau(R, K) \in (0, 1]$ such that

$$\|\vec{V}_{N,j}\|_{C_J \mathcal{H}_x^s} \leq 2C_4 \|\vec{V}_{N,j}(t_0)\|_{\mathcal{H}_x^s} + 2C_6 \eta,$$

provided that $|J| \leq \tau$. Then, by recalling that τ depends only on δ and by choosing $\eta_0 = \eta_0(\delta)$ and $\eta = \eta(\delta)$ sufficiently small, it follows from a slight modification of the iterative argument in Step 1 with (5.69) that

$$\|\vec{V}_{N,j}\|_{C_T \mathcal{H}_x^s} \leq \delta$$

on $\Omega_{j,N}(R, K, \eta_0, \eta)$, satisfying (5.79), for any $N \geq N_*$, where N_* is as in (5.80). This proves the convergence-in-probability (5.66), and therefore concludes the proof of Theorem 1.4 (i).

5.3. On the convergence rate under a higher moment assumption. In this subsection, we present a proof of Theorem 1.4 (ii) on a convergence rate under the higher moment assumption (1.30):

$$v_j \in L^6(\Omega; C([0, T]; H^s(\mathbb{T}^2))), \quad j \in \mathbb{N} \quad (5.88)$$

for some $T > 0$ and $\frac{1}{2} \leq s < 1$. Given $(v_j^{(0)}, v_j^{(1)}) \in L^6(\Omega; \mathcal{H}^s(\mathbb{T}^2))$, Proposition 4.1 guarantees (5.88) only for a short time. As in Subsection 5.2, however, the convergence argument presented below works on $[0, T]$ for any given $T > 0$, *assuming* that (5.88) (together with (1.28) and (1.29)) holds.

As compared to the presentation in Subsection 5.2, the only difference appears on how we handle the terms in (5.44) and (5.76). More precisely, we show that the terms in (5.44) and (5.76) converge in probability to 0 with a rate $N^{-\frac{1}{2}+\varepsilon_0}$ for any $\varepsilon_0 > 0$. In view of Lemma 5.1 which comes with a convergence rate of $N^{-\frac{1}{2}+\varepsilon_0}$, we only need to establish suitable replacements of Lemmas 5.2 and 5.4.

The following lemma establishes an extension of Lemma 5.4 under the higher moment assumption on v_j .

Lemma 5.5. *Let $\frac{1}{2} \leq s < 1$ and $\{(\Psi_j, v_j)\}_{j \in \mathbb{N}}$ be as in Lemma 5.2. Moreover, assume that*

$$v_j \in L^p(\Omega; C([0, T]; H^s(\mathbb{T}^2))) \quad (5.89)$$

for some $p > 2$. Then, given $T, \varepsilon_0 > 0$, we have, for each $j \in \mathbb{N}$,

$$N^{\frac{1}{2}-\varepsilon_0} \left\| \frac{1}{N} \sum_{k=1}^N \left(v_k : \Psi_k \Psi_j : - \mathbb{E}[v_k \Psi_k] \Psi_j \right) \right\|_{L_T^2 H_x^{s-1}} \longrightarrow 0 \quad (5.90)$$

in probability as $N \rightarrow \infty$, and

$$N^{\frac{1}{2}-\varepsilon_0} \left\| \frac{1}{N} \sum_{k=1}^N \left(v_k : \Psi_k \Psi_j : - \mathbb{E}[v_k \Psi_k] \Psi_j \right) \right\|_{\mathcal{A}_{N,j} L_T^2 H_x^{s-1}} \longrightarrow 0 \quad (5.91)$$

in probability as $N \rightarrow \infty$.

Proof. We only consider (5.91) since (5.90) follows from a similar consideration. From a slight modification of the proof of Lemma 5.4, it is easy to see that the contribution from the case $k = j$ satisfies (5.91). Thus, we only consider the case $k \neq j$ in the following.

Set

$$X_{k,j} = v_k \Psi_k \Psi_j - \mathbb{E}[v_k \Psi_k] \Psi_j$$

Then, from the mutual independence of $\{(\Psi_k, v_k)\}_{k \in \mathbb{N}}$, we have

$$\mathbb{E}[X_{k,j}] = \mathbb{E}[v_k \Psi_k - \mathbb{E}[v_k \Psi_k]] \cdot \mathbb{E}[\Psi_j] = 0 \quad (5.92)$$

and

$$\mathbb{E}[X_{k,j} \cdot X_{k',j}] = \mathbb{E}[(v_k \Psi_k - \mathbb{E}[v_k \Psi_k])(v_{k'} \Psi_{k'} - \mathbb{E}[v_{k'} \Psi_{k'}])] \cdot \mathbb{E}[\Psi_j^2] = 0 \quad (5.93)$$

for any $k \neq k'$ with $k, k' \neq j$.

A slight modification of (5.14) with (5.89) yields

$$\begin{aligned} \left\| \left\| v_k \Psi_k \Psi_j \right\|_{L_T^2 H_x^{s-1}} \right\|_{L_\omega^2} &\lesssim_T \|v_k\|_{L_\omega^p C_T H_x^s} \|\Psi_k \Psi_j\|_{L_\omega^{\frac{2p}{p-2}} C_T W_x^{-\varepsilon, \infty}} \\ &\lesssim 1, \end{aligned} \quad (5.94)$$

uniformly in $j, k \in \mathbb{N}$. Similarly, a minor modification of (5.15) yields

$$\left\| \mathbb{E}[v_k \Psi_k] \Psi_j \right\|_{L_T^2 H_x^{s-1}} \Big\|_{L_\omega^2} \lesssim_T 1, \quad (5.95)$$

uniformly in $j, k \in \mathbb{N}$. Hence, proceeding as in (5.7) with (5.92), (5.93), (5.94), and (5.95), we have

$$\left\| \frac{1}{N} \sum_{\substack{k=1 \\ k \neq j}}^N X_{k,j} \right\|_{L_\omega^2 \mathcal{A}_{N,j} L_T^2 H_x^{s-1}} \leq \frac{1}{N^{\frac{1}{2}}} \|X_{k,j}\|_{\mathcal{A}_N^{(2)} L_\omega^2 L_T^2 H_x^{s-1}} \lesssim_T \frac{1}{N^{\frac{1}{2}}}, \quad (5.96)$$

where $\mathcal{A}_N^{(2)} B$ is as in (2.2). Therefore, the convergence-in-probability (5.91) follows from Chebyshev's inequality with (5.96). \square

In the proof of Lemma 5.5, we crucially used the fact that the space-time norm is L^2 -based. Since the terms treated in Lemma 5.2 involve the L_x^r -base Sobolev spaces with $r > 2$, we can not proceed as in the proof of Lemma 5.5. We instead make the observation that the terms treated in Lemma 5.2 only appear in (5.49), (5.51), and (5.53):

$$\begin{aligned} & \frac{1}{N} \sum_{k=1}^N \left(v_k^2 - \mathbb{E}[v_k^2] \right) v_j, \quad \frac{2}{N} \sum_{k=1}^N \left(\Psi_k v_k - \mathbb{E}[\Psi_k v_k] \right) v_j \\ \text{and} \quad & \frac{1}{N} \sum_{k=1}^N \left(v_k^2 - \mathbb{E}[v_k^2] \right) \Psi_j \end{aligned} \quad (5.97)$$

which do not involve the difference $V_{N,j}$ defined in (5.32). The following lemma establishes convergence in probability of these terms in a direct manner.

Lemma 5.6. *Let $\frac{1}{2} \leq s < 1$ and $T > 0$. Let $\{(\Psi_j, v_j)\}_{j \in \mathbb{N}}$ be as in Lemma 5.2. Moreover, assume (5.88):*

$$v_j \in L^6(\Omega; C([0, T]; H^s(\mathbb{T}^2))). \quad (5.98)$$

Then, given $\varepsilon_0 > 0$, we have, for each $j \in \mathbb{N}$,

$$N^{\frac{1}{2}-\varepsilon_0} \left\| \frac{1}{N} \sum_{k=1}^N \left(\Psi_k v_k - \mathbb{E}[\Psi_k v_k] \right) v_j \right\|_{L_T^2 H_x^{s-1}} \longrightarrow 0, \quad (5.99)$$

$$N^{\frac{1}{2}-\varepsilon_0} \left\| \frac{1}{N} \sum_{k=1}^N \left(v_k^2 - \mathbb{E}[v_k^2] \right) v_j \right\|_{L_T^2 H_x^{s-1}} \longrightarrow 0, \quad (5.100)$$

$$N^{\frac{1}{2}-\varepsilon_0} \left\| \frac{1}{N} \sum_{k=1}^N \left(v_k^2 - \mathbb{E}[v_k^2] \right) \Psi_j \right\|_{L_T^2 H_x^{s-1}} \longrightarrow 0, \quad (5.101)$$

in probability, as $N \rightarrow \infty$, and

$$N^{\frac{1}{2}-\varepsilon_0} \left\| \frac{1}{N} \sum_{k=1}^N \left(\Psi_k v_k - \mathbb{E}[\Psi_k v_k] \right) v_j \right\|_{\mathcal{A}_{N,j} L_T^2 H_x^{s-1}} \longrightarrow 0, \quad (5.102)$$

$$N^{\frac{1}{2}-\varepsilon_0} \left\| \frac{1}{N} \sum_{k=1}^N \left(v_k^2 - \mathbb{E}[v_k^2] \right) v_j \right\|_{\mathcal{A}_{N,j} L_T^2 H_x^{s-1}} \longrightarrow 0, \quad (5.103)$$

$$N^{\frac{1}{2}-\varepsilon_0} \left\| \frac{1}{N} \sum_{k=1}^N \left(v_k^2 - \mathbb{E}[v_k^2] \right) \Psi_j \right\|_{\mathcal{A}_{N,j} L_T^2 H_x^{s-1}} \longrightarrow 0, \quad (5.104)$$

in probability, as $N \rightarrow \infty$.

Proof. Since the claim follows from a slight modification of the proof of Lemma 5.5, we keep the discussion brief. We only consider (5.103) since (5.99), (5.100), (5.101), (5.102), and (5.104) follow from a similar consideration. We first note that, by arguing as in the proof of Lemma 5.4, the contributions from the case $k = j$ can be treated easily and thus we omit details. In the following, we assume $k \neq j$.

Set

$$Y_{k,j} = \left(v_k^2 - \mathbb{E}[v_k^2] \right) v_j \quad (5.105)$$

Then, from the mutual independence of $\{(\Psi_k, v_k)\}_{k \in \mathbb{N}}$, we have

$$\mathbb{E}[Y_{k,j}] = \mathbb{E}[Y_{k,j} \cdot Y_{k',j}] = 0 \quad (5.106)$$

for any $k \neq k'$ with $k, k' \neq j$; see (5.92) and (5.93).

Proceeding as in (3.6) with (5.98), we have

$$\begin{aligned} \left\| \|v_k^2 v_j\|_{L_T^2 H_x^{s-1}} \right\|_{L_\omega^2} &\lesssim_T \|v_k\|_{L_\omega^6 C_T H_x^s}^2 \|v_j\|_{L_\omega^6 C_T H_x^s} \\ &\lesssim 1, \end{aligned} \quad (5.107)$$

uniformly in $j, k \in \mathbb{N}$. Similarly, a minor modification of (5.15) yields

$$\left\| \|\mathbb{E}[v_k^2] v_j\|_{L_T^2 H_x^{s-1}} \right\|_{L_\omega^2} \lesssim 1, \quad (5.108)$$

uniformly in $j, k \in \mathbb{N}$. Then, the claim (5.103) follows from proceeding as in the proof of Lemma 5.5 with (5.105), (5.106), (5.107), and (5.108). We omit details. \square

With Lemmas 5.5 and 5.6, Theorem 1.4 (ii) follows from a straightforward modification of the argument presented in Subsection 5.2. More precisely, given small $\varepsilon_0 > 0$, we need to prove the following two convergence results under the assumption (5.88) together with the assumed convergence of initial data (1.28) and (1.29) with the convergence rate of $N^{-\frac{1}{2}+\varepsilon_0}$:

(i) in Step 1, instead of (5.35), we prove

$$N^{\frac{1}{2}-\varepsilon_0} \|\vec{V}_{N,j}\|_{X_N^s(T)} = N^{\frac{1}{2}-\varepsilon_0} \|\vec{v}_{N,j} - \vec{v}_j\|_{X_N^s(T)} \longrightarrow 0$$

in probability as $N \rightarrow \infty$,

(ii) in Step 2, instead (5.66), we prove, for each fixed $j \in \mathbb{N}$:

$$N^{\frac{1}{2}-\varepsilon_0} \|\vec{V}_{N,j}\|_{C_T \mathcal{H}_x^s} = N^{\frac{1}{2}-\varepsilon_0} \|\vec{v}_{N,j} - \vec{v}_j\|_{C_T \mathcal{H}_x^s} \longrightarrow 0$$

in probability, as $N \rightarrow \infty$.

In Step 1, the only modification appears in (5.50), (5.52), and (5.54), where we treat the terms in (5.97) directly by Lemmas 5.5 and 5.6. Similarly, in Step 2, the only modification appears in (5.81), (5.82), and (5.83), where we once again treat the terms in (5.97) directly by Lemmas 5.5 and 5.6. Since the required modifications are straightforward, we omit details; see Subsection 6.2 for an example of an argument where we obtain a convergence rate (in a slightly different setting). This concludes the proof of Theorem 1.4 (ii).

Remark 5.7. Let $T \gg 1$. In Theorem 1.4(ii), we only assume the existence of a solution $\{(u_j, \partial_t u_j)\}_{j \in \mathbb{N}}$ to the mean-field SdNLW (1.20) on the time interval $[0, T]$ along with the higher moment assumption (1.30). In particular, we did not assume existence of a solution $\{(u_{N,j}, \partial_t u_{N,j})\}_{j=1}^N$ to HLSM $_N$ (1.10) on the time interval $[0, T]$ (whose local-in-time existence is guaranteed by Theorem 1.1(i)). We point out that an iterative application of the local-in-time convergence argument together with the blowup alternative (3.4) guarantees existence of the solution $\{(u_{N,j}, \partial_t u_{N,j})\}_{j=1}^N$ to HLSM $_N$ (1.10) on the time interval $[0, T]$ for sufficiently large $N = N(\omega, T) \gg 1$. See [27] for an argument, where existence (with a control on the norm) of a solution to a limiting equation, say on a time interval $[0, T]$, guarantees existence (and a control on the norm) of converging solutions on the same interval.

6. MEAN-FIELD LIMIT OF INVARIANT GIBBS DYNAMICS

In this section, we present a proof of Theorem 1.6(iii) on convergence of the invariant Gibbs dynamics for HLSM $_N$ (1.42) to the limiting invariant dynamics for the mean-field SdNLW (1.43), given by (1.38). In Subsection 6.1, we go over the construction of the initial data set on an appropriate probability space $(\Omega', \mathcal{F}', \mathbb{P}')$. In Subsection 6.2, we then establish convergence of the invariant Gibbs dynamics for HLSM $_N$ (1.42).

6.1. On the Gibbsian initial data. In this subsection, we go over the construction of the initial data set stated in Theorem 1.6(iii), which follows from the following proposition.

Proposition 6.1. *Let $m > 0$. Then, there exist a probability space $(\Omega_1, \mathcal{F}_1, \mathbb{P}_1)$ and random variables $\{\phi_j^{(0)}\}_{j \in \mathbb{N}}$ and $\vec{\psi}_N = \{\psi_{N,j}\}_{j=1}^N$, $N \in \mathbb{N}$, on $(\Omega_1, \mathcal{F}_1, \mathbb{P}_1)$ such that the family $\{\phi_j^{(0)}\}_{j \in \mathbb{N}}$ is independent and*

$$\text{Law}(\phi_j^{(0)}) = \mu_1 \text{ for all } j \in \mathbb{N} \quad \text{and} \quad \text{Law}(\vec{\psi}_N) = \rho_N, \quad (6.1)$$

where μ_1 is the massive Gaussian free field with mass $m > 0$ defined in (1.5) and ρ_N is the $O(N)$ linear sigma model defined in (1.35). Furthermore, we have

$$\sup_{j=1, \dots, N} \left\| \|\psi_{N,j} - \phi_j^{(0)}\|_{H_x^1} \right\|_{L^2(\Omega_1)} \lesssim N^{-\frac{1}{2}}. \quad (6.2)$$

By assuming Proposition 6.1, we first carry out the construction of the initial data sets stated in Theorem 1.6(iii.a) and (iii.b).

By extending the probability space $(\Omega_1, \mathcal{F}_1, \mathbb{P}_1)$ constructed in Proposition 6.1, let $\{\phi_j^{(1)}\}_{j \in \mathbb{N}}$ be a family of independent random variables on Ω_1 such that $\text{Law}(\phi_j^{(1)}) = \mu_0$ for each $j \in \mathbb{N}$, where μ_0 is the white noise measure on \mathbb{T}^2 defined in (1.33) with $s = 0$. Given $N \in \mathbb{N}$, we set

$$\vec{\phi}_N^{(0)} = \{\phi_{N,j}^{(0)}\}_{j=1}^N \stackrel{\text{def}}{=} \{\phi_j^{(0)}\}_{j=1}^N \quad \text{and} \quad \vec{\phi}_N^{(1)} = \{\phi_{N,j}^{(1)}\}_{j=1}^N \stackrel{\text{def}}{=} \{\phi_j^{(1)}\}_{j=1}^N,$$

where $\{\phi_j^{(0)}\}_{j \in \mathbb{N}}$ is as in Proposition 6.1. For each $N \in \mathbb{N}$, we define $\vec{v}_N^{(0)} = \{v_{N,j}^{(0)}\}_{j=1}^N$ by setting

$$v_{N,j}^{(0)} = \psi_{N,j} - \phi_{N,j}^{(0)} = \psi_{N,j} - \phi_j^{(0)}, \quad j = 1, \dots, N, \quad (6.3)$$

where $\vec{\psi}_N = \{\psi_{N,j}\}_{j=1}^N$ is as in Proposition 6.1. Then, from (6.2), we have

$$\sup_{j=1, \dots, N} \left\| \|v_{N,j}^{(0)}\|_{H_x^1} \right\|_{L^2(\Omega_1)} \lesssim N^{-\frac{1}{2}},$$

yielding (1.47).

Let $\{\xi_j\}_{j \in \mathbb{N}}$ be a family of independent space-time white noise on $\mathbb{R}_+ \times \mathbb{T}^2$ which is also independent from $\{\phi_j^{(0)}\}_{j \in \mathbb{N}}$, $\{\phi_j^{(1)}\}_{j \in \mathbb{N}}$, and $\{\vec{\psi}_N\}_{N \in \mathbb{N}}$. Namely, we may assume that ξ_j is a $\mathcal{D}'(\mathbb{R}_+ \times \mathbb{T})$ -valued random variable defined on another probability space $(\Omega_2, \mathcal{F}_2, \mathbb{P}_2)$. Then, by setting $\mathbb{P}' = \mathbb{P}_1 \otimes \mathbb{P}_2$ and $\Omega' = \Omega_1 \times \Omega_2$, we complete the construction of the initial data sets stated in Theorem 1.6 (iii).

We now turn to a proof of Proposition 6.1. Let us first introduce some notations and preliminary lemmas. Let ν_1 and ν_2 be two probability measures on a Banach space B and let $H \subset B$ be a closed subspace. We define the Wasserstein-2 distance of ν_1 and ν_2 with respect to H by setting

$$\left(W_H(\nu_1, \nu_2)\right)^2 = \inf_{X \sim \nu_1, Y \sim \nu_2} \mathbb{E}[\|X - Y\|_H^2], \quad (6.4)$$

where $X \sim \nu_1$ and $Y \sim \nu_2$ mean that X and Y are B -valued random variables with $\text{Law}(X) = \nu_1$ and $\text{Law}(Y) = \nu_2$, and the expectation is taken with respect to the probability measure associated with the coupling of X and Y . Equivalently, it can be written as

$$\left(W_H(\nu_1, \nu_2)\right)^2 = \inf_{\mathbf{p} \in \Pi(\nu_1, \nu_2)} \int_{B \times B} \|x - y\|_H^2 d\mathbf{p}(x, y), \quad (6.5)$$

where $\Pi(\nu_1, \nu_2)$ is the set of probability measures \mathbf{p} on $B \times B$ whose first and second marginals are given by ν_1 and ν_2 , namely,

$$\int_{y \in B} d\mathbf{p}(x, y) = d\nu_1(x) \quad \text{and} \quad \int_{x \in B} d\mathbf{p}(x, y) = d\nu_2(y). \quad (6.6)$$

See, for example, [17, 16], where they take (B, H, μ) to be an abstract Wiener space (for a Gaussian measure μ with its Cameron-Martin space given by H) and ν_j , $j = 1, 2$, to be probability measures on B . In this case, the infimum in (6.4) and (6.5) is indeed attained.

In our setting, we take $B = (H^{-\varepsilon}(\mathbb{T}^2))^{\otimes N}$ and consider two probability measures: the Gaussian measure $\mu_1^{\otimes N}$, where μ_1 is the massive Gaussian free field with mass $m > 0$ defined in (1.5), and the coupled Φ_2^4 -measure ρ_N in (1.35). Then, we take H to be its Cameron-Martin space of $\mu_1^{\otimes N}$, namely, $H = (H^1(\mathbb{T}^2))^{\otimes N}$ endowed with the norm:

$$\|\vec{f}\|_{H^1(\mathbb{T}^2)^{\otimes N}}^2 = \sum_{j=1}^N \|f_j\|_{H^1(\mathbb{T}^2)}^2$$

for $\vec{f} = (f_1, \dots, f_N)$.

We now state a crucial lemma from a recent work by Delgadino and Smith [14].

Lemma 6.2. *Let $m > 0$. Given $N \in \mathbb{N}$, let $\mu_1^{\otimes N}$ and ρ_N be as above. Then, we have*

$$\sup_{N \in \mathbb{N}} W_{(H^1(\mathbb{T}^2))^{\otimes N}}(\mu_1^{\otimes N}, \rho_N) < \infty.$$

Lemma 6.2 plays a key role in studying the limiting behavior of the coupled Φ_2^4 -measure ρ_N in (1.35) as $N \rightarrow \infty$. It was first proven by Shen, Smith, Zhu, and Zhu [47, Theorem 5.11; see also (5.14)] for large masses $m \gg 1$. In a recent work [14], Delgadino and Smith extended this result to arbitrarily small masses $m > 0$ by making use of Talagrand's inequality [54, 44, 17].

It follows from Lemma 6.2 with (6.4) that, for each $N \in \mathbb{N}$, there exist random vectors $\vec{\phi}_N = \{\phi_{N,j}\}_{j=1}^N \sim \mu_1^{\otimes N}$ and $\vec{\psi}_N = \{\psi_{N,j}\}_{j=1}^N \sim \rho_N$ such that

$$\mathbb{E}[\|\vec{\phi}_N - \vec{\psi}_N\|_{(H^1(\mathbb{T}^2))^{\otimes N}}^2] \lesssim 1, \quad (6.7)$$

uniformly in $N \in \mathbb{N}$. We point out that Proposition 6.1 does not immediately follow from (6.7) since $\vec{\phi}_N$ depends on $N \in \mathbb{N}$. In order to amend this issue, we need the following gluing lemma, which is an infinite-dimensional generalization of [59, Lemma 7.6]. Given a measurable space E , we denote by $\mathcal{P}(E)$ the set of probability measures on E .

Lemma 6.3. *Let I be an index set. Let E and E_j , $j \in I$, be Polish spaces. Given $\mu \in \mathcal{P}(E)$, suppose that for each $j \in I$, ν_j is a probability measure on $E \times E_j$ with marginal μ on E in the sense of (6.6). Then, there exists a probability measure $\pi \in \mathcal{P}(E \times \prod_{j \in I} E_j)$ with marginal ν_j on $E \times E_j$ for each $j \in I$.*

Proof. Let $J \subset I$ be a finite set. By the disintegration-of-measure theorem ([19, Theorem on p. 148] and [18, Proposition 452O]), for each $j \in J$, there exists a measurable map $x \in E \mapsto \pi_j^x \in \mathcal{P}(E_j)$, uniquely determined μ -almost everywhere, such that

$$\nu_j(A_j) = \int_E \pi_j^x(A_j^x) d\mu(x)$$

for any measurable set $A_j \subset E \times E_j$, where

$$A_j^x = \{y \in E_j : (x, y) \in A_j\}.$$

We then define $\pi_J \in \mathcal{P}(E \times \prod_{j \in J} E_j)$ by setting

$$\pi_J(A_J) = \int_E \left(\bigotimes_{j \in J} \pi_j^x \right) (A_J^x) d\mu(x)$$

for any measurable set $A_J \subset E \times \prod_{j \in J} E_j$, where

$$A_J^x = \left\{ y \in \prod_{j \in J} E_j : (x, y) \in A_J \right\}.$$

It is not difficult to check that π_J has marginal ν_j on $E \times E_j$ for each $j \in J$. Furthermore, one can easily see that the collection of the probability measures $\{\pi_J : J \subset I \text{ is finite}\}$ is consistent. Then, by the Kolmogorov extension theorem ([1, Appendix D]), there exists a measure $\pi \in \mathcal{P}(E \times \prod_{j \in I} E_j)$ that satisfies the desired properties. \square

We also need the following lemma; see [52, (v) on p. 52].

Lemma 6.4. *A countable product of Polish spaces is Polish.*

We are now ready to present a proof of Proposition 6.1.

Proof of Proposition 6.1. As observed above, it follows from Lemma 6.2 that, for each $N \in \mathbb{N}$, there exist $\vec{\phi}_N = \{\phi_{N,j}\}_{j=1}^N \sim \mu_1^{\otimes N}$ and $\vec{\psi}_N = \{\psi_{N,j}\}_{j=1}^N \sim \rho_N$ such that (6.7) holds. For each $N \in \mathbb{N}$, we extend $\vec{\phi}_N$ to a sequence of independent random variables $\vec{\phi}_N^\infty = \{\phi_{N,j}^\infty\}_{j \in \mathbb{N}}$ by setting $\phi_{N,j}^\infty = \phi_{N,j}$ for $j = 1, \dots, N$ and $\text{Law}(\phi_{N,j}) = \mu_1$ for $j \geq N + 1$. Then, $\vec{\phi}_N^\infty$ is an $(H^{-\varepsilon}(\mathbb{T}^2))^{\otimes \mathbb{N}}$ -valued random variable with the law $\mu_1^{\otimes \mathbb{N}}$. Note from Lemma 6.4 that the space $(H^{-\varepsilon}(\mathbb{T}^2))^{\otimes \mathbb{N}}$ is a Polish space.

We now apply Lemma 6.3 by taking $I = \mathbb{N}$, $E = (H^{-\varepsilon}(\mathbb{T}^2))^{\otimes \mathbb{N}}$, $E_N = (H^{-\varepsilon}(\mathbb{T}^2))^{\otimes N}$, $\mu = \mu_1^{\otimes \mathbb{N}}$, and $\nu_N = \text{Law}(\vec{\phi}_N^\infty, \vec{\psi}_N)$ for each $N \in \mathbb{N}$. By Lemma 6.3, there exists a probability measure π on $E \times \prod_{N \in \mathbb{N}} E_N$ with marginal ν_N on $E \times E_N$ for each $N \in \mathbb{N}$.

The probability measure π then gives rise to random variables $\vec{\phi}^{(0)} = \{\phi_j^{(0)}\}_{j \in \mathbb{N}}$ on $E = (H^{-\varepsilon}(\mathbb{T}^2))^{\otimes \mathbb{N}}$ and $\vec{\psi}_N = \{\psi_{N,j}\}_{j=1}^N$ on $E_N = (H^{-\varepsilon}(\mathbb{T}^2))^{\otimes N}$ for each $N \in \mathbb{N}$, satisfying (6.1), such that, for each $N \in \mathbb{N}$, we have

$$\sup_{N \in \mathbb{N}} \mathbb{E} \left[\left\| \{\phi_j^{(0)}\}_{j=1}^N - \vec{\psi}_N \right\|_{H^1(\mathbb{T}^2)^{\otimes N}}^2 \right] < \infty. \quad (6.8)$$

It follows from the symmetry in (1.35) that the components $\psi_{N,j}$, $j = 1, \dots, N$, have the same distribution. Hence the desired bound (6.2) follows from (6.8). \square

6.2. Convergence of invariant Gibbs dynamics. In this subsection, we prove convergence of the invariant Gibbs dynamics for HL SM_N (1.42) claimed in Theorem 1.6 (iii.c). Since the proof follows closely to that of a combination of the local well-posedness argument (the proof of Proposition 3.1) presented in Subsection 3.1 and the iterative argument presented in Subsection 5.2, we keep our discussion brief here, indicating necessary modifications.

Let $\{(\phi_j^{(0)}, \phi_j^{(1)})\}_{j \in \mathbb{N}}$ and $\vec{\psi}_N = \{\psi_{N,j}\}_{j=1}^N$, $N \in \mathbb{N}$, be as in Theorem 1.6 (iii) constructed in Subsection 6.1, and let $\vec{v}_N^{(0)} = \{v_{N,j}^{(0)}\}_{j=1}^N$ be as in (6.3). Let Φ_j , $j \in \mathbb{N}$, be the solution to (1.38).

By writing (1.41) in the Duhamel formulation, we have

$$\begin{aligned} v_{N,j} = S(t)(v_{N,j}^{(0)}, 0) - \frac{1}{N} \sum_{k=1}^N \mathcal{I} \Big(& v_{N,k}^2 v_{N,j} + 2\Phi_k v_{N,k} v_{N,j} + v_{N,k}^2 \Phi_j \\ & + : \Phi_k^2 : v_{N,j} + 2v_{N,k} : \Phi_k \Phi_j : + : \Phi_k^2 \Phi_j : \Big), \end{aligned} \quad (6.9)$$

where $S(t)$ and \mathcal{I} are as in (2.5) and (2.6), respectively. Denote by u_j the limit of $u_{N,j} = \Phi_j + v_{N,j}$. As mentioned in Subsection 1.3, the (formal) limit u_j solves (1.43) with initial data $(\phi_j^{(0)}, \phi_j^{(1)})$. Hence, recalling that $u_j \equiv \Phi_j$ is a solution to (1.43) with $(u_j, \partial_t u_j)|_{t=0} = (\phi_j^{(0)}, \phi_j^{(1)})$ and that the uniqueness of $(u_j, \partial_t u_j)$ holds in the class (1.46), we conclude that $(u_j, \partial_t u_j) = (\Phi_j, \partial_t \Phi_j)$. Namely, we have $v_j = u_j - \Phi_j \equiv 0$ for any $j \in \mathbb{N}$, which in particular (and trivially) implies

$$(v_j, \partial_t v_j) = (0, 0) \in L^p(\Omega; C([0, T]; \mathcal{H}^s(\mathbb{T}^2)))$$

for any $p \geq 1$ and $T > 0$. This is the main difference and the simplification as compared to Subsection 5.2.

• **Step 1:** In this first step, we establish the following convergence for each $T \gg 1$; for any $\varepsilon_0 > 0$,

$$N^{\frac{1}{2}-\varepsilon_0} \|\vec{v}_{N,j}\|_{X_N^s(T)} \longrightarrow 0 \quad (6.10)$$

in probability (with respect to \mathbb{P}') as $N \rightarrow \infty$, where $X_N^s(T)$ is as in (3.3).

Fix a target time $T \gg 1$, $\varepsilon_0 > 0$, and small $\delta > 0$. Let $\frac{1}{2} \leq s < 1$. In the following, we suppress dependence on T , s , ε_0 , and small $\varepsilon > 0$. Given small $\eta_0 > 0$, define the set $\Omega'_{1,N}(\eta_0) \subset \Omega'$ by setting

$$\Omega'_{1,N}(\eta_0) = \left\{ \omega' \in \Omega' : N^{\frac{1}{2}-\varepsilon_0} \|v_{N,j}^{(0)}\|_{\mathcal{A}_N H_x^s} \leq \eta_0 \right\}. \quad (6.11)$$

where $\mathcal{A}_N B$ is as in (1.11). Then, it follows from Chebyshev's inequality and (6.2) with (6.3) that there exists $N_1 = N_1(\eta_0, \delta) \in \mathbb{N}$ such that

$$\mathbb{P}'((\Omega'_{1,N}(\eta_0))^c) < \frac{\delta}{6}. \quad (6.12)$$

for any $N \geq N_1$. Next, given $K \geq 1$ define the sets $\Omega'_{2,N}(K) \subset \Omega'$ by setting

$$\Omega'_{2,N}(K) = \left\{ \omega' \in \Omega' : \|\Phi_N\|_{\mathcal{Z}_N^\varepsilon(T)} \leq K \right\}, \quad (6.13)$$

where the $\mathcal{Z}_N^\varepsilon(T)$ -norm is as in (2.17) and (2.18), and Φ_N is given by

$$\Phi_N = \{ \Phi_j, : \Phi_k \Phi_j : , \vec{0}_{j,k} \}_{j,k=1}^N$$

with $\vec{0}_{j,k} = 0$ for any $j, k = 1, \dots, N$. Then, it follows from Lemma 2.5 (iii) that there exists $K = K(\delta) \gg 1$, independent of $N \in \mathbb{N}$, such that

$$\mathbb{P}'((\Omega'_{2,N}(K))^c) < \frac{\delta}{6}. \quad (6.14)$$

Given small $\eta > 0$, define the set $\Omega'_{3,N}(\eta) \subset \Omega'$ by setting

$$\Omega'_{3,N}(\eta) \subset \Omega' = \left\{ \omega' \in \Omega' : N^{\frac{1}{2}-\varepsilon_0} \left\| \frac{1}{N} \sum_{k=1}^N : \Phi_k^2 \Phi_j : \right\|_{\mathcal{A}_{N,j} L_T^2 W_x^{-\varepsilon, \infty}} \leq \eta \right\}. \quad (6.15)$$

Then, it follows from Lemma 5.1 that there exists $N_2 = N_2(\eta, \delta) \in \mathbb{N}$ such that

$$\mathbb{P}'((\Omega'_{3,N}(\eta))^c) < \frac{\delta}{6} \quad (6.16)$$

for any $N \geq N_2$. In the following, we work on $\Omega'_N(K, \eta_0, \eta)$ defined by

$$\Omega'_N(K, \eta_0, \eta) = \Omega'_{1,N}(\eta_0) \cap \Omega'_{2,N}(K) \cap \Omega'_{3,N}(\eta). \quad (6.17)$$

Given small $\delta > 0$, by choosing $K = K(\delta)$ sufficiently large, it follows from (6.17) with (6.12), (6.14), and (6.16) that

$$\mathbb{P}((\Omega'_N(K, \eta_0, \eta))^c) < \frac{\delta}{2} \quad (6.18)$$

for any $N \geq \max(N_1, N_2)$, where $N_1(\eta_0, \delta)$ and $N_2(\eta, \delta)$ are as in (6.12) and (6.16), respectively.

We now consider (6.9) on an interval $J = [t_0, t_1] \subset [0, T]$ with $|J| \leq 1$, starting at time t_0 . By applying a slight modification of (3.6)-(3.12), where $\mathbf{Z}_N = \{Z_j^{(1)}, Z_{k,j}^{(2)}, Z_{k,j}^{(3)}\}_{j,k=1}^N$ are as in (2.20), with (6.13) and (6.15) we obtain

$$\begin{aligned} N^{\frac{1}{2}-\varepsilon_0} \|\vec{v}_{N,j}\|_{X_N^s(J)} &\leq C_1 N^{\frac{1}{2}-\varepsilon_0} \|\vec{v}_{N,j}(t_0)\|_{\mathcal{A}_N \mathcal{H}_x^s} \\ &\quad + C_2 |J| \left(\|\vec{v}_{N,k}\|_{X_N^s(J)}^2 + K^2 \right) N^{\frac{1}{2}-\varepsilon_0} \|\vec{v}_{N,j}\|_{X_N^s(J)} \\ &\quad + C_3 |J|^{\frac{1}{2}} \eta, \end{aligned} \quad (6.19)$$

where the $X_N^s(J)$ -norm is as in (5.46). Suppose that

$$\|\vec{v}_{N,j}\|_{X_N^s(J)} \leq 1$$

for some interval $J \subset [0, T]$. Then, it follows from (6.19) that there exists $\tau = \tau(K(\delta)) = \tau(\delta) \in (0, 1]$ such that

$$N^{\frac{1}{2}-\varepsilon_0} \|\vec{v}_{N,j}\|_{X_N^s(J)} \leq 2C_1 N^{\frac{1}{2}-\varepsilon_0} \|\vec{v}_{N,j}(t_0)\|_{\mathcal{A}_N \mathcal{H}_x^s} + 2C_3 \eta, \quad (6.20)$$

provided that $|J| \leq \tau$. Compare (6.20) with (5.61). Then, by noting that τ depends only on δ , and choosing $\eta_0 = \eta_0(\delta)$ and $\eta = \eta(\delta)$ sufficiently small, the desired convergence-in-probability (6.10) follows from (6.11) and a slight modification of the iterative argument in Step 1 of the proof of Theorem 1.4 (i) presented in Subsection 5.2. We omit details.

• **Step 2:** In the following, we briefly discuss how to obtain the following convergence in probability for each *fixed* $j \in \mathbb{N}$:

$$N^{\frac{1}{2}-\varepsilon_0} \|\vec{v}_{N,j}\|_{C_T \mathcal{H}_x^s} \longrightarrow 0 \quad (6.21)$$

as $N \rightarrow \infty$.

We first go over the construction of data sets indexed by $N \gg 1$. Fix $j \in \mathbb{N}$. Given small $\eta_0 > 0$, define the set $\Omega'_{4,j,N}(\eta_0) \subset \Omega'$ by setting

$$\Omega'_{4,j,N}(\eta_0) = \left\{ \omega' \in \Omega' : N^{\frac{1}{2}-\varepsilon_0} \|v_{N,j}^{(0)}\|_{H_x^s} \leq \eta_0 \right\}. \quad (6.22)$$

where $\mathcal{A}_N B$ is as in (1.11). Then, it follows from Chebyshev's inequality and (6.2) with (6.3) that there exists $N_3 = N_3(j, \eta_0, \delta) \in \mathbb{N}$ such that

$$\mathbb{P}'((\Omega'_{4,j,N}(\eta_0))^c) < \frac{\delta}{8} \quad (6.23)$$

for any $N \geq N_3$. Given small $\eta > 0$, define the $\Omega'_{5,N}(\eta) \subset \Omega'$ by setting

$$\Omega'_{5,N}(\eta) = \left\{ \omega' \in \Omega' : N^{\frac{1}{2}-\varepsilon_0} \|\vec{v}_{N,k}\|_{X_N^s(J)} \leq \eta \right\}. \quad (6.24)$$

Then, it follows from the convergence-in-probability (6.10) established in Step 1 that, given $\delta > 0$, there exists $N_4 = N_4(\eta, \delta) \in \mathbb{N}$ such that

$$\mathbb{P}((\Omega'_{5,N}(\eta))^c) < \frac{\delta}{8} \quad (6.25)$$

for any $N \geq N_4$. Given $j \in \mathbb{N}$ and $K \geq 1$, define the set $\Omega'_{6,j,N}(K) \subset \Omega'$ by setting

$$\Omega'_{6,j,N}(K) = \left\{ \omega' \in \Omega' : \|\Phi_k \Phi_j\|_{\mathcal{A}_{N,k} C_T W_x^{-\varepsilon, \infty}} \leq K \right\}. \quad (6.26)$$

Then, it follows from Lemma 2.5 (i) (for the uniform (in j, k) moment bound on the $C_T W_x^{-\varepsilon, \infty}$ -norm of $\Phi_k \Phi_j$) and Chebyshev's inequality that there exists $K = K(\delta) \gg 1$, independent of $j, N \in \mathbb{N}$, such that

$$\mathbb{P}((\Omega'_{6,j,N}(K))^c) < \frac{\delta}{8}. \quad (6.27)$$

Lastly, given small $\eta > 0$, define the set $\Omega'_{7,j,N}(\eta) \subset \Omega'$ by setting

$$\Omega'_{7,j,N}(\eta) \subset \Omega' = \left\{ \omega' \in \Omega' : N^{\frac{1}{2}-\varepsilon_0} \left\| \frac{1}{N} \sum_{k=1}^N \Phi_k^2 \Phi_j \right\|_{L_T^2 W_x^{-\varepsilon, \infty}} \leq \eta \right\}. \quad (6.28)$$

Then, it follows from Lemma 5.1 that there exists $N_5 = N_5(j, \eta, \delta) \in \mathbb{N}$ such that

$$\mathbb{P}'((\Omega'_{7,j,N}(\eta))^c) < \frac{\delta}{8} \quad (6.29)$$

for any $N \geq N_5$. In the following, we work on $\Omega'_{j,N}(K, \eta_0, \eta)$ defined by

$$\begin{aligned} \Omega'_{j,N}(K, \eta_0, \eta) &= \Omega'_N(K, \eta_0, \eta) \cap \Omega'_{4,j,N}(\eta_0) \\ &\quad \cap \Omega'_{5,N}(\eta) \cap \Omega'_{6,j,N}(K) \cap \Omega'_{7,j,N}(\eta) \end{aligned} \quad (6.30)$$

where $\Omega'_N(K, \eta_0, \eta)$ is as in (6.17). Given small $\delta > 0$, by choosing $K = K(\delta)$ sufficiently large, it follows from (6.30) with (6.18), (6.23), (6.25), (6.27), and (6.29), that

$$\mathbb{P}((\Omega'_{j,N}(K, \eta_0, \eta))^c) < \delta \quad (6.31)$$

for any $N \geq \max_{\ell=1,\dots,5} N_\ell$, where N_ℓ , $\ell = 1, \dots, 5$, are as in (6.12), (6.16), (6.23), (6.25), and (6.29), respectively.

For fixed $j \in \mathbb{N}$, we now consider (6.9) on an interval $J = [t_0, t_1] \subset [0, T]$ with $|J| \leq 1$, starting at time t_0 . By applying a slight modification of (3.6)-(3.12) (without taking the ℓ^2 -average in j ; see also the estimates in Step 2 of the proof of Theorem 1.4 (i) presented in Subsection 5.2) with (6.13), (6.24), (6.26), and (6.28), we have

$$\begin{aligned} N^{\frac{1}{2}-\varepsilon_0} \|\vec{v}_{N,j}\|_{C_T \mathcal{H}_x^s} &\leq C_1 N^{\frac{1}{2}-\varepsilon_0} \|\vec{v}_{N,j}(t_0)\|_{\mathcal{H}^s} \\ &\quad + C_2 |J| \left(\|\vec{v}_{N,k}\|_{X_N^s(J)}^2 + K^2 \right) N^{\frac{1}{2}-\varepsilon_0} \|\vec{v}_{N,j}\|_{C_T \mathcal{H}_x^s} \\ &\quad + C_3 |J|^{\frac{1}{2}} \eta (1 + K). \end{aligned} \quad (6.32)$$

Compare (6.32) with (5.87). Then, in view of (6.32) with (6.22) and (6.31), the desired convergence-in-probability (6.21) follows from a slight modification of the iterative argument in Step 1 of the proof of Theorem 1.4 (i) presented in Subsection 5.2. We omit details.

This concludes the proof of Theorem 1.6 (iii).

Acknowledgements. The authors would like to thank Leonardo Tolomeo and Jiawei Li for helpful discussions. They are also grateful to Scot Smith for telling them about the work [14]. S.L. would like to thank the School of Mathematics at the University of Edinburgh for its hospitality, where this manuscript was prepared. R.L. was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 211504053 – SFB 1060. R.L. and S.L. acknowledge support from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy – EXC-2047/1 – 390685813. R.L. and S.L. were also funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 539309657 – SFB 1720. T.O. was supported by the European Research Council (grant no. 864138 “SingStochDispDyn”) and by the EPSRC Mathematical Sciences Small Grant (grant no. EP/Y033507/1).

REFERENCES

- [1] R. Bass, *Stochastic processes*. Cambridge Series in Statistical and Probabilistic Mathematics, 33. Cambridge University Press, Cambridge, 2011. xvi+390 pp.
- [2] Á. Bényi, T. Oh, O. Pocovnicu, *Wiener randomization on unbounded domains and an application to almost sure well-posedness of NLS*, Excursions in harmonic analysis. Vol. 4, 3–25, Appl. Numer. Harmon. Anal., Birkhäuser/Springer, Cham, 2015.
- [3] Á. Bényi, T. Oh, T. Zhao, *Fractional Leibniz rule on the torus*, Proc. Amer. Math. Soc. 153 (2025), no. 1, 207–221.
- [4] J. Bourgain, *Periodic nonlinear Schrödinger equation and invariant measures*, Comm. Math. Phys. 166 (1994), no. 1, 1–26.
- [5] J. Bourgain, *Invariant measures for the 2D-defocusing nonlinear Schrödinger equation*, Comm. Math. Phys. 176 (1996), no. 2, 421–445.
- [6] B. Bringmann, *Invariant Gibbs measures for the three-dimensional wave equation with a Hartree nonlinearity II: dynamics*, J. Eur. Math. Soc. (JEMS) 26 (2024), no. 6, 1933–2089.
- [7] B. Bringmann, Y. Deng, A. Nahmod, H. Yue, *Invariant Gibbs measures for the three dimensional cubic nonlinear wave equation*, Invent. Math. 236 (2024), no. 3, 1133–1411.
- [8] D. Brydges, G. Slade, *Statistical mechanics of the 2-dimensional focusing nonlinear Schrödinger equation*, Comm. Math. Phys. 182 (1996), no. 2, 485–504.
- [9] E. Brun, G. Li, R. Liu, *Global well-posedness of the energy-critical stochastic nonlinear wave equations*, J. Differential Equations 397 (2024), 316–348.

- [10] N. Burq, N. Tzvetkov, *Probabilistic well-posedness for the cubic wave equation*, J. Eur. Math. Soc. 16 (2014), no. 1, 1–30.
- [11] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, T. Tao, *Almost conservation laws and global rough solutions to a nonlinear Schrödinger equation*, Math. Res. Lett. 9 (2002), no. 5-6, 659–682.
- [12] G. Da Prato, A. Debussche, *Strong solutions to the stochastic quantization equations*, Ann. Probab. 31 (2003), no. 4, 1900–1916.
- [13] G. Da Prato, L. Tubaro, *Wick powers in stochastic PDEs: an introduction*, Quantum and stochastic mathematical physics, 1–15. Springer Proc. Math. Stat., 377.
- [14] M. Delgadino, S.A. Smith, *An application of Talagrand’s inequality to the two dimensional linear sigma model*, in preparation.
- [15] A. Deya, *A nonlinear wave equation with fractional perturbation*, Ann. Probab. 47 (2019), no. 3, 1775–1810.
- [16] S. Fang, J. Shao, K.-T. Sturm, *Wasserstein space over the Wiener space*, Probab. Theory Related Fields 146 (2010), no. 3-4, 535–565.
- [17] D. Feyel, A.S. Üstünel, *Monge-Kantorovitch measure transportation and Monge-Ampère equation on Wiener space*, Probab. Theory Related Fields 128 (2004), no. 3, 347–385.
- [18] D.H. Fremlin, *Measure theory. Vol. 4. Topological measure spaces*. Measure theory. Vol. 4. Topological measure spaces. Part I, II. Corrected second printing of the 2003 original. Torres Fremlin, Colchester, 2006. Part I: 528 pp.; Part II: 439+19 pp. (errata).
- [19] S. Graf, R. D. Mauldin, *A classification of disintegrations of measures*, Measure and measurable dynamics (Rochester, NY, 1987), 147–158. Contemp. Math., 94 American Mathematical Society, Providence, RI, 1989.
- [20] D. Greco, T. Oh, L. Tao, L. Tolomeo, *Critical threshold for weakly interacting log-correlated focusing Gibbs measures*, Proc. Amer. Math. Soc. Ser. B 12 (2025), 150–165.
- [21] M. Gubinelli, H. Koch, T. Oh, *Renormalization of the two-dimensional stochastic nonlinear wave equations*, Trans. Amer. Math. Soc. 370 (2018), no. 10, 7335–7359.
- [22] M. Gubinelli, H. Koch, T. Oh, *Paracontrolled approach to the three-dimensional stochastic nonlinear wave equation with quadratic nonlinearity*, J. Eur. Math. Soc. (JEMS) 26 (2024), no. 3, 817–874.
- [23] M. Gubinelli, H. Koch, T. Oh, L. Tolomeo, *Global dynamics for the two-dimensional stochastic nonlinear wave equations*, Int. Math. Res. Not. IMRN. (2022), no. 21, 16954–16999.
- [24] H. Kuo, *Introduction to stochastic integration*, Universitext. Springer, New York, 2006. xiv+278 pp.
- [25] J. Lebowitz, H. Rose, E. Speer, *Statistical mechanics of the nonlinear Schrödinger equation*, J. Statist. Phys. 50 (1988), no. 3-4, 657–687.
- [26] M. Ledoux, M. Talagrand, *Probability in Banach spaces. Isoperimetry and processes*, Reprint of the 1991 edition. Classics in Mathematics. Springer-Verlag, Berlin, 2011. xii+480 pp.
- [27] G. Li, J. Li, T. Oh, N. Tzvetkov, *Probabilistic well-posedness of dispersive PDEs beyond variance blowup I: Benjamin-Bona-Mahony equation*, arXiv:2509.02344 [math.AP].
- [28] H. P. McKean, *Statistical mechanics of nonlinear wave equations. IV. Cubic Schrödinger*, Comm. Math. Phys. 168 (1995), no. 3, 479–491. Erratum: “Statistical mechanics of nonlinear wave equations. IV. Cubic Schrödinger”, Comm. Math. Phys. 173 (1995), no. 3, 675.
- [29] E. Nelson, *A quartic interaction in two dimensions*, 1966 Mathematical Theory of Elementary Particles (Proc. Conf., Dedham, Mass., 1965), pp. 69–73, M.I.T. Press, Cambridge, Mass.
- [30] D. Nualart, *The Malliavin calculus and related topics*, Second edition. Probab. Appl. (N. Y.) Springer-Verlag, Berlin, 2006. xiv+382 pp.
- [31] T. Oh, M. Okamoto, *Comparing the stochastic nonlinear wave and heat equations: a case study*, Electron. J. Probab. 26 (2021), Paper No. 9, 44 pp.
- [32] T. Oh, M. Okamoto, T. Robert, *A remark on triviality for the two-dimensional stochastic nonlinear wave equation*, Stochastic Process. Appl. 130 (2020), no. 9, 5838–5864.
- [33] T. Oh, M. Okamoto, L. Tolomeo, *Focusing Φ_3^4 -model with a Hartree-type nonlinearity*, Mem. Amer. Math. Soc. 304 (2024), no. 1529, vi+143 pp.
- [34] T. Oh, M. Okamoto, L. Tolomeo, *Stochastic quantization of the Φ_3^3 -model*, Mem. Eur. Math. Soc., 16, EMS Press, Berlin, 2025. viii+145 pp.
- [35] T. Oh, M. Okamoto, N. Tzvetkov, *Uniqueness and non-uniqueness of the Gaussian free field evolution under the two-dimensional Wick ordered cubic wave equation*, Ann. Inst. Henri Poincaré Probab. Stat. 60 (2024), no. 3, 1684–1728.
- [36] T. Oh, O. Pocovnicu, N. Tzvetkov, *Probabilistic local well-posedness of the cubic nonlinear wave equation in negative Sobolev spaces*, Ann. Inst. Fourier (Grenoble) 72 (2022) no. 2, 771–830.
- [37] T. Oh, T. Robert, N. Tzvetkov, *Stochastic nonlinear wave dynamics on compact surfaces*, Ann. H. Lebesgue 6 (2023), 161–223.

- [38] T. Oh, K. Seong, L. Tolomeo, *A remark on Gibbs measures with log-correlated Gaussian fields*, Forum Math. Sigma. 12 (2024), e50, 40 pp.
- [39] T. Oh, P. Sosoe, L. Tolomeo, *Optimal integrability threshold for Gibbs measures associated with focusing NLS on the torus*, Invent. Math. 227 (2022), no. 3, 1323–1429.
- [40] T. Oh, L. Thomann, *A pedestrian approach to the invariant Gibbs measures for the 2-d defocusing nonlinear Schrödinger equations*, Stoch. Partial Differ. Equ. Anal. Comput. 6 (2018), no. 3, 397–445.
- [41] T. Oh, L. Thomann, *Invariant Gibbs measures for the 2-d defocusing nonlinear wave equations*, Ann. Fac. Sci. Toulouse Math. 29 (2020), no. 1, 1–26.
- [42] T. Oh, L. Tolomeo, Y. Wang, G. Zheng, *Hyperbolic $P(\Phi)_2$ -model on the plane*. to appear in Comm. Math. Phys.
- [43] T. Oh, Y. Wang, Y. Zine, *Three-dimensional stochastic cubic nonlinear wave equation with almost space-time white noise*, Stoch. Partial Differ. Equ. Anal. Comput. 10 (2022), no. 3, 898–963.
- [44] F. Otto, C. Villani, *Generalization of an inequality by Talagrand and links with the logarithmic Sobolev inequality*, J. Funct. Anal. 173 (2000), no. 2, 361–400.
- [45] S. Ryang, T. Saito, K. Shigemoto, *Canonical stochastic quantization*, Progr. Theoret. Phys. 73 (1985), no. 5, 1295–1298.
- [46] H. Shen, *A stochastic PDE approach to large N problems in quantum field theory: A survey*, J. Math. Phys. 63 (2022), no. 8, 081103-1–081103-21.
- [47] H. Shen, S. A. Smith, R. Zhu, X. Zhu, *Large N limit of the $O(N)$ linear sigma model via stochastic quantization*, Ann. Probab. 50 (2022), no. 1, 131–202.
- [48] H. Shen, R. Zhu, X. Zhu, *Large N limit of the $O(N)$ linear sigma model in 3D*, Comm. Math. Phys. 394 (2022), no. 3, 953–1009.
- [49] H. Shen, R. Zhu, X. Zhu, *Large N limit and $1/N$ expansion of invariant observables in $O(N)$ linear σ -model via SPDE*, arXiv:2306.05166 [math.PR].
- [50] B. Simon, *The $P(\phi)_2$ Euclidean (quantum) field theory*, Princeton Ser. Phys. Princeton University Press, Princeton, NJ, 1974. xx+392 pp.
- [51] A.-S. Sznitman, *Topics in propagation of chaos*. École d’Été de Probabilités de Saint-Flour XIX–1989, 165–251, Lecture Notes in Math., 1464, Springer, Berlin, 1991.
- [52] S.M. Srivastava, *A course on Borel sets*. Graduate Texts in Mathematics, 180. Springer-Verlag, New York, 1998. xvi+261 pp.
- [53] C. Sun, N. Tzvetkov, W. Xu, *Weak universality results for a class of nonlinear wave equations*, to appear in Ann. Inst. Fourier. (Grenoble).
- [54] M. Talagrand, *Transportation cost for Gaussian and other product measures*, Geom. Funct. Anal. 6 (1996), no. 3, 587–600.
- [55] L. Thomann, N. Tzvetkov, *Gibbs measure for the periodic derivative nonlinear Schrödinger equation*, Nonlinearity 23 (2010), no. 11, 2771–2791.
- [56] L. Tolomeo, *Global well posedness of the two-dimensional stochastic nonlinear wave equation on an unbounded domain*, Ann. Probab. 49 (2021), no. 3, 1402–1426.
- [57] L. Tolomeo, *Ergodicity for the hyperbolic $P(\Phi)_2$ -model*, arXiv:2310.02190 [math.PR].
- [58] N. Tzvetkov, *Construction of a Gibbs measure associated to the periodic Benjamin-Ono equation*, Probab. Theory Related Fields 146 (2010), no. 3-4, 481–514.
- [59] C. Villani, *Topics in optimal transportation*. Grad. Stud. Math., 58 American Mathematical Society, Providence, RI, 2003. xvi+370 pp.
- [60] K. Wilson, *Quantum field-theory models in less than 4 dimensions*, Phys. Rev. D (3) 7 (1973), 2911–2926.
- [61] Y. Zine, *Smoluchowski-Kramers approximation for singular stochastic wave equations in two dimensions*, Electron. J. Probab. 30 (2025), Paper No. 88, 49 pp.

RUOYUAN LIU, MATHEMATICAL INSTITUTE, UNIVERSITY OF BONN, ENDENICHER ALLEE 60, 53115, BONN, GERMANY

Email address: ruoyuanl@math.uni-bonn.de

SHAO LIU, MATHEMATICAL INSTITUTE, UNIVERSITY OF BONN, ENDENICHER ALLEE 60, 53115, BONN, GERMANY

Email address: shaoliu@math.uni-bonn.de

TADAHIRO OH, SCHOOL OF MATHEMATICS, THE UNIVERSITY OF EDINBURGH, AND THE MAXWELL INSTITUTE FOR THE MATHEMATICAL SCIENCES, JAMES CLERK MAXWELL BUILDING, THE KING'S BUILDINGS, PETER GUTHRIE TAIT ROAD, EDINBURGH, EH9 3FD, UNITED KINGDOM, AND SCHOOL OF MATHEMATICS AND STATISTICS, BEIJING INSTITUTE OF TECHNOLOGY, BEIJING 100081, CHINA

Email address: `hiro.oh@ed.ac.uk`