

Forecasting Primordial Non-Gaussianity from UNIONS Lyman-Break Galaxies and *Planck* CMB lensing

Constantin Payerne¹, William d’Assignies², Christophe Yèche¹,
Hendrik Hildebrandt³, Dustin Lang^{4,5}, Thomas de Boer⁶

¹ Université Paris-Saclay, CEA, IRFU, 91191 Gif-sur-Yvette, France

² IFAE, The Barcelona Institute of Science and Technology, Campus UAB, 08193 Bellaterra Barcelona, Spain

³ Ruhr University Bochum, Faculty of Physics and Astronomy, Astronomical Institute, GCCL, 44780 Bochum, Germany

⁴ Perimeter Institute for Theoretical Physics, 31 Caroline St. North, Waterloo, ON N2L 2Y5, Canada

⁵ Department of Physics and Astronomy, University of Waterloo, 200 University Ave W, Waterloo, ON N2L 3G1, Canada

⁶ Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu HI 96822

Received –; accepted –

ABSTRACT

Context. Local primordial non-Gaussianities, characterized by the parameter $f_{\text{NL}}^{\text{loc}}$, provide a powerful window into the physics of inflation. Cross-correlating high-redshift tracer samples with the CMB lensing potential offers a particularly robust probe of $f_{\text{NL}}^{\text{loc}}$, mitigating imaging systematics that typically affect large-scale measurements from tracer auto-spectra. In this context, the Ultraviolet Near Infrared Optical Northern Survey (UNIONS) enables the selection of u -dropout high-redshift Lyman-Break Galaxies (LBGs).

Aims. We aim to forecast the expected precision on $f_{\text{NL}}^{\text{loc}}$ achievable from analyzing the cross-correlation power spectrum between the distribution of UNIONS-selected LBGs and the CMB lensing potential measured by the *Planck* satellite.

Methods. We perform a Markov Chain Monte Carlo forecast to estimate the uncertainties on $f_{\text{NL}}^{\text{loc}}$ and on a galaxy bias parameter b_0 , which captures our uncertainty in the tracer bias.

Results. We forecast $\sigma(f_{\text{NL}}^{\text{loc}}) = 34$ for an idealized photometric sample of $r < 24.3$ LBGs selected with a Random Forest classification algorithm from UNIONS-like *ugriz* imaging, with a resulting surface density of $1,100 \text{ deg}^{-2}$. This precision can be improved to $\sigma(f_{\text{NL}}^{\text{loc}}) = 20$ after spectroscopic follow-up with the Dark Energy Spectroscopic Instrument (DESI), during its next phase starting in 2029, DESI-II. We also test a more realistic selection using early UNIONS data, based on a u -dropout color–color cut over the *ugr* imaging, which yields a denser sample of $r < 24.2$ objects at $1,400 \text{ deg}^{-2}$. From this sample—covering a larger footprint and expected to have a higher large-scale galaxy bias—we forecast an improved constraint of $\sigma(f_{\text{NL}}^{\text{loc}}) = 20$, with similar precision achievable after DESI-II follow-up. In addition, we perform preliminary validation of the redshift distribution using the clustering-redshift method with DESI DR1 data, confirming the calibration from deep, small-area photometric fields. However, accounting for uncertainties in the clustering-redshift distribution significantly degrades the $f_{\text{NL}}^{\text{loc}}$ constraining power.

Key words. Galaxies: high-redshift – Cosmology: large-scale structure of Universe – Methods: statistical

1. Introduction

Lyman Break Galaxies (LBGs; Steidel et al. 1996) are young, actively star-forming galaxies at $z > 1.5$. Their rest-frame spectra exhibit a sharp flux decrement blueward of the Lyman- α transition at 1216 \AA , extending down to the Lyman limit at 912 \AA , due to absorption by neutral hydrogen in both the intergalactic medium (IGM) and within the galaxies themselves. These spectral features enable the identification of LBGs at $2.5 < z < 3.5$ through the u -dropout technique, which selects objects with a pronounced flux deficit in the u band ($3300\text{--}4000 \text{ \AA}$) relative to their g - or r -band flux (Ruhlmann-Kleider et al. 2024; Payerne et al. 2025). At higher redshifts, analogous dropout techniques (e.g., g - or r -dropouts; Malkan et al. 2017; Ono et al. 2018; Harikane et al. 2022; Ruhlmann-Kleider et al. 2024) exploit the flux decrement at wavelengths beyond 4000 \AA . Thanks to these well-defined spectral breaks, LBGs are powerful tracers for probing the early Universe using broadband photometry.

LBGs have long been central to studies of galaxy formation and evolution at high redshift (Steidel et al. 1996, 1999; Giavalisco et al. 2004; Reddy et al. 2008; Hildebrandt et al.

2009; Finkelstein et al. 2022; Harikane et al. 2023). More recently, dropout-selected galaxies have also emerged as powerful cosmological probes (see review by Wilson & White 2019). They serve as (i) highly biased tracers of large-scale structure in the high-redshift, matter-dominated Universe (Foucaud et al. 2003; Ye et al. 2025; Ruhlmann-Kleider et al. 2024), and (ii) distant background light sources for probing the intergalactic medium through Lyman- α forest absorption in their spectra (Herrera-Alcantar et al. 2025). Clustering measurements of LBGs allow constraints on the growth of structure and the evolution of dark energy within $2 < z < 6$, for example, through cross-correlation with CMB lensing (Wilson & White 2019; Miyatake et al. 2022). Large, high-redshift LBG samples spanning wide cosmic volumes further enable tests of primordial non-Gaussianity via scale-dependent bias (Schmittfull & Seljak 2018; Chaussidon et al. 2025; Payerne et al. 2025), as well as investigations of the sum of neutrino masses (Yu et al. 2018). At cluster redshifts $z > 1$, where weak-lensing shape measurements of background galaxies are challenging, the magnification of LBG samples can be exploited to estimate cluster masses (Tudorica et al. 2017) and to probe the low-redshift matter density

field through inverse galaxy–galaxy lensing (Cross & Sánchez 2024), a method largely independent of galaxy bias. In this context, dense samples of LBGs at $z > 2.5$ are particularly valuable for cosmology. Such samples are expected to be provided by current and forthcoming wide-field, multi-band imaging surveys with U -band coverage (Payerne et al. 2025; Crenshaw et al. 2025), including the ongoing Ultraviolet Near Infrared Optical Northern Survey¹ (UNIONS; Ibata et al. 2017; Chambers et al. 2016; Miyazaki et al. 2018; Gwyn et al. 2025), the Vera C. Rubin Observatory’s Legacy Survey of Space and Time² (LSST; LSST Science Collaboration et al. 2009), and the Chinese Space Station Telescope (CSST). The high number densities of LBGs expected from these surveys will enable precise cosmological studies of the high-redshift Universe, primarily through measurements of projected clustering (with LSST, *Euclid*, and CSST) and of the three-dimensional power spectrum of large-scale structure in next-generation spectroscopic programs (e.g., DESI-II, WST, Spec-S5, MUST). In this work, we focus on the constraining power of high-density LBG samples, selected from wide-area broadband imaging surveys such as UNIONS, on local-type primordial non-Gaussianity (PNG). Inflation remains the leading paradigm for the early Universe, and local PNG—quantified by the parameter $f_{\text{NL}}^{\text{loc}}$ —offers a key test of inflationary models. Multi-field inflation scenarios, in particular, predict a potentially observable level of PNG, typically $f_{\text{NL}}^{\text{loc}} \sim \mathcal{O}(1)$. While the cosmic microwave background (CMB) has already provided strong constraints, with $f_{\text{NL}}^{\text{loc}} = -0.9 \pm 5$ (Planck Collaboration et al. 2020), further improvements from CMB observations are fundamentally limited by cosmic variance. This makes large-scale structure surveys the most promising avenue for future progress.

Primordial non-Gaussianities imprint a distinctive scale dependence on the large-scale linear bias of cosmological tracers (Dalal et al. 2008; Slosar et al. 2008), such as galaxies and quasars. This feature has been widely used to constrain $f_{\text{NL}}^{\text{loc}}$, either through the three-dimensional power spectrum of tracers (Castorina et al. 2019; Rezaie et al. 2021; Mueller et al. 2022; Cagliari et al. 2024; Chaussidon et al. 2025) or via angular power spectra and cross-correlations with CMB lensing (Krolewski et al. 2024; Fabbian et al. 2025). The tightest current limit, $f_{\text{NL}}^{\text{loc}} = -3.6^{+9.0}_{-9.1}$, was obtained by Chaussidon et al. (2025) from DESI Quasar ($0.8 < z < 3.1$) and Luminous Red Galaxy ($0.6 < z < 1.1$) large-scale power spectrum measurements. In this context, high-redshift LBGs are expected to deliver independent and competitive constraints on $f_{\text{NL}}^{\text{loc}}$ thanks to their higher number densities compared to DESI quasars (Payerne et al. 2025; Crenshaw et al. 2025) and their redshift distribution spanning $z = 2.5$ – 3.5 . Probing large-scale clustering at $z > 2.5$ is particularly advantageous: LBGs sample vast cosmic volumes, are strongly biased tracers, and lie in an epoch less affected by nonlinear structure growth. Consequently, the primordial signal is less contaminated by late-time astrophysical effects, providing a cleaner window on the Universe’s initial conditions.

The paper is organized as follows: In Section 2, we introduce the formalism for the LBG angular power spectrum, the cross-angular power spectrum between the LBG population and the CMB lensing potential, and their link to local PNGs. In Section 3, we present the different datasets that we use throughout this paper to conduct PNG forecasts. In Section 4, we describe the forecasting methodology employed throughout this work, based on posterior estimation via MCMC using fiducial

data vectors. In Section 5, we present the different forecasts on the parameter $f_{\text{NL}}^{\text{loc}}$, exploring modeling choices and propagation of photometric redshift distribution uncertainties. We conclude in Section 6.

2. Formalism for the Angular Power spectrum

Under the Limber approximation (Limber 1953), the correlation function of two fields X , Y is dominated by small angular scales only (i.e., high multipoles) and the kernel varies slowly along the line of sight. The angular power spectrum simplifies to

$$C_{\ell}^{XY} \approx \int_0^{\chi_{\text{H}}} \frac{d\chi}{\chi^2} q_X(\chi) q_Y(\chi) P\left(k = \frac{\ell + 1/2}{\chi}, z(\chi)\right), \quad (1)$$

where q_x are the kernels associated to X and Y , χ is the comoving distance, and P is the matter power spectrum. From this, we discuss the auto-correlation angular power spectrum of the LBG population, and its cross-correlation with CMB lensing maps.

2.1. Galaxy density field and clustering

The observed galaxy overdensity can be written as

$$\widehat{\Delta}_{\text{g}}(\theta) = \widehat{\Delta}_{\text{g, int}}(\theta) + \widehat{\Delta}_{\text{g, RSD}}(\theta) + \widehat{\Delta}_{\text{g, mag}}(\theta), \quad (2)$$

where $\widehat{\Delta}_{\text{g, int}}(\theta)$ denotes the intrinsic clustering in the absence of primordial non-Gaussianities, $\widehat{\Delta}_{\text{g, RSD}}(\theta)$ describes the contribution from PNGs, $\widehat{\Delta}_{\text{g, mag}}(\theta)$ encodes the effect of redshift-space distortions, and $\widehat{\Delta}_{\text{g, mag}}(\theta)$ captures the contribution from lensing magnification. The kernel corresponding to the intrinsic contribution is

$$q_{\text{g}}^{\text{int}}(\chi) = b(z) n(z) \frac{dz}{d\chi}, \quad (3)$$

where $b(z)$ denotes the large-scale linear galaxy bias, and $n(z)$ the (normalized) galaxy redshift distribution. In our analysis we restrict to multipoles $\ell < 300$, which for sources at $z \simeq 2.5$ – 3 correspond to comoving wavenumbers $k \sim \ell/\chi(z) \lesssim 0.05 \, h \, \text{Mpc}^{-1}$. At these scales, the matter density field lies well within the linear regime: nonlinear clustering only becomes relevant at $k \gtrsim 0.7$ – $1.0 \, h \, \text{Mpc}^{-1}$ at these redshifts, as shown by simulations (Takahashi et al. 2012). Furthermore, galaxy bias is expected to remain scale-independent down to $k \sim 0.1$ – $0.2 \, h \, \text{Mpc}^{-1}$ (Desjacques et al. 2018). Our choice of angular scales $\ell < 300$, therefore, ensures that the clustering signal can be modeled using linear perturbation theory (e.g., the linear matter power spectrum $P(k, z)$ with a redshift- and scale-independent bias $b(z)$). Moreover, lensing magnification alters the observed galaxy number density by deflecting light from intervening structures. Depending on the survey flux limit and the slope of the luminosity function, this effect can lead to either an apparent enhancement or suppression of number counts. Redshift-space distortions arise from the peculiar velocities of galaxies along the line of sight, which modify their observed redshifts and thus their inferred distances. This induces anisotropies in the observed clustering. The corresponding kernels for these contributions are

$$q_{\text{g}}^{\text{RSD}}(\chi) = -H(z) n(z) \frac{d \ln D}{d \ln a} j''(k\chi), \quad (4)$$

$$q_{\text{g}}^{\text{mag}}(\chi) = (5s - 2) \times \frac{3 \Omega_{\text{m}} H_0^2}{2} \frac{\chi}{a(\chi)} \int_{\chi}^{\chi_{\text{H}}} d\chi' n(z(\chi')) \frac{\chi' - \chi}{\chi'}. \quad (5)$$

¹ <https://www.skysurvey.cc/>

² <https://rubinobservatory.org/about>

For the magnification term, s denotes the magnification bias, i.e., the logarithmic slope of the cumulative number counts at the survey detection limit, defined as $s(m_{\text{lim}}, z) = d \log_{10} N(< m_{\text{lim}}, z) / dm$ where $N(< m)$ is the number of objects brighter than magnitude m , and m_{lim} is the survey flux limit (Elvin-Poole et al. 2023; Euclid Collaboration: Lepori et al. 2022). We emphasize that it is a zeroth-order approximation of the magnification bias since color cuts may distort the effective selection at the limit magnitude m_{lim} . For the latter, we will consider the r band to compute the limiting magnitude of the sample. From this, the observed angular power spectrum of the galaxy density field is given by

$$C_{\ell}^{\text{gg,obs}} = \sum_{ij} C_{\ell}^{q_i q_j} + \frac{1}{\bar{n}_{\text{gal}}} \quad (6)$$

where $q_i \in \{q_{\text{g}}^{\text{int}}, q_{\text{g}}^{\text{RSD}}, q_{\text{g}}^{\text{mag}}\}$, and \bar{n}_{gal} is the surface density of the galaxy sample in steradians, denoting for the white noise contribution from the tracer shot noise.

2.2. CMB Lensing and Cross-Correlation with LSS

The temperature anisotropies and polarization patterns of the CMB are gravitationally lensed by the intervening large-scale structure between the surface of last scattering and the observer. This lensing remaps the primary CMB anisotropies and induces B-mode polarization, as well as characteristic non-Gaussian features in the observed CMB maps (Lewis & Challinor 2006). The deflections induced by lensing can be described by the lensing potential $\phi(\hat{\mathbf{n}})$, which is related to the line-of-sight integral of the gravitational potential. The associated convergence field $\kappa(\hat{\mathbf{n}}) = -\frac{1}{2} \nabla^2 \phi(\hat{\mathbf{n}})$ can then be reconstructed from high-resolution CMB temperature and polarization maps using quadratic estimators (Okamoto & Hu 2003). The lensing convergence has a projection kernel given by

$$q_{\kappa}(\chi) = \frac{3H_0^2 \Omega_{\text{m}}}{2c^2} \frac{\chi}{a(\chi)} \left(\frac{\chi_* - \chi}{\chi_*} \right), \quad (7)$$

χ_* is the comoving distance to the surface of last scattering, $a(\chi)$ is the scale factor, H_0 is the Hubble constant, and Ω_{m} is the matter density parameter today. This kernel peaks at redshift $z \sim 2$, which makes CMB lensing especially sensitive to the high-redshift Universe.

The reconstructed CMB lensing convergence power spectrum includes both signal and noise contributions, and is given by $C_{\ell}^{\kappa\kappa, \text{obs}} = C_{\ell}^{\kappa\kappa} + N_{\ell}^{\kappa}$ where $C_{\ell}^{\kappa\kappa}$ is the theoretical lensing auto-spectrum, and N_{ℓ}^{κ} is the lensing reconstruction noise, which depends on the specifications of the CMB experiment (e.g., angular resolution, instrumental noise, etc.). In this work, we adopt the noise spectrum of the CMB lensing maps derived from the Planck PR4 CMB data (Carron et al. 2022).

The CMB lensing signal can also be cross-correlated with the distribution of galaxies or other tracers of large-scale structure. Such cross-correlations provide a powerful probe of the large-scale matter distribution and the galaxy bias, and are sensitive to primordial non-Gaussianity, $f_{\text{NL}}^{\text{loc}}$, via scale-dependent effects. The observed cross-angular power spectrum between the CMB lensing field and a galaxy field is given by the sum of $C_{\ell}^{\kappa q_i}$ where the sum runs over the relevant galaxy kernel contributions q_i (i.e. intrinsic clustering, magnification, and redshift-space distortions).

2.3. PNG: impact of $f_{\text{NL}}^{\text{loc}}$ on galaxy field

Multifield inflation models can generate a small level of local PNG, parametrized by $f_{\text{NL}}^{\text{loc}}$, which modifies the statistics of the initial gravitational potential. By modifying the height of rare density peaks, the parameter $f_{\text{NL}}^{\text{loc}}$ alters the response of halo abundance to long-wavelength background modes, thereby affecting the large-scale halo bias (Dalal et al. 2008; Slosar et al. 2008; Desjacques & Seljak 2010). The tracer bias acquires an additional scale-dependent correction $b \rightarrow b + \Delta b$ in Eq. (3), with

$$\Delta b(z, k) = b_{\Phi} f_{\text{NL}}^{\text{loc}} \frac{3\Omega_{\text{m}} H_0^2}{2k^2 T(k) D(z)}. \quad (8)$$

where $T(k)$ is the matter power spectrum transfer function and $D(z)$ is the linear growth factor. Assuming a universal mass function, the bias b_{Φ} can be related to the linear tracer bias through the relation

$$b_{\Phi} = 2(b(z) - 1) \delta_c \quad (9)$$

While this relation is by far the most widely adopted in the literature, there is no compelling reason to expect it to hold for realistic tracers of large-scale structure. It is common to generalize the expression by introducing a parameter p_{Φ} , so that $b_{\Phi} = 2(b(z) - p_{\Phi}) \delta_c$ where p_{Φ} depends on the merger history of the tracer (Barreira 2020). Simulations of dark matter halos and mass-selected galaxies in IllustrisTNG (Barreira et al. 2020), as well as studies of recent mergers (Slosar et al. 2008), have shown that p_{Φ} can significantly deviate from unity. It is therefore generally accepted that $p_{\Phi} \neq 1$ must be taken into account depending on the tracer under consideration. Even if the lensing-related quantities are in principle subject to corrections from primordial non-Gaussianity (Jeong et al. 2011; Anbajagane et al. 2024), we neglect the impact of $f_{\text{NL}}^{\text{loc}}$ on the CMB lensing potential since the effects of $f_{\text{NL}}^{\text{loc}}$ are strongly suppressed by projection effects. Thus, in this work, we rely on the large-scale dependent bias effect described in Eq. (8) to constrain $f_{\text{NL}}^{\text{loc}}$.

3. Datasets

In this work, we perform a forecast study for an LBG sample that can be generated from the UNIONS data. The UNIONS is build upon a collaboration between the Hawaiian observatories: the Canada-France-Hawaii Telescope (CFHT, Mauna Kea), the Panoramic Survey Telescope and Rapid Response System (Pan-STARRS, Maui), and the Subaru Telescope (Mauna Kea). It is currently providing *ugriz* imaging over 5,000 deg² of the northern sky. The CFHT Canada-France Imaging Survey (CFHT/CFIS) targets the *u*- and *r*-bands with the Megacam imager, delivering image quality competitive with all other current large ground-based facilities. CFHT/CFIS will reach a depth of $r \simeq 25$ over 5,000 deg² and $u \simeq 24.6$ over 9,000 deg². Meanwhile, Pan-STARRS provides the *i*-band, and the Wide Imaging with Subaru Hyper Suprime-Cam of the Euclid Sky (WISHES) supplies the *z*-band. The 5σ depth of the data, measured within a 2-arcsecond diameter aperture (Gwyn et al. 2025), is $[u, g, r, i, z] = [24.45, 25.25, 24.95, 24.55, 24.05]$. Multi-band catalogs over the UNIONS footprint have been obtained using the GAAP (Gaussian Aperture and PSF; Kuijken 2008; Kuijken et al. 2015, 2019) method, which will be made publicly available upon release of the UNIONS data set. The covered surface areas are, for $\{ugr, ugri, ugriz\} = \{4760, 4630, 3730\}$ deg² (Gwyn et al. 2025). Besides the main UNIONS footprint, multi-band *ugr* catalogs are

available on the two deep fields XMM-LSS and COSMOS. We will consider using these deep fields in the next section.

In the following sections, we present the various LBG selections applied to UNIONS-like and UNIONS imaging. For each case, we derive the corresponding sample characteristics, namely the redshift distribution and the lensing magnification bias.

3.1. UNIONS-like LBG sample from a Random Forest approach by degrading deep catalogs

In this section, we present the selection of high-redshift LBGs on UNIONS-like imaging, obtained on the COSMOS field.

Photometric LBG sample: We first introduce the LBG sample that was derived in Payerne et al. (2025), obtained by applying a Random Forest selection to UNIONS-like photometry, i.e. by degrading CLAUDS³ u - and HSC⁴-PDR3 $griz$ broadband imaging on the COSMOS field. For this dataset, we get deep 5σ point source depth $[u, g, r, i, z] = [27.7, 27.4, 27.1, 26.9, 26.3]$. The combination of CLAUDS and HSC-SSP data over COSMOS (and other small deep fields such as XMM-LSS) is detailed in Desprez et al. (2023). In Payerne et al. (2025), the COSMOS CLAUDS+HSC catalog has been degraded to UNIONS depth (called UNIONS-like, with respective depth, at the time of this work $ugriz = [24.6, 25.5, 25.5, 24.2, 24.4]$) to train a Random Forest for selecting $r < 24.3$ UNIONS-like LBGs in the redshift range $z \in [2.5, 3.5]$. The LePhare (6 bands) photometric redshift distribution of this sample is shown in red full lines in the left panel of Figure 1, corresponding to a surface density of ~ 1100 photometric LBGs per deg^2 . The magnification bias factor for this sample is found to be $s(r_{\text{lim}} = 24.3) = 0.43$.

DESI-II LBG sample: This work is also the occasion to explore DESI-II scenarios, whose high-redshift target selection strategy is still under discussion. Spectroscopic redshift follow-ups will allow us to (1) remove low redshift contaminants and (2) reduce the uncertainty on the sample redshift distribution. Considering all $z > 2$ objects in the distribution in Figure 1 (left panel), the number density goes from 1100 to 880 LBGs per square degree. Moreover, the current DESI spectroscopic redshift reconstruction method⁵ has an internal efficiency, which degrades the efficiency of the recovered LBG spectroscopic sample at $z \sim 2.2$. Considering the redshift efficiency in Figure 11 of Payerne et al. (2025) or in Fig. 18 of Ruhlmann-Kleider et al. (2024) (resp. for 2-hour and 4-hour exposure) multiplying the corresponding redshift distribution, we get the corrected number densities 500 and 590 LBGs per square degrees. The corresponding 2-hour and 4-hour exposure high-redshift distributions are shown in the left panel of Figure 1.

Alternative LBG selection: We explore two alternative UNIONS-like LBG selections on the COSMOS field; Individual CLAUDS U -band images of 180-second exposures on COS-

MOS have been co-added and have undergone forced photometry using HSC g -band. Two U -band catalogs were obtained, corresponding to 180-second and 2×180 -second exposures in the u -band. For reference, the default deep CLAUDS U -band catalog uses 600-second exposures. Two other selections, namely UNIONSlike_RF_u180s and UNIONSlike_RF_u2x180s where obtained by training a RF algorithm to select a sample of 1100 deg^{-2} galaxies with $r < 24.5$ in the range $z \in [2.5, 3.5]$, where, for each galaxy sample, $s(r_{\text{lim}} = 24.5) = 0.21$ and 0.46 , respectively. The corresponding photometric redshift distributions are represented in the left panel of 1. These selections, as for the one in Payerne et al. (2025), were tested on the COSMOS field and show different selection efficiencies.

3.2. u -dropout LBG sample using UNIONS GAAP catalogs

In this section, we make use of the early GAAP UNIONS data to test different high-redshift LBG selections.

LBG color-color selection: We use the early UNIONS GAAP photometric data available in the UNIONS collaboration, to test a color-color box selection of high-redshift $r < 24.2$ LBGs referred to as [COSMOS: TMG u -dropout] in Ruhlmann-Kleider et al. (2024) (see their Table 1). For $22 < r < 24.2$, the u -dropout color-box selection is defined as

$$(i) \ u - g > 0.3, \quad (10)$$

$$(ii) \ -0.5 < g - r < 1, \quad (11)$$

$$(iii) \ [u - g > 2.2 \times (g - r) + 0.32] \quad (12)$$

$$\cup [u - g > 0.9 \cap u - g > 1.6 \times (g - r) + 0.75] \quad (13)$$

This sample is referred to as UNIONS_colorcut and yields a photometric LBG angular density of 1400 deg^{-2} and a magnification bias $s(r_{\text{lim}} = 24.2) = 0.25$. To assess the redshift distribution of these photometrically selected u -dropout LBGs, we match geometrically the selected UNIONS-LBGs to deep photometric redshift catalogs provided by CLAUDS+HSC imaging on XMM (XMM benefits from UNIONS multi-band imaging). The corresponding photometric redshift distribution is displayed in the right panel of Figure 1. In our previous Payerne et al. (2025) idealistic dataset, the resulting redshift distribution $n(z)$ is relatively narrow, peaking sharply at $z \sim 3$. In comparison to the color-box selection, which peaks at $z \sim 1.5$ with a broader shape, the difference reflects (i) the slight optimistic $ugriz$ magnitude depth used in Payerne et al. (2025) for the degradation procedure (ii) the use of a Random Forest approach (iii) the simplified and noise-free nature of the simulated conditions in Payerne et al. (2025), where measurement uncertainties, selection effects, and intrinsic galaxy diversity are minimized. The broadening of the UNIONS distribution can be attributed to observational noise in UNIONS data, all of which introduces scatter and extends the distribution away from the main peak. As a result, the realistic dataset provides a more accurate representation of the complexity encountered in real observations.

LBG random forest selection: As for the idealistic Payerne et al. (2025), an alternative is to use a Random Forest classifier to select LBGs within a specific redshift range using UNIONS imaging. However, the limited overlap between UNIONS and CLAUDS+HSC on XMM restricts the available training data (the overlap is about 2 deg^2). Instead, we construct a hybrid selection by splitting the matches into two separate samples. For the first

³ Canada-France-Hawaii Telescope Large Area U-band Deep Survey, Sawicki et al. (2019)

⁴ Hyper Suprime-Camera, Aihara et al. (2022)

⁵ First, a convolutional neural network (CNN) derived from Quasar-NET (Busca & Balland 2018) is applied to each DESI spectrum. A more precise LBG redshift is then obtained using the RedRock (RR) software (Guy et al. 2023) that uses the CNN output redshift as a prior and refines its measurement.




Fig. 1. Left: Photometric redshift distributions of the photometric LBG samples UNIONSlike_RF in red (the convolution of the UNIONSlike_RF distribution with the DESI spectroscopic efficiency is shown in dashed and dotted lines, they are not normalized to show the impact at $z \sim 2.5$). The distribution of the samples UNIONSlike_RF_u180s, and LSSTY4like_RF_u2x180s are shown in blue and dashed cyan lines. **Right:** Photometric redshift distributions of the UNIONS LBG samples UNIONS_colorcut and UNIONS_PZcut in the XMM field. The sample selected with the hybrid method à-la SOM is shown for illustration. We also show the quantity nb (the product between the large-scale bias and the normalized redshift distribution) resulting from our clustering-redshift calibration methods.

subsample, we pre-compute the mean LePhare photometric redshift in a multidimensional grid of UNIONS colors ($u - g$, $g - r$, $r - i$) and r magnitude. A UNIONS galaxy in the second subsample is considered as part of the LBG sample if the mean redshift in its associated grid cell satisfies $\langle z \rangle > 1.7$. This grid-based approach is conceptually similar to Self-Organizing Maps (SOMs, Zhang et al. 2025; Roster et al. 2025), and poorly mimics the cut-free approach of Random Forests. From this, we get a representative redshift distribution derived from a reference sample. Our method uses a fixed, axis-aligned grid, while SOMs construct an adaptive, data-driven grid that captures complex structures in color space. This hybrid SOM-like selection tests the relevance of the color-box cuts compared to a data-driven cut-free method is represented in green in the right panel of Figure 1. We observe that the distribution fairly matches the color-color box selection.

LBG photometric selection: To check the consistency of the u -dropout selection described above (color-cut), we apply an alternative selection based on the BPZ photometric redshifts (Benítez 2011) of UNIONS detections, requiring $Z_{ML} > 1.7$ in the catalog. This selection is referred to as UNIONS_PZcut, and represented in purple in the right panel of Figure 1. This validation on XMM of the color-box selection enables us to apply this selection to a wider UNIONS portion of the sky, and assess the corresponding redshift distribution through clustering-redshift methods.

Matching LBG targets with deep CLAUDS+HSC photometric catalogs is useful to assess the underlying photometric redshift distribution; however, this approach remains limited by the accuracy of the deep photometric catalogs and by the relatively low statistical significance of the inferred distribution (usually done on fields of a few square degrees). A second method relies on spectroscopic data, as high-precision spectroscopic measurements can help calibrate the less accurate photometric redshifts. For available spectroscopic subsamples of photometric datasets, direct calibration is possible (see, e.g., Lima et al. (2008); Hildebrandt et al. (2021)). Such direct calibration is challenging for LBGs, however, since obtaining a representative spectroscopic

sample for any specific photometric selection is difficult in practice (Ruhmann-Kleider et al. 2024; Payerne et al. 2025).

Measuring the redshift distribution with clustering-redshifts: In this work, we use the clustering-redshifts method (Ménard et al. 2013; Schmidt et al. 2013; d’Assignies et al. 2025b) to evaluate the redshift distribution of an arbitrary photometrically-selected dataset based on the spatial cross-correlation with a reference population, the latter with spectroscopic redshift available.

The overlap of UNIONS data we use in this work with DESI spectroscopic observations publicly available (DR1, Ross et al. 2025) is approximately $1,300 \text{ deg}^2$. The region of overlap suffers from a low completeness of DESI data ($\text{Dec} > 30 \text{ deg}$), which limits the method’s constraining power. DESI data also includes randoms. We evaluate the measurements for the different DESI tracers separately (BGs, LRG, ELG, and QSO), and then recombine them, as the tracers’ masks differ from one to another. The DESI data are then used to measure the product of the redshift distribution with galaxy bias $b_{\text{LBG}}(z) n_{\text{LBG}}(z)$ with the so-called clustering-redshifts methods (using the pipeline developed in d’Assignies et al. 2025b), over the redshift range $0 < z < 3.4$. Contrary to the majority of the previous clustering-redshift calibration methods, we do not aim to break the degeneracy between bias and redshift distribution, as the observables we consider directly depend on their product. The uncertainty $\sigma(b_{\text{LBG}}(z) n_{\text{LBG}}(z))$ can then be used directly for the Bayesian inference, and we marginalize over both bias and distribution during the same step.

For clustering-redshifts calibration, we binned the spectroscopic data in redshift bins $z_j \pm \Delta z/2$, and measure the cross correlations between each spectroscopic bin s_j and the UNIONS LBG bin, as functions of perpendicular separations r_p : $w_{s_j \text{ LBG}}(r_p)$. To maximize the signal-to-noise ratio and simplify covariance estimates, we usually reduce the data vectors to scalars given by

$$\bar{w}_{s_j \text{ LBG}} = \int_{r_{p,\min}}^{r_{p,\max}} W(r_p) w_{s_j \text{ LBG}}(r_p) dr_p, \quad (14)$$

with $W(r_p) \propto r_p^\gamma$ a normalized weighting functions. We use the scale range 1.5 – 5.5 Mpc and the weighting $\gamma = -1$, which is an excellent tradeoff between boosting the signal-to-noise ratio coming from small scales, limiting biasing due to non-linearities (d’Assignies et al. 2025b), and fiber collisions (see Choppin de Janvry et al. 2025; d’Assignies et al. 2025a). These scalars can be used to constrain $b_{\text{LBG}}(z) n_{\text{LBG}}(z)$ as

$$\bar{w}_{s_j \text{ LBG}} = b_s(z_j) b_{\text{LBG}}(z_j) n_{\text{LBG}}(z_j) \bar{w}_m(z_j), \quad (15)$$

where \bar{w}_m is a theoretical function estimated with Halofit (Takahashi et al. 2012). The spectroscopic galaxy biases $b_s(z_j)$ are measured from the auto-correlations of the spectroscopic bins, for larger scales (to limit the effect of fiber collision). The data vectors are measured with Treecorr (Jarvis 2015) and LS estimators (Landy & Szalay 1993). Moreover, the correlations of \bar{w} between different redshifts z_j can be neglected (d’Assignies et al. 2025a). Thus, we estimate the covariance matrix using a standard Jackknife estimate, and put all the off-diagonal coefficients to 0. We combined the measurements from different tracers localized at the same redshift, using an inverse weighting scheme, neglecting cross-correlations (Choppin de Janvry et al. 2025). The joint product of the LBG bias and the LBG redshift distribution, $b_{\text{LBG}}(z_j) n_{\text{LBG}}(z_j) = \bar{w}_{s_j \text{ LBG}} / b_r(z_j) \bar{w}_m(z_j)$ is shown in Figure 1 (right panel) with the corresponding error bars. For the latter, we work in a minimal scenario, where the bias and the redshift distribution cannot be disentangled. This is possible by dealing with accurate photometric redshifts. We can compute an *effective* bias of the sample by integrating $b_{\text{LBG}} n_{\text{LBG}}$, we get $b_{\text{eff}} = 1.5 \pm 0.2$, where the error is estimated from random samples of $b_{\text{LBG}} n_{\text{LBG}}$.

4. Forecasting methodology

We first detail how to parametrize uncertainties associated with our LBG sample, i.e., the large-scale bias, the outlier fraction, and the redshift distribution uncertainty, as well as accounting for clustering-redshift-derived distribution in the $f_{\text{NL}}^{\text{loc}}$ prediction. Then we present our $f_{\text{NL}}^{\text{loc}}$ forecasts using an MCMC-based approach.

4.1. Modeling of LBG properties

4.1.1. Modeling of the LBG large-scale linear bias

For the galaxy large-scale bias, we follow different prescriptions discussed in the literature. The first one we study is introduced by Wilson & White (2019),

$$b_{W19}(z, m_{\text{detect}}) = A(m_{\text{detect}})(1+z) + B(m_{\text{detect}})(1+z)^2, \quad (16)$$

with $A(m_{\text{detect}}) = -0.98(m_{\text{detect}} - 25) + 0.11$ and $B(m_{\text{detect}}) = 0.12(m_{\text{detect}} - 25) + 0.17$. Here, m_{detect} is the limiting magnitude of the LBG sample. For *u*-dropout (resp. *g*-dropout and *r*-dropout) LBGs, the limiting magnitude corresponds to the *r*-band (resp. *i*- and *z*-band). This prescription successfully reproduces the large-scale bias measurements for various dropout selections and limiting magnitudes. To allow additional flexibility and possible uncertainty in this bias prescription, we introduce a global rescaling amplitude b_0 , so that the LBG bias is modeled as

$$b(z) = b_0 b_{W19}(z, m_{\text{detect}}). \quad (17)$$

4.1.2. Fraction of LBG outlier and their galaxy bias

In this section, we expect our LBG samples to be contaminated by a fraction f_{out} of outliers, from which we can define a proper large-scale linear bias. We use a two-population model for a given sample with given total $n(z)$, with some “outliers” at redshift $z < z_{\text{mid}}$ with given redshift distribution $n_{\text{out}}(z)$, outlier fraction f_{out} and bias $b_{\text{out}}(z)$, along with a high-redshift sample at $z > z_{\text{mid}}$, with redshift distribution $n_{\text{highz}}(z)$ and bias $b_{\text{highz}}(z)$, such as

$$n(z) = f_{\text{out}} n_{\text{out}}(z) + (1 - f_{\text{out}}) n_{\text{highz}}(z). \quad (18)$$

We adopt the two-population bias model (Mergulhão et al. 2022)

$$b(z) = b_0 \times [b_1(z) + b_2(z)]. \quad (19)$$

where

$$(b_1, b_2) = \left(\frac{f_{\text{out}} n_{\text{out}}(z) b_{\text{out}}(z)}{n(z)}, \frac{(1 - f_{\text{out}}) n_{\text{highz}}(z) b_{\text{highz}}(z)}{n(z)} \right). \quad (20)$$

Moreover, we also use the model

$$b(z) = b_1(z) + b_0 \times b_2(z), \quad (21)$$

where this time, b_0 is connected *only* to the high-redshift portion of the sample, contrary to Eq. (19). As a result, only the high-redshift LBG bias is treated as unknown, and the uncertainty on $f_{\text{NL}}^{\text{loc}}$ is expected to be smaller than in the previous case, since the low-redshift outlier bias is fixed.

4.1.3. LBG redshift distribution uncertainty

Cosmological inference from the angular clustering of photometrically selected *u*-dropout LBGs relies on an accurate determination of their redshift distribution, $n(z)$ (Choi et al. 2016; Petri et al. 2025). Calibrated $n(z)$ has error bars that translate the uncertainty in calibrating the redshift distribution of LBGs. We consider a very simple toy example where the LBG sample has a $n(z)$ with error bar $\sigma(z)$. To propagate the uncertainty of the $n(z)$ on the cosmological fits, we consider random samples $\hat{n}_k \sim \mathcal{N}(n(z), \sigma(z))$, where $\sigma(z)$ is taken to be $\alpha n(z)$, i.e. a fraction of the total $n(z)$. We consider $\alpha = 0.2$ (i.e. 20% error on the recovered redshift distribution).

We explore another $n(z)$ -sampling technique, inspired from the *shift-and-stretch* standard method in the literature (Myles et al. 2021; d’Assignies et al. 2025b; Giannini et al. 2025), where we shift the high-redshift part of the Payerne et al. (2025) $n(z)$ (left panel of Figure 1), i.e. $z > 1$ part around its mean $z = 2.8$ by a factor $\Delta z \sim \mathcal{N}(0, 0.1)$, and stretch it by a factor $1 + \alpha \sim \mathcal{N}(1, 0.1)$.

4.1.4. Accounting for all with Cz estimates

The clustering-redshift method estimates the product $\langle b_{\text{LBG}} n_{\text{LBG}} \rangle(z)$, directly accounting for outliers. This inherent degeneracy between $b(z)$ and $n(z)$ cannot be disentangled without highly precise UNIONS photometric redshifts, making it practically impossible to separate the two terms $n(z)$ and $b(z)$. For standard galaxy clustering analysis, this is indeed advantageous, as marginalizing over clustering redshift uncertainties accounts for both the redshift evolution of the distribution and bias, as the degeneracy between b and n is also in the intrinsic kernel, cf. Eq. (3). As we are considering an additional term

from non-Gaussianity, we slightly modify the kernel in Eq. (8) with

$$b_{\text{LBG}} n_{\text{LBG}}(z) \left[1 + 2\delta_c \left(1 - \frac{p}{b_{\text{eff}}} \right) f_{\text{NL}}^{\text{loc}} \frac{3\Omega_m H_0^2}{2k^2 T(k) D(z)} \right], \quad (22)$$

where $b_{\text{eff}} = 1.5$, as computed previously. That way, $f_{\text{NL}}^{\text{loc}}$ is to be the only free parameter of the fit, as redshift uncertainty on bias and distribution, and outlier fraction are accounted for in the nb term. However, we are neglecting the impact of the redshift evolution on the galaxy bias in the second term. Let's note that the form of the second term is, on its own, not exact, as we are also assuming a specific (and redshift invariant) model for b_ϕ .

We further explore a more sophisticated bias modeling, adopting the prescription of [Wilson & White \(2019\)](#) used throughout this work. In this case, b_0 is no longer a free parameter but is instead fixed by the normalization constraint

$$\bar{b}_0 = \int_0^{+\infty} dz' \frac{b_{\text{LBG}} n_{\text{LBG}}(z')}{b_{W19}(z')} = 0.63, \quad (23)$$

From this, we replace b_{eff} in Eq. (22) by $\bar{b}_0 b_{W19}(z)$, incorporating redshift dependence in LBG bias.

4.2. MCMC-based forecasts

MCMC vs. Fisher: In this work, we perform forecasts to estimate the precision on $f_{\text{NL}}^{\text{loc}}$ from the angular distribution of UNIONS-like LBGs and their cross-correlation with *Planck* CMB lensing. In our forecast analysis, we consider two free parameters: (i) $f_{\text{NL}}^{\text{loc}}$, with a fiducial value of 0, and (ii) a galaxy bias-related parameter b_0 (see after), rescaled to 1 as its fiducial value. We choose to carry out these forecasts using MCMC, with the fiducial prediction serving as the mock data vector, rather than relying solely on Fisher-matrix forecasts. While Fisher methods are fast and provide a useful first-order estimate, they assume that the likelihood around the fiducial model is Gaussian and linear in the parameters. This approximation can break down in the presence of non-Gaussian posteriors, nonlinear parameter degeneracies, or physical parameter boundaries. By contrast, MCMC explores the full posterior distribution and naturally incorporates priors and parameter limits, providing a more accurate representation of the expected constraints.

MCMC setup: To forecast constraints on the amplitude of local-type primordial non-Gaussianity, we consider the observed dataset as a theoretical prediction for the clustering amplitudes of the LBG population and the CMB lensing potential, along with the corresponding theoretical covariances. In other words, the “data” we use in the forecasts are the binned theoretical predictions $\{C_b^{gg,\text{obs}}, C_b^{kg,\text{obs}}\}_{(b_0, f_{\text{NL}}^{\text{loc}})=(1,0)}$ where $C_b^{XY} = B_{b\ell} C_\ell^{XY}$, $B_{b\ell}$ is the binning matrix. We adopt a binning scheme inspired by [Krolewski et al. \(2024\)](#), with $\ell_{\text{min}} = 5$ (this lower limit is also imposed by the UNIONS sky coverage), $\ell_{\text{max}} = 300$ (safely describing galaxy clustering statistics as linear and redshift-only bias dependent), and $\Delta\ell = 5$, resulting in 60 bins. In the above equation, C_ℓ^{XY} is the *unbinned* full-sky angular power spectrum in multipole ℓ bins. Using the theoretical full-sky prediction is appropriate because it eliminates the variance of the estimator, resulting in a posterior distribution centered on the input values. However, this approach is a simplification, as it does not account for potential systematic biases in the signal estimation pipeline, which must be corrected in the analysis (e.g., via radial integral

constraints). We explore in Appendix B what could be the residual bias between the input theoretical angular power spectra used in our inference pipeline and the corresponding data vectors that would be obtained from realistic datasets.

Covariance: The covariance of the binned de-coupled angular power spectrum is given by

$$\text{Cov}(\widehat{C}_b^{XY}, \widehat{C}_{b'}^{ZW}) = B_{b\ell} \text{Cov}(\widehat{C}_\ell^{XY}, \widehat{C}_{\ell'}^{ZW}) B_{b'\ell'}, \quad (24)$$

where $B_{b\ell}$ is the binning matrix, and the covariance of the unbinned spectra is ([Brown et al. 2005](#))

$$\text{Cov}(\widehat{C}_\ell^{XY}, \widehat{C}_{\ell'}^{ZW}) = \frac{C_\ell^{XZ} C_\ell^{YW} + C_\ell^{XW} C_\ell^{YZ}}{2\ell + 1} \mathcal{M}_{\ell\ell'}^{-1}, \quad (25)$$

where $\mathcal{M}_{\ell\ell'}^{-1}$ is the mixing-mode matrix which accounts for partial sky coverage. We validate on mocks in Appendix B that this modeling choice is accurate to describe the Gaussian-disconnected part of the covariance. Not included in our mock validation pipeline are non-Gaussian contributions to the covariance, such as the super-sample covariance (SSC), which is currently considered to be the dominant non-Gaussian contribution. SSC arises from the non-linear modulation of local observables by long-wavelength density fluctuations.

Data vector: For our inference, we then adopt the theoretical “data vector” (i.e., the input prediction for the angular power spectrum), which is satisfactorily recovered by the mocks, along with the theoretical covariance that includes the full mode-coupling matrix.

Likelihood evaluation and covariance update: As stated before, we consider $f_{\text{NL}}^{\text{loc}}$ as a free parameter. We also include a free galaxy bias-related parameter, b_0 , with a default value of 1, used to rescale certain LBG bias dependencies (discussed below). For the measured LBG angular power spectrum, the cross-angular power spectrum between the LBG density and the CMB lensing, and for the combination of the two, the likelihoods (respectively, $\mathcal{L}_{gg}, \mathcal{L}_{kg}, \mathcal{L}_{gg+kg}$) are assumed to follow a multivariate Gaussian distribution. We note that the true likelihood is actually a generalized Gamma distribution ([Carron 2013](#)). It is well known that the Gamma distribution approaches a Gaussian distribution for a sufficiently large number of modes. For simplicity and computational efficiency, we compute the theoretical covariances at the fiducial values $(f_{\text{NL}}^{\text{loc}}, b_0) = (0, 1)$. A nonzero $f_{\text{NL}}^{\text{loc}}$ induces additional large-scale correlations, increasing the variance of the angular power spectrum; including this dependence in the covariance can impact parameter constraints ([Kodwani et al. 2019](#); [Eifler et al. 2009](#)). However, computing a parameter-dependent covariance at each likelihood evaluation is prohibitively expensive, and combining it with a Gaussian likelihood can artificially tighten posteriors derived from two-point statistics. As shown by [Carron \(2013\)](#), using a cosmology-dependent covariance in this way is formally inconsistent. Fixing the covariance, instead, ensures conservative information content and avoids biasing the inference. We then draw samples from the parameter posterior distribution using Bayes’ theorem:

$$\mathcal{P}(\theta|\text{data}) = \frac{\mathcal{L}_{\text{tot}}(\text{data}|\theta) \pi(\theta)}{\mathcal{L}_{\text{tot}}(\text{data})}, \quad (26)$$

where \mathcal{L}_{tot} corresponds to $\mathcal{L}_{gg}, \mathcal{L}_{kg}$, or \mathcal{L}_{gg+kg} . We use the `emcee` package ([Foreman-Mackey et al. 2013](#)) with flat priors on $f_{\text{NL}}^{\text{loc}}$ (between -500 and 500) and b_0 (between 0 and 5).

To accelerate the likelihood evaluation, we adopt the template method presented in [Fabbian et al. \(2025\)](#); Since our goal is to infer the LBG linear bias and the PNG parameter $f_{\text{NL}}^{\text{loc}}$, most contributions to the angular power spectra arise from pre-factors ($b_0, f_{\text{NL}}^{\text{loc}}$) multiplying terms that are independent of bias and PNG. This is always the case for $f_{\text{NL}}^{\text{loc}}$, and for b_0 when it simply rescales the overall bias of the considered population. We pre-compute the angular power spectra for a fiducial no-PNG cosmology ($f_{\text{NL}}^{\text{loc}} = 0$) with $b_0 = 1$, allowing only b_0 and $f_{\text{NL}}^{\text{loc}}$ to vary in the MCMC. This approach reduces the computation time of each likelihood evaluation to a few milliseconds and posterior estimations in seconds. The recovered uncertainties on $f_{\text{NL}}^{\text{loc}}$ and b_0 are taken to be the standard deviations of the recovered posterior distributions.

Marginalizing over parameters: When accounting for redshift distributions in the inference, a continuous model that can be readily marginalized over is often unavailable. On the other hand, generating realizations of $n(z)$ is usually straightforward (see Section 4.1.3 and Section 4.1.4). This is typically the situation described in [Bernstein et al. \(2025\)](#), except that we are dealing with only one bin at a time and our likelihood can be evaluated within milliseconds. Hence, we can directly compute the likelihood for every discrete realization n_k and effectively marginalize over them by stacking:

$$\mathcal{L}(x|y) \approx \frac{1}{N_{\text{real}}} \sum_k \mathcal{L}(x|y, n_k), \quad (27)$$

where N_{real} is the number of realizations, all assumed to have equal probability, independent of y .

5. Results on $f_{\text{NL}}^{\text{loc}}$ precision

The summary Table A.1 lists the forecasted precision on $f_{\text{NL}}^{\text{loc}}$ and b_0 for the various analysis cases we explore in this work.

5.1. From UNIONS-like LBG samples

The constraining power on $f_{\text{NL}}^{\text{loc}}$ from galaxy two-point statistics arises primarily from the largest clustering scales (i.e., the lowest multipoles ℓ).

The gain in $f_{\text{NL}}^{\text{loc}}$ precision enabled by the increased LBG density from UNIONS is only realized if the relevant scales of C_ℓ^{gg} are in the shot-noise-dominated regime. In the cosmic-variance-dominated regime, where $C_\ell^{\text{gg}} \gg 1/\bar{n}$ or equivalently $C_\ell^{\text{gg}} \bar{n} \gg 1$, the improvement from a higher number density saturates. For C_ℓ^{kg} , the cosmic-variance-dominated regime requires both $C_\ell^{\text{gg}} \gg 1/\bar{n}$ and $C_\ell^{\text{kk}} \gg N_\ell^{\text{kk}}$. In the lower plot of Figure 2, we show the quantities $C_\ell^{\text{gg}} \bar{n}$ and $C_\ell^{\text{kk}}/N_\ell^{\text{kk}}$. To define the cosmic-variance-dominated regime, we adopt the criteria $C_\ell^{\text{gg}} \bar{n} > 10$ and $C_\ell^{\text{kk}}/N_\ell^{\text{kk}} > 10$. We find that the LBG auto-spectrum is cosmic-variance dominated up to $\ell \lesssim 200$, implying that increasing the tracer density does not improve the signal-to-noise ratio on these scales. At smaller scales, however, a higher tracer density can still provide additional information on the tracer bias, rather than directly on the $f_{\text{NL}}^{\text{loc}}$ response, thereby helping to break the degeneracy between b_0 and $f_{\text{NL}}^{\text{loc}}$.

We first consider an idealized framework, with no outliers, no uncertainty, and with the redshift distribution set in [Payerne et al. \(2025\)](#) to evaluate the precision on $f_{\text{NL}}^{\text{loc}}$ and b_0 from a combination of the auto- and cross-spectra, C_ℓ^{gg} and C_ℓ^{kg} , respectively.




Fig. 2. Top panel shows the angular power spectra C_ℓ^{gg} , C_ℓ^{kg} , and C_ℓ^{kk} (solid lines) for the baseline sample. The cosmological-only contribution is shown as dashed lines, while the noise contribution is shown as dotted lines. Bottom panel shows the signal-to-noise ratio, defined as the ratio of the cosmological-only term to the noise, for C_ℓ^{gg} and C_ℓ^{kg} .

We obtain $\sigma(f_{\text{NL}}^{\text{loc}}) = 7$ from C_ℓ^{gg} alone. From C_ℓ^{kg} alone, the constraints are $\sigma(f_{\text{NL}}^{\text{loc}}) = 34$. The joint constraints (i.e., combining C_ℓ^{gg} and C_ℓ^{kg}) are very similar to those from C_ℓ^{gg} alone, reflecting the relatively larger uncertainties from the C_ℓ^{kg} -only case. The corresponding posterior distributions for the C_ℓ^{kg} -only and C_ℓ^{gg} -only analyses are shown in the left panel of Figure 3. We will now move to more realistic forecasts considering the different LBG uncertainties and systematics, cf. Section 4.1.

5.1.1. Impact of the angular momentum fitting range to mitigate imaging systematics

Imaging systematics, such as survey depth inhomogeneities and Galactic dust, can induce spurious density fluctuations in the tracer sample, generating artificial clustering signals. These effects are particularly problematic on the large scales most relevant for $f_{\text{NL}}^{\text{loc}}$ measurements, where they can lead to significant biases. A common mitigation strategy is to model the tracer density variations as linear functions of imaging properties and then re-weight the tracers, thereby suppressing spurious fluctuations while preserving the cosmological signal ([Chaussidon et al. 2022](#); [Krolewski et al. 2024](#)). In the context of $f_{\text{NL}}^{\text{loc}}$ inference from the tracer angular power spectrum, however, removing such systematic contributions from the auto-spectrum C_ℓ^{gg} remains challenging. For this reason, angular power spectrum-based analyses often rely primarily on the cross-spectrum with CMB lensing, C_ℓ^{kg} , since the large-scale modes of C_ℓ^{gg} are the most contaminated. It is therefore important to explore the constraining power of the LBG angular distribution when used exclusively in cross-correlation with the CMB lensing potential.

In the right panel of Figure 3, we show the precision on $f_{\text{NL}}^{\text{loc}}$ obtained from C_ℓ^{kg} only, with fitting ranges $\ell \in [\ell_{\text{min}}, 300]$ and $5 < \ell_{\text{min}} < 200$. As expected, the constraints rapidly degrade with increasing ℓ_{min} , since the sensitivity to $f_{\text{NL}}^{\text{loc}}$ resides on




Fig. 3. **Left:** Posterior distribution of the parameters $f_{\text{NL}}^{\text{loc}}$ and the large-scale linear bias (expressed in terms of relative error). **Right:** Forecasted error on $f_{\text{NL}}^{\text{loc}}$ as a function of the lower multipole cut ℓ_{min} . Solid lines show constraints using only $C_{\ell}^{\text{kg}}(\ell > \ell_{\text{min}})$, while dashed lines correspond to the combination of C_{ℓ}^{kg} and $C_{\ell}^{\text{gg}}(\ell > \ell_{\text{min}})$. Different colors indicate different LBG number densities: 800, 1100, and 2000 LBGs per deg^2 .

the largest clustering scales. In parallel, we show the constraints from a joint analysis of C_{ℓ}^{kg} and C_{ℓ}^{gg} , where the auto-spectrum is progressively truncated to $\ell \in [\ell_{\text{min}}, 300]$, under the assumption that the clustering signal of C_{ℓ}^{gg} can be reliably recovered only for $\ell > \ell_{\text{min}}$. We find that as ℓ_{min} increases, the precision on $f_{\text{NL}}^{\text{loc}}$ approaches the C_{ℓ}^{kg} -only result. The corresponding posterior distribution for $\ell_{\text{min}} = 200$ is shown in the left panel of Figure 3. While essentially no constraining power remains on $f_{\text{NL}}^{\text{loc}}$ in this case, the precision on the bias parameter b_0 is significantly improved. This indicates that relying on the small-scale clustering of the LBG population primarily constrains b_0 , rather than $f_{\text{NL}}^{\text{loc}}$, since these scales are insensitive to primordial non-Gaussianity but probe the large-scale linear bias more efficiently.

5.1.2. Impact of the $b_{\Phi}(b)$ parametrization

The sensitivity of the angular power spectrum to $f_{\text{NL}}^{\text{loc}}$ is entirely degenerate with the parameter b_{Φ} , through the combination $b_{\Phi}f_{\text{NL}}^{\text{loc}}$ in Eq. (8). In the previous section, we assumed the universality of the halo mass function, which allows a partial disentanglement of $f_{\text{NL}}^{\text{loc}}$ from the LBG bias via the $b_{\Phi}(b)$ relation in Eq. (9). For most real tracers of large-scale structure, however, this universality does not hold, and the relation must account for a parameter p_{Φ} that encodes the tracer’s merger history (Slosar et al. 2008), with $p_{\Phi} = 1$ corresponding to the universal case. The parameter p_{Φ} remains poorly constrained across tracer populations, and broad marginalization over it can degrade $f_{\text{NL}}^{\text{loc}}$ estimates (Barreira 2020). In the absence of reliable priors on p_{Φ} , only the combination $b_{\Phi}f_{\text{NL}}^{\text{loc}}$ can be constrained, although any nonzero detection of this product would still indicate the presence of local primordial non-Gaussianity. Besides the fiducial value $p_{\Phi} = 1$, we consider several alternative values inspired by Barreira (2020), namely $-1 < p_{\Phi} < 3$. We see that the accuracy on $f_{\text{NL}}^{\text{loc}}$ can improve if the parameter p is lower than unity—an aspect that can be further investigated using cosmological simulations.

5.1.3. Impact of the LBG linear bias

An essential ingredient in the modeling of $f_{\text{NL}}^{\text{loc}}$ is the tracer bias and its redshift dependence. In our baseline analysis, we adopt the prescription of Eq. (16), following Wilson & White (2019). This parametrization, however, is subject to modeling uncertainties, as current observational constraints on the LBG bias remain statistically limited (Ye et al. 2025; Ruhlmann-Kleider et al. 2024) and are highly sensitive to the adopted selection criteria.

As a simple test of modeling uncertainties, we can adopt a single effective bias parameter for the full redshift range, rather than a fully redshift-dependent prescription (see, e.g., Doux et al. (2018); Ye et al. (2025); Ruhlmann-Kleider et al. (2024)). However, to account for the presence of interlopers described with other biases, we use a two-population model (Mergulhão et al. 2022), which allows the data to self-consistently capture contamination while keeping the model minimal, with only a few free parameters: the outlier fraction f_{out} , and the outlier bias b_{out} and the high-redshift bias b_{highz} as presented in Eq. (19). For the baseline UNIONS-like RF sample, we choose the $z < 1$ (resp. $z > 1$) part of the $n(z)$ to represent the outliers (high-redshift LBGs) in our sample. We adopt the two-population bias model, which yields $f_{\text{out}} = 0.33$.

We fix the low-redshift bias to $b_{\text{out}} = \{0.5, 0.8, 1\}$ and the high-redshift bias to $b_{\text{highz}} = \{2.5, 3, 3.5\}$. Additionally, we consider $b_{\text{highz}}(z) = b_{\text{W19}}(z)$ from Eq. (16), giving an average high-redshift bias of $\langle b_{\text{W19}}(z) | z > 1 \rangle = 4.20$. The forecasted precision on $f_{\text{NL}}^{\text{loc}}$ for the different $\{b_{\text{out}}, b_{\text{highz}}\}$ choices is shown in full-lines in Figure 4. We see that, at fixed b_{out} , the constraints improve as b_{highz} increases. Conversely, at fixed b_{highz} , larger b_{out} values lead to weaker constraints. Adopting the two-population bias model with $b_{\text{out}} = \langle b_{\text{W19}}(z) | z < 1 \rangle = 1.02$ and $b_{\text{highz}} = \langle b_{\text{W19}}(z) | z > 1 \rangle = 4.20$ results in $\sigma(f_{\text{NL}}^{\text{loc}}) \simeq 35$, slightly larger than the baseline using the Wilson & White (2019) prescription for the full-redshift range bias. We also consider the two-population bias model in Eq. (21), where b_0 (the free parameter in the MCMC) is only connected to the high-redshift part of the sample, contrary to Eq. (19). The




Fig. 4. **Left:** Forecasted error on $f_{\text{NL}}^{\text{loc}}$ as a function of the high-redshift bias prescription $\bar{b}_2(z)$. Solid lines correspond to a full rescaling of the galaxy bias by b_0 in Eq. (19). In contrast, dashed lines correspond to rescaling only the high-redshift population bias by b_0 in Eq. (21). **Right:** Forecasted error on $f_{\text{NL}}^{\text{loc}}$ for different choices of $b_\phi(b)$, beyond the prescription given in Eq. (9).

results are shown in dashed lines in Figure 4, where constraining b_0 solely for the high-redshift population leads to slightly tighter $f_{\text{NL}}^{\text{loc}}$ constraints. We observe that the two-population model provides greater flexibility in modeling galaxy bias.

5.1.4. Impact of the outlier fraction

We model the Payerne et al. (2025) redshift distribution as in Eq. (18), depending on the outlier fraction at $z < 1$, (cf. Section 5.1.4). This modeling allows us to account for the effect of low-redshift outliers and to assess how $f_{\text{NL}}^{\text{loc}}$ constraints are impacted when the sample is more or less contaminated. We vary f_{out} following McDonald Mill et al. (2025), considering the values $\{0, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6\}$ while keeping the total LBG number density fixed. Maintaining the same number density ensures that the Random Forest required density budget is unchanged, but with varying efficiency in selecting high-redshift LBGs. For $f_{\text{out}} = 0$ (no contamination), we obtain the tightest constraint, $\sigma(f_{\text{NL}}^{\text{loc}}) = 16$. As f_{out} increases from 0.1 to 0.6, the precision degrades approximately linearly, with $\sigma(f_{\text{NL}}^{\text{loc}})$ increasing to $\sigma(f_{\text{NL}}^{\text{loc}}) = 161$.

For the two alternative Random Forest selections based on UNIONS-like and LSSTY4-like photometry yielding 1,100 LBG per square degrees, namely UNIONSlike_RF_u180s and LSSTY4like_RF_u2x180s (presented in Section 3), we obtain $\sigma(f_{\text{NL}}^{\text{loc}}) = 36$ and $\sigma(f_{\text{NL}}^{\text{loc}}) = 28$, respectively.

Still in the context of modifying the outlier fraction, we can assess the precision on $f_{\text{NL}}^{\text{loc}}$ for a DESI-II-like sample, i.e., when the outliers can be “removed” by spectroscopic redshift confirmation. Considering the $z > 2$ part of the sample only (and modifying the LBG density accordingly, i.e. 800 LBG per square degree) we get that $\sigma(f_{\text{NL}}^{\text{loc}}) = 15$. When accounting for the DESI spectroscopic redshift efficiency for 2-hour exposure (resp. 4-hour), we get $\sigma(f_{\text{NL}}^{\text{loc}}) = 22$ (resp. $\sigma(f_{\text{NL}}^{\text{loc}}) = 20$). We observe that a higher precision on $f_{\text{NL}}^{\text{loc}}$ can be achieved by removing outliers, which motivates enhanced sample selection from photometric datasets, and their spectroscopic follow-up, even if the current DESI spectroscopic redshift efficiency removes most of the $z < 2$ targets.

5.1.5. Impact of the redshift distribution uncertainty

Here we explore the impact of the $n(z)$ uncertainty on the PNG constraints, with two models described in Sect. 4.1.3. For the first, we consider random samples $\hat{n}_k \sim \mathcal{N}(n(z), \sigma(z))$, where $\sigma(z)$ is taken to be $0.2 n(z)$. In the second, we generate samples \hat{n}_k , with the so-called ‘shift and stretch’ method: we shift the high-redshift part of the $n(z)$ by a random factor $\Delta z \sim \mathcal{N}(0, 0.1)$, and stretch it by a random factor $1 + \alpha \sim \mathcal{N}(1, 0.1)$. To derive the error on $f_{\text{NL}}^{\text{loc}}$ accounting for the uncertainty on the $n(z)$, we stack the chains once converged and compute the mean and variance from the stacked chains (as explained in Eq. (27)).

For the first model, considering fitting C_ℓ^{kg} -only, the dispersion of $f_{\text{NL}}^{\text{loc}}$ and b_0 best-fits over the 50 MCMCs are $\text{disp}(f_{\text{NL}}^{\text{loc}}) = 6$ and $\text{disp}(b_0) = 0.01$, respectively. Marginal error on $f_{\text{NL}}^{\text{loc}}$ and b_0 are obtained after stacking the chains, and we get $\sigma(f_{\text{NL}}^{\text{loc}}) \approx 35$, that we retrieve if we consider the approximation $\sigma(f_{\text{NL}}^{\text{loc}}) = \sqrt{\sigma_{\text{fid}}^2(f_{\text{NL}}^{\text{loc}}) + \text{disp}^2(f_{\text{NL}}^{\text{loc}})}$, and $\sigma(b_0) \approx 0.08$ similar to the fiducial fit, the effect of $f_{\text{NL}}^{\text{loc}}$ precision is really small.

Using shift and stretch, we find similar results, namely that the dispersion of $f_{\text{NL}}^{\text{loc}}$ over the 50 MCMCs is roughly $\text{disp}(f_{\text{NL}}^{\text{loc}}) = 6$, leading to very few differences on the $f_{\text{NL}}^{\text{loc}}$ precision when stacking chains and taking the marginal dispersion, i.e. $\sigma(f_{\text{NL}}^{\text{loc}}) = 35$. From this simple exercise, we tested the robustness of $f_{\text{NL}}^{\text{loc}}$ constraints when accounting for the galaxy sample redshift uncertainty. We find that the effect is rather small for a well-calibrated $n(z)$.

5.2. From early UNIONS LBG samples

In this section, we go beyond our idealistic LBG samples obtained in Payerne et al. (2025) to explore how $f_{\text{NL}}^{\text{loc}}$ forecasts can be conducted using LBG selection on the early UNIONS multiband data. We first discuss the selection we use on the GAAP multiband catalog, and then we discuss the methodology to infer the LBG redshift distribution.

5.2.1. CLAUDS+HSC-calibrated $n(z)$

From the CLAUDS+HSC-calibrated redshift distribution presented in Figure 1 (right panel) and the lensing magnification bias for this sample, we forecast a precision of $\sigma(f_{\text{NL}}^{\text{loc}}) = 20$ when adopting the Wilson et al. (2017) bias model. This result is slightly better than that of the baseline photometric sample from Payerne et al. (2025), which contains a more significant fraction of low-redshift outliers. In contrast, the UNIONS selection includes a higher number of galaxies in the redshift range $1.5 < z < 2$, where the CMB lensing kernel peaks. Additionally, the UNIONS sample benefits from a higher galaxy density (increasing from 1100 to 1400 deg^{-2}), a lower magnification bias, and a higher mean bias, corresponding to a deeper r -band limit (24.2 vs. 24.3). The usable area for the forecast is also extended to the full UNIONS footprint, since the selection relies solely on ugr magnitudes. After convolving the redshift distribution with the DESI redshift reconstruction efficiency for 2-hour (resp. 4-hour) spectroscopic exposures—and adjusting for the corresponding number densities—we obtain $\sigma(f_{\text{NL}}^{\text{loc}}) = 24$ (resp. $\sigma(f_{\text{NL}}^{\text{loc}}) = 23$), assuming only the UNIONS North Galactic Cap (NGC) footprint that DESI-II could realistically survey. The significant reduction in LBG number density for the DESI-II spectroscopic follow-up with 2-hours and 4-hours exposure time (down to 140 and 170 deg^{-2}) arises from the higher proportion of UNIONS-LBGs that lie below $z < 2$ in the UNIONS selection compared to the selection in Payerne et al. (2025) peaking at $z > 3$ (down to 500 and 590 deg^{-2} from 1100 deg^{-2} initially).

5.2.2. From clustering-redshifts with DESI DR1 data

In this section, we evaluate the degradation of the PNG constraints, using the clustering-redshift estimates of the LBG large-scale bias and redshift distribution (see Section 4.1.4).

First, we fix bn to the measured $b_{\text{LBG}}n_{\text{LBG}}$ with clustering redshifts, with corresponding $b_{\text{eff}} = 1.5$, leaving $f_{\text{NL}}^{\text{loc}}$ to be the only free parameter of the fit. For this setup, we obtain a fiducial uncertainty of $\sigma(f_{\text{NL}}^{\text{loc}}) = 57$ (without accounting for uncertainties in the clustering-redshift $b(z)n(z)$ distribution), as the lower effective bias naturally reduces the constraining power.

To assess the impact of the uncertainty of the clustering-redshifts distribution, we generate random realizations $\widehat{bn}(z) \sim \mathcal{N}(b_{\text{LBG}}n_{\text{LBG}}(z), \sigma_{\text{CZ}}(z))$, and ensure that $\widehat{bn}(z)$ remains positive. For each realization, we compute an effective bias \widehat{b}_{eff} as the redshift-integrated value of $\widehat{bn}(z)$. The fiducial data vector is obtained using the reference $b_{\text{LBG}}n_{\text{LBG}}(z)$ and b_{eff} in Eq. (22), and 50 MCMC realizations are performed, accounting for the sampled $\widehat{bn}(z)$ and \widehat{b}_{eff} as input for the C_{ℓ}^{kg} modeling.

From this, the marginal uncertainty reaches $\sigma(f_{\text{NL}}^{\text{loc}}) = 134$ when stacking all converged chains. This corresponds to roughly a factor-of-two increase compared to the idealized case, reflecting a reduction in the forecasting power of the cross-correlation power spectrum. Importantly, this degradation is driven not by the dispersion in $n(z)$ itself, but by the effective bias b_{eff} , which is closely linked to the constraining power on $f_{\text{NL}}^{\text{loc}}$ —higher bias naturally leads to a better signal-to-noise ratio. Repeating the analysis with the error bars of the inferred clustering-redshift distribution reduced by a factor of $\sqrt{3}$ —corresponding to the expected improvement in clustering-redshift precision with upcoming DESI DR3 data—reduces the uncertainty to $\sigma(f_{\text{NL}}^{\text{loc}}) \approx 105$, highlighting the strong potential for improved clustering-redshift calibration with next-generation spectroscopic data.

Then we use the model which includes the impact of redshift evolution on the PNG term, and we propagate uncertainties on the clustering-redshift distribution via $\widehat{bn}(z) \sim \mathcal{N}(b_{\text{LBG}}n_{\text{LBG}}(z), \sigma_{\text{CZ}}(z))$ and \widehat{b}_0 to be the redshift integral of $\widehat{bn}(z)/b_{\text{W19}}(z)$. From this model, we obtain $\sigma(f_{\text{NL}}^{\text{loc}}) = 55$ (neglecting uncertainties), $\sigma(f_{\text{NL}}^{\text{loc}}) = 147$ when including DESI DR1 uncertainties, and $\sigma(f_{\text{NL}}^{\text{loc}}) = 120$ when assuming DESI DR3 performance. We can see that the more sophisticated model is not associated with an increase in the uncertainty in the fixed case, and creates a 15% degradation accounting for uncertainties. Thus, reducing the uncertainties associated with the clustering-redshift measurements is the main limitation of this method.

This framework offers a promising avenue for combining clustering-redshift information with realistic bias modeling, representing an important step toward fully data-driven constraints on primordial non-Gaussianity. Even though current uncertainties remain significant and are the major contribution to the uncertainty on $f_{\text{NL}}^{\text{loc}}$, this exploratory analysis demonstrates the feasibility of applying clustering-redshift methods to LBG samples over such a wide redshift range, by combining the analysis of multiple DESI tracers.

6. Summary and Conclusions

In this work, we investigated how effectively UNIONS-selected Lyman Break Galaxies (LBGs) can constrain local primordial non-Gaussianities, quantified by the parameter $f_{\text{NL}}^{\text{loc}}$. When measured from large-scale structure, $f_{\text{NL}}^{\text{loc}}$ is intrinsically degenerate with the galaxy bias, represented here by b_0 , and both parameters are jointly fitted throughout this study.

From UNIONS-like LBG sample: Using an idealized LBG sample from Payerne et al. (2025) identified from broadband $ugriz$ bands using a Random Forest algorithm, peaking at $z = 2$ and with a surface density of 1,100 deg^{-2} , we find that the scales most relevant for $f_{\text{NL}}^{\text{loc}}$ measurement ($\ell = 5\text{--}200$) are dominated by cosmic variance. Consequently, expanding the survey area—and therefore increasing the number of available modes—directly enhances the precision on $f_{\text{NL}}^{\text{loc}}$ derived from the LBG angular power spectrum. In contrast, the cross-correlation between the LBG distribution and the Planck CMB lensing map remains limited by the noise of the latter. For our fiducial model, we obtain $\sigma(f_{\text{NL}}^{\text{loc}}) = 34$, with a 9% uncertainty in the recovered galaxy bias, assuming the universality relation $b_{\phi} = 2\delta_c(b - p)$ with $p = 1$. However, the accuracy can improve if the parameter p is lower than unity—an aspect that can be further investigated using cosmological simulations. Introducing a more complex, two-population bias model degrades the precision to $\sigma(f_{\text{NL}}^{\text{loc}}) = 44$. We also examined the impact of photometric outliers on $f_{\text{NL}}^{\text{loc}}$ constraints. Tests using more realistic UNIONS-like datasets, from co-added U -band exposure on COSMOS, confirm the strong role of outliers in degrading $f_{\text{NL}}^{\text{loc}}$ constraints. Considering UNIONS as an option for DESI-II target selection naturally removes outliers below $z = 2.2$ through spectroscopic confirmation, yielding a precision of $\sigma(f_{\text{NL}}^{\text{loc}}) = 22$, accounting for DESI redshift efficiency. This clearly demonstrates that reducing the fraction of outliers—even at the cost of a lower LBG density—significantly enhances the accuracy of $f_{\text{NL}}^{\text{loc}}$ measurements. We further assessed the robustness of our forecasts to uncertainties in the underlying redshift distribution $n(z)$. Randomizing $n(z)$ at the 20% level or applying a “shift-and-stretch”

modification impacts $\sigma(f_{\text{NL}}^{\text{loc}})$ by less than 1%, confirming the stability of our results against such variations.

From UNIONS LBG sample: Finally, leveraging early UNIONS data, we tested LBG selection on real observations. The LBG sample obtained on XMM can be matched to the deep photometric catalog from CLAUDS+HSC to assess their $n(z)$, from which we found $\sigma(f_{\text{NL}}^{\text{loc}}) = 20$ with the [Wilson & White \(2019\)](#), leveraging the wider UNIONS footprint (the UNIONS selection relies only on *ugr*, not *ugriz* as in [Payerne et al. \(2025\)](#)), larger bias (lower magnitude limit) and a broader coverage of the CMB lensing kernel). In the context of DESI-II, spectroscopic follow-up would allow us to reach the high-redshift end of the distribution with no uncertainties on the redshift distribution, providing $\sigma(f_{\text{NL}}^{\text{loc}}) = 24$. We then use the clustering-redshift technique ([Ménard et al. 2013](#); [d’Assignies et al. 2025b](#)) to assess the redshift distribution of the UNIONS photometrically selected LBGs by cross-correlating them with DESI DR1 tracers over the wide range $0 < z < 3.5$. We find good agreement with the CLAUDS+HSC calibrated distribution, demonstrating the application of clustering redshifts to such an extensive redshift interval for LBGs. However, when accounting for uncertainties in the clustering-redshift $b(z)n(z)$ distribution, the constraint broadens to $\sigma(f_{\text{NL}}^{\text{loc}}) = 134$, yet is expected to improve significantly to $\sigma(f_{\text{NL}}^{\text{loc}}) = 105$ with DESI DR3 data. Overall, these exploratory results highlight the promise of the clustering-redshift method as an independent method to validate the recovered photometric redshift distribution up to $z > 2$, without relying on deep photometric catalog or spectroscopic follow-ups.

Acknowledgements. We thank David Alonso for his help in using the Core Cosmology Library (CCL, [Chisari et al. \(2019\)](#)) and NaMaster ([Alonso et al. 2019](#)). C. P. and C. Y. acknowledge support from grant ANR-22-CE31-0009 for the HZ-3D-MAP project and from grant ANR-22-CE92-0037 for the DESI-Lya project. W. d’A. acknowledges support from the MICINN projects PID2019-111317GB-C32, PID2022-141079NB-C32 as well as predoctoral program AGAUR-FI ajuts (2024 FI-1 00692) Joan Oró. IFAE is partially funded by the CERCA program of the Generalitat de Catalunya.

References

Aihara, H. et al. 2022, PASJ, 74, 247
 Alonso, D. et al. 2019, MNRAS, 484, 4127
 Anbajagane, D., Chang, C., Lee, H., & Gatti, M. 2024, JCAP, 2024, 062
 Barreira, A. 2020, JCAP, 2020, 031
 Barreira, A. et al. 2020, JCAP, 2020, 013
 Benítez, N. 2011, ASCL, ascl:1108.011
 Bernstein, G. et al. 2025, arXiv e-prints, arXiv:2506.00758
 Brown, M. L. et al. 2005, MNRAS, 360, 1262
 Busca, N. & Balland, C. 2018, arXiv e-prints, arXiv:1808.09955
 Cagliari, M. S. et al. 2024, JCAP, 2024, 036
 Carron, J. 2013, A&A, 551, A88
 Carron, J. et al. 2022, JCAP, 2022, 039
 Castorina, E. et al. 2019, JCAP, 2019, 010
 Chambers, K. C. et al. 2016, arXiv e-prints, arXiv:1612.05560
 Chaussidon, E. et al. 2022, MNRAS, 509, 3904
 Chaussidon, E. et al. 2025, JCAP, 2025, 029
 Chisari, N. E. et al. 2019, ApJS, 242, 2
 Choi, A., Heymans, C., Blake, C., et al. 2016, MNRAS, 463, 3737
 Choppin de Janvry, J. et al. 2025, arXiv e-prints, arXiv:2511.18133
 Crenshaw, J. F. et al. 2025, arXiv e-prints, arXiv:2503.06016
 Cross, D. N. & Sánchez, C. 2024, Phys. Rev. D, 110, 123534
 Dalal, N. et al. 2008, Phys. Rev. D, 77, 123514
 d’Assignies, W. et al. 2025a, arXiv e-prints, arXiv:2510.23565
 d’Assignies, W. et al. 2025b, arXiv e-prints, arXiv:2505.10416
 Desjacques, V., Jeong, D., & Schmidt, F. 2018, Physics Reports, 733, 1
 Desjacques, V. & Seljak, U. 2010, Classical and Quantum Gravity, 27, 124011
 Desprez, G., Picouet, V., Moutard, T., et al. 2023, A&A, 670, A82
 Doux, C. et al. 2018, MNRAS, 480, 5386
 Eifler, T. et al. 2009, A&A, 502, 721

Elvin-Poole, J., MacCrann, N., Everett, S., et al. 2023, MNRAS, 523, 3649
 Euclid Collaboration: Lepori, F. et al. 2022, A&A, 662, A93
 Fabbian, G. et al. 2025, arXiv e-prints, arXiv:2504.20992
 Finkelstein, S. L. et al. 2022, ApJ, 940, L55
 Foreman-Mackey, D. et al. 2013, PASP, 125, 306
 Foucaud, S. et al. 2003, A&A, 409, 835
 Giannini, G. et al. 2025, arXiv e-prints, arXiv:2509.07964
 Giavalisco, M. et al. 2004, ApJ, 600, L103
 Guy, J. et al. 2023, AJ, 165, 144
 Gwyn, S. et al. 2025, arXiv e-prints, arXiv:2503.13783
 Harikane, Y. et al. 2022, ApJS, 259, 20
 Harikane, Y. et al. 2023, ApJS, 265, 5
 Herrera-Alcantar, H. K. et al. 2025, arXiv e-prints, arXiv:2507.21852
 Hildebrandt, H. et al. 2009, arXiv e-prints, arXiv:0903.3951
 Hildebrandt, H. et al. 2021, A&A, 647, A124
 Hivon, E., Górski, K. M., Netterfield, C. B., et al. 2002, ApJ, 567, 2
 Ibata, R. A., McConnachie, A., Cuillandre, J.-C., et al. 2017, ApJ, 848, 128
 Jarvis, M. 2015, Astrophysics Source Code Library, record ascl:1508.007
 Jeong, D., Schmidt, F., & Sefusatti, E. 2011, Phys. Rev. D, 83, 123005
 Kodwani, D. et al. 2019, The Open Journal of Astrophysics, 2, 3
 Krolewski, A., Percival, W. J., Ferraro, S., et al. 2024, JCAP, 2024, 021
 Kuijken, K. 2008, A&A, 482, 1053
 Kuijken, K. et al. 2015, MNRAS, 454, 3500
 Kuijken, K. et al. 2019, A&A, 625, A2
 Landy, S. D. & Szalay, A. S. 1993, ApJ, 412, 64
 Lewis, A. & Challinor, A. 2006, Phys. Rep., 429, 1
 Lima, M., Cunha, C. E., Oyaizu, H., et al. 2008, MNRAS, 390, 118
 Limber, D. N. 1953, ApJ, 117, 134
 LSST Science Collaboration et al. 2009, arXiv e-prints, arXiv:0912.0201
 Malkan, M. A. et al. 2017, ApJ, 850, 5
 McDonald Mill, C. et al. 2025, arXiv e-prints, arXiv:2509.08052
 Ménard, B. et al. 2013, arXiv e-prints, arXiv:1303.4722
 Mergulhão, T. et al. 2022, JCAP, 2022, 021
 Miyatake, H. et al. 2022, Phys. Rev. Lett., 129, 061301
 Miyazaki, S. et al. 2018, PASJ, 70, S1
 Mueller, E.-M. et al. 2022, MNRAS, 514, 3396
 Myles, J. et al. 2021, MNRAS, 505, 4249
 Okamoto, T. & Hu, W. 2003, Phys. Rev. D, 67, 083002
 Ono, Y. et al. 2018, PASP, 70, S10
 Payerne, C. et al. 2025, JCAP, 2025, 031
 Petri, F. et al. 2025, arXiv e-prints, arXiv:2506.06475
 Planck Collaboration et al. 2020, A&A, 641, 24
 Reddy, N. A., Steidel, C. C., Pettini, M., et al. 2008, ApJS, 175, 48
 Rezaie, M. et al. 2021, MNRAS, 500, 1
 Ross, A. J. et al. 2025, JCAP, 2025, 125
 Roster, W. et al. 2025, arXiv e-prints, arXiv:2508.02779
 Ruhlmann-Kleider, V. et al. 2024, JCAP, 2024, 059
 Sawicki, M. et al. 2019, MNRAS, 489, 5202
 Schmidt, S. J. et al. 2013, MNRAS, 431, 3307
 Schmittfull, M. & Seljak, U. 2018, Phys. Rev. D, 97, 123540
 Slosar, A. et al. 2008, JCAP, 2008, 031
 Steidel, C. C. et al. 1996, ApJ, 462, L17
 Steidel, C. C. et al. 1999, ApJ, 519, 1
 Takahashi, R. et al. 2012, ApJ, 761, 152
 Tudorica, A. et al. 2017, A&A, 608, A141
 Wilson, M. J. & White, M. 2019, JCAP, 2019, 015
 Wilson, M. J. et al. 2017, MNRAS, 464, 3121
 Ye, I. et al. 2025, arXiv e-prints, arXiv:2507.02515
 Yu, B. et al. 2018, arXiv e-prints, arXiv:1809.02120
 Zhang, Y.-H. et al. 2025, arXiv e-prints, arXiv:2508.20903

Analysis case	$\sigma(f_{\text{NL}}^{\text{loc}})$	$\sigma(b_0)/b_0$
Section 5.1: Baseline sample UNIONSlike_RF		
$\bar{n}_{\text{gal}} = 1, 100 \text{ deg}^{-2}, S_{\text{survey}} = 3, 730 \text{ deg}^2$		
$C_{\ell}^{\text{gg}} + C_{\ell}^{\text{gk}}$	7	0.01
$C_{\ell}^{\text{gk}} + C_{\ell}^{\text{gg}} (\ell > 100)$	25	0.02
C_{ℓ}^{gk} -only	34	0.08
$C_{\ell}^{\text{gk}} (\ell > 25)$	61	0.09
Section 5.1.2: Impact of $b_{\phi}(b)$ modeling		
$p_{\phi} = 0.5-2$	31-62	0.08-0.14
Section 5.1.3: Impact of linear bias modeling		
$b_0 \times (1.0 + 3.5)$	44	0.08
$b_0 \times (1.0 + b_{\text{W19}})$	36	0.07
$1.0 + b_0 \times 3.5$	38	0.09
$1.0 + b_0 \times b_{\text{W19}}$	34	0.08
Section 5.1.4: Impact of the outliers fraction		
$f_{\text{out}} = 0$ (no $z < 1$ outliers)	16	0.09
$f_{\text{out}} = 0.6$	161	0.14
UNIONSlike_RF_u180s	36	0.08
LSST41like_RF_u2x180s	28	0.08
baseline high-z only	15	0.11
baseline (DESI-2h/4h exp.)	22/20	0.12
Section 5.1.5: Uncertainty on $n(z)$: baseline sample		
20% error on $n(z)$	35	0.08
high-z SSM $\Delta z, \Delta s = 0.1, 0.1$	35	0.08
Section 5.2: Color-box selection on GAAP UNIONS data		
$\bar{n}_{\text{gal}} = 1, 400 \text{ deg}^{-2}, S_{\text{survey}} = 4, 760 \text{ deg}^2$		
$n(z)$ CLAUDS+HSC calib.	20	0.07
+ (DESI-2h/4h exp.)	21/19	0.09
$n(z)b(z)$ CZ-calib.+ b_{eff}	57	-
+err DESI DR1	134	-
+err DESI DR3-like	105	-
$n(z)b(z)$ CZ-calib.+ $b_0 b_{\text{W19}}$	55	-
+err DESI DR1	147	-
+err DESI DR3-like	120	-

Table A.1. Summary of the forecasted precision on $f_{\text{NL}}^{\text{loc}}$ and the precision on the bias parameter $\sigma(b_0)/b_0$ for the different analysis cases we explore in this paper.

Appendix A: Summary of $f_{\text{NL}}^{\text{loc}}$ forecasts

We present in Table A.1 a summary of the different forecast results on $f_{\text{NL}}^{\text{loc}}$ and b_0 (when considered, only the fractional error is reported in this table) that we conducted in this paper. Results are sorted by paper sections.

Appendix B: Validation of the fiducial data vector using C_{ℓ} -based mocks

we adopt a mock-based approach:

- We generate, using Healpy, a full-sky LBG overdensity field $\bar{\delta}_i^g$ on a HEALPix grid with NSIDE=512, starting from an input C_{ℓ}^{gg} without including the shot-noise contribution (by taking only the first term in Eq. (6)).
- Each smoothed overdensity field value $\bar{\delta}_i^g$ in a given HEALPix cell⁶ is then used to generate the number of observed LBGs

⁶ For a pixelized galaxy overdensity map—where the pixel size is set by NSIDE—the finite resolution of the HEALPix grid introduces a smoothing of the underlying field. Each pixel effectively averages the signal over its solid angle, leading to a suppression of power on angular




Fig. B.1. **Upper panel:** Bias between the mean decoupled power spectra and the theoretical power spectra. **Lower panel:** Bias between the dispersion of the estimators over 100 simulations and (i) the theoretical variance computed with the full mode-coupling matrix $M_{\ell\ell'}$, and (ii) the approximation $M_{\ell\ell'} = \delta_{\ell\ell'}^k$.

in that cell, according to

$$\widehat{N}_i \sim \text{Pois}(\lambda_i = S_i \bar{n}_{\text{gal}} (1 + \bar{\delta}_i^g)), \quad (\text{B.1})$$

where \bar{n}_{gal} is the target angular number density of LBGs and S_i is the surface area of the i -th pixel. This procedure naturally incorporates the effect of shot noise.

- We compute the observed overdensity field, defined as

$$\widehat{\delta}_i^{\text{g,obs}} = \frac{\widehat{N}_i - \langle \widehat{N}_j \rangle_{\text{mask}}}{\langle \widehat{N}_j \rangle_{\text{mask}}}, \quad (\text{B.2})$$

where $\langle \widehat{N}_j \rangle_{\text{mask}}$ denotes the average number of galaxies per HEALPix pixel within a circular aperture of opening angle

$$\theta_{\text{survey}} = \arccos(1 - 2f_{\text{sky}}). \quad (\text{B.3})$$

For simplicity, we adopt a circular aperture corresponding to a sky fraction f_{sky} rather than using the actual UNIONS footprint.

- Alongside the LBG overdensity map, we generate a correlated CMB lensing convergence field,

$$\widehat{a}_{\ell m}^k = \widehat{a}_{\ell m}^{k,\text{cosmo}} + \widehat{a}_{\ell m}^{k,\text{noise}}, \quad (\text{B.4})$$

where $\widehat{a}_{\ell m}^{k,\text{cosmo}} \sim \mathcal{N}(0, C_{\ell}^{kk})$ with randomized phases, and $\widehat{a}_{\ell m}^{k,\text{noise}} \sim \mathcal{N}(0, N_{\ell}^{kk})$ is drawn from the CMB lensing noise spectrum of Carron et al. (2022).

- We use the publicly available NaMaster (Alonso et al. 2019) package to compute both the pseudo angular power spectra $\widehat{\text{PCL}}_{\ell}^{\text{gg}}, \widehat{\text{PCL}}_{\ell}^{\text{kg}}$ on the masked sky through Eq. (C.7), and we convert them to binned-decoupled angular power spectra $\widehat{C}_{\ell}^{\text{gg}}, \widehat{C}_{\ell}^{\text{kg}}$ through Eq. (C.11) (some calculation details are presented in Appendix C) by using the mixing-mode matrix

scales comparable to or smaller than the pixel size. This suppression is characterized by the pixel window function, which depends on the chosen NSIDE.

$\mathcal{M}_{\ell\ell'}^{-1}$ in Eq. (C.10) which accounts for partial sky coverage. Let's note that due to the impact of the window function, the maximum multipole trusted for the binned angular power spectrum is $l_{\max} \sim 2\text{NSIDE} = 1024$. On the other side, the maximum scale to be probed depends on the survey geometry and is such that $l_{\min} \sim \pi/\theta_{\text{survey}} \approx 5$. Finally, we consider the average over $N = 30$ realization of the fields $\widehat{C}_b^{XY} = \langle \widehat{C}_b^{XY(k)} \rangle_{k \in \{1, \dots, 30\}}$.

We compare in Figure B.1 (upper panel) the residual between the measured \widehat{C}_b^{XY} and the input C_ℓ^{XY} , and find good agreement at all considered scales, as the residuals are scattered around zero. Also, there is no impact of the pixel window function at $\ell < 300$, so we neglect its correction at the modeling level.

Next, we compare the dispersion of the estimators to the predicted covariance of the partial-sky binned, decoupled estimator. The covariance of the binned de-coupled angular power spectrum is given by

$$\text{Cov}(\widehat{C}_b^{XY}, \widehat{C}_{b'}^{ZW}) = B_{b\ell} \text{Cov}(\widehat{C}_\ell^{XY}, \widehat{C}_{\ell'}^{ZW}) B_{b'\ell'}, \quad (\text{B.5})$$

where $B_{b\ell}$ is the binning matrix, and the covariance of the unbinned spectra is (Brown et al. 2005)

$$\text{Cov}(\widehat{C}_\ell^{XY}, \widehat{C}_{\ell'}^{ZW}) = \frac{C_\ell^{XZ} C_{\ell'}^{YW} + C_\ell^{XW} C_{\ell'}^{YZ}}{2\ell + 1} \mathcal{M}_{\ell\ell'}^{-1}, \quad (\text{B.6})$$

The recovered bias is plotted in Figure B.1, showing that the variance is in reasonable agreement with the theoretical covariance when using the full mode-mixing matrix, which provides a good prescription for small sky fractions. We also compare this to the commonly used approximation $\mathcal{M}_{\ell\ell'} \sim f_{\text{sky}} \delta_{\ell\ell'}^K$, which is valid for rather uniform masks and large sky fractions f_{sky} . This approximation is not valid in our case, as it underestimates the errors by 20–50% across a large range of angular scales. Therefore, we consider the full coupling matrix for a circular sky mask.

Appendix C: Formalism for the estimation of the angular power spectrum from LBG catalog and CMB lensing map

We consider a projected field s defined by

$$s(\theta) = \int_0^{\chi_H} d\chi q(\chi) \delta(\chi\theta, \chi). \quad (\text{C.1})$$

For two full-sky fields s^X and s^Y defined as

$$\widehat{s}_X(\theta) = \int_0^{\chi_H} d\chi q_X(\chi) \widehat{\delta}(\chi\theta, \chi). \quad (\text{C.2})$$

The angular power spectrum between the two fields is computed from the spherical harmonic coefficients

$$\widehat{a}_{\ell m} = \int d^2\theta \widehat{s}(\theta) Y_{\ell m}(\theta). \quad (\text{C.3})$$

The pseudo-angular power spectrum (Alonso et al. 2019) is given by

$$\widehat{C}_\ell^{XY} = \frac{1}{2\ell + 1} \sum_{m=-\ell}^{\ell} |\widehat{a}_{\ell m}^{X,*} \widehat{a}_{\ell m}^Y|^2 \quad (\text{C.4})$$

On full-sky, the pseudo angular power spectrum averages to

$$C_\ell^{XY} = \frac{2}{\pi} \int_0^\infty dk k^2 P(k) W_\ell^X(k) W_\ell^Y(k), \quad (\text{C.5})$$

where $P(k)$ is the 3D matter power spectrum, and $W_\ell^X(k)$ and $W_\ell^Y(k)$ are the radial transfer functions of the fields, given by

$$W_\ell^X(k) = \int dz \frac{d\chi}{dz} q_X(\chi) j_\ell(k\chi). \quad (\text{C.6})$$

Following the MASTER formalism (Hivon et al. 2002), which we adopt in this work, the pseudo-angular power spectrum between two cosmological fields X and Y , denoted by PCL_ℓ^{XY} , is given by

$$\text{PCL}_\ell^{XY} = \frac{1}{2\ell + 1} \sum_{m=-\ell}^{\ell} |a_{\ell m}^{X,*} a_{\ell m}^Y|^2, \quad (\text{C.7})$$

where the spherical harmonic coefficients are computed as

$$a_{\ell m} = \int d^2\theta s(\theta) Y_{\ell m}(\theta). \quad (\text{C.8})$$

The expectation value of the pseudo-angular power spectrum is related to the true angular power spectrum via

$$\langle \text{PCL}_\ell^{XY} \rangle = \mathcal{M}_{\ell\ell'} C_\ell^{XY}, \quad (\text{C.9})$$

where $\mathcal{M}_{\ell\ell'}$ is the mode-mixing matrix that accounts for the effects of partial sky coverage (Alonso et al. 2019), and is expressed as

$$\mathcal{M}_{\ell\ell'} = \frac{2\ell' + 1}{4\pi} \sum_{\ell''} (2\ell'' + 1) \text{PCL}_{\ell''}^{\text{mask}} \begin{pmatrix} \ell & \ell' & \ell'' \\ 0 & 0 & 0 \end{pmatrix}^2, \quad (\text{C.10})$$

where $\text{PCL}_{\ell''}^{\text{mask}}$ is the angular power spectrum of the sky mask. For a homogeneous circular mask with sky fraction f_{sky} , the matrix simplifies to $\mathcal{M}_{\ell\ell'} \approx \delta_{\ell\ell'}^K f_{\text{sky}}$, leading to its inverse $\mathcal{M}_{\ell\ell'}^{-1} \approx \delta_{\ell\ell'}^K / f_{\text{sky}}$. The "decoupled" (i.e., not pseudo) angular power spectrum C_ℓ^{XY} captures the cosmological signal, and is generally estimated as a binned data vector through

$$\widehat{C}_b^{XY} = \mathcal{M}_{bb'}^{-1} \text{PCL}_{b'}^{XY} \quad (\text{C.11})$$

where the binned mode-mixing matrix is given by $\mathcal{M}_{bb'} = B_{b\ell} \mathcal{M}_{\ell\ell'} B_{b'\ell'}^{-1}$, and the binned pseudo- C_ℓ is given by $\text{PCL}_b^{XY} = B_{b\ell} \text{PCL}_\ell^{XY}$. Then,

$$\langle \widehat{C}_b^{XY} \rangle = C_b^{XY} = B_{b\ell} C_\ell^{XY} \quad (\text{C.12})$$