

Piecewise polynomial approximation on non-Lipschitz domains

D. P. Hewett^a

^aDepartment of Mathematics, University College London, London, United Kingdom

December 1, 2025

Abstract

We prove best approximation error estimates for discontinuous piecewise polynomial approximation in fractional Sobolev spaces on non-Lipschitz meshes of non-Lipschitz domains. In particular, the boundary of the domain, and the boundaries of the mesh elements, can be fractal.

MSC2020 classifications: 41A10, 41A25, 65N30, 65N38

1 Introduction

Piecewise polynomial approximation on a domain $\Omega \subset \mathbb{R}^n$ is central to many numerical methods, including the finite element method (FEM) [7, 18] and integral equation methods [2, 26] such as the boundary element method [29]. In conforming FEM, the piecewise polynomial approximation space must be globally continuous. This imposes strict constraints on the mesh geometry, and classically one tends to work with conforming simplicial meshes. However, in discontinuous Galerkin (dG) FEM, and in many integral equation methods, one works with globally discontinuous piecewise polynomials, which introduces considerably more flexibility in the type of mesh that can be used. When the underlying domain has a complicated structure, the use of meshes with non-simplicial and even non-polytopal elements offers significantly more efficient and accurate approximation of the domain geometry than can be achieved using traditional simplicial meshes. In dG-FEM this idea is well established, with the theoretical state of the art being [13], which proves optimal hp best approximation error estimates for Lipschitz domains and a class of meshes with Lipschitz elements.

In recent years there has been increasing interest in the solution of PDEs on non-Lipschitz domains with fractal boundary [1, 4, 14, 19, 23], such as the Koch snowflake illustrated in Figure 1(a). Such domains cannot be meshed with a finite number of Lipschitz elements, so the results of [13] do not apply. An obvious remedy is to first replace the domain Ω with a smoother “prefractal” approximation, and mesh this. However, this can introduce significant geometry approximation errors; furthermore, meshing the prefractal approximation may require a large number of elements, particularly if one wants to work with shape-regular simplicial meshes. For an example of the extreme mesh complexity that can arise in such geometries with conforming FEM, see, e.g., [5].




Figure 1: (a) The Koch snowflake domain Ω , an open set with fractal boundary of Hausdorff dimension $\log 4 / \log 3$. (b) A mesh of Ω comprising 13 elements with fractal boundary (7 large and 6 small), each a scaled/rotated/translated version of Ω (see [10] for details). (c) A mesh of Ω comprising 12 elements, 6 of which are triangles and 6 of which have a boundary that is the union of a line segment and a fractal curve.

To mesh a domain like that in Figure 1(a) more efficiently one might consider working with meshes containing elements with fractal boundaries, such as those illustrated in Figure 1(b),(c), so that the geometry of Ω is fully captured. Such meshes have been successfully used in the context of integral equation methods for acoustic wave scattering by fractal structures in [8–10], using the numerical quadrature rules on fractal meshes presented in [20,21]. However, aside from the results for piecewise constant approximation presented in [8–10], piecewise polynomial approximation theory on such meshes does not seem to be available in the literature.

In this paper we provide best approximation error estimates for discontinuous piecewise polynomial approximation on non-Lipschitz meshes of non-Lipschitz domains. In fact, many of our results hold for arbitrary meshes of arbitrary domains, meaning that they require no regularity assumption whatsoever on the boundary of Ω , or on the boundaries of the mesh elements. In particular, these boundaries can be fractal, with any Hausdorff dimension in the interval $[n - 1, 1]$, or can even have positive Lebesgue measure.

Our estimates are stated in the context of Sobolev spaces, and when dealing with non-smooth domains it is important to clarify the precise Sobolev space setting in which one is working. In the FEM setting one typically requires estimates in “intrinsic” Sobolev spaces such as $W^m(\Omega)$, for some $m \in \mathbb{N}$, defined in terms of square integrability of weak derivatives on Ω . In the integral equation setting one often requires estimates in “extrinsic” spaces derived from the Bessel potential Sobolev space $H^s(\mathbb{R}^n)$, for some $s \in \mathbb{R}$, such as the space $H^s(\Omega)$ of restrictions to Ω of elements of $H^s(\mathbb{R}^n)$, or the space $\tilde{H}^s(\Omega)$ defined as the closure of $C_0^\infty(\Omega)$ in $H^s(\mathbb{R}^n)$. Here we provide results in both intrinsic and extrinsic spaces.

Our results are stated in §2 and the proofs can be found in §3. Our results can be summarised thus:

- In Lemma 2.3 we provide optimal hp local best approximation error estimates based on local regularity assumptions. Lemma 2.3 generalises [13, Lemma 4.31] to the non-Lipschitz case, and its proof, like that of [13, Lemma 4.31], makes use of a “covering mesh” approach. However, as is explained before its statement, Lemma 2.3 provides a slightly stronger result than [13,

Lemma 4.31]. This strengthening means that, when using Lemma 2.3 to derive global error estimates based on global regularity assumptions, in contrast to [13], we do not need to impose any condition analogous to [13, Assumption 4.28] restricting the number of mesh elements that can intersect each covering element.

- In Theorem 2.5, which is our main result, we provide global error estimates in broken Sobolev norms for functions u in the extrinsic spaces $H^m(\Omega)$ for arbitrary $m \in \mathbb{N}$ and for arbitrary meshes of arbitrary domains.
- In Corollary 2.6 we provide a version of Theorem 2.5 valid for functions u in the intrinsic space $W^m(\Omega)$. This result holds under the assumption that Ω is a W^m extension domain, which is an implicit regularity assumption on Ω (but not on the mesh). It holds whenever Ω is an (ϵ, δ) locally uniform domain, so in particular for the Koch snowflake domain in Figure 1(a).
- In Corollary 2.7 we derive $L^2(\Omega)$ estimates for functions in the fractional order spaces $H^s(\Omega)$ for general $s \in [0, m]$, and in Corollary 2.8 we provide estimates in negative order norms. These two corollaries are both valid for arbitrary meshes of arbitrary domains.
- Finally, in Corollary 2.9 we generalise the results of Corollary 2.8 to distributions of negative smoothness. This result requires the additional assumption that $\{H^s(\Omega)\}_{s \in [0, m]}$ (equivalently, $\{\tilde{H}^{-s}(\Omega)\}_{s \in [0, m]}$) is an interpolation scale, which again is an implicit regularity assumption on Ω (but not on the mesh). Conditions under which this holds are discussed in §2; we note that it holds for any (ϵ, δ) locally uniform domain, so in particular for the Koch snowflake domain in Figure 1(a).

As hinted at above, our work is largely motivated by recent developments in the numerical solution of integral equations on fractal domains [6, 8–10]. Indeed, a simplified piecewise constant version of Corollary 2.8 was used to analyse the convergence of a piecewise constant Galerkin boundary element method for scattering by fractal screens in [10]. Furthermore, the result of Corollary 2.7 will be applied in the forthcoming paper [6] to analyse piecewise polynomial volume integral equation methods for the Lippmann-Schwinger equation for scattering by fractal inhomogeneities.

Our results are also of relevance to the analysis of dG-FEM on non-smooth domains, and in a forthcoming paper [22] we will present and analyse a dG-FEM on a fractal domain using meshes with fractal elements of the type shown in Figure 1(b). We note that, for dG-FEM applications, in addition to error estimates for approximation on the mesh elements, one also typically needs error estimates for approximation on element boundaries (cf., e.g., [13, Eqn (4.25)]). In principle, such estimates can be derived for non-Lipschitz meshes of non-Lipschitz domains, provided that the boundaries are sufficiently regular to ensure that suitable trace operators exist. This holds in particular for the mesh of Figure 1(b), because the Koch snowflake domain is a $W^m(\Omega)$ extension domain and its boundary is a d -set (in the terminology of [25]) of Hausdorff dimension $d = \log 4 / \log 3$, so by results in [25] there exists a bounded linear trace operator from $W^1(\Omega)$ to $L^2(\partial\Omega, \mathcal{H}^d)$, where \mathcal{H}^d is the d -dimensional Hausdorff measure. However, we postpone further discussion of this topic to future work.

2 Definitions and results

Let $\Omega \subset \mathbb{R}^n$ be a domain, i.e. a non-empty open set. For $m \in \mathbb{N}$ let $W^m(\Omega)$ denote the Sobolev space of functions $u \in L^2(\Omega)$ with square-integrable weak derivatives up to order m , normed by $\|u\|_{W^m(\Omega)} = \left(\sum_{j=0}^m |u|_{W^j(\Omega)}^2 \right)^{1/2}$, where $|u|_{W^j(\Omega)} = \left(\sum_{|\alpha|=j} \|D^\alpha u\|_{L^2(\Omega)}^2 \right)^{1/2}$. For $s \in \mathbb{R}$ let $H^s(\mathbb{R}^n)$ denote the Bessel potential Sobolev space of tempered distributions u on \mathbb{R}^n for which $\|u\|_{H^s(\mathbb{R}^n)} := \left(\int_{\mathbb{R}^n} (1 + |\xi|^2)^s |\hat{u}(\xi)|^2 d\xi \right)^{1/2} < \infty$. Let $\tilde{H}^s(\Omega)$ denote the closure of $C_0^\infty(\Omega)$ in $H^s(\mathbb{R}^n)$, a closed subspace of $H^s(\mathbb{R}^n)$, and let

$$H^s(\Omega) := \{U|_\Omega : U \in H^s(\mathbb{R}^n)\}, \quad (1)$$

equipped with the quotient norm

$$\|u\|_{H^s(\Omega)} := \min_{\substack{U \in H^s(\mathbb{R}^n) \\ u=U|_\Omega}} \|U\|_{H^s(\mathbb{R}^n)}, \quad u \in H^s(\Omega). \quad (2)$$

We recall that for any domain $\Omega \subset \mathbb{R}^n$ and any $s \in \mathbb{R}$ the dual space of $H^s(\Omega)$ can be unitarily realised as the space $\tilde{H}^{-s}(\Omega)$ (with $L^2(\Omega)$ as pivot space) [16, Thm 3.3]. We also recall that, for $m \in \mathbb{N}_0$, $H^m(\mathbb{R}^n) = W^m(\mathbb{R}^n)$ with equivalent norms, and $H^m(\Omega)$ is continuously embedded in $W^m(\Omega)$. In general $H^m(\Omega)$ may be a strict subset of $W^m(\Omega)$. However, if Ω is a W^m extension domain, meaning that there exists a bounded linear operator $E_\Omega^m : W^m(\Omega) \rightarrow W^m(\mathbb{R}^n)$ such that $E_\Omega^m(u)|_\Omega = u$ for all $u \in W^m(\Omega)$, then $H^m(\Omega) = W^m(\Omega)$ with equivalent norms. This holds in particular if Ω is Lipschitz [12, 28], or more generally if Ω is (ϵ, δ) locally uniform [24, 27]. The latter class includes a large subclass of domains with fractal boundary; in particular, the Koch snowflake domain pictured in Figure 1 is (ϵ, ∞) locally uniform [24], and hence is a W^m extension domain.

Definition 2.1. *Given a domain $\Omega \subset \mathbb{R}^n$, we say that a countable (e.g. finite) collection $\mathcal{T} = \{K\}$ of non-empty bounded open subsets of Ω is a mesh of Ω if $K \cap K' = \emptyset$ for $K, K' \in \mathcal{T}$, $K \neq K'$, and*

$$\left| \Omega \setminus \bigcup_{K \in \mathcal{T}} K \right| = 0. \quad (3)$$

Given such a mesh, let $h_K := \text{diam}(K)$ for each $K \in \mathcal{T}$, and let $h := \sup_{K \in \mathcal{T}} h_K$ denote the meshwidth of \mathcal{T} .

As in [13], we will make use of the notion of a *covering* of a mesh.

Definition 2.2. *Given a domain $\Omega \subset \mathbb{R}^n$, and a mesh $\mathcal{T} = \{K\}$ of Ω , we define a covering $\mathcal{T}^\# = \{\mathcal{K}\}$ of \mathcal{T} to be a set of open simplices or cubes \mathcal{K} such that for each $K \in \mathcal{T}$ there exists at least one $\mathcal{K} \in \mathcal{T}^\#$ with $K \subset \mathcal{K}$. Given a covering $\mathcal{T}^\#$ we define $h_{\mathcal{K}} := \text{diam}(\mathcal{K})$ for each $\mathcal{K} \in \mathcal{T}^\#$. We say that $\kappa : \mathcal{T} \rightarrow \mathcal{T}^\#$ is a covering choice function if $K \subset \kappa(K)$ for each $K \in \mathcal{T}$.*

The following lemma provides an optimal local hp approximation result, generalising [13, Lemma 4.31] to the non-Lipschitz case. An important difference compared to [13, Lemma 4.31] is that we control in one estimate the contributions from all of the elements K that are associated (via the function κ) with a given covering element \mathcal{K} . This allows us to avoid the need for an assumption like [13, Assumption 4.28] when deriving the global estimates presented in Theorem 2.5.

Here, and henceforth, given $p \in \mathbb{N}_0$ and $S \subset \mathbb{R}^n$ let $\mathcal{P}_p(S)$ denote the set of polynomials on S in n variables of total degree less than or equal to p .

Lemma 2.3. *Let $\Omega \subset \mathbb{R}^n$ be a domain, \mathcal{T} a mesh of Ω with covering $\mathcal{T}^\#$, and $\kappa : \mathcal{T} \rightarrow \mathcal{T}^\#$ a covering choice function. Let $h_0 > 0$ and suppose that $h_K \leq h_0$ for each $K \in \mathcal{T}^\#$. Let $u \in L^2(\Omega)$ and suppose that for each $K \in \mathcal{T}^\#$ there exists $m_K \in \mathbb{N}_0$ and $U_K \in W^{m_K}(\mathcal{K})$ such that $(U_K)|_K = u|_K$ for each $K \in \kappa^{-1}(\mathcal{K})$. Then for each $K \in \mathcal{T}^\#$ and each $p \in \mathbb{N}_0$ there exists an operator $\pi_{p,m_K,\mathcal{K}} : W^{m_K}(\mathcal{K}) \rightarrow \mathcal{P}_p(\mathcal{K})$ and a constant $C_{n,h_0,m_K,\mathcal{K}} > 0$, depending only on n , h_0 , m_K and the shape regularity of \mathcal{K} , such that, with $t_K := \min(m_K, p + 1)$,*

$$\left(\sum_{K \in \kappa^{-1}(\mathcal{K})} \|u - \pi_{p,m_K,\mathcal{K}}(U_K)\|_{W^j(K)}^2 \right)^{1/2} \leq C_{n,h_0,m_K,\mathcal{K}} \frac{h_K^{t_K-j}}{(p+1)^{m_K-j}} \sum_{r=t_K}^{m_K} |U_K|_{W^r(\mathbb{R}^n)}, \quad j = 0, 1, \dots, m_K. \quad (4)$$

Lemma 2.3 is the foundation of our main result, Theorem 2.5, which provides global best approximation error estimates in broken Sobolev norms for h and p version refinements for functions in the extrinsic spaces $H^m(\Omega)$. We emphasize that Theorem 2.5 holds for arbitrary meshes of arbitrary domains.

Definition 2.4. *Given a domain $\Omega \subset \mathbb{R}^n$, a mesh $\mathcal{T} = \{K\}$ of Ω of meshwidth $h > 0$, and $p \in \mathbb{N}_0$, we define the piecewise polynomial space*

$$V_{h,p} := \{u \in L^2(\Omega) : u|_K \in \mathcal{P}_p \text{ for each } K \in \mathcal{T}\} \subset L^2(\Omega). \quad (5)$$

Theorem 2.5. *Let $n, m \in \mathbb{N}$ and $h_0 > 0$. There exists a constant $C_{n,h_0,m} > 0$, depending only on n , h_0 and m , such that if $\Omega \subset \mathbb{R}^n$ is a domain, \mathcal{T} is a mesh of Ω with meshwidth $h \in (0, h_0]$, $u \in H^m(\Omega)$, $U \in H^m(\mathbb{R}^n) = W^m(\mathbb{R}^n)$ is such that $U|_\Omega = u$, and $p \in \mathbb{N}_0$, then, with $t := \min(m, p + 1)$,*

$$\min_{v \in V_{h,p}} \left(\sum_{K \in \mathcal{T}} \|u - v\|_{W^j(K)}^2 \right)^{1/2} \leq C_{n,h_0,m} \frac{h^{t-j}}{(p+1)^{m-j}} \sum_{r=t}^m |U|_{W^r(\mathbb{R}^n)}, \quad j = 0, 1, \dots, m. \quad (6)$$

Theorem 2.5 immediately implies the following corollary for functions in the intrinsic spaces $W^m(\Omega)$, under the assumption that $\Omega \subset \mathbb{R}^n$ is an extension domain (cf. the discussion after (2)).

Corollary 2.6. *Let $n, m \in \mathbb{N}$ and $h_0 > 0$. If $\Omega \subset \mathbb{R}^n$ is a W^m extension domain and \mathcal{T} is a mesh of Ω with meshwidth $h \in (0, h_0]$, then, for each $p \in \mathbb{N}_0$, and with $t := \min(m, p + 1)$,*

$$\min_{v \in V_{h,p}} \left(\sum_{K \in \mathcal{T}} \|u - v\|_{W^j(K)}^2 \right)^{1/2} \leq C_{m,\Omega} C_{n,h_0,m} \frac{h^{t-j}}{(p+1)^{m-j}} \|u\|_{W^m(\Omega)}, \quad j = 0, 1, \dots, m, \quad u \in W^m(\Omega), \quad (7)$$

where $C_{n,h_0,m}$ is the constant from Theorem 2.5 and $C_{m,\Omega}$ is the norm of $E_\Omega^m : W^m(\Omega) \rightarrow W^m(\mathbb{R}^n)$.

From Theorem 2.5 one can derive three further corollaries, motivated by the integral equation applications in [6, 8–10]. The first concerns L^2 approximation errors for functions in fractional order spaces. We mention here that while the proof of Corollary 2.7 (presented in §3) uses function space interpolation, it does not require the spaces $\{H^s(\Omega)\}_{s \in [0,m]}$ to be an interpolation scale;¹ it requires only that $H^{\theta m}(\Omega)$ is continuously embedded in the interpolation space $K_{\theta,2}((L^2(\Omega), H^m(\Omega)))$ for every $\theta \in (0, 1)$, which holds for all open Ω by [15, Lem. 4.2].

¹This is known not to hold for arbitrary Ω - see, e.g., [15, Lem. 4.8].

Corollary 2.7. *Let $n, m \in \mathbb{N}$ and $h_0 > 0$. Suppose that $\Omega \subset \mathbb{R}^n$ is a domain, \mathcal{T} is a mesh of Ω with meshwidth $h \in (0, h_0]$, $p \geq m - 1$, and $0 \leq s \leq m$ (with s not necessarily an integer). Then*

$$\min_{v \in V_{h,p}} \|u - v\|_{L^2(\Omega)} \leq (C_{n,h_0,m})^{s/m} \left(\frac{h}{p+1} \right)^s \|u\|_{H^s(\Omega)}, \quad u \in H^s(\Omega). \quad (8)$$

The second, presented as Corollary 2.8 below, provides best approximation error estimates in negative order norms. To aid the interpretation of Corollary 2.8, we recall that, for any domain $\Omega \subset \mathbb{R}^n$, extension by zero is a unitary isomorphism from $H^0(\Omega) = L^2(\Omega)$ to $\tilde{H}^0(\Omega)$ (its inverse being the restriction operator). The fact that $\tilde{H}^{s_1}(\Omega) \supset \tilde{H}^{s_2}(\Omega)$ for $s_1 < s_2$ then implies that extension by zero provides a continuous embedding of $H^0(\Omega) = L^2(\Omega)$ into $\tilde{H}^s(\Omega)$ for $s \leq 0$, a fact we are using implicitly in what follows.

Corollary 2.8. *Let $n, m \in \mathbb{N}$ and $h_0 > 0$. Suppose that $\Omega \subset \mathbb{R}^n$ is a domain, \mathcal{T} is a mesh of Ω with meshwidth $h \in (0, h_0]$, $p \geq m - 1$, and $-m \leq s_1 \leq 0 \leq s_2 \leq m$ (with s_1, s_2 not necessarily integers). Then*

$$\min_{v \in V_{h,p}} \|u - v\|_{\tilde{H}^{s_1}(\Omega)} \leq (C_{n,h_0,m})^{(s_2-s_1)/m} \left(\frac{h}{p+1} \right)^{s_2-s_1} \|u\|_{H^{s_2}(\Omega)}, \quad u \in H^{s_2}(\Omega). \quad (9)$$

The third extends Corollary 2.8 to functions/distributions in negative order spaces.

Corollary 2.9. *Let $n, m \in \mathbb{N}$ and $h_0 > 0$. Suppose that $\Omega \subset \mathbb{R}^n$ is a domain, and that $\{H^s(\Omega)\}_{0 \leq s \leq m}$ is an interpolation scale (equivalently, that $\{\tilde{H}^s(\Omega)\}_{-m \leq s \leq 0}$ is an interpolation scale). Suppose that \mathcal{T} is a mesh of Ω with meshwidth $h \in (0, h_0]$, $p \geq m - 1$, and that $-m \leq s_1 \leq s_2 \leq 0$ (with s_1 and s_2 not necessarily integers). Then*

$$\min_{v \in V_{h,p}} \|u - v\|_{\tilde{H}^{s_1}(\Omega)} \leq \tilde{C}(C_{n,h_0,m})^{(s_2-s_1)/m} \left(\frac{h}{p+1} \right)^{s_2-s_1} \|u\|_{\tilde{H}^{s_2}(\Omega)}, \quad u \in \tilde{H}^{s_2}(\Omega), \quad (10)$$

where \tilde{C} is the norm² of the embedding of $\tilde{H}^{s_2}(\Omega)$ into the interpolation space $K_{1-s_2/s_1,2}(\tilde{H}^{s_1}(\Omega), \tilde{H}^0(\Omega))$.

Remark 2.10. *Regarding the interpolation scale assumption in Corollary 2.9:*

- (i) *We recall from [15, Cor. 4.9] that $\{H^s(\Omega)\}_{0 \leq s \leq m}$ is an interpolation scale (in the sense of [15, Rem. 3.8]) if and only if $\{\tilde{H}^s(\Omega)\}_{-m \leq s \leq 0}$ is an interpolation scale.*
- (ii) *If $\Omega \subset \mathbb{R}^n$ is a Lipschitz hypograph or a Lipschitz domain then $\{H^s(\Omega)\}_{s \in \mathbb{R}}$ and $\{\tilde{H}^s(\Omega)\}_{s \in \mathbb{R}}$ are interpolation scales (see, e.g., [15, Corollaries 4.7 and 4.10]).*
- (iii) *If Ω is an (ϵ, δ) locally uniform domain then $\{H^s(\Omega)\}_{s \geq 0}$ and $\{\tilde{H}^s(\Omega)\}_{s \leq 0}$ are interpolation scales (see, e.g., [15, Corollaries 4.7 and 4.10]). In particular, this includes the Koch snowflake domain pictured in Figure 1.*
- (iv) *It was shown recently in [10, Cor. 3.19] that $\{H^s(\Omega)\}_{s \in \mathbb{R}}$ and $\{\tilde{H}^s(\Omega)\}_{s \in \mathbb{R}}$ are interpolation scales provided Ω is “thick” in the sense of Triebel (see [10, Def. A.3]), $\bar{\Omega} \neq \mathbb{R}^n$ and the Assouad dimension of $\partial\Omega$ (as defined in e.g. [10, Appendix D]) is less than n . In particular, these hypotheses hold if Ω is the interior of an n -attractor (in the terminology of [10]), i.e.*

²This norm is not guaranteed to equal one, since even when $\{H^s(\Omega)\}_{0 \leq s \leq m}$ is an interpolation scale, it is not in general exact - see, e.g., [15, Lem. 4.13].

the attractor of an iterated function system of contracting similarities satisfying the open set condition and having Hausdorff dimension n . This applies to all the examples in [10, Fig. 1] (including the Koch snowflake, the Gosper island and the Levy dragon). These hypotheses also hold for all the “classical snowflakes” in [11, §5.1] (which includes the Koch snowflake).

- (v) By combining [10, Prop. 3.15] with [10, Rem. 3.22] and the Wolff interpolation Theorem [32, Thm 1], one can show that $\{H^s(\Omega)\}_{s \geq 0}$ and $\{\tilde{H}^s(\Omega)\}_{s \leq 0}$ are interpolation scales whenever Ω is an open n -set (in the terminology of [10], called an interior regular domain in [11]), $\bar{\Omega} \neq \mathbb{R}^n$ and the Assouad dimension of $\partial\Omega$ is less than n .

Some further references to the literature on interpolation scale results for $H^s(\Omega)$ and $\tilde{H}^s(\Omega)$, and more general Besov and Triebel-Lizorkin spaces, can be found in [10, Rem. 3.22].

3 Proofs

3.1 Proof of Lemma 2.3

Our proof is similar to that of [13, Lemma 4.31], except that we prove a slightly stronger result by collecting together the contributions of all the mesh elements associated with a given covering element. Under the hypotheses of the lemma, for each $\mathcal{K} \in \mathcal{T}^\#$ let $\pi_{p,m_{\mathcal{K}},\mathcal{K}} : W^{m_{\mathcal{K}}}(\mathcal{K}) \rightarrow \mathcal{P}_p(\mathcal{K})$ denote a known optimal hp -version approximation operator on the simplex/cube \mathcal{K} (e.g. [3],[30], [31, Appendix C]). Since $(U_{\mathcal{K}})|_K = u$ for all $K \in \kappa^{-1}(\mathcal{K})$ and since $K \cap K' = \emptyset$ for $K \neq K'$, for $j = 0, 1, \dots, m_{\mathcal{K}}$ we can apply standard hp approximation results (e.g. [3],[30], [31, Appendix C]) to estimate, with $t_{\mathcal{K}} := \min(m_{\mathcal{K}}, p+1)$,

$$\begin{aligned} \left(\sum_{K \in \kappa^{-1}(\mathcal{K})} \|u - \pi_{p,m_{\mathcal{K}},\mathcal{K}}(U_{\mathcal{K}})\|_{W^j(K)}^2 \right)^{1/2} &= \left(\sum_{K \in \kappa^{-1}(\mathcal{K})} \|U_{\mathcal{K}} - \pi_{p,m_{\mathcal{K}},\mathcal{K}}(U_{\mathcal{K}})\|_{W^j(K)}^2 \right)^{1/2} \\ &\leq \|u - \pi_{p,m_{\mathcal{K}},\mathcal{K}}(U_{\mathcal{K}})\|_{W^j(\mathcal{K})} \\ &\leq C \frac{h_{\mathcal{K}}^{t_{\mathcal{K}}-q}}{(p+1)^{m_{\mathcal{K}}-j}} \sum_{r=t_{\mathcal{K}}}^{m_{\mathcal{K}}} |U_{\mathcal{K}}|_{W^r(\mathbb{R}^n)}, \end{aligned}$$

for some constant $C > 0$ depending only on n , $m_{\mathcal{K}}$, h_0 and the shape regularity of \mathcal{K} .

3.2 Proof of Theorem 2.5

Our approach is a generalisation of the argument used to prove [10, Proposition 3.24], which considered only the case $m = 1$, $p = 0$. The idea is to apply Lemma 2.3, using a covering formed of overlapping cubes.

Under the hypotheses of the theorem, for each $\mathbf{z} \in \mathbb{Z}^n$ let $Q_{\mathbf{z}}$ be the cube of side length h defined by $Q_{\mathbf{z}} := [0, h]^n + h\mathbf{z}$, and let $Q'_{\mathbf{z}} := [-h, 2h]^n + h\mathbf{z}$, the cube of side length $3h$ formed by taking the union of $Q_{\mathbf{z}}$ with all its neighbouring cubes, which satisfies $\text{diam}(Q'_{\mathbf{z}}) = 3n^{1/2}h$. For each $K \in \mathcal{T}$, since $\text{diam}(K) \leq h$, K intersects the cube $Q_{\mathbf{z}}$ for at least one and at most 2^n different $\mathbf{z} \in \mathbb{Z}^n$. And for each such \mathbf{z} we have $K \subset Q'_{\mathbf{z}}$. Hence $\mathcal{T}^\# := \{Q'_{\mathbf{z}}\}_{\mathbf{z} \in \mathbb{Z}^n}$ is a covering of \mathcal{T} . Let $\kappa : \mathcal{T} \rightarrow \mathcal{T}^\#$ be any covering choice function, and let $\mathbf{z}_{\kappa} : \mathcal{T} \rightarrow \mathbb{Z}^n$ be defined by $\kappa(K) = Q'_{\mathbf{z}_{\kappa}(K)}$ for $K \in \mathcal{T}$.

For each $K \in \mathcal{T}$ define $v|_K$ to be the restriction to K of $\pi_{p,m,Q'_{\mathbf{z}_K(K)}}(U|_{Q'_{\mathbf{z}_K(K)}})$, where $\pi_{p,m,Q'_{\mathbf{z}_K(K)}}$ is the operator from Lemma 2.3. This defines v almost everywhere on Ω , because of the fact that $|\Omega \setminus \bigcup_{K \in \mathcal{T}} K| = 0$. Furthermore, by Lemma 2.3 we have that

$$\sum_{K \in \mathcal{T}} \|u - v\|_{W^j(K)}^2 = \sum_{\mathbf{z} \in \mathbb{Z}^n} \sum_{K \in \mathbf{z}_K^{-1}(\mathbf{z})} \|u - v\|_{W^j(K)}^2 \leq \sum_{\mathbf{z} \in \mathbb{Z}^n} \left(C_{n,h_0,m,Q'_{\mathbf{z}}} \frac{(3n^{1/2}h)^{t-j}}{(p+1)^{m-j}} \right)^2 \sum_{r=t}^m |U|_{W^r(Q'_{\mathbf{z}})}^2.$$

Since $C_{n,h_0,m,Q'_{\mathbf{z}}}$ depends on \mathbf{z} only through the shape regularity of $Q'_{\mathbf{z}}$, which is independent of \mathbf{z} , we can replace $C_{n,h_0,m,Q'_{\mathbf{z}}}$ by a constant $\tilde{C}_{n,h_0,m}$ depending only on n, h_0 and m . Then

$$\begin{aligned} \sum_{K \in \mathcal{T}} \|u - v\|_{W^j(K)}^2 &\leq \left(\tilde{C}_{n,h_0,m} \frac{(3n^{1/2}h)^{t-j}}{(p+1)^{m-j}} \right)^2 \sum_{\mathbf{z} \in \mathbb{Z}^n} \sum_{r=t}^m |U|_{W^r(Q'_{\mathbf{z}})}^2 \\ &= 3^n \left(\tilde{C}_{n,h_0,m} \frac{(3n^{1/2}h)^{t-j}}{(p+1)^{m-j}} \right)^2 \sum_{\mathbf{z} \in \mathbb{Z}^n} \sum_{r=t}^m |U|_{W^r(Q_{\mathbf{z}})}^2 \\ &= 3^n \left(\tilde{C}_{n,h_0,m} \frac{(3n^{1/2}h)^{t-j}}{(p+1)^{m-j}} \right)^2 \sum_{r=t}^m |U|_{W^r(\mathbb{R}^n)}^2, \end{aligned}$$

where we used the fact that for each \mathbf{z} the contribution $|U|_{W^r(Q_{\mathbf{z}})}^2$ gets counted 3^n times in the sum $\sum_{\mathbf{z} \in \mathbb{Z}^n} |U|_{W^r(Q'_{\mathbf{z}})}^2$. From this, (6) follows by taking square roots, with $C_{n,h_0,m} = 3^{n/2} (3n^{1/2})^m \tilde{C}_{n,h_0,m}$. In the case where Ω is bounded, the fact that $v \in L^2(\Omega)$, and hence that $v \in V_{h,p}$, is immediate from the definition of v . In the case where Ω is unbounded the fact that $\|v\|_{L^2(\Omega)} < \infty$ follows from (6) with $j = 0$, the fact that $u \in L^2(\Omega)$, and the triangle inequality $\|v\|_{L^2(\Omega)} \leq \|u - v\|_{L^2(\Omega)} + \|u\|_{L^2(\Omega)}$.

3.3 Proof of Corollary 2.7

Let $\Pi : L^2(\Omega) \rightarrow V_{h,p}$ denote the L^2 -orthogonal projection onto $V_{h,p}$, which satisfies $\|u - \Pi u\|_{L^2(\Omega)} = \min_{v \in V_{h,p}} \|u - v\|_{L^2(\Omega)}$ for $u \in L^2(\Omega)$. For $u \in H^m(\Omega)$, taking $j = 0$ in (6) and bounding $|U|_{W^m(\mathbb{R}^n)} \leq \|U\|_{H^m(\mathbb{R}^n)}$ gives

$$\|u - \Pi u\|_{L^2(\Omega)} \leq C_{n,h_0,m} \left(\frac{h}{p+1} \right)^m \|U\|_{H^m(\mathbb{R}^n)}. \quad (11)$$

Since (11) holds for every $U \in H^m(\mathbb{R}^n)$ with $U|_{\Omega} = u$, by the definition of $\|\cdot\|_{H^m(\Omega)}$ we have

$$\|u - \Pi u\|_{L^2(\Omega)} \leq C_{n,h_0,m} \left(\frac{h}{p+1} \right)^m \|u\|_{H^m(\Omega)}. \quad (12)$$

We note also for $u \in L^2(\Omega) = H^0(\Omega)$ the trivial estimate

$$\|u - \Pi u\|_{L^2(\Omega)} \leq \|u\|_{L^2(\Omega)}. \quad (13)$$

We then deduce (8) by interpolation applied to the operator $I - \Pi$, using (12) and (13). We adapt the argument of [17, Eqn (46)], where only the special case $m = 1$ was considered. Precisely, given $s \in [0, m]$, let $\theta := s/m \in [0, 1]$ and let $\|\cdot\|_{\theta}$ denote the norm on the K -method

interpolation space $K_{\theta,2}((L^2(\Omega), H^m(\Omega)))$, with the normalisation defined in [15, (7,8)]. Then, with $\sigma := (C_{n,h_0,m})^{1/m}h/(p+1)$,

$$\|u - \Pi u\|_{L^2(\Omega)} \leq \sigma^{\theta m} \|u\|_{\theta} \leq \sigma^{\theta m} \|u\|_{H^{\theta m}(\Omega)} = \sigma^s \|u\|_{H^s(\Omega)}, \quad (14)$$

where the first inequality follows by interpolation since $K_{\theta,2}((L^2(\Omega), L^2(\Omega))) = L^2(\Omega)$ with equality of norms given the normalisation [15, (7,8)] (see [15, Lem. 2.1(iii), Thm. 2.2(i)]), the second inequality holds since, again given this normalisation, the embedding of $H^s(\Omega) = H^{\theta m}(\Omega)$ into $K_{\theta,2}((L^2(\Omega), H^m(\Omega)))$ has norm ≤ 1 [15, Lem. 4.2],³ and the final equality holds because $\theta m = s$.

3.4 Proof of Corollary 2.8

We adopt a duality argument, similar to that used in the proofs of [29, Thm. 4.1.33] and [17, Lemma A.1]. Let $-m \leq s_1 \leq 0 \leq s_2 \leq m$ and let $u \in H^{s_2}(\Omega) \subset L^2(\Omega)$. By the duality between $\tilde{H}^{-s}(\Omega)$ and $\tilde{H}^s(\Omega)$ (see, e.g., [16, Thm 3.3 & §3.1.3]), the L^2 -orthogonality of the projection Π (which we define as in the proof of Corollary 2.7 to be orthogonal projection onto $V_{h,p}$ in $L^2(\Omega)$), the Cauchy-Schwarz inequality, and two applications of (8), we have that

$$\begin{aligned} \min_{v \in V_{h,p}} \|u - v\|_{\tilde{H}^{s_1}(\Omega)} &\leq \|u - \Pi u\|_{\tilde{H}^{s_1}(\Omega)} = \sup_{0 \neq v \in H^{-s_1}(\Omega)} \frac{|(u - \Pi u, v)_{L^2(\Omega)}|}{\|v\|_{H^{-s_1}(\Omega)}} \\ &= \sup_{0 \neq v \in H^{-s_1}(\Omega)} \frac{|(u - \Pi u, v - \Pi v)_{L^2(\Omega)}|}{\|v\|_{H^{-s_1}(\Omega)}} \\ &\leq \sup_{0 \neq v \in H^{-s_1}(\Omega)} \frac{\|u - \Pi u\|_{L^2(\Omega)} \|v - \Pi v\|_{L^2(\Omega)}}{\|v\|_{H^{-s_1}(\Omega)}} \\ &\leq (C_{n,h_0,m})^{(s_2-s_1)/m} \left(\frac{h}{p+1} \right)^{s_2-s_1} \|u\|_{H^{s_2}(\Omega)}. \end{aligned}$$

3.5 Proof of Corollary 2.9

Let $m \leq s_1 \leq 0$ and let $\Pi^{s_1} : \tilde{H}^{s_1}(\Omega) \rightarrow V_{h,p}$ denote orthogonal projection onto $\tilde{V}_{h,p}$ in $\tilde{H}^{s_1}(\Omega)$. Applying (9) with $s_2 = 0$ gives

$$\|u - \Pi^{s_1} u\|_{\tilde{H}^{s_1}(\Omega)} \leq (C_{n,h_0,m})^{-s_1/m} \left(\frac{h}{p+1} \right)^{-s_1} \|u\|_{\tilde{H}^0(\Omega)}, \quad u \in \tilde{H}^0(\Omega), \quad (15)$$

and we also have the trivial estimate

$$\|u - \Pi^{s_1} u\|_{\tilde{H}^{s_1}(\Omega)} \leq \|u\|_{\tilde{H}^{s_1}(\Omega)}, \quad u \in \tilde{H}^{s_1}(\Omega). \quad (16)$$

Then (10) follows for $-m \leq s_1 \leq s_2 \leq 0$ by applying interpolation to $I - \Pi^{s_1}$, using (15) and (16).

3.6 Proof of Corollary 2.6

Just apply (6) with $U = E_{\Omega}^m u$ and bound $\|E_{\Omega}^m w\|_{W^m(\mathbb{R}^n)} \leq C_{m,\Omega} \|w\|_{W^m(\Omega)}$.

³Note that in general it does not hold that $H^{\theta m}(\Omega) = K_{\theta,2}((L^2(\Omega), H^m(\Omega)))$ - see, e.g., [15, Lem. 4.8].

4 Conclusion

We have presented best approximation error estimates for discontinuous piecewise polynomial approximation on non-Lipschitz meshes of non-Lipschitz domains. Many of our results hold for arbitrary meshes of arbitrary domains, while some require a minimal smoothness assumption on Ω , satisfied, for instance, if Ω is an (ϵ, δ) locally uniform domain. Our results are therefore valid for a large class of domains with fractal boundary, including the Koch snowflake domain.

Acknowledgements

The author thanks Simon Chandler-Wilde, Emmanuil Georgoulis, Andrea Moiola and Iain Smears for helpful discussions in relation to this work. Part of this work was carried out while the author was in residence at Institut Mittag-Leffler in Djursholm, Sweden in autumn 2025, supported by the Swedish Research Council under grant no. 2021-06594.

References

- [1] Y. ACHDOU, C. SABOT, AND N. TCHOU, *Transparent boundary conditions for the Helmholtz equation in some ramified domains with a fractal boundary*, J. Comput. Phys., 220 (2007), pp. 712–739.
- [2] K. E. ATKINSON, *The Numerical Solution of Integral Equations of the Second Kind*, Cambridge University Press, 1997.
- [3] L. BABUŠKA AND M. SURI, *The optimal convergence rate of the p -version of the finite element method*, SIAM J. Numer. Anal., 24 (1987), pp. 750–776.
- [4] P. BAGNERINI, A. BUFFA, AND E. VACCA, *Finite elements for a prefractal transmission problem*, C. R. Math., 342 (2006), pp. 211–214.
- [5] P. BAGNERINI, A. BUFFA, AND E. VACCA, *Mesh generation and numerical analysis of a Galerkin method for highly conductive prefractal layers*, Appl. Numer. Math., 65 (2013), pp. 63–78.
- [6] J. BANNISTER, A. GIBBS, AND D. P. HEWETT, *Acoustic scattering by fractal inhomogeneities via geometry-conforming Galerkin methods for the Lippmann-Schwinger equation*. In preparation.
- [7] S. C. BRENNER AND L. R. SCOTT, *The mathematical theory of finite element methods*, Springer, 2008.
- [8] A. M. CAETANO, S. N. CHANDLER-WILDE, X. CLAEYS, A. GIBBS, D. P. HEWETT, AND A. MOIOLA, *Integral equation methods for acoustic scattering by fractals*, Proc. Roy. Soc. A, 481 (2025), p. 20230650.
- [9] A. M. CAETANO, S. N. CHANDLER-WILDE, A. GIBBS, D. P. HEWETT, AND A. MOIOLA, *A Hausdorff-measure boundary element method for acoustic scattering by fractal screens*, Numer. Math., 156 (2024), p. 463–532.

- [10] A. M. CAETANO, S. N. CHANDLER-WILDE, AND D. HEWETT, *Properties of IFS attractors with non-empty interiors and related rough sets, and associated function spaces and scattering problems*. arxiv.org/abs/2511.15213.
- [11] A. M. CAETANO, D. P. HEWETT, AND A. MOIOLA, *Density results for Sobolev, Besov and Triebel-Lizorkin spaces on rough sets*, J. Funct. Anal., 281 (2021), p. 109019.
- [12] A. P. CALDERÓN, *Lebesgue spaces of differentiable functions and distributions*, Proc. Symp. Pure Math., 4 (1961), pp. 33–49.
- [13] A. CANGIANI, Z. DONG, AND E. GEORGIOULIS, *hp-version discontinuous Galerkin methods on essentially arbitrarily-shaped elements*, Math. Comp., 91 (2022), pp. 1–35.
- [14] R. CAPITANELLI AND M. A. VIVALDI, *Weighted estimates on fractal domains*, Mathematika, 61 (2015), pp. 370–384.
- [15] S. N. CHANDLER-WILDE, D. P. HEWETT, AND A. MOIOLA, *Interpolation of Hilbert and Sobolev spaces: quantitative estimates and counterexamples*, Mathematika, 61 (2015), pp. 414–443.
- [16] ———, *Sobolev spaces on non-Lipschitz subsets of \mathbb{R}^n with application to boundary integral equations on fractal screens*, Integr. Equat. Operat. Th., 87 (2017), pp. 179–224.
- [17] S. N. CHANDLER-WILDE, D. P. HEWETT, A. MOIOLA, AND J. BESSON, *Boundary element methods for acoustic scattering by fractal screens*, Numer. Math., 147 (2021), pp. 785–837.
- [18] P. G. CIARLET, *The Finite Element Method for Elliptic Problems*, North-Holland, 1978.
- [19] S. CREO, M. R. LANCIA, AND P. VERNOLE, *Transmission problems for the fractional p -Laplacian across fractal interfaces*, Discrete Contin. Dyn. Syst. - Ser. S, 15 (2022).
- [20] A. GIBBS, D. P. HEWETT, AND B. MAJOR, *Numerical evaluation of singular integrals on non-disjoint self-similar fractal sets*, Numer. Alg., 97 (2024), pp. 311–343.
- [21] A. GIBBS, D. P. HEWETT, AND A. MOIOLA, *Numerical evaluation of singular integrals on fractal sets*, Numer. Alg., 92 (2023), pp. 2071–2124.
- [22] S. GOMEZ, A. MOIOLA, AND D. P. HEWETT, *A discontinuous Galerkin method with fractal elements*. In preparation.
- [23] M. HINZ, A. ROZANOVA-PIERRAT, AND A. TEPLYAEV, *Non-Lipschitz uniform domain shape optimization in linear acoustics*, SIAM J. Control Optim., 59 (2021), pp. 1007–1032.
- [24] P. W. JONES, *Quasiconformal mappings and extendability of functions in Sobolev spaces*, Acta Math., 147 (1981), pp. 71–88.
- [25] A. JONSSON AND H. WALLIN, *Function Spaces on Subsets of \mathbb{R}^n* , Math. Rep., 2 (1984).
- [26] R. KRESS, *Linear Integral Equations*, Springer-Verlag, New York, 2nd ed., 1999.
- [27] L. G. ROGERS, *Degree-independent Sobolev extension on locally uniform domains*, J. Funct. Anal., 235 (2006), pp. 619–665.
- [28] V. S. RYCHKOV, *On restrictions and extensions of the Besov and Triebel-Lizorkin spaces with respect to Lipschitz domains*, J. London Math. Soc., 60 (1999), pp. 237–257.

- [29] S. A. SAUTER AND C. SCHWAB, *Boundary Element Methods*, Springer, 2011.
- [30] C. SCHWAB, *p- and hp-Finite Element Methods*, Clarendon Press, Oxford, 1998.
- [31] I. SMEARS, *Discontinuous Galerkin finite element approximation of Hamilton–Jacobi–Bellman equations with Cordes coefficients*, PhD thesis, University of Oxford, 2015.
- [32] T. H. WOLFF, *A note on interpolation spaces*, in Harmonic Analysis: Proceedings of a conference held at the University of Minnesota, Minneapolis, April 20–30, 1981, Springer, 1982, pp. 199–204.