

Supersymmetric zeta functions and determinants

Yu Nakayama¹ and Tadashi Okazaki²

*Center for Gravitational Physics and Quantum Information, Yukawa Institute for
Theoretical Physics, Kyoto University,
Kitashirakawa Oiwakecho, Sakyo-ku, Kyoto 606-8502, Japan
Shing-Tung Yau Center and School of Physics of Southeast University,
Yifu Architecture Building, No.2 Sipailou, Xuanwu district,
Nanjing, Jiangsu, 210096, China*

Abstract

We define supersymmetric zeta functions and supersymmetric determinants, which can reveal spectral properties complementary to those captured by the supersymmetric indices. They play a crucial role in analyzing the Cardy-like behaviors of the supersymmetric indices and the supersymmetric Casimir energies associated with the supersymmetric partition functions. We investigate a variety of examples of the supersymmetric zeta functions and determinants for two-, four-, and six-dimensional supersymmetric field theories.

¹yu.nakayama@yukawa.kyoto-u.ac.jp

²tokazaki@seu.edu.cn

Contents

1	Introduction and conclusions	4
1.1	Structure	5
1.2	Future works	6
2	Supersymmetric zeta functions	7
2.1	Definition	7
2.2	Mellin transform	9
2.3	Multiple supersymmetric zeta functions	12
2.4	Degeneracy with exponential growth	13
2.4.1	Cardy-like limit	13
2.4.2	Effective Field Theory interpretation	16
2.4.3	Asymptotic degeneracy	18
2.4.4	Supersymmetric Casimir energies	23
3	Supersymmetric determinants	25
3.1	Definition	25
3.2	Laplace transformation	25
3.3	Vacuum exponents	26
3.4	Functional equations	27
4	2d supersymmetric field theories	29
4.1	Zeta functions for 2d $\mathcal{N} = (0, 2)$ elliptic genera	29
4.1.1	2d $\mathcal{N} = (0, 2)$ chiral multiplet	31
4.1.2	2d $\mathcal{N} = (0, 2)$ Fermi multiplet	33
4.1.3	2d $\mathcal{N} = (0, 2)$ vector multiplet	35
4.1.4	2d $\mathcal{N} = (0, 2)$ SQED with chirals $(\Phi, \bar{\Phi}_a)$ and Fermis (Ψ_α, Γ_a)	36
4.1.5	2d $\mathcal{N} = (0, 2)$ $U(N_c)$ with chirals $(\Phi_s, \bar{\Phi}_a)$ and Fermis $(\Psi_\alpha, \Omega, \Gamma_{s,a})$	38
4.1.6	2d $\mathcal{N} = (0, 2)$ $SU(2)$ with 4 chirals Φ_α	39
4.1.7	2d $\mathcal{N} = (0, 2)$ $SU(N)$ with $2N$ chirals $(\Phi_\alpha, \bar{\Phi}_\beta)$	41
4.1.8	2d $\mathcal{N} = (0, 2)$ $USp(2N)$ with $2N + 2$ chirals Φ_α	43
4.1.9	2d $\mathcal{N} = (0, 2)$ $USp(2N)$ with chirals (Φ_α, A) and Fermis Γ_a	46
4.2	Zeta functions for 2d $\mathcal{N} = (2, 2)$ elliptic genera	48
4.2.1	2d $\mathcal{N} = (2, 2)$ chiral multiplet	49
4.2.2	2d $\mathcal{N} = (2, 2)$ vector multiplet	51
4.2.3	2d $\mathcal{N} = (2, 2)$ pure SYM theories	52
4.2.4	2d $\mathcal{N} = (2, 2)$ minimal models	54
4.2.5	2d $\mathcal{N} = (2, 2)$ Kazama-Suzuki models	58

4.2.6	2d $\mathcal{N} = (2, 2)$ \mathbb{CP}^{N-1} models	60
5	4d supersymmetric field theories	61
5.1	Zeta functions for 4d supersymmetric indices	61
5.1.1	4d $\mathcal{N} = 1$ chiral multiplet	63
5.1.2	4d $\mathcal{N} = 1$ vector multiplet	66
5.1.3	4d $\mathcal{N} = 1$ $SU(2)$ with 6 chirals $(\Phi_\alpha, \bar{\Phi}_\alpha)$	67
5.1.4	4d $\mathcal{N} = 1$ $SU(N)$ with $2(N+1)$ chirals $(\Phi_\alpha, \bar{\Phi}_\alpha)$	70
5.1.5	4d $\mathcal{N} = 1$ $USp(2N)$ with $2N+4$ chirals Φ_α	73
5.1.6	4d $\mathcal{N} = 2$ hypermultiplet	75
5.1.7	4d $\mathcal{N} = 2$ vector multiplet	76
5.1.8	4d $\mathcal{N} = 4$ vector multiplet	78
5.2	Zeta functions for Schur and Macdonald indices	79
5.2.1	4d $\mathcal{N} = 2$ hypermultiplet	82
5.2.2	4d $\mathcal{N} = 2$ vector multiplet	83
5.2.3	4d $\mathcal{N} = 4$ vector multiplet	85
5.2.4	4d $\mathcal{N} = 2$ pure SYM theories	86
5.2.5	$J^{h^\vee}[k]$ AD theories	90
5.2.6	The generalized (G_1, G_2) AD theories	92
5.2.7	$(J^{h^\vee}[k], F)$ AD matters	93
5.2.8	$(A_{N-1}^N[k], [N-1, 1])$ AD matter	95
5.2.9	rank 2 AD theory $AD(C_2)$	96
6	6d supersymmetric field theories	97
6.1	Zeta functions for supersymmetric indices	97
6.1.1	6d $\mathcal{N} = (1, 0)$ hypermultiplet	99
6.1.2	6d $\mathcal{N} = (1, 0)$ tensor multiplet	101
6.1.3	6d $\mathcal{N} = (1, 0)$ vector multiplet	103
6.1.4	6d $\mathcal{N} = (2, 0)$ tensor multiplet	104
6.2	Zeta functions for unrefined indices	106
6.2.1	6d $\mathcal{N} = (2, 0)$ theories	106
A	Notations	107
B	Residue formulae	108
C	Special zeta values	108
C.1	Riemann zeta function	108
C.2	Hurwitz zeta function	109

1 Introduction and conclusions

In any theory of quantum systems, the most fundamental question is to understand the spectrum of operators (e.g., the Hamiltonian), specifically their eigenvalues and distributions. From them, we can construct various spectral functions. In physics, the most familiar spectral functions are the partition function and, when the theory is equipped with nilpotent fermionic symmetries, the index. The former is of fundamental interest because it encodes the thermodynamic properties of the system, and the latter, if available, is a highly useful and, more importantly, computable quantity that can be regarded as an invariant of the system under small perturbations and can be used to analyze dualities. The main advantage of studying the partition function or index is that it can be reformulated in terms of the path integral; this path integral expression, when compared with the spectral definition, sometimes yields remarkable formulae for certain geometric and topological invariants, such as the Atiyah-Singer index theorem [1, 2] and the Jones polynomial [3].

In mathematics, other spectral functions are also of interest, such as spectral zeta functions and spectral determinants. One advantage these alternative tools offer over partition functions is their potential for larger analyticity domains. Analytic continuation of the partition function in the "temperature" is highly non-trivial. For instance, if the theory has an infinite tower of excited states, the partition function takes the form of an infinite sum, which converges on the half-plane $\text{Re}\beta > 0$ and may be singular at the high temperature limit $\beta \rightarrow 0$. On the other hand, the zeta function can have a meromorphic extension to the whole complex plane with only zeros and simple poles when the spectrum is discrete. The determinant takes the form of an infinite product with only isolated zeros and poles corresponding to the locations of the eigenvalues. Such structures can make complex analysis much easier. These different types of spectral functions can be connected to one another via a certain integral transform. Such an integral transform can map the original function space into another one, thereby providing a powerful method for finding analytical solutions to problems that might be difficult or impossible to solve by other methods. The resulting form is often particularly suitable for the saddle point approximation in the presence of large parameters. In particular, since the zeta functions and the partition functions are related via the Mellin transform, the structure of the zeros and poles of the zeta functions can encode the asymptotic behavior of the partition functions.

The asymptotic behavior of the partition functions is typically associated with

the geometry underlying the spectrum. Thus, the zeta functions encode such geometric information in a naturally regularized manner. In practice, when computing the partition functions, the quantum field theories and their path integrals are full of divergences. Nevertheless, the zeta functions provide a powerful tool for obtaining finite numbers while revealing the underlying geometry. This regularization, for instance, will be useful for understanding emergent geometries in large N theories via holography, such as the AdS/CFT correspondence [4]. The essential philosophy here is analogous to the classic question “Can one hear the shape of a drum?”, posed by Kac [5]. In fact, Martelli, Sparks and Yau [6, 7, 8] demonstrated that the volume and other geometric properties of the Sasaki-Einstein manifolds can be read off from certain spectral data of the holographically dual 4d $\mathcal{N} = 1$ superconformal field theories.

The main motivation of this work is to investigate new spectral functions for physical theories. We begin with the supersymmetric indices and introduce new types of spectral functions for the supersymmetric theories, *supersymmetric zeta functions* and *supersymmetric determinants*. Supersymmetric indices [9, 10, 11] are well-established spectral functions that encode the BPS spectra of the supersymmetric theories. Defined as a trace over the BPS states, they are used to count the BPS local operators in the theory by reading off their expansion coefficients.

In our initial exploration of these spectral functions for supersymmetric theories, we focus on the supersymmetric field theories in $d = 2, 4$ and 6 dimensions. For such even-dimensional field theories, there exist local anomalies encoded by the anomaly polynomial $(d+2)$ -form \mathcal{A}_{d+2} that is a polynomial in the Chern and Pontryagin classes of gauge and gravity fields. It is known that the Cardy-like limits (high temperature limits) of the supersymmetric indices and the supersymmetric Casimir energy (zero temperature limits) of the supersymmetric partition functions on $S^1 \times S^{d-1}$ have a universal relationship with the anomaly coefficients for the superconformal field theories. We find that these properties (e.g., the Cardy-like limits) can be understood as consequences of universal aspects of the supersymmetric zeta functions we introduce. We propose that residues and special values of the supersymmetric zeta functions can be universally expressed in terms of the anomaly coefficients or central charges for the superconformal field theories. Conversely, the anomaly coefficients and the central charges for the superconformal field theories can be calculated from the supersymmetric zeta functions.

1.1 Structure

In section 2 we define the supersymmetric zeta functions. They are linked to the supersymmetric indices via the Mellin transform. We present formulae for the Cardy-

like limit of the supersymmetric indices and the asymptotic degeneracy in terms of the supersymmetric zeta functions. Also, we propose that the special supersymmetric zeta values can encode the supersymmetric Casimir energies. In section 3 we define the supersymmetric determinants as the third variant of the spectral functions. The vacuum exponents, which appear as their special values in the asymptotic formulae, can provide criteria for identifying a supersymmetry-breaking vacuum or an unstable vacuum. In section 4, section 5 and section 6, we provide various examples of the supersymmetric zeta functions and supersymmetric determinants for 2d, 4d and 6d supersymmetric field theories. We present some conventions of the q -shifted factorials in Appendix A. Several useful formulae for the residues and the special values of the zeta functions are summarized in Appendices B and C.

1.2 Future works

- It would be interesting to explore the universal structure of the residues and special values of the supersymmetric zeta functions for other settings of supersymmetric field theories, including the theories in odd dimensions, the configurations with defect operators and the deformations such as the topological twisting and the Higgsing. We hope to report on our results in future work.
- For the supersymmetric field theories with gravity duals the large N limits of the supersymmetric indices capture the Kaluza-Klein modes of the supergravity. The associated supersymmetric zeta functions can be used to analyze the asymptotic behaviors, which typically grow much faster than those at finite N . We leave the detailed analysis of large N results to future work.
- The supersymmetric zeta functions and supersymmetric determinants also contain several other theory-dependent data, such as zeros and their behavior at large values, which are also expected to encode certain physical information. At present, these aspects are less understood, and we hope to report on them in future work.
- The supersymmetric indices may satisfy interesting relations under certain transformations, e.g. the elliptic and modular transformations [12, 13, 14], the T -reflection [15, 16]. From the perspective of the supersymmetric determinants, such properties are translated into certain functional equations. In turn, the supersymmetric zeta functions will obey certain algebraic relations. These relations will provide us with interesting avenues for further study.

- In mathematics, the arithmetic nature of the special zeta values is often discussed. The values of the Riemann zeta function at any positive even integer are transcendental due to the well-known fact that π is transcendental [17]. Although a proof is still lacking, it is conjectured that the zeta values at odd integers are algebraically independent over \mathbb{Q} and have higher transcendentality (see e.g. [18, 19, 20] for the irrationality). In physics, it is conjectured [21, 22] that the central charges such as a and c take algebraic numbers or rational numbers, depending on the number of supersymmetries. It would be interesting to further understand the arithmetic nature of our supersymmetric zeta values that encode the central charges and connections to the “zeta world” of number theory.
- While the main focus of our paper is the asymptotic behaviors of the supersymmetric indices on the first sheet, it would be interesting to study the so-called second sheet (or higher sheet) indices by studying their multi-valued structures. Such indices turn out to play an important role in accounting for the black hole microstates, see e.g. [23, 24, 25, 26, 27, 28, 29, 30, 31, 32]. We leave the analysis of the associated supersymmetric zeta functions and determinants as an interesting direction for future work.

2 Supersymmetric zeta functions

2.1 Definition

We begin with the supersymmetric index of a supersymmetric theory. It can be defined by

$$\mathbb{I}(q) = \text{Tr}_{\mathcal{H}}(-1)^F q^{\Delta}, \quad (2.1)$$

where the trace is taken over the cohomology \mathcal{H} of the chosen supercharges in the theory. F is the fermion number operator, and Δ is a certain appropriate combination of the Cartan generators of global symmetries in the theory that commute with the chosen supercharges. When one expands the index

$$\mathbb{I}(q) = \sum_n d(n) q^n, \quad (2.2)$$

the expansion coefficients $d(n)$ can be identified with the *degeneracy* of the BPS operators with charge (or “energy”) n as they count the BPS bosonic and fermionic operators or states with positive and negative signs, respectively. When we consider a supersymmetric quantum field theory with a UV Lagrangian, the supersymmetric

index can be computed as a certain supersymmetric partition function on the product space of a circle and a sphere $S^1 \times S^{d-1}$ (or more generally, on S^{d-1} bundle over S^1 when angular momentum chemical potential is introduced) up to the supersymmetric Casimir energy factor.

Here we assume that the degeneracy $d(n)$ exhibits the polynomial growth $d(n) = \mathcal{O}(n^\alpha)$ (see section 2.4 for the exponential growth). Let us write $q = e^{-\beta}$ and consider the asymptotic behavior of the number of states for the large charge n . By using the continuum approximation for the sum, one can view the expression (2.2) as the Laplace transform of the degeneracy $d(n)$. Thus, we can obtain the asymptotic behavior of the degeneracy via the inverse Laplace transform of the supersymmetric index. The inverse Laplace transform can be easily performed by taking the Laurent series expansion of the supersymmetric index of the form

$$\mathbb{I}(\beta) = \sum_k C_k \beta^k. \quad (2.3)$$

The asymptotic behavior of the degeneracy with polynomial growth $d(n) = \mathcal{O}(n^\alpha)$ arises from the negative powers of β . Performing the inverse Laplace transform, one finds

$$d(n) \sim \sum_{k=1}^{\alpha+1} \frac{C_{-k}}{(k-1)!} n^{k-1}. \quad (2.4)$$

Also, the asymptotic number of the BPS operators with charge less than or equal to x is estimated as

$$\begin{aligned} \Omega(x) &= \sum_{n \leq x} d(n) \\ &\sim \int_0^x dn \, d(n) = \sum_{k=1}^{\alpha+1} \frac{C_{-k}}{k!} x^k. \end{aligned} \quad (2.5)$$

Sometimes, $\Omega(x)$ may be better-behaved than $d(n)$ as a function of x when $d(n)$ exhibits an oscillatory behavior.

We now define our central quantities. We introduce an arithmetic generating function by defining the *supersymmetric zeta function* by

$$\begin{aligned} \mathbb{Z}(s, z) &= \text{Tr}_{\mathcal{H}} (-1)^F \frac{1}{(\Delta + z)^s} \\ &= \sum_n \frac{d(n)}{(n + z)^s}. \end{aligned} \quad (2.6)$$

It can be regarded as the generalized Dirichlet series associated with the BPS degeneracy $d(n)$ of the supersymmetric theory. For the series to be well-defined as a

function of the complex variable s , we need to consider its convergence properties. In general, if the degeneracy $d(n)$ exhibits polynomial growth $d(n) = \mathcal{O}(n^\alpha)$, the Dirichlet series (2.6) converges absolutely to an analytic function on the half plane $\text{Re}(s) > \alpha + 1$ [33]. In addition, the supersymmetric zeta functions can be analytically continued to meromorphic functions on the whole complex s -plane, which are holomorphic everywhere except for simple poles.

2.2 Mellin transform

The supersymmetric zeta function can be obtained from the supersymmetric index by applying the following integral transform:

$$\begin{aligned}
\mathbb{Z}(s, z) &= \frac{1}{\Gamma(s)} \int_1^\infty dq \, \mathbb{I}(q^{-1}) q^{-z-1} (\log q)^{s-1} \\
&= \frac{1}{\Gamma(s)} \int_1^\infty dq \sum_n d(n) q^{-(n+z)-1} (\log q)^{s-1} \\
&= \sum_n d(n) \frac{1}{\Gamma(s)} (n+z)^{-s} \int_0^\infty dt \, t^{s-1} e^{-t} \\
&= \sum_n \frac{d(n)}{(n+z)^s}.
\end{aligned} \tag{2.7}$$

Here, in the third line, we have interchanged the order of summation and integration (assuming this is valid) and made the substitution $q = e^{\frac{t}{n+z}}$. In the fourth line, we have used the Euler integral of the second kind

$$\Gamma(w) = \int_0^\infty dt \, t^{w-1} e^{-t}. \tag{2.8}$$

Alternatively, we have

$$\mathbb{Z}(s, z) = \frac{1}{\Gamma(s)} \int_0^\infty d\beta \, \mathbb{I}(\beta) e^{-\beta z} \beta^{s-1}. \tag{2.9}$$

This is the Mellin transform of the product $\mathbb{I}(\beta) e^{-\beta z}$. (Here z can be viewed as an additional contribution to the charge of the “vacuum state”.) By splitting the integral (2.9) at $\beta = 1$, we can use the Laurent expansion to perform the integration for small β , whereas the integration over the large β region gives an entire function of s . It leads to the relation

$$C_{-k} = (k-1)! \text{Res}_{s=k} \mathbb{Z}(s, 0) \quad \text{for } k > 0, \tag{2.10}$$

where $\text{Res}_{s=k} \mathbb{Z}(s, 0)$ is the residue at a simple pole $s = k$ of the supersymmetric zeta function with $z = 0$. By plugging this into (2.4) and (2.5), we get

$$d(n) \sim \sum_{k=1}^{\alpha+1} \text{Res}_{s=k} \mathbb{Z}(s, 0) n^{k-1} \tag{2.11}$$

and

$$\Omega(x) \sim \sum_{k=1}^{\alpha+1} \frac{\text{Res}_{s=k} \mathbb{Z}(s, 0)}{k} x^k. \quad (2.12)$$

The formula (2.12) is a result of the Wiener-Ikehara Tauberian theorem [34, 35] for the arithmetic generating function.

If the supersymmetric zeta function (2.6) can be analytically continued to non-positive integers, its special values give rise to the regularized “weighted moments” of the charge Δ

$$\begin{aligned} \mathbb{Z}(-k, z) &= \text{Tr}_{\mathcal{H}}(-1)^F (\Delta + z)^k \\ &= \sum_n (n + z)^k d(n). \end{aligned} \quad (2.13)$$

For superconformal field theories, the charge Δ can be identified with the (R-symmetry twisted) Hamiltonian obtained from the radial quantization so that the special supersymmetric zeta values can be viewed as the regularized energy moments. It follows that these values appear as the non-negative power coefficients in the Laurent series (2.3) for the supersymmetric index in terms of β

$$C_k = \frac{(-1)^k \mathbb{Z}(-k, 0)}{k!} \quad \text{for } k \geq 0. \quad (2.14)$$

As a simple example, let us consider the half-BPS index for 4d $\mathcal{N} = 4$ $U(N)$ super Yang-Mills (SYM) theory

$$\mathcal{I}_{\frac{1}{2}\text{BPS}}^{4\text{d } \mathcal{N}=4 \text{ } U(N)}(q) = \prod_{n=1}^N \frac{1}{(1 - q^n)} = \sum_{n_1, \dots, n_N \geq 0} q^{n_1 + 2n_2 + \dots + Nn_N}. \quad (2.15)$$

The supersymmetric spectral zeta function is given by

$$\mathbb{Z}_{\frac{1}{2}\text{BPS}}^{4\text{d } \mathcal{N}=4 \text{ } U(N)}(s, z) = \zeta_N(s, z; 1, 2, \dots, N), \quad (2.16)$$

where

$$\zeta_N(s, z; \omega_1, \dots, \omega_N) = \sum_{n_1, \dots, n_N \geq 0} (n_1 \omega_1 + \dots + n_N \omega_N + z)^{-s} \quad (2.17)$$

is the Barnes multiple zeta function [36]. It has simple poles at $s = 1, \dots, N$ with residues

$$\text{Res}_{s=N-i} \zeta_N(s, z; \omega_1, \dots, \omega_N) = \frac{F_i(e_1(\omega), \dots, e_k(\omega); z)}{\Gamma(N-i)} \prod_{j=1}^N \frac{1}{\omega_j}, \quad (2.18)$$

with $i = 0, 1, \dots, N - 1$. Here

$$F_i(e_1(\omega), \dots, e_k(\omega); z) = \sum_{k=0}^i T_k(e_1(\omega), \dots, e_k(\omega)) \frac{(-z)^{i-k}}{(i-k)!} \quad (2.19)$$

is a function of the elementary symmetric functions $e_1(\omega), \dots, e_k(\omega)$ and z . The functions $T_k(e_1(\omega), \dots, e_k(\omega))$ are the k -th Todd polynomials [37] which can be defined by

$$\prod_{i=1}^N \frac{tx_i}{1 - e^{-tx_i}} = \sum_{k=0}^{\infty} T_k(e_1(x), \dots, e_k(x)) t^k. \quad (2.20)$$

The residue formula (2.18) is obtained from the integral representation

$$\zeta_N(s, z; \omega_1, \dots, \omega_N) = \frac{1}{\Gamma(s)} \int_0^{\infty} dt \, t^{s-1} \frac{e^{-zt}}{(1 - e^{-\omega_1 t}) \dots (1 - e^{-\omega_N t})} \quad (2.21)$$

of the Barnes multiple zeta function. For example,

$$T_0 = 1, \quad (2.22)$$

$$T_1(e_1) = \frac{1}{2} e_1(x), \quad (2.23)$$

$$T_2(e_1, e_2) = \frac{1}{12} (e_1(x)^2 + e_2(x)), \quad (2.24)$$

$$T_3(e_1, e_2, e_3) = \frac{1}{24} e_1(x) e_2(x), \quad (2.25)$$

$$T_4(e_1, e_2, e_3, e_4) = \frac{1}{720} (-e_1(x)^4 + 4e_1(x)^2 e_2(x) + 3e_2(x)^2 + e_1(x) e_3(x) - e_4(x)). \quad (2.26)$$

The Barnes multiple zeta function (2.17) converges absolutely for $\text{Re}(s) > N$. It follows that the supersymmetric zeta function (2.16) has simple poles at $s = N - i$, $i = 0, \dots, N - 1$ with residues

$$\begin{aligned} & \text{Res}_{s=N-i} \mathbb{Z}_{\frac{1}{2}\text{BPS}}^{4d \, \mathcal{N}=4 \, U(N)}(s, z) \\ &= \frac{1}{\Gamma(N-i)N!} \sum_{k=0}^i T_k(|s(N+1, N)|, \dots, |s(N+1, N+1-k)|) \frac{(-z)^{i-k}}{(i-k)!}, \end{aligned} \quad (2.27)$$

where $s(n, m)$ is the Stirling number of the first kind. The residues (2.27) encode the terms with β^{-N+i} in the Laurent series expansion of the supersymmetric index (2.15).

By the inverse Laplace transform, we find the asymptotic degeneracy

$$\begin{aligned}
d_{\frac{1}{2}\text{BPS}}^{4d \mathcal{N}=4 U(N)}(n) &\sim \sum_{k=0}^{N-1} \frac{T_{N-k-1}(|s(N+1, N)|, \dots, |s(N+1, N+1-k)|)}{N!k!} n^k \\
&= \frac{1}{N!(N-1)!} n^{N-1} + \frac{N+1}{4(N-1)!(N-2)!} n^{N-2} \\
&\quad + \frac{(N+1)(9N^2+5N-2)}{288(N-1)!(N-3)!} n^{N-3} + \frac{N(N+1)^2(3N+2)}{1152(N-2)!(N-4)!} n^{N-4} + \dots
\end{aligned} \tag{2.28}$$

The number of the BPS states with charge less than or equal to x is

$$\begin{aligned}
\Omega_{\frac{1}{2}\text{BPS}}^{4d \mathcal{N}=4 U(N)}(x) &\sim \sum_{k=0}^{N-1} \frac{T_{N-k-1}(|s(N+1, N)|, \dots, |s(N+1, N+1-k)|)}{N!(k+1)!} x^{k+1} \\
&= \frac{1}{(N!)^2} x^N + \frac{N+1}{4((N-1)!)^2} x^{N-1} \\
&\quad + \frac{(N+1)(9N^2+5N-2)}{288(N-1)!(N-2)!} x^{N-2} + \frac{N(N+1)^2(3N+2)}{1152(N-2)!(N-3)!} x^{N-3} + \dots
\end{aligned} \tag{2.29}$$

The special values of the Barnes zeta function at negative integers are given by [38]

$$\zeta_N(-k, z; \omega_1, \dots, \omega_N) = \frac{(-1)^N k!}{(k+N)!} B_{k+N}(z; \omega_1, \dots, \omega_N), \tag{2.30}$$

where the multiple Bernoulli polynomial is defined by

$$\frac{x^n e^{xz}}{(e^{\omega_1 x} - 1) \dots (e^{\omega_n x} - 1)} = \sum_{k \geq 0} B_k(z; \omega_1, \dots, \omega_n) \frac{x^k}{k!}, \tag{2.31}$$

the energy moments are given by

$$\mathbb{Z}_{\frac{1}{2}\text{BPS}}^{4d \mathcal{N}=4 U(N)}(-k, 0) = \frac{(-1)^N k!}{(k+N)!} B_{k+N}(0; 1, \dots, N). \tag{2.32}$$

2.3 Multiple supersymmetric zeta functions

Our supersymmetric zeta functions can be extended to multiple supersymmetric zeta functions by introducing additional fugacities coupled to the Cartan generators of global symmetries in a straightforward fashion.

For the supersymmetric field theories in $d = 4, 5$ and 6 dimensions, there exist multiple Cartan generators of the rotational symmetry group. While these Cartan generators are coupled to independent fugacities $\{q_I\}$ for the supersymmetric indices, we first replace them with the exponential variables $\{\omega_I\}$ with the same base q for

the supersymmetric zeta functions, i.e. $q_I = q^{\omega_I}$. Then we obtain from (2.7) the multiple supersymmetric zeta functions depending on $\{\omega_I\}$ as the Mellin transform of the supersymmetric indices. We will give a variety of examples of the multiple supersymmetric zeta functions for 4d and 6d supersymmetric field theories in section 5 and section 6.

Furthermore, when the theories have other global symmetries, the supersymmetric indices can be refined by introducing equivariant parameters, also known as flavor fugacities $\{x_\alpha\}$ coupled to their Cartan generators. The resulting indices, which are referred to as the flavored supersymmetric indices, are formal multivariable Taylor series in the fugacities $\{q_i\}$ whose coefficients are Laurent series in the flavor fugacities $\{x_\alpha\}$. In this case, we obtain the flavored supersymmetric zeta functions depending on the flavor fugacities $\{x_\alpha\}$ by performing the Mellin transform (2.7) of the flavored supersymmetric indices. In this work, we specialize to the unflavored case for the supersymmetric zeta functions. We leave a full discussion on the flavored supersymmetric zeta functions to future work.

2.4 Degeneracy with exponential growth

If the supersymmetric indices have expansion coefficients of polynomial growth, the supersymmetric zeta functions can be directly defined by taking the Mellin transform (2.7) of the supersymmetric indices. However, the supersymmetric indices $\mathcal{I}(q)$ in supersymmetric field theories are typically defined as certain traces over the Fock space for a multi-particle system so that the degeneracy $d(n)$ may not exhibit polynomial growth. In such cases, we define the supersymmetric zeta functions $\mathfrak{Z}(s, z)$ by the Mellin transform of the “single-particle indices $i(q)$ ”, which we obtain by taking the plethystic logarithm of the supersymmetric indices $\mathcal{I}(q)$. As we will discuss below, the resulting supersymmetric zeta functions turn out to play a crucial role in studying the asymptotic Cardy-like behaviors and the supersymmetric Casimir energies.

2.4.1 Cardy-like limit

Consider the single-particle index $i(q)$ that is defined by the plethystic logarithm [39] of the supersymmetric index $\mathcal{I}(q)$

$$i(q) = \text{PL}[\mathcal{I}(q)] = \sum_{d \geq 1} \frac{\mu(d)}{d} \log \mathcal{I}(q^d), \quad (2.33)$$

where $\mu(d)$ is the Möbius function. Conversely, one can recover the supersymmetric index $\mathcal{I}(q)$ by taking the plethystic exponential of the single-particle index $i(q)$

$$\mathcal{I}(q) = \text{PE}[i(q)] = \exp \left[\sum_{k=1}^{\infty} \frac{i(q^k)}{k} \right]. \quad (2.34)$$

When the single-particle index $i(q)$ can be expanded as

$$i(q) = \sum_{\nu} \delta(\nu) q^{\nu}, \quad (2.35)$$

where $\delta(\nu)$ may not be integers, we define the supersymmetric zeta function as

$$\mathfrak{Z}(s, z) = \sum_{\nu} \frac{\delta(\nu)}{(\nu + z)^s}. \quad (2.36)$$

As discussed in section 2.3, this definition can be straightforwardly generalized to multiple supersymmetric zeta functions by introducing additional variables.

We assume that the supersymmetric zeta function (2.36) has simple poles at $s = \alpha_i$ and that it absolutely converges for $\text{Re}(s) > \alpha_*$, where α_* corresponds to the rightmost pole. Also, we assume that it can be analytically continued to the region $\text{Re}(s) \geq -C_0$ ($0 < C_0 < 1$). Furthermore, we assume that as $|\text{Im}(s)| \rightarrow \infty$,

$$\mathfrak{Z}(s, z) = \mathcal{O}(|\text{Im}(s)|^{C_1}) \quad (2.37)$$

uniformly on $\text{Re}(s) \geq -C_0$ for some positive number C_1 . The number C_1 can be viewed as the order of growth of the supersymmetric zeta function, which generalizes the least possible exponent (order of growth) [40, 41] of the Riemann zeta function.

We note that the logarithm of the supersymmetric index $\mathcal{I}(q)$ is given by

$$\begin{aligned} \log \mathcal{I}(\beta) &= \sum_{k=1}^{\infty} \frac{i(e^{-k\beta})}{k} \\ &= \sum_{k=1}^{\infty} \sum_{\nu} \frac{\delta(\nu)}{k} e^{-\nu k\beta}, \end{aligned} \quad (2.38)$$

where we have set $q = e^{-\beta}$. Using the Mellin inversion formula

$$e^{-\beta} = \frac{1}{2\pi i} \int_{\sigma_0 - i\infty}^{\sigma_0 + i\infty} ds \beta^{-s} \Gamma(s) \quad (2.39)$$

for $\text{Re} \beta > 0$ and $\sigma_0 > 0$, the exponential function in (2.38) can be rewritten. Consequently, we get

$$\log \mathcal{I}(\beta) = \frac{1}{2\pi i} \int_{\alpha_* + 1 - i\infty}^{\alpha_* + 1 + i\infty} ds \beta^{-s} \Gamma(s) \zeta(s+1) \mathfrak{Z}(s, 0). \quad (2.40)$$

Here we have interchanged the order of summation and integration, which is justified by the assumed absolute convergence of the series. By shifting the line of integration from $\text{Re}(s) = \alpha_* + 1$ to $\text{Re}(s) = -C_0$, we can evaluate the integral (2.40) by picking up the residues

$$\beta^{-\alpha_i} \Gamma(\alpha_i) \zeta(\alpha_i + 1) \text{Res}_{s=\alpha_i} \mathfrak{Z}(s, 0) \quad (2.41)$$

at simple poles $s = \alpha_i$ and the residue

$$\mathfrak{Z}(0, 0)' - \mathfrak{Z}(0, 0) \log \beta \quad (2.42)$$

at a double pole $s = 0$, where $\mathfrak{Z}(0, 0)' = \frac{\partial}{\partial s} \mathfrak{Z}(s, z)|_{s=0, z=0}$. Then we find that

$$\begin{aligned} \log \mathcal{I}(\beta) &= \sum_i \frac{\text{Res}_{s=\alpha_i} \mathfrak{Z}(s, 0) \Gamma(\alpha_i) \zeta(\alpha_i + 1)}{\beta^{\alpha_i}} - \mathfrak{Z}(0, 0) \log \beta + \mathfrak{Z}(0, 0)' \\ &\quad + \frac{1}{2\pi i} \int_{-C_0-i\infty}^{-C_0+i\infty} ds \beta^{-s} \Gamma(s) \zeta(s+1) \mathfrak{Z}(s, 0). \end{aligned} \quad (2.43)$$

The final integral can be bounded as follows:

$$\begin{aligned} &\left| \frac{1}{2\pi i} \int_{-C_0-i\infty}^{-C_0+i\infty} ds \beta^{-s} \Gamma(s) \zeta(s+1) \mathfrak{Z}(s, 0) \right| \\ &= \mathcal{O} \left(|\beta|^{C_0} \int_{-\infty}^{\infty} dt \exp \left(-\frac{\pi}{2} |t| \right) |t|^{C_1+C_2+C_3} \right) \\ &= \mathcal{O} (|\beta|^{C_0}), \end{aligned} \quad (2.44)$$

where we have used the assumption (2.37), the least possible exponent (order of growth) of the Riemann zeta function [40, 41]

$$\zeta(s+1) = \mathcal{O}(|\text{Im}(s)|^{C_2}) \quad (2.45)$$

and the asymptotic estimate of the gamma function [42]

$$\Gamma(s) \sim \sqrt{2\pi} |\text{Im}(s)|^{\text{Re}(s)-\frac{1}{2}} \exp \left(-\frac{\pi}{2} |\text{Im}(s)| \right), \quad (2.46)$$

which implies $\Gamma(s) = \mathcal{O} \left(\exp \left(-\frac{\pi}{2} |\text{Im}(s)| \right) |\text{Im}(s)|^{C_3} \right)$ as $|\text{Im}(s)| \rightarrow \infty$.

Hence, we obtain the formula for the asymptotic behavior as $\beta \rightarrow 0$

$$\begin{aligned} &\log \mathcal{I}(\beta) \\ &= \sum_i \frac{\text{Res}_{s=\alpha_i} \mathfrak{Z}(s, 0) \Gamma(\alpha_i) \zeta(\alpha_i + 1)}{\beta^{\alpha_i}} - \mathfrak{Z}(0, 0) \log \beta + \mathfrak{Z}(0, 0)' + \mathcal{O} (|\beta|^{C_0}). \end{aligned} \quad (2.47)$$

The asymptotic formula (2.47) applies to numerous supersymmetric indices with exponential growth. We refer to it as the Cardy-like limit of the supersymmetric index

(see [43] for the original Cardy formula for the partition function of a 2d CFT). The formula (2.47) asserts that the terms $\beta^{-\alpha_i}$ with $\alpha_i > 0$ in the Cardy-like limit of the supersymmetric indices correspond to simple poles at α_i of the supersymmetric zeta functions $\mathfrak{Z}(s, z)$, and that their coefficients are encoded by the residues at these poles with $z = 0$. For superconformal field theories in 2d, 4d, and 6d, it has been pointed out that these coefficients can be universally determined by the anomaly coefficients [44, 45] (see e.g. [46, 47, 48, 23, 24, 25, 26, 27, 32, 49] for further variants). In such cases, there exist universal relations between the residues of the supersymmetric zeta functions and the anomaly coefficients, as we will see in section 4, section 5 and section 6.

There may exist a logarithmic correction in (2.47) if the supersymmetric zeta value $\mathfrak{Z}(0, 0)$ is non-trivial. We shall call $\mathfrak{Z}(0, 0)$ the *Zeta-index*. Note that it differs from the ordinary Witten index as the former is associated with the plethystic logarithm of a supersymmetric index that exhibits exponential growth. It has been argued [45, 47] that the logarithmic term in the Cardy-like limit of the 4d $\mathcal{N} = 1$ supersymmetric index arises due to the presence of the unlifted zero modes parameterizing the quantum Coulomb branch of the 3d $\mathcal{N} = 2$ gauge theory obtained upon compactifying the 4d $\mathcal{N} = 1$ gauge theory on a circle. Accordingly, the Zeta-index for the 4d $\mathcal{N} = 1$ gauge theory is expected to encode the complex dimension of the quantum Coulomb branch of the resulting 3d theory.

Moreover, the constant term in (2.47) is given by the s -derivative of the supersymmetric zeta function evaluated at the origin ($s = 0, z = 0$). This is naturally characterized by a special value of the supersymmetric determinant discussed in section 3.

2.4.2 Effective Field Theory interpretation

At this point, it is instructive to discuss the effective field theory interpretation of the formula (2.47). The supersymmetric index \mathcal{I} is, in fact, a specific partition function on $S^1 \times S^{d-1}$ (twisted by R-symmetry, etc.). Therefore, the effective field theory analysis in the $\beta \rightarrow 0$ limit can be applied to $\mathcal{I}(\beta)$ just as it would be to a standard partition function. The idea is that the supersymmetric index, or more specifically the superconformal index, can be regarded as a type of partition function on $S^1 \times S^{d-1}$. On the other hand, we can predict the behavior of the partition function in the small β limit by using the effective field theory techniques. The comparison gives a physical interpretation of the asymptotic behavior of the free energy as a function of β^{-1} .

Suppose we want to study the partition function of a certain conformal field theory on $S^1 \times S^{d-1}$. It can be realized as a path integral over $S^1 \times S^{d-1}$ with appropriate

boundary conditions on the Euclidean time circle of radius β . Let us assume that the boundary condition is such that there is no zero mode, and the low-energy effective field theory on S^{d-1} becomes gapped. In this situation, the partition function, more precisely the free energy (i.e., the logarithm of the partition function), must be given by a local functional of the background metric $\hat{g}_{\mu\nu} = \beta^{-2}g_{\mu\nu}$ on S^{d-1} (up to Weyl anomaly). This is the consequence of the Weyl invariance because the original metric $ds^2 = \beta^2 d\tau^2 + g_{\mu\nu}dx^\mu dx^\nu$ is Weyl equivalent to $d\hat{s}^2 = d\tau^2 + \beta^{-2}g_{\mu\nu}dx^\mu dx^\nu$. In more complicated situations, we may further introduce the background Kaluza-Klein graviphoton to describe the spin and the other gauge fields to describe chemical potentials of global symmetries.

If we accept the effective field theory analysis, we should have

$$Z(\beta)_{\text{CFT}_d} = Z(\beta)_{\text{gapped}_{d-1}} \sim e^{-S_{d-1}[\hat{g}]} \quad (2.48)$$

with the effective action given by the derivative expansion:

$$S[g] = \int d^{d-1}x \sqrt{\hat{g}}(-f + c_1 \hat{R} + c_2 \hat{R}^2 + \dots). \quad (2.49)$$

Substituting the metric \hat{g} for S^{d-1} explicitly, we find

$$S[g] = \text{Vol}S^{d-1} \left(-f\beta^{1-d} + (d-2)(d-1)c_1\beta^{3-d} + (d-2)^2(d-1)^2c_2\beta^{5-d} \dots \right). \quad (2.50)$$

It should be compared with (2.47). In particular, the residues and special values of $\mathfrak{Z}(s, 0)$ are directly related to the effective field theory Wilson coefficients f , c_1 , c_2 , and so on.

For a more precise comparison, we need to consider several additional features. First of all, our target is the supersymmetric index and not the partition function. Furthermore, our definition of “time evolution” may involve extra contributions from the gauge field background. For instance, the “Hamiltonian” defining the superconformal index contains rotation and the R-symmetry chemical potential, and the extra terms in the effective action, such as the Chern-Simons terms for the background gauge field, will also contribute. In addition, from the viewpoint of the effective field theory, the supersymmetry introduces further constraints: the supersymmetry forbids the appearance of several terms in the effective action, such as the “cosmological constant” f , or the supersymmetry relates various terms, e.g., the Einstein-Hilbert term with the R-symmetry Chern-Simons term in $d = 4$.

A systematic analysis of the effective field theory for the superconformal index can be found, e.g., in the work [45] of di Pietro and Komargodski. They, for example, showed that, after taking into account the R-symmetry twisting, the Wilson coefficient

c_1 is proportional to a particular combination of the central charges $c - a$ in $d = 4$. They also studied the prediction of several coefficients in $d = 6$ from the anomaly polynomial, and further confirmation was provided in the work [50] of Chang et al. by studying the 5d effective action. We will use these facts in later sections.

If the S^1 compactification leaves the “gapless” degrees of freedom or zero modes on S^{d-1} , the effective field theory expansion can fail. Its manifestation is the presence of the $\log \beta$ term, which we called the Zeta-index. We have already mentioned the claim that the coefficient in 4d encodes the complex dimension of the quantum Coulomb branch of the compactified 3d theory. Such a contribution cannot be written as a local functional of the metric and gauge fields.

Finally, when d is even, the Weyl anomaly exists, so performing the Weyl transformation from $ds^2 = \beta^2 d\tau^2 + g_{\mu\nu} dx^\mu dx^\nu$ to $d\tilde{s}^2 = d\tau^2 + \beta^{-2} g_{\mu\nu} dx^\mu dx^\nu$ introduces extra terms proportional to β , which are related to the Casimir energy that we will discuss in section 2.4.4. In the work [51] of Benjamin et al., it was argued that if we could choose a scheme where the Casimir energy in the $d\tilde{s}^2$ metric is zero, the Casimir energy in the ds^2 metric should be entirely determined from the Weyl anomaly. We will come back to this point in the context of the superconformal index in section 2.4.4.

2.4.3 Asymptotic degeneracy

We next consider the asymptotic behavior of the degeneracy $d(n)$. Here we assume that it grows exponentially as $n \rightarrow \infty$.³ According to the Cauchy integral theorem, we have

$$d(n) = \frac{1}{2\pi i} \int_{\beta_0}^{\beta_0 + 2\pi i} d\beta \mathcal{I}(\beta) e^{n\beta}. \quad (2.51)$$

With $\beta = y + 2\pi i x$, it can be written as

$$d(n) = \int_{-\frac{1}{2}}^{\frac{1}{2}} dx \mathcal{I}(y + 2\pi i x) e^{ny + 2\pi i n x}. \quad (2.52)$$

Our strategy is to apply the saddle point approximation to the integral (2.52). As the supersymmetric index $\mathcal{I}(\beta)$ has a singularity at $\beta = 0$, we wish to evaluate the dominant saddle by making use of the Cardy-like limit (2.47) as $\beta \rightarrow 0$. The absolute value of the integrand can be approximated as

$$|\mathcal{I}(y + 2\pi i x) e^{ny + 2\pi i n x}| \sim \left| \exp \left[\text{Res}_{s=\alpha_*} \mathfrak{Z}(s, 0) \Gamma(\alpha_*) \zeta(\alpha_* + 1) (y + 2\pi i x)^{-\alpha_*} + ny \right] \right|, \quad (2.53)$$

³An important caveat is that if the degeneracy $d(n)$ does not exhibit exponential growth (say, if it alternates between positive and negative values), the following asymptotic analysis will not work in general.

where α_* is the location of the rightmost pole as discussed in section 2.4.1. Provided that the residue $\text{Res}_{s=\alpha_*} \mathfrak{Z}(s, 0)$ at the rightmost pole for $z = 0$ is positive, the maximum absolute value of the integrand in (2.52) appears when $x = 0$. Then the integrand with $x = 0$ can be well approximated by

$$\exp [\text{Res}_{s=\alpha_*} \mathfrak{Z}(s, 0) \Gamma(\alpha_*) \zeta(\alpha_* + 1) y^{-\alpha_*} + ny], \quad (2.54)$$

Applying the saddle point approximation, we should choose y by solving

$$\frac{\partial}{\partial y} \left\{ \exp [\text{Res}_{s=\alpha_*} \mathfrak{Z}(s, 0) \Gamma(\alpha_*) \zeta(\alpha_* + 1) y^{-\alpha_*} + ny] \right\} = 0. \quad (2.55)$$

Then we find

$$y = n^{-\frac{1}{\alpha_*+1}} [\text{Res}_{s=\alpha_*} \mathfrak{Z}(s, 0) \Gamma(\alpha_* + 1) \zeta(\alpha_* + 1)]^{\frac{1}{\alpha_*+1}}. \quad (2.56)$$

Substituting (2.56) into (2.52), we obtain

$$d(n) \sim e^m \int_{-\frac{1}{2}}^{\frac{1}{2}} dx \mathcal{I}(y + 2\pi ix) e^{2\pi i n x}, \quad (2.57)$$

where we have defined

$$m = n^{\frac{\alpha_*}{\alpha_*+1}} [\text{Res}_{s=\alpha_*} \mathfrak{Z}(s, 0) \Gamma(\alpha_* + 1) \zeta(\alpha_* + 1)]^{\frac{1}{\alpha_*+1}} \quad (2.58)$$

To proceed, we also note that the integral (2.57) can be further estimated by specifying a path where the angle $\arg(\beta)$ is fixed. Suppose that $|x| \leq \frac{y}{2\pi}$. Then we have the condition $|\arg(\beta)| \leq \frac{\pi}{4}$. In this case, as y is sufficiently small so that we can split the integral (2.57) into two parts i) $0 \leq |x| \leq y^\delta$ and ii) $y^\delta \leq |x| \leq \frac{1}{2}$ for some δ with $1 + \frac{\alpha_*}{3} < \delta < 1 + \frac{\alpha_*}{2}$ for which the former region is dominant in the large n limit. This process is similar to the Hardy-Ramanujan Farey dissection that divides the unit circle into specific segments or arcs in the Hardy-Ramanujan-Littlewood circle method [52]. The dominant integral takes the form

$$\begin{aligned} d(n) \sim \exp \left[m \left(1 + \frac{1}{\alpha_*} \right) + \mathfrak{Z}(0, 0)' \right] \\ \times \int_{-(\frac{m}{n})^\delta}^{(\frac{m}{n})^\delta} dx \exp \left\{ \frac{m}{\alpha_*} \left[-1 + \left(1 + \frac{2\pi i n}{m} x \right)^{-\alpha_*} \right] + 2\pi i n x - \mathfrak{Z}(0, 0) \log \left(\frac{m}{n} + 2\pi i x \right) \right\}, \end{aligned} \quad (2.59)$$

where we have used the Cardy-like limit (2.47). After the variable change $2\pi x = \frac{m}{n} \omega$, the expression can be rewritten as

$$\begin{aligned} d(n) \sim \exp \left[m \left(1 + \frac{1}{\alpha_*} \right) + \mathfrak{Z}(0, 0)' - (\mathfrak{Z}(0, 0) - 1) \log \left(\frac{m}{n} \right) - \log(2\pi) \right] \\ \times \int_{-2\pi(\frac{m}{n})^{\delta-1}}^{2\pi(\frac{m}{n})^{\delta-1}} d\omega \exp \left[m \left(\frac{1}{\alpha_* (1 + i\omega)^\alpha} - \frac{1}{\alpha_*} + i\omega \right) - \mathfrak{Z}(0, 0) \log(1 + i\omega) \right]. \end{aligned} \quad (2.60)$$

Again taking the limit $m \rightarrow \infty$ (or equivalently $n \rightarrow \infty$), the integral in the second line is approximated as

$$\begin{aligned}
& \int_{-2\pi(\frac{m}{n})^{\delta-1}}^{2\pi(\frac{m}{n})^{\delta-1}} d\omega \exp \left[m \left(\frac{1}{\alpha_*(1+i\omega)^{\alpha_*}} - \frac{1}{\alpha_*} + i\omega \right) - \mathfrak{Z}(0,0) \log(1+i\omega) \right] \\
&= \int_{-Cm^{\frac{1-\delta}{\alpha_*}}}^{Cm^{\frac{1-\delta}{\alpha_*}}} d\omega \exp \left[-\frac{m(\alpha_*+1)}{2} \omega^2 \right. \\
&\quad \left. + m \left(\frac{1}{\alpha_*(1+i\omega)^{\alpha_*}} - \frac{1}{\alpha_*} + i\omega + \frac{\alpha_*+1}{2} \omega^2 \right) - \mathfrak{Z}(0,0) \log(1+i\omega) \right] \\
&\sim \int_{-Cm^{\frac{1-\delta}{\alpha_*}}}^{Cm^{\frac{1-\delta}{\alpha_*}}} d\omega \exp \left[-\frac{m(\alpha_*+1)}{2} \omega^2 + \mathcal{O}(m|\omega|^3) - \mathfrak{Z}(0,0) \mathcal{O}(|\omega|) \right], \tag{2.61}
\end{aligned}$$

where

$$C = 2\pi [\text{Res}_{s=\alpha_*} \mathfrak{Z}(s,0) \Gamma(\alpha_*+1) \zeta(\alpha_*+1)]^{\frac{\delta-1}{\alpha_*}}. \tag{2.62}$$

As $m \rightarrow \infty$, we have $\mathcal{O}(|\omega|) = \mathcal{O}(m^{\frac{1-\delta}{\alpha_*}})$ and therefore the dominant integral is further evaluated as

$$\begin{aligned}
& \int_{-Cm^{\frac{1-\delta}{\alpha_*}}}^{Cm^{\frac{1-\delta}{\alpha_*}}} d\omega \exp \left[-\frac{m(\alpha_*+1)}{2} \omega^2 \right] \\
&= \sqrt{\frac{2}{m(\alpha_*+1)}} \int_{-C'm^{\frac{1}{2}+\frac{1-\delta}{\alpha_*}}}^{C'm^{\frac{1}{2}+\frac{1-\delta}{\alpha_*}}} d\omega' \exp(-\omega'^2) \\
&\sim \left(\frac{2\pi}{m(\alpha_*+1)} \right)^{\frac{1}{2}}. \tag{2.63}
\end{aligned}$$

Combining (2.60) and (2.63), we finally obtain

$$d(n) \sim \exp \left[\left(1 + \frac{1}{\alpha_*} \right) m + \mathfrak{Z}(0,0)' \right] \left(\frac{n}{m} \right)^{3(0,0)-1} (2\pi m(\alpha_*+1))^{-\frac{1}{2}}. \tag{2.64}$$

This asymptotic formula (2.64) provides the leading asymptotic behavior of the degeneracy for a supersymmetric zeta function $\mathfrak{Z}(s,0)$ with a simple pole at $s = \alpha_* > 0$, which can be viewed as an extension of the Meinardus theorem [53]. For example, if the supersymmetric zeta function has a simple pole at $s = 1$, we have

$$d(n) \sim \frac{1}{2} \left(\frac{R_1}{6n^3} \right)^{\frac{1}{4}} \left(\frac{6n}{\pi^2 R_1} \right)^{\frac{3(0,0)}{2}} \exp \left[2\pi \sqrt{\frac{R_1}{6}} n^{\frac{1}{2}} + \mathfrak{Z}(0,0)' \right] \tag{2.65}$$

with $R_1 = \text{Res}_{s=1} \mathfrak{Z}(s,0)$.

When the supersymmetric zeta functions have multiple simple poles, the subleading terms contribute correction terms to the asymptotic formula (2.64), although the leading term corresponding to the rightmost pole is still well approximated. A general method to evaluate the subleading terms was mathematically formulated in [54]. For equidistant simple poles at $s = i$, with $i = 0, 1, \dots, r$ and $r = \alpha_*$, the asymptotic formula for $d(n)$ will take the form

$$d(n) \sim \exp \left[n\delta_n + \sum_{j=1}^r (\text{Res}_{s=j} \Gamma(j) \zeta(j+1) \delta_n^{-j}) + \mathfrak{Z}(0,0)' \right] \left(\frac{n}{m} \right)^{\mathfrak{Z}(0,0)-1} (2\pi m(\alpha_* + 1))^{-\frac{1}{2}}, \quad (2.66)$$

where

$$\delta_n = [\text{Res}_{s=\alpha_*} \mathfrak{Z}(s,0) \Gamma(\alpha_* + 1) \zeta(\alpha_* + 1)]^{\frac{1}{\alpha_*+1}} \sum_{j=0}^r \psi_j z^{j+1} \Big|_{z \rightarrow n^{-\frac{1}{r+1}}} \quad (2.67)$$

and ψ_j can be determined by solving the recursive equation

$$h_r \left(\sum_{l=0}^s \psi_l z^l \right)^{r+1} = \sum_{k=0}^r h_k h_r^{\frac{r-k}{r+1}} z^{r-k} \left(\sum_{l=0}^s \psi_l z^l \right)^{r-k} \quad (2.68)$$

with

$$h_i = \text{Res}_{s=i} \mathfrak{Z}(s,0) \Gamma(i+1) \zeta(i+1). \quad (2.69)$$

For example, when the supersymmetric zeta function has simple poles at $s = 2$ and 1 , we find

$$d(n) \sim \frac{1}{\sqrt{6\pi}} \left(\frac{2\zeta(3)R_2}{n^4} \right)^{\frac{1}{6}} \left(\frac{n}{2\zeta(3)R_2} \right)^{\frac{\mathfrak{Z}(0,0)}{3}} \times \exp \left[\frac{3}{2} (2\zeta(3)R_2)^{\frac{1}{3}} n^{\frac{2}{3}} + \frac{R_1 \pi^2}{6} \left(\frac{1}{2\zeta(3)R_2} \right)^{\frac{1}{3}} n^{\frac{1}{3}} - \frac{\pi^4 R_1^2}{432 \zeta(3) R_2} + \mathfrak{Z}(0,0)' \right], \quad (2.70)$$

where we have defined $R_i = \text{Res}_{s=i} \mathfrak{Z}(s,0)$. Such growth appears in the large N indices of the 3d supersymmetric field theories describing a stack of N M2-branes [55, 56]. When the supersymmetric zeta function has simple poles at $s = 3, 2$ and 1 , we obtain

$$d(n) \sim \frac{1}{2\sqrt{2}} \left(\frac{R_3}{15n^5} \right)^{\frac{1}{8}} \left(\frac{15n}{\pi^4 R_3} \right)^{\frac{\mathfrak{Z}(0,0)}{4}} \times \exp \left[\frac{4\pi}{3} \left(\frac{R_3}{15} \right)^{\frac{1}{4}} n^{\frac{3}{4}} + \frac{\zeta(3)R_2}{\pi^2} \left(\frac{15}{R_3} \right)^{\frac{1}{2}} n^{\frac{1}{2}} + \frac{\pi^6 R_1 R_3 - 45 \zeta(3)^2 R_2^2}{2\pi^5} \left(\frac{5}{27 R_3^5} \right)^{\frac{1}{4}} n^{\frac{1}{4}} - \frac{5(\pi^6 \zeta(3) R_1 R_2 R_3 - 60 \zeta(3)^3 R_2^3)}{4\pi^8 R_3^2} + \mathfrak{Z}(0,0)' \right]. \quad (2.71)$$

Furthermore, when there are simple poles at $s = 4, 3, 2$ and 1 , we find

$$\begin{aligned}
d(n) \sim & \frac{1}{\sqrt{10}\pi} \left(\frac{24\zeta(5)R_4}{n^6} \right)^{\frac{1}{10}} \left(\frac{n}{24\zeta(5)R_4} \right)^{\frac{3(0,0)}{5}} \\
& \times \exp \left[\frac{5}{4} (24\zeta(5)R_4)^{\frac{1}{5}} n^{\frac{4}{5}} + \frac{R_3\pi^4}{45} \left(\frac{1}{24\zeta(5)R_4} \right)^{\frac{3}{5}} n^{\frac{3}{5}} \right. \\
& + \frac{54000\zeta(3)\zeta(5)R_2R_4 - \pi^8 R_3^2}{108000} \left(\frac{1}{18\zeta(5)^7 R_4^7} \right)^{\frac{1}{5}} n^{\frac{2}{5}} \\
& + \frac{\pi^{12} R_3^3 - 54000\pi^4 \zeta(3)\zeta(5)R_2R_3R_4 + 8100000\pi^2 \zeta(5)^2 R_1 R_4^2}{48600000} \left(\frac{1}{24\zeta(5)^{11} R_4^{11}} \right)^{\frac{1}{5}} n^{\frac{1}{5}} \\
& - \frac{1}{41990400000\zeta(5)^3 R_4^3} (\pi^{16} R_3^4 - 64800\pi^8 \zeta(3)\zeta(5)R_2 R_3^2 R_4 \\
& \left. + 3888000\pi^6 \zeta(5)^2 R_1 R_3 R_4^2 + 699840000\zeta(3)^2 \zeta(5)^2 R_2^2 R_4^2) + \mathfrak{Z}(0,0)' \right]. \quad (2.72)
\end{aligned}$$

When the supersymmetric zeta function has simple poles at $s = 5, 4, 3, 2$ and 1 , we find

$$\begin{aligned}
d(n) \sim & \frac{1}{2\sqrt{3}} \left(\frac{8R_5}{63n^7} \right)^{\frac{1}{12}} \left(\frac{63n}{8\pi^6 R_5} \right)^{\frac{3(0,0)}{6}} \\
& \times \exp \left[\frac{6\pi}{5} \left(\frac{8R_5}{63} \right)^{\frac{1}{6}} n^{\frac{5}{6}} + \frac{6\zeta(5)}{\pi^4} \left(\frac{63}{8R_5} \right)^{\frac{2}{3}} n^{\frac{2}{3}} \right. \\
& + \frac{\pi^{10} R_3 R_5 - 17010\zeta(5)^2 R_4^2}{30\pi^9} \left(\frac{7}{2R_5^3} \right)^{\frac{1}{2}} n^{\frac{1}{2}} \\
& + \frac{10\pi^{12} \zeta(3) R_2 R_5^2 - 21\pi^{10} \zeta(5) R_3 R_4 R_5 + 317520\zeta(5)^3 R_4^3}{20\pi^4} \left(\frac{63}{R_5^7} \right)^{\frac{1}{3}} n^{\frac{1}{3}} \\
& - \frac{1}{800\pi^{19}} \left(\pi^{20} R_5^2 (7R_3^2 - 400R_1 R_5) + 151200\pi^{12} \zeta(3)\zeta(5) R_2 R_4 R_5^2 \right. \\
& - 555660\pi^{10} \zeta(5)^2 R_3 R_4 R_5 + 6826283100\zeta(5)^4 R_4^4) \left(\frac{7}{648R_5^{19}} \right)^{\frac{1}{6}} n^{\frac{1}{6}} \\
& - \frac{7}{800\pi^{24} R_5^4} \left(20\pi^{22} \zeta(3) R_2 R_3 R_5^3 + 3\pi^{20} \zeta(5) R_4 R_5^2 (-21R_3^2 + 200R_1 R_5) \right. \\
& - 680400\pi^{12} \zeta(3)\zeta(5)^2 R_2 R_4^2 R_5^2 + 2857680\pi^{10} \zeta(5)^3 R_3 R_4^3 R_5 \\
& \left. \left. - 29165482080\zeta(5)^5 R_4^5 \right) + \mathfrak{Z}(0,0)' \right]. \quad (2.73)
\end{aligned}$$

Although we focus on the asymptotic BPS degeneracy for the supersymmetric theories in this work, our asymptotic formulae will be applicable to larger classes of counting functions with exponential growth by introducing appropriate zeta functions.

For example, the asymptotic density of states in d -dimensional CFT can be extracted by analyzing its partition function on $S^1 \times S^{d-1}$, which is a weighted sum over states on S^{d-1} . In such cases some of the residues of the associated zeta functions may vanish so that our formulae reduce to the formulae obtained in [51] by using the thermal effective action.⁴

It is worth noting that the above asymptotic formulae for the degeneracy $d(n)$ may not be valid if some of the residues are non-positive values. The extent of its validity and the error terms are not yet known exactly. Also we remark that the fact that the discrete spectrum may be characterized by different finite spacing needs further consideration for the determination of the exact constant term in the asymptotic degeneracy $d(n)$. Nevertheless, our asymptotic formulae for the degeneracy $d(n)$ may be used to provide a good approximation for the asymptotic number $\Omega(x)$ of the BPS operators with charge less than or equal to x , even though the formulae themselves are invalid.

2.4.4 Supersymmetric Casimir energies

The supersymmetric index \mathcal{I} for the d -dimensional supersymmetric quantum field theory is intimately related to the supersymmetric partition function $\mathcal{Z}_{S^1 \times S^{d-1}}$ on $S^1 \times S^{d-1}$. It is known that they can be distinguished by the supersymmetric Casimir energy factor

$$\mathcal{Z}_{S^1 \times S^{d-1}} = e^{-\beta E_{\text{SUSY}}} \mathcal{I}, \quad (2.74)$$

where the supersymmetric Casimir energy E_{SUSY} can be obtained from the large circle limit (zero temperature limit)

$$E_{\text{SUSY}} = - \lim_{\beta \rightarrow \infty} \frac{\partial}{\partial \beta} \log \mathcal{Z}_{S^1 \times M_{d-1}}^{\text{SUSY}}. \quad (2.75)$$

In general, the Casimir energies may depend on the regularization scheme; however, it has been argued that they are scheme-independent and enjoy the universal structure [59, 60, 61, 62, 63, 64, 65, 66, 67] (also see e.g. [68, 69, 70, 71, 72, 73] for further study). In fact, the statement was demonstrated in [65] for 4d $\mathcal{N} = 1$ superconformal field theories, for which the supersymmetric partition functions are scheme-independent. Subsequently, it was proposed by Bobev, Bullimore and Kim [66] that the supersymmetric Casimir energies for even-dimensional superconformal field theories are evaluated as the equivariant integrals of the anomaly polynomials.

⁴Also see [57, 58] for a survey of the asymptotic degeneracy of the Hilbert series by means of the effective field theory.

The main focus of our paper is the supersymmetric zeta functions, and it is not immediately obvious if we can say anything about the supersymmetric Casimir energies. However, by comparing the supersymmetric Casimir energies studied in the literature with our computation of the supersymmetric zeta values in various examples, we find that a non-trivial relation exists:

$$E_{\text{SUSY}} = \frac{1}{2} \mathfrak{Z}(-1, 0), \quad (2.76)$$

where the supersymmetric zeta function $\mathfrak{Z}(s, z)$ is associated with the plethystic logarithm of the supersymmetric indices. This is suggestive because the supersymmetric zeta values at negative integers s can be interpreted as certain regularized energy moments for superconformal field theories, but it is highly non-trivial because it implies that the supersymmetric zeta values somehow encode information about the supersymmetric partition functions rather than just the index.

We will directly compute the supersymmetric Casimir energies according to the formula (2.76) for many examples in later sections. For free theories with UV Lagrangian descriptions, we can easily check that the relation (2.76) reproduces the formulae obtained via the supersymmetric localization technique, which were originally pointed out by Kim [59], where the supersymmetric Casimir energy shift in the effective action can be formally written as

$$E_{\text{SUSY}} = \frac{1}{2} \text{Tr}(-1)^F \Delta. \quad (2.77)$$

Here the trace is taken over the single particle Hilbert space in such a way that $\text{Tr}(-1)^F q^\Delta$ is the single-particle index $i(q)$. Δ is an appropriate combination of the dilatation operator and the Cartan generators of the R-symmetry that commute with the chosen supercharges for the definition of the supersymmetric index. The summation in (2.77) should require regularization. It can be evaluated from the single-particle index $i(q)$ as [59, 60]

$$E_{\text{SUSY}} = \frac{1}{2} \lim_{q \rightarrow 1^-} q \frac{\partial}{\partial q} i(q), \quad (2.78)$$

which turns out to be compatible with (2.76).

In Lagrangian field theories where the localization computation is possible, the supersymmetric Casimir energy evaluated in [60] agrees with our computation whenever the comparison is possible. Note that beyond free theories, the way the Casimir energy was computed there is different from ours, but the final result agrees. Remarkably, our formula can also be applied to the non-Lagrangian theories. In fact, as we will see in sections 5.2.5, 5.2.7 and 5.2.8, by combining the supersymmetric Casimir energies evaluated from the formula (2.76) for the non-Lagrangian theories with the

universal formula proposed in [66] for the supersymmetric Casimir energies, we obtain the correct central charges.

We now suggest that the universal properties of the Cardy-like behaviors and the supersymmetric Casimir energies result from those of the supersymmetric zeta functions. An important byproduct of the supersymmetric zeta functions is a novel approach to extract anomaly coefficients or central charges for superconformal field theories from the BPS spectral data.

3 Supersymmetric determinants

3.1 Definition

The properties of the spectrum can also be encoded in the spectral determinants. Here we define the *supersymmetric determinant* by

$$\mathbb{D}(z) = \exp \left[\frac{\partial}{\partial s} \mathbb{Z}(s, z) \right] \Big|_{s=0}. \quad (3.1)$$

It can be written as

$$\mathbb{D}(z) = \prod_n (n + z)^{-d(n)}. \quad (3.2)$$

The supersymmetric spectral determinant is a function of the “charge variable” z , which is related to certain energy eigenvalues in the radial quantization for the superconformal field theories. The poles (resp. zeros) at $z = -n$ of order $|d(n)|$ correspond to the $|d(n)|$ bosonic (resp. $|d(n)|$ fermionic) operators of charge n in the theory.

When the degeneracy $d(n)$ does not grow polynomially, we use the supersymmetric zeta function $\mathfrak{Z}(s, z)$ given by (2.36) in the definition (3.1) and the $d(n)$ in the product expression (3.2) is replaced by $\delta(n)$. We denote the resulting supersymmetric determinant by

$$\begin{aligned} \mathfrak{D}(z) &= \exp \left[\frac{\partial}{\partial s} \mathfrak{Z}(s, z) \right] \Big|_{s=0} \\ &= \prod_n (n + z)^{-\delta(n)}. \end{aligned} \quad (3.3)$$

3.2 Laplace transformation

The spectral determinant (3.1) is related to the supersymmetric index via a Laplace transformation. From (2.9) and (3.1), the logarithm of the spectral determinant can

be written as

$$\begin{aligned}
\log \mathbb{D}(z) &= \left. \frac{\partial}{\partial s} \mathbb{Z}(s, z) \right|_{s=0} \\
&= \left. \frac{\partial}{\partial s} \left[\frac{1}{\Gamma(s)} \int_0^\infty d\beta \, \mathbb{I}(\beta) e^{-\beta z} \beta^{s-1} \right] \right|_{s=0} \\
&= \left. \left[-\frac{1}{\Gamma(s)} \int_0^\infty d\beta \, \mathbb{I}(\beta) e^{-\beta z} \beta^{s-1} \psi(s) + \frac{1}{\Gamma(s)} \int_0^\infty d\beta \, \mathbb{I}(\beta) e^{-\beta z} \beta^{s-1} \log \beta \right] \right|_{s=0} \\
&= \int_0^\infty d\beta \, \frac{\mathbb{I}(\beta)}{\beta} e^{-\beta z}^*,
\end{aligned} \tag{3.4}$$

where $\psi(s)$ is the digamma function. In the last line, we have used

$$\lim_{s \rightarrow 0} \frac{\psi(s)}{\Gamma(s)} = -1 \tag{3.5}$$

and $1/\Gamma(0) = 0$. The notation $*$ in (3.4) stands for an appropriate regularization that subtracts the divergent terms of the integrand as $\beta \rightarrow 0$. Alternatively, the Laplace transform of the supersymmetric index is equal to the log-derivative of the supersymmetric determinant

$$-\frac{\partial}{\partial z} \log \mathbb{D}(z) = \int_0^\infty d\beta \, \mathbb{I}(\beta) e^{-\beta z}^*. \tag{3.6}$$

3.3 Vacuum exponents

When the variable z is set to zero, one obtains the special value of the supersymmetric determinants $\mathfrak{D}(z)$ at $z = 0$

$$\mathfrak{D}(0) = \exp[\mathfrak{Z}(0, 0)'] = \prod_n n^{-\delta(n)}. \tag{3.7}$$

One of the important aspects of this value lies in its role in the asymptotic formulae of the supersymmetric indices. It appears in the Cardy-like limit (2.47) and the asymptotic degeneracy (2.64) as a constant term. We refer to it as the *vacuum exponent*. These asymptotic formulae are only meaningful if the vacuum exponent is a non-zero finite value. When the value $\mathfrak{D}(0)$ is zero, the asymptotic coefficient is always zero, while when the value $\mathfrak{D}(0)$ approaches infinity, the asymptotic coefficient diverges. We propose that

$$\begin{cases} \text{SUSY breaking vacuum} & \rightarrow & \mathfrak{D}(0) = 0 \\ \text{no stable vacuum} & \rightarrow & \mathfrak{D}(0) = \infty \end{cases} \tag{3.8}$$

In other words, if supersymmetry is broken, then the vacuum exponent will vanish, although the converse is not necessarily true. Similarly, if there is no normalizable vacuum due to the non-compact degrees of freedom, then the vacuum exponent will diverge.

3.4 Functional equations

We make a further remark that certain transformations and relations satisfied by the supersymmetric indices may reduce to functional equations for the supersymmetric determinants.

Suppose that the supersymmetric index satisfies the relation

$$\mathbb{I}(q^{-1}) = Cq^D \mathbb{I}(q), \quad (3.9)$$

with $C = \pm 1$ and $D \in \mathbb{R}$. Such a transformation is called the T -reflection [16] (also see [15]) as it corresponds to a reflection of the temperature. It is then tempting to explore a reflection relation for the supersymmetric determinant that relates $\mathbb{D}(z)$ to $\mathbb{D}(D - z)$. If we define a function

$$\mathbb{C}(z) := \mathbb{D}(D - z)^C \mathbb{D}(z)^{-1}, \quad (3.10)$$

this function will be crucial for understanding the behavior and analytic properties of the supersymmetric determinant, such as its poles and zeros, by relating its values on either side of the line $\text{Re}(z) = D/2$.

As an example, let us consider the half-BPS index (2.15) of 4d $\mathcal{N} = 4$ $U(N)$ SYM theory. It obeys

$$\mathcal{I}_{\frac{1}{2}\text{BPS}}^{4\text{d } \mathcal{N}=4 \text{ } U(N)}(q^{-1}) = (-1)^N q^{\frac{N(N+1)}{2}} \mathcal{I}_{\frac{1}{2}\text{BPS}}^{4\text{d } \mathcal{N}=4 \text{ } U(N)}(q). \quad (3.11)$$

The supersymmetric determinant is given by

$$\begin{aligned} \mathbb{D}_{\frac{1}{2}\text{BPS}}^{4\text{d } \mathcal{N}=4 \text{ } U(N)}(z) &= \Gamma_N(z; 1, 2, \dots, N) \\ &= \prod_{n_1, \dots, n_N \geq 0} \frac{1}{(n_1 + 2n_2 + \dots + Nn_N + z)}, \end{aligned} \quad (3.12)$$

where

$$\Gamma_r(z; \omega_1, \dots, \omega_r) := \exp \left[\left. \frac{\partial}{\partial s} \zeta_r(s, z; \omega_1, \dots, \omega_r) \right|_{s=0} \right] \quad (3.13)$$

is the Barnes multiple gamma function [36]. We have

$$\begin{aligned}
\mathbb{C}_{\frac{1}{2}\text{BPS}}^{4\text{d } \mathcal{N}=4 \text{ } U(N)}(z) &= \Gamma_N(z; 1, 2, \dots, N)^{-1} \Gamma_N\left(\frac{N(N+1)}{2} - z; 1, 2, \dots, N\right)^{(-1)^N} \\
&= S_N(z; 1, 2, \dots, N),
\end{aligned} \tag{3.14}$$

where

$$S_r(z; \omega_1, \omega_2, \dots, \omega_r) = \Gamma_r(z; \omega_1, \omega_2, \dots, \omega_r)^{-1} \Gamma_r(\omega_1 + \dots + \omega_r - z; \omega_1, \omega_2, \dots, \omega_r)^{(-1)^r} \tag{3.15}$$

is the multiple sine function [74, 75, 76]. Thus, we have the reflection formula

$$\mathbb{C}_{\frac{1}{2}\text{BPS}}^{4\text{d } \mathcal{N}=4 \text{ } U(N)}(z) = \mathbb{D}_{\frac{1}{2}\text{BPS}}^{4\text{d } \mathcal{N}=4 \text{ } U(N)}(D - z)^C \mathbb{D}_{\frac{1}{2}\text{BPS}}^{4\text{d } \mathcal{N}=4 \text{ } U(N)}(z)^{-1}, \tag{3.16}$$


where $D = N(N+1)/2$ and $C = (-1)^N$. When $N = 1$, this reduces to

$$\begin{aligned}
S_1(z; \omega_1) &= \Gamma_1(1 - z; \omega_1)^{-1} \Gamma_1(z; \omega_1)^{-1} \\
&= \frac{2\pi}{\Gamma\left(\frac{z}{\omega_1}\right) \Gamma\left(1 - \frac{z}{\omega_1}\right)} \\
&= 2 \sin\left(\frac{\pi z}{\omega_1}\right).
\end{aligned} \tag{3.17}$$

This is nothing but the Euler reflection formula

$$\Gamma(z)\Gamma(1 - z) = \frac{\pi}{\sin(\pi z)}. \tag{3.18}$$

To conclude this section, we have identified the following related families of supersymmetric spectral functions for these theories: supersymmetric indices, supersymmetric zeta functions, and supersymmetric determinants.



For theories whose degeneracies $d(n)$ may not grow polynomially, we take their plethystic logarithm $\mathbb{I}(q)$ (whose coefficients $\delta(n)$ may themselves grow polynomially) to define the supersymmetric zeta function $\mathfrak{Z}(s, z)$ and the supersymmetric determinant $\mathfrak{D}(z)$. In the following sections, we illustrate these functions by considering a variety of 2d, 4d, and 6d supersymmetric field theories.

4 2d supersymmetric field theories

4.1 Zeta functions for 2d $\mathcal{N} = (0, 2)$ elliptic genera

The elliptic genera for 2d $\mathcal{N} = (0, 2)$ supersymmetric field theories in the Ramond-Ramond (RR) sector and Neveu-Schwarz-Neveu-Schwarz (NSNS) sector are defined by

$$\mathcal{I}_{\text{R}}^{2\text{d}, (0, 2)}(x; q) = \text{Tr}_{\text{R}}(-1)^F q^{H_L} \bar{q}^{H_R} \prod_{\alpha} x_{\alpha}^{f_{\alpha}}, \quad (4.1)$$

$$\mathcal{I}_{\text{NS}}^{2\text{d}, (0, 2)}(x; q) = \text{Tr}_{\text{NS}}(-1)^F q^{H_L} \bar{q}^{H_R - \frac{R}{2}} \prod_{\alpha} x_{\alpha}^{f_{\alpha}}. \quad (4.2)$$

Here $H_L = (H + iP)/2$ and $H_R = (H - iP)/2$ are the left- and right-moving Hamiltonians in the Euclidean signature in terms of the Hamiltonian H and the momentum P , respectively. R stands for the $U(1)_R$ R-charge and f_{α} are the Cartan generators of other global symmetries. While only right-moving ground states with $H_R = 0$ contribute to the elliptic genus in the RR sector, right-moving chiral primary states with $H_R = R/2$ contribute to the elliptic genus in the NSNS sector. A prescription for computing the matrix integrals of the elliptic genera of 2d $\mathcal{N} = (0, 2)$ supersymmetric gauge theories was given in [77, 78].

One has the 4-form anomaly polynomial of the form (see e.g. [79])

$$\mathcal{A}_4 = \frac{1}{2} k_{RR} c_1(R)^2 - \frac{1}{24} k p_1(T), \quad (4.3)$$

where $c_1(R)$ is the first Chern class of the $U(1)_R$ background gauge field and $p_1(T)$ is the first Pontryagin class of the tangent bundle T of the two-dimensional spacetime. The 't Hooft anomaly coefficient k_{RR} of the $U(1)_R$ R-symmetry can be computed as

$$k_{RR} = \text{Tr} \gamma_3 U(1)_R^2, \quad (4.4)$$

where γ_3 is the two-dimensional chirality matrix that takes value $+1$ on the right-handed fermions and -1 on the left-handed fermions. For the superconformal field theories, the right-moving central charge c_R is related to the 't Hooft anomaly for the

$U(1)_R$ R-symmetry

$$c_R = 3k_{RR}. \quad (4.5)$$

The gravitational anomaly coefficient k is given by

$$k = \text{Tr } \gamma_3. \quad (4.6)$$

For the superconformal field theories, the left-moving central charge c_L can be determined by the gravitational anomaly, which yields the relation:

$$c_R - c_L = k. \quad (4.7)$$

We define the supersymmetric zeta functions $\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(s, z)$ for the 2d $\mathcal{N} = (0, 2)$ supersymmetric field theories by the Mellin transform (2.7) of the plethystic logarithms of the elliptic genera in the NSNS sector. We also define from (3.1) the supersymmetric determinants $\mathfrak{D}_{\text{NS}}^{2\text{d } (0,2)}(z)$ associated with the supersymmetric zeta functions $\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(s, z)$. The supersymmetric zeta functions $\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(s, z)$ can have a simple pole at $s = 1$.

According to the asymptotic formulae (2.47) and (2.64), we obtain

$$\log \mathcal{I}_{\text{NS}}^{2\text{d } (0,2)}(\beta) \sim \frac{\pi^2 \text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(s, 0)}{6\beta} - \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(0, 0) \log \beta + \log \mathfrak{D}_{\text{NS}}^{2\text{d } (0,2)}(0), \quad (4.8)$$

$$\begin{aligned} d_{\text{NS}}^{2\text{d } (0,2)}(n) &\sim \left(\frac{\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(s, 0)}{96n^3} \right)^{\frac{1}{4}} \left(\frac{6n}{\pi^2 \text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(s, 0)} \right)^{\frac{\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(0,0)}{2}} \\ &\times \mathfrak{D}_{\text{NS}}^{2\text{d } (0,2)}(0) \exp \left[2\pi \sqrt{\frac{\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(s, 0)}{6}} n^{\frac{1}{2}} \right], \end{aligned} \quad (4.9)$$

where $d_{\text{NS}}^{2\text{d } (0,2)}(n)$ are the expansion coefficients of the elliptic genus in the NSNS sector

$$\mathcal{I}_{\text{NS}}^{2\text{d } (0,2)}(x; q) = \sum_n d_{\text{NS}}^{2\text{d } (0,2)}(n) q^n. \quad (4.10)$$

For the superconformal field theories, we observe that the residues at a simple pole $s = 1$ of the supersymmetric zeta functions $\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(s, z)$ are given by the gravitational anomaly coefficients

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(s, 0) = 2k = 2(c_R - c_L). \quad (4.11)$$

Then the formulae (4.8) and (4.9) reduce to⁵

$$\log \mathcal{I}_{\text{NS}}^{2\text{d } (0,2)}(\beta) \sim \frac{\pi^2(c_R - c_L)}{3\beta} - \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(0,0) \log \beta + \log \mathfrak{D}_{\text{NS}}^{2\text{d } (0,2)}(0), \quad (4.12)$$

$$\begin{aligned} d_{\text{NS}}^{2\text{d } (0,2)}(n) &\sim \left(\frac{c_R - c_L}{48n^3} \right)^{\frac{1}{4}} \left(\frac{3n}{\pi^2(c_R - c_L)} \right)^{\frac{\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(0,0)}{2}} \\ &\times \mathfrak{D}_{\text{NS}}^{2\text{d } (0,2)}(0) \exp \left[2\pi \sqrt{\frac{c_R - c_L}{3}} n^{\frac{1}{2}} \right]. \end{aligned} \quad (4.13)$$

From the various examples we have examined, we find that the supersymmetric zeta values for $s = -1$ and $z = 0$ are given by

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(-1,0) = -\frac{c_L}{12}. \quad (4.14)$$

This universal structure, determined by the left-moving central charge c_L , is compatible with the supersymmetric Casimir energies of the 2d $\mathcal{N} = (0,2)$ theories.

For the superconformal field theories, the central charges are obtainable from (4.11) and (4.14) in terms of the supersymmetric zeta functions

$$c_L = -12\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(-1,0), \quad (4.15)$$

$$c_R = \frac{1}{2} \text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(s,0) - 12\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(-1,0). \quad (4.16)$$

The 4-form anomaly coefficients in (4.3) are given by

$$k = \frac{1}{2} \text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(s,0), \quad (4.17)$$

$$k_{RR} = \frac{1}{6} \text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(s,0) - 4\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2)}(-1,0). \quad (4.18)$$

4.1.1 2d $\mathcal{N} = (0,2)$ chiral multiplet

The elliptic genus of the 2d $\mathcal{N} = (0,2)$ chiral multiplet of R-charge r reads

$$\mathcal{I}_{\text{NS}}^{2\text{d } (0,2) \text{ chiral}_r}(x;q) = \frac{1}{(q^{\frac{r}{2}}x;q)_{\infty}(q^{1-\frac{r}{2}}x^{-1};q)_{\infty}}. \quad (4.19)$$

The single-particle index is

$$i_{\text{NS}}^{2\text{d } (0,2) \text{ chiral}_r}(q) = \frac{q^{\frac{r}{2}}x}{1-q} + \frac{q^{1-\frac{r}{2}}x^{-1}}{1-q}. \quad (4.20)$$

The supersymmetric zeta function is given by

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ chiral}_r}(s,z) = \zeta\left(s, \frac{r}{2} + z\right) + \zeta\left(s, 1 - \frac{r}{2} + z\right), \quad (4.21)$$

⁵A similar asymptotic formula is found in Appendix A of [44].

where

$$\zeta(s, a) = \sum_{n=0}^{\infty} \frac{1}{(n+a)^s} \quad (4.22)$$

is the Hurwitz zeta function. The supersymmetric zeta function (4.21) has a simple pole at $s = 1$ with residue

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ chiral}_r}(s, z) = 2. \quad (4.23)$$

The Zeta-index vanishes

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ chiral}_r}(0, 0) = 0. \quad (4.24)$$

The supersymmetric zeta value with $s = -1$ and $z = 0$ is given by

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ chiral}_r}(-1, 0) = -\frac{3(r-1)^2 - 1}{12}. \quad (4.25)$$

Plugging (4.23) and (4.25) into the formulae (4.17) and (4.18), we obtain the gravitational and 't Hooft anomalies

$$k = 1, \quad (4.26)$$

$$k_{RR} = (r-1)^2. \quad (4.27)$$

When the R-charge is chosen as the superconformal anomaly, they agree with the central charge of the superconformal field theories due to the 't Hooft anomaly matching. From (4.15) and (4.16), we find

$$c_L = 3(r-1)^2 - 1, \quad (4.28)$$

$$c_R = 3(r-1)^2. \quad (4.29)$$

They agree with the anomalies contributed from the 2d $\mathcal{N} = (0, 2)$ chiral multiplet of R-charge r [44].

The supersymmetric determinant is

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (0,2) \text{ chiral}_r}(z) = \frac{\Gamma\left(\frac{r}{2} + z\right) \Gamma\left(1 - \frac{r}{2} + z\right)}{2\pi}. \quad (4.30)$$

The vacuum exponent is

$$\begin{aligned} \mathfrak{D}_{\text{NS}}^{2\text{d } (0,2) \text{ chiral}_r}(0) &= \frac{\Gamma\left(\frac{r}{2}\right) \Gamma\left(1 - \frac{r}{2}\right)}{2\pi} \\ &= \frac{1}{2 \sin\left(\frac{\pi r}{2}\right)}. \end{aligned} \quad (4.31)$$




Figure 1: The vacuum exponent $\mathfrak{D}_{\text{NS}}^{2d (0,2) \text{ chiral}_r}(0)$.

It is shown in Figure 1. The vacuum exponent has a simple pole at $r = 0$, for which we have a theory of non-compact free bosons with the non-normalizable vacuum. The degeneracy $d^{2d (0,2) \text{ chiral}_r}(n)$ obeys the exponential growth for $0 < r < 2$. We obtain from (4.13) the asymptotic growth

$$d^{2d (0,2) \text{ chiral}_{r=1}}(n) \sim \frac{1}{3^{\frac{1}{4}} \cdot 4n^{\frac{3}{4}} \sin\left(\frac{\pi r}{2}\right)} \exp\left[\frac{2\pi}{3^{\frac{1}{2}}} n^{\frac{1}{2}}\right]. \quad (4.32)$$

For example, for $r = 1$ we show the exact numbers $d_{\text{exact}}^{2d (0,2) \text{ chiral}_{r=1}}(n)$ and the values $d_{\text{asyp}}^{2d (0,2) \text{ chiral}_{r=1}}(n)$ evaluated from (4.32) in the following:

n	$d_{\text{exact}}^{2d (0,2) \text{ chiral}_{r=1}}(n)$	$d_{\text{asyp}}^{2d (0,2) \text{ chiral}_{r=1}}(n)$
10	1598	3240
100	1.70092×10^{13}	3.41288×10^{13}
1000	3.5254×10^{46}	7.05732×10^{46}

(4.33)

As discussed in section 2.4.3, the asymptotic formula (4.32) generally involves ambiguity in the constant term. The numerical computation suggests that the ratio of the estimated value to the actual value approaches 2 as n becomes large.

4.1.2 2d $\mathcal{N} = (0, 2)$ Fermi multiplet

The elliptic genus of the 2d $\mathcal{N} = (0, 2)$ Fermi multiplet of R-charge r is given by

$$\mathcal{I}_{\text{NS}}^{2d (0,2) \text{ Fermi}_r}(x; q) = (q^{\frac{1+r}{2}} x; q)_{\infty} (q^{\frac{1-r}{2}} x^{-1}; q)_{\infty}. \quad (4.34)$$

The single-particle index is

$$i_{\text{NS}}^{2\text{d } (0,2) \text{ Fermi}_r}(x; q) = -\frac{q^{\frac{1+r}{2}} x}{1-q} - \frac{q^{\frac{1-r}{2}} x^{-1}}{1-q}. \quad (4.35)$$

The supersymmetric zeta function is

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ Fermi}_r}(s, z) = -\zeta\left(s, \frac{1+r}{2} + z\right) - \zeta\left(s, \frac{1-r}{2} + z\right). \quad (4.36)$$

It has a simple pole at $s = 1$ with residue

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ Fermi}_r}(s, z) = -2. \quad (4.37)$$

The Zeta-index vanishes

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ Fermi}_r}(0, 0) = 0. \quad (4.38)$$

The supersymmetric zeta value with $s = -1$ and $z = 0$ is

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ Fermi}_r}(-1, 0) = \frac{3r^2 - 1}{12}. \quad (4.39)$$

Substituting (4.37) and (4.39) into the formulae (4.17) and (4.18), we get the gravitational and 't Hooft anomalies

$$k = -1, \quad (4.40)$$

$$k_{RR} = -r^2. \quad (4.41)$$

When the theory is superconformal we find from (4.15) and (4.16) the central charges

$$c_L = 1 - 3r^2, \quad (4.42)$$

$$c_R = -3r^2. \quad (4.43)$$

They are the expected contributions from the $\mathcal{N} = (0, 2)$ Fermi multiplet of R-charge r [44].

The supersymmetric determinant is

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (0,2) \text{ Fermi}_r}(z) = \frac{2\pi}{\Gamma\left(\frac{1+r}{2} + z\right) \Gamma\left(\frac{1-r}{2} + z\right)}. \quad (4.44)$$

The vacuum exponent is

$$\begin{aligned} \mathfrak{D}_{\text{NS}}^{2\text{d } (0,2) \text{ Fermi}_r}(0) &= \frac{2\pi}{\Gamma\left(\frac{1+r}{2}\right) \Gamma\left(\frac{1-r}{2}\right)} \\ &= 2 \cos\left(\frac{\pi r}{2}\right). \end{aligned} \quad (4.45)$$

We show the vacuum exponent in Figure 2.




Figure 2: The vacuum exponent $\mathfrak{D}_{\text{NS}}^{2\text{d } (0,2) \text{ Fermi}_r}(0)$.

4.1.3 2d $\mathcal{N} = (0, 2)$ vector multiplet

The elliptic genus of the 2d $\mathcal{N} = (0, 2)$ $U(1)$ vector multiplet is

$$\mathcal{I}_{\text{NS}}^{2\text{d } (0,2) \text{ vec}}(q) = (q)_{\infty}^2. \quad (4.46)$$

The single-particle index is

$$i_{\text{NS}}^{2\text{d } (0,2) \text{ vec}}(q) = -\frac{2q}{1-q}. \quad (4.47)$$

We have the supersymmetric zeta function

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ vec}}(s, z) = -2\zeta(s, 1+z). \quad (4.48)$$

It has a simple pole at $s = 1$ with residue

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ vec}}(s, z) = -2. \quad (4.49)$$

The Zeta-index is

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ vec}}(0, 0) = 1. \quad (4.50)$$

The supersymmetric zeta value with $s = -1$ and $z = 0$ is

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ vec}}(-1, 0) = \frac{1}{6}. \quad (4.51)$$

According to the formulae (4.17) and (4.18) as well as the residue (4.49) and the zeta value (4.51), we get the gravitational and 't Hooft anomalies

$$k = -1, \quad (4.52)$$

$$k_{RR} = -1. \quad (4.53)$$

Correspondingly, we have

$$c_L = -2, \quad (4.54)$$

$$c_R = -3. \quad (4.55)$$

which are identified with the expected contributions from the 2d $\mathcal{N} = (0, 2)$ $U(1)$ vector multiplet [44].

The supersymmetric determinant is

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (0, 2) \text{ vec}}(z) = \frac{2\pi}{[\Gamma(1+z)]^2}. \quad (4.56)$$

The vacuum exponent is

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (0, 2) \text{ vec}}(0) = 2\pi. \quad (4.57)$$

4.1.4 2d $\mathcal{N} = (0, 2)$ SQED with chirals $(\Phi, \bar{\Phi}_a)$ and Fermis (Ψ_α, Γ_a)

Consider the 2d $\mathcal{N} = (0, 2)$ SQED studied in [80, 44], that is a $U(1)$ gauge theory with a chiral multiplet Φ of charge +1, $N_f - 1$ chiral multiplets $\bar{\Phi}_a$ of charge -1 , N_f Fermi multiplets Ψ_α of charge -1 and $N_f - 1$ neutral Fermi multiplets Γ_a . The theory is conjectured to be dual to a theory of N_f free Fermi multiplets Λ'_α [44]. The field content is summarized as

	$U(1)$	$SU(N_f)$	$SU(N_f - 1)$	$U(1)_R$
Φ	+1	1	1	0
$\bar{\Phi}_a$	-1	1	N_f - 1	$\frac{N_f - 2}{N_f - 1}$
Ψ_α	-1	N_f	1	0
Γ_a	0	1	N_f - 1	$\frac{1}{N_f - 1}$
Λ'_α	0	N_f	1	0

(4.58)

The elliptic genus of the $U(1)$ gauge theory in the NSNS sector is evaluated as

$$\begin{aligned} & \mathcal{I}_{\text{NS}}^{2\text{d } (0, 2) \text{ SQED } + (\Phi, \bar{\Phi}_a; \Psi_\alpha, \Gamma_a)}(x_\alpha; q) \\ &= (q)_\infty^2 \oint_{\text{JK}} \frac{ds}{2\pi i s} \frac{1}{(s; q)_\infty (qs^{-1}; q)_\infty} \prod_{\alpha=1}^{N_f} (q^{\frac{1}{2}} s^\pm x_\alpha^\mp; q)_\infty \\ & \times \prod_{a=1}^{N_f-1} \frac{(q^{\frac{1}{2} + \frac{1}{2(N_f-1)}} y_a^{-1}; q)_\infty (q^{\frac{1}{2} - \frac{1}{2(N_f-1)}} y_a; q)_\infty}{(q^{\frac{1}{2} - \frac{1}{2(N_f-1)}} s^{-1} y_a; q)_\infty (q^{\frac{1}{2} + \frac{1}{2(N_f-1)}} s y_a^{-1}; q)_\infty}. \end{aligned} \quad (4.59)$$

Here the contour integral \oint_{JK} is evaluated by applying the Jeffery-Kirwan (JK) residue prescription [81]. It agrees with the elliptic genus of N_f free Fermi multiplets

$$\mathcal{I}_{\text{NS}}^{2\text{d } (0, 2) \Lambda'_\alpha}(x_\alpha; q) = \prod_{\alpha=1}^{N_f} (q^{\frac{1}{2}} x_\alpha^\pm; q)_\infty. \quad (4.60)$$

The single-particle index is

$$i_{\text{NS}}^{2\text{d } (0,2) \text{ SQED } +(\Phi, \bar{\Phi}_a; \Psi_\alpha, \Gamma_a)}(x_\alpha; q) = - \sum_{\alpha=1}^{N_f} \frac{q^{\frac{1}{2}}(x_\alpha + x_\alpha^{-1})}{1 - q}. \quad (4.61)$$

The supersymmetric zeta function is

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ SQED } +(\Phi, \bar{\Phi}_a; \Psi_\alpha, \Gamma_a)}(s, z) = -2N_f \zeta\left(s, \frac{1}{2} + z\right). \quad (4.62)$$

It has a simple pole at $s = 1$ with residue

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ SQED } +(\Phi, \bar{\Phi}_a; \Psi_\alpha, \Gamma_a)}(s, z) = -2N_f. \quad (4.63)$$

As the theory is dual to the free Fermi multiplets, the Zeta-index vanishes

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ SQED } +(\Phi, \bar{\Phi}_a; \Psi_\alpha, \Gamma_a)}(0, 0) = 0. \quad (4.64)$$

The supersymmetric zeta value with $s = -1$ and $z = 0$ is

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ SQED } +(\Phi, \bar{\Phi}_a; \Psi_\alpha, \Gamma_a)}(-1, 0) = -\frac{N_f}{12}. \quad (4.65)$$

From the residue (4.63) and the zeta value (4.65) we find the anomalies

$$k = -N_f, \quad (4.66)$$

$$k_{RR} = 0, \quad (4.67)$$

and

$$c_L = N_f, \quad (4.68)$$

$$c_R = 0, \quad (4.69)$$

which agree with the central charges of the 2d $\mathcal{N} = (0, 2)$ SQED in [44].

The supersymmetric determinant is

$$\mathfrak{D}^{2\text{d } (0,2) \text{ SQED } +(\Phi, \bar{\Phi}_a; \Psi_\alpha, \Gamma_a)}(z) = \frac{(2\pi)^{N_f}}{\Gamma\left(\frac{1}{2} + z\right)^{2N_f}}. \quad (4.70)$$

The vacuum exponent is

$$\mathfrak{D}^{2\text{d } (0,2) \text{ SQED } +(\Phi, \bar{\Phi}_a; \Psi_\alpha, \Gamma_a)}(0) = 2^{N_f}. \quad (4.71)$$

4.1.5 2d $\mathcal{N} = (0, 2)$ $U(N_c)$ with chirals $(\Phi_s, \bar{\Phi}_a)$ and Fermis $(\Psi_\alpha, \Omega, \Gamma_{s,a})$

The 2d $\mathcal{N} = (0, 2)$ $U(N_c)$ gauge theory with N_c fundamental chiral multiplets Φ_s , $N_f + N_c$ antifundamental chiral multiplets $\bar{\Phi}_a$, N_f antifundamental Fermi multiplets Ψ_α , 2 determinant Fermi multiplets $\Omega_{1,2}$ and $N_f + N_c$ neutral Fermi multiplets $\Gamma_{s,a}$ is conjectured to be dual to the theory of $N_c N_f + 2$ free fermions $\Gamma'_{\alpha,s}$ and $\Omega'_{1,2}$ [44]. The field content and the charges are

	$U(N_c)$	$SU(N_f)$	$SU(N_c)'$	$SU(N_f + N_c)$	$SU(2)$	$U(1)_R$
Φ_s	\mathbf{N}_c	$\mathbf{1}$	$\overline{\mathbf{N}_c}$	$\mathbf{1}$	$\mathbf{1}$	0
$\bar{\Phi}_a$	$\overline{\mathbf{N}_c}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{N}_f + \mathbf{N}_c$	$\mathbf{1}$	$\frac{N_f}{N_f + N_c}$
Ψ_α	$\overline{\mathbf{N}_c}$	\mathbf{N}_f	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	0
$\Omega_{1,2}$	det	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{2}$	0
$\Gamma_{s,a}$	$\mathbf{1}$	$\mathbf{1}$	\mathbf{N}_c	$\overline{\mathbf{N}_f + \mathbf{N}_c}$	$\mathbf{1}$	$\frac{N_c}{N_f + N_c}$
$\Gamma'_{\alpha,s}$	$\mathbf{1}$	\mathbf{N}_f	$\overline{\mathbf{N}_c}$	$\mathbf{1}$	$\mathbf{1}$	0
$\Omega'_{1,2}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{2}$	0

The elliptic genus of the $U(N_c)$ gauge theory is evaluated as

$$\begin{aligned}
& \mathcal{I}_{\text{NS}}^{2d (0,2) U(N_c) + (\Phi_s, \bar{\Phi}_a; \Psi_\alpha, \Omega, \Gamma_{s,a})}(x_\alpha, u_s, v; q) \\
&= (q)^{2N_c} \oint_{\text{JK}} \prod_{i=1}^{N_c} \frac{ds_i}{2\pi i s_i} \prod_{i \neq j} (s_i^\pm s_j^\mp; q)_\infty (q s_i^\pm s_j^\mp; q)_\infty \prod_{i=1}^{N_c} \prod_{\alpha=1}^{N_f} (q^{\frac{1}{2}} s_i^\mp x_\alpha^\mp; q)_\infty \\
&\times \prod_{i=1}^{N_c} \prod_{s=1}^{N_c} \frac{1}{(s_i u_s^{-1}; q)_\infty (q s_i^{-1} u_s; q)_\infty} \prod_{a=1}^{N_f + N_c} \frac{\prod_{s=1}^{N_c} (q^{\frac{1}{2} \pm \frac{N_c}{2(N_f + N_c)}} u_s^\pm y_a^\mp; q)_\infty}{\prod_{i=1}^{N_c} (q^{\frac{N_f}{2(N_c + N_f)}} s_i^{-1} y_a; q)_\infty (q^{1 - \frac{N_f}{2(N_c + N_f)}} s_i y_a^{-1}; q)_\infty} \\
&\times (q^{\frac{1}{2}} (\prod_{i=1}^{N_c} s_i v)^\pm; q)_\infty (q^{\frac{1}{2}} (\prod_{i=1}^{N_c} s_i v^{-1})^\pm; q)_\infty.
\end{aligned} \tag{4.73}$$

It is shown [44] that the expression (4.73) is equal to the elliptic genus

$$\mathcal{I}_{\text{NS}}^{2d (0,2) \Gamma'_{\alpha,s} + \Omega'}(x_\alpha, u_s, v; q) = (q^{\frac{1}{2}} v^\pm; q)_\infty^2 \prod_{\alpha=1}^{N_f} \prod_{s=1}^{N_c} (q^{\frac{1}{2}} x_\alpha^\pm u_s^\mp; q)_\infty \tag{4.74}$$

of the theory of $N_c N_f + 2$ free fermions. The single-particle index reads

$$\begin{aligned}
& \mathcal{I}_{\text{NS}}^{2d (0,2) U(N_c) + (\Phi_s, \bar{\Phi}_a; \Psi_\alpha, \Omega, \Gamma_{s,a})}(x_\alpha, u_s, w; q) \\
&= -\frac{2q^{\frac{1}{2}}(v + v^{-1})}{1 - q} - \sum_{\alpha=1}^{N_f} \sum_{s=1}^{N_c} \frac{q^{\frac{1}{2}}(x_\alpha u_s^{-1} + x_\alpha^{-1} u_s)}{1 - q}.
\end{aligned} \tag{4.75}$$

The supersymmetric zeta function is given by

$$\mathfrak{Z}_{\text{NS}}^{2d (0,2) U(N_c) + (\Phi_s, \bar{\Phi}_a; \Psi_\alpha, \Omega, \Gamma_{s,a})}(s, z) = -2(N_c N_f + 2)\zeta\left(s, \frac{1}{2} + z\right). \tag{4.76}$$

The residue at a simple pole $s = 1$ is

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \ U(N_c) + (\Phi_s, \bar{\Phi}_a; \Psi_\alpha, \Omega, \Gamma_{s,a})}(s, z) = -4 - 2N_c N_f. \quad (4.77)$$

For $s = 0$ and $z = 0$ we have

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \ U(N_c) + (\Phi_s, \bar{\Phi}_a; \Psi_\alpha, \Omega, \Gamma_{s,a})}(0, 0) = 0. \quad (4.78)$$

When $s = -1$ and $z = 0$ we get

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \ U(N_c) + (\Phi_s, \bar{\Phi}_a; \Psi_\alpha, \Omega, \Gamma_{s,a})}(-1, 0) = -\frac{1}{6} - \frac{N_c N_f}{12}. \quad (4.79)$$

According to the formulae (4.17) and (4.18) we get the anomalies

$$k = -2 - N_c N_f, \quad (4.80)$$

$$k_{RR} = 0. \quad (4.81)$$

Also from (4.15) and (4.16), we find

$$c_L = 2 + N_c N_f, \quad (4.82)$$

$$c_R = 0. \quad (4.83)$$

They agree with the central charges obtained in [44].

The supersymmetric determinant is

$$\mathfrak{D}^{2\text{d } (0,2) \ U(N_c) + (\Phi_s, \bar{\Phi}_a; \Psi_\alpha, \Omega, \Gamma_{s,a})}(z) = \frac{(2\pi)^{N_c N_f + 2}}{\Gamma(\frac{1}{2} + z)^{2(N_c N_f + 2)}}. \quad (4.84)$$

The vacuum exponent is

$$\mathfrak{D}^{2\text{d } (0,2) \ U(N_c) + (\Phi_s, \bar{\Phi}_a; \Psi_\alpha, \Omega, \Gamma_{s,a})}(0) = 2^{N_c N_f + 2}. \quad (4.85)$$

4.1.6 2d $\mathcal{N} = (0, 2)$ $SU(2)$ with 4 chirals Φ_α

Consider the 2d $\mathcal{N} = (0, 2)$ $SU(2)$ gauge theory with 4 fundamental chiral multiplets Φ_α . The theory is conjectured to be dual to the Landau-Ginzburg (LG) theory with the Fermi multiplet Ψ' and 6 chiral multiplets Φ' interacting with the J-type Pfaffian superpotential [82, 83, 84]

$$\mathcal{W} = \Psi' \text{Pf}(\Phi'). \quad (4.86)$$

The R-charge of the J-term superpotential is required to be equal to +1 since the fermionic measure $d\theta^+$ has the R-charge -1 . The field content is summarized as

	$SU(2)$	$SU(4)$	$U(1)_x$	$U(1)_R$
Φ_α	2	4	+1	0
Φ'	1	6	+2	0
Ψ'	1	1	-4	+1

(4.87)

The elliptic genus of the $SU(2)$ gauge theory reads

$$\begin{aligned}
& \mathcal{I}_{\text{NS}}^{2\text{d } (0,2) \text{ } SU(2) + \Phi_\alpha}(x_\alpha, x; q) \\
&= \frac{1}{2}(q)_\infty^2 \oint \frac{ds}{2\pi i s} (s^{\pm 2}; q)_\infty (qs^{\pm 2}; q)_\infty \\
&\times \prod_{\alpha=1}^4 \frac{1}{(sx_\alpha x; q)_\infty (qs^{-1}x_\alpha^{-1}x^{-1}; q)_\infty (s^{-1}x_\alpha x; q)_\infty (qsx_\alpha^{-1}x^{-1}; q)_\infty}.
\end{aligned} \tag{4.88}$$

It is equal to the elliptic genus of the dual LG theory

$$\mathcal{I}_{\text{NS}}^{2\text{d } (0,2) \text{ } \Phi' + \Psi'}(x_\alpha, x; q) = (x^4; q)_\infty (qx^{-4}; q)_\infty \prod_{\alpha < \beta}^4 \frac{1}{(x_\alpha x_\beta x^2; q)_\infty (qx_\alpha^{-1}x_\beta^{-1}x^{-2}; q)_\infty}. \tag{4.89}$$

The single-particle index is

$$\begin{aligned}
& i_{\text{NS}}^{2\text{d } (0,2) \text{ } SU(2) + \Phi_\alpha}(x_\alpha, x; q) \\
&= -\frac{x^4}{1-q} - \frac{x^{-4}q}{1-q} + \sum_{\alpha < \beta}^4 \left[\frac{x_\alpha x_\beta x^2}{1-q} + \frac{x_\alpha^{-1}x_\beta^{-1}x^{-2}q}{1-q} \right].
\end{aligned} \tag{4.90}$$

When we turn off the $SU(4)$ flavor fugacities x_α and set $x = q^{R_x}$, we get

$$i_{\text{NS}}^{2\text{d } (0,2) \text{ } SU(2) + \Phi_\alpha}(x_\alpha = 1, x = q^{R_x}; q) = -\frac{q^{4R_x}}{1-q} - \frac{q^{1-4R_x}}{1-q} + \frac{6q^{2R_x}}{1-q} + \frac{6q^{1-2R_x}}{1-q}. \tag{4.91}$$

The supersymmetric zeta function is evaluated as

$$\begin{aligned}
\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ } SU(2) + \Phi_\alpha}(s, z) &= -\zeta(s, 4R_x + z) - \zeta(s, 1 - 4R_x + z) \\
&+ 6 [\zeta(s, 2R_x + z) + \zeta(s, 1 - 2R_x + z)].
\end{aligned} \tag{4.92}$$

It has a simple pole at $s = 1$ with residue

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ } SU(2) + \Phi_\alpha}(s, z) = 10. \tag{4.93}$$

The Zeta-index vanishes

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ } SU(2) + \Phi_\alpha}(0, 0) = 0. \tag{4.94}$$

For $s = -1$ and $z = 0$ the supersymmetric zeta value is evaluated as

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ } SU(2) + \Phi_\alpha}(-1, 0) = -\left(8R_x^2 - 8R_x + \frac{5}{6}\right). \tag{4.95}$$

From (4.93) and (4.95) we obtain the trial central charges

$$c_L = 96R_x^2 - 96R_x + 10, \quad (4.96)$$

$$c_R = 96R_x^2 - 96R_x + 15. \quad (4.97)$$

Applying the c -extremization [85, 79], one finds $R_x = 1/2$, which leads to $c_L = -14$ and $c_R = -9$. As the negative values violate the unitarity bound, they indicate the failure of the naive application of the c -extremization. It was proposed by Sacchi [86] that the correct central charges can be obtained by choosing the vanishing mixing coefficients $R_x = 0$ for the associated flavor currents arising from non-compact directions in the moduli space. Then one finds

$$c_L = 10, \quad (4.98)$$

$$c_R = 15. \quad (4.99)$$

We have the supersymmetric determinant

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (0,2) \text{ } SU(2) + \Phi_\alpha}(z) = \frac{\Gamma(z)^5 \Gamma(1+z)^5}{(2\pi)^5}. \quad (4.100)$$

The vacuum exponent is

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (0,2) \text{ } SU(2) + \Phi_\alpha}(0) = \infty. \quad (4.101)$$

This would imply that the theory has the non-compact free bosons.

4.1.7 2d $\mathcal{N} = (0, 2)$ $SU(N)$ with $2N$ chirals $(\Phi_\alpha, \bar{\Phi}_\beta)$

Consider the $SU(N)$ gauge theory with N fundamental chiral multiplets Φ_α and N antifundamental chiral multiplets $\bar{\Phi}_\beta$. It is argued [82, 83] that the theory is dual to the LG theory of N^2 chiral multiplets Φ' , 2 chiral multiplets (B', \bar{B}') and the Fermi multiplet Ψ' with a superpotential

$$\mathcal{W} = \Psi'(B'\bar{B}' + \det \Phi'). \quad (4.102)$$

The field content and charges are given by

	$SU(N)$	$SU(N)_x$	$SU(N)_y$	$U(1)_x$	$U(1)_y$	$U(1)_R$
Φ_α	\mathbf{N}	$\bar{\mathbf{N}}$	$\mathbf{1}$	+1	0	0
$\bar{\Phi}_\beta$	$\bar{\mathbf{N}}$	$\mathbf{1}$	\mathbf{N}	0	+1	0
Φ'	$\mathbf{1}$	$\bar{\mathbf{N}}$	\mathbf{N}	+1	+1	0
B'	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	+ N	0	0
\bar{B}'	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	0	+ N	0
Ψ'	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	- N	- N	+1

(4.103)

The elliptic genus of the $SU(N)$ gauge theory reads

$$\begin{aligned}
& \mathcal{I}_{\text{NS}}^{2d (0,2) SU(N) + (\Phi_\alpha, \bar{\Phi}_\beta)}(x_\alpha, y_\beta, x, y; q) \\
&= \frac{1}{N!} (q)_\infty^{N-1} \oint_{\text{JK}} \prod_{i=1}^{N-1} \frac{ds_i}{2\pi i s_i} \prod_{i < j} (s_i^\pm s_j^\mp; q)_\infty (q s_i^\pm s_j^\mp; q)_\infty \\
&\times \prod_{i=1}^N \prod_{\alpha=1}^N \frac{1}{(s_i x_\alpha^{-1} x; q)_\infty (q s_i^{-1} x_\alpha x^{-1}; q)_\infty} \prod_{i=1}^N \prod_{\beta=1}^N \frac{1}{(s_i^{-1} y_\beta y; q)_\infty (q s_i y_\beta^{-1} y^{-1}; q)_\infty}, \quad (4.104)
\end{aligned}$$

with $\prod_{i=1}^N s_i = 1$, $\prod_\alpha x_\alpha = 1$ and $\prod_\beta y_\beta = 1$. As discussed in [82], the expression agrees with

$$\begin{aligned}
& \mathcal{I}_{\text{NS}}^{2d (0,2) \Phi' + (B' \bar{B}') + \Psi'}(x_\alpha, y_\beta, x, y; q) \\
&= \frac{(q x^{-N} y^{-N}; q)_\infty (x^N y^N; q)_\infty}{(x^N; q)_\infty (q x^{-N}; q)_\infty (y^N; q)_\infty (q y^{-N}; q)_\infty \prod_{\alpha=1}^N \prod_{\beta=1}^N (x_\alpha^{-1} y_\beta x y; q)_\infty (q x_\alpha y_\beta^{-1} x^{-1} y^{-1}; q)_\infty}, \quad (4.105)
\end{aligned}$$

which is the elliptic genus of the dual LG theory. The single-particle index is

$$\begin{aligned}
& i_{\text{NS}}^{2d (0,2) SU(N) + (\Phi_\alpha, \bar{\Phi}_\beta)}(x_\alpha, y_\beta, x, y; q) \\
&= -\frac{x^{-N} y^{-N} q}{1-q} - \frac{x^N y^N}{1-q} + \frac{x^N}{1-q} + \frac{x^{-N} q}{1-q} + \frac{y^N}{1-q} + \frac{y^{-N} q}{1-q} \\
&+ \sum_{\alpha=1}^N \sum_{\beta=1}^N \left[\frac{x_\alpha^{-1} y_\beta x y}{1-q} + \frac{x_\alpha y_\beta^{-1} x^{-1} y^{-1} q}{1-q} \right]. \quad (4.106)
\end{aligned}$$

Setting $x_\alpha = y_\beta = 1$, $x = q^{R_x}$ and $y = q^{R_y}$, it reduces to

$$\begin{aligned}
& i_{\text{NS}}^{2d (0,2) SU(N) + (\Phi_\alpha, \bar{\Phi}_\beta)}(x_\alpha = 1, y_\beta = 1, x = q^{R_x}, y = q^{R_y}; q) \\
&= -\frac{q^{1-N(R_x+R_y)}}{1-q} - \frac{q^{N(R_x+R_y)}}{1-q} + \frac{q^{NR_x}}{1-q} + \frac{q^{1-NR_x}}{1-q} + \frac{q^{NR_y}}{1-q} + \frac{q^{1-NR_y}}{1-q} \\
&+ N^2 \left(\frac{q^{R_x+R_y}}{1-q} + \frac{q^{1-R_x-R_y}}{1-q} \right). \quad (4.107)
\end{aligned}$$

The supersymmetric zeta function is evaluated as

$$\begin{aligned}
& \mathfrak{Z}_{\text{NS}}^{2d (0,2) SU(N) + (\Phi_\alpha, \bar{\Phi}_\beta)}(s, z) \\
&= -\zeta(s, 1 - N(R_x + R_y) + z) - \zeta(s, N(R_x + R_y) + z) \\
&+ \zeta(s, NR_x + z) + \zeta(s, 1 - NR_x + z) + \zeta(s, NR_y + z) + \zeta(s, 1 - NR_y + z) \\
&+ N^2 [\zeta(s, R_x + R_y + z) + \zeta(s, 1 - R_x - R_y + z)]. \quad (4.108)
\end{aligned}$$

The residue at a simple pole $s = 1$ is

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2d (0,2) SU(N) + (\Phi_\alpha, \bar{\Phi}_\beta)}(s, z) = 2(N^2 + 1). \quad (4.109)$$

The special supersymmetric zeta value for $s = 0$ and $z = 0$ vanishes

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ } SU(N) + (\Phi_\alpha, \bar{\Phi}_\beta)}(0, 0) = 0. \quad (4.110)$$

For $s = -1$ and $z = 0$ the supersymmetric zeta value is

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ } SU(N) + (\Phi_\alpha, \bar{\Phi}_\beta)}(-1, 0) = -N^2(R_x^2 - R_x) - N^2(R_y^2 - R_y) - \frac{N^2}{6} - \frac{1}{6}. \quad (4.111)$$

Hence we get the trial central charges

$$c_L = 12(N^2 R_x^2 - N^2 R_x + N^2 R_y^2 - N^2 R_y) + 2(N^2 + 1), \quad (4.112)$$

$$c_R = 12(N^2 R_x^2 - N^2 R_x + N^2 R_y^2 - N^2 R_y) + 3(N^2 + 1). \quad (4.113)$$

Upon the c -extremization [85, 79], we get $R_x = R_y = 1/2$, which leads to the negative central charges $c_L = 2 - 4N^2$ and $c_R = 3 - 3N^2$. Again the correct central charges will be obtained by requiring no mixing anomaly coefficients [86]. Then we get

$$c_L = 2(N^2 + 1), \quad (4.114)$$

$$c_R = 3(N^2 + 1). \quad (4.115)$$

When $N = 2$, they reproduce the central charges (4.98) and (4.99) for the $SU(2)$ gauge theory with 4 fundamental chirals, as expected.

The supersymmetric determinant is

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (0,2) \text{ } SU(N) + (\Phi_\alpha, \bar{\Phi}_\beta)}(z) = \frac{\Gamma(z)^{N^2+1} \Gamma(1+z)^{N^2+1}}{(2\pi)^{N^2+1}}. \quad (4.116)$$

The vacuum exponent is

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (0,2) \text{ } SU(N) + (\Phi_\alpha, \bar{\Phi}_\beta)}(0) = \infty. \quad (4.117)$$

Again, this would imply that the theory has the non-compact free bosons.

4.1.8 2d $\mathcal{N} = (0, 2)$ $USp(2N)$ with $2N + 2$ chirals Φ_α

Next consider the $USp(2N)$ gauge theory with $2N + 2$ fundamental chiral multiplets that can be viewed as an alternative higher rank generalization of the $SU(2)$ gauge theory with 4 fundamental chiral multiplets. It is conjectured [83, 86] that the theory is dual to the LG model of a single Fermi multiplet Ψ' and $(N + 1)(2N + 1)$ chiral multiplets Φ' with a superpotential

$$\mathcal{W} = \Psi' \text{Pf} \Phi'. \quad (4.118)$$

The field content and the charges are given by

	$USp(2N)$	$SU(2N+2)$	$U(1)_x$	$U(1)_R$
Φ_α	2N	2N+2	+1	0
Φ'	1	(N+1)(2N+1)	+2	0
Ψ'	1	1	$-2(N+1)$	+1

(4.119)

The duality between these two theories results from the dimensional reduction of the Intriligator-Pouliot duality [87] for 4d $\mathcal{N} = 1$ gauge theories. When $N = 1$, the theory reduces to the $SU(2)$ gauge theory with 4 fundamental chirals.⁶

The elliptic genus of the $USp(2N)$ gauge theory reads

$$\begin{aligned}
& \mathcal{I}_{\text{NS}}^{2\text{d } (0,2) \text{ } USp(2N) + \Phi_\alpha}(x_\alpha, x; q) \\
&= \frac{1}{2^N N!} (q)_\infty^{2N} \oint_{\text{JK}} \prod_{i=1}^N \frac{ds_i}{2\pi i s_i} (s_i^{\pm 2}; q)_\infty (q s_i^{\pm 2}; q)_\infty \\
&\times \prod_{i < j} (s_i^\pm s_j^\mp; q)_\infty (s_i^\pm s_j^\pm; q)_\infty (q s_i^\pm s_j^\mp; q)_\infty (q s_i^\pm s_j^\pm; q)_\infty \\
&\times \prod_{i=1}^N \prod_{\alpha=1}^{2N+2} \frac{1}{(s_i x_\alpha x; q)_\infty (q s_i^{-1} x_\alpha^{-1} x^{-1}; q)_\infty (s_i^{-1} x_\alpha x; q)_\infty (q s_i x_\alpha^{-1} x^{-1}; q)_\infty}.
\end{aligned} \tag{4.120}$$

It coincides with the elliptic genus for the dual LG model of the form

$$\mathcal{I}_{\text{NS}}^{2\text{d } (0,2) \text{ } \Phi' + \Psi'}(x_\alpha, x; q) = \frac{(q x^{-2(N+1)}; q)_\infty (x^{2(N+1)}; q)_\infty}{\prod_{\alpha < \beta}^{2N+2} (x_\alpha x_\beta x^2; q)_\infty (q x_\alpha^{-1} x_\beta^{-1} x^{-2}; q)_\infty}. \tag{4.121}$$

The single-particle index takes the form

$$\begin{aligned}
& i_{\text{NS}}^{2\text{d } (0,2) \text{ } USp(2N) + \Phi_\alpha}(x_\alpha, x; q) \\
&= -\frac{x^{-2(N+1)} q}{1-q} - \frac{x^{2(N+1)}}{1-q} + \sum_{\alpha < \beta}^{2N+2} \left[\frac{x_\alpha x_\beta x^2}{1-q} + \frac{x_\alpha^{-1} x_\beta^{-1} x^{-2} q}{1-q} \right].
\end{aligned} \tag{4.122}$$

With $x_\alpha = 1$ and $x = q^{R_x}$, it becomes

$$\begin{aligned}
& i_{\text{NS}}^{2\text{d } (0,2) \text{ } USp(2N) + \Phi_\alpha}(x_\alpha = 1, x = q^{R_x}; q) \\
&= -\frac{q^{1-2(N+1)R_x}}{1-q} - \frac{q^{2(N+1)R_x}}{1-q} + (2N+1)(N+1) \left[\frac{q^{2R_x}}{1-q} + \frac{q^{1-2R_x}}{1-q} \right].
\end{aligned} \tag{4.123}$$

⁶In addition, it is conjectured in [88] that these theories are dual to the $SU(2N)$ gauge theory with an antisymmetric chiral, $2N+2$ fundamental chirals and a neutral Fermi interacting with a J-term superpotential.

One finds the supersymmetric zeta function

$$\begin{aligned} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ } USp(2N) + \Phi_\alpha}(s, z) \\ = -\zeta(s, 1 - 2(N+1)R_x + z) - \zeta(s, 2(N+1)R_x + z) \\ + (2N+1)(N+1) [\zeta(s, 2R_x + z) + \zeta(s, 1 - 2R_x + z)]. \end{aligned} \quad (4.124)$$

The residue at a simple pole $s = 1$ is

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ } USp(2N) + \Phi_\alpha}(s, z) = 2N(2N+3). \quad (4.125)$$

The special supersymmetric zeta value with $s = 0$ and $z = 0$ vanishes

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ } USp(2N) + \Phi_\alpha}(0, 0) = 0. \quad (4.126)$$

The special supersymmetric zeta value with $s = -1$ and $z = 0$ is

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ } USp(2N) + \Phi_\alpha}(-1, 0) = -4N(N+1)R_x^2 + 4N(N+1)R_x - \frac{1}{6}N(2N+3). \quad (4.127)$$

Accordingly, we obtain the trial central charges

$$c_L = 48N(N+1)R_x^2 - 48N(N+1)R_x + 2N(2N+3), \quad (4.128)$$

$$c_R = 48N(N+1)R_x^2 - 48N(N+1)R_x + 3N(2N+3). \quad (4.129)$$

By extremizing the trial central charges according to the c -extremization [85, 79], we get $R_x = 1/2$. However, for this value the trial central charges become negative, $c_L = -2N(4N+3)$ and $c_R = -3N(2N+1)$. Following the proposal in [86], the correct central charges will be obtained by simply setting the mixing coefficient R_x to zero. Then the expressions (4.128) and (4.129) reduce to

$$c_L = 2N(2N+3), \quad (4.130)$$

$$c_R = 3N(2N+3). \quad (4.131)$$

The supersymmetric determinant is

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (0,2) \text{ } USp(2N) + \Phi_\alpha}(z) = \frac{\Gamma(z)^{N(2N+3)} \Gamma(1+z)^{N(2N+3)}}{(2\pi)^{N(2N+3)}}. \quad (4.132)$$

As the theory has the non-compact free bosons the vacuum exponent is divergent

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (0,2) \text{ } USp(2N) + \Phi_\alpha}(0) = \infty. \quad (4.133)$$

4.1.9 2d $\mathcal{N} = (0, 2)$ $USp(2N)$ with chirals (Φ_α, A) and Fermis Γ_a

Another type of higher rank $USp(2N)$ gauge theories which generalizes the $SU(2)$ gauge theory with 4 fundamental chiral multiplets can be obtained by introducing the chiral multiplet A transforming in the antisymmetric representation. When the theory has 4 fundamental chiral multiplets Φ_α , an antisymmetric chiral multiplet A and N neutral Fermi multiplets Γ_a with a superpotential

$$\mathcal{W} = \sum_{a=1}^N \Gamma_a \text{Tr } A^a, \quad (4.134)$$

where the contractions are performed using the $USp(2N)$ invariant antisymmetric tensor $J = i\sigma_2 \otimes \mathbb{I}_N$, it is conjectured to be dual to the LG model of $6N$ chiral multiplets $\Phi'_a{}^{\alpha\beta}$ and N Fermi multiplets Ψ'_a with a superpotential [86]

$$\mathcal{W} = \sum_{a,b,c=1}^N \sum_{\alpha,\beta,\gamma,\delta=1}^4 \epsilon_{\alpha\beta\gamma\delta} \Psi'_a \Phi'_b{}^{\alpha\beta} \Phi'_c{}^{\gamma\delta} \delta_{a+b+c, 2N+1}. \quad (4.135)$$

The field content and charges are summarized as

	$USp(2N)$	$SU(4)$	$U(1)_w$	$U(1)_x$	$U(1)_R$
Φ_α	$2N$	4	$\frac{1-N}{3}$	$+1$	0
A	$N(2N-1)$	1	$+1$	0	0
Γ_a	1	1	$-a$	0	$+1$
Φ_a	1	6	$a - \frac{2N+1}{3}$	$+2$	0
Ψ_a	1	1	$a - \frac{2N+1}{3}$	-4	$+1$

(4.136)

with $a = 1, \dots, N$.

The elliptic genus of the $USp(2N)$ gauge theory reads

$$\begin{aligned} \mathcal{I}_{\text{NS}}^{2d(0,2)USp(2N)\Phi_\alpha+A+\Gamma_a}(x_\alpha, w, x; q) &= \frac{1}{2^N N!} (q)_\infty^{2N} \oint_{\text{JK}} \prod_{i=1}^N \frac{ds_i}{2\pi i s_i} (s_i^{\pm 2}; q)_\infty (qs_i^{\pm 2}; q)_\infty \\ &\times \prod_{i < j} (s_i^\pm s_j^\mp; q)_\infty (s_i^\pm s_j^\pm; q)_\infty (qs_i^\pm s_j^\mp; q)_\infty (qs_i^\pm s_j^\pm; q)_\infty \\ &\times \prod_{i=1}^N \prod_{\alpha=1}^4 \frac{1}{(s_i x_\alpha w^{\frac{1-N}{3}} x; q)_\infty (qs_i^{-1} x_\alpha^{-1} w^{\frac{N-1}{3}} x^{-1}; q)_\infty (s_i^{-1} x_\alpha w^{\frac{1-N}{3}} x; q)_\infty (qs_i x_\alpha^{-1} w^{\frac{N-1}{3}} x^{-1}; q)_\infty} \\ &\times \prod_{i,j=1}^N \frac{1}{(s_i s_j^{-1} w; q)_\infty (qs_i^{-1} s_j w; q)_\infty} \prod_{i < j} \frac{1}{(s_i^\pm s_j^\pm w; q)_\infty (qs_i^\mp s_j^\mp w^{-1}; q)_\infty} \\ &\times \prod_{a=1}^N (qw^{-a}; q)_\infty (w^a; q)_\infty. \end{aligned} \quad (4.137)$$

It agrees with

$$\begin{aligned} & \mathcal{I}_{\text{NS}}^{2\text{d } (0,2) \Phi'_a + \Psi'_a}(x_\alpha, w, x; q) \\ &= \prod_{i=1}^N \frac{(x^4 w^{-i + \frac{2N+1}{3}}; q)_\infty (q x^{-4} w^{i - \frac{2N+1}{3}}; q)_\infty}{\prod_{\alpha < \beta}^4 (x^2 w^{i - \frac{2N+1}{3}} x_\alpha x_\beta; q)_\infty (q x^{-2} w^{-i + \frac{2N+1}{3}} x_\alpha^{-1} x_\beta^{-1}; q)_\infty}. \end{aligned} \quad (4.138)$$

With $x_\alpha = 1$, $w = q^{R_w}$ and $x = q^{R_x}$, the single-particle index reads

$$\begin{aligned} & i_{\text{NS}}^{2\text{d } (0,2) \text{ } USp(2N) \Phi_\alpha + A + \Gamma_a}(x_\alpha = 1, w = q^{R_w}, x = q^{R_x}; q) \\ &= \sum_{i=1}^N \left[-\frac{q^{4R_x - (i - \frac{2N+1}{3})R_w}}{1 - q} - \frac{q^{1 - 4R_x + (i - \frac{2N+1}{3})R_w}}{1 - q} \right. \\ & \quad \left. + \frac{6q^{2R_x + (i - \frac{2N+1}{3})R_w}}{1 - q} + \frac{6q^{1 - 2R_x - (i - \frac{2N+1}{3})R_w}}{1 - q} \right]. \end{aligned} \quad (4.139)$$

The supersymmetric zeta function is evaluated as

$$\begin{aligned} & \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ } USp(2N) \Phi_\alpha + A + \Gamma_a}(s, z) \\ &= \sum_{i=1}^N \left[-\zeta \left(s, 4R_x - (i - \frac{2N+1}{3})R_w + z \right) - \zeta \left(s, 1 - 4R_x + (i - \frac{2N+1}{3})R_w + z \right) \right. \\ & \quad \left. + 6\zeta \left(s, 2R_x + (i - \frac{2N+1}{3})R_w + z \right) + 6\zeta \left(s, 1 - 2R_x - (i - \frac{2N+1}{3})R_w + z \right) \right]. \end{aligned} \quad (4.140)$$

The residue at a simple pole $s = 1$ is

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ } USp(2N) \Phi_\alpha + A + \Gamma_a}(s, z) = 10N. \quad (4.141)$$

For $s = 0$ and $z = 0$ the supersymmetric zeta value vanishes

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ } USp(2N) \Phi_\alpha + A + \Gamma_a}(0, 0) = 0. \quad (4.142)$$

The supersymmetric zeta value with $s = -1$ and $z = 0$ is given by

$$\begin{aligned} & \mathfrak{Z}_{\text{NS}}^{2\text{d } (0,2) \text{ } USp(2N) \Phi_\alpha + A + \Gamma_a}(-1, 0) \\ &= \sum_{i=1}^N \left[-\frac{5}{9}(2N - 3i + 1)^2 R_w^2 - 8R_x^2 + \frac{32}{3}(2N - 3i + 1)R_w R_x \right. \\ & \quad \left. - \frac{7}{3}(2N - 3i + 1)R_w + 8R_x - \frac{5}{6} \right] \\ &= -\frac{5}{18}N(2N^2 - N - 1)R_w^2 - 8NR_x^2 + \frac{16}{3}N(N - 1)R_w R_x \\ & \quad - \frac{7}{6}N(N - 1)R_w + 8NR_x - \frac{5}{6}N. \end{aligned} \quad (4.143)$$

Hence the trial central charges are

$$c_L = \frac{10}{3}N(2N^2 - N - 1)R_w^2 + 96NR_x^2 - 64N(N - 1)R_wR_x \\ + 14N(N - 1)R_w - 96NR_x + 10N, \quad (4.144)$$

$$c_R = \frac{10}{3}N(2N^2 - N - 1)R_w^2 + 96NR_x^2 - 64N(N - 1)R_wR_x \\ + 14N(N - 1)R_w - 96NR_x + 15N. \quad (4.145)$$

When $N = 1$, they agree with (4.96) and (4.97). Applying the c -extremization, one finds that the mixing coefficients are given by

$$R_w = -\frac{9}{2(2N - 7)}, \quad R_x = -\frac{N + 4}{2(2N - 7)}. \quad (4.146)$$

Then the values of the central charges are

$$c_L = \frac{5N(5N + 23)}{2(2N - 7)}, \quad (4.147)$$

$$c_R = \frac{45N(N + 1)}{2(2N - 7)}. \quad (4.148)$$

Note that they are negative for $N = 1, 2$ and 3 . Again the prescription for the failure of the c -extremization will be given by the vanishing two mixing coefficients R_w and R_x [86]. Consequently, we arrive at

$$c_L = 10N, \quad (4.149)$$

$$c_R = 15N. \quad (4.150)$$

The supersymmetric determinant is

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (0,2) \text{ } USp(2N) \text{ } \Phi_\alpha + A + \Gamma_a}(z) = \frac{\Gamma(z)^{5N}\Gamma(1+z)^{5N}}{(2\pi)^{5N}}. \quad (4.151)$$

We find the divergent vacuum exponent

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (0,2) \text{ } USp(2N) \text{ } \Phi_\alpha + A + \Gamma_a}(0) = \infty. \quad (4.152)$$

4.2 Zeta functions for 2d $\mathcal{N} = (2, 2)$ elliptic genera

The 2d $\mathcal{N} = (2, 2)$ supersymmetric field theories have a left-moving $U(1)$ R-symmetry J_L and a right-moving $U(1)$ R-symmetry J_R . They are related to the vectorial (R_V) and axial (R_A) R-symmetries as follows:

$$R_V = J_L + J_R, \quad R_A = J_L - J_R. \quad (4.153)$$

The elliptic genera for 2d $\mathcal{N} = (2, 2)$ supersymmetric field theories can be defined in a similar way:

$$\mathcal{I}_{\text{R}}^{2\text{d } (2,2)}(y, x; q) = \text{Tr}_{\text{R}}(-1)^F q^{H_L} \bar{q}^{H_R} y^{J_L} \prod_{\alpha} x_{\alpha}^{f_{\alpha}}, \quad (4.154)$$

$$\mathcal{I}_{\text{NS}}^{2\text{d } (2,2)}(y, x; q) = \text{Tr}_{\text{NS}}(-1)^F q^{H_L} \bar{q}^{H_R - \frac{J_R}{2}} y^{J_L} \prod_{\alpha} x_{\alpha}^{f_{\alpha}}. \quad (4.155)$$

The elliptic genus in the RR sector and that in the NSNS sector are related by spectral flow:

$$\mathcal{I}_{\text{R}}^{2\text{d } (2,2)}(y, x; q) = y^{-\frac{c}{6}} \mathcal{I}_{\text{NS}}^{2\text{d } (2,2)}(q^{-\frac{1}{2}} y, x; q), \quad (4.156)$$

where $c = c_L = c_R$.

Similarly, we define the supersymmetric zeta functions $\mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2)}(s, z)$ by the Mellin transform(2.7) of the plethystic logarithms of the NSNS sector elliptic genera with $y = x = 1$, and the associated supersymmetric determinants $\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2)}(z)$ by (3.1).

4.2.1 2d $\mathcal{N} = (2, 2)$ chiral multiplet

Consider the $\mathcal{N} = (2, 2)$ chiral multiplet of the vectorial R-charge $2r$ and of the vanishing axial R-charge. It has the left-moving R-charge r . The elliptic genus in the NSNS sector is given by

$$\begin{aligned} \mathcal{I}_{\text{NS}}^{2\text{d } (2,2) \text{ chiral}_r}(y, x; q) &= \mathcal{I}_{\text{NS}}^{2\text{d } (0,2) \text{ chiral}_r}(y^r x; q) \times \mathcal{I}_{\text{NS}}^{2\text{d } (0,2) \text{ Fermi}_r}(y^{r-1} x; q) \\ &= \frac{(q^{\frac{1}{2} + \frac{r}{2}} y^{r-1} x; q)_{\infty} (q^{\frac{1}{2} - \frac{r}{2}} y^{1-r} x^{-1}; q)_{\infty}}{(q^{\frac{r}{2}} y^r x; q)_{\infty} (q^{1 - \frac{r}{2}} y^{-r} x^{-1}; q)_{\infty}}. \end{aligned} \quad (4.157)$$

The single-particle index is

$$\begin{aligned} i_{\text{NS}}^{2\text{d } (2,2) \text{ chiral}_r}(y, x; q) &= i_{\text{NS}}^{2\text{d } (0,2) \text{ chiral}_r}(y^r x; q) + i_{\text{NS}}^{2\text{d } (0,2) \text{ Fermi}_r}(y^{r-1} x; q) \\ &= \frac{q^{\frac{r}{2}} y^r x}{1 - q} + \frac{q^{1 - \frac{r}{2}} y^{-r} x^{-1}}{1 - q} - \frac{q^{\frac{1+r}{2}} y^{r-1} x}{1 - q} - \frac{q^{\frac{1-r}{2}} y^{-r+1} x^{-1}}{1 - q}. \end{aligned} \quad (4.158)$$

The supersymmetric zeta function is evaluated as

$$\begin{aligned} \mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ chiral}_r}(s, z) &= \mathfrak{Z}_{\text{NS}}^{(0,2) \text{ chiral}_r}(s, z) + \mathfrak{Z}_{\text{NS}}^{(0,2) \text{ Fermi}_r}(s, z) \\ &= \zeta\left(s, \frac{r}{2} + z\right) + \zeta\left(s, 1 - \frac{r}{2} + z\right) - \zeta\left(s, \frac{1+r}{2} + z\right) - \zeta\left(s, \frac{1-r}{2} + z\right). \end{aligned} \quad (4.159)$$

While each term has a simple pole at $s = 1$, the total sum of the residues vanishes

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ chiral}_r}(s, z) = 0. \quad (4.160)$$

Also the Zeta-index vanishes

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ chiral}_r}(0,0) = 0. \quad (4.161)$$

For $s = -1$ and $z = 0$ the supersymmetric zeta value is

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ chiral}_r}(-1,0) = \frac{2r-1}{4}. \quad (4.162)$$

Plugging (4.160) and (4.162) into (4.17) and (4.18), we find the gravitational and 't Hooft anomalies

$$k = 0, \quad (4.163)$$

$$k_{RR} = 1 - 2r. \quad (4.164)$$

From the formulae (4.15) and (4.16), we get

$$c = c_L = c_R = 3(1 - 2r). \quad (4.165)$$

This is the expected anomaly contribution from the $\mathcal{N} = (2,2)$ chiral multiplet [78] for generic R-symmetry assignment.

The supersymmetric determinant is

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ chiral}_r}(z) = \frac{\Gamma\left(\frac{r}{2} + z\right) \Gamma\left(1 - \frac{r}{2} + z\right)}{\Gamma\left(\frac{1+r}{2} + z\right) \Gamma\left(\frac{1-r}{2} + z\right)}. \quad (4.166)$$

The vacuum exponent is

$$\begin{aligned} \mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ chiral}_r}(0) &= \frac{\Gamma\left(\frac{r}{2}\right) \Gamma\left(1 - \frac{r}{2}\right)}{\Gamma\left(\frac{1+r}{2}\right) \Gamma\left(\frac{1-r}{2}\right)} \\ &= \cot\left(\frac{\pi r}{2}\right), \end{aligned} \quad (4.167)$$

which is shown in Figure 3.

Similarly, we obtain from the expression (4.157) the elliptic genus of the $\mathcal{N} = (2,2)$ twisted chiral multiplet by replacing y with y^{-1}

$$\mathcal{I}_{\text{NS}}^{2\text{d } (2,2) \text{ tchiral}_r}(y, x; q) = \mathcal{I}_{\text{NS}}^{2\text{d } (2,2) \text{ chiral}_r}(y^{-1}, x; q). \quad (4.168)$$

Hence the supersymmetric zeta function associated with the unflavored index takes the same form as (4.159) and therefore the anomaly contribution is given by (4.165).




Figure 3: The vacuum exponent $\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ chiral}_r}(0)$.

4.2.2 2d $\mathcal{N} = (2, 2)$ vector multiplet

The $\mathcal{N} = (2, 2)$ $U(1)$ vector multiplet can be viewed as the twisted chiral multiplet of the left-moving R-charge $r = +1$.

The elliptic genus in the NSNS sector is given by combining the indices of the $\mathcal{N} = (0, 2)$ vector multiplet and the $\mathcal{N} = (0, 2)$ chiral multiplet of R-charge $+1$

$$\mathcal{I}_{\text{NS}}^{2\text{d } (2,2) \text{ vec}}(y; q) = \frac{(q)_\infty^2}{(q^{\frac{1}{2}}y; q)_\infty (q^{\frac{1}{2}}y^{-1}; q)_\infty}. \quad (4.169)$$

The single-particle index of the $\mathcal{N} = (2, 2)$ $U(1)$ vector multiplet

$$i_{\text{NS}}^{2\text{d } (2,2) \text{ vec}}(y; q) = -\frac{2q}{1-q} + \frac{(y + y^{-1})q^{\frac{1}{2}}}{1-q}. \quad (4.170)$$

The supersymmetric zeta function is given by

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ vec}}(s, z) = -2\zeta(s, 1+z) + 2\zeta\left(s, \frac{1}{2} + z\right). \quad (4.171)$$

The residue at $s = 1$ is

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ vec}}(s, z) = 0. \quad (4.172)$$

The Zeta-index is given by

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ vec}}(0, 0) = 1. \quad (4.173)$$

The supersymmetric zeta value with $s = -1$ and $z = 0$ is

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ vec}}(-1, 0) = \frac{1}{4}. \quad (4.174)$$

Substituting (4.172) and (4.174) into (4.17) and (4.18), we get the anomalies

$$k = 0, \quad (4.175)$$

$$k_{RR} = -1. \quad (4.176)$$

Also we obtain from (4.15) and (4.16)

$$c = c_L = c_R = -3. \quad (4.177)$$

This is the expected anomaly contribution from the $\mathcal{N} = (2, 2)$ $U(1)$ vector multiplet [78].

We find the supersymmetric determinant

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ vec}}(z) = \frac{\Gamma\left(\frac{1}{2} + z\right)^2}{\Gamma(1 + z)^2}. \quad (4.178)$$

The vacuum exponent is

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ vec}}(0) = \pi. \quad (4.179)$$

4.2.3 2d $\mathcal{N} = (2, 2)$ pure SYM theories

Now consider the $\mathcal{N} = (2, 2)$ pure SYM theory of gauge group G . It is conjectured [89] that when G is semisimple and simply-connected, the theory flows to a free theory of $\text{rank}(G)$ twisted chiral multiplets with the left-moving R-charges being the degrees d_i , $i = 1, \dots, \text{rank}(G)$ of the Casimir operators at low energy (see e.g. [90, 91, 92, 93] for further discussions). The degrees of the Casimirs are given by (see e.g. [94])

G	d_i	
$SU(N)$	$2, 3, \dots, N$	
$SO(2N + 1)$	$2, 4, \dots, 2N$	
$USp(2N)$	$2, 4, \dots, 2N$	
$SO(2N)$	$2, 4, \dots, 2(N - 1), N$	
G_2	$2, 6$	
F_4	$2, 6, 8, 12$	
E_6	$2, 5, 6, 8, 9, 12$	
E_7	$2, 6, 8, 10, 12, 14, 18$	
E_8	$2, 8, 12, 14, 18, 20, 24, 30$	(4.180)

For example, for $G = SU(N)$ it is dual to the theory of $(N - 1)$ free twisted chiral multiplets of the left-moving R-charges n with $n = 2, \dots, N$.

The elliptic genus in the NSNS sector can be evaluated by the matrix integral

$$\begin{aligned} & \mathcal{I}_{\text{NS}}^{2d(2,2)G}(y; q) \\ &= \frac{1}{|\text{Weyl}(G)|} \frac{(q)_\infty^{2\text{rank}(G)}}{(q^{\frac{1}{2}} y^\pm; q)_\infty^{\text{rank}(G)}} \oint_{\text{JK}} \prod_{\alpha \in \text{root}(G)} \frac{ds}{2\pi i s} \frac{(s^\alpha; q)_\infty (qs^\alpha; q)_\infty}{(q^{\frac{1}{2}} s^\alpha y; q)_\infty (q^{\frac{1}{2}} s^{-\alpha} y^{-1}; q)_\infty}, \end{aligned} \quad (4.181)$$

where $|\text{Weyl}(G)|$ is the order of the Weyl group of G . According to the dualities, it is equal to

$$\mathcal{I}_{\text{NS}}^{2d(2,2)G}(y; q) = \prod_{i=1}^{\text{rank}(G)} \frac{(q^{\frac{1+d_i}{2}} y^{1-d_i}; q)_\infty (q^{\frac{1-d_i}{2}} y^{-1+d_i}; q)_\infty}{(q^{\frac{d_i}{2}} y^{-d_i}; q)_\infty (q^{1-\frac{d_i}{2}} y^{d_i}; q)_\infty} \quad (4.182)$$

and the single-particle index is given by

$$\begin{aligned} & i_{\text{NS}}^{2d(2,2)G}(y; q) \\ &= \sum_{i=1}^{\text{rank}(G)} \left[-\frac{y^{1-d_i} q^{\frac{1+d_i}{2}}}{1-q} - \frac{y^{-1+d_i} q^{\frac{1-d_i}{2}}}{1-q} + \frac{y^{-d_i} q^{\frac{d_i}{2}}}{1-q} + \frac{y^{d_i} q^{1-\frac{d_i}{2}}}{1-q} \right]. \end{aligned} \quad (4.183)$$

The supersymmetric zeta function is evaluated as

$$\begin{aligned} & \mathfrak{Z}_{\text{NS}}^{2d(2,2)G}(s, z) \\ &= \sum_{i=1}^{\text{rank}(G)} \left[-\zeta\left(s, \frac{1+d_i}{2} + z\right) - \zeta\left(s, \frac{1-d_i}{2} + z\right) + \zeta\left(s, \frac{d_i}{2} + z\right) + \zeta\left(s, 1 - \frac{d_i}{2} + z\right) \right]. \end{aligned} \quad (4.184)$$

It follows that

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2d(2,2)G}(s, z) = 0, \quad (4.185)$$

$$\mathfrak{Z}_{\text{NS}}^{2d(2,2)G}(0, 0) = 0, \quad (4.186)$$

$$\mathfrak{Z}_{\text{NS}}^{2d(2,2)G}(-1, 0) = \frac{\dim G}{4}, \quad (4.187)$$

where we have used the fact that

$$\dim G = \sum_{i=1}^{\text{rank}(G)} (2d_i - 1). \quad (4.188)$$

According to the formulae (4.17), (4.18), (4.15) and (4.16), we find the anomalies

$$k = 0, \quad (4.189)$$

$$k_{RR} = -\dim G \quad (4.190)$$

and

$$c = c_L = c_R = -3 \dim G. \quad (4.191)$$

While the values (4.191) should not be understood as the ordinary central charges, they are the anomaly contributions of the $\mathcal{N} = (2, 2)$ vector multiplet of gauge group G [78]. They are also called the effective central charges in [93]. The pure SYM theory of gauge group G can be viewed as a sigma model on the algebraic stack $BG = [\text{point}/G]$ [95, 96, 97, 98]. The dimension of the quotient algebraic stack $[X/G]$ is given by [95]

$$\dim[X/G] = \dim X - \dim G, \quad (4.192)$$

which can be negative. For the stack $BG = [\text{point}/G]$, the dimension is $-\dim G$, which matches with $c/3$.

The supersymmetric determinant is

$$\mathfrak{D}_{\text{NS}}^{2d(2,2)G}(z) = \prod_{i=1}^{\text{rank}(G)} \frac{\Gamma(\frac{d_i}{2} + z) \Gamma(1 - \frac{d_i}{2} + z)}{\Gamma(\frac{1+d_i}{2} + z) \Gamma(\frac{1-d_i}{2} + z)}. \quad (4.193)$$

The vacuum exponent takes the form

$$\mathfrak{D}_{\text{NS}}^{2d(2,2)G}(0) = \prod_{i=1}^{\text{rank}(G)} \cot\left(\frac{\pi d_i}{2}\right). \quad (4.194)$$

It contains divergent factors associated with the degrees d_i of the Casimirs being even integers. This indicates the non-compact degrees of freedom.⁷

4.2.4 2d $\mathcal{N} = (2, 2)$ minimal models

Let us consider the $\mathcal{N} = (2, 2)$ minimal models with $c < 3$ which can be described by the LG theories of twisted chiral multiplets $\tilde{\Phi}_i$ with a quasi-homogeneous twisted superpotential $\tilde{\mathcal{W}}$. They are the rational conformal field theories with an ADE classification of the twisted superpotentials given by

minimal models	$\tilde{\mathcal{W}}$
A_{k+1}	$\tilde{\Phi}^{k+2} \quad k \geq 0$
D_{k+2}	$\tilde{\Phi}_1^{k+1} + \tilde{\Phi}_1 \tilde{\Phi}_2^2 \quad k \geq 2$
E_6	$\tilde{\Phi}_1^3 + \tilde{\Phi}_2^4$
E_7	$\tilde{\Phi}_1^3 + \tilde{\Phi}_1 \tilde{\Phi}_2^3$
E_8	$\tilde{\Phi}_1^3 + \tilde{\Phi}_2^5$

(4.195)

⁷For $SU(N)$ with N odd, $SO(2N)$ with N odd, and E_6 , the degrees d_i of the Casimirs contain odd integers corresponding to zeros so that the vacuum exponents are indeterminate.

The twisted superpotential has the left-moving R-charge +1 and the right-moving R-charge -1.

The elliptic genera of the $\mathcal{N} = (2, 2)$ minimal models in the NSNS sector are given by [99, 100]

$$\mathcal{I}_{\text{NS}}^{2d (2, 2) \min A_{k+1}}(y; q) = \frac{(q^{\frac{1}{2} + \frac{1}{2(k+2)}} y^{\frac{k+1}{k+2}}; q)_{\infty} (q^{\frac{1}{2} - \frac{1}{2(k+2)}} y^{-\frac{k+1}{k+2}}; q)_{\infty}}{(q^{\frac{1}{2(k+2)}} y^{-\frac{1}{k+2}}; q)_{\infty} (q^{1 - \frac{1}{2(k+2)}} y^{\frac{1}{k+2}}; q)_{\infty}}, \quad (4.196)$$

$$\begin{aligned} \mathcal{I}_{\text{NS}}^{2d (2, 2) \min D_{k+2}}(y; q) &= \frac{(q^{\frac{1}{2} + \frac{1}{2(k+1)}} y^{-\frac{1}{k+1} + 1}; q)_{\infty} (q^{\frac{1}{2} - \frac{1}{2(k+1)}} y^{\frac{1}{k+1} - 1}; q)_{\infty}}{(q^{\frac{1}{2(k+1)}} y^{-\frac{1}{k+1}}; q)_{\infty} (q^{1 - \frac{1}{2(k+1)}} y^{\frac{1}{k+1}}; q)_{\infty}} \\ &\times \frac{(q^{\frac{1}{2} + \frac{k}{4(k+1)}} y^{-\frac{k}{2(k+1)} + 1}; q)_{\infty} (q^{\frac{1}{2} - \frac{k}{4(k+1)}} y^{\frac{k}{2(k+1)} - 1}; q)_{\infty}}{(q^{\frac{k}{4(k+1)}} y^{-\frac{k}{2(k+1)}}; q)_{\infty} (q^{1 - \frac{k}{4(k+1)}} y^{\frac{k}{2(k+1)}}; q)_{\infty}}, \end{aligned} \quad (4.197)$$

$$\mathcal{I}_{\text{NS}}^{2d (2, 2) \min E_6}(y; q) = \frac{(q^{\frac{2}{3}} y^{\frac{2}{3}}; q)_{\infty} (q^{\frac{1}{3}} y^{-\frac{2}{3}}; q)_{\infty} (q^{\frac{5}{8}} y^{\frac{3}{4}}; q)_{\infty} (q^{\frac{3}{8}} y^{-\frac{3}{4}}; q)_{\infty}}{(q^{\frac{1}{6}} y^{-\frac{1}{3}}; q)_{\infty} (q^{\frac{5}{6}} y^{\frac{1}{3}}; q)_{\infty} (q^{\frac{1}{8}} y^{-\frac{1}{4}}; q)_{\infty} (q^{\frac{7}{8}} y^{\frac{1}{4}}; q)_{\infty}}, \quad (4.198)$$

$$\mathcal{I}_{\text{NS}}^{2d (2, 2) \min E_7}(y; q) = \frac{(q^{\frac{2}{3}} y^{\frac{2}{3}}; q)_{\infty} (q^{\frac{1}{3}} y^{-\frac{2}{3}}; q)_{\infty} (q^{\frac{11}{18}} y^{\frac{7}{9}}; q)_{\infty} (q^{\frac{7}{18}} y^{-\frac{7}{9}}; q)_{\infty}}{(q^{\frac{1}{6}} y^{-\frac{1}{3}}; q)_{\infty} (q^{\frac{5}{6}} y^{\frac{1}{3}}; q)_{\infty} (q^{\frac{1}{9}} y^{-\frac{2}{9}}; q)_{\infty} (q^{\frac{8}{9}} y^{\frac{2}{9}}; q)_{\infty}}, \quad (4.199)$$

$$\mathcal{I}_{\text{NS}}^{2d (2, 2) \min E_8}(y; q) = \frac{(q^{\frac{2}{3}} y^{\frac{2}{3}}; q)_{\infty} (q^{\frac{1}{3}} y^{-\frac{2}{3}}; q)_{\infty} (q^{\frac{3}{5}} y^{\frac{4}{5}}; q)_{\infty} (q^{\frac{2}{5}} y^{-\frac{4}{5}}; q)_{\infty}}{(q^{\frac{1}{6}} y^{-\frac{1}{3}}; q)_{\infty} (q^{\frac{5}{6}} y^{\frac{1}{3}}; q)_{\infty} (q^{\frac{1}{10}} y^{-\frac{1}{5}}; q)_{\infty} (q^{\frac{9}{10}} y^{\frac{1}{5}}; q)_{\infty}}. \quad (4.200)$$

The single-particle indices are

$$\begin{aligned} i_{\text{NS}}^{2d (2, 2) \min A_{k+1}}(q) &= -\frac{q^{\frac{1}{2} + \frac{1}{2(k+2)}} y^{\frac{k+1}{k+2}} + q^{\frac{1}{2} - \frac{1}{2(k+2)}} y^{-\frac{k+1}{k+2}}}{1 - q} \\ &+ \frac{q^{\frac{1}{2(k+2)}} y^{-\frac{1}{k+2}} + q^{1 - \frac{1}{2(k+2)}} y^{\frac{1}{k+2}}}{1 - q}, \end{aligned} \quad (4.201)$$

$$\begin{aligned} i_{\text{NS}}^{2d (2, 2) \min D_{k+2}}(q) &= -\frac{q^{\frac{1}{2} + \frac{1}{2(k+1)}} y^{-\frac{1}{k+1} + 1} + q^{\frac{1}{2} - \frac{1}{2(k+1)}} y^{\frac{1}{k+1} - 1}}{1 - q} + \frac{q^{\frac{1}{2(k+1)}} y^{-\frac{1}{k+1}} + q^{1 - \frac{1}{2(k+1)}} y^{\frac{1}{k+1}}}{1 - q} \\ &- \frac{q^{\frac{1}{2} + \frac{k}{4(k+1)}} y^{-\frac{k}{2(k+1)} + 1} + q^{\frac{1}{2} - \frac{k}{4(k+1)}} y^{\frac{k}{2(k+1)} - 1}}{1 - q} + \frac{q^{\frac{k}{4(k+1)}} y^{-\frac{k}{2(k+1)}} + q^{1 - \frac{k}{4(k+1)}} y^{\frac{k}{2(k+1)}}}{1 - q}, \end{aligned} \quad (4.202)$$

$$i_{\text{NS}}^{2d (2, 2) \min E_6}(q) = -\frac{q^{\frac{2}{3}} y^{\frac{2}{3}} + q^{\frac{1}{3}} y^{-\frac{2}{3}}}{1 - q} + \frac{q^{\frac{1}{6}} y^{-\frac{1}{3}} + q^{\frac{5}{6}} y^{\frac{1}{3}}}{1 - q} - \frac{q^{\frac{5}{8}} y^{\frac{3}{4}} + q^{\frac{3}{8}} y^{-\frac{3}{4}}}{1 - q} + \frac{q^{\frac{1}{8}} y^{-\frac{1}{4}} + q^{\frac{7}{8}} y^{\frac{1}{4}}}{1 - q}, \quad (4.203)$$

$$i_{\text{NS}}^{2d (2, 2) \min E_7}(q) = -\frac{q^{\frac{2}{3}} y^{\frac{2}{3}} + q^{\frac{1}{3}} y^{-\frac{2}{3}}}{1 - q} + \frac{q^{\frac{1}{6}} y^{-\frac{1}{3}} + q^{\frac{5}{6}} y^{\frac{1}{3}}}{1 - q} - \frac{q^{\frac{11}{18}} y^{\frac{7}{9}} + q^{\frac{7}{18}} y^{-\frac{7}{9}}}{1 - q} + \frac{q^{\frac{1}{9}} y^{-\frac{2}{9}} + q^{\frac{8}{9}} y^{\frac{2}{9}}}{1 - q}, \quad (4.204)$$

$$i_{\text{NS}}^{2d (2, 2) \min E_8}(q) = -\frac{q^{\frac{2}{3}} y^{\frac{2}{3}} + q^{\frac{1}{3}} y^{-\frac{2}{3}}}{1 - q} + \frac{q^{\frac{1}{6}} y^{-\frac{1}{3}} + q^{\frac{5}{6}} y^{\frac{1}{3}}}{1 - q} - \frac{q^{\frac{3}{5}} y^{\frac{4}{5}} + q^{\frac{2}{5}} y^{-\frac{4}{5}}}{1 - q} + \frac{q^{\frac{1}{10}} y^{-\frac{1}{5}} + q^{\frac{9}{10}} y^{\frac{1}{5}}}{1 - q}. \quad (4.205)$$

The supersymmetric zeta functions for the $\mathcal{N} = (2, 2)$ minimal models are evaluated as

$$\begin{aligned} \mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ min } A_{k+1}}(s, z) = & -\zeta\left(s, \frac{k+3}{2(k+2)} + z\right) - \zeta\left(s, \frac{k+1}{2(k+2)} + z\right) \\ & + \zeta\left(s, \frac{1}{2(k+2)} + z\right) + \zeta\left(s, \frac{2k+3}{2(k+2)} + z\right), \end{aligned} \quad (4.206)$$

$$\begin{aligned} \mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ min } D_{k+2}}(s, z) = & -\zeta\left(s, \frac{k+2}{2(k+1)} + z\right) - \zeta\left(s, \frac{k}{2(k+1)} + z\right) \\ & + \zeta\left(s, \frac{1}{2(k+1)} + z\right) + \zeta\left(s, \frac{2k+1}{2(k+1)} + z\right) \\ & - \zeta\left(s, \frac{3k+2}{4(k+1)} + z\right) - \zeta\left(s, \frac{k+2}{4(k+1)} + z\right) \\ & + \zeta\left(s, \frac{k}{4(k+1)} + z\right) + \zeta\left(s, \frac{3k+4}{4(k+1)} + z\right), \end{aligned} \quad (4.207)$$

$$\begin{aligned} \mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ min } E_6}(s, z) = & -\zeta\left(s, \frac{1}{3} + z\right) - \zeta\left(s, \frac{2}{3} + z\right) + \zeta\left(s, \frac{1}{6} + z\right) + \zeta\left(s, \frac{5}{6} + z\right) \\ & - \zeta\left(s, \frac{5}{8} + z\right) - \zeta\left(s, \frac{3}{8} + z\right) + \zeta\left(s, \frac{1}{8} + z\right) + \zeta\left(s, \frac{7}{8} + z\right), \end{aligned} \quad (4.208)$$

$$\begin{aligned} \mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ min } E_7}(s, z) = & -\zeta\left(s, \frac{1}{3} + z\right) - \zeta\left(s, \frac{2}{3} + z\right) + \zeta\left(s, \frac{1}{6} + z\right) + \zeta\left(s, \frac{5}{6} + z\right) \\ & - \zeta\left(s, \frac{11}{18} + z\right) - \zeta\left(s, \frac{7}{18} + z\right) + \zeta\left(s, \frac{1}{9} + z\right) + \zeta\left(s, \frac{8}{9} + z\right), \end{aligned} \quad (4.209)$$

$$\begin{aligned} \mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ min } E_8}(s, z) = & -\zeta\left(s, \frac{1}{3} + z\right) - \zeta\left(s, \frac{2}{3} + z\right) + \zeta\left(s, \frac{1}{6} + z\right) + \zeta\left(s, \frac{5}{6} + z\right) \\ & - \zeta\left(s, \frac{3}{5} + z\right) - \zeta\left(s, \frac{2}{5} + z\right) + \zeta\left(s, \frac{1}{10} + z\right) + \zeta\left(s, \frac{9}{10} + z\right). \end{aligned} \quad (4.210)$$

For any series of the $\mathcal{N} = (2, 2)$ minimal models, we have

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ min } G}(s, z) = 0 \quad (4.211)$$

and

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ min } G}(0, 0) = 0. \quad (4.212)$$

The special supersymmetric zeta values with $s = -1$ and $z = 0$ are given by

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ min } A_{k+1}}(-1, 0) = -\frac{k}{4(k+2)}, \quad (4.213)$$

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ min } D_{k+2}}(-1, 0) = -\frac{k}{4(k+1)}, \quad (4.214)$$

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ min } E_6}(-1, 0) = -\frac{5}{24}, \quad (4.215)$$

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ min } E_7}(-1, 0) = -\frac{2}{9}, \quad (4.216)$$

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ min } E_8}(-1, 0) = -\frac{7}{30}. \quad (4.217)$$

According to the formulae (4.15) and (4.16), we get the central charges [101]

$$c = c_L = c_R = 3 - \frac{6}{h^\vee} \quad (4.218)$$

of the $\mathcal{N} = (2, 2)$ minimal models, where h^\vee is the dual Coxeter number of the relevant Lie algebra

minimal modes	c	h^\vee
A_{k+1}	$\frac{3k}{k+2}$	$k+2$
D_{k+2}	$\frac{3k}{k+1}$	$2k+2$
E_6	$\frac{5}{2}$	12
E_7	$\frac{8}{3}$	18
E_8	$\frac{14}{5}$	30

(4.219)

The supersymmetric determinants are given by

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ min } A_{k+1}}(z) = \frac{\Gamma\left(\frac{1}{2(k+2)} + z\right) \Gamma\left(\frac{2k+3}{2(k+2)} + z\right)}{\Gamma\left(\frac{k+3}{2(k+2)} + z\right) \Gamma\left(\frac{k+1}{2(k+2)} + z\right)}, \quad (4.220)$$

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ min } D_{k+2}}(z) = \frac{\Gamma\left(\frac{1}{2(k+1)} + z\right) \Gamma\left(\frac{2k+1}{2(k+1)} + z\right) \Gamma\left(\frac{k}{4(k+1)} + z\right) \Gamma\left(\frac{3k+4}{4(k+1)} + z\right)}{\Gamma\left(\frac{k+2}{2(k+1)} + z\right) \Gamma\left(\frac{k}{2(k+1)} + z\right) \Gamma\left(\frac{3k+2}{4(k+1)} + z\right) \Gamma\left(\frac{k+2}{4(k+1)} + z\right)}, \quad (4.221)$$

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ min } E_6}(z) = \frac{\Gamma\left(\frac{1}{6} + z\right) \Gamma\left(\frac{5}{6} + z\right) \Gamma\left(\frac{1}{8} + z\right) \Gamma\left(\frac{7}{8} + z\right)}{\Gamma\left(\frac{1}{3} + z\right) \Gamma\left(\frac{2}{3} + z\right) \Gamma\left(\frac{5}{8} + z\right) \Gamma\left(\frac{3}{8} + z\right)}, \quad (4.222)$$

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ min } E_7}(z) = \frac{\Gamma\left(\frac{1}{6} + z\right) \Gamma\left(\frac{5}{6} + z\right) \Gamma\left(\frac{1}{9} + z\right) \Gamma\left(\frac{8}{9} + z\right)}{\Gamma\left(\frac{1}{3} + z\right) \Gamma\left(\frac{2}{3} + z\right) \Gamma\left(\frac{11}{18} + z\right) \Gamma\left(\frac{7}{18} + z\right)}, \quad (4.223)$$

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ min } E_8}(z) = \frac{\Gamma\left(\frac{1}{6} + z\right) \Gamma\left(\frac{5}{6} + z\right) \Gamma\left(\frac{1}{10} + z\right) \Gamma\left(\frac{9}{10} + z\right)}{\Gamma\left(\frac{1}{3} + z\right) \Gamma\left(\frac{2}{3} + z\right) \Gamma\left(\frac{3}{5} + z\right) \Gamma\left(\frac{2}{5} + z\right)}. \quad (4.224)$$

The vacuum exponents are

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ min } A_{k+1}}(0) = \cot\left(\frac{\pi}{2k+4}\right), \quad (4.225)$$

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ min } D_{k+2}}(0) = \cot\left(\frac{\pi}{2(k+1)}\right) \cot\left(\frac{\pi k}{4(k+1)}\right), \quad (4.226)$$

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ min } E_6}(0) = \sqrt{3} \cot\left(\frac{\pi}{8}\right) = 4.18154\dots, \quad (4.227)$$

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ min } E_7}(0) = \sqrt{3} \cot\left(\frac{\pi}{9}\right) = 4.75877\dots, \quad (4.228)$$

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ min } E_8}(0) = \sqrt{3} \cot\left(\frac{\pi}{10}\right) = \sqrt{15 + 6\sqrt{5}} = 5.3307\dots \quad (4.229)$$

4.2.5 2d $\mathcal{N} = (2, 2)$ Kazama-Suzuki models

Kazama-Suzuki models [102, 103] are the rational $\mathcal{N} = (2, 2)$ SCFTs that are described by the coset

$$\frac{SU(M+1)_k \times SO(2M)_1}{SU(M)_{k+1} \times U(1)_{M(M+1)(M+k+1)}}. \quad (4.230)$$

It is expected to be described by the LG theory of M twisted chiral multiplets $\tilde{\Phi}_i$ with a superpotential [104]

$$\tilde{\mathcal{W}} = \sum_{i=1}^M \frac{1}{M+k+1} \tilde{\phi}_i^{M+k+1}, \quad (4.231)$$

where

$$\sum_{i=0}^M \tilde{\Phi}_i t^i = \prod_{i=1}^M (1 + \tilde{\phi}_i t). \quad (4.232)$$

For $M = 1$ they are identical to the $\mathcal{N} = (2, 2)$ A_{k+1} minimal model.

The elliptic genus of the A_{k+1} Kazama-Suzuki model in the NSNS sector is given by [100]

$$\mathcal{I}_{\text{NS}}^{2\text{d } (2,2) \text{ KS}_{M,k}}(y; q) = \prod_{j=1}^M \frac{(q^{\frac{k+j}{2(M+k+1)}} y^{-\frac{k+j}{M+k+1}}; q)_{\infty} (q^{1-\frac{k+j}{2(M+k+1)}} y^{\frac{k+j}{M+k+1}}; q)_{\infty}}{(q^{\frac{j}{2(M+k+1)}} y^{-\frac{j}{M+k+1}}; q)_{\infty} (q^{1-\frac{j}{2(M+k+1)}} y^{\frac{j}{M+k+1}}; q)_{\infty}}. \quad (4.233)$$

Then the single-particle index reads

$$\begin{aligned} & \mathcal{I}_{\text{NS}}^{2\text{d } (2,2) \text{ KS}_{M,k}}(y; q) \\ &= \sum_{j=1}^M \left[-\frac{q^{\frac{k+j}{2(M+k+1)}} y^{-\frac{k+j}{M+k+1}}}{1-q} - \frac{q^{1-\frac{k+j}{2(M+k+1)}} y^{\frac{k+j}{M+k+1}}}{1-q} + \frac{q^{\frac{j}{2(M+k+1)}} y^{-\frac{j}{M+k+1}}}{1-q} + \frac{q^{1-\frac{j}{2(M+k+1)}} y^{\frac{j}{M+k+1}}}{1-q} \right]. \end{aligned} \quad (4.234)$$

The supersymmetric zeta function is

$$\begin{aligned} & \mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ KS}_{M,k}}(s, z) \\ &= \sum_{j=1}^M \left[-\zeta \left(s, \frac{k+j}{2(M+k+1)} + z \right) - \zeta \left(s, 1 - \frac{k+j}{2(M+k+1)} + z \right) \right. \\ & \quad \left. + \zeta \left(s, \frac{j}{2(M+k+1)} + z \right) + \zeta \left(s, 1 - \frac{j}{2(M+k+1)} + z \right) \right]. \end{aligned} \quad (4.235)$$

Although each of the Hurwitz zeta function has a simple pole at $s = 1$, the total sum of the residues vanishes

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ KS}_{M,k}}(s, z) = 0. \quad (4.236)$$

Also the special supersymmetric zeta value with $s = 0$ and $z = 0$ vanishes

$$\mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ KS}_{M,k}}(0, 0) = 0. \quad (4.237)$$

The special supersymmetric zeta value with $s = -1$ and $z = 0$ is given by

$$\begin{aligned} \mathfrak{Z}_{\text{NS}}^{2\text{d } (2,2) \text{ KS}_{M,k}}(-1, 0) &= - \sum_{j=1}^M \frac{k(2M+k-2j+2)}{4(M+k+1)^2} \\ &= - \frac{Mk}{4(M+k+1)}. \end{aligned} \quad (4.238)$$

Therefore we get from (4.15) and (4.16) the central charge

$$c = c_L = c_R = \frac{3Mk}{M+k+1}. \quad (4.239)$$

The supersymmetric determinant is

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ KS}_{M,k}}(z) = \prod_{j=1}^M \frac{\Gamma \left(\frac{j}{2(M+k+1)} + z \right) \Gamma \left(1 - \frac{j}{2(M+k+1)} + z \right)}{\Gamma \left(\frac{k+j}{2(M+k+1)} + z \right) \Gamma \left(1 - \frac{k+j}{2(M+k+1)} + z \right)}. \quad (4.240)$$

The vacuum exponent is evaluated as

$$\mathfrak{D}_{\text{NS}}^{2\text{d } (2,2) \text{ KS}_{M,k}}(0) = \prod_{i=1}^{\lfloor \frac{M+1}{2} \rfloor} \cot \left(\frac{\pi(2i-1)}{2(k+M+1)} \right) \prod_{j=1}^{\lfloor \frac{M}{2} \rfloor} \cot \left(\frac{\pi j}{k+M+1} \right), \quad (4.241)$$

where $\lfloor x \rfloor := \max\{m \in \mathbb{Z} | m \leq x\}$ is the floor function.

4.2.6 2d $\mathcal{N} = (2, 2)$ \mathbb{CP}^{N-1} models

The $\mathcal{N} = (2, 2)$ supersymmetric \mathbb{CP}^{N-1} model is described by a $U(1)$ vector multiplet with N chiral multiplets of charge $+1$ (See e.g. [105, 106, 107, 108] for earlier works on the model). It is known as an asymptotic free theory with N vacua with mass gap. So the theory is not conformal in the IR. It is mirror to the A_{N-1} affine Toda theory that is an LG theory of $N - 1$ periodic chiral multiplet fields Φ_α with a superpotential of the form

$$\mathcal{W} = \Lambda \left(\sum_{\alpha=1}^{N-1} e^{\Phi_\alpha} + \prod_{\alpha=1}^{N-1} e^{-\Phi_\alpha} \right), \quad (4.242)$$

with Λ being some constant, corresponding to the dynamical scale generated in the \mathbb{CP}^{N-1} model.

The elliptic genus is given by [77]

$$\mathcal{I}_{\text{NS}}^{2\text{d } (2,2) \mathbb{CP}^{N-1}}(y, x_\alpha; q) = \frac{(q)_\infty^2}{(q^{\frac{1}{2}} y^\pm; q)_\infty} \oint_{\text{JK}} \frac{ds}{2\pi i s} \prod_{\alpha=1}^N \frac{(q^{\frac{1}{2}} y^\mp s^\pm x_\alpha^\mp; q)_\infty}{(s x_\alpha^{-1}; q)_\infty (q s^{-1} x_\alpha; q)_\infty}, \quad (4.243)$$

where $y^N = 1$ and $\prod_\alpha x_\alpha = 1$. Picking up the JK residues at N simple poles $s = x_\beta$ with $\beta = 1, \dots, N$, we obtain

$$\mathcal{I}_{\text{NS}}^{2\text{d } (2,2) \mathbb{CP}^{N-1}}(y, x_\alpha; q) = \sum_{\beta=1}^N \mathcal{I}_{\text{NS},\beta}^{2\text{d } (2,2) \mathbb{CP}^{N-1}}(y, x_\alpha; q), \quad (4.244)$$

where

$$\mathcal{I}_{\text{NS},\beta}^{2\text{d } (2,2) \mathbb{CP}^{N-1}}(y, x_\alpha; q) = \prod_{\gamma \neq \beta} \frac{(q^{\frac{1}{2}} y^{-1} \frac{x_\beta}{x_\gamma}; q)_\infty (q^{\frac{1}{2}} y \frac{x_\gamma}{x_\beta}; q)_\infty}{(\frac{x_\beta}{x_\gamma}; q)_\infty (q \frac{x_\gamma}{x_\beta}; q)_\infty}. \quad (4.245)$$

In the unflavored limit each residue (4.245) takes the same form albeit with a singularity so that the full index is simply given by multiplying it by N . When we take the plethystic logarithm of the full index, the multiplicative factor N simply gives rise to the constant term $\log N$, which is irrelevant upon the Mellin transform. So it is enough to consider the single-particle index as the plethystic logarithm of the contribution (4.245)

$$i_{\text{NS},\beta}^{2\text{d } (2,2) \mathbb{CP}^{N-1}}(y, x_\alpha; q) = \sum_{\gamma \neq \beta} \left[-\frac{q^{\frac{1}{2}} y^{-1} \frac{x_\beta}{x_\gamma}}{1-q} - \frac{q^{\frac{1}{2}} y \frac{x_\gamma}{x_\beta}}{1-q} + \frac{\frac{x_\beta}{x_\gamma}}{1-q} + \frac{q \frac{x_\gamma}{x_\beta}}{1-q} \right]. \quad (4.246)$$

Then the supersymmetric zeta function associated with (4.246) takes the form

$$\begin{aligned} & \mathbf{3}_{\text{NS},\beta}^{2\text{d } (2,2) \mathbb{CP}^{N-1}}(s, z) \\ &= -2(N-1)\zeta\left(s, \frac{1}{2} + z\right) + (N-1)\zeta(s, z) + (N-1)\zeta(s, 1+z). \end{aligned} \quad (4.247)$$

We have

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{NS},\beta}^{2\text{d } (2,2) \text{ } \mathbb{CP}^{N-1}}(s, z) = 0 \quad (4.248)$$

and

$$\mathfrak{Z}_{\text{NS},\beta}^{2\text{d } (2,2) \text{ } \mathbb{CP}^{N-1}}(0, 0) = 0. \quad (4.249)$$

The supersymmetric zeta value with $s = -1$ and $z = 0$ is given by

$$\mathfrak{Z}_{\text{NS},\beta}^{2\text{d } (2,2) \text{ } \mathbb{CP}^{N-1}}(-1, 0) = -\frac{N-1}{4}. \quad (4.250)$$

From (4.15) and (4.16) we obtain

$$c = c_L = c_R = 3(N-1). \quad (4.251)$$

This is the 't Hooft anomaly rather than the central charge. The value $\frac{c}{3}$ is equal to the dimension $N-1$ of the \mathbb{CP}^{N-1} .

The supersymmetric determinant is

$$\mathfrak{D}_{\text{NS},\beta}^{2\text{d } (2,2) \text{ } \mathbb{CP}^{N-1}}(z) = \frac{\Gamma(z)^{N-1} \Gamma(1+z)^{N-1}}{\Gamma(\frac{1}{2}+z)^{2(N-1)}}. \quad (4.252)$$

The vacuum exponent is divergent

$$\mathfrak{D}_{\text{NS},\beta}^{2\text{d } (2,2) \text{ } \mathbb{CP}^{N-1}}(0) = \infty. \quad (4.253)$$

5 4d supersymmetric field theories

5.1 Zeta functions for 4d supersymmetric indices

The supersymmetric indices for 4d $\mathcal{N} = 1$ supersymmetric field theories can be defined as follows [10, 11]:

$$\mathcal{I}^{4\text{d } \mathcal{N}=1}(x; p; q) = \text{Tr}(-1)^F p^{j_3 + \bar{j}_3 - \frac{R}{2}} q^{j_3 - \bar{j}_3 - \frac{R}{2}} \prod_{\alpha} x_{\alpha}^{f_{\alpha}}, \quad (5.1)$$

where j_3 and \bar{j}_3 are the Cartan generators of the rotation group $SU(2)_1 \times SU(2)_2$, R is the $U(1)_R$ R-charge, and f_{α} are the Cartan generators of the other global symmetries. The supersymmetric indices of 4d $\mathcal{N} = 1$ gauge theories can be evaluated as the elliptic hypergeometric integrals [109].

The 6-form anomaly polynomial takes the form

$$\mathcal{A}_6 = \frac{1}{6} k_{RRR} c_1(R)^3 - \frac{1}{24} k_{RC_1}(R) p_1(T), \quad (5.2)$$

where $c_1(R)$ is the first Chern class for the $U(1)_R$ R-symmetry background gauge field and $p_1(T)$ is the first Pontryagin class of the tangent bundle T of the four-dimensional spacetime. The 't Hooft anomaly coefficient k_{RRR} for the $U(1)$ R-symmetry and the mixed $U(1)_R$ -gravitational anomaly coefficient k_R are given by

$$k_{RRR} = \text{Tr } U(1)_R^3, \quad k_R = \text{Tr } U(1)_R. \quad (5.3)$$

For the superconformal field theories, the c -anomaly c_{4d} and the a -anomaly a_{4d} are linearly related to the 't Hooft anomaly coefficient for the $U(1)_R$ R-symmetry and the mixed $U(1)$ -gravitational anomaly coefficient [110]

$$c_{4d} = \frac{9}{32}k_{RRR} - \frac{5}{32}k_R, \quad (5.4)$$

$$a_{4d} = \frac{9}{32}k_{RRR} - \frac{3}{32}k_R. \quad (5.5)$$

We define the multiple supersymmetric zeta functions $\mathfrak{Z}^{4d \mathcal{N}=1}(s, z; \omega_1, \omega_2)$ for the $4d \mathcal{N} = 1$ supersymmetric field theories by taking the Mellin transform (2.7) or equivalently (2.9) of the plethystic logarithms of the $4d$ supersymmetric indices (5.1) with the fugacities p and q being replaced with $p = q^{\omega_1}$ and $q = q^{\omega_2}$. Subsequently, the supersymmetric determinants $\mathfrak{D}^{4d \mathcal{N}=1}(z; \omega_1, \omega_2)$ associated with the supersymmetric zeta functions $\mathfrak{Z}^{4d \mathcal{N}=1}(s, z; \omega_1, \omega_2)$ are defined by (3.1). These supersymmetric zeta functions can be expressed in terms of the Barnes double zeta functions [36, 111], which have simple poles at $s = 2$ and $s = 1$. However, the supersymmetric zeta functions $\mathfrak{Z}^{4d \mathcal{N}=1}(s, z; \omega_1, \omega_2)$ may only have a simple pole at $s = 1$.

From the asymptotic formula (2.47), we have

$$\begin{aligned} & \log \mathcal{I}^{4d \mathcal{N}=1}(1; q^{\omega_1}; q^{\omega_2}) \\ & \sim \frac{\pi^2 \text{Res}_{s=1} \mathfrak{Z}^{4d \mathcal{N}=1}(s, 0; \omega_1, \omega_2)}{6\beta} - \mathfrak{Z}^{4d \mathcal{N}=1}(0, 0; \omega_1, \omega_2) \log \beta + \log \mathfrak{D}^{4d \mathcal{N}=1}(0; \omega_1, \omega_2). \end{aligned} \quad (5.6)$$

When we expand the $4d \mathcal{N} = 1$ supersymmetric index with $p = q$ as

$$\mathcal{I}^{4d \mathcal{N}=1}(1; q; q) = \sum_n d^{4d \mathcal{N}=1}(n) q^n, \quad (5.7)$$

the analysis of the asymptotic behavior of the expansion coefficients $d^{4d \mathcal{N}=1}(n)$ needs delicate and subtle care since the expansion coefficients $d^{4d \mathcal{N}=1}(n)$ generally do not grow exponentially.

Di Pietro and Komargodski argued [45] that the leading terms in the Cardy-like limit of the $4d$ supersymmetric indices are universally captured by the mixed $U(1)$ -gravitational anomalies and they can be expressed in terms of the $4d$ central charges

c_{4d} and a_{4d} for superconformal theories. This implies that the residues of the supersymmetric zeta functions at a simple pole $s = 1$ are given by

$$\text{Res}_{s=1} \mathfrak{Z}^{4d \mathcal{N}=1}(s, z; \omega_1, \omega_2) = 16(c_{4d} - a_{4d}) \frac{\omega_1 + \omega_2}{\omega_1 \omega_2}. \quad (5.8)$$

From the various examples we have examined, the special supersymmetric zeta values with $s = -1$ and $z = 0$ can be determined by the c - and a -anomalies

$$\begin{aligned} \mathfrak{Z}^{4d \mathcal{N}=1}(-1, 0; \omega_1, \omega_2) \\ = \frac{4(\omega_1^3 + \omega_2^3)}{9\omega_1\omega_2} c_{4d} - \frac{4(2\omega_1^3 - 3\omega_1^2\omega_2 - 3\omega_1\omega_2^2 + 2\omega_2^3)}{27\omega_1\omega_2} a_{4d}. \end{aligned} \quad (5.9)$$

Comparison with the supersymmetric Casimir energies for 4d $\mathcal{N} = 1$ supersymmetric field theories [60, 61] (also see [65, 62, 63, 112, 66]) shows that our proposed relation (2.76) holds.

For superconformal field theories, from (5.8) and (5.9) the c - and a -anomalies can be expressed in terms of the supersymmetric zeta functions

$$c_{4d} = \frac{1}{128} \text{Res}_{s=1} \mathfrak{Z}^{4d \mathcal{N}=1}(s, 0; 1, 1) + \frac{27}{32} \mathfrak{Z}^{4d \mathcal{N}=1}(-1, 0; 1, 1), \quad (5.10)$$

$$a_{4d} = -\frac{3}{128} \text{Res}_{s=1} \mathfrak{Z}^{4d \mathcal{N}=1}(s, 0; 1, 1) + \frac{27}{32} \mathfrak{Z}^{4d \mathcal{N}=1}(-1, 0; 1, 1). \quad (5.11)$$

Also, the 6-form anomaly coefficients are given by

$$k_{RRR} = -\frac{1}{4} \text{Res}_{s=1} \mathfrak{Z}^{4d \mathcal{N}=1}(s, 0; 1, 1) + 3 \mathfrak{Z}^{4d \mathcal{N}=1}(-1, 0; 1, 1), \quad (5.12)$$

$$k_R = -\frac{1}{2} \text{Res}_{s=1} \mathfrak{Z}^{4d \mathcal{N}=1}(s, 0; 1, 1). \quad (5.13)$$

5.1.1 4d $\mathcal{N} = 1$ chiral multiplet

The supersymmetric index of the 4d $\mathcal{N} = 1$ chiral multiplet with R-charge r takes the form [11]

$$\mathcal{I}^{4d \mathcal{N}=1 \text{ chiral}_r}(x; p; q) = \Gamma((pq)^{\frac{r}{2}} x; p, q), \quad (5.14)$$

where

$$\Gamma(x; p, q) = \prod_{m=0}^{\infty} \prod_{n=0}^{\infty} \frac{1 - p^{m+1} q^{n+1} x^{-1}}{1 - p^m q^n x} \quad (5.15)$$

is the elliptic gamma function. The single-particle index is given by

$$i^{4d \mathcal{N}=1 \text{ chiral}_r}(x; p; q) = \frac{(pq)^{\frac{r}{2}} x - (pq)^{\frac{2-r}{2}} x^{-1}}{(1-p)(1-q)}. \quad (5.16)$$

For $p = q$ it becomes

$$\begin{aligned} i^{4d} \mathcal{N} = 1 \text{ chiral}_r(x; q; q) &= \frac{q^r x - q^{2-r} x^{-1}}{(1 - q)^2} \\ &= \sum_{n=1}^{\infty} (nq^{n-1+r} x - nq^{n+1-r} x^{-1}). \end{aligned} \quad (5.17)$$

The supersymmetric zeta function is evaluated as

$$\begin{aligned} \mathfrak{Z}^{4d \mathcal{N} = 1 \text{ chiral}_r}(s, z; \omega_1, \omega_2) \\ = \zeta_2 \left(s, \frac{r(\omega_1 + \omega_2)}{2} + z; \omega_1, \omega_2 \right) - \zeta_2 \left(s, \frac{(2-r)(\omega_1 + \omega_2)}{2} + z; \omega_1, \omega_2 \right), \end{aligned} \quad (5.18)$$

where

$$\zeta_2(s, z; a, b) = \sum_{m, n \geq 0} (am + bn + z)^{-s} \quad (5.19)$$

is the Barnes double zeta function [36, 111] defined by the double series. When $\omega_1 = \omega_2 = 1$, it reduces to

$$\begin{aligned} \mathfrak{Z}^{4d \mathcal{N} = 1 \text{ chiral}_r}(s, z; 1, 1) &= \zeta(s-1, r+z) - \zeta(s-1, 2-r+z) \\ &\quad + (1-r-z)\zeta(s, r+z) - (-1+r-z)\zeta(s, 2-r+z). \end{aligned} \quad (5.20)$$

The Barnes double zeta function (5.19) has an analytic continuation to the complex s -plane with simple poles at $s = 1$ and $s = 2$. The residues at these poles are given by [113]

$$\text{Res}_{s=2} \zeta_2(s, z; a, b) = \frac{1}{ab}, \quad (5.21)$$

$$\text{Res}_{s=1} \zeta_2(s, z; a, b) = \frac{1}{2} \left(\frac{1}{a} + \frac{1}{b} \right) - \frac{z}{ab}. \quad (5.22)$$

Hence the residues at simple poles at $s = 2$ and $s = 1$ for the supersymmetric zeta function (5.18) are evaluated as

$$\text{Res}_{s=2} \mathfrak{Z}^{4d \mathcal{N} = 1 \text{ chiral}_r}(s, z; \omega_1, \omega_2) = 0, \quad (5.23)$$

$$\text{Res}_{s=1} \mathfrak{Z}^{4d \mathcal{N} = 1 \text{ chiral}_r}(s, z; \omega_1, \omega_2) = \frac{(1-r)(\omega_1 + \omega_2)}{\omega_1 \omega_2}. \quad (5.24)$$

The special supersymmetric zeta values can be evaluated by making use of the formula (C.18). The Zeta-index is

$$\mathfrak{Z}^{4d \mathcal{N} = 1 \text{ chiral}_r}(0, 0; \omega_1, \omega_2) = 0. \quad (5.25)$$

The supersymmetric zeta value with $s = -1$ and $z = 0$ is

$$\begin{aligned} \mathfrak{Z}^{4d \mathcal{N}=1 \text{ chiral}_r}(-1, 0; \omega_1, \omega_2) \\ = \frac{\omega_1 + \omega_2}{24\omega_1\omega_2}(r-1) [(\omega_1 + \omega_2)^2 r^2 - 2(\omega_1 + \omega_2)^2 r + 2\omega_1\omega_2]. \end{aligned} \quad (5.26)$$

Substituting (5.24) and (5.26) into the formulae (5.10), (5.11), (5.12) and (5.13), we find the anomalies

$$\begin{aligned} c_{4d} &= \frac{9}{32}k_{RRR} - \frac{5}{32}k_R \\ &= \frac{9}{32}r^3 - \frac{27}{32}r^2 + \frac{11}{16}r - \frac{1}{8}, \end{aligned} \quad (5.27)$$

$$\begin{aligned} a_{4d} &= \frac{9}{32}k_{RRR} - \frac{3}{32}k_R \\ &= \frac{9}{32}r^3 - \frac{27}{32}r^2 + \frac{3}{4}r - \frac{3}{16}, \end{aligned} \quad (5.28)$$

$$k_{RRR} = (r-1)^3, \quad (5.29)$$

$$k_R = r-1. \quad (5.30)$$

For a free chiral multiplet the R-charge is $r = 2/3$ the expressions (5.27) and (5.28) reduce to the central charges

$$c_{4d} = \frac{1}{24}, \quad (5.31)$$

$$a_{4d} = \frac{1}{48}. \quad (5.32)$$

In this case, the value $a_{4d}/c_{4d} = 1/2$ hits the lower bound for $\mathcal{N} = 1$ SCFT [114].

The supersymmetric determinant is

$$\mathfrak{D}^{4d \mathcal{N}=1 \text{ chiral}_r}(z; \omega_1, \omega_2) = \frac{\Gamma_2\left(\frac{r(\omega_1+\omega_2)}{2} + z; \omega_1, \omega_2\right)}{\Gamma_2\left(\frac{(2-r)(\omega_1+\omega_2)}{2} + z; \omega_1, \omega_2\right)}, \quad (5.33)$$

where

$$\Gamma_2(z; \omega_1, \omega_2) := \exp \left[\left. \frac{\partial}{\partial s} \zeta_2(s, z; \omega_1, \omega_2) \right|_{s=0} \right] \quad (5.34)$$

is the Barnes double gamma function [36]. The vacuum exponent with $\omega_1 = \omega_2 = 1$ is given by

$$\mathfrak{D}^{4d \mathcal{N}=1 \text{ chiral}_r}(0; 1, 1) = e^{\zeta'(-1, r) - \zeta'(-1, 2-r)} \left[\frac{\Gamma(r)\Gamma(2-r)}{2\pi} \right]^{1-r}. \quad (5.35)$$

We show a plot of the vacuum exponent (5.35) in Figure 4.




Figure 4: The vacuum exponent $\mathfrak{D}^{4d \mathcal{N}=1 \text{ chiral}_r}(0; 1, 1)$.

When one introduces a mass term $\mathcal{W} = \lambda\Phi^2$, the chiral multiplet has R-charge $r = 1$. The theory has a single supersymmetric vacuum. The supersymmetric index (5.14) is equal to 1 and the supersymmetric zeta function (5.18) vanishes. The vacuum exponent $\mathfrak{D}^{4d \mathcal{N}=1 \text{ chiral}_r}(0; 1, 1)$ is 1.

For the theory of a chiral multiplet with a linear superpotential $\mathcal{W} = \lambda\Phi$, the Polonyi model [115], the chiral multiplet has R-charge $r = 2$. There is no supersymmetric vacuum so that supersymmetry is spontaneously broken. The supersymmetric index (5.14) vanishes. It stems from the fermionic contribution -1 in the single-particle index (5.16), corresponding to the Goldstino. In this case the vacuum exponent $\mathfrak{D}^{4d \mathcal{N}=1 \text{ chiral}_r}(0; 1, 1)$ vanishes, as discussed in (3.8).

5.1.2 4d $\mathcal{N} = 1$ vector multiplet

The supersymmetric index of the 4d $\mathcal{N} = 1$ $U(1)$ vector multiplet reads [11]

$$\mathcal{I}^{4d \mathcal{N}=1 \text{ vec}}(p; q) = (p; p)_\infty (q; q)_\infty. \quad (5.36)$$

The single-particle index of the vector multiplet is

$$i^{4d \mathcal{N}=1 \text{ vec}}(p; q) = -\frac{p}{1-p} - \frac{q}{1-q}. \quad (5.37)$$

In the limit $p = q$, it reduces to

$$i^{4d \mathcal{N}=1 \text{ vec}}(q) = \frac{-2q}{1-q} = -2 \sum_{n=1}^{\infty} q^n. \quad (5.38)$$

The supersymmetric zeta function is given by

$$\mathfrak{Z}^{4d \mathcal{N}=1 \text{ vec}}(s, z; \omega_1, \omega_2) = -\omega_1^{-s} \zeta\left(s, \frac{\omega_1 + z}{\omega_1}\right) - \omega_2^{-s} \zeta\left(s, \frac{\omega_2 + z}{\omega_2}\right). \quad (5.39)$$

It has a simple pole at $s = 1$ with residue

$$\text{Res}_{s=1} \mathfrak{Z}^{4d \mathcal{N}=1 \text{ vec}}(s, z; \omega_1, \omega_2) = -\frac{1}{\omega_1} - \frac{1}{\omega_2}. \quad (5.40)$$

The Zeta-index is

$$\mathfrak{Z}^{4d \mathcal{N}=1 \text{ vec}}(0, 0; \omega_1, \omega_2) = 1. \quad (5.41)$$

The supersymmetric zeta value with $s = -1$ and $z = 0$ is

$$\mathfrak{Z}^{4d \mathcal{N}=1 \text{ vec}}(-1, 0; \omega_1, \omega_2) = \frac{\omega_1 + \omega_2}{12}. \quad (5.42)$$

Plugging (5.40) and (5.42) into the formulae (5.12) and (5.13), we get the anomalies

$$k_{RRR} = 1, \quad (5.43)$$

$$k_R = 1. \quad (5.44)$$

Also from (5.10) and (5.11), we get

$$c_{4d} = \frac{1}{8}, \quad (5.45)$$

$$a_{4d} = \frac{3}{16}, \quad (5.46)$$

which are equal to the central charges contributed from the 4d $\mathcal{N} = 1$ $U(1)$ vector multiplet. The value $a_{4d}/c_{4d} = 3/2$ hits the upper bound for $\mathcal{N} = 1$ SCFT [114].

The supersymmetric determinant is

$$\mathfrak{D}^{4d \mathcal{N}=1 \text{ vec}}(z; \omega_1, \omega_2) = \frac{2\pi \omega_1^{\frac{1}{2} - \frac{\omega_1 + z}{\omega_1}} \omega_2^{\frac{1}{2} - \frac{\omega_2 + z}{\omega_2}}}{\Gamma\left(\frac{\omega_1 + z}{\omega_1}\right) \Gamma\left(\frac{\omega_2 + z}{\omega_2}\right)}. \quad (5.47)$$

The vacuum exponent is

$$\mathfrak{D}^{4d \mathcal{N}=1 \text{ vec}}(0; \omega_1, \omega_2) = \frac{2\pi}{\sqrt{\omega_1 \omega_2}}. \quad (5.48)$$

5.1.3 4d $\mathcal{N} = 1$ $SU(2)$ with 6 chirals $(\Phi_\alpha, \bar{\Phi}_\alpha)$

Let us consider the 4d $\mathcal{N} = 1$ $SU(2)$ gauge theory with 3 fundamental chiral multiplets Φ_α and 3 antifundamental chiral multiplets $\bar{\Phi}_\alpha$. The theory is dual to a free theory of 15 chiral multiplets (Φ', B', \bar{B}') interacting with a superpotential [116]

$$\mathcal{W} = \Phi' B' \bar{B}' - \det \Phi'. \quad (5.49)$$

The field content is summarized as

	$SU(2)$	$SU(3)_x$	$SU(3)_y$	$U(1)_B$	$U(1)_R$
Φ_α	2	3	1	+1	$\frac{1}{3}$
$\bar{\Phi}_\alpha$	$\bar{\mathbf{2}}$	1	3	-1	$\frac{1}{3}$
Φ'	1	3	3	0	$\frac{2}{3}$
B'	1	$\bar{\mathbf{3}}$	1	+2	$\frac{2}{3}$
\bar{B}'	1	1	$\bar{\mathbf{3}}$	-2	$\frac{2}{3}$

(5.50)

The supersymmetric index is given by

$$\begin{aligned}
\mathcal{I}^{4d \mathcal{N}=1 \text{ } SU(2) \text{ } \Phi_\alpha}(b, x_\alpha, \tilde{x}_\alpha; p; q) \\
= \frac{(p; p)_\infty (q; q)_\infty}{2} \oint \frac{ds}{2\pi i s} \frac{1}{\Gamma(s^{\pm 2}; p, q)} \prod_{\alpha=1}^3 \Gamma((pq)^{\frac{1}{6}} b x_\alpha s^\mp; p, q) \Gamma((pq)^{\frac{1}{6}} b^{-1} \tilde{x}_\alpha s^\pm; p, q),
\end{aligned}
\tag{5.51}$$

with $\prod_\alpha x_\alpha = \prod_\alpha \tilde{x}_\alpha = 1$. It is shown to be equal to [117]

$$\prod_{\alpha=1}^3 \prod_{\beta=1}^3 \Gamma((pq)^{\frac{1}{3}} x_\alpha \tilde{x}_\beta; p, q) \prod_{\alpha=1}^3 \Gamma((pq)^{\frac{1}{3}} b^2 x_\alpha^{-1}; p, q) \prod_{\alpha=1}^3 \Gamma((pq)^{\frac{1}{3}} b^{-2} \tilde{x}_\alpha^{-1}; p, q), \tag{5.52}$$

which is identified with the index of the dual theory. The single-particle index reads

$$\begin{aligned}
i^{4d \mathcal{N}=1 \text{ } SU(2) \text{ } \Phi_\alpha}(b, x_\alpha, \tilde{x}_\alpha; p; q) \\
= \sum_{\alpha=1}^3 \sum_{\beta=1}^3 \frac{(pq)^{\frac{1}{3}} x_\alpha \tilde{x}_\beta - (pq)^{\frac{2}{3}} \tilde{x}_\alpha^{-1} x_\beta^{-1}}{(1-p)(1-q)} \\
+ \sum_{\alpha=1}^3 \frac{(pq)^{\frac{1}{3}} b^2 x_\alpha^{-1} - (pq)^{\frac{2}{3}} b^{-2} x_\alpha}{(1-p)(1-q)} + \sum_{\alpha=1}^3 \frac{(pq)^{\frac{1}{3}} b^{-2} \tilde{x}_\alpha^{-1} - (pq)^{\frac{2}{3}} b^2 \tilde{x}_\alpha}{(1-p)(1-q)}.
\end{aligned}
\tag{5.53}$$

With $p = q^{\omega_1}$, $q = q^{\omega_2}$, $x_\alpha = \tilde{x}_\alpha = 1$ and $b = q^B$, it becomes

$$\begin{aligned}
i^{4d \mathcal{N}=1 \text{ } SU(2) \text{ } \Phi_\alpha}(q^B, 1, 1; q^{\omega_1}, q^{\omega_2}) \\
= 9 \frac{q^{\frac{\omega_1+\omega_2}{3}} - q^{\frac{2(\omega_1+\omega_2)}{3}}}{(1-q^{\omega_1})(1-q^{\omega_2})} + 3 \frac{q^{\frac{\omega_1+\omega_2}{3}+2B} - q^{\frac{2(\omega_1+\omega_2)}{3}-2B}}{(1-q^{\omega_1})(1-q^{\omega_2})} + 3 \frac{q^{\frac{\omega_1+\omega_2}{3}-2B} - q^{\frac{2(\omega_1+\omega_2)}{3}+2B}}{(1-q^{\omega_1})(1-q^{\omega_2})}.
\end{aligned}
\tag{5.54}$$

The supersymmetric zeta function is evaluated as

$$\begin{aligned}
& \mathfrak{Z}^{4d \mathcal{N}=1 \text{ } SU(2) \text{ } \Phi_\alpha}(s, z; \omega_1, \omega_2; B) \\
&= 9 \left[\zeta_2 \left(s, \frac{\omega_1 + \omega_2}{3} + z; \omega_1, \omega_2 \right) - \zeta_2 \left(s, \frac{2(\omega_1 + \omega_2)}{3} + z; \omega_1, \omega_2 \right) \right] \\
&+ 3 \left[\zeta_2 \left(s, \frac{\omega_1 + \omega_2}{3} + 2B + z; \omega_1, \omega_2 \right) - \zeta_2 \left(s, \frac{2(\omega_1 + \omega_2)}{3} - 2B + z; \omega_1, \omega_2 \right) \right] \\
&+ 3 \left[\zeta_2 \left(s, \frac{\omega_1 + \omega_2}{3} - 2B + z; \omega_1, \omega_2 \right) - \zeta_2 \left(s, \frac{2(\omega_1 + \omega_2)}{3} + 2B + z; \omega_1, \omega_2 \right) \right].
\end{aligned} \tag{5.55}$$

While the Barnes double zeta function has simple poles at $s = 2$ and 1 , the total residues at these poles are

$$\text{Res}_{s=2} \mathfrak{Z}^{4d \mathcal{N}=1 \text{ } SU(2) \text{ } \Phi_\alpha}(s, z; \omega_1, \omega_2; B) = 0, \tag{5.56}$$

$$\text{Res}_{s=1} \mathfrak{Z}^{4d \mathcal{N}=1 \text{ } SU(2) \text{ } \Phi_\alpha}(s, z; \omega_1, \omega_2; B) = \frac{5(\omega_1 + \omega_2)}{\omega_1 \omega_2}. \tag{5.57}$$

Also we have

$$\mathfrak{Z}^{4d \mathcal{N}=1 \text{ } SU(2) \text{ } \Phi_\alpha}(0, 0; \omega_1, \omega_2; B) = 0 \tag{5.58}$$

and

$$\mathfrak{Z}^{4d \mathcal{N}=1 \text{ } SU(2) \text{ } \Phi_\alpha}(-1, 0; \omega_1, \omega_2; B) = -4B^2 \frac{\omega_1 + \omega_2}{\omega_1 \omega_2} + \frac{5}{36}(\omega_1 + \omega_2) + \frac{5}{27} \frac{\omega_1^3 + \omega_2^3}{\omega_1 \omega_2}. \tag{5.59}$$

From (5.57) and (5.59) as well as the formulae (5.10) and (5.11), we find the trial central charges

$$c_{4d} = \frac{5}{8} - \frac{27B^2}{(\omega_1 + \omega_2)^2}, \tag{5.60}$$

$$a_{4d} = \frac{5}{16} - \frac{27B^2}{(\omega_1 + \omega_2)^2}. \tag{5.61}$$

According to the a -maximization [21], the mixed anomaly coefficient B is required to vanish. Then we get

$$c_{4d} = \frac{5}{8}, \tag{5.62}$$

$$a_{4d} = \frac{5}{16}, \tag{5.63}$$

which agree with the central charges. In this case, $a_{4d}/c_{4d} = 1/2$ hits the lower bound for $\mathcal{N} = 1$ SCFT [114].

The supersymmetric determinant is

$$\mathfrak{D}^{4d \mathcal{N}=1 SU(2) \Phi_\alpha}(z; \omega_1, \omega_2) = \frac{\Gamma_2\left(\frac{\omega_1+\omega_2}{3} + z; \omega_1, \omega_2\right)^{15}}{\Gamma_2\left(\frac{2(\omega_1+\omega_2)}{3} + z; \omega_1, \omega_2\right)^{15}}. \quad (5.64)$$

Using Mathematica, the vacuum exponent with $\omega_1 = \omega_2 = 1$ is evaluated as

$$\mathfrak{D}^{4d \mathcal{N}=1 SU(2) \Phi_\alpha}(0; 1, 1) = \frac{1}{9\sqrt{3}} e^{\frac{5(2\pi^2 - 3\psi^{(1)}(\frac{1}{3}))}{6\sqrt{3}\pi}} = 0.0127546 \dots \quad (5.65)$$

5.1.4 4d $\mathcal{N} = 1$ $SU(N)$ with $2(N+1)$ chirals $(\Phi_\alpha, \bar{\Phi}_\alpha)$

Consider the 4d $\mathcal{N} = 1$ $SU(N)$ gauge theory with $N+1$ fundamental chiral multiplets Φ_α and $N+1$ antifundamental chiral multiplets $\bar{\Phi}_\alpha$. It is dual to a free theory consisting of $(N+1)^2 + (N+1) + (N+1)$ chiral multiplets (Φ', B', \bar{B}') interacting with a superpotential [116]

$$\mathcal{W} = \Phi' B' \bar{B}' - \det \Phi'. \quad (5.66)$$

The field content and charges are

	$SU(N)$	$SU(N+1)_x$	$SU(N+1)_y$	$U(1)_B$	$U(1)_R$
Φ_α	\mathbf{N}	$\mathbf{N+1}$	$\mathbf{1}$	+1	$\frac{1}{N+1}$
$\bar{\Phi}_\alpha$	$\bar{\mathbf{N}}$	$\mathbf{1}$	$\mathbf{N+1}$	-1	$\frac{1}{N+1}$
Φ'	$\mathbf{1}$	$\mathbf{N+1}$	$\mathbf{N+1}$	0	$\frac{2}{N+1}$
B'	$\mathbf{1}$	$\bar{\mathbf{N+1}}$	$\mathbf{1}$	+N	$\frac{N}{N+1}$
\bar{B}'	$\mathbf{1}$	$\mathbf{1}$	$\bar{\mathbf{N+1}}$	-N	$\frac{N}{N+1}$

(5.67)

The supersymmetric index is given by

$$\begin{aligned} \mathcal{I}^{4d \mathcal{N}=1 SU(N) \Phi_\alpha}(b, x_\alpha, \tilde{x}_\alpha; p, q) \\ = \frac{(p; p)_\infty^{N-1} (q; q)_\infty^{N-1}}{N!} \oint \prod_{i=1}^{N-1} \frac{ds_i}{2\pi i s_i} \prod_{i < j} \frac{1}{\Gamma(s_i^\pm s_j^\mp; p, q)} \\ \times \prod_{i=1}^N \prod_{\alpha=1}^{N+1} \Gamma((pq)^{\frac{1}{2(N+1)}} b x_\alpha s_i^\mp; p, q) \Gamma((pq)^{\frac{1}{2(N+1)}} b^{-1} \tilde{x}_\alpha s_i^\pm; p, q), \end{aligned} \quad (5.68)$$

with $\prod_\alpha x_\alpha = \prod_\alpha \tilde{x}_\alpha = 1$. It agrees with the index

$$\prod_{\alpha=1}^{N+1} \prod_{\beta=1}^{N+1} \Gamma((pq)^{\frac{1}{N+1}} x_\alpha \tilde{x}_\beta; p, q) \prod_{\alpha=1}^{N+1} \Gamma((pq)^{\frac{N}{2(N+1)}} b^N x_\alpha^{-1}; p, q) \Gamma((pq)^{\frac{N}{2(N+1)}} b^{-N} \tilde{x}_\alpha^{-1}; p, q) \quad (5.69)$$

of the dual theory. The single-particle index reads

$$\begin{aligned}
& i^{4d} \mathcal{N} = 1 \text{ } SU(N) \text{ } \Phi_\alpha(b, x_\alpha, \tilde{x}_\alpha; p; q) \\
&= \sum_{\alpha=1}^{N+1} \sum_{\beta=1}^{N+1} \frac{(pq)^{\frac{1}{N+1}} x_\alpha \tilde{x}_\beta - (pq)^{1-\frac{1}{N+1}} x_\alpha^{-1} \tilde{x}_\beta^{-1}}{(1-p)(1-q)} \\
&+ \sum_{\alpha=1}^{N+1} \frac{(pq)^{\frac{N}{2(N+1)}} b^N x_\alpha^{-1} - (pq)^{1-\frac{N}{2(N+1)}} b^{-N} x_\alpha}{(1-p)(1-q)} + \sum_{\alpha=1}^{N+1} \frac{(pq)^{\frac{N}{2(N+1)}} b^{-N} \tilde{x}_\alpha^{-1} - (pq)^{1-\frac{N}{2(N+1)}} b^N \tilde{x}_\alpha}{(1-p)(1-q)}.
\end{aligned} \tag{5.70}$$

If we set $p = q^{\omega_1}$, $q = q^{\omega_2}$ and $b = q^B$ and turn off the flavor fugacities x_α , \tilde{x}_α , it reduces to

$$\begin{aligned}
& i^{4d} \mathcal{N} = 1 \text{ } SU(N) \text{ } \Phi_\alpha(q^B, x_\alpha, \tilde{x}_\alpha; q^{\omega_1}; q^{\omega_2}) \\
&= (N+1)^2 \frac{q^{\frac{\omega_1+\omega_2}{N+1}} - q^{(1-\frac{1}{N+1})(\omega_1+\omega_2)}}{(1-q^{\omega_1})(1-q^{\omega_2})} \\
&+ (N+1) \frac{q^{\frac{N}{2(N+1)}(\omega_1+\omega_2)+BN} - q^{1-\frac{N}{2(N+1)}(\omega_1+\omega_2)-BN}}{(1-q^{\omega_1})(1-q^{\omega_2})} \\
&+ (N+1) \frac{q^{\frac{N}{2(N+1)}(\omega_1+\omega_2)-BN} - q^{1-\frac{N}{2(N+1)}(\omega_1+\omega_2)+BN}}{(1-q^{\omega_1})(1-q^{\omega_2})}.
\end{aligned} \tag{5.71}$$

The supersymmetric zeta function is

$$\begin{aligned}
& \mathfrak{Z}^{4d} \mathcal{N} = 1 \text{ } SU(N) \text{ } \Phi_\alpha(s, z; \omega_1, \omega_2) \\
&= (N+1)^2 \left[\zeta_2 \left(s, \frac{\omega_1 + \omega_2}{N+1} + z; \omega_1, \omega_2 \right) - \zeta_2 \left(s, \frac{N(\omega_1 + \omega_2)}{N+1} + z; \omega_1, \omega_2 \right) \right] \\
&+ (N+1) \left[\zeta_2 \left(s, \frac{N(\omega_1 + \omega_2)}{2(N+1)} + BN + z; \omega_1, \omega_2 \right) - \zeta_2 \left(s, \frac{(N+2)(\omega_1 + \omega_2)}{2(N+1)} - BN + z; \omega_1, \omega_2 \right) \right] \\
&+ (N+1) \left[\zeta_2 \left(s, \frac{N(\omega_1 + \omega_2)}{2(N+1)} - BN + z; \omega_1, \omega_2 \right) - \zeta_2 \left(s, \frac{(N+2)(\omega_1 + \omega_2)}{2(N+1)} + BN + z; \omega_1, \omega_2 \right) \right].
\end{aligned} \tag{5.72}$$

The residues at simple poles $s = 2$ and 1 are evaluated as

$$\text{Res}_{s=2} \mathfrak{Z}^{4d} \mathcal{N} = 1 \text{ } SU(N) \text{ } \Phi_\alpha(s, z; \omega_1, \omega_2) = 0, \tag{5.73}$$

$$\text{Res}_{s=1} \mathfrak{Z}^{4d} \mathcal{N} = 1 \text{ } SU(N) \text{ } \Phi_\alpha(s, z; \omega_1, \omega_2) = (N^2 + 1) \frac{\omega_1 + \omega_2}{\omega_1 \omega_2}. \tag{5.74}$$

For $s = 0$ and $z = 0$ the supersymmetric zeta value vanishes

$$\mathfrak{Z}^{4d} \mathcal{N} = 1 \text{ } SU(N) \text{ } \Phi_\alpha(0, 0; \omega_1, \omega_2) = 0. \tag{5.75}$$

For $s = -1$ and $z = 0$ we find

$$\begin{aligned} & \mathfrak{Z}^{4d \mathcal{N}=1 \text{ } SU(N) \text{ } \Phi_\alpha}(-1, 0; \omega_1, \omega_2) \\ &= -B^2 N^2 \frac{\omega_1 + \omega_2}{\omega_1 \omega_2} - \frac{N^4 - 4N^3 - N^2 + 2N + 1}{12(N+1)^2}(\omega_1 + \omega_2) + \frac{N^2(2N+1)(\omega_1^3 + \omega_2^3)}{12(N+1)^2 \omega_1 \omega_2}. \end{aligned} \quad (5.76)$$

According to the formulae (5.10) and (5.11), we find

$$c_{4d} = \frac{1}{16} \left[18N - \left(2 + \frac{108B^2}{(\omega_1 + \omega_2)^2} \right) N^2 - \frac{N(29N+22)+2}{(N+1)^2} \right], \quad (5.77)$$

$$a_{4d} = \frac{3}{16} \left[6N - \left(1 + \frac{36B^2}{(\omega_1 + \omega_2)^2} \right) N^2 - \frac{2N(5N+4)+1}{(N+1)^2} \right]. \quad (5.78)$$

The value a_{4d} is maximum at $B = 0$, i.e. the vanishing mixed anomaly coefficient. We then arrive at

$$c_{4d} = -\frac{9N^4}{16(N+1)^2} + \frac{7}{16}N^2 - \frac{1}{8}, \quad (5.79)$$

$$a_{4d} = -\frac{9N^4}{16(N+1)^2} + \frac{3}{8}N^2 - \frac{3}{16}. \quad (5.80)$$

For $N = 2$ they reduce to the central charges (5.62) and (5.63). For $N > 2$ they do not obey the universal bounds on c and a for $\mathcal{N} = 1$ SCFTs, whereas they match the expected anomalies of the 4d $\mathcal{N} = 1$ $SU(N)$ SQCD with $2(N+1)$ chirals.⁸

The supersymmetric determinant is

$$\begin{aligned} & \mathfrak{D}^{4d \mathcal{N}=1 \text{ } SU(N) \text{ } \Phi_\alpha}(z; \omega_1, \omega_2) \\ &= \frac{\Gamma_2\left(\frac{\omega_1+\omega_2}{N+1} + z; \omega_1, \omega_2\right)^{(N+1)^2} \Gamma_2\left(\frac{N(\omega_1+\omega_2)}{2(N+1)} + z; \omega_1, \omega_2\right)^{2(N+1)}}{\Gamma_2\left(\frac{N(\omega_1+\omega_2)}{N+1} + z; \omega_1, \omega_2\right)^{(N+1)^2} \Gamma_2\left(\frac{(N+2)(\omega_1+\omega_2)}{2(N+1)} + z; \omega_1, \omega_2\right)^{2(N+1)}}. \end{aligned} \quad (5.83)$$

It turns out that the vacuum exponents with $\omega_1 = \omega_2 = 1$ are finite values for finite $N \geq 2$. For example, for $N = 3$ we find the finite vacuum exponent with $\omega_1 = \omega_2 = 1$ given by

$$\mathfrak{D}^{4d \mathcal{N}=1 \text{ } SU(3) \text{ } \Phi_\alpha}(0; 1, 1) = \frac{1}{512} e^{-\frac{4G}{\pi}} = 0.000608467 \dots, \quad (5.84)$$

⁸For 4d $\mathcal{N} = 1$ SQCD with N_f flavors the central charges are

$$c_{4d} = -\frac{9N^4}{16N_f^2} + \frac{7}{16}N^2 - \frac{1}{8}, \quad (5.81)$$

$$a_{4d} = -\frac{9N^4}{16N_f^2} + \frac{3}{8}N^2 - \frac{3}{16}. \quad (5.82)$$

where

$$G = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)^2} = 0.915965 \dots \quad (5.85)$$

is Catalan's constant. For $N = 2, \dots, 10$ the vacuum exponents with $\omega_1 = \omega_2 = 1$ are numerically computed as follows:

N	$\mathfrak{D}^{4d} \mathcal{N} = 1 \text{ } SU(N) \text{ } \Phi_\alpha(0; 1, 1)$
2	0.0127546
3	0.000608467
4	0.0000520141
5	0.0000130731
6	0.0000142974
7	0.0000941049
8	0.00491762
9	2.599
10	17226.5

(5.86)

5.1.5 4d $\mathcal{N} = 1 \text{ } USp(2N)$ with $2N + 4$ chirals Φ_α

Another type of higher rank generalizations of the $SU(2)$ gauge theory with 6 chiral multiplets is the $\mathcal{N} = 1 \text{ } USp(2N)$ gauge theory with $2N + 4$ fundamental chiral multiplets Φ_α . It is dual to the free theory of $(N + 2)(2N + 3)$ chiral multiplets Φ' with a superpotential

$$\mathcal{W} = \text{Pf} \Phi'. \quad (5.87)$$

The duality belongs to the Intriligator-Pouliot duality [87]. The field content is summarized as

	$USp(2N)$	$SU(2N + 4)$	$U(1)_R$
Φ_α	2N	2N + 4	$\frac{1}{N+2}$
Φ'	1	(N + 2)(2N + 3)	$\frac{2}{N+2}$

(5.88)

The supersymmetric index of the $USp(2N)$ gauge theory is

$$\begin{aligned}
& \mathcal{I}^{4d} \mathcal{N} = 1 \text{ } USp(2N) \text{ } \Phi_\alpha(x_\alpha; p; q) \\
&= \frac{(p; p)_\infty^N (q; q)_\infty^N}{2^N N!} \oint \prod_{i=1}^N \frac{ds_i}{2\pi i s_i} \frac{1}{\Gamma(s_i^{\pm 2}; p, q)} \prod_{i < j} \frac{1}{\Gamma(s_i^\pm s_j^\mp; p, q) \Gamma(s_i^\pm s_j^\pm; p, q)} \\
&\times \prod_{i=1}^N \prod_{\alpha=1}^{2N+4} \Gamma((pq)^{\frac{1}{2(N+2)}} s_i^\pm x_\alpha; p, q).
\end{aligned} \quad (5.89)$$

It agrees with

$$\prod_{\alpha < \beta}^{2N+4} \Gamma((pq)^{\frac{1}{N+2}} x_\alpha x_\beta; p, q), \quad (5.90)$$

which is the index of the dual free theory. The single-particle index is

$$\begin{aligned} & i^{4d} \mathcal{N} = 1 \text{ } USp(2N) \text{ } \Phi_\alpha(x_\alpha; p; q) \\ &= \sum_{\alpha < \beta}^{2N+4} \frac{(pq)^{\frac{1}{N+2}} x_\alpha x_\beta - (pq)^{1 - \frac{1}{N+2}} x_\alpha^{-1} x_\beta^{-1}}{(1-p)(1-q)}. \end{aligned} \quad (5.91)$$

The supersymmetric zeta function is

$$\begin{aligned} & \mathfrak{Z}^{4d \text{ } \mathcal{N} = 1 \text{ } USp(2N) \text{ } \Phi_\alpha}(s, z; \omega_1, \omega_2) \\ &= (N+2)(2N+3) \left[\zeta_2 \left(s, \frac{\omega_1 + \omega_2}{N+2} + z; \omega_1, \omega_2 \right) - \zeta_2 \left(s, \frac{(N+1)(\omega_1 + \omega_2)}{N+2} + z; \omega_1, \omega_2 \right) \right]. \end{aligned} \quad (5.92)$$

The residues at simple poles $s = 2$ and 1 are

$$\text{Res}_{s=2} \mathfrak{Z}^{4d \text{ } \mathcal{N} = 1 \text{ } USp(2N) \text{ } \Phi_\alpha}(s, z; \omega_1, \omega_2) = 0, \quad (5.93)$$

$$\text{Res}_{s=1} \mathfrak{Z}^{4d \text{ } \mathcal{N} = 1 \text{ } USp(2N) \text{ } \Phi_\alpha}(s, z; \omega_1, \omega_2) = \frac{N(2N+3)(\omega_1 + \omega_2)}{\omega_1 \omega_2}. \quad (5.94)$$

The supersymmetric zeta value with $s = 0$ and $z = 0$ vanishes

$$\mathfrak{Z}^{4d \text{ } \mathcal{N} = 1 \text{ } USp(2N) \text{ } \Phi_\alpha}(0, 0; \omega_1, \omega_2) = 0. \quad (5.95)$$

For $s = -1$ and $z = 0$ the supersymmetric zeta value is given by

$$\begin{aligned} & \mathfrak{Z}^{4d \text{ } \mathcal{N} = 1 \text{ } USp(2N) \text{ } \Phi_\alpha}(-1, 0; \omega_1, \omega_2) \\ &= \frac{N(N+1)(2N+3)}{6(N+2)^2} \frac{\omega_1^3 + \omega_2^3}{\omega_1 \omega_2} - \frac{N(2N+3)(N^2 - 2N - 2)}{12(N+2)^2} (\omega_1 + \omega_2). \end{aligned} \quad (5.96)$$

Substituting (5.94) and (5.96) into formulae (5.10) and (5.11), we obtain

$$c_{4d} = - \frac{N(2N+3)(N^2 - 5N - 5)}{8(N+2)^2} \quad (5.97)$$

$$a_{4d} = - \frac{3N(2N^3 - N^2 - 10N - 6)}{16(N+2)^2}. \quad (5.98)$$

When $N = 1$ they reduce to (5.62) and (5.63). For $N > 2$ they do not obey the universal bounds on a and c , while the values (5.97) and (5.98) are the contributions to the c - and a -anomalies. As before, the reason they do not obey the universal bound

is for $N > 3$, the term in the superpotential $\text{Pf } \Phi'$ is not renormalizable, and the true “IR fixed point” is located at the zero value of the coefficient, leading to an enhanced symmetry.

The supersymmetric determinant is

$$\mathfrak{D}^{4d \mathcal{N}=1 \text{ } USp(2N) \text{ } \Phi_\alpha}(z; \omega_1, \omega_2) = \frac{\Gamma_2\left(\frac{\omega_1 + \omega_2}{N+2} + z; \omega_1, \omega_2\right)^{(N+2)(2N+3)}}{\Gamma_2\left(\frac{(N+1)(\omega_1 + \omega_2)}{N+2} + z; \omega_1, \omega_2\right)^{(N+2)(2N+3)}}. \quad (5.99)$$

We find that the vacuum exponents with $\omega_1 = \omega_2 = 1$ are finite values for finite N . For $N = 1, \dots, 8$ the vacuum exponents are numerically computed as follows:

N	$\mathfrak{D}^{4d \mathcal{N}=1 \text{ } USp(2N) \text{ } \Phi_\alpha}(0; 1, 1)$
1	0.0127546
2	0.0000610352
3	6.06434×10^{-7}
4	3.89098×10^{-8}
5	3.79592×10^{-8}
6	1.12473×10^{-6}
7	0.00181271
8	263.197

(5.100)

5.1.6 4d $\mathcal{N} = 2$ hypermultiplet

The 4d $\mathcal{N} = 2$ hypermultiplet can be constructed by a pair of two $\mathcal{N} = 1$ chiral multiplets of $r = 2/3$.

The supersymmetric index is given by

$$\begin{aligned} \mathcal{I}^{4d \mathcal{N}=2 \text{ hyp}}(y, z; p; q) &= \mathcal{I}^{4d \mathcal{N}=1 \text{ chiral}_{r=\frac{2}{3}}}(yz; p; q) \times \mathcal{I}^{4d \mathcal{N}=1 \text{ chiral}_{r=\frac{2}{3}}}(yz^{-1}; p; q) \\ &= \Gamma((pq)^{\frac{1}{3}}yz; p, q) \Gamma((pq)^{\frac{1}{3}}yz^{-1}; p, q), \end{aligned} \quad (5.101)$$

where y is the fugacity for the $SU(2) \times U(1)$ R-symmetry. The single-particle index is

$$i^{4d \mathcal{N}=2 \text{ hyp}}(y, z; p; q) = \frac{(pq)^{\frac{1}{3}}y(z + z^{-1}) - (pq)^{\frac{2}{3}}y^{-1}(z + z^{-1})}{(1-p)(1-q)}. \quad (5.102)$$

The supersymmetric zeta function is

$$\begin{aligned} \mathfrak{Z}^{4d \mathcal{N}=2 \text{ hyp}}(s, z; \omega_1, \omega_2) &= 2 \left[\zeta_2 \left(s, \frac{\omega_1 + \omega_2}{3} + z, \omega_1, \omega_2 \right) - \zeta_2 \left(s, \frac{2(\omega_1 + \omega_2)}{3} + z, \omega_1, \omega_2 \right) \right]. \end{aligned} \quad (5.103)$$

The residues at simple poles $s = 2$ and 1 are

$$\text{Res}_{s=2} \mathfrak{Z}^{4d \mathcal{N}=2 \text{ hyp}}(s, z; \omega_1, \omega_2) = 0, \quad (5.104)$$

$$\text{Res}_{s=1} \mathfrak{Z}^{4d \mathcal{N}=2 \text{ hyp}}(s, z; \omega_1, \omega_2) = \frac{2}{3} \frac{\omega_1 + \omega_2}{\omega_1 \omega_2}. \quad (5.105)$$

For $s = 0$ and $z = 0$ it vanishes

$$\mathfrak{Z}^{4d \mathcal{N}=2 \text{ hyp}}(0, 0; \omega_1, \omega_2) = 0. \quad (5.106)$$

The special value with $s = -1$ and $z = 0$ is

$$\mathfrak{Z}^{4d \mathcal{N}=2 \text{ hyp}}(-1, 0; \omega_1, \omega_2) = \frac{1}{54}(\omega_1 + \omega_2) + \frac{2}{81} \frac{\omega_1^3 + \omega_2^3}{\omega_1 \omega_2}. \quad (5.107)$$

From (5.105) and (5.107) we get

$$c_{4d} = \frac{1}{12}, \quad (5.108)$$

$$a_{4d} = \frac{1}{24}. \quad (5.109)$$

They are the 4d central charges of the $\mathcal{N} = 2$ hypermultiplet. The value $a_{4d}/c_{4d} = 1/2$ saturates the lower bound for $\mathcal{N} = 2$ SCFT [118, 114].

The supersymmetric determinant is

$$\mathfrak{D}^{4d \mathcal{N}=2 \text{ hyp}}(z; \omega_1, \omega_2) = \frac{\Gamma_2\left(\frac{(\omega_1 + \omega_2)}{3} + z; \omega_1, \omega_2\right)^2}{\Gamma_2\left(\frac{2(\omega_1 + \omega_2)}{3} + z; \omega_1, \omega_2\right)^2} \quad (5.110)$$

The vacuum exponent with $\omega_1 = \omega_2 = 1$ is given by

$$\mathfrak{D}^{4d \mathcal{N}=2 \text{ hyp}}(0; 1, 1) = \frac{1}{3^{\frac{1}{3}}} e^{\frac{2\pi^2 - 3\psi^{(1)}(\frac{1}{3})}{9\sqrt{3}\pi}} = 0.559014 \dots \quad (5.111)$$

5.1.7 4d $\mathcal{N} = 2$ vector multiplet

The 4d $\mathcal{N} = 2$ $U(1)$ vector multiplet can be realized as a combination of the $\mathcal{N} = 1$ $U(1)$ vector multiplet and the $\mathcal{N} = 1$ chiral multiplet with $r = 2/3$.

The supersymmetric index is given by

$$\begin{aligned} \mathcal{I}^{4d \mathcal{N}=2 \text{ vec}}(y; p; q) &= \mathcal{I}^{4d \mathcal{N}=1 \text{ vec}}(p; q) \times \mathcal{I}^{4d \mathcal{N}=1 \text{ chiral}_{r=\frac{2}{3}}}(y^{-2}; p; q) \\ &= (p; p)_\infty (q; q)_\infty \Gamma((pq)^{\frac{1}{3}} y^{-2}; p, q), \end{aligned} \quad (5.112)$$

where y is the fugacity for the $SU(2) \times U(1)$ R-symmetry. The single-particle index reads

$$i^{4d \mathcal{N}=2 \text{ vec}}(y; p; q) = -\frac{p}{1-p} - \frac{q}{1-q} + \frac{(pq)^{\frac{1}{3}}y^{-2} - (pq)^{\frac{2}{3}}y^2}{(1-p)(1-q)}. \quad (5.113)$$

The supersymmetric zeta function is

$$\begin{aligned} \mathfrak{Z}^{4d \mathcal{N}=2 \text{ vec}}(s, z; \omega_1, \omega_2) &= -\omega_1^{-s} \zeta\left(s, \frac{\omega_1 + z}{\omega_1}\right) - \omega_2^{-s} \zeta\left(s, \frac{\omega_2 + z}{\omega_2}\right) \\ &\quad + \zeta_2\left(s, \frac{\omega_1 + \omega_2}{3} + z; \omega_1, \omega_2\right) - \zeta_2\left(s, \frac{2(\omega_1 + \omega_2)}{3} + z; \omega_1, \omega_2\right). \end{aligned} \quad (5.114)$$

The residues at simple poles $s = 2$ and 1 are

$$\text{Res}_{s=2} \mathfrak{Z}^{4d \mathcal{N}=2 \text{ vec}}(s, z; \omega_1, \omega_2) = 0, \quad (5.115)$$

$$\text{Res}_{s=1} \mathfrak{Z}^{4d \mathcal{N}=2 \text{ vec}}(s, z; \omega_1, \omega_2) = -\frac{2}{3} \frac{\omega_1 + \omega_2}{\omega_1 \omega_2}. \quad (5.116)$$

We have the non-trivial Zeta-index

$$\mathfrak{Z}^{4d \mathcal{N}=2 \text{ vec}}(0, 0; \omega_1, \omega_2) = 1. \quad (5.117)$$

For $s = -1$ and $z = 0$ one finds

$$\mathfrak{Z}^{4d \mathcal{N}=2 \text{ vec}}(-1, 0; \omega_1, \omega_2) = \frac{5}{54}(\omega_1 + \omega_2) + \frac{1}{81} \frac{\omega_1^3 + \omega_2^3}{\omega_1 \omega_2}. \quad (5.118)$$

We obtain from (5.116) and (5.118)

$$c_{4d} = \frac{1}{6}, \quad (5.119)$$

$$a_{4d} = \frac{5}{24}. \quad (5.120)$$

They agree with the 4d central charges of the $\mathcal{N} = 2$ $U(1)$ vector multiplet. The value $a_{4d}/c_{4d} = 5/4$ hits the upper bound for $\mathcal{N} = 2$ SCFT [118, 114].

The supersymmetric determinant is

$$\mathfrak{D}^{4d \mathcal{N}=2 \text{ vec}}(z; \omega_1, \omega_2) = \frac{2\pi \omega_1^{\frac{1}{2} - \frac{\omega_1 + z}{\omega_1}} \omega_2^{\frac{1}{2} - \frac{\omega_2 + z}{\omega_2}} \Gamma_2\left(\frac{(\omega_1 + \omega_2)}{3} + z; \omega_1, \omega_2\right)}{\Gamma\left(\frac{\omega_1 + z}{\omega_1}\right) \Gamma\left(\frac{\omega_2 + z}{\omega_2}\right) \Gamma_2\left(\frac{2(\omega_1 + \omega_2)}{3} + z; \omega_1, \omega_2\right)} \quad (5.121)$$

The vacuum exponent with $\omega_1 = \omega_2 = 1$ is

$$\mathfrak{D}^{4d \mathcal{N}=2 \text{ vec}}(z; 1, 1) = \frac{2\pi}{3^{\frac{1}{6}}} e^{\frac{2\pi^2 - 3\psi^{(1)}(\frac{1}{3})}{18\sqrt{3}\pi}} = 4.69776 \dots \quad (5.122)$$

5.1.8 4d $\mathcal{N} = 4$ vector multiplet

The $\mathcal{N} = 4$ $U(1)$ vector multiplet comprises the $\mathcal{N} = 2$ $U(1)$ vector multiplet and the $\mathcal{N} = 2$ hypermultiplet.

The supersymmetric index reads

$$\begin{aligned} \mathcal{I}^{4d \mathcal{N}=4 \text{ vec}}(v, w; p; q) &= \mathcal{I}^{4d \mathcal{N}=2 \text{ vec}}\left(y = \sqrt{\frac{v}{w}}; p; q\right) \times \mathcal{I}^{4d \mathcal{N}=2 \text{ hyp}}\left(y = \sqrt{\frac{v}{w}}, z = \sqrt{vw}; p; q\right) \\ &= (p; p)_\infty (q; q)_\infty \Gamma((pq)^{\frac{1}{3}}v; p, q) \Gamma((pq)^{\frac{1}{3}}w^{-1}; p, q) \Gamma((pq)^{\frac{1}{3}}wv^{-1}; p, q). \end{aligned} \quad (5.123)$$

The single-particle index is

$$i^{4d \mathcal{N}=4 \text{ vec}}(v, w; p; q) = \frac{(pq)^{\frac{1}{3}}\left(v + \frac{1}{w} + \frac{w}{v}\right) - (pq)^{\frac{2}{3}}\left(w + \frac{1}{v} + \frac{v}{w}\right) - p - q + 2pq}{(1-p)(1-q)}. \quad (5.124)$$

The supersymmetric zeta function is

$$\begin{aligned} \mathfrak{Z}^{4d \mathcal{N}=4 \text{ vec}}(s, z; \omega_1, \omega_2) &= -\omega_1^{-s} \zeta\left(s, \frac{\omega_1 + z}{\omega_1}\right) - \omega_2^{-s} \zeta\left(s, \frac{\omega_2 + z}{\omega_2}\right) \\ &\quad + 3 \left[\zeta_2\left(s, \frac{\omega_1 + \omega_2}{3} + z; \omega_1, \omega_2\right) - \zeta_2\left(s, \frac{2(\omega_1 + \omega_2)}{3} + z; \omega_1, \omega_2\right) \right]. \end{aligned} \quad (5.125)$$

The residues at simple poles $s = 2$ and 1 vanish

$$\text{Res}_{s=2} \mathfrak{Z}^{4d \mathcal{N}=4 \text{ vec}}(s, z; \omega_1, \omega_2) = 0, \quad (5.126)$$

$$\text{Res}_{s=1} \mathfrak{Z}^{4d \mathcal{N}=4 \text{ vec}}(s, z; \omega_1, \omega_2) = 0. \quad (5.127)$$

The supersymmetric zeta value with $s = 0$ and $z = 0$ is

$$\mathfrak{Z}^{4d \mathcal{N}=4 \text{ vec}}(0, 0; \omega_1, \omega_2) = 1. \quad (5.128)$$

For $s = -1$ and $z = 0$ we have

$$\mathfrak{Z}^{4d \mathcal{N}=4 \text{ vec}}(-1, 0; \omega_1, \omega_2) = \frac{1}{9}(\omega_1 + \omega_2) + \frac{1}{27} \frac{\omega_1^3 + \omega_2^3}{\omega_1 \omega_2}. \quad (5.129)$$

Plugging (5.127) and (5.129) into the formulae (5.10) and (5.11), we get the central charges

$$c_{4d} = a_{4d} = \frac{1}{4}. \quad (5.130)$$

The supersymmetric determinant is

$$\mathfrak{D}^{4d \mathcal{N}=4 \text{ vec}}(z; \omega_1, \omega_2) = \frac{2\pi \omega_1^{\frac{1}{2} - \frac{\omega_1+z}{\omega_1}} \omega_2^{\frac{1}{2} - \frac{\omega_2+z}{\omega_2}} \Gamma_2\left(\frac{(\omega_1+\omega_2)}{3} + z; \omega_1, \omega_2\right)^3}{\Gamma\left(\frac{\omega_1+z}{\omega_1}\right) \Gamma\left(\frac{\omega_2+z}{\omega_2}\right) \Gamma_2\left(\frac{2(\omega_1+\omega_2)}{3} + z; \omega_1, \omega_2\right)^3} \quad (5.131)$$

The vacuum exponent with $\omega_1 = \omega_2 = 1$ is

$$\mathfrak{D}^{4d \mathcal{N}=4 \text{ vec}}(0; 1, 1) = \frac{2\pi}{\sqrt{3}} e^{\frac{2\pi^2 - 3\psi^{(1)}(\frac{1}{3})}{6\sqrt{3}\pi}} = 2.62611 \dots \quad (5.132)$$

5.2 Zeta functions for Schur and Macdonald indices

For $4d \mathcal{N} \geq 2$ supersymmetric field theories, one can define special types of supersymmetric indices by taking certain fugacity limits of the supersymmetric indices so that only certain short multiplets contribute to the indices [119, 120]. The $\mathcal{N} = 2$ supersymmetric index is given by

$$\mathcal{I}^{4d \mathcal{N}=2}(x; y; p; q) = \text{Tr}(-1)^F p^{j_3 + \bar{j}_3 + \frac{1}{3}(2l - \frac{r}{2})} q^{j_3 - \bar{j}_3 + \frac{1}{3}(2l - \frac{r}{2})} y^{2l+r} \prod_{\alpha} x_{\alpha}^{f_{\alpha}}, \quad (5.133)$$

where l and r stand for the Cartan generators of $SU(2)_R$ and $U(1)_r$ R-symmetry. These are related to the $\mathcal{N} = 1$ R-charge R as $R = \frac{2}{3}(-2l + \frac{r}{2})$. Replacement $y \rightarrow (pq)^{-\frac{1}{3}} \tilde{q}^{\frac{1}{2}}$ leads to

$$\mathcal{I}^{4d \mathcal{N}=2}(x; \tilde{q}; p; q) = \text{Tr}(-1)^F p^{j_3 + \bar{j}_3 - \frac{r}{2}} q^{j_3 - \bar{j}_3 - \frac{r}{2}} \tilde{q}^{l + \frac{r}{2}} \prod_{\alpha} x_{\alpha}^{f_{\alpha}}. \quad (5.134)$$

If we take the limit $p \rightarrow 0$ of the index (5.134), we obtain the Macdonald index [120]⁹

$$\mathcal{I}_{\text{Mac}}(x; \tilde{q}; q) = \mathcal{I}^{4d \mathcal{N}=2}(x; \tilde{q}; 0; q). \quad (5.135)$$

Furthermore, if we set \tilde{q} to q , the Macdonald index reduces to the Schur index

$$\mathcal{I}_{\text{Schur}}(x; q) = \mathcal{I}_{\text{Mac}}(x; q; q). \quad (5.136)$$

While the Schur index is typically associated with the Higgs branch of the theory, it also encodes some properties of the Coulomb branch [121, 122, 123, 124]. In [124] Deb and Razamat propose that the generalized Schur partition function may encode the Schur indices of the theories that are obtained from a seed theory by tuning to singular loci of its Coulomb branch. The generalized Schur partition function, which we denote by $\mathcal{I}_{\text{Schur}_{\alpha}}(x; q)$ is obtained from (5.134) by setting

$$\tilde{q} = (qp)^{1 + \frac{\alpha}{\log q} \epsilon} \quad (5.137)$$

⁹The fugacity t in [120] corresponds to \tilde{q} .

and replacing p with $1 - \epsilon$ and then taking $\epsilon \rightarrow 0$. Here α is non-negative real value.¹⁰ When $\alpha = 1$, it reduces to the ordinary Schur index.

We define the supersymmetric zeta functions $\mathfrak{Z}_{\text{Schur}}(s, z)$ (resp. $\mathfrak{Z}_{\text{Schur}_\alpha}(s, z)$) for the 4d $\mathcal{N} = 2$ supersymmetric field theories by taking the Mellin transform of the plethystic logarithms of the unflavored Schur indices (resp the unflavored generalized Schur partition functions). Similarly, we define $\mathfrak{Z}_{\text{Mac}}(s, z; \omega_1, \omega_2)$ as the Mellin transformed plethystic logarithms of the Macdonald indices with $q = q^{\omega_1}$ and $\tilde{q} = q^{\omega_2}$. Thus we have $\mathfrak{Z}_{\text{Mac}}(s, z; 1, 1) = \mathfrak{Z}_{\text{Schur}}(s, z)$. These supersymmetric zeta functions have simple pole at $s = 1$. In addition, for the Schur index, the generalized Schur partition function and Macdonald index we denote the associated supersymmetric determinants by $\mathfrak{D}_{\text{Schur}}(z)$, $\mathfrak{D}_{\text{Schur}_\alpha}(z)$ and $\mathfrak{D}_{\text{Mac}}(z; \omega_1, \omega_2)$ respectively.

Again the leading behaviors in the Cardy-like limits of the Schur and Macdonald indices are captured by the supersymmetric zeta functions. It follows from (2.47) and (2.64) that

$$\begin{aligned} & \log \mathcal{I}_{\text{Mac}}(1; q^{\omega_2}; q^{\omega_1}) \\ & \sim \frac{\pi^2 \text{Res}_{s=1} \mathfrak{Z}_{\text{Mac}}(s, 0; \omega_1, \omega_2)}{6\beta} - \mathfrak{Z}_{\text{Mac}}(0, 0; \omega_1, \omega_2) \log \beta + \log \mathfrak{D}_{\text{Mac}}(0; \omega_1, \omega_2), \end{aligned} \quad (5.138)$$

$$\begin{aligned} & d_{\text{Schur}}(n) \\ & \sim \left(\frac{\text{Res}_{s=1} \mathfrak{Z}_{\text{Schur}}(s, 0)}{96n^3} \right)^{\frac{1}{4}} \left(\frac{6n}{\pi^2 \text{Res}_{s=1} \mathfrak{Z}_{\text{Schur}}(s, 0)} \right)^{\frac{\mathfrak{Z}_{\text{Schur}}(0,0)}{2}} \\ & \quad \times \mathfrak{D}_{\text{Schur}}(0) \exp \left[2\pi \sqrt{\frac{\text{Res}_{s=1} \mathfrak{Z}_{\text{Schur}}(s, 0)}{6}} n^{\frac{1}{2}} \right]. \end{aligned} \quad (5.139)$$

For the superconformal field theories the leading Cardy-like asymptotics of the Macdonald and Schur indices are universally controlled by the anomalies. In other words, we have

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{Schur}}(s, 0) = 48(c_{4d} - a_{4d}), \quad (5.140)$$

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{Mac}}(s, 0; \omega_1, \omega_2) = \frac{48}{\omega_1} (c_{4d} - a_{4d}). \quad (5.141)$$

The relation (5.140) follows from the universal formula (2.47) and the leading terms in the Cardy-like limit of the Schur indices, which are discussed in [46, 47, 48, 125, 126, 127, 128, 22]. On the other hand, we propose the universal relation (5.141) associated with the Macdonald indices.¹¹

¹⁰The limit for integer value of α was discussed in [48].

¹¹Also see [23] for the study of the Cardy-like limits of the Macdonald indices.

Also we propose that the supersymmetric zeta values with $s = -1$ and $z = 0$ are universally given by the anomalies

$$\mathfrak{Z}_{\text{Schur}}(-1, 0) = c_{4d}, \quad (5.142)$$

$$\mathfrak{Z}_{\text{Mac}}(-1, 0; \omega_1, \omega_2) = -4\omega_1(c_{4d} - a_{4d}) + 12\omega_2(c_{4d} - a_{4d}) - \frac{\omega_2^2}{\omega_1}(7c_{4d} - 8a_{4d}). \quad (5.143)$$

From (2.76) the supersymmetric zeta values (5.142) give rise to the supersymmetric Casimir energies of the Schur indices [66]. Similarly, the supersymmetric zeta values (5.143) are expected to produce the supersymmetric Casimir energies of the Macdonald indices.

From (5.140) and (5.142) the 4d central charges of $\mathcal{N} = 2$ SCFTs can be expressed in terms of the supersymmetric zeta functions as

$$c_{4d} = \mathfrak{Z}_{\text{Schur}}(-1, 0), \quad (5.144)$$

$$a_{4d} = \mathfrak{Z}_{\text{Schur}}(-1, 0) - \frac{\text{Res}_{s=1} \mathfrak{Z}_{\text{Schur}}(s, 0)}{48}. \quad (5.145)$$

It is demonstrated [129] that there exists the 2d chiral algebra or vertex operator algebra (VOA) of central charge

$$c_{2d} = -12c_{4d}. \quad (5.146)$$

associated with any 4d $\mathcal{N} = 2$ SCFT. The unflavored Schur index is identical to the (normalized) vacuum character of the associated VOA

$$\chi_{\text{vac}}(q) = \mathcal{I}_{\text{Schur}}(q). \quad (5.147)$$

If the vacuum character is annihilated by a linear modular differential equation (LMDE), there exists a vector-valued modular form whose components $\chi_i(\tau)$ obey

$$\chi_i(\tau) = \sum_j \mathcal{S}_{ij} \chi_j(-1/\tau) \quad (5.148)$$

with $\chi_0(\tau) = \chi_{\text{vac}}(q = e^{2\pi i \tau})$. When the components $\chi_i(\tau)$ are expanded with respect to the nome $q = e^{2\pi i \tau}$, one can find the smallest exponent h_* among the expanded components. The exponent h_* is known to encode the leading asymptotic behavior and it is related to the 4d central charges as [125]

$$a_{4d} = \frac{1}{48}(24h_* - 5c_{2d}). \quad (5.149)$$

From the perspective of the associated VOA, the formulae (5.144) and (5.145) can be rewritten as

$$c_{2d} = -12\mathfrak{Z}_{\text{Schur}}(-1, 0), \quad (5.150)$$

$$h_* = -\frac{1}{2}\mathfrak{Z}_{\text{Schur}}(-1, 0) - \frac{1}{24}\text{Res}_{s=1} \mathfrak{Z}_{\text{Schur}}(s, 0). \quad (5.151)$$

5.2.1 4d $\mathcal{N} = 2$ hypermultiplet

The Schur index, the generalized Schur partition function and Macdonald index of the $\mathcal{N} = 2$ hypermultiplet are given by

$$\mathcal{I}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(x; q) = \frac{1}{(q^{\frac{1}{2}}x; q)_{\infty}(q^{\frac{1}{2}}x^{-1}; q)_{\infty}}, \quad (5.152)$$

$$\mathcal{I}_{\text{Schur}_{\alpha}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(x; q) = \frac{1}{(q^{\frac{1}{2}}x; q)_{\infty}^{\alpha}(q^{\frac{1}{2}}x^{-1}; q)_{\infty}^{\alpha}}, \quad (5.153)$$

$$\mathcal{I}_{\text{Mac}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(x; \tilde{q}; q) = \frac{1}{(\tilde{q}^{\frac{1}{2}}x; q)_{\infty}(\tilde{q}^{\frac{1}{2}}x^{-1}; q)_{\infty}}. \quad (5.154)$$

The single-particle indices are

$$i_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(x; q) = \frac{q^{\frac{1}{2}}(x + x^{-1})}{1 - q}, \quad (5.155)$$

$$i_{\text{Schur}_{\alpha}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(x; q) = \frac{\alpha q^{\frac{1}{2}}(x + x^{-1})}{1 - q}, \quad (5.156)$$

$$i_{\text{Mac}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(x; \tilde{q}; q) = \frac{\tilde{q}^{\frac{1}{2}}(x + x^{-1})}{1 - q}. \quad (5.157)$$

The supersymmetric zeta functions are

$$\mathfrak{Z}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(s, z) = 2\zeta\left(s, \frac{1}{2} + z\right), \quad (5.158)$$

$$\mathfrak{Z}_{\text{Schur}_{\alpha}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(s, z) = 2\alpha\zeta\left(s, \frac{1}{2} + z\right), \quad (5.159)$$

$$\mathfrak{Z}_{\text{Mac}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(s, z; \omega_1, \omega_2) = 2\omega_1^{-s}\zeta\left(s, \frac{\omega_2}{2\omega_1} + \frac{z}{\omega_1}\right). \quad (5.160)$$

The residues at $s = 1$ are

$$\text{Res}_{s=1}\mathfrak{Z}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(s, z) = 2, \quad (5.161)$$

$$\text{Res}_{s=1}\mathfrak{Z}_{\text{Schur}_{\alpha}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(s, z) = 2\alpha, \quad (5.162)$$

$$\text{Res}_{s=1}\mathfrak{Z}_{\text{Mac}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(s, z) = \frac{2}{\omega_1}. \quad (5.163)$$

For $s = 0$ and $z = 0$ the supersymmetric zeta values vanish

$$\mathfrak{Z}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(0, 0) = 0, \quad (5.164)$$

$$\mathfrak{Z}_{\text{Schur}_{\alpha}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(0, 0) = 0, \quad (5.165)$$

$$\mathfrak{Z}_{\text{Mac}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(0, 0) = 1 - \frac{\omega_2}{\omega_1}. \quad (5.166)$$

For $s = -1$ and $z = 0$ we have

$$\mathfrak{Z}_{\text{Schur}}^{4d \mathcal{N}=2 \text{ hyp}}(-1, 0) = \frac{1}{12}, \quad (5.167)$$

$$\mathfrak{Z}_{\text{Schur}_\alpha}^{4d \mathcal{N}=2 \text{ hyp}}(-1, 0) = \frac{\alpha}{12}, \quad (5.168)$$

$$\mathfrak{Z}_{\text{Mac}}^{4d \mathcal{N}=2 \text{ hyp}}(-1, 0) = -\frac{\omega_1}{6} + \frac{\omega_2}{2} - \frac{\omega_2^2}{4\omega_1}. \quad (5.169)$$

Plugging (5.161) and (5.167) into (5.144) and (5.145), we reproduce the 4d central charges (5.108) and (5.109). If the generalized Schur partition function for rational value α coincides with the Schur index for some SCFT, it should have the central charges $a_{4d} = \frac{\alpha}{24}$ and $c_{4d} = \frac{\alpha}{12}$ so that they hit the lower bound $a_{4d}/c_{4d} = 1/2$.

The supersymmetric determinants are

$$\mathfrak{D}_{\text{Schur}}^{4d \mathcal{N}=2 \text{ hyp}}(z) = \frac{\Gamma\left(\frac{1}{2} + z\right)^2}{2\pi}, \quad (5.170)$$

$$\mathfrak{D}_{\text{Schur}_\alpha}^{4d \mathcal{N}=2 \text{ hyp}}(z) = \frac{\Gamma\left(\frac{1}{2} + z\right)^{2\alpha}}{(2\pi)^\alpha}, \quad (5.171)$$

$$\mathfrak{D}_{\text{Mac}}^{4d \mathcal{N}=2 \text{ hyp}}(z; \omega_1, \omega_2) = \omega_1^{\frac{2z - \omega_1 + \omega_2}{\omega_1}} \frac{\Gamma\left(\frac{\omega_2}{2\omega_1} + \frac{z}{\omega_1}\right)^2}{2\pi}. \quad (5.172)$$

The vacuum exponents are

$$\mathfrak{D}_{\text{Schur}}^{4d \mathcal{N}=2 \text{ hyp}}(0) = \frac{1}{2}, \quad (5.173)$$

$$\mathfrak{D}_{\text{Schur}_\alpha}^{4d \mathcal{N}=2 \text{ hyp}}(0) = \frac{1}{2^\alpha}, \quad (5.174)$$

$$\mathfrak{D}_{\text{Mac}}^{4d \mathcal{N}=2 \text{ hyp}}(0; \omega_1, \omega_2) = \omega_1^{\frac{\omega_2}{\omega_1} - 1} \frac{\Gamma\left(\frac{\omega_2}{2\omega_1}\right)^2}{2\pi}. \quad (5.175)$$

5.2.2 4d $\mathcal{N} = 2$ vector multiplet

For the 4d $\mathcal{N} = 2$ $U(1)$ vector multiplet we have the Schur and Macdonald indices

$$\mathcal{I}_{\text{Schur}}^{4d \mathcal{N}=2 \text{ vec}}(q) = (q)_\infty^2, \quad (5.176)$$

$$\mathcal{I}_{\text{Mac}}^{4d \mathcal{N}=2 \text{ vec}}(\tilde{q}; q) = (\tilde{q}; q)_\infty (q; q)_\infty. \quad (5.177)$$

The single-particle indices are

$$i_{\text{Schur}}^{4d \mathcal{N}=2 \text{ vec}}(q) = -\frac{2q}{1-q}, \quad (5.178)$$

$$i_{\text{Mac}}^{4d \mathcal{N}=2 \text{ vec}}(\tilde{q}; q) = -\frac{\tilde{q} + q}{1-q}. \quad (5.179)$$

The supersymmetric zeta functions are

$$\mathfrak{Z}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ vec}}(s, z) = -2\zeta(s, 1+z), \quad (5.180)$$

$$\mathfrak{Z}_{\text{Mac}}^{4\text{d } \mathcal{N}=2 \text{ vec}}(s, z; \omega_1, \omega_2) = -\omega_1^{-s} \zeta\left(s, \frac{\omega_1 + z}{\omega_1}\right) - \omega_1^{-s} \zeta\left(s, \frac{\omega_2 + z}{\omega_1}\right). \quad (5.181)$$

The residues at a simple pole $s = 1$ are

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ vec}}(s, z) = -2, \quad (5.182)$$

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{Mac}}^{4\text{d } \mathcal{N}=2 \text{ vec}}(s, z; \omega_1, \omega_2) = -\frac{2}{\omega_1}. \quad (5.183)$$

The supersymmetric zeta values with $s = 0$ and $z = 0$ are

$$\mathfrak{Z}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ vec}}(0, 0) = 1, \quad (5.184)$$

$$\mathfrak{Z}_{\text{Mac}}^{4\text{d } \mathcal{N}=2 \text{ vec}}(0, 0; \omega_1, \omega_2) = \frac{\omega_2}{\omega_1}. \quad (5.185)$$

The supersymmetric zeta values with $s = -1$ and $z = 0$ are

$$\mathfrak{Z}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ vec}}(-1, 0) = \frac{1}{6}, \quad (5.186)$$

$$\mathfrak{Z}_{\text{Mac}}^{4\text{d } \mathcal{N}=2 \text{ vec}}(-1, 0; \omega_1, \omega_2) = \frac{\omega_1}{6} - \frac{\omega_2}{2} + \frac{\omega_2^2}{2\omega_1}. \quad (5.187)$$

From (5.182) and (5.186) as well as the formulae (5.144) and (5.145), the central charges (5.119) and (5.120) are reproduced.

The supersymmetric determinants are evaluated as

$$\mathfrak{D}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ vec}}(z) = \frac{2\pi}{\Gamma(1+z)^2}, \quad (5.188)$$

$$\mathfrak{D}_{\text{Mac}}^{4\text{d } \mathcal{N}=2 \text{ vec}}(z; \omega_1, \omega_2) = \frac{2\pi\omega_1^{-\frac{2z+\omega_2}{\omega_1}}}{\Gamma\left(\frac{z+\omega_1}{\omega_1}\right)\Gamma\left(\frac{z+\omega_2}{\omega_1}\right)}. \quad (5.189)$$

The vacuum exponents are

$$\mathfrak{D}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ vec}}(0) = 2\pi, \quad (5.190)$$

$$\mathfrak{D}_{\text{Mac}}^{4\text{d } \mathcal{N}=2 \text{ vec}}(0; \omega_1, \omega_2) = \frac{2\pi\omega_1^{-\frac{\omega_2}{\omega_1}}}{\Gamma\left(\frac{\omega_2}{\omega_1}\right)}. \quad (5.191)$$

5.2.3 4d $\mathcal{N} = 4$ vector multiplet

The Schur and Macdonald indices for the 4d $\mathcal{N} = 4$ $U(1)$ vector multiplet are

$$\begin{aligned}\mathcal{I}_{\text{Schur}}^{4\text{d } \mathcal{N}=4 \text{ vec}}(t; q) &= \mathcal{I}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ vec}}(q) \times \mathcal{I}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(x = t^2; q) \\ &= \frac{(q)_\infty^2}{(q^{\frac{1}{2}}t^2; q)_\infty (q^{\frac{1}{2}}t^{-2}; q)_\infty},\end{aligned}\tag{5.192}$$

$$\begin{aligned}\mathcal{I}_{\text{Mac}}^{4\text{d } \mathcal{N}=4 \text{ vec}}(t; \tilde{q}; q) &= \mathcal{I}_{\text{Mac}}^{4\text{d } \mathcal{N}=2 \text{ vec}}(\tilde{q}; q) \times \mathcal{I}_{\text{Mac}}^{4\text{d } \mathcal{N}=2 \text{ hyp}}(x = t^2; \tilde{q}; q) \\ &= \frac{(\tilde{q}; q)_\infty (q; q)_\infty}{(\tilde{q}^{\frac{1}{2}}t^2; q)_\infty (\tilde{q}^{\frac{1}{2}}t^{-2}; q)_\infty}.\end{aligned}\tag{5.193}$$

The single-particle indices read

$$i_{\text{Schur}}^{4\text{d } \mathcal{N}=4 \text{ vec}}(t; q) = \frac{-2q + q^{\frac{1}{2}}(t^2 + t^{-2})}{1 - q},\tag{5.194}$$

$$i_{\text{Mac}}^{4\text{d } \mathcal{N}=4 \text{ vec}}(t; \tilde{q}; q) = \frac{-\tilde{q} - q + \tilde{q}^{\frac{1}{2}}(t^2 + t^{-2})}{1 - q}.\tag{5.195}$$

The supersymmetric zeta functions are

$$\mathfrak{Z}_{\text{Schur}}^{4\text{d } \mathcal{N}=4 \text{ vec}}(s, z) = -2\zeta(s, 1 + z) + 2\zeta\left(s, \frac{1}{2} + z\right),\tag{5.196}$$

$$\begin{aligned}\mathfrak{Z}_{\text{Mac}}^{4\text{d } \mathcal{N}=4 \text{ vec}}(s, z; \omega_1, \omega_2) &= -\omega_1^{-s}\zeta\left(s, \frac{\omega_1 + z}{\omega_1}\right) - \omega_1^{-s}\zeta\left(s, \frac{\omega_2 + z}{\omega_1}\right) \\ &\quad + 2\omega_1^{-s}\zeta\left(s, \frac{\omega_2}{2\omega_1} + \frac{z}{\omega_1}\right).\end{aligned}\tag{5.197}$$

The residues at $s = 1$ vanish

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{Schur}}^{4\text{d } \mathcal{N}=4 \text{ vec}}(s, z) = 0,\tag{5.198}$$

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{Mac}}^{4\text{d } \mathcal{N}=4 \text{ vec}}(s, z; \omega_1, \omega_2) = 0.\tag{5.199}$$

For $s = 0$ and $z = 0$ the supersymmetric zeta values are

$$\mathfrak{Z}_{\text{Schur}}^{4\text{d } \mathcal{N}=4 \text{ vec}}(0, 0) = 1,\tag{5.200}$$

$$\mathfrak{Z}_{\text{Mac}}^{4\text{d } \mathcal{N}=4 \text{ vec}}(0, 0; \omega_1, \omega_2) = 1.\tag{5.201}$$

For $s = -1$ and $z = 0$ we have

$$\mathfrak{Z}_{\text{Schur}}^{4\text{d } \mathcal{N}=4 \text{ vec}}(-1, 0) = \frac{1}{4},\tag{5.202}$$

$$\mathfrak{Z}_{\text{Mac}}^{4\text{d } \mathcal{N}=4 \text{ vec}}(-1, 0; \omega_1, \omega_2) = \frac{\omega_2^2}{4\omega_1}.\tag{5.203}$$

Again we obtain from (5.198) and (5.202) the central charges (5.130).

The supersymmetric determinants are

$$\mathfrak{D}_{\text{Schur}}^{4\text{d } \mathcal{N}=4 \text{ vec}}(z) = \frac{\Gamma\left(\frac{1}{2} + z\right)^2}{\Gamma(1+z)^2}, \quad (5.204)$$

$$\mathfrak{D}_{\text{Mac}}^{4\text{d } \mathcal{N}=4 \text{ vec}}(z; \omega_1, \omega_2) = \omega_1^{-1} \frac{\Gamma\left(\frac{\omega_2}{2\omega_1} + \frac{z}{\omega_1}\right)^2}{\Gamma\left(\frac{z+\omega_1}{\omega_1}\right) \Gamma\left(\frac{z+\omega_2}{\omega_1}\right)}. \quad (5.205)$$

The vacuum exponents are

$$\mathfrak{D}_{\text{Schur}}^{4\text{d } \mathcal{N}=4 \text{ vec}}(0) = \pi, \quad (5.206)$$

$$\mathfrak{D}_{\text{Mac}}^{4\text{d } \mathcal{N}=4 \text{ vec}}(0; \omega_1, \omega_2) = \omega_1^{-1} \frac{\Gamma\left(\frac{\omega_2}{2\omega_1}\right)^2}{\Gamma\left(\frac{\omega_2}{\omega_1}\right)}. \quad (5.207)$$

5.2.4 4d $\mathcal{N}=2$ pure SYM theories

Consider the 4d $\mathcal{N}=2$ pure SYM theory of gauge group G . The Schur indices can be evaluated from the matrix integral of the form

$$\mathcal{I}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ } G}(q) = \frac{1}{|\text{Weyl}(G)|} (q)_{\infty}^{2\text{rank}(G)} \oint \prod_{\alpha \in \text{root}(G)} \frac{ds}{2\pi i s} (s^{\alpha}; q) (qs^{\alpha}; q)_{\infty}, \quad (5.208)$$

where $|\text{Weyl}(G)|$ is the order of the Weyl group of G . They are evaluated as [130, 131]

$$\mathcal{I}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ } SU(N)}(q) = \frac{(q^{2N}; q^{2N})_{\infty}^N}{(q^2; q^2)_{\infty}}, \quad (5.209)$$

$$\mathcal{I}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ } SO(2N+1)}(q) = \frac{(q^{4N-2}; q^{4N-2})_{\infty}^{N-1} (q^{8N-4}; q^{8N-4})_{\infty} (q^4; q^4)_{\infty}}{(q^2; q^2)_{\infty}}, \quad (5.210)$$

$$\mathcal{I}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ } USp(2N)}(q) = \frac{(q^{2N+2}; q^{2N+2})_{\infty} (q^{4N+4}; q^{4N+4})_{\infty}^{N-1} (q^4; q^4)_{\infty}}{(q^2; q^2)_{\infty}}, \quad (5.211)$$

$$\mathcal{I}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ } SO(2N)}(q) = \frac{(q^{4N-4}; q^{4N-4})_{\infty}^{N+1} (q^4; q^4)_{\infty}}{(q^{2N-2}; q^{2N-2})_{\infty} (q^2; q^2)_{\infty}}, \quad (5.212)$$

$$\mathcal{I}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ } G_2}(q) = \frac{(q^4; q^4)_{\infty} (q^6; q^6)_{\infty} (q^8; q^8)_{\infty} (q^{24}; q^{24})_{\infty}}{(q^2; q^2)_{\infty} (q^{12}; q^{12})_{\infty}}, \quad (5.213)$$

$$\mathcal{I}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ } F_4}(q) = \frac{(q^4; q^4)_{\infty} (q^6; q^6)_{\infty} (q^{18}; q^{18})_{\infty}^2 (q^{36}; q^{36})_{\infty}^2}{(q^2; q^2)_{\infty} (q^{12}; q^{12})_{\infty}}, \quad (5.214)$$

$$\mathcal{I}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ } E_6}(q) = \frac{(q^4; q^4)_{\infty} (q^6; q^6)_{\infty} (q^{24}; q^{24})_{\infty}^7}{(q^2; q^2)_{\infty} (q^8; q^8)_{\infty} (q^{12}; q^{12})_{\infty}}, \quad (5.215)$$

$$\mathcal{I}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ } E_7}(q) = \frac{(q^4; q^4)_{\infty} (q^6; q^6)_{\infty} (q^{36}; q^{36})_{\infty}^8}{(q^2; q^2)_{\infty} (q^{12}; q^{12})_{\infty} (q^{18}; q^{18})_{\infty}}, \quad (5.216)$$

$$\mathcal{I}_{\text{Schur}}^{4\text{d } \mathcal{N}=2 \text{ } E_8}(q) = \frac{(q^4; q^4)_{\infty} (q^6; q^6)_{\infty} (q^{10}; q^{10})_{\infty} (q^{60}; q^{60})_{\infty}^9}{(q^2; q^2)_{\infty} (q^{12}; q^{12})_{\infty} (q^{20}; q^{20})_{\infty} (q^{30}; q^{30})_{\infty}}. \quad (5.217)$$

The single-particle indices are

$$i_{\text{Schur}}^{4d \mathcal{N}=2 \text{ } SU(N)}(q) = -\frac{Nq^{2N}}{1-q^{2N}} + \frac{q^2}{1-q^2}, \quad (5.218)$$

$$i_{\text{Schur}}^{4d \mathcal{N}=2 \text{ } SO(2N+1)}(q) = -\frac{(N-1)q^{4N-2}}{1-q^{4N-2}} - \frac{q^{8N-4}}{1-q^{8N-4}} - \frac{q^4}{1-q^4} + \frac{q^2}{1-q^2}, \quad (5.219)$$

$$i_{\text{Schur}}^{4d \mathcal{N}=2 \text{ } USp(2N)}(q) = -\frac{q^{2N+2}}{1-q^{2N+2}} - \frac{(N-1)q^{4N+4}}{1-q^{4N+4}} - \frac{q^4}{1-q^4} + \frac{q^2}{1-q^2}, \quad (5.220)$$

$$i_{\text{Schur}}^{4d \mathcal{N}=2 \text{ } SO(2N)}(q) = -\frac{(N-1)q^{4N-4}}{1-q^{4N-4}} - \frac{q^4}{1-q^4} + \frac{q^{2N-2}}{1-q^{2N-2}} + \frac{q^2}{1-q^2}, \quad (5.221)$$

$$i_{\text{Schur}}^{4d \mathcal{N}=2 \text{ } G_2}(q) = -\frac{q^4}{1-q^4} - \frac{q^6}{1-q^6} - \frac{q^8}{1-q^8} - \frac{q^{24}}{1-q^{24}} + \frac{q^2}{1-q^2} + \frac{q^{12}}{1-q^{12}}, \quad (5.222)$$

$$i_{\text{Schur}}^{4d \mathcal{N}=2 \text{ } F_4}(q) = -\frac{q^4}{1-q^4} - \frac{q^6}{1-q^6} - \frac{2q^{18}}{1-q^{18}} - \frac{2q^{36}}{1-q^{36}} + \frac{q^2}{1-q^2} + \frac{q^{12}}{1-q^{12}}, \quad (5.223)$$

$$i_{\text{Schur}}^{4d \mathcal{N}=2 \text{ } E_6}(q) = -\frac{q^4}{1-q^4} - \frac{q^6}{1-q^6} - 7\frac{q^{24}}{1-q^{24}} + \frac{q^2}{1-q^2} + \frac{q^8}{1-q^8} + \frac{q^{12}}{1-q^{12}}, \quad (5.224)$$

$$i_{\text{Schur}}^{4d \mathcal{N}=2 \text{ } E_7}(q) = -\frac{q^4}{1-q^4} - \frac{q^6}{1-q^6} - 8\frac{q^{36}}{1-q^{36}} + \frac{q^2}{1-q^2} + \frac{q^{12}}{1-q^{12}} + \frac{q^{18}}{1-q^{18}}, \quad (5.225)$$

$$i_{\text{Schur}}^{4d \mathcal{N}=2 \text{ } E_8}(q) = -\frac{q^4}{1-q^4} - \frac{q^6}{1-q^6} - \frac{q^{10}}{1-q^{10}} - 9\frac{q^{60}}{1-q^{60}} + \frac{q^2}{1-q^2} + \frac{q^{12}}{1-q^{12}} + \frac{q^{20}}{1-q^{20}} + \frac{q^{30}}{1-q^{30}}. \quad (5.226)$$

We obtain the supersymmetric zeta functions

$$\mathfrak{Z}_{\text{Schur}}^{4d \mathcal{N}=2 \text{ } SU(N)}(s, z) = -2^{-s} N^{1-s} \zeta\left(s, \frac{2N+z}{2N}\right) + 2^{-s} \zeta\left(s, \frac{2+z}{2}\right), \quad (5.227)$$

$$\begin{aligned} \mathfrak{Z}_{\text{Schur}}^{4d \mathcal{N}=2 \text{ } SO(2N+1)}(s, z) = & -(N-1)(4N-2)^{-s} \zeta\left(s, \frac{4N-2+z}{4N-2}\right) - 4^{-s} \zeta\left(s, \frac{4+z}{4}\right) \\ & - (8N-4)^{-s} \zeta\left(s, \frac{8N-4+z}{8N-4}\right) + 2^{-s} \zeta\left(s, \frac{2+z}{2}\right), \end{aligned} \quad (5.228)$$

$$\begin{aligned} \mathfrak{Z}_{\text{Schur}}^{4d \mathcal{N}=2 \text{ } USp(2N)}(s, z) = & -(2N+2)^{-s} \zeta\left(s, \frac{2N+2+z}{2N+2}\right) - 4^{-s} \zeta\left(s, \frac{4+z}{4}\right) \\ & - (N-1)(4N+4)^{-s} \zeta\left(s, \frac{4N+4+z}{4N+4}\right) + 2^{-s} \zeta\left(s, \frac{2+z}{2}\right), \end{aligned} \quad (5.229)$$

$$\mathfrak{Z}_{\text{Schur}}^{4d \mathcal{N}=2 \text{ } SO(2N)}(s, z) = -(N+1)(4N-4)^{-s} \zeta\left(s, \frac{4N-4+z}{4N-4}\right) - 4^{-s} \zeta\left(s, \frac{4+z}{4}\right)$$

$$+ (2N - 2)^{-s} \zeta \left(s, \frac{2N - 2 + z}{2N - 2} \right) + 2^{-s} \zeta \left(s, \frac{2 + z}{2} \right), \quad (5.230)$$

$$\begin{aligned} \mathfrak{Z}_{\text{Schur}}^{4d \mathcal{N} = 2 G_2}(s, z) = & -4^{-s} \zeta \left(s, \frac{4 + z}{4} \right) - 6^{-s} \zeta \left(s, \frac{6 + z}{6} \right) - 8^{-s} \zeta \left(s, \frac{8 + z}{8} \right) \\ & - 24^{-s} \zeta \left(s, \frac{24 + z}{24} \right) + 2^{-s} \zeta \left(s, \frac{2 + z}{2} \right) + 12^{-s} \zeta \left(s, \frac{12 + z}{12} \right), \end{aligned} \quad (5.231)$$

$$\begin{aligned} \mathfrak{Z}_{\text{Schur}}^{4d \mathcal{N} = 2 F_4}(s, z) = & -2 \cdot (36)^{-s} \zeta \left(s, \frac{36 + z}{36} \right) - 2 \cdot (18)^{-s} \zeta \left(s, \frac{18 + z}{18} \right) \\ & - 6^{-s} \zeta \left(s, \frac{6 + z}{6} \right) - 4^{-s} \zeta \left(s, \frac{4 + z}{4} \right) \\ & + 12^{-s} \zeta \left(s, \frac{12 + z}{12} \right) + 2^{-s} \zeta \left(s, \frac{2 + z}{2} \right), \end{aligned} \quad (5.232)$$

$$\begin{aligned} \mathfrak{Z}_{\text{Schur}}^{4d \mathcal{N} = 2 E_6}(s, z) = & -7 \cdot (24)^{-s} \zeta \left(s, \frac{24 + z}{24} \right) - 6^{-s} \zeta \left(s, \frac{6 + z}{6} \right) - 4^{-s} \zeta \left(s, \frac{4 + z}{4} \right) \\ & + 2^{-s} \zeta \left(s, \frac{2 + z}{2} \right) + 8^{-s} \zeta \left(s, \frac{8 + z}{8} \right) + 12^{-s} \zeta \left(s, \frac{12 + z}{12} \right), \end{aligned} \quad (5.233)$$

$$\begin{aligned} \mathfrak{Z}_{\text{Schur}}^{4d \mathcal{N} = 2 E_7}(s, z) = & -8 \cdot (36)^{-s} \zeta \left(s, \frac{36 + z}{36} \right) - 6^{-s} \zeta \left(s, \frac{6 + z}{6} \right) - 4^{-s} \zeta \left(s, \frac{4 + z}{4} \right) \\ & + 2^{-s} \zeta \left(s, \frac{2 + z}{2} \right) + 12^{-s} \zeta \left(s, \frac{12 + z}{12} \right) + 18^{-s} \zeta \left(s, \frac{18 + z}{18} \right), \end{aligned} \quad (5.234)$$

$$\begin{aligned} \mathfrak{Z}_{\text{Schur}}^{4d \mathcal{N} = 2 E_8}(s, z) = & -9 \cdot (60)^{-s} \zeta \left(s, \frac{60 + z}{60} \right) - 10^{-s} \zeta \left(s, \frac{10 + z}{10} \right) - 6^{-s} \zeta \left(s, \frac{6 + z}{6} \right) \\ & - 4^{-s} \zeta \left(s, \frac{4 + z}{4} \right) + 2^{-s} \zeta \left(s, \frac{2 + z}{2} \right) + 12^{-s} \zeta \left(s, \frac{12 + z}{12} \right) \\ & + 20^{-s} \zeta \left(s, \frac{20 + z}{20} \right) + 30^{-s} \zeta \left(s, \frac{30 + z}{30} \right). \end{aligned} \quad (5.235)$$

Though each term in the supersymmetric zeta functions has a simple pole at $s = 1$, the residue vanishes

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{Schur}}^{4d \mathcal{N} = 2 G}(s, z) = 0. \quad (5.236)$$

The supersymmetric zeta value with $s = 0$ and $z = 0$ is given by

$$\mathfrak{Z}_{\text{Schur}}^{4d \mathcal{N} = 2 G}(0, 0) = \frac{\text{rank}(G)}{2}. \quad (5.237)$$

The supersymmetric zeta value with $s = -1$ and $z = 0$ is

$$\mathfrak{Z}_{\text{Schur}}^{4d \mathcal{N}=2 G}(-1, 0) = \frac{\dim G}{6}. \quad (5.238)$$

According to the formulae (5.144), we get

$$c_{4d} = \frac{\dim G}{6}. \quad (5.239)$$

This is the anomaly contribution for the 4d $\mathcal{N} = 2$ vector multiplet of gauge group G . However, the formula (5.145) does not simply give rise to the anomaly contribution $\frac{5}{24} \dim G$ for a_{4d} as the theory is not conformal.

The supersymmetric determinants are

$$\mathfrak{D}^{4d \mathcal{N}=2 SU(N)}(z) = 2^N \pi^{\frac{N-1}{2}} N^{\frac{N-z}{2}} \frac{\Gamma(1 + \frac{z}{2})}{z^N \Gamma(\frac{z}{2N})^N}, \quad (5.240)$$

$$\mathfrak{D}^{4d \mathcal{N}=2 SO(2N+1)}(z) = \frac{2^{\frac{N+1+z}{2}} (4N-2)^{-\frac{(N-1)(2N-1+z)}{4N-2}} \pi^{\frac{N-1}{2}} \Gamma(\frac{2+z}{4})}{(8N-4)^{\frac{4N-2+z}{8N-4}} (\frac{z}{4N-2})^N \Gamma(\frac{z}{8N-4}) \Gamma(\frac{z}{4N-2})^{N-1}}, \quad (5.241)$$

$$\mathfrak{D}^{4d \mathcal{N}=2 USp(2N)}(z) = \frac{2^{\frac{N(3N-1)-4+z}{2(N+1)}} \pi^{\frac{N-1}{2}} \Gamma(\frac{2+z}{4})}{(N+1)^{\frac{2N+z}{4}} (\frac{z}{N+1})^{N-1} \Gamma(1 + \frac{z}{2N+2}) \Gamma(\frac{z}{4+4N})^{N-1}}, \quad (5.242)$$

$$\mathfrak{D}^{4d \mathcal{N}=2 SO(2N)}(z) = \frac{\pi^{\frac{N-1}{2}} \Gamma(\frac{2+z}{4}) \Gamma(1 + \frac{z}{2(N-1)})}{2^{\frac{N^2+N-2+z}{2(N-1)}} (N-1)^{\frac{2N+z}{4}} \Gamma(1 + \frac{z}{4(N-1)})^{N+1}}, \quad (5.243)$$

$$\mathfrak{D}^{4d \mathcal{N}=2 G_2}(z) = \frac{\pi \Gamma(\frac{2+z}{4})}{2^{2+\frac{z}{6}} 3^{\frac{1}{2}+\frac{z}{8}} \Gamma(1 + \frac{z}{24}) \Gamma(1 + \frac{z}{8}) \Gamma(\frac{6+z}{12})}, \quad (5.244)$$

$$\mathfrak{D}^{4d \mathcal{N}=2 F_4}(z) = \frac{2^{5+\frac{z}{9}} 3^{4-\frac{5z}{12}} \pi^2 \Gamma(\frac{2+z}{4})}{z^4 \Gamma(\frac{z}{36})^2 \Gamma(\frac{z}{18})^2 \Gamma(\frac{6+z}{12})}, \quad (5.245)$$

$$\mathfrak{D}^{4d \mathcal{N}=2 E_6}(z) = \frac{\pi^3 \Gamma(1 + \frac{z}{8}) \Gamma(\frac{2+z}{4})}{2^{6+\frac{z}{6}} 3^{\frac{7}{2}+\frac{3z}{8}} \Gamma(1 + \frac{z}{24})^7 \Gamma(\frac{6+z}{12})}, \quad (5.246)$$

$$\mathfrak{D}^{4d \mathcal{N}=2 E_7}(z) = \frac{\pi^3 \Gamma(\frac{2+z}{4}) \Gamma(\frac{18+z}{36})}{16 \cdot 3^{7+\frac{5z}{12}} \Gamma(1 + \frac{z}{36})^7 \Gamma(\frac{6+z}{12})}, \quad (5.247)$$

$$\mathfrak{D}^{4d \mathcal{N}=2 E_8}(z) = \frac{\pi^4 \Gamma(\frac{2+z}{4}) \Gamma(\frac{30+z}{60})}{16 \cdot 3^{4+\frac{z}{5}} 5^{4+\frac{z}{6}} \Gamma(1 + \frac{z}{60})^8 \Gamma(\frac{6+z}{12}) \Gamma(\frac{10+z}{20})}. \quad (5.248)$$

The vacuum exponents are evaluated as

$$\mathfrak{D}^{4d \mathcal{N}=2 SU(N)}(0) = \frac{\pi^{\frac{N-1}{2}}}{N^{\frac{N}{2}}}, \quad (5.249)$$

$$\mathfrak{D}^{4d \mathcal{N}=2 SO(2N+1)}(0) = \frac{\pi^{\frac{N}{2}}}{2(2N-1)^{\frac{N}{2}}}, \quad (5.250)$$

$$\mathfrak{D}^{4d \mathcal{N}=2 \text{ } USp(2N)}(0) = \frac{\pi^{\frac{N}{2}}}{2^{\frac{N}{2}}(N+1)^{\frac{N}{2}}}, \quad (5.251)$$

$$\mathfrak{D}^{4d \mathcal{N}=2 \text{ } SO(2N)}(0) = \frac{\pi^{\frac{N}{2}}}{2^{\frac{N+2}{2}}(N-1)^{\frac{N}{2}}}, \quad (5.252)$$

$$\mathfrak{D}^{4d \mathcal{N}=2 \text{ } G_2}(0) = \frac{\pi}{4\sqrt{3}}, \quad (5.253)$$

$$\mathfrak{D}^{4d \mathcal{N}=2 \text{ } F_4}(0) = \frac{\pi^2}{162}, \quad (5.254)$$

$$\mathfrak{D}^{4d \mathcal{N}=2 \text{ } E_6}(0) = \frac{\pi^3}{1728\sqrt{3}}, \quad (5.255)$$

$$\mathfrak{D}^{4d \mathcal{N}=2 \text{ } E_7}(0) = \frac{\pi^{\frac{7}{2}}}{34992}, \quad (5.256)$$

$$\mathfrak{D}^{4d \mathcal{N}=2 \text{ } E_8}(0) = \frac{\pi^4}{810000}. \quad (5.257)$$

5.2.5 $J^{h^\vee}[k]$ AD theories

Argyres-Douglas (AD) theories [132, 133] were originally found as the IR fixed points at special singular points on the Coulomb branches of 4d $\mathcal{N}=2$ gauge theories. They typically have no Lagrangian description constructed by the $\mathcal{N}=2$ supermultiplets and contain local operators with fractional scaling dimensions. The AD theories can be engineered from 6d $\mathcal{N}=(2,0)$ theory of type $J=\{A_n, D_n, E_n\}$ on a Riemann surface Σ [134]. The Coulomb branch of the theory is described by a Hitchin system on Σ , which involves the Higgs field Φ [135]. The irregular singularity is associated with an irregular solution to Hitchin's equation with the singular form

$$\Phi = \frac{\mathfrak{t}}{z^{2+k/b}} + \cdots, \quad (5.258)$$

where \mathfrak{t} is a regular semi-simple element of the Lie algebra J . The possible values of b are listed in [136]. The resulting AD theories are labeled by the irregular singularities $J^b[k]$.

It is conjectured [137] that the single-particle Schur index of the $J^{h^\vee}[k]$ AD theory is given by¹²

$$i_{\text{Schur}}^{J^{h^\vee}[k]}(q) = \sum_{i=1}^{\text{rank}(J)} \frac{q^{d_i}}{1-q} - \frac{q^{h^\vee+k}}{1-q^{h^\vee+k}} \sum_{i=1}^{\text{rank}(J)} \sum_{j=0}^{2d_i-2} q^{1-d_i+j}, \quad (5.259)$$

where $\text{rank}(J)$ is the rank of J , h^\vee is the dual Coxeter number of J , and d_i are degrees of the Casimirs of J .

¹²Also see [121, 138] for earlier conjectures with $J=A_{N-1}$.

The supersymmetric zeta function is evaluated as

$$\mathfrak{Z}_{\text{Schur}}^{J^{h^\vee}[k]}(s, z) = \sum_{i=1}^{\text{rank}(J)} \zeta(s, d_i + z) - \sum_{i=1}^{\text{rank}(J)} \sum_{j=0}^{2d_i-2} (h^\vee + k)^{-s} \zeta\left(s, \frac{h^\vee + k + 1 - d_i + j + z}{h^\vee + k}\right). \quad (5.260)$$

It has a simple pole at $s = 1$ with residue

$$\begin{aligned} \text{Res}_{s=1} \mathfrak{Z}_{\text{Schur}}^{J^{h^\vee}[k]}(s, z) &= \text{rank}(J) - \frac{\dim J}{h^\vee + k} \\ &= \frac{(k-1)\text{rank}(J)}{k + h^\vee}. \end{aligned} \quad (5.261)$$

Here we have used the relation (4.188) as well as the relation

$$\dim J = \text{rank}(J)(h^\vee + 1) \quad (5.262)$$

for the simply-laced Lie algebra J . With $s = 0$ and $z = 0$ the supersymmetric zeta value vanishes

$$\mathfrak{Z}_{\text{Schur}}^{J^{h^\vee}[k]}(0, 0) = 0. \quad (5.263)$$

The supersymmetric zeta value for $s = -1$ and $z = 0$ is evaluated as

$$\begin{aligned} \mathfrak{Z}_{\text{Schur}}^{J^{h^\vee}[k]}(-1, 0) &= \sum_{i=1}^{\text{rank}(J)} \frac{(h^\vee + k - 2d_i + 1) [2d_i(h^\vee + k - d_i + 1) - h^\vee - k]}{12(h^\vee + k)} \\ &= \frac{k^2 \dim J}{12(h^\vee + k)} - \frac{1}{12} \text{rank}(J), \end{aligned} \quad (5.264)$$

where we have used the relations

$$\frac{1}{3} h^\vee \dim J = \sum_{i=1}^{\text{rank}(J)} d_i(d_i - 1), \quad (5.265)$$

$$\frac{4}{3} h^\vee \dim J + \text{rank}(J) = \sum_{i=1}^{\text{rank}(J)} (2d_i - 1)^2, \quad (5.266)$$

$$\frac{1}{2} (h^\vee)^2 \dim J = \sum_{i=1}^{\text{rank}(J)} d_i(2d_i - 1)(d_i - 1), \quad (5.267)$$

as well as (4.188). Substituting (5.261) and (5.264) into the formulae (5.144) and (5.145), we get

$$c_{4d} = \frac{k^2 \dim J}{12(h^\vee + k)} - \frac{1}{12} \text{rank}(J), \quad (5.268)$$

$$a_{4d} = \frac{(4k^2 + 1) \dim J}{48(h^\vee + k)} - \frac{5}{48} \text{rank}(J). \quad (5.269)$$

The explicit values are summarized as follows:

$J^{h^\vee}[k]$	$\text{rank}(J)$	h^\vee	d_i	c_{4d}	a_{4d}
$A_{N-1}^N[k]$	$N-1$	N	$i+1$	$\frac{(N-1)(k-1)(N+k+kN)}{12(N+k)}$	$\frac{(N-1)(k-1)(4kN+4N+4k-1)}{48(N+k)}$
$D_N^{2N-2}[k]$	N	$2N-2$	$2, 4, \dots, 2(N-1), N$	$\frac{N(k-1)(2kN+2N-k-2)}{12(2N+k-2)}$	$\frac{N(k-1)(8kN+8N-4k-9)}{48(2N+k-2)}$
$E_6^{12}[k]$	6	12	$2, 5, 6, 8, 9, 12$	$\frac{(k-1)(12+13k)}{2(12+k)}$	$\frac{(k-1)(47+52k)}{8(12+k)}$
$E_7^{18}[k]$	7	18	$2, 6, 8, 10, 12, 14, 18$	$\frac{7(k-1)(18+19k)}{12(18+k)}$	$\frac{7(k-1)(71+76k)}{48(18+k)}$
$E_8^{30}[k]$	8	30	$2, 8, 12, 14, 18, 20, 24, 30$	$\frac{2(k-1)(30+31k)}{3(30+k)}$	$\frac{(k-1)(119+124k)}{6(30+k)}$

(5.270)

We see that all these values of c_{4d} precisely agree with the central charges computed in [139].

The supersymmetric determinant is

$$\begin{aligned} & \mathfrak{D}_{\text{Schur}}^{J^{h^\vee}[k]}(z) \\ &= (2\pi)^{\frac{\dim J - \text{rank}(J)}{2}} (h^\vee + k)^{-\frac{(h^\vee + k + 2z) \dim J}{2(h^\vee + k)}} \frac{\prod_{i=1}^{\text{rank}(J)} \Gamma(d_i + z)}{\prod_{i=1}^{\text{rank}(J)} \prod_{j=0}^{2d_i-2} \Gamma\left(\frac{h^\vee + k + 1 - d_i + j + z}{h^\vee + k}\right)}. \end{aligned} \quad (5.271)$$

The vacuum exponent is

$$\begin{aligned} & \mathfrak{D}_{\text{Schur}}^{J^{h^\vee}[k]}(0) \\ &= (2\pi)^{\frac{\dim J - \text{rank}(J)}{2}} (h^\vee + k)^{-\frac{\dim J}{2}} \frac{\prod_{i=1}^{\text{rank}(J)} \Gamma(d_i)}{\prod_{i=1}^{\text{rank}(J)} \prod_{j=0}^{2d_i-2} \Gamma\left(\frac{h^\vee + k + 1 - d_i + j}{h^\vee + k}\right)}. \end{aligned} \quad (5.272)$$

5.2.6 The generalized (G_1, G_2) AD theories

The generalized (G_1, G_2) Argyres-Douglas theories are constructed in [140] by geometric engineering Type IIB superstring theory on the singular Calabi-Yau hypersurfaces of \mathbb{C}^4 defined as the zero locus of the polynomial associated with a pair of the ADE Lie algebras G_1 and G_2 . They are symmetric under an exchange of G_1 and G_2 .

For $\gcd(h_1^\vee, h_2^\vee) = 1$, where h_i^\vee are the dual Coxeter numbers of G_i , we have the single-particle index of the form [126]

$$i_{\text{Schur}}^{(G_1, G_2)}(q) = \frac{\left(\sum_{i=1}^{\text{rank}(G_1)} q^{d_i^{(1)}-1}\right) \left(\sum_{j=1}^{\text{rank}(G_2)} q^{d_j^{(2)}-1}\right)}{(1 - q^{h_1^\vee + h_2^\vee})}, \quad (5.273)$$

where $d_i^{(i)}$ are degrees of the Casimirs of G_i . The condition $\gcd(h_1^\vee, h_2^\vee) = 1$ is satisfied when $G_1 = J$ and $G_2 = A_{k-1}$ with even k , which is equivalent to the AD theory $J^{h^\vee}[k]$. So we can reproduce the results in the previous section.

The supersymmetric zeta function is given by

$$\mathfrak{Z}_{\text{Schur}}^{(G_1, G_2)}(s, z) = \sum_{i=1}^{\text{rank}(G_1)} \sum_{j=1}^{\text{rank}(G_2)} (h_1^\vee + h_2^\vee)^{-s} \zeta \left(s, \frac{d_i^{(1)} + d_j^{(2)} - 2 + z}{h_1^\vee + h_2^\vee} \right). \quad (5.274)$$

The residue at a simple pole $s = 1$ is evaluated as

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{Schur}}^{(G_1, G_2)}(s, z) = \frac{\text{rank}(G_1) \text{rank}(G_2)}{h_1^\vee + h_2^\vee}. \quad (5.275)$$

The Zeta-index is evaluated as

$$\begin{aligned} & \mathfrak{Z}_{\text{Schur}}^{(G_1, G_2)}(0, 0) \\ &= \frac{\text{rank}(G_1) \text{rank}(G_2) (h_1^\vee + h_2^\vee + 2) - \text{rank}(G_1) \dim G_2 - \text{rank}(G_2) \dim G_1}{2(h_1^\vee + h_2^\vee)} \\ &= 0, \end{aligned} \quad (5.276)$$

where we have used the relations (4.188) and (5.262). The special supersymmetric zeta value with $s = -1$ and $z = 0$ is evaluated as

$$\begin{aligned} & \mathfrak{Z}_{\text{Schur}}^{(G_1, G_2)}(-1, 0) \\ &= \frac{1}{12(h_1^\vee + h_2^\vee)} \left[(h_1^\vee + 3h_2^\vee + 6) \dim G_1 \text{rank}(G_2) + (h_2^\vee + 3h_1^\vee + 6) \dim G_2 \text{rank}(G_1) \right. \\ & \quad \left. - 3 \dim G_1 \dim G_2 - (h_1^\vee + h_2^\vee + 3)^2 \text{rank}(G_1) \text{rank}(G_2) \right] \\ &= \frac{(h_1^\vee + h_2^\vee + h_1^\vee h_2^\vee) \text{rank}(G_1) \text{rank}(G_2)}{12(h_1^\vee + h_2^\vee)}. \end{aligned} \quad (5.277)$$

Again we have used the relations (4.188) and (5.262). According to the formulae (5.144) and (5.145) we obtain

$$c_{4d} = \frac{(h_1^\vee + h_2^\vee + h_1^\vee h_2^\vee) \text{rank}(G_1) \text{rank}(G_2)}{12(h_1^\vee + h_2^\vee)}, \quad (5.278)$$

$$a_{4d} = \frac{(4h_1^\vee + 4h_2^\vee + 4h_1^\vee h_2^\vee - 1) \text{rank}(G_1) \text{rank}(G_2)}{48(h_1^\vee + h_2^\vee)}. \quad (5.279)$$

For $G_1 = J$ and $G_2 = A_{k-1}$ they agree with the expressions (5.268) and (5.269) for the 4d central charges as expected.

5.2.7 $(J^{h^\vee}[k], F)$ AD matters

The Argyres Douglas (AD) matters can be engineered by taking 6d $\mathcal{N} = (2, 0)$ theory on a sphere with an irregular puncture and a regular puncture. They have non-Abelian

flavor symmetries and contain the Coulomb branch operators with fractional scaling dimensions. They are labeled by $(J^b[k], Y)$ [136] where $J^b[k]$ stands for the irregular singularity and Y the regular singularity [141, 142]. When b is the dual Coxeter number h^\vee of the Lie algebra J and Y is full, the associated VOA is conjectured to be the affine Kac-Moody algebra \widehat{F}_{k_F} with level $k_F = h^\vee(-1 + \frac{1}{h^\vee+k})$ [139].

The Schur index for the AD matters $(J^{h^\vee}[k], F)$ is given by the vacuum character of the affine Kac-Moody algebra [137, 143]. The single-particle index reads

$$\begin{aligned} i_{\text{Schur}}^{(J^{h^\vee}[k], F)}(q) &= \dim F \frac{q - q^{h^\vee+k}}{(1-q)(1-q^{h^\vee+k})} \\ &= \dim F \left(\frac{1}{1-q} - \frac{1}{1-q^{h^\vee+k}} \right). \end{aligned} \quad (5.280)$$

The supersymmetric zeta function is evaluated as

$$\mathfrak{Z}_{\text{Schur}}^{(J^{h^\vee}[k], F)}(s, z) = \dim F \left[\zeta(s, z) - (h^\vee + k)^{-s} \zeta\left(s, \frac{z}{h^\vee + k}\right) \right]. \quad (5.281)$$

It has a simple pole at $s = 1$ with residue

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{Schur}}^{(J^{h^\vee}[k], F)}(s, z) = \dim F \frac{h^\vee + k - 1}{h^\vee + k}. \quad (5.282)$$

The supersymmetric zeta value vanishes for $s = 0$ and $z = 0$

$$\mathfrak{Z}_{\text{Schur}}^{J^{h^\vee}[k], F}(0, 0) = 0. \quad (5.283)$$

On the other hand, the supersymmetric zeta value at $s = -1$ is non-trivial. We find

$$\mathfrak{Z}_{\text{Schur}}^{(J^{h^\vee}[k], F)}(-1, 0) = \frac{\dim F(h^\vee + k - 1)}{12}. \quad (5.284)$$

According to the formulae (5.144) and (5.145), we get

$$c_{4d} = \frac{\dim F(h^\vee + k - 1)}{12}, \quad (5.285)$$

$$a_{4d} = \frac{\dim F(h^\vee + k - 1)(4k + 4h^\vee - 1)}{48(h^\vee + k)}. \quad (5.286)$$

The actual values (5.285) and (5.286) are listed in the following:

$(J^{h^\vee}[k], F)$	$\dim F$	h^\vee	c_{4d}	a_{4d}	k_F
$(A_{N-1}^N[k], F)$	$N^2 - 1$	N	$\frac{(N^2-1)(N+k-1)}{12}$	$\frac{(N^2-1)(N+k-1)(4N+4k-1)}{48(N+k)}$	$\frac{N(N+k-1)}{N+k}$
$(D_N^{2N-2}[k], F)$	$N(2N-1)$	$2N-2$	$\frac{N(2N-1)(2N+k-3)}{12}$	$\frac{N(2N-1)(2N+k-3)(8N+4k-9)}{48(2N+k-2)}$	$\frac{2(N-1)(2N+k-3)}{2N+k-2}$
$(E_6^{12}[k], F)$	78	12	$\frac{13(k+11)}{2}$	$\frac{13(k+1)(4k+47)}{8(k+12)}$	$\frac{12(k+11)}{k+12}$
$(E_7^{18}[k], F)$	133	18	$\frac{133(k+17)}{12}$	$\frac{133(k+17)(4k+71)}{48(k+18)}$	$\frac{18(k+17)}{k+18}$
$(E_8^{30}[k], F)$	248	30	$\frac{62(k+29)}{3}$	$\frac{31(k+29)(4k+119)}{6(k+30)}$	$\frac{30(k+29)}{k+30}$

(5.287)

We see that they precisely agree with the central charges c_{4d} computed in [139].

The supersymmetric determinant is

$$\mathfrak{D}_{\text{Schur}}^{J^{h^\vee}[k],F}(z) = (h^\vee + k)^{\dim F(\frac{1}{2} - \frac{z}{h^\vee + k})} \frac{\Gamma(z)^{\dim F}}{\Gamma\left(\frac{z}{h^\vee + k}\right)^{\dim F}} \quad (5.288)$$

The vacuum exponent is evaluated as

$$\mathfrak{D}_{\text{Schur}}^{J^{h^\vee}[k],F}(0) = (h^\vee + k)^{-\frac{1}{2} \dim F}. \quad (5.289)$$

where we have used the fact that

$$\lim_{z \rightarrow 0} \frac{\Gamma(z)}{\Gamma(z/a)} = \frac{1}{a}. \quad (5.290)$$

5.2.8 $(A_{N-1}^N[k], [N-1, 1])$ AD matter

Next consider the $(A_{N-1}^N[k], [N-1, 1])$ Argyres Douglas (AD) matter which arises from 6d $\mathcal{N} = (2, 0)$ theory of type A_{N-1} wrapped on a sphere with an irregular puncture of type $A_{N-1}^N[k]$ and a simple regular puncture associated with the Young diagram $[N-1, 1]$.

The single-particle Schur index is given by [137]

$$i_{\text{Schur}}^{A_{N-1}^N[k][N-1,1]}(q) = \sum_{i=1}^{N-1} \frac{q^i}{1-q} + \frac{2q^{\frac{N}{2}}}{1-q} - \frac{q^{N+k}}{1-q^{N+k}} \left(\sum_{i=1}^{N-1} \sum_{j=1}^{N-1} q^{j-i} + 2 \sum_{i=1}^{N-1} q^{\frac{N-2i}{2}} \right). \quad (5.291)$$

The supersymmetric zeta function is evaluated as

$$\begin{aligned} \mathfrak{Z}_{\text{Schur}}^{A_{N-1}^N[k][N-1,1]}(s, z) &= \sum_{i=1}^{N-1} \zeta(s, i+z) + 2\zeta\left(s, \frac{N}{2} + z\right) \\ &\quad - \sum_{i=1}^{N-1} \sum_{j=1}^{N-1} (k+N)^{-s} \zeta\left(s, \frac{j-i+k+N+z}{k+N}\right) \\ &\quad - 2 \sum_{i=1}^{N-1} (k+N)^{-s} \zeta\left(s, \frac{\frac{N-2i}{2} + k+N+z}{k+N}\right). \end{aligned} \quad (5.292)$$

It has a simple pole at $s = 1$ with residue

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{Schur}}^{A_{N-1}^N[k][N-1,1]}(s, z) = \frac{(N+1)(k+1)}{N+k}. \quad (5.293)$$

When $s = 0$ and $z = 0$, the supersymmetric zeta function (5.292) becomes zero

$$\mathfrak{Z}_{\text{Schur}}^{A_{N-1}^N[k][N-1,1]}(0, 0) = 0. \quad (5.294)$$

On the other hand, the supersymmetric zeta value for $s = -1$ and $z = 0$ is

$$\mathfrak{Z}_{\text{Schur}}^{A_{N-1}^N[k][N-1,1]}(-1, 0) = \frac{(N^2(k+2) - N - k)(k+1)}{12(N+k)}. \quad (5.295)$$

From (5.144) and (5.145), we find

$$c_{4d} = \frac{(N^2(k+2) - N - k)(k+1)}{12(N+k)}, \quad (5.296)$$

$$a_{4d} = \frac{(k+1)(4k(N^2 - 1) + 8N^2 - 5N - 1)}{48(N+k)}. \quad (5.297)$$

When $k = 1$, the $(A_{N-1}^N[k], [N-1, 1])$ AD matter is identified with the (A_1, A_{2N-1}) AD theory. As expected, the expressions (5.296) and (5.297) agree with the central charges for the (A_1, A_{2N-1}) AD theory [134, 144].¹³

The supersymmetric determinant reads

$$\begin{aligned} \mathfrak{D}_{\text{Schur}}^{A_{N-1}^N[k][N-1,1]}(z) &= (2\pi)^{\frac{(N+1)(N-2)}{2}} (k+N)^{-\frac{(N^2-1)(N+k+2z)}{2(N+k)}} \\ &\times \frac{\Gamma\left(\frac{N}{2} + z\right)^2 \prod_{i=1}^{N-1} \Gamma(i+z)}{\prod_{i=1}^{N-1} \Gamma\left(\frac{\frac{N-2i}{2} + N+k+z}{N+k}\right)^2 \prod_{i,j=1}^{N-1} \Gamma\left(\frac{j-i+N+k+z}{N+k}\right)}. \end{aligned} \quad (5.300)$$

We have the vacuum exponent

$$\begin{aligned} \mathfrak{D}_{\text{Schur}}^{A_{N-1}^N[k][N-1,1]}(0) &= (2\pi)^{\frac{(N+1)(N-2)}{2}} (k+N)^{-\frac{N^2-1}{2}} \\ &\times \frac{\Gamma\left(\frac{N}{2}\right)^2 \prod_{i=1}^{N-1} \Gamma(i)}{\prod_{i=1}^{N-1} \Gamma\left(\frac{\frac{N-2i}{2} + N+k}{N+k}\right)^2 \prod_{i,j=1}^{N-1} \Gamma\left(\frac{j-i+N+k}{N+k}\right)}. \end{aligned} \quad (5.301)$$

5.2.9 rank 2 AD theory $\text{AD}(C_2)$

In [128] Kaidi and Martone constructed the rank 2 Argyres Douglas theory $\text{AD}(C_2)$ that arises as a non-trivial IR fixed point on the Coulomb branch of the mass deformed Lagrangian theory with gauge group G_2 and 4 fundamental hypermultiplets.

The single-particle Schur index is given by [128]

$$\begin{aligned} i_{\text{Schur}}^{\text{AD}(C_2)}(q) &= \frac{1}{1-q} \sum_{i=0}^{\infty} (10q^{3i+1} + 4q^{3i+\frac{3}{2}} - 4q^{3i+\frac{5}{2}} - 10q^{3i+3}) \\ &= \frac{10q + 4q^{\frac{3}{2}} + 10q^2}{1-q^3}. \end{aligned} \quad (5.302)$$

¹³The 4d central charges for the (A_{k-1}, A_{kn-1}) AD theory are [134, 144]

$$c_{4d} = \frac{(k-1)(k^2n^2 + kn^2 - 2n - 2)}{12(n+1)} \quad (5.298)$$

$$a_{4d} = \frac{(k-1)(2k^2n^2 + 2kn^2 - 5n - 5)}{24n+1}. \quad (5.299)$$

We find the supersymmetric zeta function

$$\mathfrak{Z}_{\text{Schur}}^{\text{AD}(C_2)}(s, z) = 10 \cdot 3^{-s} \zeta\left(s, \frac{1+z}{3}\right) + 4 \cdot 3^{-s} \zeta\left(s, \frac{\frac{3}{2}+z}{3}\right) + 10 \cdot 3^{-s} \zeta\left(s, \frac{2+z}{3}\right). \quad (5.303)$$

The residue at a simple pole $s = 1$ is

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{Schur}}^{\text{AD}(C_2)}(s, z) = 8. \quad (5.304)$$

The Zeta-index vanishes

$$\mathfrak{Z}_{\text{Schur}}^{\text{AD}(C_2)}(0, 0) = 0. \quad (5.305)$$

The special supersymmetric zeta value with $s = -1$ and $z = 0$ is

$$\mathfrak{Z}_{\text{Schur}}^{\text{AD}(C_2)}(-1, 0) = \frac{13}{6}. \quad (5.306)$$

Plugging (5.304) and (5.306) into the formulae (5.144) and (5.145), we get

$$c_{4d} = \frac{13}{6}, \quad (5.307)$$

$$a_{4d} = 2, \quad (5.308)$$

which agree with the central charges of the AD theory $\text{AD}(C_2)$ [128].

The supersymmetric determinant is

$$\mathfrak{D}_{\text{Schur}}^{\text{AD}(C_2)}(z) = \frac{3^{5-2z} \Gamma(z)^{10} \Gamma\left(\frac{3+2z}{6}\right)^4}{4\pi^2 \Gamma\left(\frac{z}{3}\right)^{10}}. \quad (5.309)$$

The vacuum exponent is

$$\mathfrak{D}_{\text{Schur}}^{\text{AD}(C_2)}(0) = \frac{1}{972}. \quad (5.310)$$

6 6d supersymmetric field theories

6.1 Zeta functions for supersymmetric indices

The supersymmetric index for the 6d $\mathcal{N} = (1, 0)$ supersymmetric field theories is defined as follows [145, 146, 147]:

$$\mathcal{I}^{6d(1,0)}(x; p; q; t) = \text{Tr}(-1)^F p^{j_{12}+R} q^{j_{34}+R} t^{j_{56}+R} \prod_{\alpha} x_{\alpha}^{f_{\alpha}}, \quad (6.1)$$

where j_{12} , j_{34} and j_{56} are the Cartan generators of the rotation group, R is the Cartan generator of the $SU(2)_R$ R-symmetry, and f_α are the Cartan generators of other global symmetries. The index can be computed as an integral over a product of topological string amplitudes [148].

The 8-form anomaly polynomial takes the form [149]:

$$\mathcal{A}_8 = \frac{1}{4!} [\alpha c_2(R)^2 + \beta c_2(R)p_1(T) + \gamma p_1(T)^2 + \delta p_2(T)]. \quad (6.2)$$

Here $c_2(R)$ is the second Chern class of the $SU(2)_R$ R-symmetry bundle and $p_{1,2}(T)$ are the Pontryagin classes of the tangent bundle.

We define the multiple supersymmetric zeta functions $\mathfrak{Z}^{6d(1,0)}(s, z; \omega_1, \omega_2, \omega_3)$ for the 6d $\mathcal{N} = (1, 0)$ supersymmetric field theories by taking the Mellin transform (2.7) of the plethystic logarithm of the 6d supersymmetric index (6.1) with the fugacities p , q , and t being replaced with $p = q^{\omega_1}$, $q = q^{\omega_2}$, and $t = q^{\omega_3}$. We also define by (3.1) the associated supersymmetric determinants $\mathfrak{D}^{6d(1,0)}(z; \omega_1, \omega_2, \omega_3)$. These supersymmetric zeta functions $\mathfrak{Z}^{6d(1,0)}(s, z; \omega_1, \omega_2, \omega_3)$ may have simple poles at $s = 3, 2$, and 1 .

It then follows from (2.47) that

$$\begin{aligned} & \log \mathcal{I}^{6d(1,0)}(x; q^{\omega_1}; q^{\omega_2}; q^{\omega_3}) \\ & \sim \frac{\pi^4 \text{Res}_{s=3} \mathfrak{Z}^{6d(1,0)}(s, 0; \omega_1, \omega_2, \omega_3)}{45\beta^3} + \frac{\zeta(3) \text{Res}_{s=2} \mathfrak{Z}^{6d(1,0)}(s, 0; \omega_1, \omega_2, \omega_3)}{\beta^2} \\ & + \frac{\pi^2 \text{Res}_{s=1} \mathfrak{Z}^{6d(1,0)}(s, 0; \omega_1, \omega_2, \omega_3)}{6\beta} - \mathfrak{Z}^{6d(1,0)}(0, 0; \omega_1, \omega_2, \omega_3) \log \beta \\ & + \log \mathfrak{D}^{6d(1,0)}(0; \omega_1, \omega_2, \omega_3). \end{aligned} \quad (6.3)$$

Also, when we expand the indices (6.1) as

$$\mathcal{I}^{6d(1,0)}(1; q; q; q) = \sum_n d^{6d(1,0)}(n) q^n, \quad (6.4)$$

we obtain the asymptotic degeneracy (2.71)

$$\begin{aligned} & d^{6d(1,0)}(n) \\ & \sim \frac{1}{2\sqrt{2}} \left(\frac{R_3}{15n^5} \right)^{\frac{1}{8}} \left(\frac{15n}{\pi^4 R_3} \right)^{\frac{\mathfrak{Z}^{6d(1,0)}(0,0;1,1,1)}{4}} \\ & \times \mathfrak{D}^{6d(1,0)}(0; 1, 1, 1) \exp \left[\frac{4\pi}{3} \left(\frac{R_3}{15} \right)^{\frac{1}{4}} n^{\frac{3}{4}} + \frac{R_2 \zeta(3)}{\pi^2} \left(\frac{15}{R_3} \right)^{\frac{1}{2}} n^{\frac{1}{2}} \right. \\ & \left. + \frac{\pi^6 R_1 R_3 - 45 R_2^2 \zeta(3)^2}{2\pi^5} \left(\frac{15}{R_3} \right)^{\frac{5}{4}} n^{\frac{1}{4}} - \frac{5(\pi^6 R_1 R_2 R_3 \zeta(3) - 60 R_2^3 \zeta(3)^3)}{4\pi^8} \left(\frac{15}{R_3} \right)^2 \right], \end{aligned} \quad (6.5)$$

where we have introduced the abbreviations

$$R_i := \text{Res}_{s=i} \mathfrak{Z}^{6d(1,0)}(s, 0; 1, 1, 1). \quad (6.6)$$

Di Pietro and Komargodski [45] proposed that the leading and subleading terms in the Cardy-like limit (6.3) can be universally fixed by the anomaly coefficients in (6.2). The proof was presented in [50] for the superconformal field theories with pure Higgs branches, including rank- N E-string theories [150, 151] and (G, G) -type minimal conformal matter theories [152]. Assuming that the conjecture is true, the residues of the supersymmetric zeta functions are given by the anomaly coefficients

$$\text{Res}_{s=3} \mathfrak{Z}^{6d(1,0)}(s, 0; \omega_1, \omega_2, \omega_3) = \frac{40\gamma + 10\delta}{\omega_1\omega_2\omega_3}, \quad (6.7)$$

$$\text{Res}_{s=2} \mathfrak{Z}^{6d(1,0)}(s, 0; \omega_1, \omega_2, \omega_3) = 0, \quad (6.8)$$

$$\text{Res}_{s=1} \mathfrak{Z}^{6d(1,0)}(s, 0; \omega_1, \omega_2, \omega_3) = \frac{3(\sum_{i=1}^3 \omega_i)^2 \beta - 16(\sum_{i=1}^3 \omega_i^2) \gamma + 2(\sum_{i=1}^3 \omega_i^2) \delta}{6\omega_1\omega_2\omega_3}. \quad (6.9)$$

In addition, we propose that the supersymmetric zeta values with $s = -1$ and $z = 0$ can be determined by the anomaly coefficients in (6.2)

$$\begin{aligned} & \mathfrak{Z}^{6d(1,0)}(-1, 0; \omega_1, \omega_2, \omega_3) \\ &= -\frac{(\omega_1 + \omega_2 + \omega_3)^4}{192\omega_1\omega_2\omega_3} \alpha + \frac{(\omega_1 + \omega_2 + \omega_3)^2(\omega_1^2 + \omega_2^2 + \omega_3^2)}{48\omega_1\omega_2\omega_3} \beta \\ & \quad - \frac{(\omega_1^2 + \omega_2^2 + \omega_3^2)^2}{12\omega_1\omega_2\omega_3} \gamma - \frac{\omega_1^2\omega_2^2 + \omega_1^2\omega_3^2 + \omega_2^2\omega_3^2}{12\omega_1\omega_2\omega_3} \delta. \end{aligned} \quad (6.10)$$

We explicitly check the proposed formula (6.10) for the hypermultiplet, tensor multiplet, and vector multiplet. Comparing these values with the supersymmetric Casimir energies given by the equivariant integral proposed in [66] (also see [153]), We find that they are precisely related by the formula (2.76), providing another justification for the proposed relation between the supersymmetric zeta values and the supersymmetric Casimir energies.

6.1.1 6d $\mathcal{N} = (1, 0)$ hypermultiplet

The supersymmetric index of the 6d $\mathcal{N} = (1, 0)$ free hypermultiplet takes the form [145, 146]

$$\mathcal{I}^{6d(1,0)\text{ hyp}}(x; p; q; t) = \frac{1}{\Gamma((pqt)^{\frac{1}{2}}x; p, q, t)}, \quad (6.11)$$

where

$$\Gamma(z; p, q, t) = \prod_{n=0}^{\infty} \prod_{m=0}^{\infty} \prod_{l=0}^{\infty} (1 - zp^n q^m t^l)(1 - z^{-1}p^{n+1}q^{m+1}t^{l+1}) \quad (6.12)$$

is the double elliptic gamma function [154]. We have the single-particle index

$$i^{6d(1,0)\text{ hyp}}(x; p; q; t) = \frac{(pqt)^{\frac{1}{2}}(x + x^{-1})}{(1-p)(1-q)(1-t)}. \quad (6.13)$$

Setting $p = t = q$ and $x = 1$, it becomes

$$\begin{aligned} i^{6d(1,0)\text{ hyp}}(1; q; q; q) &= \frac{2q^{\frac{3}{2}}}{(1-q)^3} \\ &= \sum_{n=1}^{\infty} (n^2 - n)q^{n-\frac{1}{2}}. \end{aligned} \quad (6.14)$$

The supersymmetric zeta function is given by

$$\mathfrak{Z}^{6d(1,0)\text{ hyp}}(s, z; \omega_1, \omega_2, \omega_3) = 2\zeta_3 \left(s, \frac{1}{2}(\omega_1 + \omega_2 + \omega_3) + z; \omega_1, \omega_2, \omega_3 \right), \quad (6.15)$$

where

$$\zeta_3(s, z; \omega_1, \omega_2, \omega_3) = \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \sum_{n_3=0}^{\infty} \frac{1}{(\omega_1 n_1 + \omega_2 n_2 + \omega_3 n_3 + z)^s} \quad (6.16)$$

is the Barnes triple zeta function. When $\omega_1 = \omega_2 = \omega_3 = 1$, it reduces to

$$\begin{aligned} &\mathfrak{Z}^{6d(1,0)\text{ hyp}}(s, z; 1, 1, 1) \\ &= \zeta \left(s - 2, \frac{1}{2} + z \right) - 2z\zeta \left(s - 1, \frac{1}{2} + z \right) + \left(z^2 - \frac{1}{4} \right) \zeta \left(s, \frac{1}{2} + z \right). \end{aligned} \quad (6.17)$$

The residues at simple poles at $s = 3, 2$ and 1 are given by

$$\text{Res}_{s=3} \mathfrak{Z}^{6d(1,0)\text{ hyp}}(s, z; \omega_1, \omega_2, \omega_3) = \frac{1}{\omega_1 \omega_2 \omega_3}, \quad (6.18)$$

$$\text{Res}_{s=2} \mathfrak{Z}^{6d(1,0)\text{ hyp}}(s, z; \omega_1, \omega_2, \omega_3) = -\frac{2z}{\omega_1 \omega_2 \omega_3}, \quad (6.19)$$

$$\text{Res}_{s=1} \mathfrak{Z}^{6d(1,0)\text{ hyp}}(s, z; \omega_1, \omega_2, \omega_3) = \frac{12z^2 - \omega_1^2 - \omega_2^2 - \omega_3^2}{12\omega_1 \omega_2 \omega_3}. \quad (6.20)$$

For $s = 0$ and $z = 0$ the supersymmetric zeta value is zero

$$\mathfrak{Z}^{6d(1,0)\text{ hyp}}(0, 0; \omega_1, \omega_2, \omega_3) = 0. \quad (6.21)$$

For $s = -1$ and $z = 0$ we obtain

$$\begin{aligned} &\mathfrak{Z}^{6d(1,0)\text{ hyp}}(-1, 0; \omega_1, \omega_2, \omega_3) \\ &= -\frac{7(\omega_1^4 + \omega_2^4 + \omega_3^4) + 10(\omega_1^2 \omega_2^2 + \omega_1^2 \omega_3^2 + \omega_2^2 \omega_3^2)}{2880\omega_1 \omega_2 \omega_3}. \end{aligned} \quad (6.22)$$

The anomaly coefficients for the free hypermultiplet are [149]

$$(\alpha, \beta, \gamma, \delta) = \left(0, 0, \frac{7}{240}, -\frac{1}{60}\right). \quad (6.23)$$

Plugging these values into the formulae (6.7), (6.9) and (6.10), we reproduce the residues (6.18), (6.20) with $z = 0$ and the supersymmetric zeta value (6.22).

The supersymmetric determinant is

$$\mathfrak{D}^{6d (1,0) \text{ hyp}}(z; \omega_1, \omega_2, \omega_3) = \Gamma_3 \left(\frac{1}{2}(\omega_1 + \omega_2 + \omega_3) + z; \omega_1, \omega_2, \omega_3 \right)^2. \quad (6.24)$$

The vacuum exponent with $\omega_1 = \omega_2 = \omega_3 = 1$ is

$$\mathfrak{D}^{6d (1,0) \text{ hyp}}(0; 1, 1, 1) = 2^{\frac{1}{8}} e^{\frac{3\zeta(3)}{16\pi^2}} = 1.1157\dots \quad (6.25)$$

The degeneracy $d^{6d (1,0) \text{ hyp}}(n)$ obeys the exponential growth. From the formula (6.5) we obtain the asymptotic growth

$$d^{6d (1,0) \text{ hyp}}(n) \sim \frac{1}{2^{\frac{11}{8}} \cdot 15^{\frac{1}{8}} n^{\frac{5}{8}}} \exp \left[\frac{4\pi}{3 \cdot 15^{\frac{1}{4}}} n^{\frac{3}{4}} - \frac{5^{\frac{1}{4}} \pi}{8 \cdot 3^{\frac{3}{4}}} n^{\frac{1}{4}} + \frac{3\zeta(3)}{16\pi^2} \right]. \quad (6.26)$$

The exact numbers $d_{\text{exact}}^{6d (1,0) \text{ hyp}}(n)$ and the values $d_{\text{asympt}}^{6d (1,0) \text{ hyp}}(n)$ evaluated from (6.26) are shown as follows:

n	$d_{\text{exact}}^{6d (1,0) \text{ hyp}}(n)$	$d_{\text{asympt}}^{6d (1,0) \text{ hyp}}(n)$
10	2964	655.87
100	5.8501×10^{26}	1.19298×10^{27}
1000	1.05043×10^{161}	2.11485×10^{161}

(6.27)

6.1.2 6d $\mathcal{N} = (1, 0)$ tensor multiplet

The single-particle index for the 6d $\mathcal{N} = (1, 0)$ tensor multiplet is given by [145]

$$i^{6d (1,0) \text{ tensor}}(p; q; t) = \frac{pqt - pq - pt - qt}{(1-p)(1-q)(1-t)}. \quad (6.28)$$

When $p = t = q$, it reduces to

$$\begin{aligned} i^{6d (1,0) \text{ tensor}}(q; q; q) &= \frac{-3q^2 + q^3}{(1-q)^3} \\ &= -\sum_{n=1}^{\infty} (n^2 - 1)q^n. \end{aligned} \quad (6.29)$$

The supersymmetric zeta function is

$$\begin{aligned} & \mathfrak{Z}^{6d (1,0) \text{ tensor}}(s, z; \omega_1, \omega_2, \omega_3) \\ &= \zeta_3(s, \omega_1 + \omega_2 + \omega_3 + z; \omega_1, \omega_2, \omega_3) - \zeta_3(s, \omega_1 + \omega_2 + z; \omega_1, \omega_2, \omega_3) \\ & - \zeta_3(s, \omega_1 + \omega_3 + z; \omega_1, \omega_2, \omega_3) - \zeta_3(s, \omega_2 + \omega_3 + z; \omega_1, \omega_2, \omega_3). \end{aligned} \quad (6.30)$$

When $\omega_1 = \omega_2 = \omega_3 = 1$, it becomes

$$\begin{aligned} & \mathfrak{Z}^{6d (1,0) \text{ tensor}}(s, z) \\ &= -\zeta(s-2, 1+z) + 2z\zeta(s-1, 1+z) + (1-z^2)\zeta(s, 1+z). \end{aligned} \quad (6.31)$$

The supersymmetric zeta function (6.30) has simple poles at $s = 3, 2$ and 1 with residues

$$\text{Res}_{s=3} \mathfrak{Z}^{6d (1,0) \text{ tensor}}(s, z; \omega_1, \omega_2, \omega_3) = -\frac{1}{\omega_1 \omega_2 \omega_3}, \quad (6.32)$$

$$\text{Res}_{s=2} \mathfrak{Z}^{6d (1,0) \text{ tensor}}(s, z; \omega_1, \omega_2, \omega_3) = \frac{2z}{\omega_1 \omega_2 \omega_3}, \quad (6.33)$$

$$\text{Res}_{s=1} \mathfrak{Z}^{6d (1,0) \text{ tensor}}(s, z; \omega_1, \omega_2, \omega_3) = -\frac{6z^2 + \omega_1^2 + \omega_2^2 + \omega_3^2 - 3(\omega_1 \omega_2 + \omega_1 \omega_3 + \omega_2 \omega_3)}{6\omega_1 \omega_2 \omega_3}. \quad (6.34)$$

The Zeta-index is given by

$$\mathfrak{Z}^{6d (1,0) \text{ tensor}}(0, 0; \omega_1, \omega_2, \omega_3) = -\frac{1}{2}. \quad (6.35)$$

For $s = -1$ and $z = 0$ we have

$$\begin{aligned} & \mathfrak{Z}^{6d (1,0) \text{ tensor}}(-1, 0; \omega_1, \omega_2, \omega_3) \\ &= -\frac{1}{360\omega_1 \omega_2 \omega_3} \left[\omega_1^4 + \omega_2^4 + \omega_3^4 - 5(\omega_1^2 \omega_2^2 + \omega_1^2 \omega_3^2 + \omega_2^2 \omega_3^2) + 15\omega_1 \omega_2 \omega_3 (\omega_1 + \omega_2 + \omega_3) \right]. \end{aligned} \quad (6.36)$$

The anomaly coefficients for the 6d $\mathcal{N} = (1, 0)$ tensor multiplet is given by [149]

$$(\alpha, \beta, \gamma, \delta) = \left(1, \frac{1}{2}, \frac{23}{240}, -\frac{29}{60} \right). \quad (6.37)$$

Substituting these values into the formulae (6.7), (6.9) and (6.10), we find the residues (6.32), (6.34) for $z = 0$ and the supersymmetric zeta value (6.36).

The supersymmetric determinant is

$$\begin{aligned} & \mathfrak{D}^{6d (1,0) \text{ tensor}}(z; \omega_1, \omega_2, \omega_3) \\ &= \frac{\Gamma_3(\omega_1 + \omega_2 + \omega_3 + z; \omega_1, \omega_2, \omega_3)}{\Gamma_3(\omega_1 + \omega_2 + z; \omega_1, \omega_2, \omega_3) \Gamma_3(\omega_1 + \omega_3 + z; \omega_1, \omega_2, \omega_3) \Gamma_3(\omega_2 + \omega_3 + z; \omega_1, \omega_2, \omega_3)}. \end{aligned} \quad (6.38)$$

The vacuum exponent for $\omega_1 = \omega_2 = \omega_3 = 1$ is

$$\mathfrak{D}^{6d (1,0) \text{ tensor}}(0; 1, 1, 1) = \frac{e^{\frac{\zeta(3)}{4\pi^2}}}{\sqrt{2\pi}} = 0.411276 \dots \quad (6.39)$$

6.1.3 6d $\mathcal{N} = (1, 0)$ vector multiplet

The 6d $\mathcal{N} = (1, 0)$ free vector multiplet is not a conformal field theory while the free vector field is scale invariant [155].

The single-particle index for the 6d $\mathcal{N} = (1, 0)$ vector multiplet reads [146, 154]

$$i^{6d (1,0) \text{ vec}}(p; q; t) = \frac{pq + pt + qt - p - q - t - 2pqt}{(1-p)(1-q)(1-t)} \quad (6.40)$$

For $p = q = t$ it reduces to

$$\begin{aligned} i^{6d (1,0) \text{ vec}}(q) &= \frac{3q^2 - 3q - 2q^3}{(1-q)^3} \\ &= -\frac{3}{2} \sum_{n=1}^{\infty} (n^2 - n) q^{n-1} + \frac{3}{2} \sum_{n=1}^{\infty} (n^2 - n) q^n - \sum_{n=1}^{\infty} (n^2 - n) q^{n+1}. \end{aligned} \quad (6.41)$$

The supersymmetric zeta function is

$$\begin{aligned} \mathfrak{Z}^{6d (1,0) \text{ vec}}(s, z; \omega_1, \omega_2, \omega_3) \\ &= -\sum_{i=1}^3 \zeta_3(s, \omega_i + z; \omega_1, \omega_2, \omega_3) + \sum_{i < j}^3 \zeta_3(s, \omega_i + \omega_j + z; \omega_1, \omega_2, \omega_3) \\ &\quad - 2\zeta_3(s, \omega_1 + \omega_2 + \omega_3 + z; \omega_1, \omega_2, \omega_3). \end{aligned} \quad (6.42)$$

For $\omega_1 = \omega_2 = \omega_3 = 1$, it becomes

$$\begin{aligned} \mathfrak{Z}^{6d (1,0) \text{ vec}}(s, z) \\ &= -\frac{3}{2} [\zeta(s-2, z) - (2z-1)\zeta(s-1, z) + (z^2-z)\zeta(s, z)] \\ &\quad + \frac{3}{2} [\zeta(s-2, z+1) - (2z+1)\zeta(s-1, z+1) + (z^2+z)\zeta(s, z+1)] \\ &\quad - [\zeta(s-2, z+2) - (2z+3)\zeta(s-1, z+2) + (z^2+3z+2)\zeta(s, z+2)]. \end{aligned} \quad (6.43)$$

It has simple poles at $s = 3, 2$ and 1 with residues

$$\text{Res}_{s=3} \mathfrak{Z}^{6d (1,0) \text{ vec}}(s, z; \omega_1, \omega_2, \omega_3) = -\frac{1}{\omega_1 \omega_2 \omega_3}, \quad (6.44)$$

$$\text{Res}_{s=2} \mathfrak{Z}^{6d (1,0) \text{ vec}}(s, z; \omega_1, \omega_2, \omega_3) = \frac{2z}{\omega_1 \omega_2 \omega_3}, \quad (6.45)$$

$$\text{Res}_{s=1} \mathfrak{Z}^{6d (1,0) \text{ vec}}(s, z; \omega_1, \omega_2, \omega_3) = -\frac{6z^2 + \omega_1^2 + \omega_2^2 + \omega_3^2 + 3(\omega_1 \omega_2 + \omega_1 \omega_3 + \omega_2 \omega_3)}{6\omega_1 \omega_2 \omega_3}. \quad (6.46)$$

The Zeta-index is

$$\mathfrak{Z}^{6d (1,0) \text{ vec}}(0, 0; \omega_1, \omega_2, \omega_3) = 1. \quad (6.47)$$

For $s = -1$ and $z = 0$ the supersymmetric zeta value is

$$\begin{aligned} & \mathfrak{Z}^{6d (1,0) \text{ vec}}(-1, 0; \omega_1, \omega_2, \omega_3) \\ &= -\frac{\omega_1^4 + \omega_2^4 + \omega_3^4 - 5(\omega_1^2\omega_2^2 + \omega_1^2\omega_3^2 + \omega_2^2\omega_3^2) - 15\omega_1\omega_2\omega_3(\omega_1 + \omega_2 + \omega_3)}{360\omega_1\omega_2\omega_3}. \end{aligned} \quad (6.48)$$

The anomaly coefficients for the 6d $\mathcal{N} = (1, 0)$ vector multiplet are [149]

$$(\alpha, \beta, \gamma, \delta) = \left(-1, -\frac{1}{2}, -\frac{7}{240}, \frac{1}{60}\right). \quad (6.49)$$

We recover the residues (6.44), (6.46) for $z = 0$ and the supersymmetric zeta value (6.48) by inserting these values into the formulae (6.7), (6.9) and (6.10).

The supersymmetric determinant is

$$\begin{aligned} & \mathfrak{D}^{6d (1,0) \text{ vec}}(z; \omega_1, \omega_2, \omega_3) \\ &= \frac{\prod_{i < j}^3 \Gamma_3(\omega_i + \omega_j + z; \omega_1, \omega_2, \omega_3)}{\Gamma_3(\omega_1 + \omega_2 + \omega_3 + z; \omega_1, \omega_2, \omega_3)^2 \prod_{i=1}^3 \Gamma_3(\omega_i + z; \omega_1, \omega_2, \omega_3)}. \end{aligned} \quad (6.50)$$

When $\omega_1 = \omega_2 = \omega_3 = 1$, the vacuum exponent is

$$\mathfrak{D}^{6d (1,0) \text{ vec}}(0; 1, 1, 1) = 2\pi e^{\frac{\zeta(3)}{4\pi^2}} = 6.47744 \dots \quad (6.51)$$

6.1.4 6d $\mathcal{N} = (2, 0)$ tensor multiplet

The 6d $\mathcal{N} = (2, 0)$ free tensor multiplet consists of the 6d $\mathcal{N} = (1, 0)$ free tensor multiplet and hypermultiplet.

The single-particle index is given by [145, 147]

$$\begin{aligned} i^{6d (2,0) \text{ tensor}}(x; p, q, t) &= i^{6d (1,0) \text{ tensor}}(p; q; t) + i^{6d (1,0) \text{ hyp}}(x; p; q; t) \\ &= \frac{(pqt)^{\frac{1}{2}}(x + x^{-1}) - pq - pt - qt + pqt}{(1-p)(1-q)(1-t)}. \end{aligned} \quad (6.52)$$

With $p = t = q$ and $x = 1$, it becomes

$$\begin{aligned} i^{6d (2,0) \text{ tensor}}(1; q; q; q) &= i^{6d (1,0) \text{ tensor}}(q; q; q) + i^{6d (1,0) \text{ hyp}}(1; q; q; q) \\ &= \frac{2q^{\frac{3}{2}} - 3q^2 + q^3}{(1-q)^3}. \end{aligned} \quad (6.53)$$

The supersymmetric zeta function is

$$\begin{aligned}
& \mathfrak{Z}^{6d(2,0)\text{ tensor}}(s, z; \omega_1, \omega_2, \omega_3) \\
&= \zeta_3(s, \omega_1 + \omega_2 + \omega_3 + z; \omega_1, \omega_2, \omega_3) - \zeta_3(s, \omega_1 + \omega_2 + z; \omega_1, \omega_2, \omega_3) \\
&- \zeta_3(s, \omega_1 + \omega_3 + z; \omega_1, \omega_2, \omega_3) - \zeta_3(s, \omega_2 + \omega_3 + z; \omega_1, \omega_2, \omega_3) \\
&+ 2\zeta_3\left(s, \frac{1}{2}(\omega_1 + \omega_2 + \omega_3) + z; \omega_1, \omega_2, \omega_3\right). \tag{6.54}
\end{aligned}$$

When $\omega_1 = \omega_2 = \omega_3 = 1$, it becomes

$$\begin{aligned}
& \mathfrak{Z}^{6d(2,0)\text{ tensor}}(s, z; 1, 1, 1) \\
&= -\zeta(s-2, 1+z) + 2z\zeta(s-1, 1+z) + (1-z^2)\zeta(s, 1+z) \\
&+ \zeta\left(s-2, \frac{1}{2}+z\right) - 2z\zeta\left(s-1, \frac{1}{2}+z\right) + \left(z^2 - \frac{1}{4}\right)\zeta\left(s, \frac{1}{2}+z\right). \tag{6.55}
\end{aligned}$$

The supersymmetric zeta function (6.54) has simple poles at $s = 3, 2$ and 1 with residues

$$\text{Res}_{s=3} \mathfrak{Z}^{6d(2,0)\text{ tensor}}(s, z; \omega_1, \omega_2, \omega_3) = 0, \tag{6.56}$$

$$\text{Res}_{s=2} \mathfrak{Z}^{6d(2,0)\text{ tensor}}(s, z; \omega_1, \omega_2, \omega_3) = 0, \tag{6.57}$$

$$\text{Res}_{s=1} \mathfrak{Z}^{6d(2,0)\text{ tensor}}(s, z; \omega_1, \omega_2, \omega_3) = -\frac{\omega_1^2 + \omega_2^2 + \omega_3^2 - 2(\omega_1\omega_2 + \omega_1\omega_3 + \omega_2\omega_3)}{4\omega_1\omega_2\omega_3}. \tag{6.58}$$

We have the Zeta-index

$$\mathfrak{Z}^{6d(2,0)\text{ tensor}}(0, 0; \omega_1, \omega_2, \omega_3) = -\frac{1}{2}. \tag{6.59}$$

The supersymmetric zeta value for $s = -1$ and $z = 0$ is

$$\begin{aligned}
& \mathfrak{Z}^{6d(2,0)\text{ tensor}}(-1, 0; \omega_1, \omega_2, \omega_3) \\
&= -\frac{\omega_1^4 + \omega_2^4 + \omega_3^4 - 2(\omega_1^2\omega_2^2 + \omega_1^2\omega_3^2 + \omega_2^2\omega_3^2) + 8\omega_1\omega_2\omega_3(\omega_1 + \omega_2 + \omega_3)}{192\omega_1\omega_2\omega_3}. \tag{6.60}
\end{aligned}$$

Substituting the supersymmetric zeta value (6.60) into (2.76), we reproduce the supersymmetric Casimir energy in [147, 66]. The anomaly coefficients of the 6d $\mathcal{N} = (2, 0)$ free tensor multiplet are [156]

$$(\alpha, \beta, \gamma, \delta) = \left(1, \frac{1}{2}, \frac{1}{8}, -\frac{1}{2}\right). \tag{6.61}$$

Again the expressions (6.58), (6.56) and (6.60) can be found by plugging the anomaly coefficients into the formulae (6.7), (6.9) and (6.10).

The supersymmetric determinant is

$$\begin{aligned}
& \mathfrak{D}^{\text{6d } (2,0) \text{ tensor}}(z; \omega_1, \omega_2, \omega_3) \\
&= \mathfrak{D}^{\text{6d } (1,0) \text{ tensor}}(z; \omega_1, \omega_2, \omega_3) \mathfrak{D}^{\text{6d } (1,0) \text{ hyp}}(z; \omega_1, \omega_2, \omega_3) \\
&= \frac{\Gamma_3(\omega_1 + \omega_2 + \omega_3; \omega_1, \omega_2, \omega_3) \Gamma_3\left(\frac{1}{2}(\omega_1 + \omega_2 + \omega_3); \omega_1, \omega_2, \omega_3\right)^2}{\Gamma_3(\omega_1 + \omega_2; \omega_1, \omega_2, \omega_3) \Gamma_3(\omega_1 + \omega_3; \omega_1, \omega_2, \omega_3) \Gamma_3(\omega_2 + \omega_3; \omega_1, \omega_2, \omega_3)}. \tag{6.62}
\end{aligned}$$

The vacuum exponent with $\omega_1 = \omega_2 = \omega_3 = 1$ is

$$\mathfrak{D}^{\text{6d } (2,0) \text{ tensor}}(0; 1, 1, 1) = \frac{e^{\frac{7\zeta(3)}{16\pi^2}}}{2^{\frac{3}{8}}\sqrt{\pi}} = 0.45886\dots \tag{6.63}$$

6.2 Zeta functions for unrefined indices

6.2.1 6d $\mathcal{N} = (2, 0)$ theories

It is conjectured [157] that 6d $\mathcal{N} = (2, 0)$ theories of type $\mathfrak{g} = \{A_n, D_n, E_n\}$ admit the protected chiral algebra that is isomorphic to the W-algebra $\mathcal{W}(\mathfrak{g})$ of central charge

$$c = \text{rank}(\mathfrak{g}) + 4 \dim \mathfrak{g} \cdot h^\vee. \tag{6.64}$$

The relevant index is called the unrefined index [147]. In our conventions it is obtained from the supersymmetric index of the 6d $\mathcal{N} = (2, 0)$ theory of type \mathfrak{g} by setting $x = q^{\frac{1}{2}}$ and $p = t = q$. The unrefined index is identified with the vacuum character of the $\mathcal{W}(\mathfrak{g})$. The associated single-particle index takes the form [147, 157]

$$i_{\text{unref}}^{\text{6d } (2,0) \mathfrak{g}}(q) = \frac{\sum_{i=1}^{\text{rank}(\mathfrak{g})} q^{d_i}}{1 - q}, \tag{6.65}$$

where d_i are the degrees of the Casimir of \mathfrak{g} . This expression can be explicitly checked for type $\mathfrak{g} = \{A_n, D_n\}$ by analyzing the supersymmetric partition functions [147], while it is conjectural for type E_n .

The supersymmetric zeta function is given by

$$\mathfrak{Z}_{\text{unref}}^{\text{6d } (2,0) \mathfrak{g}}(s, z) = \sum_{i=1}^{\text{rank}(\mathfrak{g})} \zeta(s, d_i + z). \tag{6.66}$$

It has a simple pole at $s = 1$ with residue

$$\text{Res}_{s=1} \mathfrak{Z}_{\text{unref}}^{\text{6d } (2,0) \mathfrak{g}}(s, z) = \text{rank}(\mathfrak{g}). \tag{6.67}$$

The supersymmetric zeta value for $s = 0$ and $z = 0$ is given by

$$\mathfrak{Z}_{\text{unref}}^{\text{6d } (2,0) \mathfrak{g}}(0, 0) = -\frac{\dim \mathfrak{g}}{2}. \tag{6.68}$$

With $s = -1$ and $z = 0$ the supersymmetric zeta value is

$$\mathfrak{Z}_{\text{unref}}^{6d(2,0)\mathfrak{g}}(-1, 0) = -\frac{1}{12} [\text{rank}(\mathfrak{g}) + 2 \dim \mathfrak{g} \cdot h^\vee]. \quad (6.69)$$

The supersymmetric determinant is

$$\mathfrak{D}_{\text{unref}}^{6d(2,0)\mathfrak{g}}(z) = \frac{1}{(2\pi)^{\frac{\text{rank}(\mathfrak{g})}{2}}} \prod_{i=1}^{\text{rank}(\mathfrak{g})} \Gamma(d_i + z). \quad (6.70)$$

The vacuum exponent is

$$\mathfrak{D}_{\text{unref}}^{6d(2,0)\mathfrak{g}}(0) = \frac{1}{(2\pi)^{\frac{\text{rank}(\mathfrak{g})}{2}}} \prod_{i=1}^{\text{rank}(\mathfrak{g})} \Gamma(d_i). \quad (6.71)$$

where the factor $\prod_i \Gamma(d_i)$ is the classical Lie superfactorial associated with \mathfrak{g} .

As the degeneracy $d_{\text{unref}}^{6d(2,0)\mathfrak{g}}(n)$ grows exponentially, we find from (2.64) the asymptotic growth

$$\begin{aligned} d_{\text{unref}}^{6d(2,0)\mathfrak{g}}(n) &\sim \frac{1}{2} \left(\frac{\text{rank}(\mathfrak{g})}{6n^3} \right)^{\frac{1}{4}} \left(\frac{\pi^2 \text{rank}(\mathfrak{g})}{6n} \right)^{\frac{\dim \mathfrak{g}}{4}} \frac{1}{(2\pi)^{\frac{\text{rank}(\mathfrak{g})}{2}}} \prod_{i=1}^{\text{rank}(\mathfrak{g})} \Gamma(d_i) \\ &\times \exp \left[2\pi \sqrt{\frac{\text{rank}(\mathfrak{g})}{6}} n^{\frac{1}{2}} \right]. \end{aligned} \quad (6.72)$$

For example, for $\mathfrak{g} = A_4$ the exact values $d_{\text{exact}}^{6d(2,0)A_4}(n)$ and the values $d_{\text{asympt}}^{6d(2,0)A_4}(n)$ estimated from (6.72) are shown as follows:

n	$d_{\text{exact}}^{6d(2,0)A_4}(n)$	$d_{\text{asympt}}^{6d(2,0)A_4}(n)$
10	72	528251
100	4.35542×10^{12}	1.61214×10^{14}
1000	1.20684×10^{55}	4.29856×10^{55}

(6.73)

Acknowledgements

YN is in part supported by JSPS KAKENHI Grant Number 21K03581. TO was supported by the Startup Funding no. 4007012317 of Southeast University.

A Notations

We define

$$\begin{aligned} (a; q)_0 &:= 1, & (a; q)_n &:= \prod_{k=0}^{n-1} (1 - aq^k), & (q)_n &:= \prod_{k=1}^n (1 - q^k), \\ (a; q)_\infty &:= \prod_{k=0}^{\infty} (1 - aq^k), & (q)_\infty &:= \prod_{k=1}^{\infty} (1 - q^k), \end{aligned} \quad (A.1)$$

with $a, q \in \mathbb{C}$ and $|q| < 1$. We also introduce the following notations:

$$(x^\pm; q)_n := (x; q)_n (x^{-1}; q)_n, \quad (\text{A.2})$$

$$(x_1, \dots, x_k; q)_n := (x_1; q)_n \cdots (x_k; q)_n. \quad (\text{A.3})$$

B Residue formulae

The Barnes multiple zeta function $\zeta_N(s, z; \{\omega_i\})$ has simple poles at $s = 1, 2, \dots, N$. The residues at these poles are given by (2.18). Explicitly, the residues at the poles $s = N, N-1, N-2$ and $N-3$ are

$$\text{Res}_{s=N} \zeta_N(s, z; \{\omega_i\}) = \frac{1}{\Gamma(N)} \prod_{i=1}^N \frac{1}{\omega_i}, \quad (\text{B.1})$$

$$\text{Res}_{s=N-1} \zeta_N(s, z; \{\omega_i\}) = \frac{1}{\Gamma(N-1)} \frac{\sum_{i=1}^N \omega_i - 2z}{2 \prod_{i=1}^N \omega_i}, \quad (\text{B.2})$$

$$\text{Res}_{s=N-2} \zeta_N(s, z; \{\omega_i\}) = \frac{1}{\Gamma(N-2)} \frac{\sum_{i=1}^N \omega_i^2 + 3 \sum_{i < j} \omega_i \omega_j - 6 \sum_{i=1}^N \omega_i z + 6z^2}{12 \prod_{i=1}^N \omega_i}, \quad (\text{B.3})$$

$$\begin{aligned} \text{Res}_{s=N-3} \zeta_N(s, z; \{\omega_i\}) &= \frac{1}{\Gamma(N-3)} \frac{1}{24 \prod_{i=1}^N \omega_i} \\ &\times \left[\left(\sum_{i=1}^N \omega_i \right) \left(\sum_{i < j} \omega_i \omega_j \right) - 2 \sum_{i=1}^N \omega_i^2 z - 6 \sum_{i < j} \omega_i \omega_j z + 6 \sum_{i=1}^N \omega_i z^2 - 4z^3 \right]. \end{aligned} \quad (\text{B.4})$$

When $N = 1$ and $\omega_1 = 1$, the Barnes multiple zeta function becomes the Hurwitz zeta function

$$\zeta(s, z) = \sum_{n=0}^{\infty} \frac{1}{(n+z)^s}. \quad (\text{B.5})$$

Furthermore, it reduces to the Riemann zeta function when $z = 1$. The Hurwitz zeta function and the Riemann zeta function have a simple pole at $s = 1$ with residue 1.

C Special zeta values

C.1 Riemann zeta function

The zeta values for any positive even integer $2k$ are given by

$$\zeta(2k) = -\frac{B_{2k}}{2(2k)!} (2\pi i)^{2k}, \quad (\text{C.1})$$

where B_{2k} are the Bernoulli numbers. For negative integers $-k < 0$ we have

$$\zeta(-k) = -\frac{B_{k+1}}{k+1}. \quad (\text{C.2})$$

The special values of the s -derivative of the Riemann zeta function $\zeta(s)$ at $s = -1$ and $s = -2$ are given by

$$\zeta'(-1) = \frac{1}{12} - \log A = -0.165421 \dots, \quad (\text{C.3})$$

$$\zeta'(-2) = -\frac{\zeta(3)}{4\pi^2} = -0.0304485 \dots, \quad (\text{C.4})$$

where A is the Glaisher-Kinkelin constant.

C.2 Hurwitz zeta function

The special values of the Hurwitz zeta function are

$$\zeta(1-m, a) = -\frac{B_m(a)}{m}, \quad (\text{C.5})$$

where $B_j(a)$ are the Bernoulli polynomials.

For example, at $s = 0, -1$ and -2 they are

$$\zeta(0, a) = \frac{1}{2} - a, \quad (\text{C.6})$$

$$\zeta(-1, a) = -\frac{1}{2} \left(a^2 - a + \frac{1}{6} \right), \quad (\text{C.7})$$

$$\zeta(-2, a) = -\frac{1}{3} \left(a^3 - \frac{3}{2}a^2 + \frac{1}{2}a \right). \quad (\text{C.8})$$

The special value of the s -derivative of the Hurwitz zeta function at $s = 0$ is

$$\zeta'(0, a) = \log \Gamma(a) - \frac{1}{2} \log(2\pi). \quad (\text{C.9})$$

The explicit formulae for the other special values of the s -derivative of the Hurwitz zeta function are obtainable from the functional equation for the Hurwitz zeta function




Figure 5: The special value $\zeta'(-1, a)$.

(see e.g. [158, 159]). For example, we have

$$\zeta'(-1, \frac{1}{6}) = \frac{\log(12)}{144} - \frac{\pi}{12\sqrt{3}} + \frac{\psi^{(1)}(\frac{1}{3})}{8\sqrt{3}\pi} + \frac{1}{6} \left(\frac{1}{12} - \log A \right) = 0.0704526\dots, \quad (\text{C.10})$$

$$\zeta'(-1, \frac{1}{4}) = \frac{G}{4\pi} - \frac{1}{8} \left(\frac{1}{12} - \log A \right) = 0.0935679\dots, \quad (\text{C.11})$$

$$\zeta'(-1, \frac{1}{3}) = -\frac{\log(3)}{72} - \frac{\pi}{18\sqrt{3}} + \frac{\psi^{(1)}(\frac{1}{3})}{12\sqrt{3}\pi} - \frac{1}{3} \left(\frac{1}{12} - \log A \right) = 0.0937262\dots, \quad (\text{C.12})$$

$$\zeta'(-1, \frac{1}{2}) = -\frac{\log(2)}{24} - \frac{1}{2} \left(\frac{1}{12} - \log A \right) = 0.0538294\dots, \quad (\text{C.13})$$

$$\zeta'(-1, \frac{2}{3}) = -\frac{\log(3)}{72} + \frac{\pi}{18\sqrt{3}} - \frac{\psi^{(1)}(\frac{1}{3})}{12\sqrt{3}\pi} - \frac{1}{3} \left(\frac{1}{12} - \log A \right) = -0.0139624\dots, \quad (\text{C.14})$$

where $G = 0.915965\dots$ is Catalan's constant. We present numerical plots of the special values $\zeta'(-1, a)$ in Figure 5. The values at $a = 0$ and $a = 2$ are equal to (C.3)

$$\zeta'(-1, 0) = \zeta'(-1, 2) = \frac{1}{12} - \log(A) = -0.165421\dots \quad (\text{C.15})$$

C.3 Barnes multiple zeta function

For the Barnes multiple zeta function $\zeta_N(s, z; \{\omega_i\})$ with $z > 0$ let

$$z = z_1\omega_1 + \dots + z_N\omega_N \quad (\text{C.16})$$

with $z_i \in \mathbb{R}$. Then the special values of the Barnes multiple zeta function are given by

$$\begin{aligned} & \zeta_N(1-m, z; \{\omega_i\}) \\ &= (-1)^N (m-1)! \sum_{\substack{m_1+\dots+m_N=m+N-1 \\ m_1 \geq 0, \dots, m_N \geq 0}} \left(\prod_{i=1}^N \frac{B_{m_i}(z_i)}{m_i!} \right) \omega_1^{m_1-1} \dots \omega_N^{m_N-1} \end{aligned} \quad (\text{C.17})$$

for $m \in \mathbb{N}$. Here $B_j(a)$ are the Bernoulli polynomials. When $N = 1$ and $\omega_1 = 1$, it reduces to (C.5).

In particular, the special values of the Barnes double zeta function for non-positive integers can be obtained using the result in [160] (also see [161]). We have

$$\zeta_2(-m, z; a, b) = a^m \left[-\frac{B_{m+1}(\frac{z}{a})}{m+1} + (-1)^m m! \sum_{j=0}^{m+2} \frac{B_j B_{m+2-j}(1-\frac{z}{a})}{j!(m+2-j)!} \left(\frac{b}{a}\right)^{j-1} \right] \quad (\text{C.18})$$

for $m \geq 0$, where B_j are the Bernoulli numbers.

References

- [1] M. F. Atiyah and I. M. Singer, “The Index of elliptic operators. 1,” *Annals Math.* **87** (1968) 484–530.
- [2] L. Alvarez-Gaume, “Supersymmetry and the Atiyah-Singer Index Theorem,” *Commun. Math. Phys.* **90** (1983) 161.
- [3] E. Witten, “Quantum Field Theory and the Jones Polynomial,” *Commun. Math. Phys.* **121** (1989) 351–399.
- [4] J. M. Maldacena, “The Large N limit of superconformal field theories and supergravity,” *Adv. Theor. Math. Phys.* **2** (1998) 231–252, [arXiv:hep-th/9711200](https://arxiv.org/abs/hep-th/9711200).
- [5] M. Kac, “Can one hear the shape of a drum?” *Amer. Math. Monthly* **73** no. 4, part II, (1966) 1–23. <https://doi.org/10.2307/2313748>.
- [6] D. Martelli, J. Sparks, and S.-T. Yau, “The Geometric dual of a-maximisation for Toric Sasaki-Einstein manifolds,” *Commun. Math. Phys.* **268** (2006) 39–65, [arXiv:hep-th/0503183](https://arxiv.org/abs/hep-th/0503183).
- [7] D. Martelli, J. Sparks, and S.-T. Yau, “Sasaki-Einstein manifolds and volume minimisation,” *Commun. Math. Phys.* **280** (2008) 611–673, [arXiv:hep-th/0603021](https://arxiv.org/abs/hep-th/0603021).

- [8] D. Martelli and J. Sparks, “Dual Giant Gravitons in Sasaki-Einstein Backgrounds,” *Nucl. Phys. B* **759** (2006) 292–319, [arXiv:hep-th/0608060](#).
- [9] E. Witten, “Constraints on Supersymmetry Breaking,” *Nucl. Phys. B* **202** (1982) 253.
- [10] J. Kinney, J. M. Maldacena, S. Minwalla, and S. Raju, “An Index for 4 dimensional super conformal theories,” *Commun. Math. Phys.* **275** (2007) 209–254, [arXiv:hep-th/0510251](#).
- [11] C. Romelsberger, “Counting chiral primaries in $N = 1$, $d=4$ superconformal field theories,” *Nucl. Phys. B* **747** (2006) 329–353, [arXiv:hep-th/0510060](#).
- [12] T. Kawai, Y. Yamada, and S.-K. Yang, “Elliptic genera and $N=2$ superconformal field theory,” *Nucl. Phys. B* **414** (1994) 191–212, [arXiv:hep-th/9306096](#).
- [13] V. P. Spiridonov and G. S. Vartanov, “Elliptic hypergeometric integrals and ’t Hooft anomaly matching conditions,” *JHEP* **06** (2012) 016, [arXiv:1203.5677 \[hep-th\]](#).
- [14] S. S. Razamat, “On a modular property of $N=2$ superconformal theories in four dimensions,” *JHEP* **10** (2012) 191, [arXiv:1208.5056 \[hep-th\]](#).
- [15] A. Berkovich, B. M. McCoy, and W. P. Orrick, “Polynomial identities, indices, and duality for the $N=1$ superconformal model $SM(2,4n)$,” *J. Statist. Phys.* **83** (1996) 795, [arXiv:hep-th/9507072](#).
- [16] G. Basar, A. Cherman, D. A. McGady, and M. Yamazaki, “Temperature-reflection symmetry,” *Phys. Rev. D* **91** no. 10, (2015) 106004, [arXiv:1406.6329 \[hep-th\]](#).
- [17] F. Lindemann, “Ueber die Zahl π ,” *Math. Ann.* **20** no. 2, (1882) 213–225. <https://doi.org/10.1007/BF01446522>.
- [18] R. Apéry, “Irrationalité de ζ_2 et ζ_3 ,” No. 61, pp. 11–13. 1979. Luminy Conference on Arithmetic.
- [19] T. Rivoal, “La fonction zêta de Riemann prend une infinité de valeurs irrationnelles aux entiers impairs,” *C. R. Acad. Sci. Paris Sér. I Math.* **331** no. 4, (2000) 267–270. [https://doi.org/10.1016/S0764-4442\(00\)01624-4](https://doi.org/10.1016/S0764-4442(00)01624-4).

- [20] V. V. Zudilin, “One of the numbers $\zeta(5)$, $\zeta(7)$, $\zeta(9)$, $\zeta(11)$ is irrational,” *Uspekhi Mat. Nauk* **56** no. 4(340), (2001) 149–150.
<https://doi.org/10.1070/RM2001v056n04ABEH000427>.
- [21] K. A. Intriligator and B. Wecht, “The Exact superconformal R symmetry maximizes a,” *Nucl. Phys. B* **667** (2003) 183–200, [arXiv:hep-th/0304128](#).
- [22] L. Rastelli and B. C. Rayhaun, “Rationality in four dimensions,” *Phys. Rev. D* **109** no. 10, (2024) 105018, [arXiv:2308.06312 \[hep-th\]](#).
- [23] S. Choi, J. Kim, S. Kim, and J. Nahmgoong, “Large AdS black holes from QFT,” [arXiv:1810.12067 \[hep-th\]](#).
- [24] M. Honda, “Quantum Black Hole Entropy from 4d Supersymmetric Cardy formula,” *Phys. Rev. D* **100** no. 2, (2019) 026008, [arXiv:1901.08091 \[hep-th\]](#).
- [25] A. Arabi Ardehali, “Cardy-like asymptotics of the 4d $\mathcal{N} = 4$ index and AdS₅ blackholes,” *JHEP* **06** (2019) 134, [arXiv:1902.06619 \[hep-th\]](#).
- [26] J. Kim, S. Kim, and J. Song, “A 4d $\mathcal{N} = 1$ Cardy Formula,” *JHEP* **01** (2021) 025, [arXiv:1904.03455 \[hep-th\]](#).
- [27] A. Cabo-Bizet, D. Cassani, D. Martelli, and S. Murthy, “The asymptotic growth of states of the 4d $\mathcal{N} = 1$ superconformal index,” *JHEP* **08** (2019) 120, [arXiv:1904.05865 \[hep-th\]](#).
- [28] A. Amariti, I. Garozzo, and G. Lo Monaco, “Entropy function from toric geometry,” *Nucl. Phys. B* **973** (2021) 115571, [arXiv:1904.10009 \[hep-th\]](#).
- [29] A. Arabi Ardehali, J. Hong, and J. T. Liu, “Asymptotic growth of the 4d $\mathcal{N} = 4$ index and partially deconfined phases,” *JHEP* **07** (2020) 073, [arXiv:1912.04169 \[hep-th\]](#).
- [30] A. González Lezcano, J. Hong, J. T. Liu, and L. A. Pando Zayas, “Sub-leading Structures in Superconformal Indices: Subdominant Saddles and Logarithmic Contributions,” *JHEP* **01** (2021) 001, [arXiv:2007.12604 \[hep-th\]](#).
- [31] A. Amariti, M. Fazzi, and A. Segati, “The SCI of $\mathcal{N} = 4$ USp(2N_c) and SO(N_c) SYM as a matrix integral,” *JHEP* **06** (2021) 132, [arXiv:2012.15208 \[hep-th\]](#).

- [32] D. Cassani and Z. Komargodski, “EFT and the SUSY Index on the 2nd Sheet,” *SciPost Phys.* **11** (2021) 004, [arXiv:2104.01464 \[hep-th\]](#).
- [33] A. Ogg, *Modular forms and Dirichlet series*. W. A. Benjamin, Inc., New York-Amsterdam, 1969.
- [34] A. Tauber, “Ein Satz aus der Theorie der unendlichen Reihen,” *Monatsh. Math. Phys.* **8** no. 1, (1897) 273–277. <https://doi.org/10.1007/BF01696278>.
- [35] S. Ikehara, “An extension of landau’s theorem in the analytical theory of numbers,” *Journal of Mathematics and physics* **10** no. 1-4, (1931) 1–12.
- [36] E. W. Barnes, “On the theory of the multiple gamma function,” *Trans. Cambridge Philos. Soc.* **19** (1904) 374–425.
- [37] F. Hirzebruch, *Neue topologische Methoden in der algebraischen Geometrie*. Ergebnisse der Mathematik und ihrer Grenzgebiete, (N.F.), Heft 9. Springer-Verlag, Berlin-Göttingen-Heidelberg, 1956.
- [38] A. Bayad and M. Beck, “Relations for Bernoulli-Barnes numbers and Barnes zeta functions,” *Int. J. Number Theory* **10** no. 5, (2014) 1321–1335. <https://doi.org/10.1142/S1793042114500298>.
- [39] E. Getzler and M. M. Kapranov, “Modular operads,” *Compositio Math.* **110** no. 1, (1998) 65–126. <https://doi.org/10.1023/A:1000245600345>.
- [40] A. Ivić, *The Riemann zeta-function*. Dover Publications, Inc., Mineola, NY, 2003. Theory and applications, Reprint of the 1985 original [Wiley, New York; MR0792089 (87d:11062)].
- [41] E. C. Titchmarsh, *The theory of the Riemann zeta-function*. The Clarendon Press, Oxford University Press, New York, second ed., 1986. Edited and with a preface by D. R. Heath-Brown.
- [42] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, *Higher transcendental functions. Vol. I*. Robert E. Krieger Publishing Co., Inc., Melbourne, FL, 1981. Based on notes left by Harry Bateman, With a preface by Mina Rees, With a foreword by E. C. Watson, Reprint of the 1953 original.
- [43] J. L. Cardy, “Operator Content of Two-Dimensional Conformally Invariant Theories,” *Nucl. Phys. B* **270** (1986) 186–204.

- [44] A. Gadde, S. Gukov, and P. Putrov, “(0, 2) trialities,” *JHEP* **03** (2014) 076, [arXiv:1310.0818 \[hep-th\]](#).
- [45] L. Di Pietro and Z. Komargodski, “Cardy formulae for SUSY theories in $d = 4$ and $d = 6$,” *JHEP* **12** (2014) 031, [arXiv:1407.6061 \[hep-th\]](#).
- [46] M. Buican and T. Nishinaka, “On the superconformal index of Argyres–Douglas theories,” *J. Phys. A* **49** no. 1, (2016) 015401, [arXiv:1505.05884 \[hep-th\]](#).
- [47] A. Arabi Ardehali, “High-temperature asymptotics of supersymmetric partition functions,” *JHEP* **07** (2016) 025, [arXiv:1512.03376 \[hep-th\]](#).
- [48] S. Cecotti, J. Song, C. Vafa, and W. Yan, “Superconformal Index, BPS Monodromy and Chiral Algebras,” *JHEP* **11** (2017) 013, [arXiv:1511.01516 \[hep-th\]](#).
- [49] J. Nahmgoong, “6d superconformal Cardy formulas,” *JHEP* **02** (2021) 092, [arXiv:1907.12582 \[hep-th\]](#).
- [50] C.-M. Chang, M. Fluder, Y.-H. Lin, and Y. Wang, “Proving the 6d Cardy Formula and Matching Global Gravitational Anomalies,” *SciPost Phys.* **11** no. 2, (2021) 036, [arXiv:1910.10151 \[hep-th\]](#).
- [51] N. Benjamin, J. Lee, H. Ooguri, and D. Simmons-Duffin, “Universal asymptotics for high energy CFT data,” *JHEP* **03** (2024) 115, [arXiv:2306.08031 \[hep-th\]](#).
- [52] G. H. Hardy and S. Ramanujan, “Asymptotic Formulae in Combinatory Analysis,” *Proc. London Math. Soc. (2)* **17** (1918) 75–115. <https://doi.org/10.1112/plms/s2-17.1.75>.
- [53] G. Meinardus, “Asymptotische Aussagen über Partitionen,” *Math. Z.* **59** (1954) 388–398. <https://doi.org/10.1007/BF01180268>.
- [54] B. L. Granovsky and D. Stark, “A Meinardus theorem with multiple singularities,” *Comm. Math. Phys.* **314** no. 2, (2012) 329–350. <https://doi.org/10.1007/s00220-012-1526-8>.
- [55] T. Okazaki, “M2-branes and plane partitions,” *JHEP* **07** (2022) 028, [arXiv:2204.01973 \[hep-th\]](#).

- [56] H. Hayashi, T. Nosaka, and T. Okazaki, “Asymptotic Degeneracies of M2-Brane SCFTs,” *Commun. Math. Phys.* **405** no. 7, (2024) 171, [arXiv:2307.02901 \[hep-th\]](#).
- [57] B. Henning, X. Lu, T. Melia, and H. Murayama, “Operator bases, S -matrices, and their partition functions,” *JHEP* **10** (2017) 199, [arXiv:1706.08520 \[hep-th\]](#).
- [58] T. Melia and S. Pal, “EFT Asymptotics: the Growth of Operator Degeneracy,” *SciPost Phys.* **10** no. 5, (2021) 104, [arXiv:2010.08560 \[hep-th\]](#).
- [59] S. Kim, “The Complete superconformal index for N=6 Chern-Simons theory,” *Nucl. Phys. B* **821** (2009) 241–284, [arXiv:0903.4172 \[hep-th\]](#). [Erratum: *Nucl.Phys.B* 864, 884 (2012)].
- [60] H.-C. Kim and S. Kim, “M5-branes from gauge theories on the 5-sphere,” *JHEP* **05** (2013) 144, [arXiv:1206.6339 \[hep-th\]](#).
- [61] B. Assel, D. Cassani, and D. Martelli, “Localization on Hopf surfaces,” *JHEP* **08** (2014) 123, [arXiv:1405.5144 \[hep-th\]](#).
- [62] B. Assel, D. Cassani, and D. Martelli, “Supersymmetric counterterms from new minimal supergravity,” *JHEP* **11** (2014) 135, [arXiv:1410.6487 \[hep-th\]](#).
- [63] D. Cassani and D. Martelli, “The gravity dual of supersymmetric gauge theories on a squashed $S^1 \times S^3$,” *JHEP* **08** (2014) 044, [arXiv:1402.2278 \[hep-th\]](#).
- [64] J. Lorenzen and D. Martelli, “Comments on the Casimir energy in supersymmetric field theories,” *JHEP* **07** (2015) 001, [arXiv:1412.7463 \[hep-th\]](#).
- [65] B. Assel, D. Cassani, L. Di Pietro, Z. Komargodski, J. Lorenzen, and D. Martelli, “The Casimir Energy in Curved Space and its Supersymmetric Counterpart,” *JHEP* **07** (2015) 043, [arXiv:1503.05537 \[hep-th\]](#).
- [66] N. Bobev, M. Bullimore, and H.-C. Kim, “Supersymmetric Casimir Energy and the Anomaly Polynomial,” *JHEP* **09** (2015) 142, [arXiv:1507.08553 \[hep-th\]](#).
- [67] D. Martelli and J. Sparks, “The character of the supersymmetric Casimir energy,” *JHEP* **08** (2016) 117, [arXiv:1512.02521 \[hep-th\]](#).

- [68] F. Br  nner, D. Regalado, and V. P. Spiridonov, “Supersymmetric Casimir energy and $SL(3, \mathbb{Z})$ transformations,” *JHEP* **07** (2017) 041, [arXiv:1611.03831 \[hep-th\]](#).
- [69] P. Benetti Genolini, D. Cassani, D. Martelli, and J. Sparks, “The holographic supersymmetric Casimir energy,” *Phys. Rev. D* **95** no. 2, (2017) 021902, [arXiv:1606.02724 \[hep-th\]](#).
- [70] P. Benetti Genolini, D. Cassani, D. Martelli, and J. Sparks, “Holographic renormalization and supersymmetry,” *JHEP* **02** (2017) 132, [arXiv:1612.06761 \[hep-th\]](#).
- [71] I. Papadimitriou, “Supercurrent anomalies in 4d SCFTs,” *JHEP* **07** (2017) 038, [arXiv:1703.04299 \[hep-th\]](#).
- [72] A. Arabi Ardehali, F. Larsen, J. T. Liu, and P. Szepietowski, “Quantum corrections to central charges and supersymmetric Casimir energy in AdS_3/CFT_2 ,” *JHEP* **07** (2019) 071, [arXiv:1811.12367 \[hep-th\]](#).
- [73] P. Panopoulos and I. Papadimitriou, “Supersymmetric Casimir energy on $\mathcal{N} = 1$ conformal supergravity backgrounds,” *JHEP* **04** (2024) 029, [arXiv:2312.17740 \[hep-th\]](#).
- [74] N. Kurokawa, “Multiple sine functions and Selberg zeta functions,” *Proc. Japan Acad. Ser. A Math. Sci.* **67** no. 3, (1991) 61–64. <http://projecteuclid.org/euclid.pja/1195512182>.
- [75] N. Kurokawa and S.-y. Koyama, “Multiple sine functions,” *Forum Math.* **15** no. 6, (2003) 839–876. <https://doi.org/10.1515/form.2003.042>.
- [76] M. Jimbo and T. Miwa, “Quantum KZ equation with $—q— = 1$ and correlation functions of the XXZ model in the gapless regime,” *J. Phys. A* **29** (1996) 2923–2958, [arXiv:hep-th/9601135](#).
- [77] F. Benini, R. Eager, K. Hori, and Y. Tachikawa, “Elliptic genera of two-dimensional $\mathcal{N}=2$ gauge theories with rank-one gauge groups,” *Lett. Math. Phys.* **104** (2014) 465–493, [arXiv:1305.0533 \[hep-th\]](#).
- [78] F. Benini, R. Eager, K. Hori, and Y. Tachikawa, “Elliptic Genera of 2d $\mathcal{N} = 2$ Gauge Theories,” *Commun. Math. Phys.* **333** no. 3, (2015) 1241–1286, [arXiv:1308.4896 \[hep-th\]](#).

- [79] F. Benini and N. Bobev, “Two-dimensional SCFTs from wrapped branes and c-extremization,” *JHEP* **06** (2013) 005, [arXiv:1302.4451 \[hep-th\]](#).
- [80] A. Gadde, S. Gukov, and P. Putrov, “Fivebranes and 4-manifolds,” *Prog. Math.* **319** (2016) 155–245, [arXiv:1306.4320 \[hep-th\]](#).
- [81] L. C. Jeffrey and F. C. Kirwan, “Localization for nonabelian group actions,” *Topology* **34** no. 2, (1995) 291–327.
- [82] P. Putrov, J. Song, and W. Yan, “(0,4) dualities,” *JHEP* **03** (2016) 185, [arXiv:1505.07110 \[hep-th\]](#).
- [83] A. Gadde, S. S. Razamat, and B. Willett, “On the reduction of 4d $\mathcal{N} = 1$ theories on \mathbb{S}^2 ,” *JHEP* **11** (2015) 163, [arXiv:1506.08795 \[hep-th\]](#).
- [84] M. Dedushenko and S. Gukov, “IR duality in 2D $N = (0, 2)$ gauge theory with noncompact dynamics,” *Phys. Rev. D* **99** no. 6, (2019) 066005, [arXiv:1712.07659 \[hep-th\]](#).
- [85] F. Benini and N. Bobev, “Exact two-dimensional superconformal R-symmetry and c-extremization,” *Phys. Rev. Lett.* **110** no. 6, (2013) 061601, [arXiv:1211.4030 \[hep-th\]](#).
- [86] M. Sacchi, “New 2d $\mathcal{N} = (0, 2)$ dualities from four dimensions,” *JHEP* **12** (2020) 009, [arXiv:2004.13672 \[hep-th\]](#).
- [87] K. A. Intriligator and P. Pouliot, “Exact superpotentials, quantum vacua and duality in supersymmetric $SP(N(c))$ gauge theories,” *Phys. Lett. B* **353** (1995) 471–476, [arXiv:hep-th/9505006](#).
- [88] J. Jiang, S. Nawata, and J. Zheng, “2d dualities from 4d,” *SciPost Phys.* **18** no. 6, (2025) 180, [arXiv:2407.17350 \[hep-th\]](#).
- [89] O. Aharony, S. S. Razamat, N. Seiberg, and B. Willett, “The long flow to freedom,” *JHEP* **02** (2017) 056, [arXiv:1611.02763 \[hep-th\]](#).
- [90] W. Gu and E. Sharpe, “A proposal for nonabelian mirrors,” [arXiv:1806.04678 \[hep-th\]](#).
- [91] Z. Chen, W. Gu, H. Parsian, and E. Sharpe, “Two-dimensional supersymmetric gauge theories with exceptional gauge groups,” *Adv. Theor. Math. Phys.* **24** no. 1, (2020) 67–123, [arXiv:1808.04070 \[hep-th\]](#).

- [92] W. Gu, E. Sharpe, and H. Zou, “Notes on two-dimensional pure supersymmetric gauge theories,” *JHEP* **04** (2021) 261, [arXiv:2005.10845 \[hep-th\]](#).
- [93] R. Eager and E. Sharpe, “Elliptic Genera of Pure Gauge Theories in Two Dimensions with Semisimple Non-Simply-Connected Gauge Groups,” *Commun. Math. Phys.* **387** no. 1, (2021) 267–297, [arXiv:2009.03907 \[hep-th\]](#).
- [94] G. W. Schwarz, “Representations of simple Lie groups with regular rings of invariants,” *Invent. Math.* **49** no. 2, (1978) 167–191.
<https://doi.org/10.1007/BF01403085>.
- [95] T. L. Gómez, “Algebraic stacks,” *Proc. Indian Acad. Sci. Math. Sci.* **111** no. 1, (2001) 1–31. <https://doi.org/10.1007/BF02829538>.
- [96] T. Pantev and E. Sharpe, “Notes on gauging noneffective group actions,” [arXiv:hep-th/0502027](#).
- [97] T. Pantev and E. Sharpe, “GLSM’s for Gerbes (and other toric stacks),” *Adv. Theor. Math. Phys.* **10** no. 1, (2006) 77–121, [arXiv:hep-th/0502053](#).
- [98] T. Pantev and E. Sharpe, “String compactifications on Calabi-Yau stacks,” *Nucl. Phys. B* **733** (2006) 233–296, [arXiv:hep-th/0502044](#).
- [99] E. Witten, “On the Landau-Ginzburg description of N=2 minimal models,” *Int. J. Mod. Phys. A* **9** (1994) 4783–4800, [arXiv:hep-th/9304026](#).
- [100] P. Di Francesco, O. Aharony, and S. Yankielowicz, “Elliptic genera and the Landau-Ginzburg approach to N=2 orbifolds,” *Nucl. Phys. B* **411** (1994) 584–608, [arXiv:hep-th/9306157](#).
- [101] C. Vafa and N. P. Warner, “Catastrophes and the Classification of Conformal Theories,” *Phys. Lett. B* **218** (1989) 51–58.
- [102] Y. Kazama and H. Suzuki, “New N=2 Superconformal Field Theories and Superstring Compactification,” *Nucl. Phys. B* **321** (1989) 232–268.
- [103] Y. Kazama and H. Suzuki, “Characterization of N=2 Superconformal Models Generated by Coset Space Method,” *Phys. Lett. B* **216** (1989) 112–116.
- [104] D. Gepner, “Fusion rings and geometry,” *Commun. Math. Phys.* **141** (1991) 381–411.

- [105] A. D’Adda, M. Luscher, and P. Di Vecchia, “A $1/n$ Expandable Series of Nonlinear Sigma Models with Instantons,” *Nucl. Phys. B* **146** (1978) 63–76.
- [106] A. D’Adda, P. Di Vecchia, and M. Luscher, “Confinement and Chiral Symmetry Breaking in CP^{n-1} Models with Quarks,” *Nucl. Phys. B* **152** (1979) 125–144.
- [107] E. Witten, “Instantons, the Quark Model, and the $1/n$ Expansion,” *Nucl. Phys. B* **149** (1979) 285–320.
- [108] E. Witten, “Phases of $N=2$ theories in two-dimensions,” *Nucl. Phys. B* **403** (1993) 159–222, [arXiv:hep-th/9301042](https://arxiv.org/abs/hep-th/9301042).
- [109] F. A. Dolan and H. Osborn, “Applications of the Superconformal Index for Protected Operators and q-Hypergeometric Identities to $N=1$ Dual Theories,” *Nucl. Phys. B* **818** (2009) 137–178, [arXiv:0801.4947](https://arxiv.org/abs/0801.4947) [hep-th].
- [110] D. Anselmi, D. Z. Freedman, M. T. Grisaru, and A. A. Johansen, “Nonperturbative formulas for central functions of supersymmetric gauge theories,” *Nucl. Phys. B* **526** (1998) 543–571, [arXiv:hep-th/9708042](https://arxiv.org/abs/hep-th/9708042).
- [111] E. W. Barnes, “The Genesis of the Double Gamma Functions,” *Proc. Lond. Math. Soc.* **31** (1899) 358–381.
<https://doi.org/10.1112/plms/s1-31.1.358>.
- [112] A. Arabi Ardehali, J. T. Liu, and P. Szepietowski, “High-Temperature Expansion of Supersymmetric Partition Functions,” *JHEP* **07** (2015) 113, [arXiv:1502.07737](https://arxiv.org/abs/1502.07737) [hep-th].
- [113] M. Spreafico, “On the Barnes double zeta and Gamma functions,” *J. Number Theory* **129** no. 9, (2009) 2035–2063.
<https://doi.org/10.1016/j.jnt.2009.03.005>.
- [114] D. M. Hofman and J. Maldacena, “Conformal collider physics: Energy and charge correlations,” *JHEP* **05** (2008) 012, [arXiv:0803.1467](https://arxiv.org/abs/0803.1467) [hep-th].
- [115] J. Polonyi, “Generalization of the Massive Scalar Multiplet Coupling to the Supergravity,”.
- [116] N. Seiberg, “Electric - magnetic duality in supersymmetric nonAbelian gauge theories,” *Nucl. Phys. B* **435** (1995) 129–146, [arXiv:hep-th/9411149](https://arxiv.org/abs/hep-th/9411149).

- [117] V. P. Spiridonov, “On the elliptic beta function,” *Uspekhi Mat. Nauk* **56** no. 1(337), (2001) 181–182.
<https://doi.org/10.1070/rm2001v056n01ABEH000374>.
- [118] A. D. Shapere and Y. Tachikawa, “Central charges of $N=2$ superconformal field theories in four dimensions,” *JHEP* **09** (2008) 109, [arXiv:0804.1957](#) [[hep-th](#)].
- [119] A. Gadde, L. Rastelli, S. S. Razamat, and W. Yan, “The 4d Superconformal Index from q -deformed 2d Yang-Mills,” *Phys. Rev. Lett.* **106** (2011) 241602, [arXiv:1104.3850](#) [[hep-th](#)].
- [120] A. Gadde, L. Rastelli, S. S. Razamat, and W. Yan, “Gauge Theories and Macdonald Polynomials,” *Commun. Math. Phys.* **319** (2013) 147–193, [arXiv:1110.3740](#) [[hep-th](#)].
- [121] C. Cordova and S.-H. Shao, “Schur Indices, BPS Particles, and Argyres-Douglas Theories,” *JHEP* **01** (2016) 040, [arXiv:1506.00265](#) [[hep-th](#)].
- [122] C. Cordova, D. Gaiotto, and S.-H. Shao, “Infrared Computations of Defect Schur Indices,” *JHEP* **11** (2016) 106, [arXiv:1606.08429](#) [[hep-th](#)].
- [123] D. Gaiotto and H. Kim, “3D TFTs from 4d $\mathcal{N} = 2$ BPS particles,” *JHEP* **03** (2025) 173, [arXiv:2409.20393](#) [[hep-th](#)].
- [124] A. Deb and S. S. Razamat, “Generalized Schur partition functions and RG flows,” [arXiv:2506.13764](#) [[hep-th](#)].
- [125] C. Beem and L. Rastelli, “Vertex operator algebras, Higgs branches, and modular differential equations,” *JHEP* **08** (2018) 114, [arXiv:1707.07679](#) [[hep-th](#)].
- [126] D. Xie and W. Yan, “Schur sector of Argyres-Douglas theory and W -algebra,” *SciPost Phys.* **10** no. 3, (2021) 080, [arXiv:1904.09094](#) [[hep-th](#)].
- [127] D. Xie and W. Yan, “4d $\mathcal{N} = 2$ SCFTs and lisse W -algebras,” *JHEP* **04** (2021) 271, [arXiv:1910.02281](#) [[hep-th](#)].
- [128] J. Kaidi and M. Martone, “New rank-2 Argyres-Douglas theory,” *Phys. Rev. D* **104** no. 8, (2021) 085004, [arXiv:2104.13929](#) [[hep-th](#)].

- [129] C. Beem, M. Lemos, P. Liendo, W. Peelaers, L. Rastelli, and B. C. van Rees, “Infinite Chiral Symmetry in Four Dimensions,” *Commun. Math. Phys.* **336** no. 3, (2015) 1359–1433, [arXiv:1312.5344 \[hep-th\]](#).
- [130] T. Okazaki and D. J. Smith, “Chern-Simons theories with defects, Rogers-Ramanujan type functions and eta-products,” [arXiv:2408.07893 \[hep-th\]](#).
- [131] Y. Lü, “Macdonald Identities and Exact Formulas for Superconformal Indices in Super Yang-Mills Theories,” [arXiv:2511.08468 \[hep-th\]](#).
- [132] P. C. Argyres and M. R. Douglas, “New phenomena in SU(3) supersymmetric gauge theory,” *Nucl. Phys. B* **448** (1995) 93–126, [arXiv:hep-th/9505062](#).
- [133] P. C. Argyres, M. R. Plesser, N. Seiberg, and E. Witten, “New N=2 superconformal field theories in four-dimensions,” *Nucl. Phys. B* **461** (1996) 71–84, [arXiv:hep-th/9511154](#).
- [134] D. Xie, “General Argyres-Douglas Theory,” *JHEP* **01** (2013) 100, [arXiv:1204.2270 \[hep-th\]](#).
- [135] D. Gaiotto, G. W. Moore, and A. Neitzke, “Wall-crossing, Hitchin systems, and the WKB approximation,” *Adv. Math.* **234** (2013) 239–403, [arXiv:0907.3987 \[hep-th\]](#).
- [136] Y. Wang and D. Xie, “Classification of Argyres-Douglas theories from M5 branes,” *Phys. Rev. D* **94** no. 6, (2016) 065012, [arXiv:1509.00847 \[hep-th\]](#).
- [137] J. Song, D. Xie, and W. Yan, “Vertex operator algebras of Argyres-Douglas theories from M5-branes,” *JHEP* **12** (2017) 123, [arXiv:1706.01607 \[hep-th\]](#).
- [138] J. Song, “Superconformal indices of generalized Argyres-Douglas theories from 2d TQFT,” *JHEP* **02** (2016) 045, [arXiv:1509.06730 \[hep-th\]](#).
- [139] D. Xie, W. Yan, and S.-T. Yau, “Chiral algebra of the Argyres-Douglas theory from M5 branes,” *Phys. Rev. D* **103** no. 6, (2021) 065003, [arXiv:1604.02155 \[hep-th\]](#).
- [140] S. Cecotti, A. Neitzke, and C. Vafa, “R-Twisting and 4d/2d Correspondences,” [arXiv:1006.3435 \[hep-th\]](#).
- [141] D. Gaiotto, “N=2 dualities,” *JHEP* **08** (2012) 034, [arXiv:0904.2715 \[hep-th\]](#).

- [142] O. Chacaltana, J. Distler, and Y. Tachikawa, “Nilpotent orbits and codimension-two defects of 6d $N=(2,0)$ theories,” *Int. J. Mod. Phys. A* **28** (2013) 1340006, [arXiv:1203.2930](#) [hep-th].
- [143] V. G. Kac and M. Wakimoto, “A remark on boundary level admissible representations,” *C. R. Math. Acad. Sci. Paris* **355** no. 2, (2017) 128–132. <https://doi.org/10.1016/j.crma.2017.01.008>.
- [144] D. Xie and P. Zhao, “Central charges and RG flow of strongly-coupled $N=2$ theory,” *JHEP* **03** (2013) 006, [arXiv:1301.0210](#) [hep-th].
- [145] J. Bhattacharya, S. Bhattacharyya, S. Minwalla, and S. Raju, “Indices for Superconformal Field Theories in 3,5 and 6 Dimensions,” *JHEP* **02** (2008) 064, [arXiv:0801.1435](#) [hep-th].
- [146] Y. Imamura, “Perturbative partition function for squashed S^5 ,” *PTEP* **2013** no. 7, (2013) 073B01, [arXiv:1210.6308](#) [hep-th].
- [147] H.-C. Kim, J. Kim, and S. Kim, “Instantons on the 5-sphere and M5-branes,” [arXiv:1211.0144](#) [hep-th].
- [148] G. Lockhart and C. Vafa, “Superconformal Partition Functions and Non-perturbative Topological Strings,” *JHEP* **10** (2018) 051, [arXiv:1210.5909](#) [hep-th].
- [149] K. Ohmori, H. Shimizu, Y. Tachikawa, and K. Yonekura, “Anomaly polynomial of general 6d SCFTs,” *PTEP* **2014** no. 10, (2014) 103B07, [arXiv:1408.5572](#) [hep-th].
- [150] N. Seiberg and E. Witten, “Comments on string dynamics in six-dimensions,” *Nucl. Phys. B* **471** (1996) 121–134, [arXiv:hep-th/9603003](#).
- [151] O. J. Ganor and A. Hanany, “Small $E(8)$ instantons and tensionless noncritical strings,” *Nucl. Phys. B* **474** (1996) 122–140, [arXiv:hep-th/9602120](#).
- [152] M. Del Zotto, J. J. Heckman, A. Tomasiello, and C. Vafa, “6d Conformal Matter,” *JHEP* **02** (2015) 054, [arXiv:1407.6359](#) [hep-th].
- [153] S. Yankielowicz and Y. Zhou, “Supersymmetric Rényi entropy and Anomalies in 6d (1,0) SCFTs,” *JHEP* **04** (2017) 128, [arXiv:1702.03518](#) [hep-th].
- [154] V. P. Spiridonov, “Modified elliptic gamma functions and 6d superconformal indices,” *Lett. Math. Phys.* **104** (2014) 397–414, [arXiv:1211.2703](#) [hep-th].

- [155] S. El-Showk, Y. Nakayama, and S. Rychkov, “What Maxwell Theory in $D < 4$ teaches us about scale and conformal invariance,” *Nucl. Phys. B* **848** (2011) 578–593, [arXiv:1101.5385 \[hep-th\]](#).
- [156] E. Witten, “Five-brane effective action in M -theory,” *J. Geom. Phys.* **22** (1997) 103–133, [arXiv:hep-th/9610234](#).
- [157] C. Beem, L. Rastelli, and B. C. van Rees, “ \mathcal{W} symmetry in six dimensions,” *JHEP* **05** (2015) 017, [arXiv:1404.1079 \[hep-th\]](#).
- [158] V. Adamchik, “A class of logarithmic integrals,” in *Proceedings of the 1997 International Symposium on Symbolic and Algebraic Computation (Kihei, HI)*, pp. 1–8. ACM, New York, 1997. <https://doi.org/10.1145/258726.258736>.
- [159] J. Miller and V. S. Adamchik, “Derivatives of the Hurwitz zeta function for rational arguments,” *J. Comput. Appl. Math.* **100** no. 2, (1998) 201–206. [https://doi.org/10.1016/S0377-0427\(98\)00193-9](https://doi.org/10.1016/S0377-0427(98)00193-9).
- [160] K. Matsumoto, “Asymptotic series for double zeta, double gamma, and Hecke L -functions,” *Math. Proc. Cambridge Philos. Soc.* **123** no. 3, (1998) 385–405. <https://doi.org/10.1017/S0305004197002168>.
- [161] E. Friedman and S. Ruijsenaars, “Shintani-Barnes zeta and gamma functions,” *Adv. Math.* **187** no. 2, (2004) 362–395. <https://doi.org/10.1016/j.aim.2003.07.020>.