# Solving the $\partial \overline{\partial}$ for extendable currents without vanishing the boundary cohomology group

Mamadou Eramane Bodian<br/>\*  $^{1},\;$  Souhaibou Sambou $^{2},\;$  Sény Diatta<br/>  $^{3},\;$  and Salomon Sambou $^{4}$ 

<sup>1</sup>University Assane Seck of Ziguinchor, BP: 523 (Sénégal)
<sup>1</sup>m.bodian@uni-zig.sn\*
<sup>2</sup>University Gaston Berger of Saint-Louis (Sénégal)
<sup>2</sup>souhaibou.sambou@ugb.edu.sn
<sup>3</sup>University Assane Seck of of Ziguinchor, BP: 523 (Sénégal)
<sup>3</sup>s.diatta1375@zig.univ.sn
<sup>4</sup>University Assane Seck of Ziguinchor, BP: 523 (Sénégal)
<sup>4</sup>ssambou@univ-zig.sn

#### Abstract

In this paper, we consider the problem of solving the  $\partial \overline{\partial}$  equation with discribed support for differential forms in a relatively compact domain  $\Omega$  of a complex analytic manifold X. And as a consequence, we have the solution of the equation  $\partial \overline{\partial}$  for extendable currents without the annulation assumption of the De Rham cohomology group of the boundary.

**Keywords:** operator  $\partial \overline{\partial}$ , De Rham cohomology group, Dolbeault cohomology group, Bott-Chern cohomology group, Applie cohomology group, discribed support, extendable currents

Mathematics Subject Classification (2020,2010) 32D15, 32D20, 31C10, 32L20 32F32.

# Statements and Declarations

### **Competing Interests**

The authors attest that there are non-financial interests that are directly or indirectly related to the work submitted for publication.

#### Data availability statements

We have not included any data in this manuscript.

### Ethics approval and consent to participate

Ethical authors approved and consented to participate.

### Consent for publication

The authors have consented to the publication of the article.

## **Funding**

The authors have no funding.

# Authors' contributions

The authors have worked fairly in the development and work of the article.

#### Acknowledgements

The authors have no acknowledgments.

# Introduction

In this paper, we are interested in the  $\partial \overline{\partial}$  problem with discribed support for differential forms and for currents. The problem is as follows, let X be an analytic complex manifold and  $\Omega \subset X$  a relatively compact domain such that  $X \setminus \Omega$  is connected. Then, under certain assumptions on the vanishing of De Rham and Dolbeault cohomology groups with discribed support, we have

$$H^{1,1}_{BC,\overline{\Omega}}(X) = 0.$$

Our first result is the generalization of [5, Theorem 1.2] to the case where X is a Stein manifold and as an assumption about the vanishing of the 2-th De Rham cohomology group with compact support see Theorem 2.1. Secondly, we give a more general result (see Theorem 2.2), where X is an analytic complex manifold assuming that the compact De Rham and Dolbeault cohomology groups

$$H_c^2(X) = H_c^{0,1}(X) = 0.$$

The corollary 2.1 is a solution of the  $\partial \overline{\partial}$  for (1,1)-differential forms in a completely pseudoconvex smooth-boundary domain with discribed support. And by duality, we have the solution of  $\partial \overline{\partial}$  for (n-1,n-1)-extendable currents without the vanishing hypothesis on the De Rham cohomology group of the boundary see corollary 2.2. Finally, we etablish in theorem 3.1 in the last section, the extension of these results to differential (p,q)-forms under certain conditions and as a consequence we have by duality the solution of the  $\partial \overline{\partial}$ -problem for (n-p,n-q)-extendable currents see corollary 3.1.

In case of solving the  $\partial \bar{\partial}$  equations for extendable currents, vanishing the De Rahm cohomology group of the boundary was a necessary assumption for the solution with exact support. See for exemple [3,4]. However this present result removes this ambiguity.

#### 1 Notations

We fix the following notations: Let X be a complex manifold

- $\mathcal{E}^k(X) := \text{the space of } k\text{-form of class } C^{\infty} \text{ on } X;$
- $\mathcal{D}^k(X) :=$  the space of k-form of class  $C^{\infty}$  with compact support;
- $\mathcal{D}'^{k}(X) := \text{the space of } k\text{-current of } X.$

For every  $k \in \{0, 1, \dots, 2n\}$ , we have

$$\mathcal{E}^k(X) = \bigoplus_{p+q=k} \mathcal{E}^{p,q}(X)$$

and the complex structure of X induices a splitting

$$d = \partial + \overline{\partial}$$
.

For every  $p,q \in \{1,\ldots,n\}$ , we define the Bott-Chern cohomology group of smooth (p,q)-forms on X as

$$H_{BC}^{p,q}(X) = \frac{\left(\ker \partial : \mathcal{E}^{p,q}(X) \to \mathcal{E}^{p+1,q}(X)\right) \cap \left(\ker \overline{\partial} : \mathcal{E}^{p,q}(X) \to \mathcal{E}^{p,q+1}(X)\right)}{Im\partial \overline{\partial} : \mathcal{E}^{p-1,q-1}(X) \to \mathcal{E}^{p,q}(X).}$$

 $H^{p,q}_{BC,\overline{\Omega}}(X)$  is the Bott-Chern cohomology group of smooth (p,q)-forms with support in

We define the Applie cohomology group of smooth (p,q)-forms on X as

$$H_A^{p,q}(X) = \frac{\ker(\partial\overline{\partial}) : \mathcal{E}^{p,q}(X) \to \mathcal{E}^{p+1,q+1}(X)}{Im\partial : \mathcal{E}^{p-1,q}(X) \to \mathcal{E}^{p,q}(X) + Im\overline{\partial} : \mathcal{E}^{p,q-1}(X) \to \mathcal{E}^{p,q}(X).}$$

Case of either p=0 or q=0. For example, if q=0, then the (p,0) Bott-Chern cohomology group is given, from definition, by

$$H^{p,0}_{BC}(X) = \{ f \in \Gamma(X, \Omega_X^p), | \partial f := 0 \},$$

where  $\Omega_X^p$  is the sheaf of holomorphic p-forms on X. Thanks to the symmetric property of Bott-Chern cohomology group, we have

$$H^{0,p}_{BC}(X) = \overline{H^{p,0}_{BC}(X)}.$$

 $H_A^{0,0}(X)$  is the set of pluriharmonic functions on X. The Bott-Chern cohomology group of (p,q)-forms with compact support is defined as

$$H^{p,q}_{BC,c}(X) = \frac{\left(\ker \partial : \mathcal{D}^{p,q}(X) \to \mathcal{D}^{p+1,q}(X)\right) \cap \left(\ker \overline{\partial} : \mathcal{D}^{p,q}(X) \to \mathcal{D}^{p,q+1}(X)\right)}{Im\partial \overline{\partial} : \mathcal{D}^{p-1,q-1}(X) \to \mathcal{D}^{p,q}(X).}$$

# 2 Solving the $\partial \overline{\partial}$ for extendable currents

The following result generalizes [5, Theorem 1.2],

#### Theorem 2.1.

Let X be a Stein manifold of complex dimension  $n \geq 2$  such that

$$H_c^2(X) = 0.$$

Let  $\Omega \subset X$  be a relatively compact domain such that  $X \setminus \Omega$  is connected. Then

$$H^{1,1}_{BC,\overline{\Omega}}(X) = 0.$$

Proof.

Let f be a (1,1)-differential form of class  $C^{\infty}$ , d-closed with support on  $\overline{\Omega}$ . Since  $H_c^2(X)=0$ , there exists a 1-differential form g of class  $C^{\infty}$  with compact support such that dg=f. We have  $g=g_1+g_2$  where  $g_1$  and  $g_2$  are respectively a (0,1) differential form of class  $C^{\infty}$  with compact support and a (1,0) differential form of class  $C^{\infty}$  with compact support. We have  $dg=dg_1+dg_2=f$ . Since  $d=\partial+\overline{\partial}$  and for bidegree reasons we have  $\partial g_2=0$  and  $\overline{\partial}g_1=0$ . Since X is Stein, there exist functions  $w_1$  and  $w_2$  of class  $C^{\infty}$  with compact support such that  $g_1=\overline{\partial}w_1$  and  $g_2=\partial w_2$ .

$$f = \partial g_1 + \overline{\partial} g_2 = \partial \overline{\partial} w_1 + \overline{\partial} \partial w_2 = \partial \overline{\partial} (w_1 - w_2).$$

We set  $v = w_1 - w_2$ , then v is compactly supported and  $\partial \overline{\partial} v = 0$  on  $X \setminus \overline{\Omega}$ . According to [5, theorem 1.1], there exists a pluriharmonic function  $\tilde{v}$  defined on X such that  $\tilde{v}_{X \setminus \overline{\Omega}} = v$ . We set

$$\check{v} = v - \tilde{v}$$
.

Then we have  $supp(\check{v}) \subset \overline{\Omega}$  and  $\partial \overline{\partial} \check{v} = f$ . So

$$H_{BC\overline{\Omega}}^{1,1}(X) = 0.$$

More generally, we have the following result

#### Theorem 2.2.

Let X be an analytic complex manifold of complex dimension  $n \geq 2$  such that

$$H_c^2(X) = H_c^{0,1}(X) = 0,$$

and  $\Omega \subset X$  be a relatively compact domain such that  $X \setminus \Omega$  is connected. Then,

$$H^{1,1}_{BC,\overline{\Omega}}(X) = 0.$$

Proof.

Let f be a (1,1)-differential form of class  $C^{\infty}$ , d-closed with support on  $\overline{\Omega}$ . Since  $H_c^2(X) = 0$ , there exists a 1-differential form g of class  $C^{\infty}$  with compact support such that dg = f.

We have  $g = g_1 + g_2$  where  $g_1$  and  $g_2$  are respectively a (0,1) differential form of class  $C^{\infty}$  with compact support and a (1,0) differential form of class  $C^{\infty}$  with compact support. We have  $dg = dg_1 + dg_2 = f$ . Since  $d = \partial + \overline{\partial}$  and for bidegree reasons we have  $\partial g_2 = 0$  and  $\overline{\partial} g_1 = 0$ . Since  $H_c^{0,1}(X) = 0$ , there exist functions  $w_1$  and  $w_2$  of class  $C^{\infty}$  with compact support such that  $g_1 = \overline{\partial} w_1$  and  $g_2 = \partial w_2$ .

$$f = \partial q_1 + \overline{\partial} q_2 = \partial \overline{\partial} w_1 + \overline{\partial} \partial w_2 = \partial \overline{\partial} (w_1 - w_2).$$

Let  $v=w_1-w_2,\ v$  is compactly supported and  $\partial \overline{\partial} v=0$  on  $X\setminus \overline{\Omega}$ . Then v is a pluriharmonic function on  $X\setminus \overline{\Omega}$  and which cancels on  $X\setminus \mathrm{supp} v$ . Suppose that  $\overline{\Omega}$  is strictly included in  $\mathrm{supp} v$ . So v=0 on  $X\setminus \overline{\Omega}$  which is absurd. So  $\mathrm{supp} v\subset \overline{\Omega}$ . This implies that

$$H^{1,1}_{BC\,\overline{\Omega}}(X) = 0.$$

Corollary 2.1.

Let X be a complex analytic manifold of complex dimension  $n \geq 2$  and  $\Omega \subset X$  a completely pseudoconvex domain with smooth boundary. Then

$$H^{1,1}_{BC,\overline{\Omega}}(X) = 0.$$

Proof.

Since  $\Omega$  has a smooth boundary, according to [2, Theorem 2.3], there exists a completely strictly pseudoconvex domain  $\Omega'$  such that  $\Omega \subset \Omega' \subset\subset X$  and that  $H^r(\Omega') = H^r(\Omega') = 0$  for  $r \leq 2$ . By Poincaré duality  $H_c^{n-r}(\Omega') = 0 \iff H_c^j(\Omega') = 0$  for  $0 \leq j \leq n-2$ . In particular  $H_c^2(\Omega') = 0$  and  $\Omega'$  is also a Stein domain. According to theorem 2.1,

$$H^{1,1}_{BC,\overline{\Omega}}(\Omega') = H^{1,1}_{BC,\overline{\Omega}}(X) = 0.$$

**Remark 2.1.** Corollary 2.1 shows that if f is a (1.1)-differential form of class  $C^{\infty}$  with support in  $\overline{\Omega}$  and d-closed, then there exists a function u of class  $C^{\infty}$  with support in  $\overline{\Omega}$  such that  $\partial \overline{\partial} u = f$ .

By duality, we have the following result.

#### Corollary 2.2.

Under the assumptions of the corollary 2.1, if T is a (n-1,n-1)-extendable current, d-closed and defined on  $\Omega$ , then there exists a (n-2,n-2)-extendable current S such that  $\partial \overline{\partial} S = T$ .

Proof.

We have  $\check{D}^{n-1,n-1}(\Omega) = (D^{1,1}(\overline{\Omega}))'$ . Let us show that

$$\check{D}^{n-1,n-1}(\Omega) \cap \ker d = \partial \overline{\partial} \check{D}^{n-2,n-2}(\Omega).$$

Let  $T \in \check{D}^{n-1,n-1}(\Omega) \cap \ker d$ . We define

$$L_T: \partial \overline{\partial} D^{1,1}(\overline{\Omega}) \longrightarrow \mathbb{C}$$
  
 $\partial \overline{\partial} \varphi \longmapsto \langle T, \varphi \rangle,$ 

 $L_T$  is well defined because  $\partial \overline{\partial} D^{n-2,n-2}(\overline{\Omega}) = D^{n-1,n-1}(\overline{\Omega}) \cap \ker d$  (cf corollaire 2.1). If  $\varphi$  and  $\varphi'$  are two (1,1)-differential forms such that  $\partial \overline{\partial} \varphi = \partial \overline{\partial} \varphi'$ , then

$$\varphi - \varphi' = \partial \overline{\partial} \theta \quad \text{et} \quad \langle T, \partial \overline{\partial} \theta \rangle = \lim_{j \to +\infty} \langle T, \partial \overline{\partial} \theta_j \rangle = 0 \Rightarrow \langle T, \varphi \rangle = \langle T, \varphi' \rangle$$

where  $(\theta_j)_{j\in\mathbb{N}}$  is a sequence of elements of  $D^{2,2}(\Omega)$  which converge uniformly towards  $\theta$ .  $L_T$  is linear because  $\partial \overline{\partial}: D^{2,2}(\overline{\Omega}) \longrightarrow \partial \overline{\partial} D^{2,2}(\overline{\Omega})$  is a linear and continuous map because the inverse image of an open set U of  $\mathbb{C}$  by  $L_T$  is an open set. So, we have  $L_T \circ \partial \overline{\partial} = T$  since  $L_T^{-1}(U) = \partial \overline{\partial} \circ T^{-1}(U)$ . According to the Hahn-Banach theorem, we can extend  $L_T$  into a linear and continuous operator  $\tilde{L}_T: D^{1,1}(\overline{\Omega}) \longrightarrow \mathbb{C}$ , it is an extendable current and  $\partial \overline{\partial} \tilde{L}_T \varphi = T$  then

$$\langle \partial \overline{\partial} \tilde{L}_T, \varphi \rangle = \langle T, \varphi \rangle.$$

Set  $S = \tilde{L}_T$ , S is an extendable current and verifies the relationship  $\partial \overline{\partial} S = T$ .

**Remark 2.2.** The corollary 2.2 is an improvement of [1, Theorem 1.1] without the assumption  $H^j(b\Omega) = 0$  for all  $1 \le j < 2n - 1$ .

 $\Box$ 

# 3 The case of differential (p,q)-forms and extendable currents

In this section we generalize these results to differential (p,q)-forms. Then we have:

#### Theorem 3.1.

Let X be a complex analytic manifold of dimension n and  $\Omega \subset X$  a smooth boundary domain and  $1 \leq p,q \leq n$  such that  $H^{p+q}_{c,d}(X) = H^{p-1,q}_{c,\overline{\partial}}(X) = H^{p,q-1}_{c,\overline{\partial}}(X) = 0$ . If the natural mapp

$$H^{p-1,q-1}_A(X) \ \longrightarrow \ H^{p-1,q-1}_A(X\backslash\Omega)$$

is an isomorphism then  $H^{p,q}_{BC,\overline{\Omega}}(X) = 0$ .

Proof.

Let  $[f] \in H^{p,q}_{BC,\overline{\Omega}}(X) \Longrightarrow f$  is a differential (p+q)-form d-closed with compact support. Since  $H^{p+q}_{c,d}(X) = 0$ , then there exists g a differential (p+q-1)-form with compact support such that dg = f. We can decompose g into the sum of a differential (p-1,q)-form  $g_1$  and a differential (p,q-1)-form  $g_2$ ; i.e  $g = g_1 + g_2$ . The saturation of the bidegree holds  $\overline{\partial}g_1 = 0$  and  $\partial g_2 = 0$  so there are two differential forms  $h_1$  and  $h_2$  with compact support such that  $g_1 = \overline{\partial}h_1$  and  $g_2 = \partial h_2$ . Hence

$$f = \partial g_1 + \overline{\partial} g_2$$
$$= \partial \overline{\partial} h_1 + \overline{\partial} \partial h_2$$
$$= \partial \overline{\partial} (h_1 - h_2).$$

The solution  $h_1 - h_2$  we get has compact support. We end the proof by correcting this solution to new one with exact support.

• If p = q = 1, then  $h_1 - h_2$  is a pluriharmonic function defined in  $X \setminus \overline{\Omega}$ . Under the theorem's assumptions, we have a natural isomorphism map

$$H_A^{0,0}(X) \longrightarrow H_A^{0,0}(X \backslash \Omega).$$

Then, there is u a pluriharmonic function on X such that

 $[u] = [(h_1 - h_2)_{|X\setminus\overline{\Omega}}] \Longrightarrow u_{|X\setminus\overline{\Omega}} = (h_1 - h_2)_{|X\setminus\overline{\Omega}}.$  Set  $h_1 - h_2 = h_1 - h_2 - u$  which is compactly supported on  $\Omega$  and  $\partial\overline{\partial}(\widehat{h_1 - h_2}) = f$  on  $\Omega$ .

• If  $p, q \geq 2$ ,  $h_1 - h_2$  is a (p - 1, q - 1) form with compact support. We have,  $\partial \overline{\partial}(h_1 - h_2) = 0$  on  $X \setminus \overline{\Omega}$ . Under the theorem's assumptions, we have a natural isomorphism map

$$H_A^{p-1,q-1}(X) \longrightarrow H_A^{p-1,q-1}(X \backslash \Omega).$$

Then, there is u a (p-1, q-1) pluriharmonic form on X such that

 $[u] = [(h_1 - h_2)_{|X\setminus\overline{\Omega}}] \Longrightarrow u_{|X\setminus\overline{\Omega}} = (h_1 - h_2)_{|X\setminus\overline{\Omega}} + \partial\overline{\partial}v$ , where v is a (p-2, q-2) form defined in  $X\setminus\overline{\Omega}$ . Hence  $(h_1 - h_2)_{|X\setminus\overline{\Omega}} = u - \partial\overline{\partial}\tilde{v}$  where  $\tilde{v}$  is an extension of v. Set  $h_1 - h_2 = h_1 - h_2 - u + \partial\tilde{\partial}\tilde{v}$  which is compactly supported on  $\Omega$  and  $\partial\overline{\partial}(h_1 - h_2) = f$  on  $\Omega$ .

As a consequence, we have :

**Corollary 3.1.** Let X be a complex analytic manifold of complex dimension  $n \geq 2$  and  $\Omega \subset X$  a completely pseudoconvex domain with smooth boundary. If T is a (n-p,n-q)-extendable current, d-closed and defined on  $\Omega$ , then there exists a (n-p-1,n-q-1)-extendable current S such that  $\partial \overline{\partial} S = T$ .

*Proof.* The proof is identical to that of the corollary 2.2.

# 4 Conclusion

In this paper, we solve the  $\partial\bar{\partial}$ -problem by a direct method based on Bott-Chern cohomology. What is original in this case is that, unlike in papers [1], [2], [3] and [4], we did not use the null hypothesis of the boundary cohomology group of the domain, which was a necessary condition for the solution with discribed support.

# References

- [1] E. Bodian, D. Diallo, S. Sambou: Résolution du  $\partial \overline{\partial}$  pour les courants prolongeables définis sur un domaine complétement strictement pseudoconvexe d'une variété analytique complexe, Afr. J. Pure Appl. Math., vol.3,  $n^{\circ}$  1 (2018), pp. 1 4.
- [2] S. Sambou, S. Sambou: Résolution du  $\partial \overline{\partial}$  pour les formes différentielles ayant une valeur au bord au sens des courants dans un domaine strictement pseudoconvexe, Annales Mathématiques Blaise Pascal, vol.25,  $n^{\circ}$  2 (2018), p. 315 326.
- [3] S. Sambou, E Bodian and D. Diallo: Résolution du ∂∂ pour les courants prolongeables définis sur la boule euclidienne de C<sup>n</sup>. C. R. Math. Rep. Acad. Sci. Canada Vol. 38 (1) 2016, pp. 34-37.
- [4] Sambou Souhaibou, Bodian Eramane, Ossete Ingoba Winnie and Hamidine Ibrahim. Solution of the  $\partial \overline{\partial}$ -problem in a pseudoconvex domain of  $C^n$ . The Journal of Analysis. vol 30. 2022. n 4. pp 1361-1375.
- [5] XIEPING WANG: Bott-Chern cohomology and the Hartogs extension theorem for pluriharmonic functions. Ann. Sc. Norm. Super. Pisa. CI. Sci.(5) 25 (2024), n.1, pp 605-610.