

# FACET NUMBERS OF NON-CENTRALLY SYMMETRIC REFLEXIVE POLYTOPES ARISING FROM POSETS

AKI MORI, KENTA MORI AND HIDEFUMI OHSUGI

**ABSTRACT.** Twinned chain polytopes form a broad class of non-centrally symmetric reflexive polytopes and exhibit intriguing structures. In the present paper, we show that the number of facets of  $d$ -dimensional twinned chain polytopes is at most  $6^{d/2}$ . In case  $d$  is even, the equality holds if and only if the polytope is isomorphic to a free sum of  $d/2$  copies of del Pezzo polygons. This result contributes a partial answer to Nill's conjecture: the number of facets of a  $d$ -dimensional reflexive polytope is at most  $6^{d/2}$ .

## 1. INTRODUCTION

A convex polytope  $\mathcal{P} \subset \mathbb{R}^d$  is called a *lattice polytope* if any vertex of  $\mathcal{P}$  belongs to  $\mathbb{Z}^d$ . A *reflexive polytope* is a lattice polytope that contains the origin in its interior and has the property that its dual polytope is also a lattice polytope. The concept of reflexive polytopes was introduced by Batyrev [1] in the study of Gorenstein toric Fano varieties. Since then, reflexive polytopes have been extensively studied from various perspectives, including combinatorics, commutative algebra and algebraic geometry. It is known that there are only finitely many reflexive polytopes in each dimension, up to unimodular equivalence [17]. Furthermore, if the dimension  $d$  satisfies  $d \leq 4$ , reflexive polytopes can be enumerated [14, 15].

Let  $N(\mathcal{P})$  be the number of facets of a lattice polytope  $\mathcal{P}$ . Nill proposed the following conjecture:

**Conjecture 1.1** ([21, Conjecture 5.2]). Let  $\mathcal{P}$  be a  $d$ -dimensional reflexive polytope. Then  $N(\mathcal{P}) \leq 6^{d/2}$ .

In Nill's original paper, this conjecture was stated as an upper bound on the number of vertices. For dimensions  $d \leq 4$ , this conjecture is supported by the database of reflexive polytopes:  $N(\mathcal{P}) \leq 6$  if  $d = 2$ ,  $N(\mathcal{P}) \leq 14 = \lfloor 6^{3/2} \rfloor$  if  $d = 3$ , and  $N(\mathcal{P}) \leq 36$  if  $d = 4$  [16]. A rigorous proof is known only for  $d = 2$  [21, Corollary 3.3]. If  $\mathcal{P}$  is a pseudo-symmetric reflexive simplicial  $d$ -dimensional polytope, then Conjecture 1.1 holds, and the maximum  $6^{d/2}$  is attained if and only if  $\mathcal{P}$  is a free sum of  $d/2$  copies of del Pezzo polygons [22]. Additionally, Conjecture 1.1 has been proven for symmetric edge polytopes arising from join graphs and connected bipartite graphs [20].

Let  $(P, \leq_P)$  be a finite partially ordered set ("poset" for short) on  $P = \{p_1, \dots, p_d\}$ . An *antichain* of  $P$  is a subset  $W \subset P$  such that  $p_i$  and  $p_j$  are incomparable for any  $p_i, p_j \in W$  with  $i \neq j$ . In particular, the empty set  $\emptyset$  as well as each singleton  $\{p_i\}$  is an antichain of  $P$ . Let  $\mathcal{A}(P)$  denote the set of antichains of  $P$ . For each  $A \in \mathcal{A}(P)$ , we set  $\rho(A) = \sum_{p_i \in A} \mathbf{e}_i \in \mathbb{R}^d$ , where  $\mathbf{e}_1, \dots, \mathbf{e}_d$  are the canonical unit coordinate vectors of  $\mathbb{R}^d$ . For example,  $\rho(\emptyset)$  is the origin of  $\mathbb{R}^d$ . Stanley [27]

---

2020 *Mathematics Subject Classification.* 52B20, 52B12.

*Key words and phrases.* twinned chain polytopes, number of facets, partially ordered set, reflexive polytopes.

defined a  $d$ -dimensional lattice polytope called the *chain polytope* of a poset  $P$  as

$$\mathcal{C}(P) = \text{conv}(\{\rho(A) : A \in \mathcal{A}(P)\}) \subset \mathbb{R}^d.$$

For two lattice polytopes  $\mathcal{P}, \mathcal{Q} \subset \mathbb{R}^d$ , we define

$$\Gamma(\mathcal{P}, \mathcal{Q}) = \text{conv}(\mathcal{P} \cup (-\mathcal{Q})) \subset \mathbb{R}^d.$$

Let  $(Q, \leq_Q)$  be a poset on  $Q = \{q_1, \dots, q_d\}$  with  $P \cap Q = \emptyset$ . Then the polytope  $\Gamma(\mathcal{C}(P), \mathcal{C}(Q))$  is called the *twinned chain polytope* [28]. It is known [25] that  $\Gamma(\mathcal{C}(P), \mathcal{C}(Q))$  is a reflexive polytope.

The study of  $\Gamma(\mathcal{P}, \mathcal{Q})$  was inspired by research on centrally symmetric configurations of integer matrices [24]. As in the case of symmetric edge polytopes [18], many reflexive polytopes exhibit the structure of  $\Gamma(\mathcal{P}, \mathcal{Q})$ . In particular, when  $\mathcal{P}$  and  $\mathcal{Q}$  are lattice polytopes arising from posets, important properties such as normality, reflexivity, unimodular equivalence and the Ehrhart polynomials of these polytopes have been studied in [8, 9, 10, 28]. In [28], it was shown that twinned chain polytopes  $\Gamma(\mathcal{C}(P), \mathcal{C}(Q))$  are locally anti-blocking, and hence the  $h^*$ -polynomials of  $\Gamma(\mathcal{C}(P), \mathcal{C}(Q))$  are  $\gamma$ -positive. The  $\gamma$ -positivity of  $h^*$ -polynomials of locally anti-blocking reflexive polytopes is studied in [26]. On the other hand, we define

$$\Omega(\mathcal{P}, \mathcal{Q}) = \text{conv}(\mathcal{P} \times \{1\} \cup (-\mathcal{Q}) \times \{-1\}) \subset \mathbb{R}^{d+1}.$$

The polytope  $\Omega(\mathcal{P}, \mathcal{Q})$  can be regarded as a variation of the Cayley sum of  $\mathcal{P}$  and  $\mathcal{Q}$ . In particular,  $\Omega(\mathcal{P}, \mathcal{P})$  is known as the *Hansen polytope* [5] when  $\mathcal{P}$  is the *stable set polytope* of a graph. In [4], it is shown that Kalai's  $3^d$  conjecture [13] holds for Hansen polytopes of split graphs. Combinatorial properties of  $\Omega(\mathcal{C}(P), \mathcal{C}(Q))$  and  $\Omega(2\mathcal{C}(P), 2\mathcal{C}(Q))$  are studied in [11] and [3], respectively. It is known [11] that  $\Omega(\mathcal{C}(P), \mathcal{C}(Q))$  is a reflexive polytope. Moreover,  $\Gamma(\mathcal{C}(P), \mathcal{C}(Q))$  arises as a projection of  $\Omega(2\mathcal{C}(P), 2\mathcal{C}(Q))$ . In this sense, twinned chain polytopes form a natural class of reflexive polytopes reflecting certain structural features of  $\Omega(2\mathcal{C}(P), 2\mathcal{C}(Q))$  while being connected to the combinatorics of posets.

In the present paper, we investigate the number of facets of twinned chain polytopes and confirm Conjecture 1.1 for these polytopes. We recall some notation about posets. Given a subset  $W$  of  $[d] := \{1, 2, \dots, d\}$ , the *induced subposet* of  $P$  on  $W$  is the poset  $(P_W, \leq_{P_W})$  on  $P_W = \{p_i : i \in W\}$  such that  $p_i \leq_{P_W} p_j$  if and only if  $p_i \leq_P p_j$ . The *ordinal sum* of  $P$  and  $Q$  is poset  $(P \oplus Q, \leq_{P \oplus Q})$  on  $P \oplus Q = P \cup Q$  such that  $s \leq_{P \oplus Q} t$  if (a)  $s, t \in P$  and  $s \leq_P t$  (b)  $s, t \in Q$  and  $s \leq_Q t$ , or (c)  $s \in P$  and  $t \in Q$ . If  $P$  and  $Q$  are posets on *disjoint* sets, then the *disjoint union* of  $P$  and  $Q$  is the poset  $P + Q$  on the union  $P \cup Q$  such that  $s \leq t$  in  $P + Q$  if either (a)  $s, t \in P$  and  $s \leq t$  in  $P$ , or (b)  $s, t \in Q$  and  $s \leq t$  in  $Q$ . A bijective map  $\varphi : P \rightarrow Q$  is called an *order isomorphism* if for all  $x, y \in P$ , we have  $x \leq_P y$  if and only if  $\varphi(x) \leq_Q \varphi(y)$ . We write  $P \cong Q$  whenever  $P$  and  $Q$  are order isomorphic. Let  $\mathbf{1}$  be the singleton poset,  $\mathbf{C}_n$  the  $n$ -element chain poset (i.e., a poset in which any two elements are comparable), and  $\mathbf{I}_n$  the  $n$ -element antichain poset. Given a poset  $(P, \leq_P)$ , the *comparability graph*  $G_P$  of  $P$  is the graph on the vertex set  $P$  whose edge set is  $\{\{p_i, p_j\} : p_i \text{ and } p_j \text{ is comparable in } P\}$ . Throughout this paper,  $\Gamma(\mathcal{C}(P), \mathcal{C}(Q))$  will be denoted as  $\Gamma(P, Q)$ .

The main result of this paper is as follows. This result, obtained within the framework of non-centrally symmetric polytopes, adds significant value to the study of such structures.

**Theorem 1.2.** *Let  $P = \{p_1, \dots, p_d\}$  and  $Q = \{q_1, \dots, q_d\}$  be posets with  $P \cap Q = \emptyset$ . Then we have*

$$(1.1) \quad N(\Gamma(P, Q)) \leq \begin{cases} 6^{\frac{d}{2}} & \text{if } d \text{ is even,} \\ 14 \cdot 6^{\frac{d-3}{2}} \left( = \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d}{2}} \right) & \text{if } d \text{ is odd.} \end{cases}$$




FIGURE 1. Posets  $P$  and  $Q$  for which the equality holds in Theorem 1.2

Here  $7\sqrt{6}/18 \doteq 0.95$ . In case  $d$  is even, the equality holds if and only if  $P, Q \cong \mathbf{I}_2 \oplus \cdots \oplus \mathbf{I}_2$  and the map  $\varphi : P \rightarrow Q$  defined by  $\varphi(p_i) = q_i$  induces a graph isomorphism from  $G_P$  to  $G_Q$ .

For example, if  $d = 1$ , then  $N(\Gamma(P, Q)) = 2$  satisfies (1.1). The posets  $P, Q \cong \mathbf{I}_2 \oplus \mathbf{I}_2 \oplus \mathbf{I}_2$  in Figure 1 is an example of a pair of posets for which the equality holds in Theorem 1.2. Note that the condition in Theorem 1.2 is same as that in the conjecture [2, Conjecture 2] on the facets of symmetric edge polytopes.

It is worth mentioning the number of twinned chain polytopes  $\Gamma(P, Q)$ . In Table 1, we show the number of pairs of comparability graphs on  $d$  vertices [23, A123416]. The chain polytope  $\mathcal{C}(P)$  is determined by the comparability graph  $G_P$ . However, permuting the labels of the vertices of  $G_Q$  does not preserve the unimodular equivalence of  $\Gamma(P, Q)$  (see Example 2.3 for details). Hence, the number of twinned chain polytopes can be larger than the number of pairs of comparability graphs. Therefore, twinned chain polytopes form a broad class of reflexive polytopes.

number of vertices	2	3	4	5	6	7	8
comparability graphs	2	4	11	33	144	824	6,793
pairs of comparability graphs	3	10	66	561	10,440	339,900	23,075,821

TABLE 1. Number of comparability graphs / pairs of comparability graphs

The present paper is organized as follows. In Section 2, we review the equation for the number of facets of  $\Gamma(P, Q)$  and provide remarks on labeling. We show that the ordinal sum of posets corresponds to the direct sum of the associated twinned chain polytopes, and use this to determine  $N(\Gamma(P, Q))$  when  $P$  and  $Q$  are elementary posets. In Section 3, we prove several lemmas on degrees of vertices of the comparability graphs  $G_P$  and  $G_Q$ . Finally, in Section 4, after studying the 3-dimensional case in detail, we prove Theorem 1.2.

## 2. THE NUMBER OF FACETS OF TWINNED CHAIN POLYTOPES

In the present section, we review the equation of the number of facets of twinned chain polytopes and examine the number of facets based on their geometric structure when these polytopes are associated with elementary posets.

Let  $P = \{p_1, \dots, p_d\}$  and  $Q = \{q_1, \dots, q_d\}$  be posets with  $P \cap Q = \emptyset$ . For  $W \subset [d] := \{1, \dots, d\}$ , let  $(\Delta_W(P, Q), \leq_W)$  be the ordinal sum of  $P_W$  and  $Q_{\bar{W}}$ , where  $\bar{W} = [d] \setminus W$ . Note that  $|\Delta_W(P, Q)| = d$ . A subset  $C \subset P$  is called a *chain* if  $C$  is a chain as a subposet of  $P$ . Note that the empty set is a

chain of  $P$ . A chain  $C$  of  $P$  is called *maximal* if it is not contained in a larger chain of  $P$ . We write  $\mathcal{M}(P)$  for the set of all maximal chains of  $P$ . The following proposition will be frequently used throughout this paper.

**Proposition 2.1** ([28, Theorem 3.2]). *Let  $P = \{p_1, \dots, p_d\}$  and  $Q = \{q_1, \dots, q_d\}$  be posets with  $P \cap Q = \emptyset$ . Then*

$$N(\Gamma(P, Q)) = \left| \bigcup_{W \subseteq [d]} \mathcal{M}(\Delta_W(P, Q)) \right|.$$

**Remark 2.2.** Maximal chains of a poset  $P$  correspond to the maximal cliques of the comparability graph  $G_P$  of  $P$ . From Proposition 2.1,  $N(\Gamma(P, Q)) = N(\Gamma(P', Q))$  if the map  $\varphi : P = \{p_1, \dots, p_d\} \rightarrow P' = \{p'_1, \dots, p'_d\}$  defined by  $\varphi(p_i) = p'_i$  induces a graph isomorphism from  $G_P$  to  $G_{P'}$ .

The following example shows that a permutation of the labels of the vertices of  $G_Q$  does not preserve  $N(\Gamma(P, Q))$ .

**Example 2.3.** Let  $(P, \leq_P)$ ,  $(Q, \leq_Q)$  be posets on  $P = \{p_1, p_2, p_3\}$  and  $Q = \{q_1, q_2, q_3\}$  that are isomorphic to  $\mathbf{1} \oplus \mathbf{I}_2$ . Then  $G_P$  and  $G_Q$  are paths with three vertices. Let  $G_P$  be the path  $(p_1, p_3, p_2)$ . Then  $G_Q$  is one of  $(q_1, q_3, q_2)$ ,  $(q_1, q_2, q_3)$ , or  $(q_2, q_1, q_3)$ . If  $G_Q = (q_2, q_1, q_3)$ , by exchanging  $p_1, q_1$  with  $p_2, q_2$ , this case reduces to the case  $G_Q = (q_1, q_2, q_3)$ . Hence, up to symmetries, there are two possibilities for  $G_Q$  (Figure 2 shows the pairs of posets  $P$  and  $Q$  for these cases).




FIGURE 2. Pairs of posets  $P, Q \cong \mathbf{1} \oplus \mathbf{I}_2$  in Example 2.3

(a) If  $G_Q = (q_1, q_3, q_2)$ , then Table 2 is a list of  $\mathcal{M}(\Delta_W(P, Q))$ . Each set appears exactly once in the list. Hence we have  $N(\Gamma(P, Q)) = 12$ .

$W$	$\emptyset$	$\{1\}$	$\{2\}$	$\{3\}$	$\{1, 2\}$	$\{1, 3\}$	$\{2, 3\}$	$\{1, 2, 3\}$
$\mathcal{M}(\Delta_W(P, Q))$	$\{q_1, q_3\}$ $\{q_2, q_3\}$	$\{p_1, q_2, q_3\}$	$\{p_2, q_1, q_3\}$	$\{p_3, q_1\}$ $\{p_3, q_2\}$	$\{p_1, q_3\}$ $\{p_2, q_3\}$	$\{p_1, p_3, q_2\}$	$\{p_2, p_3, q_1\}$	$\{p_1, p_3\}$ $\{p_2, p_3\}$

TABLE 2. The list of  $\mathcal{M}(\Delta_W(P, Q))$  in Example 2.3 (a)

(b) If  $G_Q = (q_1, q_2, q_3)$ , then Table 3 is a list of  $\mathcal{M}(\Delta_W(P, Q))$ . Note that  $\{p_2, q_3\}$  appears twice in the list. Hence we have  $N(\Gamma(P, Q)) = 11$ .

$W$	$\emptyset$	$\{1\}$	$\{2\}$	$\{3\}$	$\{1, 2\}$	$\{1, 3\}$	$\{2, 3\}$	$\{1, 2, 3\}$
$\mathcal{M}(\Delta_W(P, Q))$	$\{q_1, q_2\}$ $\{q_2, q_3\}$	$\{p_1, q_2, q_3\}$	$\{p_2, q_1\}$ $\{p_2, q_3\}$	$\{p_3, q_1, q_2\}$	$\{p_1, q_3\}$ $\{p_2, q_3\}$	$\{p_1, p_3, q_2\}$	$\{p_2, p_3, q_1\}$	$\{p_1, p_3\}$ $\{p_2, p_3\}$

TABLE 3. The list of  $\mathcal{M}(\Delta_W(P, Q))$  in Example 2.3 (b)

Let  $\mathcal{P} \subset \mathbb{R}^d$  be a  $d$ -dimensional polytope with  $m$  facets, and  $\mathcal{P}' \subset \mathbb{R}^{d'}$  be a  $d'$ -dimensional polytope with  $m'$  facets. Suppose that  $\mathcal{P}$  and  $\mathcal{P}'$  have the origin in their interiors. Then the  $(d + d')$ -dimensional polytope

$$\mathcal{P} \oplus \mathcal{P}' = \text{conv}(\mathcal{P} \times \{0\}^{d'} \cup \{0\}^d \times \mathcal{P}') \subset \mathbb{R}^{d+d'}$$

is called the *direct sum* of  $\mathcal{P}$  and  $\mathcal{P}'$ . It is known that  $\mathcal{P} \oplus \mathcal{P}'$  has  $m \cdot m'$  facets (see [12]).

**Proposition 2.4.** *Let  $(P_1, \leq_{P_1})$ ,  $(P_2, \leq_{P_2})$ ,  $(Q_1, \leq_{Q_1})$  and  $(Q_2, \leq_{Q_2})$  be posets with  $P_1 = \{p_1, \dots, p_s\}$ ,  $P_2 = \{p_{s+1}, \dots, p_d\}$ ,  $Q_1 = \{q_1, \dots, q_s\}$ , and  $Q_2 = \{q_{s+1}, \dots, q_d\}$ . Then*

$$N(\Gamma(P_1 \oplus P_2, Q_1 \oplus Q_2)) = N(\Gamma(P_1, Q_1)) \cdot N(\Gamma(P_2, Q_2)).$$

*In particular, if  $N(\Gamma(P_1, Q_1)) \leq 6^{\frac{s}{2}}$  and  $N(\Gamma(P_2, Q_2)) \leq 6^{\frac{d-s}{2}}$ , then*

$$N(\Gamma(P_1 \oplus P_2, Q_1 \oplus Q_2)) \leq 6^{\frac{d}{2}}.$$

*Proof.* From [7, Lemma 7.2], we have

$$(2.1) \quad \mathcal{C}(P_1 \oplus P_2) \cup (-\mathcal{C}(Q_1 \oplus Q_2)) = \mathcal{C}(P_1) \oplus \mathcal{C}(P_2) \cup (-\mathcal{C}(Q_1)) \oplus (-\mathcal{C}(Q_2)).$$

Taking the convex hull of both sides of (2.1), we obtain

$$\begin{aligned} & \Gamma(P_1 \oplus P_2, Q_1 \oplus Q_2) \\ &= \text{conv}\left(\left(\mathcal{C}(P_1) \times \{0\}^{d-s} \cup \{0\}^s \times \mathcal{C}(P_2)\right) \cup \left((- \mathcal{C}(Q_1)) \times \{0\}^{d-s} \cup \{0\}^s \times (- \mathcal{C}(Q_2))\right)\right) \\ &= \text{conv}\left(\left(\mathcal{C}(P_1) \cup (- \mathcal{C}(Q_1))\right) \times \{0\}^{d-s} \cup \{0\}^s \times \left(\mathcal{C}(P_2) \cup (- \mathcal{C}(Q_2))\right)\right) \\ &= \text{conv}\left(\Gamma(P_1, Q_1) \times \{0\}^{d-s} \cup \{0\}^s \times \Gamma(P_2, Q_2)\right) \\ &= \Gamma(P_1, Q_1) \oplus \Gamma(P_2, Q_2). \end{aligned}$$

Thus we have

$$N(\Gamma(P_1 \oplus P_2, Q_1 \oplus Q_2)) = N(\Gamma(P_1, Q_1)) \cdot N(\Gamma(P_2, Q_2)).$$

If  $N(\Gamma(P_1, Q_1)) \leq 6^{\frac{s}{2}}$  and  $N(\Gamma(P_2, Q_2)) \leq 6^{\frac{d-s}{2}}$ , then  $N(\Gamma(P_1 \oplus P_2, Q_1 \oplus Q_2)) \leq 6^{\frac{s}{2}} \cdot 6^{\frac{d-s}{2}} = 6^{\frac{d}{2}}$ .  $\square$

Theorem 1.2 holds for the following posets.

**Proposition 2.5.** *Let  $(P, \leq_P)$  and  $(Q, \leq_Q)$  be posets with  $|P| = |Q| = d$ .*

- (1) *If  $P, Q \cong \mathbf{C}_d$ , then  $N(\Gamma(P, Q)) = 2^d < 14 \cdot 6^{\frac{d-3}{2}}$ .*
- (2) *If  $P, Q \cong \mathbf{I}_d$ , then  $N(\Gamma(P, Q)) = d^2 + d$ . If  $d = 2$ , then  $d^2 + d = 6^{\frac{2}{2}}$ . If  $d \neq 2$ , then  $d^2 + d < 14 \cdot 6^{\frac{d-3}{2}}$ .*
- (3) *If  $P \cong \mathbf{I}_d$ ,  $Q \cong \mathbf{C}_d$ , then  $N(\Gamma(P, Q)) = d \cdot 2^{d-1} + 1 < 14 \cdot 6^{\frac{d-3}{2}}$ .*

*Proof.* (1) From Proposition 2.4, we have

$$N(\Gamma(\mathbf{C}_d, \mathbf{C}_d)) = N(\Gamma(\mathbf{1} \oplus \dots \oplus \mathbf{1}, \mathbf{1} \oplus \dots \oplus \mathbf{1})) = N(\Gamma(\mathbf{1}, \mathbf{1}))^d = 2^d.$$

Moreover  $2^d = 2 \cdot 2^{d-1} < \frac{7}{3} \cdot \sqrt{6}^{d-1} = 14 \cdot 6^{\frac{d-3}{2}}$ .

(2) We define the partition

$$\bigcup_{W \subset [d]} \mathcal{M}(\Delta_W(P, Q)) = \mathcal{M}_1 \sqcup \mathcal{M}_2 \sqcup \mathcal{M}_3,$$

where

$$\mathcal{M}_1 = \mathcal{M}(\Delta_\emptyset(P, Q)), \quad \mathcal{M}_2 = \mathcal{M}(\Delta_{[d]}(P, Q)) \quad \text{and} \quad \mathcal{M}_3 = \bigcup_{\emptyset \neq W \subsetneq [d]} \mathcal{M}(\Delta_W(P, Q)).$$

Since  $|\mathcal{M}_1| = |\mathcal{M}_2| = d$  and  $|\mathcal{M}_3| = d(d-1)$ , the assertion follows from Proposition 2.1. Let  $\varphi(d) = d^2 + d$  and  $\psi(d) = 14 \cdot 6^{\frac{d-3}{2}}$ . Then  $\varphi(1) = 2 < 7/3 = \psi(1)$  and  $\varphi(2) = 6$ . If  $d \geq 3$ , then

$$\frac{\varphi(d+1)}{\varphi(d)} = 1 + \frac{2}{d} < \sqrt{6} = \frac{\psi(d+1)}{\psi(d)}.$$

Since  $\varphi(3) = 12 < 14 = \psi(3)$ , we have  $\varphi(d) < \psi(d)$  for  $d \geq 3$ .

(3) It is known [28, Example 3.5] that  $N(\Gamma(P, Q)) = d \cdot 2^{d-1} + 1$ . Let  $\varphi(d) = d \cdot 2^{d-1} + 1$  and  $\psi(d) = 14 \cdot 6^{\frac{d-3}{2}}$ . It is easy to check that  $\varphi(d) < \psi(d)$  for  $d \leq 5$ . If  $d \geq 5$ , then

$$\frac{\varphi(d+1)}{\varphi(d)} = \frac{(d+1) \cdot 2^d + 1}{d \cdot 2^{d-1} + 1} = 2 + \frac{2}{d} - \frac{d+2}{d(d \cdot 2^{d-1} + 1)} < 2 + \frac{2}{d} < \sqrt{6} = \frac{\psi(d+1)}{\psi(d)}.$$

Thus  $\varphi(d) < \psi(d)$  for  $d \geq 6$ . □

In particular, Theorem 1.2 holds for  $d \leq 2$ . (Figure 3 shows all  $\Gamma(P, Q)$  for  $d = 2$ .)

**Corollary 2.6.** *Let  $(P, \leq_P)$  and  $(Q, \leq_Q)$  be posets with  $|P| = |Q| = 2$ . Then we have  $N(\Gamma(P, Q)) \leq 6$ . The equality holds if and only if  $P$  and  $Q$  are isomorphic to  $\mathbf{I}_2$ .*




FIGURE 3. Twinned chain polytopes for  $d = 2$

**Corollary 2.7.** *Let  $(P, \leq_P)$  and  $(Q, \leq_Q)$  be posets with  $|P| = |Q| = d$ . If  $P, Q \cong \mathbf{I}_2 \oplus \cdots \oplus \mathbf{I}_2$ , and  $\{p_{2i-1}, p_{2i}\}, \{q_{2i-1}, q_{2i}\} \cong \mathbf{I}_2$  for  $1 \leq i \leq d/2$ , then  $N(\Gamma(P, Q)) = 6^{\frac{d}{2}}$ .*

*Proof.* It follows from Proposition 2.4 and Proposition 2.5 (2). □

The following lemma follows from a well-known fact in graph theory.

**Lemma 2.8** ([19]). *Let  $(P, \leq_P)$  be a poset with  $|P| = d$ . Then  $|\mathcal{M}(P)| \leq \lfloor 3^{\frac{d}{3}} \rfloor$ .*

From Lemma 2.8, one can show that Theorem 1.2 holds when  $P$  is a chain.

$d$	3	5	7	9	11	13	15
$\sum_{i=0}^d \binom{d}{i} \lfloor 3^{\frac{i}{3}} \rfloor$	13	82	496	2971	17756	106522	640651
$14 \cdot 6^{\frac{d-3}{2}} \left( = \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d}{2}} \right)$	14	84	504	3024	18144	108864	653184

TABLE 4. Numbers appearing in Proof of Proposition 2.9

**Proposition 2.9.** *Let  $P = \{p_1, \dots, p_d\}$  and  $Q = \{q_1, \dots, q_d\}$  be posets with  $P \cap Q = \emptyset$ . Suppose that either  $\deg_{G_P} p_i = d - 1$  or  $\deg_{G_Q} q_i = d - 1$  for each  $i = 1, 2, \dots, d$ . Then*

$$N(\Gamma(P, Q)) < \begin{cases} 6^{\frac{d}{2}} & \text{if } d \text{ is even,} \\ 14 \cdot 6^{\frac{d-3}{2}} & \text{if } d \text{ is odd.} \end{cases}$$

*Proof.* We may assume that  $d \geq 3$ . Suppose that  $\deg_{G_P} p_i = d - 1$  for  $i = 1, 2, \dots, m$  and  $\deg_{G_Q} q_j = d - 1$  for  $j = m + 1, m + 2, \dots, d$ . Let  $W \subset [d]$ . We fix an arbitrary maximal chain  $C \in \mathcal{M}(\Delta_W(P, Q))$ . Since  $\deg_{G_P} p_i = d - 1$  for any  $i \in [m]$ , we have  $P_{W \cap [m]} \subset C$ . Similarly, since  $\deg_{G_Q} q_j = d - 1$  for  $j = m + 1, m + 2, \dots, d$ , we have  $Q_{\overline{W} \cap \{m+1, \dots, d\}} \subset C$ . On the other hand,

$$C \setminus (P_{W \cap [m]} \cup Q_{\overline{W} \cap \{m+1, \dots, d\}})$$

is a maximal chain of

$$P_{W \cap \{m+1, \dots, d\}} \oplus Q_{\overline{W} \cap [m]}.$$

Therefore,

$$(2.2) \quad N(\Gamma(P, Q)) = \sum_{W \subset [d]} |\mathcal{M}(P_{W \cap \{m+1, \dots, d\}})| |\mathcal{M}(Q_{\overline{W} \cap [m]})|.$$

Let  $|\overline{W} \cap [m]| = k$  and  $|W \cap \{m + 1, \dots, d\}| = \ell$ . By Lemma 2.8, we obtain

$$N(\Gamma(P, Q)) \leq \left( \sum_{k=0}^m \binom{m}{k} \lfloor 3^{\frac{k}{3}} \rfloor \right) \left( \sum_{\ell=0}^{d-m} \binom{d-m}{\ell} \lfloor 3^{\frac{\ell}{3}} \rfloor \right) \leq (1 + \sqrt[3]{3})^d < 6^{\frac{d}{2}}$$

since  $1 + \sqrt[3]{3} < \sqrt{6}$ . Moreover, since  $(1 + \sqrt[3]{3})^{17} < \frac{7\sqrt{6}}{18} \cdot 6^{\frac{17}{2}}$ , we have  $(1 + \sqrt[3]{3})^d < \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d}{2}}$  for any  $d \geq 17$ . For odd  $d$  with  $3 \leq d \leq 15$ , one can check

$$N(\Gamma(P, Q)) \leq \left( \sum_{k=0}^m \binom{m}{k} \lfloor 3^{\frac{k}{3}} \rfloor \right) \left( \sum_{\ell=0}^{d-m} \binom{d-m}{\ell} \lfloor 3^{\frac{\ell}{3}} \rfloor \right) \leq \sum_{i=0}^d \binom{d}{i} \lfloor 3^{\frac{i}{3}} \rfloor < \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d}{2}}$$

through explicit computations (see Table 4). □

The following corollary follows immediately from equation (2.2).

**Corollary 2.10.** *If  $P \cong \mathbf{C}_d$ , then*

$$N(\Gamma(P, Q)) = \sum_{W \subset [d]} |\mathcal{M}(Q_W)| < \begin{cases} 6^{\frac{d}{2}} & \text{if } d \text{ is even,} \\ 14 \cdot 6^{\frac{d-3}{2}} & \text{if } d \text{ is odd.} \end{cases}$$



### 3. DEGREES OF VERTICES OF THE COMPARABILITY GRAPHS

In the present section, we prove several lemmas on degrees of vertices of the comparability graphs  $G_P$  and  $G_Q$ . Lemma 3.1 will be frequently used throughout this section. Let  $c(P)$  denote the number of chains in  $P$ .

**Lemma 3.1.** *Let  $P = \{p_1, \dots, p_d\}$  and  $Q = \{q_1, \dots, q_d\}$  be posets with  $P \cap Q = \emptyset$  and let*

$$\mathcal{M} = \left\{ C \in \bigcup_{W \subset [d]} \mathcal{M}(\Delta_W(P, Q)) : p_d \in C \right\}.$$

*If  $p_d$  is incomparable with  $p_{i_1}, \dots, p_{i_s}$  in  $P$  with  $0 \leq s \leq d-1$ , then we have*

$$(3.1) \quad |\mathcal{M}| \leq c(Q_{\{i_1, \dots, i_s\}}) \cdot N(\Gamma(P_{[d-1] \setminus \{i_1, \dots, i_s\}}, Q_{[d-1] \setminus \{i_1, \dots, i_s\}})).$$

*Proof.* Let

$$\begin{aligned} \mathcal{M}' &= \bigcup_{W \subset [d-1] \setminus \{i_1, \dots, i_s\}} \mathcal{M}(\Delta_W(P_{[d-1] \setminus \{i_1, \dots, i_s\}}, Q_{[d-1] \setminus \{i_1, \dots, i_s\}})), \\ \mathcal{M}'' &= \{(C_1, C') : C_1 \text{ is a chain of } Q_{\{i_1, \dots, i_s\}}, C' \in \mathcal{M}'\}. \end{aligned}$$

It is enough to show that there exists a surjective map  $\varphi : \mathcal{M}'' \rightarrow \mathcal{M}$ . Let  $(C_1, C') \in \mathcal{M}''$ . Then

$$C' \in \mathcal{M}(\Delta_W(P_{[d-1] \setminus \{i_1, \dots, i_s\}}, Q_{[d-1] \setminus \{i_1, \dots, i_s\}}))$$

for some  $W \subset [d-1] \setminus \{i_1, \dots, i_s\}$ . Let

$$C = \{p_d\} \sqcup C_1 \sqcup C_2 \sqcup C_3,$$

where

$$\begin{aligned} C_2 &= \{p \in C' \cap P : p \text{ is comparable with } p_d\}, \\ C_3 &= \{q \in C' \cap Q : \{q\} \cup C_1 \text{ is a chain of } Q\}. \end{aligned}$$

Then  $C$  belongs to  $\mathcal{M}(\Delta_{W'}(P, Q))$  where  $W' = \{d\} \sqcup W \sqcup (\{i_1, \dots, i_s\} \setminus \{i : q_i \in C_1\})$ . Hence  $C \in \mathcal{M}$ . We define  $\varphi$  by  $\varphi((C_1, C')) = C$ .

We now show that  $\varphi$  is surjective. Suppose that  $C \in \mathcal{M}$ . Then  $C \in \mathcal{M}(\Delta_{W'}(P, Q))$  for some  $d \in W' \subset [d]$ . Let

$$\begin{aligned} C_1 &:= C \cap Q_{\{i_1, \dots, i_s\}}, \\ C_2 &:= (C \cap P) \setminus \{p_d\}, \\ C_3 &:= C \cap Q_{[d-1] \setminus \{i_1, \dots, i_s\}}. \end{aligned}$$

Then  $C_1$  is a chain of  $Q_{\{i_1, \dots, i_s\}}$ , and there exists  $C' \in \mathcal{M}(\Delta_W(P_{[d-1] \setminus \{i_1, \dots, i_s\}}, Q_{[d-1] \setminus \{i_1, \dots, i_s\}}))$  where  $W = W' \cap ([d-1] \setminus \{i_1, \dots, i_s\})$  such that  $C_2 \cup C_3 \subset C'$ . It follows that  $\varphi((C_1, C')) = C$  and hence  $\varphi$  is surjective. Thus we have (3.1), as desired.  $\square$

**Lemma 3.2.** *Let  $P = \{p_1, \dots, p_d\}$  and  $Q = \{q_1, \dots, q_d\}$  be posets with  $P \cap Q = \emptyset$ . Suppose that  $p_d$  is incomparable with  $p_{i_1}, \dots, p_{i_s}$  in  $P$  and  $q_d$  is incomparable with  $q_{j_1}, \dots, q_{j_t}$  in  $Q$ . Then we have*

$$(3.2) \quad \begin{aligned} N(\Gamma(P, Q)) &\leq N(\Gamma(P_{[d-1]}, Q_{[d-1]})) \\ &\quad + c(Q_{\{i_1, \dots, i_s\}}) \cdot N(\Gamma(P_{[d-1] \setminus \{i_1, \dots, i_s\}}, Q_{[d-1] \setminus \{i_1, \dots, i_s\}})) \\ &\quad + c(P_{\{j_1, \dots, j_t\}}) \cdot N(\Gamma(P_{[d-1] \setminus \{j_1, \dots, j_t\}}, Q_{[d-1] \setminus \{j_1, \dots, j_t\}})). \end{aligned}$$



In particular, if (1.1) holds for any posets with  $\leq d-1$  elements, and if  $st \neq 0$  and  $s+t \geq 4$ , then

$$N(\Gamma(P, Q)) \begin{cases} < 6^{\frac{d}{2}} & \text{if } d \text{ is even,} \\ \leq 14 \cdot 6^{\frac{d-3}{2}} & \text{if } d \text{ is odd.} \end{cases}$$

*Proof.* We define a partition  $\bigcup_{W \subset [d]} \mathcal{M}(\Delta_W(P, Q)) = \mathcal{M}_1 \sqcup \mathcal{M}_2 \sqcup \mathcal{M}_3$  where

$$\begin{aligned} \mathcal{M}_1 &= \left\{ C \in \bigcup_{W \subset [d]} \mathcal{M}(\Delta_W(P, Q)) : p_d, q_d \notin C \right\}, \\ \mathcal{M}_2 &= \left\{ C \in \bigcup_{W \subset [d]} \mathcal{M}(\Delta_W(P, Q)) : p_d \in C \right\}, \\ \mathcal{M}_3 &= \left\{ C \in \bigcup_{W \subset [d]} \mathcal{M}(\Delta_W(P, Q)) : q_d \in C \right\}. \end{aligned}$$

It is easy to see that  $\mathcal{M}_1 \subset \bigcup_{W \subset [d-1]} \mathcal{M}(\Delta_W(P_{[d-1]}, Q_{[d-1]}))$  and hence

$$|\mathcal{M}_1| \leq N(\Gamma(P_{[d-1]}, Q_{[d-1]})).$$

From Lemma 3.1,

$$\begin{aligned} |\mathcal{M}_2| &\leq c(Q_{\{i_1, \dots, i_s\}}) \cdot N(\Gamma(P_{[d-1] \setminus \{i_1, \dots, i_s\}}, Q_{[d-1] \setminus \{i_1, \dots, i_s\}})), \\ |\mathcal{M}_3| &\leq c(P_{\{j_1, \dots, j_t\}}) \cdot N(\Gamma(P_{[d-1] \setminus \{j_1, \dots, j_t\}}, Q_{[d-1] \setminus \{j_1, \dots, j_t\}})). \end{aligned}$$

Hence we have (3.2). Thus

$$N(\Gamma(P, Q)) \leq 6^{\frac{d-1}{2}} + 2^s \cdot 6^{\frac{d-s-1}{2}} + 2^t \cdot 6^{\frac{d-t-1}{2}} = \left( 1 + \left( \frac{2}{\sqrt{6}} \right)^s + \left( \frac{2}{\sqrt{6}} \right)^t \right) 6^{\frac{d-1}{2}}.$$

Suppose that  $st \neq 0$ . Then

$$1 + \left( \frac{2}{\sqrt{6}} \right)^s + \left( \frac{2}{\sqrt{6}} \right)^t \begin{cases} \leq 1 + \frac{2}{\sqrt{6}} + \frac{2}{3} \cdot \frac{2}{\sqrt{6}} < \sqrt{6} & \text{if } s+t \geq 4, \\ \leq 1 + \frac{2}{\sqrt{6}} + \frac{4}{9} < \frac{7\sqrt{6}}{18} \cdot \sqrt{6} & \text{if } s+t \geq 5, \\ = 1 + \frac{2}{3} + \frac{2}{3} = \frac{7\sqrt{6}}{18} \cdot \sqrt{6} & \text{if } s=t=2. \end{cases}$$

If  $d$  is odd and  $(s, t) = (1, 3)$ , then

$$N(\Gamma(P, Q)) \leq 6^{\frac{d-1}{2}} + 2 \cdot \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d-2}{2}} + 2^3 \cdot \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d-4}{2}} = \left( \frac{3}{7} + \frac{1}{3} + \frac{2}{9} \right) \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d}{2}} = \frac{62}{63} \cdot \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d}{2}}.$$

□

**Lemma 3.3.** Let  $P = \{p_1, \dots, p_d\}$  and  $Q = \{q_1, \dots, q_d\}$  be posets with  $P \cap Q = \emptyset$ . Suppose that

- $p_d$  is incomparable with  $p_i$  and comparable with  $p_{i'}$  for all  $i' \in [d-1] \setminus \{i\}$ ,
- $q_d$  is incomparable with  $q_j$  and comparable with  $q_{j'}$  for all  $j' \in [d-1] \setminus \{j\}$ ,
- (1.1) holds for any posets with  $\leq d-1$  elements,

then  $N(\Gamma(P, Q))$  satisfies (1.1). The equality  $N(\Gamma(P, Q)) = 6^{\frac{d}{2}}$  holds if and only if  $d$  is even,  $i = j$ ,  $N(\Gamma(P_{[d-1] \setminus \{i\}}, Q_{[d-1] \setminus \{i\}})) = 6^{\frac{d-2}{2}}$ ,  $P = \{p_i, p_d\} \oplus P_{[d-1] \setminus \{i\}}$  and  $Q = \{q_i, q_d\} \oplus Q_{[d-1] \setminus \{i\}}$ .

*Proof.* We define

$$\begin{aligned} \mathcal{M}_1 &= \left\{ C \in \bigcup_{w \subset [d]} \mathcal{M}(\Delta_w(P, Q)) : p_d \in C \right\}, \\ \mathcal{M}_2 &= \left\{ C \in \bigcup_{w \subset [d]} \mathcal{M}(\Delta_w(P, Q)) : q_d \in C \right\}, \\ \mathcal{M}_3 &= \left\{ C \in \bigcup_{w \subset [d]} \mathcal{M}(\Delta_w(P, Q)) : p_i \in C, q_d \notin C \right\} \\ &= \left\{ C \in \bigcup_{w \subset [d-1]} \mathcal{M}(\Delta_w(P_{[d-1]}, Q_{[d-1]})) : p_i \in C \right\}, \\ \mathcal{M}_4 &= \left\{ C \in \bigcup_{w \subset [d]} \mathcal{M}(\Delta_w(P, Q)) : q_j \in C, p_d \notin C \right\} \\ &= \left\{ C \in \bigcup_{w \subset [d-1]} \mathcal{M}(\Delta_w(P_{[d-1]}, Q_{[d-1]})) : q_j \in C \right\}. \end{aligned}$$

For any  $C \in \bigcup_{w \subset [d]} \mathcal{M}(\Delta_w(P, Q))$ , if  $p_d, q_d \notin C$ , then either  $p_i$  or  $q_j$  belongs to  $C$ . Hence

$$\bigcup_{w \subset [d]} \mathcal{M}(\Delta_w(P, Q)) = \mathcal{M}_1 \sqcup \mathcal{M}_2 \sqcup (\mathcal{M}_3 \cup \mathcal{M}_4).$$

From Lemma 3.1,

$$\begin{aligned} (3.3) \quad |\mathcal{M}_1| &\leq 2 \cdot N(\Gamma(P_{[d-1] \setminus \{i\}}, Q_{[d-1] \setminus \{i\}})), \\ |\mathcal{M}_2| &\leq 2 \cdot N(\Gamma(P_{[d-1] \setminus \{j\}}, Q_{[d-1] \setminus \{j\}})), \\ |\mathcal{M}_3| &\leq N(\Gamma(P_{[d-1] \setminus \{i\}}, Q_{[d-1] \setminus \{i\}})), \\ |\mathcal{M}_4| &\leq N(\Gamma(P_{[d-1] \setminus \{j\}}, Q_{[d-1] \setminus \{j\}})). \end{aligned}$$

Thus we have

$$\begin{aligned} (3.4) \quad N(\Gamma(P, Q)) &\leq 3 \cdot N(\Gamma(P_{[d-1] \setminus \{i\}}, Q_{[d-1] \setminus \{i\}})) + 3 \cdot N(\Gamma(P_{[d-1] \setminus \{j\}}, Q_{[d-1] \setminus \{j\}})) \\ &\leq \begin{cases} 6 \cdot 6^{\frac{d-2}{2}} = 6^{\frac{d}{2}} & \text{if } d \text{ is even,} \\ 6 \cdot \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d-2}{2}} = \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d}{2}} & \text{if } d \text{ is odd.} \end{cases} \end{aligned}$$

Suppose that  $N(\Gamma(P, Q)) = 6^{\frac{d}{2}}$ . Then  $d$  is even and

- The equalities hold for the inequalities in (3.3),
- $\mathcal{M}_3 \cap \mathcal{M}_4 = \emptyset$ , and hence  $i = j$ , and
- $N(\Gamma(P_{[d-1] \setminus \{i\}}, Q_{[d-1] \setminus \{i\}})) = 6^{\frac{d-2}{2}}$ .

If  $p_i$  is incomparable with  $p_k$  for some  $k \in [d-1]$ , then

$$|\mathcal{M}_3| \leq 2 \cdot N(\Gamma(P_{[d-1] \setminus \{i,k\}}, Q_{[d-1] \setminus \{i,k\}})) \leq 2 \cdot \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d-3}{2}} = \frac{7}{9} \cdot 6^{\frac{d-2}{2}} < 6^{\frac{d-2}{2}}$$

by Lemma 3.1. This is a contradiction. Hence  $p_i$  is comparable with  $p_k$  for any  $k \in [d-1]$ . Similarly,  $q_j (= q_i)$  is comparable with  $q_k$  for any  $k \in [d-1]$ . Thus  $P = \{p_i, p_d\} \oplus P_{[d-1] \setminus \{i\}}$  and  $Q = \{q_i, q_d\} \oplus Q_{[d-1] \setminus \{i\}}$ . The converse follows from Proposition 2.4.  $\square$

**Lemma 3.4.** Let  $P = \{p_1, \dots, p_d\}$  and  $Q = \{q_1, \dots, q_d\}$  be posets with  $P \cap Q = \emptyset$ . Suppose that

- $p_d$  is incomparable with  $p_i$  and comparable with  $p_{i'}$  for all  $i' \in [d-1] \setminus \{i\}$ ,
- $q_d$  is incomparable with  $q_{j_1}, q_{j_2}$  and comparable with  $q_{j'}$  for all  $j' \in [d-1] \setminus \{j_1, j_2\}$ ,
- (1.1) holds for any posets with  $\leq d-1$  elements,

then

$$N(\Gamma(P, Q)) < \begin{cases} 6^{\frac{d}{2}} & \text{if } d \text{ is even,} \\ 14 \cdot 6^{\frac{d-3}{2}} & \text{if } d \text{ is odd.} \end{cases}$$

*Proof.* We define a partition  $\bigcup_{W \subset [d]} \mathcal{M}(\Delta_W(P, Q)) = \mathcal{M}_1 \sqcup \mathcal{M}_2 \sqcup \mathcal{M}_3 \sqcup \mathcal{M}_4$ , where

$$\begin{aligned} \mathcal{M}_1 &= \left\{ C \in \bigcup_{W \subset [d]} \mathcal{M}(\Delta_W(P, Q)) : p_d, q_d \notin C \right\}, \\ \mathcal{M}_2 &= \left\{ C \in \bigcup_{W \subset [d]} \mathcal{M}(\Delta_W(P, Q)) : p_d \in C \right\}, \\ \mathcal{M}_3 &= \left\{ C \in \bigcup_{W \subset [d]} \mathcal{M}(\Delta_W(P, Q)) : p_i, q_d \in C \right\}, \\ \mathcal{M}_4 &= \left\{ C \in \bigcup_{W \subset [d]} \mathcal{M}(\Delta_W(P, Q)) : q_d \in C, p_i \notin C \right\}. \end{aligned}$$

By the assumption for  $p_d$  and  $q_d$ , it follows that

$$\mathcal{M}_1 = \left\{ C \in \bigcup_{W \subset [d-1]} \mathcal{M}(\Delta_W(P_{[d-1]}, Q_{[d-1]})) : \{p_i, q_{j_1}, q_{j_2}\} \cap C \neq \emptyset \right\}.$$

On the other hand,  $|\mathcal{M}_4| = |\mathcal{M}|$ , where

$$\mathcal{M} = \left\{ C \in \bigcup_{W \subset [d-1]} \mathcal{M}(\Delta_W(P_{[d-1]}, Q_{[d-1]})) : \{p_i, q_{j_1}, q_{j_2}\} \cap C = \emptyset \right\}$$

since  $\mathcal{M}_4 = \{C \cup \{q_d\} : C \in \mathcal{M}\}$ . Hence we have  $|\mathcal{M}_1| + |\mathcal{M}_4| = |\mathcal{M}_1| + |\mathcal{M}| = N(\Gamma(P_{[d-1]}, Q_{[d-1]}))$ . From Lemma 3.1,

$$|\mathcal{M}_2| \leq 2 \cdot N(\Gamma(P_{[d-1] \setminus \{i\}}, Q_{[d-1] \setminus \{i\}})).$$

**Case 1.** ( $i \notin \{j_1, j_2\}$ .) First, we show

$$(3.5) \quad |\mathcal{M}_3| \leq 4 \cdot N(\Gamma(P_{[d-1] \setminus \{i, j_1, j_2\}}, Q_{[d-1] \setminus \{i, j_1, j_2\}}))$$

from a similar argument as in Proof of Lemma 3.1. Let

$$\mathcal{M}' = \bigcup_{W \subset [d-1] \setminus \{i, j_1, j_2\}} \mathcal{M}(\Delta_W(P_{[d-1] \setminus \{i, j_1, j_2\}}, Q_{[d-1] \setminus \{i, j_1, j_2\}})),$$

$$\mathcal{M}'' = \{(C_1, C') : C' \in \mathcal{M}', C_1 \text{ is a chain of } P_{\{j_1, j_2\}} \text{ such that } \{p_i\} \cup C_1 \text{ is a chain}\}.$$

It is enough to show that there exists a surjective map  $\varphi : \mathcal{M}'' \rightarrow \mathcal{M}_3$ . Let  $(C_1, C') \in \mathcal{M}''$ . Then

$$C' \in \mathcal{M}(\Delta_W(P_{[d-1] \setminus \{i, j_1, j_2\}}, Q_{[d-1] \setminus \{i, j_1, j_2\}}))$$

for some  $W \subset [d-1] \setminus \{i, j_1, j_2\}$ . Let

$$C = \{p_i, q_d\} \sqcup C_1 \sqcup C_2 \sqcup C_3,$$

where

$$C_2 = \{q \in C' \cap Q : q \text{ is comparable with } q_d\},$$

$$C_3 = \{p \in C' \cap P : \{p, p_i\} \cup C_1 \text{ is a chain of } P\}.$$

Then  $C$  belongs to  $\mathcal{M}(\Delta_{W'}(P, Q))$  where  $W' = \{i\} \sqcup W \sqcup \{k : p_k \in C_1\}$ . Hence  $C \in \mathcal{M}_3$ . We define  $\varphi$  by  $\varphi((C_1, C')) = C$ .

We now show that  $\varphi$  is surjective. Suppose that  $C \in \mathcal{M}_3$ . Then  $C \in \mathcal{M}(\Delta_{W'}(P, Q))$  for some  $W' \subset [d-1]$ . Let

$$C_1 := C \cap \{p_{j_1}, p_{j_2}\},$$

$$C_2 := (C \cap Q) \setminus \{q_d\},$$

$$C_3 := C \cap P_{[d-1] \setminus \{i, j_1, j_2\}}.$$

Then  $C_1$  is a chain of  $\{p_{j_1}, p_{j_2}\}$  such that  $\{p_i\} \cup C_1$  is a chain, and there exists

$$C' \in \mathcal{M}(\Delta_W(P_{[d-1] \setminus \{i, j_1, j_2\}}, Q_{[d-1] \setminus \{i, j_1, j_2\}})),$$

where  $W = W' \setminus \{i, j_1, j_2\}$  such that  $C_2 \cup C_3 \subset C'$ . It follows that  $\varphi((C_1, C')) = C$  and hence  $\varphi$  is surjective. Thus we have (3.5).

Therefore,

$$\begin{aligned} N(\Gamma(P, Q)) &\leq N(\Gamma(P_{[d-1]}, Q_{[d-1]})) + 2 \cdot N(\Gamma(P_{[d-1] \setminus \{i\}}, Q_{[d-1] \setminus \{i\}})) \\ &\quad + 4 \cdot N(\Gamma(P_{[d-1] \setminus \{i, j_1, j_2\}}, Q_{[d-1] \setminus \{i, j_1, j_2\}})). \end{aligned}$$

If  $d$  is even, then

$$N(\Gamma(P, Q)) \leq \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d-1}{2}} + 2 \cdot 6^{\frac{d-2}{2}} + 4 \cdot 6^{\frac{d-4}{2}} = \left(\frac{7}{18} + \frac{1}{3} + \frac{1}{9}\right) \cdot 6^{\frac{d}{2}} = \frac{5}{6} \cdot 6^{\frac{d}{2}} < 6^{\frac{d}{2}}.$$

If  $d$  is odd, then

$$\begin{aligned} N(\Gamma(P, Q)) &\leq 6^{\frac{d-1}{2}} + 2 \cdot \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d-2}{2}} + 4 \cdot \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d-4}{2}} \\ &= \left(\frac{3}{7} + \frac{1}{3} + \frac{1}{9}\right) \cdot \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d}{2}} \end{aligned}$$

$$= \frac{55}{63} \cdot \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d}{2}} < \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d}{2}}.$$

**Case 2.** ( $i = j_1$ .) From Lemma 3.1, we have

$$|\{C \in \bigcup_{W \subset [d]} \mathcal{M}(\Delta_W(P, Q)) : q_d \in C\}| \leq c(P_{j_1, j_2}) \cdot N(\Gamma(P_{[d-1] \setminus \{j_1, j_2\}}, Q_{[d-1] \setminus \{j_1, j_2\}})).$$

Since the number of chains in  $\{p_{j_1}, p_{j_2}\}$  that contains  $p_{j_1}$  is at most 2, we have

$$|\mathcal{M}_3| \leq 2 \cdot N(\Gamma(P_{[d-1] \setminus \{j_1, j_2\}}, Q_{[d-1] \setminus \{j_1, j_2\}})).$$

Therefore,

$$\begin{aligned} N(\Gamma(P, Q)) &\leq N(\Gamma(P_{[d-1]}, Q_{[d-1]})) + 2 \cdot N(\Gamma(P_{[d-1] \setminus \{i\}}, Q_{[d-1] \setminus \{i\}})) \\ &\quad + 2 \cdot N(\Gamma(P_{[d-1] \setminus \{j_1, j_2\}}, Q_{[d-1] \setminus \{j_1, j_2\}})). \end{aligned}$$

If  $d$  is even, then

$$N(\Gamma(P, Q)) \leq \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d-1}{2}} + 2 \cdot 6^{\frac{d-2}{2}} + 2 \cdot \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d-3}{2}} = \left( \frac{7}{18} + \frac{1}{3} + \frac{7}{54} \right) \cdot 6^{\frac{d}{2}} = \frac{23}{27} \cdot 6^{\frac{d}{2}} < 6^{\frac{d}{2}}.$$

If  $d$  is odd, then

$$N(\Gamma(P, Q)) \leq 6^{\frac{d-1}{2}} + 2 \cdot \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d-2}{2}} + 2 \cdot 6^{\frac{d-3}{2}} = \left( \frac{3}{7} + \frac{1}{3} + \frac{1}{7} \right) \cdot \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d}{2}} = \frac{19}{21} \cdot \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d}{2}} < \frac{7\sqrt{6}}{18} \cdot 6^{\frac{d}{2}},$$

as desired.  $\square$

#### 4. PROOF OF THEOREM 1.2

In the present section, we prove Theorem 1.2. Before that, we will show that, if  $d = 3$ , then  $N(\Gamma(P, Q))$  is less than  $14 \cdot 6^{\frac{3-3}{2}} = 14$ .

**Proposition 4.1.** *Let  $(P, \leq_P)$  and  $(Q, \leq_Q)$  be posets with  $|P| = |Q| = 3$ . Then we have*

$$N(\Gamma(P, Q)) \leq 13.$$

*The equality holds if and only if one of  $P$  and  $Q$  is isomorphic to  $\mathbf{I}_3$  and the other is isomorphic to either  $\mathbf{C}_3$  or  $\mathbf{1} \oplus \mathbf{I}_2$ .*

*Proof.* The set of all posets with three elements is  $\{\mathbf{1} + \mathbf{C}_2, \mathbf{C}_3, \mathbf{I}_3, \mathbf{1} \oplus \mathbf{I}_2, \mathbf{I}_2 \oplus \mathbf{1}\}$ . Since  $\mathbf{1} \oplus \mathbf{I}_2$  and  $\mathbf{I}_2 \oplus \mathbf{1}$  have the same comparability graph, we may assume that  $P$  and  $Q$  belong to  $\{\mathbf{1} + \mathbf{C}_2, \mathbf{C}_3, \mathbf{I}_3, \mathbf{1} \oplus \mathbf{I}_2\}$ .

**Case 1.** ( $Q = Q_{\{1\}} + Q_{\{2,3\}} \cong \mathbf{1} + \mathbf{C}_2$ .) If  $P$  is isomorphic to either  $\mathbf{1} + \mathbf{C}_2$  or  $\mathbf{1} \oplus \mathbf{I}_2$ , at least one of  $\deg_{G_P} p_2$  or  $\deg_{G_P} p_3$  is one. We may assume that  $\deg_{G_P} p_3 = 1$ . Since  $\deg_{G_Q} q_3 = 1$ , from (3.4) in Proof of Lemma 3.3, we have  $N(\Gamma(P, Q)) \leq 6 \cdot N(\Gamma(\mathbf{1}, \mathbf{1})) = 12$ . Suppose that  $P$  is isomorphic to  $\mathbf{I}_3$ . Then  $\deg_{G_P} p_1 = \deg_{G_Q} q_1 = 0$ . From Lemma 3.2, we have  $N(\Gamma(P, Q)) \leq N(\Gamma(P_{\{2,3\}}, Q_{\{2,3\}})) + c(P_{\{2,3\}}) + c(Q_{\{2,3\}}) = 5 + 3 + 4 = 12$ . If  $P$  is isomorphic to  $\mathbf{C}_3$ , then  $N(\Gamma(P, Q)) = 11$  by Corollary 2.10.

**Case 2.** ( $P, Q \not\cong \mathbf{1} + \mathbf{C}_2$ .)

From Proposition 2.5,

$$N(\Gamma(\mathbf{C}_3, \mathbf{C}_3)) = 8, N(\Gamma(\mathbf{I}_3, \mathbf{I}_3)) = 12, N(\Gamma(\mathbf{I}_3, \mathbf{C}_3)) = 13.$$

Thus we may assume that  $P \not\cong \mathbf{C}_3, \mathbf{I}_3$ . Then  $P$  is isomorphic to  $\mathbf{1} \oplus \mathbf{I}_2$ . If  $Q$  is isomorphic to  $\mathbf{1} \oplus \mathbf{I}_2$ , then  $N(\Gamma(P, Q)) = 11$  or  $12$  from Example 2.3. If  $Q \cong \mathbf{C}_3 (= \mathbf{1} \oplus \mathbf{C}_2)$ , then

$$N(\Gamma(P, Q)) = N(\Gamma(\mathbf{1}, \mathbf{1})) \cdot N(\Gamma(\mathbf{I}_2, \mathbf{C}_2)) = 2 \cdot 5 = 10$$

from Proposition 2.4. Suppose that  $Q \cong \mathbf{I}_3$ . Let  $P = \{p_1, p_2, p_3\}$  where  $p_1$  and  $p_2$  are not comparable. Then  $\bigcup_{W \subset [3]} \mathcal{M}(\Delta_W(P, Q))$  consists of

$$\begin{aligned} & \{q_1\}, \{q_2\}, \{q_3\}, \{p_1, q_2\}, \{p_1, q_3\}, \{p_2, q_1\}, \{p_2, q_3\}, \{p_3, q_1\}, \{p_3, q_2\}, \\ & \{p_1, p_3\}, \{p_1, p_3, q_2\}, \{p_2, p_3\}, \{p_2, p_3, q_1\}. \end{aligned}$$

Hence  $N(\Gamma(P, Q)) = 13$ . (Table 5 shows the values of  $N(\Gamma(P, Q))$  in Proof of Proposition 4.1.)  $\square$

$Q \backslash P$	$\mathbf{1} + \mathbf{C}_2$	$\mathbf{C}_3$	$\mathbf{I}_3$	$\mathbf{1} \oplus \mathbf{I}_2$
$\mathbf{1} + \mathbf{C}_2$	$\leq 12$	11	$\leq 12$	$\leq 12$
$\mathbf{C}_3$		8	13	10
$\mathbf{I}_3$			12	13
$\mathbf{1} \oplus \mathbf{I}_2$				11 or 12

TABLE 5. The values of  $N(\Gamma(P, Q))$  for posets with  $|P| = |Q| = 3$

We note here that the twinned chain polytope and the symmetric edge polytope are distinctly different classes of lattice polytopes.

**Remark 4.2.** Let  $G$  be a finite simple graph on the vertex set  $[d + 1]$  with the edge set  $E(G)$ . The *symmetric edge polytope*  $\mathcal{P}_G$  of  $G$  is the convex hull of  $\{\pm(\mathbf{e}_i - \mathbf{e}_j) : \{i, j\} \in E(G)\}$ , where  $\mathbf{e}_i$  is the  $i$ -th unit coordinate vector in  $\mathbb{R}^{d+1}$ . It is known that the symmetric edge polytope of a connected graph with  $d + 1$  vertices is a centrally symmetric reflexive  $d$ -dimensional polytope.

- The number of vertices of  $\Gamma(\mathbf{I}_3, \mathbf{I}_3)$  is 14 by [3, Proposition 3.3]. On the other hand, if  $d = 3$ , then the number of vertices of  $\mathcal{P}_G$  is less than or equal to  $2 \cdot \binom{4}{2} = 12$  and the equality holds if and only if  $G$  is the complete graph  $K_4$ . Thus,  $\Gamma(\mathbf{I}_3, \mathbf{I}_3)$  is not isomorphic to  $\mathcal{P}_G$  for any graph  $G$ .
- From [6, Proposition 3.5], we have  $N(\mathcal{P}_{K_4}) = 14$ . On the other hand, if  $d = 3$ , then  $N(\Gamma(P, Q)) \leq 13$  by Proposition 4.1. Hence,  $\mathcal{P}_{K_4}$  is not isomorphic to  $\Gamma(P, Q)$  for any posets  $P$  and  $Q$ .

We now prove the main theorem of this paper.

*Proof of Theorem 1.2.* By Corollary 2.6 and Proposition 4.1, the assertion holds if  $d \leq 3$ . The proof proceeds by induction. Let  $d \geq 4$ . From Proposition 2.9, we may assume that there exists  $k \in [d]$  such that  $p_k$  is incomparable with exactly  $s$  elements in  $P$  and  $q_k$  is incomparable with exactly  $t$  elements in  $Q$  with  $st \neq 0$ . Then one of the following holds:

- (i)  $s + t \geq 4$ ,
- (ii)  $(s, t) = (1, 1)$ ,
- (iii)  $(s, t) \in \{(1, 2), (2, 1)\}$ .

If either (i) or (iii) holds, then

$$N(\Gamma(P, Q)) \begin{cases} < 6^{\frac{d}{2}} & \text{if } d \text{ is even,} \\ \leq 14 \cdot 6^{\frac{d-3}{2}} & \text{if } d \text{ is odd.} \end{cases}$$

from Lemmas 3.2 and 3.4, respectively. Suppose that (ii) holds. From Lemma 3.3,  $N(\Gamma(P, Q))$  satisfies (1.1). Moreover, for even  $d$ , the equality  $N(\Gamma(P, Q)) = 6^{\frac{d}{2}}$  holds if and only if

- $N(\Gamma(P_{[d] \setminus \{i, k\}}, Q_{[d] \setminus \{i, k\}})) = 6^{\frac{d-2}{2}},$
- $P = \{p_i, p_k\} \oplus P_{[d] \setminus \{i, k\}},$
- $Q = \{q_i, q_k\} \oplus Q_{[d] \setminus \{i, k\}},$

where  $p_k$  (resp.  $q_k$ ) is incomparable with  $p_i$  (resp.  $q_i$ ). From the hypothesis of induction, we have  $P = \{p_i, p_k\} \oplus P_{[d] \setminus \{i, k\}} \cong \{p_i, p_k\} \oplus \mathbf{I}_2 \oplus \cdots \oplus \mathbf{I}_2$ ,  $Q = \{q_i, q_k\} \oplus Q_{[d] \setminus \{i, k\}} \cong \{q_i, q_k\} \oplus \mathbf{I}_2 \oplus \cdots \oplus \mathbf{I}_2$ , and the map  $\varphi : P \rightarrow Q$  defined by  $\varphi(p_i) = q_i$  induces a graph isomorphism from  $G_P$  to  $G_Q$ . (See Remark 2.2 for the statement about the map from the poset  $P$  to  $Q$ .)  $\square$

**Acknowledgment.** This work was supported by JSPS KAKENHI 24K00534.

## REFERENCES

- [1] V. Batyrev, Dual polyhedra and mirror symmetry for Calabi-Yau hypersurfaces in toric varieties, *J. Algebr. Geom.* **3** (1994), 493–535.
- [2] B. Braun and K. Bruegge, Facets of symmetric edge polytopes for graphs with few edges, *J. Integer Seq.* **26** (2023), Article 23.7.2.
- [3] T. Chappell, T. Friedl and R. Sanyal, Two double poset polytopes, *SIAM J. Discrete Math.* **31** (2017), 2378–2413.
- [4] R. Freij, M. Henze, M.W. Schmitt and G.M. Ziegler, Face numbers of centrally symmetric polytopes from split graphs, *Electron. J. Combin.* **20** (2013), 1–13.
- [5] A. B. Hansen, On a certain class of polytopes associated with independence systems, *Math. Scand.* **41** (1977), 225–241.
- [6] A. Higashitani, K. Jochemko and M. Michałek, Arithmetic aspects of symmetric edge polytopes, *Mathematika* **65** (2019), 763–784.
- [7] C. Haase, F. Kohl and A. Tsuchiya, Levelness of order polytopes, *SIAM J. Discrete Math.* **34**(2) (2020), 1261–1280.
- [8] T. Hibi and K. Matsuda, Quadratic Gröbner bases of twinned order polytopes, *Eur. J. Comb.* **54** (2016), 187–192.
- [9] T. Hibi, K. Matsuda and A. Tsuchiya, Gorenstein Fano polytopes arising from order polytopes and chain polytopes, *J. Commut. Algebra*, to appear.
- [10] T. Hibi, K. Matsuda and A. Tsuchiya, Quadratic Gröbner bases arising from partially ordered sets, *Math. Scand.* **121** (2017), 19–25.
- [11] T. Hibi and A. Tsuchiya, Facets and volume of Gorenstein Fano polytopes, *Math. Nachr.* **290**(2017), 2619–2628.
- [12] M. Henk, J. Richter-Gebert and G. M. Ziegler, Basic properties of convex polytopes, In *Handbook of discrete and computational geometry*, CRC Press Ser. Discrete Math. Appl., pages 243–270. CRC, Boca Raton, FL, 1997.
- [13] G. Kalai, The number of faces of centrally-symmetric polytopes, *Graphs Combin.* **5**(1) (1989), 389–391.
- [14] M. Kreuzer and H. Skarke, Classification of reflexive polyhedra in three-dimensions, *Adv. Theor. Math. Phys.* **2** (1998), 853–871.
- [15] M. Kreuzer and H. Skarke, Complete classification of reflexive polyhedra in four dimensions, *Adv. Theor. Math. Phys.* **4** (2000), 1209–1230.
- [16] M. Kreuzer and H. Skarke, Calabi-Yau data, Webpage, <http://hep.itp.tuwien.ac.at/~kreuzer/CY/>.
- [17] J. C. Lagarias and G. M. Ziegler, Bounds for lattice polytopes containing a fixed number of interior points in a sublattice, *Canad. J. Math.* **43** (1991), 1022–1035.
- [18] T. Matsui, A. Higashitani, Y. Nagazawa, H. Ohsugi and T. Hibi, Roots of Ehrhart polynomials arising from graphs, *J. Algebraic Combin.* **34** (2011), 721–749.
- [19] J. W. Moon and L. Moser, On cliques in graphs, *Israel J. Math.* **3** (1965), 23–28.



- [20] A. Mori, K. Mori and H. Ohsugi, Number of facets of symmetric edge polytopes arising from join graphs, *Eur. J. Comb.* **127** (2025), 104165.
- [21] B. Nill, Gorenstein toric Fano varieties, *Manuscripta Math.* **116** (2005) 183–210.
- [22] B. Nill, Classification of pseudo-symmetric simplicial reflexive polytopes, in “Algebraic and Geometric Combinatorics” (C. A. Athanasiadis et al., eds.), *Contemp. Math.* **423**, AMS, 2006, pp. 269–282.
- [23] OEIS Foundation Inc. “The On-Line Encyclopedia of Integer Sequences” at <https://oeis.org>.
- [24] H. Ohsugi and T. Hibi, Centrally symmetric configurations of integer matrices, *Nagoya Math. J.* **216** (2014), 153–170.
- [25] H. Ohsugi and T. Hibi, Reverse lexicographic squarefree initial ideals and Gorenstein Fano polytopes, *J. Commut. Algebra* **10**(2)(2018), 171–186.
- [26] H. Ohsugi and A. Tsuchiya, The  $h^*$ -polynomials of locally anti-blocking lattice polytopes and their  $\gamma$ -positivity, *Discrete Comput. Geom.* **66** (2021), 701–722.
- [27] R. P. Stanley, Two poset polytopes, *Discrete Comput. Geom.* **1** (1986), 9–23.
- [28] A. Tsuchiya, Volume, facets and dual polytopes of twinned chain polytopes, *Ann. Comb.* **22** (2018), 875–884.

AKI MORI, LEARNING CENTER, INSTITUTE FOR GENERAL EDUCATION, SETSUNAN UNIVERSITY, NEYAGAWA, OSAKA, 572-8508, JAPAN

*Email address:* aki.mori@setsunan.ac.jp

KENTA MORI, DEPARTMENT OF MATHEMATICAL SCIENCES, SCHOOL OF SCIENCE, KWANSEI GAKUIN UNIVERSITY, SANDA, HYOGO 669-1330, JAPAN

*Email address:* k-mori@kwansei.ac.jp

HIDEFUMI OHSUGI, DEPARTMENT OF MATHEMATICAL SCIENCES, SCHOOL OF SCIENCE, KWANSEI GAKUIN UNIVERSITY, SANDA, HYOGO 669-1330, JAPAN

*Email address:* ohsugi@kwansei.ac.jp