

# On the classification of $S^3$ -bundles over $\mathbb{C}P^2$

Liu wancheng

December 2, 2025

## Abstract

This paper presents a classification of the total spaces of  $S^3$ -bundles over  $\mathbb{C}P^2$  up to orientation-preserving homotopy equivalence. Our approach proceeds in two steps: we first derive the PL-homeomorphism classification for these manifolds by computing their Kreck-Stolz invariants. Then, building upon this PL classification result and through an application of surgery theory, we establish the homotopy equivalence classification.

## 1 Introduction

Examples of nondiffeomorphic homeomorphic homogeneous spaces were first discovered by Kreck and Stolz in [1]. These examples belong to a more general class of manifolds called manifolds of type  $r$ . A closed 1-connected smooth 7-dimensional manifold  $M$  is called a manifold of type  $r$  if it satisfies the following properties:

1.  $H^0(M; \mathbb{Z}) \cong H^2(M; \mathbb{Z}) \cong H^5(M; \mathbb{Z}) \cong H^7(M; \mathbb{Z}) \cong \mathbb{Z}$
2.  $H^1(M; \mathbb{Z}) \cong H^3(M; \mathbb{Z}) \cong H^6(M; \mathbb{Z}) = 0$
3.  $H^4(M; \mathbb{Z}) \cong \mathbb{Z}_r$
4.  $H^4(M; \mathbb{Z})$  is generated by  $u^2$ , where  $u$  is a generator of  $H^2(M; \mathbb{Z})$ .

In [2][3] Kreck and Stolz introduced invariants  $s_i(\bar{s}_i)$  that provide a complete smooth (topological) classification of manifolds of type  $r$ . However, the classification of manifolds of type  $r$  up to homotopy equivalence remains unsolved. Partial results can be found in Kruggel's paper [4].

The total spaces of  $S^3$ -bundles over  $\mathbb{C}P^2$  with Euler class  $e = \pm r$  are typical examples of manifolds of type  $r$ . Escher [5] obtained a classification of these manifolds up to diffeomorphism or homeomorphism using Kreck and Stolz's invariants. A partial homotopy classification based on Kruggel's analysis was also presented in the same paper. In this paper, we give a complete classification of these manifolds up to homotopy equivalence (Theorem 1.2). The proof is based on the PL-homeomorphism classification and does not depend on Kruggel's earlier work.

*Convention.* All manifolds under consideration are oriented, and all diffeomorphisms, homeomorphisms, PL-homeomorphisms and homotopy equivalences are orientation-preserving.

### 1.1 Rank 4 real vector bundles over $\mathbb{C}P^2$

In this subsection we give a brief description of 4-dimensional oriented real vector bundles over  $\mathbb{C}P^2$  and fix the notations.

Applying the functor  $[-, BSO(4)]$  to the cofiber sequence

$$S^3 \xrightarrow{h} S^2 \longrightarrow \mathbb{C}P^2 \xrightarrow{q} S^4 \quad (1)$$

associated to the Hopf map  $h: S^3 \rightarrow S^2$ , and notice that  $\pi_3(BSO(4)) = 0$ , one has a short sequence

$$0 \rightarrow \pi_4(BSO(4)) \rightarrow [\mathbb{C}P^2, BSO(4)] \rightarrow \pi_2(BSO(4)) \rightarrow 0. \quad (2)$$

We choose orientation cohomology class  $w_{\mathbb{C}P^2}$  of  $\mathbb{C}P^2$ , and  $w_{S^4}$  of  $S^4$ , respectively, such that  $q^*w_{S^4} = w_{\mathbb{C}P^2} = x^2$ , where  $x$  is a generator of  $H^2(\mathbb{C}P^2; \mathbb{Z})$ .

Let  $e$  and  $p_1$  denote the Euler class and the first Pontrjagin class of a vector bundle, respectively. We choose generators  $\alpha$  and  $\beta$  of the group  $\pi_4(BSO(4))$ , which is known to be isomorphic to  $\mathbb{Z} \oplus \mathbb{Z}$ , such that the corresponding vector bundles over  $S^4$  satisfy the following properties:

	$\alpha$	$\beta$
$e$	0	$w_{S^4}$
$p_1$	$4w_{S^4}$	$-2w_{S^4}$

Under the isomorphisms  $\pi_2(BSO(4)) \cong \mathbb{Z}_2 \cong H^2(\mathbb{C}P^2; \mathbb{Z}_2)$ , the homomorphism  $[\mathbb{C}P^2, BSO(4)] \rightarrow \pi_2(BSO(4))$  is identified with the homomorphism by taking the second Stiefel-Whitney class  $w_2$  of a vector bundle.

Therefore, from the exact sequence (2), one may deduce that 4-dimensional oriented real bundles over  $\mathbb{C}P^2$  can be obtained by the following two pullback diagrams and parametrized by a pair of integers  $(k, l)$  in each case.

$$\begin{array}{ccc} \xi_{k,l} & \longrightarrow & k\alpha + l\beta \\ \downarrow & & \downarrow \\ \mathbb{C}P^2 & \xrightarrow{q} & S^4 \end{array}$$

Spin cases

$$\begin{array}{ccc} \xi'_{k,l} & \longrightarrow & \gamma \oplus \varepsilon^2 \vee k\alpha + l\beta \\ \downarrow & & \downarrow \\ \mathbb{C}P^2 & \xrightarrow{p} & \mathbb{C}P^2 \vee S^4 \end{array}$$

Nonspin cases

Here  $\gamma$  is the tautological line bundle over  $\mathbb{C}P^2$ ,  $p$  is the map collapsing the boundary of a closed disc embedded in the interior of the 4-cell of  $\mathbb{C}P^2$  to a point. An easy calculation shows

$$\begin{aligned} p_1(\xi_{k,l}) &= (4k - 2l)w_{\mathbb{C}P^2}, \quad e(\xi_{k,l}) = lw_{\mathbb{C}P^2}, \\ p_1(\xi'_{k,l}) &= (4k - 2l + 1)w_{\mathbb{C}P^2}, \quad e(\xi'_{k,l}) = lw_{\mathbb{C}P^2}. \end{aligned} \quad (3)$$

## 1.2 Classification Results

Let  $M_{k,l}$  and  $M'_{k,l}$  denote the total spaces of the sphere bundles associated with  $\xi_{k,l}$  and  $\xi'_{k,l}$ , respectively. Noticing that  $\mathbb{C}P^2$  is nonspin, a straightforward calculation shows that  $M_{k,l}$  is nonspin and  $M'_{k,l}$  is spin. We are interested in the classification of these natural geometric objects up to diffeomorphism, homeomorphism or homotopy equivalence.

When  $l \neq 0$ , diffeomorphism and homeomorphism classifications are obtained by applying Kreck-Stolz invariants [2]. Similar methods will produce a PL-homeomorphism classification (see §2). We sum up the results in the following theorem.

**Theorem 1.1.**

1.  $M_{k,l}$  is diffeomorphic to  $M_{k',l}$  if and only if  $k$  and  $k'$  satisfy the following:
  - (a)  $k \equiv k' \pmod{6l}$ ,
  - (b)  $(k - k')(3(k + k' - l) + 1) \equiv 0 \pmod{168l}$ .
2.  $M'_{k,l}$  is diffeomorphic to  $M'_{k',l}$  if and only if  $k$  and  $k'$  satisfy the following:
  - (a)  $k \equiv k' \pmod{12l}$ ,
  - (b)  $(k - k')(k + k' - l + 2) \equiv 0 \pmod{56l}$ .
3.  $M_{k,l}$  is homeomorphic to  $M_{k',l}$  if and only if  $k \equiv k' \pmod{6l}$ .
4.  $M'_{k,l}$  is homeomorphic to  $M'_{k',l}$  if and only if  $k \equiv k' \pmod{12l}$ .
5.  $M_{k,l}$  is PL-homeomorphic to  $M_{k',l}$  if and only if  $k \equiv k' \pmod{6l}$ ,
6.  $M'_{k,l}$  is PL-homeomorphic to  $M'_{k',l}$  if and only if  $k \equiv k' \pmod{12l}$ .

Escher [5] calculated the Kreck-Stolz's invariants of these manifolds and obtained the diffeomorphism and homeomorphism classifications.

The main result of this paper is a homotopy classification of the total spaces of  $S^3$ -bundles over  $\mathbb{C}P^2$  with non-vanishing Euler classes.

**Theorem 1.2.**  $M_{k,l}$  is homotopy equivalent to  $M_{k',l}$  if and only if  $k \equiv k' \pmod{6}$ . The same holds when  $M_{k,l}$  and  $M_{k',l}$  are replaced by  $M'_{k,l}$  and  $M'_{k',l}$ , respectively.

From the exact sequence associated to the fiber bundle  $S^3 \rightarrow M'_{k,l} \rightarrow \mathbb{C}P^2$

$$\cdots \rightarrow \pi_4(S^3) \rightarrow \pi_4(M'_{k,l}) \rightarrow \pi_4(\mathbb{C}P^2) \rightarrow \cdots$$

one sees that  $\pi_4(M'_{k,l})$  is either isomorphic to  $\mathbb{Z}_2$  or vanishes. It is shown in [5] that  $\pi_4(M'_{k,l})$  is isomorphic to  $\mathbb{Z}_2$  when  $l$  is even, and conjectured that  $\pi_4(M'_{k,l})$  vanishes when  $l$  is odd, as suggested by calculations for specific examples. As a corollary of Theorem 1.2 we confirmed this conjecture.

**Corollary 1.3.** The homotopy group  $\pi_4(M'_{k,l})$  vanishes when  $l$  is odd.

We will prove Theorem 1.2 in Section 3 using the PL-homeomorphism classification in Theorem 1.1 and surgery theory.

When  $l = 0$ , the cohomology rings of both  $M_{k,l}$  and  $M'_{k,l}$  are isomorphic to  $H^*(\mathbb{C}P^2 \times S^3)$ . The classification results are obtained in [6]. For completeness we present the results here.

**Theorem 1.4** (Theorem 3.1 of [6]).

- 1 *Two manifolds  $M_{k,0}$  and  $M_{k',0}$  are homeomorphic, PL-homeomorphic or diffeomorphic if and only if  $k = k'$ ;*
- 2 *Two manifolds  $M_{k,0}$  and  $M_{k',0}$  are homotopy equivalent if and only if  $k \equiv k' \pmod{6}$ .*

*These two statements also holds when  $M_{k,0}$  and  $M_{k',0}$  are replaced by  $M'_{k,0}$  and  $M'_{k',0}$ , respectively.*

## 2 Kreck-Stolz Invariants and PL Classification

In the remaining part of this paper we always assume that  $l$  is nonzero. The smooth and homeomorphism classifications of the manifolds of type  $l$  are obtained by Kreck-Stolz in [2]. In this section we give a summary of the invariants which they used to achieved the classification, and show that after obvious modifications, the same approach also applies to the PL classification problem.

Suppose  $(M, u)$  is a manifold of type  $l$ , the invariants are defined as certain characteristic numbers of a pair  $(W, z)$  whose boundary is  $(M, u)$ . That is, we have the following:

1.  $W$  is a smooth 8-manifold with  $\partial W = M$ ;
2.  $z \in H^2(W, \mathbb{Z})$  such that  $z|_M = u$ ;
3.  $W$  is spin (nonspin) if  $M$  is spin (nonspin);
4. In the nonspin case,  $w_2(W) \equiv z \pmod{2}$ .

Notice that  $j^* : H^4(W, M; \mathbb{Q}) \rightarrow H^4(W; \mathbb{Q})$  is an isomorphism. Therefore the cohomology classes  $p_1(W)^2$ ,  $p_1(W)z^2$  and  $z^4$  can be regarded as elements in  $H^8(W, M; \mathbb{Q})$ , and we abbreviate  $p_1^2$ ,  $p_1z^2$  and  $z^4$  for the evaluations  $\langle p_1(W)^2, [W, M] \rangle$ ,  $\langle p_1(W)z^2, [W, M] \rangle$  and  $\langle z^4, [W, M] \rangle$ , respectively. The charc-

teristic numbers  $S_i(W, z) \in \mathbb{Q}$  are defined as follows:

Spin case:

$$\begin{aligned} S_1(W, z) &= -\frac{1}{2^5 \cdot 7} \text{sign}(W) + \frac{1}{2^7 \cdot 7} p_1^2, \\ S_2(W, z) &= -\frac{1}{2^4 \cdot 3} z^2 p_1 + \frac{1}{2^3 \cdot 3} z^4, \\ S_3(W, z) &= -\frac{1}{2^2 \cdot 3} z^2 p_1 + \frac{2}{3} z^4 \end{aligned}$$

Nonspin case:

$$\begin{aligned} S_1(W, z) &= -\frac{1}{2^5 \cdot 7} \text{sign}(W) + \frac{1}{2^7 \cdot 7} p_1^2 - \frac{1}{2^6 \cdot 3} z^2 p_1 + \frac{1}{2^7 \cdot 3} z^4 \\ S_2(W, z) &= -\frac{1}{2^3 \cdot 3} z^2 p_1 + \frac{5}{2^3 \cdot 3} z^4 \\ S_3(W, z) &= -\frac{1}{2^3} z^2 p_1 + \frac{13}{2^3} z^4 \end{aligned}$$

Define  $\bar{S}_i(W, z)$  as follows:  $\bar{S}_1(W, z) = 28S_1(W, z)$ ,  $\bar{S}_2(W, z) = S_1(W, z)$ , and  $\bar{S}_3(W, z) = S_3(W, z)$ .

Define  $s_i(M) \in \mathbb{Q}/\mathbb{Z}$  by  $S_i(W, z) \pmod{\mathbb{Z}}$ , and  $\bar{s}_i(M) \in \mathbb{Q}/\mathbb{Z}$  by  $\bar{S}_i(W, z) \pmod{\mathbb{Z}}$ . Kreck and Stolz's classification of manifolds of type  $l$  is as follows.

**Theorem 2.1** (Theorem 1 of [3]). *Let  $M$  and  $M'$  be two smooth manifolds of type  $l$  which are both spin or both nonspin. Then  $M$  is diffeomorphic (hemeomorphic) to  $M'$  if and only if  $s_i(M) = s_i(M')$  (resp.  $\bar{s}_i(M) = \bar{s}_i(M')$ ) for  $i = 1, 2, 3$ .*

By smoothing theory, if a closed oriented 7-dimensional PL manifold  $M$  admits a smoothing, then it admits exactly 28 smoothings up to concordance [7]. Moreover, every concordant class can be represented by  $M \# \Sigma$ , where  $\Sigma$  is an exotic sphere. Therefore, we could know  $M'$  is  $PL$ -homeomorphic to  $M$  if and only if  $M'$  is diffeomorphic to  $M \# \Sigma$  for some exotic sphere  $\Sigma$ . Consider 8-dimensional Milnor manifold  $M^8(E_8)$  which is a smooth parallelizable manifold with signature 8.  $\partial M^8(E_8)$  is a generator of  $\Theta_7$ , say  $\Sigma_1$ . We have  $s_1(\Sigma_1) = 1/28$ ,  $s_2(\Sigma_1) = s_3(\Sigma_1) = 0$ . Therefore, we have the following theorem:

**Theorem 2.2.** *Let  $M$  and  $M'$  be two smooth manifolds of type  $l$  which are both spin or both nonspin. Then  $M$  is  $PL$ -homeomorphic to  $M'$  if and only if  $28s_1(M) = 28s_1(M')$ , and  $s_i(M) = s_i(M')$  for  $i = 2, 3$ .*

*Proof of Theorem 1.1.* Let  $W_{k,l}$  be the disk bundle associated to  $\xi_{k,l}$ , and let  $\pi_{k,l} : W_{k,l} \rightarrow \mathbb{C}P^2$  be the projection. We have  $\tau(W_{k,l}) \cong \pi_{k,l}^*(\tau(\mathbb{C}P^2) \oplus \xi_{k,l})$ . Let  $z = \pi_{k,l}^*x$ , recall that  $x$  is a generator of  $H^2(\mathbb{C}P^2; \mathbb{Z})$ , then

$$\begin{aligned} p_1(W_{k,l}) &= \pi_{k,l}^*(p_1(\mathbb{C}P^2) + p_1(\xi_{k,l})) \\ &= (4k - 2l + 3)\pi_{k,l}^*w_{\mathbb{C}P^2} \\ &= (4k - 2l + 3)z^2. \end{aligned}$$

Let  $y$  be a generator of  $H^4(W_{k,l}, M_{k,l}; \mathbb{Z})$  such that  $j^*(y) = lz^2$ . Clearly,  $y$  is the Thom class of  $\xi_{k,l}$ . The orientation of  $W_{k,l}$  is defined by equation  $\langle z^2 \cup y, [W_{k,l}, M_{k,l}] \rangle = 1$ .

Since  $\langle y \cup j^*y, [W_{k,l}, M_{k,l}] \rangle = l$ ,  $\text{sign}(W)$  equals to the sign of  $l$ . By definition, we have

$$\begin{aligned} z^4 &= \langle z^2 \cup j^{*-1}z^2, [W_{k,l}, M_{k,l}] \rangle = \frac{1}{l} \langle z^2 \cup y, [W_{k,l}, M_{k,l}] \rangle = \frac{1}{l}, \\ z^2 p_1 &= \frac{4k - 2l + 3}{l}, \\ p_1^2 &= \frac{(4k - 2l + 3)^2}{l}. \end{aligned}$$

Then  $s_i$  is given by

$$\begin{aligned} s_1 &= -\frac{\text{sign}(l)}{2^5 \cdot 7} + \frac{(4k - 2l + 3)^2}{2^7 \cdot 7l} - \frac{8k - 4l + 5}{2^7 \cdot 3l}, \\ s_2 &= -\frac{2k - l - 1}{12l}, \\ s_3 &= -\frac{2k - l - 5}{4l}. \end{aligned}$$

Similarly, let  $\pi'_{k,l} : W'_{k,l} \rightarrow \mathbb{C}P^2$  be the projection. We have

$$\begin{aligned} z^4 &= \frac{1}{l}, \\ z^2 p_1 &= \frac{4k - 2l + 4}{l}, \\ p_1^2 &= \frac{(4k - 2l + 4)^2}{l}. \end{aligned}$$

And  $s_i$  is given by

$$\begin{aligned} s_1 &= -\frac{\text{sign}(l)}{2^5 \cdot 7} + \frac{(2k - l + 2)^2}{2^5 \cdot 7l}, \\ s_2 &= -\frac{2k - l + 1}{24l}, \\ s_3 &= -\frac{2k - l - 2}{6l}. \end{aligned}$$

Then Theorem 1.1 follows by comparing the invariants.  $\square$

*Proof of Corollary 1.3.* Let  $l$  be an odd integer. If  $\pi_4(M'_{k,l}) = \mathbb{Z}_2$  for some  $k$ , by a theorem of Kruggel [4, Theorem 0.1], the  $s_2$ -invariant of any type  $l$  manifold  $M$  which is homotopy equivalent to  $M'_{k,l}$  satisfies  $l(s_2(M) - s_2(M'_{k,l})) = 0 \in \mathbb{Q}/\mathbb{Z}$ . By Theorem 1.2,  $M'_{k+6,l}$  is homotopy equivalent to  $M_{k,l}$ . But the above calculation shows that  $l(s_2(M'_{k+6,l}) - s_2(M'_{k,l})) = 1/2 \in \mathbb{Q}/\mathbb{Z}$ , a contradiction.  $\square$

### 3 Fiber homotopy equivalences

**Lemma 3.1.** *There exist fiber homotopy equivalences of spherical fibrations  $f_j : M_{k+6j,l} \rightarrow M_{k,l}$  and  $f'_j : M'_{k+6j,l} \rightarrow M'_{k,l}$  for all  $j \in \mathbb{Z}$ .*

*Proof.* Let  $SG(4)$  be the topological monoid of orientation preserving self homotopy equivalences of  $S^3$ ,  $BSG(4)$  the classifying space of  $S^3$ -fibrations. Therefore, the sphere bundles of two vector bundles are fiber homotopy equivalent as  $S^3$ -fibrations if and only if the classifying maps of the vector bundles have the same image under the map  $i_* : [\mathbb{C}P^2, BSO(4)] \rightarrow [\mathbb{C}P^2, BSG(4)]$  induced by the inclusion  $i : SO(4) \rightarrow SG(4)$ .

The cofibration sequence (1) induces the following exact diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & \pi_4(BSO(4)) & \xrightarrow{q^*} & [\mathbb{C}P^2, BSO(4)] & \longrightarrow & \pi_2(BSO(4)) \longrightarrow 0 \\
& & \downarrow i_* & & \downarrow i_* & & \downarrow i_* \\
\pi_3(BSG(4)) & \xrightarrow{(\sum h)^*} & \pi_4(BSG(4)) & \xrightarrow{q^*} & [\mathbb{C}P^2, BSG(4)] & \longrightarrow & \pi_2(BSG(4)) \xrightarrow{h^*} \pi_3(BSG(4))
\end{array}$$

Diagram 1

It's known that  $i_* : \pi_4(BSO(4)) \rightarrow \pi_4(BSG(4))$  can be identified with  $\mathbb{Z}\{\alpha\} \oplus \mathbb{Z}\{\beta\} \rightarrow \mathbb{Z}_{12} \oplus \mathbb{Z}$ ,  $k\alpha + l\beta \mapsto ((k \bmod 12), l)$  [8]. To identify the homomorphism  $(\sum h)^* : \pi_3(BSG(4)) \rightarrow \pi_4(BSG(4))$  we consider the fibration  $SF(3) \rightarrow SG(4) \rightarrow S^3$ , where  $SG(4) \rightarrow S^3$  is the evaluation at the base point, and  $SF(3)$  the subspace of  $SG(4)$  consisting of orientation preserving homotopy equivalences of  $S^3$  which fix the base point. The Hopf map  $h : S^3 \rightarrow S^2$  induces the following diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & \pi_2(SF(3)) & \xrightarrow{\cong} & \pi_2(SG(4)) & \longrightarrow & \pi_2(S^3) \longrightarrow 0 \\
& & \downarrow h^* & & \downarrow h^* & & \downarrow h^* \\
0 & \longrightarrow & \pi_3(SF(3)) & \longrightarrow & \pi_3(SG(4)) & \longrightarrow & \pi_3(S^3) \longrightarrow 0
\end{array}$$

Diagram 2

By the usual adjoint correspondence,  $h^* : \pi_2(SF(3)) \rightarrow \pi_3(SF(3))$  can be identified with  $(\Sigma^3 h)^* : \pi_5(S^3) \rightarrow \pi_6(S^3)$ , which is known to be identified with the inclusion  $\mathbb{Z}_2 \hookrightarrow \mathbb{Z}_{12}$  [9]. Therefore,  $(\sum h)^* : \pi_3(BSG(4)) \rightarrow \pi_4(BSG(4))$  can be identified with the inclusion into the first summand  $\mathbb{Z}_2 \hookrightarrow \mathbb{Z}_{12} \oplus \mathbb{Z}$ .

Therefore from commutative diagram 1 one has:

$$i_*(\xi_{k+6j,l}) = q^* i_*((k+6j)\alpha + l\beta) = q^* i_*(k\alpha + l\beta) = i_*(\xi_{k,l}).$$

This shows that  $M_{k+6j,l}$  and  $M_{k,l}$  are fiber homotopy equivalent.

For the nonspin case one needs more detailed analysis since the exact sequences in diagram 1 are not exact sequences of abelian groups. We will prove the statement in two steps.

First we show that the spherical fibrations  $\xi'_{k+6j,l}$  and  $\xi'_{k,l}$  are stably fiber homotopy equivalent. The cofiber sequence (1) induces the following commutative diagram of short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \pi_4(BSO) & \longrightarrow & [\mathbb{C}P^2, BSO] & \longrightarrow & \pi_2(BSO) \longrightarrow 0 \\ & & \downarrow & & \downarrow i_* & & \downarrow \cong \\ \pi_3(BSG) & \xrightarrow{\Sigma \eta^*} & \pi_4(BSG) & \xrightarrow{q^*} & [\mathbb{C}P^2, BSG] & \longrightarrow & \pi_2(BSG) \xrightarrow{0} \pi_3(BSG) \end{array}$$

The following facts are known.

1. Taking the first Pontrjagin class induces an bijective homomorphism

$$p_1: [\mathbb{C}P^2, BSO] \rightarrow \mathbb{Z}.$$

2. There are isomorphisms  $\pi_4 BSG \cong \pi_3^s \cong \mathbb{Z}_{24}$ , and we have the following diagram

$$\begin{array}{ccc} \pi_3(BSG(4)) \cong \mathbb{Z}_2 & \longrightarrow & \pi_4(BSG(4)) \cong \mathbb{Z}_{12} \oplus \mathbb{Z} \\ \downarrow & & \downarrow \\ \pi_3(BSG) \cong \mathbb{Z}_2 & \xleftarrow{(\Sigma \eta)^*} & \pi_4(BSG) \cong \mathbb{Z}_{24} \end{array}$$

where the natural map  $\pi_4(BSG(4)) \rightarrow \pi_4(BSG)$  is the injection on the first summand and zero on the second summand, as  $\Sigma^\infty: \pi_6(S^3) \rightarrow \pi_3^s$  is identified with  $\mathbb{Z}_{12} \hookrightarrow \mathbb{Z}_{24}$  [9].

Therefore by the 5-lemma the homomorphism  $i_*: [\mathbb{C}P^2, BSO] \rightarrow [\mathbb{C}P^2, BSG]$  is surjective. Hence the group  $[\mathbb{C}P^2, BSG]$  is cyclic and isomorphic to  $\mathbb{Z}_{24}$  by counting order. Under stabilization  $BSO(4) \rightarrow BSO \rightarrow BSG$  the classifying maps for the vector bundles  $\xi'_{k,l}$  and  $\xi'_{k+6j,l}$  are homotopic, since  $p_1(\xi'_{k+6j,l}) \equiv p_1(\xi'_{k,l}) \pmod{24}$  (see formula (3)). Therefore there exists a homotopy

$$f_t: \mathbb{C}P^2 \times [0, 1] \rightarrow BSG$$

between the classifying maps for  $i_* j_*(\xi'_{k+6j,l})$  and  $i_* j_*(\xi'_{k,l})$ .

In the second step we find by induction on skeleta a lift of  $f_t$  to  $BSG(4)$

$$g_t: \mathbb{C}P^2 \times I \rightarrow BSG(4),$$

with  $g_0 = i_* \xi'_{k+6j,l}$  and  $g_1 = i_* \xi'_{k,l}$ . This shows that  $\xi'_{k+6j,l}$  and  $\xi'_{k,l}$  are fiber homotopy equivalent.

Let  $SG/SG(4)$  be the homotopy fibre of  $BSG(4) \rightarrow BSG$ , then a basic calculation shows that  $\pi_2 SG/SG(4) = 0$  and  $\pi_4 SG/SG(4) \cong \mathbb{Z}$ . By elementary obstruction theory there is a lift  $g_t^{(2)}$  of  $f_t$  on the 2-skeleton of  $\mathbb{C}P^4$

$$g_t^{(2)}: S^2 \times I \rightarrow BSG(4).$$



Let  $X = \mathbb{C}P^2 \times \partial I \cup S^2 \times I$ , then  $\mathbb{C}P^2 \times I = X \cup_m D^5$ , where we regard  $S^4 = D^4 \times \partial I \cup S^3 \times I$  and  $m: S^4 \rightarrow X$  has the form

$$\iota \times \text{id} \cup h \times \text{id}: D^4 \times \partial I \cup S^3 \times I \rightarrow \mathbb{C}P^2 \times I$$

where  $\iota: D^4 \rightarrow \mathbb{C}P^2$  is the inclusion of the top cell and  $h$  the Hopf map. Then the homotopy  $g_t^{(2)}$  extends to the desired homotopy  $g_t: \mathbb{C}P^2 \times I \rightarrow BSG(4)$  if and only if the composite

$$(g_0 \cup g_1 \cup g_t^{(2)}) \circ m: S^4 \rightarrow X = \mathbb{C}P^2 \times \partial I \cup S^2 \times I \rightarrow BSG(4)$$

is null-homotopic. By standard algebraic topology one shows that  $\pi_4(X)$  is isomorphic to  $\mathbb{Z}$ , generated by  $[m]$ . Denote  $g_0 \cup g_1 \cup g_t^{(2)}$  by  $\phi$ , it suffices to show that  $\phi_*: \pi_4(X) \rightarrow \pi_4(BSG(4))$  is trivial.

By construction, the composite  $j_* \circ \phi_*: \pi_4(X) \rightarrow \pi_4(BSG(4)) \rightarrow \pi_4(BSG)$  is trivial. On the other hand, the homomorphism  $j_*: \pi_4(BSG(4)) \rightarrow \pi_4(BSG)$  can be identified with  $\mathbb{Z}_{12} \oplus \mathbb{Z} \rightarrow \mathbb{Z}_{24}$ , where on the first summand it is the inclusion. Therefore  $\phi_*([m])$  is not in the  $\mathbb{Z}_{12}$ -summand. In the sequel we will show it is zero by showing that it is zero in  $\pi_4(BSG(4)) \otimes \mathbb{Q}$ .

Let  $X \rightarrow S^4 \vee X \vee S^4$  be the map which pinches a 4-sphere from each top cell  $D^4$  of  $\mathbb{C}P^2$ , and  $S^4 \rightarrow S^4 \vee S^4 \vee S^4$  be the map which pinches a 4-sphere from each  $D^4$  in  $S^4 = D^4 \times \partial I \cup S^3 \times I$ . Then the map  $\phi \circ m: X \rightarrow BSG(4)$  has a factorization shown in the following commutative diagram

$$\begin{array}{ccccc} S^4 & \xrightarrow{m} & X & \xrightarrow{\phi} & BSG(4) \\ \downarrow & & \downarrow & \nearrow f_0 \vee f \vee f_1 & \\ S^4 \vee S^4 \vee S^4 & \xrightarrow{\text{id} \vee m \vee \text{id}} & S^4 \vee X \vee S^4 & & \end{array}$$

where  $f_0 = i((k+6j)\alpha + l\beta)$ ,  $f_1 = i(k\alpha + l\beta)$ ,  $f = i(\eta \oplus \varepsilon^2) \times \partial I \cup g_t^{(2)}$ .

First recall that we have shown that  $[f_0] = [f_1] \in \pi_4(BSG(4)) \otimes \mathbb{Q}$ .

The only non-trivial rational homotopy group of  $BSG(4)$  is  $\pi_4(BSG(4)) \otimes \mathbb{Q} \cong \mathbb{Q}$ . There is a rational homotopy equivalence  $l: BSG(4) \rightarrow K(\mathbb{Q}, 4)$  [10]. We claim that the composite  $l \circ f: X \rightarrow BSG(4) \rightarrow K(\mathbb{Q}, 4)$  is null homotopic. This will finish the proof for the nonspin case. To prove the claim, first note that the inclusion  $\mathbb{C}P^2 \cup \mathbb{C}P^2 \subset X$  induces an isomorphism  $H^4(X; \mathbb{Q}) \rightarrow H^4(\mathbb{C}P^2; \mathbb{Q}) \oplus H^4(\mathbb{C}P^2; \mathbb{Q})$ . Therefore from the one-one correspondence between  $H^4(-; \mathbb{Q})$  and  $[-, K(\mathbb{Q}, 4)]$  it suffices to show that the map  $l \circ i \circ (\eta \oplus \varepsilon^2): \mathbb{C}P^2 \rightarrow BSG(4) \rightarrow K(\mathbb{Q}, 4)$  is null-homotopic. Notice that the map  $\eta \oplus \varepsilon^2$  factors through  $BSO(3)$ . Since the composition  $\pi_3(SO(3)) \rightarrow \pi_3(SO(4)) \rightarrow \pi_3(SG(4))$  is trivial after tensoring with  $\mathbb{Q}$ , the map  $BSO(3) \rightarrow BSO(4) \rightarrow BSG(4) \rightarrow K(\mathbb{Q}, 4)$  is trivial. This shows the claim.  $\square$

## 4 surgery exact sequence and homotopy classification

In this section we prove the homotopy classification (Theorem 1.2) by showing that every element in the PL structure set  $\mathcal{S}^{PL}(M_{k,l})$  is represented by a fiber homotopy equivalence  $f_j : M_{k+6j,l} \rightarrow M_{k,j}$ . The same also holds for the nonspin case. This approach was first used in [8].

For the convenience of the reader, we give a briefly introduction to simply connected surgery exact sequence. We work in the piecewise linear (PL) category. For more information we refer to [11].

Let  $(M, \partial M)$  be an  $n$ -dimensional compact PL manifold with possibly empty boundary. A PL manifold structure on  $(M, \partial M)$  is a triple  $(N, \partial N, g)$ , where  $N$  is a compact PL manifold and  $g : (N, \partial N) \rightarrow (M, \partial M)$  is a homotopy equivalence. Two triples  $(N_0, \partial N_0, g_0)$  and  $(N_1, \partial N_1, g_1)$  are concordant if there is a PL-homeomorphism  $f : (N_0, \partial N_0) \rightarrow (N_1, \partial N_1)$  so that the following diagram commutes up to homotopy.

$$\begin{array}{ccc} (N_0, \partial N_0) & & \\ \downarrow f & \searrow g_0 & \\ & & (M, \partial M) \\ & \nearrow g_1 & \\ (N_1, \partial N_1) & & \end{array}$$

The PL structure set  $\mathcal{S}^{PL}(M)$  is the set of concordance classes of triples  $(N, \partial N, g)$ .

Let  $G/PL$  be the homotopy fiber of  $J : BPL \rightarrow BG$ . Then the set of homotopy classes of maps  $M \rightarrow G/PL$  are in 1-1 correspondence with the set of equivalence classes of pairs  $(\eta, t)$ , where  $\eta$  is a stable PL bundle over  $M$  and  $t$  is a fiber homotopy trivialization of the associated spherical fibration  $J\eta$ .

The map of normal invariants  $\eta : \mathcal{S}^{PL}(M) \rightarrow [M, G/PL]$  is defined by sending a triple  $(N, \partial N, g)$  to  $\eta(g) := (\nu_M - (g^{-1})^* \nu_N, t(g))$ , where  $t(g)$  is a fiber homotopy trivialization of  $J(\nu_M - (g^{-1})^* \nu_N)$  defined in a standard way.

If  $\pi_1(M) \cong \pi_1(\partial M) \cong 0$  and  $n \geq 6$ , we have the following two facts [12]:

1. If  $\partial M \neq \emptyset$ ,  $\mathcal{S}^{PL}(M) \xrightarrow{\eta} [M, G/PL]$  is an isomorphism;
2. If  $\partial M = \emptyset$ , we have an exact sequence of based sets

$$0 \rightarrow \mathcal{S}^{PL}(M) \xrightarrow{\eta} [M, G/PL] \rightarrow L_n(e).$$

$$\text{where } L_n(e) \cong \pi_n(G/PL) \cong \begin{cases} \mathbb{Z}, & \text{if } n \equiv 0 \pmod{4}, \\ \mathbb{Z}_2, & \text{if } n \equiv 2 \pmod{4}, \\ 0, & \text{if } n \equiv 1 \pmod{2}. \end{cases}$$

Let  $\iota : M_{k,l} \hookrightarrow W_{k,l}$  be the canonical inclusion. Then we have the following

commutative diagram, where the normal invariants are isomorphisms by the two facts mentioned above.

$$\begin{array}{ccc} \mathcal{S}^{PL}(W_{k,l}) & \xrightarrow[\cong]{\eta} & [W_{k,l}, G/PL] \\ \downarrow & & \downarrow \\ \mathcal{S}^{PL}(M_{k,l}) & \xrightarrow[\cong]{\eta} & [M_{k,l}, G/PL] \end{array}$$

Diagram 3

We have the same diagram when  $M_{k,l}$  and  $W_{k,l}$  are replaced by  $M'_{k,l}$  and  $W'_{k,l}$ , respectively.

The infinite loop space structure on  $G/PL$  defines a generalized cohomology theory, in which  $[-, G/PL]$  is the 0-th group. Let  $M_{k,l}^0$  denote  $M_{k,l} \setminus \{\text{pt}\}$ . We have  $[M_{k,l}^0, G/PL] \cong \mathcal{S}^{PL}(M_{k,l})$  [12]. Then by diagram 3, we have  $[M_{k,l}, G/PL] \cong [M_{k,l}^0, G/PL]$ . It is easy to see that the Atiyah-Hurzebruch spectral sequence for  $M_{k,l}^0$  collapses at the  $E_2$ -page, then so does  $M_{k,l}$ . It's ready to see that the Atiyah-Hurzebruch spectral sequences for both  $\mathbb{CP}^2$  and  $M_{k,l}$  collapse at the  $E_2$ -page, therefore we have the following commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z} & \longrightarrow & [\mathbb{CP}^2, G/PL] & \longrightarrow & \mathbb{Z}_2 \longrightarrow 0 \\ & & \downarrow & & \downarrow (p_{k,l})^* & & \downarrow \cong \\ 0 & \longrightarrow & \mathbb{Z}_l & \longrightarrow & [M_{k,l}, G/PL] & \longrightarrow & \mathbb{Z}_2 \longrightarrow 0 \end{array}$$

where  $p_{k,l}$  denotes the sphere bundle projection. By the 5-lemma,  $(p_{k,l})^* : [\mathbb{CP}^2, G/PL] \rightarrow [M_{k,l}, G/PL]$  is surjective. It's known that  $[\mathbb{CP}^2, G/PL] \cong \mathbb{Z}$  ([13, Chapter 14C]), therefore  $[M_{k,l}, G/PL] \cong \mathbb{Z}_{2l}$ . Similarly, we have  $[M'_{k,l}, G/PL] \cong \mathbb{Z}_{2l}$ .

In Lemma 3.1 we have shown that there are fiber homotopy equivalences  $f_j : M_{k+6j,l} \rightarrow M_{k,l}$  and  $f'_j : M'_{k+6j,l} \rightarrow M'_{k,l}$ . They represent elements in  $\mathcal{S}^{PL}(M_{k,l})$  and  $\mathcal{S}^{PL}(M'_{k,l})$ , respectively.

**Lemma 4.1.** *Let  $f_j : M_{k+6j,l} \rightarrow M_{k,l}$  be a fiber homotopy equivalence, then  $\eta(f_j) = j \in [M_{k,l}, G/PL] \cong \mathbb{Z}_{2l}$ . The same holds when  $M_{k,l}$  and  $f_j$  are replaced by  $M'_{k,l}$  and  $f'_j$ , respectively.*

*Proof.* For any vector bundle  $\zeta$ , the disc bundle  $D(\zeta)$  is homeomorphic to the mapping cylinder of spherical bundle projection. Therefore, there exists a natural fiber homotopy equivalence  $F_j : W_{k+6j,l} \rightarrow W_{k,l}$  extending  $f_j$ . Consider the following commutative diagram

$$\begin{array}{ccccc}
\mathcal{S}^{PL}(W_{k,l}) & \xrightarrow{\eta} & [W_{k,l}, G/PL] & \xrightarrow{j_*} & [W_{k,l}, BPL] \\
\downarrow i_{k,l}^* & & \downarrow i_{k,l}^* & & \downarrow i_{k,l}^* \\
\mathcal{S}^{PL}(\mathbb{CP}^2) & \xrightarrow{\eta} & [\mathbb{CP}^2, G/PL] & \xrightarrow[\times 24]{j_*} & [\mathbb{CP}^2, BPL]
\end{array}$$

where  $i_{k,l} : \mathbb{CP}^2 \rightarrow W_{k,l}$  is the inclusion of the zero section and  $j_*$  is interpreted by forgetting the trivialization. It's well know that  $[\mathbb{CP}^2, BPL] \cong \mathbb{Z}$  and this isomorphism is determined by  $p_1$ . And  $j_* : [\mathbb{CP}^2, G/PL] \rightarrow [\mathbb{CP}^2, BPL]$  is known to be multiplication by 24 [6]. Thus we obtain

$$\begin{aligned}
i_{k,l}^* j_* \eta(F_j) &= i_{k,l}^* (\nu(W_{k,l}) - F_j^{-1*} \nu(W_{k+6j,l})) \\
&= i_{k,l}^* (\pi_{k,l}^* (-\xi_{k,l} - \tau(\mathbb{CP}^2)) - F_j^{-1*} \pi_{k+6j,l}^* (-\xi_{k+6j,l} - \tau(\mathbb{CP}^2))) \\
&= i_{k,l}^* \pi_{k,l}^* (\xi_{k+6j,l} - \xi_{k,l}) \\
&= \xi_{k+6j,l} - \xi_{k,l} \\
&= 4(k+6j) - 2l - (4k - 2l) \\
&= 24j \in \mathbb{Z}.
\end{aligned}$$

Hence  $\eta(F_j) = j \in [W_{k,l}, G/PL] \cong \mathbb{Z}$ . From the following commutative diagram

$$\begin{array}{ccc}
[W_{k,l}, G/PL] & \xleftarrow[\cong]{\pi_{k,l}^*} & [\mathbb{CP}^2, G/PL] \\
\downarrow \iota^* & \swarrow p_{k,l}^* & \\
[M_{k,l}, G/PL] & & 
\end{array}$$

we conclude that  $\eta(f_j) = \iota^* \eta(F_j) = p_{k,l}^* \pi_{k,l}^{*-1} \eta(F_j) = j \in \mathbb{Z}_{2l}$ .

The proof for  $M'_{k,l}$  is similar.  $\square$

**Proposition 4.2.** *A 7-dimensional closed nonspin (spin) PL-manifold  $M$  is homotopy equivalent to  $M_{k,l}$  ( $M'_{k,l}$ ) if and only if  $M$  is PL-homeomorphism to  $M_{k+6j,l}$  ( $M'_{k+6j,l}$ ) for some  $j \in \mathbb{Z}$ .*

*Proof.* Let  $f : M \rightarrow M_{k,l}$  be a homotopy equivalence. Then by Lemma 4.1, there exists some  $j$  such that  $\eta(f_j) = \eta(f)$ . Since  $\eta : \mathcal{S}^{PL}(M_{k,l}) \rightarrow [M_{k,l}, G/PL]$  is an isomorphism (see diagram 3), one has  $(M_{k+6j,l}, f_j) = (M, f) \in \mathcal{S}^{PL}(M_{k,l})$ . Hence  $M_{k+6j,l}$  and  $M$  are PL-homeomorphic.  $\square$

Theorem 1.2 is a consequence of Proposition 4.2 and the PL classification.

*Proof of Theorem 1.2.* Suppose  $M_{k,l}$  is homotopy equivalent to  $M_{k',l}$ , then by Proposition 4.2 there exists some  $j$  such that  $M_{k,l}$  is PL-homeomorphic to  $M_{k'+6j,l}$ . Then Theorem 1.1 (5) implies  $k \equiv k' + 6j \pmod{6}$ . Hence  $k \equiv k' \pmod{6}$ . The proof is analogous for  $M'_{k,l}$ .  $\square$

**Remark 4.3.** Combining Theorem 1.2 and Lemma 3.1, we obtain that the homotopy classification of the total spaces  $M_{k,l}$  and the homotopy classification of the pairs  $(M_{k,l}, S^3)$  coincide. The statement holds when  $M_{k,l}$  and  $(M_{k,l}, S^3)$  are replaced by  $M'_{k,l}$  and  $(M'_{k,l}, S^3)$ , respectively.

**Acknowledgement.** The author wishes to thank Yang Su for his guidance and many constructive discussions.

## References

- [1] Matthias Kreck and Stephan Stolz. A diffeomorphism classification of 7-dimensional homogeneous einstein manifolds with  $\mathrm{su}(3) \times \mathrm{su}(2) \times \mathrm{u}(1)$ -symmetry. *Annals of Mathematics*, 127(2):373–388, 1988.
- [2] Matthias Kreck and Stephan Stolz. Some nondiffeomorphic homeomorphic homogeneous 7-manifolds with positive sectional curvature. *Journal of Differential Geometry*, 33(2):465 – 486, 1991.
- [3] Matthias Kreck and Stephan Stolz. A correction on: “some nondiffeomorphic homeomorphic homogeneous 7-manifolds with positive sectional curvature”. *Journal of Differential Geometry*, 49(1):203–204, 1998.
- [4] Bernd Kruggel. Kreck-stolz invariants, normal invariants and the homotopy classification of generalised wallach spaces. *Quarterly Journal of Mathematics*, 49(4):469–485, 1998.
- [5] Christine Escher and Wolfgang Ziller. Topology of non-negatively curved manifolds. *Annals of Global Analysis and Geometry*, 46(1):23–55, 2014.
- [6] Xueqi Wang. On the classification of certain 1-connected 7-manifolds and related problems. *The Quarterly Journal of Mathematics*, 73(2):711–727, 12 2021.
- [7] Bjørn Jahren. Smoothing theory and topology of  $\mathrm{pl}/\mathrm{o}$ . *Preprint series: Pure mathematics <http://urn.nb.no/URN:NBN:no-8076>*, 1973.
- [8] Diarmuid Crowley and Christine M. Escher. A classification of  $\mathrm{s}^3$ -bundles over  $\mathrm{s}^4$ . *Differential Geometry and its Applications*, 18(3):363–380, 2003.
- [9] Hiroshi Toda. *Composition Methods in Homotopy Groups of Spheres*. Princeton University Press, Princeton, 1963.
- [10] Dennis Sullivan. *Geometric Topology: Geometric Topology Localization, Periodicity, and Galois Symmetry*. MIT Press, Cambridge, 1970.
- [11] Wolfgang Lück and Tibor Macko. *Surgery Theory: Foundations*. Springer, Cham, 2024.

- [12] Dennis Sullivan. *Triangulating and Smoothing Homotopy Equivalences and Homeomorphisms. Geometric Topology Seminar Notes*, pages 69–103. Springer Netherlands, Dordrecht, 1996.
- [13] Charles Terence Clegg Wall. *Surgery on Compact Manifolds*, volume 69 of *Mathematical Surveys and Monographs*. American Mathematical Society, second edition, 1999.