

Triplet pairing in neutron matter in a comprehensive diagrammatic approach

Panagiota Papakonstantinou^{1,*}, Eckhard Krotscheck^{2,3,**}, and Jiawei Wang^{2,***}

¹Institute for Rare Isotope Science (IRIS), Institute for Basic Science(IBS), Daejeon 34000, Korea

²Department of Physics, SUNY Buffalo, New York 14260, USA

³Institute for Theoretical Physics, Johannes Kepler University, A 4040 Linz, Austria

Abstract. We apply a large-scale summation of Feynman diagrams, including the class of parquet diagrams plus important contributions outside the parquet class, for calculating effective pairing interactions and subsequently the superfluid gap in P -wave pairing in neutron matter. We use realistic nucleon-nucleon interactions of the v_8 type and perform calculations up to a Fermi momentum of 1.8 fm^{-1} . We find that many-body correlations lead to a strong reduction of the spin-orbit interaction, and, therefore, to a radical suppression of the $^3P_2 - ^3F_2$ gap and an enhancement of the 3P_0 gap.

1 Introduction

The superfluid phase transition in neutron matter has been studied microscopically for decades; see, for example, Refs. [1, 2] for related reviews and compilations of the relevant literature. Numerous microscopic calculations predict singlet-state superfluidity consistently at sub-saturation densities. Triplet pairing is generally predicted at higher densities. However, predictions for the magnitude of the triplet-pairing gap vary widely. Whether triplet pairing develops in dense matter is particularly important in astrophysics, because it can affect the cooling of neutron stars in intricate ways, enhancing certain mechanisms and suppressing others.

In this work, we undertake the most comprehensive to-date diagrammatic approach to calculating triplet pairing in dense neutron matter by including not only ring and ladder diagrams but also diagrams beyond the parquet class, which the form of the nuclear interaction necessitates.

Next, we summarize existing microscopic approaches to the nuclear pairing gap, we briefly describe the formalism used in the present work, and we proceed to present our results.

2 Existing approaches

We may classify existing microscopic approaches to calculating the pairing gap in nuclear matter as follows: First, there are calculations of the superfluid gap at the mean-field level using bare, more or less realistic nuclear interactions. The effect of a three-body force has also been explored. Second, there are those that take into account medium polarization effects, which can strongly influence the superfluid transition temperature. Such calculations

necessarily involve assumptions about the quasiparticle interaction based on known physical observables. Third, inclusion of many-body effects in correlated basis function theory for superfluid systems, which can be mapped onto a regular Bardeen–Cooper–Schrieffer (BCS)-like theory with effective interactions. Finally, some Monte Carlo calculations exist for pairing in the singlet channel in low-density neutron matter. The present work falls into the third category. We proceed to sketch our method before we present our results.

3 Correlated wavefunctions, diagrammatic summations, and the nuclear interaction

In correlated basis functions methods, the first step is to select a trial N -body state $|\Phi_0\rangle$ that describes the system's statistics. In the case of Fermionic systems, that could be a Slater determinant of plane waves or other basis functions appropriate for the symmetries of the system. Next, one acts on the trial function with an operator \hat{F}_N that encodes dynamical correlations. Among others, it suppresses the wave function at short interparticle distances thus encoding short-range repulsion. The correlated and normalized wave function is

$$|\Psi\rangle = \frac{\hat{F}_N |\Phi_0\rangle}{\langle \Phi_0 | \hat{F}_N^\dagger \hat{F}_N | \Phi_0 \rangle^{1/2}}.$$

If $|\Phi_0\rangle$ is chosen to be a BCS state, the correlated-basis function method leads to a BCS-like theory, with the bare interaction replaced by an effective interaction which includes many-body correlations. The effective interaction is then calculated by a systematic summation of parquet diagrams of rings and ladders such as the one depicted in Fig. 1(a) and including exchange diagrams in the case of Fermion systems such as the ones depicted in Fig. 1(c).

*e-mail: ppapakon@ibs.re.kr

**e-mail: eckhardk@buffalo.edu

***e-mail: jwang97@buffalo.edu




Figure 1. Examples of diagrams discussed in the text: (a) Parquet and (b) beyond-parquet diagram; (c) exchange diagrams.

The correlation operators are determined by an unconstrained variational principle of the energy expectation value.

This well-known formalism encounters challenges in the case of the nuclear interaction which has a rich operator structure with different radial dependence for each term. Specifically, the bare nuclear interaction is strongly spin-dependent with the spin channels differing significantly. The necessary symmetrization of non-commuting operators in the correlation operator \hat{F}_N leads to terms which we can identify with non-parquet diagrams. These contributions beyond the parquet class describe processes where a spin flip can happen in the intermediate states such as the one depicted in Fig. 1(b). A systematic way to sum the beyond-parquet diagrams has been developed [3]. Here, we present results for triplet pairing in neutron matter.

4 Results

In-medium effective interactions are calculated as outlined above for different nuclear potentials of v_8 type [4, 5]. A dramatic suppression of the in-medium spin-orbit interaction is found, as shown in Fig. 2 in the case of the Reid v_8 potential. As a result, the $^3P_2 - ^3F_2$ gap is found very weak, as shown in Fig. 3(a). In addition, the 3P_0 gap is enhanced, reaching 0.4 – 0.8 MeV for the interactions used here, as shown in Fig. 3(b).




Figure 2. The in-medium spin-orbit effective interaction for the Reid v_8 interaction at different values of Fermi momentum (solid lines) compared with the bare interaction (dashed line).

5 Summary

We have carried out a comprehensive evaluation of the pairing interactions in neutron matter, including medium polarization effects and spin-flip processes, which so far have been ignored or treated in simplistic phenomenological ways. The most striking result is that many-body correlations radically suppress the 3P_2 and $^3P_2 - ^3F_2$ gap by significantly reducing the in-medium spin-orbit interaction. The 3P_0 pairing is found non-negligible. Future extensions could include the effect of three-body forces.




Figure 3. (a) The $^3P_2 - ^3F_2$ gap and (b) the 3P_0 gap calculated in this work with the indicated interactions.

Acknowledgments This work was supported by the Institute for Basic Science (2013M7A1A1075764).

References

- [1] A. Sedrakian and J. W. Clark, Superfluidity in nuclear systems and neutron stars. *Eur. Phys. J. A* **55**, 167 (2019). <https://doi.org/10.1140/epja/i2019-12863-6>
- [2] G. C. Strinati, The BCS–BEC crossover: From ultracold Fermi gases to nuclear systems. *Phys. Rep.* **738**, 1 (2018), <https://doi.org/10.1016/j.physrep.2018.02.004>
- [3] E. Krotscheck and J. Wang, Variational and parquet-diagram calculations for neutron matter. II. Twisted chain diagrams. *Phys. Rev. C* **102**, 064305 (2020). <https://doi.org/10.1103/PhysRevC.102.064305>
- [4] E. Krotscheck, P. Papakonstantinou, J. Wang, Triplet Pairing in Neutron Matter. *Astroph. J.* **955**, 76 (2023). <https://doi.org/10.3847/1538-4357/acee7c>
- [5] E. Krotscheck, P. Papakonstantinou, J. Wang, Variational and parquet-diagram calculations for neutron matter. V. Triplet pairing. *Phys. Rev. C* **109**, 015803 (2024). <https://doi.org/10.1103/PhysRevC.109.015803>

