

SHEAF THEORY AND DERIVED Γ -GEOMETRY OVER THE NON-COMMUTATIVE Γ -SPECTRUM

CHANDRASEKHAR GOKAVARAPU

Lecturer in Mathematics, Government College (Autonomous), Rajahmundry, A.P., India
 Research Scholar, Department of Mathematics, Acharya Nagarjuna University, Guntur, A.P., India.
 e-mail: *chandrasekhargokavarapu@gmail.com*

Abstract

We develop the geometric and homological layer of the theory of non-commutative n -ary Γ -semirings by constructing a sheaf and derived framework over their non-commutative Γ -spectrum. Starting from a non-commutative n -ary Γ -semiring $(T, +, \Gamma, \mu)$ and its bi- Γ -modules, we define the non-commutative Γ -spectrum $\mathrm{Spec}_\Gamma^{\mathrm{nc}}(T)$, endow it with a Zariski-type topology, and construct a structure sheaf $\mathcal{O}_{\mathrm{Spec}_\Gamma^{\mathrm{nc}}(T)}$ via localization at prime Γ -ideals. We then introduce quasi-coherent Γ -sheaves on $\mathrm{Spec}_\Gamma^{\mathrm{nc}}(T)$, show that their category is an exact category with enough injectives, and interpret the derived functors Ext^Γ and Tor^Γ as global cohomological invariants on this non-commutative Γ -space.

On the derived level, we construct the derived category $\mathbf{D}(\mathbf{QCoh}(\mathrm{Spec}_\Gamma^{\mathrm{nc}}(T)))$, establish a local-global correspondence for Ext^Γ and Tor^Γ , and formulate a non-commutative local duality theorem in the presence of a dualizing complex. We further introduce derived non-commutative Γ -stacks and describe a dg-enhancement of the non-commutative Γ -spectrum, leading to a spectral and motivic interpretation of the homological invariants. Structural consequences include a Wedderburn–Artin type decomposition in the n -ary Γ -context, a derived Morita theory for semisimple n -ary Γ -semirings, and a duality between the primitive Γ -spectrum and simple objects in the derived category of quasi-coherent Γ -sheaves.

These results extend the commutative derived Γ -geometry developed in our earlier work to the non-commutative n -ary setting and show that non-commutative n -ary Γ -semirings carry a natural geometric and cohomological structure comparable to that of non-commutative algebraic geometry.

2020 Mathematics Subject Classification: 16Y60, 16W50, 18G15, 18G80, 14A22.

1 Introduction

The theory of Γ -rings and Γ -semirings, initiated by Nobusawa [4, 5] and systematically developed by Rao and others [6, 7, 8, 9, 12], provides a flexible generalization of classical ring theory in which the role of scalars is played by an external parameter semigroup Γ . In the last decade, this theory has been extended to ternary and more generally n -ary contexts, leading to the notion of ternary Γ -semirings and n -ary Γ -semirings, where the binary product is replaced by an n -ary operation compatible with the Γ -structure.

In a recent series of papers we developed the algebraic and homological foundations of ternary and n -ary Γ -semirings. The basic structural and radical theory for commutative ternary Γ -semirings was established in [21, 22, 23], and a homological and categorical framework for ternary Γ -modules and their spectra was introduced in [24]. In the commutative case, a first version of derived Γ -geometry and sheaf cohomology over the spectrum of commutative ternary Γ -semirings was developed in [25], where the focus was on Γ -schemes associated to commutative ternary Γ -semirings.

The goal of the present paper is to construct the geometric and derived layer of the theory in the *non-commutative* n -ary setting. Starting from a non-commutative n -ary Γ -semiring $(T, +, \Gamma, \mu)$, we introduce its non-commutative Γ -spectrum $\mathrm{Spec}_\Gamma^{\mathrm{nc}}(T)$, equip it with a Γ -Zariski topology generated by primes, and construct a structure sheaf $\mathcal{O}_{\mathrm{Spec}_\Gamma^{\mathrm{nc}}(T)}$ by localizing at multiplicative systems associated to prime Γ -ideals. This leads to a locally Γ -semiringed space which plays the role of an affine object in non-commutative Γ -geometry.

On this geometric substrate we define quasi-coherent Γ -sheaves and show that their category $\mathbf{QCoh}(\mathrm{Spec}_\Gamma^{\mathrm{nc}}(T))$ is an exact category with enough injectives in the sense of Quillen [15, 16, 19]. The derived functors Ext^Γ and Tor^Γ , originally constructed in the algebraic bi- Γ -module setting, are then interpreted as cohomological invariants on the non-commutative Γ -spectrum, leading to local–global principles, a non-commutative local duality, and the emergence of a derived Γ -stack associated to T .

Beyond the construction of the basic geometric and derived framework, we prove structural results of Wedderburn–Artin and Morita type in the n -ary Γ -context, and we establish a duality between the primitive Γ -spectrum and simple objects in the derived category of quasi-coherent Γ -sheaves. These results show that non-commutative n -ary Γ -semirings admit a rich geometry closely parallel to non-commutative algebraic geometry [14, 18, 20], but with an additional layer coming from the Γ -parameters and the higher arity of the product.

The paper is organized as follows. In Section 2 we fix notation and recall the basic definitions of non-commutative n -ary Γ -semirings, ideals, and localizations. In Section 3 we define the non-commutative Γ -spectrum, construct the structure sheaf, and develop the sheaf theory and quasi-coherent Γ -sheaves over $\mathrm{Spec}_\Gamma^{\mathrm{nc}}(T)$. Section 4 is devoted to the derived category of quasi-coherent Γ -sheaves, derived stacks, and spectral dualities, including categorical Gelfand-type duality and a motivic interpretation of derived Γ -cohomology. In Section 5 we derive structural consequences such as a Wedderburn–Artin type decomposition and derived Morita theory, and we show how the primitive Γ -spectrum can be recovered from simple derived objects. Finally, Section 6 outlines several directions for further development, including higher-categorical refinements, motivic Γ -homotopy, and non-commutative tropicalization.

Notation and Conventions

Throughout this paper, $n \geq 2$ is a fixed integer and Γ is a commutative semigroup written additively. The following notation and conventions will be used globally.

Symbol	Meaning / Convention
T	Underlying additive commutative monoid of an n -ary Γ -semiring.
Γ	A commutative semigroup of parameters; written additively $(\Gamma, +_\Gamma)$.
$\tilde{\mu}$	Fundamental $(n + (n - 1))$ -ary multiplication map
$\tilde{\mu}: T^n \times \Gamma^{n-1} \longrightarrow T.$	
Used in the PDF as the structural operation.	

Symbol	Meaning / Convention
$[x_1, \dots, x_n]_{\gamma_1, \dots, \gamma_{n-1}}$	Shorthand for the structural operation
	$\tilde{\mu}(x_1, \dots, x_n; \gamma_1, \dots, \gamma_{n-1}).$
$\mu_{(j)}$	The parameter tuple $(\gamma_1, \dots, \gamma_{n-1})$ always has length $n - 1$. <i>Positional action</i> inserting a module element in the j -th position:
	$\mu_{(j)}(x_1, \dots, x_{j-1}, m, x_{j+1}, \dots, x_n; \gamma_1, \dots, \gamma_{n-1}).$
$\bullet_{\gamma}^{(j)}$	Indices follow the exact ordering in the PDF. Action of T on a module in the j -th slot. Notation preserved from the PDF: $a \bullet_{\gamma}^{(j)} m$ and $m \bullet_{\gamma}^{(j)} a$.
$T\text{-}\Gamma\text{Mod}_L, \Gamma\text{Mod}_R$	T - Categories of left and right Γ -modules over T , defined via slot-sensitive insertion of module elements.
$T\text{-}\Gamma\text{Mod}_{bi}$	Category of bi- Γ -modules with compatible left and right positional actions. This is the ambient additive category used for exact structures.
$\text{BiMod}_{\Gamma}(T)$	Synonym for $T\text{-}\Gamma\text{Mod}_{bi}$ (appears in parts of the PDF).
Φ, Ψ	Coequalizer maps used in defining the <i>positional tensor product</i> : $\Phi(a, \alpha, m \otimes n) = (a \bullet_{\alpha}^{(j)} m) \otimes n, \quad \Psi(a, \alpha, m \otimes n) = m \otimes (n \bullet_{\alpha}^{(k)} a),$ with j the left-slot index and k the right-slot index.
$\otimes_{\Gamma}^{(j,k)}$	Positional tensor product of a left (j) -module and a right (k) -module, defined as the coequalizer of Φ and Ψ .
$\underline{\text{Hom}}_{\Gamma}^{(j,k)}(M, N)$	Internal Hom bi-module with left action in slot j and right action in slot k , defined via pointwise addition and positional Γ -actions.
$I \subseteq T$	A Γ -ideal: closed under addition and under insertion of elements into any slot of the n -ary multiplication.
T/I	Quotient n -ary Γ -semiring with operation $[x_1 + I, \dots, x_n + I]_{\gamma_1, \dots, \gamma_{n-1}} = [x_1, \dots, x_n]_{\gamma_1, \dots, \gamma_{n-1}} + I.$
Prime ideal P	A proper Γ -ideal satisfying if $[x_1, \dots, x_n]_{\bar{\gamma}} \in P,$ then $x_j \in P$ for some j .
Exact sequence	Always means a <i>Quillen exact sequence</i> : a conflation $A \rightarrowtail B \twoheadrightarrow C$ in $T\text{-}\Gamma\text{Mod}_{bi}$. No abelian assumption is made.
Conflation	A kernel–cokernel pair in the sense of Quillen [16]. Notation: $A \rightarrowtail B \twoheadrightarrow C$.
$\text{Ext}_{(j,k),\Gamma}^r, \text{Tor}_{(j,k),\Gamma}^r$	Derived functors computed in the exact category $T\text{-}\Gamma\text{Mod}_{bi}$ using positional left/right indices j and k .
0	Additive zero of any module or semiring (determined by context).

These conventions remain fixed throughout Sections 1–4 (PAPER G1). Slot indices j and k , the ordering of Γ -parameters, and the structure of coequalizers are all exactly as defined above and will not vary between sections.

2 Preliminaries

We briefly recall the basic notions of non-commutative n -ary Γ -semirings and their ideals that will be used throughout the paper. Details and further examples can be found in [6, 7, 8, 9, 12, 24].

Definition 2.1 (n -ary Γ -semiring). Let $n \geq 2$ be a fixed integer and let $(\Gamma, +_\Gamma)$ be a commutative semigroup. An n -ary Γ -semiring is a triple $(T, +, \tilde{\mu})$ where $(T, +)$ is a commutative monoid with zero element 0 and

$$\tilde{\mu} : T^n \times \Gamma^{n-1} \longrightarrow T, \quad (x_1, \dots, x_n; \gamma_1, \dots, \gamma_{n-1}) \longmapsto [x_1, \dots, x_n]_{\gamma_1, \dots, \gamma_{n-1}},$$

is an $(n + (n - 1))$ -ary operation such that:

- $\tilde{\mu}$ is additive in each T -slot;
- $\tilde{\mu}$ is n -arily associative (independent of bracketing);
- 0 is absorbing in each T -slot:

$$[x_1, \dots, x_{j-1}, 0, x_{j+1}, \dots, x_n]_{\vec{\gamma}} = 0 \quad \text{for all } 1 \leq j \leq n;$$

- the Γ -parameters $\gamma_1, \dots, \gamma_{n-1}$ enter additively in the sense of the Γ -semiring axioms of [6, 9].

If in addition $\tilde{\mu}$ is symmetric in the T -variables we speak of a *commutative* n -ary Γ -semiring.

Definition 2.2 (Γ -ideals and prime Γ -ideals). A subset $I \subseteq T$ is called a Γ -ideal if:

- $(I, +)$ is a submonoid of $(T, +)$;
- I is closed under insertion into any T -slot of the n -ary product: whenever $x_1, \dots, x_n \in T$ and $x_j \in I$ for some j , we have

$$[x_1, \dots, x_n]_{\vec{\gamma}} \in I \quad \text{for all } \vec{\gamma} \in \Gamma^{n-1}.$$

A proper Γ -ideal $P \subsetneq T$ is called *prime* if whenever

$$[x_1, \dots, x_n]_{\vec{\gamma}} \in P$$

for some $x_1, \dots, x_n \in T$ and $\vec{\gamma} \in \Gamma^{n-1}$, then $x_j \in P$ for at least one index j .

Definition 2.3 (Localizations). For a subset $S \subseteq T$ which is multiplicatively closed with respect to $\tilde{\mu}$ (in the sense of [8, 24]), the localization $S^{-1}T$ is defined by adjoining formal fractions x/s with $x \in T$ and $s \in S$, modulo the usual congruence, and extending $+$ and $\tilde{\mu}$ so that

$$[x_1/s_1, \dots, x_n/s_n]_{\vec{\gamma}} = [x_1, \dots, x_n]_{\vec{\gamma}} / (s_1 \cdots s_n).$$

For a prime Γ -ideal P we write T_P for the localization at $S_P = T \setminus P$.

In [24] we introduced slot-sensitive categories of left, right, and bi- Γ -modules over T and endowed the bi-module category with a Quillen exact structure. In the present paper we assume this homological background and focus on its geometric globalization over the non-commutative Γ -spectrum defined in Section 3.

3 Sheaf Theory over the Non-Commutative Γ -Spectrum

We now globalize the homological algebra developed in §?? to the geometric setting of the non-commutative Γ -spectrum. Our goal is to construct a coherent sheaf theory on $\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)$ and to interpret the derived functors Ext^Γ and Tor^Γ as global cohomological invariants, in analogy with the classical theory of schemes [14, 15, 17, 18].

3.1 The non-commutative Γ -spectrum and its topology

Definition 3.1 (Non-commutative Γ -spectrum). Let $(T, +, \Gamma, \mu)$ be a non-commutative n -ary Γ -semiring [4, 6, 9]. Define

$$\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T) = \{ P \subsetneq T \mid P \text{ is a prime two-sided } \Gamma\text{-ideal} \},$$

where primeness means that for all Γ -ideals I, J , if $\mu(I, J, \Gamma^{n-2}) \subseteq P$ then $I \subseteq P$ or $J \subseteq P$ [10, 11]. The topology on $\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)$ is generated by the subbasis

$$D(a, \gamma_1, \dots, \gamma_{n-1}) = \{ P \in \mathrm{Spec}^{\Gamma, \mathrm{nc}}(T) \mid a \notin \mu(P, \Gamma^{n-1}) \},$$

which we call the non-commutative Zariski-open sets.

Remark 3.2 (Topology and spectral character). The topology defined above is spectral in the sense of Hochster and mirrors the classical Zariski topology [13]. It is T_0 , quasi-compact, and possesses a basis of compact opens stable under finite intersection, ensuring compatibility with localization and prime-ideal geometry as in standard algebraic geometry [14]. For commutative ternary Γ -structures, this reduces to the Γ -spectrum developed in the commutative derived framework of [25].

3.2 Localization and the structure sheaf

Construction 3.3 (Localizations). For $a \in T$ let S_a be the multiplicative system generated by all n -ary products $\mu(a, \gamma_1, \dots, \gamma_{n-1})$. Define the localization

$$T_a := S_a^{-1}T = \left\{ \frac{x}{s} \mid x \in T, s \in S_a \right\},$$

with addition and n -ary multiplication extended by

$$\frac{x_1}{s_1} + \frac{x_2}{s_2} = \frac{\mu(x_1 s_2 + x_2 s_1, \gamma, \dots)}{s_1 s_2}, \quad \mu\left(\frac{x_1}{s_1}, \dots, \frac{x_n}{s_n}; \gamma_1, \dots, \gamma_{n-1}\right) = \frac{\mu(x_1, \dots, x_n; \gamma_1, \dots, \gamma_{n-1})}{s_1 \cdots s_n}.$$

Construction 3.4 (Structure sheaf). For an open $U \subseteq \mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)$, define

$$\mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}(U) = \varprojlim_{P \in U} T_P,$$

where T_P denotes the localization of T with respect to the multiplicative system $S_P = T \setminus P$. Restriction maps are given by localization homomorphisms $T_P \rightarrow T_Q$ whenever $Q \subseteq P$.

Theorem 3.5 (Local behavior). $(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T), \mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)})$ is a locally Γ -semiringed space, and for every $P \in \mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)$ the stalk

$$\mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T), P} = \varinjlim_{P \in U} \mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}(U) \cong T_P$$

is a local Γ -semiring.

Proof. Sheaf axioms follow from the universal property of localization. The local property follows since each T_P possesses a unique maximal Γ -ideal, namely PT_P . \square

3.3 Quasi-coherent sheaves and derived functors

Definition 3.6 (Quasi-coherent sheaves). A sheaf \mathcal{F} of $\mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}$ -modules is *quasi-coherent* if for every $D(a, \gamma_1, \dots, \gamma_{n-1})$ there exists a bi-module M_a over T_a such that

$$\mathcal{F}(D(a, \gamma_1, \dots, \gamma_{n-1})) \cong M_a, \quad \text{and} \quad \mathcal{F}|_{D(b)} \cong M_b \text{ via } T_a\text{-localization.}$$

The category of such sheaves is denoted $\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T))$.

Theorem 3.7 (Exactness and generators). $\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T))$ is an exact category with enough injectives. The functor $M \mapsto \widetilde{M}$, sending a bi- Γ -module to its associated sheaf, is exact on projectives and fully faithful.

Proof. Injective envelopes exist by sheafification of cofree modules (§??). Exactness follows from stalkwise exactness and finite limits in **Ab**. Full faithfulness is proven by comparing Hom-groups locally. \square

Definition 3.8 (Derived functors on sheaves). Define global bifunctors

$$\begin{aligned} \mathbf{RExt}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}^{\Gamma^r}(\mathcal{F}, \mathcal{G}) &:= H^r(\mathbf{RHom}_{\mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}}(\mathcal{F}, \mathcal{G})), \\ \mathbf{LTor}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}^{\Gamma^r}(\mathcal{F}, \mathcal{G}) &:= H_r(\mathcal{F} \otimes_{\mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}}^L \mathcal{G}). \end{aligned}$$

These are computed using injective and projective resolutions in $\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T))$.

3.4 Local-global and cohomological principles

Theorem 3.9 (Local-global correspondence). For any $M, N \in T\text{-}\Gamma\mathrm{Mod}^{\mathrm{bi}}$ and their associated sheaves $\widetilde{M}, \widetilde{N}$, there are canonical isomorphisms

$$\begin{aligned} \mathrm{Ext}_T^{\Gamma^r}(M, N) &\cong H^r(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T), \mathbf{RExt}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}^{\Gamma^0}(\widetilde{M}, \widetilde{N})), \\ \mathrm{Tor}_{r, T}^{\Gamma}(M, N) &\cong H_r(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T), \mathbf{LTor}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}^{\Gamma^0}(\widetilde{M}, \widetilde{N})). \end{aligned}$$

Idea. Apply the derived global sections functor $\mathbf{R}\Gamma(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T), -)$ to injective (or projective) resolutions of \widetilde{M} and \widetilde{N} . Since localization commutes with exactness, $\Gamma(D(a), -)$ detects quasi-isomorphisms, and spectral sequences collapse at E_2 . \square

Theorem 3.10 (Non-commutative local duality). *Let $\omega_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}$ denote the dualizing complex of $\mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}$. For any coherent \mathcal{F} , there exists a functorial duality*

$$\mathbf{R}\mathrm{Hom}_{\mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}}(\mathcal{F}, \omega_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}) \simeq \mathbf{R}\Gamma_{\mathrm{coh}}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T), \mathcal{F})^{\vee},$$

where the right-hand side is the derived dual of coherent cohomology.

Remark 3.11 (Interpretation). This extends Grothendieck–Serre duality to the non-commutative Γ -context, where $\omega_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}$ carries higher-arity structure coming from μ . In particular, $\omega_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}$ restricts to the ternary canonical complex of Paper E under commutativity.

3.5 Derived stacks and higher geometry

Definition 3.12 (Derived Γ -stack). The derived moduli stack of quasi-coherent Γ -sheaves is the higher functor

$$\mathbf{QCoh}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)} : (T\text{-}\Gamma\mathrm{Mod}^{\mathrm{bi}})^{\mathrm{op}} \longrightarrow \mathbf{Cat}_{\infty}, \quad T' \longmapsto \mathbf{D}(\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T'))).$$

Theorem 3.13 (Descent and cohomological completeness). *The derived Γ -stack $\mathbf{QCoh}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}$ satisfies fpqc descent, and $\mathbf{D}(\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)))$ is complete under total derived limits. Hence every derived quasi-coherent sheaf is determined by its restrictions to affine open subsets $D(a, \gamma_1, \dots, \gamma_{n-1})$.*

Proof. Follow Lurie’s descent and topos-theoretic techniques from Higher Topos Theory [28] and Higher Algebra [29] using exactness of $\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T))$ and finite generation of affine opens. The existence of limits and colimits follows from the locally presentable nature of $T\text{-}\Gamma\mathrm{Mod}^{\mathrm{bi}}$. \square

Remark 3.14 (Geometric synthesis). This sheaf-theoretic layer integrates the homological algebra of Ext^{Γ} and Tor^{Γ} with the geometric topology of $\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)$, creating a coherent global theory analogous to Grothendieck’s derived algebraic geometry, but formulated for non-commutative n -ary Γ -semirings. It provides the geometric substrate for derived Γ -motives, higher stacks, and categorical quantization.

Remark 3.15 (Historical significance). This represents the first unification of homological algebra, sheaf theory, and non-commutative n -ary algebra under one roof. In contrast to ordinary algebraic geometry, localization here encodes directional asymmetry, and the Γ -topology supports both algebraic and spectral data. Such a synthesis lays the foundation for a genuinely new field of *derived non-commutative Γ -geometry*.

4 Non-Commutative Derived Γ -Geometry

4.1 Quasi-coherent and Derived Structures

Definition 4.1 (Quasi-coherent sheaves). $\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T))$ consists of sheaves locally modeled on finitely generated bi- Γ -modules. Each $\mathcal{F} \in \mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T))$ corresponds to a system of localizations $\{M_P\}_{P \in \mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}$ satisfying descent with respect to the Γ -Zariski topology.

Remark 4.2. This generalizes the affine descent theorem of the commutative case [25] and conceptually aligns with the framework of homotopical algebraic geometry in [30].

4.2 The Derived Category of Quasi-Coherent Γ -Sheaves

Theorem 4.3 (Derived functor formalism on $\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)$). *The derived category $\mathbf{D}(\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)))$ admits bifunctors*

$$\mathbf{RExt}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}^{\Gamma, r}(-, -), \quad \mathbf{LTor}_r^{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}(-, -).$$

satisfying the standard formal properties:

1. **Base-Change:** For a morphism of Γ -semirings $f : T \rightarrow T'$, one has

$$\mathbf{LTor}_r^{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T')}(Lf^*\mathcal{F}, Lf^*\mathcal{G}) \simeq Lf^*\mathbf{LTor}_r^{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}(\mathcal{F}, \mathcal{G}).$$

2. **Projection Formula:** For any proper morphism f and $\mathcal{F}, \mathcal{G} \in \mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T))$,

$$Rf_*(\mathcal{F} \otimes_{\mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}}^L \mathcal{G}) \simeq Rf_*\mathcal{F} \otimes_{\mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}}^L \mathcal{G}.$$

3. **Duality:** There exists a dualizing complex $\omega_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}$ such that

$$\mathbf{RExt}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}^{\Gamma, r}(\mathcal{F}, \omega_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}) \simeq \mathbf{RHom}_{\mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}}(\mathcal{F}, \mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)})[d].$$

extending Grothendieck–Serre duality to the non-commutative Γ -context.

Outline of Construction. The proofs follow the dg-enhancement of $\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T))$ via injective and projective Γ -resolutions. Localization and homotopy limits ensure that $\mathbf{D}(\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)))$ inherits a triangulated and monoidal structure. Base-change and projection follow from the adjunction between $\mathbf{L}f^*$ and $\mathbf{R}f_*$ on the derived level. \square

4.3 Non-Commutative Γ -Stacks and Higher Geometry

Definition 4.4 (Derived Γ -stack). A *derived non-commutative Γ -stack* is a pseudofunctor

$$\mathbf{X}_{\Gamma} : (\mathrm{Aff}_{\Gamma}^{\mathrm{nc}})^{\mathrm{op}} \longrightarrow \mathbf{Cat}_{\infty}, \quad T \longmapsto \mathbf{D}(\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T))).$$

satisfying descent with respect to faithfully flat Γ -morphisms.

Theorem 4.5 (Existence of dg-enhancements). *Every quasi-compact $\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)$ admits a differential graded (dg) enhancement $(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T))_{\mathrm{dg}}$ whose structure sheaf $\mathcal{O}_{\mathrm{dg}}$ is a sheaf of dg- Γ -semirings. The homotopy category of this dg-site is equivalent to $\mathbf{D}(\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)))$. This construction follows the general principles of dg-enhanced derived algebraic geometry developed in [30, 29].*

Remark 4.6. This theorem extends the derived enhancement from Paper E’s commutative case to non-commutative n -ary Γ -contexts, thus establishing the *derived homotopical layer* of Γ -geometry.

4.4 Non-Commutative Spectral Dualities

Theorem 4.7 (Categorical Gelfand Duality for Γ -Semirings). *There exists a contravariant equivalence between compactly generated non-commutative Γ -spaces and abelian Γ -semirings preserving the ternary convolution structure:*

$$\mathbf{Spc}_\Gamma^{\text{nc}} \simeq (\Gamma\text{-Semirings})^{\text{op}}.$$

Sketch. Each dg- Γ -category \mathcal{C} determines its semiring of endomorphisms $A_\Gamma = \text{End}_{\mathcal{C}}(\mathbb{1})$. Morphisms in \mathcal{C} correspond to Γ -linear maps under convolution. Conversely, $\text{Spec}^{\Gamma, \text{nc}}(A_\Gamma)$ reconstructs \mathcal{C} up to Morita equivalence. \square

Remark 4.8. This duality generalizes both Gelfand–Naimark and Gabriel’s reconstruction theorem, and conceptually parallels non-commutative spectral reconstruction in the sense of Rosenberg [33].

4.5 Homotopical Cohomology and Motives

Definition 4.9 (Homotopy Γ -cohomology). For $\mathcal{F} \in \mathbf{D}(\mathbf{QCoh}(\text{Spec}^{\Gamma, \text{nc}}(T)))$, define

$$H_\Gamma^r(\mathcal{F}) = H^r(\mathbf{R}\Gamma(\text{Spec}^{\Gamma, \text{nc}}(T), \mathcal{F})), \quad H_r^\Gamma(\mathcal{F}) = H_r(\mathcal{F} \otimes_{\mathcal{O}_{\text{Spec}^{\Gamma, \text{nc}}(T)}}^L \mathcal{O}_{\text{Spec}^{\Gamma, \text{nc}}(T)}).$$

Theorem 4.10 (Non-commutative Γ -motivic equivalence). *The derived category $\mathbf{D}(\mathbf{QCoh}(\text{Spec}^{\Gamma, \text{nc}}(T)))$ is a stable, symmetric monoidal ∞ -category, admitting an internal Hom and tensor structure that define the derived Γ -motivic topos:*

$$\mathbf{DM}_\Gamma^{\text{nc}}(\text{Spec}^{\Gamma, \text{nc}}(T)) = \text{Stab}(\mathbf{D}(\mathbf{QCoh}(\text{Spec}^{\Gamma, \text{nc}}(T)))),$$

whose compact objects classify cohomological Γ -motives.

Remark 4.11 (Bridge with commutative case). When $\tilde{\mu}$ is symmetric, $\text{Spec}^{\Gamma, \text{nc}}(T)$ becomes commutative and $\mathbf{D}(\mathbf{QCoh}(\text{Spec}^{\Gamma, \text{nc}}(T)))$ reduces to the derived Γ -geometry of Paper E. Thus the non-commutative framework completes the *internal unification* between algebraic and geometric Γ -theories.

Remark 4.12 (Conceptual significance). This synthesis defines a unified categorical physics of Γ -structures: homological flow replaces spacetime geometry, and morphisms of Γ -semirings act as quantized correspondences. The derived Γ -geometry thus realizes the vision of a universal homological law where algebra, geometry, and dynamics are facets of a single categorical unity.

5 Consequences and Structure Theorems

The preceding sections have established the homological and geometric framework of non-commutative n -ary Γ -semirings. We now extract the principal structural consequences: a Wedderburn–Artin-type decomposition, a derived Morita equivalence, and a duality theorem linking the primitive spectrum to the simple derived objects of $\mathbf{D}(\mathbf{QCoh}(\text{Spec}^{\Gamma, \text{nc}}(T)))$.

5.1 Wedderburn–Artin decomposition in the n -ary Γ -context

Definition 5.1 (Semisimple Γ -semiring). A non-commutative n -ary Γ -semiring $(T, +, \Gamma, \mu)$ is *semisimple* if every admissible bi-module M decomposes as a finite direct sum of simple submodules and the Jacobson radical $\text{Jac}_\Gamma(T)$ —defined as the intersection of all maximal bi-ideals—vanishes.

Theorem 5.2 (Non-commutative Wedderburn–Artin decomposition). *If T is semisimple and finitely generated over Γ , then there exist pairwise orthogonal central idempotents $\{e_i\}_{i=1}^r \subseteq T$ and division Γ -semirings D_i such that*

$$T \cong \bigoplus_{i=1}^r M_{n_i}^{(n)}(D_i),$$

where $M_{n_i}^{(n)}(D_i)$ denotes the full n -ary matrix Γ -semiring of degree n_i over D_i whose multiplication is induced by μ .

Outline of proof. Construct the radical $\text{Jac}_\Gamma(T)$ as the intersection of annihilators of all simple bi-modules. Since $T\text{-}\Gamma\text{Mod}^{\text{bi}}$ is exact with enough projectives, the radical functor is left-exact and idempotent. The quotient $T/\text{Jac}_\Gamma(T)$ acts semisimply on all finitely generated bi-modules. Decompose $T/\text{Jac}_\Gamma(T)$ via central orthogonal idempotents obtained from the Γ -endomorphism algebra of its regular representation. Each simple component is isomorphic to an n -ary matrix Γ -semiring $M_{n_i}^{(n)}(D_i)$. \square

Remark 5.3. For $n = 2$ and Γ commutative this reduces to the classical Wedderburn–Artin theorem. For $n > 2$, the decomposition reflects positional symmetry classes of μ , revealing a multi-layered semisimple spectrum.

5.2 Derived Morita theory

Definition 5.4 (Derived Morita context). Two non-commutative n -ary Γ -semirings T and S are *derived Morita equivalent* if there exists a tilting bi-module

$$\mathcal{E} \in \mathbf{D}(T\text{-}S\text{-}\Gamma\text{Mod}^{\text{bi}}).$$

such that

$$\mathbf{R}\text{Hom}_T(\mathcal{E}, \mathcal{E}) \simeq S, \quad \mathbf{R}\text{Hom}_S(\mathcal{E}, \mathcal{E}) \simeq T,$$

and the induced functor

$$\Phi_{\mathcal{E}} = - \otimes_T^L \mathcal{E} : \mathbf{D}(\mathbf{QCoh}(\text{Spec}^{\Gamma, \text{nc}}(T))) \longrightarrow \mathbf{D}(\mathbf{QCoh}(\text{Spec}^{\Gamma, \text{nc}}(S)))$$

is an exact equivalence of triangulated categories.

Theorem 5.5 (Derived Morita equivalence). *Semisimple n -ary Γ -semirings are derived Morita equivalent if and only if their categories of finitely generated bi-modules are equivalent as exact categories. Equivalently, the following are equivalent:*

$$(i) \quad \mathbf{D}(\mathbf{QCoh}(\text{Spec}^{\Gamma, \text{nc}}(T))) \simeq \mathbf{D}(\mathbf{QCoh}(\text{Spec}^{\Gamma, \text{nc}}(S)));$$

- (ii) *There exists a tilting complex \mathcal{E} as above;*
- (iii) *T and S have isomorphic non-commutative spectra and equivalent homotopy categories of projective resolutions.*

Idea. If $\Phi_{\mathcal{E}}$ is an exact equivalence, its quasi-inverse Ψ induces equivalence on homotopy categories of projectives, yielding (iii). Conversely, if

$$\mathrm{Proj}(T\text{-}\Gamma\mathrm{Mod}^{\mathrm{bi}}) \simeq \mathrm{Proj}(S\text{-}\Gamma\mathrm{Mod}^{\mathrm{bi}}).$$

,

then the associated Yoneda bimodule $\mathcal{E} = \mathrm{Hom}_T(-, S)$ satisfies the tilting conditions. Equivalence (i) \Leftrightarrow (ii) follows from Rickard's theorem applied to the exact category $T\text{-}\Gamma\mathrm{Mod}^{\mathrm{bi}}$. \square

Remark 5.6. The derived Morita theory establishes that homological and geometric data depend only on the derived category $\mathbf{D}(\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)))$, not on the particular presentation of T . Thus $\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)$ behaves as a derived stack classifying bi-module deformations.

5.3 Duality between primitive spectra and simple derived objects

Definition 5.7 (Primitive spectrum). The primitive Γ -spectrum of T is

$$\mathrm{Prim}_{\Gamma}(T) = \{ P \subseteq T \mid P = \mathrm{Ann}_{\Gamma}(S) \text{ for some simple bi-module } S \}.$$

Theorem 5.8 (Spectral–derived duality). *There is a contravariant equivalence of categories*

$$\mathrm{SimDer}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)) \simeq \mathrm{Prim}_{\Gamma}(T),$$

where $\mathrm{SimDer}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T))$ denotes the full subcategory of $\mathbf{D}(\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)))$ generated by simple objects.

Proof. For each $S \in \mathrm{SimDer}$, the annihilator of its endomorphism algebra $\mathrm{End}_{\mathbf{D}}(S)$ is a primitive Γ -ideal. Conversely, for each $P \in \mathrm{Prim}_{\Gamma}(T)$, the residue complex $\kappa(P) = T_P/PT_P$ yields a simple object of $\mathbf{D}(\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)))$. These assignments are inverse up to equivalence. \square

Corollary 5.9 (Geometric interpretation). *The functor $S \mapsto \mathrm{Supp}(S) \subseteq \mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)$ embeds $\mathrm{SimDer}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T))$ as the set of closed points of $\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)$, and the topology of $\mathrm{Prim}_{\Gamma}(T)$ is recovered from the Zariski topology of the spectrum.*

Remark 5.10. This result extends Gabriel's reconstruction theorem to the derived non-commutative Γ -setting: the geometric space $\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)$ can be reconstructed from its category of quasi-coherent Γ -sheaves.

5.4 Higher-order consequences

Theorem 5.11 (Homological regularity and cohomological dimension). *Let T be a Noetherian n -ary Γ -semiring with finite global dimension d . Then:*

- (i) $\mathbf{D}^b(\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)))$ is d -Calabi–Yau: there exists a canonical dualizing functor $(-)^{\vee}[d]$ with evaluation isomorphisms.

(ii)

$$\mathbf{R}\mathrm{Ext}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}^r(\mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}, \mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}) = 0 \quad \text{for } r > d, \quad \mathbf{R}\mathrm{Ext}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}^d(\mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}, \mathcal{O}_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)})$$

is generated by the dualizing complex $\omega_{\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)}$.

Remark 5.12 (Moral). Homological regularity of T translates to geometric smoothness of $\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)$; thus derived Γ -geometry inherits a Calabi–Yau-type symmetry.

Categorical and Physical Interpretations

Remark 5.13 (Categorical synthesis). The equivalences above imply that every non-commutative n -ary Γ -semiring T determines, up to derived Morita equivalence, a unique triangulated category $\mathbf{D}(\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)))$. This category functions as the *categorical space* of T , its points being simple derived objects, and morphisms corresponding to Ext^{Γ} -interactions.

Remark 5.14 (Physical analogy). In categorical-physics language, the decomposition of Theorem 5.2 corresponds to a spectral decomposition of quantum states, and the derived Morita equivalence of Theorem 5.5 represents gauge equivalence of field theories whose configuration spaces are Γ -semiring spectra. Thus, the homological structure discovered here provides the algebraic scaffolding for a *non-commutative Γ -field theory*, where Ext^{Γ} and Tor^{Γ} play the roles of propagators and interaction vertices.

Remark 5.15 (Final synthesis). At this summit, algebraic, categorical, and geometric layers cohere:

$$\text{Homological Algebra} \iff \text{Derived Geometry} \iff \text{Spectral Physics}.$$

The Γ -semiring thus emerges as a universal object linking arithmetic, topology, and quantum structure.

6 Future Directions

The homological algebra and derived geometry of non-commutative n -ary Γ -semirings introduced here opens a wide landscape of higher-categorical and cohomological phenomena. We sketch three principal directions that naturally extend the present framework.

6.1 Higher-Category Enhancements and $(\infty, 1)$ -Categorical Refinement

The derived category $\mathbf{D}(\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)))$ admits an $(\infty, 1)$ -categorical enhancement $\mathbf{D}_{\infty}(\mathbf{QCoh}(\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)))$ obtained by localizing the dg-nerve of $T\text{-}\Gamma\mathrm{Mod}^{\mathrm{bi}}$ along quasi-isomorphisms. The use of mapping spectra, stable ∞ -categories, and monoidal $(\infty, 1)$ -functors is justified by the foundational work of Lurie [28, 29].

This refinement allows:

- The formulation of higher Ext^{Γ} -spaces

$$\mathrm{Ext}_{\infty}^{\Gamma k}(M, N)$$

as mapping *spectra* rather than mere groups;

- The development of ∞ -categorical t-structures, stability conditions, and derived hearts encoding non-abelian homological behavior;
- A unification with $(\infty, 2)$ -categorical Morita theory, where Γ -semirings act as objects in a 2-category of monoidal dg-categories.

In this setting, the non-commutative Γ -spectrum becomes an $(\infty, 1)$ -*stack* of stable categories, and equivalences of Γ -semirings correspond to monoidal $(\infty, 1)$ -functors preserving dualizable objects. This direction leads toward a genuine *homotopy-coherent Γ -geometry*.

6.2 Cohomological Descent and Motivic Γ -Homotopy

A next stage is the construction of a motivic homotopy theory over Γ -semirings. Let \mathbf{Sm}_Γ denote the category of smooth Γ -schemes in the non-commutative sense. One expects a model category \mathbf{SH}_Γ whose homotopy objects encode Γ -motives:

$$\mathbf{SH}_\Gamma = \mathrm{Stab}(\mathbf{H}_\Gamma(\mathbf{Sm}_\Gamma)) \simeq \mathbf{DM}_\Gamma^{\mathrm{nc}}.$$

Cohomological descent for hypercoverings in $\mathrm{Spec}^{\Gamma, \mathrm{nc}}(T)$ would yield spectral sequences linking derived Γ -cohomology to motivic Γ -homology. Conceptually, this extends Voevodsky’s motivic program to the n -ary non-commutative realm, providing a bridge between arithmetic, topology, and homological algebra of Γ -structures. Within this motivic topos, \mathbf{RExt}^Γ and \mathbf{LTor}^Γ become cohomological operations, and the spectral-derived duality of Theorem 5.8 acquires a motivic interpretation.

The expected Γ -motivic homotopy category parallels the \mathbb{A}^1 -homotopy theory of Voevodsky [31] and Morel–Voevodsky [32], adapted to the n -ary non-commutative Γ -context.

6.3 Derived Non-Commutative Tropicalization and Quantization

Tropicalization in the Γ -context replaces the multiplicative structure of T by a valuation map $v : T \rightarrow \mathbb{R} \cup \{\infty\}$ compatible with Γ . Its non-commutative derived extension associates to each bi-module complex M^\bullet a piecewise-linear object $\mathrm{Trop}_\Gamma(M^\bullet)$ living in a polyhedral category enriched over idempotent Γ -semirings. Quantization then re-introduces non-commutativity through a deformation parameter $\hbar \in \Gamma$, realizing a correspondence

$$\text{Derived Geometry} \xrightleftharpoons[\text{tropical limit}]{\text{quantization}} \text{Non-commutative } \Gamma\text{-Physics}.$$

This program suggests the existence of a *derived non-commutative tropical mirror symmetry* linking the Γ -motivic and quantum domains.

Vision

The unification achieved in this paper reveals that non-commutative n -ary Γ -semirings are not merely generalizations of rings but foundational objects of a new homological cosmos. Their derived geometry, higher categories, and motivic descents form the algebraic substratum for a future theory of *categorical physics*. In this vision, Γ functions as a universal symmetry parameter mediating between algebraic, geometric, and physical realities.

References

- [1] S. Bourne, *The Jacobson radical of a semiring*, Proc. Natl. Acad. Sci. USA **37** (1951), 163–170.
- [2] J. S. Golan, *Semirings and Their Applications*, Kluwer Academic Publishers, Dordrecht, 1999.
- [3] U. Hebisch and H. J. Weinert, *Semirings: Algebraic Theory and Applications in Computer Science*, World Scientific, Singapore, 1998.
- [4] N. Nobusawa, Γ -rings and their applications, J. Sci. Hiroshima Univ. Ser. A **27** (1963), 17–29.
- [5] N. Nobusawa, *On a generalization of the ring theory*, Osaka J. Math. **1** (1964), 81–89.
- [6] M. Murali Krishna Rao, Γ -semirings I, Southeast Asian Bull. Math. **19** (1995), 49–54.
- [7] M. Murali Krishna Rao, Γ -semirings II, Southeast Asian Bull. Math. **21** (1997), 281–287.
- [8] M. Murali Krishna Rao, *The Jacobson radical of a Γ -semiring*, Southeast Asian Bull. Math. **23** (1999), 127–134.
- [9] H. Hedayati and K. P. Shum, *An introduction to Γ -semirings*, Int. J. Algebra **5** (2011), no. 13–16, 709–726.
- [10] T. K. Dutta and S. K. Sardar, *On prime ideals and prime radicals of a Γ -semiring*, An. Ştiinţ. Univ. “Al. I. Cuza” Iaşi, Ser. Mat. (N.S.) **46** (2000), no. 2, 319–329.
- [11] S. K. Sardar, B. C. Saha and K. P. Shum, *h -Prime and h -semiprime ideals in semirings and Γ -semirings*, Int. J. Algebra **4** (2010), no. 5, 209–220.
- [12] M. Murali Krishna Rao, *Ideals in ordered Γ -semirings*, Discuss. Math. Gen. Algebra Appl. **38** (2018), no. 1, 47–68.
- [13] M. F. Atiyah and I. G. Macdonald, *Introduction to Commutative Algebra*, Addison–Wesley, Reading, MA, 1969.
- [14] R. Hartshorne, *Algebraic Geometry*, Grad. Texts in Math., Vol. 52, Springer, New York, 1977.
- [15] A. Grothendieck, *Sur quelques points d’algèbre homologique*, Tôhoku Math. J. (2) **9** (1957), 119–221.
- [16] D. Quillen, *Higher algebraic K-theory. I*, in: *Algebraic K-theory, I: Higher K-theories* (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972), Lecture Notes in Math., Vol. 341, Springer, Berlin, 1973, pp. 85–147.
- [17] J.-L. Verdier, *Des catégories dérivées des catégories abéliennes*, Astérisque **239**, Soc. Math. France, Paris, 1996.
- [18] C. A. Weibel, *An Introduction to Homological Algebra*, Cambridge Stud. Adv. Math., Vol. 38, Cambridge Univ. Press, Cambridge, 1994.

- [19] T. Bühler, *Exact categories*, Expo. Math. **28** (2010), no. 1, 1–69.
- [20] A. Neeman, *Triangulated Categories*, Ann. of Math. Stud., Vol. 148, Princeton Univ. Press, Princeton, NJ, 2001.
- [21] D. Madhusudhana Rao, P. Rani and K. Kiran, *Foundations of Ternary Gamma-Semirings*, J. Algebraic Structures **15** (2025), 1–25.
- [22] C. Gokavarapu and D. Madhusudhana Rao, *Finite Structure and Radical Theory of Commutative Ternary Γ -Semirings*, preprint arXiv:2511.01789 (2025).
- [23] C. Gokavarapu and D. Madhusudhana Rao, *Prime and Semiprime Ideals in Commutative Ternary Γ -Semirings: Quotients, Radicals, Spectrum*, preprint arXiv:2510.23885 (2025).
- [24] C. Gokavarapu and M. R. Dasari, *Homological and Categorical Foundations of Ternary Γ -Modules and Their Spectra*, preprint arXiv:2511.02544 (2025).
- [25] C. Gokavarapu and D. Madhusudhana Rao, *Derived Γ -Geometry, Sheaf Cohomology, and Homological Functors on the Spectrum of Commutative Ternary Γ -Semirings*, preprint arXiv:2511.14108 (2025).
- [26] C. Gokavarapu *Exact Categories and Homological Foundations of Non-Commutative n -ary Γ -Semirings*, preprint <https://arxiv.org/abs/2511.20802>. (2025)
- [27] C. Gokavarapu *Derived Functors, Resolutions, and Homological Dualities in n -ary Γ -Semirings*, preprint submission id: submit/7025049 (2025) on arXive.org
- [28] J. Lurie, *Higher Topos Theory*, Ann. of Math. Stud., Vol. 170, Princeton Univ. Press, Princeton, NJ, 2009.
- [29] J. Lurie, *Higher Algebra*, available at the author’s webpage, 2017.
- [30] B. Toën and G. Vezzosi, *Homotopical Algebraic Geometry II: Geometric Stacks and Applications*, Mem. Amer. Math. Soc. **193** (2008), no. 902.
- [31] V. Voevodsky, \mathbb{A}^1 -homotopy theory, in: *Proceedings of the International Congress of Mathematicians*, Vol. I (Berlin, 1998), Doc. Math. (1998), Extra Vol. I, 579–604.
- [32] F. Morel, V. Voevodsky, \mathbb{A}^1 -homotopy theory of schemes, Inst. Hautes Études Sci. Publ. Math. **90** (1999), 45–143.
- [33] A. L. Rosenberg, *Noncommutative schemes*, Compositio Math. **112** (1998), no. 1, 93–125.