

A formula for the Euler characteristic of a poset through the determinant of the order-complement matrix

Pedro José Chocano Feito, Luis Felipe Prieto Martínez

November 2025

Abstract

Given a finite poset P , its zeta matrix \mathbf{Z} encode fundamental incidence-theoretic information about the order structure. In this paper we introduce and study the *order-complement matrix* $\overline{\mathbf{Z}} = \mathbf{J} - \mathbf{Z}$, where \mathbf{J} is the all-ones matrix. We prove a closed formula for its characteristic polynomial and for its determinant, showing that $\det(\overline{\mathbf{Z}}) = (-1)^n \tilde{\chi}(P)$, where $n = |P|$ and $\tilde{\chi}(P)$ is the reduced Euler characteristic of P . This provides a new, unexpectedly simple linear-algebraic expression for the Euler characteristic of a poset, complementing existing determinant formulas for matrices derived from incidence relations.

1 Introduction

A **partially ordered set**, or **poset**, is a set P equipped with a binary relation that is reflexive, antisymmetric, and transitive. This relation, usually denoted by \leq , provides a way to compare certain pairs of elements within the set, though not necessarily all of them.

In a poset (P, \leq) , a **chain** is a subset of elements $C = \{x_0, x_1, \dots, x_k\} \subseteq P$ in which every pair of elements is comparable and arranged in a strictly increasing order, that is,

$$x_0 < x_1 < \dots < x_k.$$

The **length** of a chain is defined as the number of strict comparisons, that is, in the previous case, the length of C is k .

The **Euler characteristic** $\chi(P)$ of a finite poset P , $|P| = n$, is defined as the alternating sum of the number of chains of each length in P :

$$\chi(P) = c_0 - c_1 + c_2 - \dots + (-1)^{n-1} c_{n-1}.$$

2020 Mathematics Subject Classification: 06A11, 06A07.

Keywords: reduced Euler characteristic, poset, determinant, characteristic polynomial.

Equivalently, it can be computed as the alternating sum of the ranks of the homology groups of its order complex. This invariant captures topological information about the poset and is preserved under order-theoretic equivalences such as homotopy equivalence of order complexes. The **reduced Euler characteristic** is defined as

$$\tilde{\chi}(P) = \chi(P) - 1.$$

Given a finite poset (P, \leq) , $P = \{x_1, \dots, x_n\}$, its **zeta matrix** is the $n \times n$ square matrix \mathbf{Z} whose entries are defined by

$$z_{ij} = \begin{cases} 1 & \text{if } x_i \leq x_j \\ 0 & \text{otherwise.} \end{cases}$$

This matrix is invertible.

Let us introduce another matrix which, up to our knowledge, is not so extended in the literature. We define the **order-complement matrix** $\bar{\mathbf{Z}}$ of the poset as

$$\bar{\mathbf{Z}} = \mathbf{J} - \mathbf{Z}, \tag{1}$$

where \mathbf{J} is the $n \times n$ matrix whose entries are all of them equal to 1 and \mathbf{Z} is the zeta matrix. The entries in such matrix are:

$$\bar{z}_{ij} = \begin{cases} 0 & \text{if } x_i \leq x_j \\ 1 & \text{otherwise.} \end{cases} \tag{2}$$

In this paper, we prove the following result:

Theorem 1. *Let (P, \leq) be a poset and let $\mathbf{Z}, \bar{\mathbf{Z}}$ be, respectively, its zeta matrix and its order-complement matrix.*

(a) *The characteristic polynomial of $\bar{\mathbf{Z}}$ is*

$$p(\lambda) = (-1)^n (\lambda + 1)^n - \sum_{k=0}^{n-1} (-1)^{k+n} c_k (\lambda + 1)^{n-1-k},$$

where

$$c_k = \mathbf{1}^T (\mathbf{Z} - \mathbf{I})^k \mathbf{1}$$

is the number of chains of length k and $\mathbf{1}$ is the column vector whose entries are all of them equal to 1.

(b) *Given a finite poset (P, \leq) , let $\bar{\mathbf{Z}}$ be its order-complement matrix. Then*

$$\det \bar{\mathbf{Z}} = (-1)^n \tilde{\chi}(P).$$

2 Some basic results

In this section, we present the main ingredients for the proof of Theorem 1. All of them are well-known in the literature but, for the sake of completeness, we include their proof.

The first result is a consequence of the comment after the definition of *zeta matrix* in [1]:

Lemma 1. *Let (P, \leq) be a finite poset with zeta matrix \mathbf{Z} .*

- (a) *The characteristic polynomial of \mathbf{Z} is $(\lambda - 1)^n$, 1 is the only eigenvalue of \mathbf{Z} , and $\det(\mathbf{Z}) = 1$.*
- (b) *$\text{tr}(\overline{\mathbf{Z}}) = 0$.*

Proof. Any finite poset admits a *linear extension*. A **linear extension** of a finite poset (P, \leq) is a total order \preceq on the elements of P that is compatible with the original partial order, that is, whenever $x \leq y$ in the poset, we also have $x \preceq y$. Equivalently, a linear extension is a listing x_1, x_2, \dots, x_n of the elements of P such that $x_i \leq x_j$ in P implies $i < j$. We refer the reader to [8] for more information.

So we can assume that \mathbf{Z} is upper unitriangular. From an upper triangular matrix, it is easy to obtain the characteristic polynomial, the eigenvalues and the determinant.

Finally,

$$\text{tr}(\overline{\mathbf{Z}}) = \text{tr}(\mathbf{J}) - \text{tr}(\mathbf{Z}) = n - n = 0.$$

□

Associated with a poset, we can also find the **Möbius matrix**, defined as the inverse of the zeta matrix. Its entries form the Möbius function of the poset. For more information, we refer the reader to [7, 8].

Viewing P as a category, the following result is the specialization of *Leinster's Formula* valid for any finite category with Möbius inversion (Section 2, following Definition 2.2 in [5]). Nevertheless, we will give our own proof of this result, based in the Neumann series expansion for the inverse:

Lemma 2. *Let P be a finite poset with Möbius matrix \mathbf{Z}^{-1} . Then the sum of all entries of the Möbius matrix equals the Euler characteristic of P , that is,*

$$\chi(P) = \mathbf{1}^T \mathbf{Z}^{-1} \mathbf{1},$$

where $\mathbf{1}$ is the column vector, whose entries are all of them equal to 1.

Proof. We can assume, again, that the zeta matrix \mathbf{Z} is upper unitriangular.

Write $\mathbf{Z} = \mathbf{I} + \mathbf{N}$. Then \mathbf{N} is strictly upper triangular and therefore nilpotent: there exists $m \leq |P|$ with $\mathbf{N}^m = \mathbf{0}$. Since \mathbf{N} is nilpotent, the inverse of $\mathbf{I} + \mathbf{N}$ is given by the finite Neumann series:

$$\mathbf{Z}^{-1} = \sum_{k=0}^{m-1} (-\mathbf{N})^k = \mathbf{I} - \mathbf{N} + \mathbf{N}^2 - \dots + (-1)^{m-1} \mathbf{N}^{m-1}.$$

The entry in the position (i, j) in \mathbf{N}^k counts the number of strict chains of length k with minimum x_i and maximum x_j . Hence $\mathbf{1}^T \mathbf{N}^k \mathbf{1}$ is the total number of strict chains in P having k inequalities, equivalently chains with $k + 1$ elements. Let c_k denote this number. The Euler characteristic of P is therefore

$$\chi(P) = \mathbf{1}^T \mathbf{Z}^{-1} \mathbf{1} = \sum_{k \geq 0} (-1)^k c_k.$$

□

3 Proof of Theorem 1

Proof of Statement (a)

We can assume, again, that the zeta matrix \mathbf{Z} is upper unitriangular.

First, let us prove that $\lambda = -1$ is never an eigenvalue of $\bar{\mathbf{Z}}$. Suppose that it is, then, from Equation (1), there exists a nontrivial vector $\mathbf{v} \neq \mathbf{0}$ such that

$$(\mathbf{J} - \mathbf{Z})\mathbf{v} = -\mathbf{v}$$

or, equivalently

$$(\mathbf{I} - \mathbf{Z})\mathbf{v} = \mathbf{J}\mathbf{v}.$$

All the entries in $\mathbf{J}\mathbf{v}$ are equal and, since the last row in $(\mathbf{I} - \mathbf{Z})$ is $\mathbf{0}$, all of them must be equal to 0, which is a contradiction.

Let $\mathbf{A} := \lambda \mathbf{I} + \mathbf{Z}$ (it is invertible, since we can assume $\lambda \neq -1$). Observe that $\mathbf{J} = \mathbf{1}\mathbf{1}^T$, where $\mathbf{1}$ is the column vector of all ones. Then, from Equation (1) and using the matrix determinant lemma:

$$p(\lambda) = \det(\bar{\mathbf{Z}} - \lambda \mathbf{I}) = (-1)^n \det(\mathbf{A} - \mathbf{1}\mathbf{1}^T) = (-1)^n \det(\mathbf{A}) \left(1 - \mathbf{1}^T \mathbf{A}^{-1} \mathbf{1}\right).$$

From Lemma 1, Statement (a), we know that $\det(\mathbf{A}) = (\lambda + 1)^n$. So

$$p(\lambda) = (-1)^n (\lambda + 1)^n \left(1 - \mathbf{1}^T \mathbf{A}^{-1} \mathbf{1}\right). \quad (3)$$

Since \mathbf{Z} is upper unitriangular, we can write $\mathbf{Z} = \mathbf{I} + \mathbf{N}$, where \mathbf{N} is strictly upper triangular. Then

$$\mathbf{A} = \lambda \mathbf{I} + \mathbf{Z} = (\lambda + 1)\mathbf{I} + \mathbf{N}.$$

Because \mathbf{N} is nilpotent, we can use the Neumann series to compute the inverse:

$$\mathbf{A}^{-1} = \frac{1}{\lambda + 1} \left(\mathbf{I} + \frac{-\mathbf{N}}{\lambda + 1} \right)^{-1} = \frac{1}{\lambda + 1} \sum_{k=0}^{n-1} \left(-\frac{\mathbf{N}}{\lambda + 1} \right)^k = \sum_{k=0}^{n-1} \frac{(-1)^k \mathbf{N}^k}{(\lambda + 1)^{k+1}}.$$

Substituting in Equation (3), we get the desired result.

Proof of Statement (b)

The determinant of $\overline{\mathbf{Z}}$ can be obtained from the characteristic polynomial $p(\lambda)$ evaluated at $\lambda = 0$, i.e.,

$$\det(\overline{\mathbf{Z}}) = p(0).$$

Substituting $\lambda = 0$ into $p(\lambda)$ in Equation (3), we get

$$p(0) = (-1)^n \left(1 - \mathbf{1}^T \mathbf{Z}^{-1} \mathbf{1}\right).$$

Using Lemma 2 in the previous equation, we obtain the desired result.

4 Brief literature review

To the best of our knowledge, no determinant formula involving the order-complement matrix has appeared before in the literature. Our result provides a new linear-algebraic characterization of the reduced Euler characteristic.

In the study of determinants of matrices naturally associated with posets, there is a classical and well-known work by Ballantine, Frechette, and Little, who consider the matrix

$$\mathfrak{Z} = \mathbf{Z} + \mathbf{Z}^T,$$

where \mathbf{Z} is the zeta matrix of a finite poset P . They derive a combinatorial interpretation of the determinant of \mathfrak{Z} and provide a recursive formula in the case where P is a Boolean algebra [1].

On a broader conceptual level, the notion of Euler characteristic has been extended well beyond posets and topological complexes to more abstract structures such as categories. In his influential paper [5], Leinster defines the Euler characteristic of a finite category by generalizing Rota’s Möbius inversion from posets to categories; his definition works when the category admits both a weighting and a coweighting. He then proves that this new invariant behaves in a manner entirely compatible with classical invariants (e.g., under colimits), thus situating the Euler characteristic in a very general categorical framework. Beyond Leinster’s combinatorial definition, there are further developments exploring alternative approaches: for instance, Berger and Leinster reinterpret the Euler characteristic of a category as the “summation” of a (formally divergent) alternating series; under suitable interpretations, this divergent series can be assigned a finite value that agrees with Leinster’s characteristic [2].

In the realm of posets but still in relation to the Euler characteristic, Noguchi studies the zeta series (a generating function related to the zeta function) under iterative barycentric subdivision [6]. While that work is focused on analytic properties, asymptotic behavior, and zeros of the zeta series rather than on determinants of matrices like $\mathbf{J} - \mathbf{Z}$, it nonetheless shows the richness of connections between zeta functions, poset structure, and topological invariants.

It is worth noting that the first appearance of the so-called order-complement matrix in the literature—though not under this terminology—occurs in [3]. In that paper, the author introduced this matrix independently of the works cited

above and showed, among other topological and combinatorial aspects, that its determinant and rank are homotopy and simple homotopy invariants. The author also raised the question of whether the absolute value of the determinant equals the absolute value of the reduced Euler characteristic, a question mentioned in the introduction following a referee's observation. Later, in [4], the same author analyzes eigenvalues of quadratic forms related to this matrix and their connections with the topological properties of posets.

5 Future Work

Let (P, \leq) be a finite poset. The spectrum of the zeta matrix \mathbf{Z} of the poset does not contain any information (see Lemma 1, Statement (a)). But the determinant formula proved here suggests that the spectrum of the order-complement matrix $\overline{\mathbf{Z}}$ may encode additional invariants of the poset. In particular, it would be natural to investigate how the distribution of eigenvalues of $\overline{\mathbf{Z}}$ reflects combinatorial properties of P , or whether specific families of posets give rise to recognizable spectral patterns. Understanding these eigenvalues could lead to spectral characterizations of classical order-theoretic invariants and provide new tools for comparing posets via linear-algebraic data.

Just as a first step in this direction, let us include the following:

Proposition 1. *Let (P, \leq) be a finite poset and let $\overline{\mathbf{Z}}$ be its order-complement matrix. Then:*

- (a) *-1 is not an eigenvalue of $\overline{\mathbf{Z}}$*
- (b) *The sum of the eigenvalues of $\overline{\mathbf{Z}}$ is always 0.*
- (c) *Suppose that (P, \leq) has a maximum (equiv. a minimum) x . Then 0 is an eigenvalue of $\overline{\mathbf{Z}}$.*

Moreover, if $\overline{\mathbf{Z}}'$ is the order-complement matrix corresponding to the poset $(P \setminus \{x\}, \leq)$, then the spectra of $\mathbf{Z}, \overline{\mathbf{Z}}'$ are related by

$$\sigma(\overline{\mathbf{Z}}') = \sigma\{\overline{\mathbf{Z}}\} \cup \{0\}.$$

- (d) *If (P, \leq) is a totally ordered set, then the only eigenvalue of $\overline{\mathbf{Z}}$ is 0.*

Proof. Statement (a) was already proved in Section 3. Statement (b) is a consequence of Lemma 1 (Statement (b)). In turn, Statement (c) follows from Equation (2) and the Laplace expansion for the determinant. Finally, Statement (d) easily follows from induction in the size of P (using Statement (c)). \square

References

- [1] Ballantine, C. M., Frechette, S. M., & Little, J. B. (2005). Determinants associated to zeta matrices of posets. *Linear Algebra and Its Applications*, 411, 364–370.

- [2] Berger, C., & Leinster, T. (2008). The Euler characteristic of a category as the sum of a divergent series. *Homology, Homotopy and Applications*, 10(1), 41–51.
- [3] Chocano, P. J. (2025). Matrix Invariants as Homotopy Invariants of Finite T_0 -spaces. *arXiv preprint*, arXiv:2506.16861.
- [4] Chocano, P. J. (2026). Eigenvalues and signature of quadratic forms associated with finite topological spaces. *Linear Algebra and its Applications*, 728, 263–282.
- [5] Leinster, T. (2008). The Euler characteristic of a category. *Documenta Mathematica*, 13, 21–49.
- [6] Noguchi, K. (2016). Zeros of the zeta series of a poset and iterated barycentric subdivision. *arXiv preprint*, arXiv:1603.08562 [math.CO].
- [7] Rota, G. C. (1964). On the foundations of combinatorial theory: I. Theory of Möbius functions. In J. N. Srivastava (Ed.), *Classic Papers in Combinatorics* (pp. 332–360). Boston, MA: Birkhäuser.
- [8] Stanley, R. P. (2011). *Enumerative Combinatorics, Volume 1* (2nd ed.). Cambridge: Cambridge University Press.