

Global dynamics in a reaction-diffusion competition model with edge behavior

Kuiyue Liu and Shanshan Chen*

Department of Mathematics, Harbin Institute of Technology
Weihai, Shandong 264209, P. R. China

Abstract

In this paper, we investigate a two-species competition model in a landscape consisting of a finite number of adjacent patches. For the two-patch scenario, by treating edge behavior at the interface as a strategy, it has been shown that there exists an ideal free distribution (IFD) strategy, which is a globally evolutionarily stable strategy. Specifically, when the resident species follows the IFD strategy and the mutant species does not, the mutant species is unable to invade the resident population. Building on this foundation, our work focuses on exploring the dynamics of the system when neither species can adopt the IFD strategy. We demonstrate that if the strategies of both species either exceed or fall below the IFD strategy, the mutant species can outcompete and eliminate the resident species, provided that its strategy is closer to the IFD strategy and its diffusion rates are equal to or slower than those of the resident species. Furthermore, if the strategies of the two species lie on opposite sides of the IFD strategy, the two species can coexist. This result is further extended to the case of an arbitrary but finite number of patches.

Keywords: global dynamics; invasion analysis; reaction-diffusion; edge behavior

MSC 2020: 35K57, 37C65, 92D25, 92D40.

1 Introduction

Reaction-diffusion equations are often the framework of choice to model biological invasions. For two competing species, it was demonstrated in [15, 24] that the species

*Corresponding author, Email: chenSS@hit.edu.cn

with a slower diffusion rate could outcompete and eliminate its faster counterpart if they are identical except for their diffusion rates, and one can also refer to [25, 34, 37] and references therein for the weak competition case in spatially heterogeneous environments.

As a result of habitat fragmentation, the landscape may comprise multiple patches. By ignoring the spatial structure within each patch, discrete patch models (systems of ordinary differential equations) can be used to model population dynamics. The researchers also addressed questions similar to those studied in reaction-diffusion models. For instance, one can refer to [9, 10, 11, 20, 24, 35, 54] and references therein for the dynamics of two competing species.

Taking into account the spatial structure of patches, numerous experiments have illustrated that there exists edge behavior of individuals, and individuals may have movement preference at interfaces between patches [46, 47, 48, 49, 50]. The continuous density conditions at interfaces were first proposed in [45] and [52] for finite and infinite two-patch landscapes, respectively. Further and related studies can be found in [12, 17, 39] and references therein.

The habitat preference of individuals can lead to density discontinuities at interfaces. Building on this observation and the work of Ovaskainen and Cornell [44], Maciel and Lutscher [41] first introduced the discontinuous density conditions and proposed the following single-species reaction-diffusion model with two patches:

$$\begin{cases} (u_1)_t = d_1(u_1)_{xx} + r_1 u_1 \left(1 - \frac{u_1}{k_1}\right), & 0 < x < x_1, \\ (u_2)_t = d_2(u_2)_{xx} + r_2 u_2 \left(1 - \frac{u_2}{k_2}\right), & x_1 < x < L, \\ u_2(x_1^+, t) = p u_1(x_1^-, t), \quad d_2(u_2)_x(x_1^+, t) = d_1(u_1)_x(x_1^-, t), \\ (u_1)_x(0^+) = (u_2)_x(L^-) = 0. \end{cases} \quad (1.1)$$

(They also investigated an infinite landscape consisting of periodically alternating patches of two types.) Here u_i denotes the density of the species in patch i , d_i denotes the diffusion coefficient in patch i , and r_i and k_i represent the intrinsic growth rate and carrying capacity of the species, respectively. Moreover, $p := (d_1/d_2)[\alpha/(1-\alpha)]$, where α characterizes the patch preference at the interface $x = x_1$. The density of the species is discontinuous at the interface if $p \neq 1$. Zaker et al. [59] showed that edge behavior influences the shape of the steady state of model (1.1), and total biomass could exceed total carrying capacity under certain conditions. Then Tchouaga and Lutscher [57] studied another single-species model with Allee effect and found that bi-stability can occur, characterized by two nonzero stable steady states. Moreover, one can refer to

[1, 21, 22, 23, 42, 53] for propagation phenomena in reaction-diffusion models with edge behavior.

There are also significant findings on two-species competition models similar to (1.1). For a competition model in a one-dimensional infinite landscape, Maciel and Lutscher [43] found that diffusion rate and patch preference influence competitive outcomes through multi-scale analysis and homogenization. Then Maciel et al. [40] considered the following model with $n = 2$:

$$\begin{cases} (u_i)_t = d_i(u_i)_{xx} + r_i u_i \left(1 - \frac{u_i + v_i}{k_i}\right), & x \in \Omega_i, t > 0, i = 1, 2, \dots, n, & (1.2a) \\ (v_i)_t = \hat{d}_i(v_i)_{xx} + r_i v_i \left(1 - \frac{u_i + v_i}{k_i}\right), & x \in \Omega_i, t > 0, i = 1, 2, \dots, n, & (1.2b) \\ u_{i+1}(x_i^+, t) = p_i u_i(x_i^-, t), d_{i+1}(u_{i+1})_x(x_i^+, t) = d_i(u_i)_x(x_i^-, t), & t > 0, & (1.2c) \\ v_{i+1}(x_i^+, t) = \hat{p}_i v_i(x_i^-, t), \hat{d}_{i+1}(v_{i+1})_x(x_i^+, t) = \hat{d}_i(v_i)_x(x_i^-, t), & t > 0, & (1.2d) \\ (u_1)_x(0^+, t) = (u_n)_x(L^-, t) = (v_1)_x(0^+, t) = (v_n)_x(L^-, t) = 0, & t > 0. & (1.2e) \end{cases}$$

Here the landscape is finite and composed of n adjacent patches $\{\Omega_i\}_{i=1}^n$, where $\Omega_i := (x_{i-1}, x_i)$ represents the i -th patch with length $x_i - x_{i-1} > 0$ for $i = 1, \dots, n$; $x_0 = 0$ and $x_n = L$ denote the boundaries of the landscape and the no-flux boundary conditions are imposed at $x = 0, L$; u_i and v_i denote the densities of two competing species in patch i ; and $d_i, \hat{d}_i > 0$ denote the diffusion rates of the two species in patch i . The two species admit the same per capita growth rate with intrinsic growth rate $r_i > 0$ and carrying capacity $k_i > 0$ in each patch. At the interface $x = x_i$ ($i = 1, \dots, n-1$), the discontinuous density conditions (if $p_i \neq 1$) and balanced flux conditions are imposed for the two species, where the superscripts \pm denote one-sided limits from the right and left, respectively, and $p_i, \hat{p}_i > 0$ are composite parameters with

$$p_i := \frac{\alpha_i}{1 - \alpha_i} \frac{d_i}{d_{i+1}}, \quad \hat{p}_i := \frac{\hat{\alpha}_i}{1 - \hat{\alpha}_i} \frac{\hat{d}_i}{\hat{d}_{i+1}}, \quad i = 1, \dots, n-1.$$

Here $\alpha_i, \hat{\alpha}_i \in (0, 1)$ represent the patch preference of the two species at the interface $x = x_i$. Specifically, α_i (resp. $\hat{\alpha}_i$) is the probability that the species **u** (resp. **v**) at the interface x_i moves to patch $i+1$, and $1 - \alpha_i$ (resp. $1 - \hat{\alpha}_i$) is the probability that the species **u** (resp. **v**) at the interface x_i moves to patch i . Denote

$$\bar{\mathbf{k}} = \left(\frac{k_2}{k_1}, \dots, \frac{k_n}{k_{n-1}} \right), \quad \mathbf{p} = (p_1, \dots, p_{n-1}), \quad \hat{\mathbf{p}} = (\hat{p}_1, \dots, \hat{p}_{n-1}). \quad (1.3)$$

For the two-patch case ($n = 2$), Maciel et al. [40] showed that $\bar{\mathbf{k}}$ is an IFD strategy. Moreover, if the resident species **u** adopts this IFD strategy, the mutant species **v**

with a different strategy ($\hat{\mathbf{p}} \neq \bar{\mathbf{k}}$) cannot successfully invade. This result can be easily extended to the case where n is finite but arbitrary.

A natural question arise: what is the dynamics of model (1.2) if neither \mathbf{u} nor \mathbf{v} can adopt the IFD strategy? In this paper, we aim to address this question. For the two-patch scenario ($n = 2$), $\bar{\mathbf{k}} = k_2/k_1$, $\mathbf{p} = p_1$ and $\hat{\mathbf{p}} = \hat{p}_1$, and we show that the following results hold:

- (i) If p_1 and \hat{p}_1 lie on the same side of IFD strategy (i.e., $p_1, \hat{p}_1 > k_2/k_1$ or $k_2/k_1 > p_1, \hat{p}_1$), the mutant species \mathbf{v} can outcompete and eliminate the resident species \mathbf{u} if its strategy \hat{p} is closer to k_2/k_1 and its diffusion rates are equal to or slower than those of the resident species;
- (ii) If p_1 and \hat{p}_1 lie on opposite sides of IFD strategy (i.e., $p_1 > k_2/k_1 > \hat{p}_1$ or $\hat{p}_1 > k_2/k_1 > p_1$), the two species can coexist.

The general form of this result for the case $n \geq 2$ is presented in Theorem 3.6.

The above results can be understood within the framework of adaptive dynamics. Detailed discussions on adaptive dynamics can be found in [13, 14, 19, 32, 56] and references therein. One key concept is a global (resp. local) evolutionarily stable strategy (ESS): a strategy is called a global (resp. local) ESS if a population adopting it cannot be invaded by a small population of mutants using a different (resp. different nearby) strategy [7, 56] (see Definition 2.8). A related concept is a global (resp. local) neighborhood invader strategy (NIS): a strategy is called a global (resp. local) NIS if a small population of mutants adopting it can always invade a population using a different (resp. different nearby) strategy [4, 7] (see Definition 2.9). Finally, a strategy is called a local convergent stable strategy (CSS) if, for any resident strategy in its neighborhood, a small population of mutants with a strategy closer to it can always invade. Moreover, the definition of a global CSS follows by removing the neighborhood restriction (see Definition 2.10).

In the two-patch scenario, let the composite parameters p_1 and \hat{p}_1 be the strategies of \mathbf{u} and \mathbf{v} , respectively, treating all other parameters as identical to isolate their effects. The above-mentioned result in [40] implies that the IFD strategy is a global ESS and NIS. Our result (i) strengthens this by proving that the IFD strategy is also a global CSS. Moreover, result (ii) parallels findings in reaction-diffusion-advection competition models [7], illustrating that two species can coexist when their strategies are on opposite sides of the IFD strategy.

The rest of the paper is organized as follows. Sect. 2 is devoted to the invasion analysis for model (1.2), and the local stability of the semi-trivial steady state is obtained. In Sect. 3, we studied the global dynamics of model (1.2). For simplicity, we list some notations used throughout the paper. For vectors $\mathbf{w} = (w_1, \dots, w_n) \in \mathbb{R}^n$ and $\mathbf{z} = (z_1, \dots, z_n) \in \mathbb{R}^n$, we denote

$$[\mathbf{w}]_\ell := (w_1, \dots, w_\ell), \quad [\mathbf{w}]_\ell^\# := (w_{n-\ell+1}, \dots, w_n) \quad \text{for } 1 \leq \ell \leq n, \quad (1.4)$$

and write

$$\mathbf{w} \gg (\text{resp. } \geq) \mathbf{z} \text{ if } w_i > (\text{resp. } \geq) z_i \text{ for } i = 1, \dots, n, \quad (1.5)$$

and $\mathbf{w} > \mathbf{z}$ if $\mathbf{w} \geq \mathbf{z}$ and $\mathbf{w} \neq \mathbf{z}$. Denote by $(u)_x$ the derivative of u with respect to x .

2 Local dynamics and invasion analysis

In each patch Ω_i ($i = 1, 2, \dots, n$), we view species u_i as the resident species and v_i as the mutant species. It follows from Theorem A.1 in Appendix A that (1.2) admits only two semi-trivial steady states $(\mathbf{u}^*, \mathbf{0})$ and $(\mathbf{0}, \mathbf{v}^*)$, where

$$\mathbf{u}^* = (u_1^*, \dots, u_n^*) \in C^2(\overline{\Omega}_1) \times \dots \times C^2(\overline{\Omega}_n)$$

is the unique positive solution of

$$\begin{cases} d_i(u_i^*)_{xx} + r_i u_i^* \left(1 - \frac{u_i^*}{k_i}\right) = 0, & x \in \Omega_i, \quad i = 1, 2, \dots, n, \end{cases} \quad (2.1a)$$

$$\begin{cases} u_{i+1}^*(x_i^+) = p_i u_i^*(x_i^-), \quad d_{i+1}(u_{i+1}^*)_x(x_i^+) = d_i(u_i^*)_x(x_i^-), \end{cases} \quad (2.1b)$$

$$\begin{cases} (u_1^*)_x(0^+) = (u_n^*)_x(L^-) = 0. \end{cases} \quad (2.1c)$$

In this section, we analyze the local dynamics of model (1.2) and establish a sufficient condition under which species $\mathbf{v} = (v_1, \dots, v_n)$ can invade when rare. Specifically, the mutant species $\mathbf{v} = (v_1, \dots, v_n)$ can (resp. cannot) invade when rare if the semi-trivial steady state $(\mathbf{u}^*, \mathbf{0})$ is unstable (resp. stable). The stability of $(\mathbf{u}^*, \mathbf{0})$ is determined by the principal eigenvalue of the following eigenvalue problem:

$$\begin{cases} \hat{d}_i(\phi_i)_{xx} + r_i \left(1 - \frac{u_i^*}{k_i}\right) \phi_i = \lambda \phi_i, & x \in \Omega_i, \quad i = 1, 2, \dots, n, \end{cases} \quad (2.2a)$$

$$\begin{cases} \phi_{i+1}(x_i^+) = \hat{p}_i \phi_i(x_i^-), \quad \hat{d}_{i+1}(\phi_{i+1})_x(x_i^+) = \hat{d}_i(\phi_i)_x(x_i^-), \end{cases} \quad (2.2b)$$

$$\begin{cases} (\phi_1)_x(0^+) = (\phi_n)_x(L^-) = 0. \end{cases} \quad (2.2c)$$

Proposition C.9 in Appendix C guarantees the existence of the principal eigenvalue for problem (2.2).

Lemma 2.1. *Let $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k})$ be the principal eigenvalue of the eigenvalue problem (2.2). Then, the semi-trivial steady state $(\mathbf{u}^*, \mathbf{0})$ is stable if $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) < 0$ and unstable if $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) > 0$.*

(Here we abuse the notation by denoting $\mathbf{1} - \mathbf{u}^*/\mathbf{k} := (1 - u_1^*/k_1, \dots, 1 - u_n^*/k_n)$.)

Proof. Model (1.2) can be rewritten as the abstract evolution equation (C.8) in Appendix C. Then, by Remark C.7, a steady state of (1.2) (or (C.8)) is asymptotically stable (resp., unstable) if it is linearly stable (resp., linearly unstable). Accordingly, we consider the eigenvalue problem associated with the semi-trivial steady state $(\mathbf{u}^*, \mathbf{0})$:

$$\begin{cases} d_i(\psi_i)_{xx} + r_i \left(1 - \frac{2u_i^*}{k_i}\right) \psi_i - \frac{r_i u_i^*}{k_i} \phi_i = \mu \psi_i, & x \in \Omega_i, \quad i = 1, 2, \dots, n, \\ \hat{d}_i(\phi_i)_{xx} + r_i \left(1 - \frac{u_i^*}{k_i}\right) \phi_i = \mu \phi_i, & x \in \Omega_i, \quad i = 1, 2, \dots, n, \\ \psi_{i+1}(x_i^+) = p_i \psi_i(x_i^-), \quad d_{i+1}(\psi_{i+1})_x(x_i^+) = d_i(\psi_i)_x(x_i^-), \\ \phi_{i+1}(x_i^+) = \hat{p}_i \phi_i(x_i^-), \quad \hat{d}_{i+1}(\phi_{i+1})_x(x_i^+) = \hat{d}_i(\phi_i)_x(x_i^-), \\ (\phi_1)_x(0^+) = (\phi_n)_x(L^-) = (\psi_1)_x(0^+) = (\psi_n)_x(L^-) = 0. \end{cases} \quad (2.3)$$

We first show that

$$\operatorname{Re} \mu \leq \max\{\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}), \lambda_1(\mathbf{p}, \mathbf{d}, \mathbf{1} - 2\mathbf{u}^*/\mathbf{k})\}. \quad (2.4)$$

Here, $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k})$ is the principal eigenvalue of (2.2), and $\lambda_1(\mathbf{p}, \mathbf{d}, \mathbf{1} - 2\mathbf{u}^*/\mathbf{k})$ corresponds to that of

$$\begin{cases} d_i(\psi_i)_{xx} + r_i \left(1 - \frac{2u_i^*}{k_i}\right) \psi_i = \lambda \psi_i, & x \in \Omega_i, \quad i = 1, 2, \dots, n, \\ \psi_{i+1}(x_i^+) = p_i \psi_i(x_i^-), \quad d_{i+1}(\psi_{i+1})_x(x_i^+) = d_i(\psi_i)_x(x_i^-), \\ (\psi_1)_x(0^+) = (\psi_n)_x(L^-) = 0. \end{cases} \quad (2.5)$$

The existence of the principal eigenvalue for (2.5) is also a consequence of Proposition C.9 in Appendix C. Furthermore, this proposition shows that $\operatorname{Re} \mu \leq \lambda(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k})$ when $\Phi = (\phi_1, \dots, \phi_n) \not\equiv \mathbf{0}$. If $\Phi \equiv \mathbf{0}$, then $\Psi = (\psi_1, \dots, \psi_n) \not\equiv \mathbf{0}$, and we obtain that $\operatorname{Re} \mu \leq \lambda(\mathbf{p}, \mathbf{d}, \mathbf{1} - 2\mathbf{u}^*/\mathbf{k})$. Therefore, (2.4) holds.

Now we prove that $\lambda_1(\mathbf{p}, \mathbf{d}, \mathbf{1} - 2\mathbf{u}^*/\mathbf{k}) < 0$. Multiplying the first equation of (2.5) with $(\prod_{\ell=i}^n p_\ell) u_i^*$, integrating the result over Ω_i and summing them over all i , we have

$$\begin{aligned} & \lambda_1 \left(\mathbf{p}, \mathbf{d}, \mathbf{1} - \frac{2\mathbf{u}^*}{\mathbf{k}} \right) \left(\sum_{i=1}^n \left(\prod_{\ell=i}^n p_\ell \right) \int_{x_{i-1}}^{x_i} \psi_i u_i^* dx \right) \\ &= \sum_{i=1}^n d_i \left(\prod_{\ell=i}^n p_\ell \right) \int_{x_{i-1}}^{x_i} (\psi_i)_{xx} u_i^* dx + \sum_{i=1}^n \left(\prod_{\ell=i}^n p_\ell \right) \int_{x_{i-1}}^{x_i} r_i \psi_i u_i^* \left(1 - \frac{2u_i^*}{k_i} \right) dx. \end{aligned} \quad (2.6)$$

Similarly, we deduce from (2.1a) that

$$0 = \sum_{i=1}^n d_i \left(\prod_{\ell=i}^n p_\ell \right) \int_{x_{i-1}}^{x_i} (u_i^*)_{xx} \psi_i dx + \sum_{i=1}^n \left(\prod_{\ell=i}^n p_\ell \right) \int_{x_{i-1}}^{x_i} r_i \psi_i u_i^* \left(1 - \frac{u_i^*}{k_i} \right) dx. \quad (2.7)$$

Subtracting (2.7) from (2.6) and applying integration by parts yields

$$\lambda_1(\mathbf{p}, \mathbf{d}, \mathbf{1} - 2\mathbf{u}^*/\mathbf{k}) \left(\sum_{i=1}^n \left(\prod_{\ell=i}^n p_\ell \right) \int_{x_{i-1}}^{x_i} \psi_i u_i^* dx \right) = - \sum_{i=1}^n \left(\prod_{\ell=i}^n p_\ell \right) \int_{x_{i-1}}^{x_i} \frac{r_i u_i^*}{k_i} \psi_i dx < 0.$$

This implies that $\lambda_1(\mathbf{p}, \mathbf{d}, \mathbf{1} - 2\mathbf{u}^*/\mathbf{k}) < 0$. The desired result follows from this and the fact that $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k})$ is also an eigenvalue of (2.3). \square

Lemma 2.2. *Let $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k})$ be the principal eigenvalue of (2.2) with corresponding eigenfunction $\Phi = (\phi_1(x), \dots, \phi_n(x))$, where $\phi_i > 0$ in $\bar{\Omega}_i$ for each $i = 1, \dots, n$. Then the following identity holds:*

$$\begin{aligned} & \lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) \left(\sum_{i=1}^n \left(\prod_{\ell=i}^n p_\ell \right) \int_{x_{i-1}}^{x_i} \phi_i u_i^* dx \right) \\ &= \sum_{i=1}^{n-1} \left(\prod_{\ell=i+1}^n p_\ell \right) d_i (\hat{p}_i - p_i) (u_i^*)_x(x_i^-) \phi_i(x_i^-) + \sum_{i=1}^n \left(\prod_{\ell=i}^n p_\ell \right) (d_i - \hat{d}_i) \int_{x_{i-1}}^{x_i} (\phi_i)_x (u_i^*)_x dx. \end{aligned} \quad (2.8)$$

Here we define $p_n = 1$.

Proof. Multiplying (2.2a) with $(\prod_{\ell=i}^n p_\ell) u_i^*$, integrating the result over Ω_i and summing them over all i , we have

$$\begin{aligned} & \lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) \left(\sum_{i=1}^n \left(\prod_{\ell=i}^n p_\ell \right) \int_{x_{i-1}}^{x_i} \phi_i u_i^* dx \right) \\ &= \sum_{i=1}^n \hat{d}_i \left(\prod_{\ell=i}^n p_\ell \right) \int_{x_{i-1}}^{x_i} (\phi_i)_{xx} u_i^* dx + \sum_{i=1}^n \left(\prod_{\ell=i}^n p_\ell \right) \int_{x_{i-1}}^{x_i} r_i \phi_i u_i^* \left(1 - \frac{u_i^*}{k_i} \right) dx. \end{aligned} \quad (2.9)$$

Similarly, we deduce from (2.1a) that

$$0 = \sum_{i=1}^n d_i \left(\prod_{\ell=i}^n p_\ell \right) \int_{x_{i-1}}^{x_i} (u_i^*)_{xx} \phi_i dx + \sum_{i=1}^n \left(\prod_{\ell=i}^n p_\ell \right) \int_{x_{i-1}}^{x_i} r_i \phi_i u_i^* \left(1 - \frac{u_i^*}{k_i} \right) dx. \quad (2.10)$$

Subtracting (2.10) from (2.9) and integrating by parts, we can obtain the desired result from (2.1b)-(2.1c) and (2.2b)-(2.2c). \square

The following result is a direct consequence of Lemma 2.2.

Lemma 2.3. Suppose that $\mathbf{d} = \hat{\mathbf{d}}$, and let $\bar{\mathbf{k}}$ be defined in (1.3). Then the following statements hold for model (1.2):

- (i) If $\mathbf{p} \gg \bar{\mathbf{k}}$, the semi-trivial steady state $(\mathbf{u}^*, \mathbf{0})$ is unstable for $\mathbf{p} \gg \hat{\mathbf{p}}$ and stable for $\hat{\mathbf{p}} \gg \mathbf{p}$;
- (ii) If $\bar{\mathbf{k}} \gg \mathbf{p}$, the semi-trivial steady state $(\mathbf{u}^*, \mathbf{0})$ is unstable for $\hat{\mathbf{p}} \gg \mathbf{p}$ and stable for $\mathbf{p} \gg \hat{\mathbf{p}}$.

Proof. Since $\mathbf{d} = \hat{\mathbf{d}}$, it follows from Lemma 2.2 that

$$\begin{aligned} \lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) & \left(\sum_{i=1}^n \left(\prod_{\ell=i}^n p_\ell \right) \int_{x_{i-1}}^{x_i} \phi_i u_i^* dx \right) \\ & = \sum_{i=1}^{n-1} \left(\prod_{\ell=i+1}^n p_\ell \right) d_i (\hat{p}_i - p_i) (u_i^*)_x(x_i^-) \phi_i(x_i^-). \end{aligned}$$

Here $p_n = 1$ and $(\phi_1(x), \dots, \phi_n(x))$ is the eigenfunction corresponding to $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k})$, where $\phi_i > 0$ in $\bar{\Omega}_i$ for all $i = 1, \dots, n$. Thus, the desired result follows from this and Lemma A.2. \square

Lemma 2.4. Suppose that $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) = 0$, where $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k})$ is the principal eigenvalue of (2.2) and $\Phi = (\phi_1(x), \dots, \phi_n(x))$ is the corresponding eigenfunction with $\phi_i(x) > 0$ for $x \in \bar{\Omega}_i$ and $i = 1, \dots, n$. Let $x_{\ell^*-1} \leq a < x_{\ell^*}$ and $x_{\ell^*-1} < b \leq x_{\ell^*}$ for some ℓ_* and ℓ^* with $1 \leq \ell_* < \ell^* \leq n$. Then the following two identities hold:

$$\begin{aligned} 0 & = \sum_{i=\ell_*}^{\ell^*-1} \left(\prod_{\ell=i+1}^{\ell^*} p_\ell \right) d_i (\hat{p}_i - p_i) (u_i^*)_x(x_i^-) \phi_i(x_i^-) + \hat{d}_{\ell^*} p_{\ell^*} (\phi_{\ell^*})_x(b^-) u_{\ell^*}^*(b^-) \\ & \quad - \hat{d}_{\ell_*} \left(\prod_{\ell=\ell_*}^{\ell^*} p_\ell \right) (\phi_{\ell_*})_x(a^+) u_{\ell_*}^*(a^+) + \sum_{i=\ell_*}^{\ell^*} \left(\prod_{\ell=i}^{\ell^*} p_\ell \right) (d_i - \hat{d}_i) \int_{\Omega_i \cap (a,b)} (\phi_i)_x (u_i^*)_x dx \quad (2.11) \\ & \quad - d_{\ell^*} p_{\ell^*} (u_{\ell^*}^*)_x(b^-) \phi_{\ell^*}(b^-) + d_{\ell_*} \left(\prod_{\ell=\ell_*}^{\ell^*} p_\ell \right) (u_{\ell_*}^*)_x(a^+) \phi_{\ell_*}(a^+), \end{aligned}$$

and

$$\begin{aligned} 0 & = \sum_{i=\ell_*}^{\ell^*-1} \frac{\hat{d}_i (k_i \hat{p}_i - k_{i+1}) (\phi_i)_x(x_i^-)}{\hat{p}_i \phi_i(x_i^-)} + \frac{\hat{d}_{\ell^*} k_{\ell^*} (\phi_{\ell^*})_x(b^-)}{\phi_{\ell^*}(b^-)} - \frac{\hat{d}_{\ell_*} k_{\ell_*} (\phi_{\ell_*})_x(a^+)}{\phi_{\ell_*}(a^+)} \\ & \quad + \sum_{i=\ell_*}^{\ell^*} \int_{\Omega_i \cap (a,b)} \left[\frac{\hat{d}_i k_i (\phi_i)_x^2}{\phi_i^2} + \frac{r_i (k_i - u_i^*)^2}{k_i} \right] dx - d_{\ell^*} (u_{\ell^*}^*)_x(b^-) + d_{\ell_*} (u_{\ell_*}^*)_x(a^+). \end{aligned} \quad (2.12)$$

Here we define $p_n = 1$.

Proof. Letting $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) = 0$ and replacing $(1, n)$ by (l_*, l^*) in the proof of Lemma 2.2, we can obtain that (2.11) holds. Now we show that (2.12) holds, and the proof is inspired by [40]. Dividing (2.2a) by ϕ_i/k_i , integrating the result over $\Omega_i \cap (a, b)$ and summing them from $i = \ell_*$ to $i = \ell^*$, we see from integration by parts and (2.2b) that

$$\begin{aligned} 0 = & \sum_{i=\ell_*}^{\ell^*-1} \frac{\hat{d}_i(k_i \hat{p}_i - k_{i+1})(\phi_i)_x(x_i^-)}{\hat{p}_i \phi_i(x_i^-)} + \frac{\hat{d}_{\ell^*} k_{\ell^*} (\phi_{\ell^*})_x(b^-)}{\phi_{\ell^*}(b^-)} - \frac{\hat{d}_{\ell_*} k_{\ell_*} (\phi_{\ell_*})_x(a^+)}{\phi_{\ell_*}(a^+)} \\ & + \sum_{i=\ell_*}^{\ell^*} \int_{\Omega_i \cap (a, b)} \left[\frac{\hat{d}_i k_i (\phi_i)_x^2}{\phi_i^2} + r_i k_i \left(1 - \frac{u_i^*}{k_i} \right) \right] dx. \end{aligned} \quad (2.13)$$

In addition, integrating (2.1a) over $\Omega_i \cap (a, b)$ and summing them from $i = \ell_*$ to $i = \ell^*$ gives

$$\sum_{i=\ell_*}^{\ell^*} \int_{\Omega_i \cap (a, b)} r_i u_i^* \left(1 - \frac{u_i^*}{k_i} \right) dx + d_{\ell^*} (u_{\ell^*})_x(b^-) - d_{\ell_*} (u_{\ell_*})_x(a^+) = 0. \quad (2.14)$$

Subtracting (2.14) from (2.13), we see that (2.12) holds. \square

Then we obtain two results on the signs of $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k})$. The first one is as follows.

Lemma 2.5. *Let $\bar{\mathbf{k}}$ be defined in (1.3) and let $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k})$ be the principal eigenvalue of (2.2). Then the following two statements hold:*

- (i) *If $\mathbf{p} \gg \bar{\mathbf{k}}$, then $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) > 0$ for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_1$ and $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) < 0$ for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_2$, where*

$$\mathcal{L}_1 := \{(\hat{\mathbf{p}}, \hat{\mathbf{d}}) : \mathbf{p} \gg \hat{\mathbf{p}}, \mathbf{d} \geq \hat{\mathbf{d}}\}, \quad \mathcal{L}_2 := \{(\hat{\mathbf{p}}, \hat{\mathbf{d}}) : \hat{\mathbf{p}} \gg \mathbf{p}, \hat{\mathbf{d}} \geq \mathbf{d}\}; \quad (2.15)$$

- (ii) *If $\bar{\mathbf{k}} \gg \mathbf{p}$, then $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) > 0$ for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{S}_1$ and $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) < 0$ for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{S}_2$, where*

$$\mathcal{S}_1 := \{(\hat{\mathbf{p}}, \hat{\mathbf{d}}) : \hat{\mathbf{p}} \gg \mathbf{p}, \mathbf{d} \geq \hat{\mathbf{d}}\}, \quad \mathcal{S}_2 := \{(\hat{\mathbf{p}}, \hat{\mathbf{d}}) : \mathbf{p} \gg \hat{\mathbf{p}}, \hat{\mathbf{d}} \geq \mathbf{d}\}. \quad (2.16)$$

Proof. We only prove (i), and (ii) can be treated similarly. By Lemma 2.3, we see that $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) > 0$ (resp. < 0) for $\mathbf{p} \gg \hat{\mathbf{p}}$ (resp. $\hat{\mathbf{p}} \gg \mathbf{p}$) if $\hat{\mathbf{d}} = \mathbf{d}$. Note from Proposition C.9 in Appendix C that $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k})$ is continuous with respect to $\hat{\mathbf{d}}$. Noting that \mathcal{L}_i ($i = 1, 2$) are convex, it suffices to show that $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) \neq 0$ for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_1 \cup \mathcal{L}_2$. Suppose to the contrary that $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) = 0$ for some

$(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_1 \cup \mathcal{L}_2$ and denote the corresponding eigenfunction by $\phi = (\phi_1(x), \dots, \phi_n(x))$ with $\phi_i(x) > 0$ for $x \in \bar{\Omega}_i$ and $i = 1, \dots, n$. Then we will obtain a contradiction for each of the following two cases: (i₁) $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_1$; (i₂) $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_2$. For simplicity, we only obtain a contradiction for (i₁), and (i₂) can be treated similarly.

By Lemma A.2, $u_1^*(0^+) < k_1$. Then we deduce (by (2.2a)) that $(\phi_1)_{xx}(0^+) < 0$. This combined with $(\phi_1)_x(0^+) = 0$ implies that $(\phi_1)_x(x) < 0$ for $x \in (0, \epsilon)$ with $0 < \epsilon \ll 1$. Since $(\phi_1)_x(L^-) = 0$, it follows that

$$y_1 = \inf\{y > 0 : \exists 1 \leq \ell \leq n \text{ s.t. } y \in (x_{\ell-1}, x_\ell] \text{ and } (\phi_\ell)_x(y^-) = 0\}$$

is well-defined with $y_1 \in (x_{\ell_1-1}, x_{\ell_1}]$ for some ℓ_1 with $1 \leq \ell_1 \leq n$. Then

$$(\phi_{\ell_1})_x(y_1^-) = 0 \text{ and } (\phi_i)_x(x) < 0 \text{ for } x \in \Omega_i \cap (0, y_1) \text{ with } i = 1, \dots, \ell_1. \quad (2.17)$$

Substituting $a = 0$, $b = y_1$, $\ell_* = 1$ and $\ell^* = \ell_1$ into (2.11), we obtain that

$$\begin{aligned} 0 = & \sum_{i=1}^{\ell_1-1} \left(\prod_{\ell=i+1}^{\ell_1} p_\ell \right) d_i (\hat{p}_i - p_i) (u_i^*)_x(x_i^-) \phi_i(x_i^-) + \hat{d}_{\ell_1} p_{\ell_1} (\phi_{\ell_1})_x(y_1^-) u_{\ell_1}^*(y_1^-) \\ & + \sum_{i=1}^{\ell_1} \left(\prod_{\ell=i}^{\ell_1} p_\ell \right) (d_i - \hat{d}_i) \int_{\Omega_i \cap (0, y_1)} (\phi_i)_x(u_i^*)_x dx - d_{\ell_1} p_{\ell_1} (u_{\ell_1}^*)_x(y_1^-) \phi_{\ell_1}(y_1^-) > 0, \end{aligned} \quad (2.18)$$

where we have used $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_1$, (2.17) and Lemma A.2 in the last step. This leads to a contradiction and completes the proof. \square

In the following, we show that the sign of $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k})$ can be determined without additional assumption on $\mathbf{d}, \hat{\mathbf{d}}$ if \mathbf{p} and $\hat{\mathbf{p}}$ are opposite sides of $\bar{\mathbf{k}}$ with respect to the order defined in (1.5).

Lemma 2.6. *Let $\bar{\mathbf{k}}$ be defined in (1.3) and let $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k})$ be the principal eigenvalue of (2.2). Then the following statements hold:*

(i) *If $\mathbf{p} \gg \bar{\mathbf{k}}$, then $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) > 0$ for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_3$, where*

$$\mathcal{L}_3 := \{(\hat{\mathbf{p}}, \hat{\mathbf{d}}) : \bar{\mathbf{k}} \gg \hat{\mathbf{p}}\}; \quad (2.19)$$

(ii) *If $\bar{\mathbf{k}} \gg \mathbf{p}$, then $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) > 0$ for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{S}_3$, where*

$$\mathcal{S}_3 := \{(\hat{\mathbf{p}}, \hat{\mathbf{d}}) : \hat{\mathbf{p}} \gg \bar{\mathbf{k}}\}. \quad (2.20)$$

Proof. We only prove (i), and (ii) can be treated similarly. Similar to Lemma 2.5, it suffices to show that $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) \neq 0$ for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_3$. Suppose to the contrary that $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) = 0$ for some $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_3$ and denote the corresponding eigenfunction by $\phi = (\phi_1, \dots, \phi_n)$ with $\phi_i(x) > 0$ for $x \in \bar{\Omega}_i$ and $i = 1, \dots, n$.

For $x \in (0, x_1]$, integrating (2.2a) over $(0, x)$ and noticing that $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) = 0$, we obtain that

$$\hat{d}_1(\phi_1)_x(x^-) = - \int_0^x r_1 \phi_1 \left(1 - \frac{u_1^*}{k_1}\right) dx < 0,$$

where we have used Lemma A.2 (i) in the last step. This combined with (2.2c) implies that

$$y_0 = \inf\{y : \exists 2 \leq \ell \leq n \text{ s.t. } y \in (x_{\ell-1}, x_\ell] \text{ and } (\phi_\ell)_x(y^-) = 0\}$$

is well-defined with $x_{\ell_0-1} < y_0 \leq x_{\ell_0}$ for some ℓ_0 with $2 \leq \ell_0 \leq n$. Thus,

$$(\phi_{\ell_0})_x(y_0^-) = 0, \quad (\phi_\ell)_x(x^-) < 0 \text{ for } x \in (x_{\ell-1}, x_\ell] \cap (0, y_0), \quad \ell = 1, \dots, \ell_0. \quad (2.21)$$

Noticing that $\lambda_1(\hat{\mathbf{p}}, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k}) = 0$ and substituting $a = 0$, $b = y_0$, $\ell_* = 1$ and $\ell^* = \ell_0$ into (2.12), we have

$$\begin{aligned} 0 &= \sum_{i=1}^{\ell_0-1} \frac{\hat{d}_i(k_i \hat{p}_i - k_{i+1})(\phi_i)_x(x_i^-)}{\hat{p}_i \phi_i(x_i^-)} + d_1(u_1^*)_x(0^+) \\ &\quad + \sum_{i=1}^{\ell_0} \int_{\Omega_i \cap (0, y_0)} \left[\frac{\hat{d}_i k_i (\phi_i)_x^2}{\phi_i^2} + \frac{r_i (k_i - u_i^*)^2}{k_i} \right] dx - d_{\ell_0}(u_{\ell_0}^*)_x(y_0^-) \\ &\quad + \frac{\hat{d}_{\ell_0} k_{\ell_0} (\phi_{\ell_0})_x(y_0^-)}{\phi_{\ell_0}(y_0^-)} - \frac{\hat{d}_1 k_1 (\phi_1)_x(0^+)}{\phi_1(0^+)} > 0, \end{aligned}$$

where we have used (2.1c), (2.2c), (2.21), $\bar{\mathbf{k}} \gg \hat{\mathbf{p}}$ and Lemma A.2 in the last step. This leads to a contradiction and completes the proof. \square

The following result is a direct consequence of Lemmas 2.5 and 2.6.

Theorem 2.7. *Let $\bar{\mathbf{k}}$ be defined in (1.3). Then the following statements hold for model (1.2):*

- (i) *If $\mathbf{p} \gg \bar{\mathbf{k}}$, then the semi-trivial steady state $(\mathbf{u}^*, \mathbf{0})$ is unstable for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_1 \cup \mathcal{L}_3$ and stable for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_2$, where $\mathcal{L}_1, \mathcal{L}_2$ are defined in (2.15) and \mathcal{L}_3 is defined in (2.19);*
- (ii) *If $\bar{\mathbf{k}} \gg \mathbf{p}$, then the semi-trivial steady state $(\mathbf{u}^*, \mathbf{0})$ is unstable for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{S}_1 \cup \mathcal{S}_3$ and stable for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{S}_2$, where $\mathcal{S}_1, \mathcal{S}_2$ are defined in (2.16) and \mathcal{S}_3 is defined in (2.20).*

Note that the mutant species $\mathbf{v} = (v_1, \dots, v_n)$ can (resp. cannot) invade when rare if the semi-trivial steady state $(\mathbf{u}^*, \mathbf{0})$ is unstable (resp. stable). Then, it follows from Theorem 2.7 that, for fixed (\mathbf{d}, \mathbf{q}) with $\mathbf{p} \gg \bar{\mathbf{k}}$, the mutant species \mathbf{v} can invade when rare if $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_1 \cup \mathcal{L}_3$; and for fixed (\mathbf{d}, \mathbf{q}) with $\bar{\mathbf{k}} \gg \mathbf{p}$, the mutant species \mathbf{v} can invade when rare if $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{S}_1 \cup \mathcal{S}_3$. Finally, we summarize the results on invasion analysis in the following: (see Tables 1 and 2)

Table 1: Summary of invasion analysis for $\mathbf{p} \gg \bar{\mathbf{k}}$. Here “ 1 ” (resp. “ 0 ”) represents that species \mathbf{v} can (resp. cannot) invade when rare.

Parameter condition for $\hat{\mathbf{p}}$	Parameter condition for $\hat{\mathbf{d}}$	Results
$\hat{\mathbf{p}} \gg \mathbf{p}$	$\hat{\mathbf{d}} \geq \mathbf{d}$	0
$\mathbf{p} \gg \hat{\mathbf{p}}$	$\mathbf{d} \geq \hat{\mathbf{d}}$	1
$\bar{\mathbf{k}} \gg \hat{\mathbf{p}}$	none	1

Table 2: Summary of invasion analysis for $\bar{\mathbf{k}} \gg \mathbf{p}$. Here “ 1 ” (resp. “ 0 ”) represents that species \mathbf{v} can (resp. cannot) invade when rare.

Parameter condition for $\hat{\mathbf{p}}$	Parameter condition for $\hat{\mathbf{d}}$	Results
$\hat{\mathbf{p}} \gg \mathbf{p}$	$\mathbf{d} \geq \hat{\mathbf{d}}$	1
$\mathbf{p} \gg \hat{\mathbf{p}}$	$\hat{\mathbf{d}} \geq \mathbf{d}$	0
$\hat{\mathbf{p}} \gg \bar{\mathbf{k}}$	none	1

In the following, we further develop the concepts of adaptive dynamics introduced earlier, and show that our result in the two-patch scenario is related to the concept of convergent stable strategy (CSS). For the case $n = 2$, let p_1 and \hat{p}_1 be the strategies for the resident species \mathbf{u} and mutant species \mathbf{v} , respectively, treating all other parameters as identical to isolate their effects. The invasion fitness is given by the principal eigenvalue $\lambda_1(\hat{p}_1, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k})$. Since \mathbf{u}^* depends on the resident's strategy p_1 , we denote $\lambda_1(\hat{p}_1, \hat{\mathbf{d}}, \mathbf{1} - \mathbf{u}^*/\mathbf{k})$ by $\lambda_1(p_1, \hat{p}_1)$ in the subsequent analysis.

Definition 2.8. [7, 32, 56] Suppose that $n = 2$. A strategy p_1^* is called a local evolutionarily stable strategy (ESS) if there exists $\delta > 0$ such that $\lambda_1(p_1^*, \hat{p}_1) < 0$ for all $\hat{p}_1 \in (p_1^* - \delta, p_1^* + \delta) \setminus \{p_1^*\}$. It is called a global ESS if the above condition holds for $\delta = \infty$.

Definition 2.9. [4, 7, 32] Suppose that $n = 2$. The strategy \hat{p}_1^* is called a local neighborhood invader strategy (NIS) if there exists $\delta > 0$ such that $\lambda_1(p_1, \hat{p}_1^*) > 0$ for all $p_1 \in (\hat{p}_1^* - \delta, \hat{p}_1^* + \delta) \setminus \{\hat{p}_1^*\}$. It is called a global NIS if the above condition holds for $\delta = \infty$.

Definition 2.10. [7, 31] Suppose that $n = 2$. A strategy p_1^* is called a local convergent stable strategy (CSS) if there exists $\delta > 0$ such that

$$\lambda_1(p_1, \hat{p}_1) = \begin{cases} < 0 & \text{if } p_1^* \leq p_1 < \hat{p}_1 < p_1^* + \delta \text{ or } p_1^* - \delta < \hat{p}_1 < p_1 \leq p_1^*, \\ > 0 & \text{if } p_1^* \leq \hat{p}_1 < p_1 < p_1^* + \delta \text{ or } p_1^* - \delta < p_1 < \hat{p}_1 \leq p_1^*. \end{cases}$$

It is called a global CSS if the above condition holds for $\delta = \infty$.

Remark 2.11. For the two-patch case, it was shown in [40] that the IFD strategy $\bar{\mathbf{k}} = k_2/k_1$ is a global ESS and NIS. Our result in Theorem 2.7 (or Lemma 2.3) implies that the IFD strategy is a global CSS.

3 Global dynamics

In the subsequent analysis, we investigate the global dynamics of model (1.2) using the theory of monotone dynamical systems. Subsect. 3.1 is devoted to the nonexistence of positive steady state for model (1.2), while the main result is stated in Subsect. 3.2. We note that partial proofs in this section are inspired by the method developed in [40, 60].

3.1 Nonexistence of positive steady state

Denote by (\mathbf{u}, \mathbf{v}) the positive steady state of (1.2) if it exists, where

$$\mathbf{u} = (u_1(x), \dots, u_n(x)) \gg \mathbf{0} \text{ and } \mathbf{v} = (v_1(x), \dots, v_n(x)) \gg \mathbf{0}.$$

Then (\mathbf{u}, \mathbf{v}) is a positive solution of

$$\begin{cases} d_i(u_i)_{xx} + r_i u_i \left(1 - \frac{u_i + v_i}{k_i}\right) = 0, & x \in \Omega_i, \quad i = 1, 2, \dots, n, \end{cases} \quad (3.1a)$$

$$\begin{cases} \hat{d}_i(v_i)_{xx} + r_i v_i \left(1 - \frac{u_i + v_i}{k_i}\right) = 0, & x \in \Omega_i, \quad i = 1, 2, \dots, n, \end{cases} \quad (3.1b)$$

$$u_{i+1}(x_i^+) = p_i u_i(x_i^-), \quad d_{i+1}(u_{i+1})_x(x_i^+) = d_i(u_i)_x(x_i^-), \quad (3.1c)$$

$$v_{i+1}(x_i^+) = \hat{p}_i v_i(x_i^-), \quad \hat{d}_{i+1}(v_{i+1})_x(x_i^+) = \hat{d}_i(v_i)_x(x_i^-), \quad (3.1d)$$

$$(u_1)_x(0^+) = (u_n)_x(L^-) = (v_1)_x(0^+) = (v_n)_x(L^-) = 0. \quad (3.1e)$$

In the following, we obtain some *a priori* estimates on the positive steady state (\mathbf{u}, \mathbf{v}) . The first one is for the case $n = 2$.

Lemma 3.1. *Suppose that $n = 2$. Let $\mathbf{u}, \mathbf{v} \in C^2(\overline{\Omega}_1) \times C^2(\overline{\Omega}_2)$ be the positive solution of (3.1) and let $\bar{\mathbf{k}}$ be defined in (1.3). Then the following statements hold:*

- (i) *If $\mathbf{p}, \hat{\mathbf{p}} \gg \bar{\mathbf{k}}$, then $(u_1)_x(x_1^-), (u_2)_x(x_1^+), (v_1)_x(x_1^-), (v_2)_x(x_1^+) < 0$;*
- (ii) *If $\bar{\mathbf{k}} \gg \mathbf{p}, \hat{\mathbf{p}}$, then $(u_1)_x(x_1^-), (u_2)_x(x_1^+), (v_1)_x(x_1^-), (v_2)_x(x_1^+) > 0$.*

Proof. We only prove (i), since (ii) can be proved similarly. Using arguments similar to those in the proof of (2.12), we obtain that

$$0 = \frac{d_1(k_1 p_1 - k_2)(u_1)_x(x_1^-)}{p_1 u_1(x_1^-)} + \sum_{i=1}^2 \int_{x_{i-1}}^{x_i} \left[\frac{d_i k_i (u_i)_x^2}{u_i^2} + \frac{r_i (k_i - u_i - v_i)^2}{k_i} \right] dx. \quad (3.2)$$

Suppose to the contrary that $(u_1)_x(x_1^-) \geq 0$. This combined with $\mathbf{p} \gg \bar{\mathbf{k}}$ implies that $u_i(x) + v_i(x) = k_i$ for $x \in \overline{\Omega}_i$ with $i = 1, 2$. Then, by (3.1c)-(3.1d),

$$k_2 = u_2(x_1^+) + v_2(x_1^+) = p_1 u_1(x_1^-) + \hat{p}_1 v_1(x_1^-) > k_2,$$

where we have used $\mathbf{p}, \hat{\mathbf{p}} \gg \bar{\mathbf{k}}$ in the last step. This leads to a contradiction, and consequently, $(u_1)_x(x_1^-) < 0$. By (3.1c) again, we have $(u_2)_x(x_1^+) < 0$. Similarly, we can also obtain that $(v_1)_x(x_1^-), (v_2)_x(x_1^+) < 0$. This completes the proof. \square

The second one gives some identities for later use.

Lemma 3.2. *Let $\mathbf{u}, \mathbf{v} \in C^2(\overline{\Omega}_1) \times \cdots \times C^2(\overline{\Omega}_n)$ be the positive solution of (3.1).*

- (i) *Suppose that $x_{i-1} \leq a < b \leq x_i$ with $i = 1, \dots, n$. Then*

$$\begin{aligned} \int_a^b r_i (k_i - u_i - v_i) dx &= -\frac{d_i k_i (u_i)_x(b^-)}{u_i(b^-)} + \frac{d_i k_i (u_i)_x(a^+)}{u_i(a^+)} - d_i k_i \int_a^b \frac{(u_i)_x^2}{u_i^2} dx \\ &= -\frac{\hat{d}_i k_i (v_i)_x(b^-)}{v_i(b^-)} + \frac{\hat{d}_i k_i (v_i)_x(a^+)}{v_i(a^+)} - \hat{d}_i k_i \int_a^b \frac{(v_i)_x^2}{v_i^2} dx. \end{aligned} \quad (3.3)$$

Moreover, if $[d_i(u_i)_x + \hat{d}_i(v_i)_x](a^+) \geq (\text{resp. } >)[d_i(u_i)_x + \hat{d}_i(v_i)_x](b^-)$, then

$$\int_a^b r_i (k_i - u_i - v_i) dx \geq (\text{resp. } >) 0. \quad (3.4)$$

(ii) Suppose that $x_{\ell^*-1} \leq a < x_{\ell^*}$ and $x_{\ell^*-1} < b \leq x_{\ell^*}$ for some ℓ_* and ℓ^* with $1 \leq \ell_* < \ell^* \leq n$. Then

$$\begin{aligned}
0 = & \sum_{i=\ell_*}^{\ell^*-1} \left(\prod_{\ell=i+1}^{\ell^*} p_\ell \right) d_i (\hat{p}_i - p_i) (u_i)_x (x_i^-) v_i (x_i^-) + \hat{d}_{\ell^*} p_{\ell^*} (v_{\ell^*})_x (b^-) u_{\ell^*} (b^-) \\
& - \hat{d}_{\ell_*} \left(\prod_{\ell=\ell_*}^{\ell^*} p_\ell \right) (v_{\ell_*})_x (a^+) u_{\ell_*} (a^+) + d_{\ell_*} \left(\prod_{\ell=\ell_*}^{\ell^*} p_\ell \right) (u_{\ell_*})_x (a^+) v_{\ell_*} (a^+) \\
& - d_{\ell^*} p_{\ell^*} (u_{\ell^*})_x (b^-) v_{\ell^*} (b^-) + \sum_{i=\ell_*}^{\ell^*} \left(\prod_{\ell=i}^{\ell^*} p_\ell \right) (d_i - \hat{d}_i) \int_{\Omega_i \cap (a,b)} (v_i)_x (u_i)_x dx.
\end{aligned} \tag{3.5}$$

Here we define $p_n = 1$.

Proof. Using arguments similar to those in the proof of (2.11), we can obtain that (ii) holds. Dividing (3.1a) and (3.1b) by u_i/k_i and v_i/k_i , respectively, and integrating by parts, we obtain that (3.3) holds. The proof for the remaining part of (i) can be found in [18, Lemma 4.3] (see also [40, Theorem 5.1]). \square

The third one is on the signs of $(u_i)_x$ and $(v_i)_x$.

Lemma 3.3. Let $\mathbf{u}, \mathbf{v} \in C^2(\overline{\Omega}_1) \times \cdots \times C^2(\overline{\Omega}_n)$ be the positive solution of (3.1) and suppose that $x_{i-1} \leq a < b \leq x_i$ with $i = 1, \dots, n$. Then the following two cases cannot occur:

(i) $(u_i)_x, (v_i)_x \geq 0$ with $(u_i)_x \not\equiv 0$ or $(v_i)_x \not\equiv 0$ in (a, b) , and

$$\max\{(u_i)_{xx}(a^+), (v_i)_{xx}(a^+)\} \geq 0, \quad \min\{(u_i)_{xx}(b^-), (v_i)_{xx}(b^-)\} \leq 0;$$

(ii) $(u_i)_x, (v_i)_x \leq 0$ with $(u_i)_x \not\equiv 0$ or $(v_i)_x \not\equiv 0$ in (a, b) , and

$$\min\{(u_i)_{xx}(a^+), (v_i)_{xx}(a^+)\} \leq 0, \quad \max\{(u_i)_{xx}(b^-), (v_i)_{xx}(b^-)\} \geq 0.$$

Proof. We only show that (i) cannot occur, and (ii) can be treated similarly. Suppose to the contrary that (i) occurs. Since

$$\max\{(u_i)_{xx}(a^+), (v_i)_{xx}(a^+)\} \geq 0 \quad \text{and} \quad \min\{(u_i)_{xx}(b^-), (v_i)_{xx}(b^-)\} \leq 0,$$

it follows from (3.1a) and (3.1b) that

$$u_i(a^+) + v_i(a^+) \geq k_i \quad \text{and} \quad u_i(b^-) + v_i(b^-) \leq k_i,$$

which contradicts the fact that $(u_i)_x, (v_i)_x \geq 0$ with $(u_i)_x \not\equiv 0$ or $(v_i)_x \not\equiv 0$ in (a, b) . Therefore, (i) can not occur. \square

The last one is as follows.

Lemma 3.4. *Let $\mathbf{u}, \mathbf{v} \in C^2(\overline{\Omega}_1) \times \cdots \times C^2(\overline{\Omega}_n)$ be the positive solution of (3.1). If $(u_1)_x(x_1^-), (v_1)_x(x_1^-) \leq 0$, then $(u_1)_x, (v_1)_x \leq 0$ in Ω_1 .*

Proof. Since $(u_1)_x(x_1^-), (v_1)_x(x_1^-) \leq 0$, we divide the proof into the following two cases:

$$(a_1) \ (u_1)_x(x_1^-)(v_1)_x(x_1^-) = 0; \quad (a_2) \ (u_1)_x(x_1^-), (v_1)_x(x_1^-) < 0.$$

For case (a_1) , since $0 = [d_1(u_1)_x + \hat{d}_1(v_1)_x](0^+) \geq [d_1(u_1)_x + \hat{d}_1(v_1)_x](x_1^-)$, it follows from Lemma 3.2 (i) that

$$\int_0^{x_1} r_1 (k_1 - u_1 - v_1) \geq 0. \quad (3.6)$$

In view of $(u_1)_x(x_1^-)(v_1)_x(x_1^-) = 0$, we see from (3.3) and (3.6) that $(u_1)_x = 0$ in Ω_1 or $(v_1)_x = 0$ in Ω_1 . This combined with (3.1a)-(3.1b) implies that $(u_1)_x, (v_1)_x = 0$ in Ω_1 , and the desired result holds.

For case (a_2) , since $(u_1)_x(0^+) = (v_1)_x(0^+) = 0$, it follows that

$$y_1 := \sup\{x \in [0, x_1) | (u_1)_x(x^+) = 0\} \text{ and } y_2 := \sup\{x \in [0, x_1) | (v_1)_x(x^+) = 0\}$$

are well-defined with $0 \leq y_1, y_2 < x_1$. Define $y_0 := \max\{y_1, y_2\}$. Clearly, the desired result holds for $y_0 = 0$. If $y_0 > 0$, $(u_1)_x(y_0^-), (v_1)_x(y_0^-) \leq 0$ with $(u_1)_x(y_0^-)(v_1)_x(y_0^-) = 0$. Then using similar arguments as in the proof of case (a_1) , we see that $(u_1)_x, (v_1)_x = 0$ in $[0, y_0]$, and the desired result holds. \square

Now we show the nonexistence of positive steady state for model (1.2).

Theorem 3.5. *Let $\bar{\mathbf{k}}$ be defined in (1.3). Then model (1.2) has no positive steady state if one of the following conditions holds:*

(i) $\mathbf{p} \gg \bar{\mathbf{k}}$ and $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_1^* \cup \mathcal{L}_2$, where \mathcal{L}_2 is defined in (2.15), and

$$\mathcal{L}_1^* := \{(\hat{\mathbf{p}}, \hat{\mathbf{d}}) : \mathbf{p} \gg \hat{\mathbf{p}} \gg \bar{\mathbf{k}}, \mathbf{d} \geq \hat{\mathbf{d}}\}; \quad (3.7)$$

(ii) $\bar{\mathbf{k}} \gg \mathbf{p}$ and $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{S}_1^* \cup \mathcal{S}_2$, where \mathcal{S}_2 is defined in (2.16), and

$$\mathcal{S}_1^* := \{(\hat{\mathbf{p}}, \hat{\mathbf{d}}) : \bar{\mathbf{k}} \gg \hat{\mathbf{p}} \gg \mathbf{p}, \mathbf{d} \geq \hat{\mathbf{d}}\}. \quad (3.8)$$

Proof. We only prove (i), and (ii) can be handled similarly. Since the nonlinear terms of model (1.2) are symmetric, it suffices to consider the case $\mathbf{p} \gg \bar{\mathbf{k}}$ and $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_1^*$. The rest of the proof is based on the following three claims.

Claim 1: If $(u_i)_x, (v_i)_x \leq 0$ in Ω_i and $(u_i)_x(x_i^-), (v_i)_x(x_i^-) \leq 0$ for $i = 1, \dots, n-1$, then $(u_n)_x, (v_n)_x \leq 0$ in Ω_n .

Proof of Claim: Since $(u_n)_x(L^-) = (v_n)_x(L^-) = 0$, it follows that z_1, z_2 are well-defined with $x_{n-1} \leq z_1, z_2 \leq L$, where

$$\begin{aligned} z_1 &:= \inf\{x \in [x_{n-1}, x_n] : \exists n-1 \leq i \leq n \text{ s.t. } x \in (x_{i-1}, x_i] \text{ and } (u_i)_x(y^-) = 0\}, \\ z_2 &:= \inf\{x \in [x_{n-1}, x_n] : \exists n-1 \leq i \leq n \text{ s.t. } x \in (x_{i-1}, x_i] \text{ and } (v_i)_x(y^-) = 0\}, \end{aligned}$$

and for fixed $i = 1, 2$, $z_i \in (x_{\ell_i-1}, x_{\ell_i}]$, where $\ell_i = n$ if $z_i > x_{n-1}$ and $\ell_i = n-1$ if $z_i = x_{n-1}$.

We first claim that $z_1 \leq z_2$. Suppose not, then $z_1 > z_2$. Thus, we have

$$(u_{\ell_2})_x(z_2^-) < 0 \text{ and } (u_i)_x, (v_i)_x \leq 0 \text{ in } \Omega_i \cap (0, z_2) \text{ for } i = 1, \dots, \ell_2. \quad (3.9)$$

Substituting $a = 0$, $b = z_2$, $\ell_* = 1$ and $\ell^* = \ell_2$ into (3.5) yields

$$\begin{aligned} 0 &= \sum_{i=1}^{\ell_2-1} \left(\prod_{\ell=i+1}^{\ell_2} p_\ell \right) d_i (\hat{p}_i - p_i) (u_i)_x(x_i^-) v_i(x_i^-) + \hat{d}_{\ell_2} p_{\ell_2} u_{\ell_2}(z_2^-) (v_{\ell_2})_x(z_2^-) \\ &\quad + \sum_{i=1}^{\ell_2} \left(\prod_{\ell=i}^{\ell_2} p_\ell \right) (d_i - \hat{d}_i) \int_{\Omega_i \cap (0, z_2)} (v_i)_x (u_i)_x dx - d_{\ell_2} p_{\ell_2} (u_{\ell_2})_x(z_2^-) v_{\ell_2}(z_2^-) > 0, \end{aligned} \quad (3.10)$$

where we have used (3.9), $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_1^*$ and the conditions of Claim 1 in the last step. This leads to a contradiction, and the claim $z_1 \leq z_2$ holds.

Then we claim $(u_n)_x(z_2^-) \leq 0$ if $z_2 \in (x_{n-1}, L]$. Clearly, $(u_n)_x(L^-) = 0$, and we only need to consider the case $z_2 \in (x_{n-1}, L)$. To the contrary, we have

$$(u_n)_x(z_2) = (u_n)_x(z_2^-) > 0 = (u_n)_x(L^-), \quad (v_n)_x(z_2) = (v_n)_x(z_2^-) = (v_n)_x(L^-) = 0.$$

This together with Lemma 3.2 (i) yields

$$0 < \int_{z_2}^L r_n (k_n - u_n - v_n) dx = -\hat{d}_n k_n \int_{z_2}^L \frac{(v_n)_x^2}{v_n^2} dx \leq 0,$$

which is a contradiction. Therefore, the claim holds, i.e., $(u_n)_x(z_2^-) \leq 0$ if $z_2 \in (x_{n-1}, L]$.

Now we claim that $z_1 = z_2$. Suppose not, then $x_{n-1} \leq z_1 < z_2 \leq L$. Thanks to $(u_n)_x(z_2^-) \leq 0$, we will obtain a contradiction for each of the following two cases:

$$(a_1) \ z_1 < z_2, \ (u_n)_x(z_2^-) < 0; \ (a_2) \ z_1 < z_2, \ (u_n)_x(z_2^-) = 0.$$

For case (a₁), noticing that $d_n(u_n)_x(z_1^+) = d_{\ell_1}(u_{\ell_1})_x(z_1^-) = 0$, where $\ell_1 = n$ if $z_1 > x_{n-1}$ and $\ell_1 = n-1$ if $z_1 = x_{n-1}$, we see that $z_3 = \sup\{x \in [z_1, z_2] : (u_n)_x(x^+) = 0\}$ is

well-defined with $z_1 \leq z_3 < z_2$. Thus, $(u_n)_{xx}(z_3^+) \leq 0$, $(u_n)_x(z_3^+) = 0$ and $(u_n)_x < 0$ in (z_3, z_2) . This combined the definition of z_2 implies that Lemma 3.3 (ii) occurs with $a = z_3$ and $b = z_2$, which is a contradiction.

For case (a₂), $(u_n)_x(z_2^-) = 0$. Set $\mathcal{M} := \{x \in (z_1, z_2) : (u_n)_x(x) \geq 0\}$. If $\mathcal{M} \subsetneq (z_1, z_2)$, then there exists $z_4 \in (z_1, z_2)$ such that $(u_n)_x(z_4) < 0$. Noting that $(u_n)_x(z_1^+) = (u_n)_x(z_2^-) = 0$, we have

$$z_5 := \sup\{x \in [z_1, z_4] | (u_n)_x(x^+) = 0\} \quad \text{and} \quad z_6 := \inf\{x \in [z_4, z_2] | (u_n)_x(x^-) = 0\}$$

are well-defined with $z_1 \leq z_5 < z_4$ and $z_4 < z_6 \leq z_2$. Thus, $(u_n)_{xx}(z_5^+) \leq 0$, $(u_n)_{xx}(z_6^-) \geq 0$ and $(u_n)_x < 0$ in (z_5, z_6) . This together with (3.9) implies Lemma 3.3 (ii) occurs with $a = z_5$ and $b = z_6$, which is a contradiction.

If $\mathcal{M} = (z_1, z_2)$, we have $(u_n)_x \geq 0$ in (z_1, z_2) . Then one of the following two subcases occur:

(b₁) there exists $z_* \in (z_1, z_2)$ such that $d_n(u_n)_x(z_*) + \hat{d}_n(v_n)_x(z_*) \geq 0$;

(b₂) $d_n(u_n)_x + \hat{d}_n(v_n)_x < 0$ in (z_1, z_2) .

For subcase (b₁), noticing that $(u_n)_x(z_2^-) = (v_n)_x(z_2^-) = 0$, we deduce from Lemma 3.2 (i) that

$$\int_{z_*}^{z_2} r_n (k_n - u_n - v_n) dx \geq 0. \quad (3.11)$$

In addition, since $z_2 > x_{n-1}$, we see from the definition of z_2 that

$$(v_n)_x < 0 \quad \text{in} \quad [z_*, z_2) \quad \text{and} \quad (v_n)_x(z_2^-) = 0. \quad (3.12)$$

Then it follows from (3.3) that

$$\begin{aligned} \int_{z_*}^{z_2} r_n (k_n - u_n - v_n) dx &= - \frac{\hat{d}_n k_n (v_n)_x(z_2^-)}{v_n(z_2^-)} + \frac{\hat{d}_n k_n (v_n)_x(z_*)}{v_2(z_*)} \\ &\quad - \hat{d}_n k_n \int_{z_*}^{z_2} \frac{(v_n)_x^2}{v_n^2} dx < 0, \end{aligned}$$

where we have used (3.12) in the last step. This contradicts (3.11).

For subcase (b₂), we have

$$\hat{d}_n [(u_n)_x + (v_n)_x] \leq d_n(u_n)_x + \hat{d}_n(v_n)_x < 0 \quad \text{in} \quad (z_1, z_2), \quad (3.13)$$

where we have used $\hat{d}_n \leq d_n$ and $\mathcal{M} = (z_1, z_2)$ in first step. By the definition of z_2 , $(v_n)_{xx}(z_2^-) \geq 0$, which together with (3.1b) yields $u_n(z_2^-) + v_n(z_2^-) \geq k_n$. Then it follows

from (3.13) that $u_n(x) + v_n(x) > k_n$ for $x \in (z_1, z_2)$, and consequently, we see from (3.1a) that $(u_n)_{xx} > 0$ in (z_1, z_2) , which contradicts the fact that $(u_n)_x(z_1^+) = (u_n)_x(z_2^-) = 0$. Therefore, we obtain a contradiction for each of the cases (a₁)-(a₂), and the claim $z_1 = z_2$ holds.

If $z_1 = z_2 = L$, then by the definitions of z_1 and z_2 , $(u_n)_x(x), (v_n)_x(x) \leq 0$ for $x \in \Omega_n$. Thus, Claim 1 holds for this case. If $z_1 = z_2 < L$, we have

$$(u_n)_x(z_2^+) = (u_n)_x(L^-) = (v_n)_x(z_2^+) = (v_n)_x(L^-) = 0.$$

Then, by Lemma 3.2 (i) again,

$$0 \leq \int_{z_2}^L r_n(k_n - u_n - v_n) dx = -d_2 k_n \int_{z_2}^L \frac{(u_n)_x^2}{u_n^2} dx = -\hat{d}_2 k_n \int_{z_2}^L \frac{(v_n)_x^2}{v_n^2} dx \leq 0,$$

which yields $(u_n)_x, (v_n)_x = 0$ in (z_2, L) . This combined with the definitions of z_1 and z_2 implies that Claim 1 also holds for this case.

Claim 2: We claim that model (1.2) has no positive steady state for the case $n = 2$.

Proof of Claim: Suppose to the contrary that model (1.2) has a positive steady state (\mathbf{u}, \mathbf{v}) for the case $n = 2$. It follows from Lemmas 3.1 and 3.4 and Claim 1 that

$$(u_1)_x(x_1^-), (v_1)_x(x_1^-) < 0 \text{ and } (u_i)_x, (v_i)_x \leq 0 \text{ in } \Omega_i \text{ for each } i = 1, 2. \quad (3.14)$$

Plugging $a = 0$ and $b = L$ into (3.5) yields

$$0 = d_1(\hat{p}_1 - p_1)(u_1)_x(x_1^-)v_1(x_1^-) + \sum_{i=1}^2 \left(\prod_{\ell=i}^2 p_\ell \right) (d_i - \hat{d}_i) \int_{\Omega_i} (v_i)_x(u_i)_x dx > 0, \quad (3.15)$$

where we have used (3.14), $\mathbf{p} \gg \hat{\mathbf{p}}$ and $\mathbf{d} \geq \hat{\mathbf{d}}$ in the last step. This leads to a contradiction and completes the proof for Claim 2.

Claim 3: We claim that model (1.2) has no positive steady state for the case $n = m$ with $m \geq 3$ if it has no positive steady state for $n = 2, \dots, m-1$.

Proof of Claim: Suppose to the contrary that model (1.2) has a positive steady state (\mathbf{u}, \mathbf{v}) for the case $n = m$. To obtain a contradiction, we first show that the following two cases cannot occur:

(c₁) there exists $2 \leq \ell_0 \leq m-2$ such that $(u_{\ell_0})_x(x_{\ell_0}^-)(v_{\ell_0})_x(x_{\ell_0}^-) \leq 0$ if $m \geq 4$;

(c₂) there exists $2 \leq \ell_0 \leq m-1$ and $x^* \in \Omega_{\ell_0}$ such that $(u_{\ell_0})_x(x^*)(v_{\ell_0})_x(x^*) \leq 0$.

Suppose to the contrary that (c₁) occurs. Without loss of generality, we assume that

$$(u_{\ell_0})_x(x_{\ell_0}^-) \leq 0 \text{ and } (v_{\ell_0})_x(x_{\ell_0}^-) \geq 0. \quad (3.16)$$

Construct the following auxiliary system:

$$\begin{cases} (w_i)_t = d_i(w_i)_{xx} + r_i w_i \left(1 - \frac{w_i + \xi_i}{k_i}\right), & t > 0, \quad x \in \Omega_i, \quad i = 1, \dots, \ell_0, \\ (\xi_i)_t = \hat{d}_i(\xi_i)_{xx} + r_i \xi_i \left(1 - \frac{w_i + \xi_i}{k_i}\right), & t > 0, \quad x \in \Omega_i, \quad i = 1, \dots, \ell_0, \\ w_{i+1}(x_i^+, t) = p_i w_i(x_i^-, t), \quad d_{i+1}(w_{i+1})_x(x_i^+, t) = d_i(w_i)_x(x_i^-, t), & t > 0, \\ \xi_{i+1}(x_i^+, t) = \hat{p}_i \xi_i(x_i^-, t), \quad \hat{d}_{i+1}(\xi_{i+1})_x(x_i^+, t) = \hat{d}_i(\xi_i)_x(x_i^-, t), & t > 0, \\ (w_1)_x(0^+, t) = (w_{\ell_0})_x(x_{\ell_0}^-, t) = (\xi_1)_x(0^+, t) = (\xi_{\ell_0})_x(x_{\ell_0}^-, t) = 0, & t > 0. \end{cases} \quad (3.17)$$

Then system (3.17) generates a monotone dynamical system, which is order-preserving with respect to the order “ \succeq ” defined in (B.1) (see the Appendix A). Define

$$([\mathbf{u}]_{\ell_0}, [\mathbf{v}]_{\ell_0}) = (u_1(x), \dots, u_{\ell_0}(x), v_1(x), \dots, v_{\ell_0}(x)).$$

It follows from (3.16) and Proposition B.3 that $([\mathbf{u}]_{\ell_0}, [\mathbf{v}]_{\ell_0})$ is a sub-equilibrium of (3.17).

Let $(\mathbf{w}^*, \mathbf{0})$ be the semi-trivial steady state of (3.17), where

$$\mathbf{w}^* = (w_1^*(x), \dots, w_{\ell_0}^*(x))$$

with $w_i^*(x) > 0$ for $x \in \bar{\Omega}_i$ and $i = 1, \dots, \ell_0$. It follows from Lemma 2.5 (i) that $\tilde{\lambda}_1 > 0$, where $\tilde{\lambda}_1$ is the principal eigenvalue of the following eigenvalue problem:

$$\begin{cases} \hat{d}_i(\psi_i)_{xx} + r_i \psi_i \left(1 - \frac{w_i^*}{k_i}\right) = \tilde{\lambda}_1 \psi_i, & x \in \Omega_i, \quad i = 1, \dots, \ell_0, \\ \psi_{i+1}(x_i^+) = \hat{p}_i \psi_i(x_i^-), \quad \hat{d}_{i+1}(\psi_{i+1})_x(x_i^+) = \hat{d}_i(\psi_i)_x(x_i^-), \\ (\psi_1)_x(0^+) = (\psi_{\ell_0})_x(x_{\ell_0}^-) = 0. \end{cases}$$

Let $\psi = (\psi_1(x), \dots, \psi_{\ell_0}(x))$ be the eigenfunction corresponding to $\tilde{\lambda}_1$ with $\psi_i(x) > 0$ for $x \in \bar{\Omega}_i$ and $i = 1, \dots, \ell_0$. Choose $\epsilon_1, \epsilon_2 > 0$ so that

$$\min_{x \in \bar{\Omega}_i} v_i(x) > \epsilon_2 \max_{x \in \bar{\Omega}_i} \psi_i(x) \quad \text{and} \quad \tilde{\lambda}_1 > \frac{r_i}{k_i} \left(\epsilon_1 \max_{x \in \bar{\Omega}_i} w_i^* + \epsilon_2 \max_{x \in \bar{\Omega}_i} \psi_i \right) \quad \text{for } i = 1, \dots, \ell_0. \quad (3.18)$$

A direct computation yields, for $i = 1, \dots, \ell_0$,

$$\begin{aligned} & (1 + \epsilon_1) d_i(w_i^*)_{xx} + (1 + \epsilon_1) r_i w_i^* \left(1 - \frac{w_i^* + \epsilon_1 w_i^* + \epsilon_2 \psi_i}{k_i}\right) \\ &= - \frac{(1 + \epsilon_1) r_i w_i^* (\epsilon_1 w_i^* + \epsilon_2 \psi_i)}{k_i} < 0, \end{aligned}$$

and

$$\epsilon_2 \hat{d}_i(\psi_i)_{xx} + \epsilon_2 r_i \psi_i \left(1 - \frac{(1 + \epsilon_1)w_i^* + \epsilon_2 \psi_i}{k_i} \right) = \epsilon_2 \psi_i \left(\tilde{\lambda}_1 - \frac{r_i(\epsilon_1 w_i^* + \epsilon_2 \psi_i)}{k_i} \right) > 0,$$

which implies that $((1 + \epsilon_1)\mathbf{w}^*, \epsilon_2 \psi)$ is a super-equilibrium of (3.17).

By the first inequality of (3.16), $[\mathbf{u}]_{\ell_0} = (u_1, \dots, u_{\ell_0})$ is a lower solution of

$$\begin{cases} d_i(w_i)_{xx} + r_i w_i \left(1 - \frac{w_i}{k_i} \right) = 0, & x \in \Omega_i, \quad i = 1, \dots, \ell_0, \\ w_{i+1}(x_i^+) = p_i w_i(x_i^-), \quad d_{i+1}(w_{i+1})_x(x_i^+) = d_i(w_i)_x(x_i^-), \\ (w_1)_x(0^+) = (w_{\ell_0})_x(x_{\ell_0}^-) = 0. \end{cases} \quad (3.19)$$

This implies $u_i \leq w_i^*$ in $\bar{\Omega}_i$, and consequently, $u_i < (1 + \epsilon_1)w_i^*$ in $\bar{\Omega}_i$ for $i = 1, \dots, \ell_0$. This together with (3.18) yields $((1 + \epsilon_1)\mathbf{w}^*, \epsilon_2 \psi) \succeq (\neq) ([\mathbf{u}]_{\ell_0}, [\mathbf{v}]_{\ell_0})$. Then, it follows from Theorem B.4 that the auxiliary system (3.17) admits a positive steady state, which contradicts the fact that model (1.2) has no positive steady state for $n = 2, \dots, m - 1$. Thus, (c_1) cannot occur, and case (c_2) can be handled similarly.

Since (c_1) and (c_2) cannot occur, it follows that one of the following two cases can occur:

- (d₁) $(u_i)_x, (v_i)_x > 0$ in Ω_i for each $2 \leq i \leq m - 1$ and $(u_i)_x(x_i^-), (v_i)_x(x_i^-) > 0$ for each $i = 2, \dots, m - 2$;
- (d₂) $(u_i)_x, (v_i)_x < 0$ in Ω_i for each $2 \leq i \leq m - 1$ and $(u_i)_x(x_i^-), (v_i)_x(x_i^-) < 0$ for each $i = 2, \dots, m - 2$.

Now we obtain a contradiction for each of cases (d₁) and (d₂).

For case (d₁), we see from (3.1e) that y_*, y^*, z_*, z^* are well-defined with $0 \leq y_*, z_* \leq x_1 \leq x_{m-1} \leq y^*, z^* \leq L$, where

$$\begin{aligned} y_* &= \sup\{y : \exists 1 \leq \ell \leq 2 \text{ s.t. } x \in [x_{\ell-1}, x_\ell] \text{ and } (u_\ell)_x(y^+) = 0\}, \\ y^* &= \inf\{y : \exists m - 1 \leq \ell \leq m \text{ s.t. } x \in (x_{\ell-1}, x_\ell] \text{ and } (u_\ell)_x(y^-) = 0\}, \\ z_* &= \sup\{z : \exists 1 \leq \ell \leq 2 \text{ s.t. } x \in [x_{\ell-1}, x_\ell] \text{ and } (v_\ell)_x(z^+) = 0\}, \\ z^* &= \inf\{z : \exists m - 1 \leq \ell \leq m \text{ s.t. } x \in (x_{\ell-1}, x_\ell] \text{ and } (v_\ell)_x(z^-) = 0\}. \end{aligned}$$

Let $\underline{x} = \max\{y_*, z_*\}$ and $\bar{x} = \min\{y^*, z^*\}$, and there exist $\underline{\ell}$ and $\bar{\ell}$ with $1 \leq \underline{\ell} \leq 2$ and $m - 1 \leq \bar{\ell} \leq m$ such that $\underline{x} \in [x_{\underline{\ell}-1}, x_{\underline{\ell}}]$ and $\bar{x} \in (x_{\bar{\ell}-1}, x_{\bar{\ell}}]$. Then, by (3.1a)-(3.1b), $u_{\underline{\ell}}(\underline{x}^+) + v_{\underline{\ell}}(\underline{x}^+) \geq k_{\underline{\ell}}$ and $u_{\bar{\ell}}(\bar{x}^-) + v_{\bar{\ell}}(\bar{x}^-) \leq k_{\bar{\ell}}$. This combined with (d₁) and

$\mathbf{p} \gg \hat{\mathbf{p}} \gg \bar{\mathbf{k}}$ implies that

$$\begin{aligned} k_{\bar{\ell}} &\geq u_{\bar{\ell}}(\bar{x}^-) + v_{\bar{\ell}}(\bar{x}^-) > u_{\bar{\ell}}(x_{\bar{\ell}-1}^+) + v_{\bar{\ell}}(x_{\bar{\ell}-1}^+) > \hat{p}_{\bar{\ell}-1} \left(u_{\bar{\ell}-1}(x_{\bar{\ell}-1}^-) + v_{\bar{\ell}-1}(x_{\bar{\ell}-1}^-) \right) \\ &> \cdots > \left(\prod_{\underline{\ell}}^{\bar{\ell}-1} \hat{p}_i \right) \left(u_{\underline{\ell}}(x_{\underline{\ell}}^-) + v_{\underline{\ell}}(x_{\underline{\ell}}^-) \right) > \left(\prod_{\underline{\ell}}^{\bar{\ell}-1} \hat{p}_i \right) \left(u_{\underline{\ell}}(\underline{x}^+) + v_{\underline{\ell}}(\underline{x}^+) \right) > k_{\bar{\ell}}, \end{aligned}$$

which is a contradiction.

For case (d₂), we see from Lemma 3.4 and Claim 1 that $(u_i)_x, (v_i)_x \leq 0$ in Ω_i for each $1 \leq i \leq m$ and $(u_i)_x(x_i^-), (v_i)_x(x_i^-) \leq 0$ for each $i = 1, \dots, m-1$. Then plugging $a = 0$, $b = L$, $\ell_* = 1$ and $\ell^* = m$ into (3.5) yields

$$\begin{aligned} 0 &= \sum_{i=1}^{m-1} \left(\prod_{\ell=i+1}^m p_{\ell} \right) d_i (\hat{p}_i - p_i) (u_i)_x(x_i^-) v_i(x_i^-) \\ &\quad + \sum_{i=1}^m \left(\prod_{\ell=i}^m p_{\ell} \right) (d_i - \hat{d}_i) \int_{\Omega_i} (v_i)_x (u_i)_x dx \geq 0, \end{aligned} \tag{3.20}$$

where we have used $\mathbf{p} \gg \hat{\mathbf{p}}$ and $\mathbf{d} \geq \hat{\mathbf{d}}$ in the last step. Exchanging the positions of u_i and v_i in (3.20), we can also obtain that

$$\begin{aligned} 0 &= \sum_{i=1}^{m-1} \left(\prod_{\ell=i+1}^m \hat{p}_{\ell} \right) \hat{d}_i (p_i - \hat{p}_i) (v_i)_x(x_i^-) u_i(x_i^-) \\ &\quad + \sum_{i=1}^m \left(\prod_{\ell=i}^m \hat{p}_{\ell} \right) (\hat{d}_i - d_i) \int_{\Omega_i} (v_i)_x (u_i)_x dx \leq 0, \end{aligned} \tag{3.21}$$

where we have used $\mathbf{p} \gg \hat{\mathbf{p}}$ and $\mathbf{d} \geq \hat{\mathbf{d}}$ in the last step. Then it follows from (3.20) and (3.21) that $(u_i)_x(x_i^-) = (v_i)_x(x_i^-) = 0$ for each $i = 1, \dots, m-1$, which implies that $u_i + v_i = k_i$ in Ω_i for each $i = 1, \dots, m$. Then

$$\begin{aligned} k_m &= u_m(x_{m-1}^+) + v_m(x_{m-1}^+) > \hat{p}_{m-1} [u_{m-1}(x_{m-1}^-) + v_{m-1}(x_{m-1}^-)] \\ &> \cdots > \left(\prod_{\ell=1}^{m-1} \hat{p}_{\ell} \right) [u_1(x_1^-) + v_1(x_1^-)] > k_m, \end{aligned} \tag{3.22}$$

where we have used (3.1d) and $\mathbf{p} \gg \hat{\mathbf{p}}$ in the second step and $\hat{\mathbf{p}} \gg \bar{\mathbf{k}}$ in the last step. This also leads to a contradiction and completes the proof for Claim 3.

Then the desired result follows from Claims 2-3 and induction. \square

3.2 Main result

Now we present our main result on the global dynamics of model (1.2).

Theorem 3.6. *Let $\bar{\mathbf{k}}$ be defined in (1.3). Then the following statements hold for model (1.2):*

- (i) *If $\mathbf{p} \gg \bar{\mathbf{k}}$, then the semi-trivial steady state $(\mathbf{0}, \mathbf{v}^*)$ is globally asymptotically stable for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_1^*$ and $(\mathbf{u}^*, \mathbf{0})$ is globally asymptotically stable for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_2$, where \mathcal{L}_1^* and \mathcal{L}_2 are defined in (3.7) and (2.15), respectively. Moreover, model (1.2) admits at least one stable positive steady state for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_3$, where \mathcal{L}_3 is defined in (2.19).*
- (ii) *If $\bar{\mathbf{k}} \gg \mathbf{p}$, then the semi-trivial steady state $(\mathbf{0}, \mathbf{v}^*)$ is globally asymptotically stable for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{S}_1^*$ and $(\mathbf{u}^*, \mathbf{0})$ is globally asymptotically stable for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{S}_2$, where \mathcal{S}_1^* and \mathcal{S}_2 are defined in (3.8) and (2.16), respectively. Moreover, model (1.2) admits at least one stable positive steady state for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{S}_3$, where \mathcal{S}_3 is defined in (2.20).*

Proof. We only prove (i), since (ii) can be treated similarly. We can rewrite model (1.2) as the abstract evolution equation (C.8) in Appendix C. Theorem C.8 implies that the semiflow T_t generated by (C.8) is strongly order-preserving with respect to \geq_k (defined in (C.30)) and order compact on the fractional power spaces $X_u^\alpha \times X_v^\alpha$ (defined in Remark C.3) for all $t > 0$. Consequently, the theory of monotone dynamical systems [27, 28, 33, 55] yields the global dynamics of (1.2). Specifically, Theorem 3.5 implies that model (1.2) has no positive steady state for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_1^* \cup \mathcal{L}_2$. Moreover, Theorem 2.7 shows that $(\mathbf{u}^*, \mathbf{0})$ is unstable (resp. stable) for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_1^*$ (resp. $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_2$). Since the nonlinear terms of model (1.2) are symmetric, we can also obtain that $(\mathbf{0}, \mathbf{v}^*)$ is stable (resp. unstable) for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_1^*$ (resp. $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_2$). Then, by the monotone dynamical system theory [27, 28, 33, 55], we obtain $(\mathbf{0}, \mathbf{v}^*)$ (resp. $(\mathbf{u}^*, \mathbf{0})$) is globally asymptotically stable for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_1^*$ (resp. $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_2$). By Theorem 2.7, we see that $(\mathbf{u}^*, \mathbf{0})$ is unstable for $(\hat{\mathbf{p}}, \hat{\mathbf{d}}) \in \mathcal{L}_3$. Note that the nonlinear terms of model (1.2) are symmetric. We can also obtain that $(\mathbf{0}, \mathbf{v}^*)$ is unstable. Then, by the monotone dynamical system theory again, (1.2) admits at least one stable positive steady state. \square

Finally, we summarize the global dynamics of model (1.2) in the following two tables: (see Tables 3 and 4).

4 Discussion

We study a reaction-diffusion model for two-species competition in a fragmented habitat of n adjacent patches. The model incorporates habitat preference of individuals, which

Table 3: Summary of the global dynamics for $\mathbf{p} \gg \bar{\mathbf{k}}$.

Parameter condition for $\hat{\mathbf{p}}$	Parameter condition for $\hat{\mathbf{d}}$	Results
$\hat{\mathbf{p}} \gg \mathbf{p}$	$\hat{\mathbf{d}} \geq \mathbf{d}$	$(\mathbf{u}^*, \mathbf{0})$ g.a.s.
$\mathbf{p} \gg \hat{\mathbf{p}} \gg \bar{\mathbf{k}}$	$\mathbf{d} \geq \hat{\mathbf{d}}$	$(\mathbf{0}, \mathbf{v}^*)$ g.a.s.
$\bar{\mathbf{k}} \gg \hat{\mathbf{p}}$	none	coexistence

Table 4: Summary of the global dynamics for $\bar{\mathbf{k}} \gg \mathbf{p}$.

Parameter condition for $\hat{\mathbf{p}}$	Parameter condition for $\hat{\mathbf{d}}$	Results
$\bar{\mathbf{k}} \gg \hat{\mathbf{p}} \gg \mathbf{p}$	$\mathbf{d} \geq \hat{\mathbf{d}}$	$(\mathbf{0}, \mathbf{v}^*)$ g.a.s.
$\mathbf{p} \gg \hat{\mathbf{p}}$	$\hat{\mathbf{d}} \geq \mathbf{d}$	$(\mathbf{u}^*, \mathbf{0})$ g.a.s.
$\hat{\mathbf{p}} \gg \bar{\mathbf{k}}$	none	coexistence

induces density discontinuities at interfaces. This model differs from established models, such as coupled ordinary differential equations (patch models) and classical reaction-diffusion models, which typically do not account for such discontinuities.

A central question in the study of two-species competition concerns the mechanisms that drive competitive exclusion or enable coexistence. Within a two-patch reaction-diffusion framework incorporating edge behavior, [40] demonstrated that when habitat preference is viewed as a strategy, the ideal free distribution (IFD) strategy is a global evolutionarily stable strategy (ESS). Our results extend this by proving that the IFD strategy is a global convergent stable strategy (CSS). Specifically, we show that a species with a strategy closer to the IFD can invade and outcompete the resident whose strategy is farther away, provided both strategies lie on the same side of the IFD. Conversely, two species can coexist if their strategies lie on opposite sides of the IFD. Moreover, this result is extended to an n -patch reaction-diffusion model incorporating edge behavior.

These findings are consistent with results from both coupled ordinary differential equations (patch models) and classical reaction-diffusion models, in which dispersal is typically treated as a strategy. Indeed, as demonstrated by [8, 38] and [5, 7], the IFD strategy is also a global ESS in the frameworks of patch models and classical reaction-diffusion models, respectively. Furthermore, competitive outcomes (exclusion or coexistence) are also determined by whether species' strategies reside on the same or opposite sides of the IFD, see [7, 36]. Therefore, across all three classes of models, the ideal free distribution (IFD) emerges as a universal principle that governs whether

competition results in exclusion or coexistence.

Our analysis relies crucially on the monotonicity of the species steady state for the corresponding single-species model (2.1). According to Lemma A.2, this monotonicity in solutions to system (2.1) is guaranteed if either $\mathbf{p} \gg \bar{\mathbf{k}}$ or $\bar{\mathbf{k}} \gg \mathbf{p}$ holds. Biologically, this condition requires that the discontinuities in population density at all interfaces are either uniformly larger or uniformly smaller than the corresponding discontinuities in the carrying capacity [59]. A violation of this condition may lead to non-monotonic steady states, as demonstrated in Proposition A.3. Nevertheless, the non-monotonicity of steady states poses analytical challenges that remain to be explored in future work.

Appendix A Semi-trivial steady state

In this part, we consider the positive solutions of the following system:

$$\begin{cases} d_i(u_i)_{xx} + r_i u_i \left(1 - \frac{u_i}{k_i}\right) = 0, & x \in \Omega_i, \quad i = 1, 2, \dots, n, \\ u_{i+1}(x_i^+) = p_i u_i(x_i^-), \quad d_{i+1}(u_{i+1})_x(x_i^+) = d_i(u_i)_x(x_i^-), & i = 1, \dots, n-1, \\ (u_1)_x(0^+) = (u_n)_x(L^-) = 0, \end{cases} \quad (\text{A.1})$$

where d_i, r_i, k_i are positive for each $i = 1, \dots, n$ and have the same meanings as model (1.2). Here the solution $\mathbf{u} = (u_1, \dots, u_n)$ of (A.1) is positive means that $\mathbf{u} \gg \mathbf{0}$ (i.e., $u_i > 0$ in $\bar{\Omega}_i$ for each $i = 1, \dots, n$).

Theorem A.1. *Model (A.1) admits a unique positive solution $\mathbf{u}^* \in C^2(\bar{\Omega}_1) \times \dots \times C^2(\bar{\Omega}_n)$.*

Proof. Inspired by [42, 59], we define

$$\xi = \begin{cases} x, & x \in \Omega_1, \\ \xi_{i-1} + \prod_{j=1}^{i-1} p_j (x - x_{i-1}), & x \in \Omega_i, \quad i = 2, \dots, n, \end{cases}$$

where

$$\xi_i = \begin{cases} 0, & i = 0, \\ x_1, & i = 1, \\ x_1 + \sum_{\ell=1}^{i-1} \prod_{j=1}^{\ell} p_j (x_{\ell+1} - x_{\ell}), & i = 2, \dots, n. \end{cases}$$

Clearly, $x \in \Omega_i$ if and only if $\xi \in \tilde{\Omega}_i$ with $\tilde{\Omega}_i = (\xi_{i-1}, \xi_i)$. Then using the following scaling

$$w_i = \begin{cases} u_1, & i = 1, \\ \frac{u_i}{\prod_{j=1}^{i-1} p_j}, & i = 2, \dots, n, \end{cases}$$

we convert model (A.1) into the following equivalent form:

$$\begin{cases} D_i(w_i)_{\xi\xi} + r_i w_i \left(1 - \frac{w_i}{\tilde{k}_i}\right) = 0, & \xi \in \tilde{\Omega}_i, \quad i = 1, 2, \dots, n, \\ w_{i+1}(\xi_i^+) = w_i(\xi_i^-), \quad D_{i+1}(w_{i+1})_{\xi}(\xi_i^+) = D_i(w_i)_{\xi}(\xi_i^-), & i = 1, \dots, n-1, \\ (w_1)_{\xi}(\xi_0^+) = (w_n)_{\xi}(\xi_n^-) = 0, \end{cases} \quad (\text{A.2})$$

where

$$D_i = \begin{cases} d_1, & i = 1, \\ d_i \left(\prod_{j=1}^{i-1} p_j\right)^2, & i = 2, \dots, n, \end{cases} \quad \text{and} \quad \tilde{k}_i = \begin{cases} k_1, & i = 1, \\ \frac{k_i}{\prod_{j=1}^{i-1} p_j}, & i = 2, \dots, n. \end{cases}$$

Note that system (A.2) has continuous density and flux at interface. Then it follows from [29, Theorem A.15] and the proof of [29, Theorem 3.3] that system (A.2) admits a unique positive solution $\mathbf{w}^* \in \tilde{X}$, where

$$\tilde{X} := \{\mathbf{w} = (w_1, \dots, w_n) \in C([0, \xi_n]) : w_i \in C^2([\xi_{i-1}, \xi_i])\}.$$

Consequently, system (A.1) admits a unique positive solution $\mathbf{u}^* \in \mathbf{u}^* \in C^2(\overline{\Omega}_1) \times \dots \times C^2(\overline{\Omega}_n)$. This completes the proof. \square

We now present some properties of \mathbf{u}^* , which were originally derived in [59, Theorem 1] for the case $n = 2$. Here we generalize these properties to the case where n is finite but arbitrary.

Lemma A.2. *Let $\mathbf{u} \in C^2(\overline{\Omega}_1) \times \dots \times C^2(\overline{\Omega}_n)$ be the unique positive solution of (2.1), and let $\bar{\mathbf{k}}$ be defined in (1.3). Then the following statements hold:*

(i) *If $\mathbf{p} \gg \bar{\mathbf{k}}$, then $u_1^*(0^+) < k_1$, $u_n^*(L^-) > k_n$, $(u_i^*)_x < 0$ in Ω_i for $i = 1, \dots, n$, and*

$$(u_{i+1}^*)_x(x_i^+), (u_i^*)_x(x_i^-) < 0 \quad \text{for } i = 1, \dots, n-1; \quad (\text{A.3})$$

(ii) *If $\bar{\mathbf{k}} \gg \mathbf{p}$, then $u_1^*(0^+) > k_1$, $u_n^*(L^-) < k_n$, $(u_i^*)_x > 0$ in Ω_i for $i = 1, \dots, n$, and*

$$(u_{i+1}^*)_x(x_i^+), (u_i^*)_x(x_i^-) > 0 \quad \text{for } i = 1, \dots, n-1. \quad (\text{A.4})$$

Proof. We only prove (i), and part (ii) can be addressed similarly. The proof relies on the following three claims.

Claim 1: We claim that $(u_i^*)_x \leq 0$ in Ω_i for $i = 1, \dots, n$.

Proof of Claim: If the claim is not true, then there exist $1 \leq \ell_0 \leq n$ and $y_0 \in \Omega_{\ell_0}$ such that $(u_{\ell_0}^*)_x(y_0) > 0$. It follows from (2.1c) that y_1, y_2 are well-defined with $y_1 < y_2$, where

$$y_1 = \sup\{y : \exists 1 \leq \ell \leq \ell_0 \text{ s.t. } y \in [x_{\ell-1}, x_\ell) \text{ and } (u_\ell^*)_x(y^+) = 0\},$$

$$y_2 = \inf\{y : \exists \ell_0 \leq \ell \leq n \text{ s.t. } y \in (x_{\ell-1}, x_\ell] \text{ and } (u_\ell^*)_x(y^-) = 0\},$$

and there exist ℓ_1 and ℓ_2 with $1 \leq \ell_1 \leq \ell_0$ and $\ell_0 \leq \ell_2 \leq n$ such that $y_1 \in [x_{\ell_1-1}, x_{\ell_1})$ and $y_2 \in (x_{\ell_2-1}, x_{\ell_2}]$. By the definitions of y_1 and y_2 ,

$$(u_{\ell_1}^*)_{xx}(y_1^+) \geq 0, \quad (u_{\ell_2}^*)_{xx}(y_2^-) \leq 0, \quad (u_{\ell_1}^*)_x(y_1^+) = (u_{\ell_2}^*)_x(y_2^-) = 0, \quad (\text{A.5})$$

and

$$(u_i^*)_x > 0 \text{ in } \Omega_i \cap (y_1, y_2) \text{ for } i = \ell_1, \dots, \ell_2, \quad (\text{A.6a})$$

$$(u_{i+1}^*)_x(x_i^+), (u_i^*)_x(x_i^-) > 0 \text{ for } i = \ell_1, \dots, \ell_2 - 1. \quad (\text{A.6b})$$

It follows from (2.1a) and (A.5) that

$$u_{\ell_1}^*(y_1^+) \geq k_{\ell_1} \text{ and } u_{\ell_2}^*(y_2^-) \leq k_{\ell_2}, \quad (\text{A.7})$$

which contradicts (A.6a) if $\ell_1 = \ell_2$. If $\ell_1 < \ell_2$, we deduce from (A.6)-(A.7) that $k_{\ell_2} > u_{\ell_2}^*(x_{\ell_2-1}^+)$ and $u_{\ell_1}^*(x_{\ell_1}^-) > k_{\ell_1}$, and consequently,

$$k_{\ell_2} > u_{\ell_2}^*(x_{\ell_2-1}^+) = p_{\ell_2-1} u_{\ell_2-1}^*(x_{\ell_2-1}^-) > \dots > \left(\prod_{\ell=\ell_1}^{\ell_2-1} p_\ell \right) u_{\ell_1}^*(x_{\ell_1}^-) > k_{\ell_1},$$

where we have used (2.1b) and $\mathbf{p} \gg \bar{\mathbf{k}}$ in the second and last steps, respectively. This also leads to a contradiction. Thus, Claim 1 holds.

Claim 2: We claim that (i) holds for the case $n = 2$.

Proof of Claim: We first show that (A.3) holds for $n = 2$. Suppose not, then we see from (2.1b) and Claim 1 that $d_2(u_2^*)_x(x_1^+) = d_1(u_1^*)_x(x_1^-) = 0$, which implies that $u_i^* = k_i$ in Ω_i for each $i = 1, 2$. This combined with $\mathbf{p} \gg \bar{\mathbf{k}}$ yields $k_2 = u_2(x_1^+) = p_1 u_1(x_1^-) > k_2$, which is a contradiction. Therefore, (A.3) holds with $n = 2$. Noticing that $(u_1^*)_x(0^+) = 0$, by Claim 1 again, we have $(u_1^*)_{xx}(0^+) \leq 0$, which yields $u_1^*(0^+) \leq k_1$. If $u_1^*(0^+) = k_1$, then u_1^* satisfies

$$\begin{cases} d_1(u_1^*)_{xx} + r_1 u_1^* \left(1 - \frac{u_1^*}{k_1}\right) = 0, & 0 < x < x_1, \\ (u_1^*)_x(0^+) = 0, & u_1^*(0^+) = k_1. \end{cases} \quad (\text{A.8})$$

The uniqueness of the solution of (A.8) implies that $u_1^* \equiv k_1$, which contradicts (A.3) with $i = 1$. Thus, $u_1^*(0^+) < k_1$, and $u_2^*(L^-) > k_2$ can be proved similarly. Finally, we show that $(u_i^*)_x < 0$ in Ω_i for each $i = 1, 2$. Suppose not, then there exist $1 \leq \tilde{i} \leq 2$ and $\tilde{x} \in \Omega_{\tilde{i}}$ such that $(u_{\tilde{i}}^*)_x(\tilde{x}) \geq 0$. Without loss of generality, we assume that $\tilde{i} = 1$. Then it follows from Claim 1 that $(u_1^*)_x(\tilde{x}) = 0$, and consequently, $u_1^*(x) = k_1$ for $x \in [0, \tilde{x}]$, which contradicts $u_1^*(0^+) < k_1$. This proves the claim.

Claim 3: We claim that (i) holds for $n = m$ with $m \geq 3$ if it holds for $n = 2, \dots, m-1$.

Proof of Claim: We first show that

$$(u_i^*)_x < 0 \text{ in } \Omega_i \text{ for } i = 2, \dots, m-1. \quad (\text{A.9})$$

Assume to the contrary that (A.9) does not hold. In view of Claim 1, there exist $2 \leq \ell_* \leq m-1$ and $x_* \in \Omega_{\ell_*}$ such that $(u_{\ell_*}^*)_x(x_*) = 0$. Then (2.1) can be rewritten as the following two subsystems:

$$\begin{cases} d_i(u_i^*)_{xx} + r_i u_i^* \left(1 - \frac{u_i^*}{k_i}\right) = 0, & x \in \tilde{\Omega}_i, \quad i = 1, 2, \dots, \ell_*, \\ u_{i+1}^*(x_i^+) = p_i u_i^*(x_i^-), \quad d_{i+1}(u_{i+1}^*)_x(x_i^+) = d_i(u_i^*)_x(x_i^-), \\ (u_1^*)_x(0^+) = (u_{\ell_*}^*)_x(x_*^-) = 0, \end{cases} \quad (\text{A.10})$$

and

$$\begin{cases} d_i(u_i^*)_{xx} + r_i u_i^* \left(1 - \frac{u_i^*}{k_i}\right) = 0, & x \in \hat{\Omega}_i, \quad i = \ell_*, \dots, m, \\ u_{i+1}^*(x_i^+) = p_i u_i^*(x_i^-), \quad d_{i+1}(u_{i+1}^*)_x(x_i^+) = d_i(u_i^*)_x(x_i^-), \\ (u_{\ell_*}^*)_x(x_*^+) = (u_m^*)_x(L^-) = 0, \end{cases} \quad (\text{A.11})$$

where

$$\tilde{\Omega}_i = \begin{cases} \Omega_i, & i = 1, \dots, \ell_* - 1, \\ (x_{\ell_*-1}, x_*), & i = \ell_*, \end{cases}$$

and

$$\hat{\Omega}_i = \begin{cases} (x_*, x_{\ell_*}), & i = \ell_*, \\ \Omega_i, & i = \ell_* + 1, \dots, m. \end{cases}$$

Since $\mathbf{p} \gg \bar{\mathbf{k}}$, it follows that $[\mathbf{p}]_{\ell_*-1} \gg [\bar{\mathbf{k}}]_{\ell_*-1}$ and $[\mathbf{p}]_{m-\ell_*}^\# \gg [\bar{\mathbf{k}}]_{m-\ell_*}^\#$, where $[\cdot]_{\ell_*-1}$ and $[\cdot]_{m-\ell_*}^\#$ are defined in (1.4). Note that (i) holds for $n = 2, \dots, m-1$, and subsystems (A.10) and (A.11) consist of ℓ_* and $m - \ell_* + 1$ patches, respectively, where $2 \leq \ell_*, m - \ell_* + 1 \leq m-1$. Then $u_{\ell_*}^*(x_*^-) > k_{\ell_*}$ and $u_{\ell_*}^*(x_*^+) < k_{\ell_*}$, which is a contradiction. Thus, (A.9) holds.

Then we show that (A.3) holds for $n = m$. If it is not true, by Claim 1 again, there exists $1 \leq \tilde{\ell} \leq m - 1$ such that $(u_{\tilde{\ell}}^*)_x(x_{\tilde{\ell}}^-) = 0$. Then we can also rewrite (2.1) into the following two subsystems:

$$\begin{cases} d_i(u_i^*)_{xx} + r_i u_i^* \left(1 - \frac{u_i^*}{k_i}\right) = 0, & x \in \Omega_i, \quad i = 1, \dots, \tilde{\ell}, \end{cases} \quad (\text{A.12a})$$

$$\begin{cases} u_{i+1}^*(x_i^+) = p_i u_i^*(x_i^-), & d_{i+1}(u_{i+1}^*)_x(x_i^+) = d_i(u_i^*)_x(x_i^-), \end{cases} \quad (\text{A.12b})$$

$$\begin{cases} (u_1^*)_x(0^+) = (u_{\tilde{\ell}}^*)_x(x_{\tilde{\ell}}^-) = 0, \end{cases} \quad (\text{A.12c})$$

and

$$\begin{cases} d_i(u_i^*)_{xx} + r_i u_i^* \left(1 - \frac{u_i^*}{k_i}\right) = 0, & x \in \Omega_i, \quad i = \tilde{\ell} + 1, \dots, m, \end{cases} \quad (\text{A.13a})$$

$$\begin{cases} u_{i+1}^*(x_i^+) = p_i u_i^*(x_i^-), & d_{i+1}(u_{i+1}^*)_x(x_i^+) = d_i(u_i^*)_x(x_i^-), \end{cases} \quad (\text{A.13b})$$

$$\begin{cases} (u_{\tilde{\ell}+1}^*)_x(x_{\tilde{\ell}}^+) = (u_m^*)_x(L^-) = 0. \end{cases} \quad (\text{A.13c})$$

Here we remark that, if $\tilde{\ell} = 1$, (A.12) holds except for (A.12b), and if $\tilde{\ell} = m - 1$, (A.13) holds except for (A.13b).

We first obtain a contradiction for the case $2 \leq \tilde{\ell} \leq m - 2$. Since $\mathbf{p} \gg \bar{\mathbf{k}}$, it follows that $[\mathbf{p}]_{\tilde{\ell}-1} \gg [\bar{\mathbf{k}}]_{\tilde{\ell}-1}$ and $[\mathbf{p}]_{m-\tilde{\ell}-1}^\# \gg [\bar{\mathbf{k}}]_{m-\tilde{\ell}-1}^\#$. Note that (i) holds for $n = 2, \dots, m - 1$, and subsystems (A.12) and (A.13) consist of $\tilde{\ell}$ and $m - \tilde{\ell}$ patches, respectively, where $2 \leq \tilde{\ell}, m - \tilde{\ell} \leq m - 2$. Thus, $u_{\tilde{\ell}}^*(x_{\tilde{\ell}}^-) > k_{\tilde{\ell}}$ and $u_{\tilde{\ell}+1}^*(x_{\tilde{\ell}}^+) < k_{\tilde{\ell}+1}$, and consequently,

$$\frac{u_{\tilde{\ell}+1}^*(x_{\tilde{\ell}}^+)}{u_{\tilde{\ell}}^*(x_{\tilde{\ell}}^-)} = p_{\tilde{\ell}} < \frac{k_{\tilde{\ell}+1}}{k_{\tilde{\ell}}}, \quad (\text{A.14})$$

which contradicts $\mathbf{p} \gg \bar{\mathbf{k}}$. If $\tilde{\ell} = 1$, then $u_{\tilde{\ell}}^* = k_{\tilde{\ell}}$ in $\Omega_{\tilde{\ell}}$. Note that $[\mathbf{p}]_{m-\tilde{\ell}-1}^\# \gg [\bar{\mathbf{k}}]_{m-\tilde{\ell}-1}^\#$ and subsystems (A.13) consist of $m - 1$ patches. Then $u_{\tilde{\ell}+1}^*(x_{\tilde{\ell}}^+) < k_{\tilde{\ell}+1}$, which implies that (A.14) also holds. This leads to a contradiction. Using similar arguments, we can obtain a contradiction for the case $\tilde{\ell} = m - 1$. Therefore, (A.3) holds for $n = m$. Finally, using arguments similar to those in the proof of Claim 2, we can obtain that $u_1^*(0^+) < k_1$, $u_m^*(L^-) > k_m$ and $(u_i^*)_x < 0$ in Ω_i for $i = 1, m$. This completes the proof of the claim.

Therefore, (i) holds by Claims 2-3 and induction. \square

For the two-patch case, the steady state \mathbf{u}^* is necessarily monotonic. However, this monotonicity can be lost for $n \geq 3$. While Lemma A.2 provides sufficient conditions for monotonicity, the steady state may otherwise be non-monotonic, as illustrated by the concrete example below.

Proposition A.3. *Suppose that $n = 3$ and let \mathbf{u}^* be the unique positive solution of (2.1). If $k_1 = k_3 := k$ and $p_1 p_2 = 1$ with $p_1 \neq k_2/k$ (or equivalently, $p_2 \neq k/k_2$), then \mathbf{u}^* is non-monotonic.*

Proof. Without loss of generality, we assume that $p_1 > k_2/k$. Then it follows from $p_1 p_2 = 1$ that $p_2 < k/k_2$. We claim that there exists $x_* \in (x_1, x_2)$ such that $(u_2^*)_x(x_*) = 0$. If it is not true, then $(u_2^*)_x > 0$ or $(u_2^*)_x < 0$ in (x_1, x_2) . We only derive a contradiction for the case $(u_2^*)_x > 0$ in (x_1, x_2) , as the other case is analogous.

By continuity of $(u_2^*)_x$ on $[x_1, x_2]$, we have $(u_2^*)_x(x_1^+), (u_2^*)_x(x_2^-) \geq 0$. We now divide the arguments into two cases:

$$(a_1) \ (u_2^*)_x(x_1^+), (u_2^*)_x(x_2^-) > 0; \quad (a_2) \ (u_2^*)_x(x_1^+) = 0 \text{ or } (u_2^*)_x(x_2^-) = 0.$$

For case (a_1) , we see from (2.1c) that

$$y_1 := \sup\{y \in [0, x_1] : (u_1^*)_x(y^+) = 0\} \text{ and } y_2 := \inf\{y \in (x_2, L] : (u_3^*)_x(y^-) = 0\} \quad (\text{A.15})$$

are well-defined with $0 \leq y_1 < x_1 < x_2 < y_2 \leq L$. Thus, we have

$$(u_1^*)_{xx}(y_1^+) \geq 0, \quad (u_3^*)_{xx}(y_2^-) \leq 0 \quad (\text{A.16})$$

and

$$(u_i^*)_x > 0 \text{ in } \Omega_i \cap (y_1, y_2) \text{ for } i = 1, 2, 3. \quad (\text{A.17})$$

It follows from (2.1a) and (A.16) that $u_1^*(y_1^+) \geq k_1 = k$ and $u_3^*(y_2^-) \leq k_3 = k$. This combined with (A.17) implies that

$$k > u_3^*(x_2^+) = p_2 u_2^*(x_2^-) > p_2 u_2^*(x_1^+) = p_1 p_2 u_1^*(x_1^-) > k, \quad (\text{A.18})$$

where we have used (2.1b) in the second and fourth steps. This leads to a contradiction.

Now we derive a contradiction for case (a_2) . For simplicity, we only consider the case that $(u_2^*)_x(x_1^+) = 0$ and $(u_2^*)_x(x_2^-) > 0$, as other cases can be treated similarly. Then \mathbf{u}^* satisfies

$$\begin{cases} d_1(u_1^*)_{xx} + r_1 u_1^* \left(1 - \frac{u_1^*}{k}\right) = 0, & 0 < x < x_1, \\ (u_1^*)_x(0) = (u_1^*)_x(x_1^-) = 0. \end{cases} \quad (\text{A.19})$$

The uniqueness of the solution to (A.19) implies $u_1^* \equiv k$ in $[0, x_1]$. Thus, by an argument similar to that in (A.18), we obtain

$$k \geq u_3^*(y_2^-) > u_3^*(x_2^+) = p_2 u_2^*(x_2^-) > p_2 u_2^*(x_1^+) = u_1^*(x_1^-) = k,$$

where y_2 is defined as in (A.15). This also leads to a contradiction. Therefore, the claim holds.

Then we can rewrite (2.1) into the following two subsystems:

$$\begin{cases} d_i(u_i^*)_{xx} + r_i u_i^* \left(1 - \frac{u_i^*}{k_i}\right) = 0, & x \in \tilde{\Omega}_i, \quad i = 1, 2, \\ u_2^*(x_1^+) = p_1 u_1^*(x_1^-), \quad d_2(u_2^*)_x(x_1^+) = d_1(u_1^*)_x(x_1^-), \\ (u_1^*)_x(0^+) = (u_2^*)_x(x_*^-) = 0, \end{cases}$$

and

$$\begin{cases} d_i(u_i^*)_{xx} + r_i u_i^* \left(1 - \frac{u_i^*}{k_i}\right) = 0, & x \in \hat{\Omega}_i, \quad i = 2, 3, \\ u_3^*(x_2^+) = p_2 u_2^*(x_2^-), \quad d_3(u_3^*)_x(x_2^+) = d_2(u_2^*)_x(x_2^-), \\ (u_2^*)_x(x_*^+) = (u_3^*)_x(L^-) = 0, \end{cases}$$

where

$$\tilde{\Omega}_i = \begin{cases} \Omega_1, & i = 1, \\ (x_1, x_*), & i = 2, \end{cases}$$

and

$$\hat{\Omega}_i = \begin{cases} (x_*, x_2), & i = 2, \\ \Omega_3, & i = 3. \end{cases}$$

Observing that $p_1 > k_2/k$ and $p_2 < k/k_2$, by applying Lemma A.2 to each subsystem, we see $(u_i^*)_x < 0$ in $\tilde{\Omega}_i$ for $i = 1, 2$, and $(u_i^*)_x > 0$ in $\hat{\Omega}_i$ for $i = 2, 3$. Consequently, \mathbf{u}^* is non-monotonic on $[0, L]$. \square

Appendix B Comparison principle

Throughout this section, we define an order in $(C(\bar{\Omega}_1) \times \cdots \times C(\bar{\Omega}_n))^2$: $(\mathbf{u}^1, \mathbf{v}^1) \succeq (\mathbf{u}^2, \mathbf{v}^2)$ if and only if

$$u_i^1(x) \geq u_i^2(x) \quad \text{and} \quad v_i^1(x) \leq v_i^2(x) \quad \text{for all } x \in \bar{\Omega}_i \quad \text{and } i = 1, \dots, n, \quad (\text{B.1})$$

Then we show that system (1.2) preserves the order “ \succeq ”.

Theorem B.1. *Let $\bar{\mathbf{u}}, \underline{\mathbf{v}}, \bar{\mathbf{v}}, \underline{\mathbf{u}} \in Y^+$, where*

$$\begin{aligned} Y^+ &:= \{(u_1, \dots, u_n) : u_i \in C^{2,1}(\bar{\Omega}_i \times (0, T)) \cap C(\bar{\Omega}_i \times [0, T]) \\ &\quad u_i \geq 0 \text{ in } \bar{\Omega}_i \times [0, T) \text{ for all } i = 1, \dots, n\} \end{aligned}$$

with $T \in (0, +\infty]$. Suppose that $(\bar{\mathbf{u}}, \bar{\mathbf{v}})$ and $(\underline{\mathbf{u}}, \underline{\mathbf{v}})$ satisfy, for $i = 1, \dots, n-1$,

$$\begin{aligned} u_{i+1}(x_i^+, t) &= p_i u_i(x_i^-, t), \quad d_{i+1}(u_{i+1})_x(x_i^+, t) = d_i(u_i)_x(x_i^-, t), \quad t > 0, \\ v_{i+1}(x_i^+, t) &= \hat{p}_i v_i(x_i^-, t), \quad \hat{d}_{i+1}(v_{i+1})_x(x_i^+, t) = \hat{d}_i(v_i)_x(x_i^-, t), \quad t > 0, \end{aligned} \quad (\text{B.2})$$

and for $i = 1, \dots, n$,

$$\begin{cases} (\bar{u}_i)_t - d_i(\bar{u}_i)_{xx} - f_i(\bar{u}_i, \bar{v}_i) \geq 0 \geq (\underline{u}_i)_t - d_i(\underline{u}_i)_{xx} - f_i(\underline{u}_i, \bar{v}_i), & x \in \Omega_i, \quad t > 0, \\ (\bar{v}_i)_t - \hat{d}_i(\bar{v}_i)_{xx} - \hat{f}_i(\underline{u}_i, \bar{v}_i) \geq 0 \geq (\underline{v}_i)_t - \hat{d}_i(\underline{v}_i)_{xx} - \hat{f}_i(\bar{u}_i, \underline{v}_i), & x \in \Omega_i, \quad t > 0, \\ (\bar{u}_1)_x(0^+, t) \leq 0 \leq (\underline{u}_1)_x(0^+, t), \quad (\bar{u}_n)_x(L^-, t) \geq 0 \geq (\underline{u}_n)_x(L^-, t), & t > 0, \\ (\bar{v}_1)_x(0^+, t) \leq 0 \leq (\underline{v}_1)_x(0^+, t), \quad (\bar{v}_n)_x(L^-, t) \geq 0 \geq (\underline{v}_n)_x(L^-, t), & t > 0, \end{cases} \quad (\text{B.3})$$

where

$$f_i(u_i, v_i) = r_i u_i \left(1 - \frac{u_i + v_i}{k_i} \right), \quad \hat{f}_i(u_i, v_i) = r_i v_i \left(1 - \frac{u_i + v_i}{k_i} \right). \quad (\text{B.4})$$

If $(\bar{\mathbf{u}}(\cdot, 0), \bar{\mathbf{v}}(\cdot, 0)) \succeq (\underline{\mathbf{u}}(\cdot, 0), \underline{\mathbf{v}}(\cdot, 0))$, then $(\bar{\mathbf{u}}(\cdot, t), \bar{\mathbf{v}}(\cdot, t)) \succeq (\underline{\mathbf{u}}(\cdot, t), \underline{\mathbf{v}}(\cdot, t))$ for any $t \in (0, T)$.

Proof. Fix any $T_0 \in (0, T)$, we denote

$$M := \max_{1 \leq i \leq n} (\|\bar{u}_i\|_\infty, \|\underline{u}_i\|_\infty, \|\bar{v}_i\|_\infty, \|\underline{v}_i\|_\infty),$$

where $\|w_i\|_\infty = \max_{(x,t) \in \bar{\Omega}_i \times [0, T_0]} |w_i(x, t)|$ for $w_i = \bar{u}_i, \underline{u}_i, \bar{v}_i, \underline{v}_i$, and let

$$\mu := \max_{1 \leq i \leq n} \left\{ \max_{[0, M] \times [0, M]} \left(\left| \frac{\partial f_i}{\partial u_i} \right| + \left| \frac{\partial f_i}{\partial v_i} \right| + \left| \frac{\partial \hat{f}_i}{\partial u_i} \right| + \left| \frac{\partial \hat{f}_i}{\partial v_i} \right| \right) \right\}.$$

Inspired by [22], we define, for $i = 1, \dots, n$,

$$\begin{aligned} \theta_i(x, t) &:= (\bar{u}_i(x, t) - \underline{u}_i(x, t))e^{-\mu t} + \epsilon(t+1), \\ \rho_i(x, t) &:= (\bar{v}_i(x, t) - \underline{v}_i(x, t))e^{-\mu t} + \epsilon(t+1). \end{aligned}$$

Then it follows from the mean value theorem that there exist

$$\begin{aligned} a_i(x, t) &:= \frac{\partial f_i}{\partial u_i}(\eta_i(x, t), \zeta_i(x, t)), \quad b_i(x, t) := \frac{\partial f_i}{\partial v_i}(\eta_i(x, t), \zeta_i(x, t)) \leq 0, \\ \hat{a}_i(x, t) &:= \frac{\partial \hat{f}_i}{\partial u_i}(\hat{\eta}_i(x, t), \hat{\zeta}_i(x, t)) \leq 0, \quad \hat{b}_i(x, t) := \frac{\partial \hat{f}_i}{\partial v_i}(\hat{\eta}_i(x, t), \hat{\eta}_i(x, t)), \end{aligned}$$

where $0 \leq \eta_i(x, t), \zeta_i(x, t), \hat{\eta}_i(x, t), \hat{\zeta}_i(x, t) \leq M$ for $(x, t) \in \Omega_i \times (0, T_0]$, such that

$$\begin{aligned}
\mathcal{L}_i(\theta_i, \rho_i) &:= (\theta_i)_t - d_i(\theta_i)_{xx} + (\mu - a_i(x, t))\theta_i + b_i(x, t)\rho_i \\
&= e^{-\mu t} [(\bar{u}_i - \underline{u}_i)_t - d_i(\bar{u}_i - \underline{u}_i)_{xx} - a_i(x, t)(\bar{u}_i - \underline{u}_i) + b_i(x, t)(\bar{v}_i - \underline{v}_i)] \\
&\quad + \epsilon + (\mu - a_i(x, t) + b_i(x, t))\epsilon(t+1) > 0 \text{ in } \Omega_i \times (0, T_0], \\
\hat{\mathcal{L}}_i(\theta_i, \rho_i) &:= (\rho_i)_t - \hat{d}_i(\rho_i)_{xx} + \left(\mu - \hat{b}_i(x, t)\right)\rho_i + \hat{a}_i(x, t)\theta_i \\
&= e^{-\mu t} \left((\bar{v}_i - \underline{v}_i)_t - \hat{d}_i(\bar{v}_i - \underline{v}_i)_{xx} - \hat{b}_i(x, t)(\bar{v}_i - \underline{v}_i) + \hat{a}_i(x, t)(\bar{u}_i - \underline{u}_i) \right) \\
&\quad + \epsilon + \left(\mu - \hat{b}_i(x, t) + \hat{a}_i(x, t)\right)\epsilon(t+1) > 0 \text{ in } \Omega_i \times (0, T_0].
\end{aligned} \tag{B.5}$$

Moreover,

$$d_i(\theta_i)_x(x_i^-, t) = d_{i+1}(\theta_{i+1})_x(x_i^+, t), \quad \hat{d}_i(\rho_i)_x(x_i^-, t) = \hat{d}_{i+1}(\rho_{i+1})_x(x_i^+, t), \tag{B.6a}$$

$$(\theta_1)_x(0^+, t), (\rho_1)_x(0^+, t) \leq 0, \quad (\theta_n)_x(L^-, t), (\rho_n)_x(L^-, t) \geq 0, \tag{B.6b}$$

$$\theta_i(\cdot, 0), \rho_i(\cdot, 0) > 0 \text{ in } \bar{\Omega}_i. \tag{B.6c}$$

We first prove that $\theta_i, \rho_i > 0$ in $(x, t) \in \bar{\Omega}_i \times [0, T_0]$ for $i = 1, \dots, n$. Suppose, for the sake of contradiction, that this is not true. Then, by (B.6c), at least one of the following quantities is well-defined and lies in the interval $(0, T_0]$:

$$t_1 := \inf\{t \in [0, T_0] : \exists 1 \leq i_0 \leq n \text{ and } y_0 \in \bar{\Omega}_{i_0} \text{ s.t. } \theta_{i_0}(y_0, t) = 0\},$$

$$t_2 := \inf\{t \in [0, T_0] : \exists 1 \leq i_0 \leq n \text{ and } y_0 \in \bar{\Omega}_{i_0} \text{ s.t. } \rho_{i_0}(y_0, t) = 0\}.$$

Without loss of generality, we assume that both t_1 and t_2 exist satisfying $t_1 \leq t_2$. By the definition of t_1 and t_2 , we have

$$\begin{aligned}
\theta_{i_0}(x, t) &> 0 \text{ for } x \in \Omega_{i_0}, t \in [0, t_1), \\
\rho_{i_0}(x, t) &\geq 0 \text{ for } x \in \Omega_{i_0}, t \in [0, t_1].
\end{aligned} \tag{B.7}$$

Now we claim that

$$\theta_{i_0}(x, t_1) > 0 \text{ for } x \in \Omega_{i_0}. \tag{B.8}$$

If it is not true, then there exists $y_0 \in \Omega_{i_0}$ such that $\theta_{i_0}(y_0, t_1) = 0$. Then $(\theta_{i_0})_t|_{(y_0, t_1)} \leq 0$, $(\theta_{i_0})_x|_{(y_0, t_1)} = 0$ and $(\theta_{i_0})_{xx}|_{(y_0, t_1)} \geq 0$. Thus,

$$\begin{aligned}
\mathcal{L}_{i_0}(\theta_{i_0}, \rho_{i_0})|_{(y_0, t_1)} &= (\theta_{i_0})_t|_{(y_0, t_1)} - d_{i_0}(\theta_{i_0})_{xx}|_{(y_0, t_1)} \\
&\quad + (\mu - a_{i_0}(y_0, t_1))\theta_{i_0}(y_0, t_1) + b_{i_0}(y_0, t_1)\rho_{i_0}(y_0, t_1) \leq 0,
\end{aligned}$$

which contradicts (B.5). Therefore, the claim is true and (B.8) holds. Then, by the definition of t_1 again, we see that $\theta_{i_0}(x_{i_0-1}^+, t_1) = 0$ or $\theta_{i_0}(x_{i_0}^-, t_1) = 0$. Now we obtain a

contradiction for the case $\theta_{i_0}(x_{i_0}^-, t_1) = 0$, and the other case can be treated similarly. If $i_0 = n$, it follows from (B.7)-(B.8) and the Hopf's lemma that $(\theta_{i_0})_x(x_{i_0}^-, t_0) < 0$, which contradicts (B.6b). If $i_0 < n$, by (B.7)-(B.8) and the Hopf's lemma again, we have $(\theta_{i_0})_x(x_{i_0}^-, t_0) < 0$ and $(\theta_{i_0+1})_x(x_{i_0}^+, t_0) > 0$, which contradicts (B.6a).

Thus, $\theta_i, \rho_i > 0$ in $\bar{\Omega}_i \times [0, T_0]$ for $i = 1, \dots, n$. Since $\epsilon > 0$ and $T_0 \in (0, T)$ are arbitrary, we conclude that $\bar{u}_i \geq \underline{u}_i$ and $\bar{v}_i \leq \underline{v}_i$ in $\bar{\Omega}_i \times (0, T)$ for $i = 1, \dots, n$. This completes the proof. \square

Remark B.2. We remark that if condition $(\bar{\mathbf{u}}(\cdot, 0), \bar{\mathbf{v}}(\cdot, 0)) \succeq (\underline{\mathbf{u}}(\cdot, 0), \underline{\mathbf{v}}(\cdot, 0))$ in Theorem B.1 is replaced by $(\bar{\mathbf{u}}(\cdot, 0), \bar{\mathbf{v}}(\cdot, 0)) \succeq (\neq)(\underline{\mathbf{u}}(\cdot, 0), \underline{\mathbf{v}}(\cdot, 0))$, one can obtain that $\bar{\mathbf{u}}(x, t) \gg \underline{\mathbf{u}}(x, t)$ and $\bar{\mathbf{v}}(x, t) \gg \underline{\mathbf{v}}(x, t)$ for all $x \in [0, L]$ and $t \in (0, T]$. The proof is also based on the comparison principle for classical reaction-diffusion equations, and here we omit the proof for simplicity.

By Theorem B.1 and the definitions of sub-equilibrium and super-equilibrium (see [27, Chapter I]), we can obtain the following result.

Proposition B.3. *Let $\bar{\mathbf{w}}, \bar{\mathbf{z}}, \underline{\mathbf{w}}, \underline{\mathbf{z}} \in X^+$ with*

$$X^+ := \{\mathbf{u} = (u_1, \dots, u_n) : u_i \in C^2(\bar{\Omega}_i) \text{ and } u_i \geq 0 \text{ in } \bar{\Omega}_i \text{ for all } i = 1, \dots, n\}. \quad (\text{B.9})$$

Assume that $(\bar{\mathbf{w}}, \underline{\mathbf{z}})$ and $(\underline{\mathbf{w}}, \bar{\mathbf{z}})$ satisfy (3.1c)-(3.1d), and for $i = 1, \dots, n$,

$$\begin{cases} -d_i(\bar{w}_i)_{xx} - f_i(\bar{w}_i, \bar{z}_i) \geq 0 \geq -d_i(\underline{w}_i)_{xx} - f_i(\underline{w}_i, \bar{z}_i), & x \in \Omega_i, \\ -\hat{d}_i(\bar{z}_i)_{xx} - \hat{f}_i(\underline{w}_i, \bar{z}_i) \geq 0 \geq -\hat{d}_i(\underline{z}_i)_{xx} - \hat{f}_i(\bar{w}_i, \underline{z}_i), & x \in \Omega_i, \\ (\bar{w}_1)_x(0^+) \leq 0 \leq (\underline{w}_1)_x(0^+), \quad (\bar{w}_n)_x(L^-) \geq 0 \geq (\underline{w}_n)_x(L^-), \\ (\bar{z}_1)_x(0^+) \leq 0 \leq (\underline{z}_1)_x(0^+), \quad (\bar{z}_n)_x(L^-) \geq 0 \geq (\underline{z}_n)_x(L^-). \end{cases}$$

Then $(\bar{\mathbf{w}}, \underline{\mathbf{z}})$ (resp. $(\underline{\mathbf{w}}, \bar{\mathbf{z}})$) is a super-equilibrium (resp. sub-equilibrium) of system (1.2).

Then, by [27, Lemma 1.1], we have the following result.

Theorem B.4. *Let $(\bar{\mathbf{w}}, \underline{\mathbf{z}})$ be a super-equilibrium of system (1.2), and let $(\underline{\mathbf{w}}, \bar{\mathbf{z}})$ be a sub-equilibrium of system (1.2) with $\bar{\mathbf{w}}, \bar{\mathbf{z}}, \underline{\mathbf{w}}, \underline{\mathbf{z}} \in X^+$ (defined in (B.9)). If $(\bar{\mathbf{w}}, \underline{\mathbf{z}}) \succeq (\underline{\mathbf{w}}, \bar{\mathbf{z}})$, then system (1.2) admits a steady state (\mathbf{u}, \mathbf{v}) satisfying $(\bar{\mathbf{w}}, \underline{\mathbf{z}}) \succeq (\mathbf{u}, \mathbf{v}) \succeq (\underline{\mathbf{w}}, \bar{\mathbf{z}})$.*

We end this section with a comparison principle for an elliptic equation.

Theorem B.5. Assume that, for each $i = 1, \dots, n$, the function $c_i(x)$ is positive and bounded. If $(u_1, \dots, u_n) \in C^2(\overline{\Omega}_1) \times \dots \times C^2(\overline{\Omega}_n)$ satisfies

$$\begin{cases} -d_i(u_i)_{xx} + c_i(x)u_i = f_i(x), & x \in \Omega_i, \quad i = 1, 2, \dots, n, & (\text{B.10a}) \\ u_{i+1}(x_i^+) = p_i u_i(x_i^-), & d_{i+1}(u_{i+1})_x(x_i^+) = d_i(u_i)_x(x_i^-), & (\text{B.10b}) \\ (u_1)_x(0^+) \leq 0 \leq (u_n)_x(L^-), & & (\text{B.10c}) \end{cases}$$

where $f_i \geq 0$ in $\overline{\Omega}_i$ for all $i = 1, \dots, n$, and there exists $1 \leq i^* \leq n$ such that $f_{i^*} \not\equiv 0$, then $u_i > 0$ in $\overline{\Omega}_i$ for all $i = 1, \dots, n$.

Proof. The proof follows the technique of [40, Proposition 3.3], which we generalize to the n -patch case. Let $m := \min_{1 \leq i \leq n, x \in \overline{\Omega}_i} u_i(x)$. We first prove that $m \geq 0$. Suppose, for contradiction, that $m < 0$. We claim that

$$u_i > m \text{ in } \Omega_i \text{ for each } i = 1, \dots, n. \quad (\text{B.11})$$

If this claim is false, then there exists $1 \leq i_0 \leq n$ and $y_0 \in \Omega_{i_0}$ such that $u_{i_0}(y_0) = m < 0$. Then $(u_{i_0})_{xx}(y_0) \geq 0$, which implies that

$$-d_{i_0}(u_{i_0})_{xx}(y_0) + c_{i_0}(y_0)u_{i_0}(y_0) \leq c_{i_0}(y_0)u_{i_0}(y_0) < 0,$$

contradicting (B.10a). Therefore, (B.11) holds, and consequently, there exists $1 \leq i_1 \leq n$ such that $u_{i_1}(x_{i_1-1}^-) = m$ or $u_{i_1}(x_{i_1}^+) = m$. Without loss of generality, we assume that $u_{i_1}(x_{i_1}^-) = m$. If $i_1 = n$, it follows from the Hopf's Lemma and (B.11) that $(u_n)_x(x_n^-) < 0$, which contradicts (B.10c). If $1 \leq i_1 < n$, by the Hopf's Lemma and (B.11) again, we have $(u_{i_1})_x(x_{i_1}^-) < 0$ and $(u_{i_1+1})_x(x_{i_1}^+) > 0$, which contradicts (B.10b). Therefore, $m \geq 0$.

Now we prove that $m > 0$. First, we claim that

$$u_{i^*} > 0 \text{ in } \Omega_{i^*}. \quad (\text{B.12})$$

If this is false, the strong maximum principle implies that $u_{i^*} \equiv 0$ in Ω_{i^*} . This contradicts (B.10a) since $f_{i^*} \not\equiv 0$. Therefore, (B.12) holds. Next, we show that $u_{i^*}(x_{i^*}^-) > 0$, and the proof for $u_{i^*}(x_{i^*-1}^+) > 0$ is similar. Suppose to the contrary that $u_{i^*}(x_{i^*}^-) = 0$. If $i^* = n$, it follows from the Hopf's Lemma and (B.12) that $(u_{i^*})_x(x_{i^*}^-) < 0$, which contradicts (B.10c). If $1 \leq i^* < n$, the Hopf's Lemma and (B.11) again gives $(u_{i^*})_x(x_{i^*}^-) < 0$. Since $u_{i^*+1} \geq 0$ in $\overline{\Omega}_{i^*+1}$, we have $(u_{i^*+1})_x(x_{i^*}^+) \geq 0$, which contradicts (B.10b). Therefore, $u_{i^*} > 0$ in $\overline{\Omega}_{i^*}$. Finally, we show that $u_i > 0$ in $\overline{\Omega}_i$ for all $i \neq i^*$. Without loss of generality, we assume that $1 \leq i^* < n$. The strong maximum principle gives $u_{i^*+1} > 0$

in Ω_{i^*+1} . Using arguments similar to those in the proof of $u_{i^*}(x_{i^*}^-) > 0$, we see that $u_{i^*+1}(x_{i^*+1}^-) > 0$, and consequently, $u_{i^*+1} > 0$ in $\overline{\Omega}_{i^*+1}$. The result for all remaining i follows by induction. \square

Appendix C Set-up of the problem

Following [22], we define a classical solution to model (1.2) as follows:

Definition C.1. For $T \in (0, +\infty]$, we say that a function

$$(\mathbf{u}, \mathbf{v}) = (u_1, \dots, u_n, v_1, \dots, v_n) : [0, L] \times [0, T) \rightarrow \mathbb{R}^n \times \mathbb{R}^n$$

is a classical solution of model (1.2) in $[0, L] \times [0, T)$ if:

- (i) u_i, v_i is of class $C^{2,1}(\overline{\Omega}_i \times (0, T)) \cap C(\overline{\Omega}_i \times [0, T))$ for each $i = 1, \dots, n$;
- (ii) All identities in (1.2) are satisfied pointwise for $0 < t < T$.

To show the existence of a classical solution of system (1.2), we first analyze the abstract evolution equation based on semigroup theory, inspired by [22, 40]. Since their models differ slightly from ours, the associated function spaces and operators must be adapted accordingly. For completeness, we now detail these necessary modifications. We begin by defining the function spaces

$$X_u, X_v := L^2(\overline{\Omega}_1) \times \dots \times L^2(\overline{\Omega}_n) \text{ and } X := X_u \times X_v, \quad (\text{C.1})$$

equipped with the following inner products:

$$\begin{aligned} \langle \mathbf{u}, \hat{\mathbf{u}} \rangle_{X_u} &= \sum_{i=1}^n \frac{1}{\prod_{j=1}^{i-1} p_j} \langle u_i, \hat{u}_i \rangle_{L^2(\overline{\Omega}_i)} \quad \text{for } \mathbf{u}, \hat{\mathbf{u}} \in X_u, \\ \langle \mathbf{v}, \hat{\mathbf{v}} \rangle_{X_v} &= \sum_{i=1}^n \frac{1}{\prod_{j=1}^{i-1} \hat{p}_j} \langle v_i, \hat{v}_i \rangle_{L^2(\overline{\Omega}_i)} \quad \text{for } \mathbf{v}, \hat{\mathbf{v}} \in X_v, \\ \langle \mathbf{w}, \mathbf{z} \rangle_X &= \langle \mathbf{u}, \hat{\mathbf{u}} \rangle_{X_u} + \langle \mathbf{v}, \hat{\mathbf{v}} \rangle_{X_v} \quad \text{for } \mathbf{w} = (\mathbf{u}, \mathbf{v}), \mathbf{z} = (\hat{\mathbf{u}}, \hat{\mathbf{v}}) \in X. \end{aligned}$$

These inner products induce the norms $\|\cdot\|_{X_u} = \sqrt{\langle \cdot, \cdot \rangle_{X_u}}$, $\|\cdot\|_{X_v} = \sqrt{\langle \cdot, \cdot \rangle_{X_v}}$ and $\|\cdot\|_X = \sqrt{\langle \cdot, \cdot \rangle_X}$. Equipped with these norms, X_u, X_v and X become Hilbert spaces, respectively.

Then define the function spaces

$$\begin{aligned} \mathcal{H}_u^1 &:= \{\mathbf{u} \in H^1(\overline{\Omega}_1) \times \dots \times H^1(\overline{\Omega}_n) : u_{i+1}(x_i^+) = p_i u_i(x_i^-), i = 1, \dots, n-1\}, \\ \mathcal{H}_v^1 &:= \{\mathbf{v} \in H^1(\overline{\Omega}_1) \times \dots \times H^1(\overline{\Omega}_n) : v_{i+1}(x_i^+) = \hat{p}_i v_i(x_i^-), i = 1, \dots, n-1\}, \\ \mathcal{H}^1 &= \mathcal{H}_u^1 \times \mathcal{H}_v^1, \end{aligned} \quad (\text{C.2})$$

equipped with the following norms:

$$\begin{aligned}\|\mathbf{u}\|_{\mathcal{H}_u^1} &= \sum_{i=1}^n \frac{1}{\prod_{j=1}^{i-1} p_j} \|u_i\|_{H^1(\overline{\Omega}_i)} \quad \text{for } \mathbf{u} \in \mathcal{H}_u^1, \\ \|\mathbf{v}\|_{\mathcal{H}_v^1} &= \sum_{i=1}^n \frac{1}{\prod_{j=1}^{i-1} \hat{p}_j} \|v_i\|_{H^1(\overline{\Omega}_i)} \quad \text{for } \mathbf{v} \in \mathcal{H}_v^1, \\ \|\mathbf{w}\|_{\mathcal{H}^1} &= \|\mathbf{u}\|_{\mathcal{H}_u^1} + \|\mathbf{v}\|_{\mathcal{H}_v^1} \quad \text{for } \mathbf{w} = (\mathbf{u}, \mathbf{v}) \in \mathcal{H}^1.\end{aligned}$$

By the Sobolev embedding theorem, function spaces \mathcal{H}_u^1 and \mathcal{H}_v^1 are closed subspaces of the product space $H^1(\overline{\Omega}_1) \times \cdots \times H^1(\overline{\Omega}_n)$, which implies that \mathcal{H}_u^1 , \mathcal{H}_v^1 and \mathcal{H}^1 are Banach spaces.

We further define, for any integer $k \geq 2$, the function spaces

$$\mathcal{H}_u^k, \mathcal{H}_v^k := H^k(\overline{\Omega}_1) \times \cdots \times H^k(\overline{\Omega}_n) \quad \text{and} \quad \mathcal{H}^k := \mathcal{H}_u^k \times \mathcal{H}_v^k \quad (\text{C.3})$$

equipped with the following norms, respectively:

$$\begin{aligned}\|\mathbf{u}\|_{\mathcal{H}_u^k} &= \sum_{i=1}^n \frac{1}{\prod_{j=1}^{i-1} p_j} \|u_i\|_{H^k(\overline{\Omega}_i)} \quad \text{for } \mathbf{u} \in \mathcal{H}_u^k, \\ \|\mathbf{v}\|_{\mathcal{H}_v^k} &= \sum_{i=1}^n \frac{1}{\prod_{j=1}^{i-1} \hat{p}_j} \|v_i\|_{H^k(\overline{\Omega}_i)} \quad \text{for } \mathbf{v} \in \mathcal{H}_v^k, \\ \|\mathbf{w}\|_{\mathcal{H}^k} &= \|\mathbf{u}\|_{\mathcal{H}_u^k} + \|\mathbf{v}\|_{\mathcal{H}_v^k} \quad \text{for } \mathbf{w} = (\mathbf{u}, \mathbf{v}) \in \mathcal{H}^k.\end{aligned}$$

Clearly, function spaces \mathcal{H}_u^k , \mathcal{H}_v^k and \mathcal{H}^k are Banach spaces for all $k \geq 2$.

Finally, define the function space

$$G = \{\mathbf{u} \in C^1(\overline{\Omega}_1) \times \cdots \times C^1(\overline{\Omega}_n) : u_{i+1}(x_i^+) = p_i u_i(x_i^-), i = 1, \dots, n-1\}, \quad (\text{C.4})$$

equipped with the norm

$$\|\mathbf{u}\|_G = \sum_{i=1}^n \frac{1}{\prod_{j=1}^{i-1} p_j} \|u_i\|_{C^1(\overline{\Omega}_i)}.$$

Clearly, G is a Banach space.

We now define an operator \mathcal{A} on function space X as follows:

$$\mathcal{A}\mathbf{w} = (\mathcal{A}_u \mathbf{u}, \mathcal{A}_v \mathbf{v}) \quad (\text{C.5})$$

where

$$\begin{aligned}\mathcal{A}_u \mathbf{u} &:= (-d_1(u_1)_{xx} + u_1, \dots, -d_n(u_n)_{xx} + u_n), \\ \mathcal{A}_v \mathbf{v} &:= (-\hat{d}_1(v_1)_{xx} + v_1, \dots, -\hat{d}_n(v_n)_{xx} + v_n).\end{aligned} \quad (\text{C.6})$$

The domain of \mathcal{A} is given by:

$$\begin{aligned} \mathcal{D}(\mathcal{A}) := \{ \mathbf{w} = (\mathbf{u}, \mathbf{v}) \in \mathcal{H}^2 \cap \mathcal{H}^1 : (u_1)_x(0^+) = (u_n)_x(L^-) = (v_1)_x(0^+) = (v_n)_x(L^-) = 0, \\ d_{i+1}(u_{i+1})_x(x_i^+) = d_i(u_i)_x(x_i^-), \hat{d}_{i+1}(v_{i+1})_x(x_i^+) = \hat{d}_i(v_i)_x(x_i^-) \}. \end{aligned} \quad (\text{C.7})$$

By the Sobolev embedding theorem again, $\mathcal{D}(\mathcal{A})$ is a closed subspace of \mathcal{H}^2 and thus a Banach space.

Then we formulate model (1.2) as the following abstract evolution equation:

$$\frac{d}{dt} \mathbf{w} + \mathcal{A} \mathbf{w} = F(\mathbf{w}), \quad (\text{C.8})$$

where $\mathbf{w} = (\mathbf{u}, \mathbf{v}) = (u_1, \dots, u_n, v_1, \dots, v_n) \in X$ and

$$\begin{aligned} F(\mathbf{w}) \\ := \left(f_1(u_1, v_1) + u_1, \dots, f_n(u_n, v_n) + u_n, \hat{f}_1(u_1, v_1) + v_1, \dots, \hat{f}_n(u_n, v_n) + v_n \right) \end{aligned} \quad (\text{C.9})$$

with f_i, \hat{f}_i be defined in (B.4) for $i = 1, \dots, n$.

We now show that $-\mathcal{A}$ generates an analytic semigroup on X .

Lemma C.2. *The linear operator $\mathcal{A} : \mathcal{D}(\mathcal{A}) \subset X \rightarrow X$ is symmetric maximal monotone, and $-\mathcal{A}$ is the infinitesimal generator of an analytic semigroup on X , where X , \mathcal{A} and $\mathcal{D}(\mathcal{A})$ are defined in (C.1), (C.5) and (C.7), respectively. Moreover, \mathcal{A} has a compact resolvent.*

Proof. Since \mathcal{A}_u and \mathcal{A}_v are essentially identical, it suffices to establish the properties for \mathcal{A}_u . Although the proof technique parallels that of [22, Lemma 3.3] (see also [40, Proposition 7.1]), several adaptations are required. We detail these modifications below and omit the analogous steps.

Define

$$\begin{aligned} \mathcal{D}(\mathcal{A}_u) := \{ \mathbf{u} \in \mathcal{H}_u^2 \cap \mathcal{H}_u^1 : d_{i+1}(u_{i+1})_x(x_i^+) = d_i(u_i)_x(x_i^-), i = 1, \dots, n-1, \\ (u_1)_x(0^+) = (u_n)_x(L^-) = 0 \}, \end{aligned} \quad (\text{C.10})$$

where \mathcal{H}_u^1 and \mathcal{H}_u^2 are defined in (C.2) and (C.3), respectively. For any $\mathbf{u} \in \mathcal{D}(\mathcal{A}_u)$, we use integration by parts and apply the interface and boundary conditions in (C.10) to derive:

$$\begin{aligned} \langle \mathcal{A}_u \mathbf{u}, \mathbf{u} \rangle_{X_u} &= \sum_{i=1}^n \frac{1}{\prod_{j=1}^{i-1} p_j} \left[\int_{\bar{\Omega}_i} -d_i(u_i)_{xx} u_i \, dx + \int_{\bar{\Omega}_i} (u_i)^2 \, dx \right] \\ &= \sum_{i=1}^n \frac{1}{\prod_{j=1}^{i-1} p_j} \left[\int_{\bar{\Omega}_i} d_i(u_i)_x^2 \, dx + \int_{\bar{\Omega}_i} (u_i)^2 \, dx \right] \\ &\geq \mathcal{C}_1 \|\mathbf{u}\|_{\mathcal{H}_u^1}^2 \geq 0, \end{aligned} \quad (\text{C.11})$$

where the constant $\mathcal{C}_1 > 0$ depends on $d_i, 1/p_i$ ($i = 1, \dots, n$). Using an argument similar to that in the proof of (C.11), we can prove that $\langle \mathcal{A}_u \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{u}, \mathcal{A}_u \mathbf{v} \rangle$ for all $\mathbf{u}, \mathbf{v} \in \mathcal{D}(\mathcal{A}_u)$. This implies \mathcal{A}_u is symmetric.

Next, we prove that for every $\lambda \geq 0$, the range $R(\lambda I_{X_u} + \mathcal{A}_u) = X_u$. That is, for any $\mathbf{g} = (g_1, \dots, g_n) \in X_u$, the following problem admits a solution $\mathbf{u} \in \mathcal{D}(\mathcal{A}_u)$:

$$\begin{cases} -d_i(u_i)_{xx} + (\lambda + 1)u_i = g_i, & \text{in } \Omega_i, & \text{(C.12a)} \\ u_{i+1}(x_i^+) = p_i u_i(x_i^-), & i = 1, \dots, n-1, & \text{(C.12b)} \\ d_{i+1}(u_{i+1})_x(x_i^+) = d_i(u_i)_x(x_i^-), & i = 1, \dots, n-1, & \text{(C.12c)} \\ (u_1)_x(0^+) = (u_n)_x(L^-) = 0. & & \text{(C.12d)} \end{cases}$$

We first convert this problem into a weak formulation: find $\mathbf{u} \in \mathcal{H}_u^1$ such that

$$a(\mathbf{u}, \mathbf{s}) = \langle \mathbf{g}, \mathbf{s} \rangle_{X_u} \quad \text{for all } \mathbf{s} \in \mathcal{H}_u^1, \quad \text{(C.13)}$$

where the bilinear form a on $\mathcal{H}_u^1 \times \mathcal{H}_u^1$ is defined by

$$a(\mathbf{u}, \mathbf{s}) = \sum_{i=1}^n \frac{1}{\prod_{j=1}^{i-1} p_j} \left[\int_{\bar{\Omega}_i} d_i(u_i)_x(s_i)_x dx + (\lambda + 1) \int_{\bar{\Omega}_i} u_i s_i dx \right].$$

Clearly, $a(\mathbf{u}, \mathbf{s})$ is continuous. Following a similar approach to that for (C.11), we can show that $a(\mathbf{u}, \mathbf{s})$ is coercive. Then, by the Lax–Milgram theorem, problem (C.12) admits a unique solution $\mathbf{u} \in \mathcal{H}_u^1$ with

$$\|\mathbf{u}\|_{\mathcal{H}_u^1} \leq \mathcal{C}_2 \|\mathbf{g}\|_{X_u}, \quad \text{(C.14)}$$

where the constant $\mathcal{C}_2 > 0$ depends on $d_i, p_i, 1/d_i$ and $1/p_i$ ($i = 1, \dots, n$). Combining this with the definition of \mathcal{H}_u^1 in (C.2), we conclude that (C.12b) holds.

For each $i = 1, \dots, n$, we construct a test function

$$\mathbf{v} = (v_1, \dots, v_n) \in C^\infty(\bar{\Omega}_1) \times \dots \times C^\infty(\bar{\Omega}_n)$$

such that $v_k \equiv 0$ for all $1 \leq k \leq n$ with $k \neq i$ and v_i is supported in Ω_i . Clearly, $\mathbf{v} \in \mathcal{H}_u^1$. Thus, (C.13) implies that $u_i \in H^2(\Omega_i)$ for all $i = 1, \dots, n$ (so that $\mathbf{u} \in \mathcal{H}_u^2$) and that (C.12a) holds. Then, for each $i = 1, \dots, n-1$, we construct a test function $\mathbf{v} = (v_1, \dots, v_n) \in G \subset \mathcal{H}_u^1$ with the following properties: (i) $v_k \equiv 0$ for all k such that $1 \leq k \leq i-1$ or $i+2 \leq k \leq n$; (ii) the component v_i is supported in $(x_{i-1}, x_i]$; (iii) The component v_{i+1} is supported in $[x_i, x_{i+1})$. Here G is defined in (C.4). Since

$\mathbf{u} \in \mathcal{H}_u^1 \cap \mathcal{H}_u^2$, we see from (C.13) that

$$\begin{aligned}
0 &= - \int_{x_{i-1}}^{x_i} g_i v_i dx - \frac{1}{p_i} \int_{x_i}^{x_{i+1}} g_{i+1} v_{i+1} dx + \int_{x_{i-1}}^{x_i} [d_i(u_i)_x(v_i)_x + (\lambda + 1)u_i v_i] dx \\
&\quad + \frac{1}{p_i} \int_{x_i}^{x_{i+1}} [d_{i+1}(u_{i+1})_x(v_{i+1})_x + (\lambda + 1)u_{i+1} v_{i+1}] dx \\
&= d_i(u_i)_x(x_i^-) v_i(x_i^-) - \frac{1}{p_i} d_{i+1}(u_{i+1})_x(x_i^+) v_{i+1}(x_i^+) + \int_{x_{i-1}}^{x_i} [-d_i(u_i)_{xx} + (\lambda + 1)u_i] v_i dx \\
&\quad - \int_{x_{i-1}}^{x_i} g_i v_i dx + \frac{1}{p_i} \int_{x_i}^{x_{i+1}} [-d_{i+1}(u_{i+1})_{xx} + (\lambda + 1)u_{i+1} - g_{i+1}] v_{i+1} dx \\
&= d_i(u_i)_x(x_i^-) v_i(x_i^-) - \frac{1}{p_i} d_{i+1}(u_{i+1})_x(x_i^+) v_{i+1}(x_i^+),
\end{aligned}$$

where we have used integration by parts in the second step and (C.12a) in the third step. Combining this with the fact that $\mathbf{v} \in \mathcal{H}_u^1$, we obtain (C.12c). Using similar arguments, we can deduce that $(u_1)_x(0^+) = (u_n)_x(L^-) = 0$. Thus, $\mathbf{u} \in \mathcal{D}(\mathcal{A}_u)$, and consequently, the operator \mathcal{A}_u is maximal monotone. It is then densely defined and closed (by [6, Proposition 7.1]). Then, by arguments similar to those in the proof of [22, Lemma 3.3], we can deduce that $-\mathcal{A}_u$ is the infinitesimal generator of an analytic semigroup.

Furthermore, we see from (C.12a) that

$$\|(u_i)_{xx}\|_{L_2(\Omega_i)} \leq \frac{1}{d_i} [\|g_i\|_{L_2(\Omega_i)} + (\lambda + 1)\|u_i\|_{L_2(\Omega_i)}],$$

where $\lambda \geq 0$. This combined with (C.14) implies that $\|\mathbf{u}\|_{\mathcal{H}_u^2} \leq \mathcal{C}_3 \|\mathbf{g}\|_{X_u}$, where the constant $\mathcal{C}_3 > 0$ depends on $d_i, p_i, 1/d_i$ and $1/p_i$ ($i = 1, \dots, n$). Since $R(\lambda I + \mathcal{A}_u) = X_u$ and $\lambda I + \mathcal{A}_u$ is injective (by (C.11)), it is invertible with a bounded inverse $(\lambda I + \mathcal{A}_u)^{-1} : X_u \rightarrow \mathcal{D}(\mathcal{A}_u)$. Since $\mathcal{D}(\mathcal{A}_u) \xrightarrow{c} X_u$, we see that \mathcal{A}_u has a compact resolvent. This completes the proof. \square

Remark C.3. By the proof of Lemma C.2, \mathcal{A}_u (defined in (C.6)) has a compact resolvent. An analogous property holds for the operator \mathcal{A}_v , whose domain is given by

$$\begin{aligned}
\mathcal{D}(\mathcal{A}_v) &:= \{\mathbf{v} \in \mathcal{H}_v^2 \cap \mathcal{H}_v^1 : \hat{d}_{i+1}(v_{i+1})_x(x_i^+) = \hat{d}_i(v_i)_x(x_i^-), i = 1, \dots, n-1, \\
&\quad (v_1)_x(0^+) = (v_n)_x(L^-) = 0\},
\end{aligned} \tag{C.15}$$

Because $-\mathcal{A}_u$ and $-\mathcal{A}_v$ both generate analytic semigroups, we can define their corresponding fractional power spaces X_u^α and X_v^α for $0 < \alpha < 1$. Moreover, since both \mathcal{A}_u and \mathcal{A}_v have compact resolvents, the embedding

$$X_u^\beta \times X_v^\beta \xrightarrow{c} X_u^\alpha \times X_v^\alpha \tag{C.16}$$

is compact whenever $0 < \alpha < \beta < 1$.

For later applications, we establish the following embedding result for the function spaces defined above.

Lemma C.4. *Suppose that $\alpha \in (1/4, 1)$. Then the following statements hold:*

(i) *For $s > 1/2$,*

$$W^{s,2}(\overline{\Omega}_1) \times \cdots \times W^{s,2}(\overline{\Omega}_n) \xhookrightarrow{c} C^r(\overline{\Omega}_1) \times \cdots \times C^r(\overline{\Omega}_n), \quad (\text{C.17})$$

where $0 \leq r < s - 1/2$; and

$$(X_u, \mathcal{H}_u^2)_{s,2} = (X_v, \mathcal{H}_v^2)_{s,2} = W^{2s,2}(\overline{\Omega}_1) \times \cdots \times W^{2s,2}(\overline{\Omega}_n), \quad (\text{C.18})$$

where $X_u, X_v, \mathcal{H}_u^2$ and \mathcal{H}_v^2 are defined in (C.1) and (C.3).

(ii) *For $0 < \theta < 1$,*

$$(X_u, X_u^\alpha)_{\theta,2} = (X_u, \mathcal{D}(\mathcal{A}_u))_{\alpha\theta,2}, \quad (X_v, X_v^\alpha)_{\theta,2} = (X_v, \mathcal{D}(\mathcal{A}_v))_{\alpha\theta,2}, \quad (\text{C.19})$$

and

$$X_u^\alpha, X_v^\alpha \xhookrightarrow{c} C^\nu(\overline{\Omega}_1) \times \cdots \times C^\nu(\overline{\Omega}_n), \quad (\text{C.20})$$

where $0 < \nu < 2\alpha - \frac{1}{2}$, and $X_u, X_v, \mathcal{D}(\mathcal{A}_u)$ and $\mathcal{D}(\mathcal{A}_v)$ are defined in (C.1), (C.10) and (C.15).

(Here, $(S_1, S_2)_{\theta,q}$ denotes the interpolation space between function spaces S_1 and S_2 via the K -method; see [58, Section 1.3.2].)

Proof. The proof can be derived from [51, Theorem A.1], but we include it here for completeness. The results in (C.17) and (C.18) follow directly from [2, Theorem 11.5] and [2, Theorem 11.6], respectively, while (C.19) is established in [58, Section 1.15.2].

Now we prove (C.20). Fix an arbitrary ν satisfying $0 < \nu < 2\alpha - 1/2$. We can choose $\theta \in (0, 1)$ such that $\alpha\theta \neq 1/2$ and $2\alpha\theta - 1/2 > \nu$. Then

$$X_u^\alpha \subset (X_u, X_u^\alpha)_{\theta,2} = (X_u, \mathcal{D}(\mathcal{A}_u))_{\alpha\theta,2} \subset (X_u, \mathcal{H}_u^2)_{\alpha\theta,2} = W^{2\alpha\theta,2}(\overline{\Omega}_1) \times \cdots \times W^{2\alpha\theta,2}(\overline{\Omega}_n),$$

where the second and last equalities follow from (C.19) and (C.18), respectively. Since $2\alpha\theta - 1/2 > \nu$, we see from (C.17) that the embedding (C.20) holds. This completes the proof. \square

We now state the definition of a strong solution to (C.8), following [26, Definition 3.3.1].

Definition C.5. A strong solution of (C.8) on the interval $(0, T)$ with initial condition $\mathbf{w}(0) = \mathbf{w}_0 \in X_u^\alpha \times X_v^\alpha$ is a function $\mathbf{w}: [0, T) \rightarrow X$ satisfying the following:

- (i) $\mathbf{w}(t) \in \mathcal{D}(\mathcal{A})$ and $d\mathbf{w}(t)/dt$ exists for all $t \in (0, T)$;
- (ii) The abstract equation (C.8) is satisfied on $(0, T)$ with $\mathbf{w}(0) = \mathbf{w}_0$;
- (iii) $F(\mathbf{w}(t))$ is locally Hölder continuous in t , and $\int_0^\rho \|F(\mathbf{w}(t))\|_X dt < \infty$ for some $\rho > 0$.

Theorem C.6. For any given nonnegative initial value $(\mathbf{u}_0, \mathbf{v}_0) \in X_u^\alpha \times X_v^\alpha$ with $\alpha \in (3/4, 1)$, model (1.2) admits a unique classical solution $(\mathbf{u}(x, t), \mathbf{v}(x, t))$ (defined in Definition C.1) that exists for $t \in [0, \infty)$ and satisfies $(\mathbf{u}(x, 0), \mathbf{v}(x, 0)) = (\mathbf{u}_0, \mathbf{v}_0)$. Moreover, the following properties hold:

- (i) One has

$$\begin{aligned}\mathbf{u}(\cdot, t) &\in C([0, \infty), X_u^\alpha) \cap C^1((0, \infty), X_u^\alpha), \\ \mathbf{v}(\cdot, t) &\in C([0, \infty), X_v^\alpha) \cap C^1((0, \infty), X_v^\alpha),\end{aligned}$$

where $3/4 < \alpha < 1$.

- (ii) There exist two positive constants M_u and M_v , depending on $(\mathbf{u}_0, \mathbf{v}_0)$, \mathbf{k} , \mathbf{p} and $\hat{\mathbf{p}}$, such that for all $(x, t) \in [0, L] \times [0, \infty)$,

$$\mathbf{0} \leq \mathbf{u} \leq \mathbf{M}_u \text{ and } \mathbf{0} \leq \mathbf{v} \leq \mathbf{M}_v.$$

Here $\mathbf{M}_u, \mathbf{M}_v \in \mathbb{R}^n$ are defined by

$$\mathbf{M}_u := (M_u, p_1 M_u, \dots, \prod_{j=1}^{n-1} p_j M_u) \text{ and } \mathbf{M}_v := (M_v, \hat{p}_1 M_v, \dots, \prod_{j=1}^{n-1} \hat{p}_j M_v). \quad (\text{C.21})$$

Proof. The existence of a classical solution is inspired by the approach in [51, Theorems C.1 and C.2], which studied a chemotaxis model. The details are provided below. The uniqueness follows directly from Theorem B.1 and Remark B.2.

For given $3/4 < \alpha < 1$, the embedding (C.20) implies:

$$X_u^\alpha, X_v^\alpha \xhookrightarrow{c} C^1(\overline{\Omega}_1) \times \dots \times C^1(\overline{\Omega}_n). \quad (\text{C.22})$$

As a consequence, $F: X_u^\alpha \times X_v^\alpha \rightarrow X$, defined by (C.9), is locally Lipschitz continuous. By [26, Theorem 3.3.3 and Theorem 3.5.2], for any initial value $(\mathbf{u}_0, \mathbf{v}_0) \in X_u^\alpha \times X_v^\alpha$, there exists $T_{\max} = T_{\max}(\mathbf{u}_0, \mathbf{v}_0) > 0$ such that (C.8) admits a unique strong solution in the sense of Definition C.5, and

$$\begin{aligned}(\mathbf{u}(t), \mathbf{v}(t)) &\in C([0, T_{\max}), X_u^\alpha) \times C([0, T_{\max}), X_v^\alpha), \\ (\mathbf{u}(t), \mathbf{v}(t)) &\in C^1((0, T_{\max}), X_u^\alpha) \times C^1((0, T_{\max}), X_v^\alpha),\end{aligned} \quad (\text{C.23})$$

where $3/4 < \alpha < 1$.

Let

$$\mathbf{u}(x, t) = (u_1, \dots, u_n) = \mathbf{u}(t)(x) \quad \text{and} \quad \mathbf{v}(x, t) := (v_1, \dots, v_n) = \mathbf{v}(t)(x) \quad \text{for } x \in [0, L].$$

We now show that the strong solution is indeed a classical solution of the original model (1.2). It follows from (C.22) and (C.23) that

$$u_i(x, t), v_i(x, t) \in C^{1,0}(\overline{\Omega}_i \times [0, T_{\max})) \cap C^{1,1}(\overline{\Omega}_i \times (0, T_{\max})) \quad \text{for all } i = 1, \dots, n.$$

Thus, $f_i(u_i(x, t), v_i(x, t)), \hat{f}_i(u_i(x, t), v_i(x, t)) \in C^{1,1}(\overline{\Omega}_i \times (0, T_{\max}))$. Now, for fixed $t > 0$, we rewrite (1.2a) and (1.2b) as the following form:

$$\begin{aligned} d_i(u_i)_{xx}(x, t) &= u_i(x, t) + (u_i)_t(x, t) - f_i(u_i(x, t), v_i(x, t)) \in C^1(\overline{\Omega}_i), \\ \hat{d}_i(v_i)_{xx}(x, t) &= v_i(x, t) + (v_i)_t(x, t) - \hat{f}_i(u_i(x, t), v_i(x, t)) \in C^1(\overline{\Omega}_i). \end{aligned}$$

This implies that $(\mathbf{u}(\cdot, t), \mathbf{v}(\cdot, t)) \in \mathcal{H}^3 = \mathcal{H}_u^3 \times \mathcal{H}_v^3$ for fixed $t > 0$. By (C.17), we have

$$\mathcal{H}_u^3, \mathcal{H}_v^3 \xhookrightarrow{c} C^2(\overline{\Omega}_1) \times \dots \times C^2(\overline{\Omega}_n),$$

which implies that

$$u_i(\cdot, \cdot), v_i(\cdot, \cdot) \in C^{2,1}(\overline{\Omega}_i \times (0, T_{\max})). \quad (\text{C.24})$$

Consequently, $(\mathbf{u}(x, t), \mathbf{v}(x, t))$ satisfies (1.2a)-(1.2b) pointwise. Since $(\mathbf{u}(x, t), \mathbf{v}(x, t)) \in \mathcal{D}(\mathcal{A})$ and given the regularity in (C.24), it follows that $(\mathbf{u}(x, t), \mathbf{v}(x, t))$ satisfies (1.2c)-(1.2e) pointwise. Therefore, $(\mathbf{u}(x, t), \mathbf{v}(x, t))$ is a classical solution of the original model (1.2).

Denote $\mathbf{w}(x, t) := (\mathbf{u}(x, t), \mathbf{v}(x, t))$ for simplicity. Now we show that $\{\mathbf{w}(\cdot, t) \mid t > 0\}$ is bounded in $X_u^\alpha \times X_v^\alpha$. Choose two positive constants M_u, M_v such that $(\mathbf{M}_u, \mathbf{0}) \succeq (\mathbf{u}_0, \mathbf{v}_0) \succeq (\mathbf{0}, \mathbf{M}_v)$ and ensure that the pairs $(\mathbf{M}_u, \mathbf{0})$ and $(\mathbf{0}, \mathbf{M}_v)$ (defined in (C.21)) satisfy (B.2) and (B.3). Here \succeq is defined in (B.1). Then it follows from Theorem B.1 that

$$(\mathbf{M}_u, \mathbf{0}) \succeq (\mathbf{u}(x, t), \mathbf{v}(x, t)) \succeq (\mathbf{0}, \mathbf{M}_v) \quad \text{for } (x, t) \in [0, L] \times (0, T_{\max}).$$

This implies that there exists a positive constant \mathcal{M}_1 such that

$$\|F(\mathbf{w}(x, t))\|_X \leq \mathcal{M}_1 \quad \text{for all } t > 0. \quad (\text{C.25})$$

According to [26, Lemma 3.3.2], $\mathbf{w}(\cdot, t)$ satisfies the following integral equation

$$\mathbf{w}(\cdot, t) = e^{-t\mathcal{A}}\mathbf{w}_0 + \int_0^t e^{-(t-s)\mathcal{A}}F(\mathbf{w}(\cdot, s)) \, ds. \quad (\text{C.26})$$

By Lemma C.2 and its proof, we see that \mathcal{A} has a compact resolvent, and 0 belongs to the resolvent set of \mathcal{A} . It follows from [26, Chapter I, Section 1.4] that, for any given $0 \leq \beta \leq 1$, the operator $\mathcal{A}^\beta e^{-t\mathcal{A}}$ is bounded on X (i.e., $\mathcal{A}^\beta e^{-t\mathcal{A}} \in L(X)$), and

$$\mathcal{A}^\beta e^{-t\mathcal{A}} \widetilde{\mathbf{W}} = e^{-t\mathcal{A}} \mathcal{A}^\beta \widetilde{\mathbf{W}} \text{ for all } \widetilde{\mathbf{W}} \in \mathcal{D}(\mathcal{A}^\beta) = X_u^\beta \times X_v^\beta. \quad (\text{C.27})$$

Moreover, there exists a positive constant $\widetilde{\mathcal{M}}_\beta$ such that

$$\|\mathcal{A}^\beta e^{-t\mathcal{A}}\|_{L(X)} \leq \widetilde{\mathcal{M}}_\beta t^{-\beta} e^{-\delta t}, \quad (\text{C.28})$$

where $\|\cdot\|_{L(X)}$ is the operator norm in $L(X)$. Then, for all $t > 0$, we see from (C.26) that

$$\begin{aligned} \|\mathbf{w}(\cdot, t)\|_{X_u^\alpha \times X_v^\alpha} &\leq \|e^{-t\mathcal{A}}\| \cdot \|\mathbf{w}(\cdot, 0)\|_{X_u^\alpha \times X_v^\alpha} + \int_0^t \|\mathcal{A}^\alpha e^{-(t-s)\mathcal{A}}\|_{L(X)} \|F(\mathbf{w}(\cdot, s))\|_X \, ds \\ &\leq \widetilde{\mathcal{M}}_0 \|\mathbf{w}(\cdot, 0)\|_{X_u^\alpha \times X_v^\alpha} + \int_0^t \mathcal{M}_1 \widetilde{\mathcal{M}}_\alpha (t-s)^{-\alpha} e^{-\delta(t-s)} \, ds \\ &\leq \widetilde{\mathcal{M}}_0 \|\mathbf{w}(\cdot, 0)\|_{X_u^\alpha \times X_v^\alpha} + \int_0^\infty \mathcal{M}_1 \widetilde{\mathcal{M}}_\alpha (t-s)^{-\alpha} e^{-\delta(t-s)} \, ds < \infty. \end{aligned} \quad (\text{C.29})$$

The first inequality uses (C.27), while the second follows from (C.28) with $\beta = 0$ and $\beta = \alpha$, respectively. By (C.29), we see that $\{\mathbf{w}(\cdot, t) \mid t > 0\}$ is bounded in $X_u^\alpha \times X_v^\alpha$. By (C.22) again, we see that, for every closed bounded set \mathcal{B} in $X_u^\alpha \times X_v^\alpha$, the image $F(\mathcal{B})$ is bounded in X . Then it follows from [26, Theorem 3.3.4] that $T_{\max} = \infty$. This completes the proof. \square

Remark C.7. It follows from Lemma C.2 and the proof of Theorem C.5 that $-\mathcal{A}$ generates an analytic semigroup and that $F : X_u^\alpha \times X_v^\alpha \rightarrow X$, defined by (C.9), is locally Lipschitz continuous. Then, by [26, Theorem 5.1.1], a steady state of the abstract evolution equation (C.8) (or the original model (1.2)) is asymptotically stable (resp., unstable) if it is linearly stable (resp., linearly unstable).

Note that X_u^α and X_v^α are ordered Banach spaces. Their positive cones, P_u and P_v , both have nonempty interior. The cone $K = P_u \times (-P_v)$ induces the order \geq_k as follows:

$$(u_1, v_1) \geq_k (u_2, v_2) \text{ if and only if } (u_1 - u_2, v_1 - v_2) \in K. \quad (\text{C.30})$$

Define the continuous semiflow generated by model (C.8) as follows:

$$T_t : (\mathbf{u}_0, \mathbf{v}_0) \in X_u^\alpha \times X_v^\alpha \mapsto (\mathbf{u}(\cdot, t), \mathbf{v}(\cdot, t)) \in X_u^\alpha \times X_v^\alpha. \quad (\text{C.31})$$

Theorem C.8. *Let the semiflow T_t be defined as in (C.31). Then the following statements hold:*

- (i) T_t is strongly order-preserving with respect to \geq_k for all $t > 0$, where \geq_k is defined in (C.30);
- (ii) T_t is order compact in $X_u^\alpha \times X_v^\alpha$ for all $t > 0$.

Proof. Statement (i) is a direct consequence of Theorem B.1, Remark B.2 and Theorem C.6. Next, we prove statement (ii). Note that T_t is order compact if it maps every order interval in $P_u \times P_v \subset X_u^\alpha \times X_v^\alpha$ into a relatively compact set. Consider the following order interval with respect to \geq_k :

$$I := \{(\mathbf{u}_0, \mathbf{v}_0) \in P_u \times P_v : (\bar{\mathbf{u}}_0, \mathbf{0}) \geq_k (\mathbf{u}_0, \mathbf{v}_0) \geq_k (\mathbf{0}, \bar{\mathbf{v}}_0)\}.$$

Let $(\bar{\mathbf{u}}(\cdot, t), \mathbf{0})$, $(\mathbf{0}, \bar{\mathbf{v}}(\cdot, t))$ and $(\mathbf{u}(\cdot, t), \mathbf{v}(\cdot, t))$ be the solutions of (C.8) with the initial values $(\bar{\mathbf{u}}_0, \mathbf{0})$, $(\mathbf{0}, \bar{\mathbf{v}}_0)$, and $(\mathbf{u}_0, \mathbf{v}_0)$, respectively. By Theorem C.6, these are all classical solutions to the original model (1.2). Then Theorem B.1 implies that

$$(\bar{\mathbf{u}}(\cdot, t), \mathbf{0}) \geq_k (\mathbf{u}(\cdot, t), \mathbf{v}(\cdot, t)) \geq_k (\mathbf{0}, \bar{\mathbf{v}}(\cdot, t)) \text{ for all } t > 0.$$

This combined with Theorem C.6 (ii) implies that the set

$$T_t(I) = \{(\mathbf{u}(\cdot, t), \mathbf{v}(\cdot, t)) \mid (\mathbf{u}(\cdot, 0), \mathbf{v}(\cdot, 0)) \in I\}$$

is bounded in $(C(\bar{\Omega}_1) \times \cdots \times C(\bar{\Omega}_n))^2$. This implies that there exists a positive constant \mathcal{M}_1 such that, for any fixed $t > 0$,

$$\|F(\mathbf{w}(\cdot, t))\|_X \leq \mathcal{M}_1 \text{ whenever } \mathbf{w}(\cdot, 0) \in I. \quad (\text{C.32})$$

Here $\mathbf{w}(\cdot, t) := (\mathbf{u}(\cdot, t), \mathbf{v}(\cdot, t))$, and F is defined by (C.9).

Then using arguments similar to those in the proof of Theorem C.6, we see from (C.32) that, for any initial value $\mathbf{w}(\cdot, 0) \in I$ and any fixed $t > 0$,

$$\begin{aligned} \|\mathbf{w}(\cdot, t)\|_\beta &\leq \|\mathcal{A}^{\beta-\alpha} e^{-t\mathcal{A}}\| \cdot \|\mathbf{w}_0\|_\alpha + \int_0^t \|\mathcal{A}^\beta e^{-(t-s)\mathcal{A}}\|_{L(X)} \cdot \|F(\mathbf{w}(\cdot, s))\|_X \, ds \\ &\leq \widetilde{\mathcal{M}}_{\beta-\alpha} t^{-(\beta-\alpha)} e^{-\delta t} \|\mathbf{w}_0\|_\alpha + \widetilde{\mathcal{M}}_\beta \mathcal{M}_1 \int_0^t (t-s)^{-\beta} e^{-\delta(t-s)} \, ds < \infty. \end{aligned}$$

Here $\alpha < \beta < 1$, and \widetilde{M}_β , $\widetilde{M}_{\beta-\alpha}$, δ are defined in the proof of Theorem C.6. Consequently, $T_t(I)$ is bounded in $X_u^\beta \times X_v^\beta$ for any fixed $t > 0$. This combined with the compact embedding (C.16) implies that the $T_t(I)$ is relatively compact in $X_u^\alpha \times X_v^\alpha$. This complete the proof. \square

Finally, we consider the following eigenvalue problem

$$\begin{cases} -d_i(\phi_i)_{xx} - c_i(x)\phi_i = \lambda\phi_i(x), & \text{in } \Omega_i, \quad i = 1, 2, \dots, n, \\ \phi_{i+1}(x_i^+) = p_i\phi_i(x_i^-), \quad d_{i+1}(\phi_{i+1})_x(x_i^+) = d_i(\phi_i)_x(x_i^-), \\ (\phi_1)_x(0^+) = (\phi_n)_x(L^-) = 0. \end{cases} \quad \begin{matrix} \text{(C.33a)} \\ \text{(C.33b)} \\ \text{(C.33c)} \end{matrix}$$

Proposition C.9. *Assume that $c_i(x) \in C^1(\overline{\Omega}_i)$ for all $i = 1, \dots, n$, and let $\mathbf{d} = (d_1, \dots, d_n)$. Then the eigenvalue problem (C.33) admits a simple eigenvalue $\lambda_1(\mathbf{d})$ with a corresponding positive eigenfunction Φ^* , and any other eigenvalue $\lambda \neq \lambda_1(\mathbf{d})$ of (C.33) satisfies $\text{Re}(\lambda) > \lambda_1$. Moreover, $\lambda_1(\mathbf{d})$ is continuous with respect to \mathbf{d} .*

Proof. Define

$$\Lambda := \max_{1 \leq i \leq n} \left\{ \|c_i(x)\|_{C(\overline{\Omega}_i)} + 1 \right\}.$$

For $\mathbf{g} \in G$, where G is defined in (C.4), we consider the following problem

$$\begin{cases} -d_i(\phi_i)_{xx} + [\Lambda - c_i]\phi_i = g_i, & \text{in } \Omega_i, \quad i = 1, 2, \dots, n, \\ \phi_{i+1}(x_i^+) = p_i\phi_i(x_i^-), \quad d_{i+1}(\phi_{i+1})_x(x_i^+) = d_i(\phi_i)_x(x_i^-), \\ (\phi_1)_x(0^+) = (\phi_n)_x(L^-) = 0. \end{cases} \quad \begin{matrix} \text{(C.34a)} \\ \text{(C.34b)} \\ \text{(C.34c)} \end{matrix}$$

Define

$$\tilde{\mathcal{A}}_u \mathbf{u} := (-d_1(u_1)_{xx} + (\Lambda - c_1(x))u_1, \dots, -d_n(u_n)_{xx} + (\Lambda - c_n(x))u_n), \quad \text{(C.35)}$$

and the domain of $\tilde{\mathcal{A}}_u$ is given by:

$$\begin{aligned} \mathcal{D}(\tilde{\mathcal{A}}_u) := \{ \mathbf{u} \in \mathcal{H}_u^2 \cap \mathcal{H}_u^1 : d_{i+1}(u_{i+1})_x(x_i^+) = d_i(u_i)_x(x_i^-), \quad i = 1, \dots, n-1, \\ (u_1)_x(0^+) = (u_n)_x(L^-) = 0 \}, \end{aligned} \quad \text{(C.36)}$$

Then using arguments similar to those in the proof of Lemma C.2, we conclude that $\tilde{\mathcal{A}}_u$ is symmetric and that (C.34) admits a unique solution $\Phi \in \mathcal{D}(\tilde{\mathcal{A}}_u)$. Moreover, Φ satisfies

$$\|\Phi\|_{\mathcal{H}_u^2} \leq \mathcal{C}_1 \|\mathbf{g}\|_{X_u}. \quad \text{(C.37)}$$

Here the constant $\mathcal{C}_1 > 0$ depends on d_i , p_i , $1/d_i$ and $1/p_i$ ($i = 1, \dots, n$). Since $\mathcal{H}_u^2 \xhookrightarrow{c} C(\overline{\Omega}_1) \times \dots \times C(\overline{\Omega}_n)$ and $G \xhookrightarrow{c} X_u$, it follows from (C.37) that

$$\max_{1 \leq i \leq n} \|\phi_i\|_{C^1(\overline{\Omega}_i)} \leq \mathcal{C}_2 \|\mathbf{g}\|_{X_u} \leq \mathcal{C}_3 \|\mathbf{g}\|_G,$$

where the constants $\mathcal{C}_2, \mathcal{C}_3 > 0$ depends on d_i , p_i , $1/d_i$ and $1/p_i$ ($i = 1, \dots, n$). This combined with (C.34a) implies that $\Phi \in \mathcal{H}_u^3$ and

$$\|\Phi\|_{\mathcal{H}_u^3} \leq \mathcal{C}_4 \|\mathbf{g}\|_{X_u}, \quad \text{(C.38)}$$

where the constant $\mathcal{C}_4 > 0$ depends on d_i , p_i , $1/d_i$ and $1/p_i$ ($i = 1, \dots, n$). The embedding $\mathcal{H}_u^3 \xhookrightarrow{c} C^2(\overline{\Omega}_1) \times \dots \times C^2(\overline{\Omega}_n)$ combined with (C.38) implies that

$$\max_{1 \leq i \leq n} \|\phi_i\|_{C^2(\overline{\Omega}_i)} \leq \mathcal{C}_5 \|\mathbf{g}\|_G, \quad (\text{C.39})$$

where the constant \mathcal{C}_5 depends on d_i , p_i , $1/d_i$ and $1/p_i$ ($i = 1, \dots, n$).

For fixed $\mathbf{d} \gg \mathbf{0}$, define the linear operator $T(\mathbf{d}) : G \rightarrow G$ by $T(\mathbf{d})\mathbf{g} = \Phi$, where Φ is the solution of (C.34) corresponding to \mathbf{g} . Since $\Phi \in C^2(\overline{\Omega}_1) \times \dots \times C^2(\overline{\Omega}_n)$ and $C^2(\overline{\Omega}_1) \times \dots \times C^2(\overline{\Omega}_n) \xhookrightarrow{c} G$, we can deduce that $T(\mathbf{d})$ is compact. Let

$$\mathcal{K} := \{\mathbf{w} \in G : w_i(x) \geq 0 \text{ in } \Omega_i\}$$

be the positive cone in G . This cone has nonempty interior $\mathring{\mathcal{K}}$ and satisfies $\mathcal{K} \cap (-\mathcal{K}) = \{\mathbf{0}\}$. By Theorem B.5, we obtain that, if $\mathbf{g} \in \mathcal{K} \setminus \{\mathbf{0}\}$, then $\Phi \in \mathring{\mathcal{K}}$, which implies that $T(\mathbf{d})(\mathcal{K} \setminus \{\mathbf{0}\}) \subset \mathring{\mathcal{K}}$. Then it follows from the Krein-Rutman theorem (see e.g., [16, Theorem 1.2]) that the spectral radius $r(\mathbf{d})$ is a simple eigenvalue of $T(\mathbf{d})$ with a corresponding positive eigenfunction \mathbf{g}^* . This implies that

$$-d_i(\phi_i^*)_{xx} + [\Lambda - c_i]\phi_i^* = \frac{\phi_i^*}{r(\mathbf{d})} \quad \text{in } \Omega_i, \quad i = 1, 2, \dots, n,$$

where $\Phi^* = (\phi_1^*, \dots, \phi_n^*) = T\mathbf{g}^*$ is positive. Consequently, $\lambda_1(\mathbf{d}) = 1/r(\mathbf{d}) - \Lambda$ is a simple eigenvalue of (C.33) with a positive eigenfunction Φ^* . Then, using arguments similar to those in the proof of [16, Theorem 1.4], we deduce that any eigenvalue $\lambda \neq \lambda_1$ of (C.33) satisfies $\text{Re}(\lambda) > \lambda_1$.

Finally, we will show that $\lambda_1(\mathbf{d})$ is continuous with respect to \mathbf{d} . By (C.37), the operator $T(\mathbf{d})$ is also bounded and compact on X_u . We first show that $T(\mathbf{d})$ is continuous in $L(X_u)$, where $L(X_u)$ denotes the set of bounded linear operators on X_u . That is, for any sequence $\{\mathbf{d}^k\}_{k=1}^\infty$ satisfying $\lim_{k \rightarrow \infty} \mathbf{d}^k = \mathbf{d}$, we have

$$\lim_{k \rightarrow \infty} \|T(\mathbf{d}^k) - T(\mathbf{d})\|_{L(X_u)} = 0, \quad (\text{C.40})$$

where $\|\cdot\|_{L(X_u)}$ is the operator norm in $L(X_u)$. Since the constant \mathcal{C}_4 in (C.37) depends only on d_i and $1/d_i$ ($i = 1, \dots, n$), we see from (C.37) that

$$\bigcup_{k=1}^\infty \{T(\mathbf{d}^k)\mathbf{g} : \|\mathbf{g}\|_{X_u} \leq 1\}$$

is compact in X_u , and $\{\|T(\mathbf{d}^k)\|_{L(X_u)}\}_{k=1}^\infty$ is bounded. Consequently, $\{T(\mathbf{d}^k)\}_{k=1}^\infty$ is collectively compact. Since $\tilde{\mathcal{A}}_u$ is symmetric, it follows that $T(\mathbf{d}^k)$ is self-adjoint. Then it follows from [3, Lemma 5.2 and Corollary 5.7] that (C.40) holds if and only if

$$\lim_{k \rightarrow \infty} T(\mathbf{d}^k)\mathbf{g} = T(\mathbf{d})\mathbf{g} \quad \text{in } X_u \quad \text{for any fixed } \mathbf{g} \in X_u. \quad (\text{C.41})$$

Note that $\{\|T(\mathbf{d}^k)\|_{L(X_u)}\}_{k=1}^\infty$ is bounded, and G is dense in X_u . Therefore, to prove (C.41), it suffices to show that

$$\lim_{k \rightarrow \infty} T(\mathbf{d}^k)\mathbf{g} = T(\mathbf{d})\mathbf{g} \text{ in } G \text{ for any fixed } \mathbf{g} \in G. \quad (\text{C.42})$$

Now we construct an auxiliary function for further application. Let $\mathbf{g} = (g_1, \dots, g_n) \in G$. For any given $a > 0$, we inductively solve the following ODEs. For $i = 1$:

$$\begin{cases} \frac{d\phi_1}{dx} = \psi_1, & x \in (0, x_1), \\ \frac{d\psi_1}{dx} = \frac{1}{d_1} [(\Lambda - c_1)\phi_1 - g_1], & x \in (0, x_1), \\ \phi_1(0^+) = a, \quad \psi_1(0^+) = 0. \end{cases} \quad (\text{C.43})$$

For $i = 2, \dots, n$:

$$\begin{cases} \frac{d\phi_i}{dx} = \psi_i, & x \in (x_{i-1}, x_i), \\ \frac{d\psi_i}{dx} = \frac{1}{d_i} [(\Lambda - c_i)\phi_i - g_i], & x \in (x_{i-1}, x_i), \\ \phi_i(x_{i-1}^+) = p_{i-1}\phi_{i-1}(x_{i-1}^-), \quad \psi_i(x_{i-1}^+) = \frac{d_{i-1}}{d_i}\psi_{i-1}(x_{i-1}^-). \end{cases} \quad (\text{C.44})$$

Define the auxiliary function $\mathcal{T} : (0, \infty) \times \mathbb{R}^n \rightarrow \mathbb{R}$ by $\mathcal{T}(a, \mathbf{d}) := \psi_n(L)$.

By the continuous dependence of solutions to ODEs on initial values and parameters, we see that $\mathcal{T}(a, \mathbf{d})$ is continuous with respect to (a, \mathbf{d}) . Since (C.34) has a unique solution $\Phi = (\phi_1, \dots, \phi_n) \in G$, it follows that $\mathcal{T}(a, \mathbf{d}) = 0$ admits a unique solution $a(\mathbf{d})$ for fixed \mathbf{d} . We now prove that

$$\lim_{k \rightarrow \infty} a(\mathbf{d}^k) = a(\mathbf{d}). \quad (\text{C.45})$$

Since the constant \mathcal{C}_5 in (C.39) depends only on d_i and $1/d_i (i = 1, \dots, n)$, it follows from (C.39) that $\{a(\mathbf{d}^k)\}_{k=1}^\infty$ is bounded. Consider any convergent subsequence $\{a(\mathbf{d}^{k_j})\}_{j=1}^\infty$ with limit a^* . Since $\mathcal{T}(a(\mathbf{d}^{k_j}), \mathbf{d}^{k_j}) = 0$ for all j and $\mathcal{T}(a, \mathbf{d})$ is continuous with respect to (a, \mathbf{d}) , taking the limit as $j \rightarrow \infty$ yields $\mathcal{T}(a^*, \mathbf{d}) = 0$. The uniqueness of the solution to $\mathcal{T}(a, \mathbf{d}) = 0$ then implies that $a^* = a(\mathbf{d})$. Thus, (C.45) holds. Note that

$$\left(T(\mathbf{d})\mathbf{g}, \frac{dT(\mathbf{d})\mathbf{g}}{dx} \right)$$

is the solution of (C.43)-(C.44) for $a = a(\mathbf{d})$. Similarly,

$$\left(T(\mathbf{d}^k)\mathbf{g}, \frac{dT(\mathbf{d}^k)\mathbf{g}}{dx} \right)$$

is the solution of (C.43)-(C.44) for $\mathbf{d} = \mathbf{d}^k$ and $a = a(\mathbf{d}^k)$. By the continuous dependence of solutions to ODEs on initial values and parameters, we see from (C.45) that (C.42) holds. Consequently, $T(\mathbf{d})$ is continuous in $L(X_u)$ with respect to \mathbf{d} .

Note from [30, Chapter III, Section 3.5] that a simple and isolated eigenvalue depends continuously on the operator. Since the spectral radius $r(\mathbf{d})$ is a simple and isolated eigenvalue of $T(\mathbf{d})$ on G (and hence X_u), it follows that $r(\mathbf{d})$ is continuous with respect to \mathbf{d} , and consequently $\lambda_1(\mathbf{d})$ is continuous with respect to \mathbf{d} . This completes the proof. \square

Acknowledgments

This work was supported by National Natural Science Foundation of China (No. 12171117), Taishan Scholars Program of Shandong Province (No. tsqn 202306137) and Shandong Provincial Natural Science Foundation of China (No. ZR2020YQ01).

References

- [1] Y. Alqawasmeh and F. Lutscher. Persistence and spread of stage-structured populations in heterogeneous landscapes. *J. Math. Biol.*, 78(5):1485–1527, 2019.
- [2] H. Amann. Existence and regularity for semilinear parabolic evolution equations. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 11(4):593–676, 1984.
- [3] P. M. Anselone. *Collectively Compact Operator Approximation Theory and Applications to Integral Equations*. Prentice-Hall Series in Automatic Computation. Prentice-Hall, Inc., Englewood Cliffs, NJ, 1971.
- [4] J. Apaloo. Revisiting strategic models of evolution: the concept of neighborhood invader strategies. *Theor. Popul. Biol.*, 52(1):71–77, 1997.
- [5] I. Averill, Y. Lou, and D. Munther. On several conjectures from evolution of dispersal. *J. Biol. Dyn.*, 6(2):117–130, 2012.
- [6] H. Brezis. *Functional Analysis, Sobolev Spaces and Partial Differential Equations*. Springer, New York, 2011.
- [7] R. S Cantrell, C. Cosner, and Y. Lou. Evolution of dispersal and the ideal free distribution. *Math. Biosci. Eng.*, 7(1):17–36, 2010.

- [8] R. S. Cantrell, C. Cosner, and Y. Lou. Evolutionary stability of ideal free dispersal strategies in patchy environments. *J. Math. Biol.*, 65(5):943–965, 2012.
- [9] S. Chen, J. Shi, Z. Shuai, and Y. Wu. Global dynamics of a Lotka-Volterra competition patch model. *Nonlinearity*, 35(2):817–842, 2022.
- [10] S. Chen, J. Shi, Z. Shuai, and Y. Wu. Two novel proofs of spectral monotonicity of perturbed essentially nonnegative matrices with applications in population dynamics. *SIAM J. Appl. Math.*, 82(2):654–676, 2022.
- [11] C.-Y. Cheng, K.-H. Lin, and C.-W. Shih. Coexistence and extinction for two competing species in patchy environments. *Math. Biosci. Eng.*, 16(2):909–946, 2019.
- [12] G. C. Cruywagen, P. Kareiva, M. A. Lewis, and J. D. Murray. Competition in a spatially heterogeneous environment: Modelling the risk of spread of a genetically engineered population. *Theor. Popul. Biol.*, 49(1):1–38, 1996.
- [13] U. Dieckmann and R. Law. The dynamical theory of coevolution: a derivation from stochastic ecological processes. *J. Math. Biol.*, 34(5-6):579–612, 1996.
- [14] O. Diekmann. A beginner’s guide to adaptive dynamics. *Math. Model. Popul. Dyn. Banach Cent. Publ.*, 63:47–86, 2003.
- [15] J. Dockery, V. Hutson, K. Mischaikow, and M. Pernarowski. The evolution of slow dispersal rates: a reaction diffusion model. *J. Math. Biol.*, 37(1):61–83, 1998.
- [16] Y. Du. *Order Structure and Topological Methods in Nonlinear Partial Differential Equations. Vol. 1: Maximum Principles and Applications*, volume 2 of *Series in Partial Differential Equations and Applications*. World Scientific, Hackensack, NJ, 2006.
- [17] H. I. Freedman, J. B. Shukla, and Y. Takeuchi. Population diffusion in a two-patch environment. *Math. Biosci.*, 95(1):111–123, 1989.
- [18] Q. Ge and D. Tang. Global dynamics of two-species Lotka-Volterra competition-diffusion-advection system with general carrying capacities and intrinsic growth rates. *J. Dynam. Differential Equations*, 36(2):1905–1926, 2024.

- [19] S. A. H. Geritz, E. Kisdi, G. Meszner, and J. A. J. Metz. Evolutionarily singular strategies and the adaptive growth and branching of the evolutionary tree. *Evol. Ecol.*, 12(1):35–57, 1998.
- [20] S. A. Gourley and Y. Kuang. Two-species competition with high dispersal: The winning strategy. *Math. Biosci. Eng.*, 2(2):345–362, 2005.
- [21] F. Hamel, F. Lutscher, and M. Zhang. Propagation and blocking in a two-patch reaction-diffusion model. *J. Math. Pures Appl.*, 168:213–267, 2022.
- [22] F. Hamel, F. Lutscher, and M. Zhang. Propagation phenomena in periodic patchy landscapes with interface conditions. *J. Dynam. Differential Equations*, 36:S435–S486, 2024.
- [23] F. Hamel and M. Zhang. Kpp transition fronts in a one-dimensional two-patch habitat. *To appear in Math. Med. Biol.*
- [24] A. Hastings. Can spatial variation alone lead to selection for dispersal? *Theor. Popul. Biol.*, 24(3):244–251, 1983.
- [25] X. He and W.-M. Ni. Global dynamics of the Lotka-Volterra competition-diffusion system: diffusion and spatial heterogeneity I. *Comm. Pure Appl. Math.*, 69(5):981–1014, 2016.
- [26] D. Henry. *Geometric Theory of Semilinear Parabolic Equations*, volume 840 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, 1981.
- [27] P. Hess. *Periodic-Parabolic Boundary Value Problems and Positivity*, volume 247 of *Pitman Research Notes in Mathematics Series*. Longman Scientific & Technical, Harlow; copublished in the United States with John Wiley & Sons, Inc., New York, 1991.
- [28] S.-B. Hsu, H.-L. Smith, and P. Waltman. Competitive exclusion and coexistence for competitive systems on ordered Banach spaces. *Trans. Amer. Math. Soc.*, 348(10):4083–4094, 1996.
- [29] Y. Jin, R. Peng, and J. Shi. Population dynamics in river networks. *J. Nonlinear Sci.*, 29(6):2501–2545, 2019.

- [30] T. Kato. *Perturbation Theory for Linear Operators*, volume Band 132 of *Die Grundlehren der mathematischen Wissenschaften*. Springer-Verlag New York, Inc., New York, 1966.
- [31] K.-Y. Lam and Y. Lou. Evolutionarily stable and convergent stable strategies in reaction-diffusion models for conditional dispersal. *Bull. Math. Biol.*, 76(2):261–291, 2014.
- [32] K.-Y. Lam and Y. Lou. *Introduction to Reaction-Diffusion Equations: Theory and Applications to Spatial Ecology and Evolutionary Biology*. Springer, Cham, Switzerland, 2022.
- [33] K.-Y. Lam and D. Munther. A remark on the global dynamics of competitive systems on ordered Banach spaces. *Proc. Amer. Math. Soc.*, 144(3):1153–1159, 2016.
- [34] K.-Y. Lam and W.-M. Ni. Uniqueness and complete dynamics in heterogeneous competition-diffusion systems. *SIAM J. Appl. Math.*, 72(6):1695–1712, 2012.
- [35] K.-H. Lin, Y. Lou, C.-W. Shih, and T.-H. Tsai. Global dynamics for two-species competition in patchy environment. *Math. Biosci. Eng.*, 11(4):947–970, 2014.
- [36] K. Liu, D. Tang, and S. Chen. Evolution of dispersal in a stream with better resources at downstream locations. *Stud. Appl. Math.*, 154(2):Paper No. e70017, 2025.
- [37] Y. Lou. On the effects of migration and spatial heterogeneity on single and multiple species. *J. Differential Equations*, 223(2):400–426, 2006.
- [38] Y. Lou. Ideal free distribution in two patches. *J. Nonlinear Model. Anal.*, 2(10):151–167, 2021.
- [39] F. Lutscher, M. A. Lewis, and E. McCauley. Effects of heterogeneity on spread and persistence in rivers. *Bull. Math. Biol.*, 68(8):2129–2160, 2006.
- [40] G. A. Maciel, C. Cosner, R. S. Cantrell, and F. Lutscher. Evolutionarily stable movement strategies in reaction-diffusion models with edge behavior. *J. Math. Biol.*, 80(1-2):61–92, 2020.

- [41] G. A. Maciel and F. Lutscher. How individual movement response to habitat edges affects population persistence and spatial spread. *Am. Nat.*, 182(1):42–52, 2013.
- [42] G. A. Maciel and F. Lutscher. Allee effects and population spread in patchy landscapes. *J. Biol. Dyn.*, 9(1):109–123, 2015.
- [43] G. A. Maciel and F. Lutscher. Movement behaviour determines competitive outcome and spread rates in strongly heterogeneous landscapes. *Theor. Ecol.*, 11(3):351–365, 2018.
- [44] O. Ovaskainen and S. J. Cornell. Biased movement at a boundary and conditional occupancy times for diffusion processes. *J. Appl. Probab.*, 40(3):557–580, 2003.
- [45] S. W. Pacala and J. Roughgarden. Spatial heterogeneity and interspecific competition. *Theor. Popul. Biol.*, 21(1):92–113, 1982.
- [46] J. D. Reeve and J. T. Cronin. Edge behaviour in a minute parasitic wasp. *J. Anim. Ecol.*, 79(2):483–490, 2010.
- [47] J. D. Reeve, J. T. Cronin, and K. J. Haynes. Diffusion models for animals in complex landscapes: incorporating heterogeneity among substrates, individuals and edge behaviours. *J. Anim. Ecol.*, 77(5):898–904, 2008.
- [48] L. Ries and D. M. Debinski. Butterfly responses to habitat edges in the highly fragmented prairies of central iowa. *J. Anim. Ecol.*, 70(5):840–852, 2001.
- [49] N. Schtickzelle and M. Baguette. Behavioural responses to habitat patch boundaries restrict dispersal and generate emigration-patch area relationships in fragmented landscapes. *J. Anim. Ecol.*, 72(4):533–545, 2003.
- [50] C. B. Schultz and E. E. Crone. Edge-mediated dispersal behavior in a prairie butterfly. *Ecology*, 82(7):1879–1892, 2001.
- [51] H. Shemtaga, W. Shen, and S. Sukhtaiev. Well-posedness of Keller-Segel systems on compact metric graphs. *J. Evol. Equ.*, 25(1):Paper No. 7, 2025.
- [52] N. Shigesada, K. Kawasaki, and E. Teramoto. Traveling periodic waves in heterogeneous environments. *Theor. Popul. Biol.*, 30(1):143–160, 1986.

- [53] N. Shigesada, K. Kawasaki, and H. F. Weinberger. Spreading speeds of invasive species in a periodic patchy environment: effects of dispersal based on local information and gradient-based taxis. *Jpn. J. Ind. Appl. Math.*, 32(3):675–705, 2015.
- [54] A. Slavík. Lotka-Volterra competition model on graphs. *SIAM J. Appl. Dyn. Syst.*, 19(2):725–762, 2020.
- [55] H.-L. Smith. *Monotone Dynamical Systems: an Introduction to the Theory of Competitive and Cooperative Systems*. American Mathematical Society, Providence, RI, 1995.
- [56] J.-M. Smith and G.-R. Price. The logic of animal conflict. *Nature*, 246(5427):15–18, 1973.
- [57] L. K. Tchouaga and F. Lutscher. Steady-state dynamics in a two-patch population model with and without allee effect. *Math. Appl. Sci. Eng.*, 4(3):196–226, 2023.
- [58] H. Triebel. *Interpolation theory, function spaces, differential operators*. Johann Ambrosius Barth, Heidelberg, second edition, 1995.
- [59] N. Zaker, L. Ketchemen, and F. Lutscher. The effect of movement behavior on population density in patchy landscapes. *Bull. Math. Biol.*, 82(1):Paper No. 1, 24, 2020.
- [60] P. Zhou. On a Lotka-Volterra competition system: diffusion vs advection. *Calc. Var. Partial Differential Equations*, 55(6):Paper No. 137, 2016.