

MODULUS OF ELEMENTARY DOMAINS IN THE HYPERBOLIC PLANE

IOANNIS D. PLATIS

ABSTRACT. We calculate the modulus of curve families inside a hyperbolic quadrilateral and a hyperbolic annulus.

1. INTRODUCTION

The study of curve families and their intrinsic geometric properties is a cornerstone of classical complex analysis and geometric function theory. In the Euclidean plane, the theory of conformal invariants, pioneered by Grötzsch, Ahlfors, and Beurling, provides a powerful framework for understanding the "size" or "capacity" of a family of curves connecting two sets, see e.g. [1], [7]. Central role in this framework plays the modulus of a curve family, a conformal invariant that quantifies the richness of the family and has profound applications in the theories of quasiconformal mappings, potential theory, the geometry of Riemann surfaces, as well as in sub-Riemannian geometry, see for instance [3], [2], as well as [6].

While the Euclidean theory about moduli of curve families is well-established, a comprehensive analysis within the geometry of the hyperbolic plane reveals a more nuanced landscape. The hyperbolic plane, with its constant negative curvature, offers a canonical domain for complex analysis and serves as the universal cover for all Riemann surfaces of genus greater than one. Consequently, understanding the behaviour of curve families in the hyperbolic plane is not merely a generalisation of the Euclidean case but rather a step towards solving problems in Teichmüller theory and low-dimensional geometry. The distinctive feature of hyperbolic plane is its exponential growth of area with radius, a property that fundamentally alters the analytic and geometric constraints on curve families.

In this short note we define the modulus of a curve family in the hyperbolic plane $\mathbf{H}_{\mathbb{C}}^1$ in Section 2.3 and consider two particular domains, that is, a normal hyperbolic quadrilateral (see Definition 3.1) and a hyperbolic annulus (see Definition 4.1). In the first case we calculate the modulus of the family of curves connecting the circular arcs of boundary of the normal quadrilateral (Theorem 3.2) and the modulus of the family of curves connecting the straight line segments of the boundary of the normal quadrilateral (Theorem 3.3). In the hyperbolic annulus case we calculate the modulus of the family of curves which join the two boundary components (hyperbolic circles) in Theorem 4.2 as well as the modulus of the family of curves which separate the boundary components in Theorem 4.3. All the above are the hyperbolic counterparts of their Euclidean analogues, see e.g. [1], or [7].

The author would like to thank Zoltán M. Balogh for stimulating discussions.

2. PRELIMINARIES

2.1. Hyperbolic plane. We consider the hyperbolic plane $\mathbf{H}_{\mathbb{C}}^1$ with coordinates $z = \lambda + it$, $\lambda > 0$, $t \in \mathbb{R}$. $\mathbf{H}_{\mathbb{C}}^1$ is a Riemannian manifold: the metric tensor is given by

$$g_h = \frac{d\lambda^2 + dt^2}{\lambda^2};$$

Key words and phrases. Modulus of curve families, hyperbolic plane.
2010 Mathematics Subject Classification: 30C85, 31A15.

The author was funded by the MEDICUS programme, No. 26375.

the hyperbolic distance $d_h(z_1, z_2)$ between two points $z_i = \lambda_i + it_i \in \mathbf{H}_{\mathbb{C}}^1$ is given by

$$\cosh(d_h(z_1, z_2)) = 1 + \frac{|z_1 - z_2|^2}{2\lambda_1\lambda_2}.$$

The geodesics of $(\mathbf{H}_{\mathbb{C}}^1)$ are either straight lines orthogonal to the t -axis or semicircles centred on the t -axis. Hyperbolic circles $C_h(z_0, R)$ with hyperbolic centre $z_0 = \lambda_0 + it_0$ and hyperbolic radius $R > 0$ are Euclidean circles $C_e((\lambda_0 \cosh R, t_0), \lambda_0 \sinh R)$. The line and the area elements are, respectively,

$$ds_h = \frac{\sqrt{d\lambda^2 + dt^2}}{\lambda}, \quad d\mathcal{A}_h = \frac{d\lambda dt}{\lambda^2}.$$

The group of orientation preserving isometries of $(\mathbf{H}_{\mathbb{C}}^1, d_h)$ comprises Möbius transformations of the form

$$f(z) = \frac{az + ib}{icz + d}, \quad a, b, c, d \in \mathbb{R}, \quad ad + bc = 1,$$

that is, it is isomorphic to

$$\mathrm{SU}(1, 1) = \left\{ \begin{pmatrix} a & ib \\ ic & d \end{pmatrix}, \quad a, b, c, d \in \mathbb{R}, \quad ad + bc = 1 \right\}.$$

2.2. Hyperbolic polar coordinates. In this Section 2.2 we give a detailed description of hyperbolic polar coordinates, see also [5]. We mainly use those coordinates in Section 4. To introduce polar coordinates (based on the point $(1, 0)$) we mimic the following manner of defining polar coordinates in the Euclidean plane \mathbb{R}^2 : all Euclidean geodesics passing from $(0, 0)$ are straight lines. We fix the ray $y = 0$ and given an $r > 0$ we define $x = r, y = 0$. Now any (orientation preserving) Euclidean isometry that leaves $(0, 0)$ fixed is an element of $\mathrm{SO}(2, 1)$. Thus if

$$R_\phi = \begin{pmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{pmatrix}$$

is such an element, we let it act on $(r, 0)$ on the right and we obtain

$$(x, y) = (r, 0), \quad R_\phi = (r \cos \phi, r \sin \phi).$$

In the hyperbolic case, considered as the first affine group, all geodesics passing through the neutral element $(1, 0)$ are either: a) the straight line $t = 0$ or b) semicircles centred on the t -axis. As before, we fix the ray $t = 0$ and given an $r > 0$ we define $\lambda = e^r, t = 0$. Now any orientation preserving hyperbolic isometry which fixes 1 is an element of $\mathrm{SU}(1, 1)$ of the form

$$f(z) = \frac{az - ib}{-ibz + a}, \quad a, b \in \mathbb{R}, \quad a^2 + b^2 = 1.$$

We may as well write equivalently,

$$f_\theta(z) = \frac{\cos(\theta/2)z - i \sin(\theta/2)}{-i \sin(\theta/2)z + \cos(\theta/2)}, \quad \theta \in \mathbb{R}.$$

We now take

$$\begin{aligned} z = f(e^r) &= \frac{e^r \cos(\theta/2) - i \sin(\theta/2)}{-ie^r \sin(\theta/2) + \cos(\theta/2)} \\ &= \frac{2e^r}{(1 + e^{2r}) + (1 - e^{2r}) \cos \theta} + i \frac{\sin \theta (e^{2r} - 1)}{(1 + e^{2r}) + (1 - e^{2r}) \cos \theta}. \end{aligned}$$

Therefore,

$$(\lambda, t) = \left(\frac{1}{\cosh r - \cos \theta \sinh r}, \frac{\sin \theta \sinh r}{\cosh r - \cos \theta \sinh r} \right).$$

Definition 2.1. The map $\Phi : [0, \infty) \times [0, 2\pi) \rightarrow \mathbf{H}_{\mathbb{C}}^1$ given by

$$(r, \theta) \mapsto \left(\frac{1}{\cosh r - \cos \theta \sinh r}, \frac{\sin \theta \sinh r}{\cosh r - \cos \theta \sinh r} \right),$$

is the *hyperbolic polar coordinates map* in the hyperbolic plane.

To find the inverse Φ^{-1} we calculate for each (λ, t) :

$$\frac{\lambda^2 + t^2 + 1}{2\lambda} = \cosh r, \quad \frac{2t}{\lambda^2 + t^2 - 1} = \tan \theta.$$

Therefore

$$(2.1) \quad r = \operatorname{arccosh} \left(\frac{\lambda^2 + t^2 + 1}{2\lambda} \right)$$

and the angle θ is defined uniquely in the interval $[0, 2\pi)$ by the following:

$$(2.2) \quad \theta = \begin{cases} 0 & \text{if } t = 0, \lambda \geq 1, \\ \arctan \left(\frac{2t}{\lambda^2 + t^2 - 1} \right) & \text{if } (\lambda^2 + t^2 - 1) > 0, t > 0 \\ \pi/2 & \text{if } \lambda^2 + t^2 = 1, t > 0 \\ \pi - \arctan \left(\frac{2t}{\lambda^2 + t^2 - 1} \right) & \text{if } (\lambda^2 + t^2 - 1) < 0, t > 0, \\ \pi & \text{if } t = 0, \lambda < 1, \\ \pi + \arctan \left(\frac{2t}{\lambda^2 + t^2 - 1} \right) & \text{if } (\lambda^2 + t^2 - 1) < 0, t < 0, \\ 3\pi/2 & \text{if } \lambda^2 + t^2 = 1, t < 0, \\ 2\pi - \arctan \left(\frac{2t}{\lambda^2 + t^2 - 1} \right) & \text{if } (\lambda^2 + t^2 - 1) > 0, t < 0. \end{cases}$$

Therefore, to each $(\lambda, t) \in \mathbf{H}_{\mathbb{C}}^1$, $t \neq 0$ we can assign a unique pair (r, θ) given by the equations (2.1) and (2.2), respectively. These equations define the inverse map Φ^{-1} . Straight forward computations deduce that for the Jacobian J_{Φ} of Φ we have

$$J_{\Phi} = \lambda^2 \sinh r.$$

The expression for the metric tensor is given by

$$(2.3) \quad g = dr^2 + \sinh^2 r d\theta^2.$$

In hyperbolic polar coordinates we may thus write the line and the area element respectively as

$$ds_h = \sqrt{dr^2 + \sinh^2 r d\theta^2}, \quad d\mathcal{A}_h = \sinh r dr d\theta.$$

We finally note that hyperbolic circles $C_h((1, 0), R)$ are written in hyperbolic polar coordinates as $r = R$.

2.3. Modulus of curve families. Let Γ be a family of curves in $\mathbf{H}_{\mathbb{C}}^1$; we define its modulus $\text{Mod}(\Gamma)$ by

$$(2.4) \quad \text{Mod}(\Gamma) = \inf \left\{ \iint_{\mathbf{H}_{\mathbb{C}}^1} \rho^2 d\mathcal{A}_h \mid \rho \in \text{Adm}(\Gamma) \right\},$$

where $\text{Adm}(\Gamma)$, the set of *admissible functions*, or *densities* for Γ is the set of all Borel measurable $\rho : \mathbf{H}_{\mathbb{C}}^1 \rightarrow \mathbb{R}_+$ such that

$$\int_{\gamma} \rho ds_h \geq 1, \text{ for all } \gamma \in \Gamma.$$

Here,

$$ds_h = \frac{\sqrt{d\lambda^2 + dt^2}}{\lambda}, \quad d\mathcal{A}_h = \frac{d\lambda dt}{\lambda^2}.$$

A density $\rho_0 \in \text{Adm}(\Gamma)$ is called extremal if

$$\text{Mod}(\Gamma) = \iint_{\mathbf{H}_{\mathbb{C}}^1} \rho_0^2 d\mathcal{A}_h.$$

The modulus of curve family is a conformal invariant: if f is a conformal self-mapping of $\mathbf{H}_{\mathbb{C}}^1$, then $\text{Mod}(f(\Gamma)) = \text{Mod}(\Gamma)$. And this is because any such conformal mapping has to be a Möbius transformation preserving the right half plane, i.e. it has to be of the form

$$f(z) = \frac{az + ib}{icz + d}, \quad a, b, c, d \in \mathbb{R}, \quad ad + bc > 0.$$

3. MODULUS OF THE NORMAL HYPERBOLIC QUADRILATERAL




FIGURE 3.1. A normal hyperbolic quadrilateral.

A geodesic quadrilateral in $\mathbf{H}_{\mathbb{C}}^1$ is a quadrilateral whose sides are geodesic segments. We shall consider quadrilaterals whose opposite sides are parallel and also equal (that is, they have the same hyperbolic length. Any such quadrilateral whose two opposite sides are parallel to the λ -axis may be normalised according to the following.

Definition 3.1. A *normal quadrilateral* $Q(a, b)$, $a, b > 0$ is the set

$$Q(a, b) = \{(\lambda, t) \in \mathbf{H}_{\mathbb{C}}^1 \mid -b \leq t \leq b, \sqrt{1+b^2-t^2} \leq \lambda \leq \sqrt{a^2+b^2-t^2}\}.$$

If $\mathcal{A}_e(Q(a, b))$ is the Euclidean area of $Q(a, b)$, then

$$\begin{aligned} \mathcal{A}_e(Q(a, b)) &= \int_{-b}^b \int_{\sqrt{1+b^2-t^2}}^{\sqrt{a^2+b^2-t^2}} d\lambda dt \\ &= 2 \int_0^b \left(\sqrt{a^2+b^2-t^2} - \sqrt{1+b^2-t^2} \right) dt. \end{aligned}$$

Therefore

$$(3.1) \quad \mathcal{A}_e(Q(a, b)) = (a^2 + b^2) \arctan(b/a) - (1 + b^2) \arctan b + b(a - 1).$$

Any two normal quadrilaterals $Q(a, b)$ and $Q(a', b')$ are conformally equivalent, that is, there exists an isometry that maps

$$(1, \pm b) \mapsto (1, \pm b'), \quad (a, \pm b) \mapsto (a', \pm b'),$$

if and only if $(a - 1)/b = (a' - 1)/b'$. And this happens because there exists such an isometry if and only if the following equality of cross-ratios holds:

$$[1 - ib, a - ib, a + ib, a + ib] = [1 - ib', a' - ib', a' + ib', a' + ib'].$$

Recall that if $z_i \in \mathbf{H}_{\mathbb{C}}^1$, $i = 1, \dots, 4$, are pairwise distinct points, then their cross-ratio is

$$[z_1, z_2, z_3, z_4] = \frac{(z_4 - z_2)(z_3 - z_1)}{(z_4 - z_1)(z_3 - z_2)}.$$

Therefore for the following we are going to consider that normal quadrilateral $Q_a = Q(a, 1)$. By applying (3.1) and by using the identity

$$\arctan(1/a) = (\pi/2) - \arctan a,$$

we obtain

$$(3.2) \quad \mathcal{A}_e(Q_a) = \frac{\pi}{2}(a^2 + 2) - (a^2 + 1) \arctan a + (a - 1).$$

3.1. Modulus of the family of curves connecting the circular arcs. We consider the family Γ_Q^1 of curves within Q_a connecting the circular arcs

$$L : \lambda^2 + t^2 = 2, \text{ and } R : \lambda^2 + t^2 = a^2 + 1,$$

and the subfamily $\Gamma'_Q \subset \Gamma_Q^1$ comprising curves

$$\gamma_t(\lambda) = (\lambda, t), \quad \lambda \in \left[\lambda_1 = \sqrt{2 - t^2}, \lambda_2 = \sqrt{a^2 + 1 - t^2} \right].$$

Let $\rho \in \text{Adm}(\Gamma_Q^1) \supset \text{Adm}(\Gamma'_Q)$. Then,

$$1 \leq \int_{\gamma_t} \rho ds_h = \int_{\lambda_1}^{\lambda_2} \rho(\lambda, t) \frac{d\lambda}{\lambda}.$$

By integrating with respect to t we have

$$\begin{aligned} 2 &\leq \int_{-1}^1 \int_{\lambda_1}^{\lambda_2} \rho(\lambda, t) \frac{d\lambda dt}{\lambda} \\ &\leq \left(\int_{-1}^1 \int_{\lambda_1}^{\lambda_2} \rho^2(\lambda, t) \frac{d\lambda dt}{\lambda^2} \right)^{1/2} \left(\int_{-1}^1 \int_{\lambda_1}^{\lambda_2} d\lambda dt \right)^{1/2} \end{aligned}$$

where we have applied the Cauchy-Schwarz inequality in the last step. By raising to the square and rearranging we obtain:

$$\iint_{Q_a} \rho^2 d\mathcal{A}_h \geq \frac{4}{\mathcal{A}_e(Q_a)},$$

where $\mathcal{A}_e(Q_a)$ is the Euclidean area of Q_a . By taking infima over all $\rho \in \text{Adm}(\Gamma_Q^1)$ we obtain the inequality

$$\text{Mod}(\Gamma_Q^1) \geq \frac{4}{\mathcal{A}_e(Q_a)}.$$

To show that equality holds, we show that the function $\rho_0 : \mathbf{H}_{\mathbb{C}}^1 : \mathbb{R}_+$ defined by

$$\rho_0(\lambda, t) = \frac{2\lambda}{\mathcal{A}_e(Q_a)} \chi(Q_a)(\lambda, t),$$

is an extremal density. Here, $\chi(Q_a)$ is the characteristic function of Q_a . Indeed,

$$\iint_Q \rho_0^2 d\mathcal{A}_h = \int_{-1}^1 \int_{\lambda_1}^{\lambda_2} d\lambda dt = \frac{4}{\mathcal{A}_e(Q)}.$$

We thus have

Theorem 3.2. *The modulus of the family of curves Γ_Q^1 whose elements join the circular components of the boundary of the normal hyperbolic quadrilateral Q_a is*

$$\text{Mod}(\Gamma_Q^1) = \frac{4}{\mathcal{A}_e(Q_a)},$$

where $\mathcal{A}_e(Q(a, b))$ is the Euclidean area of the quadrilateral given by (3.2).

3.2. Modulus of the family of curves connecting the line segments. We next consider the family Γ_Q^2 of curves within Q_a connecting the line segments

$$B : t = -1, \text{ and } T : t = 1,$$

and the subfamily $\Gamma_Q'' \subset \Gamma_Q^2$ comprising curves $\gamma_\lambda(s)$ defined for every fixed $\lambda \in [1, a]$ as follows:

$$\gamma_\lambda(s) = \sqrt{1 + \lambda^2}(\cos s, \sin s), \quad s \in [-s_\lambda, s_\lambda], \quad s_\lambda = \arctan(1/\lambda).$$

Let as before $\rho \in \text{Adm}(\Gamma_Q^2) \supset \text{Adm}(\Gamma_Q'')$. Then,

$$1 \leq \int_{\gamma_\lambda} \rho ds_h = \int_{-s_\lambda}^{s_\lambda} \rho(\gamma_\lambda(s)) \frac{ds}{\cos s}.$$

The hyperbolic area element is

$$d\mathcal{A}_h = \frac{\lambda d\lambda ds}{(\lambda^2 + 1) \cos^2 s}.$$

Therefore

$$\begin{aligned}
a - 1 &\leq \int_1^a \int_{-s_\lambda}^{s_\lambda} \rho(\lambda, s) \frac{ds d\lambda}{\cos s} \\
&= \int_1^a \int_{-s_\lambda}^{s_\lambda} \frac{\lambda^{1/2} \rho(\lambda, s)}{(\lambda^2 + b^2)^{1/2} \cos s} \cdot \frac{(\lambda^2 + 1)^{1/2}}{\lambda^{1/2}} ds d\lambda \\
&\leq \left(\iint_Q \rho^2 d\mathcal{A}_h \right)^{1/2} \left(\int_1^a \int_{-s_\lambda}^{s_\lambda} \frac{\lambda^2 + 1}{\lambda} ds d\lambda \right)^{1/2} \\
&\leq \left(\iint_Q \rho^2 d\mathcal{A}_h \right)^{1/2} \left(2 \int_1^a \frac{\lambda^2 + 1}{\lambda} \arctan(1/\lambda) d\lambda \right)^{1/2}.
\end{aligned}$$

The integral I inside the right parenthesis is written as

$$I = I_1 + I_2 = 2 \int_1^a \lambda \arctan(1/\lambda) d\lambda + 2 \int_1^a \frac{\arctan(1/\lambda)}{\lambda} d\lambda.$$

We have

$$I_1 = \mathcal{A}_e(Q_a), \quad I_2 = 2 (\text{Ti}_2(1) - \text{Ti}_2(1/a)),$$

where

$$(3.3) \quad \text{Ti}_2(x) = \int_0^x \frac{\arctan t}{t} dt$$

is the *inverse tangent integral*, see [4]. Since $\text{Ti}_2(1) = \mathbf{G}$, where $\mathbf{G} \approx 0.915966$ is the *Catalan constant* and $\text{Ti}_2(1/a) = \text{Ti}_2(a) - (\pi/2) \log a$ we thus have

$$I_2 = 2 \left(\mathbf{G} + \frac{\pi}{2} \log a - \text{Ti}_2(a) \right).$$

Hence

$$\text{Mod}(\Gamma_Q^2) \geq \frac{(a-1)^2}{I}.$$

Now the function $\rho_0 : \mathbf{H}_{\mathbb{C}}^1 \rightarrow \mathbb{R}_+$ defined by

$$\rho_0(\lambda, s) = \frac{a-1}{I} \frac{\lambda^2 + 1}{\lambda} \chi(Q)(\lambda, s),$$

is an extremal density:

$$\iint_Q \rho_0^2 d\mathcal{A}_h = \frac{(a-1)^2}{I} \int_1^a \int_{-s_\lambda}^{s_\lambda} \frac{\lambda^2 + b^2}{\lambda} d\lambda ds = \frac{(a-1)^2}{I}.$$

We have proved:

Theorem 3.3. *The modulus of the family of curves Γ_Q^2 whose elements join the straight line segment components of the boundary of the normal hyperbolic quadrilateral Q_a is*

$$\text{Mod}(\Gamma_Q^2) = \frac{(a-1)^2}{\mathcal{A}_e(Q_a) + 2 \left(\mathbf{G} + \frac{\pi}{2} \log a - \text{Ti}_2(a) \right)},$$

where $\mathcal{A}_e(Q_a)$ is the Euclidean area of the quadrilateral given by (3.2), Ti_2 is the inverse tangent integral as in (3.3) and \mathbf{G} is the Catalan constant.

4. MODULUS OF THE HYPERBOLIC CIRCULAR ANNULUS

Definition 4.1. A hyperbolic annulus $A(z_0, R_1, R_2)$ is the domain enclosed between the hyperbolic circles $C(z_0, R_1)$ and $C(z_0, R_2)$, that is,

$$A(z_0, R_1, R_2) = \{z \in \mathbf{H}_{\mathbb{C}}^1 \mid \cosh(R_1) \leq 1 + \frac{|z - z_0|^2}{2\Re(z)\Re(z_0)} \leq \cosh R_2\}.$$

Any hyperbolic annulus can be conformally mapped onto an annulus of the form $A((1, 0), 1, R)$, $R > 1$, which for brevity we shall denote by A_R . In hyperbolic coordinates

$$A_R = \{(r, \theta) \mid 1 \leq r \leq R\}.$$




FIGURE 4.1. The white region between the circles is the hyperbolic annulus.

We consider the domain A_R , see Figure 4.1 and two families of curves in A_R .

4.1. Modulus of the family of connecting curves. Let Γ_A^1 be the family of curves in A_R with each endpoint at each circle. Let Γ'_A be the subfamily comprising curves γ_θ , $\theta \in [0, 2\pi)$ such that

$$\gamma_\theta(s) = (s, \theta), \quad s \in [1, R].$$

If ρ is admissible for Γ_A^1 , then

$$\int_1^R \rho(s, \theta) ds \geq 1.$$

Therefore

$$\begin{aligned}
2\pi &\leq \int_1^R \int_0^{2\pi} \rho(r, \theta) dr d\theta \\
&= \int_1^R \int_0^{2\pi} \rho(r, \theta) (\sinh r)^{1/2} \cdot (\sinh r)^{-1/2} dr d\theta \\
&\leq \left(\int_1^R \int_0^{2\pi} \rho^2(r, \theta) \sinh r dr d\theta \right)^{1/2} \cdot \left(\int_1^R \int_0^{2\pi} \frac{dr d\theta}{\sinh r} \right)^{1/2},
\end{aligned}$$

where we have used Cauchy-Schwarz inequality in the last step. We take this inequality to the square and by using the fact that

$$\int_1^R \int_0^{2\pi} \frac{dr d\theta}{\sinh r} = 2\pi \log \left(\frac{\tanh(R/2)}{\tanh(1/2)} \right),$$

we obtain

$$\frac{2\pi}{\log \left(\frac{\tanh(R/2)}{\tanh(1/2)} \right)} \leq \text{Mod}(\Gamma).$$

We now show that the density ρ_0 which is such that

$$\rho_0(r, \theta) = \frac{c_0}{\sinh r} \chi_{A_R}(r, \theta), \quad c_0 = \left(\log \left(\frac{\tanh(R/2)}{\tanh(1/2)} \right) \right)^{-1},$$

is extremal. Indeed,

$$\begin{aligned}
\int_1^R \int_0^{2\pi} \rho_0^2(r, \theta) \sinh(2r) dr d\theta &= 2\pi c_0^2 \int_1^R \frac{dr}{\sinh r} \\
&= \frac{2\pi}{\log \left(\frac{\tanh(R/2)}{\tanh(1/2)} \right)}.
\end{aligned}$$

We thus have

Theorem 4.2. *The modulus of the family of curves Γ_A^1 whose elements join the boundary components of the hyperbolic circular annulus A_R is*

$$\text{Mod}(\Gamma_A^1) = \frac{2\pi}{\log \left(\frac{\tanh(R/2)}{\tanh(1/2)} \right)}.$$

4.2. Modulus of the family of separating curves. We next consider the family Γ_A^2 of closed curves inside A_R which wind around $(1, 0)$. We consider the subfamily Γ_A'' of hyperbolic circles centred at $(1, 0)$. An element of this subfamily is of the form

$$\gamma_r(s) = (r, s), \quad s \in [0, 2\pi].$$

If ρ is an admissible density for Γ_A^2 , then

$$\int_0^{2\pi} \rho(r, s) \sinh r ds \geq 1.$$

Now from

$$\frac{1}{\sinh r} \leq \int_0^{2\pi} \rho(r, \theta) d\theta$$

we have that

$$\begin{aligned} \int_1^R \frac{dr}{\sinh r} &\leq \int_1^R \int_0^{2\pi} \rho(r, \theta) (\sinh r)^{1/2} \cdot (\sinh r)^{-1/2} dr d\theta \\ &\leq \left(\int_1^R \int_0^{2\pi} \rho^2(r, \theta) \sinh r dr d\theta \right)^{1/2} \cdot \left(\int_1^R \int_0^{2\pi} \frac{1}{\sinh r} dr d\theta \right)^{1/2}. \end{aligned}$$

After raising to the square, rearranging and taking infimum over all admissible ρ we obtain

$$\text{Mod}(\Gamma_A^2) \geq \frac{1}{2\pi} \log \left(\frac{\tanh(R/2)}{\tanh(1/2)} \right).$$

We verify that the extremal density is

$$\rho'(r, \theta) = \frac{1}{2\pi \sinh r} \chi_{A_R}.$$

Indeed,

$$\begin{aligned} \iint_{A_R} (\rho')^2(r, \theta) \sinh r dr d\theta &= \frac{1}{4\pi^2} \int_1^R \int_0^{2\pi} \frac{dr d\theta}{\sinh r} \\ &= \frac{1}{2\pi} \int_1^R \frac{dr}{\sinh r} \\ &= \frac{1}{2\pi} \log \left(\frac{\tanh(R/2)}{\tanh(1/2)} \right). \end{aligned}$$

We have proved

Theorem 4.3. *The modulus of the family of closed curves Γ_A^2 whose elements separate the boundary components of the hyperbolic circular annulus A_R is*

$$\text{Mod}(\Gamma_A^2) = \frac{1}{2\pi} \log \left(\frac{\tanh(R/2)}{\tanh(1/2)} \right).$$

REFERENCES

- [1] Lars V. Ahlfors. *Lectures on quasiconformal mappings. With additional chapters by C. J. Earle and I. Kra, M. Shishikura and J. H. Hubbard*, volume 38 of *Univ. Lect. Ser.* Providence, RI: American Mathematical Society (AMS), 2nd enlarged and revised ed. edition, 2006.
- [2] Zoltán M. Balogh, Elia Bubani, and Ioannis D. Platis. Stretch maps on the affine-additive group. *Anal. Math. Phys.*, 15(3):39, 2025. Id/No 61.
- [3] Zoltán M. Balogh, Katrin Fässler, and Ioannis D. Platis. Modulus method and radial stretch map in the Heisenberg group. *Ann. Acad. Sci. Fenn., Math.*, 38(1):149–180, 2013.
- [4] Leonard Lewin. *Polylogarithms and associated functions*. New York, Oxford: North Holland. XVII, 359 p. \$ 54.75 (USA + Canada); \$ 73.25; Dfl. 150.00 (1981)., 1981.
- [5] William Ma and David Minda. Euclidean properties of hyperbolic polar coordinates. *Comput. Methods Funct. Theory*, 6(1):223–242, 2006.
- [6] Ioannis D. Platis. Modulus of revolution rings in the Heisenberg group. *Proc. Am. Math. Soc.*, 144(9):3975–3990, 2016.
- [7] A. Vasil'ev. *Moduli of families of curves for conformal and quasiconformal mappings*, volume 1788 of *Lect. Notes Math.* Berlin: Springer, 2002.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PATRAS, UNIVERSITY CAMPUS, GR-26504 RION, ACHAIA, GREECE.

Email address: idplatis@math.upatras.gr