

THE DIOPHANTINE EQUATION $(2^k - 1)(b^k - 1) = y^q$

CHANG LIU AND BO HE

ABSTRACT. In this paper, we consider the exponential Diophantine equation $(2^k - 1)(b^k - 1) = y^q$ with $k \geq 2$, odd integer b and an odd prime exponent q and obtain effective upper bounds for q in terms of b . In particular, we show that $q \leq \log_2(b + 1)$ holds apart from a finite, explicitly determined set of exceptional pairs (b, q) when $3 \leq b < 10^6$. As an application, we prove that the related equation $(2^k - 1)(b^k - 1) = x^n$, has no positive integer solution (k, x, n) for several specific odd values of b , including $b \in \{5, 7, 11, 13, 21, 23, 27, 29\}$.

1. INTRODUCTION

In 2002, Szalay [9] investigated the Diophantine equation $(2^n - 1)(3^n - 1) = x^2$ and proved that there is no solution in positive integers n and x , which is the first investigation for the Diophantine equation

$$(1) \quad (a^n - 1)(b^n - 1) = x^2.$$

Moreover, he studied similar equations such as $(2^n - 1)(5^n - 1) = x^2$, demonstrating that they have only limited solutions with specific values of n . And in the same year, Hajdu and Szalay [4] have shown that there is no solution for $(2^n - 1)(6^n - 1) = x^2$.

Cohn's [3] studied the equation $(a^n - 1)(b^n - 1) = x^2$ and explores the integer solutions for given values of a and b , deriving general results and conjectures about this equation in 2001. Subsequently, Luca and Walsh [7] performed extensive computational investigation solve this equations (1) for nearly all pairs (a, b) satisfying $2 \leq b < a \leq 100$, leaving only 70 unresolved cases in 2002. After that, Le [6] proved if $3|b$, then the equation $(2^n - 1)(b^n - 1) = x^2$ has no positive integer solution (n, x) in 2009.

Very recently, the authors complete solved [5] the first example of the exponential Diophantine equation

$$(2) \quad (a^n - 1)(b^n - 1) = y^q$$

for $(a, b) = (2, 3)$ and odd prime q . It was shown that the equation

$$(3) \quad (2^k - 1)(3^k - 1) = y^q$$

has no solutions in positive integers (x, k, q) with q is an odd prime. One of the key steps in the proof is to rewrite equation (3) in the form $(X^q - 1)(Y^q - 1) = Z^q$, by setting $X = 2^{k/q}$, $Y = 3^{k/q}$ and $Z = y$. Observe that this equation is very close in shape to $(x^k - 1)(y^k - 1) = z^k - 1$, which was completely resolved in Bennett's work [2].

In this paper, we first analyse the equation

$$(4) \quad (X^q - 1)(Y^q - 1) = Z^q,$$

as a preliminary step towards solving equation (2) in more generality.

Date: December 2, 2025.

Theorem 1. *The equation (4) has no solution in integers X, Y, Z and odd prime q with $1 < X \leq Y$.*

In view of existing results, in order to solve equation (2) for fixed pairs (a, b) , it is enough to show that $q \mid k$. In this paper we concentrate on the case $a = 2$ and b an odd integer in equation (2), and we establish the following result.

Theorem 2. *When $b \leq 10^6$ is an odd integer, the equation (15) has no solution with positive integers $k, y \geq 2$ and $q > \log_2(b+1)$ except for the following special cases.*

q	Remaining b (odd, $3 \leq b < 10^6$)
5	15, 17
11	1023, 1025
13	4095, 4097
17	$t \cdot 2^{14} \pm 1$ ($t = 1, 2, \dots, 7$)

Theorem 3. *The Diophantine equation*

$$(2^k - 1)(b^k - 1) = x^n, \quad n \geq 2$$

has no positive solution (k, x, n) for odd $b = 5, 7, 11, 13, 21, 23, 27, 29$.

2. THE DIOPHANTINE EQUATION $(X^q - 1)(Y^q - 1) = Z^q$

Let consider the diophantine equation

$$(5) \quad (X^q - 1)(Y^q - 1) = Z^q, \quad 1 < X < Y.$$

Our method is base on Bennett's works on rational approximation to algebraic numbers [1] and his proof [2] of diophantine equation $(x^k - 1)(y^k - 1) = z^k - 1$.

Let

$$(6) \quad A := X^q - 1, \quad B := Y^q - 1.$$

Then $AB = Z^q$. There exists a positive integer t such that

$$XY = Z + t.$$

Expanding the expression

$$\left((AB)^{1/q} + t\right)^q = (A+1)(B+1),$$

yields

$$(7) \quad q(AB)^{(q-1)/q}t + \binom{q}{2}(AB)^{(q-2)/q}t^2 + \dots + t^q = A + B + 1.$$

We claim that

$$A < \binom{q}{2}(AB)^{(q-2)/q}t^2.$$

If $q \geq 4$, this is immediate from $B \geq A$. If $q = 3$ and $A \geq 3t^2(AB)^{1/3}$, then $B < A^2$, it follows that

$$3(AB)^{2/3}t > 3Bt > A + B + 1.$$

which is a contradiction with equation (7). Therefore,

$$q(AB)^{(q-1)/q}t < B,$$

which is

$$(8) \quad B > q^q A^{q-1} t^q.$$

Next, observe that from (6), we have

$$\left(\frac{XY}{Z}\right)^q - \frac{A+1}{A} = \frac{A+1}{AB} < \frac{2A}{Z^q},$$

It follows that

$$(9) \quad \left| \sqrt[q]{1 + \frac{1}{A}} - \frac{XY}{Z} \right| < \frac{2A}{q \cdot Z^q}$$

Now we appeal to another result to deduce a lower bound which will contradict (9).

Lemma 1. [1] *Let $k, A \in \mathbb{Z}_{>0}$ with $k \geq 3$. Define*

$$\mu_n = \prod_{p|n} p^{1/(p-1)},$$

and suppose

$$(10) \quad \left(\sqrt{A} + \sqrt{A+1}\right)^{2(k-2)} > (k\mu_k)^k.$$

Then

$$\left| \sqrt[k]{1 + \frac{1}{A}} - \frac{p}{q} \right| > (8k\mu_k A)^{-1} \cdot q^{-\lambda},$$

with

$$(11) \quad \lambda = 1 + \frac{\log \left(k\mu_k (\sqrt{A} + \sqrt{A+1})^2 \right)}{\log \left(\frac{1}{k\mu_k} (\sqrt{A} + \sqrt{A+1})^2 \right)}.$$

and $\lambda < k$.

In our case, since $A = X^q - 1$, it is easy to verify that inequality (10) fails only when $(X, q) \in \{(2, 3), (3, 3)\}$. We suppose that (x, q) lies outside the set in $\{(2, 3), (3, 3)\}$. Combining (9) with Lemma 1 gives

$$(12) \quad Z^{q-\lambda} < 16\mu_q A^2,$$

thus,

$$(13) \quad B^{q-\lambda} < 16^q \mu_q^q A^{q+\lambda} = 16^q q^{\frac{q}{q-1}} A^{q+\lambda}$$

From $A = X^q - 1$, the expression for λ in (11) shows that λ is monotonically decreasing in $X \geq 2$ (for $q \geq 7$), so $\lambda < 3.15$. Therefore, (13) implies

$$B < 300 \cdot A^{2.7},$$

which contradicts (8).

Similarly, if $q = 5$ and $X \geq 3$, then $\lambda < 2.8$, hence

$$B < 1400 \cdot A^{3.6},$$

which is again contradicting with (8). There is no contradiction with upper bound and lower bound for b when $(X, q) = (2, q)$. We consider it separately. Let $q = 5$ and $X = 2$, then

$$(14) \quad 31(Y^5 - 1) = Z^5,$$

and therefore

$$\left| \sqrt[5]{31} - \frac{Z}{Y} \right| < \frac{31^{1/5}}{5Y^5}.$$

However, by Corollary 1.2 of [1],

$$\left| \sqrt[5]{31} - \frac{Z}{Y} \right| > \frac{0.01}{Y^{2.83}}.$$

it follows that $Y \leq 6$. Through simple verification, the equation has no solution.

For the case $q = 3$ we make a slight modification of the argument in [2]. Starting from the equation

$$(X^3 - 1)Y^3 - Z^3 = X^3 - 1$$

and writing $\alpha = (X^3 - 1)^{1/3}$, one obtains a very good rational approximation Z/Y to α , with $|\alpha - Z/Y| \ll 2Y^{-2}$. By Legendre's criterion, Z/Y must be a convergent of the simple continued fraction of α . Then, a careful inspection of the initial partial quotients of $\sqrt[3]{X^3 - 1}$ shows that any such convergent arising from a solution forces a lower bound of the shape

$$Y \geq 5X^6.$$

This contradicts the corresponding upper bound given in Lemma 1 for sufficiently large values of X . The remaining small values reduce to $X = 2, 3$, and these can be checked out by the same continued fraction estimating as in [2]. Hence the case $q = 3$ cannot occur, and the proof of Theorem 1 is complete.

3. THE FIRST UPPER BOUND OF q

In this section, we will get a relative upper bound of q by proving the following result.

Theorem 4. *For odd integer b and odd prime q , if (k, y, q) is an integer solution of Diophantine equation*

$$(15) \quad (2^k - 1)(b^k - 1) = y^q,$$

then we have $q \leq 2\sqrt{2b}$.

Let p be a prime and $n \in \mathbb{Z} \setminus \{0\}$. The p -adic valuation $\nu_p(n)$ is the largest integer $e \geq 0$ such that $p^e \mid n$. The following result, known as the Lifting-the-Exponent (LTE) Lemma, is a standard tool for estimating p -adic valuations of exponential differences.

Lemma 2 (Lifting-the-exponent Lemma [8]). *Let p be a prime, and let a and b be integers such that k is a positive integer. Suppose $p \mid (a - b)$ and $p \nmid ab$. Then, the p -adic valuation ν_p of $a^k - b^k$ is given by*

$$\nu_p(a^k - b^k) = \begin{cases} \nu_p(a - b) + \nu_p(k), & \text{if } p \text{ is odd,} \\ \nu_2(a - b), & \text{if } p = 2 \text{ and } k \text{ is odd,} \\ \nu_2(a^2 - b^2) + \nu_2\left(\frac{k}{2}\right), & \text{if } p = 2 \text{ and } k \text{ is even.} \end{cases}$$

Lemma 3. *Assume that (k, b, y, q) is a positive integer solution of equation (15) with $q > \log_2(b + 1)$. We have*

- (1) $\nu_2(k) > q - \log_2(b + 1) > 0$,
- (2) $\nu_3(k) > 0$.

Proof. For point (1), since b is odd, we have $2 \mid (b^k - 1)$, hence $2 \mid y$. Thus,

$$b^k - 1 \equiv 0 \pmod{2^q}.$$

Applying Lemma 2, we obtain

$$q \leq \nu_2(y^q) = \nu_2(b^k - 1) = \begin{cases} \nu_2(b - 1), & \text{if } k \text{ is odd,} \\ \nu_2(b^2 - 1) + \nu_2(k/2), & \text{if } k \text{ is even.} \end{cases}$$

It is a contradiction that both $q \leq \nu_2(b - 1)$ and $q > \log_2(b + 1)$. Therefore $2 \mid k$ and

$$\nu_2(k) \geq q + 1 - \nu_2(b^2 - 1) \geq q - \log_2(b + 1).$$

Here we used $\min\{\nu_2(b + 1), \nu_2(b - 1)\} = 1$.

When $p = 3$ in point (2), the condition $2^k - 1 \equiv 0 \pmod{3}$ implies that $3 \mid y$. Thus, we have

$$(2^k - 1)(b^k - 1) \equiv 0 \pmod{3^q}.$$

If $3 \mid b$, another application of Lemma 2 to $4^{k/2} - 1$ yields $3^{q-1} \mid k$. Otherwise,

$$\nu_3(b^k - 1) = \nu_3(b^2 - 1) + \nu_3(k/2) = \nu_3(b - 1) + \nu_3(b + 1) + \nu_3(k) \leq \log_3(b + 1) + \nu_3(k),$$

since odd prime p cannot divide both $b + 1$ and $b - 1$ simultaneously. Hence

$$\nu_3(k) \geq \frac{q - \log_3(b + 1) - 1}{2}.$$

The right side is positive when $b \geq 7$. The remain cases that $b = 5$ provides $\nu_3(b + 1) = 1$, one can check directly. This complete the proof of Lemma 3. \square

Lemma 4. Assume that (k, b, y, q) is a quadruple of positive integers solving equation (15). Let p be an odd prime. If $(p - 1) \mid k$, then

$$\nu_p(k) > \frac{1}{2} \left(q - p \cdot \frac{\log 2b}{2 \log p} \right).$$

In particular, if we further assume that $p \leq q$ and $q > 2\sqrt{2b}$, then $p \mid k$.

Proof. Let us begin by analyzing the p -adic valuation on both sides of equation (15). Assumed that $p - 1$ divides k . Then, by Fermat's Little Theorem, it follows that p divides both $b^{p-1} - 1$ and $b^k - 1$ for any integer b satisfying $p \nmid b$. By applying Lemma 2, we obtain

$$\nu_p(2^k - 1) = \nu_p(2^{p-1} - 1) + \nu_p\left(\frac{k}{p-1}\right) = \nu_p\left(2^{\frac{p-1}{2}} - 1\right) + \nu_p\left(2^{\frac{p-1}{2}} + 1\right) + \nu_p(k),$$

$$\nu_p(b^k - 1) = \begin{cases} \nu_p\left(b^{\frac{p-1}{2}} - 1\right) + \nu_p\left(b^{\frac{p-1}{2}} + 1\right) + \nu_p(k), & \text{if } p \nmid b, \\ 0, & \text{if } p \mid b. \end{cases}$$

Next, we observe that

$$\nu_p\left(2^{\frac{p-1}{2}} - 1\right) + \nu_p\left(2^{\frac{p-1}{2}} + 1\right) < p \cdot \frac{\log 2}{2 \log p},$$

$$\nu_p\left(b^{\frac{p-1}{2}} - 1\right) + \nu_p\left(b^{\frac{p-1}{2}} + 1\right) < p \cdot \frac{\log b}{2 \log p}.$$

As a consequence,

$$\nu_p((2^k - 1)(b^k - 1)) < p \cdot \frac{\log 2b}{2 \log p} + 2\nu_p(k).$$

On the right side of equation (15), since $p \mid y$, it implies that

$$\nu_p(y^q) \geq q.$$

Combining the inequalities derived above, we conclude

$$\nu_p(k) > \frac{1}{2} \left(q - p \cdot \frac{\log 2b}{2 \log p} \right).$$

Assume that $p \leq q$ and $q > 2\sqrt{2b}$. We have

$$\nu_p(k) > \frac{1}{2} \left(q - p \cdot \frac{\log 2b}{2 \log p} \right) \geq \frac{1}{2} \left(q - q \cdot \frac{\log 2b}{2 \log q} \right) > 0$$

which completes the proof of the lemma. \square

Lemma 5. *Assume that (k, b, y, q) is a quadruple of positive integers solving equation (15) with $q \geq 2\sqrt{2b}$, then we have $q \mid k$.*

Proof. If $q = 3$, the result follows directly from Lemma 3 with $1 + \log_2(b+1) < 2\sqrt{2b}$. Define $P_n = \{p_1 = 2, p_2 = 3, p_3, p_4, \dots, p_n\}$ to be the set of all primes up to q , where $p_3 = 5 < p_4 < \dots < p_n = q$. To prove the result, it suffices to show that $(p-1) \mid k$ for each $p \in P$, so that Lemma 4 can be applied.

Assume that for some i , we already have $(p_i - 1) \mid k$. We aim to prove $(p_{i+1} - 1) \mid k$. It suffices to establish that

$$(16) \quad \nu_r(p_{i+1} - 1) \leq \nu_r(k)$$

for all $r \in P_i$.

We first consider $r = 2, 3$. By Lemma 3 and elementary monotonicity,

$$\nu_2(p_{i+1} - 1) \leq \frac{\log(p_{i+1} - 1)}{\log 2} \leq \frac{\log(q - 1)}{\log 2} < q - \log_2(b + 1) \leq \nu_2(k).$$

$$\nu_3(p_{i+1} - 1) \leq \frac{\log(p_{i+1} - 1)}{\log 3} \leq \frac{\log(q - 1)}{\log 3} < \frac{q - \log_3(b + 1) - 1}{2} < \nu_3(k).$$

Next, we consider $r \geq 5$. We analyze two sub cases based on the size between r and $q^{1/2}$.

If $r > q^{1/2}$, it follows that

$$\nu_r(p_{i+1} - 1) \leq \frac{\log(p_{i+1} - 1)}{\log r} < \frac{\log q}{\log r} < \frac{\log r^2}{\log r} = 2.$$

Since $r \in P_i = \{2, 3, p_1, p_2, \dots, p_i\}$, then $(r - 1) \mid k$ by assumption. Thus, by Lemma 4, we deduce that $r \mid k$. Therefore

$$\nu_r(p_{i+1} - 1) \leq 1 \leq \nu_r(k).$$

If $5 \leq r < q^{1/2}$, we claim that

$$\frac{1}{2} \left(q - r \cdot \frac{\log 2b}{2 \log r} \right) \geq \frac{\log(q - 1)}{\log r}.$$

It suffices to prove

$$\frac{1}{2} \left(q - q^{\frac{1}{2}} \cdot \frac{\log \frac{q^2}{4}}{2 \log q^{\frac{1}{2}}} \right) - \frac{\log(q - 1)}{\log 5} \geq 0.$$

which indeed holds, as one verifies by examining the monotonicity of the left side. Therefore, we have shown that $(p_{i+1} - 1) \mid k$. Consequently, by Lemma 4, we

conclude $p_{i+1} \mid k$. This completes the induction, and hence every prime in the set P divides k , including q . This completes the proof. \square

Now, assume that (k, b, y, q) is a quadruple of positive integers solving equation (15) under the assumption of Theorem 4. By Lemma 5, we have deduced $q \mid k$. Define that

$$(X, Y, Z) := (2^{k/q}, b^{k/q}, y),$$

we obtain a new triple of positive integers that satisfies the Diophantine equation

$$(X^q - 1)(Y^q - 1) = Z^q.$$

Theorem 1 has shown that it is no positive integer solution. This complete the proof of Theorem 4.

4. THE SECOND UPPER BOUND OF q

In this section, we sharp the upper bound of q for some not too big value of odd integer b . By Theorem 4, we have shown that for every odd integer $3 \leq b < 10^6$ there is no solution whenever q is an odd prime with $q > 2828$.

Lemma 6. *If odd integer $b \leq 10^6$, then the corresponding results in Lemmas 3–5 hold for every prime $q \geq 23$. Moreover, under these same conditions, the Diophantine equation (15) admits no solutions.*

Proof. Since $q \geq 23 > \log_2(b+1)$, we have $\nu_2(k) > 1$ and $\nu_3(k) > 0$. Therefore, Lemma 3 applies.

Then we investigate the exponents of prime divisors in $\nu_p(2^k - 1)$ and $\nu_p(b^k - 1)$. By Lemma 2,

$$\nu_p(2^k - 1) = \nu_p(2^{p-1} - 1) + \nu_p\left(\frac{k}{p-1}\right) = \nu_p(2^{p-1} - 1) + \nu_p(k),$$

and $\nu_p(2^{p-1} - 1) = 1$ for almost every prime $p < 2828$ except for $p = 1093$ where $\nu_{1093}(2^{1092} - 1) = 2$.

Similarly,

$$\nu_p(b^k - 1) = \nu_p(b^{p-1} - 1) + \nu_p\left(\frac{k}{p-1}\right) = \nu_p(b^{p-1} - 1) + \nu_p(k),$$

through computer calculation, $\nu_p(b^{p-1} - 1) \leq 11$. Hence,

$$(17) \quad \nu_p(k) \geq \frac{q - \nu_p(b^{p-1} - 1) - \nu_p(2^{p-1} - 1)}{2} \geq 5,$$

so Lemma 4 holds.

For Lemma 5, assume that for some i , we have $(p_i - 1) \mid k$. Our goal is to show that $(p_{i+1} - 1) \mid k$. It suffices to establish that

$$\nu_r(p_{i+1} - 1) \leq \nu_r(k)$$

for all $r \in P_i$. And it has proved that $\nu_r(k) \geq 5$ by (17).

We begin with the case $r = 2, 3$. By Lemma 3,

$$q - \log_2(b+1) \leq \nu_2(k)$$

It is obvious that

$$\nu_2(p_{i+1} - 1) \leq \nu_2(q - 1) < q - \log_2(b+1).$$

for all odd prime $q \geq 23$ and $b \leq 10^6$. Similarly,

$$\nu_3(p_{i+1} - 1) \leq \nu_3(q - 1) < \frac{q - \log_3(b + 1) - 1}{2} \leq \nu_3(k).$$

for all odd prime $q \geq 23$ and $b \leq 10^6$.

For $r \geq 5$, we have

$$\nu_r(p_{i+1} - 1) \leq \nu_5(q - 1) \leq 4 \leq \nu_r(k).$$

Hence Lemma 5 is true, which means $q \mid k$. Using Theorem 1, we complete the proof of this proposition. \square

Now, we prove Theorem 2.

Proof. There are a few possibilities for $3 \leq q < 23$. By Lemma 3, we have $\nu_2(k) > 0$ and $\nu_3(k) > 0$. Let p be a prime with $3 \leq p \leq q$. We claim that if $(p - 1) \mid k$ then $p \mid k$. Since we already know that $3 \mid k$, it remains to consider $5 \leq p \leq q < 23$. Likewise,

$$\nu_p((2^k - 1)(b^k - 1)) = \nu_p((2^{p-1} - 1)(b^{p-1} - 1)) + 2\nu_p(k) \geq q$$

Hence

$$\nu_p(k) \geq \frac{q - \nu_p(2^{p-1} - 1) - \nu_p(b^{p-1} - 1)}{2}$$

Since $\nu_p(2^{p-1} - 1) = 1$, we only need to consider those b for which $\nu_p(b^{p-1} - 1) \geq 4$ (as $q \geq 5$). A finite computer calculation shows that there are no pairs (b, p) with $q - 1 - \nu_p(b^{p-1} - 1) \leq 0$ under assumption.

Therefore, $q > \log_2(b + 1)$, $3 \mid k$ and $7 \mid k$ when $q = 3$ or $q = 7$ since $2 \mid k$ and $6 \mid k$ respectively. Thus by Theorem 1 there is no solution to equation (15).

Now consider the case $q = 5$. If $\nu_2(k) \geq 2$, then $4 \mid k$ and hence $5 \mid k$. Therefore, $\nu_2(k) = 1$, which forces $\nu_2(b^2 - 1) = 5n$ where $n \in \mathbb{Z}_+$. Combination with $5 = q > \log_2(b + 1)$, the remaining unsolved cases are $b = 15, 17$.

Similarly, for $q = 11$ and $q = 13$ the remaining unsolved cases are $b = 1023, 1025$ and $b = 4095, 4097$, respectively.

Next, consider the case $q = 17$. If $\nu_2(k) \geq 4$, then $16 \mid k$ and hence $17 \mid k$. Therefore, $1 \leq \nu_2(k) \leq 3$, which implies $15 \leq \nu_2(b^2 - 1) \leq 17$. Combination with $17 = q > \log_2(b + 1)$, the remaining unsolved cases are $b = t \cdot 2^{14} \pm 1$ where $t = 1, 2, \dots, 7$.

Finally, consider $q = 19$. We have $\nu_3(k) \geq \frac{q - \log_3(b + 1) - 1}{2} = \frac{19 - \log_3(b + 1) - 1}{2} > 2$. Then $18 \mid k$ and hence $19 \mid k$. By Theorem 1, there is no solution for equation (15).

The remaining unresolved instances, with an odd prime q and odd positive $b < 10^6$ where $q > \log_2(b + 1)$ are summarized in the table. This complete the proof of Theorem 2. \square

5. PROOF OF THEOREM 3

Assume that (k, x, n) is a solution in positive integers of

$$(18) \quad (2^k - 1)(b^k - 1) = x^n, \quad k, n > 1.$$

First consider the case $n = 2$. By Theorem 3.1 in [7] there is no solution in positive integers k, x for any odd integer $b \leq 100$. Thus, we may assume that $n > 2$. Let

q be the least prime divisor of n and put $y = x^{n/q}$. Then (k, y, q) is a solution in positive integers of equation(15), namely

$$(2^k - 1)(b^k - 1) = y^q.$$

We now restrict to the four values $b = 5, 7, 11, 13, 21, 23, 27, 29$. By Theorem 2 we have

$$q \leq \log_2(b+1) < 5,$$

hence q must be 3, since q is an odd prime.

For $b = 5$, Theorem 2 yields

$$q < \log_2 6 < 3,$$

which is impossible for an odd prime q .

In the cases $b = 7, 11, 13, 21, 23, 27, 29$, comparing the 2-adic valuations on both sides of

$$(2^k - 1)(b^k - 1) = y^3$$

shows that we must have $2 \mid k$. Applying Lemma 2 with $p = 3$ gives

$$\nu_3(2^k - 1) = \nu_3(4 - 1) + \nu_3(k/2) = 1 + \nu_3(k)$$

and

$$\nu_3(b^k - 1) = \begin{cases} \nu_3(b - 1) + \nu_3(k) & \text{if } b = 7, 13, 29; \\ \nu_3(b^2 - 1) + \nu_3(k) & \text{if } b = 11, 23, 29; \\ 0 & \text{if } b = 21, 27. \end{cases}$$

Consequently,

$$\nu_3((2^k - 1)(b^k - 1)) = \begin{cases} 2 + 2\nu_3(k) & \text{if } b = 7, 13; \\ 2 + 2\nu_3(k) & \text{if } b = 11, 23, 29; \\ 1 + \nu_3(k) & \text{if } b = 21, 27. \end{cases}$$

On the other hand,

$$\nu_3(y^q) = \nu_3(y^3) = 3\nu_3(y),$$

and the equality of both sides implies

$$3 = q \leq \nu_3(y^q) = \begin{cases} 2 + 2\nu_3(k) & \text{if } b = 7, 13; \\ 2 + 2\nu_3(k) & \text{if } b = 11, 23, 29; \\ 1 + \nu_3(k) & \text{if } b = 21, 27. \end{cases},$$

so $3 \mid k$. This contradicts Theorem 1, which is the equation $(X^q - 1)(Y^q - 1) = Z^q$ has no integer solution with $1 < X \leq Y$ and q an odd prime.

Combining all the above cases, we conclude that for $b = 5, 7, 11, 13, 21, 23, 27, 29$ the Diophantine equation

$$(2^k - 1)(b^k - 1) = x^n, \quad n \geq 2,$$

admits no solution in positive integers (k, x, n) . This completes the proof of Theorem 3.

ACKNOWLEDGMENTS

The first author acknowledges the support of the China Scholarship Council program (Project ID: 202106310023). The second author was supported by Natural Science Foundation of China (Grant No. 12161001).

REFERENCES

- [1] M. A. Bennett, *Explicit lower bounds for rational approximation to algebraic numbers*, Proceedings of the London Mathematical Society, **75**(1)(1997), 63–78.
- [2] M. A. Bennett, *The Diophantine equation $(x^k - 1)(y^k - 1) = (z^k - 1)^t$* , Indagationes Mathematicae, **18**(4)(2007), 507–525.
- [3] J. H. E. Cohn, *The diophantine equation $(a^n - 1)(b^n - 1) = x^2$* , Periodica Mathematica Hungarica, **44**(2) (2002), 169–175.
- [4] L. Hajdu and L. Szalay, *On the Diophantine Equations $(2^n - 1)(6^n - 1) = x^2$ and $(a^n - 1)(a^{kn} - 1) = x^2$* , Periodica Mathematica Hungarica, **40** (2000), 141–145.
- [5] B. He, C. Liu, *The diophantine equation $(2^k - 1)(3^k - 1) = x^n$* , Periodica Mathematica Hungarica, **91** (2025), 582–587.
- [6] M. H. Le, *A note on the exponential Diophantine equation $(2^n - 1)(b^n - 1) = x^2$* , Publicationes Mathematicae Debrecen **74**(3-4) (2009): 453–455.
- [7] F. Luca and P. G. Walsh, *The product of like-indexed terms in binary recurrences*, Journal of Number Theory, **96**(1) (2002), 152–173.
- [8] A. H. Parvardi, *Lifting the exponent lemma (LTE)*. available at <https://artofproblemsolving.com/community/c6h393335p2198886>.
- [9] L. Szalay, *On the Diophantine equation $(2^n - 1)(3^n - 1) = x^2$* , Publicationes Mathematicae Debrecen, **57**(1-2) (2000), 1–9.

(Chang Liu) MATHEMATISCHES INSTITUT DER UNIVERSITÄT GÖTTINGEN, BUNSENSTR. 3-5, DE-37073, GÖTTINGEN, GERMANY

Email address, Chang Liu: `chang.liu@mathematik.uni-goettingen.de`

(Bo He) 1. MATHEMATISCHES INSTITUT DER UNIVERSITÄT GÖTTINGEN, BUNSENSTR. 3-5, DE-37073, GÖTTINGEN, GERMANY; 2. APPLIED MATHEMATICS INSTITUTE OF ABA TEACHERS UNIVERSITY, WENCHUAN, SICHUAN, 623002, P. R. CHINA

Email address, Bo He: `bo.he@mathematik.uni-goettingen.de`, `bhe@live.cn`