

HYPONORMALITY AND QUASINORMALITY OF UNBOUNDED TOEPLITZ OPERATORS WITH NON-HARMONIC SYMBOL ON THE FOCK-SOBOLEV SPACE

Anuradha Gupta and Kajal Negi

Abstract

In this paper, we establish the essential criteria for the hyponormality and quasinormality of the unbounded Toeplitz operator T_φ with non-harmonic symbol, acting on the Fock-Sobolev space $F^{2,m}(\mathbb{C})$. The study shows that quasinormality does not inherently imply hyponormality of unbounded Toeplitz operator with non-harmonic symbols.

Mathematics Subject Classification: Primary 47B35, Secondary 46E22, 47B20, 47A05.

Keywords: Fock-Sobolev space, Hyponormality, Quasinormality, Toeplitz operator.

1 Introduction

The Fock space denoted by \mathbb{F}^2 , also referred as the Bargmann space, is a Hilbert space consisting of entire functions defined on the complex plane \mathbb{C} and endowed with the standard Lebesgue measure dA . In \mathbb{F}^2 , the norm of a function f belonging to \mathbb{F}^2 , is denoted by $\|f\|_2$. This norm is defined as $\|f\|_2^2 = \frac{1}{\pi} \int |f(z)|^2 e^{-|z|^2} dA(z)$. For a non-negative integer m , let $L_m^2(\mathbb{C})$ be the space of Lebesgue measurable functions f on \mathbb{C} such that $\frac{1}{\pi m!} \int |f(z)|^2 |z|^{2m} e^{-|z|^2} dA(z) < \infty$. The Fock-Sobolev space $F^{2,m}(\mathbb{C})$ comprises entire functions f satisfying $f \in L_m^2(\mathbb{C})$. H.R. Cho and K. Zhu established in their work [2] that a function $f \in F^{2,m}(\mathbb{C}^n)$ if and only if $z^\alpha f(z) \in F^2(\mathbb{C}^n)$ for all multi-indices α with $|\alpha| = m$. The space $F^{2,m}(\mathbb{C})$ is a Hilbert space equipped with the inner product inherited from $L_m^2(\mathbb{C})$, which is defined as follows:

$$\langle f, g \rangle = \frac{1}{\pi m!} \int f(z) \overline{g(z)} |z|^{2m} e^{-|z|^2} dA(z) \text{ where } f, g \in L_m^2(\mathbb{C}).$$

It is well-established that $F^{2,m}(\mathbb{C})$ constitutes a closed subspace of $L_m^2(\mathbb{C})$. The orthogonal projection $P : L_m^2(\mathbb{C}) \rightarrow F^{2,m}(\mathbb{C})$ is defined as

$$Pf(z) = \frac{1}{\pi m!} \int f(w) K_m(z, w) |w|^{2m} e^{-|w|^2} dA(w), \quad f \in L_m^2(\mathbb{C}) \quad (1.1)$$

where $K_m(z, w)$ represents the reproducing kernel function of the Fock-Sobolev space $F^{2,m}(\mathbb{C})$ given by the series expression $K_m(z, w) = \sum_{n=0}^{\infty} \frac{m!}{(n+m)!} (z\bar{w})^n$. In $F^{2,m}(\mathbb{C})$, polynomials form a dense subset.

Additionally, monomials are mutually orthogonal. This implies that the set $\left\{ \frac{z^k}{\sqrt{\langle z^k, z^k \rangle}} \right\}$ forms an

orthonormal basis for $F^{2,m}(\mathbb{C})$. Let φ be a Lebesgue measurable function on \mathbb{C} that satisfies the condition

$$\int |\varphi(w)| |w|^{2m} |K_m(z, w)| e^{-|w|^2} dA(w) < \infty.$$

We can define the unbounded Toeplitz operator T_φ on $F_m^2(\mathbb{C})$ with symbol φ as

$$T_\varphi f(z) = \langle (\varphi f)(w), K_m(w, z) \rangle_m = \frac{1}{\pi m!} \int \varphi(w) f(w) K_m(z, w) |w|^{2m} e^{-|w|^2} dA(w).$$

A linear operator T in a complex Hilbert space \mathcal{H} is called densely defined if its domain $D(T)$ is dense in \mathcal{H} . The adjoint of the operator T is denoted by T^* . If the domain $D(T^*)$ consists of all $y \in \mathcal{H}$ for which there exists a unique $y^* \in \mathcal{H}$ such that $\langle Tx, y \rangle = \langle x, y^* \rangle$ for every $x \in D(T)$ then the adjoint operator $T^* : D(T^*) \rightarrow \mathcal{H}$ is defined by $y^* = T^*y$ for every $y \in D(T^*)$. Let S denote the subset of all polynomials in $F^{2,m}(\mathbb{C})$. Clearly, S forms a dense subspace of $F^{2,m}(\mathbb{C})$. The operator T_φ , with φ as a non-harmonic symbol, is therefore densely defined on S . The hyponormality of operators in the Hardy space has been studied by C.C. Cowen [3], while T. Nakazi and K. Takahashi [10] investigated hyponormality in the same space using a different approach. In the weighted Bergman space, K. Sumin and L. Jongrak [7] examined the necessary or sufficient conditions for hyponormality with non-harmonic symbols. Motivated by their research, we have studied the hyponormality and quasinormality of unbounded Toeplitz operators in the Fock-Sobolev space $F^{2,m}(\mathbb{C})$ with non-harmonic symbols. These operators are essential tools in signal processing, where they are employed for purposes such as noise reduction and image enhancement. Moreover, they hold significant significance in prediction theory, enabling the generation of forecasts based on historical data. Additionally, they are instrumental in wavelet analysis for extracting features and in solving boundary value problems encountered in differential equations. It is generally accepted that any bounded quasinormal operator on a Hilbert space is hyponormal, this conventional implication does not universally hold for unbounded operators. This paper presents the necessary conditions for hyponormality and quasinormality of unbounded Toeplitz operators in the Fock-Sobolev space. We begin with the following known property of Toeplitz operators which is instrumental in the subsequent results:

Proposition 1.1. [11] *If $f, g \in L^\infty(\mathbb{C})$ then the following equivalences hold:*

1. $T_{f+g} = T_f + T_g$.
2. $T_f^* = T_{\bar{f}}$.
3. $T_f T_g = T_{fg}$ if f or g is analytic.

Lemma 1.2. [5] *Let t and s be non-negative integers and $w \in \mathbb{C}$, then*

$$P(\bar{w}^t w^s) = \begin{cases} \frac{(s+m)!}{(s+m-t)!} w^{(s-t)} & \text{for } s \geq t, \\ 0 & \text{for } s < t, \end{cases}$$

where P is orthogonal projection from $L_m^2(\mathbb{C})$ onto $F^{2,m}(\mathbb{C})$.

Remark 1.3. [5] Let $s, t \in \mathbb{N}$ and $z \in \mathbb{C}$

$$\langle z^s, z^t \rangle = \begin{cases} \pi(s+m)!, & \text{if } s = t, \\ 0, & \text{otherwise.} \end{cases} \quad (1.2)$$

2 Necessary conditions for hyponormality

Let T be a densely defined linear operator on a complex Hilbert space \mathcal{H} . The operator T is considered to be hyponormal if $D(T) \subseteq D(T^*)$ and the self-commutator $[T^*, T] := T^*T - TT^*$, is positive. In this section, the necessary conditions for the hyponormality of the unbounded Toeplitz operator T_φ on the subspace S of $F^{2,m}$, where φ is a non-harmonic symbol, are being examined.

Theorem 2.1. *Let $\varphi(z) = az^p\bar{z}^n + bz^s\bar{z}^t$, where $p, n, s, t \in \mathbb{N}$, $z, a, b \in \mathbb{C}$, $p \geq n$, $t \geq s$ and $|p - n| \neq |t - s|$. The Toeplitz operator T_φ is hyponormal then*

1. For $t > s + 1$, $p = n + 1$

$$|a|^2 \left(\frac{(p+k+m)!^2}{(k+m+1)!} - \frac{(p+k+m-1)!^2}{(k+m-1)!} \right) \geq |b|^2 \left(\frac{(t+k+m)!^2}{(t+k+m-s)!} - \frac{(s+k+m)!^2}{(s+k+m-t)!} \right).$$

2. For $t > s + 1$, $p > n + 1$

$$|a|^2 \left(\frac{(p+k+m)!^2}{(p+k+m-n)!} - \frac{(n+k+m)!^2}{(n+k+m-p)!} \right) \geq |b|^2 \left(\frac{(t+k+m)!^2}{(t+k+m-s)!} - \frac{(s+k+m)!^2}{(s+k+m-t)!} \right).$$

3. For $t = s + 1$, $p > n + 1$

$$(p+k+m-1)!^2((p+k+m)^2 - 1) \geq (s+k+m)!^2((s+k+m+1)^2 - (k+m+1)(k+m)).$$

Proof. Let $p, n, s, t \in \mathbb{N}$, $z, a, b \in \mathbb{C}$ with $p \geq n$, $t \geq s$, $|p - n| \neq |t - s|$ and $\varphi(z) = az^p\bar{z}^n + bz^s\bar{z}^t$, let the Toeplitz operator T_φ be hyponormal then for $k \in \mathbb{N}$, $k > p, t$ and $z^k \in S$

$$\|T_\varphi(z^k)\|^2 - \|T_\varphi^*(z^k)\|^2 \geq 0. \quad (2.1)$$

By using Lemma 1.2

$$\begin{aligned} T_\varphi(z^k) &= P(az^p\bar{z}^n + bz^s\bar{z}^t)(z^k) \\ &= aP(z^{p+k}\bar{z}^n) + bP(z^{s+k}\bar{z}^t) \\ &= a \frac{(p+k+m)!}{(p+k+m-n)!} z^{p+k-n} + b \frac{(s+k+m)!}{(s+k+m-t)!} z^{s+k-t}. \end{aligned} \quad (2.2)$$

Using $|p - n| \neq |t - s|$, 2.2 and 1.2.

$$\begin{aligned} \langle T_\varphi z^k, T_\varphi z^k \rangle &= \left\langle a \frac{(p+k+m)!}{(p+k+m-n)!} z^{p+k-n} + b \frac{(s+k+m)!}{(s+k+m-t)!} z^{s+k-t}, \right. \\ &\quad \left. a \frac{(p+k+m)!}{(p+k+m-n)!} z^{p+k-n} + b \frac{(s+k+m)!}{(s+k+m-t)!} z^{s+k-t} \right\rangle \\ &= \pi \left(|a|^2 \frac{(p+k+m)!^2}{(p+k+m-n)!} + |b|^2 \frac{(s+k+m)!^2}{(s+k+m-t)!} \right). \end{aligned} \quad (2.3)$$

Consider

$$\begin{aligned} T_\varphi^*(z^k) &= P(\bar{a}\bar{z}^p z^{n+k} + \bar{b}\bar{z}^s z^{t+k}) \\ &= \bar{a}P(z^{n+k}\bar{z}^p) + \bar{b}P(z^{t+k}\bar{z}^s) \\ &= \bar{a} \frac{(n+k+m)!}{(n+k+m-p)!} z^{n+k-p} + \bar{b} \frac{(t+k+m)!}{(t+k+m-s)!} z^{t+k-s}. \end{aligned} \quad (2.4)$$

Using 2.4 and 1.2

$$\begin{aligned}
\langle T_\varphi^* z^k, T_\varphi^* z^k \rangle &= \langle \bar{a} \frac{(n+k+m)!}{(n+k+m-p)!} z^{n+k-p} + \bar{b} \frac{(t+k+m)!}{(t+k+m-s)!} z^{t+k-s}, \bar{a} \frac{(n+k+m)!}{(n+k+m-p)!} z^{n+k-p} \\
&\quad + \bar{b} \frac{(t+k+m)!}{(t+k+m-s)!} z^{t+k-s} \rangle \\
&= \pi \left(|a|^2 \frac{(n+k+m)!^2}{(n+k+m-p)!} + |b|^2 \frac{(t+k+m)!^2}{(t+k+m-s)!} \right). \tag{2.5}
\end{aligned}$$

On putting 2.3 and 2.5 in 2.1.

$$\pi \left(|a|^2 \left(\frac{(p+k+m)!^2}{(p+k+m-n)!} - \frac{(n+k+m)!^2}{(n+k+m-p)!} \right) + |b|^2 \left(\frac{(s+k+m)!^2}{(s+k+m-t)!} - \frac{(t+k+m)!^2}{(t+k+m-s)!} \right) \right) \geq 0. \tag{2.6}$$

Case 1. If $p > n + 1$, $t = s + 1$. Let $\mathbb{T} = \{z \in \mathbb{C} \mid |z| = 1\}$ and $a, b \in \mathbb{T}$ then 2.6 becomes

$$\frac{(p+k+m)!^2}{(p+k+m-n)!} + \frac{(s+k+m)!^2}{(k+m-1)!} \geq \frac{(n+k+m)!^2}{(n+k+m-p)!} + \frac{(t+k+m)!^2}{(k+m+1)!} \tag{2.7}$$

As $p > n$ and $(p+k+m-n)! \geq (k+m+1)!$ and $(k+m+1)! \geq (n+k+m-p)!$. Therefore,

$$\frac{(p+k+m)!^2}{(k+m+1)!} \geq \frac{(p+k+m)!^2}{(p+k+m-n)!} \text{ and } \frac{(n+k+m)!^2}{(k+m+1)!} \leq \frac{(n+k+m)!^2}{(n+k+m-p)!}.$$

$$\text{Therefore, } \frac{(p+k+m)!^2}{(k+m+1)!} - \frac{(n+k+m)!^2}{(k+m+1)!} \geq \frac{(p+k+m)!^2}{(p+k+m-n)!} - \frac{(n+k+m)!^2}{(n+k+m-p)!}$$

Therefore, 2.7 becomes,

$$\frac{(p+k+m)!^2}{(k+m+1)!} - \frac{(n+k+m)!^2}{(k+m+1)!} + \frac{(s+k+m)!^2(k+m+1)(k+m)}{(k+m+1)!} - \frac{(t+k+m)!^2}{(k+m+1)!} \geq 0 \tag{2.8}$$

$$\implies (p+k+m)!^2 - (n+k+m)!^2 + (s+k+m)!^2(k+m+1)(k+m) - (t+k+m)!^2 \geq 0. \tag{2.9}$$

As $p > n$ therefore $-(p+k+m+1)!^2 < -(n+k+m)!^2$.

$$\implies (p+k+m)!^2 + (s+k+m)!^2(k+m+1)(k+m) \geq (p+k+m-1)!^2 + (s+k+m+1)!^2.$$

$$\implies (p+k+m-1)!^2((p+k+m)^2 - 1) \geq (s+k+m)!^2((s+k+m+1)^2 - (k+m+1)(k+m)).$$

Case 2: If $t > s + 1$, $p = n + 1$ then from 2.6

$$|a|^2 \left(\frac{(p+k+m)!^2}{(k+m+1)!} - \frac{(p+k+m-1)!^2}{(k+m-1)!} \right) \geq |b|^2 \left(\frac{(t+k+m)!^2}{(t+k+m-s)!} - \frac{(s+k+m)!^2}{(s+k+m-t)!} \right). \tag{2.10}$$

Case 3: If $t > s + 1$, $p > n + 1$ then from 2.6

$$|a|^2 \left(\frac{(p+k+m)!^2}{(p+k+m-n)!} - \frac{(n+k+m)!^2}{(n+k+m-p)!} \right) \geq |b|^2 \left(\frac{(t+k+m)!^2}{(t+k+m-s)!} - \frac{(s+k+m)!^2}{(s+k+m-t)!} \right). \quad (2.11)$$

□

The above conditions are only necessary but not sufficient which is shown in the following example:

Example 2.2. Let $\varphi(z) = z\bar{z}^3 + z^2\bar{z}$ where $z \in \mathbb{C}$ and let $m = 1$ as it satisfies given condition (1) in Theorem 2.1. Using Proposition 1.1, Lemma 1.2 and $z - z^4 \in S$, $T_\varphi(z - z^4) = -116z^2 - 7z^5$, $T_\varphi^*(z - z^4) = 6 - 25z^3 - 8z^6$, $\langle T_\varphi^*T_\varphi(z - z^4), (z - z^4) \rangle = 116016\pi$ and $\langle T_\varphi T_\varphi^*(z - z^4), (z - z^4) \rangle = 337596\pi$. Hence, $(\langle T_\varphi^*T_\varphi - T_\varphi T_\varphi^* \rangle(z - z^4), (z - z^4)) = -221580 \pi < 0$. Thus, T_φ is not hyponormal.

Example 2.3. Let $\varphi(z) = 4z^3\bar{z} + 6z\bar{z}^4$ where $z \in \mathbb{C}$ and let $m = 1$ as it satisfies given condition (2) in Theorem 2.1. Using Proposition 1.1, Lemma 1.2 and $z - z^4 \in S$, $T_\varphi(z - z^4) = -2160z + 20z^3 - 32z^6$, $T_\varphi^*(z - z^4) = -480z^2 + 36z^4 - 54z^7$, $\langle T_\varphi^*T_\varphi(z - z^4), (z - z^4) \rangle = 14501760\pi$ and $\langle T_\varphi T_\varphi^*(z - z^4), (z - z^4) \rangle = 119111040\pi$. Hence, $(\langle T_\varphi^*T_\varphi - T_\varphi T_\varphi^* \rangle(z - z^4), (z - z^4)) = -104609280 \pi < 0$. Thus, T_φ is not hyponormal.

Example 2.4. Let $\varphi(z) = 4z^3\bar{z} + 6z^2\bar{z}^3$ where $z \in \mathbb{C}$ and let $m = 1$ as it satisfies given condition (3) in Theorem 2.1. Using Proposition 1.1 Lemma 1.2 and $z - z^4 \in S$, $T_\varphi(z - z^4) = 144 - 1240z^3 - 32z^6$, $T_\varphi^*(z - z^4) = -360z^2 - 336z^5$, $\langle T_\varphi^*T_\varphi(z - z^4), (z - z^4) \rangle = 43293696\pi$ and $\langle T_\varphi T_\varphi^*(z - z^4), (z - z^4) \rangle = 82753920\pi$. Hence, $(\langle T_\varphi^*T_\varphi - T_\varphi T_\varphi^* \rangle(z - z^4), (z - z^4)) = -39460224 \pi < 0$. Thus, T_φ is not hyponormal.

If we put $p - n = 1 = t - s$ and $k = 1$ in 2.6 then the following remark follows :-

Remark 2.5. Let $\varphi(z) = az^p\bar{z}^n + bz^s\bar{z}^t$, where $p, n, s, t \in \mathbb{N}$, $a, b \in \mathbb{C}$, $p \geq n \geq 0$, $t \geq s \geq 0$ and $p - n = 1 = t - s$. Then, T_φ is hyponormal if

$$|a|^2(p+m)!^2((1+p+m)^2 - (1+m)(2+m)) \geq |b|^2(t+m)!^2((1+t+m)^2 - (1+m)(2+m)).$$

Remark 2.6. Let T and S be operators on complex Hilbert space, we will be using next formula from [4] to check hyponormality

$$\begin{aligned} \langle ((T+S)^*, T+S)u, u \rangle &= \langle Tu, Tu \rangle - \langle T^*u, T^*u \rangle \\ &\quad + 2\operatorname{Re}(\langle Tu, Su \rangle - \langle T^*u, S^*u \rangle) + \langle Su, Su \rangle - \langle S^*u, S^*u \rangle. \end{aligned} \quad (2.12)$$

Theorem 2.7. Suppose $\mathbf{C} \in \mathbb{C}$, $s \in \mathbb{N}$ and $n \in \mathbb{N} \cup \{0\}$ such that $n \leq N$. If $T_{z^n + C|z|^{2s}}$ is hyponormal operator then

$$\begin{aligned} \pi \left(\sum_{k=n}^N |a_k|^2 \left((k+n+m)! - \frac{((k+m)!)^2}{(k+m-n)!} \right) + \right. \\ \left. 2 \sum_{k=0}^N \operatorname{Re}(Ca_{k+n}\bar{a}_k) \frac{(k+m+n+s)!(k+m)! - (k+m+n)!(k+m+s)!}{(k+m)!} \right) \geq 0. \end{aligned}$$

Proof. Let the operator T_φ be hyponormal where the symbol $\varphi(z) = az^n + C|z|^{2s}$ then for $f(z) = \sum_{k=0}^N a_k z^k \in S$. The Toeplitz operator T_φ satisfies 2.12.

$$\text{Consider } T_{z^n}(f(z)) = P(z^n) \left(\sum_{k=0}^N a_k z^k \right) = \sum_{k=0}^N a_k z^{k+n}. \quad (2.13)$$

$$\text{By using 1.2, } \langle T_{z^n}(f(z)), T_{z^n}(f(z)) \rangle = \pi \sum_{k=0}^N |a_k|^2 (k+n+m)!. \quad (2.14)$$

$$\text{Consider } T_{z^n}^*(f(z)) = P(\bar{z}^n) \left(\sum_{k=0}^N (a_k z^k) \right) = \sum_{k=0}^N a_k P(z^k \bar{z}^n) = \sum_{k=n}^N a_k \frac{(k+m)!}{(k+m-n)!} z^{k-n}. \quad (2.15)$$

From 1.2

$$\langle T_{z^n}^*(f(z)), T_{z^n}^*(f(z)) \rangle = \sum_{k=n}^N |a_k|^2 \frac{(k+m)!^2}{(k+m-n)!^2} \langle z^{k-n}, z^{k-n} \rangle = \pi \sum_{k=n}^N |a_k|^2 \frac{(k+m)!^2}{(k+m-n)!}. \quad (2.16)$$

$$\begin{aligned} \text{Using Lemma 1.2. } T_{C|z|^{2s}}(f(z)) &= P(C|z|^{2s} \sum_{k=0}^N a_k z^k) = C \sum_{k=0}^N a_k P(|z|^{2s} z^k) \\ &= C \sum_{k=0}^N a_k \frac{(k+s+m)!}{(k+m)!} z^k. \end{aligned} \quad (2.17)$$

Using 1.2, we have

$$\begin{aligned} \langle T_{z^n}(f(z)), T_{C|z|^{2s}}(f(z)) \rangle &= \left\langle \sum_{k=0}^N a_k z^{k+n}, C \sum_{k=0}^N a_k \frac{(k+s+m)!}{(k+m)!} z^k \right\rangle \\ &= \pi \bar{C} \sum_{k=0}^N a_k \overline{a_{k+n}} (k+m+n+s)! \end{aligned} \quad (2.18)$$

$$\begin{aligned} \text{and } \langle T_{\bar{z}^n}(f(z)), T_{\bar{C}|z|^{2s}}(f(z)) \rangle &= C \left\langle \sum_{k=n}^N a_k \frac{(k+m)!}{(k+m-n)!} z^{k-n}, \sum_{k=0}^N a_k \frac{(k+m+s)!}{(k+m)!} z^k \right\rangle \\ &= \pi C \sum_{k=0}^N \bar{a}_k a_{k+n} \frac{(k+m+n)!(k+m+s)!}{(k+m)!}. \end{aligned} \quad (2.19)$$

Using 2.18 and 2.19

$$\begin{aligned} &\text{Re} \left(\langle T_{z^n}(f(z)), T_{C|z|^{2s}}(f(z)) \rangle - \langle T_{\bar{z}^n}(f(z)), T_{\bar{C}|z|^{2s}}(f(z)) \rangle \right) \\ &= \pi \sum_{k=0}^N \text{Re}(C a_{k+n} \bar{a}_k) \frac{(k+m+n+s)!(k+m)! - (k+m+n)!(k+m+s)!}{(k+m)!}. \end{aligned} \quad (2.20)$$

If the operator T_φ is hyponormal then from 2.12

$$\begin{aligned}
& \pi \left(\sum_{k=0}^N |a_k|^2 (k+n+m)! - \sum_{k=n}^N |a_k|^2 \frac{((k+m)!)^2}{(k+m-n)!} + \right. \\
& \quad \left. 2 \sum_{k=0}^N \operatorname{Re}(C a_{k+n} \overline{a_k}) \frac{(k+m+n+s)!(k+m)! - (k+m+n)!(k+m+s)!}{(k+m)!} \right) \geq 0. \\
& = \pi \left(\sum_{k=0}^{n-1} |a_k|^2 (k+n+m)! + \sum_{k=n}^N |a_k|^2 \left((k+m+n)! - \frac{((k+m)!)^2}{(k+m-n)!} \right) + \right. \\
& \quad \left. 2 \sum_{k=0}^N \operatorname{Re}(C a_{k+n} \overline{a_k}) \frac{(k+m+n+s)!(k+m)! - (k+m+n)!(k+m+s)!}{(k+m)!} \right) \geq 0.
\end{aligned} \tag{2.21}$$

Hence, if the operator T_φ where $\varphi(z) = z^n + C|z|^{2s}$ is hyponormal then,

$$\begin{aligned}
& \pi \left(\sum_{k=n}^N |a_k|^2 \left((k+n+m)! - \frac{((k+m)!)^2}{(k+m-n)!} \right) + \right. \\
& \quad \left. 2 \sum_{k=0}^N \operatorname{Re}(C a_{k+n} \overline{a_k}) \frac{(k+m+n+s)!(k+m)! - (k+m+n)!(k+m+s)!}{(k+m)!} \right) \geq 0.
\end{aligned} \tag{2.22}$$

□

Remark 2.8. Let $f(z) = a_0 + a_1 z \in S$ and $\phi(z) = z^n + C|z|^{2s}$ where $z, a_0, a_1 \in \mathbb{C}$ and $n, s \in \mathbb{N}$. If the operator T_φ is hyponormal then for $n > 1$ 2.21 reduces to

$$\pi \left(|a_0|^2 (n+m)! + |a_1|^2 (1+n+m)! \right) \geq 0.$$

If $n = 0$ then 2.21 becomes $\pi \left(|a_0|^2 (m)! \right)$ which is trivially positive. Now, consider $n = 1$ in 2.21, thus

$$\begin{aligned}
& \pi \left(|a_0|^2 (1+m)! + |a_1|^2 (1+m)! + 2 \operatorname{Re}(C \overline{a_0} a_1) s(m+s)! \right) \geq 0. \\
& \implies \left(|a_0|^2 (1+m)! + |a_1|^2 (1+m)! + 2|C||a_0||a_1|s(m+s)! \right) \geq 0.
\end{aligned}$$

If $a_0 \neq 0 \neq a_1$ then,

$$\left(\left| \frac{a_0}{a_1} \right|^2 (1+m)! + (1+m)! + 2|C| \left| \frac{a_0}{a_1} \right| s(m+s)! \right) \geq 0.$$

The above inequality is a quadratic inequality in terms of $\left| \frac{a_1}{a_2} \right|$ and takes only non-negative values, so its discriminant will be non-positive. Therefore, $|C|^2 \leq \frac{(1+m)!^2}{s^2(m+s)!^2}$. As $(1+m)!^2 \leq s^2(m+s)!^2$, we conclude that $|C| \leq 1$.

3 Necessary conditions for quasinormality

A densely defined linear operator T on a Hilbert space \mathcal{H} is termed as quasinormal if $D(T) \subseteq D(T^*)$ and satisfies the relationship $T(T^*T) = (T^*T)T$ which implies that T commutes with T^*T . In this section, we explore the necessary conditions for the quasinormality of unbounded Toeplitz operators T_φ on the subspace S of the Fock–Sobolev space, where φ is a non-harmonic symbol.

Theorem 3.1. *Let $\varphi(z) = az^p\bar{z}^n + bz^s\bar{z}^t$, $z, a, b \in \mathbb{C}$ with $p, n, s, t \in \mathbb{N}$ and $p \geq n$, $t \geq s$. If the Toeplitz operator T_φ is quasinormal then*

1. *If $p \neq n$ and $s \neq t$ then the operator T_φ is quasinormal.*

2. *If $p = n$ and $s \neq t$ then*

$$\frac{(p+k+m)!(t+k+m)!^2}{(k+m)!(t+k+m+s)!} = \frac{(s+k+m)!^2(s+k+m-p-t)!}{(s+k+m-t)!^2}.$$

3. *If $p \neq n$ and $s = t$ then*

$$\frac{(s+k+m)!(k+m+n)!^2}{(k+m)!(k+n+m-p)!} = \frac{(p+k+m)!^2(p+k+m+s-n)!}{(p+k+m-n)!^2}.$$

Proof. Let the operator T_φ is quasinormal where the symbol $\varphi(z) = az^p\bar{z}^n + bz^s\bar{z}^t$. Then for $k \in \mathbb{N}$, $k > 2p, 2t$ and $z^k \in S$.

$$\langle (T_\varphi^*T_\varphi^2 - T_\varphi T_\varphi^*T_\varphi)(z^k), (z^k) \rangle = 0. \quad (3.1)$$

Consider

$$T_\varphi(z^k) = P(az^p\bar{z}^n + bz^s\bar{z}^t)(z^k) = a \frac{(p+k+m)!}{(p+k+m-n)!} z^{p+k-n} + b \frac{(s+k+m)!}{(s+k+m-t)!} z^{s+k-t} \quad (3.2)$$

$$\text{and } T_\varphi^*(z^k) = P(\bar{a}\bar{z}^p z^n + \bar{b}\bar{z}^s z^t)(z^k) = \bar{a} \frac{(n+k+m)!}{(n+k+m-p)!} z^{n+k-p} + \bar{b} \frac{(t+k+m)!}{(t+k+m-s)!} z^{t+k-s}. \quad (3.3)$$

Using 3.2 and Lemma 1.2 we get

$$\begin{aligned} T_\varphi^2(z^k) &= T_\varphi(T_\varphi(z^k)) \\ &= P(az^p\bar{z}^n + bz^s\bar{z}^t) \left(a \frac{(p+k+m)!}{(p+k+m-n)!} z^{p+k-n} + b \frac{(s+k+m)!}{(s+k+m-t)!} z^{s+k-t} \right) \\ &= a^2 \frac{(p+k+m)!(2p+k+m-n)!}{(p+k+m-n)!(2p+k+m-2n)!} z^{2p+k-2n} \\ &\quad + ab \left(\frac{(s+k+m)!(s+k+m+p-t)!}{(s+k+m-t)!(s+k+m+p-t-n)!} z^{s+k+p-t-n} \right. \\ &\quad \left. + \frac{(p+k+m)!(p+k+m+s-n)!}{(p+k+m-n)!(p+k+m+s-n-t)!} z^{p+k+s-n-t} \right) \\ &\quad + b^2 \frac{(s+k+m)!(2s+k+m-t)!}{(s+k+m-t)!(2s+k+m-2t)!} z^{2s+k-2t} \end{aligned} \quad (3.4)$$

$$\begin{aligned}
\text{and } T_\varphi^* T_\varphi(z^k) &= P(\bar{a}z^p z^n + \bar{b}\bar{z}^s z^t) \left(a \frac{(p+k+m)!}{(p+k+m-n)!} z^{p+k-n} + b \frac{(s+k+m)!}{(s+k+m-t)!} z^{s+k-t} \right) \\
&= |a|^2 \frac{(p+k+m)!^2}{(p+k+m-n)!(k+m)!} z^k + a\bar{b} \frac{(p+k+m)!(p+k+m+t-n)!}{(p+k+m-n)!(p+k+m+t-n-s)!} \\
&\quad z^{p+k+t-n-s} + \bar{a}b \frac{(s+k+m)!(s+k+m+n-t)!}{(s+k+m-t)!(s+k+m+n-t-p)!} z^{s+k+n-t-p} \\
&\quad + |b|^2 \frac{(s+k+m)!^2}{(s+k+m-t)!(k+m)!} z^k.
\end{aligned} \tag{3.5}$$

Using Lemma 1.2 and 3.3 , 3.5.

$$\begin{aligned}
\langle T_\varphi^* T_\varphi z^k, T_\varphi^* z^k \rangle &= \left\langle |a|^2 \frac{(p+k+m)!^2}{(p+k+m-n)!(k+m)!} z^k + a\bar{b} \frac{(p+k+m)!(p+k+m+t-n)!}{(p+k+m-n)!(p+k+m+t-n-s)!} \right. \\
&\quad z^{p+k+t-n-s} + \bar{a}b \frac{(s+k+m)!(s+k+m+n-t)!}{(s+k+m-t)!(s+k+m+n-t-p)!} z^{s+k+n-t-p} \\
&\quad + |b|^2 \frac{(s+k+m)!^2}{(s+k+m-t)!(k+m)!} z^k, \bar{a} \frac{(k+n+m)!}{(k+n+m-p)!} z^{k+n-p} \\
&\quad \left. + \bar{b} \frac{(t+k+m)!}{(t+k+m-s)!} z^{t+k-s} \right\rangle
\end{aligned} \tag{3.6}$$

Case 1. If $p = n$ and $s = t$ then 3.6 becomes

$$\begin{aligned}
&\pi \left(a|a|^2 \frac{(p+k+m)!^3}{(k+m)!^2} + a^2 \bar{b} \frac{(p+k+m)!^2 (s+k+m)!}{(k+m)!^2} + 2|a|^2 b \frac{(p+k+m)!^2 (s+k+m)!}{(k+m)!^2} \right. \\
&\quad + |b|^2 a \frac{(s+k+m)!^2 (p+k+m)!}{(k+m)!^2} + a|b|^2 \frac{(p+k+m)!(s+k+m)!^2}{(k+m)!^2} \\
&\quad \left. + \bar{a}b^2 \frac{(s+k+m)!^2 (p+k+m)!}{(k+m)!^2} + |b|^2 b \frac{(s+k+m)!^3}{(k+m)!^2} \right).
\end{aligned} \tag{3.7}$$

Case 2. If $p = n$ and $s \neq t$ then 3.6 becomes

$$\pi \left(a|a|^2 \frac{(p+k+m)!^3}{(k+m)!^2} + a|b|^2 \frac{(p+k+m)!(s+k+m)!^2}{(k+m)!(s+k+m-t)!} + a|b|^2 \frac{(p+k+m)!(t+k+m)!^2}{(k+m)!(t+k+m-s)!} \right). \tag{3.8}$$

Case 3. If $p \neq n$ and $s = t$ then 3.6 becomes

$$\pi \left(|a|^2 b \frac{(s+k+m)!(k+m+n)!^2}{(k+m)!(k+m+n-p)!} + |a|^2 b \frac{(p+k+m)!^2 (s+k+m)!}{(p+k+m-n)!(k+m)!} + |b|^2 b \frac{(s+k+m)!^3}{(k+m)!^2} \right). \tag{3.9}$$

Case 4. If $p \neq n$ and $s \neq t$ then 3.6 becomes

$$\langle T_\varphi^* T_\varphi z^k, T_\varphi^* z^k \rangle = 0. \quad (3.10)$$

Consider,

$$\begin{aligned} & \langle T_\varphi^* T_\varphi^2 z^k, z^k \rangle \\ &= \left\langle a^2 \frac{(p+k+m)!(2p+k+m-n)!}{(p+k+m-n)!(2p+k+m-2n)!} z^{2p+k-2n} + ab \left(\frac{(s+k+m)!(s+k+m+p-t)!}{(s+k+m-t)!(s+k+m+p-t-n)!} \right. \right. \\ & \quad \left. \left. z^{s+k+p-t-n} + \frac{(p+k+m)!(p+k+m+s-n)!}{(p+k+m-n)!(p+k+m+s-n-t)!} z^{p+k+s-n-t} \right) \right. \\ & \quad \left. + b^2 \frac{(s+k+m)!(2s+k+m-t)!}{(s+k+m-t)!(2s+k+m-2t)!} z^{2s+k-2t}, a \frac{(p+k+m)!}{(p+k+m-n)!} z^{p+k-n} + b \frac{(s+k+m)!}{(s+k+m-t)!} z^{s+k-t} \right\rangle. \end{aligned} \quad (3.11)$$

Case i. If $p = n$ and $s = t$ then 3.11 becomes

$$\begin{aligned} & \pi \left(\bar{a}b^2 \frac{(s+k+m)!^2(p+k+m)!}{(k+m)!^2} + b|b|^2 \frac{(s+k+m)!^3}{(k+m)!^2} + |a|^2 a \frac{(p+k+m)!^3}{(k+m)!^2} \right. \\ & \quad + a^2 \bar{b} \frac{(p+k+m)!^2(s+k+m)!}{(k+m)!^2} + 2|a|^2 b \frac{(s+k+m)!(p+k+m)!^2}{(k+m)!^2} \\ & \quad + a|b|^2 \frac{(s+k+m)!^2(p+k+m)!}{(k+m)!^2} \\ & \quad \left. + a|b|^2 \frac{(p+k+m)!(s+k+m)!^2}{(k+m)!^2} \right). \end{aligned} \quad (3.12)$$

Case ii. If $p = n$ and $s \neq t$ then 3.11 becomes

$$\pi \left(|a|^2 a \frac{(p+k+m)!^3}{(k+m)!^2} + a|b|^2 \frac{(s+k+m)!^2(s+k+m+p-t)!}{(s+k+m-t)!^2} + a|b|^2 \frac{(s+k+m)!^2(p+k+m)!}{(k+m)!(k+s+m-t)!} \right). \quad (3.13)$$

Case iii. If $p \neq n$ and $s = t$ then 3.11 becomes

$$\begin{aligned} & \pi \left(b|a|^2 \frac{(s+k+m)!(p+k+m)!^2}{(k+m)!(p+k+m-n)!} + |a|^2 b \frac{(p+k+m)!^2(p+k+m+s-n)!}{(p+k+m-n)!^2} \right. \\ & \quad \left. + |b|^2 b \frac{(s+k+m)!^3}{(k+m)!^2} \right). \end{aligned} \quad (3.14)$$

Case iv. If $p \neq n$ and $s \neq t$ then 3.11 becomes

$$\langle T_\varphi^* T_\varphi^2 z^k, z^k \rangle = 0. \quad (3.15)$$

As the operator T_φ is quasinormal where the symbol $\varphi(z) = az^p\bar{z}^n + bz^s\bar{z}^t$, therefore, on comparing the corresponding cases, we get,

Case a. If $p = n$ and $s = t$ then from 3.6 and 3.11

$$\langle T_\varphi^* T_\varphi^2 z^k, z^k \rangle = \langle T_\varphi T_\varphi^* T_\varphi z^k, z^k \rangle. \quad (3.16)$$

Therefore, the operator T_φ is quasinormal.

Case b. If $p = n$ and $s \neq t$ then from 3.8 and 3.13,

$$a|b|^2 \left(\frac{(p+k+m)!(t+k+m)!^2}{(k+m)!(t+k+m-s)!} - \frac{(s+k+m)!^2(s+k+m+p-t)!}{(s+k+m-t)!^2} \right) = 0.$$

If $a \neq 0 \neq b$ then,

$$\frac{(p+k+m)!(t+k+m)!^2}{(k+m)!(t+k+m-s)!} = \frac{(s+k+m)!^2(s+k+m-p-t)!}{(s+k+m-t)!}.$$

Case c. If $p \neq n$ and $s = t$ then from 3.9 and 3.14,

$$|a|^2 b \left(\frac{(s+k+m)!(k+m+n)!^2}{(k+m)!(k+n+m-p)!} - \frac{(p+k+m)!^2(p+k+m+s-n)!}{(p+k+m-n)!^2} \right) = 0.$$

If $a \neq 0 \neq b$ then

$$\frac{(s+k+m)!(k+m+n)!^2}{(k+m)!(k+n+m-p)!} = \frac{(p+k+m)!^2(p+k+m+s-n)!}{(p+k+m-n)!^2}$$

Case d. If $p \neq n$ and $s \neq t$ then from 3.10 and 3.15 the operator T_φ is quasinormal. \square

The conditions presented in Theorem 3.1 are necessary but not sufficient. The following example verifies the same:

Example 3.2. Let $\varphi(z) = 4\bar{z}^3 z^2 + 6z^3 \bar{z}$ where $z \in \mathbb{C}$ and let $m = 1$ as it satisfies given condition (1) in Theorem 3.1. From example 2.2 T_φ is not hyponormal. To check for quasinormality T_φ must satisfy, 3.1. Consider $T_\varphi T_\varphi(z - z^4) = 3840z + 417600z^2 + 576z^4 - 60228z^5 - 2880z^8$, $T_\varphi^* T_\varphi(z - z^4) = 2304 + 303300z + 1920z^3 + 49392z^4 - 17280z^7$, $\langle T_\varphi^* T_\varphi(z - z^4), T_\varphi^*(z - z^4) \rangle = 0$ and $\langle T_\varphi^2(z - z^4), T_\varphi(z - z^4) \rangle = 5529600\pi$. Hence, $\langle (T_\varphi^* T_\varphi^2 - T_\varphi T_\varphi^* T_\varphi)(z - z^4), (z - z^4) \rangle = 5529600\pi \neq 0$. Thus T_φ is not quasinormal.

Remark 3.3. In general, it is known that every bounded quasinormal operator on a Hilbert space is hyponormal, but in case of the unbounded Toeplitz operator, the conventional implication does not hold. The following example verifies the same:

Example 3.4. Let $\varphi(z) = 2z^3 + 2z^3\bar{z} + \bar{z}^3 + 3z\bar{z}^3$ where $z \in \mathbb{C}$ and $m = 1$. Then by Lemma 1.2 $T_\varphi(z) = 2z^4 + 10z^3$, $T_\varphi^*(z) = z^4 + 15z^3$, $T_\varphi^2(z) = T_\varphi(2z^4 + 10z^3) = 4z^7 + 52z^6 + 140z^5 + 720z^2 + 1920z + 240$ and $T_\varphi^* T_\varphi(z) = 2z^7 + 68z^6 + 210z^5 + 360z^2 + 1440z + 480$.

And by 1.2 $\langle T_\varphi^* T_\varphi^2 z, z \rangle = 0$ and $\langle T_\varphi T_\varphi^* T_\varphi z, z \rangle = 0$. Therefore, $\langle (T_\varphi^* T_\varphi^2 - T_\varphi T_\varphi^* T_\varphi)z, z \rangle = 0$. Therefore, the operator T_φ is quasinormal. For being hyponormal T_φ should satisfy, $\langle (T_\varphi^* T_\varphi - T_\varphi T_\varphi^*)z, z \rangle \geq 0$. By using 1.2 $\langle T_\varphi^* T_\varphi z, z \rangle = 2880\pi$ and $\langle T_\varphi T_\varphi^* z, z \rangle = 5520\pi$. Therefore, $\langle (T_\varphi^* T_\varphi - T_\varphi T_\varphi^*)z, z \rangle = -2640\pi < 0$. Hence, the operator T_φ is quasinormal but not hyponormal.

Acknowledgments

The second author is thankful to the Council of Scientific and Industrial Research (CSIR) [grant number 09/0045(13796)/2022 – I] for their support in the form of a grant-in-aid for this research.

Author contributions

All authors contributed equally to the manuscript and read and approved the final manuscript.

Financial disclosure

None reported.

Conflict of interest

The authors declare no potential conflict of interests.

References

- [1] Amita Aggarwal and Anuradha Gupta. Hyponormality of quasi-homogeneous Toeplitz operators on the Fock space. *J. Math. Anal. Appl.*, 542(1):Paper No. 128842, 13, 2025.
- [2] Hong Rae Cho and Kehe Zhu. Fock-Sobolev spaces and their Carleson measures. *J. Funct. Anal.*, 263(8):2483–2506, 2012.
- [3] Carl Claudius Cowen. Hyponormality of Toeplitz operators. *Proc. Amer. Math. Soc.*, 103(3):809–812, 1988.
- [4] Matthew Fleeman and Constanze Liaw. Hyponormal Toeplitz operators with non-harmonic symbol acting on the Bergman space. *arXiv preprint arXiv:1707.09677*, 2017.
- [5] A. Gupta and K. Negi. Hyponormality and quasinormality of unbounded toeplitz operators on the fock-sobolev space. *Complex Analysis and Operator Theory*, 19:78, 2025.
- [6] J. Janas. On unbounded hyponormal operators iii. *Studia Mathematica*, 112(1):75–82, 1994.
- [7] Sumin Kim and Jongrak Lee. Hyponormal Toeplitz operators with non-harmonic symbols on the weighted Bergman spaces. *Ann. Funct. Anal.*, 14(1):Paper No. 14, 14, 2023.
- [8] Sumin Kim and Jongrak Lee. Remarks on hyponormal Toeplitz operators with nonharmonic symbols. *Open Math.*, 21(1):Paper No. 20230114, 8, 2023.
- [9] Jongrak Lee. Normal Toeplitz operators on the Fock spaces. *Symmetry* 12(10), page 1615, 2020.
- [10] Takahiko Nakazi and Katsutoshi Takahashi. Hyponormal Toeplitz operators and extremal problems of Hardy spaces. *Trans. Amer. Math. Soc.*, 338(2):753–767, 1993.
- [11] Houcine Sadraoui. *Hyponormality of Toeplitz operators and composition operators*. ProQuest LLC, Ann Arbor, MI, 1992. Thesis (Ph.D.)–Purdue University.

[12] Noriaki Setô. Bargmann's inequalities in spaces of arbitrary dimension. *Publ. Res. Inst. Math. Sci.*, 9:429–461, 1973/74.

[13] Kehe Zhu. *Analysis on Fock spaces*, volume 263. Springer Science & Business Media, 2012.

Anuradha Gupta

Department of Mathematics, Delhi College of Arts and Commerce,
University of Delhi, Netaji Nagar,
New Delhi-110023, India.
email: dishna2@yahoo.in

Kajal Negi

Department of Mathematics,
University of Delhi,
New Delhi-110007, India.
email: kajalnegi1109@gmail.com