

A guide to topological reconstruction on endomorphism monoids and polymorphism clones

Paolo Marimon¹ and Michael Pinsker²

¹Institut für Diskrete Mathematik & Geometrie, TU Wien. Vienna, Austria.

²Institut für Diskrete Mathematik & Geometrie, TU Wien. Vienna, Austria.

December 2, 2025

Abstract

Various spaces of symmetries of a structure are naturally endowed with both an algebraic and a topological structure. For example, the automorphism group of a structure is, on top of being a group, a topological group when equipped with the topology of pointwise convergence. In some cases, the algebraic structure of such space alone is sufficiently rich to determine its topology (under some requirements on the topology). For automorphism groups, the problem of when this happens has been actively pursued over the last 40 years. With the exception of some early work of Lascar, the analogue of this problem for endomorphism monoids and polymorphism clones has only received attention in the past 15 years. In this guide, we survey the current state of affairs in this relatively young line of research. We moreover use this opportunity to polish several existing results and to extend them beyond what was hitherto known.

Keywords: endomorphism monoid, polymorphism clone, topological reconstruction, Zariski topology, pointwise convergence topology, bi-interpretation, ω -categoricity.

1 Introduction

Given a structure \mathbb{A} with domain Ω , we may associate with it various spaces of symmetries: its automorphism group $\text{Aut}(\mathbb{A})$, its monoids of elementary self-embeddings $\text{EEmb}(\mathbb{A})$, of self-embeddings $\text{Emb}(\mathbb{A})$, and of endomorphisms $\text{End}(\mathbb{A})$, and its polymorphism clone $\text{Pol}(\mathbb{A})$ (the space of homomorphisms from some finite power of \mathbb{A} into itself, see Section 2.1).

Each space of symmetries of \mathbb{A} is naturally endowed with the topology of pointwise convergence τ_{pw} . For the spaces which are subsets of Ω^Ω (which include all but the last space above), this is just the subspace topology induced by the product topology on Ω^Ω , where Ω is endowed with the discrete topology; in the case of $\text{Pol}(\mathbb{A})$ it is the topology induced in the sum space $\mathcal{O} := \bigcup_{n \in \mathbb{N}} \Omega^{\Omega^n}$ of the spaces Ω^{Ω^n} which again bear the product topology. It is folklore that the subgroups of the full symmetric group S_Ω on Ω which are topologically closed in it correspond to automorphism groups of first-order structures with domain Ω . Analogously, closed submonoids of the full transformation monoid Ω^Ω are precisely the endomorphism monoids of first-order structures, and closed subclones of the full clone $\mathcal{O} := \bigcup_{n \in \mathbb{N}} \Omega^{\Omega^n}$ the polymorphism clones of first-order structures.

When Ω is countable, as we will be assuming throughout this paper, closed submonoids of Ω^Ω are Polish topological semigroups. That is, they are separable and completely metrizable (i.e. Polish) and the semigroup operation is continuous. Similarly, closed subgroups

of S_Ω are Polish topological groups (that is, both group multiplication and inversion are continuous), and closed subclones of \mathcal{O} are Polish topological clones (that is, composition of functions is continuous).

The picture described above raises two natural questions:

Question 1.1. What information can we recover about the original structure \mathbb{A} from a given space of symmetries (thought of as a topological group / monoid / clone)?

Question 1.2. To what extent does the algebraic structure of a space of symmetries determine its topological structure?

Regarding Question 1.1, we have an extremely satisfying answer when \mathbb{A} belongs to the class of ω -categorical structures, which includes e.g. $(\mathbb{N}; =)$, $(\mathbb{Q}; <)$, the random graph, countable vector spaces over finite fields, and the countable atomless Boolean algebra B_∞ . In that case, its spaces of symmetries recover the original structure up to certain notions of bi-interpretation [1, 14, 16]. In particular, $\text{Aut}(\mathbb{A})$ and $\text{EEmb}(\mathbb{A})$ recover \mathbb{A} up to bi-interpretability, $\text{Pol}(\mathbb{A})$ recovers \mathbb{A} up to the stronger notion of primitive positive bi-interpretability, and $\text{End}(\mathbb{A})$ recovers it up to the intermediate notion of existential positive bi-interpretability (modulo some minor caveats, see Theorem 2.8). Moreover, by a result of [51], restricting our attention to ω -categorical structures with no algebraicity, their spaces of symmetries with the topologies of pointwise convergence recover the original structure even up to bi-definability, see [27].

This fact makes Question 1.2 even more inviting in the ω -categorical context. There are several ways of making Question 1.2 precise, which gives rise to different notions of “topological reconstruction”. Let us focus for a moment on a closed transformation monoid S and on a class of topological semigroups \mathcal{C} . We say that S is **reconstructible** with respect to \mathcal{C} if whenever there is an algebraic isomorphism $\alpha : S \rightarrow T \in \mathcal{C}$, there is also a topological isomorphism $\xi : S \rightarrow T$. If, instead, we ask the topological isomorphism between S and T is already given by α , then we say that S has **automatic homeomorphicity** (AH). Automatic homeomorphicity with respect to the class of Polish semigroups is known as the **Unique Polish Property** (UPP) since it is equivalent to there being a unique Polish semigroup topology on S . Finally, one might ask that all semigroup homomorphism from S to a member T of \mathcal{C} are continuous. This property is known as **automatic continuity** (AC). All notions above can be defined analogously for topological groups and topological clones.

For automorphism groups of ω -categorical structures, these notions of reconstruction have been heavily studied in model theory, group theory, and topological dynamics. Reconstruction of $\text{Aut}(\mathbb{A})$ with respect to the class $\mathcal{C}_G^{\text{cl}}$ of automorphism groups of ω -categorical structures (also known as closed oligomorphic permutation groups) is equivalent to being able to recover \mathbb{A} from $\text{Aut}(\mathbb{A})$ up to bi-interpretation. And there are well-established (though not fully general) methods for proving properties that imply automatic homeomorphicity of $\text{Aut}(\mathbb{A})$ with respect to the class $\mathcal{C}_G^{\text{cl}}$ of closed subgroups of S_Ω . A notable fact for groups is that by the Open Mapping Theorem for Polish groups [6, Theorem 1.2.6], if $\text{Aut}(\mathbb{A})$ has automatic continuity with respect to a given class of Polish groups, then it also has automatic homeomorphicity with respect to it; in particular, this applies to $\mathcal{C}_G^{\text{cl}}$. Automatic continuity with respect to $\mathcal{C}_G^{\text{cl}}$ can be equivalently characterised by the **small index property** (SIP): a Polish group has the small index property if every subgroup of index $\leq \aleph_0$ is open¹. The small index property has been proved for several natural classes of ω -categorical structures: S_Ω , $\text{Aut}(\mathbb{Q}, <)$, the automorphism group of the countable atomless Boolean algebra, the automorphism groups of all ω -categorical ω -stable structures and of the random graph [23, 55, 35, 30]. With the exception of $\text{Aut}(\mathbb{Q}, <)$, the mentioned groups have been shown to have a stronger property than SIP, known as **ample generics**,

¹Some authors define the SIP by asking that subgroups of index $< 2^{\aleph_0}$ are open [39]. By Baire category, every open subgroup of a Polish group has index $\leq \aleph_0$, which is why we use the former definition for the purposes of automatic continuity.




Figure 1: Implications amongst automatic continuity (AC) and automatic homeomorphicity (AH) notions with respect to different classes, with the relevant citations. Above, G indicates the automorphism group of an ω -categorical structure, \bar{G} its monoid of elementary embeddings, and S its endomorphism monoid.

which implies automatic continuity with respect to the class \mathcal{C}_G^2 of second countable topological groups [34]. Finally, showing that $\text{Aut}(\mathbb{A})$ has a **weak $\forall\exists$ -interpretation** (in the sense of [51]) also implies that $\text{Aut}(\mathbb{A})$ has automatic homeomorphicity with respect to $\mathcal{C}_G^{\text{cl}}$. This latter method applies to several structures for which the SIP is not known (such as the generic poset), or does not hold (such as the Cherlin-Hrushovski structure, cf. Definition 2.6) [2].

Moving from automorphism groups to monoids of elementary embeddings, endomorphism monoids, and polymorphism clones, the way various notions of reconstruction interact takes several unexpected turns. The point of this survey is to help the reader navigate this sea of notions of topological reconstruction and the techniques that have been developed in this field for spaces of symmetries other than automorphism groups. In the process of doing so, we polished several existing results in the literature to clarify the picture of the current state of the art, and we also prove the significant transfer result (Theorem 4.7) that, for \mathbb{A} a countable ω -categorical structure, automatic homeomorphicity of $\text{Aut}(\mathbb{A})$ with respect to $\mathcal{C}_G^{\text{cl}}$ implies automatic homeomorphicity of $\text{EEmb}(\mathbb{A})$ with respect to the class $\mathcal{C}_{\bar{T}}^{\text{cl}}$ of closed transformation monoids on Ω .

Roughly, the picture we will paint for monoids of elementary embeddings and endo-

morphism monoids is as follows: very little is known about reconstructibility (wrt $\mathcal{C}_f^{\text{cl}}$) as a separate notion from automatic homeomorphicity (wrt $\mathcal{C}_f^{\text{cl}}$). The only known ω -categorical counterexample to reconstruction is the same for automorphism groups, monoids of elementary embeddings, endomorphism monoids, and polymorphism clones [13], and comes from a classical construction by [26]. Regarding automatic homeomorphicity and automatic continuity, we summarise the current situation in Figure 1. Notably, there is no Open Mapping Theorem for Polish semigroups, and hence automatic continuity and homeomorphicity are formally incomparable notions. The high connectivity of the top part of the figure displays the main result of this paper (Theorem 4.7). Nevertheless, the reader should not be fooled by what is omitted from the figure: no monoid of elementary embeddings of an ω -categorical structure has UPP or automatic continuity with respect to $\mathcal{C}_f^{\text{cl}}$. Meanwhile, the bottom of the figure is highly disconnected from the top half, with the only implications being the obvious ones. In particular, no matter how strong the reconstruction properties of $\text{Aut}(\mathbb{A})$ are, $\text{End}(\mathbb{A})$ may fail to have automatic homeomorphicity even with respect to itself. This is the result of our efforts in polishing previous results of [18].

Despite the disconnect between the bottom and top part of Figure 1, several techniques have been developed to prove versions of automatic homeomorphicity for endomorphism monoids (and polymorphism clones). On one hand, there is a body of “bottom-up” results which rely on transferring upwards automatic homeomorphicity from $\text{Aut}(\mathbb{A})$ to $\text{End}(\mathbb{A})$, passing through $\text{EEmb}(\mathbb{A})$. This in particular includes work of [7, 56, 8, 25] and previous work on monoids of elementary embeddings now subsumed by Theorem 4.7 [47, 18, 7, 56]. On the other hand, the seminal paper [25] introduced several new ideas and techniques which are especially suited to the study of semigroups and clones and which do not (necessarily) rely on upwards transfer from the study of some underlying topological group. In particular, the authors introduce **Property X**, a frequently occurring property for semigroups that can be used to deduce maximality of the topology of pointwise convergence amongst Polish semigroup topologies. Property X can then be mixed with the study of the **Zariski topology** τ_Z , a topology defined through purely intrinsic information on an algebraic semigroup, which, if shown to coincide with τ_{pw} , implies that τ_{pw} is minimal amongst Hausdorff semigroup topologies. This yields a route to prove UPP without relying on any form of reconstruction for an underlying group which is often fruitful since $\tau_{\text{pw}} = \tau_Z$ on $\text{End}(\mathbb{A})$ under fairly general conditions [49, 24, 42]. The latter fact is notable since it stands in stark contrast with the situation for groups, where the Zariski topology tends to be less well-behaved and usually distinct from τ_{pw} [28, 3].

Indeed, the work of [49] in combination with [51] (see [27]) yields another remarkable fact that we note in this survey: monoids of elementary embeddings of ω -categorical structures with no algebraicity have **automatic action compatibility** in the sense that any algebraic isomorphism between two of them is induced from a bijection between their underlying domains. The analogue of this statement for automorphism groups is wide open, with substantial works in the field proving it under additional assumptions [51, 44, 2].

In spite of the growing body of available techniques, we are still far from any form of complete picture for topological reconstruction of endomorphism monoids or polymorphism clones. Throughout this guide, we include several open questions which we believe may help the field grow.

2 Preliminaries

We will assume some basic knowledge of model theory, such as the first five chapters of [53]. For more on ω -categorical structures and topological reconstruction in the context of automorphism groups, we refer the reader to the survey [39].

2.1 Spaces of symmetries and their topologies

Let Ω be a countable set. We endow Ω^Ω with the topology of pointwise convergence τ_{pw} given by the product topology on Ω^Ω , where Ω is assigned the discrete topology. We write $\mathcal{O}^{(1)} := \Omega^\Omega$ for the **full transformation monoid** on Ω , where the monoid operation is composition of functions. When we talk about the topology of pointwise convergence τ_{pw} on any subspace of $\mathcal{O}^{(1)}$ we mean the induced subspace topology. The monoid $\mathcal{O}^{(1)}$ with the topology τ_{pw} is a **topological semigroup** in the sense that the semigroup operation is continuous. Moreover, $\mathcal{O}^{(1)}$ is Polish, i.e. separable and completely metrizable. Closed submonoids of $\mathcal{O}^{(1)}$ are still topological semigroups with τ_{pw} . The full symmetric group S_Ω with τ_{pw} is a Polish **topological group** (i.e. composition and inversion are continuous), and any of its subgroups which is closed in S_Ω is a Polish topological group.

For $n \geq 1$, let $\mathcal{O}^{(n)} = \Omega^{\Omega^n}$ be the set of functions $f : \Omega^n \rightarrow \Omega$, and $\mathcal{O} := \bigcup_{n \in \mathbb{N}} \mathcal{O}^{(n)}$. We call \mathcal{O} the **full clone** on Ω . Analogously to the previous paragraph, we endow each $\mathcal{O}^{(n)}$ with the product topology and endow \mathcal{O} with the sum topology. This defines the topology of pointwise convergence τ_{pw} on \mathcal{O} . Note that τ_{pw} has a sub-basis of open sets $\{\mathcal{U}_{(\bar{a},b)} \mid \bar{a} \in \Omega^n, b \in \Omega, n \geq 1\}$, where for $\bar{a} \in \Omega^n, b \in \Omega$,

$$\mathcal{U}_{(\bar{a},b)} := \{f \in \mathcal{O}^{(n)} \mid f(\bar{a}) = b\}.$$

Similarly, $\{\mathcal{U}_{(a,b)} \mid a, b \in \Omega\}$ forms a subbasis of open sets of τ_{pw} on $\mathcal{O}^{(1)}$. We often find it convenient to think of the notion of topological closure induced by τ_{pw} : given $\mathcal{F} \subseteq \mathcal{O}$ and $f \in \mathcal{O}$, we say that $f \in \overline{\mathcal{F}}$ if for all finite $C \subseteq \Omega$ there is some $g \in \mathcal{F}$ such that $g|_C = f|_C$. Topological closure on $\mathcal{O}^{(1)}$ is identical.

Definition 2.1. We call $\mathcal{C} \subseteq \mathcal{O}$ a **function clone** over B if

- \mathcal{C} contains all projections: for each $1 \leq i \leq k \in \mathbb{N}$, \mathcal{C} contains the k -ary projection to the i th coordinate $\pi_i^k \in \mathcal{O}^{(k)}$, given by $(x_1, \dots, x_k) \mapsto x_i$;
- \mathcal{C} is closed under composition: for all $f \in \mathcal{C} \cap \mathcal{O}^{(n)}$ and all $g_1, \dots, g_n \in \mathcal{C} \cap \mathcal{O}^{(m)}$, $f(g_1, \dots, g_n)$, given by

$$(x_1, \dots, x_m) \mapsto f(g_1(x_1, \dots, x_m), \dots, g_n(x_1, \dots, x_m)),$$

is in $\mathcal{C} \cap \mathcal{O}^{(m)}$.

We say that a function clone is **closed** if it is closed in τ_{pw} on \mathcal{O} . Given a set of functions $\mathcal{S} \subseteq \mathcal{O}$, we denote by $\langle \mathcal{S} \rangle$ the smallest closed function clone containing \mathcal{S} . A closed function clone \mathcal{C} together with τ_{pw} is a (Polish) **topological clone** in the sense that each $\mathcal{C} \cap \mathcal{O}^{(n)}$ is clopen and each operation $\text{comp}_m^n : \mathcal{O}^{(n)} \times (\mathcal{O}^{(m)})^n \rightarrow \mathcal{O}^{(m)}$ given by $(f, g_1, \dots, g_n) \mapsto f(g_1, \dots, g_n)$ is continuous.

Definition 2.2. Let \mathcal{C} and \mathcal{D} be function clones. A **clone homomorphism** $\xi : \mathcal{C} \rightarrow \mathcal{D}$ is a map which preserves arities of functions, is constant on the projections $\{\pi_i^k \mid 1 \leq i \leq k\}$ and preserves composition of functions.

Definition 2.3. Let \mathbb{A} and \mathbb{B} be first-order structures in the same relational language. A map $f : \mathbb{A} \rightarrow \mathbb{B}$ is a **homomorphism** if it sends tuples in a relation of \mathbb{A} to tuples in the corresponding relation of \mathbb{B} ; it is an **embedding** if it is moreover injective and sends only tuples in a relation of \mathbb{A} to tuples in the corresponding relation of \mathbb{B} ; and it is an **elementary embedding** if it preserves the truth of arbitrary first-order formulas. We write $\text{Aut}(\mathbb{A})$, $\text{Emb}(\mathbb{A})$, $\text{EEmb}(\mathbb{A})$, and $\text{End}(\mathbb{A})$ for, respectively, the automorphism group, the monoid of self-embeddings, the monoid of elementary self-embeddings, and the endomorphism monoid of \mathbb{A} . By a **polymorphism** of \mathbb{A} we mean a map $\mathbb{A}^n \rightarrow \mathbb{A}$ for some $n \geq 1$ which

preserves each relation R of \mathbb{A} :

$$\text{if } \begin{pmatrix} a_1^1 \\ \vdots \\ a_k^1 \end{pmatrix}, \dots, \begin{pmatrix} a_1^n \\ \vdots \\ a_k^n \end{pmatrix} \in R, \text{ then } \begin{pmatrix} f(a_1^1, \dots, a_1^n) \\ \vdots \\ f(a_k^1, \dots, a_k^n) \end{pmatrix} \in R.$$

Note that unary polymorphisms correspond to endomorphisms and that the polymorphisms of a relational structure form a closed subclone of \mathcal{O} . We write $\text{Pol}(\mathbb{A})$ for the polymorphism clone of \mathbb{A} .

The following correspondences are easy to prove (allowing relational structures to be in an infinite language):

- closed subgroups of S_Ω correspond to automorphism groups of relational structures on Ω ;
- closed submonoids of $\mathcal{O}^{(1)}$ correspond to endomorphism monoids of relational structures on Ω ;
- closed function clones on Ω correspond to polymorphism clones of relational structures on Ω .

2.2 ω -categorical structures and related notions

Definition 2.4. Let Ω be a countable set (we allow Ω to be finite). We say that a permutation group $G \curvearrowright \Omega$ is **oligomorphic** if $G \curvearrowright \Omega^n$ has finitely many orbits for each $n \in \mathbb{N}$ in its natural action $g \cdot (x_1, \dots, x_n) = (gx_1, \dots, gx_n)$ for $(x_1, \dots, x_n) \in \Omega^n$. A structure \mathbb{A} is **ω -categorical** if its automorphism group is oligomorphic.

With our definition, all finite structures are ω -categorical. A large class of countably infinite ω -categorical structures is given by homogeneous structures in a finite relational language (**finitely homogeneous** structures):

Definition 2.5. A countably infinite structure \mathbb{A} in a relational language is **homogeneous** if every isomorphism between finite substructures extends to an automorphism of \mathbb{A} .

Homogeneous structures are in one-to-one correspondence with certain well-behaved classes of finite relational structures known as Fraïssé classes. Given a Fraïssé class \mathcal{K} , the homogeneous structure whose **age** (i.e. class of finite substructures) is \mathcal{K} is known as its **Fraïssé limit**. We refer the reader to [39] for a survey on ω -categorical and homogeneous structures. We usually name homogeneous structure after their age (for which they are the **generic** countable object). Thus, the generic K_n -free graph \mathcal{H}_n is the countable homogeneous graph whose age is the class of finite graphs which do not embed a copy of the complete graph K_n , the generic poset is the Fraïssé limit of finite partially ordered sets, and so on.

Spaces of symmetries of ω -categorical structures have natural characterisations in terms of their actions: closed oligomorphic subgroups of S_Ω correspond to the automorphism groups of ω -categorical structures; closed transformation monoids on Ω^Ω are called oligomorphic if they contain an oligomorphic permutation group, and correspond to endomorphism monoids of ω -categorical structures. Moreover, monoids of elementary embeddings of ω -categorical structures correspond precisely to closed oligomorphic transformation monoids with a dense set of invertibles. Similarly, polymorphism clones of ω -categorical structures correspond to closed oligomorphic clones, i.e., closed clones whose unary operations are an oligomorphic transformation monoid.

In some sections, we will be working with countable **saturated** structures. These are countable structures \mathbb{A} such that for each of their finite subsets B , \mathbb{A} realises all 1-types

over B . In particular, ω -categorical structures are saturated. In countable saturated structures we have that $\text{EEmb}(\mathbb{A}) = \overline{\text{Aut}(\mathbb{A})}$.

The latter fact provides one motivation for reconstructing the topology on the class of closed submonoids \mathcal{S} of Ω^Ω whose subgroup G of invertibles is dense. Further, monoids of this class are an essential intermediate step when lifting reconstruction results from automorphism groups to endomorphism monoids. Finally, if G is even oligomorphic, such monoids are called **model-complete cores**, and they appear as an essential concept in various applications e.g. in Ramsey theory or Constraint Satisfaction (see [41, 4, 10, 11]). The name of model-complete core was originally given to ω -categorical structures whose endomorphism monoid has a dense set of invertibles; it was shown in [10] that every ω -categorical structure \mathbb{A} has a substructure \mathbb{C} which is a model-complete core and such that \mathbb{A} has a homomorphism into \mathbb{C} , and \mathbb{C} is ω -categorical and unique up to isomorphism [10]. Closed clones whose monoid of unary functions are a model-complete core are called the same.

Given a group $G \curvearrowright \Omega$ and $B \subseteq \Omega$ finite, the **algebraic closure** of B , $\text{acl}(B)$, is the set of all elements of Ω which have finite orbit under the action $G_{(B)} \curvearrowright \Omega$ of the pointwise stabiliser of B . We say that $G \curvearrowright \Omega$ has **locally finite algebraic closure** if the algebraic closure of any finite set is finite; this is the case for all oligomorphic actions. Moreover, $G \curvearrowright \Omega$ has **no algebraicity** if for all finite $B \subseteq \Omega$, $\text{acl}(B) = B$. It is easy to see that this implies that the center $Z(G)$ of G is trivial.

Some remarkable results of [36, 37] on monoids of elementary embeddings rely on the assumption that the underlying group G is **G-finite**: for every open subgroup $U \leq G$, the intersection of all closed subgroups of finite index in U still has finite index in U . As far as we know, there are no known examples of finitely homogeneous structures which are not G-finite. However, allowing for an infinite language it is easy to build such structures [38]:

Definition 2.6. By the Cherlin-Hrushovski structure \mathbb{A}_{CH} we mean the homogeneous structure obtained as the Fraïssé limit of finite structures with, for each $n \in \mathbb{N}$, a $2n$ -ary equivalence relation E_n on n -tuples, such that each E_n has at most two equivalence classes.

The Cherlin-Hrushovski structure also does not have the small index property [36].

2.3 Interpretations

One motivation for reconstructing the topology of various spaces of symmetries of an ω -categorical structure is that one can then further reconstruct the structure itself up to various notions of interpretation.

Definition 2.7. Let \mathbb{A} and \mathbb{B} be first-order structures, respectively in the signatures \mathcal{L} and \mathcal{L}' . A **first-order interpretation (fo-interpretation)** of \mathbb{B} in \mathbb{A} is a partial surjection $I : A^d \rightarrow B$ for some $d \in \mathbb{N}$ such that for each relation R of \mathbb{B} defined by an atomic \mathcal{L}' -formula $\phi(x_1, \dots, x_n)$, the relation

$$I^{-1}(R) := \{(a_1^1, \dots, a_1^d, \dots, a_n^1, \dots, a_n^d) \in A^{nd} \mid \mathbb{B} \models \phi(I(a_1^1, \dots, a_1^d), \dots, I(a_n^1, \dots, a_n^d))\}$$

is first-order definable in \mathbb{A} . We say that d is the **dimension** of the interpretation I . Below, we are mainly interested in stronger notions of interpretations. In particular, we say that I is an

- **existential interpretation** (e-interpretation) if each $I^{-1}(R)$ is definable in \mathbb{A} as an existential formula;
- **existential positive interpretation** (ep-interpretation) if each $I^{-1}(R)$ is existentially positively definable in \mathbb{A} ;

- **primitive positive interpretation** (pp-interpretation) if each $I^{-1}(R)$ is primitively positively definable in \mathbb{A} .

It is easy to see that any interpretation I of \mathbb{B} in \mathbb{A} naturally gives rise to a continuous group homomorphism $\text{Aut}(I) : \text{Aut}(\mathbb{A}) \rightarrow \text{Aut}(\mathbb{B})$. Moreover, for ω -categorical \mathbb{A} and \mathbb{B} , a continuous group homomorphism $f : \text{Aut}(\mathbb{A}) \rightarrow \text{Aut}(\mathbb{B})$ is of the form $\text{Aut}(I)$ for some interpretation I if and only if its range acts oligomorphically on B [1]. Analogous statements hold for e-interpretations with respect to embedding monoids [14, Theorem 3.2], for ep-interpretations with respect to endomorphism monoids [14, Theorem 3.6], and for pp-interpretations with respect to polymorphism clones [16, Proposition 22]. Whilst this already motivates understanding when homomorphism between oligomorphic spaces of symmetries are continuous, our main interest in this survey lies in topological isomorphisms. We will see below these encode bi-interpretations of varying degrees of strength.

Note that interpretations compose naturally. In particular, if I is an (fo/e/ep/pp)-interpretation of \mathbb{B} in \mathbb{A} and J is an (fo/e/ep/pp)-interpretation of \mathbb{A} in \mathbb{B} , we say that \mathbb{A} and \mathbb{B} are (fo/e/ep/pp)-**bi-interpretable** if the induced partial functions $I \circ J$ and $J \circ I$ are (fo/e/ep/pp)-definable in \mathbb{B} and \mathbb{A} respectively. See e.g. [1, 14, 16] for details.

Theorem 2.8. *Let \mathbb{A} and \mathbb{B} be ω -categorical structures of size > 1 . Then:*

1. *\mathbb{A} and \mathbb{B} are bi-interpretable if and only if $\text{Aut}(\mathbb{A})$ and $\text{Aut}(\mathbb{B})$ are isomorphic as topological groups (Coquand, Corollary 1.4 in [1]).*
2. *If $\text{Emb}(\mathbb{A})$ and $\text{Emb}(\mathbb{B})$ are isomorphic as topological monoids, then \mathbb{A} and \mathbb{B} are existentially bi-interpretable [14, Theorem 3.14]. The converse does not hold [14, Example 4.4].*
3. *If \mathbb{A} and \mathbb{B} are existentially positively bi-interpretable, then $\text{End}(\mathbb{A})$ and $\text{End}(\mathbb{B})$ are isomorphic as topological monoids. The converse holds as long as neither of \mathbb{A} and \mathbb{B} have a constant endomorphism (by [14, Theorem 3.12]).*
4. *\mathbb{A} and \mathbb{B} are primitively positively bi-interpretable if and only if $\text{Pol}(\mathbb{A})$ and $\text{Pol}(\mathbb{B})$ are isomorphic as topological clones [16].*

As mentioned, (3) does not hold in full generality: let \mathbb{A} be an ω -categorical structure with a constant endomorphism. Let \mathbb{A}' be the structure obtained by adding a new element c to \mathbb{A} as well as unary predicates for $\{c\}$ and its complement in \mathbb{A}' (which is \mathbb{A}). Then $\text{End}(\mathbb{A})$ and $\text{End}(\mathbb{A}')$ are topologically isomorphic. However, any structure existentially positively interpretable in \mathbb{A} also has a constant endomorphism [14, Lemma 3.9], so \mathbb{A} and \mathbb{A}' cannot be bi-interpretable. Nevertheless, one can get a fully general correspondence looking at clones and using [16]:

Lemma 2.9. *Let \mathbb{A} and \mathbb{B} be ω -categorical structures. Then, \mathbb{A} and \mathbb{B} are existentially positively bi-interpretable if and only if $\langle \text{End}(\mathbb{A}) \rangle$ and $\langle \text{End}(\mathbb{B}) \rangle$ are isomorphic as topological clones.*

Proof. One direction is a direct consequence of (3). For the other, suppose that the two clones are isomorphic. These clones are the polymorphism clones of the structures \mathbb{A}', \mathbb{B}' obtained by enriching \mathbb{A}, \mathbb{B} by the 4-ary relation defined by $x = y \vee u = v$. By (4), \mathbb{A}', \mathbb{B}' are pp-bi-interpretable, and hence \mathbb{A}, \mathbb{B} are ep-bi-interpretable. The reason constant endomorphisms do not pose a problem in this setting is that they are detected by the clone structure: an endomorphism f is constant if and only if $f \circ \pi_1^2 = f \circ \pi_2^2$ for the two binary projections π_1^2, π_2^2 . \square

Studying monoids of elementary embeddings is also relevant to the problem of reconstructing structures up to first-order bi-interpretability. The following folklore result is a direct consequence of Proposition 3.1 and Theorem 2.8.

Proposition 2.10. *Let \mathbb{A} and \mathbb{B} be ω -categorical structures of size > 1 . Then, \mathbb{A} and \mathbb{B} are first-order bi-interpretable if and only if $\text{EEmb}(\mathbb{A})$ and $\text{EEmb}(\mathbb{B})$ are isomorphic as topological monoids.*

While this characterization of first-order bi-interpretability might seem less natural than the one with automorphism groups, it may actually be easier to verify than the latter. For example, by Theorem 5.6 [49], if two ω -categorical structures with no algebraicity have isomorphic endomorphism monoids of elementary embeddings, then this isomorphism will be a topological one. No analogue of this result is known for automorphism groups.

2.4 Notions of reconstruction

Definition 2.11. Let S be a topological semigroup and \mathcal{C} be a class of topological semigroups. We say that

1. S is **reconstructible** with respect to \mathcal{C} if whenever there is a semigroup isomorphism from S to a member \mathcal{T} of \mathcal{C} , there is also a semigroup isomorphism which is moreover a homeomorphism between S and \mathcal{T} ;
2. S has **automatic homeomorphicity** with respect to \mathcal{C} if every semigroup isomorphism between S and a member \mathcal{T} of \mathcal{C} is a homeomorphism;
3. S has **automatic continuity** with respect to \mathcal{C} if every semigroup homomorphism from S to a member \mathcal{T} of \mathcal{C} is continuous.

Analogous definitions exist for topological groups and clones and the corresponding notions of homomorphism.

Definition 2.12. We define several classes of topological groups, semigroups and clones \mathcal{C}_X^Y depending on the parameters X and Y as follows: $X \in \{\mathcal{G}, \bar{\mathcal{G}}, \mathcal{T}, \mathcal{C}\}$ indicates whether the class we are looking at is a class of topological groups (\mathcal{G}), topological monoids with a dense group of invertible elements ($\bar{\mathcal{G}}$), semigroups (\mathcal{T}), or clones (\mathcal{C}). Meanwhile, we use $Y \in \{2, \text{cl}, \text{sat}, \text{ol}\}$ to indicate whether we are looking at the class of (in the order of decreasing generality):

1. second countable topological groups, semigroups, clones, which we denote, respectively, by $\mathcal{C}_{\mathcal{G}}^2, \mathcal{C}_{\bar{\mathcal{G}}}^2, \mathcal{C}_{\mathcal{C}}^2$.
2. closed subgroups of S_{Ω} , closed submonoids of Ω^{Ω} with a dense set of invertibles, closed submonoids of Ω^{Ω} , or closed subclones of \mathcal{O} , which we denote, respectively, by $\mathcal{C}_{\mathcal{G}}^{\text{cl}}, \mathcal{C}_{\bar{\mathcal{G}}}^{\text{cl}}, \mathcal{C}_{\mathcal{T}}^{\text{cl}}, \mathcal{C}_{\mathcal{C}}^{\text{cl}}$.
3. the classes in item 2 restricted to those elements whose underlying group of invertible elements is the automorphism group of a countable saturated structure. We denote these by $\mathcal{C}_{\mathcal{G}}^{\text{sat}}, \mathcal{C}_{\bar{\mathcal{G}}}^{\text{sat}}, \mathcal{C}_{\mathcal{T}}^{\text{sat}}, \mathcal{C}_{\mathcal{C}}^{\text{sat}}$, respectively.
4. the classes in item 2 restricted to those elements whose underlying group of invertible elements is oligomorphic. We denote these by $\mathcal{C}_{\mathcal{G}}^{\text{ol}}, \mathcal{C}_{\bar{\mathcal{G}}}^{\text{ol}}, \mathcal{C}_{\mathcal{T}}^{\text{ol}}, \mathcal{C}_{\mathcal{C}}^{\text{ol}}$, respectively.

Another established notion of reconstruction is when a semigroup (group, clone) has a unique Polish compatible topology. This is known as the **Unique Polish Property** (UPP). Note that UPP is equivalent to automatic homeomorphicity with respect to the class of Polish topological semigroups (groups, clones).

3 Reconstruction

The following is implicit in [36, 37]:

Proposition 3.1 (Proposition 11 in [18]). *Let G and H be closed permutation groups on Ω . Let $\xi : G \rightarrow H$ be a continuous homomorphism. Then,*

1. ξ extends to a continuous homomorphism $\bar{\xi} : \bar{G} \rightarrow \bar{H}$;
2. if ξ is an isomorphism then $\bar{\xi}$ is an isomorphism and a homeomorphism.

Consequently, G and H are isomorphic as topological groups if and only if \overline{G} and \overline{H} are isomorphic as topological semigroups.

Proposition 3.1 was an ingredient for the following result of Lascar, which already suggests that monoids of elementary embeddings yield more information towards reconstruction than automorphism groups:

Theorem 3.2 ([37]). *Let $G \in \mathcal{C}_G^{\text{ol}}$ be G -finite, and let $H \in \mathcal{C}_G^{\text{cl}}$. Let $\eta : \overline{G} \rightarrow \overline{H}$ be an algebraic isomorphism. Then, $\eta|_G : G \rightarrow H$ is a topological isomorphism.*

Note that Proposition 3.1 yields the following upwards reconstruction result:

Corollary 3.3. *Let $G \in \mathcal{C}_G^{\text{cl}}$ have reconstruction with respect to $\mathcal{C}_G^{\text{cl}}$. Then, \overline{G} has reconstruction with respect to $\mathcal{C}_G^{\text{cl}}$.*

The result above also yields transfer of reconstruction from G with respect to $\mathcal{C}_G^{\text{ol}}$ to \overline{G} with respect to $\mathcal{C}_G^{\text{ol}}$. Very little is known about other forms of transfer of reconstructibility:

Question 3.4. Let $G \in \mathcal{C}_G^{\text{cl}}$.

- (reconstruction lifting) Does G having reconstruction with respect to $\mathcal{C}_G^{\text{cl}}$ imply that \overline{G} has reconstruction with respect to $\mathcal{C}_G^{\text{cl}}$?
- (reconstruction descent) Does \overline{G} having reconstruction with respect to $\mathcal{C}_G^{\text{cl}}$ imply that G has reconstruction with respect to $\mathcal{C}_G^{\text{cl}}$?

The main issue with the first question is that we may not be able to ensure that any topological semigroup isomorphic to \overline{G} is such that its subgroup of invertible elements is dense. For the second question, an issue is that algebraic isomorphisms between groups may not lift to algebraic isomorphisms of their closures. Both questions are open even if G is oligomorphic, cf. [13, Questions 4.1 & 4.2].

It is important to keep in mind throughout this survey that we do have a counterexample to reconstruction at all levels of symmetries of an ω -categorical structure; however, surprisingly it is the only one we are aware of.

Theorem 3.5 ([26, 13]). *There are $G, H \in \mathcal{C}_G^{\text{ol}}$ such that*

- *the groups G and H are algebraically, but not topologically isomorphic;*
- *the monoids \overline{G} and \overline{H} are algebraically, but not topologically isomorphic;*
- *the clones $\langle \overline{G} \rangle$ and $\langle \overline{H} \rangle$ are algebraically, but not topologically isomorphic.*

The structures in Theorem 3.5 rely on the existence of separable profinite groups P and Q which are algebraically but not topologically isomorphic [58]. By a generalisation of the Cherlin-Hrushovski structure, these can then be encoded as the quotients by an oligomorphic normal subgroup of two oligomorphic groups Σ_P and Σ_Q [13, Proposition 3.4]. These are then “perturbed” to obtain G and H . We note that the latter groups have a non-trivial center (and in particular, algebraicity) and are not G -finite.

4 Automatic homeomorphicity

4.1 Automatic homeomorphicity for monoids of elementary embeddings

In this subsection, we prove that for any countable saturated structure, automatic homeomorphicity lifts from its automorphism group (with respect to $\mathcal{C}_G^{\text{cl}}$) to its monoid of elementary embeddings (with respect to $\mathcal{C}_G^{\text{cl}}$). This is an original result, completing the answer to a problem which is central to several papers on topological reconstruction for monoids [18, 7,

56, 47]. In particular, [47] answer this problem positively in the context of saturated structures whose automorphism group has a trivial centre. Uncovering the key idea behind their proof is the essential tool to answer this question in full generality.

We begin from the following key lemma for reconstruction:

Lemma 4.1 ([18, Lemma 12]). *Let $G \in \mathcal{C}_G^{\text{cl}}$ have automatic homeomorphicity with respect to $\mathcal{C}_G^{\text{cl}}$. Suppose that the only injective $\Phi \in \text{End}(\overline{G})$ that fixes G pointwise is the identity. Then \overline{G} has automatic homeomorphicity with respect to $\mathcal{C}_G^{\text{cl}}$.*

The majority of this section is dedicated to proving the assumption on $\text{End}(\overline{G})$ is always satisfied when G is the automorphism group of a saturated structure.

Definition 4.2. Let $G \in \mathcal{C}_G^{\text{cl}}$. We call a subset $S \subseteq \Omega$ **free** if the orbits of tuples under the action $G_{\{S\}} := \{\alpha \in G \mid \alpha(S) = S\} \curvearrowright S$ are precisely the restrictions of orbits of $G \curvearrowright \Omega$ to S . We say that S is **Galois-closed** if the pointwise stabilizer G_S of S in G acts without fixed points on $\Omega \setminus S$. Finally, $f \in \overline{G}$ is free Galois-closed if its image is.

We remark that this notion is a translation of the notion of superhomogeneity used in [47] from structures to permutation groups. We shall now apply some results of that paper which are formulated there in terms of structures. For any countable saturated structure \mathbb{A} , there is a structure \mathbb{A}' , which is saturated, homogeneous (in a countable language), and such that $\text{Aut}(\mathbb{A}) = \text{Aut}(\mathbb{A}')$ (see, e.g., [47]), allowing us to speak in terms of $G := \text{Aut}(\mathbb{A})$. When G is oligomorphic, this is known as the **canonical structure** for G . In this case \mathbb{A}' is obtained by adding a relation for every orbit of tuples under G .

Proposition 4.3 ([47]). *Let $G \in \mathcal{C}_G^{\text{cl}}$ and let $\Phi \in \text{End}(\overline{G})$ fix G pointwise. Then for all free Galois-closed $h \in \overline{G}$ there exists $g_h \in Z(G)$, such that $\Phi(h) = h \circ g_h$.*

Proof. For $f \in \overline{G}$, let

$$f^* := \{(\alpha, \beta) \mid \alpha, \beta \in G, \alpha \circ f = f \circ \beta\}.$$

Clearly, for any $h \in \overline{G}$ we have $\Phi(h)^* = h^*$. By Corollary 3.12 and Proposition 3.13 of [47] this implies that there is some $g_h \in Z(G)$ such that $\Phi(h) = h \circ g_h$. \square

Our key observation is that the main results of [47] rely on a correspondence between maps in $\text{End}(\overline{G})$ which fix G pointwise and homomorphisms $\phi : \overline{G} \rightarrow Z(G)$ mapping G to 1:

Proposition 4.4. *Let $G \in \mathcal{C}_G^{\text{sat}}$ and let $\Phi \in \text{End}(\overline{G})$ fix G pointwise. Then, there is a semigroup homomorphism $\phi : \overline{G} \rightarrow Z(G)$ such that $\phi(G) = \{1\}$ and for each $f \in \overline{G}$*

$$\Phi(f) = \phi(f) \circ f.$$

Proof. For any $f \in \overline{G}$, by [47, Proposition 5.1], we can take free Galois-closed $g, h \in \overline{G}$ such that $g = h \circ f$. By Proposition 4.3, there are $z_g, z_h \in Z(G)$ such that $\Phi(g) = g \circ z_g$ and $\Phi(h) = h \circ z_h$. Now, since Φ is a homomorphism,

$$h \circ f \circ z_g = g \circ z_g = \Phi(g) = \Phi(h \circ f) = \Phi(h)\Phi(f) = h \circ z_h \circ \Phi(f).$$

Since h is injective and $z_h, z_g \in Z(\overline{G})$ (being in $Z(G)$), we get

$$\Phi(f) = z_h^{-1} \circ z_g \circ f.$$

In particular, we can set $\phi(f) = z_h^{-1} \circ z_g$. Since Φ fixes G pointwise, we must have that $\phi(G) = 1$. We now claim that $\phi : \overline{G} \rightarrow Z(G)$ is a homomorphism. To see this, note that for $f, g \in \overline{G}$, since $\phi(f), \phi(g) \in Z(G)$,

$$(f \circ g) \circ (\phi(f) \circ \phi(g)) = (\phi(f) \circ f) \circ (\phi(g) \circ g) = \Phi(f \circ g) = (f \circ g) \circ (\phi(f \circ g)),$$

which by injectivity of $f \circ g$ implies $\phi(f \circ g) = \phi(f) \circ \phi(g)$ as required. \square

The proof of the following lemma will use some basic model-theoretic techniques.

Lemma 4.5. *Let $G \in \mathcal{C}_G^{\text{sat}}$. Let $h \in \overline{G}$. Then there are $f \in \overline{G}$ and $\alpha, \beta \in G$ such that*

$$\alpha \circ f \circ h = f \circ \beta.$$

Proof. Let $h \in \overline{G}$. Let \mathbb{A} be a saturated structure with automorphism group G . Write \mathbb{A}_1 for \mathbb{A} and \mathbb{A}_0 for $h(\mathbb{A}_1)$. Since h is an elementary self-embedding, its inverse is an isomorphism $\alpha_0 : \mathbb{A}_0 \rightarrow \mathbb{A}_1$. We first construct an elementary extension $\mathbb{A}_2 \succeq \mathbb{A}_1$ together with an isomorphism $\alpha_1 : \mathbb{A}_1 \rightarrow \mathbb{A}_2$ extending α_0 . To do so, take an enumeration of \mathbb{A}_1 . Consider the type $p(z; A_0) := \text{tp}(\mathbb{A}_1 / A_0)$ in a variable z of length ω . Let \mathbb{A}_2 be a realisation of $p(x; \alpha_0(A_0))$ and define α_1 to be the map sending every element of \mathbb{A}_1 to its realisation in \mathbb{A}_2 . This map is an isomorphism and extends α_0 . By repeating this process, we get an elementary chain

$$\mathbb{A}_0 \preceq \mathbb{A}_1 \preceq \cdots \preceq \mathbb{A}_i \preceq \mathbb{A}_{i+1} \preceq \cdots$$

where, at each stage $i \geq 1$ we have an isomorphism $\alpha_i : \mathbb{A}_i \rightarrow \mathbb{A}_{i+1}$ extending α_{i-1} . Write \mathbb{A}' for the union of this elementary chain and α for $\bigcup_{i \in \mathbb{N}} \alpha_i$. By construction, α is an automorphism of \mathbb{A}' . Since it is a union of a countable chain of countable saturated models, \mathbb{A}' is still a countable saturated model of the first-order theory of \mathbb{A} . In particular, there is an isomorphism $f : \mathbb{A}' \rightarrow \mathbb{A}_1$ (cf. [53, Lemma 5.2.8]). Define $h' : \mathbb{A}' \rightarrow \mathbb{A}'$ as $f^{-1}hf$. Note that, for $a, b \in \mathbb{A}'$,

$$h'(a) = b \Leftrightarrow f^{-1}hf(a) = b \Leftrightarrow hf(a) = f(b).$$

Hence, since f is an isomorphism between \mathbb{A}' and \mathbb{A} , it yields an isomorphism between the structures $(\mathbb{A}'; h') \cong (\mathbb{A}; h)$ (i.e., the expansions of \mathbb{A}' and \mathbb{A} by, respectively, h' and h). In particular, to prove our lemma, it is sufficient to prove its conclusion for $(\mathbb{A}'; h')$ rather than for (\mathbb{A}, h) . Now, $fh' = hf$ yields an isomorphism $\mathbb{A}' \rightarrow \mathbb{A}_0$. Moreover, recall that α is an isomorphism $\mathbb{A}' \rightarrow \mathbb{A}'$ such that $\alpha(\mathbb{A}_0) = \alpha_0(\mathbb{A}_0) = \mathbb{A}_1$. Thus, since fh' , α , and f^{-1} are all isomorphisms, $f^{-1}\alpha fh' := \beta \in \text{Aut}(\mathbb{A}')$. Hence,

$$\alpha \circ f \circ h' = f \circ \beta,$$

as desired. \square

Corollary 4.6. *Let $G \in \mathcal{C}_G^{\text{sat}}$, and let H be any group. Then, any homomorphism $\phi : \overline{G} \rightarrow H$ such that $\phi(G) = \{1\}$ is constant.*

Proof. Let $h \in \overline{G}$. By Lemma 4.5 there are $f \in \overline{G}$ and $\alpha, \beta \in G$ such that

$$\alpha \circ f \circ h = f \circ \beta.$$

Since ϕ is a homomorphism,

$$\phi(\alpha) \circ \phi(f) \circ \phi(h) = \phi(f) \circ \phi(\beta).$$

Since $\phi(\alpha) = \phi(\beta) = 1$ and $\phi(f) \in H$, this implies that $\phi(h) = 1$. \square

Theorem 4.7. *Let $G \in \mathcal{C}_G^{\text{sat}}$ have automatic homeomorphicity with respect to $\mathcal{C}_G^{\text{cl}}$. Then, \overline{G} has automatic homeomorphicity with respect to $\mathcal{C}_T^{\text{cl}}$.*

Moreover, suppose $\theta : \overline{G} \rightarrow \Omega^\Omega$ is an injective homomorphism and that $\theta(G)$ is closed in S_Ω . Then θ is a homeomorphism onto its image, which is an element of $\mathcal{C}_G^{\text{cl}}$.

Proof. By Proposition 4.4 and Corollary 4.6, any $\Phi \in \text{End}(\overline{G})$ that fixes every element of G is the identity function on \overline{G} . Hence \overline{G} has automatic homeomorphicity with respect to $\mathcal{C}_T^{\text{cl}}$ by Lemma 4.1. Given the first part, the proof of the second part is identical to that of [7, Lemma 4.1]. \square

Note that the same proof yields that if G has automatic homeomorphicity with respect to a subclass \mathcal{C} of $\mathcal{C}_G^{\text{cl}}$ (e.g. $\mathcal{C}_G^{\text{ol}}$), then \overline{G} has automatic homeomorphicity with respect to the subclass of those monoids in $\mathcal{C}_T^{\text{cl}}$ whose group of invertibles belongs to \mathcal{C} (e.g. $\mathcal{C}_T^{\text{ol}}$). The corollary below now follows by the same argument as in [47, Theorem 2.4]:

Corollary 4.8. *Let $G \in \mathcal{C}_G^{\text{ol}}$ be G -finite. Then \overline{G} has automatic homeomorphicity with respect to $\mathcal{C}_{\overline{G}}^{\text{cl}}$.*

4.2 Automatic homeomorphicity for endomorphism monoids

Surprisingly, automatic homeomorphicity may fail for the endomorphism monoids of structures whose automorphism groups have automatic homeomorphicity even in very tame cases. The following is an adaptation of [18, Theorem 8]. Our main contribution is that the structure in the counterexample below is extremely tame. Firstly, it is model-theoretically tame being ω -stable and a reduct of a finitely homogeneous structure. These structures are very well understood [22]. Secondly, being ω -categorical and ω -stable, its automorphism group has ample generics [30], and so our example is also tame from the point of view of reconstruction of its automorphism group.

Theorem 4.9. *There is an ω -categorical structure \mathbb{A} which is ω -stable and first-order definable from a finitely homogeneous structure such that $\text{Aut}(\mathbb{A})$ has automatic continuity with respect to \mathcal{C}_G^2 , but $\text{End}(\mathbb{A})$ (and hence also the clone $\langle \text{End}(\mathbb{A}) \rangle$) has a non-continuous automorphism. In particular, $\text{End}(\mathbb{A})$ (and $\langle \text{End}(\mathbb{A}) \rangle$) do not have automatic homeomorphicity with respect to $\mathcal{C}_T^{\text{ol}}$ (and $\mathcal{C}_C^{\text{ol}}$).*

Proof. Let \mathbb{B} be the structure with an equivalence relation E with infinitely many classes of size 2. Let \mathbb{A}' consist of two disjoint copies $\mathbb{B}_1, \mathbb{B}_2$ of \mathbb{B} plus an additional new element \star which is (only) related to itself in E , and which has additional unary predicates B_1, B_2 for the domains of $\mathbb{B}_1, \mathbb{B}_2$. It is easy to see that \mathbb{A}' is finitely homogeneous (hence, ω -categorical) and ω -stable. In particular, it has ample generics by [30] and so $\text{Aut}(\mathbb{A}')$ has automatic continuity with respect to \mathcal{C}_G^2 . Let

$$\mathcal{F} := \{f : A' \rightarrow A' \mid f(B_1) \subseteq B_2 \cup \{\star\}, f(B_2 \cup \{\star\}) = \{\star\}, f \text{ preserves } E\}.$$

Consider the closed monoid $\mathcal{T} := \overline{\text{Aut}(\mathbb{A}')} \cup \mathcal{F}$; its invertibles are precisely the elements of $\text{Aut}(\mathbb{A}')$. Let \mathcal{T}_∞ consist of those elements of \mathcal{F} which miss an infinite number of E -classes in B_2 . Let $\alpha \in \text{Aut}(\mathbb{A}')$ flip the two elements of any E -class (and fix \star). We define the map $\eta : \mathcal{T} \rightarrow \mathcal{T}$ by

$$\eta(f) := \begin{cases} f & f \notin \mathcal{T}_\infty; \\ \alpha f & f \in \mathcal{T}_\infty. \end{cases}$$

Using that α commutes with all elements of $\overline{\text{Aut}(\mathbb{A}')}$, it is easy to see that η is an automorphism of \mathcal{T} . However, taking a sequence $(f_n)_{n \in \mathbb{N}}$ outside of \mathcal{T}_∞ converging to a non-constant element g in \mathcal{T}_∞ , $(\eta(f_n))_{n \in \mathbb{N}}$ converges to g but $\eta(g) = \alpha g \neq g$. Hence η is not continuous. \square

In spite of the above negative result, there are still reasonable situations where one may hope to lift automatic homeomorphicity results from $\text{Aut}(\mathbb{A})$ to $\text{End}(\mathbb{A})$. In particular, [7, 56] developed a sophisticated technique which they use to show automatic homeomorphicity for the endomorphism monoids of (\mathbb{Q}, \leq) and some structures first-order definable in it with different automorphism groups. They make use of the small index property for $\text{Aut}(\mathbb{A})$ and some additional information on the structure of its open subgroups, which may be provided by some (possibly weak) form of elimination of imaginaries.

Their main idea is as follows: consider an isomorphism $\theta : \text{End}(\mathbb{A}) \rightarrow \mathcal{S} \in \mathcal{C}_T^{\text{cl}}$. This can be viewed as a faithful monoid action of $\text{End}(\mathbb{A})$ on Ω , the domain of \mathcal{S} ; in the following

discussion, to distinguish this action from the original action of $\text{End}(\mathbb{A})$, we write A for the domain of the original action. The aim is to directly show that θ is a homeomorphism by studying how the induced action of $G := \text{Aut}(\mathbb{A})$ on Ω can extend to θ . By the orbit-stabilizer theorem, for any $x \in \Omega$ there is a bijection between the orbit $X := \text{Orb}(x)$ of x under $G \curvearrowright \Omega$ and the left-cosets in G of the stabilizer $G_x = \{g \in G \mid \theta(g)(x) = x\}$. Since X is countable, G_x has countable index. Thus, if G has the SIP we know that G_x is open. Since the open subgroups of G are precisely those containing the pointwise stabilizer G_B of a finite set B under the action $G \curvearrowright A$, we can conclude that $G_x \supseteq G_B$ for some finite B . With additional information on the open subgroups of G , however, we can draw stronger conclusions.

We say that G has **elimination of imaginaries** if every open subgroup of G is equal to G_B for some finite B , and that it has **weak elimination of imaginaries** if every open subgroup of G is sandwiched between G_B and the setwise stabiliser $G_{\{B\}}$ for some finite B . Several ω -categorical structures have weak elimination of imaginaries, e.g., $(\mathbb{N}, =)$, the random graph, and infinite vector spaces over finite fields. It is easy to prove that if G has weak elimination of imaginaries, then for any open subgroup $H \leq G$ there is a finite algebraically closed $B \subseteq A$ and $L \leq G_{\{B\}} \curvearrowright B$ such that

$$H = G_{(B,L)} := \{f \in G \mid f|_B \in L\}.$$

Groups with the SIP and weak elimination of imaginaries are said to have the **strong small index property** (SSIP) [19].

Let us now return to analysing an isomorphism $\theta : \text{End}(\mathbb{A}) \rightarrow \mathcal{S} \in \mathcal{C}_{\mathcal{T}}^{\text{cl}}$ assuming that G has the SSIP. In that situation, for any $x \in \Omega$ we have $G_x = G_{(B,L)}$ as above. In particular, we can identify the orbit $X = \text{Orb}(x)$ with the orbit of the pair (B, L) under the natural action of G where for $g \in G$, $g(B, L) = (gB, gL)$; here, gL is a subgroup of $G_{\{gB\}} \curvearrowright gB$. Doing this for all G -orbits, we can rewrite the action of G on Ω as the action of G on a disjoint union of G -orbits of pairs $(B_i, L_i)_{i \in I}$, where I is a countable set. Using Theorem 4.7 and the proof of [7, Lemma 4.2], we can then also describe the action of \bar{G} on Ω : for any $f \in \bar{G}$ and any (gB_i, gL_i) , where $g \in G$ and $i \in I$, we obtain $f(gB_i, gL_i) = (hgB_i, hgL_i)$ where $h \in G$ is an arbitrary element which agrees with f on gB_i .

For the final step, which requires the most work, one needs to describe how $\text{End}(\mathbb{A})$ acts on the orbits of the action $G \curvearrowright \Omega$ induced by θ and deduce that this action yields that θ is a homeomorphism. Whilst this cannot always be done successfully (as demonstrated by Theorem 4.9), it already yielded substantial results for various structures with a first-order definition in (\mathbb{Q}, \leq) (see [20]; the relations in the statement below are all taken to be reflexive):

Theorem 4.10 ([7, 56]). *Let \mathbb{A} be any of: (\mathbb{Q}, \leq) , (\mathbb{Q}, sep) , $(\mathbb{Q}, \text{betw})$, $(\mathbb{Q}, \text{circ})$. Then, $\text{End}(\mathbb{A})$ has automatic homeomorphicity with respect to $\mathcal{C}_{\mathcal{T}}^{\text{cl}}$. Moreover, $\text{Pol}(\mathbb{A})$ has automatic homeomorphicity with respect to $\mathcal{C}_{\mathcal{C}}^{\text{cl}}$.*

5 On the number of Polish topologies

In this subsection, we study Polish topologies on closed oligomorphic monoids. Whilst all monoids in $\mathcal{C}_{\mathcal{G}}^{\text{ol}}$ have at least two Polish semigroup topologies, several interesting monoids in $\mathcal{C}_{\mathcal{T}}^{\text{ol}}$ have the Unique Polish Property.

5.1 Multiple Polish topologies on monoids of elementary embeddings

Lemma 5.1 ([24, Proposition 5.1]). *Let $\mathcal{S} \in \mathcal{C}_{\mathcal{T}}^{\text{cl}}$ be a monoid of injective functions whose group of invertible elements is not closed in Ω^{Ω} . Then, there are at least two distinct Polish semigroup topologies on \mathcal{S} . In particular, this is the case for any $\mathcal{S} \in \mathcal{C}_{\mathcal{G}}^{\text{ol}}$.*

Proof idea. Consider the topology \mathcal{I}_4 on the monoid of injective functions $\text{Inj}(\Omega)$ generated by τ_{pw} and the open sets $(\mathcal{W}_a)_{a \in \Omega}$ of the form

$$\mathcal{W}_a := \{f \in \text{Inj}(\Omega) \mid a \notin \text{Im}(f)\}.$$

This is a Polish topology on $\text{Inj}(\Omega)$ [25, Theorem 5.15 & Section 5.5]. It is easy to see that the induced topology τ_c on \mathcal{S} is a Polish semigroup topology in which the group of invertibles is closed, yielding that $\tau_c \neq \tau_{\text{pw}}$. \square

Similar ideas were used by [52] to prove, for every n , failure of UPP for the endomorphism monoids of the generic graph without independent sets of size n and of the complete multipartite graph with ω -many blocks of size n . Whilst not all endomorphisms of these structures are injective, there is some $m \in \mathbb{N}$ such that, for any endomorphism, the preimage of any point has size $\leq m$, and this is sufficient to build a similar topology to τ_c which is distinct from τ_{pw} .

Definition 5.2. Let $G \curvearrowright \Omega$ have locally finite algebraic closure. We say that acl forms a **pregeometry** if it satisfies the exchange property: for all $a, b \in \Omega$ and $C \subseteq \Omega$ finite, if $a \in \text{acl}(C \cup \{b\}) \setminus \text{acl}(C)$, then $b \in \text{acl}(C \cup \{a\})$.

When acl forms a pregeometry, we can adequately define the **dimension** $\dim(B)$ of a finite set $B \subseteq \Omega$ as the smallest n such that there is $C \subseteq \Omega$ with $|C| = n$ and $\text{acl}(C) = \text{acl}(B)$ [57] (we assign dimension \aleph_0 to infinite subsets of Ω). Note that acl forms a pregeometry in any oligomorphic permutation group with no algebraicity, or in the automorphism group of any vector space over a finite field.

Lemma 5.3. Let $G \curvearrowright \Omega$ be such that acl forms a pregeometry. Then, there are at least countably many Polish semigroup topologies on \overline{G} .

Proof. This is essentially the proof of [24, Theorem 5.22 (i) & (vi)]. Note that for any $f \in \overline{G}$, the image $\text{Im}(f)$ is algebraically closed. Let $\mathcal{S}_0 := \tau_c$ from the proof of Lemma 5.1. For each $n \in \mathbb{N}$ let

$$F_n := \{f \in \overline{G} \mid \dim(\Omega \setminus \text{Im}(f)) = n\}.$$

For all $n \in \mathbb{N}$, we define \mathcal{S}_n to be the topology generated by \mathcal{S}_{n-1} and F_{n-1} .

Claim 1: For each $n \in \mathbb{N}$, \mathcal{S}_n is a semigroup topology.

Proof of Claim 1. We need to prove that composition is continuous. Note that for all $f, g \in \overline{G}$,

$$\dim(\Omega \setminus \text{Im}(fg)) = \dim(\Omega \setminus \text{Im}(f)) + \dim(\Omega \setminus \text{Im}(g)).$$

In particular, for any $l \leq n$, choosing $m, k \in \mathbb{N}$ such that $m + k = l$, we get that $F_m \cdot F_k \subseteq F_l$. This yields continuity as desired. \square

Claim 2: For all $l < n$ the topologies \mathcal{S}_l and \mathcal{S}_n are distinct.

Proof of Claim 2. Firstly, using the fact that acl is a pregeometry (and Neumann's Lemma [29, Corollary 4.2.2]), it is easy to see that the sets F_l are non-empty for all $l \in \mathbb{N}$. Let $f \in F_n$ and let \mathcal{B} be an open neighbourhood of f in \mathcal{S}_n ; it suffices to show that $\mathcal{B} \neq F_n$. Since f does not belong to any F_m for any $m < n$, we must have that $\mathcal{B} \in \mathcal{S}_0$. Hence, there exist a basic open neighbourhood $\mathcal{U}_{(\bar{a}, \bar{b})} \in \tau_{\text{pw}}$ and some set

$$\mathcal{W}_{\{c_1, \dots, c_k\}} = \{f \in \overline{G} \mid \forall i \ c_i \notin \text{Im}(f)\}$$

such that $f \in \mathcal{U}_{(\bar{a}, \bar{b})} \cap \mathcal{W}_{\{c_1, \dots, c_k\}} \subseteq \mathcal{B}$. By a back & forth argument, using Neumann's Lemma and the fact that algebraic closure forms a pregeometry, it is easy to construct some $g \in \overline{G}$ such that $g \in \mathcal{B}$ but $g \notin F_n$ by building g so that it agrees with f on \bar{a} and so that its image avoids an algebraically closed subset of Ω of dimension $> n$ containing $\{c_1, \dots, c_k\}$. This implies that $F_n \neq \mathcal{B}$, as we wanted to show. \square

Claim 3: For each $n \in \mathbb{N}$, the topology \mathcal{S}_n is Polish.

Proof of Claim 3. By [33, Lemma 13.2] it is sufficient to prove that for each n , F_n is closed in \overline{G} . Note that

$$\overline{G} \setminus F_n := \{f \in \overline{G} \mid \dim(\Omega \setminus \text{Im}(f)) > n\} \cup \{f \in \overline{G} \mid \dim(\Omega \setminus \text{Im}(f)) < n\}.$$

The first of the two sets in this union is open in \mathcal{S}_0 and the second is just $\bigcup_{i < n} F_i$. Hence this set is open in \mathcal{S}_n , as desired. \square

\square

It is natural to wonder whether the technique of Lemma 5.3 can be extended to all monoids in $\mathcal{C}_G^{\text{ol}}$. We point out that there are ω -categorical structures for which every non-surjective elementary embedding has co-infinite image, meaning that one cannot define different topologies by open sets of the form of the sets F_n . In particular, using a classical construction of Hrushovski [31], one can build an ω -categorical graph such that every vertex has infinite valency and such that for any two vertices and any path of length two connecting them, the vertex in the middle of the path belongs to their algebraic closure (cf. [40, Theorem 5.1]). From this, it is easy to prove that any of its non-surjective elementary embedding has co-infinite image. Thus, we ask the following:

Question 5.4. Does every $\overline{G} \in \mathcal{C}_G^{\text{ol}}$ have infinitely many Polish semigroup topologies?

5.2 Minimality of τ_{pw} and the Zariski topology

Definition 5.5. Let \mathcal{S} be a monoid. The **semigroup Zariski topology** τ_Z , introduced in [25], is the topology on \mathcal{S} for which a subbasis is given by all sets

$$M_{\phi, \psi} := \{s \in \mathcal{S} \mid \phi(s) \neq \psi(s)\},$$

where $\phi, \psi : \mathcal{S} \rightarrow \mathcal{S}$ are functions of the form

$$s \mapsto t_k s t_{k-1} s \dots t_1 s t_0 \text{ and } s \mapsto q_l s q_{l-1} s \dots q_1 s q_0,$$

for fixed $k, l \geq 1$ and $t_0, \dots, t_k, q_0, \dots, q_l \in \mathcal{S}$.

When \mathcal{S} is a group, a potentially finer topology can be defined by allowing inverses of s to appear in the definitions of ϕ and ψ ; this topology is the **group Zariski topology**, and we shall not consider it here.

Contrary to the pointwise convergence topology τ_{pw} on a transformation monoid \mathcal{S} , the semigroup Zariski topology τ_Z is defined in purely algebraic terms rather than from the action of \mathcal{S} . In particular, any isomorphism between monoids is a homeomorphism with respect to this topology. In [25], it was pointed out that although τ_Z need not be Hausdorff or a semigroup topology, it is contained in any topology enjoying these properties; in particular, $\tau_Z \subseteq \tau_{\text{pw}}$. In fact, in [25, 24] it was shown that $\tau_Z = \tau_{\text{pw}}$ for the endomorphism monoids of several ω -categorical structures, and a general condition for this to happen was provided in [49]. An ω -categorical structure \mathbb{A} has a **mobile core** if any of its elements is contained in the image of an endomorphism into a substructure which is a model-complete core. Trivially, this is satisfied if \mathbb{A} is itself a model-complete core.

Theorem 5.6 ([49]). *Let \mathbb{A} be an ω -categorical structure without algebraicity which has a mobile core. Then $\tau_Z = \tau_{\text{pw}}$ on $\text{End}(\mathbb{A})$ if one of the following two conditions holds:*

- *the model-complete core of \mathbb{A} is finite;*
- *the model-complete core of \mathbb{A} is infinite and does not have algebraicity.*

In [49], also the first example of an ω -categorical structure \mathbb{A} for which $\tau_Z \neq \tau_{pw}$ on $\text{End}(\mathbb{A})$ was provided, answering a question from [24]. We remark that another albeit less natural example could actually be obtained from Theorem 3.5 [13]: clearly, if $\text{End}(\mathbb{A})$ and $\text{End}(\mathbb{B})$ are isomorphic but not topologically isomorphic, $\tau_{pw} \neq \tau_Z$ must hold for one of the two monoids. This example also answers positively [49, Question 4.1] asking about the existence of an ω -categorical structure with a Hausdorff semigroup topology on its endomorphism monoid which is not finer than τ_{pw} . Indeed, using a non-topological isomorphism between $\text{End}(\mathbb{A})$ and $\text{End}(\mathbb{B})$ to send τ_{pw} from one monoid onto the other, one sees that one of the two structures must be a positive example (although we do not know which one). However, the following strengthening of the conclusion of the question remains open:

Question 5.7. Is there $S \in \mathcal{C}_T^{\text{ol}}$ such that there is a Hausdorff semigroup topology on S which is strictly coarser than τ_{pw} ?

Meanwhile, vastly generalizing the example from [49], a large number of further instances of $\tau_Z \neq \tau_{pw}$ has been announced recently:

Theorem 5.8 ([42]). *Let $G \in \mathcal{C}_G^{\text{el}}$ have locally finite algebraic closure and non-trivial center. Then τ_Z is not Hausdorff on \overline{G} , and hence in particular $\tau_Z \neq \tau_{pw}$.*

At least in the case of transitive clones, minimality of τ_{pw} can be lifted from the monoid its unary functions. Call a semigroup S acting on Ω **weakly directed** if for all $a, b \in \Omega$ there are $f, g \in S$ and $c \in \Omega$ such that $f(c) = a$ and $g(c) = b$. Any transformation monoid containing a transitive permutation group is weakly directed. A clone is weakly directed if the monoid of its unary functions is.

Lemma 5.9 ([24, Lemma 6.3]). *Let $C \in \mathcal{C}_C^{\text{el}}$ be weakly directed. Suppose that τ_{pw} is minimal amongst Hausdorff semigroup topologies on the monoid of unary operations of C . Then, τ_{pw} is minimal amongst Hausdorff clone topologies on C .*

5.3 Maximality of τ_{pw} , Property X, and automatic continuity

Definition 5.10. Let S be a topological semigroup and G a subset of S . We say that S has **Property X** with respect to G if the following holds: for every $s \in S$ there are $f_s, g_s \in S$ and $t_s \in G$ such that $s = f_s t_s g_s$ and for every neighbourhood \mathcal{U} of t_s , the set $f_s(\mathcal{U} \cap G)g_s$ is a neighbourhood of s .

Theorem 5.11 ([25, Theorem 3.1]). *Let S be a topological semigroup with topology τ and $G \subseteq S$. If S has Property X with respect to G , then:*

1. *if τ is Polish and G is a Polish subgroup of S then τ is maximal among the Polish semigroup topologies for S ;*
2. *if G is a subsemigroup of S with automatic continuity with respect to a class \mathcal{C} of topological semigroups, then S has automatic continuity with respect to \mathcal{C} .*

The following notion was introduced in [21]:

Definition 5.12. A countably infinite relational structure \mathbb{A} is **homomorphism-homogeneous** (H-homogeneous) if every homomorphism between finite substructures of \mathbb{A} extends to an endomorphism of the whole structure.

Definition 5.13. A Fraïssé class \mathcal{K} in a relational language has the **strong amalgamation property with homomorphism gluing** (SAPHG) if for any $A, B_1, B_2 \in \mathcal{K}$ and embeddings $f_i : A \rightarrow B_i$ for $i \in \{1, 2\}$, there is a strong amalgam, i.e. $C \in \mathcal{K}$ and embeddings $g_i : B_i \rightarrow C$ for $i \in \{1, 2\}$ such that $g_1 \circ f_1 = g_2 \circ f_2$ and $g_1(B_1) \cap g_2(B_2) = g_1 f_1(A)$, with the property that whenever h is a map from a subset of C to some $D \in \mathcal{K}$ containing $g_1 \circ f_1(A)$ in its domain and $h \circ g_1$ and $h \circ g_2$ are homomorphisms, then h is also a homomorphism.

We remark that the SAPHG implies that the amalgam C in the definition is “canonical” in a sense that agrees with several notions of canonical amalgamation in the literature [54, 45, 32]; the SAPHG is strictly stronger than the existence of a canonical amalgam.

Theorem 5.14 ([24]). *Let \mathbb{A} be a homogeneous and H -homogeneous structure whose age has SAPHG. Then, $\text{End}(\mathbb{A})$ equipped with τ_{pw} has Property X with respect to $\text{Aut}(\mathbb{A})$.*

Example 5.15. The assumptions of Theorem 5.14 apply to the following structures:

1. the random graph \mathcal{R} ;
2. the generic directed graph;
3. the disjoint union ωK_n of countably many copies of the complete graph K_n ;
4. the generic strict poset;
5. any of the structures above including loops;
6. the generic equivalence relation with n -many countably infinite classes $E_{n,\omega}$;
7. the Cherlin-Hrushovski structure \mathbb{A}_{CH} .

All structures above except the generic strict (or non-strict) poset and \mathbb{A}_{CH} are such that their automorphism groups are known to have ample generics, and so automatic continuity with respect to $\mathcal{C}_{\mathcal{G}}^2$ [30]. With the exception of \mathbb{A}_{CH} these examples are already mentioned in [24]; the claimed properties are easy to prove for \mathbb{A}_{CH} .

[25] also show that the endomorphism monoid of the countable atomless Boolean algebra B_∞ has Property X with respect to its automorphism group, and the latter is known to have automatic continuity with respect to $\mathcal{C}_{\mathcal{G}}^2$.

The following proposition can be applied to deduce that the examples above whose automorphism group has automatic continuity with respect to $\mathcal{C}_{\mathcal{G}}^2$ are such that their endomorphism monoids have automatic continuity with respect to $\mathcal{C}_{\mathcal{T}}^2$:

Proposition 5.16 ([25, Proposition 4.1]). *Let $G \in \mathcal{C}_{\mathcal{G}}^2$. Then, G has automatic continuity with respect to $\mathcal{C}_{\mathcal{G}}^2$ if and only if it has automatic continuity with respect to $\mathcal{C}_{\mathcal{T}}^2$.*

While Property X is a powerful tool for showing maximality of τ_{pw} and lifting of automatic continuity from subgroups of a monoid, [50] note that there is some leeway in Definition 5.10: the decomposition of $s \in S$ could be of arbitrary length without changing the truth of Theorem 5.11; they call this generalization Property \bar{X} . Similarly, in the same definition s can be composed from the left with a left-invertible element of S , leading to **pseudo-Property \bar{X}** . They apply this notion to $\text{End}(\mathbb{Q}, \leq)$ equipped with τ_{rich} , a finer topology than τ_{pw} , and the subgroup $\text{Aut}(\mathbb{Q}, \leq)$ equipped with τ_{pw} to conclude that any Polish semigroup topology on $\text{End}(\mathbb{Q}, \leq)$ must be contained in τ_{rich} . By a series of further arguments, of which some can be recast as statements about pseudo-Property \bar{X} , and others involve different techniques such as Baire category, they obtain:

Theorem 5.17 ([50]). *$\text{End}(\mathbb{Q}, \leq)$ has the UPP.*

[50] note that most of the results on the UPP in Example 5.15 do not use the full strength of Polishness: they have unique second countable metrisable semigroup topologies. This occurs whenever the UPP can be shown applying the second item of Theorem 5.11. The considerably more complex proof of Theorem 5.17, which only partially fits into this framework, additionally requires the space to be Baire.

For the endomorphism monoids of some well-studied structures the status of the UPP is still open; among homogeneous graphs, the only one for which whether the UPP holds is not known is the following (see [52]):

Question 5.18. Does the endomorphism monoid of the complete multipartite graph $\mathbb{K}_{\omega,\omega}$ (with ω -many blocks of size ω) in the language of graphs have UPP?

Property X can also be applied in the context of clones. Given a topological clone \mathcal{C} with a topology τ , we can associate with it a topological semigroup $\mathcal{S}_{\mathcal{C}}$ with domain \mathcal{C} , topology τ , and the operation $*$ given by $f * g = f(g, \dots, g)$. Whilst the algebraic structure of $\mathcal{S}_{\mathcal{C}}$ is weaker than that of \mathcal{C} in the sense that not every semigroup homomorphism with respect to $*$ is a clone homomorphism, it is often sufficiently rich to restrict the possible clone topologies on \mathcal{C} :

Lemma 5.19 ([25, Lemma 7.1]). *Let \mathcal{C} be a topological clone and let $\mathcal{S}_{\mathcal{C}}$ have automatic continuity with respect to $\mathcal{C}_{\mathcal{T}}^2$. Then, \mathcal{C} has automatic continuity with respect to $\mathcal{C}_{\mathcal{T}}^2$.*

In [25], the above lemma and Property X are used to show that τ_{pw} is the unique second countable Hausdorff clone topology on the polymorphism clone of the countable atomless Boolean algebra. In [24, Theorem 6.2], it is shown that homogeneous, H-homogeneous structures \mathbb{A} with the SAPHG and whose age is closed under finite non-empty direct products are such that $\mathcal{S}_{\text{Pol}(\mathbb{A})}$ has Property X with respect to $\text{Aut}(\mathbb{A})$. This is then applied to prove UPP and automatic continuity with respect to $\mathcal{C}_{\mathcal{C}}^2$ for various structures, such as the random graph and ωK_{ω} with loops. We invite the reader to compare this result also to [46, Theorem 4.1]. In the latter theorem, the authors introduce the **amalgamated extension property** (AEP) [46, Definition 3.23], a weaker amalgamation property than the SAPHG. They show that homogeneous, H-homogeneous structures \mathbb{A} with the AEP whose age is closed under finite non-empty direct products, and whose endomorphisms contain all constant functions, are such that $\text{Pol}(\mathbb{A})$ has automatic homeomorphicity with respect to $\mathcal{C}_{\mathcal{C}}^{\text{cl}}$. The latter result is based on “gate covering” techniques of [18], but, unlike the original paper, does not rely on upwards transfers of automatic homeomorphicity from the underlying automorphism group.

5.4 More on automatic continuity and open mappings

Preceding the positive instances in Example 5.15, automatic continuity for semigroups was believed to be a less fruitful notion than for groups due to the following result:

Theorem 5.20 ([18, Corollary 10]). *No $\overline{G} \in \mathcal{C}_{\mathcal{G}}^{\text{cl}}$ has automatic continuity with respect to $\mathcal{C}_{\mathcal{T}}^{\text{cl}}$.*

The small index property for groups can be meaningfully extended to semigroups. A **right congruence** on a semigroup \mathcal{S} is an equivalence relation $\mathcal{E} \subseteq \mathcal{S} \times \mathcal{S}$ such that whenever $(g, f) \in \mathcal{E}$ and $h \in \mathcal{S}$, then $(hg, hf) \in \mathcal{E}$. We define a left-congruence analogously but with respect to composition from the other side.

Definition 5.21. A topological semigroup \mathcal{S} has the **right small index property** (rSIP) if every right congruence with countably many classes is open in $\mathcal{S} \times \mathcal{S}$. We define analogously the **left small index property** (lSIP).

Proposition 5.22 ([25, Proposition 4.3]). *A topological monoid \mathcal{S} has the rSIP if and only if it has automatic continuity with respect to $\mathcal{C}_{\mathcal{T}}^{\text{cl}}$.*

Proposition 5.23 ([25, Corollary 4.5]). *If a topological monoid \mathcal{S} has automatic continuity with respect to $\mathcal{C}_{\mathcal{T}}^2$, then it has both the rSIP and the lSIP.*

Automatic continuity with respect to $\mathcal{C}_{\mathcal{T}}^{\text{cl}}$ and with respect to $\mathcal{C}_{\mathcal{T}}^2$ are distinct notions for Polish semigroups, and even for closed oligomorphic transformation monoids [25, Proposition 5.18]. Moreover, contrasting the Open Mapping Theorem for Polish groups, automatic continuity with respect to $\mathcal{C}_{\mathcal{T}}^{\text{cl}}$ does not imply automatic homeomorphicity with respect to $\mathcal{C}_{\mathcal{T}}^{\text{cl}}$ for Polish semigroups (cf. [25, Proposition 5.15]); however, we do not know an example of this phenomenon within the class of endomorphism monoids.

Question 5.24. Is there $\mathcal{S} \in \mathcal{C}_{\mathcal{T}}^{\text{cl}}$ with automatic continuity with respect to $\mathcal{C}_{\mathcal{T}}^{\text{cl}}$ but without automatic homeomorphicity with respect to $\mathcal{C}_{\mathcal{T}}^{\text{cl}}$? If so, is there an oligomorphic example?

Regarding the openness of clone homomorphisms, we have both a lifting result from semigroups as well as a result specific to clones. The former implies that for isomorphisms from a transitive clone \mathcal{C} , it is sufficient to check openness on its unary functions.

Theorem 5.25 ([18]). *Let $\mathcal{C}, \mathcal{D} \in \mathcal{C}_C^{\text{cl}}$. Let $\xi : \mathcal{C} \rightarrow \mathcal{D}$ be a clone homomorphism and let ξ' be its restriction to unary functions.*

- *Suppose \mathcal{D} is weakly directed. If ξ' is continuous, then so is ξ ;*
- *Suppose \mathcal{C} is weakly directed and ξ is an isomorphism. If ξ' is open, so is ξ .*

The following trick for clones is not available for monoids, since homomorphisms of the latter do not necessarily preserve constant functions:

Proposition 5.26 ([18]). *Let $\mathcal{C} \in \mathcal{C}_C^{\text{cl}}$ contain all constant functions. Then, any isomorphism $\xi : \mathcal{C} \rightarrow \mathcal{D} \in \mathcal{C}_C^{\text{cl}}$ is open.*

6 Rubin's dream: stronger notions of reconstruction

The group S_Ω acts continuously on Ω^Ω by conjugation: $(\theta, f) \mapsto f^\theta := \theta f \theta^{-1}$ for $\theta \in S_\Omega$ and $f \in \Omega^\Omega$. Hence, for any closed subsemigroup \mathcal{S} of Ω^Ω and any $\theta \in S_\Omega$, we obtain a new closed subsemigroup $\mathcal{S}^\theta := \{f^\theta \mid f \in \mathcal{S}\}$ which is topologically isomorphic to it by fixing θ in the above map. We call such homeomorphisms **set-induced**. This idea leads to a stronger notion of reconstruction:

Definition 6.1. Let $\mathcal{S} \in \mathcal{C}_T^{\text{cl}}$ and $\mathcal{C} \subseteq \mathcal{C}_T^{\text{cl}}$. We say that \mathcal{S} has **automatic action compatibility** with respect to \mathcal{C} if any algebraic isomorphism from \mathcal{S} onto an element of \mathcal{C} is set-induced.

Again, the analogous notion can be defined also in the context of permutation groups and clones. In the latter case, the action on Ω^{Ω^n} is

$$(\theta, f) \mapsto f^\theta(x_1, \dots, x_n) := \theta f(\theta^{-1}(x_1), \dots, \theta^{-1}(x_n)).$$

These notions have a structural interpretation: we call structures \mathbb{A}, \mathbb{B} on Ω **(fo/e/ep/pp)-bi-definable** if they are (fo/e/ep/pp)-bi-interpretable with $d = 1$ and a bijection $I : \Omega \rightarrow \Omega$ (using the notation from Definition 2.7). In the situation where they are ep-bi-definable, for $\theta := I$ we have $\text{End}(\mathbb{B}) = \text{End}(\mathbb{A})^\theta$. Analogous statements hold for the other notions of definability. Moreover, for all of these notions the converse (i.e. the existence of a permutation θ moving one space of symmetries onto the other by conjugation) holds in the oligomorphic case. This follows from the theorem of Ryll-Nardzewski, Engeler, and Svenonius for automorphism groups, is due to [15] for pp-bi-definability, and is folklore for the other cases.

For the above form of reconstruction to hold, we need to exclude algebraicity: for any oligomorphic permutation group G we can construct a topologically isomorphic oligomorphic G' whose action is not isomorphic to that of G by adding a fixed point. Thus, the interesting classes are $\mathcal{C}_G^{\text{na}}, \mathcal{C}_G^{\text{na}}, \mathcal{C}_T^{\text{na}}$, and $\mathcal{C}_C^{\text{na}}$: spaces of symmetries of ω -categorical structures without algebraicity. In this context, we already have the following consequence of Theorem 5.6 regarding automatic homeomorphicity:

Corollary 6.2. *Let $\overline{G} \in \mathcal{C}_G^{\text{na}}$. Then, \overline{G} has automatic homeomorphicity with respect to $\mathcal{C}_G^{\text{na}}$.*

Proof. By Theorem 5.6, $\tau_Z = \tau_{\text{pw}}$ for any structure in $\mathcal{C}_G^{\text{na}}$, yielding that any isomorphism between two structures in the class is also topological. \square

However, much more is true. Namely, it is a result of [51] that any fo-bi-interpretation of \mathbb{B} in \mathbb{A} , where \mathbb{A}, \mathbb{B} are ω -categorical structures without algebraicity, gives rise to a bi-definition: this is essentially Lemma 2.11 in [51]. In some of the literature on reconstruction one gets the impression that this fact has been overlooked, perhaps due to the complex mathematical language used in Rubin's paper and the belief that the statement holds only

in the presence of a weak $\forall\exists$ -interpretation (the main topic of Rubin’s article). Be that as it may, the same statement holds for all stronger (i.e. e/ep/pp) notions of bi-interpretation by analysing Rubin’s proof, and a new proof has been announced in [27]:

Theorem 6.3. *Let $\mathcal{S}, \mathcal{T} \in \mathcal{C}_{\mathcal{T}}^{\text{na}}$. Then any topological isomorphism from \mathcal{S} onto \mathcal{T} is set-induced. Similar statements hold in $\mathcal{C}_{\mathcal{G}}^{\text{na}}$ and $\mathcal{C}_{\mathcal{C}}^{\text{na}}$.*

Theorem 6.3 has, in the case of groups, been exploited to show that the equivalence relation of topological isomorphism on the class $\mathcal{C}_{\mathcal{G}}^{\text{na}}$ of closed oligomorphic groups without algebraicity is **smooth** [43]. By [27], the same is true for monoids (i.e. the class $\mathcal{C}_{\mathcal{T}}^{\text{na}}$) and clones (i.e. the class $\mathcal{C}_{\mathcal{C}}^{\text{na}}$). Both works contain generalisations beyond the context of no algebraicity. As a consequence of Theorem 6.3, automatic homeomorphicity and automatic action compatibility coincide in the context of $\mathcal{C}_{\mathcal{G}}^{\text{na}}$. Together with Corollary 6.2 this yields:

Corollary 6.4. *Let $\overline{G} \in \mathcal{C}_{\mathcal{G}}^{\text{na}}$. Then, \overline{G} has automatic action compatibility with respect to $\mathcal{C}_{\mathcal{G}}^{\text{na}}$.*

Some specific cases of Corollary 6.4 were proven in [8]. The following proposition allows to lift being set-induced from semigroup homomorphisms to clone homomorphisms.

Proposition 6.5 ([8, Theorem 4.3]). *Let $\mathcal{C}, \mathcal{D} \in \mathcal{C}_{\mathcal{C}}^{\text{cl}}$ with \mathcal{C} being weakly directed. Let $\xi : \mathcal{C} \rightarrow \mathcal{D}$ be a surjective clone homomorphism and let ξ' be its restriction to unary functions. If ξ' is set-induced, then so is ξ .*

We can thus extend Corollary 6.4 to transitive clones.

Corollary 6.6. *Let $\mathcal{C}, \mathcal{D} \in \mathcal{C}_{\mathcal{C}}^{\text{na}}$ be model-complete core clones. If \mathcal{C} is transitive, then any clone isomorphism $\xi : \mathcal{C} \rightarrow \mathcal{D}$ is set-induced.*

7 Final remarks on clones

In this final section, we summarise some of the literature on reconstruction for polymorphism clones mentioned throughout this survey and list some final questions on the topic. Current techniques for clones usually rely on lifting reconstruction properties from the underlying group or monoid. Whilst this was originally the case also for the study of reconstruction for monoids, now we have several powerful direct techniques. We hope that the study of clones will develop in a similar direction. The original “gate covering” techniques of [18] relied on lifting to prove automatic homeomorphicity with respect to $\mathcal{C}_{\mathcal{C}}^{\text{cl}}$ of the Horn clone, the polymorphism clone of the random graph, and all minimal closed clones above the automorphism group of the random graph. Meanwhile, [7, 56] use Proposition 5.26 and other lifting tricks to deduce automatic continuity of the polymorphism clones of (\mathbb{Q}, \leq) and the other structures mentioned in Theorem 4.10. Although Property X for $\mathcal{S}_{\text{Pol}(\mathbb{A})}$ can be used to deduce UPP and automatic continuity for $\text{Pol}(\mathbb{A})$ [25, 24], current proofs of UPP do rely on lifting minimality of τ_{pw} using Lemma 5.9, and automatic continuity still relies on lifting from $\text{Aut}(\mathbb{A})$. The only existing technique that seems to not rely on lifting is [46, Theorem 4.1], which utilises gate coverings to show automatic homeomorphicity with respect to $\mathcal{C}_{\mathcal{C}}^{\text{cl}}$ for the polymorphism clones of various homogeneous structures, such as the generic (non-strict) poset.

One challenge we face is that current techniques often rely on specific structural properties such as particular forms of amalgamation. However, most structures do not enjoy these properties themselves, although in many cases they are first-order definable, or interpretable, in another structure that does. For example, amongst the continuum of closed clones whose underlying permutation group is S_{Ω} (presented in [12]), we only know automatic homeomorphicity for those containing the full monoid $\mathcal{O}^{(1)}$, for the Horn clone [18], and for $\langle S_{\Omega} \rangle$ [7].

Question 7.1. Does the polymorphism clone of every structure which is first-order definable in $(\mathbb{N}, =)$ have automatic homeomorphicity with respect to $\mathcal{C}_{\mathcal{C}}^{\text{cl}}$? In general, are there

conditions which guarantee that automatic homeomorphicity of $\text{End}(\mathbb{A})$ (or $\text{Pol}(\mathbb{A})$) for a structure \mathbb{A} imply the same for the structures first-order definable or interpretable in it?

Note that Theorem 4.9 shows that some non-trivial conditions must be imposed. We are not aware of any model-complete core clone without automatic homomorphicity whose monoid of unary functions has automatic homeomorphicity.

Of particular interest, especially in applications to Constraint Satisfaction [9, 48], is when an ω -categorical structure \mathbb{A} pp-interprets all finite structures. This is the case if and only if there is a continuous clone homomorphism from $\text{Pol}(\mathbb{A})$ to the clone $\mathcal{P}_{\{0,1\}}$ of projections on a two-element set. Naturally, one then wishes to know whether all such homomorphisms must be continuous, or whether at least the existence of a homomorphism implies the existence of a continuous one. A structure first-order definable in the Cherlin-Hrushovski structure provides a counterexample to the former statement [17]. However, the proof requires the ultrafilter lemma, and the structure is only homogenisable in an infinite language. The truth of the latter statement is unknown, cf. [17, Questions 7.3 & 7.4].

Question 7.2. Let $\mathcal{C} \in \mathcal{C}_{\mathcal{C}}^{\text{ol}}$ have a clone homomorphism to $\mathcal{P}_{\{0,1\}}$. Does it then always also have a continuous such homomorphism? Can the existence of a discontinuous homomorphism from some $\mathcal{C} \in \mathcal{C}_{\mathcal{C}}^{\text{ol}}$ to $\mathcal{P}_{\{0,1\}}$ be proven in ZF? Can it be proven for the polymorphism clone of a structure which is first-order definable in a finitely homogeneous structure?

A positive answer to the first question is made plausible by the result from [5, Theorem 1.6] that for an ω -categorical model complete core, $\text{Pol}(\mathbb{A}, c_1, \dots, c_m)$ has a continuous homomorphism to $\mathcal{P}_{\{0,1\}}$ for some constants c_1, \dots, c_m if and only if $\text{Pol}(\mathbb{A}, d_1, \dots, d_n)$ has a homomorphism to $\mathcal{P}_{\{0,1\}}$ for some constants d_1, \dots, d_n .

Acknowledgements: We thank Luna Elliott and Roman Feller for helpful comments. Funded by the European Union (ERC, POCOCOP, 101071674). Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or the European Research Council Executive Agency. Neither the European Union nor the granting authority can be held responsible for them. This research was funded in whole or in part by the Austrian Science Fund (FWF) [I 5948]. For the purpose of Open Access, the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript (AAM) version arising from this submission.

References

- [1] Gisela Ahlbrandt and Martin Ziegler. “Quasi-finitely axiomatizable totally categorical theories”. In: *Annals of Pure and Applied Logic* 30.1 (1986), pp. 63–82.
- [2] Silvia Barbina and Dugald Macpherson. “Reconstruction of Homogeneous Relational Structures”. In: *Journal of Symbolic Logic* 72.3 (2007), pp. 792–802.
- [3] S Bardyla, L Elliott, JD Mitchell, and Y Péresse. “A note on intrinsic topologies of groups”. In: *arXiv preprint arXiv:2506.11500* (2025).
- [4] Libor Barto, Michael Kompatscher, Miroslav Olšák, Trung Van Pham, and Michael Pinsker. “Equations in oligomorphic clones and the constraint satisfaction problem for ω -categorical structures”. In: *Journal of Mathematical Logic* 19.2 (2019), #1950010.
- [5] Libor Barto and Michael Pinsker. “Topology is irrelevant”. In: *SIAM Journal on Computing* 49.2 (2020), pp. 365–393.
- [6] Howard Becker and Alexander Kechris. *The Descriptive Set Theory of Polish Group Actions*. LMS Lecture Note Series 232. Cambridge University Press, 1996.
- [7] Mike Behrisch, John Truss, and Edith Vargas-García. “Reconstructing the topology on monoids and polymorphism clones of the rationals”. In: *Studia Logica* 105.1 (2017), pp. 65–91.

- [8] Mike Behrisch and Edith Vargas-García. “On a stronger reconstruction notion for monoids and clones”. In: *Forum Mathematicum* 33.6 (2021), pp. 1487–1506.
- [9] Manuel Bodirsky. *Complexity of Infinite-Domain Constraint Satisfaction*. Lecture Notes in Logic (52). Cambridge, United Kingdom; New York, NY: Cambridge University Press, 2021.
- [10] Manuel Bodirsky. “Cores of countably categorical structures”. In: *Logical Methods in Computer Science (LMCS)* 3.1 (2007), pp. 1–16.
- [11] Manuel Bodirsky. “Ramsey Classes: Examples and Constructions”. In: *Surveys in Combinatorics. London Mathematical Society Lecture Note Series* 424. Cambridge University Press, 2015.
- [12] Manuel Bodirsky, Hubie Chen, and Michael Pinsker. “The Reducts of Equality up to Primitive Positive Interdefinability”. In: *Journal of Symbolic Logic* 75.4 (2010), pp. 1249–1292.
- [13] Manuel Bodirsky, David Evans, Michael Kompatscher, and Michael Pinsker. “A counterexample to the reconstruction of ω -categorical structures from their endomorphism monoids”. In: *Israel Journal of Mathematics* 224.1 (2018), pp. 57–82.
- [14] Manuel Bodirsky and Markus Junker. “ \aleph_0 -categorical structures: interpretations and endomorphisms”. In: *Algebra Universalis* 64.3-4 (2011), pp. 403–417.
- [15] Manuel Bodirsky and Jaroslav Nešetřil. “Constraint Satisfaction with Countable Homogeneous Templates”. In: *Journal of Logic and Computation* 16.3 (2006), pp. 359–373.
- [16] Manuel Bodirsky and Michael Pinsker. “Topological Birkhoff”. In: *Transactions of the American Mathematical Society* 367 (2015), pp. 2527–2549.
- [17] Manuel Bodirsky, Michael Pinsker, and András Pongrácz. “Projective clone homomorphisms”. In: *Journal of Symbolic Logic* 86.1 (2021), pp. 148–161.
- [18] Manuel Bodirsky, Michael Pinsker, and András Pongrácz. “Reconstructing the Topology of Clones”. In: *Transactions of the American Mathematical Society* 369 (2017), pp. 3707–3740.
- [19] Peter J. Cameron. *Oligomorphic permutation groups*. Cambridge: Cambridge University Press, 1990.
- [20] Peter J. Cameron. “Transitivity of permutation groups on unordered sets”. In: *Mathematische Zeitschrift* 148 (1976), pp. 127–139.
- [21] Peter J. Cameron and Jaroslav Nešetřil. “Homomorphism-homogeneous relational structures”. In: *Combinatorics, Probability and Computing* 15.1-2 (2006), pp. 91–103.
- [22] G. Cherlin, L. Harrington, and A.H. Lachlan. “ \aleph_0 -categorical, \aleph_0 -stable Structures”. In: *Annals of Pure and Applied Logic* 28 (1985), pp. 103–135.
- [23] John Dixon, Peter M. Neumann, and Simon Thomas. “Subgroups of small index in infinite symmetric groups”. In: *Bulletin of the London Mathematical Society* 18.6 (1986), pp. 580–586.
- [24] L. Elliott, J. Jonušas, J.D. Mitchell, Y. Péresse, and M. Pinsker. “Polish topologies on endomorphism monoids of relational structures”. In: *Advances in Mathematics* 431 (2023b), p. 109214.
- [25] Luke Elliott, Julius Jonušas, Zachary Mesyan, J Mitchell, Michał Morayne, and Yann Peresse. “Automatic continuity, unique Polish topologies, and Zariski topologies on monoids and clones”. In: *Transactions of the American Mathematical Society* 376.11 (2023a), pp. 8023–8093.
- [26] David M. Evans and Paul R. Hewitt. “Counterexamples to a conjecture on relative categoricity”. In: *Annals of Pure and Applied Logic* 46.2 (1990), pp. 201–209.
- [27] Roman Feller and Michael Pinsker. “Decidability of interpretability”. announced.

- [28] Zaniar Ghadernezhad and Javier de la Nuez González. “Group topologies on automorphism groups of homogeneous structures”. In: *Pacific Journal of Mathematics* 327.1 (2024), pp. 83–105.
- [29] Wilfrid Hodges. *Model theory*. Cambridge: Cambridge University Press, 1993.
- [30] Wilfrid Hodges, Ian Hodkinson, Daniel Lascar, and Saharon Shelah. “The Small Index Property for ω -Stable ω -Categorical Structures and for the Random Graph”. In: *Journal of the London Mathematical Society* S2-48.2 (1993), pp. 204–218.
- [31] Ehud Hrushovski. *A stable \aleph_0 -categorical pseudo-plane*. Unpublished notes. 1987.
- [32] Itay Kaplan and Pierre Simon. “Automorphism groups of finite topological rank”. In: *Transactions of the American Mathematical Society* 372.3 (2019), pp. 2011–2043.
- [33] Alexander S. Kechris. *Classical descriptive set theory*. Vol. 156. Graduate Texts in Mathematics. Springer, 1995.
- [34] Alexander S. Kechris and Christian Rosendal. “Turbulence, amalgamation and generic automorphisms of homogeneous structures”. In: *Proceedings of the London Mathematical Society* 94.3 (2007), pp. 302–350.
- [35] Aleksandra Kwiatkowska. “The group of homeomorphisms of the Cantor set has ample generics”. In: *Bulletin of the London Mathematical Society* 44.6 (2012), pp. 1132–1146.
- [36] Daniel Lascar. “Autour de la propriété du petit indice”. In: *Proceedings of the London Mathematical Society* 62.1 (1991), pp. 25–53.
- [37] Daniel Lascar. “Le demi-groupe des endomorphismes d’une structure \aleph_0 -catégorique”. In: *Actes de la Journée Algèbre Ordonnée (Le Mans, 1987)* (1989), pp. 33–43.
- [38] Daniel Lascar. “On the category of models of a complete theory”. In: *The Journal of Symbolic Logic* 47.2 (1982), pp. 249–266.
- [39] Dugald Macpherson. “A survey of homogeneous structures”. In: *Discrete Mathematics* 311.15 (2011), pp. 1599–1634.
- [40] Paolo Marimon. “Invariant Keisler measures for ω -categorical structures”. In: *The Journal of Symbolic Logic* 90.2 (2025), pp. 791–807.
- [41] Antoine Mottet and Michael Pinsker. “Cores over Ramsey structures”. In: *Journal of Symbolic Logic* 86.1 (2021), pp. 352–361.
- [42] Javi de la Nuez Gonzalez, Zaniar Ghadernezhad, Paolo Marimon, and Michael Pinsker. “Minimal and intrinsic topologies on monoids of elementary embeddings”. announced.
- [43] Gianluca Paolini and Andre Nies. “Two classes of oligomorphic groups with smooth topological isomorphism relation”. In: *arXiv preprint* (2024), arXiv–2410.
- [44] Gianluca Paolini and Saharon Shelah. “Reconstructing Structures with the Strong Small Index Property up to Bi-Definability”. In: *Fundamenta Mathematicae* 247 (2019), pp. 25–35.
- [45] Gianluca Paolini and Saharon Shelah. “The Strong Small Index Property For Free Homogeneous Structures”. In: *Submitted* (2018). Preprint available at ArXiv 1703.10517.
- [46] Christian Pech and Maja Pech. “Polymorphism clones of homogeneous structures: gate coverings and automatic homeomorphicity”. In: *Algebra universalis* 79.2 (2018), p. 35.
- [47] Christian Pech and Maja Pech. “Reconstructing the Topology of the Elementary Self-embedding Monoids of Countable Saturated Structures”. In: *Studia Logica* 106.3 (2018), pp. 595–613.
- [48] Michael Pinsker. “Current challenges in infinite-domain constraint satisfaction: Dilemmas of the infinite sheep”. In: *2022 IEEE 52nd International Symposium on Multiple-Valued Logic (ISMVL)*. IEEE. 2022, pp. 80–87.

- [49] Michael Pinsker and Clemens Schindler. “On the Zariski topology on endomorphism monoids of omega-categorical structures”. In: *The Journal of Symbolic Logic* (2023), pp. 1–19.
- [50] Michael Pinsker and Clemens Schindler. “The semigroup of increasing functions on the rational numbers has a unique Polish topology”. In: *arXiv preprint arXiv:2305.04921* (2023).
- [51] Matatyahu Rubin. “On the reconstruction of ω -categorical structures from their automorphism groups”. In: *Proceedings of the London Mathematical Society* 3.69 (1994), pp. 225–249.
- [52] Clemens Schindler. “Semigroup topologies on endomorphism monoids of omega-categorical structures”. PhD thesis. Technische Universität Wien, 2023.
- [53] Katrin Tent and Martin Ziegler. *A course in model theory*. Lecture Notes in Logic. Cambridge University Press, 2012.
- [54] Katrin Tent and Martin Ziegler. “On the isometry group of the Urysohn space”. In: *Journal of the London Mathematical Society* 87.1 (2013), pp. 289–303.
- [55] John Truss. “Infinite permutation groups. II. Subgroups of small index”. In: *Journal of Algebra* 120.2 (1989), pp. 494–515.
- [56] John Truss and Edith Vargas-García. “Reconstructing the topology on monoids and polymorphism clones of reducts of the rationals”. In: *Contributions to Discrete Mathematics* 16.2 (2021), pp. 1–22.
- [57] Neil White. *Theory of matroids*. 26. Cambridge University Press, 1986.
- [58] Ernst Witt. “Über die Kommutatorgruppe kompakter Gruppen”. In: *Rendiconti di Matematica e delle sue Applicazioni. Serie V*. 14 (1954), pp. 125–129.