

Structural and Spectral Properties of Strictly Interval Graphs

Lilian Markenzon¹, Claudia M. Justel²

¹ PPGI - Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil.

² PGSC - Departamento de Engenharia de Computação, Instituto Militar de Engenharia, Rio de Janeiro, Brazil.

Abstract

In this paper we deal with a subclass of chordal graphs, which are simultaneously strictly chordal and interval, the *strictly interval graphs*. We present a new characterization of the class that leads to a simple linear recognition algorithm. Next we introduce a new subclass of strictly interval graphs, the *SI-core* graphs, that are non-split and non-cograph graphs and show that several elements of the new class are Laplacian integral.

Keywords: strictly interval graphs, Laplacian eigenvalues.

2000 MSC: 05C50, 05C75

1. Introduction

Strictly chordal graphs (also called block duplicate graphs), a subclass of chordal graphs, were first introduced by [7]; in this paper a forbidden subgraph characterization was given by the authors. The same class was defined in terms of hypergraphs in [10]. In 2010, [3] characterized strictly chordal graphs by their critical clique graph structure. This characterization leads to several structural properties of the class.

In this paper we deal with a subclass of chordal graphs, which are simultaneously strictly chordal and interval, the *strictly interval graphs*. They were introduced by Markenzon and Waga [15], who proved that they are (2-net, bipartite claw)-free (see Figure 1).

In this paper we focus on the relation between classic invariants of graphs and their integer Laplacian eigenvalues. It is well known in the literature

the quest for classes of graphs with elements which are Laplacian integral graphs. The cographs are the most important example of these classes, but the quest goes on. We present here a new characterization of strictly interval graphs by using their critical clique graph that leads to a simpler recognition algorithm. Next we introduce a subclass of strictly interval graphs, the *SI*-core graphs. Based on their structure, we show that several elements of the class are Laplacian integral.




Figure 1: 2-net and bipartite claw graphs

2. Graph Basic Concepts

All graphs mentioned in this paper are connected graphs.

Basic concepts about chordal graphs are assumed to be known and can be found in Blair and Peyton [2] and Golumbic [6]. Let $G = (V, E)$ be a graph, with $|E| = m$, $|V| = n > 0$. The *set of neighbors* of a vertex $v \in V$ is denoted by $N(v) = \{w \in V \mid \{v, w\} \in E\}$ and its *closed neighborhood* by $N[v] = N(v) \cup \{v\}$. Two vertices u and v are *true twins* in G if $N[u] = N[v]$ and they are *false twins* in G if $N(u) = N(v)$. A vertex v is said to be *simplicial* in G if $N(v)$ is a clique in G . Given a chordal graph G , the *derived graph* $D(G)$ is the induced subgraph obtained by deleting all the simplicial vertices of G .

A subset $S \subset V$ is a *vertex separator* for non-adjacent vertices u and v (a *uv*-separator) if the removal of S from the graph separates u and v into distinct connected components. If no proper subset of S is a *uv*-separator then S is a *minimal uv*-separator. When the pair of vertices remains unspecified, we refer to S as a *minimal vertex separator*.

A *clique-tree* of G is defined as a tree T whose vertices are the maximal cliques of G such that for every two maximal cliques Q and Q' each clique in the path from Q to Q' in T contains $Q \cap Q'$. The set of maximal cliques of G is denoted by \mathbb{Q} . The *leafage* of a chordal graph G , $\ell(G)$, is the minimum number of leaves among the clique-trees of G . Blair and Peyton [2] proved that, for a

clique-tree $T = (V_T, E_T)$, a set $S \subset V$ is a minimal vertex separator of G if and only if $S = Q' \cap Q''$ for some edge $\{Q', Q''\} \in E_T$. Moreover, the multiset \mathbb{M} of the minimal vertex separators of G is the same for every clique-tree of G . The algorithm presented in Markenzon and Pereira [13] computes the set of minimal vertex separators, \mathbb{S} , of a chordal graph G in linear time.

It is worth mentioning special kinds of cliques. A *simplicial clique* is a maximal clique containing at least one simplicial vertex. A simplicial vertex appears in only one maximal clique of the graph. A simplicial clique Q is called a *boundary clique* if there exists a maximal clique Q' such that $Q \cap Q' = X$ is the set of non-simplicial vertices of Q . Shibata [18] showed that if a maximal clique Q is a leaf in some clique-tree then Q is a boundary clique. The converse of this fact was proved by Hara and Takemura [8]. These results can be summarized in the next theorem.

Theorem 1. [8] *A maximal clique is a leaf in some clique-tree if and only if it is a boundary clique.*

A maximal clique such that all vertices are true twins is called a *critical clique* [12]. The set \mathbb{C} of critical cliques of G forms a partition of vertices of V .

The *critical clique graph* of a chordal graph G , $CC(G)$, is the connected graph whose vertices are the critical cliques of G and whose edges connect vertices corresponding to cliques Q_i and Q_j such that some vertex of Q_i is neighbor in G of a vertex in Q_j . Note that $CC(G)$ has no true twins [3], then the max cliques of $CC(G)$ have at most one simplicial vertex.

3. Subclasses of chordal graphs

In this section we present the definitions and some characterizations of the subclasses of chordal graphs that somehow will be used in this paper.

An *interval graph* is the intersection graph of a family I of intervals on the real line. There are several characterizations of this class. The following characterization theorem guarantees that for an interval graph, its maximal cliques can be linearly ordered.

Theorem 2. [5] *Let $G = (V, E)$ be a non-complete chordal graph. Then, G is an interval graph if and only if G has a clique-tree that is a path.*

Strictly chordal graphs, defined in [10] based on hypergraph properties, were firstly introduced as *block duplicate graphs* by Golombic and Peled [7]. A *strictly chordal graph* is a graph obtained by adding zero or more true twins to each vertex of a block graph G (a graph is a *block graph* [9] if it is connected and every block is a clique). The class was proved to be chordal, gem-free and dart-free ([7], [10]), see Figure 2.



Figure 2: Gem and dart graphs.

Theorem 3. [3] *Let G be a connected graph. G is strictly chordal iff $CC(G)$ is a block graph.*

Strictly chordal graphs can also be characterized in terms of the structure of their minimal vertex separators.

Theorem 4. [14] *Let $G = (V, E)$ be a chordal graph. The following statements are equivalent:*

1. G is a strictly chordal graph.
2. For any distinct $S, S' \in \mathbb{S}$, $S \cap S' = \emptyset$.
3. G is a $\{\text{gem}, \text{dart}\}$ -free graph.

Corollary 4.1. *Let $G = (V, E)$ be a strictly chordal graph and \mathbb{S} its set of minimal vertex separators.*

- (a) $S \in \mathbb{S}$ is a minimal separator of G .
- (b) S belongs to exactly $\mu(S) + 1$ maximal cliques of G , for every $S \in \mathbb{S}$.
- (c) boundary cliques of G contain only one mvs.

Based on this characterization, the recognition algorithm for strictly chordal graphs is extremily simple and relies only on the determination of \mathbb{S} .

The critical clique graph of strictly chordal graphs have interesting properties.

Lemma 5. [12] *Let $G = (V, E)$ a chordal graph. Then, the set of critical cliques in G forms a partition of V .*

By Theorem 4, the proof of the next lemma becomes immediate.

Lemma 6. *Let G be a strictly chordal graph. A critical clique of G is composed by:*

1. *all the simplicial vertices of a maximal clique of G or*
2. *all the vertices of a minimal vertex separator of G .*

4. Strictly Interval Graphs: new characterization and recognition algorithm

In this section we restudy a subclass of interval graphs, the *strictly interval graphs* (*SIG*), graphs which are at the same time strictly chordal and interval graphs. A first characterization of the class was presented in [15]. We present structural results about the class, leading to a new characterization and a simple recognition algorithm.

Theorem 7. [15] *Let $G = (V, E)$ be a non-complete strictly chordal graph. The following statements are equivalent:*

1. *G is a strictly interval graph.*
2. *G is a $\{2\text{-net, bipartite claw}\}$ -free graph (see Fig. 1).*
3. *$\text{leafage}(G) = 2$.*

Corollary 7.1. *Let G be a non-complete strictly interval graph. Then every maximal clique of G contains at most two minimal vertex separators.*

As it was seen in Theorem 3, strictly chordal graphs are characterized by their critical clique graphs. The next theorem presents a new characterization of strictly interval graphs, also based in their critical clique graphs, that supports a more efficient recognition algorithm.

Theorem 8. *Let G be a non-complete strictly chordal graph. G is a strictly interval graph iff $D(CC(G))$ is a path.*

Proof. \rightarrow Let G be a strictly interval graph. By Theorem 3, $CC(G)$ is a block graph. So, all mvs of G are represented by one vertex in $CC(G)$.

In $D(CC(G))$ there are not simplicial vertices of G . Suppose that S_1, S_2 and S_3 , mvs of G , form a path $P = [s_1, s_2, s_3]$ in $D(CC(G))$. Let us consider a vertex s' that does not belong to this path but, together with P , establish a connected induced subgraph of $D(CC(G))$. Vertex s' can be linked to P in three ways:

1. forming the edge (s', s_1) (or (s', s_3)): in this case it forms a path in $D(CC(G))$.
2. forming the edge (s', s_2) : in this case G has a bipartite claw.
3. with two edges, forming a triangle: in this case G has a 2-net.
4. with three edges: in this case G has a dart and it is not strictly chordal.

Item 1 satisfies the hypothesis. With itens 2, 3 and 4, by Theorem 7, we do not have a strictly interval graph. So, $D(CC(G))$ is a path.

\leftarrow Let $D(CC(G))$ be a path. Since G is strictly chordal, we need to prove that G is an interval graph, or, equivalently, G has a clique-tree that is a path (Theorem 2) or the maximal cliques of G can be linearly ordered. Since $D(CC(G))$ is a path, the maximal cliques of G represented in $D(CC(G))$ admit a linear ordering.

By Lemma 6, each mvs of G establish a maximal clique of $CC(G)$. Each edge of $D(CC(G))$ belongs to a maximal clique of G . So, these maximal cliques also form a path in the clique tree.

Let us consider the addition of simplicial vertices of $CC(G)$ to $D(CC(G))$. This addition produces three different cases:

- the construction of boundary cliques of $CC(G)$, containing the minimal vertex separators that are vertices of $D(CC(G))$. It does not matter how many boundary cliques contain the same mvs ; they will appear sequentially in the clique-tree of G and the resulting graph is an interval graph.
- The addition of a simplicial vertex adjacent to an edge of $D(CC(G))$; it results in a max clique with three vertices, one of them being a simplicial vertex. In the clique-tree, this vertex corresponds to an increase

of the cardinality of one maximal clique and it does not affect the clique-tree. A new addition over this same edge of $D(CC(G))$ cannot be performed: the new vertex cannot be a true twin of the first vertex added because the max cliques of $CC(G)$ have at most one simplicial vertex. It also cannot be a false twin because the result would be a new maximal clique producing a dart in G - impossible since G is a strictly chordal graph.

- two (or more) false twins as simplicial vertices adjacent to a maximal clique results in a dart induced subgraph in G . It is not a valid addition.

Therefore, there are only two possible additions of simplicial vertices to $D(CC(G))$ (by hypothesis, a path) and both maintain the linear order of the maximal cliques of G . Then, G is an interval graph and the result follows. \square

Theorem 8 provides directly a recognition algorithm. The steps are:

Step 1: Recognition of G as strictly chordal;

Step 2: Determination of $D(CC(G))$:

- Lemma 6 provides a simple determination of the vertices of $CC(G)$;
- removal of simplicial vertices of $CC(G)$;

Step 3: Test if the result is a path.

It is important to highlight that all these steps are trivially simple to implement and have linear time complexity.

5. Spectral Basic Concepts

The *Laplacian matrix* of a graph G is defined as $L(G) = \mathcal{D}(G) - A(G)$, where $\mathcal{D}(G)$ denotes the diagonal degree matrix and $A(G)$ the adjacency matrix of G . As $L(G)$ is symmetric, there are n real eigenvalues and as $L(G)$ is positive semidefinite, all these eigenvalues are non-negatives and 0 is an eigenvalue. We denote the eigenvalues of $L(G)$, the *Laplacian eigenvalues* of G , by $\mu_1(G) \geq \mu_2(G), \dots, \geq \mu_n(G) = 0$.

All different Laplacian eigenvalues of G together with their multiplicities form the *Laplacian spectrum* of G , denoted by $\text{Spec}L(G) = [\bar{\mu}_1^{m_1}, \dots, \bar{\mu}_d^{m_d}]$, where $1 \leq d \leq n$ is the number of distinct eigenvalues of $L(G)$ and m_i the multiplicity of $\bar{\mu}_i$, $1 \leq i \leq d$. A graph is called *Laplacian integral* if $\text{Spec}L(G)$ consists of integers.

The next results about linear algebra will be used later. Given a $n \times p$ matrix A and a $m \times q$ matrix B , the *Kronecker product* (or tensor product) $A \otimes B$ is the $nm \times pq$ block matrix:

$$A \otimes B = \begin{pmatrix} a_{1,1}B & a_{1,2}B & \dots & a_{1,p}B \\ \vdots & \vdots & & \vdots \\ a_{n,1}B & a_{n,2}B & \dots & a_{n,p}B \end{pmatrix}$$

Let M be a real $n \times n$ matrix in the following block form

$$M = \begin{pmatrix} M_{1,1} & \dots & M_{1,t} \\ \vdots & \ddots & \vdots \\ M_{t,1} & \dots & M_{t,t} \end{pmatrix}$$

where $M_{i,j}$ are $n_i \times n_j$ matrices, $1 \leq i, j \leq t$ and $n = \sum_{r=1}^t n_r$. If $b_{i,j}$ the average row sum of $M_{i,j}$, then $B = B(M) = (b_{i,j})$ is the *quotient matrix* of M . Moreover, if for each pair i, j , $M_{i,j}$ has constant row sum, then B is the *equitable quotient matrix* of M . You et al. [19] showed the following result.

Theorem 9. [19] *Given a matrix M and B a equitable quotient matrix of M , the spectrum of B is included in the spectrum of M ($\text{Spec}B \subseteq \text{Spec}M$).*

6. A new subclass: *SI*-core graphs

We define a new subclass of strictly chordal graphs, the *strictly interval-core graphs*, or simply *SI-core graphs*.

G is a *SI-core graph* iff it is a strictly interval graph with exactly two minimal vertex separators; these separators have the same cardinality s , $s \geq 2$, their union defines a maximal clique of size $2s$ and each separator has the same number $p \geq 2$ of simplicial vertices adjacent to it. Given fixed integers $s \geq 2$ and $p \geq 2$, we denote $\mathcal{G}(s, p)$ the set of *SI-core graphs* with these parameters. Figure 3 shows an example of a *SI-core graph* $G \in \mathcal{G}(2, 3)$.




Figure 3: A *SI*-core graph in $\mathcal{G}(2, 3)$




Figure 4: The *min* and *max* *SI*-core graphs in $\mathcal{G}(2, 3)$

For the new class we will show that an element can have many integer Laplacian eigenvalues and even be an integral graph. It is interesting to observe that the class of *SI*-core graphs is P_5 -free, it is not a subclass of cographs and it is not a subclass of split graphs. Actually the *SI*-core graph is a multi-core graph [16].

Two special graphs belonging to $\mathcal{G}(s, p)$ can be considered: the graphs with minimum and maximum number of edges, for now on denoted *min* *SI*-core graph, $G_{min}(s, p)$, and *max* *SI*-core graph, $G_{max}(s, p)$.

Figure 4 shows examples of the *min* and the *max* *SI*-core graphs in $\mathcal{G}(2, 3)$. Observe that the *min* *SI*-core graph in Figure 4 is the "carioca" graph, that belongs to the class defined by Kirkland et al. in [11], restudied in [1] as (k, t) -split graphs. The *min* *SI*-core graph of $\mathcal{G}(s, p)$, for any s, p , constitute the intersection between the two classes.

Property 10. *Let s, p be fixed integer values. The min *SI*-core graph has $\frac{s(s-1)}{2} + 2ps$ edges; the max *SI*-core graph has $\frac{s(s-1)}{2} + 2ps + p(p-1)$ edges.*

Observe that it is possible to establish, for s and p fixed integers, the number of non-isomorphic *SI*-core graphs. In order to do that we need the concept of the partition function $a(n)$.

A *partition of a positive integer n* , also called an integer partition, is a way of writing n as a sum of positive integers. The number of partitions of n is given by the partition function $a(n)$, entry A000041 in [17].

Theorem 11. *Let s, p be fixed integer values. There are $\frac{a(p)(a(p)+1)}{2}$ non-isomorphic SI -core graphs.*

Proof. Let $G \in \mathcal{G}(s, p)$. Each partition of the p simplicial vertices adjacent to a mv s can be combined with a partition of the other p simplicial vertices. Hence, with the first partition we have p different combinations. With the second partition we have $p-1$ different combinations and so on. So, the number of combinations without repetition is given by the sum of the elements of an arithmetic progression from 1 to p . \square

From the results above it is not difficult to see that there are SI -core graphs for any even number of vertices greater or equal 8. Moreover, we can determine the number of elements of the chosen class. For instance, there are 66 non-isomorphic SI -core graphs in $\mathcal{G}(4, 6)$.

7. Laplacian spectrum of SI -core graphs

Theorem 12. *Let $G_{min}(s, p)$ be the min SI -core graph in $\mathcal{G}(s, p)$. Then the number of non integer Laplacian eigenvalues of $G_{min}(s, p)$ with $s \geq 2$ and $p \geq 2$, ℓ , is $\ell = 0$ or $\ell = 2$.*

Proof. Let $G(s, p)$ be the min SI -core graph in $\mathcal{G}(s, p)$. Then, $L(G_{min}(s, p))$ can be written as

$$\begin{pmatrix} (2s+p)\mathbf{I}_{2s} - \mathbf{J}_{2s} & -\mathbf{X} \\ -\mathbf{X}^T & \mathbf{J}_{2p} \end{pmatrix},$$

where $\mathbf{X} = \mathbf{I}_2 \otimes \mathbf{J}_{s,p}$ and \otimes denotes the Kronecker product of matrices.

It is easy to conclude that $2s+p$ is an eigenvalue of $L(G_{min}(s, p))$ with multiplicity at least $2(s-1)$ and s is an eigenvalue of $L(G_{min}(s, p))$ with multiplicity at least $2(p-1)$ (eigenvalues corresponding to true and false twins).

By Theorem 9, the spectrum of the 4×4 equitable quotient matrix M is included in the spectrum of $L(G_{min}(s, p))$:

$$M = \begin{pmatrix} (2s+p)\mathbf{I}_2 - \mathbf{J}_2 & -p\mathbf{I}_2 \\ -s\mathbf{I}_2 & s\mathbf{J}_2 \end{pmatrix}.$$

Observe that the eigenvalues of M are:

- 0, corresponding to eigenvector $[\mathbf{1}_2, \mathbf{1}_2]$;
 - $s + p$, corresponding to eigenvector $[p\mathbf{1}_2, -s\mathbf{1}_2]$;
 - λ , corresponding to eigenvector $[\alpha z, \beta z]$, where z is ortogonal to vector $\mathbf{1}_2$ and λ is an eigenvalue of matrix M_1 correspndig to eigenvector $[\alpha, \beta]$,
- $$M_1 = \begin{pmatrix} (2s + p) & -p \\ -s & s \end{pmatrix}.$$

Matrix M_1 has eigenvalues given by $\frac{2s+p+s \pm \sqrt{(2s+p-s)^2 + 4sp}}{2}$. Since $2s + p + s$ e $2s + p - s$ are stricly positive integer numbers with the same parity, M_1 matrix eigenvalues are integer if and only if $(2s + p - s)^2 + 4ps$ is a perfect square. Then we can conclude that all eigenvalues of $L(G_{min}(s, p))$ are integer numbers if and only if $(2s + p - s)^2 + 4ps$ is a perfect square ($\ell = 0$) or $L(G_{min}(s, p))$ has exactly two non integer Laplacian eigenvalues ($\ell = 2$). \square

Corollary 12.1. *Let $G_{min}(s, p)$ be the min SI-core graph in $\mathcal{G}(s, p)$. The Laplacian spectrum of $G_{min}(s, p)$ with $s \geq 2$ and $p \geq 2$ is*

$$\left[\frac{3s + p + \sqrt{(s + p)^2 + 4sp}}{2}; (2s + p)^{2(s-1)}; (s + p); s^{2(p-1)}; \frac{3s + p - \sqrt{(s + p)^2 + 4sp}}{2}; 0 \right].$$

$G_{min}(s, p)$ is Laplacian integral when $(s + p)^2 + 4sp$ is a perfect square. For instance, $G_{min}(2, 3)$, $G_{min}(3, 10)$, $G_{min}(4, 6)$ are Laplacian integral. Next, by using the concepts given in [4], it will be proved that if for fixed $s, p \geq 2$, $G_{min}(s, p)$ is Laplacian integral, then all SI-core graphs in $\mathcal{G}(s, p)$ are Laplacian integral.

Given a connected graph $G = (V(G), E(G))$, let (F, S) be a cluster in G , where F is the set of false twin vertices and S is the set of the shared neighbors. Let $H = (V(H), E(H))$ be any graph such that $V(H) = F$. The graph $G(H) = (V(G(H)), E(G(H)))$ is defined having $V(G(H)) = V(G)$ and $E(G(H)) = E(G) \cup E(H)$.

Now, let be a connected graph G with a family of $k \geq 1$ pairwise disjoint clusters $(F_1, S_1), \dots, (F_k, S_k)$, with $|S_j| = s_j$, $|F_j| = c_j$, $1 \leq j \leq k$, and consider a family of graphs H_1, \dots, H_k with $|V(H_j)| = c_j$ for all j . The graph $G(H_1, \dots, H_k)$ is obtained from G by adding the edges in H_1, \dots, H_k as before. The Laplacian eigenvalues of H_1, \dots, H_k are denoted by $\mu_1(H_j) \geq \dots \geq \mu_{c_j}(H_j) = 0$, for $j = 1, \dots, k$.

Corollary 4.3 in [4] shows that the Laplacian eigenvalues of $G(H_1, \dots, H_k)$ remain the same, independently of the graphs H_1, \dots, H_k , with the exception of $|F_1| + \dots + |F_k| - k$ of them, and

$$\det(\lambda I - L(G(H_1, \dots, H_k))) = p_L(\lambda) \prod_{j=1}^k (\prod_{i=1}^{c_j-1} (\lambda - (s_j + \mu_i(L(H_j))))).$$

Therefore, $s_j + \mu_i(H) \in \text{spec} L(G(H_1, \dots, H_k))$, for $1 \leq j \leq k$ and $1 \leq i \leq c_j - 1$ and the remaining eigenvalues of $G(H_1, \dots, H_k)$ are the roots of the polynomial $p_L(\lambda)$ of degree $n - \sum_{j=1}^k c_j + k$, where $\det(\lambda I - L(G)) = p_L(\lambda) \prod_{j=1}^k (\lambda - s_j)^{c_j-1}$.

Lemma 13. *Given $s, p \geq 2$, let $G(s, p) \neq G_{\min}(s, p)$ be a SI -core graph in $\mathcal{G}(s, p)$, and H the subgraph of G induced by $E(H) = E(G(s, p)) \setminus E(G_{\min}(s, p))$. Then it exists $H = H_1 \cup H_2$ such that $|V(H_1)| = |V(H_2)| = p$ and $G(s, p) = G_{\min}(s, p)(H_1, H_2)$.*

Proof. Since $G(s, p) \neq G_{\min}(s, p)$, by Property 10 it can be concluded that

$$\frac{s(s-1)}{2} + 2ps < |E(G(s, p))| \leq \frac{s(s-1)}{2} + 2ps + p(p-1).$$

Moreover, by definition of SI -core graphs, G has $2p$ simplicial vertices, p of them adjacent to each mvs . As simplicial vertices can be true twins or false twins, we can conclude that the edges in $E(H)$ connect true twins corresponding to simplicial vertices adjacent to one minimal vertex separator of $G(s, p)$. Then, it exist $H = H_1 \cup H_2$ where $H_j = \cup_{l=p_1^j, \dots, p_{r_j}^j} K_l$, and $p_1^j, \dots, p_{r_j}^j$ is a partition of p for $j = 1, 2$, such that $G(s, p) = G_{\min}(H_1, H_2)$. \square

Theorem 14. *Given $s, p \geq 2$, let $G(s, p) \neq G_{\min}(s, p)$ be a SI -core graph in $\mathcal{G}(s, p)$. Then the number of non integer Laplacian eigenvalues of $G(s, p)$, ℓ , is $\ell = 0$ or $\ell = 2$.*

Proof. Consider $G_{\min}(s, p)$ in $\mathcal{G}(s, p)$, $s \geq 2$ and $p \geq 2$. Let H be the graph in the hypothesis of Lemma 13. By using the notation in [4], $G_{\min}(s, p)$ has 2 clusters (F_1, S_1) and (F_2, S_2) where $|F_1| = |F_2| = p$ and $|S_1| = |S_2| = s$ are the number of false twins and the number of shared neighbors (mvs), respectively. Furthermore, by Lemma 13, $G_{\min}(s, p)(H_1, H_2)$ with $H_1 \cup H_2 = H$, and each H_j is the union of the complete graphs K_l , $l = p_1^j, \dots, p_{r_j}^j$ containing simplicial vertices adjacent to the minimal vertex separator S_j , $j = 1, 2$. By

using Corollary 4.3 in [4] for $G_{\min}(s, p)(H_1, H_2)$, its Laplacian eigenvalues coincide with all but $|F_1| + |F_2| - 2 = 2(p - 1)$ eigenvalues of $G_{\min}(s, p)$. Among the new eigenvalues, when $p_i^j \geq 2$, $(s_j + p_i^j)$ has multiplicity $(p_i^j - 1)$; the remaining eigenvalues are equal to s_j . Then, the number of non integer Laplacian eigenvalues of $G(s, p)$ is the same of that for $G_{\min}(s, p)$ and the result follows. \square

Corollary 14.1. *Given $s, p \geq 2$, let $G_{\max}(s, p)$ be the max SI-core graph in $\mathcal{G}(s, p)$. The Laplacian spectrum of $G_{\max}(s, p)$ is*

$$\left[\frac{3s + p + \sqrt{(s + p)^2 + 4sp}}{2}; (2s + p)^{2(s-1)}; s + p; (s + p)^{2(p-1)}; \frac{3s + p - \sqrt{(s + p)^2 + 4sp}}{2}; 0 \right].$$

Proof. Since $G_{\max}(s, p)$ verifies the hypothesis of Theorem 14, and $G_{\max}(s, p) = G_{\min}(s, p)(H_1, H_2)$ where $H_1 = H_2 = K_p$ as defined below, the only Laplacian eigenvalues of $G_{\max}(s, p)$ different from those of $G_{\min}(s, p)$ are $(s + p)^{2(p-1)}$. By Corollary 12.1, the result follows. \square

Corollary 14.2. *Given $s, p \geq 2$, let $G(s, p) \neq G_{\min}(s, p)$ and $\neq G_{\max}$ be a SI-core graph in $\mathcal{G}(s, p)$ and H the subgraph of G induced by $E(H) = E(G(s, p)) \setminus E(G_{\min}(s, p))$. The Laplacian spectrum of $G(s, p)$ is*

$$\left[\frac{3s + p + \sqrt{(s + p)^2 + 4sp}}{2}; (2s + p)^{2(s-1)}; s + p; (s + \mu_1(L(H))); \dots; (s + \mu_{2(p-1)}(L(H))); \frac{3s + p - \sqrt{(s + p)^2 + 4sp}}{2}; 0 \right],$$

where $H = H_1 \cup H_2$, $H_j = \cup_{l=p_1^j, \dots, p_{r_j}^j} K_l$, with $p_1^j, \dots, p_{r_j}^j$ a partition of p for $j = 1, 2$, such that $G(s, p) = G_{\min}(H_1, H_2)$, and $\mu_i(L(H))$, $1 \leq i \leq 2(p - 1)$ are all but two smallest eigenvalues of $L(H)$.

Proof. By Lemma 13, it exists $H = H_1 \cup H_2$ such that $G(s, p) = G_{\min}(H_1, H_2)$. By the proof of Theorem 14, the Laplacian eigenvalues are determined and the Corollary follows. \square

8. Combinatorial and Spectral Results

In this section we present some structural properties of the SI -core graphs, establishing conditions for defining a multi-set Z of real numbers that correspond to the spectrum of the graph. The next property follows from Corollaries 12.1 and 14.1.

Property 15. *Let s, p be fixed integer values.*

- *The Laplacian eigenvalues $\mu_1, \mu_2, \dots, \mu_{2s-1}, \mu_{2s}, \mu_{n-1}, \mu_n$ are the same for any graph $G \in \mathcal{G}(s, p)$.*
- *For $2s + 1 \leq i \leq n - 2$ the Laplacian eigenvalues μ_i are integers and they belong to the interval $[s, s + p]$ for any graph $G \in \mathcal{G}(s, p)$.*

Let us consider an ordered multi-set with n elements

$$Z = \{c_1, d_1, \dots, d_{2(s-1)}, c_2, x_1, \dots, x_{p-1}, y_1, \dots, y_{p-1}, c_3, c_4\}.$$

Considering two subsets $X = \{x_1, \dots, x_{p-1}\}$, and $Y = \{y_1, \dots, y_{p-1}\}$ and representing the repeated elements by x^{exp} we obtain

$$Z = \{c_1, d_1^{2(s-1)}, c_2, x_1^{exp_1}, \dots, x_j^{exp_j}, y_1^{exp_1}, \dots, y_\ell^{exp_\ell}, c_3, c_4\},$$

subject to the following conditions:

- $c_1 = \frac{3s+p+\sqrt{(s+p)^2+4sp}}{2};$
- $d_1 = 2s + p;$
- $c_2 = s + p;$
- for $1 \leq i \leq j$, $s \leq x_i \leq s + p;$

If $x_i > s$ then

- $x_i \geq s + 2;$
- $exp_i = k_i(x_i - s - 1)$ times, $k_i = 1, 2, \dots;$
- $n_i = k_i(x_i - s);$

- else $x_i = s$ and $n_i = 0$;
- $\sum_1^j n_i \leq p$;
- for $1 \leq i \leq \ell$, $s \leq y_i \leq s + p$;
 If $y_i > s$ then
 - $y_i \geq s + 2$;
 - $\exp_i = k'_i(y_i - s - 1)$ times, $k_i = 1, 2, \dots$;
 - $n'_i = k'_i(y_i - s)$;
- else $y_i = s$ and $n'_i = 0$;
- $\sum_1^j n'_i \leq p$;
- $c_3 = \frac{3s+p-\sqrt{(s+p)^2+4sp}}{2}$;
- $c_4 = 0$.

Theorem 16. *Let Z be an ordered multi-set. Then Z corresponds to the Laplacian eigenvalues of a SI-core graph.*

Proof. A SI-core graph $G(s, p)$ must be built and its Laplacian eigenvalues must be equal to Z . First, we determine s and p . Let consider the elements $d_1 = (2s + p)^{2(s-1)}$. Let $e = 2(s - 1)$; hence $s = \frac{e+2}{2}$. As $d_1 = 2^{\frac{e+2}{2}} + p$, we have $p = d_1 - 2s$.

Knowing s and p , it is easy to see, by Corollaries 12.1 and 14.1, that

- $c_1 = \mu_1 = \frac{3s+p+\sqrt{(s+p)^2+4sp}}{2}$,
- $d_i, 1 \leq i \leq 2(s - 1) = \mu_j, 2 \leq j \leq 2s - 1 = 2s + p$.
- $c_2 = \mu_{2s} = s + p$,
- $c_3 = \mu_{n-1} = \frac{3s+p-\sqrt{(s+p)^2+4sp}}{2}$ and
- $c_4 = \mu_n = 0$.

These Laplacian eigenvalues are the same for any $G \in \mathcal{G}(s, p)$.

Now, the remaining values of Z allow us to determine which SI -core graph in $\mathcal{G}(s, p)$ correspond to this ordered multi-set.

Let consider now the elements x_i , $1 \leq i \leq j$ of Z . Suppose, w.l.g, that the eigenvalues of G corresponding to simplicial elements all adjacent to the same mv s S_1 of G are considered together, $x_i = \mu_k$, $2s+1 \leq k \leq 2s+p-1$. Apply the same reasoning to the elements y_i , $1 \leq i \leq \ell$ of Z ; they correspond to the same mv s S_2 of G considered together, $y_i = \mu_k$, $2s+p \leq k \leq 2s+2p-2$. By the ordered multi-set Z definition, $s \leq x_i \leq s+p$. For all $1 \leq i \leq j$, if $x_i = s$ then all p vertices adjacent to S_1 are false twins; if $x_i = s+p$ then all p vertices adjacent to S_1 are true twins. The same reasoning can be applied to y_i , $1 \leq i \leq \ell$. In this case Z corresponds to the Laplacian eigenvalues of $G_{min}(s, p)$ and $G_{max}(s, p)$.

Suppose now that $x_i = \mu_k$ different from s and $s+p$. The eigenvalue μ_k corresponds to a simplicial vertex v . If $\mu_k > s$, the eigenvalue μ_k appears because v has true twins. So, there is a complete graph K_l of cardinality $l = \mu_k - s$, composed by simplicial vertices. The multiplicity of μ_k originated by K_l , $v \in K_l$, is $l - 1 = \mu_k - s - 1$, see Theorem 14. As more than one set of simplicial vertices being true twins can have the same cardinality, the corresponding eigenvalue can appear also more times. The value exp_k corresponds to the multiplicity of μ_k . The total number of simplicial vertices being true twins adjacent to S_1 and S_2 , respectively, must be at most p . These results correspond to the restrictions presented in the definition of the ordered multi-set Z . \square

For instance, let us consider the ordered multi-set $\{18, 16^4, 13, 12^8, 6^6, 3^4, 1, 0\}$. As it obeys to the description of multi-set Z , it corresponds to a graph G , by Theorem 16. Observe the element $d_1 = 16$ with multiplicity 4. So, $4 = 2(s-1)$ and $s = 3$. As $c_2 = 13$, $p = 13 - 3 = 10$. Graph G has 26 vertices: two separators of cardinality 3 and 20 simplicial vertices, 10 adjacent to each separator. Moreover, since 12 has multiplicity 8 in Z , it can be concluded that exists $\mu_i = 12$ eigenvalue of G corresponding to a clique of size $12 - 3 = 9$. Furthermore, since 6 has multiplicity 6 in Z it can be concluded that exists $\mu_i = 6$, eigenvalue of G corresponding to 3 cliques of simplicial vertices of size 3. The remaining eigenvalues are equal to 3. See in Figure 5 a SI -core graph whose Laplacian spectrum is $[18, 16^4, 13, 12^8, 6^6, 3^4, 1, 0]$.




Figure 5: A *SI*-core graph in $\mathcal{G}(3, 10)$

9. Answering some questions

Question 1: Given n , is it possible to build a *SI*-core graph with n vertices?

Firstly n must be even; with any pair $s \geq 2$ and $p \geq 2$ such that $2s + 2p = n$ it is possible to build $G \in \mathcal{G}(s, p)$. The $2s$ vertices form the maximal clique of *mvss* S_1 and S_2 ; p vertices are adjacent to S_1 and p vertices are adjacent to S_2 . These vertices can form mutually exclusive cliques.

Question 2: Are there *SI*-core graphs determined by their Laplacian spectrum?

Given an ordered multi-set Z it is possible to build a *SI*-core graph. However, some spectra can correspond to one graph (the *min* and *max* *SI*-core graphs, for instance) and some other spectra can correspond to more than one non-isomorphic graph. For instance, graphs G_1 and G_2 in Figure 6 have the same spectrum,

$$[10.21699, 9^2, 7, 5^4, 2^4, 0.78301, 0],$$

however they are not isomorphic.

Question 3: Given $a \in \mathbb{N}$, fixed s and p , is there a *SI*-core graph $G(S, p)$ whose Laplacian spectrum contains a ?

Since s and p are already known, several eigenvalues of the graph have their value determined in advance. However, there are $2(p - 1)$ values that are possible of be modified. By the ordered multi-set Z definition, we know the restrictions that must be applied.

Firstly $s \leq a \leq s + p$.




Figure 6: Two *SI*-core graphs in $\mathcal{G}(2, 5)$

If $a = s$, the solution is immediate. Actually, the value $a = s$ can appear $2(p - 1)$ times in the spectra of G . Let us consider $a \neq s$, that is $a > s$.

- a must be greather or equal $s + 2$;
- a must appear $k(a - s - 1)$ times, $k = 1, 2, \dots$;
- each eigenvalue corresponds to a clique of simplicial vertices of size $a - s$; the sum of these elements must be less or equal p .

Actually, it is quite easy to include certain values as a Laplacian eigenvalue of $G(s, p)$. Let start with $G_{min}(s, p)$. Its spectrum is already known:

$$\left[\frac{3s + p + \sqrt{(s + p)^2 + 4sp}}{2}; (2s + p)^{2(s-1)}; (s + p); s^{2(p-1)}; \frac{3s + p - \sqrt{(s + p)^2 + 4sp}}{2}; 0 \right].$$

The term $s^{2(p-1)}$ corresponds to: s^{p-1}, s^{p-1} . One (or two) of these terms can be rewritten as $a^{a-s-1}, s^{p-a+s+1}$. For instance, we want that the value 7 appears as an eigenvalue of $G(3, 10)$. As 7 satisfies the condition $3 \leq 7 \leq 13$, we know that 7 must appear at least 3 times (or 6) as an eigenvalue. It is interesting that, althout 9 times seems to be ok, it is not, since $3(7 - 3)$ is not less or equal 10 (the last condition).

Based on these results, we can conclude that there are k cliques of $a - s$ simplicial vertices adjacent to the same mus in G .

10. Acknowledgment

The second author was partially supported by CNPq, Grant 405552/2023-8 and FAPERJ, Process SEI 260003/001228/2023.

References

- [1] Abreu, N., Justel, C.M., Markenzon, L. *Strictly chordal graphs: Structural properties and integer Laplacian eigenvalues*, Linear Algebra and its Applications **682** (2024) 351-362.
- [2] Blair, J.R.S., Peyton, B., *An introduction to chordal graphs and clique trees*, In: George, J.A., Gilbert, J. R., Liu, J. W. H. (Eds.), *Graph theory and sparse matrix computation*. Springer Verlag, IMA **56**, 1-30, 1993.
- [3] Brandstädt, A., Wagner, P., *Characterising (k,ℓ) -leaf powers*, Discrete Applied Mathematics **158** (2010) 110-122.
- [4] Cardoso, D.M., Rojo, O. *Edge perturbation on graphs with clusters: Adjacency, Laplacian and signless Laplacian eigenvalues*, Linear Algebra and its Applications **512** (2017) 113-128.
- [5] Gilmore, P.C., Hoffman, A.J., *A characterization of comparability graphs and of interval graphs*, Canad. J. Math. **16** (1964) 539-548.
- [6] Golumbic, M.C., *Algorithmic graph theory and perfect graphs* (2nd edition), Academic Press, New York, 2004.
- [7] Golumbic, M.C., Peled, U.N., *Block duplicate graphs and a hierarchy of chordal graphs*, Discrete Appl. Math. **124** (2002) 67-71.
- [8] Hara, H., Takemura, A., *Boundary cliques, clique trees and perfect sequences of maximal cliques of a chordal graph*, METR 2006-41, Department of Mathematical Informatics, University of Tokyo, 2006.
- [9] Harary, F., *A characterization of block graphs*, Canad. Math. Bull. **6**(1) (1963) 1-6.
- [10] Kennedy, W., *Strictly chordal graphs and phylogenetic roots*, Master Thesis, University of Alberta, 2005.

- [11] Kirkand, S., de Freitas, M. A. A., Del Vecchio, R., Abreu, N.M.M., *Split Nonthreshold Laplacian Integral Graphs* Linear and Multilinear Algebra **58**(2) (2010) 221-233.
- [12] Lin, G., Kearney, P.E., Jiang, T. *Phylogenetic k -Root and Steiner k -Root*, In Lee, D.T., Teng, S.H. (Eds.) Algorithms and Computation (ISAAC 2000) *LNCS* **1969** (2000) 539-551.
- [13] Markenzon, L., Pereira, P.R.C., *One-phase algorithm for the determination of minimal vertex separators of chordal graphs*, Int. Trans. Oper. Res. **17** (2010) 683-690.
- [14] Markenzon, L., Waga, C.F.E.M., *New results on ptolemaic graphs*, Discrete Appl. Math. **196** (2015), 135-140.
- [15] Markenzon, L., Waga, C. F. E. M., *Strictly interval graphs: characterization and linear time recognition*, Electr. Notes Discrete Math. **52** (2016), 181–188.
- [16] Markenzon, L., Paciornik, N., *Multicore Graphs: Characterization and Properties*. In: Hoffman, F., Holliday, S., Rosen, Z., Shahrokhi, F., Wierman, J. (eds) Combinatorics, Graph Theory and Computing. SE-ICCGTC 2021. Springer Proceedings in Mathematics & Statistics, **448** (2024) Springer.
- [17] OEIS Foundation Inc. (2025), Entry A000041 in The On-Line Encyclopedia of Integer Sequences, <https://oeis.org/A000041>.
- [18] Shibata, Y., *On the tree representation of chordal graphs*, J. Graph Theory **12** (1988) 421-428.
- [19] You, L., Yang, M., W. So, W., Xi, W., *On the spectrum of an equitable quotient matrix*, Linear Algebra Appl. **577** (2019), 21-40.