

ON THE FUTURE COVER OF A SOFIC SHIFT

KLAUS THOMSEN

ABSTRACT. The paper contains a new proof of the theorem by Krieger which establishes the canonicity of the future cover of a sofic shift. In addition the paper describes a method to produce new canonical covers from a given one, resulting in canonical covers related to, but generally different from the future cover.

1. INTRODUCTION

A cover of a sofic shift Y is a subshift of finite type X together with a factor map $\pi : X \rightarrow Y$ from X onto Y . Every sofic shift has a wealth of different covers, so the least one can do to simplify considerations regarding covers is to identify two covers $\pi : X \rightarrow Y$ and $\pi' : X' \rightarrow Y$ of the same sofic shift Y when they are isomorphic in the sense that there is a conjugacy $\chi : X \rightarrow X'$ such that $\pi' \circ \chi = \pi$. Recall that a subshift of finite type can be realized as the edge shift X_G of a finite directed graph G and that a labeling of the set of edges E_G is a map $L_G : E_G \rightarrow A$, where A is a finite set. If $Y = L_G(X_G)$, where $L_G : X_G \rightarrow Y \subseteq A^{\mathbb{Z}}$ is the map obtained by reading the labels of elements in X_G , we say that (G, L_G) is a presentation of Y . Then any cover of Y is isomorphic to one coming from a presentation of Y , see Theorem 3.2.1 in [LM]. But even after having reduced covers to labeled graphs, every sofic shift has an uncanny host of different presentations. Fortunately the future cover of Wolfgang Krieger, [Kr1], and its associated labeled graph which we denote by $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$, stands out because it is canonical in the sense that a conjugacy $\psi : Y \rightarrow Z$ of sofic shifts lifts to a conjugacy $\psi_{\mathbb{K}} : X_{\mathbb{K}(Y)} \rightarrow X_{\mathbb{K}(Z)}$ such that

$$\begin{array}{ccc} X_{\mathbb{K}(Y)} & \xrightarrow{\psi_{\mathbb{K}}} & X_{\mathbb{K}(Z)} \\ L_{\mathbb{K}(Y)} \downarrow & & \downarrow L_{\mathbb{K}(Z)} \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes. Thus the subshift of finite type $X_{\mathbb{K}(Y)}$ and the factor map $L_{\mathbb{K}(Y)}$ are both conjugacy invariants for Y . The same is of course also true for any other cover which is canonical in the same way. It must be emphasized that also components of a canonical cover that are not of maximal entropy as well as the wandering parts of the cover, which often arise even when Y is mixing, also give rise to conjugacy invariants for Y . This observation is the main motivation for the work and the results that are presented below.

We start by giving a new proof of Krieger's theorem. While his proof is a tour de force in handling sliding block codes, the proof given here is more conceptual and leads to an abstract characterization of the future cover analogous to the abstract characterization of the canonical cover of an irreducible sofic shift known as the minimal right-resolving cover or

2020 *Mathematics Subject Classification.* Primary ; Secondary .
Key words and phrases. Sofic shifts, canonical covers.

Fischer cover, [F]. Alternative proofs of parts of Krieger's theorem have been given before; first by Boyle, Kitchens and Marcus in [BKM] and later by Nasu in [N]. The proof presented here is also different from those.

The main new result of the paper is a theorem which out of a given canonical cover produces a new one. To appreciate this, recall that Krieger proved the uniqueness of the conjugacy $\psi_{\mathbb{K}}$ in the diagram above. Without this additional requirement it is obvious that a canonical cover is not unique up to isomorphism; for example the disjoint union of two copies of the future cover will again be canonical in the sense used so far, but not with unique lifts of conjugacies. In the following we shall therefore refer to a construction of covers as *weakly canonical* when it gives lifts of conjugacies, but possibly lifts that are not unique, while a construction which gives rise to unique lifts of conjugacies, as the future cover does, will be called a *strongly canonical* cover. The main result of the paper, Theorem 4.8, describes a method to produce a strongly canonical cover out of a given weakly canonical one.

An important virtue of the construction is that it produces new strongly canonical covers of sofic shifts. Up to now only two strongly canonical covers for sofic shifts have been known; at least when we for simplicity ignore the doubling coming from the passage from right-resolving to left-resolving covers that can be achieved by 'time-reversal'. Besides the future cover it is the minimal right-resolving cover, and only for irreducible sofic shifts. It is therefore interesting to compare the strongly canonical covers that can be produced using Theorem 4.8 with the future cover and, in the irreducible case, the minimal right-resolving cover. For this we introduce a method to find the future cover of a sofic shift starting from an arbitrary given presentation. This is based on a variation of the so-called subset construction described in §3.3 of [LM]. More specifically, it is shown that a restriction to a subset of the vertexes in the subset construction, applied to an arbitrary presentation of the sofic shift, gives rise to a labeled graph whose merged graph is isomorphic to the future cover. Thus the future cover is the factor obtained from a subgraph of the subset construction by identifying vertexes with the same follower set. This fact is used to produce the examples occurring in the paper, and makes it easy to show by examples that the new strongly canonical covers that can be obtained from Theorem 4.8 in general are different both from the future cover, and in the irreducible case, also the minimal right-resolving cover.

Acknowledgement: I thank Brian Marcus, Tom Meyerovitch and Chengyu Wu for the collaboration that produced [MMTW]. I'm particularly grateful to Brian for renewing my interest in sofic shifts, and I thank him also for showing me his solution to Exercise 3.2.8 (e) in [LM].

2. KRIEGER'S FUTURE COVER AND ITS CANONICITY

2.1. Setup and notation. Let A be a finite set. The set $A^{\mathbb{Z}}$ of bi-infinite sequences

$$x = (x_i)_{i \in \mathbb{Z}} = \cdots x_{-3}x_{-2}x_{-1}x_0x_1x_2x_3 \cdots$$

of elements from A is a compact metric space; a metric can be defined for example as in Example 6.1.10 of [LM]. The shift σ is the homeomorphism of $A^{\mathbb{Z}}$ defined such that

$$\sigma(x)_i = x_{i+1}.$$

A closed subset $X \subseteq A^{\mathbb{Z}}$ is shift-invariant when $\sigma(X) = X$ and it is then called a *subshift*. We shall mainly consider *subshifts of finite type* (abbreviated to SFT) and *sofic shifts*. See Definition 2.1.1 and Definition 3.1.3 in [LM].

The set $\mathbb{W}(X)$ of *words* in a subshift X consists of the finite strings $a_1 a_2 \cdots a_n \in A^n$ with the property that $a_1 a_2 \cdots a_n = x_1 x_2 \cdots x_n$ for some element $x = (x_i)_{i \in \mathbb{Z}} \in X$. Given an element $x \in X$ and an integer $j \in \mathbb{Z}$ we denote by $x_{(-\infty, j]}$ the element

$$\cdots x_k x_{k+1} x_{k+2} \cdots x_j \in A^{(-\infty, j]},$$

and by $X(-\infty, j]$ the set

$$X(-\infty, j] := \{x_{(-\infty, j]} : x \in X\}.$$

The symbols $x_{[j, \infty)}$ and $X[j, \infty)$ have a similar meaning. The same goes for expressions like $x_{[i, j)}$ and $x_{(i, j]}$ when $i < j$ or $x_{[i, j]}$ when $i \leq j$. The set of σ -periodic points of a subshift X will be denoted by $\text{Per}(X)$.

Let G be a finite directed graph with edges (or arrows) E_G and vertexes V_G . For $e \in E_G$ let $s_G(e) \in V_G$ be the start vertex and $t_G(e)$ the terminal vertex of e . We shall always assume that G does not contain sinks or sources; i.e. that $s_G^{-1}(v) \neq \emptyset$ and $t_G^{-1}(v) \neq \emptyset$ for all $v \in V_G$. We extend the definition of t_G to finite and left-infinite paths in G , and the definition of s_G to finite or right-infinite paths in G , in the obvious way. We denote the edge shift of G by X_G . Thus X_G consists of the bi-infinite sequences

$$(e_i)_{i \in \mathbb{Z}} = \cdots e_{-3} e_{-2} e_{-1} e_0 e_1 e_2 e_3 \cdots \in E_G^{\mathbb{Z}}$$

of edges in G such that $s_G(e_{i+1}) = t_G(e_i)$ for all $i \in \mathbb{Z}$, and the words in $\mathbb{W}(X_G)$ are the finite paths in G . We shall sometimes refer to the elements of X_G as *rays* in G . The set X_G is an SFT and every SFT is conjugate to X_G for some graph G by Theorem 2.3.2 in [LM].

Let A be a finite set (the alphabet) and $L_G : E_G \rightarrow A$ a map considered as a labelling of the edges in G . We extend L_G to finite or infinite paths in the natural way. In particular, when $x := (e_i)_{i \in \mathbb{Z}} \in X_G$,

$$L_G(x) := (L_G(e_i))_{i \in \mathbb{Z}} = \cdots L_G(e_{-3}) L_G(e_{-2}) L_G(e_{-1}) L_G(e_0) L_G(e_1) L_G(e_2) L_G(e_3) \cdots \in A^{\mathbb{Z}}.$$

Then (G, L_G) is a *labeled graph*, $Y := L_G(X_G) \subseteq A^{\mathbb{Z}}$ is a sofic shift and (G, L_G) is a *presentation* of Y , cf. §3.1 in [LM].

The first lemma deals with factor codes, right-resolving graphs and right-closing factor codes. We refer to page 18, Definition 3.3.1 and Definition 8.1.8 in [LM], respectively.

Lemma 2.1. *Let X be an SFT and $\pi : X \rightarrow Y$ a right-closing factor code. There is a right-resolving labeled graph (G, L_G) and a conjugacy $\psi : X \rightarrow X_G$ such that*

$$\begin{array}{ccc} X & \xrightarrow{\psi} & X_G \\ & \searrow \pi \quad \swarrow L_G & \\ & Y & \end{array}$$

commutes.

Proof. Recode, first using (the proof of) Theorem 3.2.1 of [LM] to get a labeled graph (H, L_H) and a conjugacy $\psi_0 : X \rightarrow X_H$ such that

$$\begin{array}{ccc}
X & \xrightarrow{\psi_0} & X_H \\
& \searrow \pi & \swarrow L_H \\
& Y &
\end{array}$$

commutes. Note that $L_H : X_H \rightarrow Y$ is a right-closing factor code since π is. It follows therefore from Proposition 8.1.9 in [LM] that (H, L_H) is right-closing with delay as defined in Definition 5.1.4 of [LM]. Consequently Proposition 5.1.11 in [LM] gives us a right-resolving labeled graph (G, L_G) and a conjugacy $\psi_1 : X_H \rightarrow X_G$ such that

$$\begin{array}{ccc}
X_H & \xrightarrow{\psi_1} & X_G \\
& \searrow L_H & \swarrow L_G \\
& Y &
\end{array}$$

commutes. Set $\psi := \psi_1 \circ \psi_0$.

□

2.2. Regular vertexes and rays. Let (H, L_H) be a labeled graph and $Y = L_H(X_H)$ the sofic shift it presents. A vertex $v \in V_H$ is *regular* when there is an element $z \in X_H$ such that $t_H(z_{(-\infty, -1]}) = v$ and

$$(1) \quad \{L_H(x) : x \in X_H[0, \infty), s_H(x) = v\} = \{y \in Y[0, \infty) : L_H(z_{(-\infty, -1]})y \in Y\}.$$

For any element $z \in X_H$ with $t_H(z_{(-\infty, -1]}) = v$, the inclusion

$$\{L_H(x) : x \in X_H[0, \infty), s_H(x) = v\} \subseteq \{y \in Y[0, \infty) : L_H(z_{(-\infty, -1]})y \in Y\}$$

always holds, so the real condition in (1) is that the opposite inclusion,

$$\{L_H(x) : x \in X_H[0, \infty), s_H(x) = v\} \supseteq \{y \in Y[0, \infty) : L_H(z_{(-\infty, -1]})y \in Y\},$$

also holds.

In order to work with regularity of vertexes we introduce the following notation. Given an element $y \in Y$, set

$$(2) \quad F(y) := \{w \in Y[0, \infty) : y_{(-\infty, -1]}w \in Y\},$$

and given a vertex $v \in V_H$, set

$$f_H(v) := \{L_H(x) : x \in X_H[0, \infty), s_H(x) = v\}.$$

Then $v \in V_H$ is regular when there is an element $z \in X_H$ such that $t_H(z_{(-\infty, -1]}) = v$ and

$$(3) \quad f_H(v) = F(L_H(z)).$$

The set $f_H(v)$ will be called the *follower set* of v and $F(y)$ the *follower set* of y .

Lemma 2.2. *Assume that (H, L_H) is right-resolving. Let γ be a finite path in H , and assume that $s_H(\gamma)$ is regular. Then $t_H(\gamma)$ is regular.*

Proof. Let $z \in X_H$ be an element such that $t_H(z_{(-\infty, -1]}) = s_H(\gamma)$ and

$$(4) \quad \{L_H(\mu) : \mu \in X_H[0, \infty), s_H(\mu) = s_H(\gamma)\} = \{y \in Y[0, \infty) : L_H(z_{(-\infty, -1]})y \in Y\}.$$

Set $\gamma' := z_{(-\infty, -1]}\gamma$ and note that $\gamma' = z'_{(-\infty, -1]}$ for some $z' \in X_H$ with $t_H(z'_{(-\infty, -1]}) = t_H(\gamma)$. If $y \in Y[0, \infty)$ and $L_H(z'_{(-\infty, -1]})y = L_H(z_{(-\infty, -1]})L_H(\gamma)y \in Y$, then (4) implies

that $L_H(\gamma)y = L_H(\nu)$ for some $\nu \in X_H[0, \infty)$ with $s_H(\nu) = s_H(\gamma)$. Since (H, L_H) is right-resolving, $\nu = \gamma\nu'$ for some $\nu' \in X_H[0, \infty)$ with $s_H(\nu') = t_H(\gamma)$. Note that $L_H(\nu') = y$. \square

A labeled graph will be called *regular* when all vertexes are regular. Adapting Definition 3.3.7 in [LM] we say that a labeled graph (H, L_H) is *follower-separated* when $f_H(v) = f_H(w) \Rightarrow v = w$.

Example 2.3. The following labeled graph (G, L_G) is right-resolving and follower-separated. But it is not regular; the vertex 1 is not regular.




FIGURE 1. A right-resolving follower-separated labeled graph which is not regular.

Remark 2.4. Every synchronized graph in the sense of Jonoska, cf. §4 in [Jo], is regular. By Lemma 3.3.15, Proposition 3.3.16 and Corollary 3.3.19 in [LM] the minimal right-resolving presentation of an irreducible sofic shift is synchronizing in the sense of Jonoska and therefore also regular. The labeled graph in Figure 3 depicts the future cover of the even shift and it is regular, but it is not synchronizing. To give an irreducible example consider the following labeled graph which is strongly connected, right-resolving, and presents the even shift.




FIGURE 2. A strongly connected labeled graph which is right-resolving and regular, but not synchronizing.

Let $x \in X_H$. We are interested in the condition that

$$(5) \quad \{L_H(\mu) : \mu \in X_H[0, \infty), s_H(\mu) = t_H(x_{(-\infty, k]})\} = \{y \in Y[0, \infty) : L_H(x_{(-\infty, k]})y \in Y\},$$

or

$$(6) \quad f_H(t_H(x_{(-\infty, k]})) = F(\sigma^{k+1}(L_H(x))),$$

for all $k \in \mathbb{Z}$. Let X be a subshift and d a metric for the topology of X . For $x \in X$, set

$$\begin{aligned} \mathbb{U}(x) &:= \left\{ z \in X : \lim_{i \rightarrow -\infty} d(\sigma^i(z), \sigma^i(x)) = 0 \right\} \\ &= \{z \in X : z_i = x_i, i \leq N, \text{ for some } N \in \mathbb{Z}\}. \end{aligned}$$

This set is sometimes called 'the unstable manifold' of x .

Lemma 2.5. *Assume that (H, L_H) is right-resolving. Let $x \in X_H$. Then $L_H : \mathbb{U}(x) \rightarrow \mathbb{U}(L_H(x))$ is surjective if and only if (5) holds for all $k \in \mathbb{Z}$.*

Proof. Assume first that (5) holds for all $k \in \mathbb{Z}$ and consider an element $y \in \mathbb{U}(L_H(x))$. Then $y_{(-\infty, L]} = L_H(x_{(-\infty, L]})$ for some $L \in \mathbb{Z}$. Thanks to (5) this implies that $y_{[L+1, \infty)} \in f_H(t_H(x_{(-\infty, L]}))$, and hence also that there is an element $z \in X_H$ such that $z_{(-\infty, L]} = x_{(-\infty, L]}$ and $L_H(z_{[L+1, \infty)}) = y_{[L+1, \infty)}$. Then $z \in \mathbb{U}(x)$ and $L_H(z) = y$.

Assume then that $L_H : \mathbb{U}(x) \rightarrow \mathbb{U}(L_H(x))$ is surjective, and consider $k \in \mathbb{Z}$ and an element $y \in Y[0, \infty)$ such that $L_H(x_{(-\infty, k]})y \in Y$. Then $L_H(x_{(-\infty, k]})y \in \mathbb{U}(L_H(x))$ and there is therefore an element $z \in \mathbb{U}(x)$ such that $L_H(z) = L_H(x_{(-\infty, k]})y$. Choose $L < k$ such that $z_{(-\infty, L]} = x_{(-\infty, L]}$. Since (H, L_H) is right-resolving, it follows that $z_{(-\infty, k]} = x_{(-\infty, k]}$. Hence

$$y = L_H(z_{[k+1, \infty)}) \in f_H(t_H(x_{(-\infty, k]})).$$

□

An element $x \in X_H$ will be called *regular* when $L_H : \mathbb{U}(x) \rightarrow \mathbb{U}(L_H(x))$ is surjective. Then the following clearly holds.

Lemma 2.6. *Assume that $x \in X_H$ is regular. Then $s_H(x_k) \in V_H$ is regular for all $k \in \mathbb{Z}$.*

We denote in the following the set of regular rays in X_H by $\mathcal{R}(L_H)$.

Example 2.7. In the labeled graph H depicted in Figure 3 the vertex c is regular and hence all the vertexes are regular by Lemma 2.2. However, X_H contains non-regular rays; the two elements of X_H of period 2 are not regular.




FIGURE 3. A regular labeled graph with non-regular rays.

Lemma 2.8. *Assume that (X_H, L_H) is right-resolving and follower-separated. Then L_H is injective on $\mathcal{R}(L_H)$.*

Proof. If $x, y \in \mathcal{R}(L_H)$ and $L_H(x) = L_H(y)$, it follows first from (6) that $t_H(x_k) = t_H(y_k)$ for all $k \in \mathbb{Z}$ because we assume that (X_H, L_H) is follower-separated. Since (H, L_H) is right-resolving it follows that $x = y$. □

2.3. The future cover and Krieger's theorem. Let A be a finite set and $Y \subseteq A^{\mathbb{Z}}$ a sofic subshift. To define the future cover of Y we use the following notation. When $D \subseteq A^{\mathbb{N}}$ is a subset of $A^{\mathbb{N}}$ and $a \in A$, we let $\frac{D}{a}$ denote the set

$$\frac{D}{a} := \{x_1x_2x_3 \cdots \in A^{\mathbb{N}} : x_0x_1x_2 \cdots \in D, x_0 = a\} = \{x \in A^{\mathbb{N}} : ax \in D\};$$

i.e. $\frac{D}{a}$ is the set of elements of $A^{\mathbb{N}}$ that are obtained from the elements of D that start with a by deleting the first coordinate. More generally, when $w \in A^m$, we set

$$\frac{D}{w} = \{x \in A^{\mathbb{N}} : wx \in D\}.$$

Let $y \in Y$. Recall from (2) that

$$F(y) = \{z_{[0,\infty)} : z \in Y, z_j = y_j, j \leq -1\} \subseteq A^{\mathbb{N}}.$$

Set

$$V_{\mathbb{K}(Y)} := \{F(y) : y \in Y\}.$$

This is a finite set because Y is sofic. $V_{\mathbb{K}(Y)}$ is the set of vertexes in a labeled graph $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ where there is a labeled arrow

$$F(y) \xrightarrow{a} F(z)$$

from $F(y)$ to $F(z)$ in $\mathbb{K}(Y)$ precisely when $F(z) = \frac{F(y)}{a}$. The labeled graph $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ is then clearly right-resolving. Since

$$F(\sigma^{i+1}(y)) = \frac{F(\sigma^i(y))}{y_i},$$

there is an element $\alpha_Y(y) \in X_{\mathbb{K}(Y)}$ determined by the two conditions:

- a) $s_{\mathbb{K}(Y)}(\alpha_Y(y)_i) = F(\sigma^i(y))$, $i \in \mathbb{Z}$, and
- b) $L_{\mathbb{K}(Y)}(\alpha_Y(y)) = y$.

The map $\alpha_Y : Y \rightarrow X_{\mathbb{K}(Y)}$ commutes with the shift, i.e. $\alpha_Y \circ \sigma = \sigma \circ \alpha_Y$, but it is not continuous.

Lemma 2.9. $L_{\mathbb{K}(Y)}(X_{\mathbb{K}(Y)}) = Y$.

Proof. Let $x = (x_i)_{i=-\infty}^{\infty} \in X_{\mathbb{K}(Y)}$ and consider an $N \in \mathbb{N}$. By definition $s_{\mathbb{K}(Y)}(x_{-N}) = F(y)$ for some $y \in Y$ and

$$L_{\mathbb{K}(Y)}(x_{-N})L_{\mathbb{K}(Y)}(x_{-N+1})L_{\mathbb{K}(Y)}(x_{-N+2}) \cdots \in F(y).$$

In particular,

$$L_{\mathbb{K}(Y)}(x_{-N})L_{\mathbb{K}(Y)}(x_{-N+1}) \cdots L_{\mathbb{K}(Y)}(x_N) \in \mathbb{W}(Y)$$

for all $N \in \mathbb{N}$, implying that $L_{\mathbb{K}(Y)}(X_{\mathbb{K}(Y)}) \subseteq Y$. The reversed inclusion, $Y \subseteq L_{\mathbb{K}(Y)}(X_{\mathbb{K}(Y)})$, follows from the observation b) above. \square

Thus $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ is a presentation of Y , and it is known as *the future cover* of Y , cf. Exercise 3.2.8 on page 75 of [LM]. ¹

¹Krieger introduced the word 'future' in [Kr1], and called an equivalent version of $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ 'the future state chain'. The terminology "future cover" was used in [BKM], but only for a component of $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$, and only for transitive sofic shifts. In Exercise 3.2.8 of [LM] the quotation marks were deleted and $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ was called the future cover for general sofic shifts.

We can now formulate Krieger's theorem on the canonicity of the future cover. In our notation Theorem 2.14 in [Kr1] reads as follows:

Lemma 2.10. (Theorem 2.14 of [Kr1]) *Let Y and Z be sofic shifts and $\psi : Y \rightarrow Z$ a conjugacy. There is a unique conjugacy $\psi_{\mathbb{K}} : X_{\mathbb{K}(Y)} \rightarrow X_{\mathbb{K}(Z)}$ such that*

- (a) $\psi \circ L_{\mathbb{K}(Y)} = L_{\mathbb{K}(Z)} \circ \psi_{\mathbb{K}}$, and
- (b) $\psi_{\mathbb{K}} \circ \alpha_Y = \alpha_Z \circ \psi$.

When Y (and Z) are transitive, Krieger was able to partly dispense with condition (b) in Lemma 2.10, cf. Corollary 2.16 in [Kr1]. And in Theorem 2.2 of [Kr2] he removed condition (b) entirely by showing that it follows from (a). In this way he proved

Theorem 2.11. (W. Krieger, 1984, 1987) *Let Y and Z be sofic shifts and $\psi : Y \rightarrow Z$ a conjugacy. There is a unique conjugacy $\psi_{\mathbb{K}} : X_{\mathbb{K}(Y)} \rightarrow X_{\mathbb{K}(Z)}$ such that*

$$\begin{array}{ccc} X_{\mathbb{K}(Y)} & \xrightarrow{\psi_{\mathbb{K}}} & X_{\mathbb{K}(Z)} \\ L_{\mathbb{K}(Y)} \downarrow & & \downarrow L_{\mathbb{K}(Z)} \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes. This conjugacy $\psi_{\mathbb{K}}$ has the property that also

$$(7) \quad \begin{array}{ccc} X_{\mathbb{K}(Y)} & \xrightarrow{\psi_{\mathbb{K}}} & X_{\mathbb{K}(Z)} \\ \alpha_Y \uparrow & & \uparrow \alpha_Z \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes.

Note that (7) says that the right-inverse α_Y of $L_{\mathbb{K}(Y)}$ is canonical in the same sense as $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$. It can therefore also be used to get conjugacy invariants for Y .

Before we can give the proof of Theorem 2.11 we need some preparations. First of all we need the relation

$$(8) \quad f_{\mathbb{K}(Y)}(F(y)) = F(y)$$

in which $F(y)$ is first considered as a vertex in $\mathbb{K}(Y)$ and next as a subset of $A^{\mathbb{N}}$. In more detail,

Lemma 2.12. $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ *is follower-separated. In fact, for all $y \in Y$,*

$$\{L_{\mathbb{K}(Y)}(w) : w \in X_{\mathbb{K}(Y)}[0, \infty), s_{\mathbb{K}(Y)}(w) = F(y)\} = F(y).$$

Proof. This follows directly from the definition of $\mathbb{K}(Y)$. □

Lemma 2.13. $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ *is regular.*

Proof. Let $F(y) \in V_{\mathbb{K}(Y)}$ and consider the element $\alpha_Y(y) \in X_{\mathbb{K}(Y)}$. Then $t_{\mathbb{K}(Y)}(\alpha_Y(y)_{(-\infty, -1]}) = F(y)$. Note that $L_{\mathbb{K}(Y)}(\alpha_Y(y)_{(-\infty, -1]}) = y_{(-\infty, -1]}$, which implies that

$$\{y' \in Y[0, \infty) : L_{\mathbb{K}(Y)}(\alpha_Y(y)_{(-\infty, -1]})y' \in Y\} = F(y).$$

Thanks to (8) this is all we need to show, cf. (3). □

Lemma 2.14. $\alpha_Y(y)$ is regular in $X_{\mathbb{K}(Y)}$ for all $y \in Y$, i.e. $\alpha_Y(Y) \subseteq \mathcal{R}(L_{\mathbb{K}(Y)})$.

Proof. We must check that

$$f_{\mathbb{K}(Y)}(t_{\mathbb{K}(Y)}(\alpha_Y(y)_{(-\infty, k]}) = F(\sigma^{k+1}(\alpha_Y(y)))$$

for $k \in \mathbb{Z}$. This is a direct check, using (8):

$$\begin{aligned} f_{\mathbb{K}(Y)}(t_{\mathbb{K}(Y)}(\alpha_Y(y)_{(-\infty, k]}) &= f_{\mathbb{K}(Y)}(s_{\mathbb{K}(Y)}(\alpha_Y(y)_{k+1})) \\ &= f_{\mathbb{K}(Y)}(F(\sigma^{k+1}(y))) = F(\sigma^{k+1}(y)) = F(\sigma^{k+1}(\alpha_Y(y))). \end{aligned}$$

□

Lemma 2.15. Let $\gamma : Y \rightarrow X_{\mathbb{K}(Y)}$ be a map such that $\gamma(Y) \subseteq \mathcal{R}(L_{\mathbb{K}(Y)})$ and $L_{\mathbb{K}(Y)} \circ \gamma = \text{id}_Y$. Then $\gamma = \alpha_Y$.

Proof. By definition and by Lemma 2.14 α_Y has the same properties as γ . Since $L_{\mathbb{K}(Y)}(\alpha_Y(y)) = y = L_{\mathbb{K}(Y)}(\gamma(y))$ for all $y \in Y$, and since $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ is follower-separated by Lemma 2.12, it follows from Lemma 2.8 that $\gamma = \alpha_Y$. □

Given an element $w \in A^m$ we denote in the following by w^∞ the m -periodic element w^∞ in $\text{Per}(A^\mathbb{Z})$ with the property that $(w^\infty)_{[0, m-1]} = w$.

Lemma 2.16. Let d be a metric for the topology of $X_{\mathbb{K}(Y)}$, let $x \in X_{\mathbb{K}(Y)}$ and let $N \in \mathbb{Z}$. There is a periodic point $p \in \text{Per}(Y)$ and an element $x' \in X_{\mathbb{K}(Y)}$ such that $x'_i = x_i$, $i \geq N$, and $\lim_{i \rightarrow -\infty} d(\sigma^i(x'), \sigma^i(\alpha_Y(p))) = 0$. In particular, $x' \in \mathcal{R}(L_{\mathbb{K}(Y)})$.

Proof. Choose $y \in Y$ such that $F(y) = s_{\mathbb{K}(Y)}(x_N)$. By the pigeon hole principle there are integers $\{n_i\}_{i \in \mathbb{N}}$ such that

$$\dots < n_3 < n_2 < n_1 < n_0 < 0$$

and $s_{\mathbb{K}(Y)}(\alpha_Y(y)_{n_j}) = s_{\mathbb{K}(Y)}(\alpha_Y(y)_{n_0})$ for all j . Let $q_j := L_{\mathbb{K}(Y)}(\alpha_Y(y)_{[n_j, n_0-1]}) = y_{[n_j, n_0-1]}$ and note that $p^j := q_j^\infty \in Y$. We claim that $F(\sigma^{n_0}(y)) \subseteq F(p^j)$ for all $j \in \mathbb{N}$. To see this, let $z \in F(\sigma^{n_0}(y))$. Since $F(\sigma^{n_0}(y)) = s_{\mathbb{K}(Y)}(\alpha_Y(y)_{n_j}) = s_{\mathbb{K}(Y)}(\alpha_Y(y)_{n_0}) = F(\sigma^{n_0}(y))$ and $L_{\mathbb{K}(Y)}(\alpha_Y(y)_{[n_j, n_0-1]}) = q_j$, it follows that $F(\sigma^{n_0}(y)) = \frac{F(\sigma^{n_0}(y))}{q_j}$, and hence also that

$$F(\sigma^{n_0}(y)) = \frac{F(\sigma^{n_0}(y))}{(q_j)^k}$$

for all $k \in \mathbb{N}$. Hence

$$F(\sigma^{n_0}(y)) = \{w \in F(\sigma^{n_0}(y)) : (q_j)^k w \in F(\sigma^{n_0}(y)) \ \forall k \in \mathbb{N}\},$$

which implies that $p_{(-\infty, -1]}^j z \in Y$, i.e. $z \in F(p^j)$. We claim next that $F(\sigma^{n_0}(y)) = F(p^l)$ for some $l \in \mathbb{N}$. By passing to a subsequence we may assume for this that $F(p^j) = F(p^1)$ for all j . If $F(p^1) \setminus F(\sigma^{n_0}(y)) \neq \emptyset$, there is an element $z \in Y[0, \infty)$ such that $y_{[n_j, n_0-1]} z \in Y[0, \infty)$ for all $j > 1$, but $y_{(-\infty, n_0-1]} z \notin Y$, which is impossible. This proves the claim. It follows that $\alpha_Y(y)_{[n_0, -1]}$ is a path, labeled by $y_{[n_0, -1]}$, which starts at $F(\sigma^{n_0}(y)) = F(p^l) = s_{\mathbb{K}(Y)}(\alpha_Y(p^l)_0)$ and ends at $s_{\mathbb{K}(Y)}(\alpha_Y(y)_0) = F(y) = s_{\mathbb{K}(Y)}(x_N)$. Set $p := \sigma^{-n_0}(p^l)$ and

$$x' := \alpha_Y(p)_{(-\infty, n_0-1]} \alpha_Y(y)_{[n_0, -1]} x_{[N, \infty)} \in X_{\mathbb{K}(Y)}.$$

Then p and x' have the desired properties, and x' is regular because $\mathbb{U}(x') = \mathbb{U}(\alpha_Y(p))$ and $\alpha_Y(p) \in \mathcal{R}(L_{\mathbb{K}(Y)})$ by Lemma 2.14. □

Corollary 2.17. *For every element $x \in X_{\mathbb{K}(Y)}$ and every $N \in \mathbb{Z}$ there is a regular ray $x' \in \mathcal{R}(L_{\mathbb{K}(Y)})$ such that $x'_i = x_i$ for $i \geq N$.*

Proof. This follows from Lemma 2.16. \square

In the following we shall work with labeled-graph homomorphisms, cf. Definition 3.1.2 of [LM]. Let (G, L_G) and (H, L_H) be labeled graphs. A *labeled-graph homomorphism* $\theta : (G, L_G) \rightarrow (H, L_H)$ is a graph homomorphism $\theta : G \rightarrow H$ such that $L_H \circ \theta(e) = L_G(e)$ for all $e \in E_G$. The sliding block code $X_G \rightarrow X_H$ induced by θ , which we also denote by θ , has the property that $L_H \circ \theta = L_G$ on X_G .

Lemma 2.18. *Assume that (G, L_G) is a right-resolving and regular presentation of Y . It follows that there is a labeled-graph homomorphism $\theta : (G, L_G) \rightarrow (\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ such that*

$$\theta(\mathcal{R}(L_G)) \subseteq \mathcal{R}(L_{\mathbb{K}(Y)}).$$

If (G, L_G) also is follower-separated, $\theta : G \rightarrow \mathbb{K}(Y)$ is injective.

Proof. Let $v \in V_G$. Since v is regular there is an $x \in X_G$ such that $t_G(x_{-\infty, -1]) = v$ and $f_G(v) = F(L_G(x))$. Define $\theta(v)$ such that

$$\theta(v) := F(L_G(x)) \in V_{\mathbb{K}(Y)}.$$

This is well-defined: If x' has the same properties as x , it follows that $F(L_G(x')) = f_G(v) = F(L_G(x))$.

Let $e \in E_G$. Choose $x \in X_G$ such that $t_G(x_{-\infty, -1]) = s_G(e)$ and $f_G(s_G(e)) = F(L_G(x))$. Set $x'_i = x_{i+1}$, $i \leq -2$, $x'_{-1} = e$ and let $x'_i, i \geq 0$, be arbitrary subject to the condition that $x' \in X_G$. Then $t_G(x'_{(-\infty, -1]}) = t_G(e)$ and by using that (G, L_G) is right-resolving it follows that

$$\begin{aligned} f_G(t_G(e)) &= \frac{f_G(s_G(e))}{L_G(e)} = \frac{\{y \in Y[0, \infty) : L_G(x_{(-\infty, -1]})y \in Y\}}{L_G(e)} \\ &= \{y \in Y[0, \infty) : L_G(x'_{(-\infty, -1]})y \in Y\}. \end{aligned}$$

Hence $\theta(t_G(e)) = F(L_G(x'))$ and $F(L_G(x')) = \frac{F(L_G(x))}{L_G(e)}$. There is therefore an edge e' in $\mathbb{K}(Y)$ from $\theta(s_G(e)) = F(L_G(x))$ to $\theta(t_G(e)) = F(L_G(x'))$ with label $L_{\mathbb{K}(Y)}(e') = L_G(e)$. Since $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ is right-resolving this arrow is unique and we set $\theta(e) = e'$. Then $\theta : G \rightarrow \mathbb{K}(Y)$ is a labeled-graph morphism.

When $x \in \mathcal{R}(L_G)$, it follows from the definition of θ and the defining relation (5) for regular rays that $\theta(x) \in X_{\mathbb{K}(Y)}$ is the path

$$\dots \xrightarrow{L_G(x_{k-1})} F(L_G(\sigma^k(x))) \xrightarrow{L_G(x_k)} F(L_G(\sigma^{k+1}(x))) \dots$$

Thus $\theta(x) = \alpha_Y(L_G(x))$ and hence $\theta(x) \in \mathcal{R}(L_{\mathbb{K}(Y)})$ by Lemma 2.14.

Finally, assume that $v, w \in V_G$ are such that $\theta(v) = \theta(w)$. There are elements $x, x' \in X_G$ such that $t_G(x_{(-\infty, -1]}) = v$ and $t_G(x'_{(-\infty, -1]}) = w$, $f_G(v) = F(L_G(x))$ and $f_G(w) = F(L_G(x'))$. By definition of θ this implies that $f_G(v) = \theta(v) = \theta(w) = f_G(w)$ and hence that $v = w$ when (G, L_G) is follower-separated. \square

2.4. Proof of Krieger's theorem, Theorem 2.11. Let $\psi : Y \rightarrow Z$ be a conjugacy of sofic shifts. The future cover $(X_{\mathbb{K}(Y)}, L_{\mathbb{K}(Y)})$ of Y is isomorphic to the cover $(X_{\mathbb{K}(Y)}, \psi \circ L_{\mathbb{K}(Y)})$ of Z in the sense that

$$\begin{array}{ccc} X_{\mathbb{K}(Y)} & \xlongequal{\quad} & X_{\mathbb{K}(Y)} \\ L_{\mathbb{K}(Y)} \downarrow & & \downarrow \psi \circ L_{\mathbb{K}(Y)} \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes. Note that $\psi \circ L_{\mathbb{K}(Y)}$ is right-closing because $L_{\mathbb{K}(Y)}$ is right-resolving and ψ is a conjugacy. By Lemma 2.1 there is therefore a right-resolving labeled graph (G, L_G) and a conjugacy $\phi : X_{\mathbb{K}(Y)} \rightarrow X_G$ such that

$$\begin{array}{ccc} X_{\mathbb{K}(Y)} & \xrightarrow{\phi} & X_G \\ \psi \circ L_{\mathbb{K}(Y)} \downarrow & & \downarrow L_G \\ Z & \xlongequal{\quad} & Z \end{array}$$

commutes. Hence also

$$\begin{array}{ccc} X_{\mathbb{K}(Y)} & \xrightarrow{\phi} & X_G \\ L_{\mathbb{K}(Y)} \downarrow & & \downarrow L_G \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes. If $L_{\mathbb{K}(Y)} : \mathbb{U}(x) \rightarrow \mathbb{U}(L_{\mathbb{K}(Y)}(x))$ is surjective, then so is $L_G : \mathbb{U}(\phi(x)) \rightarrow \mathbb{U}(L_G(\phi(x))) = \mathbb{U}(\psi(L_{\mathbb{K}(Y)}(x)))$, showing that

$$(9) \quad \phi(\mathcal{R}(L_{\mathbb{K}(Y)})) \subseteq \mathcal{R}(L_G).$$

Let $v \in V_G$ and consider an element $x \in X_G$ with $s_G(x_0) = v$. Then $x = \phi(z)$, where $z = \phi^{-1}(x)$. Since ϕ is a conjugacy of shift spaces, it is given by a sliding block code defined on blocks of a given size, say $D \in \mathbb{N}$, cf. §1.5 in [LM]. It follows that $v = s_G(\phi(z')_0)$ for all $z' \in X_{\mathbb{K}(Y)}$ with the property that $z'_i = z_i$ for $i > -D$. By Corollary 2.17 there is a regular ray $z' \in \mathcal{R}(L_{\mathbb{K}(Y)})$ such that $z'_i = z_i$, $i > -D$. Then $\phi(z') \in \mathcal{R}(L_G)$ by (9) and hence $v = s_G(\phi(z)_0) = s_G(\phi(z')_0)$ is regular by Lemma 2.6. This shows that (G, L_G) is regular. It follows then from Lemma 2.18 that there is a labeled-graph homomorphism $\theta_0 : (G, L_G) \rightarrow (\mathbb{K}(Z), L_{\mathbb{K}(Z)})$ such that $\theta_0(\mathcal{R}(L_G)) \subseteq \mathcal{R}(L_{\mathbb{K}(Z)})$ and

$$\begin{array}{ccc} X_G & \xrightarrow{\theta_0} & X_{\mathbb{K}(Z)} \\ L_G \downarrow & & \downarrow L_{\mathbb{K}(Z)} \\ Z & \xlongequal{\quad} & Z \end{array}$$

commutes. All together we get a sliding block code $\phi' : X_{\mathbb{K}(Y)} \rightarrow X_{\mathbb{K}(Z)}$ such that

$$\phi'(\mathcal{R}(L_{\mathbb{K}(Y)})) \subseteq \mathcal{R}(L_{\mathbb{K}(Z)})$$

and

$$\begin{array}{ccc} X_{\mathbb{K}(Y)} & \xrightarrow{\phi'} & X_{\mathbb{K}(Z)} \\ L_{\mathbb{K}(Y)} \downarrow & & \downarrow L_{\mathbb{K}(Z)} \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes.

Now apply this with ψ replaced by ψ^{-1} . This gives us a sliding block code $\phi'' : X_{\mathbb{K}(Z)} \rightarrow X_{\mathbb{K}(Y)}$ such that

$$\phi''(\mathcal{R}(L_{\mathbb{K}(Z)})) \subseteq \mathcal{R}(L_{\mathbb{K}(Y)})$$

and

$$\begin{array}{ccc} X_{\mathbb{K}(Z)} & \xrightarrow{\phi''} & X_{\mathbb{K}(Y)} \\ L_{\mathbb{K}(Z)} \downarrow & & \downarrow L_{\mathbb{K}(Y)} \\ Z & \xrightarrow{\psi^{-1}} & Y \end{array}$$

commutes. Then

$$\phi'' \circ \phi'(\mathcal{R}(L_{\mathbb{K}(Y)})) \subseteq \mathcal{R}(L_{\mathbb{K}(Y)})$$

and

$$\begin{array}{ccc} X_{\mathbb{K}(Y)} & \xrightarrow{\phi'' \circ \phi'} & X_{\mathbb{K}(Y)} \\ L_{\mathbb{K}(Y)} \downarrow & & \downarrow L_{\mathbb{K}(Y)} \\ Y & \xlongequal{\quad} & Y \end{array}$$

commutes. Let $x \in \mathcal{R}(L_{\mathbb{K}(Y)})$. Then $L_{\mathbb{K}(Y)}(x) = L_{\mathbb{K}(Y)}(\phi'' \circ \phi'(x))$ and $\phi'' \circ \phi'(x) \in \mathcal{R}(L_{\mathbb{K}(Y)})$. Since $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ is right-resolving and follower-separated, Lemma 2.8 implies that $x = \phi'' \circ \phi'(x)$. Since $\mathcal{R}(L_{\mathbb{K}(Y)})$ is dense in $X_{\mathbb{K}(Y)}$ by Corollary 2.17, it follows that $\phi'' \circ \phi' = \text{id}$. By exchanging the roles of Y and Z it follows that also $\phi' \circ \phi'' = \text{id}$. Set $\psi_{\mathbb{K}} := \phi'$.

Uniqueness: If $\phi_i : X_{\mathbb{K}(Y)} \rightarrow X_{\mathbb{K}(Z)}$, are conjugacies such that $L_{\mathbb{K}(Z)} \circ \phi_i = \psi \circ L_{\mathbb{K}(Y)}$, $i = 1, 2$, then

$$\phi_i(\mathcal{R}(L_{\mathbb{K}(Y)})) = \mathcal{R}(L_{\mathbb{K}(Z)}).$$

Let $x \in \mathcal{R}(L_{\mathbb{K}(Y)})$. Since $L_{\mathbb{K}(Z)} \circ \phi_1(x) = \psi \circ L_{\mathbb{K}(Y)}(x) = L_{\mathbb{K}(Z)} \circ \phi_2(x)$ and $L_{\mathbb{K}(Z)}$ is injective on $\mathcal{R}(L_{\mathbb{K}(Z)})$ by Lemma 2.8, it follows that $\phi_1(x) = \phi_2(x)$. Since $\mathcal{R}(L_{\mathbb{K}(Y)})$ is dense in $X_{\mathbb{K}(Y)}$ by Corollary 2.17, it follows that $\phi_1 = \phi_2$.

Finally, to get the commuting diagram (7), set $\gamma := \psi_{\mathbb{K}}^{-1} \circ \alpha_Z \circ \psi$. By using Lemma 2.14, and that $\psi_{\mathbb{K}}^{-1}$ and ψ both are conjugacies, we find that $\gamma(Y) \subseteq \mathcal{R}(L_{\mathbb{K}(Y)})$. Since

$$L_{\mathbb{K}(Y)} \circ \gamma = \psi^{-1} \circ L_{\mathbb{K}(Z)} \circ \alpha_Z \circ \psi = \psi^{-1} \circ \psi = \text{id}_Y = L_{\mathbb{K}(Y)} \circ \alpha_Y,$$

it follows from Lemma 2.15 shows that $\gamma = \alpha_Y$. \square

As noted in the introduction an alternative proof of Theorem 2.11, without the uniqueness part and the diagram (7), was given by Nasu in [N].

2.5. The terminal component in the future cover of an irreducible sofic shift. The material in this section is probably folklore to experts, but I haven't found it in the literature.

Let H be a finite directed graph without sinks or sources. A vertex $v \in V_H$ is *recurrent* when there is a path γ in H such that $s_H(\gamma) = t_H(\gamma) = v$, and it is called *transient* otherwise. Let V_H^r denote the set of recurrent vertexes and V_H^t the set of transient vertexes. Two vertexes $v, w \in V_H$ *communicate*, and we write $v \rightsquigarrow w$, when there is a path γ in H with $s_H(\gamma) = v$ and $t_H(\gamma) = w$. When $v, w \in V_H^r$, $v \rightsquigarrow w$ and $w \rightsquigarrow v$, we write $v \sim w$. This an equivalence relation \sim on V_H^r and the elements $C \in V_H^r / \sim$ define a strongly connected subgraph H_C of H with $V_{H_C} = C$ and $E_{H_C} = \{e \in E_H : s_H(e), t_H(e) \in C\}$. The elements of V_H^r / \sim and the corresponding graphs H_C , $C \in V_H^r / \sim$, will be referred to as the *components* in H and the irreducible subshifts $X_C := X_{H_C}$, $C \in V_H^r / \sim$, as the components in X_H .

The components $C \in V_H^r / \sim$ are vertexes in a directed graph C_H where there is an arrow from C to C' when $v \rightsquigarrow w$ for some elements $v \in C$, $w \in C'$. While there are no sinks or sources in H by assumption, the graph C_H of its communicating components always have both. A *terminal component* (*source component*) in H is a sink (source) C in C_H and the associated subshift X_C is a *terminal component* (*source component*) in X_H .

Lemma 2.19. *Let (H, L_H) be a right-resolving and follower-separated labeled graph and C a terminal component in H . Then (H_C, L_H) is isomorphic, as a labeled graph, to the minimal right-resolving presentation of $L_H(X_C)$*

Proof. Note that (H_C, L_H) is right-resolving and follower-separated. The lemma follows therefore from Corollary 3.3.19 in [LM]. \square

Let Y be a sofic subshift. When $w \in \mathbb{W}(Y)$, set

$$F(w) := \{z \in Y[0, \infty) : wz \in Y[0, \infty)\}.$$

A dual version of the following lemma has appeared as Lemma 4.7 of [J].

Lemma 2.20. *For every $y \in Y$ there is an $N \in \mathbb{N}$ such that*

$$F(y_{[-n, -1]}) = F(y)$$

for all $n \geq N$.

Proof. Note that $F(y_{[-n-1, -1]}) \subseteq F(y_{[-n, -1]})$ for all n and that

$$F(y) = \bigcap_n F(y_{[-n, -1]}).$$

Since $\{F(w) : w \in \mathbb{W}(Y)\}$ is a finite set there must be a sequence $n_1 < n_2 < \dots$ in \mathbb{N} such that $F(y_{[-n_i, -1]}) = F(y_{[-n_1, -1]})$ for all i . Set $N := n_1$. \square

A word $w \in \mathbb{W}(Y)$ is *synchronizing* when $u, v \in \mathbb{W}(Y)$, $uw, wv \in \mathbb{W}(Y) \Rightarrow uwwv \in \mathbb{W}(Y)$.

Lemma 2.21. *Let (H, L_H) be a right-resolving, regular and follower-separated labeled graph and C a terminal component in H . For each $v \in C$ there is a finite path γ in H such that $t_H(\gamma) = v$ and $L_H(\gamma)$ is synchronizing for $L_H(X_H)$.*

Proof. Let $v \in C$. Since (H, L_H) is regular there is an element $z \in X_H$ such that $t_H(z_{(-\infty, -1]}) = v$ and $f_H(v) = F(L_H(z))$. By Lemma 2.20 there is an $N \in \mathbb{N}$ such that $F(L_H(z_{[-N, -1]})) = F(L_H(z))$. By Proposition 3.3.16 in [LM] there is a word $u \in \mathbb{W}(L_H(X_H))$ such that $L_H(z_{[-N, -1]})u$ is synchronizing for $L_H(X_H)$. Since u is a prefix of an element from $F(L_H(z)) = f_H(v)$ there is a path γ_1 in H such that $s_H(\gamma_1) = v$ and $L_H(\gamma_1) = u$. Note that $t_H(\gamma_1) \in C$ since C is a terminal component. Since H_C is strongly connected there is a path γ_2 in H_C such that $s_H(\gamma_2) = t_H(\gamma_1)$ and $t_H(\gamma_2) = v$. Set $\gamma := z_{[-N, -1]}\gamma_1\gamma_2$ and note that $L_H(\gamma)$ is synchronizing for $L_H(X_H)$ because $L_H(z_{[-N, -1]})u$ is a prefix of $L_H(\gamma)$. \square

Proposition 2.22. *Let Y be an irreducible sofic shift space. There is only one terminal component C in $\mathbb{K}(Y)$ and $(\mathbb{K}(Y)_C, L_{\mathbb{K}(Y)})$ is isomorphic to the minimal right-resolving presentation of Y .*

Proof. Let v, w both be vertexes of a terminal component in $\mathbb{K}(Y)$. By Lemma 2.21 there are synchronizing words s, t for Y such that $v = F(s)$ and $w = F(t)$. Let $u \in \mathbb{W}(Y)$. Since Y is irreducible there is a word $x \in \mathbb{W}(Y)$ such that $u = x_{[i,j]}$ for some $i \leq j$ and $sxt \in \mathbb{W}(Y)$. By definition of $\mathbb{K}(Y)$ there is therefore a path in $\mathbb{K}(Y)$ with label xt from $v = F(s)$ to $w = F(t)$. It follows that v and w lie in the same component C of $\mathbb{K}(Y)$. Furthermore, since the word u was arbitrary it follows that $L_{\mathbb{K}(Y)}(X_C) = Y$. It follows then from Lemma 2.19 that $(\mathbb{K}(Y)_C, L_{\mathbb{K}(Y)})$ is isomorphic to the minimal right-resolving presentation of Y . \square

Lemma 2.23. *Let G and H be directed graphs with unique terminal components, C and C' , respectively. Assume that $\chi : X_G \rightarrow X_H$ is a conjugacy. It follows that $\chi(X_C) = X_{C'}$.*

Proof. The periodic points p of X_C are characterized by the property that for all $x \in X_G$ there is an element $y \in X_G$ and a natural numbers $k \in \mathbb{N}$ such that $\lim_{i \rightarrow \infty} d(\sigma^{-i}(x), \sigma^{-i}(y)) = 0$ and $\lim_{i \rightarrow \infty} d(\sigma^i(y), \sigma^{i+k}(p)) = 0$. There is a similar characterization of the periodic elements of $X_{C'}$ and we conclude therefore that $\chi(\text{Per } X_C) \subseteq \text{Per}(X_{C'})$. Since the periodic points of X_C are dense in X_C it follows that $\chi(X_C) \subseteq X_{C'}$. Hence, by symmetry, $\chi(X_C) = X_{C'}$. \square

Corollary 2.24. *Let $\psi : Y \rightarrow Z$ be a conjugacy of irreducible sofic subshifts. Let C and C' be the terminal components in $\mathbb{K}(Y)$ and $\mathbb{K}(Z)$, respectively, and let $\psi_{\mathbb{K}} : X_{\mathbb{K}(Y)} \rightarrow X_{\mathbb{K}(Z)}$ be the conjugacy from Theorem 2.11. Then $\psi_{\mathbb{K}}(X_C) = X_{C'}$.*

Corollary 2.25. *(Corollary (2.16) in [Krl]) Let $\psi : Y \rightarrow Z$ be a conjugacy of irreducible sofic subshifts. Let (G, L_G) and (H, L_H) be the minimal right-resolving presentations of Y and Z , respectively. It follows that there is unique conjugacy $\phi : X_G \rightarrow X_H$ such that*

$$(10) \quad \begin{array}{ccc} X_G & \xrightarrow{\phi} & X_H \\ L_G \downarrow & & \downarrow L_H \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes.

Proof. The existence of ϕ follows from Corollary 2.24, Proposition 2.22 and Theorem 2.11. The uniqueness follows here by the same argument which was used by Krieger in [Krl]: The elements of Y that contain synchronized words arbitrarily far 'to the left' have unique pre-image under L_G because they are magic for (G, L_G) by Exercise 3.3.3 in [LM] or Lemma 4.3 in [MMTW]. (These elements were called F -finitary in [Krl].) Since G is strongly connected it follows from Proposition 3.3.16 in [LM] that they are dense in X_G . Thus the elements $x \in X_G$ for which $L_G^{-1}(L_G(x)) = \{x\}$ are dense in X_G . Denote this set by \mathcal{X}_G . If $\phi : X_G \rightarrow X_H$ is a conjugacy such that (10) commutes, $\phi(\mathcal{X}_G) = \mathcal{X}_H$. If ϕ_1 and ϕ_2 are both conjugacies that make the diagram (10) commute, and $x \in \mathcal{X}_G$, we have that $\phi_i(x) \in \mathcal{X}_H$ and

$$L_H(\phi_1(x)) = \psi(L_G(x)) = L_H(\phi_2(x)),$$

implying that $\phi_1(x) = \phi_2(x)$. Since \mathcal{X}_G is dense in X_G , it follows that $\phi_1 = \phi_2$. \square

As noted in the introduction, an alternative proof of Corollary 2.25, without the uniqueness part, was given in [BKM].

3. ADDITIONAL PROPERTIES OF THE FUTURE COVER

3.1. A universal property of the future cover $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$. Let (H, L_H) be a labeled graph. A subset $U \subseteq V_H$ of vertexes in H is a *hereditary* subset when $e \in E_H$, $s_H(e) \in U \Rightarrow t_H(e) \in U$. When this holds we let H_U denote the graph with $V_{H_U} := U$ and $E_{H_U} := \{e \in E_H : t_H(e) \in U\}$. By restriction the labeling L_H defines a labeling of H_U and we denote the resulting labeled graph by (H_U, L_H) . We call this a *hereditary labeled subgraph* of (H, L_H) . Lemma 2.18 can then be improved to the following.

Proposition 3.1. *Assume that (G, L_G) is a right-resolving and regular presentation of the sofic shift Y . It follows that there is a labeled graph homomorphism $\theta : (G, L_G) \rightarrow (\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ such that*

$$\theta(\mathcal{R}(L_G)) \subseteq \mathcal{R}(L_{\mathbb{K}(Y)}),$$

and $\theta(V_G)$ is a hereditary subset of $V_{\mathbb{K}(Y)}$. If, furthermore, (G, L_G) is also follower-separated, θ is a labeled graph isomorphism of (G, L_G) onto the hereditary labeled subgraph

$$(\mathbb{K}(Y)_{\theta(V_G)}, L_{\mathbb{K}(Y)}).$$

Proof. By Lemma 2.18 and its proof there is a labeled-graph homomorphism $\theta : (G, L_G) \rightarrow (\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ such that $\theta(v) = F(L_G(x))$ where $x \in X_G$, $t_G(x_{(-\infty, -1]}) = v$ and $f_G(v) = F(L_G(x))$. Let $F(\theta(v)) \xrightarrow{a} F(z)$ be an arrow in $\mathbb{K}(Y)$. Then a is the first letter in an element $w \in F(L_G(x)) = f_G(v)$, and there is therefore a right-infinite path γ in G such that $s_G(\gamma) = v$ and $L_G(\gamma) = w$. The first arrow in γ is an arrow $e \in E_G$ with $s_G(e) = v$ and $L_G(e) = a$. There is therefore an arrow $F(\theta(v)) \xrightarrow{a} \theta(t_G(e))$ in $\mathbb{K}(Y)$. Since $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ is right-resolving this implies that $F(z) = \theta(t_G(e))$, proving that $\theta(V_G)$ is a hereditary subset of $V_{\mathbb{K}(Y)}$. The remaining assertions follow now from Lemma 2.18. \square

Corollary 3.2. *Let Y be a sofic shift. Any labeled graph which presents Y and is right-resolving, regular and follower-separated is isomorphic to a hereditary labeled subgraph of $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$.*

Corollary 3.3. *Let Y be a sofic shift space. For any presentation (G, L_G) of Y which is right-resolving, regular and follower-separated, the number $\#V_G$ of vertexes is less than or equal to the number of follower sets in Y . The future cover $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ is the only presentation of Y , up to isomorphism of labeled graphs, which is right-resolving, regular and follower-separated, and for which the number of vertexes is equal to the number of follower sets in Y .*

The future cover has one more property which seems worthwhile to point out; it is predecessor-separated: Consider a labeled graph (H, L_H) , and $v \in V_H$ a vertex in H . The *predecessor set* of v is the set

$$p_H(v) := \{L_H(z) : z \in X_H(-\infty, -1], t_H(z) = v\}.$$

(H, L_H) is *predecessor-separated* when $p_H(v) = p_H(w) \Rightarrow v = w$.

Lemma 3.4. *Let (H, L_H) be a right-resolving, regular and follower-separated labeled graph. Then (H, L_H) is predecessor-separated.*

Proof. Assume $v, w \in V_H$ and $p_H(v) = p_H(w)$. Since v is regular there is a $z \in X_H$ such that $t_H(z_{(-\infty, -1]}) = v$ and $f_H(v) = F(L_H(z))$. Then $L_H(z_{(-\infty, -1]}) \in p_H(v) = p_H(w)$, so there is

a $u \in X_H(-\infty, -1]$ such that $t_H(u) = w$ and $L_H(u_{(-\infty, -1]}) = L_H(z_{(-\infty, -1]})$. It follows that $f_H(w) \subseteq F(L_H(z)) = f_H(v)$. By symmetry $f_H(v) \subseteq f_H(w)$ and hence $f_H(v) = f_H(w)$, which implies that $v = w$ since we assume that (H, L_H) is follower-separated. \square

Corollary 3.5. *The labeled graph $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ is both follower-separated and predecessor separated.*

Proof. Combine Lemma 2.12, Lemma 2.13 and Lemma 3.4. \square

Example 3.6. If we denote the sofic shift presented by the labeled graph in Figure 1 by Y the future cover $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ of Y is isomorphic to the following labeled graph.




FIGURE 4. The future cover of the sofic shift presented by the labeled graph in Figure 1.

The labeled graph in Figure 1 is not isomorphic to that in Figure 4 although the two graphs have the same number of vertexes. Thus regularity is a crucial assumption both in Corollary 3.3, and indeed in Proposition 3.1.

Example 3.7. The following labeled graph (G, L_G) is right-resolving, follower-separated and regular (synchronizing, in fact). It is the minimal right-resolving presentation of the even shift.




FIGURE 5. A right-resolving, regular and follower-separated presentation of the even shift.

The future cover of the even shift is depicted in Figure 3. It has one vertex more than the minimal right-resolving presentation, showing that the condition of maximality of the number of vertexes in Corollary 3.3 can not be dropped.

3.2. A universal property of the future cover $L_G : X_{\mathbb{K}(Y)} \rightarrow Y$. By Proposition 4 in [BKM] the minimal right-resolving cover of an irreducible sofic shift has the following universal property:

Proposition 3.8. *Let Y be an irreducible sofic shift and (G, L_G) the minimal right-resolving presentation of Y . Let X be an irreducible SFT and $\pi : X \rightarrow Y$ a right-closing factor map. There is a sliding block code $\rho : X \rightarrow X_G$ such that*

$$\begin{array}{ccc} X & \xrightarrow{\rho} & X_G \\ \pi \downarrow & \swarrow L_G & \\ Y & & \end{array}$$

commutes.

This property can be considered as a minimality condition with respect to irreducible and right-closing covers. In general, or more specifically when Y is not of almost finite type, the minimal right-resolving cover is not minimal with respect to arbitrary irreducible covers by Corollary 13 in [BKM]. It is also not minimal with respect to general right-resolving covers; for example the future cover of the even shift which is depicted in Figure 3 does not factor through the minimal right-resolving cover of the even shift. However, its minimality property can be used to characterise it up to isomorphism.

Proposition 3.9. *Let Y be an irreducible SFT and $\pi : X \rightarrow Y$ a factor map with X an SFT. Then π is isomorphic to the minimal right-resolving cover of Y if and only if*

- a) X is irreducible,
- b) π is right-closing,
- c) π is injective on the set of doubly transitive points in X , and
- d) $\pi : X \rightarrow Y$ is minimal with respect to irreducible right-closing covers of Y .

Proof. The following arguments are all implicitly contained in [BKM]. First, it is well-known from [BKM] and [LM] that the minimal right-resolving cover has all four properties, and hence so has any cover isomorphic to it. So assume that $\pi : X \rightarrow Y$ is a cover for which they all hold. We must show that π is then isomorphic to the minimal right-resolving cover $L_G : X_G \rightarrow Y$ of Y . For this note that there are sliding block codes $\rho : X \rightarrow X_G$ and $\rho' : X_G \rightarrow X$ such that

$$\begin{array}{ccccc} X_G & \xrightarrow{\rho'} & X & \xrightarrow{\rho} & X_G \\ & \searrow L_G & \downarrow \pi & \swarrow L_G & \\ & & Y & & \end{array}$$

and

$$\begin{array}{ccccc} X & \xrightarrow{\rho} & X_G & \xrightarrow{\rho'} & X \\ & \searrow \pi & \downarrow L_G & \swarrow \pi & \\ & & Y & & \end{array}$$

both commute. An application of Lemma 9.1.13 in [LM] shows that ρ and ρ' both maps doubly transitive points to doubly transitive points, so condition c) implies that $\rho \circ \rho' = \text{id}_{X_G}$ and $\rho' \circ \rho = \text{id}_X$, first on doubly transitive points and then all over by continuity. \square

We need some preparations in order to discuss a similar characterization of the future cover.

Lemma 3.10. *Let (H, L_H) be a regular right-resolving presentation of the sofic shift Y . For $x \in X_H$ and $N \in \mathbb{N}$ there is a $y \in \mathcal{R}(L_H)$ such that $y_i = x_i$ for $i \geq -N$.*

Proof. Let $N \in \mathbb{N}$. Since $s_H(x_{[-N, \infty)})$ is regular there is a $z \in X_H$ such that $t_H(z_{(-\infty, -1]}) = s_H(x_{[-N, \infty)})$ and $f_H(s_H(x_{[-N, \infty)})) = F(L_H(z))$. It follows from Lemma 2.20 that there is an $M \in \mathbb{N}$ such that $F(L_H(z)) = F(L_H(z_{[-M, -1]}))$. Set $w_1 := z_{[-M, -1]}$. Since $s_H(w_1)$ is regular we can repeat the argument to get a finite path w_2 in H such that $t_H(w_2) = s_H(w_1)$ and $F(L_H(w_2)) = f_H(s_H(w_1))$. Repeat this argument inductively and set

$$y := \cdots w_3 w_2 w_1 x_{[-N, \infty)}.$$

For $j \in \{-N - \sum_{k=1}^n |w_k| : n \in \mathbb{N}\}$ we have by construction that

$$f_H(t_H(y_{(-\infty, j-1]})) = F(\sigma^j(y)).$$

As in the proof of Lemma 2.2, by using that (H, L_H) is right-resolving it follows from this that the same holds for all $j \in \mathbb{Z}$; i.e. $y \in X_H$ is regular. \square

Lemma 3.11. *Let (H, L_H) be a right-resolving presentation of the sofic shift Y . Then (H, L_H) is regular if and only if $\mathcal{R}(L_H)$ is dense in X_H .*

Proof. One implication follows from Lemma 3.10. The other from Lemma 2.6. \square

Definition 3.12. *Let $\pi : X \rightarrow Y$ be a sliding block code between subshifts. An element $x \in X$ is regular for π when $\pi : \mathbb{U}(x) \rightarrow \mathbb{U}(\pi(x))$ is surjective. The set of elements $x \in X$ that are regular for π will be denoted by $\mathcal{R}(\pi)$. We say that π is regular when $\mathcal{R}(\pi)$ is dense in X .*

Proposition 3.13. *Let $\pi : X \rightarrow Y$ be a factor code where X is an SFT. Assume that π is right-closing and regular. There is a sliding block code $\rho : X \rightarrow X_{\mathbb{K}(Y)}$ such that*

$$\begin{array}{ccc} X & \xrightarrow{\rho} & X_{\mathbb{K}(Y)} \\ \pi \downarrow & \swarrow L_{\mathbb{K}(Y)} & \\ Y & & \end{array}$$

commutes.

Proof. By Lemma 2.1 there is a right-resolving labeled graph (G, L_G) and a conjugacy $\psi : X \rightarrow X_G$ such that

$$\begin{array}{ccc} X & \xrightarrow{\psi} & X_G \\ \pi \downarrow & \swarrow L_G & \\ Y & & \end{array}$$

commutes. Note that $\psi(\mathcal{R}(\pi)) = \mathcal{R}(L_G)$. It follows from Lemma 3.11 that (G, L_G) is regular since π is. It follows then from Lemma 2.18 that there is a sliding block code $\mu : X_G \rightarrow X_{\mathbb{K}(Y)}$ such that

$$\begin{array}{ccc} X_G & \xrightarrow{\mu} & X_{\mathbb{K}(Y)} \\ L_G \downarrow & \swarrow L_{\mathbb{K}(Y)} & \\ Y & & \end{array}$$

commutes. Set $\rho := \mu \circ \psi$. \square

In the language of [BKM] which we have adopted above, Proposition 3.13 shows that the future cover is minimal with respect to regular and right-closing factor maps. The regularity condition on π can not be removed: The right-resolving labeled graph (G, L_G) in Figure 7 presents a sofic shift Y whose future cover $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ is obtained by removing the vertex 10 in Figure 7, and L_G does not factor through $L_{\mathbb{K}(Y)}$. The future cover is therefore not minimal with respect to arbitrary right-resolving factor maps in general.

Lemma 3.14. *Let $\pi : X \rightarrow Y$, $\pi' : X' \rightarrow Y$ and $\rho : X \rightarrow X'$ be sliding block codes such that $\pi' \circ \rho = \pi$. Then $\rho(\mathcal{R}(\pi)) \subseteq \mathcal{R}(\pi')$.*

Proof. Let $x \in \mathcal{R}(\pi)$ and consider an element $y \in \mathbb{U}(\pi'(\rho(x))) = \mathbb{U}(\pi(x))$. There is then an element $z \in \mathbb{U}(x)$ such that $\pi(z) = y$. Since $\rho(z) \in \mathbb{U}(\rho(x))$ and $\pi'(\rho(z)) = y$, it follows that $\rho(x) \in \mathcal{R}(\pi')$. \square

Proposition 3.15. *Let $\pi : X \rightarrow Y$ be a factor map with X an SFT. Then π is isomorphic to the future cover $L_{\mathbb{K}(Y)} : X_{\mathbb{K}(Y)} \rightarrow Y$ of Y if and only if*

- a) π is regular,
- b) π is right-closing,
- c) π is injective on the set $\mathcal{R}(\pi)$ of regular points, and
- d) $\pi : X \rightarrow Y$ is minimal with respect to regular and right-closing factor maps.

Proof. $L_{\mathbb{K}(Y)}$ is regular by Lemma 3.11 and Lemma 2.13, and right-closing because the future cover is right-resolving. By Proposition 3.13 it is minimal with respect to regular and right-closing factor maps. That $L_{\mathbb{K}(Y)}$ is injective on $\mathcal{R}(L_{\mathbb{K}(Y)})$ follows from Lemma 2.8.

Assume $\pi : X \rightarrow Y$ is a factor map with X an SFT such that a), b), c) and d) hold. It follows that there are sliding block codes $\rho : X \rightarrow X_{\mathbb{K}(Y)}$ and $\rho' : X_{\mathbb{K}(Y)} \rightarrow X$ such that

$$\begin{array}{ccccc} X_{\mathbb{K}(Y)} & \xrightarrow{\rho'} & X & \xrightarrow{\rho} & X_{\mathbb{K}(Y)} \\ & \searrow L_{\mathbb{K}(Y)} & \downarrow \pi & \swarrow L_{\mathbb{K}(Y)} & \\ & & Y & & \end{array}$$

and

$$\begin{array}{ccccc} X & \xrightarrow{\rho} & X_{\mathbb{K}(Y)} & \xrightarrow{\rho'} & X \\ & \searrow \pi & \downarrow L_{\mathbb{K}(Y)} & \swarrow \pi & \\ & & Y & & \end{array}$$

both commute. It follows from Lemma 3.14 that ρ and ρ' both map regular points to regular points. Since π and $L_{\mathbb{K}(Y)}$ are regular it follows that $\rho \circ \rho' = \text{id}_{X_{\mathbb{K}(Y)}}$ and $\rho' \circ \rho = \text{id}_X$, first on regular points and then all over by continuity. \square

3.3. The relation between the future cover and the follower set graph. In this section we show how the future cover is related to another commonly used presentation of sofic shifts, the follower set graph, cf. page 73 in [LM].

Given a right-resolving labeled graph (H, L_H) , the set of its regular vertexes is a hereditary subset by Lemma 2.2 and we denote the corresponding hereditary labeled subgraph by (H_{reg}, L_H) . As shown by the example depicted in Figure 1, the sofic shift presented by (H_{reg}, L_H) can be smaller than that presented by (H, L_H) .

Proposition 3.16. *Let Y be a sofic shift and (G, \mathcal{L}) the follower set graph of Y . The future cover $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ of Y is isomorphic to $(G_{\text{reg}}, \mathcal{L})$ as a labeled graph.*

Proof. The proof uses the observation that two words $w_1, w_2 \in \mathbb{W}(Y)$ have the same follower sets as defined in [LM] if and only if $F(w_1) = F(w_2)$. Let $y \in Y$. By Lemma 2.20 there is an $N \in \mathbb{N}$ such that $F(y) = F(y_{[-n, -1]})$ for all $n \geq N$. If $z \in Y$ and $F(y) = F(z)$ there is an $M \in \mathbb{N}$ such that $F(z_{[-n, -1]}) = F(y_{[-n, -1]})$ for all $n \geq M$. It follows that we can define a map $\chi : V_{\mathbb{K}(Y)} \rightarrow V_G$ such that $\chi(F(y)) = F(y_{[-n, -1]})$ for all large $n \in \mathbb{N}$. This map, χ , is clearly injective. For $y, z \in Y$ there is a labeled arrow $F(y) \xrightarrow{a} F(z)$ in $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ if and only if $F(z) = \frac{F(y)}{a}$ if and only if $F(z_{[-n, -1]}) = \frac{F(y_{[-n, -1]})}{a}$ for all large n if and only if there is a labeled arrow $\chi(F(y)) \xrightarrow{a} \chi(F(z))$ in G . This shows that χ is an embedding of labeled graphs. To show that $\chi(V_{\mathbb{K}(Y)})$ is a hereditary subset, let $w \in \mathbb{W}(Y)$ and assume that $\chi(F(y)) \xrightarrow{a} F(w)$ in G . Then $F(w) = \frac{F(y)}{a}$, and hence $F(w) = F(z)$, where $z \in Y$ is any element with $z_{(-\infty, -1]} = y_{(-\infty, -1]}a$. It follows that $F(w) = \chi(F(z))$.

Since $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ is regular by Lemma 2.13 and since the range of χ is a hereditary subgraph of (G, \mathcal{L}) it follows that $\chi(V_{\mathbb{K}(Y)})$ consist only of regular vertexes in G . Conversely, if $w \in \mathbb{W}(Y)$ and $F(w)$ is a regular vertex in G , it follows from the defining relation (1) that $F(w) = F(z)$ for some $z \in Y$, proving that $F(w) = \chi(F(z)) \in \chi(V_{\mathbb{K}(Y)})$. \square

Example 3.17. Consider the following labeled graph.




FIGURE 6. The minimal right-resolving presentation of a mixing sofic shift whose follower set graph is larger than its future cover.

The follower sets of the vertexes a, b, c, d, e are also follower sets of (synchronizing) words in $\mathbb{W}(Y)$. Besides these the follower sets of words from $\mathbb{W}(Y)$ include only the follower sets of the words 10 and 0, resulting in the following labeled graph depicting the follower set graph of Y .




FIGURE 7. The follower set graph of the sofic shift presented by Figure 6.

The vertex 10 is not regular while all the other are, and hence the future cover of Y is the labeled graph obtained from the graph in Figure 7 by deleting the vertex 10 and the arrows it emits.

4. AN ADDENDUM TO KRIEGER'S THEOREM

4.1. Subset constructions. In this section by a *dynamical system* we mean a pair (X, ϕ) where X is a compact metrizable space and $\phi : X \rightarrow X$ is a homeomorphism of X onto itself.

Let (X, ϕ) be a dynamical system and d a metric for the topology of X . The set of non-empty compact subsets K of X can be equipped with a metric, the *Hausdorff metric* d_{Haus} , defined by

$$d_{Haus}(K, K') := \max \left\{ \sup_{x \in K} \left(\inf_{y \in K'} d(x, y) \right), \sup_{y \in K'} \left(\inf_{x \in K} d(x, y) \right) \right\}.$$

This turns the set of non-empty compact subsets of X into a compact metric space

$$(X^\#, d_{Haus}).$$

See for example Section 2.6 in [B].

When (X, d) and (X', d') are two compact metric spaces and $\psi : X \rightarrow X'$ is a continuous map we get a continuous map $\psi^\# : X^\# \rightarrow X'^\#$ defined such that

$$\psi^\#(K) := \psi(K)$$

for $K \in X^\#$. In particular, we get from (X, ϕ) a new dynamical system $(X^\#, \phi^\#)$. More generally when (X, ϕ) and (Y, ϕ') are dynamical systems and $\pi : X \rightarrow Y$ is a map of dynamical systems in the sense that $\pi \circ \phi = \phi' \circ \pi$, the set

$$X^\pi := \{K \in X^\# : \pi(x) = \pi(y) \forall x, y \in K\}$$

is a closed $\phi^\#$ -invariant subset of $X^\#$ and $(X^\pi, \phi^\#)$ is also a dynamical system. When π is surjective and hence a factor map, then so is $\pi^\# : (X^\pi, \phi^\#) \rightarrow (Y, \phi')$.

The following is now a straightforward observation.

Lemma 4.1. *Consider two conjugacies $\psi' : (X, \phi) \rightarrow (X', \phi')$ and $\psi : (Y, \mu) \rightarrow (Z, \mu')$ of dynamical systems, and factor maps $\pi : (X, \phi) \rightarrow (Y, \mu)$ and $\pi' : (X', \phi') \rightarrow (Z, \mu')$ such that*

$$\begin{array}{ccc} X & \xrightarrow{\psi'} & X' \\ \pi \downarrow & & \downarrow \pi' \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes. Then $\psi'^{\#} : X^{\pi} \rightarrow X'^{\pi'}$ is a conjugacy, and

$$\begin{array}{ccc} X^{\pi} & \xrightarrow{\psi'^{\#}} & X'^{\pi'} \\ \pi^{\#} \downarrow & & \downarrow \pi'^{\#} \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes.

4.2. Subset constructions for sofic shifts. Let (H, L_H) be a labeled graph and $Y := L_H(X_H)$ the presented sofic shift. We define a labeled graph $(H'', L_{H''})$ as follows. The set $V_{H''}$ of vertexes consists of the non-empty subsets F of vertexes from V_H . For $F, F' \in V_{H''}$ there is an edge $e' \in E_{H''}$ with $s_{H''}(e') = F$ and $t_{H''}(e') = F'$ when there is a symbol $a \in A$, the alphabet of Y , such that

$$F \subseteq \{s_H(e) : e \in E_H, L_H(e) = a\}$$

and

$$F' = \{t_H(e) : e \in E_H, s_H(e) \in F, L_H(e) = a\}.$$

We set then $L_{H''}(e') = a$. While the graph H'' in principle contains all subsets of V_H as vertexes, it is clear from the definition of arrows in $E_{H''}$ that typically only a much smaller collection of such subsets will occur in elements of $X_{H''}$. In particular, when we consider H'' in the following it will be assumed that H'' has been trimmed by repeatedly removing sources and sinks to obtain a graph that does not contain neither.

Lemma 4.2. *Assume that (H, L_H) is right-resolving. There is a conjugacy $\kappa : (X_H^{L_H}, \sigma^{\#}) \rightarrow (X_{H''}, \sigma)$ such that*

$$\begin{array}{ccc} X_H^{L_H} & \xrightarrow{\kappa} & X_{H''} \\ & \searrow L_H^{\#} \quad \swarrow L_{H''} & \\ & Y & \end{array}$$

commutes.

Proof. Let $K \in X_H^{L_H}$. Note that K is a finite set since (H, L_H) is right-resolving. Define

$$F_i := \{s_H(k_i) : k \in K\}$$

for $i \in \mathbb{Z}$. For each $i \in \mathbb{Z}$ there is a unique edge $e_i(K)$ in $E_{H''}$ from F_i to F_{i+1} with label $L_{H''}(e_i(K)) = L_H(k_i)$ for all $k \in K$, and we set

$$\kappa(K) := (e_i(K))_{i \in \mathbb{Z}} \in X_{H''}.$$

Since $e_i(\sigma(K)) = e_{i+1}(K)$ we find that

$$\kappa \circ \sigma^\#(K) = \kappa(\sigma(K)) = (e_i(\sigma(K)))_{i \in \mathbb{Z}} = (e_{i+1}(K))_{i \in \mathbb{Z}} = (\sigma(\kappa(K)))_{i \in \mathbb{Z}},$$

proving that $\kappa \circ \sigma^\# = \sigma \circ \kappa$. Note also that $L_{H''}(\kappa(K)) = L_H^\#(K)$.

To construct an inverse map $\kappa^{-1} : X_{H''} \rightarrow X_H^{L_H}$ to κ , let $x = (x_i)_{i \in \mathbb{Z}} \in X_{H''}$. Since (H, L_H) is right-resolving, $\#_{s_{H''}}(x_i) \geq \#_{s_{H''}}(x_{i+1})$, and we choose $N \in \mathbb{N}$ such that $\#_{s_{H''}}(x_i) = \#_{s_{H''}}(x_N)$ for all $i \leq N$. For each $v \in s_{H''}(x_N)$ there is a unique path $\gamma^v \in X_H$ such that $s_H(\gamma_N^v) = v$ and $L_H(\gamma_v) = L_{H''}(x)$. Setting

$$\kappa^{-1}(x) := \{\gamma_v : v \in s_{H''}(x_N)\}$$

we obtain a well-defined map $\kappa^{-1} : X_{H''} \rightarrow X_H^{L_H}$. It is straightforward to check that κ^{-1} is the inverse of κ .

Since $X_H^{L_H}$ and $X_{H''}$ are compact Hausdorff spaces it suffices now to show that κ is continuous. For this we use the metric d on X_H defined such that $d(x, y) = \frac{1}{n+1}$, where $n = \infty$ when $x = y$ and otherwise $n = \min \{j \in \mathbb{N} \cup \{0\} : x_{[-j, j]} \neq y_{[-j, j]}\}$. Consider then two elements $K, K' \in X_H^{L_H}$. If $d_{Haus}(K, K') < \frac{1}{N}$, it follows that

$$\{k_i : k \in K\} = \{k_i : k \in K'\}$$

for $-N \leq i \leq N$, and hence also that $\kappa(K)_i = e_i(K) = e_i(K') = \kappa(K')_i$ for $-N \leq i \leq N$. This shows that κ is continuous. \square

Corollary 4.3. *Let Y and Z be sofic shifts and (G, L_G) , (H, L_H) right-resolving presentations of Y and Y' , respectively. Assume that there are conjugacies $\psi : Y \rightarrow Z$ and $\phi : X_G \rightarrow X_H$ such that*

$$\begin{array}{ccc} X_G & \xrightarrow{\phi} & X_H \\ L_G \downarrow & & \downarrow L_H \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes. There is a conjugacy $\phi'' : X_{G''} \rightarrow X_{H''}$ such that

$$\begin{array}{ccc} X_{G''} & \xrightarrow{\phi''} & X_{H''} \\ L_{G''} \downarrow & & \downarrow L_{H''} \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes.

Proof. Combine Lemma 4.1 and Lemma 4.2. \square

Thus $(X_{G''}, L_{G''})$ is a weakly canonical cover when (G, L_G) is. However, the conjugacy ϕ'' may not be unique and there is no analogue to the right-inverse for the labeling map as in Krieger's theorem. We continue therefore to identify a labeled subgraph of $(G'', L_{G''})$ which does give rise to a strongly canonical cover.

Since (H, L_H) is right-resolving, $L_H^{-1}(y)$ is a finite set and hence an element of $X_H^{L_H}$ for each $y \in Y$. We define $\beta_H : Y \rightarrow X_H^{L_H}$ such that

$$\beta_H(y) := \kappa(L_H^{-1}(y)),$$

where $\kappa : X_H^{L_H} \rightarrow X_{H''}$ is the conjugacy from Lemma 4.2. Note that

$$L_{H''} \circ \beta_Y = \text{id}_Y.$$

From the proof of Lemma 4.2 we get the following.

Lemma 4.4. *Let $y \in Y$. Then $\beta_H(y) \in X_{H''}$ is the unique element with the properties that $L_{H''}(\beta_H(y)) = y$ and $s_{H''}(\beta_H(y)_j) = \{s_H(x_j) : x \in L_H^{-1}(y)\}$ for all $j \in \mathbb{Z}$.*

To proceed we need some terminology. Let X be a subshift. We say that two elements $x, y \in X$ are *backward asymptotic* and that x is *backward asymptotic to y* when there is an $N \in \mathbb{Z}$ such that $x_i = y_i$ for all $i \leq N$. When $A \subseteq X$ is a subset and $x \in X$ we say that x is *backward asymptotic to A* when x is backward asymptotic to an element of A . When A is a subshift of X we say that $x \in X$ is *backward transitive to A* when x is backward asymptotic to A and every word $w \in \mathbb{W}(A)$ appears infinitely often to the left in x , in the sense that for all $N \in \mathbb{Z}$ there are $i \leq j \leq N$ such that $w = x_{[i,j]}$.

Lemma 4.5. *Let C be a component in H'' such that there is an element $y \in Y$ with $\beta_H(y)$ backward asymptotic to X_C , and let $z \in X_{H''}$ be an element backward transitive to X_C . Then $z = \beta_H(L_{H''}(z))$.*

Proof. Set $y' := L_{H''}(z)$. Then y' is backward transitive to $L_{H''}(X_C)$. Let C' be the component in H'' with the property that $\beta_H(y')$ is backward asymptotic to $X_{C'}$. Note that the number

$$M(C) := \#s_{H''}(x_j)$$

is the same for all elements $x \in X_C$ and all $j \in \mathbb{Z}$. Since $L_{H''}(z) = y'$ we have that

$$(11) \quad s_{H''}(z_j) \subseteq s_{H''}(\beta_H(y')_j)$$

for all $j \in \mathbb{Z}$, implying that $M(C) \leq M(C')$ since z is backward asymptotic to X_C and $\beta_H(y')$ to $X_{C'}$. On the other hand, if w is a word in $L_{H''}(X_C)$ it occurs infinitely often to the left in y' . Since $L_{H''}^{-1}(w)$ is a finite set and $L_{H''}(\beta_H(y')) = y'$ it follows that $w = L_{H''}(w')$ for some word w' appearing infinitely often to the left in $\beta_H(y')$. Then $w' \in \mathbb{W}(X_{C'})$ and we conclude therefore that $L_{H''}(X_C) \subseteq L_{H''}(X_{C'})$. It follows that there is a $z' \in X_{C'}$ such that y is backward asymptotic to $L_{H''}(z')$, and there is therefore a $N \in \mathbb{Z}$ such that $s_{H''}(z'_j) \subseteq s_{H''}(\beta_H(y)_j)$ for all $j \leq N$. It follows that $M(C') \leq M(C)$ and therefore also that $M(C) = M(C')$. Then (11) implies that $z = \beta_H(y') = \beta_H(L_{H''}(z))$. \square

Definition 4.6. *We denote by H' the hereditary subgraph of H'' generated by the components C of H'' with the property that there is a point $y \in Y$ such that $\beta_H(y)$ is backward asymptotic to X_C .*

To simplify the notation, set $L_{H'} := L_{H''}|_{H'}$.

Lemma 4.7. $X_{H'} = \overline{\beta_H(Y)}$. In particular, $(X_{H'}, L_{H'})$ is a right-resolving presentation of Y .

Proof. By definition of H' every element of $X_{H'}$ can be approximated by elements z with the property that z is backward transitive to X_C , where C is the component in H'' with the property that $\beta_H(y)$ is backward asymptotic to X_C for some $y \in Y$. Since $z \in \beta_H(Y)$ by Lemma 4.5, it follows that $X_{H'} \subseteq \overline{\beta_H(Y)}$. On the other hand, every element of $\beta_H(Y)$ is backward asymptotic to some X_C , where C , by definition, is a component in H' . Hence $\beta_H(Y) \subseteq X_{H'}$. \square

Theorem 4.8. *Let (G, L_G) and (H, L_H) be right-resolving presentations of the sofic shifts Y and Z , respectively. Assume that there are conjugacies $\psi : Y \rightarrow Z$ and $\phi : X_G \rightarrow X_H$ such that*

$$\begin{array}{ccc} X_G & \xrightarrow{\phi} & X_H \\ L_G \downarrow & & \downarrow L_H \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes. There is a unique conjugacy $\phi' : X_{G'} \rightarrow X_{H'}$ such that

$$\begin{array}{ccc} X_{G'} & \xrightarrow{\phi'} & X_{H'} \\ L_{G'} \downarrow & & \downarrow L_{H'} \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes.

Proof. It follows from Corollary 4.3 that there is a conjugacy $\phi'' : X_{G''} \rightarrow X_{H''}$ such that

$$\begin{array}{ccc} X_{G''} & \xrightarrow{\phi''} & X_{H''} \\ L_{G''} \downarrow & & \downarrow L_{H''} \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes. When we use the same symbol κ for the two conjugacies $\kappa : X_G^{L_G} \rightarrow X_{G''}$ and $\kappa : X_H^{L_H} \rightarrow X_{H''}$ from Lemma 4.2, we note that $\phi'' = \kappa \circ \phi^\# \circ \kappa^{-1}$. Since $\phi^\#(L_G^{-1}(y)) = L_H^{-1}(\psi(y))$ it follows that $\phi''(\beta_G(y)) = \beta_H(\psi(y))$ for all $y \in Y$. By Lemma 4.7 this implies that $\phi''(X_{G'}) = X_{H'}$, and we set $\phi' := \phi''|_{X_{G'}}$. This proves the existence part. To prove uniqueness, consider a conjugacy $\lambda : X_{G'} \rightarrow X_{H'}$ such that

$$\begin{array}{ccc} X_{G'} & \xrightarrow{\lambda} & X_{H'} \\ L_{G'} \downarrow & & \downarrow L_{H'} \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes. Set

$$\lambda' := \kappa^{-1} \circ \lambda \circ \kappa|_{\kappa^{-1}(X_{G'})}.$$

By Lemma 4.2 also

$$\begin{array}{ccc} X_G^{L_G} \cap \kappa^{-1}(X_{G'}) & \xrightarrow{\lambda'} & X_H^{L_H} \cap \kappa^{-1}(X_{H'}) \\ L_G^\# \downarrow & & \downarrow L_H^\# \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes. Note that $L_G^{-1}(y) \in X_G^{L_G} \cap \kappa^{-1}(X_{G'})$ by Lemma 4.7. It follows that $\lambda'(L_G^{-1}(y)) \subseteq L_H^{-1}(\psi(y))$ and $\lambda'^{-1}(L_H^{-1}(\psi(y))) \subseteq L_G^{-1}(y)$, implying that $\lambda'(L_G^{-1}(y)) = L_H^{-1}(\psi(y))$ for all $y \in Y$ and hence $\lambda(\beta_G(y)) = \beta_H(\psi(y)) = \phi'(\beta_G(y))$ for all $y \in Y$. By Lemma 4.7 this implies that $\lambda = \phi'$. □

We note that in analogy with (7) in Krieger's theorem, the diagram

$$(12) \quad \begin{array}{ccc} X_{G'} & \xrightarrow{\phi'} & X_{H'} \\ \beta_G \uparrow & & \uparrow \beta_H \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes.

One of the possible applications of Theorem 4.8 is to the minimal right-resolving presentations of two conjugate irreducible sofic shifts. The paper finishes with three examples where the canonical cover $(G', L_{G'})$ arising in this setup is exhibited and compared to the future cover. For this we need a general procedure to construct the future cover.

5. CONSTRUCTING THE FUTURE COVER

In this section we present a method to construct the future cover of a sofic shift from any given presentation of it.

5.1. On the merging of a labeled graph. Let (H, L_H) be a labeled graph. The *merged graph* $([H], L_{[H]})$ of (H, L_H) is then defined as follows, cf. page 78 of [LM]. The set of vertexes $V_{[H]}$ is the set

$$V_{[H]} := \{f_H(v) : v \in V_H\}$$

and for $a \in A$ there is a labeled arrow $f_H(v) \xrightarrow{a} f_H(w)$ when there are vertexes $v', w' \in V_H$ such that $f_H(v') = f_H(v)$, $f_H(w') = f_H(w)$ and there is a labeled arrow $v' \xrightarrow{a} w'$ in H . When (H, L_H) is right-resolving we can define a map $f_H : E_H \rightarrow E_{[H]}$ such that $f_H(e)$ is the arrow $f_H(e) \in E_{[H]}$ with the properties

- $s_{[H]}(f_H(e)) = f_H(s_H(e))$, and
- $L_{[H]}(e) = L_H(e)$.

Then $f_H : H \rightarrow [H]$ is a labeled-graphs homomorphism.

Lemma 5.1. *Assume that (H, L_H) is right-resolving. Let $v, w \in V_H$. There is a labeled arrow $f_H(v) \xrightarrow{a} f_H(w)$ in $[H]$ if and only if there is a labeled arrow $v \xrightarrow{a} u$ in H such that $f_H(u) = f_H(w)$.*

Proof. Assume that $f_H(v) \xrightarrow{a} f_H(w)$ in $[H]$ and let $v_1, w_1 \in V_H$ be such that $v_1 \xrightarrow{a} w_1$ in H , $f_H(v_1) = f_H(v)$ and $f_H(w_1) = f_H(w)$. Since $f_H(v) = f_H(v_1)$ there is an arrow e in E_H labeled a which leaves v . Set $u := t_H(e)$. By using that (H, L_H) is right-resolving we find then that

$$f_H(u) = \frac{f_H(v)}{a} = \frac{f_H(v_1)}{a} = f_H(w_1).$$

This establishes the 'only if' part. The 'if' part is trivial. \square

Lemma 5.2. *Assume that (H, L_H) is right-resolving. Then $([H], L_{[H]})$ is right-resolving and follower-separated, and there is a labeled-graph homomorphism $f_H : H \rightarrow [H]$ such that $f_H : X_H \rightarrow X_{[H]}$ is a factor map.*

Proof. $([H], L_{[H]})$ is right-resolving by Lemma 3.3.8 of [LM]. To see that $f_H : X_H \rightarrow X_{[H]}$ is surjective, consider a finite path

$$(13) \quad f_H(v_1) \xrightarrow{a_1} f_H(v_2) \xrightarrow{a_2} f_H(v_3) \xrightarrow{a_3} \dots \xrightarrow{a_{n-1}} f_H(v_n)$$

in $[H]$. Repeated applications of Lemma 5.1 give a path

$$v_1 \xrightarrow{a_1} u_2 \xrightarrow{a_2} u_3 \xrightarrow{a_3} \dots \xrightarrow{a_{n-1}} u_n$$

in H which is mapped to (13) under f_H . The surjectivity of f_H follows therefore from compactness of X_H . The argument shows also that the follower set of $f_H(v_1)$ in $([H], L_{[H]})$ is the same as the follower set of v_1 in (H, L_H) ; in symbols

$$f_{[H]}(f_H(v_1)) = f_H(v_1).$$

In particular, $([H], L_{[H]})$ is follower-separated \square

We note that the morphism $f_H : H \rightarrow [H]$ is injective on V_H if and only if it is an isomorphism of labeled graphs, and this happens if and only if (H, L_H) is follower-separated.

5.2. A cover of the future cover. Let (G, L_G) be a labeled graph and $Y = L_G(X_G)$ the sofic shift presented by (G, L_G) . When $D \subseteq V_G$ and $a \in A$, set

$$[D, a] := \{t_G(e) : e \in E_G, s_G(e) \in D, L_G(e) = a\};$$

the set of vertexes that can be reached from an element of D using an edge in G labeled by a . We consider the collection 2^{V_G} of non-empty subsets of V_G as the set of vertexes in a graph \overline{G} with edges

$$E_{\overline{G}} := \{(D, a) \in 2^{V_G} \times A : [D, a] \neq \emptyset\}.$$

The start vertex $s_{\overline{G}}(D, a)$ of $(D, a) \in E_{\overline{G}}$ is $D \in 2^{V_G}$ and the terminal vertex $t_{\overline{G}}(D, a)$ is $[D, a]$. With this definition, there is a graph morphism $G \rightarrow \overline{G}$ sending a vertex $v \in V_G$ to $\{v\} \in 2^{V_G}$ and $e \in E_G$ to $(\{s_G(e)\}, L_G(e)) \in E_{\overline{G}}$, and we extend L_G from E_G to a labeling $L_{\overline{G}}$ of $E_{\overline{G}}$ such that

$$L_{\overline{G}}(D, a) := a$$

when $(D, a) \in E_{\overline{G}}$. Then $(\overline{G}, L_{\overline{G}})$ is a right-resolving labeled graph and $L_{\overline{G}}(X_{\overline{G}}) = Y$.

For $y \in Y$, set

$$D^y := \{t_G(x) : x \in X_G(-\infty, -1], L_G(x) = y_{(-\infty, -1]}\}.$$

The set of vertexes $V_{\underline{G}} := \{D^y : y \in Y\}$ is a hereditary subset of vertexes in $V_{\overline{G}}$; that is, if $e \in E_{\overline{G}}$ and $s_{\overline{G}}(e) \in V_{\underline{G}}$, then $s_{\overline{G}}(e) \in V_{\underline{G}}$. Hence, when we set $E_{\underline{G}} := \{e \in E_{\overline{G}} : s_{\overline{G}}(e) \in V_{\underline{G}}\}$, we have defined a graph \underline{G} , and hence also a labeled graph $(\underline{G}, L_{\underline{G}})$, when $L_{\underline{G}} := L_{\overline{G}}|_{E_{\underline{G}}}$.

Note that we can define a shift-commuting map $\alpha_G : Y \rightarrow X_{\underline{G}}$ such that

- a) $s_{\underline{G}}(\alpha_G(y)_k) = D^{\sigma^k(y)}$ for all $k \in \mathbb{Z}$, and
- b) $L_{\underline{G}}(\alpha_G(y)) = y$.

In particular, $L_{\underline{G}}(X_{\underline{G}}) = Y$ and $(\underline{G}, L_{\underline{G}})$ is thus a presentation of Y . We show next that the future cover of Y is the merged graph of $(\underline{G}, L_{\underline{G}})$.

Let $([\underline{G}], L_{[\underline{G}]})$ be the merged graph of $(\underline{G}, L_{\underline{G}})$. Thus the set of vertexes $V_{[\underline{G}]}$ consists of the set

$$V_{[\underline{G}]} := \{f_{\underline{G}}(v) : v \in V_{\underline{G}}\}.$$

By Lemma 5.2 $([\underline{G}], L_{[\underline{G}]})$ is right-resolving and the morphism $f_{\underline{G}} : \underline{G} \rightarrow [\underline{G}]$ of labeled graphs induces a factor map $f_{\underline{G}} : X_{\underline{G}} \rightarrow X_{[\underline{G}]}$.

Lemma 5.3. $F(y) = f_{\underline{G}}(D^y)$ for all $y \in Y$.

Proof. It follows from the definitions that

$$\begin{aligned} f_{\underline{G}}(D^y) &= \{L_{\underline{G}}(x) : x \in X_{\underline{G}}[0, \infty), s_{\underline{G}}(x) = D^y\} \\ &= \{L_G(x') : x' \in X_G[0, \infty), s_G(x') \in D^y\} \\ &= \{z_{[0, \infty)} : z \in Y, z_j = y_j, j \leq -1\} = F(y). \end{aligned}$$

□

Lemma 5.4. *Let $y, z \in Y$, $a \in A$. There is a labeled arrow*

$$(14) \quad F(y) \xrightarrow{a} F(z)$$

in $\mathbb{K}(Y)$ if and only if there is a labeled arrow

$$(15) \quad f_{\underline{G}}(D^y) \xrightarrow{a} f_{\underline{G}}(D^z)$$

in $[\underline{G}]$.

Proof. By Lemma 5.3 an arrow as in (14) exists if and only if $f_{\underline{G}}(D^z) = \frac{f_{\underline{G}}(D^y)}{a}$, or if and only if

$$\{L_{\underline{G}}(x) : x \in X_{\underline{G}}[0, \infty), s_{\underline{G}}(x) = D^z\} = \frac{\{L_{\underline{G}}(x') : x' \in X_{\underline{G}}[0, \infty), s_{\underline{G}}(x') = D^y\}}{a}.$$

The latter equality holds exactly when there is a labeled arrow $D^y \xrightarrow{a} D$ in \underline{G} for some $D \in V_{\underline{G}}$ with $f_{\underline{G}}(D) = f_{\underline{G}}(D^z)$. By Lemma 5.1 this happens if and only if there is a labeled arrow as in (15). □

It follows from the last two lemmas that there is an isomorphism of labeled graphs $(\mathbb{K}(Y), L_{\mathbb{K}(Y)}) \simeq ([\underline{G}], L_{[\underline{G}]})$ defined such that


$$V_{\mathbb{K}(Y)} \ni F(y) \mapsto f_{\underline{G}}(D^y),$$

allowing us to make the following identification:


Corollary 5.5. $(\mathbb{K}(Y), L_{\mathbb{K}(Y)}) = ([\underline{G}], L_{[\underline{G}]})$ as labeled graphs.

6. FOUR EXAMPLES

It seems to be a well-established tradition that examples of sofic shifts start with the even shift, and we have no reason to deviate from this. The minimal right-resolving presentation of the even shift is shown in Figure 5 and the future cover in Figure 3. When (G, L_G) is the labeled graph from Figure 5, the corresponding labeled graph $(G', L_{G'})$ from Theorem 4.8 is




If instead (G, L_G) is the labeled graph from Figure 3, the labeled graph $(G', L_{G'})$ becomes




Thus we see, already from the even shift, that the minimal right-resolving cover, the future cover, the labeled graph $(G', L_{G'})$ from Theorem 4.8 applied first to the minimal right-resolving cover and then to the future cover, are all different covers.

We end the paper by giving two examples where the future cover of a mixing sofic shift is the same as the minimal right-resolving cover, while the canonical cover from Theorem 4.8 applied to the minimal right-resolving cover is different. The examples also show that in some cases the labeled graph $(\underline{G}, L_{\underline{G}})$ from Section 5.2 is the same as $(G', L_{G'})$ and in others it isn't.


Example 6.1. This following labeled graph (G, L_G) is irreducible, right-resolving and follower-separated and hence the minimal right resolving graph of the sofic shift Y it represents. Note that Y is mixing.



The graph $(\underline{G}, L_{\underline{G}})$ is the following.




Here the vertex $\{a, b\}$ has the same follower set as $\{a\}$ and the merged graph, which is the future cover $(\mathbb{K}(Y), L_{\mathbb{K}(Y)})$ by Corollary 5.5, is the graph (16) we started with. The future cover is therefore the same as the minimal right-resolving cover. In this case the labeled graph G' is




Thus $(G', L_{G'})$ is a proper subgraph of $(\underline{G}, L_{\underline{G}})$ in this case.

Example 6.2. The following labeled graph (G, L_G) is irreducible, right resolving and follower-separated and hence the minimal right resolving graph of the sofic shift Y it represents. Note that Y is mixing.



(17)

In this case the graphs $(\underline{G}, L_{\underline{G}})$ and $(G', L_{G'})$ are the same:



Since the vertex $\{a, b\}$ has the same follower set as $\{a\}$, the future cover is again the labeled graph (17) we started with, and hence equal to the minimal right-resolving cover, while the canonical cover $(G', L_{G'})$ is bigger.

REFERENCES

- [B] M. Barnsley, *Fractals everywhere*, Academic Press, 1988.
- [BKM] M. Boyle, B. Kitchens and B.H. Marcus, *A note on minimal covers for sofic systems*, Proc. Amer. Math. Soc. **95** (1985), 403-411.
- [F] R. Fischer, *Sofic systems and graphs*, Monatsh. Math. **80** (1975), 179-186.
- [J] R. Johansen, *On the structure of covers of sofic shifts*, Documenta Mathematica **16** (2011), 111-131.
- [Jo] N. Jonoska, *Sofic shifts with synchronizing presentations*, Theoret. Comput. Sci. **158** (1996), 81-115.
- [Kr1] W. Krieger, *On sofic systems I*, Israel J. Math. **48** (1984), 305-330.
- [Kr2] W. Krieger, *On sofic systems II*, Israel J. Math. **60** (1987), 167-176.
- [LM] D. Lind and B. Marcus, *An introduction to symbolic dynamics and coding*, Cambridge University Press, 1995.
- [MMTW] B. Marcus, T. Meyerovitch, K. Thomsen and C. Wu, *Factorizable embeddings and the period of an irreducible sofic shift*, arXiv:2508.02554
- [N] M. Nasu, *Topological conjugacy for sofic systems*, Ergod. Th. & Dynam. Sys. **6** (1986), 265-280.