

# Digamma-Function Representation of the Ground-State Energy in Antiferromagnetic Heisenberg $XXX_s$ Spin Chains

Reaz Shafqat

Department of Mathematics and Natural Sciences, BRAC University,  
Dhaka, Bangladesh

December 2025

## Abstract

The antiferromagnetic Heisenberg spin chain remains a central framework for exploring exactly solvable models within quantum integrable systems. For the isotropic XXX chain, the ground-state energy per site of the spin- $\frac{1}{2}$  system is famously given by  $\ln 2$ . Extending the classic formulations to arbitrary spin- $s$ , Takhtajan and Babujian derived two separate finite-series expressions for integer and half-integer spin representations. Current work introduces a unified analytical expression for the ground-state energy density in terms of the digamma function. This compact formulation reproduces both the Takhtajan Babujian series.

## Introduction<sup>1</sup>

The study of quantum spin chains occupies a central role in condensed-matter physics, integrable models, and mathematical physics. Among these systems, the antiferromagnetic Heisenberg model stands out as a paradigmatic exactly solvable many-body model, capturing key features of strong correlations, quantum criticality, and universal behaviour in one dimension. Since Bethe's original solution of the spin- $\frac{1}{2}$  XXX chain [1], the model has remained a cornerstone in the development of integrable systems, inspiring powerful frameworks such as the coordinate and Algebraic Bethe Ansatz [3, 13, 14], the quantum inverse scattering method [12], and deep connections to quantum groups and conformal field theory (CFT) [9, 10, 24, 23].

For higher-spin generalisations of the isotropic XXX chain, the exact analytic structure becomes richer. Independently, Takhtajan [6] and Babujian [8] constructed closed-form thermodynamic Bethe–Ansatz (TBA) expressions for the ground-state energy density for

---

<sup>1</sup>This work is based on the master's thesis of Reaz Shafqat submitted to the Department of Theoretical Physics of the University of Dhaka, Session: 21-22.

arbitrary spin  $s$ . Their analyses established the now-famous distinction between integer and half-integer spin chains, later connected to the emergence of topological  $\theta$ -terms in the low-energy nonlinear sigma model (NLSM) description [9, 10, 11]. These results revealed that the ground-state energy admits two finite-series expansions that differ structurally for integer and half-integer  $s$ , a reflection of the fundamental dichotomy between gapped (integer) and critical (half-integer) chains predicted by the Haldane conjecture[9, 10].

Despite the elegance and completeness of the Takhtajan–Babujian formalism, the resulting expressions do not provide a single, analytically continuous formula valid for all real  $s$ . Moreover, modern developments—such as anyonic spin chains, fractionalised excitations, and extensions via quantum-group symmetry [21, 22, 18, 19]—motivate a unified perspective on the ground-state energy that treats  $s$  as a continuous variable. This work proposes a compact analytic representation of the antiferromagnetic vacuum energy density in terms of the digamma functions. This formulation reproduces the Takhtajan–Babujian results exactly. The proposed expression offers an analytically tractable, physically transparent, and computationally convenient alternative to traditional finite-series expansions, thereby simplifying both theoretical analysis and numerical evaluation.

The remainder of this paper is organised as follows. The following section reviews the essential theoretical foundations of the Heisenberg XXX chain: the Takhtajan–Babujian generalisation to arbitrary spin- $s$ , and the resulting integer/half-integer series structure for the ground-state energy density. Afterwards, the digamma-function representation of the ground state energy density with a detailed derivation is outlined.

## Babujian’s thermodynamic Bethe ansatz for XXX <sub>$s$</sub> chain

A brief overview of Babujian’s thermodynamic Bethe ansatz (TBA) for the antiferromagnetic spin- $s$  Heisenberg XXX chain is presented, where  $s$  may take either integer or half-integer values [7, 8]. This formulation generalizes the earlier spin- $\frac{1}{2}$  treatment developed by Takahashi and Yang–Yang [5, 3]. The analysis starts from the Bethe equations for  $N$  sites,  $M$  down spins (magnons) with periodic boundary conditions. The rapidities shift as  $\lambda_j \rightarrow i\lambda_j + \frac{1}{2}$ , one obtains

$$\left(\frac{\lambda_j - is}{\lambda_j + is}\right)^N = \prod_{\substack{\ell=1 \\ \ell \neq j}}^M \frac{\lambda_j - \lambda_\ell - i}{\lambda_j - \lambda_\ell + i}. \quad (1)$$

In the thermodynamic limit, solutions are assumed to form “strings” of length  $n$  in the complex plane,

$$\lambda_j^{n,\alpha} = \lambda_j^n + i(n+1-2\alpha), \quad \alpha = 1, \dots, n, \quad (2)$$

with real string centres  $\lambda_j^n$ . The corresponding Bethe ansatz equation will take on the following form:

$$\left(\frac{\lambda_j + is}{\lambda_j - is}\right)^N = \prod_{\substack{k=1 \\ k \neq j}}^{\xi_1} \frac{\lambda_j - \lambda_k + i}{\lambda_j + \lambda_k - i} \prod_{\substack{k=1 \\ k \neq j}}^{\xi_2} \frac{\lambda_j - \lambda_k + 2i}{\lambda_j - \lambda_k - 2i} \dots \prod_{\substack{k=1 \\ k \neq j}}^{\xi_{2s}} \frac{\lambda_j - \lambda_k + 2si}{\lambda_j - \lambda_k - 2si}. \quad (3)$$

If  $\xi_n$  denotes the number of  $n$ -strings, then

$$M = \sum_{n=1}^{\infty} n \xi_n. \quad (4)$$

Multiplying (3) over the members of a given string and taking the logarithm yields Bethe equations for the string centres,

$$N \theta_{n,2s}(\lambda_j^n) = 2\pi J_j^n + \sum_{k=1}^{\infty} \sum_{\ell=1}^{\xi_k} \Xi_{n,k}(\lambda_j^n - \lambda_\ell^k), \quad (5)$$

where  $J_j^n$  are integer or half-integer quantum numbers and

$$\theta_n(\lambda) = 2 \arctan \frac{\lambda}{n}, \quad \theta_{m,p}(\lambda) = \sum_{\ell=1}^{\min(m,p)} \theta_{m+p+1-2\ell}(\lambda) \quad (6)$$

are basic scattering phases. The  $\Xi_{m,k}$  are fixed linear combinations of the  $\theta_\ell$  that encode scattering between  $m$ - and  $k$ -strings. The multi-string scattering phase can be expressed through the kernels  $\Xi_{m,p}(\lambda)$  using the convolution identity

$$\theta_{m,p}(\lambda) = \hat{p}(\Xi_{m,p}(\lambda) + \pi \operatorname{sgn} \lambda \delta_{m,p}), \quad (7)$$

where

$$\hat{p}g(\lambda) = \int_{-\infty}^{\infty} p(\lambda - \lambda') g(\lambda') d\lambda', \quad p(\lambda) = \left(4 \cosh \frac{\pi \lambda}{2}\right)^{-1}. \quad (8)$$

In the thermodynamic limit, the discrete roots are described by continuous densities of occupied and unoccupied quantum numbers,  $\rho_n(\lambda)$  and  $\tilde{\rho}_n(\lambda)$ , defined such that  $\rho_n(\lambda) d\lambda$  gives the number of  $n$ -strings with centres in  $[\lambda, \lambda + d\lambda]$  (and similarly for  $\tilde{\rho}_n$ ). Differentiating (5) and converting sums into integrals, one obtains the basic linear relation

$$2\pi [\rho_n(\lambda) + \tilde{\rho}_n(\lambda)] = \theta'_{n,2s}(\lambda) - \sum_{k=1}^{\infty} \int_{-\infty}^{\infty} K_{n,k}(\lambda - \lambda') \rho_k(\lambda') d\lambda', \quad (9)$$

where  $K_{n,k}(\lambda) = \frac{d}{d\lambda} \Xi_{n,k}(\lambda)/2\pi$  is the TBA kernel. In the thermodynamic limit, the relation between particle and hole densities takes the form

$$\tilde{\rho}_m(\lambda) = - \sum_{n=1}^{\infty} \hat{A}_{m,n}[\rho_n(\lambda) - p(\lambda) \delta_{n,2s}], \quad (10)$$

where  $\hat{A}_{m,n}$  denotes the integral operator built from the kernel  $A_{m,n}(\lambda)$ . The energy per site can be expressed directly in terms of the  $\rho_n$ ,

$$\frac{E_s}{N} = -\frac{1}{2} \sum_{n=1}^{\infty} \int \theta'_{n,2s}(\lambda) \rho_n(\lambda) d\lambda \quad (11)$$

The equilibrium distributions at temperature  $T$  and magnetic field  $H$  follow from minimizing the free energy  $F = E_s - TS - HS^z$  at  $T = 0$  and  $H = 0$  with respect to  $\rho_n$  and  $\tilde{\rho}_n$ , where the Entropy has the standard TBA form [3, 16]. Introducing the pseudoenergies

$$\varepsilon_n(\lambda) = T \ln \frac{\tilde{\rho}_n(\lambda)}{\rho_n(\lambda)}, \quad (12)$$

The stationarity condition yields coupled non-linear integral equations of the form

$$\varepsilon_n(\lambda) = Hn - \frac{1}{2}\theta'_{n,2s}(\lambda) - T \sum_{k=1}^{\infty} (K_{n,k} * \ln[1 + e^{-\varepsilon_k/T}])(\lambda), \quad (13)$$

with  $*$  denoting convolution. These are Babujian's TBA equations for the spin- $s$  chain. Finally, the zero-temperature limit  $T \rightarrow 0$  of (13) determines the ground state. One finds that only strings of length  $2s$  are occupied in the antiferromagnetic vacuum, with pseudoenergy

$$\varepsilon_{2s}(\lambda) = -\frac{\pi}{4 \cosh(\frac{\pi}{2}\lambda)}, \quad \varepsilon_{n \neq 2s}(\lambda) \geq 0. \quad (14)$$

Substituting this into the expression for the free energy gives the ground-state energy

$$\frac{E_s^0}{N} = -\sum_{k=1}^{2s} \beta(k), \quad \beta(k) = \int_0^1 \frac{x^{k-1}}{1+x} dx. \quad (15)$$

The integral in (15) is evaluated by using the recursive reduction of power in the numerator. So for integer and half-integer, one gets two separate reduction formulas. Thus, evaluating the sum, Babujian obtains the compact formulas

$$\text{integer } s : \quad \frac{E_s^0}{N} = -\sum_{k=1}^s \frac{1}{2k-1}, \quad (16)$$

$$\text{half-integer } s : \quad \frac{E_s^0}{N} = -\left( \ln 2 + \sum_{k=1}^{s-\frac{1}{2}} \frac{1}{2k} \right), \quad (17)$$

which display the characteristic difference between integer and half-integer spins, in line with Haldane's conjecture on the low-energy physics of Heisenberg chains [9, 10].

## Proposed Expression

Following [25, 5], it can simply be stated that the ground state of the antiferromagnetic Heisenberg spin chain contains no holes. Hence in (10)  $\tilde{\rho}_m(\lambda) = 0$ . So,

$$\rho(\lambda) = p(\lambda)\delta_{n,2s} = \frac{-\delta_{n,2s}}{4 \cosh(\pi\lambda/2)}$$

The above equation suggests that the ground state only contains strings of length:  $2s$ . Now, in order to determine the ground state energy (11) is considered. Since the ground

state contains only strings of length  $2s$ , the choice  $n = 2s$  follows directly. Hence,

$$\theta_{2s,2s}(\lambda) = \sum_{\ell=1}^{2s} \theta_{4s+1-2\ell}(\lambda) = 2 \sum_{\ell=1}^{2s} \arctan \left( \frac{\lambda}{4s+1-2\ell} \right)$$

Now differentiating w.r.t  $\lambda$ :

$$\theta'_{2s,2s}(\lambda) \sum_{\ell=1}^{2s} \frac{4s+1-2\ell}{\lambda^2 + (4s+1-2\ell)^2}$$

Plugging the expression of  $\rho(\lambda)$  and  $\theta'_{2s,2s}(\lambda)$  in (11), giving:

$$\frac{E(s)}{N} = \sum_{l=1}^{2s} \frac{-1}{4} \int_{-\infty}^{\infty} \frac{4s+1-2l}{\lambda^2 + (4s+1-2l)^2} \frac{d\lambda}{\cosh(\pi\lambda/2)}, \quad (18)$$

Integrals involving the hyperbolic secant kernel arise naturally in integrable spin chains and thermodynamic Bethe ansatz calculations. For  $4s+1-2l > 0$  with  $0 \leq l \leq 2s$  the standard Laplace representation is employed:

$$\frac{4s+1-2l}{\lambda^2 + (4s+1-2l)^2} = \int_0^{\infty} e^{-(4s+1-2l)t} \cos(\lambda t) dt. \quad (19)$$

Substituting (19) into (18) and interchanging the order of integration gives

$$\begin{aligned} \frac{E(s)}{N} &= \frac{-1}{4} \int_0^{\infty} e^{-(4s+1-2l)t} \left( \int_{-\infty}^{\infty} \frac{\cos(\lambda t)}{\cosh(\pi\lambda/2)} d\lambda \right) dt \\ &= \frac{-1}{4} \int_0^{\infty} e^{-(4s+1-2l)t} J(t) dt, \end{aligned} \quad (20)$$

where

$$J(t) = \int_{-\infty}^{\infty} \frac{\cos(\lambda t)}{\cosh(\pi\lambda/2)} d\lambda. \quad (21)$$

To compute  $J(t) = \int_{-\infty}^{\infty} \frac{\cos(\lambda t)}{\cosh(\pi\lambda/2)} d\lambda$ , first this Fourier cosine transform is converted to the Fourier transform:

$$\int_{-\infty}^{\infty} \frac{\cos(\lambda t)}{\cosh(\pi\lambda/2)} d\lambda = \Re \left( \int_{-\infty}^{\infty} \frac{e^{-i\lambda t}}{(\cosh \pi\lambda/2)} d\lambda \right) \quad (22)$$

First, the following substitution has been made:

$$\mu = \frac{\pi\lambda}{2}, \quad \lambda = \frac{2\mu}{\pi}, \quad d\lambda = \frac{2}{\pi} d\mu,$$

to rewrite (21) as

$$J(t) = \frac{2}{\pi} \int_{-\infty}^{\infty} \frac{\exp(i\frac{2t}{\pi}\mu)}{\cosh \mu} d\mu. \quad (23)$$

Expressing  $\cosh(\mu) = \frac{2}{e^\mu + e^{-\mu}}$ :

$$J(t) = \frac{2}{\pi} \int_{-\infty}^{\infty} \frac{\exp(i\frac{2t}{\pi}\mu)}{e^\mu + e^{-\mu}} d\mu$$

Introducing the substitution  $x = e^\mu$ , which implies  $\mu = \ln x$ , the integration limits are transformed accordingly:

$$J(t) = \frac{4}{\pi} \int_{-\infty}^{\infty} \frac{\exp\left(i\frac{2t}{\pi}\mu\right)}{e^\mu + e^{-\mu}} d\mu = \frac{4}{\pi} \int_0^{\infty} \frac{x^{(i\frac{2t}{\pi})}}{1+x^2} dx = \frac{4}{\pi} \int_0^{\infty} \frac{x^{(i\frac{2t}{\pi}+1)-1}}{1+x^2} dx$$

Setting  $\alpha = i\frac{2t}{\pi} + 1$  and  $\beta = 2$  gives:

$$J(t) = \frac{4}{\pi} \int_0^{\infty} \frac{x^{\alpha-1}}{1+x^\beta} dx$$

Now, letting  $y = (1+x^\beta)^{-1}$ , since  $0 < \Re(\alpha) < \beta$ , the integral  $J(t)$  becomes:

$$\frac{4}{\pi} \int_0^{\infty} \frac{x^{\alpha-1}}{1+x^\beta} dx = \frac{4}{\pi\beta} \int_0^1 y^{-\alpha/\beta} (1-y)^{\alpha/\beta-1} dy = \frac{4}{\pi\beta} B\left(1 - \frac{\alpha}{\beta}, \frac{\alpha}{\beta}\right)$$

The expression reduces to the Beta function. Using its representation in terms of Gamma functions, one obtains:

$$= \frac{4}{\pi\beta} \Gamma\left(1 - \frac{\alpha}{\beta}\right) \Gamma\left(\frac{\alpha}{\beta}\right) = \frac{4\pi}{\pi\beta} \csc\left(\frac{\alpha\pi}{\beta}\right).$$

Applying values of  $\alpha$  &  $\beta$ :

$$J(t) = 2 \csc\left\{\left(i\frac{2t}{\pi} + 1\right) \frac{\pi}{2}\right\} = \sec(it) = 2\operatorname{sech}(t) \quad (24)$$

Substituting (24) into (20) gives the simpler representation

$$\frac{E(s)}{N} = \frac{-2}{4} \int_0^{\infty} e^{-(4s+1-2l)t} \operatorname{sech}(t) dt. \quad (25)$$

Use the standard expansion

$$\operatorname{sech}(t) = \frac{2}{e^t + e^{-t}} = \frac{2e^{-t}}{1 + e^{-2t}}, \quad (26)$$

Now the integral interval in (25) is 0 to  $\infty$ . On this interval  $0 < e^{-2t} \leq 1$ . So, the binomial expansion is taken into account (26).

$$\operatorname{sech}(t) = \frac{2e^{-t}}{1 + e^{-2t}} = 2 \sum_{n=1}^{\infty} (-1)^n e^{-(2n+1)t} \quad (27)$$

which, when inserted into (25), yields

$$\begin{aligned} \frac{E(s)}{N} &= \frac{-4}{4} \sum_{n=0}^{\infty} (-1)^n \int_0^{\infty} e^{-(4s+1-2l+2n+1)t} dt \\ &= - \sum_{n=0}^{\infty} \frac{(-1)^n}{4s+1-2l+2n+1}. \end{aligned} \quad (28)$$

One defines:

$$A = \frac{4s+1-2l+1}{2}, \quad 4s+1-2l+2n+1 = 2(n+A),$$

so that (28) becomes

$$\frac{E(s)}{N} = - \sum_{n=0}^{\infty} \frac{(-1)^n}{n+A}. \quad (29)$$

(29) can be represented by Lerch transcendence,  $\Phi(z, s, a) = \sum_{k=0}^{\infty} \frac{z^k}{(a+k)^s}$  by setting,  $z = -1, s = 1, k = n$  and  $a = A$ . Therefore, the following equation is obtained,

$$\Phi(-1, 1, A) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(A+n)} \quad (30)$$

It is known that the Lerch transcendent[20] has a digamma function representation expressed as:

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{n+z} = \frac{1}{2} \left[ \psi\left(\frac{z+1}{2}\right) - \psi\left(\frac{z}{2}\right) \right], \quad (31)$$

where  $\psi$  denotes the digamma function. From (29) and (31),

$$\frac{E(s)}{N} = \frac{-1}{2} \left( \psi\left(\frac{A+1}{2}\right) - \psi\left(\frac{A}{2}\right) \right) \quad (32)$$

Putting the value of  $A$ :

$$\begin{aligned} \frac{E(s)}{N} &= \frac{-1}{2} \left( \psi\left(\frac{4s+1-2l+1+2}{4}\right) - \psi\left(\frac{4s+1-2l+1}{4}\right) \right) \\ \Rightarrow \frac{E(s)}{N} &= \frac{-1}{2} \left( \psi\left(\frac{4s+4-2l}{4}\right) - \psi\left(\frac{4s+2-2l}{4}\right) \right) \end{aligned} \quad (33)$$

Reintroducing the summation in the integral (18) admits the closed form

$$\frac{E(s)}{N} = \frac{-1}{2} \sum_{l=1}^{2s} \left( \psi\left(\frac{4s+4-2l}{4}\right) - \psi\left(\frac{4s+2-2l}{4}\right) \right) \quad (34)$$

Now evaluating different terms of the summation in (34). Hence, for  $l = 1$ :

$$\frac{-1}{2} \left( \psi\left(s + \frac{1}{2}\right) - \psi\left(\frac{s}{2}\right) \right)$$

For,  $l = 2$ :

$$\frac{-1}{2} \left( \psi(s) - \psi\left(s - \frac{1}{2}\right) \right)$$

So the summation in (34) is a telescopic sum, hence,

$$\begin{aligned} \frac{E(s)}{N} &= \frac{-1}{2} \left( \psi\left(s + \frac{1}{2}\right) - \psi\left(\frac{s}{2}\right) \cdots - \psi(1) + \psi(1) - \psi\left(\frac{1}{2}\right) \right) \\ \frac{E(s)}{N} &= \frac{-1}{2} \left( \psi\left(s + \frac{1}{2}\right) - \psi\left(\frac{1}{2}\right) \right) \end{aligned} \quad (35)$$

However, no step in the derivation of (35) relies on whether  $s$  is an integer or half-integer. Therefore, for integer  $s$ ,

$$\psi\left(s + \frac{1}{2}\right) = -\gamma - 2\ln(2) + \sum_{k=1}^s \frac{2}{2k-1}$$

and

$$\psi\left(\frac{1}{2}\right) = -\gamma - 2\ln(2)$$

Where  $\gamma$  = Euler-Mascheroni Constant. Now, from (35) one arrives at,

$$\frac{E(s)}{N} = -\sum_{k=1}^s \frac{1}{2k-1} \quad (36)$$

Now for  $s$ : half integers,

$$\psi\left(s + \frac{1}{2}\right) = -\gamma + \sum_{k=1}^{s-\frac{1}{2}} \frac{1}{k}$$

Furthermore, upon inserting the known value of  $\psi(\frac{1}{2})$ , the expression becomes:

$$\frac{E(s)}{N} = -\left(\ln 2 + \sum_{k=1}^{s-\frac{1}{2}} \frac{1}{2k}\right) \quad (37)$$

So, (36) and (37) exactly match the proposed expression of Babujian (17).

## Conclusion

In this work, a compact analytical formula (35) has been proposed for the antiferromagnetic ground-state energy density of the isotropic Heisenberg spin- $s$  chain using the digamma function. This representation unifies the treatment of integer and half-integer spin, reproduces known results from the Takhtajan–Babujian solution, and bypasses the need for separate finite-series expansions. A promising direction for future work is to explore whether this digamma-based expression can be extended to anyonic spin chains governed by  $SU(2)_k$  fusion rules [21, 26]. Such systems admit fractionalized spin degrees of freedom and exhibit rich topological and conformal structure. Developing an analogous exact formula in this setting could significantly advance the understanding of energy scaling, quantum group symmetries, and topological sectors in one-dimensional quantum systems.

## Acknowledgements

The author would like to thank Dr. Tibra Ali for his supervision and constant support throughout the work.



## References

- [1] H. Bethe, “Zur Theorie der Metalle. I. Eigenwerte und Eigenfunktionen der linearen Atomkette,” *Zeitschrift für Physik* **71** (1931) 205–226.
- [2] L. Hulthén, “Über das Austauschproblem eines Kristalles,” *Arkiv Mat. Astron. Fysik* **26A** (1938) 1.
- [3] C. N. Yang and C. P. Yang, “Thermodynamics of a One-Dimensional System of Bosons with Repulsive Delta-Function Interaction,” *J. Math. Phys.* **10** (1969) 1115–1122.
- [4] C. N. Yang and C. P. Yang, “One-Dimensional Chain of Anisotropic Spin-Spin Interactions. I. Proof of Bethe’s Hypothesis for Ground State in a Finite System,” *Phys. Rev.* **150** (1966) 321–327.
- [5] M. Takahashi, “One-Dimensional Heisenberg Model at Finite Temperature,” *Prog. Theor. Phys.* **46** (1971) 401–415.
- [6] L. A. Takhtajan, “The picture of low-lying excitations in the isotropic Heisenberg chain of arbitrary spin  $S$ ,” *Phys. Lett. A* **87** (1982) 479–482.
- [7] H. M. Babujian, “Exact solution of the one-dimensional isotropic Heisenberg chain with arbitrary spins  $S$ ,” *Phys. Lett. A* **90** (1982) 479–482.
- [8] H. M. Babujian, “Exact solution of the one-dimensional isotropic Heisenberg chain with arbitrary spins  $S$ ,” *Nucl. Phys. B* **215** (1983) 317–336.
- [9] F. D. M. Haldane, “Nonlinear field theory of large-spin Heisenberg antiferromagnets: Semiclassically quantized solitons of the one-dimensional easy-axis Néel state,” *Phys. Rev. Lett.* **50** (1983) 1153–1156.
- [10] F. D. M. Haldane, “Continuum dynamics of the 1-D Heisenberg antiferromagnet: Identification with the  $O(3)$  nonlinear sigma model,” *Phys. Lett. A* **93** (1983) 464–468.
- [11] S. C. Zhang, H. J. Schulz and T. Ziman, “Ground-state energies of the nonlinear sigma model and the Heisenberg spin chains,” *Phys. Rev. Lett.* **63** (1989) 1110–1113.
- [12] N. Yu. Reshetikhin, “A method of functional equations in the theory of exactly solvable quantum systems,” *Lett. Math. Phys.* **7** (1983) 205–213.
- [13] V. E. Korepin, N. M. Bogoliubov and A. G. Izergin, *Quantum Inverse Scattering Method and Correlation Functions*, Cambridge University Press (1993).
- [14] A. Klümper, “Integrability of quantum chains: Theory and applications to the spin-1/2 XXZ chain,” in *Quantum Magnetism*, Lecture Notes in Physics **645**, Springer (2004) 349–379.
- [15] H.-J. Mikeska and A. K. Kolezhuk, “One-Dimensional Magnetism,” in *Quantum Magnetism*, Lecture Notes in Physics **645**, Springer (2004).

- [16] M. Takahashi, *Thermodynamics of One-Dimensional Solvable Models*, Cambridge University Press, Cambridge (1999).
- [17] C. J. S. Langer (ed.), “Spin Chains and Exactly Solvable Models in Low-Dimensional Quantum Magnetism,” *J. Phys. A: Math. Gen.* **38** (2005) R289–R332.
- [18] M. Mourigal *et al.*, “Fractional spinon excitations in the quantum Heisenberg anti-ferromagnetic chain,” *arXiv:1306.4678*.
- [19] K.-T. Chen, “Duality at lattice scale around the Takhtajan–Babujian point in the bilinear–biquadratic spin-1 chain,” *arXiv:1512.00967*.
- [20] I. S. Gradshteyn and I. M. Ryzhik, *Tables of Integrals, Series, and Products*, 6th ed., Academic Press, San Diego (2000), see §9.55 “The Lerch Function  $\Phi(z, s, v)$ ”, p. 1029.
- [21] A. E. Feiguin, S. Trebst, A. W. W. Ludwig and M. Troyer, “Interacting Anyons in Topological Quantum Liquids: The Golden Chain,” *arXiv:cond-mat/0612341* (2007).
- [22] P. Lecheminant, “Topological phase transitions in  $SU(N)$  spin chains,” *Nucl. Phys. B* **901**, 510–535 (2015).
- [23] I. Affleck, “Universal term in the free energy at a critical point and the conformal anomaly,” *Phys. Rev. Lett.* **56**, 746–748 (1986).
- [24] A. B. Zamolodchikov and A. B. Zamolodchikov, “Factorized S-matrices and integrable field theories,” *Ann. Phys.* **120**, 253–291 (1979).
- [25] A. Doikou, “Simplified Calculation of Boundary S Matrices,” *hep-th/9709069* (1997).
- [26] P. Fendley, F. Lesage, and H. Saleur, “Exact results for integrable  $SU(2)_k$  quantum chains,” *J. Stat. Phys.* **79**, 799–827 (1995).