

Parity results for the reciprocals of false theta functions

Jing Jin¹, Huan Xu² and Olivia X.M. Yao³

¹College of Agricultural Information,
Jiangsu Agri-animal Husbandry Vocational College,
Taizhou, 225300, Jiangsu, P. R. China

^{2,3}School of Mathematical Sciences,
Suzhou University of Science and Technology,
Suzhou, 215009, Jiangsu, P. R. China

Email: ¹jinjing19841@126.com, ²xu_huanz@163.com, ³yaoxiangmei@163.com

Abstract. Recently, Keith investigated arithmetic properties for the reciprocals of some false theta functions and posed several conjectures. In this paper, we prove some parity results for the reciprocals of some false theta functions by using some identities on Ramanujan's general theta function. In particular, our results imply some conjectures of Keith.

Keywords: congruences, false theta functions, parity, Ramanujan's general theta functions.

AMS Subject Classification: 11P83, 05A17

1 Introduction

The aim of this paper is to investigate parity results for the reciprocals of some false theta functions and confirm some conjectures given by Keith [11] by utilizing several identities on Ramanujan's general theta function. Let us begin with some definitions.

Ramanujan's general theta function is defined by

$$f(a, b) = \sum_{n=-\infty}^{\infty} a^{n(n+1)/2} b^{n(n-1)/2}, \quad |ab| < 1. \quad (1.1)$$

In Ramanujan's notation, the Jacobi triple product identity can be rewritten as

$$f(a, b) = f(b, a) = (-a, -b, ab; ab)_{\infty}. \quad (1.2)$$

Here and throughout this paper, we use the following notation:

$$(a; q)_{\infty} := \prod_{k=0}^{\infty} (1 - aq^k), \quad (a; q)_n := \frac{(a; q)_{\infty}}{(aq^n; q)_{\infty}},$$

$$(a_1, a_2, \dots, a_k; q)_{\infty} := (a_1; q)_{\infty} (a_2; q)_{\infty} \cdots (a_k; q)_{\infty},$$

where q is a complex number with $|q| < 1$. In addition, for all positive integers m , define

$$f_m := (q^m; q^m)_\infty.$$

Three important special cases of (1.1) are defined by,

$$f(-q, -q^2) = \sum_{n=-\infty}^{\infty} (-1)^n q^{n(3n-1)/2} = f_1, \quad (1.3)$$

$$f(-q, -q) = \sum_{n=-\infty}^{\infty} (-1)^n q^{n^2} = \frac{f_1^2}{f_2}, \quad (1.4)$$

$$f(-q, -q^3) = \sum_{n=0}^{\infty} (-q)^{n(n+1)/2} = \frac{f_1 f_4}{f_2}. \quad (1.5)$$

Note that the reciprocals of the right sides of (1.3)–(1.5) are the generating functions of the ordinary partition function $p(n)$ [1], the overpartition function $\bar{p}(n)$ [7], and the partition with odd parts distinct function [9], respectively, which are three of the most important types of partition functions. A number of congruences for the three partition functions have been proved; see for example [8, 9, 12].

In 1917, Rogers [15] introduced false theta functions, which are series that would be classical theta functions except for changes in signs of an infinite number of terms. False theta functions closely resemble ordinary theta functions; however, they do not have the modular transformation properties that theta functions have. In his notebooks [13] and lost notebook [14], Ramanujan recorded many identities involving false theta functions $\Psi(a, b)$ which is defined by

$$\Psi(a, b) := \sum_{n=0}^{\infty} a^{n(n+1)/2} b^{n(n-1)/2} - \sum_{n=-\infty}^{-1} a^{n(n+1)/2} b^{n(n-1)/2}. \quad (1.6)$$

In recent years, there has been an increase in interest in false theta functions. They appear in a variety of contexts including modular and quantum modular forms, meromorphic Jacobi forms, quantum knot invariants, and many more; see for example [2, 3, 5, 6, 10, 16].

Very recently, Keith [11] studies reciprocals of false theta functions and proved some interesting results on congruences, asymptotic bounds, and combinatorial identities. In particular, he showed that for $n \geq 0$,

$$\begin{aligned} c_{5,1}(8n+5) &\equiv 0 \pmod{2}, \\ c_{5,1}(32n+31) &\equiv 0 \pmod{4}, \end{aligned}$$

where $c_{r,s}(n)$ is defined by

$$\sum_{n=0}^{\infty} c_{r,s}(n) q^n := \frac{1}{\Psi(-q^r, q^s)}. \quad (1.7)$$

In addition, he proved that $c_{9,1}(8n+4) \equiv 0 \pmod{2}$ if n cannot be represented in the form $n = 10k^2 - 4k$ for integer k .

At the end of his paper, Keith [11] posed the following conjecture on congruences modulo 2 for $c_{13,1}(n)$:

Conjecture 1.1 [11] For $n \geq 0$,

$$c_{13,1}(32n + 23) \equiv 0 \pmod{2}, \quad (1.8)$$

$$c_{13,1}(64n + 63) \equiv 0 \pmod{2}, \quad (1.9)$$

$$c_{13,1}(72n + j) \equiv 0 \pmod{2}, \quad (1.10)$$

where $j \in \{15, 21, 39, 69\}$.

In this paper, using some identities on Rmanujan's general theta function $f(a, b)$, we prove some parity results for $c_{r,s}(n)$ where $(r, s) \in \{(9, 1), (7, 3), (13, 1), (11, 3), (9, 5)\}$. In particular, we establish several infinite families of congruences modulo 2 which imply the congruences given in Conjecture 1.1.

2 Parity results for $c_{9,1}(n)$

Theorem 2.1 Let p be a prime with $p \equiv 11, 19 \pmod{20}$. For $n, k \geq 0$ with $p \nmid n$,

$$c_{9,1}\left(8p^{2k+1}n + \frac{6p^{2k+2} + 4}{5}\right) \equiv 0 \pmod{2} \quad (2.1)$$

and

$$c_{9,1}\left(8p^{2k+1}n + \frac{26p^{2k+2} + 4}{5}\right) \equiv 0 \pmod{2}. \quad (2.2)$$

Proof. Using (1.2) and the fact that

$$(1 - q)^2 \equiv (1 - q^2) \pmod{2},$$

one can prove that for all positive integers r and s ,

$$f(q^r, q^s)^2 \equiv f(q^{2r}, q^{2s}) \pmod{2}. \quad (2.3)$$

In view of (1.6) and (1.7), we see that for all positive integers r and s ,

$$\begin{aligned} \sum_{n=0}^{\infty} c_{r,s}(n)q^n &= \frac{1}{\sum_{n=0}^{\infty} (-1)^{n(n+1)/2} q^{(r+s)n^2/2 + (r-s)n/2} - \sum_{n=-\infty}^{-1} (-1)^{n(n+1)/2} q^{(r+s)n^2/2 + (r-s)n/2}} \\ &\equiv \frac{1}{\sum_{n=-\infty}^{\infty} (-1)^{n(n+1)/2} q^{(r+s)n^2/2 + (r-s)n/2}} \\ &\equiv \frac{1}{f(q^r, q^s)} \\ &\equiv \frac{f(q^s, q^r)}{f(q^{2s}, q^{2r})} \pmod{2}. \quad (\text{by (2.3)}) \end{aligned} \quad (2.4)$$

It follows from [4, Entry 30 (ii) and (iii)] that

$$f(a, b) = f(a^3b, ab^3) + af(b/a, a^5b^3). \quad (2.5)$$

Taking $a = q$ and $b = q^9$ in (2.5) yields

$$f(q, q^9) = f(q^{12}, q^{28}) + qf(q^8, q^{32}). \quad (2.6)$$

Setting $(r, s) = (9, 1)$ in (2.4), we obtain

$$\sum_{n=0}^{\infty} c_{9,1}(n)q^n \equiv \frac{f(q, q^9)}{f(q^2, q^{18})} \pmod{2}. \quad (2.7)$$

Substituting (2.6) into (2.7) and extracting those terms in which the power of q is congruent to 0 modulo 2, then replacing q^2 by q , we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} c_{9,1}(2n)q^n &\equiv \frac{f(q^6, q^{14})}{f(q, q^9)} \\ &\equiv \frac{f(q, q^9)f(q^6, q^{14})}{f(q^2, q^{18})} \pmod{2}. \end{aligned} \quad (\text{by (2.3)}) \quad (2.8)$$

If we substitute (2.6) into (2.8) and pick out those terms in which the power of q is congruent to 1 modulo 2, then divided by q and replace q^2 by q , we deduce that

$$\begin{aligned} \sum_{n=0}^{\infty} c_{9,1}(4n+2)q^n &\equiv \frac{f(q^4, q^{16})f(q^3, q^7)}{f(q, q^9)} \\ &\equiv \frac{f(q^4, q^{16})}{f(q^2, q^{18})} f(q, q^9)f(q^3, q^7) \pmod{2}. \end{aligned} \quad (\text{by (2.3)}) \quad (2.9)$$

It follows from [4, Entry 29] that if $ab = cd$, then

$$f(a, b)f(c, d) = f(ac, bd)f(ad, bc) + af(b/c, ac^2d)f(b/d, acd^2). \quad (2.10)$$

Taking $(a, b, c, d) = (q, q^9, q^3, q^7)$ in (2.10), we arrive at

$$f(q, q^9)f(q^3, q^7) = f(q^4, q^{16})f(q^8, q^{12}) + qf(q^2, q^{18})f(q^6, q^{14}). \quad (2.11)$$

If we substitute (2.11) into (2.9) and extract those terms in which the power of q is congruent to 0 modulo 2, then replace q^2 by q , we arrive at

$$\begin{aligned} \sum_{n=0}^{\infty} c_{9,1}(8n+2)q^n &\equiv \frac{f(q^2, q^8)^2 f(q^4, q^6)}{f(q, q^9)} \\ &\equiv \frac{f(q, q^4)^4 f(q^2, q^3)^2}{f(q, q^9)} \pmod{2}. \end{aligned} \quad (\text{by (2.3)}) \quad (2.12)$$

It is easy to check that

$$\begin{aligned} \frac{f(q, q^4)^4 f(q^2, q^3)^2}{f(q, q^9)} &= \frac{f(q, q^4)^3 f(q^2, q^3)^2 f(q^4, q^6) f_5}{f_{10}^2} \\ &\equiv \frac{f(q, q^4)^3 f(q^2, q^3)^4 f_5}{f_{10}^2} \pmod{2}. \end{aligned} \quad (\text{by (2.3)}) \quad (2.13)$$

Combining (2.12) and (2.13) yields

$$\sum_{n=0}^{\infty} c_{9,1}(8n+2)q^n \equiv \frac{f(q, q^4)^3 f(q^2, q^3)^4 f_5}{f_{10}^2} \pmod{2}. \quad (2.14)$$

It is easy to verify that

$$f(q, q^4)f(q^2, q^3) = \frac{f_2 f_5^3}{f_1 f_{10}}. \quad (2.15)$$

In light of (2.14) and (2.15),

$$\begin{aligned} \sum_{n=0}^{\infty} c_{9,1}(8n+2)q^n &\equiv \frac{f_2^3 f_5^{10}}{f_1^3 f_{10}^5} f(q^2, q^3) \\ &\equiv \frac{f_2^2}{f_1} f(q^2, q^3) \pmod{2}, \end{aligned} \quad (2.16)$$

where we have used the fact that for any positive integer k ,

$$f_k^2 \equiv f_{2k} \pmod{2}. \quad (2.17)$$

It is well known that

$$\sum_{m=0}^{\infty} q^{m(m+1)/2} = \frac{f_2^2}{f_1}. \quad (2.18)$$

In view of (1.1), (2.16) and (2.18),

$$\sum_{n=0}^{\infty} c_{9,1}(8n+2)q^n \equiv \sum_{m=0}^{\infty} q^{m(m+1)/2} \sum_{k=-\infty}^{\infty} q^{(5k^2+k)/2} \pmod{2},$$

which can be rewritten as

$$\sum_{n=0}^{\infty} c_{9,1}(8n+2)q^{40n+6} \equiv \sum_{m=0}^{\infty} \sum_{k=-\infty}^{\infty} q^{5(2m+1)^2 + (10k+1)^2} \pmod{2}.$$

Therefore, if $40n+6$ is not of the form $5x^2+y^2$, then $c_{9,1}(8n+2) \equiv 0 \pmod{2}$. Note that if N is of the form $5x^2+y^2$ and p is a prime with $p \equiv 11, 19 \pmod{20}$, then $\nu_p(N)$ is even since $\left(\frac{-5}{p}\right) = -1$. Here $\nu_p(N)$ denotes the highest power of p dividing N and $\left(\frac{\cdot}{p}\right)$ denotes the Legendre symbol. It is easy to check that if $p \nmid n$, then for any $\alpha \in \mathbb{N}$,

$$\nu_p \left(40 \left(p^{2\alpha+1}n + \frac{3(p^{2\alpha+2}-1)}{20} \right) + 6 \right) = \nu_p(40p^{2\alpha+1}n + 6p^{2\alpha+2}) = 2\alpha + 1$$

is odd and $40 \left(p^{2\alpha+1}n + \frac{3(p^{2\alpha+2}-1)}{20} \right) + 6$ is not of the form $5x^2+y^2$. Therefore, if p is a prime with $p \equiv 11, 19 \pmod{20}$ and $p \nmid n$, then for any $\alpha \in \mathbb{N}$,

$$c_{9,1} \left(8 \left(p^{2\alpha+1}n + \frac{3(p^{2\alpha+2}-1)}{20} \right) + 2 \right) \equiv 0 \pmod{2},$$

from which, we arrive at (2.1).

Substituting (2.11) into (2.9) and picking out those terms in which the power of q is congruent to 1 modulo 2, then dividing by q and replacing q^2 by q , we get

$$\sum_{n=0}^{\infty} c_{9,1}(8n+6)q^n \equiv f(q^2, q^8)f(q^3, q^7) \pmod{2}. \quad (2.19)$$

In view of (1.1) and (2.19)

$$\sum_{n=0}^{\infty} c_{9,1}(8n+6)q^n \equiv \sum_{m,k=-\infty}^{\infty} q^{5m^2+3m+5k^2+2k} \pmod{2}. \quad (2.20)$$

It follows from (2.20) that

$$\sum_{n=0}^{\infty} c_{9,1}(8n+6)q^{20n+13} \equiv \sum_{m,k=-\infty}^{\infty} q^{(10m+3)^2+(10k+2)^2} \pmod{2}.$$

Thus, if $20n+13$ is not of the form x^2+y^2 , then $c_{9,1}(8n+6)$ is even. If N is of the form x^2+y^2 and p is a prime with $p \equiv 11, 19 \pmod{20}$, then $\nu_p(N)$ is even because $\left(\frac{-1}{p}\right) = -1$. It is easy to check that if $p \nmid n$, then for any $\alpha \in \mathbb{N}$,

$$\nu_p \left(20 \left(p^{2\alpha+1}n + \frac{13(p^{2\alpha+2}-1)}{20} \right) + 13 \right) = \nu_p(20p^{2\alpha+1}n + 13p^{2\alpha+2}) = 2\alpha + 1$$

is odd and $20 \left(p^{2\alpha+1}n + \frac{13(p^{2\alpha+2}-1)}{24} \right) + 13$ is not of the form x^2+y^2 . Thus, if p is a prime with $p \equiv 11, 19 \pmod{20}$ and $p \nmid n$, then for any $\alpha \in \mathbb{N}$,

$$c_{9,1} \left(8 \left(p^{2\alpha+1}n + \frac{13(p^{2\alpha+2}-1)}{20} \right) + 6 \right) \equiv 0 \pmod{2},$$

which yields (2.2). This completes the proof of Theorem 2.1. ■

3 Parity results for $c_{7,3}(n)$

Theorem 3.1 (1) For $n \geq 0$,

$$c_{7,3}(16n+5) \equiv 0 \pmod{2}. \quad (3.1)$$

(2) For $n \geq 0$, $c_{7,3}(16n+13) \equiv 0 \pmod{2}$ if and only if n can not be represented in the form $n = 5k^2 + 4k$ for some integer k .

(3) Let p be a prime with $p \equiv 11, 19 \pmod{20}$. For $n, k \geq 0$ with $p \nmid n$,

$$c_{7,3} \left(8p^{2k+1}n + \frac{34p^{2k+2}+1}{5} \right) \equiv 0 \pmod{2}. \quad (3.2)$$

Remark. Note that the second part (2) of Theorem 3.1 is analogous to the parity results for $c_{9,1}(16n+14)$ proved by Keith [11].

Proof. Setting $(r, s) = (7, 3)$ in (2.4) yields

$$\sum_{n=0}^{\infty} c_{7,3}(n)q^n \equiv \frac{f(q^3, q^7)}{f(q^6, q^{14})} \pmod{2}. \quad (3.3)$$

Taking $(a, b) = (q^3, q^7)$ in (2.5) yields

$$f(q^3, q^7) = f(q^{16}, q^{24}) + q^3 f(q^4, q^{36}). \quad (3.4)$$

Substituting (3.4) into (3.3) and extracting those terms in which the power of q is congruent to 1 modulo 2, then dividing by q and replacing q^2 by q , we obtain

$$\begin{aligned}\sum_{n=0}^{\infty} c_{7,3}(2n+1)q^n &\equiv q \frac{f(q^2, q^{18})}{f(q^3, q^7)} \\ &\equiv q \frac{f(q^2, q^{18})}{f(q^6, q^{14})} f(q^3, q^7) \pmod{2}. \quad (\text{by (2.3)})\end{aligned}\quad (3.5)$$

Substituting (3.4) into (3.5), we obtain

$$\sum_{n=0}^{\infty} c_{7,3}(2n+1)q^n \equiv q \frac{f(q^2, q^{18})}{f(q^6, q^{14})} (f(q^{16}, q^{24}) + q^3 f(q^4, q^{36})) \pmod{2}.$$

Thus, we can immediately read off the following:

$$\begin{aligned}\sum_{n=0}^{\infty} c_{7,3}(4n+1)q^n &\equiv q^2 \frac{f(q, q^9)}{f(q^3, q^7)} f(q^2, q^{18}) \\ &\equiv q^2 \frac{f(q^2, q^{18})}{f(q^6, q^{14})} f(q, q^9) f(q^3, q^7) \pmod{2} \quad (\text{by (2.3)})\end{aligned}\quad (3.6)$$

and

$$\begin{aligned}\sum_{n=0}^{\infty} c_{7,3}(4n+3)q^n &\equiv \frac{f(q^8, q^{12})}{f(q^3, q^7)} f(q, q^9) \\ &\equiv \frac{f(q^8, q^{12})}{f(q^6, q^{14})} f(q, q^9) f(q^3, q^7) \pmod{2}. \quad (\text{by (2.3)})\end{aligned}\quad (3.7)$$

If we substitute (2.11) into (3.6) and pick out those terms in which the power of q is congruent to 1 modulo 2, then divided by q and replace q^2 by q , we obtain

$$\begin{aligned}\sum_{n=0}^{\infty} c_{7,3}(8n+5)q^n &\equiv q f(q, q^9)^2 \\ &\equiv q f(q^2, q^{18}) \pmod{2}, \quad (\text{by (2.3)})\end{aligned}$$

which yields (3.1) and

$$\sum_{n=0}^{\infty} c_{7,3}(16n+13)q^n \equiv f(q, q^9) \pmod{2}.$$

In addition, it follows from (1.1) and the above congruence that

$$\sum_{n=0}^{\infty} c_{7,3}(16n+13)q^n \equiv \sum_{k=-\infty}^{\infty} q^{5k^2+4k} \pmod{2}.$$

The second part (2) of Theorem 3.1 follows from the above congruence.

Substituting (2.11) into (3.7) and picking out those terms in which the power of q is congruent to 1 modulo 2, then dividing by q and replacing q^2 by q , we obtain

$$\sum_{n=0}^{\infty} c_{7,3}(8n+7)q^n \equiv f(q, q^9) f(q^4, q^6) \pmod{2}. \quad (3.8)$$

In light of (1.1) and (3.8),

$$\sum_{n=0}^{\infty} c_{7,3}(8n+7)q^n \equiv \sum_{m,k=-\infty}^{\infty} q^{5k^2+4k+5m^2+m} \pmod{2}. \quad (3.9)$$

We can rewrite (3.9) as

$$\sum_{n=0}^{\infty} c_{7,3}(8n+7)q^{20n+17} \equiv \sum_{m,k=-\infty}^{\infty} q^{(10k+4)^2+(10m+1)^2} \pmod{2}. \quad (3.10)$$

Using (3.10) and the same method for proving (2.2), we can get (3.2). This completes the proof of Theorem 3.1. \blacksquare

4 Parity results for $c_{13,1}(n)$

In this section, we prove some parity results for $c_{13,1}(n)$. In particular, we confirm Conjecture 1.1.

The following theorem generalizes (1.8) and (1.9).

Theorem 4.1 *For $n, k \geq 0$,*

$$c_{13,1}\left(2^{3k+5}n + \frac{19 \times 2^{3k+3} + 9}{7}\right) \equiv 0 \pmod{2}, \quad (4.1)$$

$$c_{13,1}\left(2^{3k+6}n + \frac{27 \times 2^{3k+4} + 9}{7}\right) \equiv 0 \pmod{2}, \quad (4.2)$$

$$c_{13,1}\left(2^{3k+7}n + \frac{3 \times 2^{3k+5} + 9}{7}\right) \equiv 0 \pmod{2}. \quad (4.3)$$

Setting $k = 0$ in (4.1) and (4.2), we obtain the following corollary.

Corollary 4.2 *Congruences (1.8) and (1.9) are true.*

Proof. Taking $(r, s) = (13, 1)$ in (2.4) yields

$$\sum_{n=0}^{\infty} c_{13,1}(n)q^n \equiv \frac{f(q, q^{13})}{f(q^2, q^{26})} \pmod{2}. \quad (4.4)$$

Setting $(a, b) = (q, q^{13})$ in (2.5), we arrive at

$$f(q, q^{13}) = f(q^{16}, q^{40}) + qf(q^{12}, q^{44}). \quad (4.5)$$

Substituting (4.5) into (4.4) and extracting those terms in which the power of q is congruent to 1 modulo 2, then dividing by q and replacing q^2 by q , we deduce that

$$\sum_{n=0}^{\infty} c_{13,1}(2n+1)q^n \equiv \frac{f(q^6, q^{22})}{f(q, q^{13})}$$

$$\equiv \frac{f(q^6, q^{22})}{f(q^2, q^{26})} f(q, q^{13}) \pmod{2}. \quad (\text{by (2.3)}) \quad (4.6)$$

Substituting (4.5) into (4.6) and picking out those terms in which the power of q is congruent to 1 modulo 2, then dividing by q and replacing q^2 by q , we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} c_{13,1}(4n+3)q^n &\equiv \frac{f(q^3, q^{11})}{f(q, q^{13})} f(q^6, q^{22}) \\ &\equiv \frac{f(q^6, q^{22})}{f(q^2, q^{26})} f(q, q^{13}) f(q^3, q^{11}) \pmod{2}. \quad (\text{by (2.3)}) \end{aligned} \quad (4.7)$$

Setting $(a, b, c, d) = (q, q^{13}, q^3, q^{11})$ in (2.10) yields

$$f(q, q^{13}) f(q^3, q^{11}) = f(q^4, q^{24}) f(q^{12}, q^{16}) + q f(q^{10}, q^{18}) f(q^2, q^{26}). \quad (4.8)$$

If we substitute (4.8) into (4.7) and pick out those terms in which the power of q is congruent to 1 modulo 2, then divided by q and replace q^2 by q , we have

$$\sum_{n=0}^{\infty} c_{13,1}(8n+7)q^n \equiv f(q^3, q^{11}) f(q^5, q^9) \pmod{2}. \quad (4.9)$$

Putting $(a, b, c, d) = (q^3, q^{11}, q^5, q^9)$ in (2.10) yields

$$f(q^3, q^{11}) f(q^5, q^9) = f(q^8, q^{20}) f(q^{12}, q^{16}) + q^3 f(q^6, q^{22}) f(q^2, q^{26}). \quad (4.10)$$

In light of (4.9) and (4.10),

$$\sum_{n=0}^{\infty} c_{13,1}(8n+7)q^n \equiv f(q^8, q^{20}) f(q^{12}, q^{16}) + q^3 f(q^6, q^{22}) f(q^2, q^{26}) \pmod{2}, \quad (4.11)$$

which yields

$$\sum_{n=0}^{\infty} c_{13,1}(16n+7)q^n \equiv f(q^4, q^{10}) f(q^6, q^8) \pmod{2}, \quad (4.12)$$

and

$$\sum_{n=0}^{\infty} c_{13,1}(16n+15)q^n \equiv q f(q, q^{13}) f(q^3, q^{11}) \pmod{2}. \quad (4.13)$$

It follows from (4.12) that for $n \geq 0$,

$$c_{13,1}(32n+23) \equiv 0 \pmod{2}. \quad (4.14)$$

Thanks to (4.8) and (4.13),

$$\sum_{n=0}^{\infty} c_{13,1}(16n+15)q^n \equiv q f(q^4, q^{24}) f(q^{12}, q^{16}) + q^2 f(q^{10}, q^{18}) f(q^2, q^{26}) \pmod{2}, \quad (4.15)$$

which yields

$$\sum_{n=0}^{\infty} c_{13,1}(32n+15)q^n \equiv q f(q, q^{13}) f(q^5, q^9) \pmod{2} \quad (4.16)$$

and

$$\sum_{n=0}^{\infty} c_{13,1}(32n+31)q^n \equiv f(q^2, q^{12})f(q^6, q^8) \pmod{2}. \quad (4.17)$$

It follows from (4.17) that for $n \geq 0$,

$$c_{13,1}(64n+63) \equiv 0 \pmod{2}. \quad (4.18)$$

Setting $(a, b, c, d) = (q, q^{13}, q^5, q^9)$ in (2.10) yields

$$f(q, q^{13})f(q^5, q^9) = f(q^6, q^{22})f(q^{10}, q^{18}) + qf(q^8, q^{20})f(q^4, q^{24}). \quad (4.19)$$

In view of (4.16) and (4.19),

$$\sum_{n=0}^{\infty} c_{13,1}(32n+15)q^n \equiv qf(q^6, q^{22})f(q^{10}, q^{18}) + q^2f(q^8, q^{20})f(q^4, q^{24}) \pmod{2}, \quad (4.20)$$

from which, we arrive at

$$\sum_{n=0}^{\infty} c_{13,1}(64n+15)q^n \equiv qf(q^4, q^{10})f(q^2, q^{12}) \pmod{2} \quad (4.21)$$

and

$$\sum_{n=0}^{\infty} c_{13,1}(64n+47)q^n \equiv f(q^3, q^{11})f(q^5, q^9) \pmod{2}. \quad (4.22)$$

It follows from (4.21) that for $n \geq 0$,

$$c_{13,1}(128n+15) \equiv 0 \pmod{2}. \quad (4.23)$$

In light of (4.9) and (4.22),

$$c_{13,1}(64n+47) \equiv c_{13,1}(8n+7) \pmod{2}. \quad (4.24)$$

By (4.24) and mathematical induction,

$$c_{13,1} \left(2^{3k+3}n + \frac{5 \times 2^{3k+3} + 9}{7} \right) \equiv c_{13,1}(8n+7) \pmod{2}. \quad (4.25)$$

Replacing n by $4n+2$, $8n+7$ and $16n+1$ in (4.25) and using (4.14), (4.18) and (4.23), we arrive at (4.1), (4.2) and (4.3), respectively. This completes the proof of Theorem 4.1. \blacksquare

Theorem 4.3 *Let p be a prime with $p \equiv 15, 27 \pmod{28}$. For $m, n, k \geq 0$ with $p \nmid n$,*

$$c_{13,1} \left(2^{3k+3}p^{2m+1}n + \frac{5 \times 2^{3k+3}p^{2m+2} + 9}{7} \right) \equiv 0 \pmod{2}. \quad (4.26)$$

Proof. In view of (1.1) and (4.9),

$$\sum_{n=0}^{\infty} c_{13,1}(8n+7)q^n \equiv \sum_{m,k=-\infty}^{\infty} q^{7m^2+4m+7k^2+2k} \pmod{2}. \quad (4.27)$$

Using (4.27) and the same method for proving (2.2), we can prove that if p is a prime with $p \equiv 15, 27 \pmod{28}$, then for $m, n \geq 0$ with $p \nmid n$,

$$c_{13,1} \left(8p^{2m+1}n + \frac{40p^{2m+2} + 9}{7} \right) \equiv 0 \pmod{2}. \quad (4.28)$$

Replacing n by $p^{2m+1}n + \frac{5(p^{2m+2}-1)}{7}$ ($p \nmid n$) in (4.25) and utilizing (4.28), we arrive at (4.26). The proof is complete. \blacksquare

Theorem 4.4 *Let p be a prime with $p \equiv 15, 27 \pmod{28}$. For $n, k \geq 0$ with $p \nmid n$,*

$$c_{13,1} \left(8p^{2k+1}n + \frac{26p^{2k+2} + 9}{7} \right) \equiv 0 \pmod{2}. \quad (4.29)$$

Proof. Substituting (4.5) into (4.6) and picking out those terms in which the power of q is congruent to 0 modulo 2, then replacing q^2 by q , we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} c_{13,1}(4n+1)q^n &\equiv \frac{f(q^3, q^{11})}{f(q, q^{13})} f(q^8, q^{20}) \\ &\equiv \frac{f(q^8, q^{20})}{f(q^2, q^{26})} f(q, q^{13}) f(q^3, q^{11}) \pmod{2}. \quad (\text{by (2.3)}) \end{aligned} \quad (4.30)$$

If we substitute (4.8) into (4.30) and pick out those terms in which the power of q is congruent to 1 modulo 2, then divided by q and replace q^2 by q , we deduce that

$$\sum_{n=0}^{\infty} c_{13,1}(8n+5)q^n \equiv f(q^4, q^{10}) f(q^5, q^9) \pmod{2}. \quad (\text{by (2.3)}) \quad (4.31)$$

Based on (1.1) and (4.31),

$$\sum_{n=0}^{\infty} c_{13,1}(8n+5)q^n \equiv \sum_{m,k=-\infty}^{\infty} q^{7m^2+3m+7k^2+2k} \pmod{2}. \quad (4.32)$$

Using (4.32) and the same method for proving (2.2), we can prove Theorem 4.4. This completes the proof. \blacksquare

To conclude this section, we give a proof of (1.10).

Corollary 4.5 *Congruence (1.10) is true.*

Proof. It is easy to check that

$$\sum_{m=-\infty}^{\infty} q^{7m^2+4m} = \sum_{m=-\infty}^{\infty} q^{63m^2+12m} + \sum_{m=-\infty}^{\infty} q^{63m^2+54m+11} + \sum_{m=-\infty}^{\infty} q^{63m^2+96m+36} \quad (4.33)$$

and

$$\sum_{k=-\infty}^{\infty} q^{7k^2+2k} = \sum_{k=-\infty}^{\infty} q^{63k^2+6k} + \sum_{k=-\infty}^{\infty} q^{63k^2+48k+9} + \sum_{k=-\infty}^{\infty} q^{63k^2+90k+32}. \quad (4.34)$$

Substituting (4.33) and (4.34) into (4.27) and picking out those terms in which the power of q is congruent to 1 modulo 3, then dividing by q and replacing q^3 by q , we obtain

$$\sum_{n=0}^{\infty} c_{13,1}(24n+15)q^n \equiv q^{14} \sum_{m,k=-\infty}^{\infty} q^{21m^2+18m+21k^2+30k} \pmod{2},$$

from which, we deduce that for $n \geq 0$,

$$c_{13,1}(72n+15) \equiv c_{13,1}(72n+39) \equiv 0 \pmod{2}. \quad (4.35)$$

In addition,

$$\sum_{m=-\infty}^{\infty} q^{7m^2+3m} = \sum_{m=-\infty}^{\infty} q^{63m^2+9m} + \sum_{m=-\infty}^{\infty} q^{63m^2+51m+10} + \sum_{m=-\infty}^{\infty} q^{63m^2+93m+34}. \quad (4.36)$$

If we substitute (4.34) and (4.36) into (4.32) and pick out those terms in which the power of q is congruent to 2 modulo 3, then divided by q^2 and replace q^3 by q , we obtain

$$\sum_{n=0}^{\infty} c_{13,1}(24n+21)q^n \equiv q^{10} \sum_{m,k=-\infty}^{\infty} q^{21m^2+3m+21k^2+30k} \pmod{2},$$

which implies that for $n \geq 0$,

$$c_{13,1}(72n+21) \equiv c_{13,1}(72n+69) \equiv 0 \pmod{2}. \quad (4.37)$$

Congruence (1.10) follows from (4.35) and (4.37). The proof is complete. \blacksquare

5 Parity results for $c_{11,3}(n)$

Theorem 5.1 For $n, k \geq 0$,

$$c_{11,3}\left(2^{3k+5}n + \frac{3 \times 2^{3k+3} + 4}{7}\right) \equiv 0 \pmod{2}, \quad (5.1)$$

$$c_{11,3}\left(2^{3k+6}n + \frac{19 \times 2^{3k+4} + 4}{7}\right) \equiv 0 \pmod{2}, \quad (5.2)$$

$$c_{11,3}\left(2^{3k+7}n + \frac{27 \times 2^{3k+5} + 4}{7}\right) \equiv 0 \pmod{2}. \quad (5.3)$$

Proof. Setting $(r, s) = (11, 3)$ in (2.4), we obtain

$$\sum_{n=0}^{\infty} c_{11,3}(n)q^n \equiv \frac{f(q^3, q^{11})}{f(q^6, q^{22})} \pmod{2}. \quad (5.4)$$

Taking $(a, b) = (q^3, q^{11})$ in (2.5) yields

$$f(q^3, q^{11}) = f(q^{20}, q^{36}) + q^3 f(q^8, q^{48}). \quad (5.5)$$

Substituting (5.5) into (5.4) and extracting those terms in which the power of q is congruent to 0 modulo 2, then replacing q^2 by q , we arrive at

$$\begin{aligned}\sum_{n=0}^{\infty} c_{11,3}(2n)q^n &\equiv \frac{f(q^{10}, q^{18})}{f(q^3, q^{11})} \\ &\equiv \frac{f(q^{10}, q^{18})}{f(q^6, q^{22})} f(q^3, q^{11}) \pmod{2}. \quad (\text{by (2.3)})\end{aligned}\quad (5.6)$$

Substituting (5.5) into (5.6) and extracting those terms in which the power of q is congruent to 0 modulo 2, then replacing q^2 by q , we arrive at

$$\begin{aligned}\sum_{n=0}^{\infty} c_{11,3}(4n)q^n &\equiv \frac{f(q^5, q^9)}{f(q^3, q^{11})} f(q^{10}, q^{18}) \\ &\equiv \frac{f(q^{10}, q^{18})}{f(q^6, q^{22})} f(q^3, q^{11}) f(q^5, q^9) \pmod{2}. \quad (\text{by (2.3)})\end{aligned}\quad (5.7)$$

Substituting (4.10) into (5.7) and extracting those terms in which the power of q is congruent to 1 modulo 2, then dividing them by q and replacing q^2 by q , we have

$$\sum_{n=0}^{\infty} c_{11,3}(8n+4)q^n \equiv q f(q, q^{13}) f(q^5, q^9) \pmod{2}. \quad (5.8)$$

It follows from (4.16) and (5.8) that for $n \geq 0$,

$$c_{11,3}(8n+4) \equiv c_{13,1}(32n+15) \pmod{2}. \quad (5.9)$$

Replacing n by $2^{3k+2}n + \frac{3(2^{3k}-1)}{7}$ in (5.9) and utilizing (4.3), we obtain (5.1).

Replacing n by $2^{3k+3}n + \frac{19 \times 2^{3k+1}-3}{7}$ in (5.9) yields

$$c_{11,3}\left(2^{3k+6}n + \frac{19 \times 2^{3k+4} + 4}{7}\right) \equiv c_{13,1}\left(2^{3k+8}n + \frac{19 \times 2^{3k+6} + 9}{7}\right) \pmod{2}. \quad (5.10)$$

Replacing k by $k+1$ in (4.1) yields

$$c_{13,1}\left(2^{3k+8}n + \frac{19 \times 2^{3k+6} + 9}{7}\right) \equiv 0 \pmod{2},$$

from which with (5.10), we arrive at (5.2).

Replacing n by $2^{3k+4}n + \frac{27 \times 2^{3k+2}-3}{7}$ in (5.9) yields

$$c_{11,3}\left(2^{3k+7}n + \frac{27 \times 2^{3k+5} + 4}{7}\right) \equiv c_{13,1}\left(2^{3k+9}n + \frac{27 \times 2^{3k+7} + 9}{7}\right) \pmod{2}. \quad (5.11)$$

It follows from (4.2) that

$$c_{13,1}\left(2^{3k+9}n + \frac{27 \times 2^{3k+7} + 9}{7}\right) \equiv 0 \pmod{2}.$$

Congruence (5.3) follows from (5.11) and the above congruence. The proof of Theorem 5.1 is complete. \blacksquare

Theorem 5.2 Let p be a prime with $p \equiv 15, 27 \pmod{28}$. For $m, n, k \geq 0$ with $p \nmid n$,

$$c_{11,3} \left(2^{3k+3} p^{2m+1} n + \frac{3 \times 2^{3k+3} p^{2m+2} + 4}{7} \right) \equiv 0 \pmod{2}. \quad (5.12)$$

Proof. Replacing n by $4n + 1$ in (4.25) yields

$$c_{13,1}(32n + 15) \equiv c_{13,1} \left(2^{3k+5} n + \frac{3 \times 2^{3k+5} + 9}{7} \right) \pmod{2}. \quad (5.13)$$

Replacing n by $2^{3k} n + \frac{3(2^{3k}-1)}{7}$ in (5.9), we get

$$c_{11,3} \left(2^{3k+3} n + \frac{3 \times 2^{3k+3} + 4}{7} \right) \equiv c_{13,1} \left(2^{3k+5} n + \frac{3 \times 2^{3k+5} + 9}{7} \right) \pmod{2}. \quad (5.14)$$

Combining (5.9), (5.13) and (5.14) yields

$$c_{11,3} \left(2^{3k+3} n + \frac{3 \times 2^{3k+3} + 4}{7} \right) \equiv c_{11,3}(8n + 4) \pmod{2}. \quad (5.15)$$

In view of (1.1) and (5.8),

$$\sum_{n=0}^{\infty} c_{11,3}(8n + 4)q^{n-1} \equiv \sum_{m,k=-\infty}^{\infty} q^{7m^2+6m+7k^2+2k} \pmod{2}. \quad (5.16)$$

Using (5.16) and the same method for proving (2.2), we can prove that if p is a prime with $p \equiv 15, 27 \pmod{28}$, then for $m, n \geq 0$ with $p \nmid n$,

$$c_{11,3} \left(8p^{2m+1} n + \frac{24p^{2m+2} + 4}{7} \right) \equiv 0 \pmod{2}. \quad (5.17)$$

Replacing n by $p^{2m+1} n + \frac{3(p^{2m+2}-1)}{7}$ in (5.15) and using (5.17), we arrive at (5.12). This completes the proof of Theorem 5.2. \blacksquare

Theorem 5.3 Let p be a prime with $p \equiv 15, 27 \pmod{28}$. For $n, k \geq 0$ with $p \nmid n$,

$$c_{11,3} \left(8p^{2k+1} n + \frac{122p^{2k+2} + 4}{7} \right) \equiv 0 \pmod{2}. \quad (5.18)$$

Proof. Substituting (5.5) into (5.6) and extracting those terms in which the power of q is congruent to 1 modulo 2, then dividing by q and replacing q^2 by q , we arrive at

$$\begin{aligned} \sum_{n=0}^{\infty} c_{11,3}(4n + 2)q^n &\equiv q \frac{f(q^5, q^9)}{f(q^3, q^{11})} f(q^4, q^{24}) \\ &\equiv q \frac{f(q^4, q^{24})}{f(q^6, q^{22})} f(q^3, q^{11}) f(q^5, q^9) \pmod{2}. \quad (\text{by (2.3)}) \end{aligned} \quad (5.19)$$

If we substitute (4.10) into (5.19) and pick out those terms in which the power of q is congruent to 0 modulo 2, then replace q^2 by q , we arrive at

$$\sum_{n=0}^{\infty} c_{11,3}(8n + 2)q^n \equiv q^2 f(q^2, q^{12}) f(q, q^{13}) \pmod{2}. \quad (5.20)$$

In light of (1.1) and (5.20),

$$\sum_{n=0}^{\infty} c_{11,3}(8n+2)q^n \equiv \sum_{m,k=-\infty}^{\infty} q^{7m^2+6m+7k^2+5k+2} \pmod{2}.$$

Utilizing the above congruence and the same method for proving (2.2), we can prove (5.18). The proof of Theorem 5.3 is complete. \blacksquare

6 Parity results for $c_{9,5}(n)$

Theorem 6.1 For $n, k \geq 0$,

$$c_{9,5} \left(2^{3k+5}n + \frac{27 \times 2^{3k+3} + 1}{7} \right) \equiv 0 \pmod{2}, \quad (6.1)$$

$$c_{9,5} \left(2^{3k+6}n + \frac{3 \times 2^{3k+4} + 1}{7} \right) \equiv 0 \pmod{2}, \quad (6.2)$$

$$c_{9,5} \left(2^{3k+7}n + \frac{19 \times 2^{3k+5} + 1}{7} \right) \equiv 0 \pmod{2}. \quad (6.3)$$

Proof. Setting $(r, s) = (9, 5)$ in (2.4) yields

$$\sum_{n=0}^{\infty} c_{9,5}(n)q^n \equiv \frac{f(q^5, q^9)}{f(q^{10}, q^{18})} \pmod{2}. \quad (6.4)$$

Taking $(a, b) = (q^5, q^9)$ in (2.5) yields

$$f(q^5, q^9) = f(q^{24}, q^{32}) + q^5 f(q^4, q^{52}). \quad (6.5)$$

Substituting (6.5) into (6.4) and extracting those terms in which the power of q is congruent to 1 modulo 2, then dividing by q and replacing q^2 by q , we arrive at

$$\begin{aligned} \sum_{n=0}^{\infty} c_{9,5}(2n+1)q^n &\equiv q^2 \frac{f(q^2, q^{26})}{f(q^5, q^9)} \\ &\equiv q^2 \frac{f(q^2, q^{26})}{f(q^{10}, q^{18})} f(q^5, q^9) \pmod{2}. \quad (\text{by (2.3)}) \end{aligned} \quad (6.6)$$

If we substitute (6.5) into (6.6) and extract those terms in which the power of q is congruent to 1 modulo 2, then divided by q and replace q^2 by q , we arrive at

$$\begin{aligned} \sum_{n=0}^{\infty} c_{9,5}(4n+3)q^n &\equiv q^3 \frac{f(q, q^{13})}{f(q^5, q^9)} f(q^2, q^{26}) \\ &\equiv q^3 \frac{f(q^2, q^{26})}{f(q^{10}, q^{18})} f(q, q^{13}) f(q^5, q^9) \pmod{2}. \quad (\text{by (2.3)}) \end{aligned} \quad (6.7)$$

Substituting (4.19) into (6.7) and extracting those terms in which the power of q is congruent to 1 modulo 2, then dividing by q and replacing q^2 by q , we arrive at

$$\sum_{n=0}^{\infty} c_{9,5}(8n+7)q^n \equiv q f(q, q^{13}) f(q^3, q^{11}) \pmod{2}. \quad (6.8)$$

In light of (4.13) and (6.8),

$$c_{9,5}(8n+7) \equiv c_{13,1}(16n+15) \pmod{2}. \quad (6.9)$$

Replacing n by $2^{3k+2}n + \frac{27 \times 2^{3k} - 6}{7}$ in (6.9) and using (4.2), we get (6.1).

Replacing n by $2^{3k+3}n + \frac{3 \times 2^{3k+1} - 6}{7}$ in (6.9) and using (4.3), we get (6.2).

Replacing n by $2^{3k+4}n + \frac{19 \times 2^{3k+2} - 6}{7}$ in (6.9) yields

$$c_{9,5} \left(2^{3k+7}n + \frac{19 \times 2^{3k+5} + 1}{7} \right) \equiv c_{13,1} \left(2^{3k+8}n + \frac{19 \times 2^{3k+6} + 9}{7} \right) \pmod{2}. \quad (6.10)$$

It follows from (4.1) that

$$c_{13,1} \left(2^{3k+8}n + \frac{19 \times 2^{3k+6} + 9}{7} \right) \equiv 0 \pmod{2}. \quad (6.11)$$

Congruence (6.3) follows from (6.10) and (6.11). ■

Theorem 6.2 *Let p be a prime with $p \equiv 15, 27 \pmod{28}$. For $m, n, k \geq 0$ with $p \nmid n$,*

$$c_{9,5} \left(2^{3k+3}p^{2m+1}n + \frac{3 \times 2^{3k+4}p^{2m+2} + 1}{7} \right) \equiv 0 \pmod{2}. \quad (6.12)$$

Proof. Replacing n by $2n+1$ in (4.25), we have

$$c_{13,1} \left(2^{3k+4}n + \frac{3 \times 2^{3k+5} + 9}{7} \right) \equiv c_{13,1}(16n+15) \pmod{2}. \quad (6.13)$$

Replacing n by $2^{3k}n + \frac{3 \times 2^{3k+1} - 6}{7}$ in (6.9) yields

$$c_{9,5} \left(2^{3k+3}n + \frac{3 \times 2^{3k+4} + 1}{7} \right) \equiv c_{13,1} \left(2^{3k+4}n + \frac{3 \times 2^{3k+5} + 9}{7} \right) \pmod{2}. \quad (6.14)$$

Combining (6.9), (6.13) and (6.14), we get

$$c_{9,5} \left(2^{3k+3}n + \frac{3 \times 2^{3k+4} + 1}{7} \right) \equiv c_{9,5}(8n+7) \pmod{2}. \quad (6.15)$$

In light of (1.1) and (6.8),

$$\sum_{n=0}^{\infty} c_{9,5}(8n+7)q^n \equiv \sum_{m,k=-\infty}^{\infty} q^{7m^2+6m+7k^2+4k+1} \pmod{2}. \quad (6.16)$$

Using (6.16) and the same method for proving (2.2), we can prove that if p is a prime with $p \equiv 15, 27 \pmod{28}$, then for $m, n \geq 0$ with $p \nmid n$,

$$c_{9,5} \left(8p^{2m+1}n + \frac{48p^{2m+2} + 1}{7} \right) \equiv 0 \pmod{2}. \quad (6.17)$$

Replacing n by $p^{2m+1}n + \frac{6(p^{2m+2}-1)}{7}$ in (6.15) and using (6.17), we get (6.12). This completes the proof of Theorem 6.2. ■

Theorem 6.3 *Let p be a prime with $p \equiv 15, 27 \pmod{28}$. For $n, k \geq 0$ with $p \nmid n$,*

$$c_{9,5} \left(8p^{2k+1}n + \frac{34p^{2k+2} + 1}{7} \right) \equiv 0 \pmod{2}. \quad (6.18)$$

Proof. If we substitute (6.5) into (6.6) and pick out those terms in which the power of q is congruent to 0 modulo 2, then replace q^2 by q , we arrive at

$$\begin{aligned} \sum_{n=0}^{\infty} c_{9,5}(4n+1)q^n &\equiv q \frac{f(q, q^{13})}{f(q^5, q^9)} f(q^{12}, q^{16}) \\ &\equiv q \frac{f(q^{12}, q^{16})}{f(q^{10}, q^{18})} f(q, q^{13}) f(q^5, q^9) \pmod{2}. \quad (\text{by (2.3)}) \end{aligned} \quad (6.19)$$

Substituting (4.19) into (6.19) and extracting those terms in which the power of q is congruent to 1 modulo 2, then replacing q^2 by q , we arrive at

$$\sum_{n=0}^{\infty} c_{9,5}(8n+5)q^n \equiv f(q^6, q^8) f(q^3, q^{11}) \pmod{2}.$$

Thanks to (1.1) and the above congruence,

$$\sum_{n=0}^{\infty} c_{9,5}(8n+5)q^n \equiv \sum_{m,k=-\infty}^{\infty} q^{7m^2+m+7k^2+4k} \pmod{2}. \quad (6.20)$$

Using (6.20) and the same method for proving (2.2), we can arrive at (6.18). This completes the proof. ■

Statements and Declarations

Funding. This work was supported by the National Natural Science Foundation of China (grant no. 12371334) and the Qinlan project of Jiangsu province of China and the research project of Jiangsu Agri-animal Husbandry Vocational College (No. NSF2025CB22).

Author Contributions. The authors contributed equally to the preparation of this article. All authors read and approved the final manuscript.

Competing Interests. The authors declare that they have no conflict of interest.

Data Availability Statements. Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

References

- [1] G.E. Andrews, The Theory of Partitions, Addison-Wesley, Reading, Mass, 1976; reprinted, Cambridge University Press, 1998.
- [2] G.E. Andrews, Ramanujan's "lost" notebook. I. partial θ -functions, Adv. Math. 41 (1981) 137–172.

- [3] G.E. Andrews and S.O. Warnaar, The Bailey transform and false theta functions, *Ramanujan J.* 14 (2007) 173–188.
- [4] B.C. Berndt, *Ramanujan’s Notebooks, Part III*, Springer, New York, 1991.
- [5] K. Bringmann and A. Mila, \mathcal{W} -algebras, false theta functions and quantum modular forms, I, *Int. Math. Res. Not.* 21 (2015) 11351–11387.
- [6] H. Burson, A bijective proof of a false theta function identity from Ramanujan’s Lost Notebook, *Ann. Combin.* 23 (2019) 579–588.
- [7] S. Corteel and J. Lovejoy, Overpartitions, *Trans. Amer. Math. Soc.* 356 (2004) 1623–1635.
- [8] M.D. Hirschhorn and J.A. Sellers, Arithmetic relations for overpartitions, *J. Combin. Math. Combin. Comput.* 53 (2005) 65–73.
- [9] M.D. Hirschhorn and J.A. Sellers, Arithmetic properties of partitions with odd parts distinct, *Ramanujan J.* 22 (2010) 273–284.
- [10] C. Jennings-Shaffer and A. Milas, On q -series identities for false theta series, *Adv. Math.* 375 (2020) 107411.
- [11] W.J. Keith, Reciprocals of false theta functions, *Ramanujan J.* 68 (2025) 27.
- [12] S. Ramanujan, Some properties of $p(n)$, the number of partitions of n , *Proc. Cambridge Philos. Soc.* 19 (1919) 214–216.
- [13] S. Ramanujan, *Notebooks, vol. 2*. Tata Institute of Fundamental Research, Bombay, 1957.
- [14] S. Ramanujan. *The Lost Notebook and Other Unpublished Papers*. Narosa Pub. House, New Delhi, 1988.
- [15] L.J. Rogers, On two theorems of combinatory analysis and some allied identities, *Proc. London Math. Soc.* 16 (1917) 315–336.
- [16] L.Q. Wang, New proofs of Ramanujan’s identities on false theta functions, *Ramanujan J.* 50 (2019) 423–431.