

SL_4 -KLOOSTERMAN SUM VIA THE BRUHAT DECOMPOSITION

SUZUHO OSONOE, MAKI NAKASUJI

ABSTRACT. We define the Kloosterman sum for SL_4 over the Kloosterman set via the Bruhat decomposition and stratify the Kloosterman set using the reduced word decomposition of the Weyl group element. The Kloosterman sum for an SL_4 -long word is decomposed into finer parts (called the fine Kloosterman sum), and can be written as a finite sum of a product of two classical Kloosterman sums.

1. INTRODUCTION

The classical Kloosterman sum is defined by

$$S(m, n; c) = \sum_{\substack{a, d \pmod{c} \\ ad \equiv 1 \pmod{c}}} e\left(\frac{ma + nd}{c}\right),$$

where $m, n, c \in \mathbb{Z}$ with $c > 0$, and $e(x) = \exp(2\pi i x)$. This exponential sum was first introduced by H. D. Kloosterman in 1926. Since then, the Kloosterman sum have been deeply studied in such fields as spectral theory, modular forms and Poincaré series. For example, the Kloosterman sum appears as an exponential sum over $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z})$ in the computation of the Fourier coefficients of the classical Poincaré series.

An important result concerning this sum is the Weil bound:

$$(1) \quad |S(m, n; c)| \leq (m, n, c)^{\frac{1}{2}} c^{\frac{1}{2}} \tau(c),$$

where (m, n, c) denotes the greatest common divisor, and $\tau(c)$ is the number of positive divisors of c . This bound enabled the effective use of Kloosterman sums in the study of various areas such as the Kuznetsov trace formula, Fourier expansions and L-functions.

The classical Kloosterman sum may be defined as a sum over the Kloosterman set arising from the Bruhat decomposition of the group $SL_2(\mathbb{Z})$. In recent years, there have been attempts to generalize the Kloosterman sum to higher rank groups such as GL_n and SL_n . In these generalizations, it is necessary to replace the original number-theoretic approach with a group-theoretic one, for which Bruhat decompositions and

2020 *Mathematics Subject Classification.* Primary 11L05; Secondary 11F22.

Key words and phrases. Kloosterman sums, Weyl group, Reduced word decomposition.

This work was supported by Japan Society for the Promotion of Science, Grant-in-Aid for Scientific Research No. 22K03274(M. Nakasuji).

Weyl groups play key roles. In the case of SL_3 , the definition and properties of the generalized Kloosterman sum have been studied, and its analytic structure has been clarified. Kiral and Nakasuji ([1]) have provided a stratification of the Kloosterman sum for SL_3 using the reduced word decomposition of the Weyl group element. Additionally, the Kloosterman sum for an SL_3 -long word is decomposed into finer parts (and called the fine Kloosterman sum), which can be written as a finite sum of a product of two classical Kloosterman sums [1, Theorem 5.10].

However, the structure of the Kloosterman sum for SL_4 has not yet been sufficiently studied. In the present paper, we attempt to construct and analyze the Kloosterman sum for an SL_4 -long word, following the group-theoretic approach used for the SL_3 case in [1].

The paper is organized as follows. In Section 2, we review the Bruhat decomposition and the Weyl groups, which are essential for the group-theoretic approach to the Kloosterman sum. In particular, we give a preliminary discussion of the reduced word decomposition of the long word that will be used in the later sections. In Section 3, we define the Kloosterman sum for SL_4 and state the resulting properties.

2. PRELIMINARIES

The Bruhat decomposition plays a key role in the definition and structure of a Kloosterman sum for SL_n . For $G = SL_n$, let $B \subset G$ denote the Borel subgroup consisting of upper triangular matrices. The Bruhat decomposition can be expressed using elements of the Weyl group W as

$$G = \bigsqcup_{w \in W} BwB.$$

In the case of SL_n , the Weyl group W can be identified with the symmetric group S_n . Additionally, any element $w \in W \cong S_n$ can be written as a product of simple transposition $s_i = (i, (i+1))$, for $1 \leq i \leq n-1$.

Let

$$w_0 = \begin{pmatrix} 1 & 2 & 3 & \cdots & (n-1) & n \\ \downarrow & \downarrow & \downarrow & \cdots & \downarrow & \downarrow \\ n & (n-1) & (n-2) & \cdots & 2 & 1 \end{pmatrix} \in W.$$

In the present paper, we stratify the Kloosterman set using the reduced word decomposition of the long word:

$$w_0 = s_1(s_2s_1)(s_3s_2s_1) \cdots (s_{n-1} \cdots s_1).$$

In terms of the matrix representation of SL_n , this is expressed as

$$(2) \quad w_0 = \begin{pmatrix} & & & & (-1)^{n-1} \\ & & & \ddots & \\ & & -1 & & \\ & & & 1 & \\ & -1 & & & \\ 1 & & & & \end{pmatrix}_{n \times n}.$$

Let U be the group of unipotent matrices and T be the group of diagonal matrices. For a given $A \in UwTU$, we write its decomposition as follows.

$$(3) \quad A = \begin{pmatrix} 1 & a_{12} & a_{13} & \cdots & a_{1n} \\ & 1 & a_{23} & \cdots & a_{2n} \\ & & \ddots & \ddots & \vdots \\ & & & & 1 \end{pmatrix} w \begin{pmatrix} t_1 & & & & \\ & t_2 & & & \\ & & \ddots & & \\ & & & & t_n \end{pmatrix} \begin{pmatrix} 1 & b_{12} & b_{13} & \cdots & b_{1n} \\ & 1 & b_{23} & \cdots & b_{2n} \\ & & \ddots & \ddots & \vdots \\ & & & & 1 \end{pmatrix}.$$

The above decomposition is a finer version of the Bruhat decomposition, where a_{ij}, b_{ij} and t_i can be expressed in terms of the entries of A ([1, Proposition 3.1]). For subsets $I = \{i_1 < i_2 < \cdots < i_k\} \subseteq \{1, \dots, n\}$ and $J = \{j_1 < j_2 < \cdots < j_k\} \subseteq \{1, \dots, n\}$, let $M_{I,J}$ be the determinant of the submatrix obtained by removing the I -th row and the J -th column from the original matrix, which we write as $M_{I,J} = \langle \mathbf{e}_I^*, A\mathbf{e}_J \rangle$. For example, let

$$A = \begin{pmatrix} A_{11} & A_{12} & A_{13} & A_{14} \\ A_{21} & A_{22} & A_{23} & A_{24} \\ A_{31} & A_{32} & A_{33} & A_{34} \\ A_{41} & A_{42} & A_{43} & A_{44} \end{pmatrix}.$$

Then, the minor $M_{234,123} (= \langle \mathbf{e}_{2,3,4}^*, A\mathbf{e}_{1,2,3} \rangle)$ means

$$\det \begin{pmatrix} A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \\ A_{41} & A_{42} & A_{43} \end{pmatrix}.$$

For a given positive root α , we define a map $\iota_\alpha : SL_2(\mathbb{R}) \rightarrow SL_n(\mathbb{R})$ such that its image lies in the rank-one subgroup associated with α .

In the case of SL_4 , for the simple roots $\alpha = (1, -1, 0, 0)$, $\beta = (0, 1, -1, 0)$, $\gamma = (0, 0, 1, -1)$ and $x = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL_2$, we define

$$(4) \quad \iota_\alpha(x) = \begin{pmatrix} a & b & & \\ c & d & & \\ & & 1 & \\ & & & 1 \end{pmatrix}, \quad \iota_\beta(x) = \begin{pmatrix} 1 & & & \\ & a & b & \\ & c & d & \\ & & & 1 \end{pmatrix}, \quad \iota_\gamma(x) = \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & a & b \\ & & c & d \end{pmatrix}.$$

Using ι_* as above, we use s_* to denote the following:

$$(5) \quad s_\alpha = \iota_\alpha \left(\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \right), \quad s_\beta = \iota_\beta \left(\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \right), \quad s_\gamma = \iota_\gamma \left(\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \right).$$

These s_* terms correspond to the simple transpositions $s_\alpha = (12)$, $s_\beta = (23)$ and $s_\gamma = (34)$ in the symmetric group S_4 .

3. SL_4 -LONG WORD KLOOSTERMAN SUM

3.1. Kloosterman set and Kloosterman sum. In this section, following [1, Section 4], we define the Kloosterman sum for $SL_4(\mathbb{Z})$.

For a vector $\mathbf{c} = \{c_1, c_2, c_3\} \in \mathbb{Z}^3$, we define

$$(6) \quad t(\mathbf{c}) = \begin{pmatrix} c_1 & & & \\ & \frac{c_2}{c_1} & & \\ & & \frac{c_3}{c_2} & \\ & & & \frac{1}{c_3} \end{pmatrix},$$

and

$$(7) \quad \Omega_\omega(\mathbf{c}) = \{A \in SL_4(\mathbb{Z}) \mid A = u_L \omega t(\mathbf{c}) u_R \in B \omega B, \ u_L, u_R \in U\}.$$

Let $\mathbf{n} = \{n_1, n_2, n_3\} \in \mathbb{Z}^3$. We use the character $\psi_{\mathbf{n}}$ to denote

$$\psi_{\mathbf{n}}(u) = e(n_1 u_{1,2} + n_2 u_{2,3} + n_3 u_{3,4}),$$

where $u = (u_{i,j})$ is a unipotent matrix, and $e(x) = \exp(2\pi i x)$.

For $\mathbf{m}, \mathbf{n}, \mathbf{c} \in \mathbb{Z}^3$, we define the usual (coarse) Kloosterman sum for $SL_4(\mathbb{Z})$ as

$$(8) \quad S_\omega(\mathbf{m}, \mathbf{n}; \mathbf{c}) = \sum_{\substack{A \in U(\mathbb{Z}) \backslash \Omega_\omega(\mathbf{c}) / U(\mathbb{Z}) \\ A = u_L \omega t(\mathbf{c}) u_R}} \psi_{\mathbf{m}}(u_L) \psi_{\mathbf{n}}(u_R).$$

From here on, we consider the Kloosterman sum for SL_4 for the case of $w = w_0$ with (2) which is the long word in the Weyl group, appearing in the decomposition of A in (7).

For a given $A \in SL_4(\mathbb{Z}) \cap B w_0 B$, we write

$$(9) \quad A = \begin{pmatrix} A_{11} & A_{12} & A_{13} & A_{14} \\ A_{21} & A_{22} & A_{23} & A_{24} \\ A_{31} & A_{32} & A_{33} & A_{34} \\ A_{41} & A_{42} & A_{43} & A_{44} \end{pmatrix}.$$

In addition, for a given $A \in SL_4 \cap Uw_0TU$, we write A as in (3):

$$(10) \quad A = \begin{pmatrix} 1 & a_{12} & a_{13} & a_{14} \\ & 1 & a_{23} & a_{24} \\ & & 1 & a_{34} \\ & & & 1 \end{pmatrix} w_0 \begin{pmatrix} t_1 & & & \\ & t_2 & & \\ & & t_3 & \\ & & & t_4 \end{pmatrix} \begin{pmatrix} 1 & b_{12} & b_{13} & b_{14} \\ & 1 & b_{23} & b_{24} \\ & & 1 & b_{34} \\ & & & 1 \end{pmatrix}.$$

The diagonal matrix T in (10) becomes $t(\mathbf{c})$ in (6) when

$$(11) \quad c_1 = t_1, \quad c_2 = t_1 t_2, \quad c_3 = t_1 t_2 t_3.$$

From [1, Proposition 3.1], we obtain

$$(12) \quad t_1 = A_{41}, \quad t_1 t_2 = M_{34,12} = \begin{vmatrix} A_{31} & A_{32} \\ A_{41} & A_{42} \end{vmatrix}, \quad t_1 t_2 t_3 = M_{234,123} = \begin{vmatrix} A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \\ A_{41} & A_{42} & A_{43} \end{vmatrix}.$$

Lemma 3.1. *For a given $A \in SL_4(\mathbb{Z}) \cap Bw_0B$ as above, we have*

$$\gcd(A_{41}, A_{42}, A_{43}) = \gcd(M_{234,123}, M_{134,123}, M_{124,123}).$$

Proof. Let $f_1 = \gcd(A_{41}, A_{42}, A_{43})$ and $f_2 = \gcd(M_{234,123}, M_{134,123}, M_{124,123})$. Each minor of f_2 includes $[A_{41} \ A_{42} \ A_{43}]$. This proves $f_1 | f_2$.

For the converse, since $|A| = 1$, we can write

$$A_{41} = \begin{vmatrix} M_{234,123} & M_{234,134} & M_{234,124} \\ M_{134,123} & M_{134,134} & M_{134,124} \\ M_{124,123} & M_{124,134} & M_{124,124} \end{vmatrix}, \quad A_{42} = \begin{vmatrix} M_{234,123} & M_{234,234} & M_{234,124} \\ M_{134,123} & M_{134,234} & M_{134,124} \\ M_{124,123} & M_{124,234} & M_{124,124} \end{vmatrix},$$

$$A_{43} = \begin{vmatrix} M_{234,123} & M_{234,234} & M_{234,134} \\ M_{134,123} & M_{134,234} & M_{134,134} \\ M_{124,123} & M_{124,234} & M_{124,134} \end{vmatrix}.$$

Each minor of f_2 appears in the first column of each determinant. This means that $f_2 | f_1$. \square

We set

$$(13) \quad f = \gcd(A_{41}, A_{42}, A_{43}) = \gcd(M_{234,123}, M_{134,123}, M_{124,123}).$$

We define $D_1, D_2, D_3 \in \mathbb{Z}_{>0}$ as follows using (11) and (12),

$$(14) \quad c_1 = A_{41} = D_1 f, \quad c_2 = M_{34,12} = D_2 f, \quad c_3 = M_{234,123} = D_3 f.$$

Furthermore, we define the following set as the fine Kloosterman set

$$\Omega(D_1, D_2, D_3, f) = \{A \in SL_4(\mathbb{Z}) \mid \gcd(A_{41}, A_{42}, A_{43}) = f, \ A_{41} = D_1 f, \\ M_{34,12} = D_2 f, \ M_{234,123} = D_3 f\}.$$

Definition 3.2. For $\mathbf{m}, \mathbf{n} \in \mathbb{Z}^3$, we define the fine Kloosterman sum for SL_4 as follows, using D_1, D_2, D_3 and f defined above.

$$(15) \quad \mathcal{S}_{\omega_0}(\mathbf{m}, \mathbf{n}; D_1, D_2, D_3, f) = \sum_{\substack{A \in U(\mathbb{Z}) \backslash \Omega(D_1, D_2, D_3, f) / U(\mathbb{Z}) \\ A = u_L \omega t(D_1 f, D_2 f, D_3 f) u_R}} \psi_{\mathbf{m}}(u_L) \psi_{\mathbf{n}}(u_R).$$

Similarly as before, the (usual) coarse Kloosterman sum for SL_4 can thus be written as the sum

$$(16) \quad S_{\omega_0}(\mathbf{m}, \mathbf{n}; (c_1, c_2, c_3)) = \sum_{f | (c_1, c_2, c_3)} \mathcal{S}_{\omega_0} \left(\mathbf{m}, \mathbf{n}; \frac{c_1}{f}, \frac{c_2}{f}, \frac{c_3}{f}, f \right).$$

From (16), the properties of the fine Kloosterman sum are expected to give the properties of (usual) coarse Kloosterman sum.

3.2. Properties of fine Kloosterman sum. In this subsection, we consider the parametrization of the fine Kloosterman cells, and discuss the properties of the fine Kloosterman sum. Using the reduced word decomposition $\omega_0 = s_{\alpha} s_{\beta} s_{\alpha} s_{\gamma} s_{\beta} s_{\alpha}$ with the notation in (5), we parametrize the fine Kloosterman sets $\Omega(D_1, D_2, D_3, f)$:

$$(17) \quad A = \iota_{\alpha}(\gamma_1) \iota_{\beta}(\gamma_3) \iota_{\alpha}(\gamma_2) \iota_{\gamma}(\gamma_6) \iota_{\beta}(\gamma_5) \iota_{\alpha}(\gamma_4),$$

where

$$\gamma_i = \begin{pmatrix} x_i & b_i \\ d_i & y_i \end{pmatrix} \text{ (if } 1 \leq i \leq 5), \quad \gamma_6 = \begin{pmatrix} x_6 & b_6 \\ f & y_6 \end{pmatrix}.$$

Then we can write

$$(18) \quad D_1 = d_4 d_5, \quad D_2 = d_2 d_3 d_5, \quad D_3 = d_1 d_3.$$

The calculation of (17) leads to

$$A_{41} = d_4 d_5 f, \quad A_{42} = d_5 f y_4, \quad A_{43} = f y_5,$$

$$M_{234,123} = d_1 d_3 f, \quad M_{134,123} = d_3 f x_1, \quad M_{124,123} = f x_3.$$

If

$$\gcd(A_{41}, A_{42}) = d_5 f, \quad \gcd(M_{234,123}, M_{134,123}) = d_3 f,$$

then $\gcd(y_4, d_4) = 1$, $\gcd(x_1, d_1) = 1$, $\gcd(y_5, d_5) = 1$ and $\gcd(x_3, d_3) = 1$. In other words, y_4 is defined up to modulo d_4 , x_1 is defined up to modulo d_1 , y_5 is defined up to modulo d_5 and x_3 is defined up to modulo d_3 .

Lemma 3.3. For $A \in SL_4(\mathbb{Z})$, we have $A_{44} M_{123,123} \equiv 1 \pmod{f}$. In particular, $\gcd(A_{44}, f) = \gcd(M_{123,123}, f) = 1$.

Proof. Expanding the determinant of A along the last row, we obtain

$$\begin{aligned} |A| &= -A_{41}M_{123,234} + A_{42}M_{123,134} - A_{43}M_{123,134} + A_{44}M_{123,123} \\ &= f\{-d_1'M_{123,234} + (A_{42}/f)M_{123,134} - (A_{43}/f)M_{123,124}\} + A_{44}M_{123,123}. \end{aligned}$$

□

The calculation of (17) leads to $A_{44} = y_6$ and $M_{123,123} = x_6$. Therefore, by Lemma 3.3, we have $x_6y_6 \equiv 1 \pmod{f}$.

From [1, Proposition 3.1], for $A \in SL_4(\mathbb{Z}) \cap u_L\omega_0Tu_R$, we know that

$$u_L = \begin{pmatrix} 1 & \frac{\langle e_{1,3,4}^*, Ae_{1,2,3} \rangle}{t_1t_2t_3} & \frac{\langle e_{1,4}^*, Ae_{1,2} \rangle}{t_1t_2} & \frac{\langle e_1^*, Ae_1 \rangle}{t_1} \\ & 1 & \frac{\langle e_{2,4}^*, Ae_{1,2} \rangle}{t_1t_2} & \frac{\langle e_2^*, Ae_1 \rangle}{t_1} \\ & & 1 & \frac{\langle e_3^*, Ae_1 \rangle}{t_1} \\ & & & 1 \end{pmatrix},$$

and

$$u_R = \begin{pmatrix} 1 & \frac{\langle e_4^*, Ae_2 \rangle}{t_1} & \frac{\langle e_4^*, Ae_3 \rangle}{t_1} & \frac{\langle e_4^*, Ae_4 \rangle}{t_1} \\ & 1 & \frac{\langle e_{3,4}^*, Ae_{1,3} \rangle}{t_1t_2} & \frac{\langle e_{3,4}^*, Ae_{1,4} \rangle}{t_1t_2} \\ & & 1 & \frac{\langle e_{2,3,4}^*, Ae_{1,2,4} \rangle}{t_1t_2t_3} \\ & & & 1 \end{pmatrix}.$$

Defining $u_1 = d_1x_2 + d_2x_3y_1$, $u_2 = d_2x_4 + d_4x_5y_2$, $v_1 = d_1x_2y_3 + d_2y_1$, $v_2 = d_2x_4y_5 + d_4y_2$, we have the following proposition.

Proposition 3.4. *For a given $A \in SL_4(\mathbb{Z})$, we choose $d_1, d_2, d_3, d_4, d_5, f$ as in (14) and (18). Then for $u_1, u_2, v_1, v_2 \in \mathbb{Z}$ as above, the Bruhat decomposition of A can be written as*

$$(19) \quad A = u_L \begin{pmatrix} & & -1 \\ & 1 & \\ -1 & & \\ 1 & & \end{pmatrix} \begin{pmatrix} d_4d_5f & & & \\ & \frac{d_2d_3}{d_4} & & \\ & & \frac{d_1}{d_2d_5} & \\ & & & \frac{1}{d_1d_3f} \end{pmatrix} u_R,$$

where

$$u_L = \begin{pmatrix} 1 & \frac{x_1}{d_1} & \frac{x_1 u_1 - d_2 x_3}{d_1 d_2 d_3} & \frac{x_1 S - d_2 x_3 T + d_2 d_4 d_5 x_6}{d_1 d_2 d_3 d_4 d_5 f} \\ & 1 & \frac{u_1}{d_2 d_3} & \frac{S}{d_2 d_3 d_4 d_5 f} \\ & & 1 & \frac{T}{d_4 d_5 f} \\ & & & 1 \end{pmatrix}$$

and

$$u_R = \begin{pmatrix} 1 & \frac{y_4}{d_4} & \frac{y_5}{d_4 d_5} & \frac{y_6}{d_4 d_5 f} \\ & 1 & \frac{v_2}{d_2 d_5} & \frac{d_4 d_5 y_3 + d_3 y_6 u_2}{d_2 d_3 d_5 f} \\ & & 1 & \frac{d_5 v_1 + d_1 d_3 x_5 y_6}{d_1 d_3 f} \\ & & & 1 \end{pmatrix},$$

in which $T = d_3 u_2 + d_4 d_5 x_6 y_3$, $S = d_2 d_4 d_5 x_6 y_1 (x_3 y_3 - 1) + d_3 (u_1 u_2 - d_1 d_4 x_5)$.

Proposition 3.5. *For given nonzero integers $d_1, d_2, d_3, d_4, d_5, f$ and $x_1 \pmod{d_1}$, $y_4 \pmod{d_4}$, $x_3 \pmod{d_3}$, $y_5 \pmod{d_5}$, the product in (19) gives rise to an integral matrix if and only if the following congruence conditions are satisfied.*

$$(20) \quad d_2 d_4 d_5 x_6 - T d_2 x_3 + S x_1 \equiv 0 \pmod{d_1 d_2 d_3}$$

$$(21) \quad S \equiv 0 \pmod{d_2 d_3}$$

$$(22) \quad y_4 (d_2 d_4 d_5 x_6 - T d_2 x_3 + S x_1) + d_2 d_3 (d_2 x_3 - x_1 u_1) \equiv 0 \pmod{d_1 d_2 d_3 d_4}$$

$$(23) \quad S y_4 - d_2 d_3 u_1 \equiv 0 \pmod{d_2 d_3 d_4}$$

$$(24) \quad T y_4 - d_2 d_3 \equiv 0 \pmod{d_4}$$

$$(25) \quad y_5 (d_2 d_4 d_5 x_6 - T d_2 x_3 + S x_1) + v_2 d_3 (d_2 x_3 - x_1 u_1) + d_1 d_3 d_4 x_1 \equiv 0 \pmod{d_1 d_2 d_3 d_4 d_5}$$

$$(26) \quad d_1 d_3 d_4 - d_3 u_1 v_2 + S y_5 \equiv 0 \pmod{d_2 d_3 d_4 d_5}$$

$$(27) \quad T y_5 - d_3 v_2 \equiv 0 \pmod{d_4 d_5}$$

$$(28) \quad -(d_3 u_2 y_6 + d_4 d_5 y_3) + T y_6 \equiv 0 \pmod{d_4 d_5 f}$$

$$(29) \quad d_1 d_3 d_4 x_5 y_6 - u_1 (d_3 u_2 y_6 + d_4 d_5 y_3) + d_4 d_5 v_1 + S y_6 \equiv 0 \pmod{d_2 d_3 d_4 d_5 f}$$

$$(30) \quad x_1 (d_1 d_3 d_4 x_5 y_6 - u_1 (d_3 u_2 y_6 + d_4 d_5 y_3) + d_4 d_5 v_1 + S y_6) + d_2 d_4 d_5 x_3 y_3 + d_2 d_4 d_5 \times (x_6 y_6 - 1) + d_2 d_3 u_2 x_3 y_6 - T d_2 x_3 y_6 \equiv 0 \pmod{d_1 d_2 d_3 d_4 d_5 f}$$

In what follows, we will assume x_1 and y_4 are chosen to be relatively prime to $d_1d_2d_3d_4d_5f$ and we will denote the inverses of x_1 and y_4 by \bar{x}_1 and \bar{y}_4 , respectively.

Lemma 3.6. *For x_1, x_3, y_4 relatively prime to $d_1d_2d_3d_4d_5f$ such that $x_1\bar{x}_1 \equiv 1$, $y_4\bar{y}_4 \equiv 1$ and $x_3y_3 \equiv 1$, the following congruence conditions are satisfied.*

$$(31) \quad d_2d_3u_1 \equiv d_2^2d_3\bar{x}_1x_3 \pmod{d_1d_2d_3}$$

$$(32) \quad d_3u_2 + d_4d_5x_6y_3 \equiv d_2d_3\bar{y}_4 \pmod{d_4}$$

$$(33) \quad d_3v_2 \equiv d_2d_3\bar{y}_4y_5 \pmod{d_4}$$

$$(34) \quad d_2d_3d_4(d_5v_1 + d_1d_3x_5y_6) \equiv d_2^2d_3d_4d_5\bar{x}_1 \pmod{d_2d_3}$$

Proof. Substituting (20) into (22) and multiplying both sides by \bar{x}_1 , we obtain (31). For (32), it follows from multiplying both sides of (24) by \bar{y}_4 .

Substituting the obtained conditions on d_3u_2 into (27) yields (33) as follows.

$$\begin{aligned} d_3u_2y_5 + d_4d_5x_6y_3 - d_3v_2 &\equiv 0 \pmod{d_4d_5} \\ d_3v_2 &\equiv d_2d_3\bar{y}_4y_5 \pmod{d_4}. \end{aligned}$$

For (34), we use the assumption $x_3y_3 \equiv 1 \pmod{d_1d_2d_3d_4d_5f}$. Substituting (21) into (29) and multiplying both side by d_2 , the term $d_2d_3u_2$ is expressed as

$$(35) \quad d_2d_4(d_5v_1 + d_1d_3x_5y_6) \equiv d_2d_3u_2 \cdot y_6u_1 + d_2d_4d_5y_3u_1 \pmod{d_2d_3}.$$

Next, we consider the conditions on $d_2d_3u_2$. Substituting (21) into (20) and applying the assumption yields the following:

$$\begin{aligned} d_2d_4d_5x_6 - d_2d_3x_3u_2 - d_2d_4d_5x_3x_6y_3 &\equiv 0 \pmod{d_2d_3} \\ d_2d_3u_2 &\equiv 0 \pmod{d_2d_3}. \end{aligned}$$

Substituting the above condition into (35), and then applying $d_2d_3u_1$, we obtain (34) as

$$\begin{aligned} d_2d_3d_4(d_5v_1 + d_1d_3x_5y_6) &\equiv d_2d_3u_1 \cdot d_4d_5y_3 \pmod{d_2d_3} \\ d_2d_3d_4(d_5v_1 + d_1d_3x_5y_6) &\equiv d_2^2d_3d_4d_5\bar{x}_1 \pmod{d_2d_3}. \end{aligned}$$

□

Theorem 3.7. *Let $\mathbf{m} = \{m_1, m_2, m_3\}$, $\mathbf{n} = \{n_1, n_2, n_3\} \in \mathbb{Z}^3$. The fine Kloosterman sum $\mathcal{S}_{\omega_0}(\mathbf{m}, \mathbf{n}; D_1, D_2, D_3, f) = \mathcal{S}_{\omega_0}(\mathbf{m}, \mathbf{n}; d_4d_5, d_2d_3d_5, d_1d_3, f)$ with the notation in (18) becomes*

$$\begin{aligned} &d_1^3d_2^2d_3^2d_4^2d_5^4f^4 \\ &\times \sum_{\substack{x_3 \pmod{d_3} \\ y_5 \pmod{d_5}}} S(m_1, (m_2d_1fx_3 + n_3d_2d_5)/d_3f; d_1) \cdot S(n_1, (n_2d_4fy_5 + m_3d_2d_3)/d_5f; d_4), \end{aligned}$$

when it satisfies $m_2 d_1 \equiv 0 \pmod{d_2 d_3}$, $m_3 \equiv 0 \pmod{d_5 f}$, $n_2 d_4 \equiv 0 \pmod{d_2 d_3 d_5}$ and $n_3 \equiv 0 \pmod{d_1 d_3 d_4 f}$.

Proof. We calculate using the definition of the fine Kloosterman sum along with the notation in (18) as follows:

$$\begin{aligned} \mathcal{S}_{\omega_0}(\mathbf{m}, \mathbf{n}; D_1, D_2, D_3, f) &= \mathcal{S}_{\omega_0}(\mathbf{m}, \mathbf{n}; d_4 d_5, d_2 d_3 d_5, d_1 d_3, f) \\ &= \sum_{\substack{A \in U(\mathbb{Z}) \setminus \Omega(d_4 d_5, d_2 d_3 d_4, d_1 d_3, f) / U(\mathbb{Z}) \\ A = u_L \omega(d_4 d_5 f, d_2 d_3 d_5 f, d_1 d_3 f) u_R}} \psi_{\mathbf{m}}(u_L) \psi_{\mathbf{n}}(u_R). \end{aligned}$$

Let us express superdiagonal elements in the unipotent factors of the Bruhat decomposition in terms of these coordinates as in Proposition 3.4. Then $\psi_{\mathbf{m}}(u_L) \psi_{\mathbf{n}}(u_R)$ becomes

$$e \left(\frac{m_1 x_1}{d_1} + \frac{m_2 u_1}{d_2 d_3} + \frac{m_3 (d_3 u_2 + d_4 d_5 x_6 y_3)}{d_4 d_5 f} + \frac{n_1 y_4}{d_4} + \frac{n_2 v_2}{d_2 d_5} + \frac{n_3 (d_5 v_1 + d_1 d_3 x_5 y_6)}{d_1 d_3 f} \right).$$

Then using Lemma 3.6, we substitute the values of u and v into the above to compute

$$\begin{aligned} \mathcal{S}_{\omega_0}(\mathbf{m}, \mathbf{n}; d_4 d_5, d_2 d_3 d_5, d_1 d_3, f) &= \sum_{\substack{x_3 \pmod{d_3} \\ y_5 \pmod{d_5}}} \sum_{\substack{x_1 \pmod{d_1} \\ x_1 \bar{x}_1 \equiv 1 \pmod{d_1}}} e \left(\frac{m_1 x_1 + \{(m_2 d_1 f x_3 + n_3 d_2 d_5) / d_3 f\} \bar{x}_1}{d_1} \right) \\ &\quad \times \sum_{\substack{y_4 \pmod{d_4} \\ y_4 \bar{y}_4 \equiv 1 \pmod{d_4}}} e \left(\frac{n_1 y_4 + \{(n_2 d_4 f y_5 + m_3 d_2 d_3) / d_5 f\} \bar{y}_4}{d_4} \right) \\ &\quad \times \sum_{k_1=0}^{d_4 d_5 f-1} e \left(\frac{m_2 d_1}{d_2 d_3} k_1 \right) \sum_{k_2=0}^{d_1 d_2 d_3 d_5 f-1} e \left(\frac{m_3}{d_5 f} k_2 \right) \\ &\quad \times \sum_{k_3=0}^{d_1 d_2 d_3 d_5 f-1} e \left(\frac{n_2 d_4}{d_2 d_3 d_5} k_3 \right) \sum_{k_4=0}^{d_1 d_4 d_5 f-1} e \left(\frac{n_3}{d_1 d_3 d_4 f} k_4 \right). \end{aligned}$$

Here, the congruence conditions on Lemma 3.6 is arranged so that the modulus is $d_1 d_2 d_3 d_4 d_5 f$. Summation over k_i gives the congruence condition is $m_2 d_1 \equiv 0 \pmod{d_2 d_3}$, $m_3 \equiv 0 \pmod{d_5 f}$, $n_2 d_4 \equiv 0 \pmod{d_2 d_3 d_5}$, $n_3 \equiv 0 \pmod{d_1 d_3 d_4 f}$, and the product of two classical Kloosterman sums as follows:

$$\begin{aligned} &d_1^3 d_2^2 d_3^2 d_4^2 d_5^4 f^4 \\ &\times \sum_{\substack{x_3 \pmod{d_3} \\ y_5 \pmod{d_5}}} S(m_1, (m_2 d_1 f x_3 + n_3 d_2 d_5) / d_3 f; d_1) \cdot S(n_1, (n_2 d_4 f y_5 + m_3 d_2 d_3) / d_5 f; d_4). \end{aligned}$$

□

From (16), using Theorem 3.7 and the Weil bound in (1), we can obtain a bound on the SL_4 -long word Kloosterman sum.

Proposition 3.8. *For given $\mathbf{m}, \mathbf{n} \in \mathbb{Z}^3$, and $c_1, c_2, c_3 > 0$, we may bound the long-word Kloosterman sum as*

$$|S_{\omega_0}(\mathbf{m}, \mathbf{n}; (c_1, c_2, c_3))| \leq c_1^3 c_2^2 c_3^2 (c_2 + 1) (m_1, c_3)^{\frac{1}{2}} (n_1, c_1)^{\frac{1}{2}} \sqrt{c_1 c_3} \tau((c_1, c_2, c_3)) \tau(c_1) \tau(c_3),$$

where $\tau(c)$ is the divisor function.

Proof. For the given decomposition of a fine Kloosterman sum as a sum of classical Kloosterman sums in Theorem 3.7, using the Weil bound in (1) on the individual sums with attention to (16), we obtain

$$|S_{\omega_0}(\mathbf{m}, \mathbf{n}; (c_1, c_2, c_3))| \leq \sum_{f|(c_1, c_2, c_3)} d_1^3 d_2^2 d_3^2 d_4^2 d_5^4 f^4 \times \sum_{\substack{x_3 \pmod{d_3} \\ y_5 \pmod{d_5}}} (m_1, d_1)^{\frac{1}{2}} (n_1, d_4)^{\frac{1}{2}} \sqrt{d_1 d_4} \tau(d_1) \tau(d_4)$$

Each d can be bounded above by c with (14) and (18), and the number of divisors f is bounded by the divisor function. □

3.3. Parametrization of other group. We now consider the case of the symplectic group $Sp(4, \mathbb{Z})$. The symplectic group is the group consisting of all $2n \times 2n$ symplectic matrices that preserve the standard symplectic form, with matrix multiplication as the group operation, and is defined as follows:

$$(36) \quad Sp(2n, \mathbb{Z}) = \{A \in GL_{2n}(\mathbb{Z}) \mid A^\top J A = J\},$$

where A^\top is the transpose of A . J is defined as

$$J = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix},$$

and I_n is the identity matrix.

For example, $Sp(2, \mathbb{Z})$ is the group of 2×2 symplectic matrices, which coincides with the special linear group $SL_2(\mathbb{Z})$. That is,

$$Sp(2, \mathbb{Z}) = SL_2(\mathbb{Z}) = \{A \in GL_2(\mathbb{Z}) \mid \det(A) = 1\}.$$

For $n \geq 2$, it is convenient to write matrices in block form. For $n = 2$ this block form is given by

$$(37) \quad A = \begin{pmatrix} P & Q \\ R & S \end{pmatrix},$$

where $P, Q, R, S \in M_2(\mathbb{Z})$, and the symplectic condition $A^\top J A = J$ with $J = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$ is equivalent to the following conditions:

$$P^\top R = R^\top P, \quad Q^\top S = S^\top Q, \quad P^\top S - R^\top Q = I_2.$$

By applying (37) to (9) we obtain the following conditions on the minors of A . In the case of $Sp(4, \mathbb{Z})$, we have following conditions:

$$\begin{aligned} M_{13,12} + M_{24,12} &= 0, & M_{13,34} + M_{24,34} &= 0, \\ M_{13,13} + M_{24,13} &= 1, & M_{13,14} + M_{24,14} &= 0, \\ M_{13,23} + M_{24,23} &= 0, & M_{13,24} + M_{24,24} &= 1. \end{aligned}$$

We now consider the orthogonal group $O(n)$. It has a subgroup consisting of all matrices with determinant 1, called the special orthogonal group, denoted by $SO(n)$. It is defined as

$$SO(n) = \{A \in GL_n(\mathbb{Z}) \mid A^\top A = I_n, \det(A) = 1\}.$$

From this definition, we can derive the conditions on the entries of a matrix A . In the case of $SO(4, \mathbb{Z})$, we have following conditions:

$$\begin{aligned} A_{1i}^2 + A_{2i}^2 + A_{3i}^2 + A_{4i}^2 &= 1, & (1 \leq i \leq 4), \\ A_{11}A_{1i} + A_{21}A_{2i} + A_{31}A_{3i} + A_{41}A_{4i} &= 0, & (2 \leq i \leq 4), \\ A_{12}A_{1i} + A_{22}A_{2i} + A_{32}A_{3i} + A_{42}A_{4i} &= 0, & (3 \leq i \leq 4), \\ A_{13}A_{14} + A_{23}A_{24} + A_{33}A_{34} + A_{43}A_{44} &= 0. \end{aligned}$$

Although both $Sp(4, \mathbb{Z})$ and $SO(4)$ have determinant 1, but additional conditions are imposed on the minors and entries of A . Therefore, the definition of the Kloosterman set and its parametrization as in (17) are expected to be more complicated than in the case of $SL_4(\mathbb{Z})$.

4. SL_5 -LONG WORD KLOOSTERMAN SUM

In this section, we outline a partial study of the fine Kloosterman sum for $SL_5(\mathbb{Z})$ following the cases of $SL_3(\mathbb{Z})$ and $SL_4(\mathbb{Z})$.

4.1. Kloosterman set and Kloosterman sum. For a vector $\mathbf{c} = \{c_1, c_2, c_3, c_4\} \in \mathbb{Z}^4$, we define

$$(38) \quad t(\mathbf{c}) = \begin{pmatrix} c_1 & & & & \\ & \frac{c_2}{c_1} & & & \\ & & \frac{c_3}{c_2} & & \\ & & & \frac{c_4}{c_3} & \\ & & & & \frac{1}{c_4} \end{pmatrix},$$

and

$$\Omega_\omega(\mathbf{c}) = \{A \in SL_5(\mathbb{Z}) \mid A = u_L \omega t(\mathbf{c}) u_R \in B\omega B, u_L, u_R \in U\}.$$

Let $\mathbf{n} = \{n_1, n_2, n_3, n_4\} \in \mathbb{Z}^4$. We use the character $\psi_{\mathbf{n}}$ to denote

$$\psi_{\mathbf{n}}(u) = e(n_1 u_{1,2} + n_2 u_{2,3} + n_3 u_{3,4} + n_4 u_{4,5}).$$

For $\mathbf{m}, \mathbf{n}, \mathbf{c} \in \mathbb{Z}^4$, we define the usual (coarse) Kloosterman sum for $SL_5(\mathbb{Z})$ as

$$S_\omega(\mathbf{m}, \mathbf{n}; \mathbf{c}) = \sum_{\substack{A \in U(\mathbb{Z}) \backslash \Omega_\omega(\mathbf{c}) / U(\mathbb{Z}) \\ A = u_L \omega t(\mathbf{c}) u_R}} \psi_{\mathbf{m}}(u_L) \psi_{\mathbf{n}}(u_R).$$

For the case of $\omega = \omega_0$ with (2), we write $A \in SL_5(\mathbb{Z}) \cap Bw_0B$ as

$$A = \begin{pmatrix} A_{11} & A_{12} & A_{13} & A_{14} & A_{15} \\ A_{21} & A_{22} & A_{23} & A_{24} & A_{25} \\ A_{31} & A_{32} & A_{33} & A_{34} & A_{35} \\ A_{41} & A_{42} & A_{43} & A_{44} & A_{45} \\ A_{51} & A_{52} & A_{53} & A_{54} & A_{55} \end{pmatrix}.$$

In addition, considering the decomposition $A \in SL_5(\mathbb{Z}) \cap Uw_0TU$ in (3), the diagonal matrix T becomes $t(\mathbf{c})$ in (38) when

$$(39) \quad c_1 = t_1, \quad c_2 = t_1 t_2, \quad c_3 = t_1 t_2 t_3, \quad c_4 = t_1 t_2 t_3 t_4.$$

From [1, Proposition 3.1], we obtain

$$(40) \quad t_1 = A_{51}, \quad t_1 t_2 = M_{45,12}, \quad t_1 t_2 t_3 = M_{345,123}, \quad t_1 t_2 t_3 t_4 = M_{2345,1234}.$$

Lemma 4.1. *For a given $A \in SL_5(\mathbb{Z}) \cap Bw_0B$ as above, we have*

$$\gcd(A_{51}, A_{52}, A_{53}, A_{54}) = \gcd(M_{2345,1234}, M_{1235,1234}, M_{1245,1234}, M_{1345,1234}).$$

We set

$$f = \gcd(A_{51}, A_{52}, A_{53}, A_{54}) = \gcd(M_{2345,1234}, M_{1235,1234}, M_{1245,1234}, M_{1345,1234}).$$

We define $D_1, D_2, D_3, D_4 \in \mathbb{Z}_{>0}$ as follows using (39) and (40),

$$(41) \quad c_1 = A_{51} = D_1 f, \quad c_2 = M_{45,12} = D_2 f, \quad c_3 = M_{345,123} = D_3 f, \quad c_4 = M_{2345,1234} = D_4 f.$$

Furthermore, we define the following set as the fine Kloosterman set

$$\Omega(D_1, D_2, D_3, D_4, f) = \{A \in SL_5(\mathbb{Z}) \mid \gcd(A_{51}, A_{52}, A_{53}, A_{54}) = f, A_{51} = D_1 f, \\ M_{45,12} = D_2 f, M_{345,123} = D_3 f, M_{2345,1234} = D_4 f\}.$$

Definition 4.2. For $\mathbf{m}, \mathbf{n} \in \mathbb{Z}^4$, we define the fine Kloosterman sum for SL_5 as follows, using D_1, D_2, D_3, D_4 and f defined above.

$$(42) \quad \mathcal{S}_{\omega_0}(\mathbf{m}, \mathbf{n}; D_1, D_2, D_3, D_4, f) = \sum_{\substack{A \in U(\mathbb{Z}) \backslash \Omega(D_1, D_2, D_3, D_4, f) / U(\mathbb{Z}) \\ A = u_L \omega_t(D_1 f, D_2 f, D_3 f, D_4 f) u_R}} \psi_{\mathbf{m}}(u_L) \psi_{\mathbf{n}}(u_R).$$

Similarly as before, the (usual) coarse Kloosterman sum for SL_5 can thus be written as the sum

$$S_{\omega_0}(\mathbf{m}, \mathbf{n}; (c_1, c_2, c_3, c_4)) = \sum_{f | (c_1, c_2, c_3, c_4)} \mathcal{S}_{\omega_0} \left(\mathbf{m}, \mathbf{n}; \frac{c_1}{f}, \frac{c_2}{f}, \frac{c_3}{f}, \frac{c_4}{f}, f \right).$$

4.2. Parametrization of fine Kloosterman cells. In this subsection, we consider the parametrization of the fine Kloosterman cells. In the case of SL_5 , for the simple roots $\alpha = (1, -1, 0, 0, 0)$, $\beta = (0, 1, -1, 0, 0)$, $\gamma = (0, 0, 1, -1, 0)$, $\delta = (0, 0, 0, 1, -1)$ and $x = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL_2$, we define

$$\begin{aligned} \iota_{\alpha}(x) &= \begin{pmatrix} a & b & & & \\ c & d & & & \\ & & 1 & & \\ & & & 1 & \\ & & & & 1 \end{pmatrix}, & \iota_{\beta}(x) &= \begin{pmatrix} 1 & & & & \\ & a & b & & \\ & c & d & & \\ & & & 1 & \\ & & & & 1 \end{pmatrix}, \\ \iota_{\gamma}(x) &= \begin{pmatrix} 1 & & & & \\ & 1 & & & \\ & & a & b & \\ & & c & d & \\ & & & & 1 \end{pmatrix}, & \iota_{\delta}(x) &= \begin{pmatrix} 1 & & & & \\ & 1 & & & \\ & & 1 & & \\ & & & a & b \\ & & & c & d \end{pmatrix}. \end{aligned}$$

Using ι_* as above, we use s_* to denote the following:

$$(43) \quad s_{\alpha} = \iota_{\alpha} \left(\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \right), \quad s_{\beta} = \iota_{\beta} \left(\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \right), \quad s_{\gamma} = \iota_{\gamma} \left(\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \right), \quad s_{\delta} = \iota_{\delta} \left(\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \right).$$

Using the reduced word decomposition $\omega_0 = s_{\alpha} s_{\beta} s_{\alpha} s_{\gamma} s_{\beta} s_{\alpha} s_{\delta} s_{\gamma} s_{\beta} s_{\alpha}$ with the notation in (43), we parametrize the fine Kloosterman sets $\Omega(D_1, D_2, D_3, D_4, f)$:

$$(44) \quad A = \iota_{\alpha}(\gamma_1) \iota_{\beta}(\gamma_3) \iota_{\alpha}(\gamma_2) \iota_{\gamma}(\gamma_6) \iota_{\beta}(\gamma_5) \iota_{\alpha}(\gamma_4) \iota_{\delta}(\gamma_{10}) \iota_{\gamma}(\gamma_9) \iota_{\beta}(\gamma_8) \iota_{\alpha}(\gamma_7),$$

where

$$\gamma_i = \begin{pmatrix} x_i & b_i \\ d_i & y_i \end{pmatrix} \quad (\text{if } 1 \leq i \leq 9), \quad \gamma_{10} = \begin{pmatrix} x_{10} & b_{10} \\ f & y_{10} \end{pmatrix}.$$

Then we can write

$$(45) \quad D_1 = d_7 d_8 d_9, \quad D_2 = d_4 d_5 d_6 d_8 d_9, \quad D_3 = d_2 d_3 d_5 d_6 d_9, \quad D_4 = d_1 d_3 d_6.$$

From [1, Proposition 3.1], for $A \in SL_5(\mathbb{Z}) \cap u_L \omega_0 T u_R$, we know that

$$u_L = \begin{pmatrix} 1 & \frac{\langle e_{1,3,4,5}^*, Ae_{1,2,3,4} \rangle}{t_1 t_2 t_3 t_4} & \frac{\langle e_{1,4,5}^*, Ae_{1,2,3} \rangle}{t_1 t_2 t_3} & \frac{\langle e_{1,5}^*, Ae_{1,2} \rangle}{t_1 t_2} & \frac{\langle e_1^*, Ae_1 \rangle}{t_1} \\ & 1 & \frac{\langle e_{2,4,5}^*, Ae_{1,2,3} \rangle}{t_1 t_2 t_3} & \frac{\langle e_{2,5}^*, Ae_{1,2} \rangle}{t_1 t_2} & \frac{\langle e_2^*, Ae_1 \rangle}{t_1} \\ & & 1 & \frac{\langle e_{3,5}^*, Ae_{1,2} \rangle}{t_1 t_2} & \frac{\langle e_3^*, Ae_1 \rangle}{t_1} \\ & & & 1 & \frac{\langle e_4^*, Ae_1 \rangle}{t_1} \\ & & & & 1 \end{pmatrix}$$

and

$$u_R = \begin{pmatrix} 1 & \frac{\langle e_5^*, Ae_2 \rangle}{t_1} & \frac{\langle e_5^*, Ae_3 \rangle}{t_1} & \frac{\langle e_5^*, Ae_4 \rangle}{t_1} & \frac{\langle e_5^*, Ae_5 \rangle}{t_1} \\ & 1 & \frac{\langle e_{4,5}^*, Ae_{1,3} \rangle}{t_1 t_2} & \frac{\langle e_{4,5}^*, Ae_{1,4} \rangle}{t_1 t_2} & \frac{\langle e_{4,5}^*, Ae_{1,5} \rangle}{t_1 t_2} \\ & & 1 & \frac{\langle e_{3,4,5}^*, Ae_{1,2,4} \rangle}{t_1 t_2 t_3} & \frac{\langle e_{3,4,5}^*, Ae_{1,2,5} \rangle}{t_1 t_2 t_3} \\ & & & 1 & \frac{\langle e_{2,3,4,5}^*, Ae_{1,2,3,5} \rangle}{t_1 t_2 t_3 t_4} \\ & & & & 1 \end{pmatrix}.$$

Here, we define

$$u_1 = d_1 x_2 + d_2 x_3 y_1, \quad u_2 = d_2 x_4 + d_4 x_5 y_2, \quad u_3 = d_4 x_7 + d_7 x_8 y_4, \quad u_4 = d_6 x_9 y_5 + d_9 x_{10} y_6,$$

$$v_1 = d_1 x_2 y_3 + d_2 y_1, \quad v_2 = d_2 x_4 y_5 + d_4 y_2, \quad v_3 = d_4 x_7 y_8 + d_7 y_4, \quad v_4 = d_6 x_9 y_{10} + d_9 x_5 y_6.$$

Computing the minors and the entries of A using (44), for d_i ($1 \leq i \leq 9$), f as in (41) and (45), the Bruhat decomposition of $A \in SL_5(\mathbb{Z}) \cap u_L \omega_0 T u_R$ can be written as

$$T = \begin{pmatrix} d_7 d_8 d_9 f & & & & \\ & \frac{d_4 d_5 d_6}{d_7} & & & \\ & & \frac{d_2 d_3}{d_4 d_8} & & \\ & & & \frac{d_1}{d_2 d_5 d_9} & \\ & & & & \frac{1}{d_1 d_3 d_6 f} \end{pmatrix},$$

$$(46) \quad u_L = \begin{pmatrix} 1 & \frac{x_1}{d_1} & \frac{x_1 u_1 - d_2 x_3}{d_1 d_2 d_3} & \frac{\langle e_{1,5}^*, Ae_{1,2} \rangle}{d_4 d_5 d_6 d_8 d_9 f} & \frac{\langle e_1^*, Ae_1 \rangle}{d_7 d_8 d_9 f} \\ & 1 & \frac{u_1}{d_2 d_3} & \frac{\langle e_{2,5}^*, Ae_{1,2} \rangle}{d_4 d_5 d_6 d_8 d_9 f} & \frac{\langle e_2^*, Ae_1 \rangle}{d_7 d_8 d_9 f} \\ & & 1 & \frac{d_3 u_2 + d_4 d_5 x_6 y_3}{d_4 d_5 x_6} & \frac{\langle e_3^*, Ae_1 \rangle}{d_7 d_8 d_9 f} \\ & & & 1 & \frac{d_5 d_6 u_3 + d_7 d_8 u_4}{d_7 d_8 d_9 f} \\ & & & & 1 \end{pmatrix}$$

and

$$(47) \quad u_R = \begin{pmatrix} 1 & \frac{y_7}{d_7} & \frac{y_8}{d_7 d_8} & \frac{y_9}{d_7 d_8 d_9} & \frac{y_{10}}{d_7 d_8 d_9 f} \\ & 1 & \frac{v_3}{d_4 d_8} & \frac{d_5 y_9 u_3 + d_7 d_8 y_5}{d_4 d_5 d_8 d_9} & \frac{\langle e_{4,5}^*, Ae_{1,5} \rangle}{d_4 d_5 d_6 d_8 d_9 f} \\ & & 1 & \frac{d_8 v_2 + d_2 d_5 x_8 y_9}{d_2 d_5 d_9} & \frac{\langle e_{3,4,5}^*, Ae_{1,2,5} \rangle}{d_2 d_3 d_5 d_6 d_9 f} \\ & & & 1 & \frac{d_5 d_9 v_1 + d_1 d_3 v_4}{d_1 d_3 d_6 f} \\ & & & & 1 \end{pmatrix}.$$

Now we consider the fine Kloosterman sum

$$\mathcal{S}_{\omega_0}(\mathbf{m}, \mathbf{n}; D_1, D_2, D_3, D_4 f) = \mathcal{S}_{\omega_0}(\mathbf{m}, \mathbf{n}; d_7 d_8 d_9, d_4 d_5 d_6 d_8 d_9, d_2 d_3 d_5 d_6 d_9, d_1 d_3 d_6, f)$$

with the notation in (45). From (46) and (47), the superdiagonal elements in the unipotent factors of the Bruhat decomposition $\psi_{\mathbf{m}}(u_L)\psi_{\mathbf{n}}(u_R)$ in (42) becomes

$$e \left(\frac{m_1 x_1}{d_1} + \frac{m_2 u_1}{d_2 d_3} + \frac{m_3 (d_3 u_2 + d_4 d_5 x_6 y_3)}{d_4 d_5 d_6} + \frac{m_4 (d_5 d_6 u_3 + d_7 d_8 u_4)}{d_7 d_8 d_9 f} \right. \\ \left. + \frac{n_1 y_7}{d_7} + \frac{n_2 v_3}{d_4 d_8} + \frac{n_3 (d_8 v_2 + d_2 d_5 x_8 y_9)}{d_2 d_5 d_9} + \frac{n_4 (d_5 d_9 v_1 + d_1 d_3 v_4)}{d_1 d_3 d_6 f} \right).$$

ACKNOWLEDGMENT

We are also sincerely grateful to Professor Eren Mehmet Kiral for his helpful comment and careful reading.

REFERENCES

- [1] E. M. Kiral and M. Nakasuji. Parametrization of Kloosterman sets and SL_3 -Kloosterman sums, *Advances in Mathematics* 403 (2022) 108392.
- [2] H. Iwaniec. *Topics in Classical Automorphic Forms (Graduate Studies in Mathematics, volume 17)*, American Mathematical Society, (1997).

- [3] D. Bump. *Automorphic Forms on $GL(3, \mathbb{R})$* , Lecture Notes in Mathematics, Springer-Verlag, Berlin Heidelberg New York Tokyo (1984).

SUZUHO OSONOE

Graduate School of Science and Technology

Sophia University,

7-1 Kioi-cho, Chiyoda-ku, Tokyo, 102-8554, Japan

`s-osonoe-7m6@eagle.sophia.ac.jp`

MAKI NAKASUJI

Department of Information and Communication Science, Faculty of Science,

Sophia University,

7-1 Kioi-cho, Chiyoda-ku, Tokyo, 102-8554, Japan

`nakasuji@sophia.ac.jp`

and

Mathematical Institute,

Tohoku University,

6-3 Aoba, Aramaki, Aoba-ku, Sendai, 980-8578, Japan