

SOME TOPOLOGICAL PROPERTIES OF THE INTRINSIC VOLUME METRIC

STEVEN HOEHNER

ABSTRACT. We prove that for fixed $j \in \{2, \dots, d-1\}$, the j th intrinsic volume metric δ_j is not uniformly bounded, and the metric space $(\mathcal{K}_j^d, \delta_j)$ is incomplete. In particular, this proves that for $j \in \{2, \dots, d-1\}$, the Hausdorff and j th intrinsic volume metrics do not induce the same topology on \mathcal{K}_j^d .

1. INTRODUCTION AND MAIN RESULTS

Fix $d \in \mathbb{N}$. Let \mathcal{K}^d denote the set of all convex, compact sets in \mathbb{R}^d . For $K \in \mathcal{K}^d$, we let $\dim(K)$ denote the dimension of the affine hull of K . For $j \in \{1, \dots, d\}$, we set $\mathcal{K}_j^d := \{K \in \mathcal{K}^d : \dim(K) \geq j\}$. For $K, L \in \mathcal{K}^d$, the Hausdorff distance $d_H(K, L)$ may be defined by

$$d_H(K, L) = \inf \{ \lambda \geq 0 : K \subset L + \lambda B_d, L \subset K + \lambda B_d \},$$

where B_d is the d -dimensional Euclidean unit ball centered at the origin. The symmetric difference metric is defined by $d_S(K, L) = \text{vol}_d(K \triangle L)$, where vol_d denotes the d -dimensional volume functional and $K \triangle L = (K \setminus L) \cup (L \setminus K)$. It is well-known that the metric spaces (\mathcal{K}^d, d_H) and $(\mathcal{K}_d^d \cup \{\emptyset\}, d_S)$ are complete. As explained in [1], in the latter case the empty set is appended to include limits of sequences of convex bodies $\{K_i\} \subset \mathcal{K}_d^d$ which converge in the Hausdorff metric to a set $K_0 \in \mathcal{K}^d$ with $\dim(K_0) < d$. More specifically, if $K_i \rightarrow K_0 \in \mathcal{K}^d$ and $\dim(K_0) < d$, then for such a sequence we have $d_S(K_i, \emptyset) = \text{vol}_d(K_i) \rightarrow 0$, so $K_i \rightarrow \emptyset$ with respect to d_S . It is also known that on uniformly bounded subsets of \mathcal{K}_d^d , the metrics d_H and d_S are equivalent. For more background on the Hausdorff and symmetric difference metrics, see the article [6] by Shephard and Webster.

For $j \in \{1, \dots, d\}$ and $K, L \in \mathcal{K}_j^d$, the j th *intrinsic volume metric* $\delta_j(K, L)$ was defined in [1] by

$$(1) \quad \delta_j(K, L) := \begin{bmatrix} d \\ j \end{bmatrix} \int_{\text{Gr}(d, j)} \text{vol}_j((P_H K) \triangle (P_H L)) d\nu_j(H).$$

Date: December 3, 2025.

2020 Mathematics Subject Classification: 52A27 (52A20, 52A39)

Key words and phrases: convex body, intrinsic volume, metric, quermassintegral

Here and throughout the paper, $\text{Gr}(d, j)$ is the Grassmannian manifold of all j -dimensional subspaces of \mathbb{R}^d , ν_j is the unique Haar probability measure on $\text{Gr}(d, j)$, P_H denotes the orthogonal projection of \mathbb{R}^d into the subspace $H \in \text{Gr}(d, j)$, and $\begin{bmatrix} d \\ j \end{bmatrix} = \binom{d}{j} \frac{\text{vol}_d(B_d)}{\text{vol}_j(B_j) \text{vol}_{d-j}(B_{d-j})}$ is the flag coefficient of Klain and Rota [5]. The intrinsic volume metric has found natural applications to the approximation of convex bodies by polytopes; see [1, 2, 4].

It was conjectured in [1] that d_H and δ_j induce the same topology on \mathcal{K}_j^d . In Open Question 2.2 of [1], they asked the following

Open Question 1.1 (uniform bound). *Let $j \in \{2, \dots, d-1\}$. If $\{K_i\}$ is a sequence in \mathcal{K}_j^d that converges to $K \in \mathcal{K}_j^d$ with respect to δ_j , does it follow that $\{K_i\}$ is uniformly bounded, that is, does there exist $R > 0$ such that $\bigcup_{i \in \mathbb{N}} K_i \subset RB_d$?*

As remarked in [1], a positive answer to this question would imply that a sequence $\{K_i\}$ which converges with respect to δ_j also converges with respect to d_H . Moreover, as noted in [1], for the special case $j = d$, this question was answered affirmatively by Shephard and Webster in [6, Lemma 11], and for $j = 1$ see [7, Theorem 3] and [3, Theorem 2].

We show that, interestingly, when $2 \leq j \leq d-1$, the answers to these questions are negative.

Theorem 1.2. *Let $2 \leq j \leq d$ be given. There exists a sequence $\{K_i\} \subset \mathcal{K}_j^d$ such that*

$$\delta_j(K_i, K) \rightarrow 0 \quad \text{but} \quad \sup_{x \in K_i} \|x\| \rightarrow \infty \quad \text{as} \quad i \rightarrow \infty.$$

Consequently, d_H and δ_j induce different topologies on \mathcal{K}_j^d if $2 \leq j \leq d$.

Remark 1.3. For $j = d$, note that $\delta_d = d_S$ is the symmetric difference metric. The nonequivalence in Theorem 1.2 is on \mathcal{K}_d^d without a uniform boundedness restriction; on uniformly bounded families, the topologies induced by d_H and d_S do coincide.

Theorem 1.4. *Let $d \geq 3$ and $2 \leq j \leq d-1$. The metric space $(\mathcal{K}_j^d, \delta_j)$ is incomplete.*

It is thus natural to ask if appending the empty set to \mathcal{K}_j^d will make $(\mathcal{K}_j^d \cup \{\emptyset\}, \delta_j)$ complete. While this is true for $j \in \{1, d\}$ (see [3, 7] for the case $j = 1$), as we show below, it is still false for intermediate $j \in \{2, \dots, d-1\}$.

Theorem 1.5. *Let $d \geq 3$ and $2 \leq j \leq d-1$. The metric space $(\mathcal{K}_j^d \cup \{\emptyset\}, \delta_j)$ is incomplete.*

Throughout the paper, we use c, C, C_1, C_2 , etc. to denote positive absolute constants whose value may change from line to line. If a positive constant depends on some parameter, the dependence will always be made explicit, e.g., $c(d)$ is a constant that depends only on d .

2. PROOFS

2.1. Proof of Theorem 1.2. Let $E := \text{span}\{e_1, \dots, e_j\} \subset \mathbb{R}^d$ and fix a j -dimensional convex body $K \subset E$ (so $K \in \mathcal{K}_j^d$). Choose $x_0 \in \text{relint}(K)$ (relative to E) and a unit vector $u \in E$.

Shrinking ε if necessary, we may assume that $x_0 + \varepsilon B_{u^\perp \cap E}^{j-1} \subset K$. For parameters $L > 0$ and $\varepsilon > 0$, define the following thin cylinder inside E :

$$(2) \quad N(L, \varepsilon) := \{x_0 + tu + v : t \in [0, L], v \in \varepsilon B_{u^\perp \cap E}^{j-1}\} \subset E.$$

Here $B_{u^\perp \cap E}^{j-1} := \{x \in u^\perp \cap E : \|x\| \leq 1\}$. Now set

$$(3) \quad K(L, \varepsilon) := \text{conv}(K \cup N(L, \varepsilon)) \subset E.$$

Note that $\dim K(L, \varepsilon) = j$ and $K(L, \varepsilon) \in \mathcal{K}_j^d$ for all $L, \varepsilon > 0$.

Fix $H \in \text{Gr}(d, j)$. Let $u_H := (P_H u) / \|P_H u\|$ when $P_H u \neq 0$, and decompose H as the direct sum $H = \mathbb{R}u_H \oplus E_H$ where $E_H := (P_H u)^\perp \cap H$. Let π_H denote the orthogonal projection along u_H . Then for every $y \in E_H$, the fiber $(P_H(K(L, \varepsilon)))_y \setminus (P_H K)_y$ is either empty or is a single segment of length at most $L\|P_H u\|$, and this can only occur when $y \in \pi_H(P_H(\varepsilon B_{u^\perp \cap E}^{j-1}))$. Hence, with $\ell_H := \|P_H u\|$ we get

$$\text{vol}_j([P_H(K(L, \varepsilon))] \triangle (P_H K)) \leq L \ell_H \text{vol}_{j-1}(\pi_H(P_H(\varepsilon B_{u^\perp \cap E}^{j-1}))).$$

Therefore, integrating over H we obtain

$$\delta_j(K(L, \varepsilon), K) \leq \begin{bmatrix} d \\ j \end{bmatrix} \int_{\text{Gr}(d, j)} L \ell_H \text{vol}_{j-1}(P_H(\varepsilon B_{u^\perp \cap E}^{j-1})) d\nu_j(H).$$

The orthogonal projection is 1-Lipschitz on H and thus does not increase the $(j-1)$ -dimensional volume. Hence,

$$\text{vol}_{j-1}(P_H(\varepsilon B_{u^\perp \cap E}^{j-1})) \leq \text{vol}_{j-1}(\varepsilon B_{u^\perp \cap E}^{j-1}) = \text{vol}_{j-1}(B_{u^\perp \cap E}^{j-1}) \varepsilon^{j-1} = \text{vol}_{j-1}(B_{j-1}) \varepsilon^{j-1}.$$

Thus, using also $\ell_H \leq 1$, we get

$$\delta_j(K(L, \varepsilon), K) \leq L \begin{bmatrix} d \\ j \end{bmatrix} \text{vol}_{j-1}(B_{j-1}) \varepsilon^{j-1} = C(d, j) L \varepsilon^{j-1}$$

where $C(d, j) := \begin{bmatrix} d \\ j \end{bmatrix} \text{vol}_{j-1}(B_{j-1})$.

Now choose a sequence $\{L_i\}$ with $L_i \rightarrow \infty$, and set $\varepsilon_i := L_i^{-2}$. For any fixed $j \geq 2$, we have

$$\delta_j(K(L_i, \varepsilon_i), K) \leq C(d, j) L_i \varepsilon_i^{j-1} = C(d, j) L_i^{1-2(j-1)} = C(d, j) L_i^{3-2j} \rightarrow 0 \quad \text{as } i \rightarrow \infty.$$

and $\sup_{x \in K(L_i, \varepsilon_i)} \|x\| \rightarrow \infty$ as $i \rightarrow \infty$. Thus, we have proved the first assertion of the theorem with $K_i := K(L_i, L_i^{-2}) \in \mathcal{K}_j^d$.

To prove the second assertion, consider the point $x_i := x_0 + L_i u \in N(L_i, \varepsilon_i) \subset K(L_i, \varepsilon_i)$. Let

$$R_K := \sup\{\|x\| : x \in K\} < \infty.$$

Observe that by the reverse triangle inequality,

$$\|x_i\| = \|x_0 + L_i u\| \geq \|L_i u\| - \|x_0\| = L_i - \|x_0\|,$$

and, for any $x \in K$,

$$\|x_i - x\| \geq \|x_i\| - \|x\| \geq L_i - \|x_0\| - R_K.$$

Therefore,

$$d_H(K(L_i, \varepsilon_i), K) \geq \text{dist}(x_i, K) \geq L_i - \|x_0\| - R_K \longrightarrow \infty \quad \text{as } i \rightarrow \infty.$$

□

2.2. Proof of Theorem 1.4. Fix a j -dimensional subspace $E \in \text{Gr}(d, j)$ and a unit vector $u \in E$. Given $x \in E$, $L > 0$ and $\varepsilon > 0$, define the following thin cylinder (a j -dimensional convex body inside E):

$$N(x; L, \varepsilon) := \text{conv}(x + [-L, L]u, x + \varepsilon B_{u^\perp \cap E}^{j-1}) \subset E,$$

where $B_{u^\perp \cap E}^{j-1}$ is the Euclidean unit ball in $u^\perp \cap E$. For any $K \in \mathcal{K}_j^d$, set

$$K^+(x; L, \varepsilon) := \text{conv}(K \cup N(x; L, \varepsilon)).$$

There exists a constant $C(d, j) > 0$ (depending only on d, j) such that for all K , all $x \in E$, and all $L, \varepsilon > 0$,

$$(4) \quad \delta_j(K^+(x; L, \varepsilon), K) \leq C(d, j)L\varepsilon^{j-1}.$$

Indeed, for each $H \in \text{Gr}(d, j)$, the j -dimensional volume in H contributed by the added cylinder is bounded above by

$$\begin{aligned} & (\text{length of } P_H\text{-image of the segment}) \times ((j-1)\text{-volume of the transverse cross-section}) \\ & \leq 2L \text{vol}_{j-1}(B_{j-1})\varepsilon^{j-1}, \end{aligned}$$

and we integrate this bound over H (the projection is 1-Lipschitz on E). Multiplying both sides of this inequality by the fixed factor $\begin{bmatrix} d \\ j \end{bmatrix}$ and setting $C(d, j) := 2 \begin{bmatrix} d \\ j \end{bmatrix} \text{vol}_{j-1}(B_{j-1})$, we obtain (4).

Next, we construct a δ_j -Cauchy sequence whose projections escape to infinity. Fix any $K_0 \in \mathcal{K}_j^d$ and let E and u be as above. Choose a sequence $\{L_m\}_{m \geq 0}$ with $L_m \rightarrow \infty$. Define $\varepsilon_m > 0$ by

$$(5) \quad \varepsilon_m^{j-1} L_m = \frac{2^{-(m+1)}}{C(d, j)}.$$

We inductively define

$$K_{m+1} := \text{conv}(K_m \cup N(x_m; L_m, \varepsilon_m)),$$

where $x_m := x_0 + T_m u$ and $T_m \rightarrow \infty$ is arbitrary (e.g., $T_m = m$). Then for a.e. H with $\|P_H u\| > 0$, we have $\|P_H x_m\| > \|P_H u\| T_m - \|P_H x_0\| \rightarrow \infty$ as $m \rightarrow \infty$. Hence, for any prescribed $R_m \rightarrow \infty$, we eventually have $P_H x_m \in H \setminus B_H(0, R_m)$. In particular, the projected cylinder $P_H N(x_m; L_m, \varepsilon_m)$ lies outside $B_H(0, R_m)$ for all sufficiently large m . By (4) and (5), we have

$$(6) \quad \delta_j(K_{m+1}, K_m) \leq C(d, j)L_m \varepsilon_m^{j-1} = 2^{-(m+1)}.$$

Hence $\sum_m \delta_j(K_{m+1}, K_m) < \infty$, and $\{K_m\}$ is δ_j -Cauchy.

Suppose by way of contradiction that $\{K_m\}$ converges in the metric δ_j to some $K_\infty \in \mathcal{K}_j^d$:

$$\delta_j(K_m, K_\infty) = \left[\begin{matrix} d \\ j \end{matrix} \right] \int_{\text{Gr}(d,j)} \text{vol}_j((P_H K_m) \triangle (P_H K_\infty)) d\nu_j(H) \longrightarrow 0 \quad \text{as } m \rightarrow \infty.$$

Then with $f_m(H) := \text{vol}_j((P_H K_m) \triangle (P_H K_\infty))$, this implies $f_m \rightarrow 0$ in $L^1(H)$. By the definition of δ_j , since $f_m \rightarrow 0$ in $L^1(H)$ there exists a subsequence (still denoted $\{f_m\}$) such that for ν_j -almost every $H \in \text{Gr}(d, j)$,

$$(7) \quad f_m(H) = \text{vol}_j((P_H K_m) \triangle (P_H K_\infty)) \longrightarrow 0 \quad \text{as } m \rightarrow \infty.$$

Next, we show that for a.e. H , each new cylinder contributes a fixed positive j -dimensional volume outside $B_H(0, R_m) = \{y \in H : \|y\| \leq R_m\}$. We will need the following lemma, proved below.

Lemma 2.1. *Fix $E \in \text{Gr}(d, j)$ and a unit vector $u \in E$. For ν_j -almost every $H \in \text{Gr}(d, j)$, the map $P_H|_E : E \rightarrow H$ has full rank j , and the linear map*

$$T_H := (\pi_H \circ P_H)|_{u^\perp \cap E} : u^\perp \cap E \longrightarrow E_H,$$

where $E_H := \{x \in H : \langle x, u_H \rangle = 0\}$ and $u_H := \frac{P_H u}{\|P_H u\|}$, is an isomorphism with $(j-1)$ -Jacobian $J_{j-1}(T_H) > 0$.

Fix an H that satisfies Lemma 2.1, and set

$$\ell_H := \|P_H u\| > 0, \quad b(H) := J_{j-1}(T_H) \text{vol}_{j-1}(B_{j-1}) > 0, \quad c(H) := 2\ell_H b(H) > 0.$$

Since P_H is linear and T_H is an isomorphism, the H -projection of the cylinder contains a rectangular block,

$$P_H N(x_m; L_m, \varepsilon_m) \supset \left(P_H x_m + [-L_m, L_m] \cdot P_H u \right) \times \pi_H \left(P_H (\varepsilon_m B_{u^\perp \cap E}^{j-1}) \right),$$

whose j -dimensional volume is at least

$$(8) \quad \text{vol}_j(P_H N(x_m; L_m, \varepsilon_m)) \geq 2L_m \|P_H u\| \varepsilon_m^{j-1} J_{j-1}(T_H) \text{vol}_{j-1}(B_{j-1}) = c(H) \varepsilon_m^{j-1} L_m.$$

By (5), the right-hand side equals $c(H)C(d, j)^{-1}2^{-(m+1)}$.

Since $P_H K_m \subset B_H(0, \rho_m)$ for some $\rho_m < \infty$, choose $R_m > \rho_m$. Thus, for all large enough m , $P_H N(x_m; L_m, \varepsilon_m) \subset H \setminus B_H(0, R_m)$ and hence is disjoint from $P_H K_m$. Thus, since $P_H K_{m+1} = \text{conv}(P_H K_m \cup P_H N(x_m; L_m, \varepsilon_m))$, and since we choose R_m so that $P_H K_m \subset B_H(0, \rho_m) \subset B_H(0, R_m)$, for ν_j -a.e. $H \in \text{Gr}(d, j)$, there exists $m_0 \in \mathbb{N}$ such that for all $m \geq m_0$,

$$(P_H K_{m+1} \setminus P_H K_m) \cap (H \setminus B_H(0, R_m)) \supset P_H N(x_m; L_m, \varepsilon_m).$$

and

$$(9) \quad \begin{aligned} \text{vol}_j \left((P_H K_{m+1} \setminus P_H K_m) \cap (H \setminus B_H(0, R_m)) \right) &\geq \text{vol}_j(P_H N(x_m; L_m, \varepsilon_m)) \\ &\geq c(H)C(d, j)^{-1}2^{-(m+1)}. \end{aligned}$$

In the final step, we will obtain a contradiction using $L^1(H)$ -convergence to a bounded projection. Fix the above H . The sets $P_H K_\infty$ are bounded in H , so there exists $R > 0$ such

that $P_H K_\infty \subset B_H(0, R)$. From (7), for the subsequence (which we still denote by $\{K_m\}$), we have

$$\text{vol}_j((P_H K_m) \triangle (P_H K_\infty)) \longrightarrow 0 \quad \text{as } m \rightarrow \infty.$$

In particular, for this fixed R , $\text{vol}_j((P_H K_m) \setminus B_H(0, R)) \rightarrow 0$ as $m \rightarrow \infty$. But $R_m \rightarrow \infty$, so eventually $R_m > R$, and then (9) yields a uniform positive lower bound

$$\text{vol}_j((P_H K_{m+1}) \setminus B_H(0, R)) \geq c(H)C(d, j)^{-1}2^{-(m+1)} > 0$$

for infinitely many m . This contradicts the fact that $\text{vol}_j((P_H K_m) \setminus B_H(0, R)) \rightarrow 0$ as $m \rightarrow \infty$. Therefore, no $K_\infty \in \mathcal{K}_j^d$ can be the δ_j -limit of $\{K_m\}$, so $(\mathcal{K}_j^d, \delta_j)$ is incomplete.

Proof of Lemma 2.1. We first show the statement on the full rank of $P_H|_E$. For $v \in E \setminus \{0\}$, the set $\mathcal{A}_v := \{H \in \text{Gr}(d, j) : H \subset v^\perp\}$ identifies with $\text{Gr}(d-1, j)$, a proper smooth submanifold of strictly lower dimension. Hence, $\nu_j(\mathcal{A}_v) = 0$. Let $\mathbb{S}(E) := \{v \in E : \|v\| = 1\}$ denote the unit sphere in E centered at the origin. Since $\mathbb{S}(E)$ is compact, it contains a countable dense subset $Q \subset \mathbb{S}(E)$. For $q \in Q$, define

$$\mathcal{A}_q := \{H \in \text{Gr}(d, j) : H \subset q^\perp\}.$$

Each \mathcal{A}_q is naturally identified with $\text{Gr}(d-1, j)$, so $\dim \mathcal{A}_q = j(d-j-1) < j(d-j) = \dim \text{Gr}(d, j)$, and therefore $\nu_j(\mathcal{A}_q) = 0$. If $H \in \text{Gr}(d, j)$ is such that $H \subset v^\perp$ for some nonzero $v \in E$, then $v/\|v\| \in \mathbb{S}(E) \cap H^\perp$, a nonempty closed subset of $\mathbb{S}(E)$. Since Q is dense in $\mathbb{S}(E)$, there exists $q \in Q \cap H^\perp$. Thus $H \subset q^\perp$, so $H \in \mathcal{A}_q$. This yields

$$\{H \in \text{Gr}(d, j) : \exists v \in E \setminus \{0\} \text{ with } H \subset v^\perp\} \subset \bigcup_{q \in Q} \mathcal{A}_q,$$

which is a countable union of ν_j -null sets and thus is a ν_j -null set. Thus, for ν_j -a.e. H , $E \cap H^\perp = \{0\}$ and $P_H|_E$ is bijective.

Next, we show that T_H is $(j-1)$ -isomorphism on $u^\perp \cap E$. Fix any H in which $P_H|_E$ has full rank, so $P_H u \neq 0$ and u_H is defined. By the first part, this set has ν_j measure 1. If $T_H w = 0$ for some $w \in u^\perp \cap E$, then $P_H w \in \text{span}\{u_H\}$, say $P_H w = \lambda u_H = \lambda P_H u / \|P_H u\|$ for some $\lambda \in \mathbb{R}$. Then $P_H(w - \beta u) = 0$ where $\beta = \lambda / \|P_H u\|$, and since $w - \beta u \in E$ and $P_H|_E$ is injective, we get $w - \beta u = 0$. Hence $0 = \langle u, w \rangle = \beta \langle u, u \rangle = \beta$, so $w = 0$. Therefore, T_H is injective. Since $\dim(u^\perp \cap E) = \dim(E_H) = j-1$, T_H is an isomorphism, and its $(j-1)$ -Jacobian is positive. \square

2.3. Proof of Theorem 1.5. Fix a j -dimensional subspace $E \in \text{Gr}(d, j)$ with $2 \leq j \leq d-1$, a unit vector $u \in E$, and a nonempty compact convex set $K_0 \subset E$ with $\dim K_0 = j$. For $L > 0$, $\varepsilon > 0$ and a base point $x_0 \in K_0$, define the thin cylinder

$$N(x_0; L, \varepsilon) := \text{conv}(x_0 + [-L, L]u, x_0 + \varepsilon B_{u^\perp \cap E}^{j-1}),$$

and set $K^+(x_0; L, \varepsilon) := \text{conv}(K_0 \cup N(x_0; L, \varepsilon))$. As in the proof of Theorem 1.4, for all $L, \varepsilon > 0$,

$$(10) \quad \delta_j(K^+(x_0; L, \varepsilon), K_0) \leq C(d, j)L\varepsilon^{j-1}.$$

where $C(d, j) := 2 \left[\frac{d}{j} \right] \text{vol}_{j-1}(B_{j-1}) > 0$.

We will construct a δ_j -Cauchy sequence with uniformly positive distance to \emptyset . Let $a_0 := \delta_j(K_0, \emptyset)$, which is positive since $K_0 \neq \emptyset$. Choose any sequence $L_m \rightarrow \infty$, and define

$$(11) \quad \varepsilon_m := \left(\frac{2^{-(m+1)} a_0}{4C(d, j)L_m} \right)^{\frac{1}{j-1}}.$$

Define inductively $K_{m+1} := \text{conv}(K_m \cup N(x_m; L_m, \varepsilon_m))$, where the choice of the base point $x_m \in E$ will be specified later. Then by (10) and (11),

$$(12) \quad \delta_j(K_{m+1}, K_m) \leq C(d, j)L_m \varepsilon_m^{j-1} = \frac{a_0}{4} \cdot 2^{-(m+1)}.$$

Hence, $\sum_m \delta_j(K_{m+1}, K_m) \leq a_0/4$, so $\{K_m\}$ is δ_j -Cauchy. Moreover, by the triangle inequality,

$$|\delta_j(K_{m+1}, \emptyset) - \delta_j(K_m, \emptyset)| \leq \delta_j(K_{m+1}, K_m) \leq \frac{a_0}{4} \cdot 2^{-(m+1)},$$

so $\{\delta_j(K_m, \emptyset)\}$ is Cauchy in \mathbb{R} and thus converges to some $\alpha \in [0, \infty)$. We now show that $\alpha > 0$. Note that for every m ,

$$\delta_j(K_m, \emptyset) \geq \delta_j(K_0, \emptyset) - \delta_j(K_0, K_m) \geq a_0 - \sum_{i=0}^{m-1} \delta_j(K_{i+1}, K_i) \geq a_0 - \frac{a_0}{4} = \frac{3}{4}a_0 > 0.$$

Passing to the limit, we thus obtain

$$(13) \quad \alpha = \lim_{m \rightarrow \infty} \delta_j(K_m, \emptyset) \geq \frac{3}{4}a_0 > 0.$$

Therefore, $\{K_m\}$ does not converge to \emptyset in $(\mathcal{K}_j^d \cup \{\emptyset\}, \delta_j)$.

Let $\mathcal{G} \subset \text{Gr}(d, j)$ denote the set of “good” subspaces in Lemma 2.1. Then by Lemma 2.1, $\nu_j(\mathcal{G}) = 1$. For fixed $H \in \mathcal{G}$, define

$$\ell_H := \|P_H u\| > 0, \quad b(H) := J_{j-1}(T_H) \text{vol}_{j-1}(B_{j-1}) > 0, \quad c(H) := 2\ell_H b(H) > 0.$$

We now choose the base points so that projected mass appears arbitrarily far from the origin. Let $\{R_m\}_{m \geq 0}$ be any sequence with $R_m \rightarrow \infty$. Since $P_H|_E$ is onto, for each m we can choose the base point $x_m \in E$ so that $P_H x_m$ lies sufficiently far in the $+u_H$ -direction to guarantee

$$(14) \quad P_H N(x_m; L_m, \varepsilon_m) \subset H \setminus B_H(0, R_m).$$

Since the projection is linear and T_H is an isomorphism, the projected cylinder contains a rectangular block in H whose j -volume is bounded below by

$$(15) \quad \text{vol}_j(P_H N(x_m; L_m, \varepsilon_m)) \geq 2L_m \|P_H u\| \varepsilon_m^{j-1} J_{j-1}(T_H) \text{vol}_{j-1}(B_{j-1}) = c(H) \varepsilon_m^{j-1} L_m.$$

By (11), the right-hand side equals $\frac{c(H)a_0}{4C(d, j)} \cdot 2^{-(m+1)}$. Moreover, since $P_H K_m$ is contained in some ball $B_H(0, \rho_m)$ with $\rho_m < \infty$ (as by continuity, projections of compact sets are compact),

choosing $R_m > \rho_m$ makes $P_H N(x_m; L_m, \varepsilon_m)$ disjoint from $P_H K_m$, and from (14) and (15) we get the lower bound

$$(16) \quad \text{vol}_j \left((P_H K_{m+1} \setminus P_H K_m) \cap (H \setminus B_H(0, R_m)) \right) \geq \frac{c(H)a_0}{4C(d, j)} \cdot 2^{-(m+1)} > 0.$$

Finally, we show that the sequence $\{K_m\}$ has no limit in \mathcal{K}_j^d . Suppose by way of contradiction that $K_m \rightarrow K_\infty$ in δ_j for some $K_\infty \in \mathcal{K}_j^d$. Following along the same lines as before, there exists a subsequence (again, not relabeled) such that for ν_j -a.e. H ,

$$(17) \quad \text{vol}_j \left((P_H K_m) \triangle (P_H K_\infty) \right) \longrightarrow 0 \quad \text{as } m \rightarrow \infty.$$

Fix the good $H \in \mathcal{G}$ chosen above, so that (17) holds for these H . Since K_∞ is compact, $P_H K_\infty \subset B_H(0, R)$ for some $R > 0$. Since $R_m \rightarrow \infty$, for all sufficiently large m we have $R_m > R$, and hence

$$\text{vol}_j \left((P_H K_m) \setminus B_H(0, R) \right) \longrightarrow 0.$$

But (16) yields, for infinitely many m ,

$$\text{vol}_j \left((P_H K_{m+1}) \setminus B_H(0, R) \right) \geq \frac{c(H)a_0}{4C(d, j)} \cdot 2^{-(m+1)} > 0,$$

a contradiction. Therefore, $\{K_m\}$ has no limit in \mathcal{K}_j^d .

To conclude the proof, note that by (13), the sequence does not converge to \emptyset , and by the last step, it does not converge to any $K_\infty \in \mathcal{K}_j^d$ either. Hence, $(\mathcal{K}_j^d \cup \{\emptyset\}, \delta_j)$ is not complete. \square

REFERENCES

- [1] F. Besau and S. Hoehner. An intrinsic volume metric for the class of convex bodies in \mathbb{R}^n . *Communications in Contemporary Mathematics*, 26(3):2350006, 2024. 1, 2
- [2] F. Besau, S. Hoehner, and G. Kur. Intrinsic and dual volume deviations of convex bodies and polytopes. *International Mathematics Research Notices*, 22:17456–17513, 2021. 2
- [3] A. Florian. On a metric for the class of compact convex sets. *Geometriae Dedicata*, 30:69–80, 1989. 2
- [4] S. Hoehner, C. Schütt, and E. Werner. Approximation of the Euclidean ball by polytopes with a restricted number of k -faces. *arXiv:2510.22771*, 2025. 2
- [5] D. A. Klain and G.-C. Rota. *Introduction to Geometric Probability*. Cambridge University Press, 1997. 2
- [6] G. C. Shephard and R. J. Webster. Metrics for sets of convex bodies. *Mathematika*, 12:73–88, 1965. 1, 2
- [7] R. A. Vitale. L_p metrics for compact, convex sets. *Journal of Approximation Theory*, 45:280–287, 1985. 2

DEPARTMENT OF MATHEMATICS & COMPUTER SCIENCE, LONGWOOD UNIVERSITY,
U.S.A. *E-mail address*: hoehnersd@longwood.edu