

Signed Mahonian Polynomials on Colored Derangements

Hasan Arslan^{*,a}, Nazmiye Alemdar^a

^a*Department of Mathematics, Erciyes University, 38039, Kayseri, Turkey*

Abstract

The polynomial $\sum_{\pi \in W} q^{maj(\pi)}$ of major index over a classical Weyl group W with a generating set S is called the Mahonian polynomial over W , and also the polynomial $\sum_{\pi \in W} (-1)^{l(\pi)} q^{maj(\pi)}$ of major index together with sign over the group W is called the signed Mahonian polynomial over the group W , where l is the length function on W defined in terms of the generating set S . We concern with the signed Mahonian polynomial

$$\sum_{\pi \in D_n^{(c)}} (-1)^{L(\pi)} q^{fmaj(\pi)}$$

on the set $D_n^{(c)}$ of colored derangements in the group $G_{c,n}$ of colored permutations, where L denotes the length function defined by means of a complex root system described by Bremke and Malle in $G_{c,n}$ and $fmaj$ defined by Adin and Roichman in $G_{c,n}$ represents the *flag-major index*, which is a Mahonian statistic. As an application of the formula for signed Mahonian polynomials on the set of colored derangements, we will derive a formula to count colored derangements of even length in $G_{c,n}$ when c is an even number. Finally, we conclude by providing a formula for the difference between the number of derangements of even and odd lengths in $G_{c,n}$ when c is even.

Keywords: Derangement, Mahonian statistic, Mahonian polynomials, length function.

2020 Mathematics Subject Classification: 05A05, 05A15, 05A19.

Email Adressess: hasanarslan@erciyes.edu.tr (H. Arslan), nakari@erciyes.edu.tr (N. Alemdar)

*Corresponding Author: Hasan Arslan

1 Introduction

In this paper, we deal with the q -enumeration of the Mahonian polynomial of colored derangements in the group of colored permutations $G_{c,n}$ based on the flag major index. Let \mathbb{N} be the set of natural numbers. For $n \in \mathbb{N}$, let $[n] := \{1, 2, \dots, n\}$ (where $[0] := \emptyset$). The cardinality of a set X will be denoted by $|X|$. Let c and n be positive integers. The group of colored permutations $G_{c,n}$ may be described as follows: The group $G_{c,n}$ of n letters with c colors may be regarded as a split group extension $G_{c,n} = \mathfrak{S}_n \rtimes \mathbb{Z}_c^n$ of \mathbb{Z}_c^n by \mathfrak{S}_n , where $\mathbb{Z}_c := \{0, 1, \dots, c-1\}$, \mathbb{Z}_c^n is the direct product of n copies of \mathbb{Z}_c and \mathfrak{S}_n is the symmetric group on $[n]$. A much more natural manner to introduce $G_{c,n}$ is in the following way: Now consider the set $I = \{i^{[k]} : i = 1, \dots, n; k = 0, \dots, c-1\}$ as the set $[n]$ colored by the colors $0, 1, \dots, c-1$. Then, the colored permutations group $G_{c,n}$ consists of all bijections $\pi : I \mapsto I$ providing that $\pi(i^{[k]}) = (\pi(i))^{[k]}$ for all $i = 1, \dots, n; k = 0, \dots, c-1$. Therefore, any element π of the group $G_{c,n}$ is written as follows:

$$\pi = \left(\begin{array}{cccc} 1 & 2 & \dots & n \\ \pi_1^{[t_1]} & \pi_2^{[t_2]} & \dots & \pi_n^{[t_n]} \end{array} \right),$$

where $\pi_i \in [n]$ and $0 \leq t_i \leq c-1$ for all $i = 1, \dots, n$. We shall represent π by the word by removing its first row:

$$\pi = \pi_1^{[t_1]} \pi_2^{[t_2]} \dots \pi_n^{[t_n]}.$$

The *color* number of π is defined as the sum $\text{col}(\pi) = \sum_{i=1}^n t_i$. In particular, the underlying permutation of a colored permutation $\pi = \pi_1^{[t_1]} \pi_2^{[t_2]} \dots \pi_n^{[t_n]} \in G_{c,n}$ is denoted by $|\pi| = \pi_1 \pi_2 \dots \pi_n \in \mathfrak{S}_n$.

As is well-known from [21], the inversion number and the descent set of $\sigma \in \mathfrak{S}_n$ are respectively defined in the following way:

$$\begin{aligned} \text{inv}(\sigma) &= |\{(i, j) \in [n] \times [n] : i < j \text{ and } \sigma_i > \sigma_j\}| \\ \text{Des}(\sigma) &= \{i \in [n-1] : \sigma_i > \sigma_{i+1}\}. \end{aligned}$$

The usual (type A) major index, a statistic derivable from $\text{Des}(\sigma)$, is identified as $\text{maj}(\sigma) = \sum_{i \in \text{Des}(\sigma)} i$. MacMahon proved in [20] that the the number of inversions inv is equidistributed with the major index maj over the symmetric group \mathfrak{S}_n , that is,

$$\sum_{\sigma \in \mathfrak{S}_n} q^{\text{inv}(\sigma)} = \sum_{\sigma \in \mathfrak{S}_n} q^{\text{maj}(\sigma)} = [n]!_q$$

where q is an indeterminate, $[i]_q := \frac{1-q^i}{1-q}$ for all $i = 1, \dots, n$ is a q -integer and $[n]_q! := [1]_q [2]_q \dots [n-1]_q [n]_q$ is the usual q -analogue of $n!$. If a permutation statistic is equidistributed with the length function, that is, the number of inversions, then it is called *Mahonian*.

Given $\pi = \left(\begin{smallmatrix} 1 & 2 & \cdots & n \\ \pi_1^{[t_1]} & \pi_2^{[t_2]} & \cdots & \pi_n^{[t_n]} \end{smallmatrix} \right) \in G_{c,n}$, the length of π is defined by

$$L(\pi) := \text{col}(\pi) + c \cdot \sum_{t_j \neq 0} |\{(i, j) : i < j \text{ and } \pi_i < \pi_j\}| + \text{inv}(|\pi|) \quad (1)$$

due to [10], where $\text{inv}(|\pi|)$ corresponds to the classical inversion in \mathfrak{S}_n . Clearly, $L(|\pi|) = \text{inv}(|\pi|)$ for all $\pi \in G_{c,n}$. For example, the length of $\pi = 2^{[3]} 1^{[1]} 3 4^{[2]} 5 \in G_{4,5}$ is $L(\pi) = 6 + 4 \cdot 3 + 1 = 19$. It is well known from [5] that for each $\pi \in G_{c,n}$ the length $L(\pi)$ corresponds to the colored inversion $\text{inv}_c(\pi)$. The flag-major index of any colored element can be practically computed by using the next theorem.

Theorem 1.1 (Adin-Roichman [1]). *Let $\pi \in G_{c,n}$. Then*

$$f\text{maj}(\pi) = c \cdot \text{maj}(\pi) + \text{col}(\pi) \quad (2)$$

where maj is computed with respect to the following total order:

$$1^{[c-1]} < \cdots < n^{[c-1]} < \cdots < 1^{[2]} < \cdots < n^{[2]} < 1^{[1]} < \cdots < n^{[1]} < 1 < \cdots < n. \quad (3)$$

In Theorem 1.1, $\text{maj}(\pi)$ equals the sum $\sum_{i \in \text{Des}_c(\pi)} i$, where $\text{Des}_c(\pi) := \{i \in [n-1] : \pi_i > \pi_{i+1}\}$ is determined by considering the total ordering in (3).

To illustrate Theorem 1.1, we will continue with our running example. Thus, we obtain that the flag-major index of $\pi = 2^{[3]} 1^{[1]} 3 4^{[2]} 5 \in G_{4,5}$ is $f\text{maj}(\pi) = 4 \cdot 3 + 6 = 18$. Adin and Roichman in [1] showed that the flag major index in the case $c = 1$ (i.e., for the symmetric group \mathfrak{S}_n) coincides with the usual major index. It is well known from [5] that the flag major index is a Mahonian statistic, that is,

$$\sum_{\pi \in G_{c,n}} q^{f\text{maj}(\pi)} = \prod_{i=1}^n [ci]_q = \prod_{i=1}^n q^{L(w)}. \quad (4)$$

Although the statistics inv and maj have symmetric joint distribution

$$\sum_{\pi \in \mathfrak{S}_n} t^{\text{inv}(\pi)} q^{\text{maj}(\pi)} = \sum_{\pi \in \mathfrak{S}_n} q^{\text{inv}(\pi)} t^{\text{maj}(\pi)} \quad (5)$$

over \mathfrak{S}_n [15], the statistics $f\text{maj}$ and inv_c do not generally have a symmetric joint distribution over $G_{c,n}$. For instance, observation of Table 1 shows that the joint symmetric distribution condition in question is not satisfied for $G_{4,2}$. In [3], Ahmia, Ramir ez and Villamizar introduced an inversion-like statistic $\widetilde{\text{inv}}$ on $G_{c,n}$ by

$$\widetilde{\text{inv}}(\pi) = c \cdot \text{inv}(|\pi|) + \text{col}(\pi) \quad (6)$$

for each $\pi \in G_{c,n}$. It easily follows that \widetilde{inv} is equidistributed with $fmaj$ over $G_{c,n}$. Clearly, $\widetilde{inv}(\pi) = \widetilde{inv}(\pi^{-1})$ for each $\pi \in G_{c,n}$. Furthermore, we will assert in Conjecture 2.6 that the joint distribution of the statistics \widetilde{inv} and $fmaj$ is symmetric over $G_{c,n}$ in the sense of (5).

For $n \geq 1$, let $D_n^{(c)} = \{\pi \in G_{c,n} : \pi(i) \neq i \text{ for all } i \in [n]\}$ be the set of all colored derangements in $G_{c,n}$. Specifically, we investigate the Mahonian polynomials defined on the set $D_n^{(c)}$ of all colored derangements in the group $G_{c,n}$:

$$d_n^{(c)}(q) := \sum_{\pi \in D_n^{(c)}} q^{fmaj(\pi)} \quad (7)$$

The use of the length function L together with $fmaj$, which we have been unable to find in the existing literature, leads to some new formulas for the signed Mahonian polynomials on the set $D_n^{(c)}$ of colored derangements in the colored permutations group $G_{c,n}$. In addition, Bagno [7] introduced a length function ℓ on $G_{c,n}$. Signed Mahonian and Euler-Mahonian polynomials studied in the literature are mainly based on the length function ℓ (see [11, 14]). We remark that the length function L , which will be used throughout the paper, is completely different from ℓ . Utilizing equations (4) and (7) together, we will produce an explicit formula for the signed Mahonian polynomials over $D_n^{(c)}$

$$\bar{d}_n^{(c)}(q) := \sum_{\pi \in D_n^{(c)}} (-1)^{L(\pi)} q^{fmaj(\pi)} \quad (8)$$

when c is even. Gessel [17] found a remarkable formula for the Mahonian polynomial over the symmetric group \mathfrak{S}_n , which is also known as $G_{1,n}$, by major index. Subsequently, another elegant proof was given by Wachs [22] in a combinatorial way. Chow [13] extended Wachs' method to produce a formula for the Mahonian polynomial over the group of signed permutations B_n (a Weyl group of type B), which can be viewed as the group $G_{2,n}$, by employing the flag major index.

In a recent work, Ji and Zhang [18] derived a formula for the Mahonian polynomial of derangements in the even-signed permutations group (a Weyl group of type D) by considering the D-major index introduced by Biagioli and Caselli in [8], extending Wachs' approach and utilizing a refinement of Stanley's shuffle theorem produced by themselves in [19]. Moreover, Ji and Zhang [18] established an explicit formula for signed Mahonian polynomial on the set of derangements in each classical Weyl group W .

2 The signed Mahonian polynomial over $D_n^{(c)}$

In this section, our primary focus will be on deriving a formula for the sum

$$\sum_{\pi \in G_{c,n}} (-1)^{L(\pi)} q^{fmaj(\pi)} \quad (9)$$

where L stands for the length function described by the formula (1). We will prove equation (9) depending on the parity of c . To this end, we will use the formula

$$\sum_{\pi \in G_{c,n}} (-1)^{L(|\pi|)} q^{fmaj(\pi)} = [c]_q [2c]_{-q} [3c]_q [4c]_{-q} \cdots [nc]_{(-1)^{n-1}q}. \quad (10)$$

proven by Biagioli and Caselli in [9].

Theorem 2.1. *If c is an even number, then we have*

$$\sum_{\pi \in G_{c,n}} (-1)^{L(\pi)} q^{fmaj(\pi)} = [c]_{-q} [2c]_q [3c]_{-q} [4c]_q \cdots [nc]_{(-1)^n q}. \quad (11)$$

Proof. Since c is even and considering the length formula given in equation (1), we deduce that $L(\pi) \equiv \text{col}(\pi) + \text{inv}(|\pi|) \pmod{2}$ for any $\pi \in G_{c,n}$. From equations (2) and (10), we obtain

$$\begin{aligned} \sum_{\pi \in G_{c,n}} (-1)^{L(\pi)} q^{fmaj(\pi)} &= \sum_{\pi \in G_{c,n}} (-1)^{\text{col}(\pi) + \text{inv}(|\pi|)} q^{fmaj(\pi)} \\ &= \sum_{\pi \in G_{c,n}} (-1)^{L(|\pi|)} (-q)^{fmaj(\pi)} \\ &= [c]_{-q} [2c]_q [3c]_{-q} [4c]_q \cdots [nc]_{(-1)^n q} \end{aligned}$$

as desired. □

Equation (11) can also be expressed as

$$\sum_{\pi \in G_{c,n}} (-1)^{L(\pi)} q^{fmaj(\pi)} = \left(\frac{1-q}{1+q} \right)^{\lfloor \frac{n+1}{2} \rfloor} \prod_{i=1}^n [ci]_q \quad (12)$$

when c is even. In Table 1, one can see the distribution of the length function L and flag major index $fmaj$ on $G_{4,2}$. We will illustrate the formula given in (11) in the following example.

Example 2.2. Taking Table 1 into account, it can be checked that the formula (11) holds for $G_{4,2}$. Namely, we have

$$\sum_{\pi \in G_{4,2}} (-1)^{L(\pi)} q^{f\text{maj}(\pi)} = 1 + q^2 - q^8 - q^{10} = [4]_{-q} [8]_q.$$

When we set $c = 2$, Theorem 2.1 corresponds to

$$\sum_{\pi \in B_n} (-1)^{l_B(\pi)} q^{f\text{maj}(\pi)} = [2]_{-q} [4]_q \cdots [2n]_{(-1)^n q}$$

which is a result of Adin, Gessel, and Roichman for the hyperoctahedral group B_n (see Theorem 5.1 in [2]). In [1], Adin and Roichman proved that the flag major index in the case $c = 2$ (i.e., for the hyperoctahedral group B_n) is identical with the flag major index in the group B_n . It should be also noted here that in the case of $c = 2$ the length function L agrees with the canonical length function l_B of the Weyl group B_n (see [10]).

Table 1: Distribution of the length, flag major index and \widetilde{inv} on $G_{4,2}$

$\pi =$ $\pi_1 \pi_2$	$L(\pi)$	$f\text{maj}(\pi)$	$\widetilde{inv}(\pi)$	$\pi =$ $\pi_1 \pi_2$	$L(\pi)$	$f\text{maj}(\pi)$	$\widetilde{inv}(\pi)$
1 2	0	0	0	2 1 ^[3]	4	7	7
1 ^[1] 2	1	1	1	2 ^[1] 1 ^[3]	5	8	8
1 ^[2] 2	2	2	2	2 ^[2] 1 ^[3]	6	9	9
1 ^[3] 2	3	3	3	2 ^[3] 1 ^[3]	7	10	10
2 1	1	4	4	1 2 ^[1]	5	5	1
2 ^[1] 1	2	1	5	1 ^[1] 2 ^[1]	6	2	2
2 ^[2] 1	3	2	6	1 ^[2] 2 ^[1]	7	3	3
2 ^[3] 1	4	3	7	1 ^[3] 2 ^[1]	8	4	4
2 1 ^[1]	2	5	5	1 2 ^[2]	6	6	2
2 ^[1] 1 ^[1]	3	6	6	1 ^[1] 2 ^[2]	7	7	3
2 ^[2] 1 ^[1]	4	3	7	1 ^[2] 2 ^[2]	8	4	4
2 ^[3] 1 ^[1]	5	4	8	1 ^[3] 2 ^[2]	9	5	5
2 1 ^[2]	3	6	6	1 2 ^[3]	7	7	3
2 ^[1] 1 ^[2]	4	7	7	1 ^[1] 2 ^[3]	8	8	4
2 ^[2] 1 ^[2]	5	8	8	1 ^[2] 2 ^[3]	9	9	5
2 ^[3] 1 ^[2]	6	5	9	1 ^[3] 2 ^[3]	10	6	6

When c is an odd number, a brute-force computation for $c \leq 7$ and $n \leq 3$

shows that the polynomial

$$\sum_{\pi \in G_{c,n}} (-1)^{L(\pi)} q^{fmaj(\pi)}$$

does not have a nice factorial-type product formula as in (11). Similarly to Theorem 2.1, we state the following result without proof, based on equations (2), (4) and (6).

Corollary 2.3. *When c is an even number, we have*

$$\sum_{\pi \in G_{c,n}} (-1)^{\widetilde{inv}(\pi)} q^{fmaj(\pi)} = \sum_{\pi \in G_{c,n}} (-1)^{fmaj(\pi)} q^{\widetilde{inv}(\pi)} = \prod_{i=1}^n [ci]_{-q}.$$

Table 2: Distribution of the length, flag major index and \widetilde{inv} on $G_{3,2}$

$\pi =$ $\pi_1 \pi_2$	$L(\pi)$	$fmaj(\pi)$	$\widetilde{inv}(\pi)$	$\pi =$ $\pi_1 \pi_2$	$L(\pi)$	$fmaj(\pi)$	$\widetilde{inv}(\pi)$
2 1	1	3	3	1 2	0	0	0
2 ^[1] 1	2	1	4	1 ^[1] 2	1	1	1
2 ^[2] 1	3	2	5	1 ^[2] 2	2	2	2
2 1 ^[1]	2	4	4	1 2 ^[1]	4	4	1
2 ^[1] 1 ^[1]	3	5	5	1 ^[1] 2 ^[1]	5	2	2
2 ^[2] 1 ^[1]	4	3	6	1 ^[2] 2 ^[1]	6	3	3
2 1 ^[2]	3	5	5	1 2 ^[2]	5	5	2
2 ^[1] 1 ^[2]	4	6	6	1 ^[1] 2 ^[2]	6	6	3
2 ^[2] 1 ^[2]	5	7	7	1 ^[2] 2 ^[2]	7	4	4

The next result immediately follows from the equations (13), (14) and (15) which occur in the proof of Theorem 4.4 in [9].

Corollary 2.4. *When c is an odd number, we have*

$$\sum_{\pi \in G_{c,n}} (-1)^{\widetilde{inv}(\pi)} q^{fmaj(\pi)} = \begin{cases} [c]_q [2c]_{-q} [3c]_q [4c]_{-q} \cdots [nc]_{(-1)^{n-1}q}, & \text{if } n \text{ is even} \\ [c]_{-q} [c]_q [2c]_{-q} [3c]_q [4c]_{-q} \cdots [(n-2)c]_q [(n-1)c]_{-q} [n]_{q^c}, & \text{if } n \text{ is odd.} \end{cases}$$

Now, we will give the following example to illustrate Corollary 2.4 based on Table 2.

Example 2.5. *Considered Table 2, we get*

$$\sum_{\pi \in G_{3,2}} (-1)^{\widetilde{inv}(\pi)} q^{fmaj(\pi)} = 1 + q^2 - q^3 + q^4 - q^5 - q^7 = [3]_q [6]_{-q}.$$

If we substitute $c \rightarrow 1$ in Corollary 2.4, then we encounter the polynomial

$$\sum_{\pi \in \mathfrak{S}_n} (-1)^{inv(\pi)} q^{maj(\pi)} = [1]_q [2]_{-q} \cdots [n]_{(-1)^{n-1}q}$$

which is nothing else but the result constructed by Gessel and Simon for the symmetric group \mathfrak{S}_n (see Corollary 2 in [23]). We also realized that for some c odd numbers the polynomial $\sum_{\pi \in G_{c,n}} (-1)^{fmaj(\pi)} q^{\widetilde{inv}(\pi)}$ has the same formula as in Corollary 2.4. For instance, this situation may be checked for each group $G_{3,2}$, $G_{5,2}$, $G_{7,2}$ and $G_{3,3}$. In addition, we observed that the joint distribution of the permutation statistics \widetilde{inv} and $fmaj$ on $G_{c,n}$ is symmetric for some values of c and n . Therefore, the above observation allows us to pose the following conjecture, which is computationally confirmed for every group $G_{c,n}$ with $1 \leq c \leq 7$ and $n = 2$ as well as for the groups $G_{3,3}$ and $G_{4,3}$ (see Appendix A).

Conjecture 2.6. *The statistic \widetilde{inv} has a symmetric joint distribution with $fmaj$ over $G_{c,n}$. That is,*

$$\sum_{\pi \in G_{c,n}} t^{\widetilde{inv}(\pi)} q^{fmaj(\pi)} = \sum_{\pi \in G_{c,n}} q^{\widetilde{inv}(\pi)} t^{fmaj(\pi)}.$$

It is important to note here that Conjecture 2.6 may be considered as a generalization of the equality (5). The best way to prove that the statistics \widetilde{inv} and $fmaj$ on $G_{c,n}$ have a symmetric joint distribution is to find, if possible, an explicit bijection $\psi : G_{c,n} \rightarrow G_{c,n}$ such that for all $\pi \in G_{c,n}$

$$\widetilde{inv}(\pi) = fmaj(\psi(\pi)) \quad \text{and} \quad fmaj(\pi) = \widetilde{inv}(\psi(\pi)).$$

In other words, one needs to show that ψ interchanges the two statistics \widetilde{inv} and $fmaj$.

Following Wachs' work [22], for any colored permutation $\pi \in G_{c,A}$, where $A = \{0 < a_1 < a_2 < \cdots < a_k\}$, the *reduction* of π to the non-fixed points is defined to be the colored permutation $dp(\pi) \in G_{c,k}$ by replacing each letter $\pi_j^{[r_j]}$ by $i^{[r_j]}$ if $\pi_j = a_i$, $i = 1, 2, \dots, k$. Thus, the *derangement part* $dp(\pi)$ of a colored permutation $\pi \in G_{c,n}$ is the reduction of subword of non-fixed points of π . For example, $dp(2^{[3]} 1^{[1]} 3 4^{[2]} 5) = 2^{[3]} 1^{[1]} 3^{[2]}$. It should be noted here that the derangement part of a colored permutation is a colored derangement, and

conversely, any derangement in $D_n^{(c)}$ and a subset having $n - k$ elements of $[n]$ give rise to a unique colored permutation in $G_{c,n}$ with $n - k$ fixed points.

Now let $\pi = \pi_1^{[t_1]} \pi_2^{[t_2]} \cdots \pi_n^{[t_n]} \in G_{c,n}$. A letter $\pi_i^{[t_i]}$ is called a *subcedant* (respectively an *excedant*) of π if $\pi_i < i$ (respectively, $\pi_i > i$) with respect to the total ordering in (3). It is obvious that all excedants of π are positive. Let $s(\pi)$ and $e(\pi)$ denote the number of subcedants and excedants of π , respectively. Let $\tilde{\pi}$ be a colored permutation having k letters obtained from $\pi \in G_{c,k}$ with $k \leq n$ in the following order:

- Replace i th smallest (in absolute value) subcedant $\pi_j^{[t_j]}$ of π by $i^{[t_j]}$, $i = 1, 2, \dots, s(\pi)$.
- Assign i th smallest fixed point of π to $s(\pi) + 1$, $i = 1, 2, \dots, k - s(\pi) - e(\pi)$.
- Write $n - i + 1$ instead of i th largest excedant of π , where i ranges from 1 to $e(\pi)$.

Note that if $k = n$, then $\tilde{\pi} \in G_{c,n}$. Moreover, if $\pi \in D_n^{(c)}$ then $\tilde{\pi} \in D_A^{(c)}$, where $A = \{1, 2, \dots, s(\pi)\} \cup \{n - e(\pi) + 1, n - e(\pi) + 2, \dots, n\}$. For instance, the collections of subcedants, excedants and fixed points of $\pi = 2 \ 1^{[1]} \ 3 \ 4^{[2]} \ 5 \ 7 \ 6^{[2]}$ are, respectively, $\{1^{[1]}, 4^{[2]}, 6^{[2]}\}$, $\{2, 7\}$ and $\{3, 5\}$. Since the above construction, π corresponds to $\tilde{\pi} = 6 \ 3^{[1]} \ 4 \ 2^{[2]} \ 5 \ 7 \ 1^{[2]}$. One can observe that descent set of π and $\tilde{\pi}$ are the same. This is not an accident.

Lemma 2.7. *Let $\pi \in G_{c,k}$ with $k \leq n$. Then $Des_c(\pi) = Des_c(\tilde{\pi})$ and $col(\pi) = col(\tilde{\pi})$.*

Proof. It is easy to see from the construction of $\tilde{\pi}$ that the colors remain the same. Based on the total ordering in (3), we achieved $Des_c(\pi) = Des_c(\tilde{\pi})$ by following the same arguments as in the proof of Lemma 1 in [22]. \square

Given two disjoint permutations $\tau = \tau_1 \cdots \tau_j$ and $\sigma = \sigma_1 \cdots \sigma_k$, a *shuffle* of τ and σ is a permutation $w = w_1 \cdots w_{j+k}$ such that τ and σ appear as subsequences in w . The collection of all shuffles of τ and σ is denoted by $sh(\tau, \sigma)$. Garsia and Gessel [16] showed that the weight of the major index on $sh(\tau, \sigma)$ is

$$\sum_{w \in sh(\tau, \sigma)} q^{maj(w)} = q^{maj(\tau) + maj(\sigma)} \begin{bmatrix} j + k \\ k \end{bmatrix}_q. \quad (13)$$

Equation (13) remains true if τ and σ are colored words of different letters from the alphabet in equation (3). It is apparent that if $\pi \in sh(\tau, \sigma)$ we have $col(\pi) =$

$col(\tau) + col(\sigma)$. Considering the definition of flag major index in (2), it is not hard to see that the following identity is true:

$$\sum_{\pi \in sh(\tau, \sigma)} q^{fmaj(w)} = q^{fmaj(\tau) + fmaj(\sigma)} \begin{bmatrix} j+k \\ k \end{bmatrix}_{q^c}. \quad (14)$$

The above construction of $\tilde{\pi}$ from π yields to following bijection, proved by Assaf in [6], is a generalized version of Theorem 2 in [22] for the $G_{c,n}$. In the proof of the following lemma, Assaf used the total ordering

$$n^{[c-1]} < \dots < 1^{[c-1]} < \dots < n^{[2]} < \dots < 1^{[2]} < n^{[1]} < \dots < 1^{[1]} < 1 < \dots < n$$

instead of the total ordering in equation (3). However, we note that Lemma 2.8 still remains true when used the total ordering in equation (3).

Lemma 2.8 (Assaf [6]). *Let $\alpha \in D_k^{(c)}$ with $k \leq n$ and $\gamma = (s(\alpha) + 1) \dots (s(\alpha) + k)$. Then the map $\phi : \{\pi \in G_{c,n} : dp(\pi) = \alpha\} \rightarrow sh(\tilde{\alpha}, \gamma)$ defined to be $\phi(\pi) = \tilde{\pi}$ is a bijection such that $Des_c(\phi(\pi)) = Des_c(\pi)$ and $col(\phi(\pi)) = col(\pi)$.*

Thus, the bijection ϕ given in Lemma 2.8 preserves the flag major index, i.e., $fmaj(\phi(\pi)) = fmaj(\pi)$ for all $\pi \in \{\pi \in G_{c,n} : dp(\pi) = \alpha\}$. In the following corollary, we will extend Wachs' formula

$$\sum_{\substack{dp(\pi)=\sigma \\ \pi \in \mathfrak{S}_n}} q^{maj(\pi)} = q^{maj(\sigma)} \begin{bmatrix} n \\ k \end{bmatrix}_q$$

given in [22] to colored case by employing Lemma 2.7, Lemma 2.8 and equation (14).

Corollary 2.9. *Let $0 \leq k \leq n$ and $\sigma \in D_k^{(c)}$. Then we have*

$$\sum_{\substack{dp(\pi)=\sigma \\ \pi \in G_{c,n}}} q^{fmaj(\pi)} = q^{fmaj(\sigma)} \begin{bmatrix} n \\ k \end{bmatrix}_{q^c}$$

where $\begin{bmatrix} n \\ k \end{bmatrix}_{q^c} = \frac{[n]_{q^c}!}{[k]_{q^c}! [n-k]_{q^c}!}$ is the q -binomial coefficient.

Proposition 2.10. *Let $0 \leq k \leq n$ and $\sigma \in D_k^{(c)}$. If c is even, then we have*

$$\sum_{\substack{dp(\pi)=\sigma \\ \pi \in G_{c,n}}} (-1)^{L(\pi)} q^{fmaj(\pi)} = (-1)^{L(\sigma)} q^{fmaj(\sigma)} \begin{bmatrix} n \\ k \end{bmatrix}_{q^c}.$$

Proof. Viewed through the lens of Corollary 2.9, it suffices to prove that

$$L(\pi) \equiv L(dp(\pi)) \pmod{2}$$

for each colored permutation $\pi \in G_{c,n}$. Suppose that the fixed points of $\pi = \pi_1^{[t_1]} \pi_2^{[t_2]} \dots \pi_n^{[t_n]}$ are in increasing order $p_1 < p_2 < \dots < p_k$. Since $L(\pi) = inv_c(\pi)$, the contribution of inversions at exactly each fixed point p_j to the total number of inversions $inv_c(\pi)$ can be obtained from the proof of Theorem 4.1 in [3] as $2(p_j - j)(n - k - p_j + j)$ and so the contribution of this case to the inversion number $inv_c(\pi)$ is even. On the other hand, any $\pi_r^{[t_r]}$ with $t_r \neq 0$ appearing after the position of p_j and greater than p_j contributes the number c to $inv_c(\pi)$. Thus, removing all fixed points from π , $inv_c(\pi)$ decreases by an even number since c is even. In conclusion, we deduce that $L(\pi) \equiv L(dp(\pi)) \pmod{2}$. \square

Theorem 2.11. *For any $n \geq 1$, we have*

$$d_n^{(c)}(q) = \sum_{\pi \in D_n^{(c)}} q^{maj(\pi)} = [c]_q [2c]_q \dots [nc]_q \sum_{k=0}^n \frac{(-1)^k q^{c \binom{k}{2}}}{[c]_q [2c]_q \dots [kc]_q}. \quad (15)$$

Proof. Considering Corollary 2.9 and equation (4), we have

$$\begin{aligned} [c]_q [2c]_q \dots [nc]_q &= \sum_{\pi \in G_{c,n}} q^{maj(\pi)} \\ &= \sum_{k=0}^n \sum_{\alpha \in D_k^{(c)}} \sum_{dp(\pi)=\alpha} q^{maj(\pi)} \\ &= \sum_{k=0}^n \sum_{\alpha \in D_k^{(c)}} q^{maj(\alpha)} \begin{bmatrix} n \\ k \end{bmatrix}_{q^c} \\ &= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_{q^c} d_k^{(c)}(q). \end{aligned}$$

Hence, Gauss inversion (i.e., q -binomial inversion) given in Corollary 3.38 of [4] on the resulting equation leads to the desired formula (15). \square

For any $n \geq 1$, we denote by $d_n^{(c)}$ the number of all colored derangements in $D_n^{(c)}$. Then

$$d_n^{(c)} = n! \sum_{k=0}^n \frac{(-1)^k c^{n-k}}{k!}. \quad (16)$$

It should be emphasized that Assaf [6] established the formula for $d_n^{(c)}$ in (16) by using Principle of Inclusion and Exclusion. Next result of this paper is the

following theorem, giving us the formula for the signed Mahonian polynomial over $D_n^{(c)}$.

Theorem 2.12. *Let c be even and $n \geq 1$. Then, we have*

$$\bar{d}_n^{(c)}(q) = \sum_{\alpha \in D_n^{(c)}} (-1)^{L(\alpha)} q^{f_{maj}(\alpha)} = [c]_q [2c]_q \dots [nc]_q \sum_{k=0}^n \frac{(-1)^k q^{c \binom{k}{2}}}{[c]_q [2c]_q \dots [kc]_q} \left(\frac{1-q}{1+q} \right)^{\lfloor \frac{n-k+1}{2} \rfloor}.$$

Proof. Summing over all colored derangements $\alpha \in D_k^{(c)}$ with $0 \leq k \leq n$, we can deduce from Proposition 2.10 and equation (8) that

$$\begin{aligned} \sum_{\pi \in G_{c,n}} (-1)^{L(\pi)} q^{f_{maj}(\pi)} &= \sum_{k=0}^n \sum_{\alpha \in D_k^{(c)}} \sum_{dp(\pi)=\alpha, \pi \in G_{c,n}} (-1)^{L(\pi)} q^{f_{maj}(\pi)} \\ &= \sum_{k=0}^n \sum_{\alpha \in D_k^{(c)}} (-1)^{L(\alpha)} q^{f_{maj}(\alpha)} \left[\begin{matrix} n \\ k \end{matrix} \right]_{q^c} \\ &= \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_{q^c} \bar{d}_k^{(c)}. \end{aligned}$$

Taking the formula in (12) into account, we get

$$\sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_{q^c} \bar{d}_k^{(c)} = \left(\frac{1-q}{1+q} \right)^{\lfloor \frac{n+1}{2} \rfloor} \prod_{i=1}^n [ci]_q. \quad (17)$$

It is also well known from the proof of Theorem 2.2 in [3] that $[ci]_q = [c]_q [i]_{q^c}$ for any positive integer i . Applying Gauss inversion to equation (17) yields to

$$\begin{aligned} \bar{d}_n^{(c)} &= \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_{q^c} (-1)^{n-k} q^{c \binom{n-k}{2}} \left(\frac{1-q}{1+q} \right)^{\lfloor \frac{k+1}{2} \rfloor} \prod_{i=1}^k [ci]_q \\ &= [c]_q [2c]_q \dots [nc]_q \sum_{k=0}^n \frac{(-1)^k q^{c \binom{k}{2}}}{[c]_q [2c]_q \dots [kc]_q} \left(\frac{1-q}{1+q} \right)^{\lfloor \frac{n-k+1}{2} \rfloor} \end{aligned}$$

which is equivalent to the desired formula. \square

Let $D_n^{+(c)}$ denote the sets of colored derangements with even length (according to the length function L) in $G_{c,n}$. Define $d_n^{+(c)} := |D_n^{+(c)}|$. We infer from Theorem 2.11 and Theorem 2.12 that

$$d_n^{+(c)}(q) = [c]_q [2c]_q \dots [nc]_q \sum_{k=0}^n \frac{(-1)^k q^{c \binom{k}{2}}}{[c]_q [2c]_q \dots [kc]_q} \left(\frac{1}{2} + \frac{1}{2} \left(\frac{1-q}{1+q} \right)^{\lfloor \frac{n-k+1}{2} \rfloor} \right) \quad (18)$$

when c is even. In similar vein, we indicate by $D_n^{-(c)}$ the set of odd colored derangements in $G_{c,n}$. Let $d_n^{-(c)} := |D_n^{-(c)}|$. Substituting $q \rightarrow 1$ in equation (18), we obtain

$$d_n^{+(c)} = \frac{n!}{2} \sum_{k=0}^{n-1} \frac{(-1)^k c^{n-k}}{k!} + (-1)^n.$$

Moreover, we conclude that $d_n^{-(c)} = \frac{n!}{2} \sum_{k=0}^{n-1} \frac{(-1)^k c^{n-k}}{k!}$. Thus, the difference between the number of even and odd colored derangements in $G_{c,n}$ is

$$d_n^{+(c)} - d_n^{-(c)} = (-1)^n \quad (19)$$

when c is even. As a matter of the fact, the formula (19) is actually a general form of the difference, which can be easily obtained from [18], for the hyperoctahedral group corresponding to the case $c = 2$. Since the number $d_n^{+(2)}$ of even length derangements in the hyperoctahedral group B_n has been already computed by Ji and Zhang in [18], one can also count the number of odd derangements in the group B_n . In the case of symmetric group \mathfrak{S}_n , Chapman [12] provided a bijective proof that the difference between the number of even and odd derangements in \mathfrak{S}_n is $d_n^+ - d_n^- = (-1)^{n-1}(n-1)$. Due to [18], it can easily be seen that the difference in question is equal to zero for the even-signed permutation group.

Question:

- *It is a natural question to ask here what the structure of the signed Mahonian polynomial $\tilde{d}_n^{(c)}(q)$ is exactly when c is odd? In this case, what is the concrete formula for the difference $d_n^{+(c)} - d_n^{-(c)}$?*

3 Appendix A

In this section, we will present Table 3 and Table 4, which show the distributions of $fmaj$ and \widetilde{inv} on the groups $G_{3,3}$ and $G_{4,3}$, respectively. These tables helps one to satisfy that $fmaj$ and \widetilde{inv} have a symmetric joint distribution on both $G_{3,3}$ and $G_{4,3}$.

References

- [1] R.M. Adin, Y. Roichman, The flag major index and group actions on polynomial rings, European J. Combin. **22** (2001) 431-446.
- [2] R.M. Adin, I.M. Gessel, Y. Roichman, Signed Mahonians, J. Combin. Theory Ser. A **109**(1) (2005) 25-43.

- [3] M. Ahmia, J.L. Ramírez, D. Villamizar, Inversions in colored permutations, derangements, and involutions, arXiv:2505.01550, (2025).
- [4] M. Aigner, Combinatorial Theory. Springer-Verlag, New York, USA, (1979).
- [5] H. Arslan, A. Altoum, M. Zaarour, An inversion statistic on the generalized symmetric groups, Adv. in Appl. Math. **154** (2024) 102655.
- [6] S.H. Assaf, Cyclic derangements, Electron. J. Combin. **17** (2010) R163.
- [7] E. Bagno, Euler-Mahonian parameters on colored permutation groups, Sémin. Lothar. Combin. **51** (2004), Art. B51f, 16 pp.
- [8] R. Biagioli, F. Caselli, Invariant algebras and major indices for classical Weyl groups, Proc. London Math. Soc. **88**(3) (2004) 603-631.
- [9] R. Biagioli, F. Caselli, Weighted enumerations on projective reflection groups, Adv. in Appl. Math. **48**(1) (2012) 249-268.
- [10] K. Bremke, G. Malle, Reduced words and a length function for $G(e, 1, n)$, Indag. Math. **8** (1997) 453-469.
- [11] H. Chang, S.-P. Eu, S. Fu, Z. Lin, Y.-H. Lo, Signed Mahonian polynomials for major and sorting indices, J. Algebraic Combin. **53** (2021) 201-226.
- [12] R. Chapman, An involution on derangements, Discrete Math. **231** (2001) 121-122.
- [13] C.-O. Chow, On derangement polynomials of type B , Sémin. Lothar. Combin. **55** (2005/07/2007), Art. B55b, 6 pp.
- [14] S.-P. Eu, Z.-C. Lin, Y.-H. Lo, Signed Euler-Mahonian identities, European J. Combin. **91** (2021) 103209.
- [15] D. Foata, M.P. Schützenberger, Major index and inversion number of permutations, Math. Nachr. **83** (1978) 143-159.
- [16] A.M. Garsia, I.M. Gessel, Permutation statistics and partitions, Adv. in Math. **31** (1979) 288-305.
- [17] I.M. Gessel, C. Reutenauer, Counting permutations with given cycle structure and descent set, J. Combin. Theory Ser. A **64** (1993) 189-215.
- [18] K.Q. Ji, D.T.X. Zhang, Signed Mahonian polynomials on derangements in classical Weyl groups, European J. Combin. **124** (2025) 104083.

- [19] K.Q. Ji, D.T.X. Zhang, Some refinements of Stanley's shuffle theorem, J. Combin. Theory Ser. A **203** (2024) 105830.
- [20] P.A. MacMahon, Combinatory analysis Volume I, Cambridge University Press (1915).
- [21] R.P. Stanley, Enumerative combinatorics Volume I, Cambridge University Press (2011).
- [22] M.L. Wachs, On q -derangement numbers, Proc. Amer. Math. Soc. **106** (1989) 273-278.
- [23] M.L. Wachs, An involution for signed Eulerian numbers, Discrete Math. **99** (1992) 59-62.

Table 3: Distribution of $f\text{maj}$ and \widetilde{inv} on $G_{3,3}$

Rank	π_1	π_2	π_3	$f\text{maj}(\pi)$	$\widetilde{inv}(\pi)$	Rank	π_1	π_2	π_3	$f\text{maj}(\pi)$	$\widetilde{inv}(\pi)$	Rank	π_1	π_2	π_3	$f\text{maj}(\pi)$	$\widetilde{inv}(\pi)$
1	1	2	3	0	0	55	1	2	3 ^[1]	7	1	109	1	2	3 ^[2]	8	2
2	1 ^[1]	2	3	1	1	56	1 ^[1]	2	3 ^[1]	8	2	110	1 ^[1]	2	3 ^[2]	9	3
3	1 ^[2]	2	3	2	2	57	1 ^[2]	2	3 ^[1]	9	3	111	1 ^[2]	2	3 ^[2]	10	4
4	1	2 ^[1]	3	4	1	58	1	2 ^[1]	3 ^[1]	5	2	112	1	2 ^[1]	3 ^[2]	12	3
5	1 ^[1]	2 ^[1]	3	2	2	59	1 ^[1]	2 ^[1]	3 ^[1]	3	3	113	1 ^[1]	2 ^[1]	3 ^[2]	10	4
6	1 ^[2]	2 ^[1]	3	3	3	60	1 ^[2]	2 ^[1]	3 ^[1]	4	4	114	1 ^[2]	2 ^[1]	3 ^[2]	11	5
7	1	2 ^[2]	3	5	2	61	1	2 ^[2]	3 ^[1]	6	3	115	1	2 ^[2]	3 ^[2]	7	4
8	1 ^[1]	2 ^[2]	3	6	3	62	1 ^[1]	2 ^[2]	3 ^[1]	7	4	116	1 ^[1]	2 ^[2]	3 ^[2]	8	5
9	1 ^[2]	2 ^[2]	3	4	4	63	1 ^[2]	2 ^[2]	3 ^[1]	5	5	117	1 ^[2]	2 ^[2]	3 ^[2]	6	6
10	2	1	3	3	3	64	2	1	3 ^[1]	10	4	118	2	1	3 ^[2]	11	5
11	2 ^[1]	1	3	1	4	65	2 ^[1]	1	3 ^[1]	8	5	119	2 ^[1]	1	3 ^[2]	9	6
12	2 ^[2]	1	3	2	5	66	2 ^[2]	1	3 ^[1]	9	6	120	2 ^[2]	1	3 ^[2]	10	7
13	2	1 ^[1]	3	4	4	67	2	1 ^[1]	3 ^[1]	5	5	121	2	1 ^[1]	3 ^[2]	12	6
14	2 ^[1]	1 ^[1]	3	5	5	68	2 ^[1]	1 ^[1]	3 ^[1]	6	6	122	2 ^[1]	1 ^[1]	3 ^[2]	13	7
15	2 ^[2]	1 ^[1]	3	3	6	69	2 ^[2]	1 ^[1]	3 ^[1]	4	7	123	2 ^[2]	1 ^[1]	3 ^[2]	11	8
16	2	1 ^[2]	3	5	5	70	2	1 ^[2]	3 ^[1]	6	6	124	2	1 ^[2]	3 ^[2]	7	7
17	2 ^[1]	1 ^[2]	3	6	6	71	2 ^[1]	1 ^[2]	3 ^[1]	7	7	125	2 ^[1]	1 ^[2]	3 ^[2]	8	8
18	2 ^[2]	1 ^[2]	3	7	7	72	2 ^[2]	1 ^[2]	3 ^[1]	8	8	126	2 ^[2]	1 ^[2]	3 ^[2]	9	9
19	1	3	2	6	3	73	1	3	2 ^[1]	7	4	127	1	3	2 ^[2]	8	5
20	1 ^[1]	3	2	7	4	74	1 ^[1]	3	2 ^[1]	8	5	128	1 ^[1]	3	2 ^[2]	9	6
21	1 ^[2]	3	2	8	5	75	1 ^[2]	3	2 ^[1]	9	6	129	1 ^[2]	3	2 ^[2]	10	7
22	1	3 ^[1]	2	4	4	76	1	3 ^[1]	2 ^[1]	11	5	130	1	3 ^[1]	2 ^[2]	12	6
23	1 ^[1]	3 ^[1]	2	2	5	77	1 ^[1]	3 ^[1]	2 ^[1]	9	6	131	1 ^[1]	3 ^[1]	2 ^[2]	10	7
24	1 ^[2]	3 ^[1]	2	3	6	78	1 ^[2]	3 ^[1]	2 ^[1]	10	7	132	1 ^[2]	3 ^[1]	2 ^[2]	11	8
25	1	3 ^[2]	2	5	5	79	1	3 ^[2]	2 ^[1]	6	6	133	1	3 ^[2]	2 ^[2]	13	7
26	1 ^[1]	3 ^[2]	2	6	6	80	1 ^[1]	3 ^[2]	2 ^[1]	7	7	134	1 ^[1]	3 ^[2]	2 ^[2]	14	8
27	1 ^[2]	3 ^[2]	2	4	7	81	1 ^[2]	3 ^[2]	2 ^[1]	5	8	135	1 ^[2]	3 ^[2]	2 ^[2]	12	9
28	3	1	2	3	6	82	3	1	2 ^[1]	10	7	136	3	1	2 ^[2]	11	8
29	3 ^[1]	1	2	1	7	83	3 ^[1]	1	2 ^[1]	8	8	137	3 ^[1]	1	2 ^[2]	9	9
30	3 ^[2]	1	2	2	8	84	3 ^[2]	1	2 ^[1]	9	9	138	3 ^[2]	1	2 ^[2]	10	10
31	3	1 ^[1]	2	4	7	85	3	1 ^[1]	2 ^[1]	5	8	139	3	1 ^[1]	2 ^[2]	12	9
32	3 ^[1]	1 ^[1]	2	5	8	86	3 ^[1]	1 ^[1]	2 ^[1]	6	9	140	3 ^[1]	1 ^[1]	2 ^[2]	13	10
33	3 ^[2]	1 ^[1]	2	3	9	87	3 ^[2]	1 ^[1]	2 ^[1]	4	10	141	3 ^[2]	1 ^[1]	2 ^[2]	11	11
34	3	1 ^[2]	2	5	8	88	3	1 ^[2]	2 ^[1]	6	9	142	3	1 ^[2]	2 ^[2]	7	10
35	3 ^[1]	1 ^[2]	2	6	9	89	3 ^[1]	1 ^[2]	2 ^[1]	7	10	143	3 ^[1]	1 ^[2]	2 ^[2]	8	11
36	3 ^[2]	1 ^[2]	2	7	10	90	3 ^[2]	1 ^[2]	2 ^[1]	8	11	144	3 ^[2]	1 ^[2]	2 ^[2]	9	12
37	2	3	1	6	6	91	2	3	1 ^[1]	7	7	145	2	3	1 ^[2]	8	8
38	2 ^[1]	3	1	7	7	92	2 ^[1]	3	1 ^[1]	8	8	146	2 ^[1]	3	1 ^[2]	9	9
39	2 ^[2]	3	1	8	8	93	2 ^[2]	3	1 ^[1]	9	9	147	2 ^[2]	3	1 ^[2]	10	10
40	2	3 ^[1]	1	4	7	94	2	3 ^[1]	1 ^[1]	11	8	148	2	3 ^[1]	1 ^[2]	12	9
41	2 ^[1]	3 ^[1]	1	2	8	95	2 ^[1]	3 ^[1]	1 ^[1]	9	9	149	2 ^[1]	3 ^[1]	1 ^[2]	10	10
42	2 ^[2]	3 ^[1]	1	3	9	96	2 ^[2]	3 ^[1]	1 ^[1]	10	10	150	2 ^[2]	3 ^[1]	1 ^[2]	11	11
43	2	3 ^[2]	1	5	8	97	2	3 ^[2]	1 ^[1]	6	9	151	2	3 ^[2]	1 ^[2]	13	10
44	2 ^[1]	3 ^[2]	1	6	9	98	2 ^[1]	3 ^[2]	1 ^[1]	7	10	152	2 ^[1]	3 ^[2]	1 ^[2]	14	11
45	2 ^[2]	3 ^[2]	1	4	10	99	2 ^[2]	3 ^[2]	1 ^[1]	5	11	153	2 ^[2]	3 ^[2]	1 ^[2]	12	12
46	3	2	1	9	9	100	3	2	1 ^[1]	10	10	154	3	2	1 ^[2]	11	11
47	3 ^[1]	2	1	7	10	101	3 ^[1]	2	1 ^[1]	8	11	155	3 ^[1]	2	1 ^[2]	9	12
48	3 ^[2]	2	1	8	11	102	3 ^[2]	2	1 ^[1]	9	12	156	3 ^[2]	2	1 ^[2]	10	13
49	3	2 ^[1]	1	4	10	103	3	2 ^[1]	1 ^[1]	11	11	157	3	2 ^[1]	1 ^[2]	12	12
50	3 ^[1]	2 ^[1]	1	5	11	104	3 ^[1]	2 ^[1]	1 ^[1]	12	12	158	3 ^[1]	2 ^[1]	1 ^[2]	13	13
51	3 ^[2]	2 ^[1]	1	3	12	105	3 ^[2]	2 ^[1]	1 ^[1]	10	13	159	3 ^[2]	2 ^[1]	1 ^[2]	11	14
52	3	2 ^[2]	1	5	11	106	3	2 ^[2]	1 ^[1]	6	12	160	3	2 ^[2]	1 ^[2]	13	13
53	3 ^[1]	2 ^[2]	1	6	12	107	3 ^[1]	2 ^[2]	1 ^[1]	7	13	161	3 ^[1]	2 ^[2]	1 ^[2]	14	14
54	3 ^[2]	2 ^[2]	1	7	13	108	3 ^[2]	2 ^[2]	1 ^[1]	8	14	162	3 ^[2]	2 ^[2]	1 ^[2]	15	15

Table 4: Distribution of $fmaj$ and \widetilde{inv} on $G_{4,3}$

Rank	π_1	π_2	π_3	$fmaj(\pi)$	$\widetilde{inv}(\pi)$	Rank	π_1	π_2	π_3	$fmaj(\pi)$	$\widetilde{inv}(\pi)$	Rank	π_1	π_2	π_3	$fmaj(\pi)$	$\widetilde{inv}(\pi)$	Rank	π_1	π_2	π_3	$fmaj(\pi)$	$\widetilde{inv}(\pi)$
1	1	2	3	0	0	97	1	2	3[1]	9	5	193	1	2	3[2]	10	2	289	1	2	3[3]	11	5
2	1[1]	2	3	1	1	98	1[1]	2	3[1]	10	2	194	1[1]	2	3[2]	11	3	290	1[1]	2	3[3]	12	4
3	1[2]	2	3	2	2	99	1[2]	2	3[1]	11	3	195	1[2]	2	3[2]	12	4	291	1[2]	2	3[3]	13	5
4	1[3]	2	3	3	3	100	1[3]	2	3[1]	12	4	196	1[3]	2	3[2]	13	5	292	1[3]	2	3[3]	14	6
5	1	2[1]	3	5	1	101	1	2[1]	3[1]	6	2	197	1	2[1]	3[2]	15	3	293	1	2[1]	3[3]	16	4
6	1[1]	2[1]	3	2	2	102	1[1]	2[1]	3[1]	3	3	198	1[1]	2[1]	3[2]	12	4	294	1[1]	2[1]	3[3]	13	5
7	1[2]	2[1]	3	3	3	103	1[2]	2[1]	3[1]	4	4	199	1[2]	2[1]	3[2]	13	5	295	1[2]	2[1]	3[3]	14	6
8	1[3]	2[1]	3	4	4	104	1[3]	2[1]	3[1]	5	5	200	1[3]	2[1]	3[2]	14	6	296	1[3]	2[1]	3[3]	15	7
9	1	2[2]	3	6	2	105	1	2[2]	3[1]	7	3	201	1	2[2]	3[2]	8	4	297	1	2[2]	3[3]	17	5
10	1[1]	2[2]	3	7	3	106	1[1]	2[2]	3[1]	8	4	202	1[1]	2[2]	3[2]	9	5	298	1[1]	2[2]	3[3]	18	6
11	1[2]	2[2]	3	4	4	107	1[2]	2[2]	3[1]	5	5	203	1[2]	2[2]	3[2]	6	6	299	1[2]	2[2]	3[3]	15	7
12	1[3]	2[2]	3	5	5	108	1[3]	2[2]	3[1]	6	6	204	1[3]	2[2]	3[2]	7	7	300	1[3]	2[2]	3[3]	16	8
13	1	2[3]	3	7	3	109	1	2[3]	3[1]	8	4	205	1	2[3]	3[2]	9	5	301	1	2[3]	3[3]	10	6
14	1[1]	2[3]	3	8	4	110	1[1]	2[3]	3[1]	9	5	206	1[1]	2[3]	3[2]	10	6	302	1[1]	2[3]	3[3]	11	7
15	1[2]	2[3]	3	9	5	111	1[2]	2[3]	3[1]	10	6	207	1[2]	2[3]	3[2]	11	7	303	1[2]	2[3]	3[3]	12	8
16	1[3]	2[3]	3	6	6	112	1[3]	2[3]	3[1]	7	7	208	1[3]	2[3]	3[2]	8	8	304	1[3]	2[3]	3[3]	9	9
17	2	1	3	4	4	113	2	1	3[1]	13	5	209	2	1	3[2]	14	6	305	2	1	3[3]	15	7
18	2[1]	1	3	1	5	114	2[1]	1	3[1]	10	6	210	2[1]	1	3[2]	11	7	306	2[1]	1	3[3]	12	8
19	2[2]	1	3	2	6	115	2[2]	1	3[1]	11	7	211	2[2]	1	3[2]	12	8	307	2[2]	1	3[3]	13	9
20	2[3]	1	3	3	7	116	2[3]	1	3[1]	12	8	212	2[3]	1	3[2]	13	9	308	2[3]	1	3[3]	14	10
21	2	1[1]	3	5	5	117	2	1[1]	3[1]	6	6	213	2	1[1]	3[2]	15	7	309	2	1[1]	3[3]	16	8
22	2[1]	1[1]	3	6	6	118	2[1]	1[1]	3[1]	7	7	214	2[1]	1[1]	3[2]	16	8	310	2[1]	1[1]	3[3]	17	9
23	2[2]	1[1]	3	3	7	119	2[2]	1[1]	3[1]	4	8	215	2[2]	1[1]	3[2]	13	9	311	2[2]	1[1]	3[3]	14	10
24	2[3]	1[1]	3	4	8	120	2[3]	1[1]	3[1]	5	9	216	2[3]	1[1]	3[2]	14	10	312	2[3]	1[1]	3[3]	15	11
25	2	1[2]	3	6	6	121	2	1[2]	3[1]	7	7	217	2	1[2]	3[2]	8	8	313	2	1[2]	3[3]	17	9
26	2[1]	1[2]	3	7	7	122	2[1]	1[2]	3[1]	8	8	218	2[1]	1[2]	3[2]	9	9	314	2[1]	1[2]	3[3]	18	10
27	2[2]	1[2]	3	8	8	123	2[2]	1[2]	3[1]	9	9	219	2[2]	1[2]	3[2]	10	10	315	2[2]	1[2]	3[3]	19	11
28	2[3]	1[2]	3	5	9	124	2[3]	1[2]	3[1]	6	10	220	2[3]	1[2]	3[2]	7	11	316	2[3]	1[2]	3[3]	16	12
29	2	1[3]	3	7	7	125	2	1[3]	3[1]	8	8	221	2	1[3]	3[2]	9	9	317	2	1[3]	3[3]	10	10
30	2[1]	1[3]	3	8	8	126	2[1]	1[3]	3[1]	9	9	222	2[1]	1[3]	3[2]	10	10	318	2[1]	1[3]	3[3]	11	11
31	2[2]	1[3]	3	9	9	127	2[2]	1[3]	3[1]	10	10	223	2[2]	1[3]	3[2]	11	11	319	2[2]	1[3]	3[3]	12	12
32	2[3]	1[3]	3	10	10	128	2[3]	1[3]	3[1]	11	11	224	2[3]	1[3]	3[2]	12	12	320	2[3]	1[3]	3[3]	13	13
33	1	2	3	8	4	129	1	2	3[1]	9	2	225	1	2	3[2]	10	6	321	1	2	3[3]	11	7
34	1[1]	2	3	9	5	130	1[1]	2	3[1]	10	6	226	1[1]	2	3[2]	11	7	322	1[1]	2	3[3]	12	8
35	1[2]	2	3	10	6	131	1[2]	2	3[1]	11	7	227	1[2]	2	3[2]	12	8	323	1[2]	2	3[3]	13	9
36	1[3]	2	3	11	7	132	1[3]	2	3[1]	12	8	228	1[3]	2	3[2]	13	9	324	1[3]	2	3[3]	14	10
37	1	2[1]	2	5	5	133	1	2[1]	2[1]	14	6	229	1	2[1]	2[2]	15	7	325	1	2[1]	2[3]	16	8
38	1[1]	2[1]	2	2	6	134	1[1]	2[1]	2[1]	11	7	230	1[1]	2[1]	2[2]	12	8	326	1[1]	2[1]	2[3]	13	9
39	1[2]	2[1]	2	3	7	135	1[2]	2[1]	2[1]	12	8	231	1[2]	2[1]	2[2]	13	9	327	1[2]	2[1]	2[3]	14	10
40	1[3]	2[1]	2	4	8	136	1[3]	2[1]	2[1]	13	9	232	1[3]	2[1]	2[2]	14	10	328	1[3]	2[1]	2[3]	15	11
41	1	2[2]	2	6	6	137	1	2[2]	2[1]	7	7	233	1	2[2]	2[2]	16	8	329	1	2[2]	2[3]	17	9
42	1[1]	2[2]	2	7	7	138	1[1]	2[2]	2[1]	8	8	234	1[1]	2[2]	2[2]	17	9	330	1[1]	2[2]	2[3]	18	10
43	1[2]	2[2]	2	4	8	139	1[2]	2[2]	2[1]	5	9	235	1[2]	2[2]	2[2]	14	10	331	1[2]	2[2]	2[3]	15	11
44	1[3]	2[2]	2	5	9	140	1[3]	2[2]	2[1]	6	10	236	1[3]	2[2]	2[2]	15	11	332	1[3]	2[2]	2[3]	16	12
45	1	2[3]	2	7	7	141	1	2[3]	2[1]	8	8	237	1	2[3]	2[2]	9	9	333	1	2[3]	2[3]	18	10
46	1[1]	2[3]	2	8	8	142	1[1]	2[3]	2[1]	9	9	238	1[1]	2[3]	2[2]	10	10	334	1[1]	2[3]	2[3]	19	11
47	1[2]	2[3]	2	9	9	143	1[2]	2[3]	2[1]	10	10	239	1[2]	2[3]	2[2]	11	11	335	1[2]	2[3]	2[3]	20	12
48	1[3]	2[3]	2	6	10	144	1[3]	2[3]	2[1]	7	11	240	1[3]	2[3]	2[2]	8	12	336	1[3]	2[3]	2[3]	17	13
49	3	1	2	4	8	145	3	1	2[1]	13	9	241	3	1	2[2]	14	10	337	3	1	2[3]	15	11
50	3[1]	1	2	1	9	146	3[1]	1	2[1]	10	10	242	3[1]	1	2[2]	11	11	338	3[1]	1	2[3]	12	12
51	3[2]	1	2	2	10	147	3[2]	1	2[1]	11	11	243	3[2]	1	2[2]	12	12	339	3[2]	1	2[3]	13	13
52	3[3]	1	2	3	11	148	3[3]	1	2[1]	12	12	244	3[3]	1	2[2]	13	13	340	3[3]	1	2[3]	14	14
53	3	1[1]	2	5	9	149	3	1[1]	2[1]	6	10	245	3	1[1]	2[2]	15	11	341	3	1[1]	2[3]	16	12
54	3[1]	1[1]	2	6	10	150	3[1]	1[1]	2[1]	7	11	246	3[1]	1[1]	2[2]	16	12	342	3[1]	1[1]	2[3]	17	13
55	3[2]	1[1]	2	3	11	151	3[2]	1[1]	2[1]	4	12	247	3[2]	1[1]	2[2]	13	13	343	3[2]	1[1]	2[3]	14	14
56	3[3]	1[1]	2	4	12	152	3[3]	1[1]	2[1]	5	13	248	3[3]	1[1]	2[2]	14	14	344	3[3]	1[1]	2[3]	15	15
57	3	1[2]	2	6	10	153	3	1[2]	2[1]	7	11	249	3	1[2]	2[2]	8	12	345	3	1[2]	2[3]	17	13
58	3[1]	1[2]	2	7	11	154	3[1]	1[2]	2[1]	8	12	250	3[1]	1[2]	2[2]	9	13	346	3[1]	1[2]	2[3]	18	14
59	3[2]	1[2]	2	8	12	155	3[2]	1[2]	2[1]	9	13	251	3[2]	1[2]	2[2]	10	14	347	3[2]	1[2]	2[3]	19	15
60	3[3]	1[2]	2	5	13	156	3[3]	1[2]	2[1]	6	14	252	3[3]	1[2]	2[2]	7	15	348	3[3]	1[2]	2[3]	16	16
61	3	1[3]	2	7	11	157	3	1[3]	2[1]	8	12	253	3	1[3]	2[2]	9	13	349	3	1[3]	2[3]	10	14
62	3[1]	1[3]	2	8	12	158	3[1]	1[3]	2[1]	9	13	254	3[1]	1[3]	2[2]	10	14	350	3[1]	1[3]	2[3]	11	15
63	3[2]	1[3]	2	9	13	159	3[2]	1[3]	2[1]	10	14	255	3[2]	1[3]	2[2]	11	15	351	3[2]	1[3]	2[3]	12	16
64	3[3]	1[3]	2	10	14	160	3[3]	1[3]	2[1]	11	15	256	3[3]	1[3]	2[2]	12	16	352	3[3]	1[3]	2[3]	13	17
65	2	3	1	8	8	161	2	3	1[1]	9	9	257	2	3	1[2]	10	10	353	2	3	1[3]	11	11
66	2[1]	3	1	9	9	162	2[1]	3	1[1]	10	10	258	2[1]	3	1[2]	11	11	354	2[1]	3	1[3]	12	12
67	2[2]	3	1	10	10	163	2[2]	3	1[1]	11	11	259	2[2]	3	1[2]	12	12	355	2[2]	3	1[3]	13	13
68	2[3]	3	1	11	11	164	2[3]	3	1[1]	12	12	260	2[3]	3									