

Swarming by curvature control in arbitrary dimension

P. Degond⁽¹⁾, A. Diez⁽²⁾, A. Frouvelle⁽³⁾

(1) Institut de Mathématiques de Toulouse ; UMR5219
Université de Toulouse ; CNRS
UPS, F-31062 Toulouse Cedex 9, France
email: pierre.degond@math.univ-toulouse.fr

(2) RIKEN Center for Interdisciplinary Theoretical and Mathematical Sciences (iTHEMS),
RIKEN iTHEMS, Wako, Saitama 351-0198, Japan
email: antoine.diez@riken.jp

(3) CEREMADE, CNRS ; UMR 7534
Université Paris Dauphine – PSL, 75016 Paris, France
email: frouvelle@ceremade.dauphine.fr

Abstract

We consider an interacting particle system proposed in the literature to model fish behavior. In this model, the agents move at constant speed and control the curvature of their trajectory (i.e. the time-derivative of their velocity) so as to align their velocity with that of their neighbors, up to some noise. We provide a novel n -dimensional formulation of this model for any $n \geq 3$ and derive its mean-field kinetic formulation using bundle geometry concepts. The target of the paper is the derivation of a fluid model in the hydrodynamic limit. We show that this fluid model is the “self-organized hydrodynamic” (SOH) model already found in earlier work pertaining to the Vicsek model. The derivation is based on the introduction of appropriate “generalized collision invariants” (GCI). The action of the n -dimensional orthogonal group is used to reduce the expression of the GCI to a set of two functions satisfying a system of equations which is solved by means of a variational formulation. This leads to explicit formulas for the coefficients of the SOH model in terms of those of the original interacting particle system.

Acknowledgements: PD holds a visiting professor association with the Department of Mathematics, Imperial College London, UK.

Data statement: No new data was generated in the course of this research.

Conflict of interest: The authors declare no conflict of interest.

Key words: Interacting particle system, kinetic equation, hydrodynamic model, tangent bundle, generalized collision invariant, group action, orthogonal group, variational formulation

AMS Subject classification: 35Q70, 35Q84, 36Q92, 82C31, 82C21, 82C22, 53B50, 57M60, 92D50

1 Introduction

Swarming is a feature shared by many living systems such as locust swarms [67], fish schools [61] or semen [28] to cite only a few. Systems of interacting motile individuals produce large scale structures such as density or orientation waves, cluster formation or oscillations. Such structures have large sizes in comparison with the interaction range of each agent. The mechanisms that generate them from local-scale agent interactions have been the object of intense scrutiny in the last three decades or so [73]. In spite of this, a lot of questions remain unanswered. Mathematical modelling provides a useful tool to unravel these mechanisms.

An assembly of interacting particles or agents can be modelled at the microscopic scale by a system of coupled ordinary or stochastic differential equations (ODEs, or SDEs) for the agents' locations and velocities for instance. However, particle systems do not provide much qualitative insight into the formation of large-scale structures except by means of intensive numerical simulations [26, 72]. The situation is drastically different with continuum (or macroscopic, or fluid) models which encode average properties of the system such as the local particle density or mean velocity. Such fluid models can be analyzed through e.g. stability analysis which provides orders of magnitudes of pattern sizes (see an application of this in e.g. [10]). However, for the analysis to be relevant, a rigorous link between the particle system and the fluid model must be available, including a “dictionary” linking the physical constants of the particle system and those of the fluid model. Such a dictionary is seldom provided.

One way to produce this dictionary is by means of the “hydrodynamic limit” (see e.g. [23, Sect. 3.8] in the case of rarefied gas dynamics). This method is based on a scale separation assumption, assuming that the time and space scales of the evolution of the system as a whole are larger than those associated to particle interactions. These assumptions align well with the goal of understanding the emergence of the large-scale structures appearing in collective dynamics. With such scaling assumptions, the particle system, or rather, its statistical description through mean-field kinetic equations, becomes singularly perturbed by a small parameter and the hydrodynamic model is obtained in the limit of this parameter tending to zero. Other methods of derivation of hydrodynamic models exist, such as the moment method (see e.g. [60]) but this method requires a closure assumption which is either arbitrary or phenomenological. By contrast, the hydrodynamic limit carries no assumption, but for the scaling assumptions made initially, which amount to select the applicability regime of the so-obtained hydrodynamic model.

Among models of collective dynamics, the Vicsek model [73] occupies a prominent position with many follow-ups (see among others, [12, 13, 26, 27, 40, 42, 47] and the review [73]). In this model, the particle speeds are supposed constant as a way to account for self-propulsion. Particles control their velocity (which is thus, up to scaling, a normalized vector) so as to align it with the average velocity of their neighbors, up to some noise reflecting the uncertainty in the control. A macroscopic version of the Vicsek model was proposed by Toner and Tu in [69, 70, 71]. This phenomenological model does not provide any link between its coefficients and those of the Vicsek model, thus failing to establish the expected “dictionary”. The hydrodynamic limit of the Vicsek model was performed for the first time in [35] (see also [1, 31, 32]) and leads to a novel type of fluid models called “Self-Organized Hydrodynamics (SOH)” [33] different from the Toner and Tu model [69].

The swarming model by curvature-control proposed in the present paper is an elaboration of the Vicsek model. Instead of the velocity, the variable by which the particles control their

motion is the time-derivative of the velocity (which is, up to scaling, equal to the curvature of the particles' trajectories, hence the terminology). Like in the Vicsek model, agents control their curvature so as to align their velocity with the average velocity of their neighbors. Again, noise is added to this curvature control to reflect uncertainties in the control. A preliminary version of this model for a single particle was originally proposed in 2 dimensions (2D) in [44] based on the analysis of fish trajectories. This single particle model was shown to behave like a diffusion at large scales in [36]. The complete, many-particle version of the curvature-control model was first proposed in 2D and its hydrodynamic limit, derived, in [37]. Strikingly, its hydrodynamic limit appeared to be the same SOH model as for the Vicsek model, except for the expression of its coefficients which took into account the specificities of curvature-control. Later, A three-dimensional (3D) version of the particle curvature-control model was proposed in [22].

In this paper, our goal is twofold. We first propose a n -dimensional (n D) version of the curvature-control model for arbitrary $n \geq 3$ (the case $n = 2$ being already treated in [37]). Compared with [37], this extension is, by far, non-trivial. Indeed, in 2D, the curvature is a scalar and the model is posed in standard flat space. In n D, ($n \geq 3$), the curvature is a normal vector to the velocity, so that the pair (velocity, curvature) lies on the tangent bundle to the sphere. It is thus necessary to use bundle geometry, which greatly complexifies the framework. In [22], this geometric complexity was overcome by the use of specific concepts to 3D (namely cross product), not generalizable to higher dimensions. The second goal is to perform the hydrodynamic limit of the curvature-control model in n D, to show that, like in 2D, the corresponding fluid model is the SOH model (in its n D version), and to give an explicit link between its coefficients and the particle curvature-control model. A fluid model based on the curvature-control model of [22] was proposed in [75] using truncated moment expansions. It is significantly different from the SOH model (these differences will be outlined in Section 4.3). While, in the curvature-control model, particle interactions are of mean-field type, a 2D model built from data [43] has features reminiscent to Boltzmann-like interactions, but otherwise similar in spirit.

Beside the Vicsek model and its variants, many other collective dynamics models have been proposed, such as the three-zones model [5, 20] or the Cucker-Smale model [1, 8, 21, 29, 50, 51, 52, 64] or the soft-core interaction model [39] to name only a few. Cucker-Smale and Vicsek type models can be connected via singular limits [9, 15, 16].

As already outlined, the curvature-control model in n D requires a bundle geometric framework. Indeed, the particle system must be appropriately formulated in the tangent bundle to the sphere. The mean-field kinetic description of the particle system is a partial differential equation (PDE) posed on this bundle, which also requires appropriate definitions of bundle differential operators. Classically, hydrodynamic limits strongly rely on appropriate conservation relations satisfied at the level of the mean-field kinetic model. However, here, there are not enough such conservation relations to conduct the hydrodynamic limit in this classical way. The same problem was already found in the Vicsek case, and was overcome by the introduction of a weaker form of conservation relation called “generalized collision invariant” (GCI) [1, 31, 32, 33, 35]. Here, the equations for the GCI are more complicated and do not lend themselves easily to analytic resolution. However, we show that the n -dimensional orthogonal group acts on the GCI and that the latter are invariant by one of its subgroups. This enables us to derive a reduced form of the GCI and to deduce explicit formulas for the coefficients of the resulting SOH model, hence producing the “dictionary” between the particle and fluid models as claimed. The GCI and their reduced form satisfy systems of PDE's which have novel variational formulations and

which are shown to be uniquely solvable.

To summarize, the main innovations contained in this work are

- a new particle curvature-control model in arbitrary dimension and its associated mean-field kinetic equations formulated in a bundle geometric framework,
- the first establishment of its associated hydrodynamic limit,
- the use of group action to reduce the complexity of a key object involved in the hydrodynamic limit, the “generalized collision invariant” (GCI),
- the derivation and unique solvability of a novel kind of variational formulation for the reduced GCI.

The outline of the paper is as follows. The particle curvature control model is presented in Section 2 and its mean-field kinetic formulation in Section 3. Then, the main result of the paper, the establishment of the hydrodynamic limit, is stated in Section 4. The hydrodynamic model consists of two equations. The first one, the continuity equation, is easily established in the same section. The second one, the velocity equation, requires the introduction of the GCI. This is the object of Section 5, where an existence and uniqueness theorem for the GCI is given. An action of the orthogonal group on the GCI is introduced in Section 6. Thanks to this action, a reduced form of the GCI consisting of two scalar functions (the reduced GCI pair) is derived in Section 7, and a variational formulation for the reduced GCI pair is introduced and studied. This finally enables us to derive the velocity equation in Section 8, which completes the proof of the main theorem. A conclusion is drawn in section 9. Appendix B provides the derivation of the reduced GCI pair, while Appendices A and C give proofs of auxiliary results stated in the main text. Bundle geometry concepts are gradually introduced when needed, in separate subsections at the beginning of Sections 2, 3, 4, 6 and 7.

2 The particle curvature-control (PCC) model

In this section, we derive the particle curvature-control (PCC) model which is the starting point of our study. We begin with an elementary presentation of the PCC model. Then, we introduce some geometric concepts which allow us to reformulate the PCC model in a geometric framework. The model is then scaled in dimensionless form.

2.1 Elementary presentation of the PCC system

We start by presenting the PCC model for a single particle before moving to a many-particle system. Then, we motivate the introduction of the geometric framework which will be fully developed in the forthcoming sections.

2.1.1 Single-particle PCC model

We start with the single-particle noiseless PCC model and describe the motion of a particle under a “force” that acts on the curvature of its trajectory. Let a particle have position $X(t) \in \mathbb{R}^n$ at time $t \in [0, \infty)$, where $n \geq 3$ is the spatial dimension (we exclude the case $n = 1$, for which the model does not make sense, and the case $n = 2$ which has been already treated in [37]). We assume

that the particle has given constant speed c_0 , i.e. its velocity dX/dt satisfies $|(dX/dt)(t)| = c_0$ at all times t , where $|\cdot|$ denotes the Euclidean norm in \mathbb{R}^n . Thus, we can write $dX/dt = c_0 \mathcal{V}$ where \mathcal{V} is the velocity direction (or equivalently, the unit vector tangent to the trajectory of the particle, oriented towards increasing times) satisfying

$$|\mathcal{V}| = 1. \quad (2.1)$$

In other words $\mathcal{V}(t) \in B$, $\forall t \in [0, \infty)$, where $B =: \mathbb{S}^{n-1}$ is the $n - 1$ -dimensional sphere. Now, we introduce the curvature of the particle trajectory $K(t)$ at time t . By definition, $K = d\mathcal{V}/ds$ is the derivative of the tangent vector \mathcal{V} with respect to the arclength s of the trajectory. Since the particle speed is c_0 and is constant, we have, up to a constant, $s = c_0 t$. Hence,

$$K = \frac{1}{c_0} \frac{d\mathcal{V}}{dt}. \quad (2.2)$$

The vector $K(t)$ belongs to \mathbb{R}^n but differentiating (2.1) with respect to t , we get

$$\mathcal{V} \cdot K = 0, \quad (2.3)$$

where \cdot denotes the Euclidean inner-product in \mathbb{R}^n . Hence, $K(t)$ is at all times orthogonal to the velocity $\mathcal{V}(t)$. Differentiating this relation once more, we get

$$\frac{dK}{dt} \cdot \mathcal{V} = -c_0 |K|^2. \quad (2.4)$$

This means that variations of the curvature cannot be arbitrary. Their projection along \mathcal{V} must equal $-c_0 |K|^2$. In particular, only the component of dK/dt normal to \mathcal{V} can be controlled by external phenomena.

Thus, let a time-dependent “force” $T(t)$ (where T stands for “torsion” of the trajectory, i.e. the derivative of its curvature) be such that

$$T \cdot \mathcal{V} = 0. \quad (2.5)$$

Then the single-particle motion is written

$$\begin{cases} \frac{dX}{dt} = c_0 \mathcal{V}, \end{cases} \quad (2.6)$$

$$\begin{cases} \frac{d\mathcal{V}}{dt} = c_0 K, \end{cases} \quad (2.7)$$

$$\begin{cases} \frac{dK}{dt} = T - c_0 |K|^2 \mathcal{V}. \end{cases} \quad (2.8)$$

We note that the constraints (2.1) and (2.3) are satisfied at all times by the solutions of this system, as soon as they are satisfied at time $t = 0$.

Now, we specify the choice of the force vector T in the single-particle PCC model. We assume that we are given a target direction \mathcal{J} (possibly depending on space and time) and that T is a relaxation force towards a fixed scalar multiple $\lambda > 0$ of the projection $P_{\mathcal{V}^\perp} \mathcal{J}$ of \mathcal{J} on the hyperplane orthogonal to \mathcal{V} , i.e.

$$T = -\nu (K - \lambda P_{\mathcal{V}^\perp} \mathcal{J}). \quad (2.9)$$

We recall that

$$P_{\mathcal{V}^\perp} \mathcal{J} = \mathcal{J} - (\mathcal{J} \cdot \mathcal{V}) \mathcal{V}. \quad (2.10)$$

The coefficient $\nu > 0$ is the relaxation rate. We notice that T given by (2.9) satisfies (2.5) as it should. We claim that the PCC model for single-particle dynamics (2.6)-(2.9) models the alignment of the velocity \mathcal{V} with the target vector \mathcal{J} . To show this, let us assume that \mathcal{J} is constant and uniform (i.e. does not depend on (x, t)). Then, the equation for (\mathcal{V}, K) decouples from that of X and we focus on the former. We get

$$\frac{d\mathcal{V}}{dt} = c_0 K, \quad \frac{dK}{dt} = -c_0 |\mathcal{V}|^2 \mathcal{V} - \nu(K - \lambda P_{\mathcal{V}^\perp} \mathcal{J}).$$

We further simplify the problem by considering a 2-dimensional motion (for illustration only, the core of the paper being concerned with dimensions $n \geq 3$). Hence, in an appropriate frame, we have

$$\mathcal{V}(t) = \begin{pmatrix} \cos \theta(t) \\ \sin \theta(t) \end{pmatrix}, \quad K(t) = \begin{pmatrix} -\sin \theta(t) \\ \cos \theta(t) \end{pmatrix} \kappa(t), \quad \mathcal{J} = \begin{pmatrix} |\mathcal{J}| \\ 0 \end{pmatrix},$$

thus defining $\theta(t)$ and $\kappa(t)$. Then, (2.47) reduces to

$$\frac{d\theta}{dt} = c_0 \kappa, \quad \frac{d\kappa}{dt} = -\nu(\kappa + \lambda |\mathcal{J}| \sin \theta), \quad (2.11)$$

or equivalently

$$\frac{d^2 \theta}{dt^2} + \nu \frac{d\theta}{dt} + \nu c_0 \lambda |\mathcal{J}| \sin \theta = 0.$$

This is the equation of a damped pendulum. In particular, if θ is small (i.e. if \mathcal{V} is almost aligned with \mathcal{J}), we can write

$$\frac{d^2 \theta}{dt^2} + \nu \frac{d\theta}{dt} + \nu c_0 \lambda |\mathcal{J}| \theta = 0,$$

whose solutions, according to the sign of $\nu - 4c_0 \lambda |\mathcal{J}|$, are combinations of real exponentials or damped sinusoids both tending to 0 as $t \rightarrow \infty$ (see Fig. 1 for illustrations). Hence, the large time behavior of the system shows that \mathcal{V} aligns to \mathcal{J} , as claimed. It also shows that, when ν is not large, the particles are wiggling a long time around the target direction before reaching it asymptotically.

Finally, we introduce noise in the single-particle PCC model. We assume that some Brownian noise with mean-square variance $2D$, with D a given constant positive diffusion coefficient, acts on the curvature K only. The reason for not adding Brownian noise to the velocity \mathcal{V} is that it would make \mathcal{V} non time-differentiable hence making it impossible to define the curvature K . The precise definition of what Brownian noise in curvature only means (given that the curvature depends on \mathcal{V} through the constraint (2.3)) depends on some geometric framework which will be described in forthcoming sections. At this stage, let us formally write the noisy single-particle PCC model as the following stochastic differential system:

$$\begin{cases} dX = c_0 \mathcal{V} dt, & (2.12) \\ d\mathcal{V} = c_0 K dt, & (2.13) \\ dK = (-\nu(K - \lambda P_{\mathcal{V}^\perp} \mathcal{J}) - c_0 |\mathcal{V}|^2 \mathcal{V}) dt + \sqrt{2D} dB_t, & (2.14) \end{cases}$$

where dB_t is a Brownian noise.




Figure 1: Examples of single-particle trajectories in space in 2D (blue curve) and vector \mathcal{J} (red arrow): (a) large alignment frequency $\nu > 4c_0\lambda|\mathcal{J}|$; (b) small alignment frequency $\nu < 4c_0\lambda|\mathcal{J}|$. Note that the Vicsek case corresponds to the overdamped limit of (2.11) and that the Vicsek trajectories would thus resemble case (a).

2.1.2 Many-particle PCC model

We now introduce the many-particle PCC model. Let N be the number of particles, and let the position, velocity direction and curvature of the i -th particle be given by the triple $(X_i(t), \mathcal{V}_i, K_i)$, where \mathcal{V}_i and K_i are constrained by (2.1), (2.3). In the many-particle PCC model (or in short, PCC model), each particle evolves independently through the single-particle PCC model introduced in the previous section, except that all the particle trajectories are coupled through the definition of the alignment vector \mathcal{J} . So, the trajectories of the particles are the solutions of the following system

$$\begin{cases} dX_i = c_0 \mathcal{V}_i dt, & (2.15) \\ d\mathcal{V}_i = c_0 K_i dt, & (2.16) \\ dK_i = \left(-\nu (K_i - \lambda P_{\mathcal{V}_i^\perp} \mathcal{J}(X_i(t), t)) - c_0 |K_i|^2 \mathcal{V}_i \right) dt + \sqrt{2D} dB_t^i, & (2.17) \end{cases}$$

for $i = 1, \dots, N$, where now $(dB_t^i)_{i=1, \dots, N}$ are N independent Brownian motions in K_i only (the precise meaning of this will be given below).

The vector $\mathcal{J}(x, t)$, which provides the coupling between the particles, is defined as the mean of the velocity directions of the particles located in a neighborhood of point x at time t . Hence, in (2.17), $\mathcal{J}(X_i(t), t)$ will describe the mean of the velocity directions of the neighboring particles to Particle i . This feature turns the PCC model into an interacting particle system. There are two variants of the model, according to whether \mathcal{J} is further normalized or not:

- **Non-normalized PCC model.** In the non-normalized version, the vector \mathcal{J} is simply the mean of the velocity directions \mathcal{V}_j of the Particles j located in a neighborhood of x at time t . It is defined by

$$\mathcal{J}(x, t) = \frac{1}{R^n} \sum_{\ell=1}^N \mathcal{K}\left(\frac{|X_\ell(t) - x|}{R}\right) \mathcal{V}_\ell(t), \quad (2.18)$$

where $\mathcal{K}: [0, \infty) \rightarrow [0, \infty)$ is a sensing function satisfying

$$\int_{\mathbb{R}^n} \mathcal{K}(|x|) dx = 1, \quad (2.19)$$

and $R > 0$ is the sensing radius. This sensing function weights the importance of the particles in the neighborhood of x . For instance, the simplest sensing function would be (up to normalization) the indicator function $\chi_{B(0,1)}$ of the ball $B(0, 1)$ centered at 0 with radius 1: $\mathcal{K}(s) \sim \chi_{B(0,1)}$. In this case, the sensing function \mathcal{K} in (2.18) would just select those particles located in the ball centered at $X_i(t)$ with radius R . However, we could imagine more sophisticated sensing functions, where the importance of particle j would be weighted by a function of the distance $|X_i(t) - X_j(t)|$.

- **Normalized PCC model.** In the normalized version, the vector \mathcal{J} is the normalized mean of the velocity directions \mathcal{V}_j of the Particles j located in a neighborhood of x at time t . In particular, this assumes that this mean is nowhere zero. The vector \mathcal{J} is given by:

$$\mathcal{J}(x, t) = \frac{\sum_{\ell=1}^N \mathcal{K}\left(\frac{|X_\ell(t) - x|}{R}\right) \mathcal{V}_\ell(t)}{\left| \sum_{\ell=1}^N \mathcal{K}\left(\frac{|X_\ell(t) - x|}{R}\right) \mathcal{V}_\ell(t) \right|}, \quad (2.20)$$

with \mathcal{K} satisfying the same assumptions as in the non-normalized case. Here, we must assume that the dynamics is defined only up to the first time (if it exists) where the denominator vanishes.

The non-normalized PCC model consists of Eqs. (2.15)-(2.17), (2.18), while the normalized PCC model, of Eqs. (2.15)-(2.17), (2.20). The two systems are solved supposing that an initial condition $(X_{i0}, \mathcal{V}_{i0}, K_{i0})_{i=1,\dots,N}$ is given.

We now restrict to the three-dimensional case $n = 3$ and establish that the non-normalized PCC model coincides with the model of [22]. Introduce an arbitrary coefficient $\delta > 0$ and define the new particle variables S_i (for “spin”) by

$$S_i = -\frac{1}{\delta} K_i \times \mathcal{V}_i, \quad (2.21)$$

where \times denotes the vector product in three dimensions. Then, we have the relations

$$K_i = \delta S_i \times \mathcal{V}_i, \quad S_i \cdot \mathcal{V}_i = 0. \quad (2.22)$$

From (2.21), (2.22), (2.16), (2.17) and (2.18), we deduce that the triples $(X_i, V_i, S_i)_{i=1}^N$ satisfy the system

$$\begin{cases} \frac{dX_i}{dt} = c_0 \mathcal{V}_i, \\ \frac{dV_i}{dt} = \frac{1}{\chi} S_i \times \mathcal{V}_i, \\ \frac{dS_i}{dt} = \mathcal{V}_i \times \left[J \sum_{j=1}^N n_{ij} \mathcal{V}_j - \eta \frac{d\mathcal{V}_i}{dt} + \xi_i \right], \end{cases} \quad (2.23)$$

$$(2.24)$$

$$(2.25)$$

with

$$\xi_i = \sqrt{(2d)\eta T} \frac{dB_t^i}{dt}, \quad (2.26)$$

and

$$\frac{1}{\chi} = c_0 \delta, \quad J = \frac{\nu \lambda}{\delta}, \quad n_{ij} = \frac{1}{R^3} \mathcal{K} \left(\frac{|X_i - X_j|}{R} \right), \quad \eta = \frac{\nu}{c_0 \delta}, \quad (2d)\eta T = \frac{2D}{\delta}.$$

Now, we can compare Equations (2.24)-(2.26) with Equations (1), (2) of [22] and notice that they are identical, apart from the scaling factor c_0 in (2.23), which can be easily removed by introducing $\mathcal{W}_i = c_0 \mathcal{V}_i$. We conclude that the normalized PCC model is identical with the model of [22] in dimension three. Hence, it can be viewed as an extension of [22] to any dimensions, noting that the former was already an extension of the two-dimensional model of [37].

2.1.3 A geometric view of the single-particle noiseless PCC model

We return to System (2.6)-(2.8) and introduce some geometric concepts. Suppose that $t \mapsto K(t)$, $[0, \infty) \rightarrow \mathbb{R}^n$ satisfies (2.3) but is not necessarily given by (2.2). Denote by $\Gamma = d\mathcal{V}/dt$ the particle acceleration and by $\mathcal{T} = dK/dt$ the time derivative of K . If we repeat the considerations of Section 2.1.1, we find that

$$\mathcal{V} \cdot \Gamma = 0 \quad \text{and} \quad \mathcal{T} \cdot \mathcal{V} + K \cdot \Gamma = 0. \quad (2.27)$$

Of course, if K is given by (2.2), we recover (2.4). We notice that, for a given pair (\mathcal{V}, K) satisfying (2.1), (2.3), the set of pairs $(\Gamma, \mathcal{T}) \in \mathbb{R}^{2n}$ satisfying (2.27) is a $2(n-1)$ -dimensional vector sub-space. Back to System (2.6) - (2.8), we can decompose

$$\begin{pmatrix} c_0 K \\ T - c_0 |K|^2 \mathcal{V} \end{pmatrix} = \begin{pmatrix} c_0 K \\ -c_0 |K|^2 \mathcal{V} \end{pmatrix} + \begin{pmatrix} 0 \\ T \end{pmatrix}. \quad (2.28)$$

We notice that both vectors at the right-hand side satisfy (2.27) (it is obvious for the former, while the latter requires (2.5)).

We now introduce some elements of geometry. The $(n-1)$ -dimensional sphere to which \mathcal{V} belongs is a differentiable manifold and for each \mathcal{V} , a vector K satisfying (2.3) is an element of the tangent space $T_{\mathcal{V}}B$ to B at \mathcal{V} . The set of pairs (\mathcal{V}, K) where $\mathcal{V} \in B$ and $K \in T_{\mathcal{V}}B$ is called the tangent bundle of B and will be denoted by M . The tangent space of M at (\mathcal{V}, K) , denoted by $T_{(\mathcal{V}, K)}M$ consists of those pairs (Γ, \mathcal{T}) satisfying (2.27). A vector field on B is a map $k: B \rightarrow M$ such that, for any $\mathcal{V} \in B$, $k(\mathcal{V}) \in T_{\mathcal{V}}B$. Like in classical analysis, vector fields can be interpreted as particle velocities which, after integration, give rise to trajectories on B . Likewise, a vector field on M is a map $\xi: M \rightarrow N$ (where N is the tangent bundle to M) such that for all $(\mathcal{V}, K) \in M$, $\xi(\mathcal{V}, K) \in T_{(\mathcal{V}, K)}M$. Hence, we see that the left-hand side of (2.28) is a vector field on M and that System (2.6)-(2.8) can be seen as a differential equation on $\mathbb{R}^n \times M$.

In the present work, it will be important to distinguish motions on M (i.e. vector fields on M) which are “parallel to \mathcal{V} ” or “parallel to K ”. For this purpose, we notice that both vectors at the right-hand side of (2.28) are vector fields on M . Furthermore, each of them depends on a

vector field on B (namely K for the first one and T for the second one). More generally, for any vector field Γ on B , we can construct the following vector field on M :

$$\mathcal{L}_{(\mathcal{V}, K)}^H \Gamma = \begin{pmatrix} \Gamma(\mathcal{V}) \\ -(\Gamma(\mathcal{V}) \cdot K)\mathcal{V} \end{pmatrix}. \quad (2.29)$$

We note that the right-hand side of (2.29) satisfies (2.27) and defines a valid vector field on M . This vector field will be called the horizontal lift of Γ at (\mathcal{V}, K) . Loosely speaking, horizontal lifts define motions on M in a direction “parallel to \mathcal{V} ”. We see that, moving parallel to \mathcal{V} imposes K to vary, as the second component of the vector (2.29) shows. This is because, as \mathcal{V} varies, K must vary too because of the constraint (2.3).

Likewise, for any vector field T on B , we can construct the following vector field on M :

$$\mathcal{L}_{(\mathcal{V}, K)}^V T = \begin{pmatrix} 0 \\ T(\mathcal{V}) \end{pmatrix}. \quad (2.30)$$

Again, we remark that the right-hand side of (2.30) satisfies (2.27) and defines a valid vector field on M . This vector field will be referred to as the vertical lift of T . Again, loosely speaking, vertical lifts define motions on M in a direction “parallel to K ”. By contrast to motions in the \mathcal{V} -direction, motions in the K -direction do not involve any variation in \mathcal{V} , as the vanishing first component of the vector (2.30) shows. This is because the tangent space $T_{\mathcal{V}}B$ along which K varies is a vector space, and moving along the tangent space $T_{\mathcal{V}}B$ does not change its base-point \mathcal{V} . Hence, we conclude that motions along \mathcal{V} and motions along K do not play a symmetric role, and this will be one of the difficulties which requires the introduction of a rigorous geometric framework.

Finally, with these definitions, (2.7)-(2.8) can be re-written

$$\frac{d}{dt} \begin{pmatrix} \mathcal{V} \\ K \end{pmatrix} = c_0 \mathcal{L}_{(\mathcal{V}, K)}^H K + \mathcal{L}_{(\mathcal{V}, K)}^V T. \quad (2.31)$$

In the next subsection, we introduce this geometric framework more rigorously.

2.2 Geometric preliminaries

This subsection recalls the geometric concepts that were mentioned in the previous section. We refer to [74] for an introduction to differential geometry, to [38, 41] for Riemannian geometry and to [46] for the geometry of bundles. We refer to Figure 2 for a depiction of the various sets and maps involved in the forthcoming description.

2.2.1 The tangent bundle to the sphere

We recall that we assume $n \geq 3$. We denote by $B = \mathbb{S}^{n-1}$ the $(n - 1)$ -dimensional sphere and by $F = \mathbb{R}^{n-1}$. We let $TB = (M, \pi, B, F)$ be the *tangent bundle* to B . In this quadruple, $M = \sqcup_{v \in B} T_v B$ is the support manifold to the vector bundle and consists of the disjoint union of the tangent spaces $T_v B$ to B at v , B is the base, F is the typical fiber and $\pi: M \rightarrow B$ is the base map. Since B is embedded in \mathbb{R}^n , M is embedded in \mathbb{R}^{2n} and we can write:

$$\alpha \in M \iff (\alpha = (v, \kappa) \in \mathbb{R}^{2n} \text{ such that } |v| = 1 \text{ and } \kappa \cdot v = 0), \quad (2.32)$$

where the symbols “ \cdot ” and “ $||$ ” are the Euclidean inner product and norm on \mathbb{R}^n . The notations v for elements of B and κ for elements of $T_v B$ stand for “velocity” and “curvature” respectively.

Now, we introduce the tangent bundle $TM = (N, \bar{\pi}, M, G)$ of M , where N is the support manifold of TM , $G = F^2 = \mathbb{R}^{2n-2}$ is the typical fiber and $\bar{\pi}: N \rightarrow M$ is the base map. Now, N is embedded in \mathbb{R}^{4n} . Let $\alpha = (v, \kappa) \in M$ (with (v, κ) satisfying (2.32)). Then, we have

$$\begin{aligned} \xi \in T_\alpha M &\iff (\xi = (a, \tau) \in \mathbb{R}^{2n} \text{ such that } a \cdot v = 0 \text{ and } \tau \cdot v + a \cdot \kappa = 0) \\ &\iff (\xi = (a, \bar{\tau} - (a \cdot \kappa)v) \text{ such that } a, \bar{\tau} \in \{v\}^\perp). \end{aligned} \quad (2.33)$$

The notations (a, τ) for the elements of $T_\alpha M$ respectively stand for “acceleration” and “torsion”.

The Euclidean structure of \mathbb{R}^n induces a *Riemannian structure* on B : the metric on $T_v B$ is defined by

$$\langle \kappa_1, \kappa_2 \rangle_v = \kappa_1 \cdot \kappa_2, \quad (2.34)$$

for all $\kappa_i \in T_v B = \{v\}^\perp$, $i = 1, 2$. We will drop the subscript v when the context is clear. The Riemannian structure on B determines the *Levi-Civita connection*, i.e. the unique torsion-free linear connection on B compatible with the Riemannian structure (here, torsion has its differential geometric meaning, and should not be confused with the torsion of a curve which has been used so far). Denote by $\mathcal{X}(B)$ the set of smooth vector fields on B and let $k, h \in \mathcal{X}(B)$. The Levi-Civita connection generates a new vector field $\nabla_k^B h$ (the gradient of h along the field k). In general, we will assume that $k \in \mathcal{X}(B)$ can be extended into a smooth function $\bar{k}: U_B \rightarrow \mathbb{R}^n$, where U_B is an open set of \mathbb{R}^n containing B . Then, thanks to [38, Chapt. 2, Exercise 3], the Levi-Civita connection is given by

$$\nabla_k^B h(v) = P_{v^\perp}((\bar{k} \cdot \nabla_v) \bar{h})(v), \quad (2.35)$$

We have denoted by ∇_v the usual nabla operator operating on functions of $v \in \mathbb{R}^n$ and by P_{v^\perp} the orthogonal projection $\mathbb{R}^n \rightarrow \{v\}^\perp$. Matrix-wise, $P_{v^\perp} = I_n - v \otimes v$ (where I_n is the identity matrix of \mathbb{R}^n and \otimes stands for the tensor product).

2.2.2 Vertical and horizontal bundle

Let $\alpha = (v, \kappa) \in M$. The space $T_\alpha M$ is decomposed into two components, which respectively represent motions “in the direction of v ” and “in the direction of κ ”. We first start with motions in the direction of κ and define the *vertical subspace* $V_\alpha = \ker(d\pi)_\alpha$, where $(d\pi)_\alpha: T_\alpha M \rightarrow T_v B$ is the differential of the base map π . For any $\xi = (a, \tau) \in T_\alpha M$ (with (a, τ) satisfying (2.33)), we have

$$(d\pi)_\alpha(\xi) = a, \quad (2.36)$$

and thus

$$\xi \in V_\alpha \iff (\xi = (0, \tau) \in \mathbb{R}^{2n} \text{ such that } \tau \cdot v = 0). \quad (2.37)$$

In particular, $\dim V_\alpha = n - 1$. For any $\alpha \in M$, the map $\mathcal{L}_\alpha^V: T_v B \rightarrow V_\alpha$ defined by

$$\mathcal{L}_\alpha^V(\tau) = (0, \tau), \quad \forall \tau \in T_v B, \quad (2.38)$$

is called the *vertical lift*. We note that

$$(d\pi)_\alpha \circ \mathcal{L}_\alpha^V = 0. \quad (2.39)$$

Now, we define directions of motion “in the v direction”. This requires the connection. We first define the horizontal lift. In general, a *horizontal lift* is a linear map $\mathcal{L}_\alpha^H: T_v B \rightarrow T_\alpha M$ such that

$$(d\pi)_\alpha \circ \mathcal{L}_\alpha^H = I_{T_v B}, \quad \forall \alpha = (v, \kappa) \in M, \quad (2.40)$$

where $I_{T_v B}$ is the identity of $T_v B$. Then, the *horizontal subspace* H_α associated with the horizontal lift \mathcal{L}_α^H is $H_\alpha = \text{Im } \mathcal{L}_\alpha^H$. The horizontal and vertical spaces are supplementary subspaces of $T_\alpha M$:

$$V_\alpha \oplus H_\alpha = T_\alpha M. \quad (2.41)$$

When B is endowed with a linear connection ∇^B , there exists a *unique horizontal map* \mathcal{L}^H associated with this connection [46, Vol. 2, Chapt. VII, Sect. 6, Prop. IX]. It can be shown that, for the sphere endowed with the Levi-Civita connection 2.35, we have

$$\mathcal{L}_\alpha^H(a) = (a, -(\kappa \cdot a)v), \quad \forall a \in T_v B. \quad (2.42)$$

and

$$\xi \in H_\alpha \iff (\xi = (a, \tau) \in \mathbb{R}^{2n} \text{ with } \tau = -(\kappa \cdot a)v) \quad (2.43)$$




Figure 2: Diagram of the sets and maps involved in the geometric framework. Arrows with a triple head mean surjections and arrows with an anchor are for injections.

2.3 Geometric presentation of the PCC system

In this subsection, we revisit the derivation of the PCC model introduced in Subsection 2.1 in light of the geometric concepts recalled in Subsection 2.2. Many different constructions of stochastic differential equations on manifolds and bundles have been proposed in the literature (see [18] for the deterministic case and [4, 11, 45, 48, 57, 59, 68] for the stochastic case). Our construction relies on [54] (see also [53]).

We first consider the noiseless single-particle PCC model. Let a single particle be characterized by its position $X \in \mathbb{R}^n$, and the pair $\mathcal{A} =: (\mathcal{V}, K) \in M$ consisting of the direction of its velocity $\mathcal{V} \in \mathbb{S}^{n-1}$ and the curvature vector of its trajectory $K \in \mathbb{R}^n$ such that (2.3) holds. The assumptions of the model have been detailed in Section 2.1. The single-particle noiseless PCC model is written according to (2.31), which can be re-written in terms of $\mathcal{A} \in M$ according to

$$\begin{cases} \frac{dX}{dt} = c_0 \mathcal{V}, \\ \frac{d\mathcal{A}}{dt} = c_0 \mathcal{L}_{\mathcal{A}}^H K - \nu \mathcal{L}_{\mathcal{A}}^V (K - \lambda P_{\mathcal{V}^\perp}(\mathcal{J}(X, t))) \end{cases} \quad (2.44)$$

$$\frac{d\mathcal{A}}{dt} = c_0 \mathcal{L}_{\mathcal{A}}^H K - \nu \mathcal{L}_{\mathcal{A}}^V (K - \lambda P_{\mathcal{V}^\perp}(\mathcal{J}(X, t))) \quad (2.45)$$

To pass to the noisy single-particle PCC model, we need to generate Brownian noise on M which acts on the curvature only. For this purpose, we generate Brownian noise on B and then, lift it with the vertical lift to generate Brownian noise on the vertical subspace $V_{\mathcal{A}}$ of $T_{\mathcal{A}}M$. We know that Brownian noise on B at \mathcal{V} can be obtained by projecting Brownian noise on \mathbb{R}^n onto $T_{\mathcal{V}}B$ using the Stratonovich construction [53, 54]. Let $(\mathbf{e}_1, \dots, \mathbf{e}_n)$ be the canonical basis of \mathbb{R}^n . Then, Brownian noise in the K variable only with variance $2D$ is written $(2D)^{1/2} \sum_{j=1}^n \mathcal{L}_{\mathcal{A}}^V(P_{\mathcal{V}^\perp} \mathbf{e}_j) \circ dB_{j,t}$ where $(dB_{j,t})_{j=1, \dots, n}$ are n independent Brownian motions on \mathbb{R} and the symbol \circ indicates that the equation is meant in the Stratonovich sense. Hence, the noisy single-particle PCC model reads:

$$\begin{cases} dX = c_0 \mathcal{V} dt, \\ d\mathcal{A} = c_0 \mathcal{L}_{\mathcal{A}}^H K dt - \nu \mathcal{L}_{\mathcal{A}}^V (K - \lambda P_{\mathcal{V}^\perp}(\mathcal{J}(X, t))) dt + \sqrt{2D} \sum_{j=1}^n \mathcal{L}_{\mathcal{A}}^V (P_{\mathcal{V}^\perp} \mathbf{e}_j) \circ dB_{j,t}. \end{cases} \quad (2.46)$$

$$d\mathcal{A} = c_0 \mathcal{L}_{\mathcal{A}}^H K dt - \nu \mathcal{L}_{\mathcal{A}}^V (K - \lambda P_{\mathcal{V}^\perp}(\mathcal{J}(X, t))) dt + \sqrt{2D} \sum_{j=1}^n \mathcal{L}_{\mathcal{A}}^V (P_{\mathcal{V}^\perp} \mathbf{e}_j) \circ dB_{j,t}. \quad (2.47)$$

Finally, the noisy many-particle PCC model (or PCC model in short) is a direct generalization of the previous model. Using the notations of Section 2.1.2, this system reads

$$\begin{cases} dX_i = c_0 \mathcal{V}_i dt, \\ d\mathcal{A}_i = c_0 \mathcal{L}_{\mathcal{A}_i}^H K_i dt - \nu \mathcal{L}_{\mathcal{A}_i}^V (K_i - \lambda P_{\mathcal{V}_i^\perp}(\mathcal{J}(X_i, t))) dt + \sqrt{2D} \sum_{j=1}^n \mathcal{L}_{\mathcal{A}_i}^V (P_{\mathcal{V}_i^\perp} \mathbf{e}_j) \circ dB_{j,t}^i, \end{cases} \quad (2.48)$$

$$d\mathcal{A}_i = c_0 \mathcal{L}_{\mathcal{A}_i}^H K_i dt - \nu \mathcal{L}_{\mathcal{A}_i}^V (K_i - \lambda P_{\mathcal{V}_i^\perp}(\mathcal{J}(X_i, t))) dt + \sqrt{2D} \sum_{j=1}^n \mathcal{L}_{\mathcal{A}_i}^V (P_{\mathcal{V}_i^\perp} \mathbf{e}_j) \circ dB_{j,t}^i, \quad (2.49)$$

where now $(dB_{j,t}^i)_{j=1, \dots, n}^{i=1, \dots, N}$ are nN independent Brownian motions in \mathbb{R} . The vector \mathcal{J} is defined by (2.18) in the non-normalized case and by (2.20) in the normalized case.

2.4 Scaling of the PCC model

2.4.1 Scaling in the normalized case

In this case, \mathcal{J} is dimensionless, so, the parameter λ has the physical dimension of the inverse of a length. We will use λ^{-1} as a spatial scale. Since c_0 is a speed, $(c_0 \lambda)^{-1}$ will be used as a time scale. We introduce the following changes of variables and unknowns:

$$X_i = \lambda^{-1} \tilde{X}_i, \quad t = (c_0 \lambda)^{-1} \tilde{t}, \quad K_i = \lambda \tilde{K}_i, \quad (2.50)$$

which defines dimensionless variables and unknowns $\tilde{X}_i, \tilde{t}, \tilde{K}_i$. We note that the unknown \mathcal{V}_i is already dimensionless and does not require any scaling. The scaling on K_i requires that we introduce a scaled variable $\tilde{\mathcal{A}}_i = (\tilde{\mathcal{V}}_i, \tilde{K}_i)$ which is obtained from the unscaled variable $\mathcal{A}_i = (\mathcal{V}_i, K_i)$ by setting

$$\mathcal{A}_i = \Sigma_\lambda(\tilde{\mathcal{A}}_i) \quad \text{or equivalently} \quad \tilde{\mathcal{A}}_i = \Sigma_{\lambda^{-1}}(\mathcal{A}_i). \quad (2.51)$$

where for $\lambda > 0$, the scaling map $\Sigma_\lambda: M \rightarrow M$, is defined by $\Sigma_\lambda(v, \kappa) = (v, \lambda\kappa)$, $\forall \alpha = (v, \kappa) \in M$. The scaling map satisfies

$$(d\Sigma_\lambda)_\alpha \circ \mathcal{L}_\alpha^H = \mathcal{L}_{\Sigma_\lambda \alpha}^H, \quad (d\Sigma_\lambda)_\alpha \circ \mathcal{L}_\alpha^V = \lambda \mathcal{L}_{\Sigma_\lambda \alpha}^V. \quad (2.52)$$

We now let $(\tilde{X}_i, \tilde{\mathcal{A}}_i)$ be functions of the rescaled time \tilde{t} and define

$$\tilde{\mathcal{J}}(\tilde{x}, t) = \mathcal{J}(\lambda^{-1}\tilde{x}, (c_0\lambda)^{-1}\tilde{t}).$$

We introduce three scaled constants

$$\bar{v} = \frac{v}{c_0\lambda}, \quad \bar{D} = \frac{D}{c_0\lambda^3}, \quad \bar{R} = \lambda R. \quad (2.53)$$

Then, using (2.52), we find that the normalized PCC model in the scaled variable $(\tilde{X}_i, \tilde{\mathcal{A}}_i)$ and the scaled time \tilde{t} is given by (dropping the tildes and the bars for clarity) :

$$\begin{cases} dX_i = \mathcal{V}_i dt, \end{cases} \quad (2.54)$$

$$\begin{cases} d\mathcal{A}_i = \mathcal{L}_{\mathcal{A}_i}^H K_i dt - v \mathcal{L}_{\mathcal{A}_i}^V (K_i - P_{\mathcal{V}_i^\perp}(\mathcal{J}(X_i, t))) dt + \sqrt{2D} \sum_{j=1}^n \mathcal{L}_{\mathcal{A}_i}^V (P_{\mathcal{V}_i^\perp} e_j) \odot dB_{j,t}^i, \end{cases} \quad (2.55)$$

with \mathcal{J} given by

$$\mathcal{J}(x, t) = \frac{\sum_{\ell=1}^N \mathcal{K}\left(\frac{|X_\ell(t) - x|}{R}\right) \mathcal{V}_\ell(t)}{\left| \sum_{\ell=1}^N \mathcal{K}\left(\frac{|X_\ell(t) - x|}{R}\right) \mathcal{V}_\ell(t) \right|}. \quad (2.56)$$

2.4.2 Scaling in the non-normalized case

We assume that \mathcal{K} is dimensionless. Let x_0 be a spatial scale. \mathcal{J} has the dimension of the inverse of a volume, i.e., x_0^{-n} while K_i has dimension x_0^{-1} . From the second term at the right-hand side of (2.49), λ has the dimension of x_0^{n-1} . Thus, we can set the length scale $x_0 = \lambda^{1/(n-1)}$. So, we introduce the scaling

$$X_i = \lambda^{\frac{1}{n-1}} \tilde{X}_i, \quad t = \lambda^{\frac{1}{n-1}} c_0^{-1} \tilde{t}, \quad K_i = \lambda^{-\frac{1}{n-1}} \tilde{K}_i, \quad \mathcal{J} = \lambda^{-\frac{n}{n-1}} \tilde{\mathcal{J}}, \quad (2.57)$$

so defining the dimensionless variables and unknowns $\tilde{X}_i, \tilde{t}, \tilde{K}_i, \tilde{\mathcal{J}}$. We also define the following dimensionless constants:

$$\bar{v} = v \lambda^{\frac{1}{n-1}} c_0^{-1}, \quad \bar{D} = D \lambda^{\frac{3}{n-1}} c_0^{-1}, \quad \bar{R} = R \lambda^{-\frac{1}{n-1}}. \quad (2.58)$$

Then, the non-normalized PCC model in the scaled variables $(\tilde{X}_i, \tilde{\mathcal{A}}_i)$ and the scaled time \tilde{t} is given by (2.54), (2.55) (again dropping the tildes and the bars for clarity) with \mathcal{J} given by

$$\mathcal{J}(x, t) = \frac{1}{R^n} \sum_{\ell=1}^N \mathcal{K}\left(\frac{|X_\ell(t) - x|}{R}\right) \mathcal{V}_\ell(t). \quad (2.59)$$

3 The kinetic curvature-control (KCC) model

In this section, we formally establish the kinetic mean-field model, below referred to as the kinetic curvature-control model (KCC model), associated with the PCC model. We first develop additional geometric concepts that are needed to achieve this goal. We then proceed to the derivation of the KCC model and conclude this section by presenting its hydrodynamic rescaling.

3.1 Geometric preliminaries

Again, we refer to Figure 2 for an illustration of the concepts described in this section and to [38, 41, 46, 74] for geometric background.

3.1.1 Connection map

In this section, we define additional operators linked to the horizontal and vertical lifts. The vertical (resp. horizontal) projection is the projection \mathcal{P}_α^V (resp. \mathcal{P}_α^H) of $T_\alpha M$ on V_α parallel to H_α (resp. on H_α parallel to V_α). They are given by

$$\mathcal{P}_\alpha^V \xi = (0, \tau + (\kappa \cdot a)v) = (0, \tau - (v \cdot \tau)v) = (0, P_{v^\perp} \tau), \quad (3.1)$$

$$\mathcal{P}_\alpha^H \xi = (a, -(\kappa \cdot a)v), \quad (3.2)$$

for all $\xi = (a, \tau)$ in $T_\alpha M$ (i.e. satisfying (2.33)). Next, we note that the vertical and horizontal lifts are isomorphism: $\mathcal{L}_\alpha^V: T_{\pi\alpha} B \rightarrow V_\alpha$ and $\mathcal{L}_\alpha^H: T_{\pi\alpha} B \rightarrow H_\alpha$. Their inverses are denoted by $(\mathcal{L}_\alpha^V)^{-1}$ and $(\mathcal{L}_\alpha^H)^{-1}$ and given by

$$(\mathcal{L}_\alpha^V)^{-1} \xi = \tau, \quad \forall \xi \in V_\alpha \quad \text{i.e.} \quad \forall \xi = (0, \tau) \quad \text{with} \quad \tau \cdot v = 0. \quad (3.3)$$

$$(\mathcal{L}_\alpha^H)^{-1} \xi = a, \quad \forall \xi \in H_\alpha \quad \text{i.e.} \quad \forall \xi = (a, -(a \cdot \kappa)v) \quad \text{with} \quad a \cdot v = 0. \quad (3.4)$$

Then, we define B_α^H and $B_\alpha^V: T_\alpha M \rightarrow T_v B$ by

$$B_\alpha^V = (\mathcal{L}_\alpha^V)^{-1} \circ \mathcal{P}_\alpha^V, \quad B_\alpha^H = (\mathcal{L}_\alpha^H)^{-1} \circ \mathcal{P}_\alpha^H. \quad (3.5)$$

Thanks to (2.40), we have

$$B_\alpha^H = (d\pi)_\alpha, \quad (3.6)$$

and thus, B_α^H is given by (2.36). We note that B_α^H does not depend on the connection. On the other hand, the map B_α^V does, and is called the connection map. We have

$$B_\alpha^V \xi = P_{v^\perp} \tau, \quad \forall \xi = (a, \tau) \quad \text{satisfying} \quad (2.33). \quad (3.7)$$

3.1.2 Riemannian structure on M

We endow M with the Riemannian structure induced by the *Sasaki metric* [49, 66] and [38, Chap. 3, Exercise 2]. For $\alpha \in M$ and $\xi_i \in T_\alpha M$, $i = 1, 2$, the Sasaki metric on $T_\alpha M$ is defined by

$$\langle\langle \xi_1, \xi_2 \rangle\rangle_\alpha = \langle B_\alpha^H \xi_1, B_\alpha^H \xi_2 \rangle_{\pi\alpha} + \langle B_\alpha^V \xi_1, B_\alpha^V \xi_2 \rangle_{\pi\alpha}. \quad (3.8)$$

Writing $\alpha = (v, \kappa) \in \mathbb{R}^{2n}$ and $\xi_i = (a_i, \tau_i) \in \mathbb{R}^{2n}$, with (v, κ) satisfying (2.32) and (a_i, τ_i) satisfying (2.33), the Sasaki metric is written:

$$\langle\langle \xi_1, \xi_2 \rangle\rangle_\alpha = a_1 \cdot a_2 + P_{v^\perp} \tau_1 \cdot P_{v^\perp} \tau_2. \quad (3.9)$$

We note that this metric is **not** the metric induced by the embedding $M \rightarrow \mathbb{R}^{2n}$. Indeed, the latter would add a term $(a_1 \cdot \kappa)(a_2 \cdot \kappa)$ to the right-hand side of (3.9). The vertical and horizontal subspaces are orthogonal for the Sasaki metric:

$$V_\alpha^\perp \oplus H_\alpha = T_\alpha M, \quad \forall \alpha \in M.$$

and thus, the projection operators \mathcal{P}_α^V and \mathcal{P}_α^H are self-adjoint with respect to the Sasaki metric. The Sasaki metric on M is linked with the metric on B by

$$\langle\langle \mathcal{L}_\alpha^H \kappa, \xi \rangle\rangle_\alpha = \langle \kappa, B_\alpha^H \xi \rangle_{\pi\alpha}, \quad \langle\langle \mathcal{L}_\alpha^V \kappa, \xi \rangle\rangle_\alpha = \langle \kappa, B_\alpha^V \xi \rangle_{\pi\alpha}, \quad (3.10)$$

for all $\alpha = (v, \kappa) \in M$, $\xi \in T_\alpha M$ and $\kappa \in T_v B$. Consequently, B_α^H and \mathcal{L}_α^H are mutually adjoint and so are B_α^V and \mathcal{L}_α^V .

3.1.3 Integration on M

We define the positive orientation of \mathbb{R}^n as that given by the canonical basis $(\mathbf{e}_1, \dots, \mathbf{e}_n)$, with $\mathbf{e}_i = (0, \dots, 0, 1, 0, \dots, 0)$, where the 1 is the i -th position. We choose the positive orientation of B such that an orthogonal basis $(\varepsilon_1, \dots, \varepsilon_{n-1})$ of $T_v B$ is direct if and only if $(v, \varepsilon_1, \dots, \varepsilon_{n-1})$ is a direct basis of \mathbb{R}^n . The volume form Vol_B associated with the Riemannian structure induced by the embedding of B into \mathbb{R}^n is given as follows: Let $(\varepsilon_1, \dots, \varepsilon_{n-1})$ be a local, positively oriented orthonormal frame. Then, the volume form Vol_B on B has expression

$$\text{Vol}_B(v)(\kappa_1, \dots, \kappa_{n-1}) = \det(\langle \varepsilon_i, \kappa_j \rangle_v), \quad \forall (\kappa_1, \dots, \kappa_{n-1}) \in (T_v B)^{n-1}. \quad (3.11)$$

We define the following $n-1$ -form on M for all $(\xi_1, \dots, \xi_{n-1}) \in (T_\alpha M)^{n-1}$ by

$$\Omega(\alpha)(\xi_1, \dots, \xi_{n-1}) = \text{Vol}_B(\pi\alpha)(B_\alpha^V(\xi_1), \dots, B_\alpha^V(\xi_{n-1})). \quad (3.12)$$

Then,

$$\text{Vol}_M = \pi^* \text{Vol}_B \wedge \Omega. \quad (3.13)$$

(where $\pi^* \text{Vol}_B$ denotes the pullback of the form Vol_B of B to a form on M by the bundle base map π) is a volume form on M associated with the Sasaki metric and we choose the orientation of M provided by this form. We will denote by $|\text{Vol}_B|$ and $|\text{Vol}_M|$ the densities associated with the volume forms on B and M respectively.

Following [46, Vol. 1, Ch. 7], for an integrable function $f: M \rightarrow \mathbb{R}$, we can integrate $f\Omega$ along the fibers \mathbb{R}^{n-1} as follows

$$\left(\int_{\mathbb{R}^{n-1}} f \Omega \right)(v) =: \int_{T_v B} f \Omega_v, \quad (3.14)$$

where Ω_v is a $n - 1$ -form on $T_v B$ called the retrenchment of Ω on $T_v B$ and defined for all $\kappa, \eta_1, \dots, \eta_{n-1}$ in $T_v B$ by

$$\Omega_v(\kappa)(\eta_1, \dots, \eta_{n-1}) = \Omega(\alpha)(\mathcal{L}_\alpha^V \eta_1, \dots, \mathcal{L}_\alpha^V \eta_{n-1}) = \text{Vol}_B(v)(\eta_1, \dots, \eta_n),$$

with $\alpha = (v, \kappa)$ and using (3.12). Then, the following Fubini theorem holds true

$$\int_M f \text{Vol}_M = \int_B \left(\int_{\mathbb{R}^{n-1}} f \Omega \right) \text{Vol}_B. \quad (3.15)$$

The same relations are true replacing the volume forms by their densities. Let $(\varepsilon_1, \dots, \varepsilon_{n-1})$ be a local, positively oriented orthonormal frame of B . Define a coordinate map $T_v B \rightarrow \mathbb{R}^{n-1}$, $\kappa \mapsto (\kappa^1, \dots, \kappa^{n-1})$ with $\kappa^i = \langle \kappa, \varepsilon_i \rangle_v$. It maps isometrically $T_v B$ onto \mathbb{R}^{n-1} . Then, in the coordinates $(\kappa^1, \dots, \kappa^{n-1})$, the retrenchment Ω_v on $T_v B$ has expression

$$\Omega_v = d\kappa^1 \wedge \dots \wedge d\kappa^{n-1}. \quad (3.16)$$

3.1.4 Differential operators on M

Let $f: M \rightarrow \mathbb{R}$ be a smooth function. We recall that the gradient of f is the vector field $\nabla_\alpha f \in \mathcal{X}(M)$ on M defined by

$$\langle \nabla_\alpha f(\alpha), \xi \rangle_\alpha = (df)_\alpha(\xi), \quad \forall \alpha \in M, \quad \forall \xi \in T_\alpha M. \quad (3.17)$$

Below, we will define partial gradients of functions on M with respect to v or to κ separately, denoted by $\nabla_v f$ and $\nabla_\kappa f$ respectively. The index α permits to remember that $\nabla_\alpha f$ is the gradient of f in the classical sense (i.e. with respect to the pair $\alpha = (v, \kappa)$). Let $\Xi \in \mathcal{X}(M)$ be a smooth vector field. The divergence of Ξ is the scalar field $\nabla_\alpha \cdot \Xi$ defined by

$$\int_M \nabla_\alpha \cdot \Xi f |\text{Vol}_M| = - \int_M \langle \Xi, \nabla_\alpha f \rangle |\text{Vol}_M|, \quad (3.18)$$

for all smooth functions $f: M \rightarrow \mathbb{R}$. In other words, gradient and divergence are formal anti-adjoint to each other with respect to the L^2 inner-product. The index α in $\nabla_\alpha \cdot \Xi$ serves the same purpose as for the gradient: it will help us to distinguish divergences with respect to the pair $\alpha = (v, \kappa)$ from partial divergences with respect to v or to κ separately, denoted by $\nabla_v \cdot \Xi$ and $\nabla_\kappa \cdot \Xi$ respectively. We recall an alternate formula for the divergence [41, §4.8]. Let $\Xi \in \mathcal{X}(\mathcal{M})$ and $D\Xi$ be the linear map $\mathcal{X}(\mathcal{M}) \rightarrow \mathcal{X}(\mathcal{M})$, $Y \mapsto \nabla_Y^M \Xi$, where ∇^M is the Levi-Civita connection associated with the Sasaki metric. Then,

$$\nabla_\alpha \cdot \Xi = \text{Tr } D\Xi. \quad (3.19)$$

We now define the horizontal (i.e. “in the v direction”) and vertical (i.e. “in the κ direction”) gradients and divergences which we will denote by using the symbols ∇_v and ∇_κ respectively. The gradients map smooth functions $f: M \rightarrow \mathbb{R}$ to smooth vector fields Ξ such that $\Xi(\alpha) \in T_{\pi\alpha} B$. Such vector fields are smooth sections of the pullback bundle π^*TB . Specifically, the pullback bundle $\pi^*TB = (\tilde{M}, \tilde{\pi}, M, F)$ is such that $\tilde{M} = \sqcup_{\alpha \in M} T_{\pi\alpha} B$ and for any $(\alpha, \kappa) \in \tilde{M}$

with $\kappa \in T_{\pi\alpha}B$, we have $\tilde{\pi}(\alpha, \kappa) = \alpha$. The space of smooth sections of π^*TB will be denoted by $\mathcal{Y}(M)$. The horizontal (resp. vertical) gradients $\nabla_v f$ (resp. $\nabla_\kappa f$) are defined by

$$\nabla_v f(\alpha) = B_\alpha^H(\nabla_\alpha f(\alpha)), \quad \nabla_\kappa f(\alpha) = B_\alpha^V(\nabla_\alpha f(\alpha)), \quad (3.20)$$

and $\nabla_v f, \nabla_\kappa f \in \mathcal{Y}(M)$. The horizontal (resp. vertical) divergences $\nabla_v \cdot \Xi$ (resp. $\nabla_\kappa \cdot \Xi$) of a field $\Xi \in \mathcal{Y}(M)$ are smooth functions $M \rightarrow \mathbb{R}$ defined by

$$\nabla_v \cdot \Xi = \nabla_\alpha \cdot (\mathcal{L}^H \Xi), \quad \nabla_\kappa \cdot \Xi = \nabla_\alpha \cdot (\mathcal{L}^V \Xi). \quad (3.21)$$

Thanks to (3.10), gradient and divergence are formal L^2 -anti-duals, i.e.

$$\int_M \nabla_v \cdot \Xi f |\text{Vol}_M| = - \int_M \langle \Xi, \nabla_v f \rangle_{\pi\alpha} |\text{Vol}_M|, \quad (3.22)$$

and similarly by replacing ∇_v by ∇_κ . Classical vector calculus identities carry over to ∇_v and ∇_κ , i.e.

$$\nabla_v(fg) = f\nabla_v g + g\nabla_v f, \quad \nabla_v \cdot (f\Xi) = (\nabla_v f, \Xi)_v + f\nabla_v \cdot \Xi. \quad (3.23)$$

and similarly by replacing ∇_v by ∇_κ .

3.1.5 Remarkable formulas

In this section, we give various useful lemmas whose proofs can be found in Appendix A.

Lemma 3.1 (Divergences of a field which does not depend on κ).

Let $k \in \mathcal{X}(B)$ and denote by $\nabla \cdot k$ its divergence. We define an element of $\mathcal{Y}(M)$ by $\tilde{k} = k \circ \pi$. Such elements of $\mathcal{Y}(M)$ can be viewed as “not depending on κ ”. Then, we have

$$\nabla_v \cdot \tilde{k} = \nabla \cdot k, \quad \nabla_\kappa \cdot \tilde{k} = 0. \quad (3.24)$$

Lemma 3.2 (Gradient of a function which only depends on v).

(i) Let $\varphi: B \rightarrow \mathbb{R}$ be smooth and let $\nabla \varphi \in \mathcal{X}(B)$ be its gradient. Define $\tilde{\varphi}: M \rightarrow \mathbb{R}$ by $\tilde{\varphi} = \varphi \circ \pi$. Then, we have

$$\nabla_v \tilde{\varphi}(\alpha) = \nabla \varphi(\pi\alpha), \quad \nabla_\kappa \tilde{\varphi}(\alpha) = 0, \quad \forall \alpha \in M. \quad (3.25)$$

(ii) Conversely, let $\psi: M \rightarrow \mathbb{R}$ be smooth and such that $\nabla_\kappa \psi \equiv 0$. Then, there exists $\varphi: B \rightarrow \mathbb{R}$, smooth, such that $\psi = \varphi \circ \pi$.

Lemma 3.3 (Gradient of $\langle \kappa, \kappa \rangle_v$).

We consider the function $M \rightarrow \mathbb{R}$, $\alpha \mapsto \langle \kappa, \kappa \rangle_v$, with $\alpha = (v, \kappa)$. Then, we have

$$\nabla_\kappa \langle \kappa, \kappa \rangle_v = 2\kappa, \quad \nabla_v \langle \kappa, \kappa \rangle_v = 0. \quad (3.26)$$

3.1.6 Symplectic structure, Hamiltonian and Poisson bracket

There is a symplectic structure on M which combines the symplectic structure on the cotangent bundle of B (see e.g. [6]) with the metric. We define a one-form $\omega^{(1)}$ on M as follows. Let $\alpha = (v, \kappa) \in M$. Let $\xi \in T_\alpha M$. Then,

$$\omega^{(1)}(\alpha)(\xi) = \langle \kappa, (d\pi)_\alpha(\xi) \rangle_v. \quad (3.27)$$

The symplectic form $\omega^{(2)}$ on TM is given by

$$\omega^{(2)} = d\omega^{(1)}. \quad (3.28)$$

Thanks to the non-degeneracy of $\omega^{(2)}$, we can define a linear isomorphism $I_\alpha: T_\alpha M^* \rightarrow T_\alpha M$, $\sigma \rightarrow I_\alpha(\sigma)$ as follows:

$$\sigma(\xi) = \omega^{(2)}(\alpha)(\xi, I_\alpha(\sigma)), \quad \forall \xi \in T_\alpha M. \quad (3.29)$$

Let $\mathbf{H}: M \times (0, \infty) \rightarrow \mathbb{R}$, $(\alpha, t) \mapsto \mathbf{H}(\alpha, t)$ be a smooth function which will be called the Hamiltonian. The Hamiltonian vector field with Hamiltonian \mathbf{H} is the vector field $I(d_\alpha \mathbf{H})$, where d_α stands for the derivative with respect to α for fixed t . A Hamiltonian flow is the flow of a Hamiltonian vector field, i.e. the flow associated with a differential equation of the form $d\alpha/dt = I(d_\alpha \mathbf{H}(\cdot, t))(\alpha(t))$.

As an example, let us consider the Hamiltonian $\mathbf{H}_J: M \rightarrow \mathbb{R}$ given by

$$\mathbf{H}_J(\alpha) = -v \cdot \mathcal{J} + \frac{1}{2} \langle \kappa, \kappa \rangle_v, \quad \alpha = (v, \kappa) \in M, \quad (3.30)$$

where $\mathcal{J} \in \mathbb{R}^n$ is a given vector. Then, we have

$$\mathcal{L}_{\mathcal{A}_i}^H K_i + v \mathcal{L}_{\mathcal{A}_i}^V P_{v_i^\perp} \mathcal{J} = I(d\mathbf{H}_{J(X_i(t), t)})(\mathcal{A}_i(t)), \quad (3.31)$$

where $d\mathbf{H}_{J(X_i(t), t)}$ denotes the derivative of the function $M \rightarrow \mathbb{R}$, $\alpha \mapsto \mathbf{H}_{J(X_i(t), t)}$, for fixed t . Thus, (3.31) is a Hamiltonian vector field with Hamiltonian $\mathbf{H}_{J(X_i(t), t)}$. Hence, Equation (2.55) for \mathcal{A}_i in the PCC model can be written

$$d\mathcal{A}_i = I(d\mathbf{H}_{J(X_i(t), t)})(\mathcal{A}_i(t)) dt - v \mathcal{L}_{\mathcal{A}_i}^V K_i dt + \sqrt{2\bar{D}} \sum_{j=1}^n \mathcal{L}_{\mathcal{A}_i}^V (P_{v_i^\perp} e_j) \circ dB_{j,t}^i. \quad (3.32)$$

In this form, we notice that the first term at the right-hand side of (3.32) is Hamiltonian while the last two terms are dissipative. We will notice a similar structure on the kinetic model and this structure will give rise to several important properties of the model.

For this purpose, it is convenient to introduce the Poisson bracket associated with the symplectic form $\omega^{(2)}$. It is defined for two smooth functions f and $h: M \rightarrow \mathbb{R}$ by

$$\{f, h\} = \omega^{(2)}(I(dh), I(df)) = df(I(dh)), \quad (3.33)$$

and is again a smooth function $M \rightarrow \mathbb{R}$. The Poisson bracket is an anticommutative bilinear form on the space of smooth functions that satisfies the Jacobi identity and the Leibnitz rule. We can show that

$$I(d\mathbf{H}) = \mathbb{J} \nabla_\alpha \mathbf{H}, \quad \mathbb{J} = \mathcal{L}^H \circ \mathcal{B}^V - \mathcal{L}^V \circ \mathcal{B}^H. \quad (3.34)$$

and consequently

$$\{f, h\} = \langle \nabla_v f, \nabla_\kappa h \rangle - \langle \nabla_\kappa f, \nabla_v h \rangle. \quad (3.35)$$

By Liouville's theorem, $\omega^{(2)}$ is preserved by a Hamiltonian flow (including the case where this Hamiltonian is time-dependent) and so, any exterior power of $\omega^{(2)}$ is preserved as well. The n -th exterior power $(\omega^{(2)})^{\wedge n}$ is equal to the Riemannian volume form Vol_M up to a constant, namely

$$(\omega^{(2)})^{\wedge n} = n! (-1)^{\frac{n(n+1)}{2}} \text{Vol}_M. \quad (3.36)$$

Thus, the flow of the vector field $\mathbb{J}\nabla_\alpha \mathbf{H}$ on M , where \mathbf{H} is any smooth Hamiltonian: $M \times (0, \infty) \rightarrow \mathbb{R}$, $(\alpha, t) \mapsto \mathbf{H}(\alpha, t)$, preserves the Sasaki volume form. It implies that $\mathbb{J}\nabla_\alpha \mathbf{H}$ is divergence free. Hence, we have

$$\nabla_v \cdot \nabla_\kappa \mathbf{H} - \nabla_\kappa \cdot \nabla_v \mathbf{H} = 0. \quad (3.37)$$

It results alternate expressions of the bracket of two smooth functions f and h :

$$\{f, h\} = \nabla_v \cdot (f \nabla_\kappa h) - \nabla_\kappa \cdot (f \nabla_v h) = \nabla_\kappa \cdot (h \nabla_v f) - \nabla_v \cdot (h \nabla_\kappa f). \quad (3.38)$$

From this and (3.22) (and its analog when ∇_v is changed in ∇_κ), we deduce that if g is another smooth function $M \rightarrow \mathbb{R}$, we have

$$\int_M \{f, h\} g |\text{Vol}_M| = \int_M f \{h, g\} |\text{Vol}_M|. \quad (3.39)$$

In particular, if $g = 1$, we find

$$\int_M \{f, h\} |\text{Vol}_M| = 0. \quad (3.40)$$

3.2 Kinetic model

We first derive the KCC model in the single-particle case. We then successively present the many-particle KCC models in the non-normalized and normalized cases.

3.2.1 Single particle KCC model

For the single-particle PCC model, applying [54, Theorem V.1.2], we get

Theorem 3.4 (Kinetic model for the single particle PCC model).

Suppose $\mathcal{J}: \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}^n$, $(x, t) \mapsto \mathcal{J}(x, t)$ is given and smooth. Consider the single-particle PCC model, given by (2.54), (2.55) with $i = N = 1$ associated with this choice of \mathcal{J} . Let $(X(t), \mathcal{A}(t))$ be a trajectory of this stochastic differential system with initial condition (X_0, \mathcal{A}_0) drawn according to the probability distribution $f_0 |dx \wedge \text{Vol}_M|$, where $f_0: \mathbb{R}^n \times M \rightarrow [0, \infty)$ is a given integrable function. Then, the probability distribution $f(\cdot, \cdot, t) |dx \wedge \text{Vol}_M|$ of $(X(t), \mathcal{A}(t))$ at time t (where $f(\cdot, \cdot, t): \mathbb{R}^n \times M \rightarrow [0, \infty)$ is an integrable function) satisfies the following kinetic equation:

$$\partial_t f + \nabla_x \cdot (v f) + \nabla_v \cdot (\kappa f) - v \nabla_\kappa \cdot ((\kappa - P_{v^\perp} \mathcal{J}) f) - D \Delta_\kappa f = 0, \quad (3.41)$$

with $\Delta_\kappa f =: \nabla_\kappa \cdot (\nabla_\kappa f)$ being the so-called vertical Laplacian of f on M . Equation (3.41) will be referred to as the kinetic curvature-control (KCC) model.

Remark 3.1. Eq. (3.41) has the same form as if (v, κ) belonged to a standard linear space. However, we should keep in mind that the meaning of the operators $\nabla_v \cdot$ and $\nabla_\kappa \cdot$ is given by (3.21).

Let \mathbf{H}_J be the Hamiltonian given by (3.30). Thanks to (3.25) and (3.26), we have

$$\nabla_v \mathbf{H}_J = -v P_{v^\perp} J, \quad \nabla_\kappa \mathbf{H}_J = \kappa. \quad (3.42)$$

Thus, in view of (3.35), the kinetic equation (3.41) can be written

$$\partial_t f + \nabla_x \cdot (v f) + \{f, \mathbf{H}_J\} - v \nabla_\kappa \cdot (\kappa f) - D \Delta_\kappa f = 0. \quad (3.43)$$

Eq. (3.41) is of Fokker-Planck type, i.e. the second-order operator is a partial diffusion which operates on the κ -variable only and not on the x and v variables. Fokker-Planck equations on manifolds have previously been considered e.g. in [14, 19, 48, 58, 65]. However, the specific Fokker-Planck equation (3.41) posed on the cartesian product of \mathbb{R}^n and the tangent bundle to the sphere \mathbb{S}^{n-1} has not been studied previously, to the best of our knowledge.

3.2.2 Many particle KCC (non-normalized case)

We consider the PCC model (2.54), (2.55), (2.59) in the non-normalized case. We associate it with its empirical measure μ_t^N on $\mathbb{R}^n \times M$ such that

$$\mu_t^N(x, \alpha) = \frac{1}{N} \sum_{i=1}^N \delta_{(X_i(t), \mathcal{A}_i(t))}(x, \alpha), \quad x \in \mathbb{R}^n, \quad \alpha \in M,$$

and we denote by μ_0^N the empirical measure associated with the initial data $((X_{i0}, \mathcal{A}_{i0}))_{i=1, \dots, N}$. We now consider μ_t^N and μ_0^N as random measures. An adaptation of classical results on mean-field limits (see e.g. the review [24, 25]) would show the following:

Theorem 3.5 (Kinetic model for the many-particle PCC model (non-normalized case)).

Let μ_0^N converge as $N \rightarrow \infty$ in probability sense towards a deterministic absolutely continuous probability measure $f_0 |dx \wedge \text{Vol}_M|$. Then, for any $t \in [0, \infty)$, μ_t^N converges in probability sense towards a deterministic absolutely continuous probability measure $f_t |dx \wedge \text{Vol}_M|$ and $f(x, \alpha, t) \equiv f_t(x, \alpha)$ satisfies (3.41) with $J = J_f(x, t)$ given by:

$$J_f(x, t) = \frac{1}{R^n} \int_{\mathbb{R}^n \times M} \mathcal{K}\left(\frac{|x - y|}{R}\right) v f(y, \alpha, t) |dx \wedge \text{Vol}_M|(y, \alpha). \quad (3.44)$$

We do not precise the functional setting and refer to [24, 25] for details.

3.2.3 Many particle KCC (normalized case)

We consider the particle system (2.54), (2.55), (2.56) in the normalized case. The normalization brings a major difficulty to prove a rigorous convergence result from the particle dynamics to the kinetic model and has not been solved yet, even for the Vicsek model, except when the singularity brought by the normalization is regularized [17]. Here, we will assume that the following is true

Conjecture 3.1 (Kinetic model for the many-particle PCC model (normalized case)).

With the same assumptions as for Theorem 3.5, and in the generic case where the denominator of (2.56) does not vanish, μ_t^N converges towards a deterministic absolutely continuous probability measure $f_t |dx \wedge \text{Vol}_M|$ and $f(x, \alpha, t) \equiv f_t(x, \alpha)$ satisfies (3.41) with $\mathcal{J} = \mathcal{J}_f(x, t)$ given by:

$$\mathcal{J}_f(x, t) = \frac{\int_{\mathbb{R}^{n \times M}} \mathcal{K}\left(\frac{|x-y|}{R}\right) v f(y, \alpha, t) |dx \wedge \text{Vol}_M|(y, \alpha)}{\left| \int_{\mathbb{R}^{n \times M}} \mathcal{K}\left(\frac{|x-y|}{R}\right) v f(y, \alpha, t) |dx \wedge \text{Vol}_M|(y, \alpha) \right|}. \quad (3.45)$$

3.3 Hydrodynamic scaling

In the remainder of this paper, **we focus on the normalized KCC model** and derive its hydrodynamic limit. Indeed, the non-normalized model leads to additional difficulties stemming from the existence of phase transitions which will be dealt with in future work.

In the hydrodynamic scaling, temporal and spatial phenomena are supposed to occur at large scales compared with the dynamics of the α variable. To express this, as classical in kinetic theory [23, Sect. 3.8], we change the time and space scales and introduce the following change of variables:

$$\hat{x} = \varepsilon x, \quad \hat{t} = \varepsilon t, \quad \text{with} \quad \varepsilon \ll 1.$$

At the same time, we assume that the trajectory curvatures are small at the microscopic scale, so that they are order unity at the macroscopic scale. For that reason, we do not rescale κ . In the new variables \hat{x}, \hat{t} , the unknowns f and \mathcal{J} are parametrized by ε and will be denoted by $f^\varepsilon, \mathcal{J}^\varepsilon$. Now, the equations for $f^\varepsilon, \mathcal{J}^\varepsilon$ are given in the normalized case by (dropping the hats on \hat{x}, \hat{t}):

$$\partial_t f^\varepsilon + \nabla_x \cdot (v f^\varepsilon) + \frac{1}{\varepsilon} (\{f^\varepsilon, \mathbf{H}_{\mathcal{J}^\varepsilon}\} - v \nabla_\kappa \cdot (\kappa f^\varepsilon) + D \nabla_\kappa f^\varepsilon) = 0, \quad (3.46)$$

with $\mathcal{J}_f^\varepsilon$ given for any $f(x, \alpha, t)$ by

$$\mathcal{J}_f^\varepsilon(x, t) = \frac{\int_{\mathbb{R}^{n \times M}} \mathcal{K}\left(\frac{|x-y|}{\varepsilon R}\right) v f(y, \alpha, t) |dx \wedge \text{Vol}_M|(y, \alpha)}{\left| \int_{\mathbb{R}^{n \times M}} \mathcal{K}\left(\frac{|x-y|}{\varepsilon R}\right) v f(y, \alpha, t) |dx \wedge \text{Vol}_M|(y, \alpha) \right|}, \quad (3.47)$$

We expand (3.47) in powers of ε . In view of (2.19), straightforward calculations give

$$\mathcal{J}_f^\varepsilon = \mathbf{u}_f + \mathcal{O}(\varepsilon^2), \quad (3.48)$$

with

$$\mathbf{u}_f(x, t) = \frac{\int_M v f(x, \alpha, t) |\text{Vol}_M|(\alpha)}{\left| \int_M v f(x, \alpha, t) |\text{Vol}_M|(\alpha) \right|}, \quad (3.49)$$

We only keep the leading order in (3.48) as the following orders are not involved in the limit $\varepsilon \rightarrow 0$ which we are interested in.

Finally, we are left with the following perturbation problem:

$$\partial_t f^\varepsilon + \nabla_x \cdot (v f^\varepsilon) + \frac{1}{\varepsilon} \left(\{f^\varepsilon, \mathbf{H}_{\mathbf{u}_{f^\varepsilon}}\} - \nu \nabla_\kappa \cdot (\kappa f^\varepsilon) + D \nabla_\kappa f^\varepsilon \right) = 0, \quad (3.50)$$

with \mathbf{u}_f given by (3.49). We will refer to this system as the scaled KCC model. Letting $\varepsilon \rightarrow 0$ in (3.50) is referred to as the hydrodynamic limit. The main result of this paper is the determination of the limit. It will be useful to introduce the collision operator

$$Q(f) = -\{f, \mathbf{H}_{\mathbf{u}_f}\} + \nu \nabla_\kappa \cdot (\kappa f) + D \Delta_\kappa f, \quad (3.51)$$

so that (3.50) can be simply written

$$\partial_t f^\varepsilon + \nabla_x \cdot (v f^\varepsilon) = \frac{1}{\varepsilon} Q(f^\varepsilon). \quad (3.52)$$

We note that Q is nonlinear due to the dependence of \mathbf{u} on f .

4 The hydrodynamic model

In this section, we derive the equilibria of the collision operator. Then, we state the main theorem of this paper which establishes that the scaled KCC model (3.52) converges to a hydrodynamic-type model (referred to as the “self-organized hydrodynamic” model (SOH)) in the limit $\varepsilon \rightarrow 0$. Finally, we show that the first equation of the SOH model, the continuity equation is easily derived from the KCC model. But first of all, we introduce a useful geometric change of variables.

4.1 Geometric preliminary: a change of variables in B

We denote by $B_{\mathbf{u}} = B \setminus \{\pm \mathbf{u}\}$. We identify $\mathbb{S}^{n-2} \approx B \cap \{\mathbf{u}\}^\perp$. We introduce the diffeomorphism $\mathbf{V}_{\mathbf{u}}: (0, \pi) \times \mathbb{S}^{n-2} \rightarrow B_{\mathbf{u}}$, $(\theta, w) \mapsto \mathbf{V}_{\mathbf{u}}(\theta, w)$, defined by

$$\mathbf{V}_{\mathbf{u}}(\theta, w) = v = \cos \theta \mathbf{u} + \sin \theta w, \quad (4.1)$$

where

$$\cos \theta = \mathbf{u} \cdot v, \quad |w| = 1, \quad \mathbf{u} \cdot w = 0. \quad (4.2)$$

The diffeomorphism $\mathbf{V}_{\mathbf{u}}$ is depicted in dimension $n = 3$ in Fig. 3 of Section 7.1.1 below. We orient \mathbb{S}^{n-2} in such a way that $(\varepsilon_1, \dots, \varepsilon_{n-2})$ is a direct orthonormal basis of $T_w \mathbb{S}^{n-2}$ if and only if $(e_\theta, \varepsilon_1, \dots, \varepsilon_{n-2})$ is a direct orthonormal basis of $T_v \mathbb{S}^{n-1}$, where

$$e_\theta = -\sin \theta \mathbf{u} + \cos \theta w. \quad (4.3)$$

The volume form $\text{Vol}_{\mathbb{S}^{n-2}}$ on \mathbb{S}^{n-2} is defined analogously to Vol_B , i.e. it is the Riemannian volume form for the Riemannian structure induced by the embedding of \mathbb{S}^{n-2} in \mathbb{R}^n . Now, we have (the proof is left to the reader)

$$|\text{Vol}_B| = \sin^{n-2} \theta d\theta |\text{Vol}_{\mathbb{S}^{n-2}}|. \quad (4.4)$$

4.2 Equilibria

Let $\mathbf{u} \in \mathbb{S}^{n-1}$. We define the (generalized) von Mises distribution $\mathbf{M}_{\mathbf{u}}: M \rightarrow \mathbb{R}$ by

$$\mathbf{M}_{\mathbf{u}} = \frac{1}{Z} \exp\left(-\frac{\nu}{D} \mathbf{H}_{\mathbf{u}}\right), \quad (4.5)$$

with Z such that

$$\int_M \mathbf{M}_{\mathbf{u}} |\text{Vol}_M| = 1. \quad (4.6)$$

Lemma 4.1. *The normalization factor Z is given by*

$$Z = \left(\frac{2\pi D}{\nu}\right)^{\frac{n-1}{2}} |\mathbb{S}^{n-2}| \int_0^\pi \exp\left(\frac{\nu^2}{D} \cos \theta\right) \sin^{n-2} \theta \, d\theta, \quad (4.7)$$

(where $|\mathbb{S}^{n-2}|$ denotes the $n-2$ -dimensional measure of \mathbb{S}^{n-2}) and does not depend on \mathbf{u} .

Proof. Using Fubini's theorem (3.15), together with (3.14), (3.16) and (3.30), we have

$$\begin{aligned} Z &= \int_M \exp\left(-\frac{\nu}{D} \mathbf{H}_{\mathbf{u}}(\alpha)\right) |\text{Vol}_M|(\alpha) \\ &= \int_B \exp\left(\frac{\nu^2}{D} (v \cdot \mathbf{u})\right) \left(\int_{\mathbb{R}^{n-1}} \exp\left(-\frac{\nu}{2D} \sum_{i=1}^{n-1} (\kappa^i)^2\right) d\kappa^1 \dots d\kappa^{n-1} \right) |\text{Vol}_B|(v) \\ &= \left(\frac{2\pi D}{\nu}\right)^{\frac{n-1}{2}} \int_B \exp\left(\frac{\nu^2}{D} v \cdot \mathbf{u}\right) |\text{Vol}_B|(v). \end{aligned}$$

Now, we use the diffeomorphism described in Section 4.1 and thanks to (4.4), we get

$$\int_B \exp\left(\frac{\nu^2}{D} v \cdot \mathbf{u}\right) |\text{Vol}_B|(v) = |\mathbb{S}^{n-2}| \int_0^\pi \exp\left(\frac{\nu^2}{D} \cos \theta\right) \sin^{n-2} \theta \, d\theta,$$

which leads to (4.7). ■

Thanks to the von Mises distribution, we can write the collision operator as follows

$$Q(f) = -\{f, \mathbf{H}_{\mathbf{u}_f}\} + D \nabla_\kappa \cdot \left(\mathbf{M}_{\mathbf{u}_f} \nabla_\kappa \left(\frac{f}{\mathbf{M}_{\mathbf{u}_f}} \right) \right). \quad (4.8)$$

Lemma 4.2 (Negativity of Q).

Let $f: M \rightarrow \mathbb{R}$ be smooth and decaying fast enough at infinity in κ . Then, we have

$$\int_M Q(f) \frac{f}{\mathbf{M}_{\mathbf{u}_f}} |\text{Vol}_M| = -D \int_M \left\langle \nabla_\kappa \left(\frac{f}{\mathbf{M}_{\mathbf{u}_f}} \right), \nabla_\kappa \left(\frac{f}{\mathbf{M}_{\mathbf{u}_f}} \right) \right\rangle \mathbf{M}_{\mathbf{u}_f} |\text{Vol}_M| \leq 0. \quad (4.9)$$

Proof. Using (3.22), we have

$$\int_M \nabla_\kappa \cdot \left(\mathbf{M}_{\mathbf{u}_f} \nabla_\kappa \left(\frac{f}{\mathbf{M}_{\mathbf{u}_f}} \right) \right) \frac{f}{\mathbf{M}_{\mathbf{u}_f}} |\text{Vol}_M| = - \int_M \left\langle \nabla_\kappa \left(\frac{f}{\mathbf{M}_{\mathbf{u}_f}} \right), \nabla_\kappa \left(\frac{f}{\mathbf{M}_{\mathbf{u}_f}} \right) \right\rangle \mathbf{M}_{\mathbf{u}_f} |\text{Vol}_M|.$$

On the other hand, using the fact that the Poisson bracket is a derivation as well as (4.5), we can write

$$\begin{aligned} \int_M \{f, \mathbf{H}_{\mathbf{u}_f}\} \frac{f}{\mathbf{M}_{\mathbf{u}_f}} |\text{Vol}_M| &= \frac{Z}{2} \int_M \{f^2, \mathbf{H}_{\mathbf{u}_f}\} \exp\left(\frac{\nu}{D} \mathbf{H}_{\mathbf{u}_f}\right) |\text{Vol}_M| \\ &= \frac{DZ}{2\nu} \int_M \left\{ f^2, \exp\left(\frac{\nu}{D} \mathbf{H}_{\mathbf{u}_f}\right) \right\} |\text{Vol}_M| = 0, \end{aligned}$$

thanks to (3.40). Inserting the expression (4.8) for the collision operator at the left-hand side of (4.9) and using the above computation leads to the right-hand side of (4.9) and ends the proof. \blacksquare

We introduce the coefficient c_1 called the “order parameter”, defined by:

$$c_1 = \frac{\int_B \exp\left(\frac{\nu^2}{D} v \cdot \mathbf{u}\right) (v \cdot \mathbf{u}) |\text{Vol}_B|(v)}{\int_B \exp\left(\frac{\nu^2}{D} v \cdot \mathbf{u}\right) |\text{Vol}_B|(v)} = \frac{\int_0^\pi \exp\left(\frac{\nu^2}{D} \cos \theta\right) \cos \theta \sin^{n-2} \theta d\theta}{\int_0^\pi \exp\left(\frac{\nu^2}{D} \cos \theta\right) \sin^{n-2} \theta d\theta}, \quad (4.10)$$

which does not depend on \mathbf{u} .

Lemma 4.3 (Properties of the order parameter).

(i) $c_1 \in [0, 1)$.

(ii) For $\mathbf{u} \in \mathbb{S}^{n-1}$, we have

$$\int_M \mathbf{M}_{\mathbf{u}}(\alpha) v |\text{Vol}_M|(\alpha) = c_1 \frac{\mathbf{u}}{|\mathbf{u}|}. \quad (4.11)$$

Proof. This is a classical result which has been proved in the literature [31]. \blacksquare

Proposition 4.4 (Equilibria).

Let $f: M \rightarrow \mathbb{R}$ be smooth and decaying fast enough at infinity in κ . Then, we have

$$Q(f) = 0, \quad (4.12)$$

if and only if $\exists \rho \geq 0, \exists \mathbf{u} \in \mathbb{S}^{n-1}$ such that

$$f = \rho \mathbf{M}_{\mathbf{u}}, \quad (4.13)$$

Distributions of the type (4.13) are called “equilibria”.

Proof. Let f be such that $Q(f) = 0$. Then, by (4.9), we have

$$0 = \int_M Q(f) \frac{f}{\mathbf{M}_{\mathbf{u}_f}} |\text{Vol}_M| = -D \int_M \left\langle \nabla_\kappa \left(\frac{f}{\mathbf{M}_{\mathbf{u}_f}} \right), \nabla_\kappa \left(\frac{f}{\mathbf{M}_{\mathbf{u}_f}} \right) \right\rangle \mathbf{M}_{\mathbf{u}_f} |\text{Vol}_M|.$$

Since the integrated quantity in the last integral is nonnegative and $\mathbf{M}_{\mathbf{u}_f} > 0$, this implies $\nabla_\kappa(f/\mathbf{M}_{\mathbf{u}_f}) = 0$, hence, by Lemma 3.2 (ii), there exists a function $\varphi: B \rightarrow \mathbb{R}$ such that

$$f = \mathbf{M}_{\mathbf{u}_f} \varphi \circ \pi.$$

Then, we compute, thanks to (3.25),

$$\begin{aligned} \nabla_v f &= \mathbf{M}_{\mathbf{u}_f} (\nabla_v \varphi) \circ \pi - \frac{\nu}{D} \mathbf{M}_{\mathbf{u}_f} \varphi \circ \pi \nabla_v \mathbf{H}_{\mathbf{u}_f}, \\ \nabla_\kappa f &= -\frac{\nu}{D} \mathbf{M}_{\mathbf{u}_f} \varphi \circ \pi \nabla_\kappa \mathbf{H}_{\mathbf{u}_f}. \end{aligned}$$

Thus,

$$Q(f) = -\{f, \mathbf{H}_{\mathbf{u}_f}\} = -\mathbf{M}_{\mathbf{u}_f} \langle \kappa, (\nabla_v \varphi) \circ \pi \rangle.$$

Since $Q(f) = 0$, we have $\langle \kappa, (\nabla_v \varphi) \circ \pi \rangle = 0$, $\forall \alpha \in M$, which implies $\nabla_v \varphi = 0$, hence, φ is a constant on B . Therefore, there exists a constant $\rho \in \mathbb{R}$ such that $f = \rho \mathbf{M}_{\mathbf{u}_f}$. Since $f \geq 0$, $\rho \in [0, \infty)$. Consequently, f is of the form (4.13). Conversely, suppose f is given by (4.13) for arbitrary $\mathbf{u} \in \mathbb{S}^{n-1}$ and $\rho \in [0, \infty)$. Then, computing \mathbf{u}_f using (3.49) with the help of (4.10), (4.11) and of the fact that c_1 is non-negative leads to the fact that any $\mathbf{u}_f = \mathbf{u}$, which shows that such f is a solution of (4.12). This ends the proof. \blacksquare

4.3 Hydrodynamic limit: main result

The main result of this paper is the following

Theorem 4.5 (Hydrodynamic limit of the KCC model (formal result)).

Suppose f^ε is a smooth solution of the scaled KCC model (3.52), which converges as $\varepsilon \rightarrow 0$ towards f as smoothly as needed. Then, there exist two functions $\rho: \mathbb{R}^n \times [0, \infty) \rightarrow (0, \infty)$, $(x, t) \mapsto \rho(x, t)$ and $\mathbf{u}: \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{S}^{n-1}$, $(x, t) \mapsto \mathbf{u}(x, t)$ such that

$$f(x, \alpha, t) = \rho(x, t) \mathbf{M}_{\mathbf{u}(x, t)}(\alpha), \quad \forall (x, \alpha, t) \in \mathbb{R}^n \times M \times [0, \infty). \quad (4.14)$$

Furthermore, ρ and \mathbf{u} satisfy the following “self-organized hydrodynamic (SOH)” system

$$\partial_t \rho + \nabla_x \cdot (c_1 \rho \mathbf{u}) = 0. \quad (4.15)$$

$$\rho (\partial_t \mathbf{u} + c_2 (\mathbf{u} \cdot \nabla_x) \mathbf{u}) + c_3 P_{\mathbf{u}^\perp} \nabla_x \rho = 0, \quad (4.16)$$

with c_1 given by (4.10), c_3 , by

$$c_3 = \frac{D}{\nu^2}. \quad (4.17)$$

To define c_2 , we first introduce $\mathbf{M}_{\mathbf{u}}^\natural: [0, \pi] \times \mathbb{R} \times [0, \infty) \rightarrow \mathbb{R}$, given by

$$\mathbf{M}_{\mathbf{u}}^\natural = \mathbf{M}_{\mathbf{u}}^\natural(\theta, \kappa_\parallel, \kappa_\perp) = \frac{1}{Z} \exp \left(-\frac{\nu}{D} \left(-\nu \cos \theta + \frac{1}{2} (\kappa_\parallel^2 + \sin^2 \theta \kappa_\perp^2) \right) \right), \quad (4.18)$$

and the solution $(\zeta_w, \zeta_r), [0, \pi] \times \mathbb{R} \times [0, \infty) \rightarrow \mathbb{R}^2$, of the following system:

$$\mathcal{U}\zeta_w - \kappa_\perp \zeta_r = \sin \theta, \quad (4.19)$$

$$\mathcal{U}\zeta_r - \frac{(n-3)D}{\sin^2 \theta \kappa_\perp^2} \zeta_r + \kappa_\perp \zeta_w = 0. \quad (4.20)$$

where the operator \mathcal{U} operates on any smooth function $\zeta: (0, \pi) \times \mathbb{R} \times [0, \infty) \rightarrow \mathbb{R}$, $(\theta, \kappa_\parallel, \kappa_\perp) \rightarrow \zeta$ as follows:

$$\mathcal{U} = \mathcal{U}_1 + \mathcal{U}_2, \quad (4.21)$$

$$\mathcal{U}_1 \zeta = \kappa_\parallel \partial_\theta \zeta + (\cos \theta \sin \theta \kappa_\perp^2 - \nu \sin \theta) \partial_{\kappa_\parallel} \zeta - 2 \frac{\cos \theta}{\sin \theta} \kappa_\parallel \kappa_\perp \partial_{\kappa_\perp} \zeta, \quad (4.22)$$

$$\mathcal{U}_2 \zeta = D \left(\frac{1}{\mathbf{M}_u^\flat} \partial_{\kappa_\parallel} (\mathbf{M}_u^\flat \partial_{\kappa_\parallel} \zeta) + \frac{1}{\sin^2 \theta \mathbf{M}_u^\flat \kappa_\perp^{n-3}} \partial_{\kappa_\perp} (\mathbf{M}_u^\flat \kappa_\perp^{n-3} \partial_{\kappa_\perp} \zeta) \right). \quad (4.23)$$

Then, c_2 is given by

$$c_2 = \frac{\int_{(0, \pi) \times \mathbb{R} \times (0, \infty)} \cos \theta (\mathbf{M}_u^\flat \zeta_w)(\theta, \kappa_\parallel, \kappa_\perp) \sin^{2n-3} \theta \kappa_\perp^{n-3} d\theta d\kappa_\parallel d\kappa_\perp}{\int_{(0, \pi) \times \mathbb{R} \times (0, \infty)} (\mathbf{M}_u^\flat \zeta_w)(\theta, \kappa_\parallel, \kappa_\perp) \sin^{2n-3} \theta \kappa_\perp^{n-3} d\theta d\kappa_\parallel d\kappa_\perp}. \quad (4.24)$$

Remark 4.1. (i) System (4.19), (4.20) will be given a variational formulation and shown to be well-posed below (see (7.35) for the variational formulation in dimension $n \geq 4$ and (7.46) in dimension $n = 3$). Boundary conditions associated with this variational formulation will follow from the variational formulation. Only the component ζ_w is needed to compute the coefficient c_2 . However, ζ_w and ζ_r are coupled and ζ_w cannot be found independently of ζ_r .

(ii) \mathbf{M}_u^\flat is nothing but the expression of the von Mises M_u in an appropriate change of variables which will be introduced below.

In dimension 2, it was already proved in [37] that the hydrodynamic limit of the KCC model is the SOH System. Theorem 4.5 extends the validity of this result to any dimension. This extension is far from obvious as it brings geometric intricacies that are absent in dimension 2. Indeed, in dimension 2, the PCC and KCC models can be formulated in standard flat spaces and the function ζ_w is obtained by solving a single equation instead of a system like (4.19), (4.20). Previously, the SOH model was derived as the hydrodynamic limit of the kinetic Vicsek model in [35] (in the normalized case) and in [31, 32] (in the non-normalized case). It also appears as the hydrodynamic limit of a Cucker-Smale model in [1]. All the previous results are formal but rigorous local convergence results were proved in [55, 56]. Local existence of smooth solutions of the SOH model was proved in [33] and in [76] in the viscous case. Its numerical resolution has been performed in [63].

A fluid model has been derived from the curvature-control model of [22] in [75]. This model is quite different from the SOH model. It is an extension of the Toner-Tu model in which \mathbf{u} is not normalized but instead, a relaxation term of the Ginzburg-Landau type is added at the right-hand side of the momentum equation (8.9). In addition, the model of [75] involves another variable, the locally-averaged curvature, and thus takes the form of three coupled PDEs.

The model of [75] is derived through a moment expansion of the kinetic equation and adequate moment closure assumptions. We do not think that such a model could be derived from the kinetic equation (3.52) by a hydrodynamic limit, even modifying the underpinning scaling assumptions. It seems however possible to derive a model involving an additional equation for the average curvature at the price of slightly modifying the initial PCC model (2.48), (2.49) and its associated kinetic model (3.52). This will be investigated in future work.

The remainder of this paper is devoted to the proof of Theorem 4.5. The establishment of the continuity equation (4.15) is easy and is done in the next subsection. All the forthcoming developments will be formal except otherwise stated.

4.4 The continuity equation

In this section, we start the proof of Theorem 4.5 by determining the limit of f^ε and showing the validity of the continuity equation (4.15).

Lemma 4.6 (Equilibrium in the spatially nonhomogeneous case).

Suppose $f^\varepsilon \rightarrow f$ as $\varepsilon \rightarrow 0$ as smoothly as needed. Then, f is given by (4.14).

Proof. From (3.52), we have $Q(f^\varepsilon) = \mathcal{O}(\varepsilon)$, so, when $\varepsilon \rightarrow 0$, we get $Q(f) = 0$. Thus, we can apply Prop. 4.4 and obtain that for any given (x, t) , the function $\alpha \mapsto f(x, \alpha, t)$ satisfies (4.13), which shows (4.14). ■

Proposition 4.7 (Continuity equation).

The functions ρ and \mathbf{u} satisfy the continuity equation (4.15).

Proof. Integrating (4.8) with respect to $\alpha \in M$, we have

$$\int_M Q(f) |\text{Vol}_M| = 0.$$

Hence, integrating (3.52) with respect to α leads to

$$\partial_t \left(\int_M f^\varepsilon |\text{Vol}_M| \right) + \nabla_x \cdot \left(\int_M f^\varepsilon v |\text{Vol}_M| \right) = 0. \quad (4.25)$$

As $\varepsilon \rightarrow 0$, we have

$$\begin{aligned} \int_M f^\varepsilon |\text{Vol}_M| &\longrightarrow \rho \int_M \mathbf{M}_u |\text{Vol}_M|(\alpha) = \rho, \\ \int_M f^\varepsilon v |\text{Vol}_M| &\longrightarrow \rho \int_M \mathbf{M}_u v |\text{Vol}_M| = c_1 \rho \mathbf{u}, \end{aligned}$$

by (4.6) and (4.11) respectively. Hence, letting $\varepsilon \rightarrow 0$ in (4.25) leads to (4.15). ■

4.5 The velocity equation

The establishment of the velocity equation (4.16) is far less straightforward. The main difficulty is the lack of non-trivial collision invariants, i.e. functions $\zeta, M \rightarrow \mathbb{R}$ other than the constants, such that

$$\int_M Q(f) \zeta |\text{Vol}_M| = 0, \quad \forall f : M \rightarrow \mathbb{R},$$

(note that the constant collision invariants give rise to the continuity equation established in the previous subsection).

In rarefied gas dynamics [23], non-trivial collision invariants are the microscopic momentum and energy. Integrating the Boltzmann equation against them gives rise to the momentum and energy conservation equations in the macroscopic model. Here, microscopic momentum or energy is not preserved, and there is no substitute that would allow us to easily obtain an evolution equation for \mathbf{u} . This problem was first encountered in the case of the the Vicsek model in [35], where it was resolved by introducing the concept of generalized collision invariant (GCI), which is a weakened form of collision invariant. It was applied to the 2D curvature control model in [37]. However, the passage from 2D to arbitrary dimension requires crossing a conceptual gap, due to the fact that the GCI in the 2D curvature control model is a scalar, while it is a vector in higher dimensions. The components of the GCI in higher dimensions do not simply follow from a simple scaling up formula from the scalar GCI in the 2D model.

Below, we split the derivation of the velocity equation in four sections. First, in Section 5, we introduce the concept of GCI (and more precisely, of vector GCI). We prove its existence and uniqueness thanks to the transcription of the vector GCI equation into a variational formulation in appropriate spaces. We show that this variational formulation falls in the framework developed in [3], from which existence and uniqueness are easily deduced.

However, a more detailed form of the vector GCI is necessary to derive an explicit form of the velocity equation. A first step towards this goal is to exploit the action of the orthogonal group O_n on the vector GCI. Section 6 is devoted to showing that the vector GCI is invariant under the action of the isotropy subgroup of O_n which fixes \mathbf{u} (which is classically isomorphic to O_{n-1}). This suggests that the vector GCI is the solution of a variational formulation posed on spaces of functions that are invariant under O_{n-1} . In this section, we derive such a variational formulation by averaging the previously found variational formulation over the Haar measure of O_{n-1} , and show the existence and uniqueness of its solution.

We further exploit the invariance of the vector GCI under the action of O_{n-1} in Section 7, by showing that it is necessarily expressed in terms of two scalar functions called the reduced GCI pair, which are nothing but the pair (ζ_w, ζ_r) introduced in Theorem 4.5. We then derive a variational formulation for the reduced GCI pair from the variational formulation for the vector GCI posed on spaces of O_{n-1} -invariant functions. Existence and uniqueness of the former is guaranteed by those of the latter. The variational formulation for the GCI pair can be interpreted formally as the PDE system (4.19), (4.20).

The final step, performed in Section 8, is to derive the velocity equation (4.16) from the knowledge of the GCI pair and to show the expression (4.24) for c_2 . The computations involved in this step heavily rely on the previous sections.

5 Generalized collision invariants (GCI)

In this section, we introduce the concept of generalized collision invariant (GCI) and show the existence and uniqueness (up to constants) of the GCI.

5.1 GCI: definition

We introduce a linear operator constructed from the collision operator by freezing the \mathbf{u} variable: let $\mathbf{u} \in \mathbb{S}^{n-1}$ be given and fixed. For a smooth function $f: M \rightarrow \mathbb{R}$, we define

$$\mathcal{Q}_{\mathbf{u}}(f) = -\{f, \mathbf{H}_{\mathbf{u}}\} + D\nabla_{\kappa} \cdot \left(\mathbf{M}_{\mathbf{u}} \nabla_{\kappa} \left(\frac{f}{\mathbf{M}_{\mathbf{u}}} \right) \right). \quad (5.1)$$

By (4.8), for any smooth function $f: M \rightarrow \mathbb{R}$, we have

$$\mathcal{Q}(f) = \mathcal{Q}_{\mathbf{u}_f}(f). \quad (5.2)$$

Since the bracket is a derivation, the formal L^2 -adjoint of $\mathcal{Q}_{\mathbf{u}}$ is $\mathcal{Q}_{\mathbf{u}}^*$ given on any smooth function $\zeta: M \rightarrow \mathbb{R}$ by

$$\mathcal{Q}_{\mathbf{u}}^*(\zeta) = \{\zeta, \mathbf{H}_{\mathbf{u}}\} + \frac{D}{\mathbf{M}_{\mathbf{u}}} \nabla_{\kappa} \cdot (\mathbf{M}_{\mathbf{u}} \nabla_{\kappa} \zeta). \quad (5.3)$$

Now, we define a GCI associated with \mathbf{u} as follows:

Definition 5.1 (GCI).

Let $\mathbf{u} \in \mathbb{S}^{n-1}$. A GCI associated with \mathbf{u} is a function $\zeta: M \rightarrow \mathbb{R}$ such that

$$\int_M \mathcal{Q}_{\mathbf{u}}(f) \zeta |\text{Vol}_M| = 0, \quad \forall f \text{ such that } P_{\mathbf{u}^\perp} \left(\int_M f v |\text{Vol}_M| \right) = 0. \quad (5.4)$$

Proposition 5.2 (Equation solved by the GCI).

Let $\mathbf{u} \in \mathbb{S}^{n-1}$. A smooth function $\zeta: M \rightarrow \mathbb{R}$ is a GCI associated with \mathbf{u} if and only if there exists a vector $A \in \mathbb{R}^n$ such that $A \cdot \mathbf{u} = 0$ and

$$\mathcal{Q}_{\mathbf{u}}^*(\zeta)(\alpha) = v \cdot A, \quad \forall \alpha = (v, \kappa) \in M, \quad (5.5)$$

Proof. In the space $L^2(M, \mathbb{R})$, consider the finite-dimensional linear subspace $\mathcal{F}_{\mathbf{u}}$ consisting of all functions of the form $\alpha \mapsto (v \cdot A)$ (with $\alpha = (v, \kappa)$), where $A \in \mathbb{R}^n$ is any vector such that $A \cdot \mathbf{u} = 0$. Eq. (5.4) can be equivalently written

$$f \in \mathcal{F}_{\mathbf{u}}^\perp \implies f \in \text{Span}\{\mathcal{Q}_{\mathbf{u}}^*(\zeta)\}^\perp.$$

Since both $\mathcal{F}_{\mathbf{u}}$ and $\text{Span}\{\mathcal{Q}_{\mathbf{u}}^*(\zeta)\}$ are finite dimensional, hence closed, this statement is again equivalent to

$$\text{Span}\{\mathcal{Q}_{\mathbf{u}}^*(\zeta)\} \subset \mathcal{F}_{\mathbf{u}},$$

which is exactly what (5.5) means. ■

Now, we define a “vector GCI” as follows. Let $\mathbf{u} \in \mathbb{S}^{n-1}$ be given and let (e_1, \dots, e_{n-1}) be an orthonormal basis of $\{\mathbf{u}\}^\perp$. For each $k \in \{1, \dots, n-1\}$, denote by ζ_k a GCI associated with \mathbf{u} and the choice $A = e_k$ in Prop. 5.2. Define a vector GCI $\vec{\zeta}$ associated with \mathbf{u} by

$$\vec{\zeta} = \sum_{k=1}^{n-1} \zeta_k e_k. \quad (5.6)$$

Applying Prop. 5.2, we find that $\vec{\zeta}$ satisfies the vector equation

$$\mathcal{Q}_{\mathbf{u}}^*(\vec{\zeta})(\alpha) = P_{\mathbf{u}^\perp} v, \quad \forall \alpha = (v, \kappa) \in M, \quad (5.7)$$

componentwise. A vector GCI has the following important property, which justifies the introduction of the GCI concept:

Lemma 5.3 (A vector GCI cancels the collision operator).

For any smooth enough function $f: M \rightarrow \mathbb{R}$, let $\vec{\zeta}$ be a vector GCI associated with \mathbf{u}_f . Then, we have

$$\int_M \mathcal{Q}(f) \vec{\zeta} |\text{Vol}_M| = 0. \quad (5.8)$$

Proof. f satisfies

$$P_{\mathbf{u}_f^\perp} \left(\int_M f v |\text{Vol}_M| \right) = \left| \int_M f v |\text{Vol}_M| \right| P_{\mathbf{u}_f^\perp} \mathbf{u}_f = 0.$$

Since $\vec{\zeta}$ is a vector GCI associated with \mathbf{u}_f , according to Def. 5.1, we deduce that

$$\int_M \mathcal{Q}_{\mathbf{u}_f}(f) \vec{\zeta} |\text{Vol}_M| = 0.$$

With (5.2), we get (5.8). ■

5.2 Existence and uniqueness (up to constants) of the vector GCI

We now prove the existence of solutions of the stationary linear Fokker-Planck equation (5.7). In [30, 34], a variational framework for the linear time-dependent Fokker-Planck equation was proposed. It relied on [7] which made essential use of a result of Lions [62]. This method was used again in [37], this time in the case of the stationary Fokker-Planck equation, but there is a mistake in the proof. Recently, [3] revisited this variational framework and showed a Poincaré estimate and the existence of solutions for both the stationary and time-dependent Fokker-Planck equations. In this section, we adapt the results of [3] to the present geometric setting. Since the proofs involve only minor adaptations of [3], they are omitted. We first introduce the following functional spaces.

5.2.1 Spaces of functions and distributions on M

The following list of spaces is tedious but is needed for the sake of notations. We begin with definitions of spaces of scalar functions on M .

(i) The space $L_v^2(L_\kappa^2)$ is defined as the closure of the space of smooth functions $M \rightarrow \mathbb{R}$ for the norm

$$\|f\|_{L_v^2(L_\kappa^2)}^2 = \int_M |f|^2 \mathbf{M}_u |\text{Vol}_M|. \quad (5.9)$$

We denote by $\langle\langle f, g \rangle\rangle_{L_v^2(L_\kappa^2)}$ the associated inner-product. We also define the closed hyperplane

$$L_v^2(L_\kappa^2)_0 = \left\{ f \in L_v^2(L_\kappa^2) \mid \int_M f \mathbf{M}_u |\text{Vol}_M| = 0 \right\}.$$

Both $L_v^2(L_\kappa^2)$ and $L_v^2(L_\kappa^2)_0$ are Hilbert spaces.

(ii) The space $\tilde{L}_v^2(L_\kappa^2)$ is the space of sections k of the pullback bundle π^*TB such that the map $M \rightarrow \mathbb{R}, (v, \kappa) \rightarrow \langle k, k \rangle_v$ belongs to $L_v^2(L_\kappa^2)$. It is a Hilbert space for the inner product

$$\langle\langle k, h \rangle\rangle_{\tilde{L}_v^2(L_\kappa^2)} = \int_M \langle k, h \rangle \mathbf{M}_u |\text{Vol}_M|. \quad (5.10)$$

and we denote by $\|k\|_{\tilde{L}_v^2(L_\kappa^2)}^2$ the associated norm.

(iii) The space $L_v^2(H_\kappa^1)$ is the closure of the space of smooth functions $M \rightarrow \mathbb{R}$ for the norm

$$\|f\|_{L_v^2(H_\kappa^1)}^2 = \|f\|_{L_v^2(L_\kappa^2)}^2 + \|\nabla_\kappa f\|_{\tilde{L}_v^2(L_\kappa^2)}^2, \quad (5.11)$$

and we denote by $\langle\langle f, g \rangle\rangle_{L_v^2(H_\kappa^1)}$ the associated inner-product. It is a Hilbert space.

(iv) The space $\mathcal{D}(M)$ is the space of smooth (infinitely differentiable) functions $M \rightarrow \mathbb{R}$ with compact support. Its dual $\mathcal{D}'(M)$ is the space of distributions on M . For $T \in \mathcal{D}'(M)$ and $\varphi \in \mathcal{D}(M)$, the duality bracket $\langle\langle T, \varphi \rangle\rangle_{\mathcal{D}', \mathcal{D}}$ is defined to extend the inner product of $L_v^2(L_\kappa^2)$. The weight \mathbf{M}_u in the duality bracket must be taken into account when defining the distributional derivative. Hence, for $T \in \mathcal{D}'$, we define $\nabla_v T$ and $\nabla_\kappa T$ by duality against an element $\varphi \in \mathcal{Y}(M)$ with compact support as follows

$$\langle\langle \nabla_v T, \varphi \rangle\rangle_{\mathcal{D}', \mathcal{D}} = -\langle\langle T, \frac{1}{\mathbf{M}_u} \nabla_v \cdot (\mathbf{M}_u \varphi) \rangle\rangle_{\mathcal{D}', \mathcal{D}},$$

and similarly replacing v by κ . We note that

$$\frac{1}{\mathbf{M}_u} \nabla_v \cdot (\mathbf{M}_u \varphi) = \nabla_v \cdot \varphi + \frac{v^2}{D} P_{v^\perp} \mathbf{u} \cdot \varphi, \quad \frac{1}{\mathbf{M}_u} \nabla_\kappa \cdot (\mathbf{M}_u \varphi) = \nabla_\kappa \cdot \varphi - \frac{v}{D} \kappa \cdot \varphi,$$

are smooth so that the above definitions make sense. Thanks to this definition, gradients in the usual and distributional senses coincide when applied to smooth functions. Since the bracket is a derivation, and \mathbf{M}_u is a function of \mathbf{H}_u , we have

$$\langle\langle \{T, \mathbf{H}_u\}, \varphi \rangle\rangle_{\mathcal{D}', \mathcal{D}} = -\langle\langle T, \{\varphi, \mathbf{H}_u\} \rangle\rangle_{\mathcal{D}', \mathcal{D}}. \quad (5.12)$$

(v) The dual space $L_v^2(H_\kappa^{-1})$ of $L_v^2(H_\kappa^1)$ is a subspace of $\mathcal{D}'(M)$, namely, the subspace of continuous linear forms on $\mathcal{D}(M)$ that extend to elements of $L_v^2(H_\kappa^1)$. We denote by $\langle\langle T, \varphi \rangle\rangle_{L_v^2(H_\kappa^{-1}), L_v^2(H_\kappa^1)}$ the duality between $T \in L_v^2(H_\kappa^{-1})$ and $\varphi \in L_v^2(H_\kappa^1)$. The space $L_v^2(H_\kappa^{-1})$ is normed by the usual operator norm of linear forms, namely

$$\|T\|_{L_v^2(H_\kappa^{-1})} = \sup_{\varphi \in L_v^2(H_\kappa^1), \|\varphi\|_{L_v^2(H_\kappa^1)} \leq 1} \left| \langle\langle T, \varphi \rangle\rangle_{L_v^2(H_\kappa^{-1}), L_v^2(H_\kappa^1)} \right|.$$

$L_v^2(H_\kappa^{-1})$ is a Hilbert space.

Now, we extend these definitions to spaces of vector-valued functions and distributions. Let (e_1, \dots, e_{n-1}) be an orthonormal basis of $\{\mathbf{u}\}^\perp$.

(i) The space $L_v^2(L_\kappa^2)_\mathbf{u}$ is the set of vector valued functions $\vec{\psi}: M \rightarrow \{\mathbf{u}\}^\perp$ such that $\vec{\psi}_i = \vec{\psi} \cdot e_i \in L_v^2(L_\kappa^2)$, $\forall i \in \{1, \dots, n-1\}$, endowed with the norm

$$\|\vec{\psi}\|_{L_v^2(L_\kappa^2)_\mathbf{u}}^2 = \sum_{i=1}^{n-1} \|\vec{\psi}_i\|_{L_v^2(L_\kappa^2)}^2, \quad (5.13)$$

and associated inner product

$$\langle\langle \vec{\psi}, \vec{\varphi} \rangle\rangle_{L_v^2(L_\kappa^2)_\mathbf{u}} = \sum_{i=1}^{n-1} \langle\langle \vec{\psi}_i, \vec{\varphi}_i \rangle\rangle_{L_v^2(L_\kappa^2)} = \int_M \vec{\psi} \cdot \vec{\varphi} \mathbf{M}_\mathbf{u} |\text{Vol}_M|,$$

where $\vec{\psi} \cdot \vec{\varphi} = \sum_{i=1}^{n-1} \vec{\psi}_i \vec{\varphi}_i$ is the usual inner product in $\{\mathbf{u}\}^\perp$. Both the norm and inner-product are independent of the choice of the orthonormal basis of $\{\mathbf{u}\}^\perp$. We also define the closed hyperplane

$$L_v^2(L_\kappa^2)_{\mathbf{u}0} = \left\{ \vec{\psi} \in L_v^2(L_\kappa^2)_\mathbf{u} \mid \vec{\psi}_i \in L_v^2(L_\kappa^2)_0, \forall i \in \{1, \dots, n-1\} \right\}.$$

Both $L_v^2(L_\kappa^2)_\mathbf{u}$ and $L_v^2(L_\kappa^2)_{\mathbf{u}0}$ are Hilbert spaces.

(ii) Let \vec{k} be a section of the bundle $\pi^*(TB) \otimes (M \times \{\mathbf{u}\}^\perp)$. We can write $\vec{k} = \sum_{i=1}^{n-1} \vec{k}_i \otimes e_i$, with \vec{k}_i being a section of the pullback bundle $\pi^*(TB)$, i.e. $\vec{k}_i(\alpha) \in T_{\pi\alpha}B$, $\forall \alpha \in M$. The space $\vec{L}_v^2(L_\kappa^2)_\mathbf{u}$ is the space of such sections such that $\vec{k}_i \in \vec{L}_v^2(L_\kappa^2)$, $\forall i \in \{1, \dots, n-1\}$. This space is a Hilbert space when endowed with norm

$$\|\vec{k}\|_{\vec{L}_v^2(L_\kappa^2)_\mathbf{u}}^2 = \sum_{i=1}^{n-1} \|\vec{k}_i\|_{\vec{L}_v^2(L_\kappa^2)}^2, \quad (5.14)$$

and associated inner product $\langle\langle \vec{k}, \vec{h} \rangle\rangle_{\vec{L}_v^2(L_\kappa^2)_\mathbf{u}}$. Again, both the norm and inner-product are independent of the choice of the orthonormal basis of $\{\mathbf{u}\}^\perp$.

(iii) Let $\vec{\psi}: M \rightarrow \{\mathbf{u}\}^\perp$. Then, $\nabla_\kappa \vec{\psi}$, when it exists, is a section of $\pi^*(TB) \otimes (M \times \{\mathbf{u}\}^\perp)$ and we can write $\nabla_\kappa \vec{\psi} = \sum_{i=1}^{n-1} \nabla_\kappa \vec{\psi}_i \otimes e_i$. The space $L_v^2(H_\kappa^1)_\mathbf{u}$ is given by

$$L_v^2(H_\kappa^1)_\mathbf{u} = \left\{ \vec{\psi} \in L_v^2(L_\kappa^2)_\mathbf{u} \mid \nabla_\kappa \vec{\psi} \in \vec{L}_v^2(L_\kappa^2)_\mathbf{u} \right\}.$$

It is a Hilbert space for the norm

$$\|\vec{\psi}\|_{L_v^2(H_\kappa^1)_\mathbf{u}}^2 = \|\vec{\psi}\|_{L_v^2(L_\kappa^1)_\mathbf{u}}^2 + \|\nabla_\kappa \vec{\psi}\|_{L_v^2(L_\kappa^1)_\mathbf{u}}^2 = \sum_{i=1}^{n-1} \|\vec{\psi}_i\|_{L_v^2(H_\kappa^1)}^2,$$

and associated inner-product $(\vec{\psi}, \vec{\phi})_{L_v^2(H_\kappa^1)_\mathbf{u}}$, which again, do not depend on the choice of the orthonormal basis of $\{\mathbf{u}\}^\perp$.

(iv) The space $D_\mathbf{u}(M)$ is the space of smooth (infinitely differentiable) functions $M \rightarrow \{\mathbf{u}\}^\perp$ with compact support. Its dual $D'_\mathbf{u}(M)$ is the space of distributions on M with values in $\{\mathbf{u}\}^\perp$. Let $\vec{T} \in D'_\mathbf{u}(M)$. We can write $\vec{T} = \sum_{i=1}^{n-1} \vec{T}_i e_i$, with $\vec{T}_i \in D'(M)$, $\forall i \in \{1, \dots, n-1\}$. For $\vec{T} \in D'_\mathbf{u}(M)$ and $\vec{\psi} = \sum_{i=1}^{n-1} \vec{\psi}_i e_i \in D_\mathbf{u}(M)$, the duality bracket $(\vec{T}, \vec{\psi})_{D'_\mathbf{u}, D_\mathbf{u}}$ is defined to extend the inner product of $L_v^2(L_\kappa^1)_\mathbf{u}$. We have

$$(\vec{T}, \vec{\psi})_{D'_\mathbf{u}, D_\mathbf{u}} = \sum_{i=1}^{n-1} (\vec{T}_i, \vec{\psi}_i)_{D', D}, \quad (5.15)$$

(v) The dual space $L_v^2(H_\kappa^{-1})_\mathbf{u}$ of $L_v^2(H_\kappa^1)_\mathbf{u}$ is a subspace of $D'_\mathbf{u}(M)$, namely, the subspace of continuous linear forms on $D_\mathbf{u}(M)$ that extend to elements of $L_v^2(H_\kappa^1)_\mathbf{u}$. We have

$$\vec{T} = \sum_{i=1}^{n-1} \vec{T}_i e_i \in L_v^2(H_\kappa^{-1})_\mathbf{u} \iff \vec{T}_i \in L_v^2(H_\kappa^{-1}), \quad \forall i \in \{1, \dots, n-1\}.$$

We denote by $(\vec{T}, \vec{\psi})_{L_v^2(H_\kappa^{-1})_\mathbf{u}, L_v^2(H_\kappa^1)_\mathbf{u}}$ the duality between $\vec{T} \in L_v^2(H_\kappa^{-1})_\mathbf{u}$ and $\vec{\psi} \in L_v^2(H_\kappa^1)_\mathbf{u}$. The identity (5.15) is still valid for the duality between $L_v^2(H_\kappa^{-1})_\mathbf{u}$ and $L_v^2(H_\kappa^1)_\mathbf{u}$ at the left-hand side and between $L_v^2(H_\kappa^{-1})$ and $L_v^2(H_\kappa^1)$ at the right-hand side. The space $L_v^2(H_\kappa^{-1})_\mathbf{u}$ is a Hilbert space when normed by the usual operator norm of linear forms, namely

$$\|\vec{T}\|_{L_v^2(H_\kappa^{-1})_\mathbf{u}} = \sup_{\vec{\psi} \in L_v^2(H_\kappa^1)_\mathbf{u}, \|\vec{\psi}\|_{L_v^2(H_\kappa^1)_\mathbf{u}} \leq 1} |(\vec{T}, \vec{\psi})_{L_v^2(H_\kappa^{-1})_\mathbf{u}, L_v^2(H_\kappa^1)_\mathbf{u}}|.$$

(vi) For any smooth function $\mathbf{H}: M \rightarrow \mathbb{R}$ and any $\vec{T} = \sum_{i=1}^{n-1} \vec{T}_i e_i \in D'_\mathbf{u}(M)$, we define $\{\vec{T}, \mathbf{H}\} = \sum_{i=1}^{n-1} \{\vec{T}_i, \mathbf{H}\} e_i$. Like in the scalar case, for any $\vec{\psi} \in D_\mathbf{u}(M)$, we have

$$(\{\vec{T}, \mathbf{H}_\mathbf{u}\}, \vec{\psi})_{D'_\mathbf{u}, D_\mathbf{u}} = -(\vec{T}, \{\vec{\psi}, \mathbf{H}_\mathbf{u}\})_{D'_\mathbf{u}, D_\mathbf{u}}. \quad (5.16)$$

The space $\mathbb{Y}_\mathbf{u}$ is defined by

$$\mathbb{Y}_\mathbf{u} = \left\{ \vec{\psi} \in L_v^2(H_\kappa^1)_\mathbf{u}, \quad \{\vec{\psi}, \mathbf{H}_\mathbf{u}\} \in L_v^2(H_\kappa^{-1})_\mathbf{u} \right\},$$

where $\{\vec{\psi}, \mathbf{H}_\mathbf{u}\}$ is defined in the distributional sense by (5.16). The space $\mathbb{Y}_\mathbf{u}$ is normed by

$$\|\vec{\psi}\|_{\mathbb{Y}_\mathbf{u}}^2 = \|\vec{\psi}\|_{L_v^2(H_\kappa^1)_\mathbf{u}}^2 + \|\{\vec{\psi}, \mathbf{H}_\mathbf{u}\}\|_{L_v^2(H_\kappa^{-1})_\mathbf{u}}^2,$$

and is a Hilbert space.

The following lemmas provide some properties of the space $\mathbb{Y}_\mathbf{u}$. Their proofs are fairly straightforward adaptations of corresponding results proved in [3], and are omitted.

Lemma 5.4 (Density of $D_{\mathbf{u}}(M)$ in $\mathbb{Y}_{\mathbf{u}}$).

$D_{\mathbf{u}}(M)$ is dense in $\mathbb{Y}_{\mathbf{u}}$.

Lemma 5.5 (Poincaré inequality in $\mathbb{Y}_{\mathbf{u}}$).

There exists a constant $C > 0$ such that for any $\vec{\psi} \in \mathbb{Y}_{\mathbf{u}} \cap L_v^2(L_\kappa^2)_{\mathbf{u}0}$ we have

$$\|\vec{\psi}\|_{L_v^2(L_\kappa^2)_{\mathbf{u}}} \leq C(\|\nabla_\kappa \vec{\psi}\|_{\tilde{L}_v^2(L_\kappa^2)_{\mathbf{u}}} + \|\{\vec{\psi}, \mathbf{H}_{\mathbf{u}}\}\|_{L_v^2(H_\kappa^{-1})_{\mathbf{u}}}). \quad (5.17)$$

5.2.2 Proof of existence of the vector GCI

Let $\vec{\chi} \in L_v^2(L_\kappa^2)_{\mathbf{u}}$. We are going to prove the existence of solutions to the following equation:

$$\mathcal{Q}_{\mathbf{u}}^*(\vec{\psi}) = \vec{\chi}, \quad (5.18)$$

provided that $\vec{\chi} \in L_v^2(L_\kappa^2)_{\mathbf{u}0}$, and we will characterize the space of such solutions. For that purpose, we transform (5.18) into a variational formulation. Let $\vec{\phi} \in D_{\mathbf{u}}(M)$. Taking the inner-product of (5.18) (in the space $\{\mathbf{u}\}^\perp$) with $\mathbf{M}_{\mathbf{u}} \vec{\phi}$, integrating over M and using (3.22) supposing that $\vec{\psi}$ has all the requested regularity for this to be possible, we get

$$-\int_M \{\vec{\psi}, \mathbf{H}_{\mathbf{u}}\} \cdot \vec{\phi} \mathbf{M}_{\mathbf{u}} |\text{Vol}_M| + D \int_M \sum_{i=1}^{n-1} \langle \nabla_\kappa \vec{\phi}_i, \nabla_\kappa \vec{\psi}_i \rangle \mathbf{M}_{\mathbf{u}} |\text{Vol}_M| = - \int_M \vec{\chi} \cdot \vec{\phi} \mathbf{M}_{\mathbf{u}} |\text{Vol}_M|.$$

Now, we see that if both $\vec{\phi}$ and $\vec{\psi}$ belong to $\mathbb{Y}_{\mathbf{u}}$, this formula has a meaning by interpreting the first integral as a duality bracket between $L_v^2(H_\kappa^{-1})_{\mathbf{u}}$ and $L_v^2(H_\kappa^1)_{\mathbf{u}}$. This leads to the variational formulation

$$\begin{cases} \vec{\psi} \in \mathbb{Y}_{\mathbf{u}}, \\ -(\langle \{\vec{\psi}, \mathbf{H}_{\mathbf{u}}\}, \vec{\phi} \rangle_{L_v^2(H_\kappa^{-1})_{\mathbf{u}}, L_v^2(H_\kappa^1)_{\mathbf{u}}} + D(\langle \nabla_\kappa \vec{\psi}, \nabla_\kappa \vec{\phi} \rangle_{\tilde{L}_v^2(L_\kappa^2)_{\mathbf{u}}}) = -(\langle \vec{\chi}, \vec{\phi} \rangle_{L_v^2(L_\kappa^2)_{\mathbf{u}}}), \quad \forall \vec{\phi} \in \mathbb{Y}_{\mathbf{u}}, \end{cases} \quad (5.19)$$

By testing (5.19) with $\vec{\phi} = e_i$ for $i = 1, \dots, n-1$ (which belong to $\mathbb{Y}_{\mathbf{u}}$) and noting that $\nabla_\kappa \vec{\phi} = 0$ and $\{\vec{\phi}, \mathbf{H}_{\mathbf{u}}\} = 0$, we see that the existence of a solution to (5.19) requires that $\vec{\chi} \in L_v^2(L_\kappa^2)_{\mathbf{u}0}$. The following theorem, whose proof is a fairly straightforward adaptation of [3] and is omitted, shows that, for $\vec{\chi} \in L_v^2(L_\kappa^2)_{\mathbf{u}}$, $\vec{\chi}$ being in $L_v^2(L_\kappa^2)_{\mathbf{u}0}$ is a necessary and sufficient condition for the existence of a solution to (5.19).

Theorem 5.6 (Existence of a solution to (5.19)).

Let $\vec{\chi} \in L_v^2(L_\kappa^2)_{\mathbf{u}}$. Then, the variational formulation (5.19) has a solution if and only if $\vec{\chi} \in L_v^2(L_\kappa^2)_{\mathbf{u}0}$ and in that case, the solution is unique in $\mathbb{Y}_{\mathbf{u}} \cap L_v^2(L_\kappa^2)_{\mathbf{u}0}$.

From Theorem 5.6, we immediately deduce the following

Corollary 5.7 (Existence and space of GCI).

(i) There exists a unique variational solution (in the sense of (5.19) with χ given by $\vec{\chi}(v, \kappa) = P_{\mathbf{u}^\perp} v$, $\forall \alpha = (v, \kappa) \in M$) in $\tilde{\mathbb{Y}}_{\mathbf{u}} \cap L_v^2(L_\kappa^2)_{\mathbf{u}0}$ to Equation (5.7). This unique solution is denoted by $\vec{\zeta}_{\mathbf{u}}$. The set of solutions to (5.7) in $\mathbb{Y}_{\mathbf{u}}$ is given by $\{\vec{\zeta}_{\mathbf{u}} + \vec{C} \mid \vec{C} \in \{\mathbf{u}\}^\perp\}$.

(ii) The space of (scalar) GCI associated with a given direction $\mathbf{u} \in \mathbb{S}^{n-1}$ is given by

$$\mathcal{G}_{\mathbf{u}} = \left\{ \vec{\zeta}_{\mathbf{u}} \cdot A + C \mid A \in \{\mathbf{u}\}^{\perp} \text{ and } C \in \mathbb{R} \right\},$$

and is an n -dimensional vector space.

Proof. (i) The function $M \rightarrow \{\mathbf{u}\}^{\perp}$, $(v, \kappa) \mapsto \vec{\chi}(v, \kappa) = P_{\mathbf{u}^{\perp}} v$ belongs to $L_v^2(L_{\kappa}^2)_{\mathbf{u}0}$ because for each $i = 1, \dots, n-1$, we have $P_{\mathbf{u}^{\perp}} v \cdot e_i = v \cdot e_i$ and

$$\int_B (v \cdot e_i) \exp\left(\frac{v}{D} \cdot \mathbf{u}\right) |\text{Vol}_B| = 0,$$

by antisymmetry. Theorem 5.6 ensures the existence of $\vec{\zeta}_{\mathbf{u}}$, the unique solution in $\mathbb{Y}_{\mathbf{u}} \cap L_v^2(L_{\kappa}^2)_{\mathbf{u}0}$ of (5.5) in the sense of the variational formulation (5.19).

(ii) Follows immediately from Point (i) and Prop. 5.2. \blacksquare

Remark 5.1 (Back to construction (5.6)).

The uniqueness (up to constants) of the vector GCI shows that the construction (5.6) does not depend on the choice of the orthonormal basis $\{e_1, \dots, e_{n-1}\}$ of $\{\mathbf{u}\}^{\perp}$.

6 Action of the orthogonal group on the vector GCI

In this section, we show that the orthogonal group O_n acts on the vector GCI $\vec{\zeta}_{\mathbf{u}}$ and that O_{n-1} (identified with the isotropy subgroup of O_n which fixes \mathbf{u}) fixes $\vec{\zeta}_{\mathbf{u}}$. Then, thanks to a suitably defined projection on subspaces of invariant functions under O_{n-1} , we show that we can restrict the variational formulation satisfied by $\vec{\zeta}_{\mathbf{u}}$ to such spaces of invariant functions. We first review some geometric background about the action of O_n on the various objects involved in this study.

6.1 Geometric preliminaries

In this section, we review how the orthogonal group O_n acts on B , M , the space $S(M, \mathbb{R})$ of smooth functions $M \rightarrow \mathbb{R}$, the spaces $\mathcal{X}(M)$ and $\mathcal{Y}(M)$ of smooth vector fields on M and of smooth sections of the pullback bundle $\pi^*(TB)$ respectively, and the various spaces of vector-valued functions and distributions defined in Section 5.2.1.

6.1.1 Action of O_n on B and M

We recall that O_n acts on the left on \mathbb{R}^n by the action $O_n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$, $(R, v) \mapsto Rv$ (meaning that the orthogonal matrix R acts on the n -tuple of coordinates of v in the canonical basis of \mathbb{R}^n). This action can be extended to a left action on \mathbb{R}^{2n} by $(R, (v, \kappa)) \mapsto R(v, \kappa) =: (Rv, R\kappa)$ and to a left action on \mathbb{R}^{4n} by $(R, (v, \kappa, a, \tau)) \mapsto R(v, \kappa, a, \tau) =: (Rv, R\kappa, Ra, R\tau)$. Now, since $|Rv| = |v|$, the action of O_n on \mathbb{R}^n can be restricted to an action of O_n on B . Similarly, since $Rv \cdot R\kappa = v \cdot \kappa$, the action of O_n on \mathbb{R}^{2n} can be restricted to an action of O_n on M . Finally, because $Ra \cdot Rv = a \cdot v$ and $Ra \cdot R\kappa + Rv \cdot R\tau = a \cdot \kappa + v \cdot \tau$, the action of O_n on \mathbb{R}^{4n} can be restricted to an action of O_n on N , the support manifold of the tangent bundle TM .

The action of O_n is compatible with the metrics on B : let $v \in B$ and $\kappa_1, \kappa_2 \in T_v B$. Then,

$$\langle R\kappa_1, R\kappa_2 \rangle_{Rv} = \langle \kappa_1, \kappa_2 \rangle_v, \quad \forall R \in O_n, \quad (6.1)$$

i.e. the action of O_n on B leads to orthogonal transformations from $T_v B$ to $T_{Rv} B$. Similarly, it is compatible with the metric on M : let $\alpha \in M$ and $\xi_1, \xi_2 \in T_\alpha M$. Then,

$$\langle\langle R\xi_1, R\xi_2 \rangle\rangle_{R\alpha} = \langle\langle \xi_1, \xi_2 \rangle\rangle_\alpha, \quad \forall R \in O_n. \quad (6.2)$$

In other words, the action of R on M leads to orthogonal transformations from $T_\alpha M$ to $T_{R\alpha} M$.

Now, the action of O_n commutes with the lifts and connection maps: let $\alpha \in M$ and $\xi \in T_\alpha M$. We have

$$B_{R\alpha}^V(R\xi) = RB_\alpha^V \xi, \quad B_{R\alpha}^H(R\xi) = RB_\alpha^H \xi, \quad \forall R \in O_n, \quad (6.3)$$

and likewise, let $\alpha \in M$, $\alpha = (v, \kappa)$ and let $a, \tau \in T_v B$. We have

$$\mathcal{L}_{R\alpha}^V(R\tau) = R\mathcal{L}_\alpha^V \tau, \quad \mathcal{L}_{R\alpha}^H(Ra) = R\mathcal{L}_\alpha^H a, \quad \forall R \in O_n. \quad (6.4)$$

6.1.2 Action of O_n on functions and vector fields

The left action of O_n on M induces a right action of O_n on $S(M, \mathbb{R})$ by $S(M, \mathbb{R}) \times O_n \rightarrow S(M, \mathbb{R})$, $(f, R) \rightarrow f^R$ where $f^R(\alpha) = f(R\alpha)$, $\forall \alpha \in M$. Similarly, there is a right action of O_n on $\mathcal{X}(M)$ by $\mathcal{X}(M) \times O_n \rightarrow \mathcal{X}(M)$, $(\Xi, R) \rightarrow \Xi^R$ where $\Xi^R(\alpha) = R^T \Xi(R\alpha)$, $\forall \alpha \in M$. Finally, there is a right action of O_n on $\mathcal{Y}(M)$ by $\mathcal{Y}(M) \times O_n \rightarrow \mathcal{Y}(M)$, $(\Xi, R) \rightarrow \Xi^R$ where $\Xi^R(\alpha) = R^T \Xi(R\alpha)$, $\forall \alpha \in M$. Simply we note that, in the case of $\mathcal{X}(M)$, the symbol $R^T \Xi$ means the action of O_n on N while in the case of $\mathcal{Y}(M)$ the same symbol means the action of O_n on M .

For $R \in O_n$, define $\sigma_R: M \rightarrow M$, $\alpha \mapsto R\alpha$. Then, $(d\sigma_R)_\alpha: T_\alpha M \rightarrow T_{R\alpha} M$ is given by $(d\sigma_R)_\alpha(\xi) = R\xi$. We note that $f^R = f \circ \sigma_R$. Now, the Sasaki volume form $|\text{Vol}_M|$ is invariant by the action of O_n on M in the sense that

$$\sigma_R^* \text{Vol}_M = \text{Vol}_M, \quad \forall R \in O_n. \quad (6.5)$$

In particular, the maps σ_R preserve the orientation. Obviously, this implies that

$$|\sigma_R^* \text{Vol}_M| = |\text{Vol}_M|, \quad \forall R \in O_n.$$

As a consequence, for any integrable function $f: M \rightarrow \mathbb{R}$ and for any $R \in O_n$, we have

$$\int_M f^R(\alpha) |\text{Vol}_M|(\alpha) = \int_M f(\alpha) |\text{Vol}_M|(\alpha). \quad (6.6)$$

Gradients and divergence transform under the action of O_n under the following rules. Suppose $f \in S(M, \mathbb{R})$ and $\Xi \in \mathcal{X}(M)$. Then, we have, for any $R \in O_n$:

$$\nabla_\alpha f^R = (\nabla_\alpha f)^R, \quad \nabla_\alpha \cdot \Xi^R = (\nabla_\alpha \cdot \Xi)^R. \quad (6.7)$$

Now, suppose that $f \in S(M, \mathbb{R})$ and $\Xi \in \mathcal{Y}(M)$. Then, we have, for any $R \in O_n$:

$$\nabla_v f^R = (\nabla_v f)^R, \quad \nabla_\kappa f^R = (\nabla_\kappa f)^R, \quad (6.8)$$

$$\nabla_v \cdot \Xi^R = (\nabla_v \cdot \Xi)^R, \quad \nabla_\kappa \cdot \Xi^R = (\nabla_\kappa \cdot \Xi)^R. \quad (6.9)$$

Suppose that $\Xi, \Upsilon \in \mathcal{Y}(M)$. Then, we have, for any $R \in O_n$:

$$\langle \Xi, \Upsilon \rangle^R = \langle \Xi^R, \Upsilon^R \rangle. \quad (6.10)$$

Finally, Let $\mathbf{u} \in \mathbb{R}^n$. We can easily check that

$$(\mathbf{H}_{\mathbf{u}})^R = \mathbf{H}_{R^T \mathbf{u}}. \quad (6.11)$$

6.1.3 Action of O_n on vector-valued functions and distributions

Let $S(M, \mathbb{R}^n)$ be the space of smooth vector-valued function $\vec{\varphi}: M \rightarrow \mathbb{R}^n$. We define a right action of O_n on this space by: $S(M, \mathbb{R}^n) \times O_n \rightarrow S(M, \mathbb{R}^n)$, $(\vec{\varphi}, R) \mapsto \vec{\varphi}^R$ where $\vec{\varphi}^R(\alpha) = R^T \vec{\varphi}(R\alpha)$, $\forall \alpha \in M$. Let $\mathbf{u} \in \mathbb{S}^{n-1}$. We note that if $\vec{\varphi}(\alpha) \in \{\mathbf{u}\}^\perp$, then $\vec{\varphi}^R(\alpha) \in \{R^T \mathbf{u}\}^\perp$. In this section, we review how this action affects the various spaces of vector-valued functions and distributions defined in Section 5.2.1. The proofs are straightforward and are omitted.

(i) Let $R \in O_n$. Then, $(L_v^2(L_\kappa^2)_{\mathbf{u}})^R = L_v^2(L_\kappa^2)_{R^T \mathbf{u}}$ and the inner-product is equivariant:

$$((\vec{\psi}^R, \vec{\varphi}^R))_{L_v^2(L_\kappa^2)_{R^T \mathbf{u}}} = ((\vec{\psi}, \vec{\varphi}))_{L_v^2(L_\kappa^2)_{\mathbf{u}}}, \quad \forall \vec{\psi}, \vec{\varphi} \in L_v^2(L_\kappa^2)_{\mathbf{u}}, \quad \forall R \in O_n. \quad (6.12)$$

(ii) We define a group action of O_n on the bundle $\pi^*(TB) \otimes (M \times \mathbb{R}^n)$ as follows: let (e_1, \dots, e_n) be an orthonormal basis of \mathbb{R}^n and let \vec{k} be a section of this bundle. We can write $\vec{k} = \sum_{i=1}^n \vec{k}_i \otimes e_i$, with \vec{k}_i being a section of π^*B , for all $i \in \{1, \dots, n\}$. We let $\vec{k}^R = \sum_{i=1}^n \vec{k}_i^R \otimes R^T e_i$, where we recall that $\vec{k}_i^R(\alpha) = R^T \vec{k}_i(R\alpha)$, $\forall \alpha \in M$. We easily verify that this group action does not depend on the choice of the orthonormal basis (e_1, \dots, e_n) . Under this group action, we have $(\pi^*(TB) \otimes (M \times \{\mathbf{u}\}^\perp))^R = (\pi^*(TB) \otimes (M \times \{R^T \mathbf{u}\}^\perp))$ and the $\tilde{L}_v^2(L_\kappa^2)_{\mathbf{u}}$ inner-product is equivariant by O_n :

$$((\vec{k}^R, \vec{h}^R))_{\tilde{L}_v^2(L_\kappa^2)_{R^T \mathbf{u}}} = ((\vec{k}, \vec{h}))_{\tilde{L}_v^2(L_\kappa^2)_{\mathbf{u}}}, \quad \forall \vec{k}, \vec{h} \in \tilde{L}_v^2(L_\kappa^2)_{\mathbf{u}}, \quad \forall R \in O_n, \quad (6.13)$$

(iii) Under the group action defined at Point (i), we have $(L_v^2(H_\kappa^1)_{\mathbf{u}})^R = L_v^2(H_\kappa^1)_{R^T \mathbf{u}}$ and for any $\vec{\psi} \in L_v^2(H_\kappa^1)_{\mathbf{u}}$, $\nabla_\kappa(\vec{\psi}^R) = (\nabla_\kappa \vec{\psi})^R$ (where at the right-hand side, the group action is that defined at Point (ii)). Furthermore, the $L_v^2(H_\kappa^1)_{\mathbf{u}}$ inner-product is equivariant by O_n :

$$((\nabla_\kappa \vec{\psi}^R, \nabla_\kappa \vec{\varphi}^R))_{\tilde{L}_v^2(L_\kappa^2)_{R^T \mathbf{u}}} = ((\nabla_\kappa \vec{\psi}, \nabla_\kappa \vec{\varphi}))_{\tilde{L}_v^2(L_\kappa^2)_{\mathbf{u}}}, \quad \forall \vec{\psi}, \vec{\varphi} \in L_v^2(H_\kappa^1)_{\mathbf{u}}, \quad \forall R \in O_n, \quad (6.14)$$

together with (6.12).

(iv) We define an action of O_n on the space of \mathbb{R}^n -valued distribution $\mathcal{D}'(M, \mathbb{R}^n)$ (the dual of the space $\mathcal{D}(M, \mathbb{R}^n)$ of smooth functions $M \rightarrow \mathbb{R}^n$ with compact support) by

$$((\vec{T}^R, \vec{\psi}))_{\mathcal{D}'(M, \mathbb{R}^n), \mathcal{D}(M, \mathbb{R}^n)} = ((\vec{T}, \vec{\psi}^{R^T}))_{\mathcal{D}'(M, \mathbb{R}^n), \mathcal{D}(M, \mathbb{R}^n)}, \quad \forall \vec{T} \in \mathcal{D}'(M, \mathbb{R}^n), \quad \forall \vec{\psi} \in \mathcal{D}(M, \mathbb{R}^n). \quad (6.15)$$

More explicitly, let (e_1, \dots, e_n) be an orthonormal basis of \mathbb{R}^n and let $\vec{T} \in \mathcal{D}'(M, \mathbb{R}^n)$. Write $\vec{T} = \sum_{i=1}^n \vec{T}_i e_i$ with $\vec{T}_i \in \mathcal{D}'(M)$ for all $i \in \{1, \dots, n\}$. Then, we have

$$\vec{T}^R = \sum_{i=1}^n \vec{T}_i^R R^T e_i. \quad (6.16)$$

Hence, we have $(D'_u(M))^R = (D'_{R^T u}(M))$ and the duality bracket between $D'_u(M)$ and $D_u(M)$ is equivariant:

$$((\vec{T}^R, \vec{\psi}^R))_{D'_{R^T u}(M), D_{R^T u}(M)} = ((\vec{T}, \vec{\psi}))_{D'_u(M), D_u(M)}, \quad \forall \vec{T} \in D'_u(M), \quad \forall \vec{\psi} \in D_u(M). \quad (6.17)$$

We deduce that the duality bracket between $L_v^2(H_\kappa^{-1})_u$ and $L_v^2(H_\kappa^1)_u$ is also equivariant:

$$((\vec{T}^R, \vec{\psi}^R))_{L_v^2(H_\kappa^{-1})_{R^T u}, L_v^2(H_\kappa^1)_{R^T u}} = ((\vec{T}, \vec{\psi}))_{L_v^2(H_\kappa^{-1})_u, L_v^2(H_\kappa^1)_u}, \quad \forall \vec{T} \in L_v^2(H_\kappa^{-1})_u, \quad \forall \vec{\psi} \in L_v^2(H_\kappa^1)_u. \quad (6.18)$$

Then, thanks to (5.15), (6.8), (6.9), (6.11), one easily verifies that for all $R \in O_n$, we have

$$\{\vec{T}^R, \mathbf{H}_{R^T u}\} = \{\vec{T}, \mathbf{H}_u\}^R, \quad \forall T \in D'_u(M). \quad (6.19)$$

We conclude that $(Y_u)^R = Y_{R^T u}$, $\forall R \in O_n$.

6.2 Action of the orthogonal group on the vector GCI

We have the following

Lemma 6.1 (Action of O_n on the vector GCI).

Let $\mathbf{u} \in \mathbb{S}^{n-1}$ and let $\vec{\zeta}_u$ be the vector GCI associated with \mathbf{u} . Then, we have

$$\vec{\zeta}_u^R = \vec{\zeta}_{R^T u}, \quad \forall R \in O_n. \quad (6.20)$$

Proof. As seen in the proof of Corollary 5.7, $\vec{\zeta}_u$ is the unique solution of variational formulation (5.19) with $\vec{\chi}$ given by $\vec{\chi}(\alpha) = P_{u^\perp} v$, $\forall \alpha = (v, \kappa) \in M$. Then, the results of Section 6.1.3 show that $\vec{\zeta}_u^R$ is a solution of the same variational formulation but with $\vec{\chi}$ replaced by $\vec{\chi}^R$. We compute, for all $\alpha = (v, \kappa) \in M$:

$$\vec{\chi}^R(\alpha) = R^T \vec{\chi}(R\alpha) = R^T P_{u^\perp}(Rv) = R^T (Rv - (Rv \cdot \mathbf{u})\mathbf{u}) = v - (v \cdot R^T \mathbf{u})R^T \mathbf{u} = P_{R^T u} v.$$

Hence, by uniqueness of the solution to the variational formulation (5.19), we obtain (6.20), which ends the proof. \blacksquare

With (6.20), it is enough to determine $\vec{\zeta}_u$ for a single vector \mathbf{u} , for instance the last vector \mathbf{e}_n of the canonical basis of \mathbf{R}^n . Indeed, since O_n acts transitively on B , there exists a (non-unique) orthogonal transformation $R \in O_n$ such that $R^T \mathbf{e}_n = \mathbf{u}$. Then, $\vec{\zeta}_u = \vec{\zeta}_{\mathbf{e}_n}^R$, which allows us to define $\vec{\zeta}_u$ from $\vec{\zeta}_{\mathbf{e}_n}$. Thus, in all the developments about the determination of $\vec{\zeta}_u$, we will specify $\mathbf{u} = \mathbf{e}_n$ and will use indifferently \mathbf{u} or \mathbf{e}_n to designate it.

By (6.20), $\vec{\zeta}_u$ satisfies

$$\vec{\zeta}_u^R = \vec{\zeta}_u, \quad \forall R \in O_n \text{ such that } R\mathbf{u} = \mathbf{u}. \quad (6.21)$$

The subgroup of O_n consisting of orthogonal transformations R such that $R\mathbf{u} = \mathbf{u}$ is the isotropy subgroup of \mathbf{u} and is isomorphic to O_{n-1} . We will identify below these two groups. We will use this invariance relation to obtain a simplified form of $\vec{\zeta}_u$. To start with, we first show that we can restrict variational formulation (5.19) to spaces of invariant functions under the action of O_{n-1}

6.3 Variational formulation in spaces of invariant functions under the action of O_{n-1}

We first define appropriate spaces of invariant functions and distributions under the action of O_{n-1} . Then, we define rotational averages, which are projections from functions or distribution spaces onto the corresponding subspaces of invariant objects. This eventually enables us to show that the variational formulation for the vector GCI (5.19) can be restricted to these invariant spaces.

6.3.1 Spaces of invariant vector-valued functions and distributions on M

In this section, we define spaces of invariant vector-valued functions and distributions on M and state some properties which are direct consequences of Section 6.1.3.

(i) We define the subspace of invariant functions $L_v^2(L_\kappa^2)_{\text{inv}}$ of $L_v^2(L_\kappa^2)_u$ by

$$L_v^2(L_\kappa^2)_{\text{inv}} = \{ \vec{\psi} \in L_v^2(L_\kappa^2)_u \mid \vec{\psi}^R = \vec{\psi}, \forall R \in O_{n-1} \}.$$

The space $L_v^2(L_\kappa^2)_{\text{inv}}$ is a closed subspace of $L_v^2(L_\kappa^2)_u$ and hence, is a Hilbert space normed by $\| \cdot \|_{L_v^2(L_\kappa^2)_u}$.

(ii) We define

$$\vec{L}_v^2(L_\kappa^2)_{\text{inv}} = \{ \vec{k} \in \vec{L}_v^2(L_\kappa^2)_u \mid \vec{k}^R = \vec{k}, \forall R \in O_{n-1} \}.$$

Again, $\vec{L}_v^2(L_\kappa^2)_{\text{inv}}$ is a closed subspace of $\vec{L}_v^2(L_\kappa^2)_u$ and hence, is a Hilbert space.

(iii) We define

$$L_v^2(H_\kappa^1)_{\text{inv}} = \{ \vec{\psi} \in L_v^2(H_\kappa^1)_u \mid \vec{\psi}^R = \vec{\psi}, \forall R \in O_{n-1} \}.$$

We note that, for $\vec{\psi} \in L_v^2(H_\kappa^1)_{\text{inv}}$, $\nabla_\kappa \vec{\psi} \in \vec{L}_v^2(L_\kappa^2)_{\text{inv}}$. The space $L_v^2(H_\kappa^1)_{\text{inv}}$ is a closed subspace of $L_v^2(H_\kappa^1)_u$ and hence, a Hilbert space, normed by $\| \cdot \|_{L_v^2(H_\kappa^1)_u}$.

(iv) The space $\mathcal{D}_{\text{inv}}(M)$ is the subspace of $\mathcal{D}_u(M)$ of elements $\vec{\phi}$ such that $\vec{\phi}^R = \vec{\phi}$. The space $\mathcal{D}'_{\text{inv}}(M)$ is the subspace of $\mathcal{D}'_u(M)$ of elements \vec{T} such that $\vec{T}^R = \vec{T}$. We note that $\mathcal{D}'_{\text{inv}}(M)$ is the dual of $\mathcal{D}_{\text{inv}}(M)$.

(v) The subspace of invariant elements of $L_v^2(H_\kappa^{-1})_u$ by the group O_{n-1} is given by

$$L_v^2(H_\kappa^{-1})_{\text{inv}} = \{ \vec{T} \in L_v^2(H_\kappa^{-1})_u \mid \vec{T}^R = \vec{T}, \forall R \in O_{n-1} \}.$$

It is the dual of $L_v^2(H_\kappa^1)_{\text{inv}}$. It is again a closed subspace of $L_v^2(H_\kappa^{-1})_u$ and hence, a Hilbert space, still normed by $\| \cdot \|_{L_v^2(H_\kappa^{-1})_u}$.

(vi) The subspace of invariant elements of \mathbb{Y}_u by the action of O_{n-1} is

$$\mathbb{Y}_{\text{inv}} = \{ \vec{\psi} \in \mathbb{Y}_u \mid \vec{\psi}^R = \vec{\psi}, \forall R \in O_{n-1} \}.$$

For $\vec{\psi} \in \mathbb{Y}_{\text{inv}}$, we have $\nabla_\kappa \vec{\psi} \in \vec{L}_v^2(L_\kappa^2)_{\text{inv}}$ and $\{ \vec{\psi}, \mathbf{H}_u \} \in L_v^2(H_\kappa^{-1})_{\text{inv}}$. The space \mathbb{Y}_{inv} is a closed subspace of \mathbb{Y}_u and hence, is a Hilbert space, still normed by $\| \cdot \|_{\mathbb{Y}_u}$.

Lemma 6.2. *We have*

$$\vec{\psi} \in L_v^2(L_\kappa^2)_{\text{inv}} \implies \int_M \vec{\psi} M_{\mathbf{u}} |\text{Vol}_M| = 0. \quad (6.22)$$

Proof. Let $\vec{\psi} \in L_v^2(L_\kappa^2)_{\text{inv}}$. Thanks to (6.6) and to the fact that $M_{\mathbf{u}}^R = M_{\mathbf{u}}$, $\forall R \in O_{n-1}$, we have

$$\int_M \vec{\psi} M_{\mathbf{u}} |\text{Vol}_M| = R \int_M \vec{\psi}^R M_{\mathbf{u}} |\text{Vol}_M| = R \int_M \vec{\psi} M_{\mathbf{u}} |\text{Vol}_M|, \quad \forall R \in O_{n-1}.$$

Hence, $\int_M \vec{\psi} M_{\mathbf{u}} |\text{Vol}_M|$ is a vector in $\{\mathbf{u}\}^\perp$, which is invariant by O_{n-1} . The only such vector is 0. \blacksquare

6.3.2 Rotational averages

We define projection maps from spaces of vector-valued functions and distributions on M to corresponding spaces of invariant functions. These projection maps are nothing but rotational averages.

(i) For $\vec{\varphi} \in \mathcal{D}_{\mathbf{u}}(M)$, we define its average $\vec{\varphi}_{\text{av}}$ over O_{n-1} by

$$\vec{\varphi}_{\text{av}} = \int_{O_{n-1}} \vec{\varphi}^R dR, \quad (6.23)$$

where dR is the normalized Haar measure on O_{n-1} . We have $\vec{\varphi}_{\text{av}} \in \mathcal{D}_{\text{inv}}(M)$. Furthermore, the map $\mathcal{D}_{\mathbf{u}}(M) \rightarrow \mathcal{D}_{\text{inv}}(M)$, $\vec{\varphi} \mapsto \vec{\varphi}_{\text{av}}$ is continuous for the topology of $\mathcal{D}_{\mathbf{u}}(M)$ and

$$\nabla_v \vec{\varphi}_{\text{av}} = (\nabla_v \vec{\varphi})_{\text{av}}, \quad \nabla_\kappa \vec{\varphi}_{\text{av}} = (\nabla_\kappa \vec{\varphi})_{\text{av}}.$$

(ii) Let $\vec{T} \in \mathcal{D}'_{\mathbf{u}}(M)$. Then, \vec{T}_{av} is the element of $\mathcal{D}'_{\text{inv}}(M)$ such that

$$(\vec{T}_{\text{av}}, \vec{\varphi})_{\mathcal{D}'_{\text{inv}}, \mathcal{D}_{\text{inv}}} = (\vec{T}, \vec{\varphi}_{\text{av}})_{\mathcal{D}'_{\mathbf{u}}, \mathcal{D}_{\mathbf{u}}} = \int_{O_{n-1}} (\vec{T}^R, \vec{\varphi})_{\mathcal{D}'_{\mathbf{u}}, \mathcal{D}_{\mathbf{u}}} dR = \int_{O_{n-1}} (\vec{T}, \vec{\varphi}^R)_{\mathcal{D}'_{\mathbf{u}}, \mathcal{D}_{\mathbf{u}}} dR, \quad (6.24)$$

where the last two equalities come from the linearity and continuity of \vec{T} . Thanks to the invariance of the Haar measure by matrix inversion, the definitions (6.24) and (6.23) are identical when $T \in \mathcal{D}_{\mathbf{u}}(M)$. The map $\mathcal{D}'_{\mathbf{u}}(M) \rightarrow \mathcal{D}'_{\text{inv}}(M)$, $\vec{T} \mapsto \vec{T}_{\text{av}}$ is continuous for the topology of $\mathcal{D}_{\mathbf{u}}(M)$ and we have

$$\nabla_v \vec{T}_{\text{av}} = (\nabla_v \vec{T})_{\text{av}}, \quad \nabla_\kappa \vec{T}_{\text{av}} = (\nabla_\kappa \vec{T})_{\text{av}}. \quad (6.25)$$

Lemma 6.3 (Properties of rotational averages).

(i) *We have*

$$\vec{\varphi} \in \mathbb{Y}_{\mathbf{u}} \implies \vec{\varphi}_{\text{av}} \in \mathbb{Y}_{\text{inv}}, \quad \text{and} \quad \|\vec{\varphi}_{\text{av}}\|_{\mathbb{Y}_{\text{inv}}} \leq \|\vec{\varphi}\|_{\mathbb{Y}_{\mathbf{u}}}. \quad (6.26)$$

(ii) For all $\vec{\varphi} \in \mathbb{Y}_{\mathbf{u}}$ and $\vec{\psi} \in \mathbb{Y}_{\text{inv}}$, we have

$$\int_{\text{O}_{n-1}} \left((\vec{\psi}, \vec{\varphi}^R) \right)_{L_v^2(L_\kappa^2)_{\mathbf{u}}} dR = \left((\vec{\psi}, \vec{\varphi}_{\text{av}}) \right)_{L_v^2(L_\kappa^2)_{\mathbf{u}}}, \quad (6.27)$$

$$\int_{\text{O}_{n-1}} \left((\nabla_\kappa \vec{\psi}, \nabla_\kappa \vec{\varphi}^R) \right)_{\tilde{L}_v^2(L_\kappa^2)_{\mathbf{u}}} dR = \left((\nabla_\kappa \vec{\psi}, \nabla_\kappa \vec{\varphi}_{\text{av}}) \right)_{\tilde{L}_v^2(L_\kappa^2)_{\mathbf{u}}}, \quad (6.28)$$

$$\int_{\text{O}_{n-1}} \left((\{\vec{\psi}, \mathbf{H}_{\mathbf{u}}\}, \vec{\varphi}^R) \right)_{L_v^2(H_\kappa^{-1})_{\mathbf{u}}, L_v^2(H_\kappa^1)_{\mathbf{u}}} dR = \left((\{\vec{\psi}, \mathbf{H}_{\mathbf{u}}\}, \vec{\varphi}_{\text{av}}) \right)_{L_v^2(H_\kappa^{-1})_{\mathbf{u}}, L_v^2(H_\kappa^1)_{\mathbf{u}}}. \quad (6.29)$$

Proof of Lemma 6.3. (i) *Step 1.* By Cauchy-Schwarz inequality, Fubini's theorem and (6.12), we have

$$\begin{aligned} \|\vec{\varphi}_{\text{av}}\|_{L_v^2(L_\kappa^2)_{\mathbf{u}}}^2 &= \int_M \left| \int_{\text{O}_{n-1}} \vec{\varphi}^R dR \right|^2 \mathbf{M}_{\mathbf{u}} |\text{Vol}_M| \leq \int_M \left(\int_{\text{O}_{n-1}} |\vec{\varphi}^R|^2 dR \right) \mathbf{M}_{\mathbf{u}} |\text{Vol}_M|, \\ &= \int_{\text{O}_{n-1}} \|\vec{\varphi}^R\|_{L_v^2(L_\kappa^2)_{\mathbf{u}}}^2 dR = \int_{\text{O}_{n-1}} \|\vec{\varphi}\|_{L_v^2(L_\kappa^2)_{\mathbf{u}}}^2 dR = \|\vec{\varphi}\|_{L_v^2(L_\kappa^2)_{\mathbf{u}}}^2. \end{aligned}$$

Here, for an element of X of $\{\mathbf{u}\}^\perp$, we have denoted by $|X|^2 = \sum_{i=1}^{n-1} (X \cdot e_i)^2$.

Step 2. Using Section 6.1.3 (iii) and the same arguments as just above, we can also prove

$$\|\nabla_\kappa \vec{\varphi}_{\text{av}}\|_{\tilde{L}_v^2(L_\kappa^2)_{\mathbf{u}}}^2 \leq \|\nabla_\kappa \vec{\varphi}\|_{\tilde{L}_v^2(L_\kappa^2)_{\mathbf{u}}}^2.$$

Step 3. We take $\vec{\varphi} \in L_v^2(H_\kappa^{-1})_{\mathbf{u}}$ and show that $\vec{\varphi}_{\text{av}} \in L_v^2(H_\kappa^{-1})_{\mathbf{u}}$. Indeed, for all $\vec{\psi} \in L_v^2(H_\kappa^1)_{\mathbf{u}}$, we have, thanks to the previous two steps

$$\begin{aligned} \left| \left((\vec{\varphi}_{\text{av}}, \vec{\psi}) \right)_{D'_{\mathbf{u}}, D_{\mathbf{u}}} \right| &= \left| \left((\vec{\varphi}, \vec{\psi}_{\text{av}}) \right)_{L_v^2(H_\kappa^{-1})_{\mathbf{u}}, L_v^2(H_\kappa^1)_{\mathbf{u}}} \right| \\ &\leq \|\vec{\varphi}\|_{L_v^2(H_\kappa^{-1})_{\mathbf{u}}} \|\vec{\psi}_{\text{av}}\|_{L_v^2(H_\kappa^1)_{\mathbf{u}}} \leq \|\vec{\varphi}\|_{L_v^2(H_\kappa^{-1})_{\mathbf{u}}} \|\vec{\psi}\|_{L_v^2(H_\kappa^1)_{\mathbf{u}}}, \end{aligned}$$

which shows that $\vec{\varphi}_{\text{av}} \in L_v^2(H_\kappa^{-1})_{\mathbf{u}}$ and

$$\|\vec{\varphi}_{\text{av}}\|_{L_v^2(H_\kappa^{-1})_{\mathbf{u}}} \leq \|\vec{\varphi}\|_{L_v^2(H_\kappa^{-1})_{\mathbf{u}}}. \quad (6.30)$$

Step 4. For $\vec{\varphi} \in D'_{\mathbf{u}}(M)$, we have, thanks to (6.25),

$$\{\vec{\varphi}_{\text{av}}, \mathbf{H}_{\mathbf{u}}\} = \{\vec{\varphi}, \mathbf{H}_{\mathbf{u}}\}_{\text{av}}.$$

From (6.30), it follows that if $\{\vec{\varphi}, \mathbf{H}_{\mathbf{u}}\} \in L_v^2(H_\kappa^{-1})_{\mathbf{u}}$, then, $\{\vec{\varphi}_{\text{av}}, \mathbf{H}_{\mathbf{u}}\} \in L_v^2(H_\kappa^{-1})_{\mathbf{u}}$ and

$$\|\{\vec{\varphi}_{\text{av}}, \mathbf{H}_{\mathbf{u}}\}\|_{L_v^2(H_\kappa^{-1})_{\mathbf{u}}} \leq \|\{\vec{\varphi}, \mathbf{H}_{\mathbf{u}}\}\|_{L_v^2(H_\kappa^{-1})_{\mathbf{u}}}.$$

Final step. Clearly, $\vec{\varphi}_{\text{av}}$ is invariant by O_{n-1} and collecting all the previous estimates, we get the estimate in the last statement of (6.26). It follows that $\vec{\varphi}_{\text{av}} \in \mathbb{Y}_{\text{inv}}$ which completes the proof of item (i).

(ii) Eqs. (6.27) and (6.28) follow from Fubini's theorem and (6.29), from (6.24). ■

6.3.3 Variational formulation for the vector GCI on invariant spaces

In this section, we show that the variational formulation for the vector GCI (5.19) can be equivalently formulated within spaces of invariant functions.

Lemma 6.4 (Variational formulation involving invariant spaces).

Let $\vec{\chi} \in L_v^2(L_\kappa^2)_{\text{inv}}$. Then, $\vec{\psi}$ is the unique solution of the variational formulation (5.19) if and only if it is the unique solution of the following variational formulation

$$\left\{ \begin{array}{l} \vec{\psi} \in \mathbb{Y}_{\text{inv}}, \\ -((\{\vec{\psi}, \mathbf{H}_u\}, \vec{\phi}))_{L_v^2(H_\kappa^{-1})_u, L_v^2(H_\kappa^1)_u} + D((\nabla_\kappa \vec{\psi}, \nabla_\kappa \vec{\phi}))_{\tilde{L}_v^2(L_\kappa^2)_u} \\ \qquad \qquad \qquad = -((\vec{\chi}, \vec{\phi}))_{L_v^2(L_\kappa^2)_u}, \quad \forall \vec{\phi} \in \mathbb{Y}_{\text{inv}}. \end{array} \right. \quad (6.31)$$

Proof. We first note that the variational formulation (6.31) has a unique solution because we can reproduce the proof of Theorem 5.6 using the invariant vector-valued version of the spaces used in the proof. We just note that, because of Lemma 6.2, an element of $\vec{\chi} \in L_v^2(L_\kappa^2)_{\text{inv}}$ or \mathbb{Y}_{inv} automatically satisfies $\int_M \vec{\chi} M_u |\text{Vol}_M| = 0$.

Now, let $\vec{\psi}$ be the unique solution of (5.19). Then, $\vec{\psi} \in \mathbb{Y}_{\text{inv}}$. Indeed, using $\vec{\phi}^R$ as a test function in the variational formulation (5.19) and applying the results of Section 6.1.3 and the fact that $\vec{\chi} \in L_v^2(L_\kappa^2)_{\text{inv}}$, we find that $\vec{\psi}^R$ is another solution of the same variational formulation. Since this variational formulation has a unique solution, we deduce that $\vec{\psi}^R = \vec{\psi}$. Furthermore, (5.19) is still valid by testing with function $\vec{\phi}$ in \mathbb{Y}_{inv} . So, $\vec{\psi}$ is also the unique solution of (6.31).

Conversely, Suppose $\vec{\psi}$ is the unique solution of (6.31). Then we obviously get $\vec{\psi} \in \mathbb{Y}_u$. Now, let us take $\vec{\phi} \in \mathbb{Y}_u$ and use $\vec{\phi}_{\text{av}} \in \mathbb{Y}_{\text{inv}}$ as a test function in (6.31). Then, by Lemma 6.3 (ii), we get

$$\int_{O_{n-1}} \left\{ -((\{\vec{\psi}, \mathbf{H}_u\}, \vec{\phi}^R))_{L_v^2(H_\kappa^{-1})_u, L_v^2(H_\kappa^1)_u} + D((\nabla_\kappa \vec{\psi}, \nabla_\kappa \vec{\phi}^R))_{\tilde{L}_v^2(L_\kappa^2)_u} + ((\vec{\chi}, \vec{\phi}^R))_{L_v^2(L_\kappa^2)_u} \right\} dR = 0.$$

But, using the results of Section 6.1.3 to transfer the exponent R onto $\vec{\psi}$ and the fact that $\vec{\psi}$ is invariant under the action of O_{n-1} , we end up with $\vec{\psi}$ being a solution of (5.19), which proves the claim. \blacksquare

7 Reduced GCI pair

In this section, we first show that, by the invariance property (6.21), the vector GCI can be reduced to two scalar quantities named the reduced GCI pair. Then from the variational formulation for the vector GCI in invariant spaces (6.31), we derive a variational formulation for the reduced GCI pair in appropriate function spaces.

7.1 From vector GCI to reduced GCI pair

In this subsection, we state the theorem establishing that the vector GCI can be expressed in terms of the reduced GCI pair. The proof of this theorem will be deferred to Appendix B. We start by introducing a convenient change of variables.

7.1.1 A change of variables in M

We consider the derivative of the diffeomorphism $\mathbf{V}_{\mathbf{u}}$ introduced in Section 4.1. We recall that we denote by $B_{\mathbf{u}} = B \setminus \{\pm \mathbf{u}\}$ and we denote by $M_{\mathbf{u}}$ the support manifold of $T B_{\mathbf{u}}$. We also recall that we identify $\mathbb{S}^{n-2} \approx B \cap \{\mathbf{u}\}^{\perp}$. We consider the tangent bundle $T(0, \pi)$ whose support manifold will be denoted by M_V (we note that $M_V \approx (0, \pi) \times \mathbb{R}$) and its elements by $(\theta, \kappa_{\parallel})$. Likewise, we consider the tangent bundle $T\mathbb{S}^{n-2}$ whose support manifold will be denoted by M_T and its elements, by (w, κ_T) . We now introduce the bundle diffeomorphism $\mathbf{A}_{\mathbf{u}}: M_V \times M_T \rightarrow M_{\mathbf{u}}$ given by $\mathbf{A}_{\mathbf{u}} = d\mathbf{V}_{\mathbf{u}}$, where $\mathbf{V}_{\mathbf{u}}$ is given by (4.1). Specifically, $\mathbf{A}_{\mathbf{u}}$ is given by

$$\mathbf{A}_{\mathbf{u}}((\theta, \kappa_{\parallel}), (w, \kappa_T)) = \alpha = (v, \kappa) = (\cos \theta \mathbf{u} + \sin \theta w, \kappa_{\parallel} e_{\theta} + \sin \theta \kappa_T), \quad (7.1)$$

with e_{θ} given by (4.3) and where

$$\kappa_T \cdot w = \kappa_T \cdot \mathbf{u} = 0, \quad (7.2)$$

the latter being a consequence of the fact that $(w, \kappa_T) \in M_T$. We note that

$$w = \frac{1}{\sin \theta}(v - \cos \theta \mathbf{u}), \quad e_{\theta} = \frac{1}{\sin \theta}(\cos \theta v - \mathbf{u}), \quad \kappa_{\parallel} = \kappa \cdot e_{\theta}, \quad \kappa_T \cdot e_{\theta} = 0. \quad (7.3)$$

The diffeomorphism $\mathbf{A}_{\mathbf{u}}$ is depicted in dimension $n = 3$ in Fig. 3. Now, thanks to the diffeomorphism $\mathbf{A}_{\mathbf{u}}$, for any smooth map $\varphi: M \rightarrow \mathbb{R}$, we can define $\varphi^{\dagger}: M_V \times M_T \rightarrow \mathbb{R}$ by

$$\varphi^{\dagger} = \varphi \circ \mathbf{A}_{\mathbf{u}}. \quad (7.4)$$

To compute the jacobian of $\mathbf{A}_{\mathbf{u}}$, we first need to specify the volume form on $M_V \times M_T$ and its orientation. We start with M_T and denote by $\alpha_T = (w, \kappa_T)$ a generic element of M_T and by $\xi^T = (\sigma^w, \sigma^{\kappa_T})$ a generic element of $T_{\alpha_T} M_T$, i.e. $(\sigma^w, \sigma^{\kappa_T}) \in \mathbb{R}^{2n}$ with

$$\sigma^w \cdot \mathbf{u} = 0, \quad \sigma^w \cdot w = 0, \quad (7.5)$$

$$\sigma^{\kappa_T} \cdot \mathbf{u} = 0, \quad \sigma^{\kappa_T} \cdot w + \sigma^w \cdot \kappa_T = 0. \quad (7.6)$$

We define a $n - 2$ -form ω on M_T in way similar to Ω (see (3.12)): we denote by π_T the bundle projection $M_T \rightarrow \mathbb{S}^{n-2}$ and, for $\xi_i^T \in T_{\alpha_T} M_T$, $i \in \{1, \dots, n - 2\}$, we let

$$\omega(\alpha_T)(\dots, \xi_i^T, \dots) = \text{Vol}_{\mathbb{S}^{n-2}}(\pi_T \alpha_T) \left(\dots, \bar{B}_{\alpha_T}^V(\xi_i^T), \dots \right), \quad (7.7)$$

where $\bar{B}_{\alpha_T}^V$ is the connection map $T_{\alpha_T} M_T \rightarrow T_{\pi_T \alpha_T} \mathbb{S}^{n-2}$ (the bar highlighting that it is different from the connection map for M). Like in the case of B , we have $\bar{B}_{\alpha_T}^V(\sigma^w, \sigma^{\kappa_T}) = P_{w^{\perp}} \sigma^{\kappa_T}$. Again,

$$\text{Vol}_{M_T} = \pi_T^* \text{Vol}_{\mathbb{S}^{n-2}} \wedge \omega. \quad (7.8)$$

is a volume form on M_T and we orient M_T by this volume form.

For $M_V \approx (0, \pi) \times \mathbb{R}$, we can proceed analogously, but since M_V is included in the flat space \mathbb{R}^2 , we can more simply define the volume form on M_V as $\text{Vol}_{M_V} = d\theta \wedge d\kappa_{\parallel}$, whose density is simply $d\theta d\kappa_{\parallel}$. We deduce that

$$|\text{Vol}_{M_V \times M_T}| = |\text{Vol}_{M_T}| d\theta d\kappa_{\parallel}. \quad (7.9)$$




Figure 3: The diffeomorphism $\mathbf{A}_{\mathbf{u}}$ depicted in dimension $n = 3$. The velocity v is changed into the pair (θ, w) where θ is the angle between \mathbf{u} and v and where w belongs to the unit circle \mathbb{S}^1 in the plane orthogonal to \mathbf{u} (diffeomorphism $\mathbf{V}_{\mathbf{u}}$ introduced in Section 4.1). The tangent vector κ to \mathbb{S}^2 at v is changed into the pair $(\kappa_{\parallel}, \kappa_T)$ where κ_{\parallel} is the projection of κ on e_{θ} , the director of the line normal to v in the plane $\text{Span}(\mathbf{u}, v)$, and κ_T is a tangent vector to \mathbb{S}^1 at w .

A computation using spherical coordinates leads to

$$|\mathbf{A}_{\mathbf{u}}^* \text{Vol}_M| = \sin^{2(n-2)} \theta |\text{Vol}_{M_T}| d\theta d\kappa_{\parallel}. \quad (7.10)$$

The retrenchment ω_w of ω on $T_w \mathbb{S}^{n-2}$ is computed similarly as that of Ω (see (3.16)). Let $(\varepsilon_1, \dots, \varepsilon_{n-2})$ be a local, positively oriented orthonormal frame in an open subset of \mathbb{S}^{n-2} containing w . We define a coordinate system on $T_w \mathbb{S}^{n-2}$ by $(\kappa_T^1, \dots, \kappa_T^{n-2})$ where $\kappa_T^i = \kappa_T \cdot \varepsilon_i$. Thus, we can apply (3.16) except that here, we are concerned with \mathbb{S}^{n-2} instead of \mathbb{S}^{n-1} . Hence, we have

$$\omega_w = d\kappa_T^1 \wedge \dots \wedge d\kappa_T^{n-2}. \quad (7.11)$$

7.1.2 Reduction of the vector GCI and reduced GCI pair

We state the main theorem establishing the expression of the vector GCI in terms of the reduced GCI pair. Its proof is given in Appendix B.

Theorem 7.1 (Reduction of the vector GCI).

Let $\vec{\psi} \in S(M, \{\mathbf{u}\}^{\perp})$ (the space of smooth functions $M \rightarrow \{\mathbf{u}\}^{\perp}$) satisfy the invariance relation (6.21) and let $\vec{\psi}^{\dagger}$ be obtained from $\vec{\psi}$ through the change of variables (7.4). Then, there exists a pair of functions: $(0, \pi) \times \mathbb{R} \times [0, \infty) \rightarrow \mathbb{R}^2$, $(\theta, \kappa_{\parallel}, \kappa_{\perp}) \rightarrow (\psi_w, \psi_r)$ such that

$$\begin{aligned} \vec{\psi}^{\dagger}(\theta, \kappa_{\parallel}, w, \kappa_T) &= \psi_w(\theta, \kappa_{\parallel}, |\kappa_T|) w + \psi_r(\theta, \kappa_{\parallel}, |\kappa_T|) \frac{\kappa_T}{|\kappa_T|}, \\ \forall ((\theta, \kappa_{\parallel}), (w, \kappa_T)) &\in M_v \times M_T \text{ such that } \kappa_T \neq 0, \end{aligned} \quad (7.12)$$

and

$$\vec{\psi}^\dagger(\theta, \kappa_\parallel, w, 0) = \psi_w(\theta, \kappa_\parallel, 0) w, \quad \forall ((\theta, \kappa_\parallel), w) \in M_V \times \mathbb{S}^{n-2}. \quad (7.13)$$

When $\vec{\psi} = \vec{\xi}_u$ is the vector GCI, the associated pair (ξ_w, ξ_r) is called the reduced GCI pair.

Remark 7.1 (Consistency of (7.12)).

From Section 6.2, we know that $\vec{\psi}$, w , κ_T all belong to $\{\mathbf{u}\}^\perp$, so that (7.12) makes sense.

Example 7.1 (Reduced pair for the function $\chi(v, \kappa) = P_{u^\perp} v$).

The function $\chi(v, \kappa) = P_{u^\perp} v$ satisfies the invariance relation (6.21). Its reduced pair is:

$$\chi_w(\theta, \kappa_\parallel, \kappa_\perp) = \sin \theta, \quad \chi_r(\theta, \kappa_\parallel, \kappa_\perp) = 0. \quad (7.14)$$

7.2 Variational formulation for the reduced GCI pair

Our goal is now to derive a variational formulation for the reduced GCI pair from the variational formulation for $\vec{\xi}_u$ involving invariant spaces (6.31). We will have to distinguish two cases: the generic case $n \geq 4$ and the special case $n = 3$. In the generic case, (ξ_w, ξ_r) are functions of $(\theta, \kappa_\parallel, \kappa_\perp) \in \mathcal{N} =: (0, \pi) \times \mathbb{R} \times (0, \infty)$. So, the variational formulation for the reduced GCI pair will involve appropriate function or distribution spaces over \mathcal{N} which we first introduce. Then, we establish this variational formulation and eventually derive the PDE and boundary conditions satisfied by the reduced GCI pair. The case $n = 3$ will require a similar yet slightly different treatment which is summarized at the end.

7.2.1 Spaces of functions on \mathcal{N} and properties

In this subsection, we assume that $n \geq 4$. We start with the definition of appropriate function spaces on \mathcal{N} and then, establish their link with spaces of invariant functions on M .

(i) For $p, q \in [0, \infty)$, we define the space

$$L_{p,q}^2 = \left\{ \psi : \mathcal{N} \rightarrow \mathbb{R}, \mid \|\psi\|_{L_{p,q}^2}^2 =: \int_{\mathcal{N}} |\psi|^2 \mathbf{M}_u^\dagger \kappa_\perp^q \sin^p \theta d\theta d\kappa_\parallel d\kappa_\perp < \infty \right\},$$

endowed with the Hilbert norm $\|\cdot\|_{L_{p,q}^2}$ and associated inner-product $(\cdot, \cdot)_{L_{p,q}^2}$. We recall that \mathbf{M}_u^\dagger is given by (4.18). We denote by $\mathbb{L} = L_{2(n-2), n-3}^2$ and the associated norm and inner product by $\|\cdot\|_{\mathbb{L}}$ and $(\cdot, \cdot)_{\mathbb{L}}$ respectively.

(ii) We define the space

$$\mathbb{H} = \left\{ \psi \in \mathbb{L} \text{ such that } \|\psi\|_{\mathbb{H}}^2 =: \|\psi\|_{\mathbb{L}}^2 + \|\partial_{\kappa_\parallel} \psi\|_{\mathbb{L}}^2 + \|\partial_{\kappa_\perp} \psi\|_{L_{2(n-3), n-3}^2}^2 < \infty \right\},$$

endowed with the Hilbert norm $\|\cdot\|_{\mathbb{H}}$ and associated inner-product $(\cdot, \cdot)_{\mathbb{H}}$. We define the semi-norm

$$\|\psi\|_{\mathbb{H}}^2 = \|\partial_{\kappa_\parallel} \psi\|_{\mathbb{L}}^2 + \|\partial_{\kappa_\perp} \psi\|_{L_{2(n-3), n-3}^2}^2,$$

so that

$$\|\psi\|_{\mathbb{H}}^2 = \|\psi\|_{\mathbb{L}}^2 + \|\psi\|_{\mathbb{H}}^2.$$

We denote by $((\cdot, \cdot))_{\mathbb{H}}^\bullet$ the associated bilinear form.

We denote by $\mathcal{D}(\mathcal{N})$ the space of C^∞ functions with compact support on \mathcal{N} . We note the following important property, whose proof uses classical functional analytic arguments and is omitted:

Lemma 7.2 (Density of $\mathcal{D}(\mathcal{N})$ in \mathbb{H} when $n \geq 4$).

Suppose $n \geq 4$. Then, $\mathcal{D}(\mathcal{N})$ is dense in \mathbb{H} .

Remark 7.2 (About Lemma 7.2).

The domain \mathcal{N} has a boundary at $\kappa_\perp = 0$ and the definition of \mathbb{H} involves derivatives in κ_\perp . If $n \geq 4$ the inner product in the definition of \mathbb{H} involves the weight κ_\perp^{n-3} with an exponent $n-3 \geq 1$. Because of this weight, we can prove that $\mathcal{D}(\mathcal{N})$ is dense in \mathbb{H} . If $n = 3$, this weight disappears and \mathbb{H} is akin to a classical Sobolev space H^1 (i.e. without any weight in the inner-product) with respect to the κ_\perp variable. Thus, in this case, $\mathcal{D}(\mathcal{N})$ is not dense in \mathbb{H} . Whether $\mathcal{D}(\mathcal{N})$ is dense in \mathbb{H} or not impacts the structure of its dual space. Indeed, dual spaces of classical H^1 spaces cannot be identified with spaces of distributions when the domain involves boundaries. By contrast, duals of H_0^1 -type spaces (which are the closures of \mathcal{D} for the Sobolev norm) can (see [2, Sections 3.7-3.14]). In the forthcoming sections, we will use the dual space of \mathbb{H} and it will be of critical importance that it is a space of distributions, hence the restriction to $n \geq 4$. The case $n = 3$ will require additional developments to circumvent the non-density of $\mathcal{D}(\mathcal{N})$ in \mathbb{H} . There is no such problem with the other variables: the domain in k_\parallel is \mathbb{R} and has no boundary; the domain in θ is $(0, \pi)$ and has boundaries but the definition of \mathbb{H} does not involve any derivative in θ .

(iii) We define the space $\mathbb{K} = \mathbb{H} \cap L_{2(n-3), n-5}^2$, endowed with the norm

$$\|\psi\|_{\mathbb{K}}^2 = \|\psi\|_{\mathbb{H}}^2 + (n-3) \|\psi\|_{L_{2(n-3), n-5}^2}^2.$$

The associated inner-product is denoted by $((\cdot, \cdot))_{\mathbb{K}}$. We define the semi-norm

$$\|\psi\|_{\mathbb{K}}^\bullet = \|\psi\|_{\mathbb{H}}^\bullet + (n-3) \|\psi\|_{L_{2(n-3), n-5}^2}^\bullet,$$

so that

$$\|\psi\|_{\mathbb{K}}^2 = \|\psi\|_{\mathbb{L}}^2 + \|\psi\|_{\mathbb{K}}^\bullet.$$

We denote by $((\cdot, \cdot))_{\mathbb{K}}^\bullet$ the associated bilinear form. Like for \mathbb{H} , we have the

Lemma 7.3 (Density of $\mathcal{D}(\mathcal{N})$ in \mathbb{K} when $n \geq 4$).

Suppose $n \geq 4$. Then, $\mathcal{D}(\mathcal{N})$ is dense in \mathbb{K} .

(iv) We define

$$\mathcal{L}^2(\mathcal{L}^2) = \mathbb{L}^2,$$

and denote its elements by (ψ_w, ψ_r) . It is endowed with the norm

$$\|(\psi_w, \psi_r)\|_{\mathcal{L}^2(\mathcal{L}^2)}^2 = \|\psi_w\|_{\mathbb{L}}^2 + \|\psi_r\|_{\mathbb{L}}^2.$$

(v) We define

$$\mathcal{L}^2(\mathcal{H}^1) = \mathbb{H} \times \mathbb{K}$$

endowed with the norm

$$\|(\psi_w, \psi_r)\|_{\mathcal{L}^2(\mathcal{H}^1)}^2 = \|\psi_w\|_{\mathbb{H}}^2 + \|\psi_r\|_{\mathbb{K}}^2.$$

and we have

$$\|(\psi_w, \psi_r)\|_{\mathcal{L}^2(\mathcal{H}^1)}^2 = \|(\psi_w, \psi_r)\|_{\mathcal{L}^2(\mathcal{L}^2)}^2 + \|\psi_w\|_{\mathbb{H}}^2 + \|\psi_r\|_{\mathbb{K}}^2.$$

In view of Lemmas 7.2 and 7.3, since $n \geq 4$, the space $\mathcal{D}(\mathcal{N})^2$ is dense in $\mathcal{L}^2(\mathcal{H}^1)$.

We now establish the link between invariant function spaces on M and function spaces on \mathcal{N} .

Proposition 7.4 (Link between $L_v^2(L_\kappa^2)_{\text{inv}}$ (resp. $L_v^2(H_\kappa^1)_{\text{inv}}$) and $\mathcal{L}^2(\mathcal{L}^2)$ (resp. $\mathcal{L}^2(\mathcal{H}^1)$)).

(i) The function $\tilde{\psi}$ belongs to $L_v^2(L_\kappa^2)_{\text{inv}}$ if and only if $\tilde{\psi}^\dagger$ (related to $\tilde{\psi}$ through (7.4)) is written

$$\begin{aligned} \tilde{\psi}^\dagger(\theta, \kappa_\parallel, w, \kappa_T) &= \psi_w(\theta, \kappa_\parallel, |\kappa_T|) w + \psi_r(\theta, \kappa_\parallel, |\kappa_T|) \frac{\kappa_T}{|\kappa_T|}, \\ &\text{for } (\theta, \kappa_\parallel, w, \kappa_T) \in M_V \times M_T \text{ a.e.}, \end{aligned} \quad (7.15)$$

(where a.e. stands for “almost everywhere”) and $(\psi_w, \psi_r) \in \mathcal{L}^2(\mathcal{L}^2)$. Furthermore, defining c as

$$c = |\mathbb{S}^{n-2}| |\mathbb{S}^{n-3}|, \quad (7.16)$$

we have

$$\|\tilde{\psi}\|_{L_v^2(L_\kappa^2)_u}^2 = c \|(\psi_w, \psi_r)\|_{\mathcal{L}^2(\mathcal{L}^2)}^2. \quad (7.17)$$

(ii) The function $\tilde{\psi}$ belongs to $L_v^2(H_\kappa^1)_{\text{inv}}$ if and only if $\tilde{\psi}^\dagger$ is given by (7.15) and $(\psi_w, \psi_r) \in \mathcal{L}^2(\mathcal{H}^1)$. Furthermore,

$$\|\tilde{\psi}\|_{L_v^2(H_\kappa^1)_u}^2 = c \|(\psi_w, \psi_r)\|_{\mathcal{L}^2(\mathcal{H}^1)}^2, \quad (7.18)$$

and

$$\|\nabla_\kappa \tilde{\psi}\|_{L_v^2(L_\kappa^2)_u}^2 = c \left(\|\psi_w\|_{\mathbb{H}}^2 + \|\psi_r\|_{\mathbb{K}}^2 \right). \quad (7.19)$$

Proof. (i) Let $\tilde{\psi} \in \mathcal{D}_{\text{inv}}(M)$. Then, Theorem 7.1 applies and we get (7.15). We have

$$|\tilde{\psi}^\dagger|^2(\theta, \kappa_\parallel, w, \kappa_T) = (|\psi_w|^2 + |\psi_r|^2)(\theta, \kappa_\parallel, |\kappa_T|), \quad \text{for } (\theta, \kappa_\parallel, w, \kappa_T) \in M_V \times M_T \text{ a.e.}$$

Hence, using (7.10), and the Fubini theorem for fiber bundles (3.15), we get

$$\begin{aligned} \int_M |\tilde{\psi}|^2 \mathbf{M}_u |\text{Vol}_M| &= \int_{M_V} \left(\int_{M_T} |\tilde{\psi}^\dagger|^2 \mathbf{M}_u^\dagger |\text{Vol}_{M_T}| \right) \sin^{2(n-2)} \theta d\theta d\kappa_\parallel \\ &= \int_{M_V} \left(\int_{M_T} (|\psi_w|^2 + |\psi_r|^2) \mathbf{M}_u^\dagger |\text{Vol}_{M_T}| \right) \sin^{2(n-2)} \theta d\theta d\kappa_\parallel \\ &= \int_{M_V} \left[\int_{\mathbb{S}^{n-2}} \left(\int_{T_w \mathbb{S}^{n-2}} (|\psi_w|^2 + |\psi_r|^2) \mathbf{M}_u^\dagger |\omega_w| \right) |\text{Vol}_B| \right] \sin^{2(n-2)} \theta d\theta d\kappa_\parallel, \end{aligned}$$

where ω_w is the retrenchment of ω on $T_w \mathbb{S}^{n-2}$, given by (7.11). Recalling the expression (8.8) for \mathbf{M}_u^\dagger , we see that the expression inside the inner integral only depends on $|\kappa_T|$. Hence, we

can use spherical coordinates in $T_w \mathbb{S}^{n-2} \approx \mathbb{R}^{n-2}$ for the inner integral and find that the result is independent of w . Thus, the integral over \mathbb{S}^{n-2} is just that of a constant and we get

$$\int_M |\vec{\psi}|^2 \mathbf{M}_u |\text{Vol}_M| = c \int_{M_V} \left(\int_0^\infty (|\psi_w|^2 + |\psi_r|^2) \mathbf{M}_u^\natural)(\theta, \kappa_\parallel, \kappa_\perp) \kappa_\perp^{n-3} d\kappa_\perp \right) \sin^{2(n-2)} \theta d\theta d\kappa_\parallel,$$

with \mathbf{M}_u^\natural given by (4.18). This is nothing but (7.17). Now, (7.15) and (7.17) extend to $\vec{\psi} \in L_v^2(L_\kappa^2)_{\text{inv}}$ by density of $\mathcal{D}_{\text{inv}}(M)$ in $L_v^2(L_\kappa^2)_{\text{inv}}$. Conversely, if $(\psi_w, \psi_r) \in \mathcal{L}^2(\mathcal{L}^2)$, then, $\vec{\psi}$ given in terms of $\vec{\psi}^\dagger$ by (7.15) is invariant by O_{n-1} and belongs to $L_v^2(L_\kappa^2)_u$ by virtue of (7.17). Thus it is an element of $L_v^2(L_\kappa^2)_{\text{inv}}$.

(ii) Let $\vec{\psi} \in \mathcal{D}_{\text{inv}}(M)$. Then, the following formula is proven in Appendix C, Proposition C.4(i)

$$|(\nabla_\kappa \vec{\psi})^\dagger|^2 = |\partial_{\kappa_\parallel} \psi_w|^2 + |\partial_{\kappa_\parallel} \psi_r|^2 + \frac{1}{\sin^2 \theta} (|\partial_{\kappa_\perp} \psi_w|^2 + |\partial_{\kappa_\perp} \psi_r|^2) + \frac{n-3}{\sin^2 \theta |\kappa_T|^2} |\psi_r|^2. \quad (7.20)$$

Now, inserting this into the integral over M , we can reproduce the calculation made in the proof of item (i) and we easily obtain (7.18) showing that $(\psi_w, \psi_r) \in \mathcal{L}^2(\mathcal{H}^1)$. As in the proof of item (i), the converse is obvious. ■

7.2.2 Spaces of distributions on \mathcal{N} and properties

In this subsection, we still assume that $n \geq 4$. We follow the same steps as in Subsection 7.2.1: we define distribution spaces on \mathcal{N} and then, explore their link with spaces of invariant distributions on M .

(i) We denote by $\mathcal{D}'(\mathcal{N})$ the dual of $\mathcal{D}(\mathcal{N})$, i.e. the space of distributions on \mathcal{N} . We define the duality bracket $\langle\langle T, \varphi \rangle\rangle_{\mathcal{D}', \mathcal{D}}$ between $T \in \mathcal{D}'(\mathcal{N})$ and $\varphi \in \mathcal{D}(\mathcal{N})$ so as to extend the inner-product of \mathbb{L} .

(ii) We define the space \mathbb{H}' as the dual of \mathbb{H} . By Lemma 7.2, $\mathcal{D}(\mathcal{N})$ is dense in \mathbb{H} and so, we can identify \mathbb{H}' to the space of distributions $T \in \mathcal{D}'(\mathcal{N})$ such that

$$\|T\|_{\mathbb{H}'} =: \sup_{\varphi \in \mathcal{D}(\mathcal{N}), \varphi \neq 0} \frac{|\langle\langle T, \varphi \rangle\rangle_{\mathcal{D}', \mathcal{D}}|}{\|\varphi\|_{\mathbb{H}}} < \infty,$$

endowed with $\|\cdot\|_{\mathbb{H}'}$ (see [2, Sections 3.7-3.14]).

(iii) We define the space \mathbb{K}' as the dual of \mathbb{K} . For the same reason as for \mathbb{H}' , we can identify \mathbb{K}' with the space of distribution $T \in \mathcal{D}'(\mathcal{N})$ such that

$$\|T\|_{\mathbb{K}'} =: \sup_{\varphi \in \mathcal{D}(\mathcal{N}), \varphi \neq 0} \frac{|\langle\langle T, \varphi \rangle\rangle_{\mathcal{D}', \mathcal{D}}|}{\|\varphi\|_{\mathbb{K}}} < \infty,$$

endowed with $\|\cdot\|_{\mathbb{K}'}$.

(iv) Let $\vec{T} \in \mathcal{D}'_{\text{inv}}(M)$. We define two distributions T_w and T_r in $\mathcal{D}'(\mathcal{N})$ such that for all φ in $\mathcal{D}(\mathcal{N})$,

$$\langle\langle T_w, \varphi \rangle\rangle_{\mathcal{D}', \mathcal{D}} = c^{-1} \langle\langle \vec{T}, (w\varphi) \circ \mathbf{A}_{\mathbf{u}}^{-1} \rangle\rangle_{\mathcal{D}'_{\mathbf{u}}, \mathcal{D}_{\mathbf{u}}}, \quad (7.21)$$

$$\langle\langle T_r, \varphi \rangle\rangle_{\mathcal{D}', \mathcal{D}} = c^{-1} \langle\langle \vec{T}, (\frac{\kappa_T}{|\kappa_T|} \varphi) \circ \mathbf{A}_{\mathbf{u}}^{-1} \rangle\rangle_{\mathcal{D}'_{\mathbf{u}}, \mathcal{D}_{\mathbf{u}}}, \quad (7.22)$$

where we refer to Section 6.2 for the definition of $\mathbf{A}_{\mathbf{u}}$, and to Eq. (7.16) for that of c . At the right-hand sides of these formulas, φ is considered as a function on $M_V \times M_T$ rather than just a function on \mathcal{N} . We note that since φ has compact support on \mathcal{N} , κ_T is never zero on the support of φ , so that $(w\varphi) \circ \mathbf{A}_{\mathbf{u}}^{-1}$ and $((\kappa_T/|\kappa_T|)\varphi) \circ \mathbf{A}_{\mathbf{u}}^{-1}$ are elements of $\mathcal{D}_{\text{inv}}(M)$ and the definitions of T_w and T_r make sense. Furthermore, if $\vec{T} \in \mathcal{D}_{\text{inv}}(M)$, it can be written $\vec{T}^\dagger = T_w w + T_r (\kappa_T/|\kappa_T|)$ and, thanks to the definition of the duality bracket for $\mathcal{D}'(\mathcal{N})$, this definition of (T_w, T_r) coincides with (7.21), (7.22). Now, for $\vec{\varphi} \in \mathcal{D}_{\text{inv}}(M)$, we have the characterization $\vec{\varphi}^\dagger = \varphi_w w + \varphi_r (\kappa_T/|\kappa_T|)$. So, we can write

$$\langle\langle \vec{T}, \vec{\varphi} \rangle\rangle_{\mathcal{D}'_{\mathbf{u}}, \mathcal{D}_{\mathbf{u}}} = c \left(\langle\langle T_w, \varphi_w \rangle\rangle_{\mathcal{D}', \mathcal{D}} + \langle\langle T_r, \varphi_r \rangle\rangle_{\mathcal{D}', \mathcal{D}} \right). \quad (7.23)$$

(v) We define

$$\mathcal{L}^2(\mathcal{H}^{-1}) = \mathbb{H}' \times \mathbb{K}'.$$

It is the dual of $\mathcal{L}^2(\mathcal{H}^1)$. Since $n \geq 4$, the space $\mathcal{D}(\mathcal{N})^2$ is dense in $\mathcal{L}^2(\mathcal{H}^1)$ and consequently, $\mathcal{L}^2(\mathcal{H}^{-1})$ can be identified with the space of pairs of distributions (T_w, T_r) such that

$$\|(T_w, T_r)\|_{\mathcal{L}^2(\mathcal{H}^{-1})} \quad (7.24)$$

$$=: \sup_{(\varphi_w, \varphi_r) \in \mathcal{D}(\mathcal{N})^2, (\varphi_w, \varphi_r) \neq 0} \frac{|\langle\langle T_w, \varphi_w \rangle\rangle_{\mathbb{H}', \mathbb{H}} + \langle\langle T_r, \varphi_r \rangle\rangle_{\mathbb{K}', \mathbb{K}}|}{(\|\varphi_w\|_{\mathbb{H}}^2 + \|\varphi_r\|_{\mathbb{K}}^2)^{1/2}} \quad (7.25)$$

$$= (\|T_w\|_{\mathbb{H}'}^2 + \|T_r\|_{\mathbb{K}'}^2)^{1/2} < \infty. \quad (7.26)$$

The space $\mathcal{L}^2(\mathcal{H}^{-1})$ is normed by $\|\cdot\|_{\mathcal{L}^2(\mathcal{H}^{-1})}$. We have

$$\vec{T} \in L_v^2(H_{\kappa}^{-1})_{\text{inv}} \iff (T_w, T_r) \in \mathcal{L}^2(\mathcal{H}^{-1}), \quad (7.27)$$

and

$$\|\vec{T}\|_{L_v^2(H_{\kappa}^{-1})_{\mathbf{u}}}^2 = c \|(T_w, T_r)\|_{\mathcal{L}^2(\mathcal{H}^{-1})}^2, \quad (7.28)$$

(vi) We refer to (4.21) for the definition of operator \mathcal{U}_1 . The space $\mathbb{Y}_{\mathcal{N}}$ is defined by

$$\mathbb{Y}_{\mathcal{N}} = \{(\psi_w, \psi_r) \in \mathcal{L}^2(\mathcal{H}^1) \text{ such that } (\mathcal{U}_1 \psi_w - \kappa_{\perp} \psi_r, \mathcal{U}_1 \psi_r + \kappa_{\perp} \psi_w) \in \mathcal{L}^2(\mathcal{H}^{-1})\},$$

where $\mathcal{U}_1 \psi_w$ and $\mathcal{U}_1 \psi_r$ are defined in the distributional sense (see (7.31) below), endowed with the norm

$$\|(\psi_w, \psi_r)\|_{\mathbb{Y}_{\mathcal{N}}}^2 = \|(\psi_w, \psi_r)\|_{\mathcal{L}^2(\mathcal{H}^1)}^2 + \|(\mathcal{U}_1 \psi_w - \kappa_{\perp} \psi_r, \mathcal{U}_1 \psi_r + \kappa_{\perp} \psi_w)\|_{\mathcal{L}^2(\mathcal{H}^{-1})}^2. \quad (7.29)$$

We now explore the Link between \mathbb{Y}_{inv} and $\mathbb{Y}_{\mathcal{N}}$. We first explore some properties of the operator \mathcal{U}_1 .

Lemma 7.5 (\mathcal{U}_1 is formally skew-adjoint).

(i) For any φ, ψ in $\mathcal{D}(\mathcal{N})$, we have

$$((\mathcal{U}_1 \varphi, \psi))_{\mathbb{L}} = -((\varphi, \mathcal{U}_1 \psi))_{\mathbb{L}}. \quad (7.30)$$

(ii) Let $T \in \mathcal{D}'(\mathcal{N})$. Then the operator \mathcal{U}_1 applied to T in the distributional sense satisfies

$$((\mathcal{U}_1 T, \varphi))_{\mathcal{D}', \mathcal{D}} = -((T, \mathcal{U}_1 \varphi))_{\mathcal{D}', \mathcal{D}}, \quad \forall \varphi \in \mathcal{D}(\mathcal{N}). \quad (7.31)$$

(iii) For $\vec{T} \in \mathcal{D}'_{\text{inv}}(M)$, we have

$$\{\vec{T}, \mathbf{H}_{\mathbf{u}}\}^{\dagger} = \left(w(\mathcal{U}_1 T_w - |\kappa_T| T_r) + \frac{\kappa_T}{|\kappa_T|} (\mathcal{U}_1 T_r + |\kappa_T| T_w) \right), \quad (7.32)$$

where the operator \mathcal{U}_1 is applied to T_w and T_r in the distributional sense, by virtue of (7.31)

Proof. (ii) is an immediate consequence of (i). To show (i), we first remark that for $\vec{\psi} \in \mathcal{D}_{\text{inv}}$, we have (see Appendix C, Proposition C.4(ii))

$$\{\vec{\psi}, \mathbf{H}_{\mathbf{u}}\}^{\dagger} = w(\mathcal{U}_1 \psi_w - |\kappa_T| \psi_r) + \frac{\kappa_T}{|\kappa_T|} (\mathcal{U}_1 \psi_r + |\kappa_T| \psi_w), \quad (7.33)$$

Now, let φ, ψ in $\mathcal{D}(\mathcal{N})$ and define $\vec{\varphi} = (w\varphi) \circ \mathbf{A}_{\mathbf{u}}^{-1}$ and define $\vec{\psi}$ in a similar way. Then, $\vec{\varphi}, \vec{\psi}$ belong to $\mathcal{D}_{\text{inv}}(M)$ and, thanks to (7.33), we have

$$\{\vec{\varphi}, \mathbf{H}_{\mathbf{u}}\} = (w\mathcal{U}_1 \varphi + \kappa_T \varphi) \circ \mathbf{A}_{\mathbf{u}}^{-1},$$

and similarly for $\{\vec{\psi}, \mathbf{H}_{\mathbf{u}}\}$. Now, from (5.16) and given the regularity of φ and ψ , we deduce that

$$(((w\mathcal{U}_1 \varphi + \kappa_T \varphi) \circ \mathbf{A}_{\mathbf{u}}^{-1}, (w\psi) \circ \mathbf{A}_{\mathbf{u}}^{-1}))_{L_v^2(L_k^2)} = -(((w\varphi) \circ \mathbf{A}_{\mathbf{u}}^{-1}, (w\mathcal{U}_1 \psi + \kappa_T \psi) \circ \mathbf{A}_{\mathbf{u}}^{-1}))_{L_v^2(L_k^2)},$$

which leads to (7.30) since w and κ_T are orthogonal.

(iii) Let $\vec{T} \in \mathcal{D}'_{\text{inv}}(M)$ and $\vec{\psi} \in \mathcal{D}_{\text{inv}}(M)$. Thanks to (5.16), (7.23), (7.33) and (7.31), we have

$$\begin{aligned} ((\{\vec{T}, \mathbf{H}_{\mathbf{u}}\}, \vec{\psi}))_{\mathcal{D}'_{\mathbf{u}}, \mathcal{D}_{\mathbf{u}}} &= -((\vec{T}, \{\vec{\psi}, \mathbf{H}_{\mathbf{u}}\}))_{\mathcal{D}'_{\mathbf{u}}, \mathcal{D}_{\mathbf{u}}} \\ &= -c \left[((T_w, \mathcal{U}_1 \psi_w - \kappa_{\perp} \psi_r))_{\mathcal{D}', \mathcal{D}} + ((T_r, \mathcal{U}_1 \psi_r + \kappa_{\perp} \psi_w))_{\mathcal{D}', \mathcal{D}} \right] \\ &= c \left[((\mathcal{U}_1 T_w - \kappa_{\perp} T_r, \psi_w))_{\mathcal{D}', \mathcal{D}} + ((\mathcal{U}_1 T_r + \kappa_{\perp} T_w, \psi_r))_{\mathcal{D}', \mathcal{D}} \right]. \end{aligned}$$

Thanks to (7.23), this last expression is equivalent to (7.32). ■

Remark 7.3 (Direct computation).

Identity (7.30) can be cross-checked by a straightforward, albeit tedious, direct computation.

With this lemma, we can show the

Proposition 7.6 (Link between \mathbb{Y}_{inv} and $\mathbb{Y}_{\mathcal{N}}$).

The function $\vec{\psi}$ belongs to \mathbb{Y}_{inv} if and only if $\vec{\psi}$ is written according to (7.15) and (ψ_w, ψ_r) belongs to $\mathbb{Y}_{\mathcal{N}}$. Furthermore,

$$\|\vec{\psi}\|_{\mathbb{Y}_{\text{inv}}}^2 = c \|(\psi_w, \psi_r)\|_{\mathbb{Y}_{\mathcal{N}}}^2. \quad (7.34)$$

Proof. Let $\vec{\psi} \in \mathbb{Y}_{\text{inv}}$. By Proposition 7.4, $\vec{\psi}$ is given by (7.15) with $(\psi_w, \psi_r) \in \mathcal{L}^2(\mathcal{H}^1)$. Now, thanks to (7.27) and to (7.32), $\{\vec{\psi}, \mathbf{H}_u\} \in L_v^2(H_\kappa^{-1})$ if and only if $(\mathcal{U}_1\psi_w - \kappa_\perp\psi_r, \mathcal{U}_1\psi_r + \kappa_\perp\psi_w) \in \mathcal{L}^2(\mathcal{H}^{-1})$, which shows that $(\psi_w, \psi_r) \in \mathbb{Y}_{\mathcal{N}}$. We can move backwards this chain of implications, which shows the claimed equivalence. The equality (7.34) follows from (7.29), (7.17) and (7.28). ■

7.2.3 Expression of the variational formulation for the reduced GCI pair

Again, in this subsection, we assume that $n \geq 4$. We establish a variational formulation solved by the reduced GCI pair and derive the corresponding system of PDE (4.19), (4.20) in strong form.

Theorem 7.7 (Variational formulation).

Suppose $n \geq 4$. Let $\vec{\chi} \in L_v^2(L_\kappa^2)_{\text{inv}}$ and define $(\chi_w, \chi_r) \in \mathcal{L}^2(\mathcal{L}^2)$ as in (7.15). Then, $\vec{\psi}$ is the unique solution of the variational formulation (6.31) if and only if (ψ_w, ψ_r) linked to $\vec{\psi}$ by (7.15) is the unique solution of the variational formulation

$$\left\{ \begin{array}{l} (\psi_w, \psi_r) \in \mathbb{Y}_{\mathcal{N}}, \\ -\langle (\mathcal{U}_1\psi_w - \kappa_\perp\psi_r, \varphi_w) \rangle_{\mathbb{H}', \mathbb{H}} - \langle (\mathcal{U}_1\psi_r + \kappa_\perp\psi_w, \varphi_r) \rangle_{\mathbb{K}', \mathbb{K}} + D((\psi_w, \varphi_w))_{\mathbb{H}}^\bullet \\ \quad + D((\psi_r, \varphi_r))_{\mathbb{K}}^\bullet = -\langle (\chi_w, \varphi_w) \rangle_{\mathbb{L}} - \langle (\chi_r, \varphi_r) \rangle_{\mathbb{L}}, \quad \forall (\varphi_w, \varphi_r) \in \mathbb{Y}_{\mathcal{N}}. \end{array} \right. \quad (7.35)$$

Proof. This is a direct consequence of (7.32), (7.23) and the counterparts for bilinear forms of (7.19) and (7.17). The variational formulation (7.35) has a unique solution since (6.31) has a unique solution. ■

Remark 7.4 (About the variational formulation (7.35)).

The condition $(\varphi_w, \varphi_r) \in \mathbb{Y}_{\mathcal{N}}$ does not imply that $(\varphi_w, 0)$ nor $(0, \varphi_r)$ separately belong to $\mathbb{Y}_{\mathcal{N}}$. This is because φ_w and φ_r are entangled in the condition $(\mathcal{U}_1\varphi_w - \kappa_\perp\varphi_r, \mathcal{U}_1\varphi_r + \kappa_\perp\varphi_w) \in \mathcal{L}^2(\mathcal{H}^{-1})$ of the definition of $\mathbb{Y}_{\mathcal{N}}$. Thus, if we separate (7.35) in two equations by taking either $\varphi_r = 0$ or $\varphi_w = 0$, we obtain a different variational formulation, which may be ill-posed.

From Theorem 7.7, we immediately get the following

Corollary 7.8 (Variational formulation satisfied by the reduced GCI pair).

Let (ζ_w, ζ_r) be the reduced GCI pair. Then, (ζ_w, ζ_r) satisfies the variational formulation (7.35), with (ψ_w, ψ_r) replaced by (ζ_w, ζ_r) and (χ_w, χ_r) given by (7.14).

In the following theorem, we provide a strong formulation of the problem associated with the variational formulation (7.35). It involves the derivation of both the PDE and its boundary conditions. The latter are obtained formally assuming that the solution of the PDE is smooth, a property that is still open.

Theorem 7.9 (PDE system associated with the variational formulation (7.35) and boundary conditions).

We assume that $n \geq 4$.

(i) Let $(\zeta_w, \zeta_r) \in \mathbb{Y}_{\mathcal{N}}$ be the unique solution to the variational formulation (7.35), with (ψ_w, ψ_r) replaced by (ζ_w, ζ_r) and (χ_w, χ_r) given by (7.14). Then, (ζ_w, ζ_r) satisfies System (4.19), (4.20) in the distributional sense.

(ii) (formal) Supposing that (ζ_w, ζ_r) is smooth. Then the boundary conditions associated to the variational formulation (7.35) are:

- no boundary condition at $\theta = 0$ nor $\theta = \pi$ on neither ζ_w nor ζ_r ;
- dimension $n = 4$: no boundary condition on ζ_w and homogeneous Dirichlet condition on ζ_r at $\kappa_{\perp} = 0$:

$$\zeta_r(\theta, \kappa_{\parallel}, 0) = 0; \quad (7.36)$$

- dimension $n \geq 5$: no boundary condition at $\kappa_{\perp} = 0$ on neither ζ_w nor ζ_r .

Proof. (i) Let $\varphi \in \mathcal{D}(\mathcal{N})$. Then, $(\varphi, 0) \in \mathbb{Y}_{\mathcal{N}}$ and $(0, \varphi) \in \mathbb{Y}_{\mathcal{N}}$. Thus, using these two pairs as test functions in (7.35) and interpreting the various bilinear forms as duality between distributions and test functions, we are led straightforwardly to (4.19), (4.20).

(ii) Define $\mathcal{D}(\bar{\mathcal{N}})$ as the set of C^{∞} functions with compact support on an open set containing the closure $\bar{\mathcal{N}}$ of \mathcal{N} . We now investigate under which conditions on $\varphi \in \mathcal{D}(\bar{\mathcal{N}})$ we have $(\varphi, 0) \in \mathbb{Y}_{\mathcal{N}}$ or $(0, \varphi) \in \mathbb{Y}_{\mathcal{N}}$. We recall that we assume $n \geq 4$ and we will treat the case $n = 3$ later on.

- Since $\sin^2 \theta \leq 1$ and φ is compactly supported in $\bar{\mathcal{N}}$, we rather straightforwardly have φ and $\partial_{\kappa_{\parallel}} \varphi \in \mathbb{L}$ and $\partial_{\kappa_{\perp}} \varphi \in L^2_{2(n-3), n-3}$, so that $\varphi \in \mathbb{H}$ for all values of $n \geq 4$. On the other hand, for $n = 4$, the function $\kappa_{\perp}^{n-5} = 1/\kappa_{\perp}$ is not integrable and $\varphi \notin L^2_{2(n-3), n-5}$. By contrast, if $n \geq 5$, $\varphi \in L^2_{2(n-3), n-5}$ and consequently, $\varphi \in \mathbb{K}$.
- Now, we clearly have $\mathcal{U}_1 \varphi = \tilde{\varphi} / \sin \theta$ with $\tilde{\varphi} \in \mathcal{D}(\bar{\mathcal{N}})$. So, $\|\mathcal{U}_1 \varphi\|_{\mathbb{L}} = \|\tilde{\varphi}\|_{L^2_{2(n-3), n-3}} < \infty$ if $n \geq 4$. It follows that $\mathcal{U}_1 \varphi$ belongs to \mathbb{L} and consequently, to \mathbb{H}' and \mathbb{K}' . Similarly, since φ is compactly supported, $\kappa_{\perp} \varphi$ belongs to \mathbb{L} and so, to \mathbb{H}' and \mathbb{K}' as well.

It follows that $(\varphi, 0) \in \mathbb{Y}_{\mathcal{N}}$ for $n \geq 4$ and that $(0, \varphi) \in \mathbb{Y}_{\mathcal{N}}$ for $n \geq 5$.

Now, we take $\varphi \in \mathcal{D}(\bar{\mathcal{N}})$ and assume that the solution (φ_w, φ_r) of the variational formulation (7.35) is smooth. We use $(\varphi, 0) \in \mathbb{Y}_{\mathcal{N}}$ as a test function in (7.35) for all $n \geq 4$. By using Green's formula, and point (i) of the present lemma, we end up with

$$\left(\int_{(0, \pi) \times \mathbb{R}} (\partial_{\kappa_{\perp}} \psi_w \varphi \mathbf{M}_{\mathbf{u}}^{\dagger} \sin^{2(n-3)} \theta \kappa_{\perp}^{n-3}) d\theta d\kappa_{\parallel} \right) \Big|_{\kappa_{\perp}=0} = 0,$$

which is a tautology when $n \geq 4$. Performing the same computation using $(0, \varphi) \in \mathbb{Y}_{\mathcal{N}}$ as a test function in (7.35) for all $n \geq 5$, we get a tautology again. Now, in the case $n = 4$, for $\varphi \in \mathcal{D}(\tilde{\mathcal{N}})$ to be in \mathbb{K} , we need $\varphi(\theta, \kappa_{\parallel}, 0) = 0$ to compensate for the weight $1/\kappa_{\perp}$, which leads to the Dirichlet condition (7.36). This ends the proof. \blacksquare

7.2.4 Case $n = 3$

In dimension $n = 3$, we use spherical coordinates. Let (e_1, e_2, \mathbf{u}) be a direct orthonormal basis and let (θ, φ) be the spherical coordinates in this basis. We let $(v, e_{\theta}, e_{\varphi})$ be the local spherical coordinates basis. Both θ and e_{θ} have been introduced in (4.2) and (4.3) respectively, and we have

$$w = \cos \varphi e_1 + \sin \varphi e_2, \quad e_{\varphi} = -\sin \varphi e_1 + \cos \varphi e_2.$$

Because $T_w \mathbb{S}^{n-2}$ is one-dimensional and spanned by e_{φ} , we can write

$$\kappa_T = \bar{\kappa}_T e_{\varphi}, \quad \text{so that} \quad \bar{\kappa}_T = \begin{cases} |\kappa_T| & \text{if } \bar{\kappa}_T \geq 0, \\ -|\kappa_T| & \text{if } \bar{\kappa}_T \leq 0, \end{cases}$$

and $\bar{\kappa}_T \in \mathbb{R}$.

For any function $\tilde{\psi} \in \mathcal{S}(M, \{\mathbf{u}\}^{\perp})$ satisfying the invariance relation (6.21), we define the functions $\tilde{\psi}_w, \tilde{\psi}_r: \tilde{\mathcal{N}} =: (0, \pi) \times \mathbb{R}^2 \rightarrow \mathbb{R}$, such that

$$\tilde{\psi}_w(\theta, \kappa_{\parallel}, \bar{\kappa}_T) = \psi_w(\theta, \kappa_{\parallel}, |\bar{\kappa}_T|), \quad \tilde{\psi}_r(\theta, \kappa_{\parallel}, \bar{\kappa}_T) = \text{Sign}(\bar{\kappa}_T) \psi_r(\theta, \kappa_{\parallel}, |\bar{\kappa}_T|), \quad (7.37)$$

for all $(\theta, \kappa_{\parallel}, \bar{\kappa}_T) \in \tilde{\mathcal{N}}$. In view of (7.12), any such function $\tilde{\psi}$ can be written in the form

$$\tilde{\psi}^{\dagger}(\theta, \kappa_{\parallel}, w, \kappa_T) = \tilde{\psi}_w(\theta, \kappa_{\parallel}, \bar{\kappa}_T) w + \tilde{\psi}_r(\theta, \kappa_{\parallel}, \bar{\kappa}_T) e_{\varphi}, \quad \forall ((\theta, \kappa_{\parallel}), (w, \kappa_T)) \in M_V \times M_T. \quad (7.38)$$

If $\tilde{\psi} = \tilde{\zeta}_{\mathbf{u}}$, the associated $(\tilde{\zeta}_w, \tilde{\zeta}_r)$ is the reduced GCI pair in dimension $n = 3$.

By analogy with the spaces defined in Subsections 7.2.1 and 7.2.2, we introduce the following spaces (notations that are a direct adaptation of those of the former sections are not repeated).

(i) For $p \in \mathbb{N}$, we define the Hilbert space:

$$\tilde{L}_p^2 = \left\{ \psi : \tilde{\mathcal{N}} \rightarrow \mathbb{R}, \mid \|\psi\|_{\tilde{L}_p^2}^2 =: \int_{\tilde{\mathcal{N}}} |\psi|^2 \mathbf{M}_{\mathbf{u}}^{\dagger} \sin^p \theta d\theta d\kappa_{\parallel} d\bar{\kappa}_T < \infty \right\}.$$

We denote by $\tilde{\mathbb{L}} = \tilde{L}_2^2$.

(ii) We define

$$\tilde{\mathbb{H}} = \left\{ \psi \in \tilde{\mathbb{L}} \mid \|\psi\|_{\tilde{\mathbb{H}}}^2 =: \|\psi\|_{\tilde{\mathbb{L}}}^2 + \|\partial_{\kappa_{\parallel}} \psi\|_{\tilde{\mathbb{L}}}^2 + \|\partial_{\kappa_{\perp}} \psi\|_{\tilde{L}_0^2}^2 < \infty \right\},$$

and recall that

$$\|\psi\|_{\tilde{\mathbb{H}}}^2 = \|\partial_{\kappa_{\parallel}} \psi\|_{\tilde{\mathbb{L}}}^2 + \|\partial_{\kappa_{\perp}} \psi\|_{\tilde{L}_0^2}^2.$$

Because $\tilde{\mathcal{N}}$ has no boundary in the $\bar{\kappa}_T$ direction, the following density theorem holds true (this would not be the case if we used κ_{\perp} instead of $\bar{\kappa}_T$, see Remark 7.2):

Lemma 7.10 (Density of $\mathcal{D}(\widetilde{\mathcal{N}})$ in $\widetilde{\mathbb{H}}$).
 $\mathcal{D}(\widetilde{\mathcal{N}})$ is dense in $\widetilde{\mathbb{H}}$.

(iii) We define $\widetilde{\mathcal{L}^2(\mathcal{L}^2)} = \widetilde{\mathbb{L}^2}$ and denote its elements by $(\bar{\psi}_w, \bar{\psi}_r)$.

(iv) We define $\widetilde{\mathcal{L}^2(\mathcal{H}^1)} = \widetilde{\mathbb{H}^2}$. Thanks to Lemma 7.10, $\mathcal{D}(\widetilde{\mathcal{N}})^2$ is dense in $\widetilde{\mathcal{L}^2(\mathcal{H}^1)}$.

Proposition 7.11 (Link between $L_v^2(L_\kappa^2)_{\text{inv}}$ (resp. $L_v^2(H_\kappa^1)_{\text{inv}}$) and $\widetilde{\mathcal{L}^2(\mathcal{L}^2)}$ (resp. $\widetilde{\mathcal{L}^2(\mathcal{H}^1)}$)).
We suppose $n = 3$. The function $\bar{\psi}$ belongs to $L_v^2(L_\kappa^2)_{\text{inv}}$ (resp. $L_v^2(H_\kappa^1)_{\text{inv}}$) if and only if $\bar{\psi}^\dagger$ is written according to (7.38) (in the a.e. sense) and $(\bar{\psi}_w, \bar{\psi}_r) \in \widetilde{\mathcal{L}^2(\mathcal{L}^2)}$ (resp. $\widetilde{\mathcal{L}^2(\mathcal{H}^1)}$). Furthermore,

$$\|\bar{\psi}\|_{L_v^2(L_\kappa^2)_{\text{u}}}^2 = 2\pi \|(\bar{\psi}_w, \bar{\psi}_r)\|_{\widetilde{\mathcal{L}^2(\mathcal{L}^2)}}^2, \quad (7.39)$$

$$\|\bar{\psi}\|_{L_v^2(H_\kappa^1)_{\text{u}}}^2 = 2\pi \|(\bar{\psi}_w, \bar{\psi}_r)\|_{\widetilde{\mathcal{L}^2(\mathcal{H}^1)}}^2, \quad (7.40)$$

and

$$\|\nabla_\kappa \bar{\psi}\|_{\widetilde{L}_v^2(L_\kappa^2)_{\text{u}}}^2 = 2\pi \left(\|\bar{\psi}_w\|_{\widetilde{\mathbb{H}}}^2 + \|\bar{\psi}_r\|_{\widetilde{\mathbb{H}}}^2 \right). \quad (7.41)$$

Remark 7.5 (About the factor 2π in (7.39) and (7.40)).

From (7.16), it seems at first glance that the factor appearing in (7.39) and (7.40) should be 4π . However, here, we integrate over $\bar{\kappa}_T \in \mathbb{R}$ instead of over $\kappa_\perp \in [0, \infty)$. Hence a factor 2 is absorbed in the duplication of the integration domain.

Proof. The proof of Prop. 7.11 is similar to that of Prop. 7.4 and is omitted. Let us just mention that the analog of (7.20) is

$$|(\nabla_\kappa \bar{\psi})^\dagger|^2 = |\partial_{\kappa_\parallel} \bar{\psi}_w|^2 + |\partial_{\kappa_\parallel} \bar{\psi}_r|^2 + \frac{1}{\sin^2 \theta} (|\partial_{\bar{\kappa}_T} \bar{\psi}_w|^2 + |\partial_{\bar{\kappa}_T} \bar{\psi}_r|^2). \quad (7.42)$$

■

(v) We define $\widetilde{\mathbb{H}}'$ as the dual of $\widetilde{\mathbb{H}}$. By Lemma 7.10, $\widetilde{\mathbb{H}}'$ is a distribution space.

(vi) We define

$$\widetilde{\mathcal{L}^2(\mathcal{H}^{-1})} = (\widetilde{\mathbb{H}}')^2.$$

Like in Subsection 7.2.2, we can identify this space with the space of pairs of distributions (T_w, T_r) satisfying an analog of identities (7.26) to (7.28) (with $c = 2\pi$), the precise writing of which are left to the reader.

(vii) We define the operator $\widetilde{\mathcal{U}}_1$ acting on $\bar{\psi} \in \widetilde{\mathbb{H}}$ as follows:

$$\widetilde{\mathcal{U}}_1 \bar{\psi} = \kappa_\parallel \partial_\theta \bar{\psi} + (\cos \theta \sin \theta \bar{\kappa}_T^2 - \nu \sin \theta) \partial_{\kappa_\parallel} \bar{\psi} - 2 \frac{\cos \theta}{\sin \theta} \kappa_\parallel \bar{\kappa}_T \partial_{\bar{\kappa}_T} \bar{\psi}. \quad (7.43)$$

The space $\widetilde{\mathbb{Y}}_{\mathcal{N}}$ is defined by

$$\widetilde{\mathbb{Y}}_{\mathcal{N}} = \{(\bar{\psi}_w, \bar{\psi}_r) \in \widetilde{\mathcal{L}^2(\mathcal{H}^1)} \text{ such that } (\widetilde{\mathcal{U}}_1 \bar{\psi}_w - \bar{\kappa}_T \bar{\psi}_r, \widetilde{\mathcal{U}}_1 \bar{\psi}_r + \bar{\kappa}_T \bar{\psi}_w) \in \widetilde{\mathcal{L}^2(\mathcal{H}^{-1})}\},$$

where $\widetilde{\mathcal{U}}_1\psi_w$ and $\widetilde{\mathcal{U}}_1\psi_r$ are defined in the distributional sense. Then, Lemma 7.5 is satisfied, “mutatis mutandis”. Let us just make the analog of (7.33) explicit:

$$\{\vec{\psi}, \mathbf{H}_u\}^\dagger = w \left((\widetilde{\mathcal{U}}_1\vec{\psi}_w - \bar{\kappa}_T\vec{\psi}_r) + e_\varphi (\widetilde{\mathcal{U}}_1\vec{\psi}_r + \bar{\kappa}_T\vec{\psi}_w) \right). \quad (7.44)$$

This proof of this identity is analog to that of (7.33) and is omitted. Then, we get the analog of Prop. 7.6:

Proposition 7.12 (Link between \mathbb{Y}_{inv} and $\widetilde{\mathbb{Y}}_{\mathcal{N}}$).

We assume $n = 3$. The function $\vec{\psi}$ belongs to \mathbb{Y}_{inv} if and only if $\vec{\psi}$ is written according to (7.38) and $(\vec{\psi}_w, \vec{\psi}_r)$ belongs to $\widetilde{\mathbb{Y}}_{\mathcal{N}}$. Furthermore,

$$\|\vec{\psi}\|_{\mathbb{Y}_{\text{inv}}}^2 = 2\pi \|(\psi_w, \psi_r)\|_{\widetilde{\mathbb{Y}}_{\mathcal{N}}}^2. \quad (7.45)$$

We can now derive the variational formulation satisfied by $(\vec{\psi}_w, \vec{\psi}_r)$, i.e. perform the analog of Subsection 7.2.3. We have the

Theorem 7.13 (Variational formulation; case $n = 3$).

We assume $n = 3$. Let $\vec{\chi} \in L_v^2(L_\kappa^2)_{\text{inv}}$ and define $(\vec{\chi}_w, \vec{\chi}_r) \in \widetilde{\mathcal{L}^2(\mathcal{L}^2)}$ as in (7.38). Then, $\vec{\psi}$ is the unique solution of the variational formulation (6.31) if and only if $(\vec{\psi}_w, \vec{\psi}_r)$ linked to $\vec{\psi}$ by (7.38) is the unique solution of the variational formulation

$$\left\{ \begin{array}{l} (\vec{\psi}_w, \vec{\psi}_r) \in \widetilde{\mathbb{Y}}_{\mathcal{N}}, \\ -\left(\left(\widetilde{\mathcal{U}}_1\vec{\psi}_w - \bar{\kappa}_T\vec{\psi}_r, \vec{\varphi}_w\right)_{\widetilde{\mathbb{H}'}, \widetilde{\mathbb{H}}} - \left(\widetilde{\mathcal{U}}_1\vec{\psi}_r + \bar{\kappa}_T\vec{\psi}_w, \vec{\varphi}_r\right)_{\widetilde{\mathbb{H}'}, \widetilde{\mathbb{H}}} + D\left(\vec{\psi}_w, \vec{\varphi}_w\right)_{\widetilde{\mathbb{H}}} \right. \\ \left. + D\left(\vec{\psi}_r, \vec{\varphi}_r\right)_{\widetilde{\mathbb{H}}} \right)_{\widetilde{\mathbb{H}}} = -\left(\vec{\chi}_w, \vec{\varphi}_w\right)_{\widetilde{\mathbb{L}}} - \left(\vec{\chi}_r, \vec{\varphi}_r\right)_{\widetilde{\mathbb{L}}}, \quad \forall (\vec{\varphi}_w, \vec{\varphi}_r) \in \widetilde{\mathbb{Y}}_{\mathcal{N}}. \end{array} \right. \quad (7.46)$$

followed by the

Corollary 7.14 (Variational formulation satisfied by the reduced GCI pair; case $n = 3$).

Let $(\vec{\xi}_w, \vec{\xi}_r)$ be the reduced GCI pair in dimension $n = 3$. Then, $(\vec{\xi}_w, \vec{\xi}_r)$ satisfies the variational formulation (7.46), with $(\vec{\psi}_w, \vec{\psi}_r)$ replaced by $(\vec{\xi}_w, \vec{\xi}_r)$ and $(\vec{\chi}_w, \vec{\chi}_r) = (\sin \theta, 0)$.

The strong form associated with the variational formulation (7.46) is given in

Theorem 7.15 (PDE system associated with the variational formulation (7.46) and boundary conditions, case $n = 3$).

(i) Let $(\vec{\xi}_w, \vec{\xi}_r) \in \widetilde{\mathbb{Y}}_{\mathcal{N}}$ be the unique solution to the variational formulation (7.46) (with $(\vec{\psi}_w, \vec{\psi}_r)$ replaced by $(\vec{\xi}_w, \vec{\xi}_r)$ and $(\vec{\chi}_w, \vec{\chi}_r) = (\sin \theta, 0)$). Then, (ξ_w, ξ_r) linked to $(\vec{\xi}_w, \vec{\xi}_r)$ through (7.37) satisfies System (4.19), (4.20) in the distributional sense.

(ii) (formal) Supposing that $(\vec{\xi}_w, \vec{\xi}_r)$ is smooth. Then the boundary conditions implied by (7.37) and the variational formulation (7.46) are

- no boundary condition at $\theta = 0$ nor $\theta = \pi$ on neither ζ_w nor ζ_r ;
- Neumann boundary conditions on ζ_w and homogeneous Dirichlet condition on ζ_r at $\kappa_\perp = 0$:

$$\partial_{\kappa_\perp} \zeta_w(\theta, \kappa_\parallel, 0) = 0, \quad \zeta_r(\theta, \kappa_\parallel, 0) = 0; \quad (7.47)$$

Proof. The same proof as for Theorem 7.9 leads to a system of equation for $(\bar{\zeta}_w, \bar{\zeta}_r)$ posed on $\widetilde{\mathcal{N}}$. By restricting it to \mathcal{N} , and using the fact that, on \mathcal{N} , $\bar{\kappa}_T = \kappa_\perp$, we recover System (4.19), (4.20). We get no boundary condition for the system posed on \mathcal{N} . However, when restricting it to \mathcal{N} , the mere smoothness of $(\bar{\zeta}_w, \bar{\zeta}_r)$ implies Neumann boundary conditions on ζ_w and Dirichlet boundary conditions on ζ_r at $\kappa_\perp = 0$. This ends the proof. ■

Remark 7.6 (About the variational formulation and the strong form).

Although the strong form can be restricted to a system of PDE posed on \mathcal{N} , this is not possible for the variational formulation. Indeed, the definition of the space $\widetilde{\mathcal{V}}_{\mathcal{N}}$ involves the use of the dual space $\widetilde{\mathcal{H}}'$ the definition of which requires a domain without boundaries, i.e. $\widetilde{\mathcal{N}}$. The variational formulation is the only formulation of the problem for which existence and uniqueness of solutions is proved. So, the reduction to a PDE posed on \mathcal{N} as formulated in Theorem 7.15 is not useful for a rigorous mathematical use. Its only merit is to highlight the analogies and differences between dimension 3 and higher dimensions.

8 Evolution equation for \mathbf{u}

We now use the reduced GCI pair to derive the equation for \mathbf{u} . This is the final step of the formal proof of Theorem 4.5.

Lemma 8.1 (Evolution equation for \mathbf{u} : first form).

The functions ρ and \mathbf{u} defined by (4.14) satisfy the following equation

$$\int_M (\partial_t + v \cdot \nabla_x) (\rho \mathbf{M}_{\mathbf{u}}) \vec{\zeta}_{\mathbf{u}} |\text{Vol}_M| = 0. \quad (8.1)$$

Proof. We multiply (3.52) by $\vec{\zeta}_{\mathbf{u}_{f^\varepsilon}}$ and integrate over M . Thanks to (5.8), we have

$$\int_M Q(f^\varepsilon) \vec{\zeta}_{\mathbf{u}_{f^\varepsilon}} |\text{Vol}_M| = 0.$$

Hence, we get

$$\int_M (\partial_t + v \cdot \nabla_x) f^\varepsilon \vec{\zeta}_{\mathbf{u}_{f^\varepsilon}} |\text{Vol}_M| = 0. \quad (8.2)$$

Now, when $\varepsilon \rightarrow 0$, we have formally $f^\varepsilon \rightarrow \rho \mathbf{M}_{\mathbf{u}}$ and $\mathbf{u}_{f^\varepsilon} \rightarrow \mathbf{u}_{\rho \mathbf{M}_{\mathbf{u}}} = \mathbf{u}$ by the proof of Prop. 4.4. Now, we show that, for any $\alpha \in M$, the map $\psi: \mathbf{u} \mapsto \zeta_{\mathbf{u}}(\alpha), \mathbb{S}^{n-1} \rightarrow \mathbb{R}$ is smooth. We fix $\mathbf{u}_0 \in \mathbb{S}^{n-1}$. We first recall that \mathbb{S}^{n-1} is diffeomorphic to the coset space $\text{SO}_n / \text{SO}_{n-1}$ where SO_{n-1}

denotes the isotropy subgroup of \mathbf{u}_0 . Hence, the projection $\pi: \text{SO}_n \rightarrow \mathbb{S}^{n-1}$ has the structure of a quotient manifold. Now, thanks to (6.20), the map $\varphi, R \mapsto \zeta_{R\mathbf{u}_0}(\alpha) = R\zeta_{\mathbf{u}_0}(R^T\alpha)$, $\text{SO}_n \rightarrow \mathbb{R}$ is smooth. Since $\varphi = \psi \circ \pi$, it follows from [46, Vol 1, Sect. 3.9, Prop. V] that ψ is smooth. Hence, we have a pointwise convergence $\vec{\zeta}_{\mathbf{u}_{f^\varepsilon}} \rightarrow \vec{\zeta}_{\mathbf{u}}$ as $\varepsilon \rightarrow 0$. Thus letting $\varepsilon \rightarrow 0$ in (8.2) and supposing that the convergence of f^ε to f is strong enough, we are led to (8.1), which ends the proof. \blacksquare

Now, we define the following tensors:

$$\mathbb{A} = \int_{M_V \times M_T} \sin^2 \theta \left((\mathbf{M}_{\mathbf{u}} \vec{\zeta}_{\mathbf{u}})^\dagger(\theta, \kappa_{\parallel}, w, \kappa_T) \otimes w \otimes w \right) |\mathbf{A}_{\mathbf{u}}^*(\text{Vol}_M)|, \quad (8.3)$$

$$\mathbb{B} = \int_{M_V \times M_T} \sin \theta \left((\mathbf{M}_{\mathbf{u}} \vec{\zeta}_{\mathbf{u}})^\dagger(\theta, \kappa_{\parallel}, w, \kappa_T) \otimes w \right) |\mathbf{A}_{\mathbf{u}}^*(\text{Vol}_M)|, \quad (8.4)$$

$$\mathbb{C} = \int_{M_V \times M_T} \sin \theta \cos \theta \left((\mathbf{M}_{\mathbf{u}} \vec{\zeta}_{\mathbf{u}})^\dagger(\theta, \kappa_{\parallel}, w, \kappa_T) \otimes w \right) |\mathbf{A}_{\mathbf{u}}^*(\text{Vol}_M)|, \quad (8.5)$$

$$\mathbb{D} = \int_{M_V \times M_T} (\mathbf{M}_{\mathbf{u}} \vec{\zeta}_{\mathbf{u}})^\dagger(\theta, \kappa_{\parallel}, w, \kappa_T) |\mathbf{A}_{\mathbf{u}}^*(\text{Vol}_M)|, \quad (8.6)$$

$$\mathbb{E} = \int_{M_V \times M_T} \cos \theta (\mathbf{M}_{\mathbf{u}} \vec{\zeta}_{\mathbf{u}})^\dagger(\theta, \kappa_{\parallel}, w, \kappa_T) |\mathbf{A}_{\mathbf{u}}^*(\text{Vol}_M)|, \quad (8.7)$$

where we recall that $|\mathbf{A}_{\mathbf{u}}^*(\text{Vol}_M)|$ is given by (7.10). Here, w and $\vec{\zeta}_{\mathbf{u}}^\dagger$ are considered as vectors of \mathbb{R}^n and \mathbb{A} is a rank-3 tensor, \mathbb{B} and \mathbb{C} are rank-2 tensors (i.e. matrices) and \mathbb{D} and \mathbb{E} are vectors. These tensors are useful to transform the left-hand side of (8.2) into explicit expressions, as shown next. We note that

$$\mathbf{M}_{\mathbf{u}}^\dagger(\theta, \kappa_{\parallel}, w, \kappa_T) = \frac{1}{Z} \exp \left(-\frac{\nu}{D} \left(-\nu \cos \theta + \frac{1}{2}(\kappa_{\parallel}^2 + \sin^2 \theta |\kappa_T|^2) \right) \right). \quad (8.8)$$

The right-hand side of (8.8) does not depend on w and defines the function $\mathbf{M}_{\mathbf{u}}^\natural = \mathbf{M}_{\mathbf{u}}^\natural(\theta, \kappa_{\parallel}, \kappa_T)$.

Lemma 8.2 (Evolution equation for \mathbf{u} : second form).

The functions ρ and \mathbf{u} defined by (4.14) satisfy the following equation

$$\frac{D^2}{\nu} (\mathbb{D} \partial_t \rho + \mathbb{E}(\mathbf{u} \cdot \nabla_x) \rho) + \mathbb{B} \left(\rho \partial_t \mathbf{u} + \frac{D}{\nu^2} \nabla_x \rho \right) + \rho \mathbb{C}(\mathbf{u} \cdot \nabla_x) \mathbf{u} + \rho \mathbb{A} : \nabla_x \mathbf{u} = 0, \quad (8.9)$$

where the $:$ in the last term indicates contraction of the last two indices of \mathbb{A} with the two indices of $\nabla_x \mathbf{u}$ while the second and third terms involve the matrix multiplication of \mathbb{B} and \mathbb{C} with the vector to their right.

Proof. We compute $(\partial_t + \nu \cdot \nabla_x)(\rho \mathbf{M}_{\mathbf{u}})$. This computation is straightforward but is reproduced here for the readers' convenience. We have

$$(\partial_t + \nu \cdot \nabla_x)(\rho \mathbf{M}_{\mathbf{u}}) = \mathbf{M}_{\mathbf{u}} \left((\partial_t + \nu \cdot \nabla_x) \rho + \rho (d \log \mathbf{M}_{\mathbf{u}})_{\mathbf{u}} ((\partial_t + \nu \cdot \nabla_x) \mathbf{u}) \right).$$

We have $\log \mathbf{M}_{\mathbf{u}} = -(\nu/D) \mathbf{H}_{\mathbf{u}} - \log Z$. Hence, for $\sigma \in T_{\mathbf{u}} \mathbb{S}^{n-1}$, we have

$$(d \log \mathbf{M}_{\mathbf{u}})_{\mathbf{u}}(\sigma) = -(\nu/D) (d \mathbf{H}_{\mathbf{u}})_{\mathbf{u}}(\sigma) = \frac{\nu^2}{D} P_{\mathbf{u}^\perp} \nu \cdot \sigma.$$

Thus, we have

$$(\partial_t + v \cdot \nabla_x)(\rho \mathbf{M}_u) = \mathbf{M}_u \left((\partial_t + v \cdot \nabla_x) \rho + \frac{v^2}{D} \rho P_{u^\perp} v \cdot ((\partial_t + v \cdot \nabla_x) \mathbf{u}) \right).$$

We use the change of variables (7.1) and get

$$\begin{aligned} \frac{D}{v^2} \left((\partial_t + v \cdot \nabla_x)(\rho \mathbf{M}_u) \right)^\dagger &= \mathbf{M}_u^\dagger \left\{ \frac{D}{v^2} (\partial_t \rho + \cos \theta \mathbf{u} \cdot \nabla_x \rho) + \sin \theta w \cdot \left(\rho \partial_t \mathbf{u} + \frac{D}{v^2} \nabla_x \rho \right) \right. \\ &\quad \left. + \rho \sin \theta \cos \theta w \cdot ((\mathbf{u} \cdot \nabla_x) \mathbf{u}) + \rho \sin^2 \theta (w \otimes w) : (\nabla_x \mathbf{u}) \right\}. \end{aligned} \quad (8.10)$$

Now, tensorizing on the left by $\vec{\zeta}_u^\dagger$ and integrating the result with respect to $|\mathbf{A}_u^*(\text{Vol}_M)|$, we get (8.9), which ends the proof. ■

Now, thanks to the explicit expression (7.12) of $\vec{\zeta}_u^\dagger$, we can compute the tensors \mathbb{A} to \mathbb{E} explicitly.

Lemma 8.3 (Expressions of the tensors \mathbb{A} to \mathbb{E}).

There exist two real numbers \mathbb{b} and \mathbb{c} such that

$$\mathbb{A} = 0, \quad (8.11)$$

$$\mathbb{B} = \mathbb{b} P_{u^\perp}, \quad \mathbb{C} = \mathbb{c} P_{u^\perp}, \quad (8.12)$$

$$\mathbb{D} = \mathbb{E} = 0. \quad (8.13)$$

Furthermore, we have

$$\mathbb{b} = (-1)^n \frac{c}{n-1} \int_{\mathcal{N}} (\mathbf{M}_u^\dagger \zeta_w)(\theta, \kappa_\parallel, \kappa_\perp) \sin^{2n-3} \theta \kappa_\perp^{n-3} d\theta d\kappa_\parallel d\kappa_\perp, \quad (8.14)$$

$$\mathbb{c} = (-1)^n \frac{c}{n-1} \int_{\mathcal{N}} (\mathbf{M}_u^\dagger \zeta_w)(\theta, \kappa_\parallel, \kappa_\perp) \cos \theta \sin^{2n-3} \theta \kappa_\perp^{n-3} d\theta d\kappa_\parallel d\kappa_\perp, \quad (8.15)$$

where c is given by (7.16).

Proof. We use the same methodology as in the proof of Proposition 7.4(i). We first consider \mathbb{B} given by (8.4). Thanks to (7.10), we have

$$\mathbb{B} = \int_{M_V \times M_T} ((\mathbf{M}_u \zeta_u)^\dagger(\theta, \kappa_\parallel, w, |\kappa_T|) \otimes w) \sin^{2n-3} \theta |\text{Vol}_{M_T}| d\theta d\kappa_\parallel.$$

Then, we apply the Fubini theorem and get

$$\begin{aligned} \mathbb{B} &= \int_{M_V} \left(\int_{M_T} ((\mathbf{M}_u \zeta_u)^\dagger(\theta, \kappa_\parallel, w, |\kappa_T|) \otimes w) |\text{Vol}_{M_T}| \right) \sin^{2n-3} \theta d\theta d\kappa_\parallel \\ &= \int_{M_V} B(\theta, \kappa_\parallel) \sin^{2n-3} \theta d\theta d\kappa_\parallel. \end{aligned} \quad (8.16)$$

With (7.12), we have

$$\begin{aligned} B &= B_1 + B_2, \\ B_1 &= \int_{M_T} (\mathbf{M}_{\mathbf{u}}^{\natural} \zeta_w)(\theta, \kappa_{\parallel}, |\kappa_T|) (w \otimes w) |\text{Vol}_{M_T}|, \\ B_2 &= \int_{M_T} (\mathbf{M}_{\mathbf{u}}^{\natural} \zeta_r)(\theta, \kappa_{\parallel}, |\kappa_T|) \left(\frac{\kappa_T}{|\kappa_T|} \otimes w \right) |\text{Vol}_{M_T}|. \end{aligned}$$

Thanks to (7.8), (3.14) and to the Fubini theorem for fiber bundles (3.15), we can write

$$B_1 = \int_{\mathbb{S}^{n-2}} (w \otimes w) \left(\int_{T_w \mathbb{S}^{n-2}} (\mathbf{M}_{\mathbf{u}}^{\natural} \zeta_w)(\theta, \kappa_{\parallel}, |\kappa_T|) |\omega_w| \right) |\text{Vol}_{\mathbb{S}^{n-2}}|,$$

where ω_w is the retrenchment of ω on $T_w \mathbb{S}^{n-2}$. Now, we can follow the same computation as in the proof of Proposition 7.4(i) and get

$$B_1 = |\mathbb{S}^{n-3}| \left(\int_0^\infty (\mathbf{M}_{\mathbf{u}}^{\natural} \zeta_w)(\theta, \kappa_{\parallel}, \kappa_{\perp}) \kappa_{\perp}^{n-3} d\kappa_{\perp} \right) \left(\int_{\mathbb{S}^{n-2}} (w \otimes w) |\text{Vol}_{\mathbb{S}^{n-2}}| \right).$$

It is readily seen that

$$\int_{\mathbb{S}^{n-2}} (w \otimes w) |\text{Vol}_{\mathbb{S}^{n-2}}| = \frac{|\mathbb{S}^{n-2}|}{n-1} P_{\mathbf{u}^{\perp}}. \quad (8.17)$$

Hence, we have

$$B_1 = \frac{|\mathbb{S}^{n-2}| |\mathbb{S}^{n-3}|}{n-1} \int_0^\infty (\mathbf{M}_{\mathbf{u}}^{\natural} \zeta_w)(\theta, \kappa_{\parallel}, \kappa_{\perp}) \kappa_{\perp}^{n-3} d\kappa_{\perp} P_{\mathbf{u}^{\perp}}.$$

To compute B_2 , we consider a direct orthonormal basis $(\mathbf{e}_1, \dots, \mathbf{e}_{n-1}, \mathbf{u})$ of \mathbb{R}^n and we compute the (i, j) -th entry of B_2 in this basis, i.e.

$$(B_2)_{ij} = \int_{M_T} (\mathbf{M}_{\mathbf{u}}^{\natural} \zeta_r)(\theta, \kappa_{\parallel}, |\kappa_T|) \frac{\kappa_T \cdot \mathbf{e}_i}{|\kappa_T|} w_j |\text{Vol}_{M_T}|.$$

By the same manipulations as for B_1 , we get

$$(B_2)_{ij} = \int_{\mathbb{S}^{n-2}} w_j \left(\int_{T_w \mathbb{S}^{n-2}} (\mathbf{M}_{\mathbf{u}}^{\natural} \zeta_r)(\theta, \kappa_{\parallel}, |\kappa_T|) \frac{\kappa_T \cdot \mathbf{e}_i}{|\kappa_T|} |\omega_w| \right) |\text{Vol}_{\mathbb{S}^{n-2}}|.$$

Let $(\varepsilon_1, \dots, \varepsilon_{n-2})$ be a local, positively oriented orthonormal frame of \mathbb{S}^{n-2} . Define a coordinate map $T_w \mathbb{B} \rightarrow \mathbb{R}^{n-1}$, $\kappa \mapsto (\kappa^1, \dots, \kappa^{n-2})$ with $\kappa^i = \langle \kappa, \varepsilon_i \rangle_v$. Then, $|\omega_w| = d\kappa^1 \dots d\kappa^{n-2}$ and we have

$$\begin{aligned} & \int_{T_w \mathbb{S}^{n-2}} (\mathbf{M}_{\mathbf{u}}^{\natural} \zeta_r)(\theta, \kappa_{\parallel}, |\kappa_T|) \frac{\kappa_T \cdot \mathbf{e}_i}{|\kappa_T|} |\omega_w| \\ &= \sum_{k=1}^{n-2} (\varepsilon_k \cdot \mathbf{e}_i) \int_{\mathbb{R}^{n-2}} (\mathbf{M}_{\mathbf{u}}^{\natural} \zeta_r) \left(\theta, \kappa_{\parallel}, \left(\sum_{\ell=1}^{n-2} |\kappa_T^{\ell}|^2 \right)^{1/2} \right) \frac{\kappa_T^k}{|\kappa_T|} d\kappa_T^1 \dots d\kappa_T^{n-2}. \end{aligned}$$

Now, by the change of variables $(\kappa_T^1, \dots, \kappa_T^k, \dots, \kappa_T^{n-2}) \rightarrow (\kappa_T^1, \dots, -\kappa_T^k, \dots, \kappa_T^{n-2})$, we realize that the integral in the k -th term of the sum at the right-hand side of the previous formula vanishes by antisymmetry. Hence, $B_2 = 0$. It follows that $\mathbb{B} = \mathbb{b}P_{\mathbf{u}}$ as it should, with \mathbb{b} given by (8.14). Going back to the formula (8.5), we realize that the same formula holds for \mathbb{C} , simply multiplying the integrated function by $\cos \theta$.

Now, we consider \mathbb{A} . Reproducing the same computation as for \mathbb{B} , we would get an analog to (8.16) with B replaced by A such that

$$\begin{aligned} A &= A_1 + A_2, \\ A_1 &= \int_{M_T} (\mathbf{M}_{\mathbf{u}}^{\natural} \zeta_w)(\theta, \kappa_{\parallel}, |\kappa_T|) (w \otimes w \otimes w) |\text{Vol}_{M_T}|, \\ A_2 &= \int_{M_T} (\mathbf{M}_{\mathbf{u}}^{\natural} \zeta_r)(\theta, \kappa_{\parallel}, |\kappa_T|) \left(\frac{\kappa_T}{|\kappa_T|} \otimes w \otimes w \right) |\text{Vol}_{M_T}|. \end{aligned}$$

Now, $A_2 = 0$ by the same argument as for B_2 , and $A_1 = 0$ because we have an even number of w -factors inside the integral corresponding to (8.17) and the integral vanishes by antisymmetry. Hence, we get (8.11). To get (8.13), the reasoning is the same, except that now there is only one w -factor inside the integral corresponding to (8.17) but the integral vanishes as well for the same reason. This ends the proof. ■

Remark 8.1 (About the expressions of the tensors \mathbb{A} to \mathbb{E}).

Expressions (8.11)-(8.13) for the tensors \mathbb{A} to \mathbb{E} could be obtained by simple symmetry considerations using (6.21). However, the expressions (8.14), (8.15) of \mathbb{b} and \mathbb{c} require Theorem 7.1 and the use of the reduced GCI pair.

We are now able to state the final form of the equation for \mathbf{u} . The following lemma completes the proof of Theorem 4.5.

Lemma 8.4 (Evolution equation for \mathbf{u} : final form).

The functions ρ and \mathbf{u} defined by (4.14) satisfy (4.16), where c_2 is given by (4.24) and c_3 by (4.17).

Proof. We insert the expressions of the tensors \mathbb{A} to \mathbb{E} found in Lemma 8.3 into (8.9) and get

$$\mathbb{b} P_{\mathbf{u}^{\perp}} \left(\rho \partial_t \mathbf{u} - \frac{D}{v^2} \nabla_x \rho \right) + \mathbb{c} \rho P_{\mathbf{u}^{\perp}} ((\mathbf{u} \cdot \nabla_x) \mathbf{u}) = 0.$$

Noting that both vectors $\partial_t \mathbf{u}$ and $(\mathbf{u} \cdot \nabla_x) \mathbf{u}$ belong to $\{\mathbf{u}\}^{\perp}$, and dividing by \mathbb{b} , we find (4.16) with $c_2 = \mathbb{c}/\mathbb{b}$ and c_3 given by (4.17). Now, thanks to (8.14), (8.15), we find that the expression of c_2 is precisely given by (4.24), which ends the proof. ■

9 Conclusion and perspectives

In this paper, we have provided a new formulation of the particle curvature-control model proposed in [22, 37] which enables us to extend it to arbitrary dimensions $n \geq 3$ (previously the

model was stated in dimensions 2 in [37] and 3 in [22]). We have also established the associated mean-field kinetic equations. Both the particle and kinetic models require a bundle geometric framework. Then, the main result of this paper is the derivation of the hydrodynamic limit of the kinetic model. We have shown that it coincides with the SOH model already found in the context of the Vicsek model. It requires the introduction of generalized collision invariants (GCI), which are solutions to a Fokker-Planck type equation. The action of the orthogonal group permits to express the GCI in terms of a pair of functions which satisfy a system of PDEs. Unique solvability of this system has been shown by means of a novel type of variational formulation.

This work opens many directions of research. First of all, the derivation of the SOH model is formal and lacks a mathematically rigorous proof. Numerical simulations should be developed to assess the accuracy and applicability range of the SOH model. We would also like to extend the present work to the non-normalized case. In this case, the model exhibits phase transitions in a similar way to the Vicsek model. For the latter, the multiplicity and stability of the spatially-homogeneous equilibria has been uncovered in [31, 32]. The corresponding results for the curvature-control model are still open. Some modifications of the model could be investigated, such as, for instance, restricting the curvature to a unit vector, to mimic the fact that animals have probably a range of preferences for their curvature. Another modification could be to replace the mean-field type of interaction by a Boltzmann like interaction, like in [43]. Finally one could also try to modify the model such that its hydrodynamic limit involves an additional equation for the locally-averaged curvature, like in [75]. This non-exhaustive list highlights the wealth of new research directions opened by the study of the curvature-control model of swarming behavior.

References

- [1] P. Aceves-Sánchez, M. Bostan, J.-A. Carrillo, and P. Degond. Hydrodynamic limits for kinetic flocking models of Cucker-Smale type. *Math. Biosci. Eng.*, 16:7883–7910, 2019.
- [2] R. A. Adams and J. J. Fournier. *Sobolev spaces*, volume 140. Elsevier, 2003.
- [3] D. Albritton, S. Armstrong, J.-C. Mourrat, and M. Novack. Variational methods for the kinetic Fokker-Planck equation. *Anal. PDE*, 17:1953–2010, 2024.
- [4] J. Angst, I. Bailleul, and C. Tardif. Kinetic brownian motion on riemannian manifolds. *Electron. J. Probab.*, 20(110):1–40, 2015.
- [5] I. Aoki. A simulation study on the schooling mechanism in fish. *Bull. Jpn. Soc. Sci. Fish*, 48(8):1081–1088, 1982.
- [6] V. I. Arnol’d. *Mathematical methods of classical mechanics*, volume 60. Springer Science & Business Media, 2013.
- [7] M. Baouendi and P. Grisvard. Sur une équation d’évolution changeant de type. *J. Funct. Anal.*, 2(3):352–367, 1968.
- [8] A. B. Barbaro, J. A. Canizo, J. A. Carrillo, and P. Degond. Phase transitions in a kinetic flocking model of Cucker–Smale type. *Multiscale Model. Simul.*, 14(3):1063–1088, 2016.
- [9] A. B. Barbaro and P. Degond. Phase transition and diffusion among socially interacting self-propelled agents. *Discrete Contin. Dyn. Syst. Ser. B*, 19:1249–1278, 2014.
- [10] J. Barré, P. Degond, D. Peurichard, and E. Zatorska. Modelling pattern formation through differential repulsion. *Netw. Heterog. Media*, 15:307–352, 2020.
- [11] F. Baudoin and C. Tardif. Hypocoercive estimates on foliations and velocity spherical Brownian motion. *Kinet. Relat. Models*, 11(2):1–23, 2018.
- [12] E. Bertin, M. Droz, and G. Grégoire. Boltzmann and hydrodynamic description for self-propelled particles. *Phys. Rev. E*, 74(2):022101, 2006.
- [13] E. Bertin, M. Droz, and G. Grégoire. Hydrodynamic equations for self-propelled particles: microscopic derivation and stability analysis. *Journal of Physics A: Mathematical and Theoretical*, 42(44):445001, 2009.
- [14] J.-M. Bismut. The hypoelliptic Laplacian on the cotangent bundle. *J. Amer. Math. Soc.*, 18(2):379–476, 2005.
- [15] M. Bostan and J. A. Carrillo. Asymptotic fixed-speed reduced dynamics for kinetic equations in swarming. *Math. Models Methods Appl. Sci.*, 23(13):2353–2393, 2013.
- [16] M. Bostan and J. A. Carrillo. Reduced fluid models for self-propelled particles interacting through alignment. *Math. Models Methods Appl. Sci.*, 27(07):1255–1299, 2017.

- [17] M. Briant, A. Diez, and S. Merino-Aceituno. Cauchy theory for general kinetic Vicsek models in collective dynamics and mean-field limit approximations. *SIAM J. Math. Anal.*, 54(1):1131–1168, 2022.
- [18] I. Bucataru, O. Constantinescu, and M. F. Dahl. A geometric setting for systems of ordinary differential equations. *Int. J. Geom. Methods Mod. Phys.*, 8(06):1291–1327, 2011.
- [19] S. Calogero. Exponential convergence to equilibrium for kinetic Fokker-Planck equations. *Comm. Partial Differential Equations*, 37(8):1357–1390, 2012.
- [20] F. Cao, S. Motsch, A. Reamy, and R. Theisen. Asymptotic flocking for the three-zone model. *Math. Biosci. Eng.*, 17(6):7692–7707, 2020.
- [21] J. A. Carrillo, M. Fornasier, J. Rosado, and G. Toscani. Asymptotic flocking dynamics for the kinetic Cucker–Smale model. *SIAM J. Math. Anal.*, 42(1):218–236, 2010.
- [22] A. Cavagna, L. Del Castello, I. Giardina, T. Grigera, A. Jelic, S. Melillo, T. Mora, L. Parisi, E. Silvestri, M. Viale, et al. Flocking and turning: a new model for self-organized collective motion. *J. Stat. Phys.*, 158:601–627, 2015.
- [23] C. Cercignani, R. Illner, and M. Pulvirenti. *The mathematical theory of dilute gases*, volume 106. Springer Science & Business Media, 2013.
- [24] L.-P. Chaintron and A. Diez. Propagation of chaos: a review of models, methods and applications. i. models and methods. *Kinet. Relat. Models*, 15:895–1015, 2022.
- [25] L.-P. Chaintron and A. Diez. Propagation of chaos: a review of models, methods and applications. ii. applications. *Kinet. Relat. Models*, 15:1017–1173, 2022.
- [26] H. Chaté, F. Ginelli, G. Grégoire, and F. Raynaud. Collective motion of self-propelled particles interacting without cohesion. *Phys. Rev. E*, 77(4):046113, 2008.
- [27] A. Costanzo and C. Hemelrijk. Spontaneous emergence of milling (vortex state) in a Vicsek-like model. *Journal of Physics D: Applied Physics*, 51(13):134004, 2018.
- [28] A. Creppy, F. Plouraboué, O. Praud, X. Druart, S. Cazin, H. Yu, and P. Degond. Symmetry-breaking phase transitions in highly concentrated semen. *Journal of The Royal Society Interface*, 13(123):20160575, 2016.
- [29] F. Cucker and S. Smale. Emergent behavior in flocks. *IEEE Trans. Automat. Control*, 52(5):852–862, 2007.
- [30] P. Degond. Global existence of smooth solutions for the Vlasov-Fokker-Planck equation in 1 and 2 space dimensions. In *Ann. Sci. Éc. Norm. Super. (4)*, volume 19, pages 519–542, 1986.
- [31] P. Degond, A. Frouvelle, and J.-G. Liu. Macroscopic limits and phase transition in a system of self-propelled particles. *J. Nonlinear Sci.*, 23:427–456, 2013.

- [32] P. Degond, A. Frouvelle, and J.-G. Liu. Phase transitions, hysteresis, and hyperbolicity for self-organized alignment dynamics. *Arch. Ration. Mech. Anal.*, 216:63–115, 2015.
- [33] P. Degond, J.-G. Liu, S. Motsch, and V. Panferov. Hydrodynamic models of self-organized dynamics: derivation and existence theory. *Methods Appl. Anal.*, 20:089–114, 2013.
- [34] P. Degond and S. Mas-Gallic. Existence of solutions and diffusion approximation for a model Fokker-Planck equation. *Transport Theory and Statistical Physics*, 16(4-6):589–636, 1987.
- [35] P. Degond and S. Motsch. Continuum limit of self-driven particles with orientation interaction. *Math. Models Methods Appl. Sci.*, 18(supp01):1193–1215, 2008.
- [36] P. Degond and S. Motsch. Large scale dynamics of the persistent turning walker model of fish behavior. *J. Stat. Phys.*, 131:989–1021, 2008.
- [37] P. Degond and S. Motsch. A macroscopic model for a system of swarming agents using curvature control. *J. Stat. Phys.*, 143:685–714, 2011.
- [38] M. P. do Carmo. *Riemannian Geometry*. Birkhäuser, 1992.
- [39] M. R. D’Orsogna, Y.-L. Chuang, A. L. Bertozzi, and L. S. Chayes. Self-propelled particles with soft-core interactions: patterns, stability, and collapse. *Phys. Rev. Lett.*, 96(10):104302, 2006.
- [40] A. Figalli, M.-J. Kang, and J. Morales. Global well-posedness of the spatially homogeneous Kolmogorov–Vicsek model as a gradient flow. *Arch. Ration. Mech. Anal.*, 227:869–896, 2018.
- [41] S. Gallot, D. Hulin, and J. Lafontaine. *Riemannian geometry*, volume 2. Springer, 1990.
- [42] I. M. Gamba and M.-J. Kang. Global weak solutions for Kolmogorov–Vicsek type equations with orientational interactions. *Arch. Ration. Mech. Anal.*, 222:317–342, 2016.
- [43] J. Gautrais, F. Ginelli, R. Fournier, S. Blanco, M. Soria, H. Chaté, and G. Theraulaz. Deciphering interactions in moving animal groups. *PLoS Comput. Biol.*, 8:e1002678, 2012.
- [44] J. Gautrais, C. Jost, M. Soria, A. Campo, S. Motsch, R. Fournier, S. Blanco, and G. Theraulaz. Analyzing fish movement as a persistent turning walker. *J. Math. Biol.*, 58:429–445, 2009.
- [45] M. Girolami and B. Calderhead. Riemann manifold Langevin and Hamiltonian Monte-Carlo methods. *J. R. Stat. Soc. Ser. B. Stat. Methodol.*, 73(2):123–214, 2011.
- [46] W. Greub, S. Halperin, and R. Vanstone. *Connections, Curvature, and Cohomology*. Academic press, 1972.
- [47] Q. Griette and S. Motsch. Kinetic equations and self-organized band formations. *Active Particles, Volume 2: Advances in Theory, Models, and Applications*, pages 173–199, 2019.

- [48] M. Grothaus and M. C. Mertin. Hypocoercivity of Langevin-type dynamics on abstract smooth manifolds. *Stochastic Process. Appl.*, 146:22–59, 2022.
- [49] S. Gudmundsson and E. Kappos. On the geometry of tangent bundles. *Expo. Math.*, 20(1):1–41, 2002.
- [50] S.-Y. Ha and J.-G. Liu. A simple proof of the Cucker-Smale flocking dynamics and mean-field limit. *Commun. Math. Sci.*, 7:297–325, 2009.
- [51] S.-Y. Ha and E. Tadmor. From particle to kinetic and hydrodynamic descriptions of flocking. *Kinet. Relat. Mod.*, 1:415–435, 2008.
- [52] J. Haskovec. A simple proof of asymptotic consensus in the Hegselmann–Krause and Cucker–Smale models with normalization and delay. *SIAM J. Appl. Dyn. Syst.*, 20(1):130–148, 2021.
- [53] E. P. Hsu. *Stochastic analysis on manifolds*. American Mathematical Soc., 2002.
- [54] N. Ikeda and S. Watanabe. *Stochastic differential equations and diffusion processes*. Elsevier, 2014.
- [55] N. Jiang, Y.-L. Luo, and T.-F. Zhang. From kinetic flocking model of Cucker–Smale type to self-organized hydrodynamic model. *Math. Models Methods Appl. Sci.*, 34(13):2395–2467, 2024.
- [56] N. Jiang, L. Xiong, and T.-F. Zhang. Hydrodynamic limits of the kinetic self-organized models. *SIAM J. Math. Anal.*, 48(5):3383–3411, 2016.
- [57] E. Joergensen. Construction of the Brownian motion and the Ornstein-Uhlenbeck process in a Riemannian manifold on basis of the Gangolli-Mc. Kean injection scheme. *Probab. Theory Related Fields*, 44(1):71–87, 1978.
- [58] G. Lebeau. Geometric Fokker-Planck equations. *Port. Math.*, 62(4):469–530, 2005.
- [59] T. Lelièvre, M. Rousset, and G. Stoltz. Langevin dynamics with constraints and computation of free energy differences. *Math. Comp.*, 81(280):2071–2125, 2012.
- [60] C. D. Levermore. Moment closure hierarchies for kinetic theories. *J. Stat. Phys.*, 83:1021–1065, 1996.
- [61] G. Lin, R. Escobedo, X. Li, T. Xue, Z. Han, C. Sire, V. Guttal, and G. Theraulaz. Experimental evidence of stress-induced critical state in schooling fish. *bioRxiv*, pages 2025–02, 2025.
- [62] J. L. Lions. *Equations différentielles opérationnelles et problèmes aux limites*, volume 111. Springer-Verlag, 2013.
- [63] S. Motsch and L. Navoret. Numerical simulations of a nonconservative hyperbolic system with geometric constraints describing swarming behavior. *Multiscale Model. Simul.*, 9(3):1253–1275, 2011.

- [64] S. Motsch and E. Tadmor. A new model for self-organized dynamics and its flocking behavior. *J. Stat. Phys.*, 144:923–947, 2011.
- [65] F. Nier, X. Sang, and F. White. Global subelliptic estimates for geometric Kramers-Fokker-Planck operators on closed manifolds. *arXiv preprint arXiv:2402.07511*, 2024.
- [66] S. Sasaki. On the differential geometry of tangent bundles of riemannian manifolds. *Tohoku Math. J. (2)*, 10(3):338–354, 1958.
- [67] S. Sayin, E. Couzin-Fuchs, I. Petelski, Y. Günzel, M. Salahshour, C.-Y. Lee, J. M. Graving, L. Li, O. Deussen, G. A. Sword, et al. The behavioral mechanisms governing collective motion in swarming locusts. *Science*, 387(6737):995–1000, 2025.
- [68] M. Soloveitchik. Fokker-Planck equation on a manifold. effective diffusion and spectrum. *Potential Anal.*, 4:571–593, 1995.
- [69] J. Toner and Y. Tu. Long-range order in a two-dimensional dynamical XY model: how birds fly together. *Phys. Rev. Lett.*, 75(23):4326, 1995.
- [70] J. Toner and Y. Tu. Flocks, herds, and schools: A quantitative theory of flocking. *Phys. Rev. E*, 58(4):4828, 1998.
- [71] J. Toner, Y. Tu, and S. Ramaswamy. Hydrodynamics and phases of flocks. *Annals of Physics*, 318(1):170–244, 2005.
- [72] T. Vicsek, A. Czirók, E. Ben-Jacob, I. Cohen, and O. Shochet. Novel type of phase transition in a system of self-driven particles. *Phys. Rev. Lett.*, 75(6):1226, 1995.
- [73] T. Vicsek and A. Zafeiris. Collective motion. *Physics reports*, 517(3-4):71–140, 2012.
- [74] F. W. Warner. *Foundations of Differentiable Manifolds and Lie Groups*. Springer, 1983.
- [75] X. Yang and M. C. Marchetti. Hydrodynamics of turning flocks. *Phys. Rev. Lett.*, 115(25):258101, 2015.
- [76] T.-F. Zhang and N. Jiang. A local existence of viscous self-organized hydrodynamic model. *Non linear Anal. Real World Appl.*, 34:495–506, 2017.

Appendix

A Proofs of remarkable formulas (Section 3.1.5)

A.1 Proof of Lemma 3.1

Read from here

For $k \in \mathcal{X}(B)$, we will denote $\mathcal{L}^V \tilde{k}$ by $\mathcal{L}^V k$ for simplicity, and similarly with V replaced by H . Let $(\varepsilon_1, \dots, \varepsilon_{n-1})$ be a local orthonormal frame on B . Then, it is readily checked that $(\mathcal{L}^H \varepsilon_1, \dots, \mathcal{L}^H \varepsilon_{n-1}, \mathcal{L}^V \varepsilon_1, \dots, \mathcal{L}^V \varepsilon_{n-1})$ is a local orthonormal frame on M . Now, using (3.19), we can write

$$\nabla_v \cdot \tilde{k} = \nabla_a \cdot (\mathcal{L}^H k) = \sum_{i=1}^{n-1} \left(\langle \nabla_{\mathcal{L}^H \varepsilon_i}^M \mathcal{L}^H k, \mathcal{L}^H \varepsilon_i \rangle + \langle \nabla_{\mathcal{L}^V \varepsilon_i}^M \mathcal{L}^H k, \mathcal{L}^V \varepsilon_i \rangle \right). \quad (\text{A.1})$$

By the torsionfreeness of the connection, we have

$$\nabla_{\mathcal{L}^H \varepsilon_i}^M \mathcal{L}^H k = \nabla_{\mathcal{L}^H k}^M \mathcal{L}^H \varepsilon_i + [\mathcal{L}^H \varepsilon_i, \mathcal{L}^H k],$$

and since $\langle \mathcal{L}^H \varepsilon_i, \mathcal{L}^H \varepsilon_i \rangle = 1$ and the connection is compatible with the metric, we have

$$\langle \nabla_{\mathcal{L}^H k}^M \mathcal{L}^H \varepsilon_i, \mathcal{L}^H \varepsilon_i \rangle = 0,$$

and similarly with $\mathcal{L}^H \varepsilon_i$ replaced by $\mathcal{L}^V \varepsilon_i$. Thus, we get

$$\nabla_v \cdot \tilde{k} = \sum_{i=1}^{n-1} \left(\langle [\mathcal{L}^H \varepsilon_i, \mathcal{L}^H k], \mathcal{L}^H \varepsilon_i \rangle + \langle [\mathcal{L}^V \varepsilon_i, \mathcal{L}^H k], \mathcal{L}^V \varepsilon_i \rangle \right).$$

Now, for any two vector fields $k, h \in \mathcal{X}(B)$, we have [49, Prop. 5.1]:

$$[\mathcal{L}^H h, \mathcal{L}^H k] = \mathcal{L}^H ([h, k]) - \mathcal{L}^V (\mathcal{R}(h, k)\kappa), \quad (\text{A.2})$$

$$[\mathcal{L}^H h, \mathcal{L}^V k] = \mathcal{L}^V (\nabla_h^B k), \quad (\text{A.3})$$

where the Levi-Civita connection ∇^B on B was used and \mathcal{R} is its Riemann curvature tensor. Thus, we have

$$\begin{aligned} \nabla_v \cdot \tilde{k} &= \sum_{i=1}^{n-1} \left(\langle \mathcal{L}^H ([\varepsilon_i, k]) - \mathcal{L}^V (\mathcal{R}(\varepsilon_i, k)\kappa), \mathcal{L}^H \varepsilon_i \rangle - \langle \mathcal{L}^V (\nabla_k^B \varepsilon_i), \mathcal{L}^V \varepsilon_i \rangle \right) \\ &= \sum_{i=1}^{n-1} \left(\langle \mathcal{L}^H ([\varepsilon_i, k]), \mathcal{L}^H \varepsilon_i \rangle - \langle \mathcal{L}^V (\nabla_k^B \varepsilon_i), \mathcal{L}^V \varepsilon_i \rangle \right), \end{aligned}$$

where the term involving the Riemann curvature tensor has been cancelled due to orthogonality of the horizontal and vertical subspaces. Now, by the definition of the Sasaki metric, we can remove the operators \mathcal{L}^H in the first term and \mathcal{L}^V in the second term provided we replace the metric on M by the metric on B . This leads to

$$\nabla_v \cdot \tilde{k} = \sum_{i=1}^{n-1} \left(\langle [\varepsilon_i, k], \varepsilon_i \rangle - \langle \nabla_k^B \varepsilon_i, \varepsilon_i \rangle \right).$$

We have $\langle \nabla_k^B \varepsilon_i, \varepsilon_i \rangle = 0$ again because $\langle \varepsilon_i, \varepsilon_i \rangle = 1$. Using the same manipulations as previously, we also notice that

$$\sum_{i=1}^{n-1} \langle [\varepsilon_i, k], \varepsilon_i \rangle = \sum_{i=1}^{n-1} \langle \nabla_{\varepsilon_i}^B k, \varepsilon_i \rangle = \nabla \cdot k.$$

Hence, we have just proved the first formula (3.24).

For the second formula (3.24), we proceed analogously. We have

$$\begin{aligned} \nabla_\kappa \cdot \tilde{k} &= \nabla_\alpha \cdot (\mathcal{L}^V k) = \sum_{i=1}^{n-1} \left(\langle \nabla_{\mathcal{L}^H \varepsilon_i}^M \mathcal{L}^V k, \mathcal{L}^H \varepsilon_i \rangle + \langle \nabla_{\mathcal{L}^V \varepsilon_i}^M \mathcal{L}^V k, \mathcal{L}^V \varepsilon_i \rangle \right). \\ &= \sum_{i=1}^{n-1} \left(\langle [\mathcal{L}^H \varepsilon_i, \mathcal{L}^V k], \mathcal{L}^H \varepsilon_i \rangle + \langle [\mathcal{L}^V \varepsilon_i, \mathcal{L}^V k], \mathcal{L}^V \varepsilon_i \rangle \right). \end{aligned} \quad (\text{A.4})$$

Now, for any two vector fields $k, h \in \mathcal{X}(B)$, we have [49, Prop. 5.1]:

$$[\mathcal{L}^V h, \mathcal{L}^V k] = 0. \quad (\text{A.5})$$

We also recall (A.3) as well as the orthogonality of the horizontal and vertical subspaces. This leads to

$$\nabla_\kappa \cdot \tilde{k} = \sum_{i=1}^{n-1} \langle \mathcal{L}^V (\nabla_{\varepsilon_i}^B k), \mathcal{L}^H \varepsilon_i \rangle = 0.$$

Thus proving the second formula (3.24) and ending the proof. \blacksquare

A.2 Proof of Lemma 3.2

(i) Let $\alpha \in M$ and $v = \pi\alpha$. Let $a \in T_v B$. We have

$$\begin{aligned} \langle \nabla_v \tilde{\varphi}(\alpha), a \rangle_v &= \langle B_\alpha^H \nabla_\alpha \tilde{\varphi}(\alpha), a \rangle_v = \langle \nabla_\alpha \tilde{\varphi}(\alpha), \mathcal{L}_\alpha^H a \rangle_\alpha = (d\tilde{\varphi})_\alpha (\mathcal{L}_\alpha^H a) \\ &= ((d\varphi)_{\pi\alpha} \circ (d\pi)_\alpha) (\mathcal{L}_\alpha^H a) = (d\varphi)_v(a) = \langle \nabla_v \varphi(v), a \rangle_v, \end{aligned}$$

where we have used (2.40) to say that $(d\pi)_\alpha (\mathcal{L}_\alpha^H a) = a$.

Similarly we get, for any $\tau \in T_v B$:

$$\begin{aligned} \langle \nabla_\kappa \tilde{\varphi}(\alpha), \tau \rangle_v &= \langle B_\alpha^V \nabla_\alpha \tilde{\varphi}(\alpha), \tau \rangle_v = \langle \nabla_\alpha \tilde{\varphi}(\alpha), \mathcal{L}_\alpha^V \tau \rangle_\alpha = (d\tilde{\varphi})_\alpha (\mathcal{L}_\alpha^V \tau) \\ &= ((d\varphi)_{\pi\alpha} \circ (d\pi)_\alpha) (\mathcal{L}_\alpha^V \tau) = 0, \end{aligned}$$

thanks to (2.39).

(ii) Let $\alpha = (v, \kappa)$ be given. We define $\theta: \mathbb{R} \rightarrow \mathbb{R}$ such that $\theta(t) = \psi(v, \kappa t)$. Then,

$$\begin{aligned} \frac{d\theta}{dt}(t) &= (d\psi)_{(v, \kappa t)}(0, \kappa) = (d\psi)_{(v, \kappa t)}(\mathcal{L}_{(v, \kappa t)}^V \kappa) \\ &= \langle \nabla_\alpha \psi(v, \kappa t), \mathcal{L}_{(v, \kappa t)}^V \kappa \rangle_{(v, \kappa t)} = \langle B_{(v, \kappa t)}^V \nabla_\alpha \psi(v, \kappa t), \kappa \rangle_v \\ &= \langle \nabla_\kappa \psi(v, \kappa t), \kappa \rangle_v = 0. \end{aligned}$$

Hence, $\theta(0) = \theta(1)$ i.e. $\psi(v, 0) = \psi(v, \kappa)$. Defining $\varphi: B \rightarrow \mathbb{R}$ by $\varphi(v) = \psi(v, 0)$, we deduce that $\psi = \varphi \circ \pi$, which ends the proof. \blacksquare

A.3 Proof of Lemma 3.3

Let $\tau \in T_v B$. Then, we have

$$\langle \nabla_\kappa \langle \kappa, \kappa \rangle_v, \tau \rangle = \langle B_\alpha^V \nabla_\alpha \langle \kappa, \kappa \rangle_v, \tau \rangle = \langle \nabla_\alpha \langle \kappa, \kappa \rangle_v, \mathcal{L}_\alpha^V \tau \rangle_\alpha \quad (\text{A.6})$$

$$= (d\langle \kappa, \kappa \rangle_v)_\alpha (\mathcal{L}_\alpha^V \tau) = \frac{d}{dt} \langle \tilde{\kappa}(t), \tilde{\kappa}(t) \rangle_{\tilde{v}(t)} \Big|_{t=0}, \quad (\text{A.7})$$

where $\tilde{\alpha}(t) = (\tilde{v}(t), \tilde{\kappa}(t))$ is the following path in M : $\tilde{v}(t) = v$, $\tilde{\kappa}(t) = \kappa + t\tau$. Indeed, we notice that $\tilde{\alpha}(0) = (v, \kappa) = \alpha$ and

$$\frac{d\tilde{\alpha}}{dt} \Big|_{t=0} = (0, \tau) = \mathcal{L}_\alpha^V \tau.$$

Then, we straightforwardly get that

$$\frac{d}{dt} \langle \tilde{\kappa}(t), \tilde{\kappa}(t) \rangle_{\tilde{v}(t)} \Big|_{t=0} = \langle 2\kappa, \tau \rangle_v,$$

which, inserted into (A.7), leads to the first equation (3.26).

Let now $a \in T_v B$. Similarly to the previous computation, we have

$$\langle \nabla_v \langle \kappa, \kappa \rangle_v, a \rangle = \langle B_\alpha^H \nabla_\alpha \langle \kappa, \kappa \rangle_v, a \rangle = \langle \nabla_\alpha \langle \kappa, \kappa \rangle_v, \mathcal{L}_\alpha^H a \rangle_\alpha \quad (\text{A.8})$$

$$= (d\langle \kappa, \kappa \rangle_v)_\alpha (\mathcal{L}_\alpha^H a) = \frac{d}{dt} \langle \tilde{\kappa}(t), \tilde{\kappa}(t) \rangle_{\tilde{v}(t)} \Big|_{t=0}, \quad (\text{A.9})$$

where $\tilde{\alpha}(t) = (\tilde{v}(t), \tilde{\kappa}(t))$ is a path such that

$$(\tilde{v}(0), \tilde{\kappa}(0)) = (v, \kappa), \quad \frac{d}{dt} (\tilde{v}, \tilde{\kappa}) \Big|_{t=0} = \mathcal{L}_\alpha^H a = (a, -(a \cdot \kappa)v). \quad (\text{A.10})$$

Let (e_1, \dots, e_n) be an orthonormal frame of \mathbb{R}^n such that $v = e_n$, $a = |a|e_{n-1}$ and $\kappa = \kappa_{n-2}e_{n-2} + \kappa_{n-1}e_{n-1}$. For given α and $a \in T_v B$, such a frame always exists. Now, define the path

$$\tilde{v}(t) = \sin(|a|t)e_{n-1} + \cos(|a|t)e_n, \quad \tilde{\kappa}(t) = \kappa_{n-2}e_{n-2} + \kappa_{n-1}e_{n-1} - \kappa_{n-1} \tan(|a|t)e_n.$$

We easily see that $\tilde{\alpha} = (\tilde{v}, \tilde{\kappa})$ is a path in M (i.e. $|\tilde{v}| = 1$, $\tilde{\kappa} \cdot \tilde{v} = 0$) defined in a neighborhood of $t = 0$ and satisfying (A.10). Now, we have

$$\langle \tilde{\kappa}(t), \tilde{\kappa}(t) \rangle_{\tilde{v}(t)} = \kappa_{n-2}^2 + \kappa_{n-1}^2 (1 + \tan^2(|a|t)).$$

Thus,

$$\frac{d}{dt} \langle \tilde{\kappa}(t), \tilde{\kappa}(t) \rangle_{\tilde{v}(t)} = 2|a|\kappa_{n-1}^2 \tan(|a|t)(1 + \tan^2(|a|t)),$$

which leads to

$$\frac{d}{dt} \langle \tilde{\kappa}(t), \tilde{\kappa}(t) \rangle_{\tilde{v}(t)} \Big|_{t=0} = 0.$$

Inserting it in (A.9), we get the second equation of (3.26), which ends the proof. \blacksquare

B Reduced GCI pair: proof of Theorem 7.1

B.1 Geometric preliminaries

We first remark that $\vec{\psi}^\dagger$ satisfies the following invariance under action of O_{n-1} :

$$\vec{\psi}^\dagger(\theta, \kappa_\parallel, R\omega, R\kappa_T) = R\vec{\psi}^\dagger(\theta, \kappa_\parallel, \omega, \kappa_T), \quad \forall R \in O_{n-1}. \quad (\text{B.1})$$

We remark that $(R\omega, R\kappa_T)$ is the image of (ω, κ_T) by the left group action of O_{n-1} on M_T . We decompose $\vec{\psi}^\dagger$ according to

$$\vec{\psi}^\dagger(\theta, \kappa_\parallel, \omega, \kappa_T) = \psi_w^\dagger(\theta, \kappa_\parallel, \omega, \kappa_T) \omega + \vec{\psi}_T^\dagger(\theta, \kappa_\parallel, \omega, \kappa_T), \quad (\text{B.2})$$

with

$$\psi_w^\dagger(\theta, \kappa_\parallel, \omega, \kappa_T) \in \mathbb{R} \quad \text{and} \quad \vec{\psi}_T^\dagger(\theta, \kappa_\parallel, \omega, \kappa_T) \in T_\omega \mathbb{S}^{n-2}. \quad (\text{B.3})$$

Straightforwardly, from (B.1), we get

$$\psi_w^\dagger(\theta, \kappa_\parallel, R\omega, R\kappa_T) = \psi_w^\dagger(\theta, \kappa_\parallel, \omega, \kappa_T), \quad \forall R \in O_{n-1}, \quad (\text{B.4})$$

$$\vec{\psi}_T^\dagger(\theta, \kappa_\parallel, R\omega, R\kappa_T) = R\vec{\psi}_T^\dagger(\theta, \kappa_\parallel, \omega, \kappa_T), \quad \forall R \in O_{n-1}, \quad (\text{B.5})$$

Thus, we need to prove that

$$\psi_w^\dagger(\theta, \kappa_\parallel, \omega, \kappa_T) = \psi_w(\theta, \kappa_\parallel, |\kappa_T|), \quad \vec{\psi}_T^\dagger(\theta, \kappa_\parallel, \omega, \kappa_T) = \psi_r(\theta, \kappa_\parallel, |\kappa_T|) \frac{\kappa_T}{|\kappa_T|}. \quad (\text{B.6})$$

We use the spherical coordinates on B with pole \mathbf{u} . In this section (only) and for the sake of simplicity, we list the canonical basis of \mathbb{R}^n in a different order, namely $(\mathbf{u}, \mathbf{e}_1, \dots, \mathbf{e}_{n-1})$ with associated coordinates written $(x_u, x_1, \dots, x_{n-1})$. Instead of the diffeomorphism (4.1), we consider $\tilde{\mathbf{V}}_{\mathbf{u}}: (0, \pi)^{n-2} \times \mathbb{R}(\text{mod. } 2\pi) \rightarrow \tilde{B}_{\mathbf{u}}, \quad (\theta, \varphi_1, \varphi_2, \dots, \varphi_{n-2}) \mapsto v = \tilde{\mathbf{V}}_{\mathbf{u}}(\theta, \varphi_1, \varphi_2, \dots, \varphi_{n-2}) \in \mathbb{S}^{n-1}$ such that

$$\tilde{\mathbf{V}}_{\mathbf{u}} = \cos \theta \mathbf{u} + \sin \theta w(\varphi_1, \varphi_2, \dots, \varphi_{n-2}),$$

with

$$w(\varphi_1, \varphi_2, \dots, \varphi_{n-2}) = \begin{pmatrix} 0 \\ \cos \varphi_1 \\ \sin \varphi_1 \cos \varphi_2 \\ \sin \varphi_1 \sin \varphi_2 \cos \varphi_3 \\ \vdots \\ \sin \varphi_1 \sin \varphi_2 \dots \sin \varphi_{n-3} \cos \varphi_{n-2} \\ \sin \varphi_1 \sin \varphi_2 \dots \sin \varphi_{n-3} \sin \varphi_{n-2} \end{pmatrix},$$

and

$$\tilde{B}_{\mathbf{u}} = B \setminus \{(x_u, x_1, x_2, \dots, x_{n-3}, 0, 0), \mid x_u^2 + x_1^2 + \dots + x_{n-3}^2 = 1\},$$

being the excluded set corresponding to the points where $\tilde{\mathbf{V}}_{\mathbf{u}}$ is not uniquely invertible and thus, is not a diffeomorphism. We deduce that

$$\frac{\partial \tilde{\mathbf{V}}_{\mathbf{u}}}{\partial \theta} = -\sin \theta \mathbf{u} + \cos \theta w =: e_\theta(\theta, \varphi_1, \dots, \varphi_{n-2}),$$

and for $i \in \{1, \dots, n-3\}$,

$$\begin{aligned} \frac{\partial \tilde{\mathbf{V}}_{\mathbf{u}}}{\partial \varphi_i} &= \sin \theta \sin \varphi_1 \dots \sin \varphi_{i-1} \begin{pmatrix} 0 \\ \vdots \\ 0 \\ -\sin \varphi_i \\ \cos \varphi_i \cos \varphi_{i+1} \\ \cos \varphi_i \sin \varphi_{i+1} \cos \varphi_{i+2} \\ \vdots \\ \cos \varphi_i \sin \varphi_{i+1} \dots \sin \varphi_{n-3} \cos \varphi_{n-2} \\ \cos \varphi_i \sin \varphi_{i+1} \dots \sin \varphi_{n-3} \sin \varphi_{n-2} \end{pmatrix} \\ &:= \sin \theta \sin \varphi_1 \dots \sin \varphi_{i-1} e_{\varphi_i}, \end{aligned}$$

where the last 0 is on the i -th line, while, for $i = n-2$,

$$\frac{\partial \tilde{\mathbf{V}}_{\mathbf{u}}}{\partial \varphi_{n-2}} = \sin \theta \sin \varphi_1 \dots \sin \varphi_{n-3} \begin{pmatrix} 0 \\ \vdots \\ 0 \\ -\sin \varphi_{n-2} \\ \cos \varphi_{n-2} \end{pmatrix} := \sin \theta \sin \varphi_1 \dots \sin \varphi_{n-3} e_{\varphi_{n-2}},$$

The tuple $(e_{\varphi_1}, \dots, e_{\varphi_{n-2}})$ form an orthonormal basis of $T_w \mathbb{S}^{n-2}$ while $(e_{\theta}, e_{\varphi_1}, \dots, e_{\varphi_{n-2}})$ form an orthonormal basis of $T_v \mathbb{S}^{n-1}$. We note that e_{φ_i} only depends on $(\varphi_i, \dots, \varphi_{n-2})$ but we will still write $e_{\varphi_i}(\varphi_1, \dots, \varphi_{n-2})$ or, by abuse of notation, $e_{\varphi_i}(w)$. In view of (7.1), a tangent vector $\kappa \in T_v B$ can be decomposed according to

$$\begin{aligned} \kappa &= \kappa_{\parallel} e_{\theta}(\theta, \varphi_1, \dots, \varphi_{n-2}) + \sin \theta \kappa_T(\varphi_1, \dots, \varphi_{n-2}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}}), \\ \kappa_T(\varphi_1, \dots, \varphi_{n-2}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}}) &= \sum_{i=1}^{n-2} \kappa_{\varphi_i} e_{\varphi_i}(\varphi_1, \dots, \varphi_{n-2}) \in T_w(\mathbb{S}^2), \end{aligned} \quad (\text{B.7})$$

with $\kappa_{\varphi_i} \in \mathbb{R}$, $\forall i \in \{1, \dots, n-2\}$. We stress that $(\kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}})$ denotes the components of κ_T in $T_w \mathbb{S}^{n-2}$ and not those of κ . We now define $\vec{\psi}^b$ such that

$$\vec{\psi}^{\dagger}(\theta, \kappa_{\parallel}, w, \kappa_T) = \vec{\psi}^b(\theta, \kappa_{\parallel}, \varphi_1, \dots, \varphi_{n-2}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}}). \quad (\text{B.8})$$

and define $\psi_w^b, \vec{\psi}_T^b$ accordingly. Furthermore, we decompose $\vec{\psi}_T^b$ as follows:

$$\begin{aligned} \vec{\psi}_T^b(\theta, \kappa_{\parallel}, \varphi_1, \dots, \varphi_{n-2}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}}) \\ = \sum_{k=1}^{n-2} \psi_k^b(\theta, \kappa_{\parallel}, \varphi_1, \dots, \varphi_{n-2}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}}) e_{\varphi_k}(\varphi_1, \dots, \varphi_{n-2}), \end{aligned} \quad (\text{B.9})$$

with $\psi_k^b(\theta, \kappa_{\parallel}, \varphi_1, \dots, \varphi_{n-2}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}}) \in \mathbb{R}$.

We will use (B.4), (B.5), with R ranging in a generating set of O_{n-1} . We remind that we identify O_{n-1} with the isotropy subgroup of \mathbf{u} in O_n , meaning that all orthogonal transformations of \mathbb{R}^{n-1} are meant being orthogonal transformations of \mathbb{R}^n that leave \mathbf{u} invariant. We will mainly use three families of orthogonal transformation, which are successively presented in the following subsections.

B.2 Action of orthogonal transformations

B.2.1 First family of orthogonal transformations

Fix $(\theta, \varphi_1, \dots, \varphi_j, \dots, \varphi_{n-2}) \in (0, \pi)^{n-2} \times \mathbb{R} \pmod{2\pi}$. Define the rotation $R_{j,\beta}$ of \mathbb{R}^{n-1} which maps $(w, e_{\varphi_1}, \dots, e_{\varphi_{n-2}})(\theta, \varphi_1, \dots, \varphi_j, \dots, \varphi_{n-2})$ to $(w, e_{\varphi_1}, \dots, e_{\varphi_{n-2}})(\theta, \varphi_1, \dots, \varphi_j + \beta, \dots, \varphi_{n-2})$, for some $j \in \{1, \dots, n-2\}$ and some $\beta \in \mathbb{R}$ such that $(\theta, \varphi_1, \dots, \varphi_j + \beta, \dots, \varphi_{n-2}) \in (0, \pi)^{n-2} \times \mathbb{R} \pmod{2\pi}$. This rotation, which maps a direct orthonormal basis of \mathbb{R}^{n-1} to another one, is perfectly defined. For $\kappa \in T_v B$ written as in (B.7), we have

$$\begin{aligned} R_{j,\beta} \kappa &= \kappa_{\parallel} e_{\theta}(\theta, \varphi_1, \dots, \varphi_j + \beta, \dots, \varphi_{n-2}) + \sin \theta R_{j,\beta} \kappa_T, \\ R_{j,\beta} \kappa_T &= \sum_{i=1}^{n-2} \kappa_{\varphi_i} e_{\varphi_i}(\varphi_1, \dots, \varphi_j + \beta, \dots, \varphi_{n-2}). \end{aligned}$$

Therefore, $(\kappa_{\parallel}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}})$ are unchanged through the action of $R_{j,\beta}$.

- *Action of $R_{j,\beta}$ on ψ_w^b .* We have

$$\psi_w^{\dagger}(\theta, \kappa_{\parallel}, R_{j,\beta} w, R_{j,\beta} \kappa_T) = \psi_w^b(\theta, \kappa_{\parallel}, \varphi_1, \dots, \varphi_j + \beta, \dots, \varphi_{n-2}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}}).$$

Thus, thanks to (B.4), we get that ψ_w^b does not depend on $\varphi_j, \forall j \in \{1, \dots, n-2\}$, i.e.

$$\psi_w^b = \psi_w^b(\theta, \kappa_{\parallel}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}}). \quad (\text{B.10})$$

- *Action of $R_{j,\beta}$ on $\vec{\psi}_T^{\dagger}$.* On the one hand, we have

$$\begin{aligned} \vec{\psi}_T^{\dagger}(R_{j,\beta} v, R_{j,\beta} \kappa) &= \vec{\psi}_T^b(\theta, \kappa_{\parallel}, \varphi_1, \dots, \varphi_j + \beta, \dots, \varphi_{n-2}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}}) \\ &= \sum_{k=1}^{n-2} \psi_k^b(\theta, \kappa_{\parallel}, \varphi_1, \dots, \varphi_j + \beta, \dots, \varphi_{n-2}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}}) e_{\varphi_k}(\varphi_1, \dots, \varphi_j + \beta, \dots, \varphi_{n-2}), \end{aligned}$$

and on the other hand

$$\begin{aligned} R_{j,\beta} \vec{\psi}_T^{\dagger}(v, \kappa) &= R_{j,\beta} \vec{\psi}_T^b(\theta, \kappa_{\parallel}, \varphi_1, \dots, \varphi_j, \dots, \varphi_{n-2}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}}) \\ &= \sum_{k=1}^{n-2} \psi_k^b(\theta, \kappa_{\parallel}, \varphi_1, \dots, \varphi_j, \dots, \varphi_{n-2}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}}) e_{\varphi_k}(\varphi_1, \dots, \varphi_j + \beta, \dots, \varphi_{n-2}). \end{aligned}$$

Thus, thanks to (B.5), we get that for all k , ψ_k^b is independent of $\varphi_j, \forall j \in \{1, \dots, n-2\}$, i.e.

$$\psi_k^b = \psi_k^b(\theta, \kappa_{\parallel}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}}), \quad \forall k \in \{1, \dots, n-2\}. \quad (\text{B.11})$$

B.2.2 Second family of orthogonal transformations

We return to $\vec{\psi}_T^{\dagger}$ and when $\kappa_T \neq 0$, we decompose

$$\vec{\psi}_T^{\dagger}(\theta, \kappa_{\parallel}, w, \kappa_T) = \psi_r^{\dagger}(\theta, \kappa_{\parallel}, w, \kappa_T) \frac{\kappa_T}{|\kappa_T|} + \vec{\psi}_C^{\dagger}(\theta, \kappa_{\parallel}, w, \kappa_T), \quad \vec{\psi}_C^{\dagger}(\theta, \kappa_{\parallel}, w, \kappa_T) \cdot \kappa_T = 0. \quad (\text{B.12})$$

Taking the inner product of the first equation (B.12) with κ_T , we note that

$$\begin{aligned} \psi_r^\dagger\left(\theta, \kappa_\parallel, w, \sum_{m=1}^{n-2} \kappa_{\varphi_m} e_{\varphi_m}(w)\right) &= \frac{1}{|\kappa_T|} \sum_{k=1}^{n-2} \kappa_{\varphi_k} \psi_k^b(\theta, \kappa_\parallel, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}}) \\ &=: \psi_r^b(\theta, \kappa_\parallel, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}}), \end{aligned} \quad (\text{B.13})$$

thus defining ψ_r^b and emphasizing that ψ_r^b does not depend on $\varphi_1, \dots, \varphi_{n-2}$. Now, we fix $\kappa_{T0} \neq 0$ given in $T_w \mathbb{S}^{n-2}$ and decompose $\kappa_{T0} = \sum_{k=1}^{n-2} (\kappa_0)_{\varphi_k} e_{\varphi_k}$. Suppose, without loss of generality, that $(\kappa_0)_{\varphi_1} > 0$ (if $(\kappa_0)_{\varphi_1} < 0$, we will just change the sign of f_1 below; if $(\kappa_0)_{\varphi_1} = 0$, we just pick up a non-zero component $(\kappa_0)_{\varphi_j}$ as there always exists one since $\kappa_T \neq 0$). Then, define $f_1 = \kappa_{T0}/|\kappa_{T0}|$ and construct an orthonormal basis $(f_1, f_2, \dots, f_{n-2})$ of $T_w \mathbb{S}^{n-2}$ by applying the Gram-Schmidt orthonormalization principle to the basis $(f_1, e_{\varphi_2}, \dots, e_{\varphi_{n-2}})$. It is readily seen that (w, f_1, \dots, f_{n-2}) is a direct orthonormal basis of \mathbb{R}^{n-1} . Then, any vector $\kappa_T \in T_w \mathbb{S}^{n-2}$ can be decomposed in this basis according to $\kappa_T = \sum_{k=1}^{n-2} \tilde{\kappa}_i f_i$ and $(\tilde{\kappa}_1, \dots, \tilde{\kappa}_{n-2})$ depends smoothly of $(\kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}})$. In particular, κ_{T0} is decomposed into $\kappa_{T0} = \sum_{k=1}^{n-2} (\tilde{\kappa}_0)_i f_i$ with $(\tilde{\kappa}_0)_1 = |\kappa_{T0}|$ and $(\tilde{\kappa}_0)_i = 0$ for $i \in \{2, \dots, n-2\}$. Then, we can define a change of variables of $\vec{\psi}_T^\dagger(\theta, \kappa_\parallel, w, \cdot)$ as follows (we stress that $(\theta, \kappa_\parallel, w)$ are kept fixed):

$$\vec{\psi}_T^\dagger(\theta, \kappa_\parallel, w, \kappa_T) = \vec{\psi}_T^\sharp(\tilde{\kappa}_1, \dots, \tilde{\kappa}_{n-2}),$$

and we define ψ_r^\sharp and $\vec{\psi}_C^\sharp$ accordingly. Now, we decompose $\vec{\psi}_C^\sharp$ in the basis (f_1, \dots, f_{n-2}) :

$$\vec{\psi}_C^\sharp(\tilde{\kappa}_1, \dots, \tilde{\kappa}_{n-2}) = \sum_{k=1}^{n-2} \psi_k^\sharp(\tilde{\kappa}_1, \dots, \tilde{\kappa}_{n-2}) f_k. \quad (\text{B.14})$$

We note that the second condition of (B.12) implies that

$$\sum_{k=1}^{n-2} \psi_k^\sharp(\tilde{\kappa}_1, \dots, \tilde{\kappa}_{n-2}) \tilde{\kappa}_k = 0. \quad (\text{B.15})$$

Hence,

$$\vec{\psi}_T^\sharp(\tilde{\kappa}_1, \dots, \tilde{\kappa}_{n-2}) = \sum_{k=1}^{n-2} \left[\psi_r^\sharp(\tilde{\kappa}_1, \dots, \tilde{\kappa}_{n-2}) \frac{\tilde{\kappa}_k}{|\kappa_T|} + \psi_k^\sharp(\tilde{\kappa}_1, \dots, \tilde{\kappa}_{n-2}) \right] f_k.$$

Now, we apply (B.5) with the orthogonal reflection S_j in the hyperplane $\text{Span}\{w, f_1, \dots, \hat{f}_j, \dots, f_{n-2}\}$ of \mathbb{R}^{n-1} where \hat{f}_j means that f_j is omitted in the list and $j \in \{2, \dots, n-2\}$. Since $S_j w = w$, the angles $(\varphi_1, \dots, \varphi_{n-2})$ are unmodified by the operation of S_j . On the other hand,

$$S_j \kappa_T = \sum_{i \neq j} \tilde{\kappa}_i f_i - \tilde{\kappa}_j f_j,$$

meaning that $(\tilde{\kappa}_1, \dots, \tilde{\kappa}_{j-1}, \tilde{\kappa}_j, \tilde{\kappa}_{j+1}, \dots, \tilde{\kappa}_{n-2})$ is changed into $(\tilde{\kappa}_1, \dots, \tilde{\kappa}_{j-1}, -\tilde{\kappa}_j, \tilde{\kappa}_{j+1}, \dots, \tilde{\kappa}_{n-2})$.

We have

$$\begin{aligned}
\vec{\psi}_T^\dagger(\theta, \kappa_\parallel, S_j w, S_j \kappa_T) &= \vec{\psi}_T^\dagger(\theta, \kappa_\parallel, w, S_j \kappa_T) = \vec{\psi}_T^\dagger(\tilde{\kappa}_1, \dots, -\tilde{\kappa}_j, \dots, \tilde{\kappa}_{n-2}) \\
&= \sum_{k \neq j} \left(\psi_r^\dagger(\tilde{\kappa}_1, \dots, -\tilde{\kappa}_j, \dots, \tilde{\kappa}_{n-2}) \frac{\tilde{\kappa}_k}{|\kappa_T|} + \psi_k^\dagger(\tilde{\kappa}_1, \dots, -\tilde{\kappa}_j, \dots, \tilde{\kappa}_{n-2}) \right) f_k \\
&\quad + \left(-\psi_r^\dagger(\tilde{\kappa}_1, \dots, -\tilde{\kappa}_j, \dots, \tilde{\kappa}_{n-2}) \frac{\tilde{\kappa}_j}{|\kappa_T|} + \psi_j^\dagger(\tilde{\kappa}_1, \dots, -\tilde{\kappa}_j, \dots, \tilde{\kappa}_{n-2}) \right) f_j.
\end{aligned}$$

On the other hand, we have

$$\begin{aligned}
S_j \vec{\psi}_T^\dagger(\theta, \kappa_\parallel, w, \kappa_T) &= S_j \vec{\psi}_T^\dagger(\tilde{\kappa}_1, \dots, \tilde{\kappa}_j, \dots, \tilde{\kappa}_{n-2}) \\
&= \sum_{k \neq j} \left(\psi_r^\dagger(\tilde{\kappa}_1, \dots, \tilde{\kappa}_{n-2}) \frac{\tilde{\kappa}_k}{|\kappa_T|} + \psi_k^\dagger(\tilde{\kappa}_1, \dots, \tilde{\kappa}_{n-2}) \right) f_k \\
&\quad - \left(\psi_r^\dagger(\tilde{\kappa}_1, \dots, \kappa_{n-2}) \frac{\tilde{\kappa}_j}{|\kappa_T|} + \psi_j^\dagger(\tilde{\kappa}_1, \dots, \tilde{\kappa}_{n-2}) \right) f_j.
\end{aligned}$$

Now, we evaluate these two expressions at κ_{T0} . By construction (see (B.15)), we have

$$\kappa_{T0} = (\tilde{\kappa}_0)_1 f_1, \quad \psi_1^\dagger((\tilde{\kappa}_0)_1, 0, \dots, 0) = 0,$$

with $(\tilde{\kappa}_0)_1 = |\kappa_{T0}|$, and we remind that $j \in \{2, \dots, n-2\}$. Hence, thanks to (B.5), we get

$$\begin{aligned}
\psi_r^\dagger((\tilde{\kappa}_0)_1, 0, \dots, 0) \frac{(\tilde{\kappa}_0)_1}{|\kappa_T|} + \sum_{k=1}^{n-2} \psi_k^\dagger((\tilde{\kappa}_0)_1, 0, \dots, 0) f_k &= \psi_r^\dagger((\tilde{\kappa}_0)_1, 0, \dots, 0) \frac{(\tilde{\kappa}_0)_1}{|\kappa_T|} \\
&\quad + \sum_{k \neq j} \psi_k^\dagger((\tilde{\kappa}_0)_1, 0, \dots, 0) f_k - \psi_j^\dagger((\tilde{\kappa}_0)_1, 0, \dots, 0) f_j.
\end{aligned}$$

Hence, $\psi_j^\dagger((\tilde{\kappa}_0)_1, 0, \dots, 0) = 0, \forall j \in \{2, \dots, n-2\}$. With (B.14), (B.15), it follows that $\psi_C^\dagger((\tilde{\kappa}_0)_1, 0, \dots, 0) = 0$, i.e. $\vec{\psi}_C^\dagger(\theta, \kappa_\parallel, w, \kappa_{T0}) = 0$. Since, this equality is valid for all $\kappa_{T0} \neq 0$, we conclude that

$$\vec{\psi}_C^\dagger(\theta, \kappa_\parallel, w, \kappa_T) = 0, \quad \forall \kappa_T \in T_w \mathbb{S}^{n-2}, \quad \kappa_T \neq 0.$$

Hence, we can write

$$\vec{\psi}_T^\dagger(\theta, \kappa_\parallel, w, \kappa_T) = \psi_r^\dagger(\theta, \kappa_\parallel, w, \kappa_T) \frac{\kappa_T}{|\kappa_T|}, \quad \forall \kappa_T \in T_w \mathbb{S}^{n-2}, \quad \kappa_T \neq 0, \quad (\text{B.16})$$

with ψ_r^\dagger not depending on the angles $(\varphi_1, \dots, \varphi_{n-2})$ as shown by (B.13).

In the case where $\kappa_T = 0$, then, for all element of O_{n-1} such that $Rw = w$, (B.5) gives

$$\vec{\psi}_T^\dagger(\theta, \kappa_\parallel, w, 0) = R \vec{\psi}_T^\dagger(\theta, \kappa_\parallel, w, 0).$$

Such orthogonal transformations span O_{n-2} , the set of orthogonal transformations of $T_w \mathbb{S}^{n-2}$ and $\vec{\psi}_T^\dagger(\theta, \kappa_\parallel, w, 0)$ is an element of $T_w \mathbb{S}^{n-2}$ which is a fixed point of all elements of O_{n-2} . Such element must be equal to zero. Hence we get

$$\vec{\psi}_T^\dagger(\theta, \kappa_\parallel, w, 0) = 0. \quad (\text{B.17})$$

B.2.3 Third family of orthogonal transformations

We now use the planar rotation $R_{j,\ell,\beta}$ of \mathbb{R}^{n-1} of angle β in the plane $\text{Span}\{e_{\varphi_j}, e_{\varphi_\ell}\}$. In this rotation, no angle is changed because w is unchanged, being orthogonal to both e_{φ_j} and e_{φ_ℓ} . For this reason, we drop the dependences of e_θ , w and e_{φ_i} on $(\theta, \varphi_1, \dots, \varphi_{n-2})$ since no confusion is possible. Now, we get

$$R_{j,\ell,\beta} e_{\varphi_j} = \cos \beta e_{\varphi_j} + \sin \beta e_{\varphi_\ell}, \quad R_{j,\ell,\beta} e_{\varphi_\ell} = -\sin \beta e_{\varphi_j} + \cos \beta e_{\varphi_\ell}.$$

Hence, For $\kappa \in T_v B$ written as in (B.7), we have

$$\begin{aligned} R_{j,\ell,\beta} \kappa &= \kappa_{\parallel} e_\theta + \sin \theta R_{j,\ell,\beta} \kappa_T, \\ R_{j,\ell,\beta} \kappa_T &= \sum_{i \notin \{j,\ell\}} \kappa_{\varphi_i} e_{\varphi_i} + (\kappa_{\varphi_j} \cos \beta - \kappa_{\varphi_\ell} \sin \beta) e_{\varphi_j} + (\kappa_{\varphi_j} \sin \beta + \kappa_{\varphi_\ell} \cos \beta) e_{\varphi_\ell}. \end{aligned}$$

Therefore, $(\kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}})$ is changed into $(\kappa_{\varphi_1}, \dots, \kappa_{\varphi_j} \cos \beta - \kappa_{\varphi_\ell} \sin \beta, \dots, \kappa_{\varphi_j} \sin \beta + \kappa_{\varphi_\ell} \cos \beta, \dots, \kappa_{\varphi_{n-2}})$. We return to the coordinates (B.8) which define $\psi_w^b(\theta, \kappa_{\parallel}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}})$ and $\psi_r^b(\theta, \kappa_{\parallel}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_{n-2}})$.

- Action of $R_{j,\ell,\beta}$ on ψ_w^b . Eq. (B.4) gives

$$\psi_w^\dagger(\theta, \kappa_{\parallel}, R_{j,\ell,\beta} w, R_{j,\ell,\beta} \kappa_T) = \psi_w^\dagger(\theta, \kappa_{\parallel}, w, \kappa_T). \quad (\text{B.18})$$

We have

$$\begin{aligned} \psi_w^\dagger(\theta, \kappa_{\parallel}, R_{j,\ell,\beta} w, R_{j,\ell,\beta} \kappa_T) \\ = \psi_w^b(\theta, \kappa_{\parallel}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_j} \cos \beta - \kappa_{\varphi_\ell} \sin \beta, \dots, \kappa_{\varphi_j} \sin \beta + \kappa_{\varphi_\ell} \cos \beta, \dots, \kappa_{\varphi_{n-2}}). \end{aligned}$$

Thus, (B.18) leads to

$$\begin{aligned} \psi_w^b(\theta, \kappa_{\parallel}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_j} \cos \beta - \kappa_{\varphi_\ell} \sin \beta, \dots, \kappa_{\varphi_j} \sin \beta + \kappa_{\varphi_\ell} \cos \beta, \dots, \kappa_{\varphi_{n-2}}) \\ = \psi_w^b(\theta, \kappa_{\parallel}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_j}, \dots, \kappa_{\varphi_\ell}, \dots, \kappa_{\varphi_{n-2}}), \quad \forall \beta \in \mathbb{R}. \end{aligned} \quad (\text{B.19})$$

Define

$$\kappa_{j\ell} = (\kappa_{\varphi_j}^2 + \kappa_{\varphi_\ell}^2)^{1/2}, \quad \omega_j = \frac{\kappa_{\varphi_j}}{\kappa_{j\ell}} =: \cos \gamma, \quad \omega_\ell = \frac{\kappa_{\varphi_\ell}}{\kappa_{j\ell}} =: \sin \gamma,$$

with $\gamma \in \mathbb{R}$. Then, (B.19) can be written

$$\begin{aligned} \psi_w^b(\theta, \kappa_{\parallel}, \kappa_{\varphi_1}, \dots, \kappa_{j\ell} \cos(\gamma + \beta), \dots, \kappa_{j\ell} \sin(\gamma + \beta), \dots, \kappa_{\varphi_{n-2}}) \\ = \psi_w^b(\theta, \kappa_{\parallel}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_j}, \dots, \kappa_{\varphi_\ell}, \dots, \kappa_{\varphi_{n-2}}), \quad \forall \beta \in \mathbb{R}. \end{aligned}$$

Taking $\beta = -\gamma$, we get

$$\begin{aligned} \psi_w^b(\theta, \kappa_{\parallel}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_j}, \dots, \kappa_{\varphi_\ell}, \dots, \kappa_{\varphi_{n-2}}) \\ = \psi_w^b(\theta, \kappa_{\parallel}, \kappa_{\varphi_1}, \dots, \kappa_{j\ell}, \dots, 0, \dots, \kappa_{\varphi_{n-2}}). \end{aligned}$$

Proceeding successively with $(j, \ell) = (n-3, n-2), (n-4, n-3), \dots, (2, 3), (1, 2)$, we get

$$\psi_w^b(\theta, \kappa_{\parallel}, \kappa_{\varphi_1}, \dots, \kappa_{\varphi_j}, \dots, \kappa_{\varphi_\ell}, \dots, \kappa_{\varphi_{n-2}}) = \psi_w^b(\theta, \kappa_{\parallel}, |\kappa_T|, 0, \dots, 0).$$

Therefore, defining ψ_w by,

$$\psi_w(\theta, \kappa_{\parallel}, |\kappa_T|) = \psi_w^b(\theta, \kappa_{\parallel}, |\kappa_T|, 0, \dots, 0),$$

we get the first equality (B.6).

- *Action of $R_{j,\ell,\beta}$ on ψ_r^b .*

Using (B.16), we have

$$\vec{\psi}_T^\dagger(\theta, \kappa_{\parallel}, R_{j,\ell,\beta} w, R_{j,\ell,\beta} \kappa_T) = \psi_r^\dagger(\theta, \kappa_{\parallel}, R_{j,\ell,\beta} w, R_{j,\ell,\beta} \kappa_T) \frac{R_{j,\ell,\beta} \kappa_T}{|R_{j,\ell,\beta} \kappa_T|}.$$

On the other hand we have

$$R_{j,\ell,\beta} \vec{\psi}_T^\dagger(\theta, \kappa_{\parallel}, w, \kappa_T) = \psi_r^\dagger(\theta, \kappa_{\parallel}, w, \kappa_T) \frac{R_{j,\ell,\beta} \kappa_T}{|\kappa_T|}.$$

Thus, thanks to (B.5) and the fact that $|R_{j,\ell,\beta} \kappa_T| = |\kappa_T|$, we get

$$\psi_r^\dagger(\theta, \kappa_{\parallel}, R_{j,\ell,\beta} w, R_{j,\ell,\beta} \kappa_T) = \psi_r^\dagger(\theta, \kappa_{\parallel}, w, \kappa_T),$$

which is the same equation as (B.18) for ψ_w^\dagger . Therefore, the same conclusion applies. Hence, with

$$\psi_r(\theta, \kappa_{\parallel}, |\kappa_T|) = \psi_r^b(\theta, \kappa_{\parallel}, |\kappa_T|, 0, \dots, 0),$$

we get the second equality (B.6).

Conversely, we easily check that (7.12) satisfies the invariance relation (6.21). This ends the proof of Theorem 7.1. \blacksquare

C Auxilliary computations for Subsection 7.2

In all this section, we assume that $\vec{\psi} \in \mathcal{D}_{\text{inv}}(M)$.

C.1 Main statements

We let $(\mathbf{e}_1, \dots, \mathbf{e}_{n-1}, \mathbf{u})$ be the canonical basis of \mathbb{R}^n ordered in the natural order (which is different from the order used in the proof of Theorem 7.1). Thus, $\{\mathbf{u}\}^\perp = \text{Span}(\mathbf{e}_1, \dots, \mathbf{e}_{n-1})$ and we denote by $\psi_i = \vec{\psi} \cdot \mathbf{e}_i$. From Section 6.2, we know that $\kappa_T \in \{\mathbf{u}\}^\perp$ and we define $\kappa_{Ti} = \kappa_T \cdot \mathbf{e}_i$, $i = 1, \dots, n-1$ (note that κ_{Ti} is different from κ_T^i defined in Subsection 7.1.1). Finally, we obviously have $\mathbf{u}_i =: \mathbf{u} \cdot \mathbf{e}_i = 0$. Now, we decompose $\vec{\psi}$ according to (7.15). We first state the following auxilliary two lemmas, which will be proved in Subsections C.2 and C.3 respectively.

Lemma C.1 (gradients of θ , κ_{\parallel} and $|\kappa_T|$).

We have

$$\nabla_v \theta = e_{\theta}, \quad \nabla_v \kappa_{\parallel} = \cos \theta \kappa_T, \quad \nabla_v |\kappa_T| = -\frac{\cos \theta}{\sin^2 \theta} \kappa_{\parallel} \frac{\kappa_T}{|\kappa_T|} - \frac{\cos \theta}{\sin \theta} |\kappa_T| e_{\theta}, \quad (\text{C.1})$$

$$\nabla_{\kappa} \theta = 0, \quad \nabla_{\kappa} \kappa_{\parallel} = e_{\theta}, \quad \nabla_{\kappa} |\kappa_T| = \frac{1}{\sin \theta} \frac{\kappa_T}{|\kappa_T|}. \quad (\text{C.2})$$

Lemma C.2 (Gradients of w_i and $\kappa_{Ti}/|\kappa_T|$).

For all $i \in \{1, \dots, n-1\}$, we have

$$\nabla_v w_i = \frac{1}{\sin \theta} (P_{v^{\perp}} \mathbf{e}_i - \cos \theta w_i e_{\theta}), \quad \nabla_{\kappa} w_i = 0, \quad (\text{C.3})$$

$$\nabla_v \frac{\kappa_{Ti}}{|\kappa_T|} = \frac{\cos^2 \theta}{\sin^2 \theta} \frac{\kappa_{\parallel}}{|\kappa_T|} w_i e_{\theta} + \left(-\frac{w_i}{\sin \theta} + \frac{\cos \theta}{\sin^2 \theta} \frac{\kappa_{\parallel} \kappa_{Ti}}{|\kappa_T|^2} \right) \frac{\kappa_T}{|\kappa_T|} - \frac{\cos \theta}{\sin^2 \theta} \frac{\kappa_{\parallel}}{|\kappa_T|} P_{v^{\perp}} \mathbf{e}_i, \quad (\text{C.4})$$

$$\nabla_{\kappa} \frac{\kappa_{Ti}}{|\kappa_T|} = -\frac{\cos \theta}{\sin \theta} \frac{w_i}{|\kappa_T|} e_{\theta} - \frac{1}{\sin \theta} \frac{\kappa_{Ti}}{|\kappa_T|^2} \frac{\kappa_T}{|\kappa_T|} + \frac{1}{\sin \theta |\kappa_T|} P_{v^{\perp}} \mathbf{e}_i. \quad (\text{C.5})$$

Thanks to these lemmas, we can state the

Proposition C.3 (gradients of ψ_i in terms of ψ_w and ψ_r). We have

$$\begin{aligned} (\nabla_v \psi_i)^{\dagger} &= e_{\theta} \left(w_i \left(\partial_{\theta} \psi_w - \frac{\cos \theta}{\sin \theta} |\kappa_T| \partial_{\kappa_{\perp}} \psi_w - \frac{\cos \theta}{\sin \theta} \psi_w + \frac{\cos^2 \theta}{\sin^2 \theta} \frac{\kappa_{\parallel}}{|\kappa_T|} \psi_r \right) \right. \\ &\quad \left. + \frac{\kappa_{Ti}}{|\kappa_T|} \left(\partial_{\theta} \psi_r - \frac{\cos \theta}{\sin \theta} |\kappa_T| \partial_{\kappa_{\perp}} \psi_r \right) \right) \\ &\quad + \frac{\kappa_T}{|\kappa_T|} \left(w_i \left(\cos \theta |\kappa_T| \partial_{\kappa_{\parallel}} \psi_w - \frac{\cos \theta}{\sin^2 \theta} \kappa_{\parallel} \partial_{\kappa_{\perp}} \psi_w - \frac{1}{\sin \theta} \psi_r \right) \right. \\ &\quad \left. + \frac{\kappa_{Ti}}{|\kappa_T|} \left(\cos \theta |\kappa_T| \partial_{\kappa_{\parallel}} \psi_r - \frac{\cos \theta}{\sin^2 \theta} \kappa_{\parallel} \partial_{\kappa_{\perp}} \psi_r + \frac{\cos \theta}{\sin^2 \theta} \frac{\kappa_{\parallel}}{|\kappa_T|} \psi_r \right) \right) \\ &\quad + P_{v^{\perp}} \mathbf{e}_i \left(\frac{1}{\sin \theta} \psi_w - \frac{\cos \theta}{\sin^2 \theta} \frac{\kappa_{\parallel}}{|\kappa_T|} \psi_r \right), \end{aligned} \quad (\text{C.6})$$

$$\begin{aligned} (\nabla_{\kappa} \psi_i)^{\dagger} &= e_{\theta} \left(w_i \left(\partial_{\kappa_{\parallel}} \psi_w - \frac{\cos \theta}{\sin \theta} \frac{1}{|\kappa_T|} \psi_r \right) + \frac{\kappa_{Ti}}{|\kappa_T|} \partial_{\kappa_{\parallel}} \psi_r \right) \\ &\quad + \frac{\kappa_T}{|\kappa_T|} \frac{1}{\sin \theta} \left(w_i \partial_{\kappa_{\perp}} \psi_w + \frac{\kappa_{Ti}}{|\kappa_T|} \left(\partial_{\kappa_{\perp}} \psi_r - \frac{1}{|\kappa_T|} \psi_r \right) \right) \\ &\quad + P_{v^{\perp}} \mathbf{e}_i \frac{1}{\sin \theta |\kappa_T|} \psi_r. \end{aligned} \quad (\text{C.7})$$

Proof. By the chain rule, we get

$$\begin{aligned} \nabla_v \psi_i &= w_i \left(\partial_{\theta} \psi_w \nabla_v \theta + \partial_{\kappa_{\parallel}} \psi_w \nabla_v \kappa_{\parallel} + \partial_{\kappa_{\perp}} \psi_w \nabla_v (|\kappa_T|) \right) + \psi_w \nabla_v w_i \\ &\quad + \frac{\kappa_{Ti}}{|\kappa_T|} \left(\partial_{\theta} \psi_r \nabla_v \theta + \partial_{\kappa_{\parallel}} \psi_r \nabla_v \kappa_{\parallel} + \partial_{\kappa_{\perp}} \psi_r \nabla_v (|\kappa_T|) \right) + \psi_r \nabla_v \left(\frac{\kappa_{Ti}}{|\kappa_T|} \right), \end{aligned} \quad (\text{C.8})$$

and similarly for $\nabla_\kappa \psi_i$ by changing v into κ everywhere. Now, inserting the formulas found in Lemmas C.1 and C.2, in these expressions, we get the result after some straightforward computations. ■

As a consequence of this proposition, we get

Proposition C.4 (Expression of $|\nabla_\kappa \vec{\psi}|^2$ and $\{\vec{\psi}, \mathbf{H}_u\}$).

(i) $|\nabla_\kappa \vec{\psi}|^2$ is given by (7.20).

(ii) $\{\vec{\psi}, \mathbf{H}_u\}^\dagger$ is given by (7.33).

Proof. (i) We take the squared norm of (C.7). We note that both vectors e_θ and $\kappa_T/|\kappa_T|$ are normed and they are orthogonal to each other. On the other hand, we have

$$\begin{aligned} |P_{v^\perp} e_i|^2 &= |e_i|^2 - (e_i \cdot v)^2 = 1 - \sin^2 \theta w_i^2, \\ e_\theta \cdot P_{v^\perp} e_i &= e_\theta \cdot e_i = \cos \theta w_i, \\ \frac{\kappa_T}{|\kappa_T|} \cdot P_{v^\perp} e_i &= \frac{\kappa_T}{|\kappa_T|} \cdot e_i = \frac{\kappa_{Ti}}{|\kappa_T|}. \end{aligned}$$

It results, after some cancellations:

$$\begin{aligned} |\nabla_\kappa \vec{\psi}|^2 &= \left(w_i \partial_{\kappa_\parallel} \psi_w + \frac{\kappa_{Ti}}{|\kappa_T|} \partial_{\kappa_\parallel} \psi_r \right)^2 + \frac{1}{\sin^2 \theta} \left(w_i \partial_{\kappa_\perp} \psi_w + \frac{\kappa_{Ti}}{|\kappa_T|} \partial_{\kappa_\perp} \psi_r \right)^2 \\ &\quad + \left(1 - w_i^2 - \frac{\kappa_{Ti}^2}{|\kappa_T|^2} \right) \frac{1}{\sin^2 \theta |\kappa_T|^2} \psi_r^2. \end{aligned}$$

Then, using that w and $\kappa_T/|\kappa_T|$ are normed and orthogonal to each other, summing over $i \in \{1, \dots, n-1\}$, we get (7.20).

(ii) We have

$$\{\vec{\psi}, \mathbf{H}_u\} = \langle \kappa, \nabla_v \vec{\psi} \rangle + v \langle P_{v^\perp} \mathbf{u}, \nabla_\kappa \vec{\psi} \rangle. \quad (\text{C.9})$$

We compute:

$$\begin{aligned} \langle \kappa, e_\theta \rangle &= \kappa_\parallel, \\ \langle \kappa, \frac{\kappa_T}{|\kappa_T|} \rangle &= \sin \theta |\kappa_T|, \\ \langle \kappa, P_{v^\perp} \mathbf{e}_i \rangle &= \langle \kappa, \mathbf{e}_i \rangle = \kappa_\parallel e_\theta \cdot \mathbf{e}_i + \sin \theta \kappa_{Ti} = \cos \theta \kappa_\parallel w_i + \sin \theta |\kappa_T| \frac{\kappa_{Ti}}{|\kappa_T|}. \end{aligned}$$

and

$$\begin{aligned} \langle P_{v^\perp} \mathbf{u}, e_\theta \rangle &= \mathbf{u} \cdot e_\theta = -\sin \theta, \\ \langle P_{v^\perp} \mathbf{u}, \frac{\kappa_T}{|\kappa_T|} \rangle &= \mathbf{u} \cdot \frac{\kappa_T}{|\kappa_T|} = 0, \\ \langle P_{v^\perp} \mathbf{u}, P_{v^\perp} \mathbf{e}_i \rangle &= -(\mathbf{u} \cdot v)(\mathbf{e}_i \cdot v) = -\cos \theta \sin \theta w_i. \end{aligned}$$

Inserting the first set of expressions into (C.6), multiplying the second set by v and insrtng it into (C.7), and adding the two resulting expressions, after some cancellations, we get

$$\{\psi_i, \mathbf{H}_u\}^\dagger = w_i (\mathcal{U}_1 \psi_w - |\kappa_T| \psi_r) + \frac{\kappa_{Ti}}{|\kappa_T|} (\mathcal{U}_1 \psi_r + |\kappa_T| \psi_r). \quad (\text{C.10})$$

Then, multiplying this expression by \mathbf{e}_i and summing over i , we get (7.33). ■

C.2 Proof of Lemma C.1

We first show the following:

$$\theta = \cos^{-1}(v \cdot \mathbf{u}), \quad \kappa_{\parallel} = -\frac{\kappa \cdot \mathbf{u}}{\sin \theta}, \quad |\kappa_T| = \frac{(|\kappa|^2 - \kappa_{\parallel}^2)^{1/2}}{\sin \theta}. \quad (\text{C.11})$$

Indeed, the first formula (C.11) directly comes from the first formula (4.2). The second formula (C.11) comes from the second and third formulas (7.3) and from the fact that $\kappa \cdot v = 0$. Finally, the third formula (C.11) comes from (7.1) and the fourth formula (7.3).

Then, taking the gradient with respect to v of the first expression in (C.11) and using the first expression (3.25) and the second equation (7.3), we get

$$\nabla_v \theta = -\left(1 - (v \cdot \mathbf{u})^2\right)^{-1/2} \nabla_v (v \cdot \mathbf{u}) = -\frac{1}{\sin \theta} P_{v^\perp} \mathbf{u} = e_\theta,$$

which is the first formula (C.1).

Now, by the second expression in (C.11), and thanks to the previous identity, we get

$$\nabla_v \kappa_{\parallel} = \frac{\cos \theta}{\sin^2 \theta} (\kappa \cdot \mathbf{u}) \nabla_v \theta - \frac{1}{\sin \theta} \nabla_v (\kappa \cdot \mathbf{u}) = -\frac{\cos \theta}{\sin \theta} \kappa_{\parallel} e_\theta - \frac{1}{\sin \theta} \nabla_v (\kappa \cdot \mathbf{u}). \quad (\text{C.12})$$

Now, let $a \in T_v B$ and denote by α the pair (v, κ) as usual. We compute

$$\begin{aligned} \langle \nabla_v (\kappa \cdot \mathbf{u}), a \rangle_v &= \langle B_\alpha^H \nabla_\alpha (\kappa \cdot \mathbf{u}), a \rangle_v = \langle \nabla_\alpha (\kappa \cdot \mathbf{u}), \mathcal{L}_\alpha^H a \rangle_\alpha \\ &= (d(\kappa \cdot \mathbf{u}))_\alpha (\mathcal{L}_\alpha^H a) = \frac{d}{dt} (\tilde{\kappa}(t) \cdot \mathbf{u}) \Big|_{t=0}, \end{aligned}$$

where $\tilde{\alpha}(t) = (\tilde{v}(t), \tilde{\kappa}(t))$ is a path in M such that $\tilde{\alpha}(0) = (v, \kappa)$ and $(d/dt)(\tilde{v}(t), \tilde{\kappa}(t))|_{t=0} = \mathcal{L}_\alpha^H a = (a, -(a \cdot \kappa)v)$, by (2.42). Let (f_1, \dots, f_n) be an orthonormal basis such that $v = f_n$, $a = |a|f_{n-1}$, $\kappa = \tilde{\kappa}_{n-2}f_{n-2} + \tilde{\kappa}_{n-1}f_{n-1}$ with $(\tilde{\kappa}_{n-2}, \tilde{\kappa}_{n-1}) \in \mathbb{R}^2$. Then define the path $\tilde{\alpha}(t)$ as follows:

$$\tilde{v}(t) = \sin(|a|t) f_{n-1} + \cos(|a|t) f_n, \quad \tilde{\kappa}(t) = \tilde{\kappa}_{n-2}f_{n-2} + \tilde{\kappa}_{n-1}f_{n-1} - \tilde{\kappa}_{n-1} \tan(|a|t) f_n.$$

Then, this path satisfies the above requirements and we have

$$\tilde{\kappa}(t) \cdot \mathbf{u} = \tilde{\kappa}_{n-2}f_{n-2} \cdot \mathbf{u} + \tilde{\kappa}_{n-1}f_{n-1} \cdot \mathbf{u} - \tilde{\kappa}_{n-1} \tan(|a|t) f_n \cdot \mathbf{u}.$$

So,

$$\frac{d}{dt} (\tilde{\kappa}(t) \cdot \mathbf{u}) \Big|_{t=0} = -|a| \tilde{\kappa}_{n-1} f_n \cdot \mathbf{u} = -(a \cdot \kappa)(v \cdot \mathbf{u}) = \langle -\cos \theta \kappa, a \rangle_v.$$

Thus,

$$\nabla_v (\kappa \cdot \mathbf{u}) = -\cos \theta \kappa.$$

Inserting this into (C.12), we get

$$\nabla_v \kappa_{\parallel} = \frac{\cos \theta}{\sin \theta} (-\kappa_{\parallel} e_\theta + \kappa) = \cos \theta \kappa_T,$$

which is the second formula (C.1).

Thanks to the third equation (C.11), we have

$$\nabla_v |\kappa_T| = \frac{1}{2(|\kappa|^2 - \kappa_{\parallel}^2)^{1/2} \sin \theta} (\nabla_v |\kappa|^2 - 2\kappa_{\parallel} \nabla_v \kappa_{\parallel}) - (|\kappa|^2 - \kappa_{\parallel}^2)^{1/2} \frac{\cos \theta}{\sin^2 \theta} \nabla_v \theta. \quad (\text{C.13})$$

Now, in view of the second formula in (3.26), we have $\nabla_v |\kappa|^2 = 0$. Inserting the first two equations of (C.1) into (C.13), we get the third equation (C.1).

The first equation (C.2) follows directly from the second equation (3.25).

By the second equation (3.25) and the second equation (C.11), we get

$$\nabla_{\kappa} \kappa_{\parallel} = -\frac{1}{\sin \theta} \nabla_{\kappa} (\kappa \cdot \mathbf{u}). \quad (\text{C.14})$$

Now, let $\tau \in T_v B$. We compute

$$\begin{aligned} \langle \nabla_{\kappa} (\kappa \cdot \mathbf{u}), \tau \rangle_v &= \langle B_{\alpha}^V \nabla_{\alpha} (\kappa \cdot \mathbf{u}), \tau \rangle_v = \langle \nabla_{\alpha} (\kappa \cdot \mathbf{u}), \mathcal{L}_{\alpha}^V \tau \rangle_{\alpha} \\ &= (d(\kappa \cdot \mathbf{u}))_{\alpha} (\mathcal{L}_{\alpha}^V \tau) = \frac{d}{dt} (\tilde{\kappa}(t) \cdot \mathbf{u}) \Big|_{t=0}, \end{aligned}$$

where $\tilde{\alpha}(t) = (\tilde{v}(t), \tilde{\kappa}(t))$ is a path in M such that $\tilde{\alpha}(0) = (v, \kappa)$ and $(d/dt)(\tilde{v}(t), \tilde{\kappa}(t))|_{t=0} = \mathcal{L}_{\alpha}^V \tau = (0, \tau)$, by (2.38). The path $\tilde{\alpha}(t) = (v, \kappa + t\tau)$ satisfies these requirements. We have $\tilde{\kappa}(t) \cdot \mathbf{u} = (\kappa \cdot \mathbf{u}) + t(\tau \cdot \mathbf{u})$, hence $(d/dt)(\tilde{\kappa}(t) \cdot \mathbf{u})|_{t=0} = (\tau \cdot \mathbf{u}) = \langle \tau, P_{v^{\perp}} \mathbf{u} \rangle_v$, showing that

$$\nabla_{\kappa} (\kappa \cdot \mathbf{u}) = P_{v^{\perp}} \mathbf{u} = -\sin \theta e_{\theta}.$$

Inserting this formula into (C.14) leads to the second formula (C.2).

Finally, thanks to the first formula in (3.26), we get

$$\nabla_{\kappa} |\kappa_T| = \frac{1}{\sin \theta (|\kappa|^2 - \kappa_{\parallel}^2)^{1/2}} (\kappa - \kappa_{\parallel} \nabla_{\kappa} \kappa_{\parallel}),$$

and thanks to the second equation (C.2), this leads to the third equation (C.2). This ends the proof of Lemma C.1. \blacksquare

C.3 Proof of Lemma C.2

With the first equation (7.3), we have $w_i = (1 - (v \cdot \mathbf{u})^2)^{-1/2} v_i$. So,

$$\begin{aligned} \nabla_v w_i &= (1 - (v \cdot \mathbf{u})^2)^{-1/2} \nabla_v (v \cdot \mathbf{e}_i) + (1 - (v \cdot \mathbf{u})^2)^{-3/2} v_i (v \cdot \mathbf{u}) \nabla_v (v \cdot \mathbf{u}), \\ &= \frac{1}{\sin \theta} P_{v^{\perp}} \mathbf{e}_i + \frac{\cos \theta}{\sin^3 \theta} v_i P_{v^{\perp}} \mathbf{u} = \frac{1}{\sin \theta} P_{v^{\perp}} \mathbf{e}_i - \frac{\cos \theta}{\sin \theta} w_i e_{\theta}, \end{aligned}$$

which is the first formula (C.3).

The second formula (C.3) follows directly from the second formula of (3.25).

With the first equation (7.2), we get

$$e_\theta \cdot \mathbf{e}_i = \cos \theta w_i. \quad (\text{C.15})$$

Then, thanks to (7.1) and the third equation (C.11), we have

$$\frac{\kappa_{Ti}}{|\kappa_T|} = \frac{\kappa_i - \kappa_\parallel \cos \theta w_i}{(|\kappa|^2 - \kappa_\parallel^2)^{1/2}}. \quad (\text{C.16})$$

We deduce that

$$\begin{aligned} \nabla_v \frac{\kappa_{Ti}}{|\kappa_T|} &= \frac{1}{(|\kappa|^2 - \kappa_\parallel^2)^{1/2}} (\nabla_v \kappa_i - \cos \theta w_i \nabla_v \kappa_\parallel + \kappa_\parallel w_i \sin \theta \nabla_v \theta - \kappa_\parallel \cos \theta \nabla_v w_i) \\ &\quad - \frac{1}{2} \frac{1}{(|\kappa|^2 - \kappa_\parallel^2)^{3/2}} (\nabla_v |\kappa|^2 - 2\kappa_\parallel \nabla_v \kappa_\parallel) (\kappa_i - \kappa_\parallel \cos \theta w_i). \end{aligned}$$

By the same proof as for $\nabla_v(\kappa \cdot \mathbf{u})$ in the previous lemma, we have

$$\nabla_v \kappa_i = \nabla_v(\kappa \cdot \mathbf{e}_i) = -(v \cdot \mathbf{e}_i) \kappa = -\sin \theta w_i (\kappa_\parallel e_\theta + \sin \theta \kappa_T).$$

Then, the second equation (3.26), the first equation (C.3) and Lemma C.1 lead to

$$\begin{aligned} \nabla_v \frac{\kappa_{Ti}}{|\kappa_T|} &= \frac{1}{\sin \theta |\kappa_T|} \left(-\sin \theta w_i (\kappa_\parallel e_\theta + \sin \theta \kappa_T) - \cos \theta^2 w_i \kappa_T + \kappa_\parallel w_i \sin \theta e_\theta \right. \\ &\quad \left. - \kappa_\parallel \frac{\cos \theta}{\sin \theta} (P_{v^\perp} \mathbf{e}_i - \cos \theta w_i e_\theta) \right) + \frac{\cos \theta}{\sin^3 \theta} \frac{\kappa_\parallel}{|\kappa_T|^2} (\kappa_i - \kappa_\parallel \cos \theta w_i) \frac{\kappa_T}{|\kappa_T|}, \end{aligned}$$

which, after rearrangement, leads to (C.4).

Similarly, thanks to (C.16), we obtain

$$\begin{aligned} \nabla_\kappa \frac{\kappa_{Ti}}{|\kappa_T|} &= \frac{1}{(|\kappa|^2 - \kappa_\parallel^2)^{1/2}} (\nabla_\kappa \kappa_i - \cos \theta w_i \nabla_\kappa \kappa_\parallel) \\ &\quad - \frac{1}{2} \frac{1}{(|\kappa|^2 - \kappa_\parallel^2)^{3/2}} (\nabla_\kappa |\kappa|^2 - 2\kappa_\parallel \nabla_\kappa \kappa_\parallel) (\kappa_i - \kappa_\parallel \cos \theta w_i) \end{aligned}$$

By the same proof as for $\nabla_\kappa(\kappa \cdot \mathbf{u})$ in the previous lemma, we have

$$\nabla_\kappa \kappa_i = \nabla_\kappa(\kappa \cdot \mathbf{e}_i) = P_{v^\perp} \mathbf{e}_i.$$

Then, the first equation (3.26) and Lemma C.1 lead to

$$\nabla_\kappa \frac{\kappa_{Ti}}{|\kappa_T|} = \frac{1}{\sin \theta |\kappa_T|} (P_{v^\perp} \mathbf{e}_i - \cos \theta w_i e_\theta) - \frac{1}{\sin^3 \theta |\kappa_T|^3} (\kappa - \kappa_\parallel e_\theta) (\kappa_i - \kappa_\parallel \cos \theta w_i),$$

which, after rearrangement, yields (C.5). ■