

# MIRI spectrophotometry of GN-z11: Detection and nature of an optical red continuum component

A. Crespo Gómez<sup>1</sup>, L. Colina<sup>2</sup>, P. G. Pérez-González<sup>2</sup>, J. Álvarez-Márquez<sup>2</sup>, M. García-Marín<sup>3</sup>, A. Alonso-Herrero<sup>4</sup>, M. Annunziatella<sup>2</sup>, A. Bik<sup>5</sup>, S. Bosman<sup>6,7</sup>, A. J. Bunker<sup>8</sup>, A. Labiano<sup>4,9</sup>, D. Langeroodi<sup>10</sup>, P. Rinaldi<sup>11</sup>, G. Östlin<sup>5</sup>, L. Boogaard<sup>15</sup>, S. Gillman<sup>12,13</sup>, G. Barro<sup>14</sup>, S. L. Finkelstein<sup>16</sup>, and G. C. K. Leung<sup>17</sup>

(Affiliations can be found after the references)

December 3, 2025

## ABSTRACT

We present new MIRI F560W, F770W and F1000W imaging of the galaxy GN-z11 at a redshift of 10.603. We report a significant detection ( $14\sigma$ ) in the F560W and F770W images, and a marginal detection ( $3.2\sigma$ ) in the F1000W filter. The new MIRI observations cover the optical-red spectral range, and significantly extend previous NIRCам wavelength coverage from rest-frame  $0.38\mu\text{m}$  up to  $0.86\mu\text{m}$ . In this work, we analyse the spectral energy distribution (SED) combining this new MIRI imaging data with archival NIRSpect/Prism and MRS spectroscopy, and NIRCам imaging, i.e. covering the rest-frame  $0.12 - 0.86\mu\text{m}$ . New constraints such as the equivalent widths of the strong optical lines ([O III] 5008, H $\beta$  and H $\alpha$ ) and the continuum emission at rest-frame  $0.48\mu\text{m}$ ,  $0.66\mu\text{m}$ , and  $0.86\mu\text{m}$ , free from emission line contributions, are presented. The continuum emission shows a flat energy distribution, in  $f_\nu$ , up to  $0.5\mu\text{m}$ , compatible with the presence of a mixed stellar population of young ( $4 \pm 1$  Myr) and mature ( $63 \pm 23$  Myr) stars that also account for the [O III], H $\beta$  and H $\alpha$  emission lines. The continuum at rest-frame  $0.66\mu\text{m}$  shows a  $36 \pm 3\%$  flux excess above the predicted flux for a mixed stellar population, pointing to the presence of an additional source contributing at these wavelengths. This excess increases to  $91 \pm 28\%$  at rest-frame  $0.86\mu\text{m}$ , although with a large uncertainty due to the marginal detection in the F1000W filter. We consider that hot dust emission in the dusty torus around a type 2 AGN could be responsible for the observed excess. Alternatively, this excess could be due to hot dust emission or to a photoluminescence dust process (Extended Red Emission, ERE) under the extreme UV radiation field, as observed in some local metal-poor galaxies and in young compact starbursts. The presence of a type 1 AGN is not supported by the observed SED as the hot dust emission in luminous high- $z$  QSOs contributes at wavelengths above rest-frame  $1\mu\text{m}$ , and an additional ad-hoc red source would be required to explain the observed flux excess at  $0.66$  and  $0.86\mu\text{m}$ . Additional deep MIRI imaging covering the rest-frame near-IR are needed to confirm the flux detection at  $10\mu\text{m}$  with higher significance, and to discriminate between the different hot dust emission in the extreme starburst and AGN scenarios with MIRI imaging at longer wavelengths.

**Key words.** galaxies: high-redshift – galaxies: individual: GN-z11 – galaxies: starburst – infrared: galaxies

## 1. Introduction

GN-z11 was firstly identified as a  $z > 10$  Lyman-break galaxy candidate using multi-colour imaging from HST CANDELS (Bouwens et al. 2010). A subsequent HST grism spectroscopic analysis revealed a flux dropout identified as the Lyman break by Oesch et al. (2016), yielding a redshift value of  $z \sim 11.09$ . The arrival of JWST has enabled the study of this object at optical and near-IR wavelength with unprecedented sensitivity and spatial resolution. In particular, the JWST Advanced Deep Extragalactic Survey (JADES; Eisenstein et al. 2023) derived a redshift  $z = 10.6034 \pm 0.0013$  using multiple emission lines based on NIRSpect/Prism data (Bunker et al. 2023). Besides, this NIRSpect spectrum has revealed the presence of a high ionization field ( $\log(U) \sim -2.3$ ) and a super-solar abundance of nitrogen ( $[\text{N}/\text{O}] > -0.49$ ). Recent JWST NIRCам imaging have shown that GN-z11 UV-emission can be depicted as the combination of a point-source and an extended component, showing an extremely compact morphology (combined  $R_{\text{eff}}$  of  $64 \pm 20$  pc) and large UV luminosity ( $M_{\text{UV}} = -21.58 \pm 0.02$ ; Tacchella et al. 2023a).

JWST multi-wavelength photometry and spectroscopy result on star-formation rates (SFRs) and stellar masses between  $20 - 30 M_\odot \text{ yr}^{-1}$  and  $\log(M_*/M_\odot) = 8.7 - 9.1$  (Bunker et al. 2023; Tacchella et al. 2023a), respectively. The relatively large stellar mass at only  $\sim 420$  Myr after the Big Bang suggests a

rapid build-up of stellar mass, which has been also found in other JWST spectroscopically confirmed galaxies at  $z > 10$  (Robertson et al. 2023; Curtis-Lake et al. 2023; Carniani et al. 2024; Naidu et al. 2025). The exceptionally UV brightness of this object allows its observation using ground-based facilities. Recent NOEMA observations (Fudamoto et al. 2024) have determined upper limits on the [C II]  $158\mu\text{m}$  emission and  $160\mu\text{m}$  dust continuum yielding  $\log(M_{\text{mol, [C II]}}/M_\odot) < 9.3$  and  $\log(M_{\text{dust}}/M_\odot) < 6.9$ . These results are consistent with a negligible dust attenuation and the blue colour ( $\beta = -2.4$ ) obtained from a SED-fitting analysis (Tacchella et al. 2023a).

Using medium resolution ( $R = 1000$ ) NIRSpect observations, Maiolino et al. (2024) present several pieces of evidences favouring the scenario where GN-z11 is a Narrow Line Seyfert 1 (NLS1) with a black hole mass of  $\log(M_{\text{BH}}/M_\odot) = 6.2 \pm 0.3$  accreting material at a large Eddington ratio ( $\lambda_{\text{Edd}} = L_{\text{bol}}/L_{\text{Edd}} \sim 5.5$ ). The authors found the presence of high-excitation ( $> 60$  eV) emission lines in the UV spectrum (e.g., [Ne IV]  $\lambda 2422, 2244$ ) which are usually present in AGN (Le Fèvre et al. 2019; Terao et al. 2022). In addition, the high critical density ( $n_{\text{e, N III}} > 10^5 \text{ cm}^{-3}$ ) and broad ( $\text{FWHM}_{\text{He II } \lambda 1640} \sim 1200 \text{ km s}^{-1}$ ) emission lines revealed by the NIRSpect data are typically observed in the broad-line regions (BLRs) of AGNs. The presence of an outflow powered by this AGN was also inferred from a high equivalent width and blue-

shifted absorption of a C iv doublet (Maiolino et al. 2024). The small and compact size of the BLR linked to this potential AGN would justify the large [N/O] of GN-z11 through a very rapid chemical enrichment driven by only a few SNe. In addition, a blue bump has been observed in the UV using NIR-Spec/Prism data (Ji et al. 2025). This bump has been proposed to be linked to dense Fe II-emitting clouds ( $n_{\text{H}} \sim 10^{11} \text{ cm}^{-3}$ ) infalling at  $\sim 3000 \text{ km s}^{-1}$ . GN-z11 hosting an AGN and the presence of tentative nearby companions detected as Ly $\alpha$  blobs make this region a possible candidate for being the core of a protocluster with a halo mass  $\log(M_{\text{halo}}/M_{\odot}) = 10.4$  (Scholtz et al. 2024).

However, alternative scenarios have been proposed to justify the overabundance of nitrogen in GN-z11, such as the presence of Wolf-Rayet (WR) stars (Senchyna et al. 2024) or super-massive stars in a proto-globular cluster (Charbonnel et al. 2023). In addition, Bhatt et al. (2024) show that cosmological simulations do not support a super-Eddington type 1 AGN scenario for GN-z11. The analysis of the H $\alpha$  and [O III] 5008 lines observed with MIRI/MRS carried out by Álvarez-Márquez et al. (2025) does not support the presence of an accreting black hole dominating the optical emission lines and continuum in GN-z11. In the case H $\alpha$  and the optical continuum are entirely produced by the BLR, the accretion rate and bolometric luminosity would be unrealistically large, but the black hole mass would be smaller ( $\log(M_{\text{BH}}/M_{\odot}) = 5.77$ ) than the one inferred by UV lines in Maiolino et al. (2024). In addition,  $L_{\text{H}\alpha}/L_{\text{X},2-10 \text{ keV}}$  and  $L_{\text{H}\alpha}/L_{5100}$  were found inconsistent with the low- $z$  AGN relations (Ho et al. 2001a; Greene & Ho 2005).

To date, the majority of studies on GN-z11 have relied on UV data. However, we still lack important information at optical wavelengths, where the contribution from the young stellar populations begin to decrease and those from more evolved populations, or redder emission sources, is expected to become more prominent. So far, only Álvarez-Márquez et al. (2025) has covered the optical wavelengths, using MIRI/MRS data, detecting [O III] 5008 and H $\alpha$  emission consistent with star formation. This paper presents the first mid-infrared F560W and F770W images of GN-z11, tracing its optical emission at rest-frame 0.5 and  $0.66 \mu\text{m}$ , respectively. We combine these images with ancillary spectro-photometric data covering from the rest- 0.15 up to  $0.86 \mu\text{m}$  to trace the UV to optical emission continuum of this object and extend the analysis of its SED up to optical-red wavelengths. The paper is organised as follows. Section 2 presents the new JWST/MIRI images used in this work and the ancillary JWST data. Section 3 describes the continuum-only extraction and SED fitting analysis carried out. The main results drawn from the analysis and their discussion are presented in Section 4. Finally, Section 5 summarises the main conclusions and results of this work. Throughout this paper we assume a Chabrier initial mass function (IMF, Chabrier 2003) and a flat  $\Lambda$ CDM cosmology, with  $\Omega_{\text{m}} = 0.31$  and  $H_0 = 67.7 \text{ km s}^{-1} \text{ Mpc}^{-1}$  (Planck Collaboration et al. 2020). For this cosmology, 1 arcsec corresponds to 4.08 kpc at  $z = 10.603$  while the luminosity distance is  $D_{\text{L}} = 113.4 \text{ Gpc}$ .

## 2. Observation and data processing

### 2.1. JWST/MIRI data and calibration

The JWST images of the GN-z11 were obtained on December 8, 2023 using the MIRI imager (MIRIM, Bouchet et al. 2015) with the F560W and F770W filters, as part of the cycle 1 JWST program ID 1264 (PI: Colina, L.). The observations were performed with the FASTR1 read-out mode in a five- and

seven-point dither, for the F560W and F770W respectively, with a medium-size cycling pattern. The total on-source integration time corresponds to 2012 and 5866 seconds distributed in 10 and 28 integrations, respectively for F560W and F770W.

These MIRIM images have been calibrated using the JWST pipeline (v1.12.0) with the context 1170 of the Calibration Reference Data System (CRDS). This CRDS includes photometric calibrations considering the temporal evolution and aperture corrections taking into account the point-spread function (PSF) cruciform. In addition to the general procedure, further steps have been applied to correct for striping and background gradients (see more details in Álvarez-Márquez et al. 2023; Pérez-González et al. 2024a; Östlin et al. 2025). The process concluded with the creation of final MIRI images dithered to a pixel scale of  $0.06''/\text{pixel}$ . According to the JWST documentation<sup>1</sup>, the spatial resolutions for these images are FWHM =  $0.207''$  and  $0.269''$ , for the F560W and F770W, respectively. Following this calibration, we obtain a  $3\sigma$  depth of 26.67 and 26.96 mag for F560W and F770W for unresolved point sources, respectively.

### 2.2. Ancillary JWST data: photometry and spectroscopy

In addition to the MIRI F560W and F770W images, this work takes advantage of the large amount of high-quality JWST data available for GN-z11 covering a large wavelength range. To trace the near-IR counterpart of this galaxy, we use the F1000W and F2100W images available from the MEOw survey (PID 5407, PI: G. Leung) which covers up to rest-frame  $\sim 2 \mu\text{m}$ . These observations were performed with a total exposure time of 722 and 3086 s reaching a  $3\sigma$  level of 230 nJy and  $2.6 \mu\text{Jy}$  (i.e., 25.5 and 22.9 mag, respectively). In addition, the 11 NIRCcam images available from JADES DR3 (Eisenstein et al. 2023; Bunker et al. 2024; D'Eugenio et al. 2025), with median  $3\sigma$  depths of  $\sim 5 \text{ nJy}$  (i.e.,  $\sim 29.6 \text{ mag}$ , D'Eugenio et al. 2025), are used to trace the UV continuum from rest-frame 0.1 up to  $\sim 0.4 \mu\text{m}$ . We performed aperture photometry as explained in Sect. 3.1 for all the NIRCcam and MIRI images available.


We have also incorporated JWST ancillary spectroscopic data for this object. In particular, we consider the fluxes from the UV lines detected on the NIRSpec/Prism data presented in Bunker et al. (2023) after a re-normalization to match the NIRCcam photometry (see Sect. 3.1 and App. A). In addition to the UV lines from NIRSpec, we also use the flux from the optical lines present in the MRS data and extracted by Álvarez-Márquez et al. (2025). The MRS spectra contain the H $\beta$  + [O III] and H $\alpha$  lines which fall into the MIRI/F560W and F770W wavelength ranges, respectively. We do not apply any normalization as the MRS line fluxes were corrected for aperture losses using the MRS PSFs (Argyriou et al. 2023).

## 3. Analysis

### 3.1. Aperture photometry and NIRSpec renormalization

We extract the GN-z11 photometry from the available NIRCcam and MIRI images considering circular apertures, centred on the F560W emission peak. The size of these apertures (i.e.,  $r = 0.5''$ ) has been chosen to fully encompass both the UV and the optical emission. Before extracting the fluxes from NIRCcam, we matched the PSF of these images to the spatial resolution of MIRI/F560W. At MIRI resolution, we con-

<sup>1</sup> <https://jwst-docs.stsci.edu/jwst-mid-infrared-instrument/miri-performance/miri-point-spread-functions>



**Fig. 1.** GN-z11 images in the NIRCam/F200W, MIRI/F560W, MIRI/F770W and MIRI/F1000W filters. The NIRCam/F200W image was convolved to match the MIRI/F560W resolution (see Sect. 3.1). Dashed circles represent the 0.5'' aperture used for the photometry extraction. White circles on the lower right corners of each panel represent the FWHM for each filter.

**Table 1.** NIRCam and MIRI photometry

Filter	units	Obs.	Cont. level
NIRCam/F115W	nJy	$31 \pm 9$	-
NIRCam/F150W	nJy	$118 \pm 9$	-
NIRCam/F182M	nJy	$177 \pm 11$	$154 \pm 12$
NIRCam/F200W	nJy	$201 \pm 7$	$185 \pm 8$
NIRCam/F210M	nJy	$175 \pm 14$	$164 \pm 14$
NIRCam/F277W	nJy	$161 \pm 3$	$161 \pm 3$
NIRCam/F335M	nJy	$130 \pm 7$	$123 \pm 7$
NIRCam/F356W	nJy	$153 \pm 3$	$150 \pm 3$
NIRCam/F410M	nJy	$133 \pm 6$	$129 \pm 6$
NIRCam/F444W	nJy	$163 \pm 4$	$140 \pm 4$
MIRIM/F560W	nJy	$368 \pm 26$	$156 \pm 31$
MIRIM/F770W	nJy	$274 \pm 20$	$212 \pm 22$
MIRIM/F1000W	nJy	$249 \pm 77$	$237 \pm 77^\dagger$
MIRIM/F2100W	$\mu$ Jy	$<2.6$	$<2.6$

**Notes.** Third and fourth columns display the observed and continuum level, i.e. the measured photometry and the continuum fluxes after subtracting the emission line contribution (see Sect 3.2), respectively. We adopt  $3\sigma$  upper limits for F2100W.  $^\dagger$ : The continuum level for F1000W is derived assuming a  $[\text{S II}] 9071,9533/\text{H}\alpha = 0.17$  (see Sect. 4.1).

sider GN-z11 as a point-like source and therefore we applied the aperture corrections derived from the empirical PSFs presented in Libralato et al. (2024). At  $r = 0.5''$ , these corrections are 1.29, 1.33, 1.54 for MIRI/F560W and the NIRCam images, MIRI/F770W and MIRI/F2100W, respectively. To optimise the S/N on the F1000W detection we have assumed an aperture of  $r = 0.3''$ , applying an aperture correction factor of 1.71. The associated uncertainty of each photometric measurement is defined as the standard deviation of the local background around GN-z11 ( $1'' < r < 3''$ ), masking all the additional objects present in the field-of-view (FoV). In this work, we have re-scaled the MIRI background noise by a factor of 2.24 to take into account the correlated noise induced when drizzling the individual observations (see Östlin et al. 2025, for details). We adopt a  $3\sigma$  upper limit for the F2100W image, which does not present significant detection. Table 1 presents the flux values measured from the NIRCam and MIRI images. The S/N is generally high ( $>10$ ) for

all photometric values, with the exception of F1000W, in which GN-z11 is detected at a  $\sim 3\sigma$  level.

We also use the NIRCam fluxes to renormalise the NIRSpec spectra, correcting from potential slit losses. In brief, we matched the NIRSpec/Prism spectra to the NIRCam photometry multiplying by a factor of 1.325. A detailed description of the process is presented in Appendix A.

### 3.2. Continuum-only fluxes and EWs

Since the optical emission lines detected in the MRS (i.e.,  $[\text{O III}] 5008$ ,  $\text{H}\alpha$ ) fall within the F560W and F770W filters, respectively, their contribution can be subtracted from the broadband fluxes, allowing us to estimate the underlying continuum at 5.6 and 7.7  $\mu\text{m}$ , and derive the equivalent width (EW) of these emission lines.

In this work, we have used the  $[\text{O III}] 5008$  and  $\text{H}\alpha$  fluxes presented by Álvarez-Márquez et al. (2025) and the transmission curves for the F560W and F770W filters<sup>2</sup>. For accuracy, we have included the  $[\text{O III}] 4960$  and  $\text{H}\beta$  contributions using  $[\text{O III}] 5008/[\text{O III}] 4960 = 3$  and  $\text{H}\alpha/\text{H}\beta = 2.86$ , considering case B and with negligible extinction (Bunker et al. 2023; Álvarez-Márquez et al. 2025). Assuming  $\log(U) = -3$  and  $0.2 Z_\odot$ , the potential contributions from  $[\text{N II}]\lambda\lambda 6549,6585$  and  $[\text{S II}]\lambda\lambda 6718,6732$  to the F770W flux are minimal ( $<5\%$ , Curti et al. 2020).

In addition, we have extended the derivation of the continuum levels to UV wavelengths by using the NIRCam fluxes and the NIRSpec/Prism re-normalised emission lines presented in Bunker et al. (2023). Finally, we estimate the continuum in MIRI/F1000W based on the  $[\text{S II}] 9071,9533/\text{H}\alpha = 0.17$  ratio derived from Kewley & Dopita (2002), assuming  $0.2 Z_\odot$  and  $\log(U) = -3$ . Although F1000W is slightly narrower than F770W, the  $[\text{S II}] 9071,9533$  flux ( $\sim 12$  nJy) would contribute only up to  $\sim 5\%$  of the total F1000W photometry. The computed continuum-only fluxes for the NIRCam and MIRI filters are listed in Table 1.

Álvarez-Márquez et al. (2025) estimate an upper limit of a possible broad component in  $\text{H}\alpha$  of  $2 - 3 \times 10^{-18} \text{ erg s}^{-1} \text{ cm}^{-2}$ . If we consider this broad component, the F560W and F770W

<sup>2</sup> Transmission curves are obtained from the SVO Filter Profile Service



continuum levels would decrease by  $6 - 9$  nJy and  $18 - 27$  nJy, respectively, assuming a ratio  $H\alpha/H\beta = 2.86$  for the broad component. Based on the MIRI/F560W and F770W continuum-only fluxes, we derive rest-frame  $EW([O\text{ III}] 5008) = 786 \pm 81 \text{ \AA}$  and  $EW(H\alpha) = 547 \pm 72 \text{ \AA}$ , considering only the narrow line contribution on  $H\alpha$ .

### 3.3. SED fitting

In order to investigate the nature of GN-z11, we have analysed its SED based on the available spectro-photometric data. In particular, the SED constructed with the NIRCам+MIRI photometry and the NIRSpec+MRS spectroscopy was fitted with two codes, CIGALE v2025.0 (Burgarella et al. 2005; Noll et al. 2009; Boquien et al. 2019) and SYNTHESIZER-AGN (Pérez-González et al. 2003, 2008). Since the F1000W flux shows a lower significance compared to the other MIRI values, we exclude F1000W (and the upper limit for F2100W) from the SED fitting to prevent biasing the results. However, we will extract the modelled fluxes at these filters to evaluate their consistency.

The SED fitting from CIGALE makes use of the UV+optical emission lines and the MIRI fluxes. In addition, as the current version of CIGALE does not use spectroscopic data, we have constructed pseudo-continuum fluxes by integrating the re-normalised NIRSpec/Prism spectrum in  $750 \text{ \AA}$  bins. The emission lines were assigned separate bins to avoid contaminating the pseudo-continuum values. These UV pseudo-continuum, which isolate the effect of the emission lines and trace the galaxy continuum emission, are considered during the SED fitting along with the MIRI fluxes. In our CIGALE model, the stellar component is fitted using the stellar population templates from the Binary Population and Spectral Synthesis library (BPASSv2.2, Stanway & Eldridge 2018) with  $Z = 0.2 Z_{\odot}$  (following Álvarez-Márquez et al. 2025) and a Chabrier initial mass function (IMF, Chabrier 2003) with an upper stellar mass limit of  $100 M_{\odot}$ . Nebular emission is included via Cloudy models (Ferland et al. 2017), assuming  $Z = 0.2 Z_{\odot}$ , an electron density of  $1000 \text{ cm}^{-3}$ , and an ionisation parameter of  $\log(U) = -2.3$  (following Bunker et al. 2023). Dust extinction is assumed to follow the attenuation curve from Calzetti et al. (2000) with  $A_V < 0.3$  mag, constrained by the Balmer line ratios derived from previous NIRSpec and MRS analysis (Bunker et al. 2023; Álvarez-Márquez et al. 2025). We have also considered the potential presence of a non-stellar component (i.e., type 1 or 2 AGN) by adding an AGN component modelled as described by Fritz et al. (2006). This model assumes a broken power-law point source as the ionization source and the presence of a smooth dusty torus. Since our dataset covers up to rest-frame  $0.86 \mu\text{m}$ , we cannot fully constrain the full model parameter space. Therefore, for simplicity, in this work we fixed an opening angle of  $60^\circ$  and a  $R_{\text{max}}/R_{\text{min}} = 10$ , following the results found for high- $z$  luminous and massive QSOs (Bosman et al. 2025) while we allow the radial and angular indices range between  $\beta = [-1, -0.5]$  and  $\gamma = [0, 6]$ , respectively. We tested both type 1 and type 2 scenarios by forcing the angle of the line of sight to the AGN polar axis be  $\theta < 20^\circ$  and  $> 70^\circ$ , respectively.

For the SED fitting carried out with SYNTHESIZER-AGN, we used the re-normalised NIRSpec/Prism spectrum, the MRS optical lines and the MIRI fluxes. Throughout the fitting, we consider the stellar population models described by Bruzual & Charlot (2003), hereafter BC03, assuming a Chabrier (2003) IMF with stellar masses between  $0.1$  and  $100 M_{\odot}$ . The nebular continuum and emission lines are modelled using Cloudy c23.01 (Gunasekera et al. 2023) predictions for a variety of ion-

izing SEDs (parametrised by a range of effective temperatures ( $8000 - 120000 \text{ K}$ ) and ionizing photon flux ( $10^{37} - 10^{60} \text{ s}^{-1}$ )), and a variety of electron densities ( $\log(n_e/\text{cm}^{-3}) = 2 - 4$ ) and metallicities ( $0.02 - 0.2 Z_{\odot}$ , with solar abundances for individual elements). The dust extinction is described by the Calzetti et al. (2000) attenuation law where we constrained the  $A_V$  to be lower than  $0.30$  mag, as for the CIGALE fitting. The AGN emission is included following the QSO composite spectrum created by combining the UV+optical and near-IR templates from Selsing et al. (2016) (up to rest-  $1.13 \mu\text{m}$ ) and Glikman et al. (2006) (rest-  $1.13 - 3.5 \mu\text{m}$ ), respectively. This empirical template is added to the fitting procedure to represent average accretion disk and dust emission in luminous type 1 AGNs. Additionally, we also consider the presence, without energy balance, of an additional dusty torus to reproduce the observed red SED shape. This component is defined by using self-consistent AGN torus model from Polletta et al. (2006, 2007), which is a highly obscured ( $A_V = 4$ ) QSO with UV emission coming from scattered light and IR emission coming from hot/warm dust.

To quantify the dependence of the physical parameters when using different stellar population models, we compared the CIGALE results considering both the BC03 and BPASSv2.2 stellar templates. The BPASSv2.2 models predicts stronger UV emission due to binary evolution, allowing stellar populations older than  $10 \text{ Myr}$  to still produce significant ionizing flux. This yields typically older ages, higher SFRs, and lower stellar masses, while BC03 favours younger and more massive solutions for GN-z11. However, these differences are  $< 10 - 15\%$  and, in all the cases, smaller than those introduced by assuming different star-formation histories (SFHs). We have also examined the potential effect of adopting a different extinction law during the SED fitting. For this comparison, we have tested the Charlot & Fall (2000) attenuation curve, in which the young stars and the nebular emission suffer an additional attenuation compared with the older stars ( $> 10 \text{ Myr}$ ). We find that, in our case, the SED fitting using Charlot & Fall (2000) lead to stellar masses only  $\sim 0.1$  dex smaller than those assuming Calzetti et al. (2000) while variations in other physical parameters (e.g.,  $\text{SFR}_{100 \text{ Myr}}, M_{\odot}$ ) are lower than  $10\%$ . We therefore argue that neither the choice of the stellar population templates or the attenuation curve affects any conclusion drawn from our analysis.


In an effort to discover the nature of GN-z11, in Sections 4.2 and 4.3 we tested different SFHs and the possible presence of an AGN for both SED fitting codes. Differences in the SFHs and AGN modelling are presented and discussed in the following sections.

## 4. Results and discussion

### 4.1. Continuum-only fluxes and colour-colour diagrams

The large spectro-photometric dataset available for GN-z11 combining NIRCам, NIRSpec, MIRI and MRS allows us to trace both the nebular lines and the continuum from rest- UV up to  $0.86 \mu\text{m}$ . By removing the nebular line contributions from the aperture photometry, we obtain continuum-only fluxes, which allow us to trace the GN-z11 underlying continuum emission (see Sect. 3.2).

The continuum-only results show (see Table 1) how, although the MIRI/F560W value agree within the uncertainties with the continuum level expected from previous SED fittings based on UV-only data (Jiang et al. 2021; Tacchella et al. 2023a), our F770W photometric value exceeds their predicted flux. In particular, once we take into account the difference with the pho-




**Fig. 2.** Continuum-only colour-colour diagrams. Black squares represent the GN-z11 colours based on the NIRCам and MIRI photometry after the subtraction of the emission lines contribution measured with NIRSpect and MRS (i.e., continuum-only colours) presented in Table 1. Continuum (dashed) lines display the continuum-only colours derived considering a stellar population colour-coded by their age assuming an instantaneous (continuum) SFH. These colours include the contribution from the stellar continuum calculated using the stellar models from BPASSv2.2 and its associated nebular continuum derived with Cloudy (see Sect. 3.2). Thin and thick lines represent the 0.05 and 0.2  $Z_{\odot}$  tracks, respectively. Orange, green and purple diamonds show the derived continuum-only colours from the composed  $z \sim 1$  QSO (Glikman et al. 2006; Selsing et al. 2016, see Sect. 3.3), the Seyfert 2 Mrk3 (Spinelli et al. 2006) and a  $z = 7.1$  QSO (Bosman et al. 2024, 2025). Red, blue and brown diamonds represent the values derived from a stacked spectra of  $\sim 150$  LRDs (Pérez-González et al. 2024b, Pérez-González et al. in prep.), the COSMOS-Webb median-stacked LRD (Akins et al. 2025) and the unique LRD known as ‘The Cliff’ (de Graaff et al. 2025), respectively. In the left and middle panels the brown diamond lie outside the colour range displayed. Black arrows in the lower right corners represent the vector magnitude for  $A_V = 0.3$  mag.

tometry from Tacchella et al. (2023a) (see App. A), our F770W continuum level ( $212 \pm 20$  nJy) is  $>70\%$  larger ( $>5\sigma$ ) than the SED fitting expected value when using UV-only photometry ( $\sim 120$  nJy). This excess at  $\sim 0.66 \mu\text{m}$  rest-frame could be hinting at the presence of a non-negligible old stellar population or a red component that would be strongly emitting at optical-red and near-IR wavelengths.

A first step to investigate the nature of GN-z11, and this possible red excess, is to use classic colour-colour diagrams covering from rest-frame UV ( $0.17 \mu\text{m}$ ) to optical-red ( $0.66 - 0.86 \mu\text{m}$ ) wavelengths. Figure 2 displays the diagrams for the UV (F200W – F444W), optical (F444W – F560W) and optical-red (F560W – F770W and F770W – F1000W) colours derived from the continuum-only fluxes. In this figure we compare these continuum-only colours to those expected from stellar populations models and those from intrinsically red objects (e.g., QSO, AGNs and LRDs). In particular, we have considered the stellar populations templates from BPASSv2.2 for continuum and instantaneous SFHs and a range of stellar ages limited to the age of the Universe at  $z = 10.6$  (i.e., 440 Myr). Their associated nebular continua are also included using the corresponding Cloudy models for those ages and SFHs (Ferland et al. 2017). For completeness, we displayed the age evolution tracks for metallicities 0.05 and 0.2  $Z_{\odot}$ . Green diamonds represent the colours of the nuclear ultraviolet-optical spectrum of Mrk 3. This low- $z$  galaxy is classified as a heavily obscured Seyfert 2 nucleus (Guainazzi et al. 2016), with a hidden BLR identified by the presence of broad Balmer lines in polarised light (Miller & Goodrich 1990). Its spectrum, obtained with the HST/STIS spectrograph using an aperture size of  $0.2''$  (i.e., 50 pc at the distance of Mrk3, Spinelli et al. 2006), represents the SED of a luminous type 2 AGN while minimizing the potential contamination due to the stellar light that could be affecting larger apertures. Moreover, its line ratios ( $\text{O3H}\gamma = -0.1$ ,  $\text{Ne3O2} = -0.2$  and  $\text{O32} = 0.9$ ), agree with those found in high- $z$  AGNs (Mazzolari et al. 2024,

2025). Orange diamonds correspond to the colours measured for the template of luminous blue QSOs at intermediate redshifts ( $1 < z < 2$ ) constructed from the simultaneous ultraviolet to near-infrared spectroscopy (Glikman et al. 2006; Selsing et al. 2016). Purple diamonds represent the colours derived for the NIRSpect+MRS spectrum of the  $z = 7.1$  QSO J1120+0641, whose BLR and dust emission has been found to be consistent with those of lower-redshift quasars (Bosman et al. 2024, 2025). We have filled the gap between the NIRSpect and MRS data (i.e.,  $3.12 - 4.9 \mu\text{m}$ ) following its best-fit SED shape ( $f_{\lambda} \propto \lambda^{-0.25}$ ; Bosman et al. 2025). Brown diamonds show the colour derived from the spectra of the singular little red dot (LRD) presented by de Graaff et al. (2025), known as ‘The Cliff’. This object presents the strongest Balmer break among LRDs and shows  $\text{H}\alpha$  absorption, which are explained under the ‘black hole star’ (BH\*) scenario in which dense gas surrounds a powerful ionising source (see also Naidu et al. 2025 and Taylor et al. 2025). Finally, the red and blue diamonds display the colour derived for an updated stacked spectra from Pérez-González et al. (2024b), combining  $>150$  nearby and high- $z$  LRDs, and the COSMOS-Webb median-stacked LRD from Akins et al. (2025), respectively. The spectra of all these sources together with the GN-z11 JWST photometry are presented in Figure 3.

The continuum-only fluxes show an SED blue in the rest-UV (F200W – F444W =  $-0.30 \pm 0.05$  mag) with a flattening in  $f_{\nu}$  up to rest-frame  $0.5 \mu\text{m}$  (F444W – F560W =  $0.12 \pm 0.21$  mag). Beyond  $[\text{O III}] 5008$ , the SED appears to rise, with colours F560W – F770W =  $0.33 \pm 0.24$  mag and F770W – F1000W =  $0.18 \pm 0.40$  mag ( $0.26 \pm 0.24$  and  $0.33 \pm 0.40$  mag, respectively, if we consider the impact of the potential broad hydrogen Balmer lines components, see Sect. 3.2). Although the rest-UV and optical colours can be reproduced by a young ( $\sim 10$  Myr) stellar burst, it fails to explain the optical-red colours (e.g., F560W – F770W and F770W – F1000W, see Figure 2). On the other hand, the Seyfert



**Fig. 3.** NIRCам and MIRI photometry compared with different red SED objects. Grey squares and black circles represent the NIRCам+MIRI observed and continuum-only (see Sect. 4.1) fluxes, respectively. Coloured lines represent the SEDs for different AGNs types and high- $z$  objects with optical-red colours presented in Sect. 4.1. Red line represents the stacked spectra for  $\sim 150$  local and high- $z$  LRDs (Pérez-González et al. 2024b, Pérez-González et al. in prep.). Brown line displays the spectra from the exotic  $z \sim 3.5$  LRD ‘The Cliff’ while brown diamonds represent its re-normalised fluxes from de Graaff et al. (2025) at rest- 0.9 and 1.9  $\mu\text{m}$ . Blue line shows the COSMOS-Webb median-stacked LRD SED model presented in Akins et al. (2025). Orange and purple lines show the  $z \sim 1$  QSO composite spectrum, created by combining the UV+optical and near-IR spectral templates from Selsing et al. (2016) and Glikman et al. (2006), and the NIRSpect+MRS spectrum for the  $z = 7.1$  QSO J1120+0641 (Bosman et al. 2024, 2025), respectively. The HST/STIS spectrum for the Sy2 Mrk3 presented by Spinelli et al. (2006) is displayed as a green line. All observed spectra have been redshifted to  $z = 10.6$  and normalised at 4800 Å.

2 spectrum can produce the optical-red continuum-only colours observed with MIRI but fails to explain the UV-blue colour F200W – F444W. Finally, the LRD stacked and ‘The Cliff’ spectra are too red, specially beyond 0.4  $\mu\text{m}$ , and they are only marginally consistent with F770W – F1000W due to the large uncertainty in the F1000W flux. These results, based on the traditional colour-colour diagrams, suggest that while the UV emission from GN-z11 can be produced by a young star-formation, the presence of an extra red source is needed to explain its rest-frame optical-red colours. Note however that in these colour-colour diagrams, the stellar plus nebular continuum models do not contain any contribution due to small amounts of hot dust emitting at optical-red wavelengths. Likewise, mixed stellar populations, that would produce optical colours different from those of (single age) instantaneous bursts are not represented.

In the following sections we perform a more detailed modelling using different SED fitting codes with different scenarios (e.g., SFHs, AGN; see Sect. 3.3) to investigate the origin behind these UV, optical and optical-red colours. Besides, SED fitting will allow us to model both the continuum and the emission lines

simultaneously, providing a more complete view of the physical mechanisms driving the emission in GN-z11.

#### 4.2. SF-only SED modelling

The colour-colour analysis carried out suggests an increase in the continuum emission beyond 0.66  $\mu\text{m}$ . In this section we perform an SED analysis including both the emission lines and the photometric points to dig into the nature of this enhancement.

As a first approach for the CIGALE SED fitting, we considered that GN-z11 can be modelled by a stellar population following a tau-decaying SFH with  $\tau$  varying in the [1, 5000 Myr] range, and with ages up to 440 Myr (i.e., age of Universe at  $z = 10.603$ ). Figure 4 shows the best-fit SED obtained from these adopted model parameters and the comparison with the observed NIRCам+MIRI fluxes. The Bayesian analysis results in an almost constant SFH ( $\tau = 2500 \pm 1500$  Myr) and a stellar population of  $M_* = (5.6 \pm 0.6) \times 10^8 M_\odot$  and an age of  $t = 21 \pm 3$  Myr. Although for this model, we obtain H $\alpha$  and [O III] 5008 that are in agreement with those measured with the MRS, the modelled F770W flux (i.e.,  $176 \pm 1$  nJy) is underestimated by  $\sim 45\%$ ,  $5\sigma$

**Table 2.** Modelled fluxes derived from the SED fitting


Line/Filter	Observed	Single pop	Mixed pop.	Mix.+ty1 (CIG.)	Mix.+AGN (SYN.)	Mix.+ty2 (CIG.)
H $\beta$	<44	22 $\pm$ 1	24 $\pm$ 0	22 $\pm$ 1	26 $\pm$ 4	22 $\pm$ 1
[O III] 4960	<44	44 $\pm$ 1	52 $\pm$ 1	44 $\pm$ 1	43 $\pm$ 10	46 $\pm$ 1
[O III] 5008	136 $\pm$ 14	132 $\pm$ 4	156 $\pm$ 4	133 $\pm$ 3	128 $\pm$ 30	136 $\pm$ 1
H $\alpha$	68 $\pm$ 9	63 $\pm$ 2	70 $\pm$ 2	61 $\pm$ 1	63 $\pm$ 10 <sup>†</sup>	63 $\pm$ 1
F560W	368 $\pm$ 26	326 $\pm$ 4	382 $\pm$ 7	335 $\pm$ 5	357 $\pm$ 50	360 $\pm$ 2
F770W	274 $\pm$ 20	176 $\pm$ 1	202 $\pm$ 5	185 $\pm$ 5	261 $\pm$ 30	254 $\pm$ 5
F1000W	249 $\pm$ 77	111 $\pm$ 3	130 $\pm$ 6	126 $\pm$ 6	274 $\pm$ 40	285 $\pm$ 11

**Notes.** CIG. and SYN. acronyms stand for CIGALE and SYNTHESIZER-AGN (See Sects. 3.3, 4.2 and 4.3). Units are in  $10^{-19}$  erg s $^{-1}$  cm $^{-2}$  and nJy for the lines and photometric points, respectively. <sup>†</sup>: We derive a  $1 \times 10^{-18}$  erg s $^{-1}$  cm $^{-2}$  flux for the broad component in H $\alpha$ .

**Table 3.** Physical parameters from SED fitting

	Observed	Single pop	Mixed pop.	Mix.+ty1 (CIG.)	Mix.+AGN (SYN.)	Mix.+ty2 (CIG.)
sSFR [Gyr $^{-1}$ ]		53 $\pm$ 6	33 $\pm$ 12	83 $\pm$ 20	85 $\pm$ 19	56 $\pm$ 4
$M_*$ [ $10^8 M_\odot$ ]		5.6 $\pm$ 0.6	14.7 $\pm$ 1.2	3.7 $\pm$ 0.8	2.1 $\pm$ 0.9	5.3 $\pm$ 0.3
$M_*^{\text{young}}$ [ $10^8 M_\odot$ ]		-	1.8 $\pm$ 0.2	1.7 $\pm$ 0.2	0.5 $\pm$ 0.2	1.8 $\pm$ 0.1
$t_{\text{young}}$ [Myr]		-	4 $\pm$ 1	6 $\pm$ 1	0.32 $\pm$ 0.15	6 $\pm$ 1
$t_{\text{old}}$ [Myr]		21 $\pm$ 3	63 $\pm$ 23	39 $\pm$ 21	6 $\pm$ 2	20 $\pm$ 2
SFR $_{10\text{Myr}}$ [ $M_\odot$ yr $^{-1}$ ]	24 $\pm$ 3 <sup>†</sup>	30 $\pm$ 2	18 $\pm$ 3	18 $\pm$ 3	18 $\pm$ 5	18 $\pm$ 2
log( $L_{\text{AGN}}$ ) [erg s $^{-1}$ ]		-	-	44.57 $\pm$ 0.04	44.6 $\pm$ 0.5	44.77 $\pm$ 0.02
$A_V$ [mag]	<0.2 <sup>†</sup>	0.15 $\pm$ 0.01	0.13 $\pm$ 0.02	0.04 $\pm$ 0.01	0.10 $\pm$ 0.03	0.03 $\pm$ 0.01
$Z^{\dagger\dagger}$ [ $Z_\odot$ ]	0.17 $\pm$ 0.03 <sup>†</sup>	-	-	-	0.16 $\pm$ 0.03	-
EW(H $\alpha$ ) [Å]	547 $\pm$ 72	983 $\pm$ 55	932 $\pm$ 24	870 $\pm$ 38	540 $\pm$ 110	572 $\pm$ 17
EW(H $\beta$ ) [Å]	140 $\pm$ 19	175 $\pm$ 10	169 $\pm$ 5	166 $\pm$ 6	170 $\pm$ 40	146 $\pm$ 3
EW([O III] 5008) [Å]	786 $\pm$ 81	1144 $\pm$ 63	1159 $\pm$ 35	1080 $\pm$ 41	840 $\pm$ 230	932 $\pm$ 19

**Notes.** <sup>†</sup>: Values derived from MIRI/MRS data presented in Álvarez-Márquez et al. (2025). <sup>††</sup>: Fixed metallicity value was adopted during the CIGALE fits (see Sect. 3.3).



**Fig. 4.** CIGALE SED models derived from the single and mixed stellar population scenarios described in Sect. 4.2. Blue (orange) line and squares represent the SED models and the expected fluxes for NIRCам/MIRI filters derived using a single  $\tau$ -decaying (mixed) stellar population. Black line and triangles represent the NIRSpec/Prism spectra and the NIRCам+MIRI photometry, respectively. The blue and orange lines have been convolved to match the NIRSpec/Prism resolution ( $R=100$ ).



below the value observed with MIRI (see Table 1). The blue optical continuum of this young stellar population makes this difference more pronounced in F1000W, modelling a flux  $\sim 44\%$  fainter ( $2\sigma$  considering the large F1000W uncertainties). For completeness, we have tested an exponentially growing SFH (negative  $\tau$ ) obtaining consistent results.

Since a single young stellar population is not able to justify the UV and optical continuum and the nebular emission of this galaxy, we consider the possibility of having a mix of different stellar populations. In recent years, a number of works have proposed that stellar populations of different ages are required to fit the UV to optical SED of high- $z$  galaxies. Tacchella et al. (2023b) found the presence of old stellar populations in  $z > 7$  galaxies using SED fitting based on NIRSpec+NIRCam data. Underlying  $\sim 100$  Myr stellar populations have been also found in five  $5 < z < 9$  galaxies in which the SEDs are dominated by blue emission produced in young bursty star-forming regions (Giménez-Arteaga et al. 2023). Some authors have found that a double power-law describes better the SFHs in high- $z$  galaxies, as single population models can be out-shone by young UV-bright stars, yielding to lower stellar masses (Giménez-Arteaga et al. 2024). In addition, stellar populations older than  $\sim 10$  Myr produce a redder optical continuum, in accordance with the MIRI colours found in GN-z11 (see Fig. 2).

Therefore, we have also considered a mixed stellar component consisting of a young population ( $t < 10$  Myr) with a constant star formation and a more mature one ( $20 < t < 440$  Myr) formed in a past burst. In this scenario, GN-z11 can be interpreted as a young compact burst in a mature host galaxy. Figure 4 displays the best-fit SED obtained under this approach and the residuals with the observed NIRCam+MIRI fluxes. Under this mixed stellar population scenario, GN-z11 is composed of a young ( $4 \pm 1$  Myr) stellar population that represents the 12% of the total stellar mass ( $1.5 \pm 0.1 \times 10^9 M_\odot$ ), while the more mature one has an average age of  $63 \pm 23$  Myr. These values are consistent with the double-component SED fitting carried out in Tacchella et al. (2023a), where the authors found a point-source with a young stellar population ( $\sim 11$  Myr) along with a more massive and mature one ( $\sim 10^9 M_\odot$  and  $\sim 35$  Myr) dominating the extended emission. Although the near-IR emission produced by the mature stellar population increases the modelled flux in F770W up to  $202 \pm 5$  nJy, it is still  $3\sigma$  fainter than the measured value (see Table 2). Even if we add the contribution of the upper limit of the potential broad component of H $\alpha$  ( $< 2 - 3 \times 10^{18}$  erg s $^{-1}$  cm $^{-2}$ , Álvarez-Márquez et al. 2025), the flux in F770W would increase only by 18–27 nJy and would be still incompatible with the MIRI data by almost  $2\sigma$  (see Table 2). Despite the increase in the modelled flux in F1000W (i.e.,  $130 \pm 6$  nJy) driven by the optical and near-IR emission from the older stellar population, we are not recovering the observed MIRI flux (i.e.,  $249 \pm 77$  nJy).

Although assuming either a single or mixed stellar population provides a good SED fit up to rest-frame  $0.5 \mu\text{m}$ , both approaches fail to reproduce the F770W and F1000W photometry. As discussed in Sect. 4.1, the significant increase in the SED of GN-z11 beyond  $0.66 \mu\text{m}$  cannot be reproduced solely by a combination of different stellar populations. This result suggests the presence of an additional, intrinsically red, component in GN-z11.

### 4.3. SF + AGN SED modelling


In addition to the young starburst (Bunker et al. 2023; Tacchella et al. 2023a; Álvarez-Márquez et al. 2025), some studies indicate the presence of high critical density lines and broad emission lines in the ultraviolet suggesting that GN-z11 could host a type 1 active black hole (Maiolino et al. 2024). To evaluate the presence of a potential AGN in GN-z11, we also consider the combination of star-formation and AGN (Type 1 or 2). The potential presence of an AGN would naturally produce a red component associated to the hot dust emission from the torus around an accreting black hole. The best-fit SED models for the different SF+AGN scenarios are presented in the top panel of Fig. 5. Specifically, CIGALE results including a type 1 and 2 AGNs are represented in blue and green, respectively, while SYNTHESIZER-AGN result is displayed as a red line. Tables 2 and 3 summarise the photometric and physical results of these models.

Adding a type 1 AGN to the mixed stellar population modelled by CIGALE does not significantly improve the SED fitting beyond  $0.66 \mu\text{m}$ . Top and middle panels from Fig. 5 show how the type 1 AGN starts to dominate the emission only above  $0.9 \mu\text{m}$  and the best-fit modelled F770W and F1000W fluxes are well below the observed values by factors 1.5 and 2, respectively. The type 1 AGN is modelled in SYNTHESIZER-AGN by a QSO composite template (see Sect. 3.3) which includes the contribution from broad emission lines and the dust emission in the optical-red and near-infrared spectral range. Top and bottom panels in Fig. 5 shows that the QSO template is almost flat until rest-frame  $\sim 1 \mu\text{m}$  and it is not able to reproduce the observed fluxes in the MIRI reddest filters. Only when including an ad-hoc extra dusty torus (i.e., highly obscured QSO, see Sect. 3.3) the modelled F770W and F1000W fluxes (257 and 258 nJy, respectively) are compatible with the observed ones. However, the addition of this extra hot dust emission to the empirical spectrum of the QSO is not physically justified. As shown in Figure 3, the  $> 0.7 \mu\text{m}$  emission from the empirical  $z \sim 1$  QSO SED agrees with the spectrum of the  $z = 7.1$  luminous QSO J1120+0641, for which JWST/MIRI spectroscopy has revealed a dust emission in the dusty torus with a temperature of about 1400 K (Bosman et al. 2024). This agreement in the SED between the low-intermediate- $z$  and high- $z$  QSOs difficulties the physical justification for the presence of such an extra dusty torus emission.

We consider next the presence of a type 2 AGN, whose UV and optical emission is expected to be obscured by the torus. According to the CIGALE best-fit model, the steep optical and near-IR continuum from the type 2 AGN emission increases the SED modelled fluxes up to 254 nJy for the F770W filter, being compatible with the MIRI measured fluxes (see Table 2). Despite the F1000W photometry not being taken into account during the SED fitting due to its low significance, the type 2 scenario yields a F1000W flux (285 nJy) compatible within the uncertainties, whereas the SED best-fit lies below the upper limit for F2100W. Although the available AGN models in CIGALE do not include the contribution from the emission lines produced by the AGN, the analysis carried out indicates that assuming a type 2 AGN yields a better agreement with the observed MIRI fluxes (i.e., optical to red continuum SED slope) than using different type 1 scenarios.

Regardless of the AGN type considered, we derive similar ages for the different stellar populations. Specifically, we estimate ages of  $\sim 4 - 6$  and  $\sim 20 - 40$  Myr for the young and mature stellar populations, respectively. Ages for the stellar pop-





**Fig. 5.** SED best-fit models considering the star formation and the different AGN scenarios. Top panel show the best-fit SEDs obtained with CIGALE assuming a type 1 and type 2 AGNs as blue and green lines, respectively. Red line displays the best-fit derived with SYNTHESIZER-AGN using a QSO template and an additional dusty torus. Black line and black circles represent the NIRSpec/Prism data presented in Bunker et al. (2023) and the NIRC and MIRI photometry as derived in Sect. 3.1, respectively. Coloured squares show the modelled NIRC and MIRI fluxes from the different best-fit models, while their residuals with the measured photometry are displayed below. Green, blue and red lines have been convolved to match the spectral resolution from NIRSpec/Prism (i.e.,  $R=100$ ). Middle panel shows the CIGALE best-fit models (in black) and the contributions of the nebular and stellar components (in blue) and the Fritz et al. (2006) torus emission (in brown). Dashed and continuous lines differentiate the results drawn from the SED fitting assuming a type 1 and type 2 AGN, respectively. Bottom panel displays the SYNTHESIZER-AGN best-fit (in black) along with the contributions of the nebular and stellar components (in blue), the QSO template (in magenta) and the dusty torus (in brown).

ulations have been derived independently for the spatially resolved point-source and extended stellar components ( $\sim 11$  and  $35$  Myr, respectively) identified in NIRCarn images (Tacchella et al. 2023a). Although the resolution of the MIRI images (i.e., FWHM =  $0.27''$  for F770W) does not allow us to spatially resolve these two components (rest-UV  $r_{\text{eff}} = 0.05''$  for the extended one), our derived ages are compatible the previous NIRCarn photometry.

To ensure the robustness of these results, we replicated the CIGALE SED fits using the SKIRTOR AGN models (Stalevski et al. 2012, 2016), which include a clumpy dust distribution for the torus. These AGN models yield similar results, where only a type 2 AGN can explain the F770W and F1000W fluxes. In next section, we will discuss the general physical scenario drawn from this SED best-fit along with other non-AGN possibilities.

#### 4.4. Origin of the red excess emission

As already shown in the previous sections, the flux in the F770W filter, and in the low-significance ( $3\sigma$ ) F1000W value, shows an excess that cannot be fully explained by the combination of a young and intermediate age stellar populations, even when including the contribution by the nebular emission (i.e., the narrow H $\alpha$  line detected with the MRS). This excess hints to the presence of a source radiating at (rest-) wavelengths of  $0.66$  to  $0.86 \mu\text{m}$ . The excess at these wavelengths can be interpreted as the emission produced by either the presence of a dusty torus of an AGN or dust associated with the extremely compact starburst identified in GN-z11.

##### 4.4.1. Dust in AGN: Dusty torus emission

Although GN-z11 has not been detected in X-rays, a  $3\sigma$  upper limit of  $L_X(2-10 \text{ keV}) < 3 \times 10^{43} \text{ erg s}^{-1}$  has been derived assuming the typical photon index for narrow-line Seyfert 1 galaxies (Maiolino et al. 2024). In addition, a value  $< 2.9 \times 10^{43} \text{ erg s}^{-1}$ , based on its H $\alpha$  luminosity (Ho et al. 2001b; Álvarez-Márquez et al. 2025), could also be consistent with hosting a type 2 AGN. These upper limits agree well with the predicted  $L_X(2-10 \text{ keV})$  based on the SED-fitting assuming the presence of type 1 and 2 AGNs. Using the X-ray extension of the CIGALE models (X-CIGALE, Yang et al. 2020), we obtain a  $L_X(2-10 \text{ keV}) = (0.9 \pm 0.1)$  and  $(2.1 \pm 0.3) \times 10^{43} \text{ erg s}^{-1}$  for the best-fit SED considering the type 1 and 2 AGNs, respectively. Similar X-ray 2-10 keV luminosities (i.e.,  $1.8 \times 10^{43}$  and  $2.6 \times 10^{43} \text{ erg s}^{-1}$ , respectively) are obtained when applying the  $L_X(2-10 \text{ keV})-L_{\text{bol}}$  relation from Marconi et al. (2004) and the bolometric luminosity derived from the SED fitting (see Table 3). In summary, the predicted X-ray luminosities of the best-fit stellar plus AGN (both type 1 and 2) SEDs are compatible with the no detection of X-ray emission in GN-z11.

However, the scenario of a type 1 AGN is not favoured by the present results. The potential presence of an undetected, weak broad H $\alpha$  line of  $2-3 \times 10^{-18} \text{ erg s}^{-1} \text{ cm}^{-2}$  tracing a type 1 AGN (Álvarez-Márquez et al. 2025) would contribute to about  $18-27 \text{ nJy}$  to the F770W flux, not sufficient to explain the excess of  $91 \text{ nJy}$  found during the SED fitting (see Sect. 4.3). In addition, Figure 3 displays how luminous intermediate  $z \sim 1$  (Glikman et al. 2006; Selsing et al. 2016) and  $z > 7$  QSOs (Bosman et al. 2024) have a spectral energy distribution at rest-wavelengths  $0.6-0.8 \mu\text{m}$  flatter than otherwise detected at F770W and F1000W in GN-z11. The SED of these QSOs are consistent with hot dust from the black hole surrounding torus

emitting at a temperature of about  $1400 \text{ K}$ , contributing at wavelengths of  $1 \mu\text{m}$  and above (e.g., Bosman et al. 2024). Moreover, our SYNTHESIZER-AGN best SED fit combining the stellar populations and type 1 AGN requires an additional ad-hoc red source (see Sect 4.3) in addition to the hot ( $\sim 1300-1400 \text{ K}$ ) dust component to explain the observed flux excess at  $0.66$  and  $0.86 \mu\text{m}$ . As already mentioned in the previous section, this extra component is not physically motivated, and therefore the scenario of a standard type 1 AGN is not supported by the present data.

Under the scenario of a Seyfert 2 contribution, the AGN bolometric luminosity derived by CIGALE can be used to estimate the size and mass of the dusty torus assuming its physical parameters (i.e., radial extent and inclination of the torus, number of clouds along the equator of the torus, angular and radial distribution of clouds, optical depth per cloud) similar to those derived for low- $z$  Seyfert 2 galaxies. Following the clumpy dusty torus (Nenkova et al. 2008a,b; García-Bernete et al. 2019), the inner radius of the torus ( $R_{\text{sub}}$ ) is given as

$$R_{\text{sub}}[\text{pc}] = 0.4 \times \left( \frac{1500 \text{ K}}{T_{\text{sub}}} \right)^{2.6} \times \left( \frac{L_{\text{AGN}}}{10^{45} \text{ erg s}^{-1}} \right)^{0.5} \quad (1)$$

where  $L_{\text{AGN}}$  is the AGN bolometric luminosity and  $T_{\text{sub}}$  is the dust sublimation temperature. Assuming the radial extent of the torus ( $R_T$ ) to be about  $10-30$  times  $R_{\text{sub}}$  and a dust sublimation temperature of  $1500 \text{ K}$ , we obtain that the radius of the dusty torus would be in the  $3-9 \text{ pc}$  range for the derived AGN luminosity of  $5.9 \times 10^{44} \text{ erg s}^{-1}$  (see Table 3). Following the  $M_{\text{dust}}$  expression by García-Bernete et al. (2019) and assuming the typical number of clouds, their opacity, and their radial and angular distribution in Seyfert 2 galaxies (i.e.,  $N_0 = 14$ ,  $\tau_V = 70$ ,  $q = 0$  and  $\sigma = 60^\circ$ , respectively), we derive a torus mass of  $0.5-1.5 \times 10^5 M_\odot$ . Thus, the size and mass of the potential torus in GN-z11 would be in the range of those detected in low- $z$  Seyfert 2 galaxies with median size and mass of  $3.5 \pm 3.9 \text{ pc}$  and  $(3.9 \pm 5.1) \times 10^5 M_\odot$ , respectively (García-Bernete et al. 2019). Although the SED fitting and colour analysis of the red excess are compatible with the presence of type 2 AGN, the true nature of this potential AGN along with its resemblance to low- $z$  counterparts are beyond the scope of this work and needs to await the confirmation of the  $10 \mu\text{m}$  flux and the extension of the photometry into redder wavelengths with additional MIRI imaging.

##### 4.4.2. Dust emission in compact starbursts

The large star-formation rate and stellar mass along with the compact size of GN-z11 indicate that this galaxy has extreme stellar mass and star formation surface densities. ( $\Sigma_{\text{SFR}} = 9.33 \times 10^2 M_\odot \text{ yr}^{-1} \text{ kpc}^{-2}$ , Álvarez-Márquez et al. 2025). In addition, the UV and optical line ratios have shown that this target is a low-metallicity galaxy with low dust extinction (Bunker et al. 2023; Álvarez-Márquez et al. 2025). This is in agreement with scenarios of UV bright galaxies at  $z > 10$  where the narrow line component of the nebular emission is barely extinct as the dust would be evacuated outside these extreme compact starbursts by the outflows (Fiore et al. 2023; Ziparo et al. 2023; Dekel et al. 2023; Li et al. 2024). In fact, recent studies have identified the existence of dusty outflows in RXCJ2248-ID ( $z = 6.1$ ; Crespo Gómez et al. 2025), and J0217-0208 ( $z = 6.2$ ; Marques-Chaves et al. 2025) as predicted in the Attenuation-Free models (AFM, Ferrara et al. 2023; Fiore et al. 2023). As GN-z11, these galaxies are also identified as N-emitters (Topping et al. 2024; Marques-Chaves et al. 2025). These two galaxies show large extinctions in the outflowing ionised gas traced

by the broad emission line components identified in their optical spectra. Extinctions of  $A_V = 1.5 - 2.5$  magnitudes are measured while the blue compact starburst appears almost extinction-free as traced by the narrow Balmer line components and the rest-UV continuum emission. These results show that, despite a low dust-extinction in the UV-bright starburst nucleus, a considerable amount of dust is present in these compact N-emitters. The detection of similar dusty outflows in GN-z11 would require extremely deep spectroscopy with MIRI in order to detect the presence of broad line components in the optical hydrogen and oxygen lines.

Moreover, complex dust and gas structures in the interstellar medium around young compact massive clusters have been detected in low metallicity galaxies. Recent NIRCам and MIRI imaging of the low- $z$  galaxy IZw18 (Hirschauer et al. 2024) has shown in detail the complexity of the dusty star formation in this extremely metal-poor galaxy (gas-phase oxygen abundance of 3% solar, see Hirschauer et al. 2024 and references therein). The presence of massive evolved stars, dust-enshrouded stellar objects as well as shells and filaments of dust/gas due to supernovae has been identified. As shown in the images of IZw18 presented in Hirschauer et al. (2024), the dust in and around the stellar clusters could be heated by the strong radiation field, re-radiating in the optical-red and near-infrared wavelength range, and resulting in a flux excess above the stellar+nebular continuum at these wavelengths. Flux excess at wavelengths  $0.66 - 0.86 \mu\text{m}$  as measured in GN-z11 has already been claimed in the low- $z$  metal-poor galaxies SBS0335-052E (Reines et al. 2008; Adamo et al. 2010b) and Haro11 (Adamo et al. 2010a). Notably, Adamo et al. (2010b) found that many of the young and massive stellar clusters present in SBS0335-052E show flux excesses amounting to 0.2 to 0.5 magnitudes above their predicted stellar continuum at rest-frame  $\sim 0.8 \mu\text{m}$ . These galaxies have a hard radiation field ( $> 54.4 \text{ eV}$ ) as demonstrated by the presence of strong He II 1640 and He II 4686 emission lines (Kehrig et al. 2018; Sirressi et al. 2022; Wofford et al. 2021), and even more recently demonstrated by the detection with MIRI spectroscopy of extended mid-infrared [Ne V]  $14.32 \mu\text{m}$  line emission, with an ionization potential of 97 eV, in SBS0335-052E (Mingozzi et al. 2025) and IZw18 (Arroyo-Polonio et al. 2025).

A photoluminescence mechanism (Extended Red Emission, or ERE, see Witt & Viji 2004) has been hypothesised to explain this excess at  $0.7 - 0.9 \mu\text{m}$  in these low-metallicity galaxies (e.g. Reines et al. 2008; Adamo et al. 2010b). This non-thermal mechanism originates from UV-excited small dust grains or complex molecules in dense circumstellar environments. However, if the excess flux observed in MIRI extends beyond one micron, it would not be consistent with the ERE scenario. If the SED continues to rise, the most likely explanation would be thermal emission of dust close to the sublimation temperature (i.e., 1000 – 1500 K). Although uncommon, some dense nucleus and young clusters in low metallicity galaxies show near-IR excess compatible with the presence of dust emitting at 600 – 1100 K (Mrk996, Thuan et al. 2008; NGC5253, Vanzì & Sauvage 2004; Calzetti et al. 2015). The presence of  $> 1000 \text{ K}$  dust has been also observed around Wolf-Rayet stars (Marchenko & Moffat 2007; Lau et al. 2023), whose existence has been proposed as a potential origin of the unusual large N/O ratio in GN-z11 (Cameron et al. 2023; Gunawardhana et al. 2025).

Recent NIRSpect and MIRI spectroscopy has clearly detected PAH emission, including the strong  $3.3 \mu\text{m}$  feature, in the dominant super-cluster (SSC-N) of the nearby galaxy IIZw40 (Lai et al. 2025). This indicates that dust and complex molecules are

produced and survive in low-metallicity ( $0.1 - 0.2 Z_\odot$ , Leitherer et al. 2018) extreme radiation environments as those present in SSC-N, with a radiation field ( $G$ )  $8.6 \times 10^4$  times higher than the Habing radiation field ( $G_0 = 1.6 \times 10^{-3} \text{ erg s}^{-1} \text{ cm}^{-2}$ , Habing 1968). The radiation field in GN-z11, derived from the extreme UV luminosity ( $L_{1500} = 1.7 \times 10^{29} \text{ erg s}^{-1} \text{ Hz}^{-1}$ , Bunker et al. 2023) and small size of the galaxy ( $r_{\text{eff}} = 64 \text{ pc}$ , Tacchella et al. 2023a), is a factor 5 higher (i.e.,  $G/G_0 = 4.3 \times 10^5$ ) than that of SSC-N. Therefore, dust and complex molecules could also be present in the low-metallicity ( $0.17 Z_\odot$ , Álvarez-Márquez et al. 2025) interstellar medium of GN-z11 as in IZw40. The presence and emission processes of dust in very extreme radiation environments such as those present in GN-z11 have to be explored further with MIRI imaging at longer wavelengths covering the near-IR rest-frame.

## 5. Summary and conclusions

This paper presents new imaging of GN-z11 with the MIRI F560W, F770W and F1000W filters, which extend the rest-frame UV-optical spectral coverage up to  $0.86 \mu\text{m}$ . In these MIRI images, GN-z11 shows a compact structure, compatible with previous results based on NIRCам imaging (Tacchella et al. 2023a). We have combined these new MIRI images with archival NIRSpect/Prism and MIRI/MRS spectroscopy, and NIRCам imaging to analyse the SED of GN-z11 covering the rest-frame  $0.12 - 0.86 \mu\text{m}$ . The complete spectral coverage of the NIRSpect and MRS spectroscopy allowed us to evaluate, and remove, the emission line contribution from the NIRCам and MIRI broadband photometry, deriving the continuum-only SED of GN-z11. The main results drawn from this analysis can be summarised as follows:

- Classical colour-colour diagrams have been used to compare the observed colours with several SED templates including stellar populations of different ages, AGNs and high- $z$  LRDs. None of the templates can explain the colours of GN-z11 over the entire spectral range covered by our data. The rest-frame UV-optical colours (i.e. F200W – F444W versus F444W – F560W) of GN-z11 are compatible with the SEDs of unobscured young and intermediate age instantaneous bursts (including nebular emission), while not consistent with their optical and optical-red colours (i.e. F444W – F560W, F560W – F770W and F770W – F1000W). The SEDs of high- $z$  LRDs and QSOs are not consistent with any of the UV-optical-red colours. The spectrum of Mrk 3, classified as a classical Seyfert 2, is compatible with the optical-red colours but fails to explain the observed UV-optical colours.
- The GN-z11 SED beyond  $\sim 0.66 \mu\text{m}$  cannot be entirely explained by the stellar and associated nebular emission alone. The observed flux in the F770W filter (rest-frame  $0.66 \mu\text{m}$ ) shows an excess of  $36 \pm 3\%$  above the predicted flux for a mixed stellar population of young (4 Myr) and intermediate (63 Myr) age stars. The flux marginally detected in the F1000W filter (rest-frame  $0.86 \mu\text{m}$ ) represents an even larger excess, although with larger uncertainties (i.e.,  $91 \pm 28\%$ ).
- Both the continuum-only colours and the SED analysis covering the entire  $0.12 - 0.86 \mu\text{m}$  spectral range identify an optical-red continuum excess at wavelengths  $0.66 - 0.86 \mu\text{m}$ . This hints to a red source which appears undetected in the ultraviolet and represents a minor contribution to the emission in the optical range (i.e.,  $< 0.6 \mu\text{m}$ ). The origin of this



excess is unclear. One scenario proposes that it can be produced by the emission from the dusty torus of a type 2 AGN characterised by a radius of 3–9 pc with a mass of  $(0.5 - 1.5) \times 10^5 M_{\odot}$ . Alternatively, the excess could be related to hot dust emission or photoluminescence of dust (Extended Red Emission) in the presence of the extreme radiation field ( $G/G_0 = 4.3 \times 10^5$ ) produced by the compact and massive starburst in GN-z11, as already hypothesised in some low- $z$  low-metallicity starbursts. The presence of a standard type 1 AGN is not supported by the present data as an additional ad-hoc red source is required to explain the observed flux excess.

In summary, flux excess has been identified in the MIRI/F770W (rest-  $0.66 \mu\text{m}$ ) and also, although with lower significance, in MIRI/F1000W (rest-  $0.86 \mu\text{m}$ ) images of GN-z11. This excess is likely due to the dust emission associated to an AGN dusty torus or to dust in the environment of an extremely compact and young starburst. However, the current wavelength coverage is still limited and the true nature of the heating source is still unclear. Deep imaging with MIRI/F1000W and redder filters are required to confirm the  $0.86 \mu\text{m}$  detection and extend the SED coverage beyond  $H\alpha$ , characterizing its near-IR spectral shape up to at least  $1.1 \mu\text{m}$ . Imaging with deep integrations using MIRI F1000W and F1280W filters will be executed with JWST in the near future (PID 7605). This will help to discriminate between the two proposed scenarios. In addition, deeper than existing MIRI/MRS spectroscopy would also be required to detect or rule out the presence of a weak broad  $H\alpha$  associated with a  $\sim 10^6 M_{\odot}$  type 1 AGN, or to identify the existence of dusty outflows.

**Acknowledgements.** A.C.G. acknowledges support by JWST contract B0215/JWST-GO-02926. L.C. and J.A.-M. acknowledge support by grants PIB2021-127718NB-I00 and PID2024-158856NA-I00, A.A.-H. by grant PID2021-124665NB-I00, and P.G.-P. by grant PID2022-139567NB-I00 funded by the Spanish Ministry of Science and Innovation and the State Agency of Research MCIN/AEI/10.13039/501100011033 and ERDF "A way of making Europe". M.A. acknowledges financial support from Comunidad de Madrid under Atracción de Talento grant 2020-T2/TIC-19971. D.L. was supported by research grants (VIL16599, VIL54489) from VILLUM FONDEN. A.B. and G.O. acknowledge support from the Swedish National Space Administration (SNSA). A.B. acknowledges funding from the "FirstGalaxies" Advanced Grant from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant agreement No. 789056). L.A.B. acknowledges support from the Dutch Research Council (NWO) under grant VI.Veni.242.055 (<https://doi.org/10.61686/LAJVP77714>) and the ERC Consolidator grant 101088676 ("VOYAJ"). This work is based on observations made with the NASA/ESA/CSA James Webb Space Telescope. The data were obtained from the Mikulski Archive for Space Telescopes at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-03127 for JWST; and from the [European JWST archive \(eJWST\)](#) operated by the ESDC.

## References

- Adamo, A., Östlin, G., Zackrisson, E., et al. 2010a, MNRAS, 407, 870  
 Adamo, A., Zackrisson, E., Östlin, G., & Hayes, M. 2010b, ApJ, 725, 1620  
 Akins, H. B., Casey, C. M., Lambrides, E., et al. 2025, ApJ, 991, 37  
 Álvarez-Márquez, J., Crespo Gómez, A., Colina, L., et al. 2025, A&A, 695, A250  
 Álvarez-Márquez, J., Crespo Gómez, A., Colina, L., et al. 2023, A&A, 671, A105  
 Argyriou, I., Glasse, A., Law, D. R., et al. 2023, A&A, 675, A111  
 Arroyo-Polonio, A., Kehrig, C., Vílchez, J. M., et al. 2025, ApJ, 987, L36  
 Bhatt, M., Gallerani, S., Ferrara, A., et al. 2024, A&A, 686, A141  
 Boquien, M., Burgarella, D., Roehlly, Y., et al. 2019, A&A, 622, A103  
 Bosman, S. E. I., Álvarez-Márquez, J., Colina, L., et al. 2024, Nature Astronomy, 8, 1054  
 Bosman, S. E. I., Álvarez-Márquez, J., Davies, F. B., et al. 2025, sub. ApJ, arXiv:2511.02902  
 Bouchet, P., García-Marín, M., Lagage, P. O., et al. 2015, PASP, 127, 612  
 Bouwens, R. J., Illingworth, G. D., González, V., et al. 2010, ApJ, 725, 1587  
 Bruzual, G. & Charlot, S. 2003, MNRAS, 344, 1000  
 Bunker, A. J., Cameron, A. J., Curtis-Lake, E., et al. 2024, A&A, 690, A288  
 Bunker, A. J., Saxena, A., Cameron, A. J., et al. 2023, A&A, 677, A88  
 Burgarella, D., Buat, V., & Iglesias-Páramo, J. 2005, MNRAS, 360, 1413  
 Calzetti, D., Armus, L., Bohlin, R. C., et al. 2000, ApJ, 533, 682  
 Calzetti, D., Johnson, K. E., Adamo, A., et al. 2015, ApJ, 811, 75  
 Cameron, A. J., Saxena, A., Bunker, A. J., et al. 2023, A&A, 677, A115  
 Carniani, S., Hainline, K., D'Eugenio, F., et al. 2024, Nature, 633, 318  
 Chabrier, G. 2003, PASP, 115, 763  
 Charbonnel, C., Schaerer, D., Prantzos, N., et al. 2023, A&A, 673, L7  
 Charlot, S. & Fall, S. M. 2000, ApJ, 539, 718  
 Crespo Gómez, A., Tamura, Y., Colina, L., et al. 2025, sub. A&A, arXiv:2511.14658  
 Curti, M., Mannucci, F., Cresci, G., & Maiolino, R. 2020, MNRAS, 491, 944  
 Curtis-Lake, E., Carniani, S., Cameron, A., et al. 2023, Nature Astronomy, 7, 622  
 de Graaff, A., Rix, H.-W., Naidu, R. P., et al. 2025, A&A, 701, A168  
 Dekel, A., Sarkar, K. C., Birnboim, Y., Mandelker, N., & Li, Z. 2023, MNRAS, 523, 3201  
 D'Eugenio, F., Cameron, A. J., Scholtz, J., et al. 2025, ApJS, 277, 4  
 Eisenstein, D. J., Johnson, B. D., Robertson, B., et al. 2023, Accepted to ApJ Supplement, arXiv:2310.12340  
 Ferland, G. J., Chatzikos, M., Guzmán, F., et al. 2017, Rev. Mexicana Astron. Astrofis., 53, 385  
 Ferrara, A., Pallottini, A., & Dayal, P. 2023, MNRAS, 522, 3986  
 Fiore, F., Ferrara, A., Bischetti, M., Feruglio, C., & Travascio, A. 2023, ApJ, 943, L27  
 Fritz, J., Franceschini, A., & Hatziminaoglou, E. 2006, MNRAS, 366, 767  
 Fudamoto, Y., Oesch, P. A., Walter, F., et al. 2024, MNRAS, 530, 340  
 García-Bernete, I., Ramos Almeida, C., Alonso-Herrero, A., et al. 2019, MNRAS, 486, 4917  
 Giménez-Arteaga, C., Fujimoto, S., Valentino, F., et al. 2024, A&A, 686, A63  
 Giménez-Arteaga, C., Oesch, P. A., Brammer, G. B., et al. 2023, ApJ, 948, 126  
 Glikman, E., Helfand, D. J., & White, R. L. 2006, ApJ, 640, 579  
 Greene, J. E. & Ho, L. C. 2005, ApJ, 630, 122  
 Guainazzi, M., Risaliti, G., Awaki, H., et al. 2016, MNRAS, 460, 1954  
 Gunasekera, C. M., van Hoof, P. A. M., Chatzikos, M., & Ferland, G. J. 2023, Research Notes of the American Astronomical Society, 7, 246  
 Gunawardhana, M. L. P., Brinchmann, J., Croom, S., et al. 2025, MNRAS, 543, 3172  
 Habing, H. J. 1968, Bull. Astron. Inst. Netherlands, 19, 421  
 Hirschauer, A. S., Crouzet, N., Habel, N., et al. 2024, AJ, 168, 23  
 Ho, L. C., Feigelson, E. D., Townsley, L. K., et al. 2001a, ApJ, 549, L51  
 Ho, L. C., Feigelson, E. D., Townsley, L. K., et al. 2001b, ApJ, 549, L51  
 Ji, X., Maiolino, R., Ferland, G., et al. 2025, MNRAS, 541, 2134  
 Jiang, L., Kashikawa, N., Wang, S., et al. 2021, Nature Astronomy, 5, 256  
 Kehrig, C., Vílchez, J. M., Guerrero, M. A., et al. 2018, MNRAS, 480, 1081  
 Kewley, L. J. & Dopita, M. A. 2002, ApJS, 142, 35  
 Lai, T. S. Y., Duval, S., Smith, J. D. T., et al. 2025, ApJ, 991, L56  
 Lau, R. M., Wang, J., Hankins, M. J., et al. 2023, ApJ, 951, 89  
 Le Fèvre, O., Lemaux, B. C., Nakajima, K., et al. 2019, A&A, 625, A51  
 Leitherer, C., Byler, N., Lee, J. C., & Levesque, E. M. 2018, ApJ, 865, 55  
 Li, Z., Dekel, A., Sarkar, K. C., et al. 2024, A&A, 690, A108  
 Libralato, M., Argyriou, I., Dicken, D., et al. 2024, PASP, 136, 034502  
 Maiolino, R., Scholtz, J., Witsstok, J., et al. 2024, Nature, 627, 59  
 Marchenko, S. V. & Moffat, A. F. J. 2007, in Astronomical Society of the Pacific Conference Series, Vol. 367, Massive Stars in Interactive Binaries, ed. N. St.-Louis & A. F. J. Moffat, 213  
 Marconi, A., Risaliti, G., Gilli, R., et al. 2004, MNRAS, 351, 169  
 Marques-Chaves, R., Schaerer, D., Dessauges-Zavadsky, M., et al. 2025, arXiv e-prints, arXiv:2510.12411  
 Mazzolari, G., Scholtz, J., Maiolino, R., et al. 2025, A&A, 700, A12  
 Mazzolari, G., Übler, H., Maiolino, R., et al. 2024, A&A, 691, A345  
 Miller, J. S. & Goodrich, R. W. 1990, ApJ, 355, 456  
 Mingozi, M., García del Valle-Espinosa, M., James, B. L., et al. 2025, ApJ, 985, 253  
 Naidu, R. P., Matthee, J., Katz, H., et al. 2025, arXiv e-prints, arXiv:2503.16596  
 Nenkova, M., Sirocky, M. M., Ivezić, Ž., & Elitzur, M. 2008a, ApJ, 685, 147  
 Nenkova, M., Sirocky, M. M., Nikutta, R., Ivezić, Ž., & Elitzur, M. 2008b, ApJ, 685, 160  
 Noll, S., Burgarella, D., Giovannoli, E., et al. 2009, A&A, 507, 1793  
 Oesch, P. A., Brammer, G., van Dokkum, P. G., et al. 2016, ApJ, 819, 129  
 Östlin, G., Pérez-González, P. G., Melinder, J., et al. 2025, A&A, 696, A57  
 Pérez-González, P. G., Barro, G., Rieke, G. H., et al. 2024a, ApJ, 968, 4  
 Pérez-González, P. G., Barro, G., Rieke, G. H., et al. 2024b, ApJ, 968, 4

- Pérez-González, P. G., Gil de Paz, A., Zamorano, J., et al. 2003, *MNRAS*, 338, 508
- Pérez-González, P. G., Rieke, G. H., Villar, V., et al. 2008, *ApJ*, 675, 234
- Planck Collaboration, Aghanim, N., Akrami, Y., et al. 2020, *A&A*, 641, A6
- Polletta, M., Tajer, M., Maraschi, L., et al. 2007, *ApJ*, 663, 81
- Polletta, M. d. C., Wilkes, B. J., Siana, B., et al. 2006, *ApJ*, 642, 673
- Reines, A. E., Johnson, K. E., & Hunt, L. K. 2008, *AJ*, 136, 1415
- Robertson, B. E., Tacchella, S., Johnson, B. D., et al. 2023, *Nature Astronomy*, 7, 611
- Scholtz, J., Witten, C., Laporte, N., et al. 2024, *A&A*, 687, A283
- Selsing, J., Fynbo, J. P. U., Christensen, L., & Krogager, J. K. 2016, *A&A*, 585, A87
- Senchyna, P., Plat, A., Stark, D. P., et al. 2024, *ApJ*, 966, 92
- Sirressi, M., Adamo, A., Hayes, M., et al. 2022, *AJ*, 164, 208
- Spinelli, P. F., Storch-Bergmann, T., Brandt, C. H., & Calzetti, D. 2006, *ApJS*, 166, 498
- Stalevski, M., Fritz, J., Baes, M., Nakos, T., & Popović, L. Č. 2012, *MNRAS*, 420, 2756
- Stalevski, M., Ricci, C., Ueda, Y., et al. 2016, *MNRAS*, 458, 2288
- Stanway, E. R. & Eldridge, J. J. 2018, *MNRAS*, 479, 75
- Tacchella, S., Eisenstein, D. J., Hainline, K., et al. 2023a, *ApJ*, 952, 74
- Tacchella, S., Johnson, B. D., Robertson, B. E., et al. 2023b, *MNRAS*, 522, 6236
- Taylor, A. J., Kokorev, V., Kocevski, D. D., et al. 2025, *ApJ*, 989, L7
- Terao, K., Nagao, T., Onishi, K., et al. 2022, *ApJ*, 929, 51
- Thuan, T. X., Hunt, L. K., & Izotov, Y. I. 2008, *ApJ*, 689, 897
- Topping, M. W., Stark, D. P., Senchyna, P., et al. 2024, *MNRAS*, 529, 3301
- Vanzi, L. & Sauvage, M. 2004, *A&A*, 415, 509
- Witt, A. N. & Vijh, U. P. 2004, in *Astronomical Society of the Pacific Conference Series*, Vol. 309, *Astrophysics of Dust*, ed. A. N. Witt, G. C. Clayton, & B. T. Draine, 115
- Wofford, A., Vidal-García, A., Feltre, A., et al. 2021, *MNRAS*, 500, 2908
- Yang, G., Boquien, M., Buat, V., et al. 2020, *MNRAS*, 491, 740
- Ziparo, F., Ferrara, A., Sommovigo, L., & Kohandel, M. 2023, *MNRAS*, 520, 2445

<sup>1</sup> Space Telescope Science Institute (STScI), 3700 San martin Drive, Baltimore, MD 21218, USA  
e-mail: acrespo@stsci.edu

<sup>2</sup> Centro de Astrobiología (CAB), CSIC-INTA, Ctra. de Ajalvir km 4, Torrejón de Ardoz, E-28850, Madrid, Spain

<sup>3</sup> European Space Agency, Space Telescope Science Institute, Baltimore, Maryland, USA

<sup>4</sup> Centro de Astrobiología (CAB), CSIC-INTA, Camino Viejo del Castillo s/n, 28692 Villanueva de la Cañada, Madrid, Spain

<sup>5</sup> Department of Astronomy, Stockholm University, Oscar Klein Centre, AlbaNova University Centre, 106 91 Stockholm, Sweden

<sup>6</sup> Institute for Theoretical Physics, Heidelberg University, Philosophenweg 12, 69120 Heidelberg, Germany

<sup>7</sup> Max-Planck-Institut für Astronomie, Königstuhl 17, 69117 Heidelberg, Germany

<sup>8</sup> Department of Physics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX13RH, U.K.

<sup>9</sup> Telespazio UK for the European Space Agency, ESAC, Camino Bajo del Castillo s/n, 28692 Villanueva de la Cañada, Spain

<sup>10</sup> DARK, Niels Bohr Institute, University of Copenhagen, Jagtvej 128, 2200 Copenhagen, Denmark

<sup>11</sup> Steward Observatory, University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721, USA

<sup>12</sup> Cosmic Dawn Center, DTU Space, Technical University of Denmark, Elektrovej 327, 2800 Kgs. Lyngby, Denmark

<sup>13</sup> Cosmic Dawn Centre, Copenhagen, Denmark

<sup>14</sup> University of the Pacific, Stockton, CA 90340, USA

<sup>15</sup> Leiden Observatory, Leiden University, PO Box 9513, NL-2300 RA Leiden, The Netherlands

<sup>16</sup> Department of Astronomy, The University of Texas at Austin, Austin, TX, USA

<sup>17</sup> MIT Kavli Institute for Astrophysics and Space Research, 77 Massachusetts Ave., Cambridge, MA 02139, USA

## Appendix A: NIRCcam-NIRSpec-MIRI normalization

As mentioned in Sect. 3.1, we used the NIRCcam photometry to renormalise the NIRSpec spectra. In this appendix we introduce the main steps followed during this renormalization.

First, we PSF-matched the NIRCcam images to the MIRI/F560W resolution. This is done by deriving empirical PSFs (ePSFs) based on the stars available in the FoV (i.e., between 3 and 6 non-saturated in each filter). We then create the corresponding kernel for each NIRCcam image using the MIRI PSF models from Libralato et al. (2024), as there are not enough available stars in the MIRI FoV. Once the convolution has been applied, we extracted the NIRCcam and MIRI photometry using circular apertures of  $r = 0.5''$  (see Sect. 1 for details). As a consistency check, we extracted the photometry at the native spatial resolution of each filter, applying the corresponding aperture corrections. The resulting fluxes are consistent with those obtained from the PSF-matched data within uncertainties.

We noticed that our NIRCcam photometry is slightly larger (20 – 25%) than the one derived in Tacchella et al. 2023a for a smaller aperture ( $r = 0.35''$ ). In our analysis, we have used a more recent data release from JADES (i.e., DR3) where the NIRCcam images were reduced with improved calibration files from JWST. In addition, we have used a larger and more distant from GN-z11 annular aperture to derive the background level. The combined effect of using a more distant aperture to derive the background level and more recent calibration files and pipeline favour the detection of diffuse emission that was missed in the original calibrated images.

Finally, we convolved the NIRSpec/Prism spectra with the NIRCcam filter transmission curves to derive the expected fluxes in each filter. These NIRSpec continuum values were therefore compared with the NIRCcam photometry, measuring a median 32.5% offset that remains nearly constant with wavelength, showing variations of less than 5%. We had also compared the NIRCcam photometry with the NIRSpec pseudo-continuum fluxes obtained in 750Å bins (see Sect. 3.3), obtaining similar results. We therefore multiply the NIRSpec/Prism spectra by 1.325 to match the NIRCcam and NIRSpec data. Bunker et al. (2023) reported an agreement between the NIRSpec/Prism reduction, which included slit-loss corrections for point-like source, and the  $r = 0.1''$  NIRCcam photometry from Tacchella et al. 2023a. This difference is produced by as GN-z11 UV morphology deviates from an ideal point-like source, which is assumed during the data reduction. In fact, even larger differences between NIRCcam and NIRSpec/MSA (i.e., 30 – 50%) have been found in other high- $z$  galaxies (e.g., JADES-GS-z14-0, Carniani et al. 2024).

For consistency, we tested whether this difference in the UV flux would lead to any noticeable difference during the SED fitting. Therefore, we replicated the SED analysis described in Sect. 4.3, with a double stellar population and the presence of an AGN, considering the NIRSpec pseudo-filters prior normalization. As in this work we assumed a uniform re-normalization, most of the properties derived with the UV slope remains unaltered. However, having a lower UV continuum results in a slightly older (i.e., 6 Myr vs 4 Myr) and more massive young stellar population (i.e., 13% larger). As the  $H\alpha$  and the optical and near-IR fluxes do not change, we do not observe significant differences either in the SFR, the total stellar mass or the AGN luminosity.