

# Computing the Hopf invariant

Oleg R. Musin and Timur Shamazov

## Abstract

We consider Whitehead's integral formula and propose an algorithm for computing the Hopf invariant for simplicial mappings.

## 1 Introduction

In 1931 Heinz Hopf [4] defined the map  $h : S^3 \rightarrow S^2$  and proved that  $h$  is not homotopic to the constant map. The *Hopf invariant* of a smooth or simplicial map  $f : S^3 \rightarrow S^2$  is the linking number

$$H(f) := \text{lk}(f^{-1}(x), f^{-1}(y)) \in \mathbb{Z},$$

where  $x \neq y \in S^2$  are generic points, i.e.  $f^{-1}(x)$  and  $f^{-1}(y)$  are circles or unions of circles.

Actually,  $H(f)$  is the element  $[f]$  of the group  $\pi_3(S^2) = \mathbb{Z}$ . The Hopf invariant completely classifies the homotopy classes of maps from  $S^3$  to  $S^2$ . Therefore, in order to understand whether a map is null-homotopic, it is sufficient to check whether its Hopf invariant is equal to zero.

A generalization of the Hopf invariant for continuous mappings  $f : S^{2n-1} \rightarrow S^n$  is known. Steenrod [14] defined the Hopf invariant using the cup product in the cohomology ring, we consider this definition in Section 2. In the smooth case, there is *Whitehead's integral formula* [16] that expresses the Hopf invariant through an integral of a special differential form over  $S^{2n-1}$ . Let  $\omega$  be a  $n$ -form on  $S^n$  such that  $\int_{S^n} \omega = 1$ . Then we may consider its pullback  $f^*\omega$ , which is a  $n$ -form on  $S^{2n-1}$ . Consider a  $(n-1)$ -form  $\theta$  on  $S^{2n-1}$  such that  $d\theta = f^*\omega$ . Then the Hopf invariant is given by

$$H(f) = \int_{S^{2n-1}} \theta \wedge f^*\omega \tag{1}$$

In 1984, S. P. Novikov [12] proposed a generalization of (1) for multivalued mappings.

If  $f : S^m \rightarrow S^n$  is a simplicial map, then this map is uniquely defined by

$$L : V(T_1) \rightarrow V(T_2),$$

where  $T_1$  and  $T_2$  are triangulations of  $S^m$  and  $S^n$ , and there is no  $n$ -dimensional simplex in  $T_1$  such that the  $L$ -image of its vertices is  $n + 1$  distinct vertices in  $T_2$ . Since  $L$  can be extended linearly to all  $k$ -dimensional,  $k \geq 1$ , faces of  $T_1$ , we obtain a simplicial map

$$f_L : S^m \rightarrow S^n.$$

It is clear, that  $f_L$  coincides with  $f$ . Suppose that a coloring (labeling)  $L$  is given. The question is how to find  $[f_L] \in \pi_m(S^n)$ ? This problem has been solved only for some special cases, see [1, 5–7, 9–11].

Let  $T_2$  be the boundary triangulation of  $(n + 1)$ -simplex  $\Delta^{n+1}$ . We denote it by  $S_{n+2}^n$ . Then

$$L : V(T_1) \rightarrow V(S_{n+2}^n) = \{1, \dots, n + 2\}$$

be a coloring (labeling) of the vertex set of  $T_1$ .

Motivation comes from obtaining estimates of the invariant  $\mu(d)$ , which appears in the quantitative Sperner theorem from [11]. In that work, it is shown that the number of fully labeled 3-simplices for a map  $f$  from a triangulated  $D^3$  into  $\{1, 2, 3, 4\}$ , having no fully labeled 2-simplices on the boundary, must be at least  $\mu([f])$ , where  $[f]$  is the homotopy class of the map  $f$ .

In this paper, we consider an extension of (1) for simplicial maps  $f : S^{2n-1} \rightarrow S^n$  and propose an algorithm for computing the Hopf invariant.

## 2 Hopf invariant for simplicial maps

Steenrod [14], see also [2, 15, 17], considered differential forms, the Hopf invariant, and the integral Whitehead formula for triangulations, simplicial and cell complexes.

Let  $f : S^{2n-1} \rightarrow S^n$  be a simplicial map. To calculate the Hopf invariant, take an  $n$ -cocycle  $\omega$  representing a generator of  $H^n(S^n)$ , and consider its pullback  $f^*\omega \in C^n(S^{2n-1})$ . Since  $H^n(S^{2n-1})$  is trivial, there exists an  $(n - 1)$ -cochain  $\theta$  such that  $\delta\theta = f^*\omega$ . It is easy to verify that the cochain  $\theta \smile f^*\omega$  is a cocycle. Indeed,

$$\delta(\theta \smile f^*\omega) = \delta\theta \smile f^*\omega - (-1)^{|\theta|}\theta \smile \delta f^*\omega = f^*\omega \smile f^*\omega - 0 = 0,$$

because  $\delta f^*\omega = f^*(\delta\omega) = 0$  and the cup product  $f^*\omega \smile f^*\omega \in C^{2n}(S^{2n-1})$  is identically zero for dimensional reasons.

Here, the **coboundary operator**  $\delta : C^k(X) \rightarrow C^{k+1}(X)$  is defined on a  $k$ -cochain  $\alpha$  by

$$(\delta\alpha)([v_0, \dots, v_{k+1}]) = \sum_{i=0}^{k+1} (-1)^i \alpha([v_0, \dots, \hat{v}_i, \dots, v_{k+1}]),$$

where  $[v_0, \dots, v_{k+1}]$  is an oriented  $(k + 1)$ -simplex and  $\hat{v}_i$  denotes omission of the vertex  $v_i$ .

The **cup product**  $\smile : C^k(X) \times C^\ell(X) \rightarrow C^{k+\ell}(X)$  is defined on oriented simplices by

$$(\alpha \smile \beta)([v_0, \dots, v_{k+\ell}]) = \alpha([v_0, \dots, v_k]) \cdot \beta([v_{k+1}, \dots, v_{k+\ell}]).$$

This operation induces the cup product in cohomology, which is associative, graded-commutative, and natural with respect to continuous maps.

Therefore, we can consider the cohomology class of the cocycle  $\theta \smile f^*(\omega)$ . This class lies in the group  $H^{2n-1}(S^{2n-1})$ , so it is proportional to the generator of the group  $H^{2n-1}(S^{2n-1})$ . The coefficient of proportionality is the Hopf invariant  $H(f)$ .

**Lemma 1.** *The Hopf invariant  $H(f)$  of  $f$  is given by the pairing of the cochain  $\theta \smile f^*\omega$  with the fundamental class  $\Delta$  of  $S^{2n-1}$ :*

$$H(f) = \langle \theta \smile f^*\omega, \Delta \rangle. \quad (2)$$

**Lemma 2.** *Let  $T$  be a finite simplicial triangulation of  $S^{2n-1}$ . A cochain  $\theta \in C^{n-1}(S^{2n-1}; \mathbb{Q})$  satisfying  $\delta\theta = f^*\omega$  can be obtained by solving a linear system over  $\mathbb{Q}$  with the following structure:*

1. *The variables are the values of  $\theta$  on the  $(n-1)$ -simplices of  $T$ .*
2. *For each  $n$ -simplex  $\sigma$  of  $T$ , the equation*

$$(\delta\theta)(\sigma) = f^*\omega(\sigma) \quad (3)$$

*provides one linear equation.*


3. *In each such equation, exactly  $n+1$  variables appear (corresponding to the  $(n-1)$ -faces of  $\sigma$ ), and their coefficients are  $\pm 1$ .*

*The number of equations equals the number of  $n$ -simplices in  $T$ . Because  $H^n(S^{2n-1}; \mathbb{Q}) = 0$ , this system is consistent and admits a rational solution.*

In case  $n = 2$  each equation has the form:

$$\langle \theta, v_0v_1 \rangle + \langle \theta, v_1v_2 \rangle + \langle \theta, v_2v_0 \rangle = \langle f^*\omega, v_0v_1v_2 \rangle$$

for every 2-simplex  $v_0v_1v_2$  in  $S^3$ . That is, the simplicial analog of Stokes' formula must be satisfied.



A solution to this system of linear equations exists but it is not unique.

**Theorem 3.** *The dimension of the solution space is*

$$f_{n-1} - f_{n-2} + f_{n-3} + \dots + (-1)^n f_0 + (-1)^{n+1}$$

Where  $f_i$  is the number of simplices of dimension  $i$  in the sphere  $S^{2n-1}$ .

*Proof.* We will prove this by induction on  $n$ . Indeed, the dimension of the solution space is equal to  $\dim \ker(\delta : C^{n-1} \rightarrow C^n)$ . Since  $H^n(S^{2n-1}) = 0$ , we have

$$\begin{aligned} \dim \ker(\delta : C^{n-1} \rightarrow C^n) &= \dim \operatorname{im}(\delta : C^{n-2} \rightarrow C^{n-1}) = \\ &= \dim C^{n-1} - \dim \ker(\delta : C^{n-2} \rightarrow C^{n-1}) \end{aligned}$$

And since  $\dim C^{n-1} = f_{n-1}$  we can apply the induction hypothesis.  $\square$

As a consequence, for the computation of the Hopf invariant on  $S^{2n-1}$ , one may choose a spanning subcomplex in the  $n$ -skeleton of  $S^{2n-1}$ , assign values of  $\theta$  arbitrarily on it, and then find a unique extension to the remaining  $(n-1)$ -simplices such that

$$\delta\theta = f^*\omega,$$

where  $f^*\omega \in H^n(S^{2n-1}; \mathbb{Z})$  is the pullback of the generator of  $H^n(S^n; \mathbb{Z})$ .

### 3 Algorithm and its computational complexity for computing the Hopf invariant

Let  $S_{n+2}^n$  be boundary triangulation of  $(n+1)$ -simplex  $\Delta^{n+1}$  sphere  $S^n$ , and let  $T_1$  be a finite simplicial triangulation of the sphere  $S^{2n-1}$ . Let a simplicial map  $f : T_1 \rightarrow S_{n+2}^n$  be defined by a vertex labelling: to each vertex  $v$  of  $T_1$  we assign a label  $\ell(v) \in \{1, 2, \dots, n+2\}$ . The Hopf invariant  $H(f)$  can be computed as follows:

1. Fix an orientation of  $S_{n+2}^n$  and select a single oriented  $n$ -simplex  $\bar{\sigma} = [w_0, \dots, w_n]$  of  $S_{n+2}^n$ . Define the function  $\omega$  on  $n$ -simplices of  $S_{n+2}^n$  by

$$\omega(\sigma) = \begin{cases} +1, & \text{if } \sigma = \bar{\sigma}, \\ -1, & \text{if } \sigma = -\bar{\sigma} \text{ (the oppositely oriented simplex),} \\ 0, & \text{otherwise.} \end{cases}$$

2. For each oriented  $n$ -simplex  $\sigma = [v_0, \dots, v_n]$  of  $T_1$ , set

$$f^*\omega(\sigma) = \omega([\ell(v_0), \dots, \ell(v_n)]).$$

By construction,  $f^*\omega(\sigma) \in \{-1, 0, +1\}$ . The value is nonzero exactly when the labelled simplex  $[\ell(v_0), \dots, \ell(v_n)]$  coincides with  $\bar{\sigma}$  or its opposite.

3. Introduce one integer variable  $x_\tau$  for each oriented  $(n-1)$ -simplex  $\tau$  of  $T_1$ , this variable represents the value  $\theta(\tau)$ . The coboundary equation on an oriented  $n$ -simplex  $\sigma = [v_0, \dots, v_n]$  becomes

$$\sum_{i=0}^n (-1)^i x_{[v_0, \dots, \widehat{v_i}, \dots, v_n]} = f^* \omega(\sigma),$$

where  $\widehat{v_i}$  denotes omission. This yields a linear system over  $\mathbb{Z}$  with  $|T_1^{(n)}|$  equations (one per  $n$ -simplex of  $T_1$ ) and  $|T_1^{(n-1)}|$  unknowns. Compute any rational solution  $\theta$  (e.g., using Gaussian elimination over  $\mathbb{Q}$ ).

4. Choose a fundamental cycle  $\Delta = \sum_{\sigma \in T_1^{(2n-1)}} \varepsilon_\sigma \sigma$  of  $T_1$  (where  $\varepsilon_\sigma = \pm 1$  according to the orientation). For each oriented  $(2n-1)$ -simplex  $\sigma = [u_0, \dots, u_{2n-1}]$ , compute

$$(\theta \smile f^* \omega)(\sigma) = \theta([u_0, \dots, u_{n-1}]) \cdot f^* \omega([u_{n-1}, \dots, u_{2n-1}]).$$

The Hopf invariant is the sum

$$H(f) = \langle \theta \smile f^* \omega, \Delta \rangle = \sum_{\sigma \in T_1^{(2n-1)}} \varepsilon_\sigma (\theta \smile f^* \omega)(\sigma).$$

**Theorem 4.** *This algorithm computes the Hopf invariant of a simplicial map  $f : S^{2n-1} \rightarrow S^n$  defined by a vertex labeling  $\ell : T_1^{(0)} \rightarrow (S_{n+2}^n)^{(0)}$ . The result is an integer  $H(f)$  that depends only on the homotopy class of  $f$ .*

**Theorem 5.** *The complexity of the algorithm is  $O(f_n^3)$ .*

*Proof.* The most computationally expensive step of our algorithm is solving the system of linear equations with  $f_n$  equations. Solving it in the general case requires  $O(f_n^3)$  time.  $\square$

We observe that the matrix of our system is sparse, with each row containing only  $n+1$  non-zero entries. The number of columns corresponds to the number of  $n$ -simplices in the triangulation of  $S^{2n-1}$ , denoted by  $f_n$ . Such systems can be solved using various heuristic approaches. For specific methods, we refer to [13] and [3].

In the special case when  $n$  is odd, it is known that every map has Hopf invariant equal to zero, so in that case the computation reduces to  $O(1)$  complexity.

We believe that no local combinatorial formula for the Hopf invariant exists, so this algorithm is essentially the fastest known method for computing it.

**Acknowledgments.** We thank Dennis Sullivan for suggesting the idea of computing the Hopf invariant via Whitehead's formula for simplicial maps.

Timur Shamazov is supported by the "Priority 2030" strategic academic leadership program. The author thanks the Summer Research Programme at MIPT - LIPS-25 for the opportunity to work on this and other problems

# References

- [1] K. Apolonskaya and O. R. Musin, Minimal simplicial spherical mappings with a given degree, preprint, arXiv:2511.10870
- [2] R. Bott and L.W. Tu, Differential Forms in Algebraic Topology, Springer, 1982
- [3] T. A. Davis, Direct Methods for Sparse Linear Systems, SIAM, 2006.
- [4] H. Hopf, (1931), Über die Abbildungen der dreidimensionalen Sphäre auf die Kugelfläche, *Mathematische Annalen*, **104** (1931) 637–665.
- [5] K. V. Madahar, Simplicial maps from the 3–sphere to the 2–sphere, *Adv. Geom.*, **2** (2012), 99–106.
- [6] K. V. Madahar and K. S. Sarkaria, A minimal triangulation of the Hopf map and its application, *Geom. Dedicata*, **82** (2000), 105–114.
- [7] K. V. Madahar and K. S. Sarkaria, Minimal simplicial self-maps of the 2-sphere *Geom. Dedicata*, **84** (2001), 25–31.
- [8] J. W. Milnor, Topology from the differentiable viewpoint, The University Press of Virginia, Charlottesville, Virginia, 1969.
- [9] O. R. Musin, Around Sperner’s lemma, preprint, arXiv:1405.7513
- [10] O. R. Musin, Homotopy invariants of covers and KKM–type lemmas, *Algebraic & Geometric Topology*, **16:3** (2016) 1799–1812.
- [11] O. R. Musin, Homotopy groups and quantitative Sperner–type lemma, arXiv:2007.08715
- [12] S. P. Novikov, The analytic generalized Hopf invariant. Many-valued functionals, *Russian Mathematical Surveys*, **39:5** (1984), 113–124.
- [13] Y. Saad, Iterative Methods for Sparse Linear Systems, SIAM, 2003.
- [14] N. E. Steenrod, Cohomology invariants of mappings, *Ann. of Math. (2)*, **50** (1949), 954–988.
- [15] D.T. Sullivan, Infinitesimal calculations in topology, *Inst. des Hautes et Sci. Publ. Math.*, **4** (1977), 269–331.
- [16] J. H. C. Whitehead, An expression of Hopf’s invariant as an integral, *Proc. Nat. Acad. Sci. USA*, **33**, (1947), 117–123.
- [17] H. Whitney, Geometric Integration Theory, Princeton University Press, 1957

O. R. Musin, University of Texas Rio Grande Valley, School of Mathematical and Statistical Sciences, One West University Boulevard, Brownsville, TX, 78520, USA.

*E-mail address:* oleg.musin@utrgv.edu

T. Shamazov, Saint Petersburg State University, Department of Mathematics and Computer Science

*E-mail address:* shamazovt@inbox.ru