

UNIQUENESS PROBLEM FOR ACCRETIVE SCHRÖDINGER OPERATORS WITH COMPLEX SINGULAR COEFFICIENTS

VLADIMIR MIKHAIETS AND VOLODYMYR MOLYBOGA

ABSTRACT. The paper studies the uniqueness problem for the one-dimensional Schrödinger operator associated with the formal differential expression

$$l[u] = -u'' + qu + i[(ru)' + ru'],$$

in the complex Hilbert space $L^2(\mathbb{R})$. The coefficients of the expression are complex-valued and satisfy

$$q = s + Q', \quad s \in L^1_{loc}(\mathbb{R}) \quad \text{and} \quad Q, r \in L^2_{loc}(\mathbb{R}),$$

where the derivative is understood in the sense of distributions. In particular, the potential q can be a Radon measure on the line. With the help of specially selected quasi-derivatives, the expression l is treated as quasi-differential. The domains of the minimal L_0 and maximal L operators associated with the expression l in the space $L^2(\mathbb{R})$ are described. We find constructive conditions on the behaviour of $\text{Im } r$ near $\pm\infty$ that guarantee that $L_0 = L$ if the operator L_0 is accretive.

1. INTRODUCTION

Let H be a complex Hilbert space with scalar product (\cdot, \cdot) . A linear operator A in the space H is called accretive [14, 28] if

$$\text{Re}(Ax, x) \geq 0, \quad \forall x \in \text{Dom}(A),$$

and maximal accretive if it has no nontrivial accretive extensions in the Hilbert space H . An operator A is called m -accretive if the left complex half-plane belongs to the resolvent set $\rho(A)$ of A . Such operators play an important role in the theory of semigroups because the condition that the strongly continuous operator semigroup e^{-At} , $t \geq 0$, is contractive is equivalent to the m -accretivity of the operator A . The class of m -accretive operators is invariant with respect to the mapping $A \mapsto A^*$ [14, Problem V-3.31].

It follows from Phillips' theorem [23, Theorem 1.1.1] that a maximal accretive operator is either m -accretive or nondensely defined and nonclosable. Therefore, every densely defined accretive operator has a closure. A densely defined operator A in H is symmetric (self-adjoint) if and only if the operators iA and $-iA$ are accretive (m -accretive). Thus, the problem of the m -accretivity of a linear operator encompasses the problem of the self-adjointness of a symmetric operator.

The problem of self-adjointness of symmetric Schrödinger operators in Hilbert spaces $L^2(\mathbb{R}^n)$, $n \in \mathbb{N}$, is inspired by problems of quantum mechanics. It has a long history and a huge bibliography (see monographs [25, 1, 36, 37, 6] and works [19, 30, 2, 12, 9, 15] and the references therein). The obtained results are, in many cases, complete and form the basis of the spectral analysis of such operators and the scattering theory. In mathematical models of real physical processes in strongly inhomogeneous media, differential operators with strongly singular coefficients naturally arise. As a rule, they can be interpreted as distributions from certain classes.

The study of properties of such operators is a complex mathematical problem due to difficulties of a fundamental character. They naturally arise even in the problem of the proper definition of such operators. Such a definition needs a description of functions on which the formal differential expression is given, and also the mapping set by the expression.

In this paper, we introduce and study the class of Schrödinger operators in the complex Hilbert space $L^2(\mathbb{R})$, which are associated with the formal differential expression

$$(*) \quad l[u] := -u'' + au' + bu,$$

with complex-valued coefficients a and b satisfying the following conditions:

$$b = s + B', \quad s \in L^1_{loc}(\mathbb{R}) \quad B, a \in L^2_{loc}(\mathbb{R}),$$

2010 *Mathematics Subject Classification.* 34B20, 34B24, 34L40.

Key words and phrases. Hilbert space, Schrödinger operator, accretive operator, singular coefficients, quasi-differential operator, uniqueness problem.

with the derivative being understood in the sense of distributions.

Using the substitutions

$$r := \frac{1}{2i}a, \quad Q := B - ir,$$

the formal differential expression $(*)$ can be written in the more convenient form

$$(1) \quad l[u] = -u'' + qu + i[(ru)' + ru'],$$

where the complex-valued coefficients q and r satisfy the conditions

$$(2) \quad q = s + Q', \quad s \in L_{loc}^1(\mathbb{R}) \quad \text{and} \quad Q, r \in L_{loc}^2(\mathbb{R}).$$

From now on, we assume without loss of generality that the formal differential expression becomes the form (1), and its coefficients satisfy conditions (2).

The article is structured as follows. Section 2 gives the regularization of the formal differential expression (1) using properly selected quasi-derivatives and describes the maximal, pre-minimal, and minimal operators associated with expressions l and l^+ in the Hilbert space $L^2(\mathbb{R})$. Section 3 presents Theorem 2, which contains sufficient conditions for the m -accretivity of minimal operators for l and l^+ , provided that the pre-minimal operators in the space $L^2(\mathbb{R})$ are accretive. Section 4 contains Theorem 3, which gives other conditions for the m -accretivity of minimal operators. These conditions are imposed only on the behaviour of $\text{Im } r$ on some sequences of finite intervals that tend to $\pm\infty$ as $n \rightarrow \pm\infty$.

2. 1D SCHRÖDINGER OPERATORS WITH SINGULAR COEFFICIENTS

Let us consider the differential expression (1) with complex-valued coefficients q and r on the line \mathbb{R} . If $q \in L_{loc}^2(\mathbb{R})$ and $r \in W_{2,loc}^1(\mathbb{R})$, then the differential expression l is well posed on the set $C_{comp}^\infty(\mathbb{R})$ of infinitely smooth compactly supported functions on \mathbb{R} (this set is dense in $L^2(\mathbb{R})$). Therefore, with expression (1), we associate the pre-minimal operator in the Hilbert space $L^2(\mathbb{R})$:

$$L_{00} : u \rightarrow l[u], \quad \text{Dom}(L_{00}) = C_{comp}^\infty(\mathbb{R}),$$

and the maximal operator

$$L : u \rightarrow l[u], \quad \text{Dom}(L) = \{u \in L^2(\mathbb{R}) \cap W_{2,loc}^2(\mathbb{R}) \mid l[u] \in L^2(\mathbb{R})\}.$$

It is clear that $L_{00} \subset L$. The operator L is closed in the space $L^2(\mathbb{R})$ [4], therefore, the operator L_{00} is closable. Its closure $\tilde{L}_{00} := L_0$ is called the minimal operator associated with expression (1). The formally adjoint differential expression to l becomes

$$l^+[v] = -v'' + \bar{q}v + i[(\bar{r}v)' + \bar{r}v'],$$

where the overline denotes complex conjugation of the coefficients. Therefore, if the coefficients q and r in (1) are real-valued, then $l = l^+$ and the operators L_{00} and L_0 are symmetric. As is known, in this case (see [20, 28])

$$L = L_0^* = L_{00}^*.$$

Similarly, we define the operators L_{00}^+ , L_0^+ and L^+ generated by the differential expression l^+ in the space $L^2(\mathbb{R})$.

The operators L_{00} , L_0 , and L can also be defined in the case where $q \in L_{loc}^1(\mathbb{R})$ and $r \in AC_{loc}(\mathbb{R})$. They also act on the functions as a differential expression l [20, 32, 8].

In this paper, we investigate the case when the coefficients of the expression l are complex-valued and satisfy (2). In particular, if the function Q has locally bounded variation, then the coefficient q is a complex-valued Radon measure on a locally compact space \mathbb{R} . Schrödinger operators with such potentials appear in many problems of modern mathematical physics (see, for example, [1, 2, 9, 18] and the bibliography given there). In this case, first of all, we should study the question about a reasonable definition of the operators generated by the formal differential expression l . The most natural, in our opinion, is the definition based on the regularization of the singular differential expression with the help of the Shin-Zettl quasi-derivatives [29, 35, 10]. The most studied here is the case where $r \equiv 0$ and $q = \bar{q}$, see [2, 12, 17, 9] and references therein. The more general case is considered in [16]. In this case, the operator $L_0 = L_0^+$ is symmetric and $L = L_0^* = (L_0^+)^*$.

In this paper, we investigate the general case of singular complex-valued coefficients based on the regularization of the formal differential expression l using quasi-derivatives.

Let us put

$$G_1 := Q + ir \quad \text{and} \quad G_2 := Q - ir.$$

Then expression (1) can be presented in a more convenient form

$$(3) \quad l[u] = -u'' + (G_1 u)' - G_2 u' + su$$

since

$$(G_1 u)' - G_2 u' = Q' u + i[(ru)' + ru'].$$

Under conditions (2), the Shin-Zettl matrix function is well posed $A(\cdot) \in L_{loc}^1$:

$$(4) \quad A(\cdot) = \begin{pmatrix} G_1 & 1 \\ -G_1 G_2 + s & -G_2 \end{pmatrix}.$$

The Shin-Zettl matrix for l defines the quasi-derivatives as follows [29, 35, 10]:

$$(5) \quad u^{[0]} := u, \quad u^{[1]} := u' - G_1 u, \quad u^{[2]} := \left(u^{[1]}\right)' + G_2 u^{[1]} + (G_1 G_2 - s) u.$$

Using quasi-derivatives (5), we properly define the formal Schrödinger differential expression (1) as a quasi-differential one by the formulas:

$$l[u] := -u^{[2]} \quad \text{with} \quad \text{Dom}(l) := \left\{ u : \mathbb{R} \rightarrow \mathbb{C} \mid u, u^{[1]} \in \text{AC}_{loc}(\mathbb{R}) \right\}.$$

Now, in the complex Hilbert space $L^2(\mathbb{R})$, we define the Schrödinger operator L generated by formal differential expression (1) as a maximal quasi-differential operator

$$Lu := l[u], \quad \text{Dom}(L) := \left\{ u \in L^2(\mathbb{R}) \mid u, u^{[1]} \in \text{AC}_{loc}(\mathbb{R}), l[u] \in L^2(\mathbb{R}) \right\}.$$

The restriction of the maximal operator L to compactly supported functions generates a pre-minimal operator:

$$L_{00}u := Lu, \quad \text{Dom}(L_{00}) := \{u \in \text{Dom}(L) \mid \text{supp } u \Subset \mathbb{R}\}.$$

Shin-Zettl matrix (4) defines the formally Lagrange adjoint quasi-differential expression l^+ as follows [29, 35, 10]:

$$(6) \quad v^{\{0\}} := v, \quad v^{\{1\}} := v' - \overline{G_2} v, \quad v^{\{2\}} := \left(v^{\{1\}}\right)' + \overline{G_1} v^{\{1\}} + (\overline{G_1 G_2} - \overline{s}) v,$$

and

$$l^+[v] := -v^{\{2\}}, \quad \text{Dom}(l^+) := \left\{ v : \mathbb{R} \rightarrow \mathbb{C} \mid v, v^{\{1\}} \in \text{AC}_{loc}(\mathbb{R}) \right\}.$$

The quasi-differential expression l^+ generates the following maximal and pre-minimal operators:

$$L^+ v := l^+[v], \quad \text{Dom}(L^+) := \left\{ v \in L^2(\mathbb{R}) \mid v, v^{\{1\}} \in \text{AC}_{loc}(\mathbb{R}), l^+[v] \in L^2(\mathbb{R}) \right\},$$

and

$$L_{00}^+ v := l^+[v], \quad \text{Dom}(L_{00}^+) := \{v \in \text{Dom}(L^+) \mid \text{supp } v \Subset \mathbb{R}\}.$$

Let us introduce the notation

$$\begin{aligned} [u, v](x) &:= u(x) \overline{v^{\{1\}}(x)} - u^{[1]}(x) \overline{v(x)}, \\ [u, v]_a^b &:= [u, v](b) - [u, v](a), \quad -\infty \leq a < b \leq \infty. \end{aligned}$$

The main result of this section is the following theorem.

Theorem 1. *The following statements hold:*

- 1⁰. *The pre-minimal operators L_{00} and L_{00}^+ are densely defined in the Hilbert space $L^2(\mathbb{R})$.*
- 2⁰. *$(L_{00})^* = L^+$ and $(L_{00}^+)^* = L$.*
- 3⁰. *The operators L , L^+ are closed, and the pre-minimal operators L_{00} , L_{00}^+ admit a closure.*
- 4⁰. *Domains of the minimal operators L_0 , L_0^+ allow the following descriptions:*

$$\text{Dom}(L_0) = \left\{ u \in \text{Dom}(L) \mid [u, v]_{-\infty}^\infty = 0 \quad \forall v \in \text{Dom}(L^+) \right\}$$

and

$$\text{Dom}(L_0^+) = \left\{ v \in \text{Dom}(L^+) \mid [u, v]_{-\infty}^\infty = 0 \quad \forall u \in \text{Dom}(L) \right\},$$

with

- i) $\text{Dom}(L) \subset W_{2,loc}^1(\mathbb{R})$ and $\text{Dom}(L^+) \subset W_{2,loc}^1(\mathbb{R})$,
- ii) $\text{Dom}(L_{00}) \subset W_{2,comp}^1(\mathbb{R})$ and $\text{Dom}(L_{00}^+) \subset W_{2,comp}^1(\mathbb{R})$.

5⁰. The quadratic form associated with the pre-minimal operator is expressed as follows

$$(\mathbf{L}_{00}u, u)_{L^2(\mathbb{R})} = \int_{\mathbb{R}} |u'|^2 dx - \int_{\mathbb{R}} (G_1 u \bar{u}' + G_2 u' \bar{u}) dx + \int_{\mathbb{R}} s |u|^2 dx, \quad u \in \text{Dom}(\mathbf{L}_{00}).$$

The proof of Theorem 1 relies on the following statements.

Lemma 1 (Generalized Lagrange identity [35, Corollary 1]). *For arbitrary functions $u \in \text{Dom}(\mathbf{L})$ and $v \in \text{Dom}(\mathbf{L}^+)$, the following relation holds:*

$$(7) \quad \int_a^b l[u] \bar{v} dx - \int_a^b \overline{ul^+[v]} dx = [u, v]_a^b$$

whenever $-\infty < a < b < +\infty$.

Lemma 2. *For arbitrary functions $u \in \text{Dom}(\mathbf{L})$ and $v \in \text{Dom}(\mathbf{L}^+)$ the following finite limits exist:*

$$[u, v](-\infty) := \lim_{x \rightarrow -\infty} [u, v](x), \quad [u, v](\infty) := \lim_{x \rightarrow \infty} [u, v](x).$$

Proof. Let us fix b in relation (7), and proceed to the limit at $a \rightarrow -\infty$. Since $u, v, l[u], l^+[v] \in L^2(\mathbb{R})$ by the assumptions of the lemma, the limit $[u, v](-\infty)$ exists and is finite. The existence and finiteness of the limit $[u, v](\infty)$ are proved similarly. \square

Combining Lemmas 1 and 2 yields the following integral identity.

Statement 1. *For arbitrary $u \in \text{Dom}(\mathbf{L})$ and $v \in \text{Dom}(\mathbf{L}^+)$, the following identity holds:*

$$(8) \quad \int_{-\infty}^{\infty} l[u] \bar{v} dx - \int_{-\infty}^{\infty} \overline{ul^+[v]} dx = [u, v]_{-\infty}^{\infty},$$

where the limits

$$[u, v](-\infty) := \lim_{x \rightarrow -\infty} [u, v](x), \quad [u, v](\infty) := \lim_{x \rightarrow \infty} [u, v](x),$$

exist and are finite.

Let $\Delta = (a, b)$ with $-\infty < a < b < \infty$ be a finite interval on \mathbb{R} .

In the complex Hilbert space $L^2(\Delta)$, let us consider the operators generated by the formal Schrödinger differential expression (1). Basic assumptions (2) regarding the coefficients on a finite interval take the form

$$q = s + Q', \quad Q, r \in L^2(\Delta) \quad \text{and} \quad s \in L^1(\Delta).$$

Therefore, the corresponding Shin-Zettl matrix-valued function (4) is well posed, and the corresponding quasi-derivatives (5) and (6) are well defined.

Let us consider the following quasi-differential expressions l and l^+ on a finite interval Δ :

$$l_{\Delta}[u] := -u^{[2]}, \quad \text{Dom}(l_{\Delta}) := \left\{ u : \overline{\Delta} \rightarrow \mathbb{C} \mid u, u^{[1]} \in \text{AC}(\overline{\Delta}) \right\},$$

and

$$l_{\Delta}^+[v] := -v^{\{2\}}, \quad \text{Dom}(l_{\Delta}^+) := \left\{ v : \overline{\Delta} \rightarrow \mathbb{C} \mid v, v^{\{1\}} \in \text{AC}(\overline{\Delta}) \right\}.$$

They naturally generate the minimal and maximal operators in the space $L^2(\Delta)$:

$$\mathbf{L}_{\Delta} u := l_{\Delta}[u], \quad \text{Dom}(\mathbf{L}_{\Delta}) := \left\{ u \in L^2(\Delta) \mid u, u^{[1]} \in \text{AC}(\overline{\Delta}), l_{\Delta}[u] \in L^2(\Delta) \right\},$$

$$\mathbf{L}_{0\Delta} u := \mathbf{L}_{\Delta} u, \quad \text{Dom}(\mathbf{L}_{0\Delta}) := \left\{ u \in \text{Dom}(\mathbf{L}_{\Delta}) \mid u^{[j]}(a) = u^{[j]}(b) = 0, j = 0, 1 \right\},$$

and

$$\mathbf{L}_{\Delta}^+ v := l_{\Delta}^+[v], \quad \text{Dom}(\mathbf{L}_{\Delta}^+) := \left\{ v \in L^2(\Delta) \mid v, v^{\{1\}} \in \text{AC}(\overline{\Delta}), l_{\Delta}^+[v] \in L^2(\Delta) \right\},$$

$$\mathbf{L}_{0\Delta}^+ v := \mathbf{L}_{\Delta}^+ v, \quad \text{Dom}(\mathbf{L}_{0\Delta}^+) := \left\{ v \in \text{Dom}(\mathbf{L}_{\Delta}^+) \mid v^{\{1\}}(a) = v^{\{1\}}(b) = 0, j = 0, 1 \right\}.$$

Let us recall a well-known result.

Statement 2 ([35, Theorem 10, 11]). *The minimal operators $\mathbf{L}_{0\Delta}$ and $\mathbf{L}_{0\Delta}^+$ are densely defined in the Hilbert space $L^2(\Delta)$, and the following statements hold:*

- (a) $(\mathbf{L}_{0\Delta})^* = \mathbf{L}_{\Delta}^+$, (c) $\mathbf{L}_{0\Delta} = (\mathbf{L}_{\Delta}^+)^*$,
- (b) $(\mathbf{L}_{0\Delta}^+)^* = \mathbf{L}_{\Delta}$, (d) $\mathbf{L}_{0\Delta}^+ = (\mathbf{L}_{\Delta})^*$.

Proof of Theorem 1. The proof of the properties $1^0 - 4^0$ is similar to the case of the semi-axis and formally self-adjoint differential expressions $l = l^+$ [35, 20]. In their proof, the properties of operators defined on a finite interval are used. For the reader's convenience, we will give the proof.

1^0 . Let us prove the density of the domain of the pre-minimal operator $\text{Dom}(L_{00})$ in the Hilbert space $L^2(\mathbb{R})$.

Suppose that a function $h \in L^2(\mathbb{R})$ is orthogonal to $\text{Dom}(L_{00})$, and show that $h \equiv 0$, which will mean the density of $\text{Dom}(L_{00})$ in the space $L^2(\mathbb{R})$.

Let Δ be an arbitrary fixed finite interval. The quasi-differential expression l defines operators $L_{0\Delta}$ and L_Δ . The space $L^2(\Delta)$ can be embedded in the space $L^2(\mathbb{R})$ if we assume that outside the interval $\overline{\Delta}$, the function $u \in L^2(\Delta)$ is zero. In this way, the domain $\text{Dom}(L_{0\Delta})$ of the operator $L_{0\Delta}$ becomes part of $\text{Dom}(L)$. Moreover, the function thus extended belongs to $\text{Dom}(L_{00})$.

Thus, the function h is orthogonal to $\text{Dom}(L_{0\Delta})$. Then, by Statement 2, the set $\text{Dom}(L_{0\Delta})$ is dense in the space $L^2(\Delta)$. Therefore, $h|_{\overline{\Delta}} = 0$ almost everywhere.

Since the interval $\Delta \subset \mathbb{R}$ is chosen arbitrarily, we have $h = 0$ almost everywhere on \mathbb{R} , which is equivalent to the density of $\text{Dom}(L_{00})$ in $L^2(\mathbb{R})$.

The density of the domain of the pre-minimal operator $\text{Dom}(L_{00}^+)$ in $L^2(\mathbb{R})$ is proved similarly.

2^0 . Let us prove the first relation

$$(L_{00})^* = L^+.$$

The second relation is proved similarly.

It follows from 1^0 that for the operator L_{00} , there exists the adjoint operator $(L_{00})^*$.

Let $u \in \text{Dom}(L_{00})$ and $v \in \text{Dom}(L^+)$. Then, using the generalized Lagrange identity (8), we obtain

$$(L_{00}u, v)_{L^2(\mathbb{R})} = (u, L^+v)_{L^2(\mathbb{R})}.$$

Therefore $L^+ \subset (L_{00})^*$. Let us prove the inverse inclusion.

Let v be an arbitrary element of $\text{Dom}((L_{00})^*)$, and let $\Delta = (\alpha, \beta)$ be a fixed finite interval on \mathbb{R} . Then

$$((L_{00})^*v, u)_{L^2(\mathbb{R})} = (v, L_{00}u)_{L^2(\mathbb{R})} \quad \text{for every } u \in \text{Dom}(L_{0\Delta}).$$

Since $u|_{\mathbb{R} \setminus \overline{\Delta}} = 0$, the scalar products are represented as integrals over the interval $\overline{\Delta}$, that is, they are scalar products in $L^2(\Delta)$, and

$$(9) \quad (((L_{00})^*v)_\Delta, u)_{L^2(\Delta)} = (v_\Delta, L_{0\Delta}u)_{L^2(\Delta)} \quad \text{for every } u \in \text{Dom}(L_{0\Delta}).$$

Here $((L_{00})^*v)_\Delta$ and v_Δ are restrictions of the functions to the interval $\overline{\Delta}$.

Using (9) and applying the generalized Lagrange identity (7), we obtain

$$((L_{00})^*v)_\Delta = L_\Delta^+ v_\Delta = (l^+[v])_\Delta.$$

Since the interval Δ can be chosen arbitrarily, then

$$v \in \text{Dom}(L^+) \quad \text{and} \quad (L_{00})^*v = l^+[v] = L^+v,$$

which is what had to be proved.

3^0 . This statement is a direct consequence of property 2^0 .

4^0 . Since the operator L_0 is closed, and because

$$(L_0)^* = L^+ \quad \text{and} \quad L_0 = (L^+)^*,$$

it is obvious that the set $\text{Dom}(L_0)$ consists of those and only those functions $u \in \text{Dom}(L)$ that satisfy the relation

$$(u, L^+v)_{L^2(\mathbb{R})} = (Lu, v)_{L^2(\mathbb{R})} \quad \text{for every } v \in \text{Dom}(L^+),$$

which, taking into account the generalized Lagrange identity on \mathbb{R} , is equivalent to statement 4^0 .

5⁰. By assumptions (2), we have $G_1, G_2 \in L^2_{loc}(\mathbb{R})$ and $\text{Dom}(L_{00}) \subset W^1_{2,comp}(\mathbb{R})$. Hence, integrating by parts, we obtain for any $u \in \text{Dom}(L_{00})$ the following:

$$\begin{aligned} (L_{00}u, u)_{L^2(\mathbb{R})} &= \int_{\mathbb{R}} \left(-(u^{[1]})' \bar{u} - G_2 u^{[1]} \bar{u} - G_1 G_2 |u|^2 + s |u|^2 \right) dx \\ &= \int_{\mathbb{R}} \left((u^{[1]} \bar{u})' - G_2 u^{[1]} \bar{u} - G_1 G_2 |u|^2 + s |u|^2 \right) dx \\ &= \int_{\mathbb{R}} \left((u' - G_1 u) \bar{u}' - G_2 (u' - G_1 u) \bar{u} - G_1 G_2 |u|^2 + s |u|^2 \right) dx \\ &= \int_{\mathbb{R}} \left(|u'|^2 - (G_1 u \bar{u}' + G_2 u' \bar{u}) + s |u|^2 \right) dx. \end{aligned}$$

The proof of Theorem 1 is complete. \square

3. UNIQUENESS PROBLEM

The uniqueness problem for a differential expression l in the Hilbert space $L^2(\mathbb{R})$ consists in finding conditions on its coefficients q and r under which the equality $L_0 = L$ holds. If these functions are real-valued, then the operator L_0 is symmetric, and $L = L_0^*$. We thus deal with the self-adjointness conditions for the symmetric operator L_0 in the Hilbert space $L^2(\mathbb{R})$. In the case when $r \equiv 0$ and $q \in L^1_{loc}(\mathbb{R})$, this topic was studied in detail by many analysts. In particular, H. Weyl [33] was the first to prove the self-adjointness of the operator L_0 in the case where $r \equiv 0$ and when the function q is continuous and bounded from below on \mathbb{R} . It is easy to see that the operator L_0 is bounded below in this case. In this connection, the hypothesis arose that this abstract condition already guarantees the self-adjointness of the operator L_0 . It turned out to be true even for the more general Sturm–Liouville differential expression

$$l[u] = -(pu')' + qu + i[(ru)' + ru'].$$

Hartman [11] and Rellich [26] showed that $L_0 = L$ if

$$\int_0^\infty p^{-1/2}(t) dt = \int_{-\infty}^0 p^{-1/2}(t) dt = \infty,$$

$r \equiv 0$, $0 < p \in C^2(\mathbb{R})$ and the function q is piecewise continuous on \mathbb{R} . Under more general assumptions

$$q = Q' + s, \quad \frac{1}{\sqrt{|p|}}, \frac{Q}{\sqrt{|p|}}, \frac{r}{\sqrt{|p|}} \in L^2_{loc}(\mathbb{R}), \quad s \in L^1_{loc}(\mathbb{R}), \quad 0 < p \in W^1_{2,loc}(\mathbb{R}),$$

this fact was proven by the authors in [16].

Similar theorems have been proved for second-order elliptic differential operators in the spaces $L^2(\mathbb{R}^n)$, with $n \in \mathbb{N}$, and other classes of operators (see, for example, [21, 24, 30, 31, 34] and the references therein).

The case where the symmetric operator L_0 is not bounded from below was also studied quite thoroughly. In particular, Sears [27] showed that if $r \equiv 0$, $q \in C(\mathbb{R})$, and if there exists an even continuous function $Q(\cdot) \geq \delta > 0$ on \mathbb{R} such that

$$q(x) \geq -Q(x), \quad \int_0^\infty Q^{-1/2}(x) dx = \infty,$$

furthermore, either the function Q is monotone on $[0; \infty)$ or

$$\limsup_{x \rightarrow \infty} \frac{Q'(x)}{\{Q(x)\}^{3/2}} < \infty,$$

then the operator L_0 is self-adjoint. See also papers [11, 13, 22] and references therein.

The case when the coefficients of the differential expression l are complex-valued is more complicated because the operator L_0 in general is non-symmetric. The question about the conditions for its m -accretivity, to the best of the authors' knowledge, has not been previously studied, even in the case of continuous coefficients q and r .

In this paper, we investigate the case when these coefficients are singular and satisfy assumption (2). To this end, we develop a methodology of [16], where symmetric Sturm–Liouville operators in $L^2(\mathbb{R})$ were studied. The main result of this section is Theorem 2. Its proof is based on the results of Section 2.

Let us introduce some notation. Given a real-valued function $f \in L^1_{loc}(\mathbb{R})$, we let

$$f^+ := \frac{|f| + f}{2} \geq 0, \quad f^- := \frac{|f| - f}{2} \geq 0, \quad f = f^+ - f^-,$$

$$r = r_0 + ir_1, \quad \text{where} \quad r_0 := \operatorname{Re} r, \quad r_1 := \operatorname{Im} r.$$

Theorem 2. *Let the coefficients of the formal differential Schrödinger expression (1) satisfy conditions (2), and suppose that the pre-minimal operators L_{00} and L_{00}^+ are accretive and*

$$r_1^\pm(x) = O(m(x)) \quad \text{as} \quad x \rightarrow \mp\infty,$$

where the function $m(x)$ satisfies

$$(m_1) \quad 1 \leq m(x) \in W^1_{2,loc}(\mathbb{R}),$$

$$(m_2) \quad \int_{-\infty}^0 \frac{ds}{m(s)} = \int_0^{+\infty} \frac{ds}{m(s)} = +\infty.$$

Then the minimal operators L_0 and L_0^* are m -accretive, and $L_0 = L$ and $L_0^* = L^+$.

Remark 1. In the case where $q \in L^2_{loc}(\mathbb{R})$ and $r \in W^1_{2,loc}(\mathbb{R})$, the operators L_{00} and L_{00}^+ are accretive if and only if both are accretive. Therefore, the accretivity assumption for L_{00}^+ in Theorem 2 can be removed in this case. The question about whether this can be done in the general case remains open.

Corollary 1. *Let Λ be an arbitrary closed sector of the half-plane $\operatorname{Re} \lambda \geq 0$ with vertex at the point 0, and let assumptions (m_1) , (m_2) of Theorem 2 be satisfied. If*

$$(L_{00}u, u)_{L^2(\mathbb{R})} \in \Lambda \quad \text{for every} \quad u \in \operatorname{Dom}(L_{00}), \quad \text{and if} \quad (L_{00}^+v, v)_{L^2(\mathbb{R})} \in \overline{\Lambda} \quad \text{for every} \quad v \in \operatorname{Dom}(L_{00}^+),$$

then $\rho(L_0) \supset \mathbb{C} \setminus \Lambda$ and $\rho(L_0^+) \supset \mathbb{C} \setminus \overline{\Lambda}$.

First, we will prove some necessary technical results.

Lemma 3. *Let initial assumptions (2) be satisfied. Then, for an arbitrary compactly supported function $\varphi \in W^2_{2,comp}(\mathbb{R})$ and for all $u \in \operatorname{Dom}(L)$ and $v \in \operatorname{Dom}(L^+)$, the following statements hold:*

- (i) $\varphi u \in \operatorname{AC}_{comp}(\mathbb{R})$,
 $\varphi v \in \operatorname{AC}_{comp}(\mathbb{R})$;
- (ii) $(\varphi u)^{[1]} = \varphi' u + \varphi u^{[1]} \in \operatorname{AC}_{comp}(\mathbb{R})$,
 $(\varphi v)^{[1]} = \varphi' v + \varphi v^{\{1\}} \in \operatorname{AC}_{comp}(\mathbb{R})$;
- (iii) $l[\varphi u] = \varphi l[u] - \varphi'' u - 2\varphi' u' + (G_1 - G_2) \varphi' u \in L^2_{comp}(\mathbb{R})$,
 $l^+[\varphi v] = \varphi l^+[v] - \varphi'' v - 2\varphi' v' + (\overline{G_2} - \overline{G_1}) \varphi' v \in L^2_{comp}(\mathbb{R})$;
- (iv) $\operatorname{supp} \varphi u$ is a compact set,
 $\operatorname{supp} \varphi v$ is a compact set,

and

$$\varphi u \in \operatorname{Dom}(L_{00}) \subset W^1_{2,comp}(\mathbb{R}) \quad \text{and} \quad \varphi v \in \operatorname{Dom}(L_{00}^+) \subset W^1_{2,comp}(\mathbb{R}).$$

In particular, if $a' < a < b < b'$ and if the function $\varphi \in W^2_{2,comp}(\mathbb{R})$ satisfy the conditions

- (a) $0 \leq \varphi(x) \leq 1, \quad x \in \mathbb{R}$,
- (b) $\varphi(x) = \begin{cases} 1, & x \in [a; b]; \\ 0, & x \in (\infty; a') \cup (b'; \infty), \end{cases}$

then the conclusions of the lemma hold.

Proof. It can be proved by direct calculations. □

Proof of Theorem 2. Let the assumptions of the theorem be satisfied. Without loss of generality, we assume that

$$(10) \quad \operatorname{Re} (L_{00}u, u)_{L^2(\mathbb{R})} \geq (u, u)_{L^2(\mathbb{R})} \quad \text{for every} \quad u \in \operatorname{Dom}(L_{00}),$$

and that

$$(11) \quad \operatorname{Re} (v, L_{00}^+ v)_{L^2(\mathbb{R})} \geq (v, v)_{L^2(\mathbb{R})} \quad \text{for every } v \in \operatorname{Dom}(L_{00}^+).$$

To prove that the minimal operator L_0 is m -accretive, it suffices to show that the null space $\mathcal{N}(L^+)$ of the maximal operator L^+ contains only the zero element, namely

$$\dim(\mathcal{N}(L^+)) = \dim(\operatorname{rank} L_0)^\perp = \operatorname{def} L_0 = 0.$$

Then, an arbitrary point of the half-plane $\operatorname{Rm} \lambda < 1$ is also regular for the operator L_0 because the defect numbers of the closed operator are constant in each connected component of a set of points of regular type of this operator (see, for example, [5] and [14, Theorem 3.2]).

Let $v \in L^2(\mathbb{R})$ be a solution of the equation

$$(12) \quad L^+ v = 0.$$

Let us show that $v \equiv 0$.

Lemma 4. *Let the general assumptions (2) be satisfied, and let $v \in L^2(\mathbb{R})$ be a solution of the equation*

$$L^+ v = 0.$$

Then, for an arbitrary compactly supported function $\varphi \in W_{2,comp}^2(\mathbb{R})$ with $\operatorname{Im} \varphi = 0$ the following inequality holds:

$$(13) \quad \operatorname{Re} (\varphi v, L_{00}^+ \varphi v)_{L^2(\mathbb{R})} = \int_{\mathbb{R}} (\varphi')^2 |v|^2 dx + 2 \int_{\mathbb{R}} r_1 \varphi' \varphi |v|^2 dx \geq \int_{\mathbb{R}} \varphi^2 |v|^2 dx.$$

Proof. Given a function $\varphi \in W_{2,comp}^2(\mathbb{R})$ with $\operatorname{Im} \varphi = 0$ and taking into account that

$$\varphi v \in \operatorname{Dom}(L_{00}^+) \subset W_{2,comp}^1(\mathbb{R}),$$

and that $L^+[v] = 0$, we obtain by direct calculations (see Lemma 3) the following:

$$(\varphi v, L_{00}^+ \varphi v)_{L^2(\mathbb{R})} = (\varphi v, L^+[\varphi v])_{L^2(\mathbb{R})} = \int_{\mathbb{R}} (-\varphi'' \varphi |v|^2 - 2\varphi' \varphi v \bar{v}' + (G_2 - G_1) \varphi' \varphi |v|^2) dx.$$

Then, applying the integration by parts formula to the first term (recall that the function φ has compact support), we obtain

$$\begin{aligned} (\varphi v, L_{00}^+ \varphi v)_{L^2(\mathbb{R})} &= \int_{\mathbb{R}} (\varphi' \varphi' |v|^2 + \varphi' \varphi v \bar{v}' + \varphi' \varphi v \bar{v}' - 2\varphi' \varphi v \bar{v}' + (G_2 - G_1) \varphi' \varphi |v|^2) dx \\ &= \int_{\mathbb{R}} ((\varphi')^2 |v|^2 + \varphi' \varphi (v \bar{v}' - v \bar{v}') + (G_2 - G_1) \varphi' \varphi |v|^2) dx. \end{aligned}$$

Therefore, for any $\varphi \in W_{2,comp}^2(\mathbb{R})$ with $\operatorname{Im} \varphi = 0$ and every solution v of the equation $L^+ v = 0$, we have

$$(\varphi v, L_{00}^+ \varphi v)_{L^2(\mathbb{R})} = \int_{\mathbb{R}} (\varphi')^2 |v|^2 dx + \int_{\mathbb{R}} \varphi' \varphi (v \bar{v}' - v \bar{v}') dx - 2i \int_{\mathbb{R}} r \varphi' \varphi |v|^2 dx,$$

and (recall that $r = r_0 + ir_1$)

$$(14) \quad \begin{aligned} (\varphi v, L_{00}^+ \varphi v)_{L^2(\mathbb{R})} &= \int_{\mathbb{R}} (\varphi')^2 |v|^2 dx + 2 \int_{\mathbb{R}} r_1 \varphi' \varphi |v|^2 dx \\ &\quad + \int_{\mathbb{R}} \varphi' \varphi (v \bar{v}' - v \bar{v}') dx - 2i \int_{\mathbb{R}} r_0 \varphi' \varphi |v|^2 dx. \end{aligned}$$

Since

$$\operatorname{Re} \int_{\mathbb{R}} \varphi' \varphi (v \bar{v}' - v \bar{v}') dx = 0 \quad \text{and} \quad \operatorname{Re} 2i \int_{\mathbb{R}} r_0 \varphi' \varphi |v|^2 dx = 0,$$

equality (14) allows us to write condition (11) in the required form

$$\operatorname{Re} (\varphi v, L_{00}^+ \varphi v)_{L^2(\mathbb{R})} = \int_{\mathbb{R}} (\varphi')^2 |v|^2 dx + 2 \int_{\mathbb{R}} r_1 \varphi' \varphi |v|^2 dx \geq \int_{\mathbb{R}} \varphi^2 |v|^2 dx.$$

□

Let us consider the following sequence of intervals $\{\Delta_n\}_{n \in \mathbb{Z}}$:

$$\Delta_0 := [-1; 1], \quad \Delta_n := [n; n+1], \quad \Delta_{-n} := [-n-1; -n] \quad \text{for each } n \in \mathbb{N}.$$

Let $\{\varphi_n\}_{n \in \mathbb{N}}$ be a sequence of real-valued functions from $W_{2,comp}^2(\mathbb{R})$ that satisfy the following conditions:

- (a₁) $0 \leq \varphi_n(x) \leq 1, x \in \mathbb{R};$
- (a₂) $\varphi_n(x) = 1, x \in [-n; n];$
- (a₃) $\text{supp } \varphi_n \subset [-n-1; n+1];$
- (a₄) $\varphi'_n(x) \geq 0, x \in \Delta_{-n}$ and $\varphi'_n(x) \leq 0, x \in \Delta_n;$
- (a₅) there exists a $K > 0$ such that

$$|\varphi'_n(x)| \leq K \quad \text{for every } x \in \mathbb{R}.$$

We now substitute the functions φ_n into (13) and note that

$$\text{supp } \varphi'_n \subset \Delta_{-n} \cup \Delta_n, \quad n \in \mathbb{N}.$$

We then obtain

$$\begin{aligned}
\int_{-n}^n |v|^2 dx &\leq \int_{\mathbb{R}} \varphi_n^2 |v|^2 dx \leq \int_{\mathbb{R}} (\varphi'_n)^2 |v|^2 dx + 2 \int_{\mathbb{R}} r_1 \varphi'_n \varphi_n |v|^2 dx \\
&= \int_{\Delta_{-n} \cup \Delta_n} (\varphi'_n)^2 |v|^2 dx + 2 \int_{\Delta_{-n}} r_1 \varphi'_n \varphi_n |v|^2 dx + 2 \int_{\Delta_n} r_1 \varphi'_n \varphi_n |v|^2 dx \\
&\leq K^2 \int_{\Delta_{-n} \cup \Delta_n} |v|^2 dx + 2 \int_{\Delta_{-n}} (r_1^+ - r_1^-) \varphi'_n \varphi_n |v|^2 dx + 2 \int_{\Delta_n} (r_1^+ - r_1^-) \varphi'_n \varphi_n |v|^2 dx \\
&= K^2 \int_{\Delta_{-n} \cup \Delta_n} |v|^2 dx + 2 \int_{\Delta_{-n}} r_1^+ \varphi'_n \varphi_n |v|^2 dx - 2 \int_{\Delta_n} r_1^- \varphi'_n \varphi_n |v|^2 dx \\
&\quad - 2 \int_{\Delta_{-n}} r_1^- \varphi'_n \varphi_n |v|^2 dx + 2 \int_{\Delta_n} r_1^+ \varphi'_n \varphi_n |v|^2 dx \\
&\leq K^2 \int_{\Delta_{-n} \cup \Delta_n} |v|^2 dx + 2 \int_{\Delta_{-n}} r_1^+ \varphi'_n \varphi_n |v|^2 dx - 2 \int_{\Delta_n} r_1^- \varphi'_n \varphi_n |v|^2 dx.
\end{aligned}$$

Here, we used

$$-2 \int_{\Delta_{-n}} r_1^- \varphi'_n \varphi_n |v|^2 dx \leq 0 \quad \text{and} \quad 2 \int_{\Delta_n} r_1^+ \varphi'_n \varphi_n |v|^2 dx \leq 0.$$

Thus,

$$(15) \quad \int_{-n}^n |v|^2 dx \leq K^2 \int_{\Delta_{-n} \cup \Delta_n} |v|^2 dx + 2 \int_{\Delta_{-n}} r_1^+ \varphi'_n |v|^2 dx - 2 \int_{\Delta_n} r_1^- \varphi'_n |v|^2 dx.$$

According to assumptions (m₁) of Theorem 2, there exist numbers $N_0 \in \mathbb{N}$ and $C > 0$ such that

$$(16) \quad r_1^+(x) \leq Cm(x) \quad \text{for every } x \in (-\infty; -N_0],$$

$$(17) \quad r_1^-(x) \leq Cm(x) \quad \text{for every } x \in [N_0; \infty).$$

Then, taking into account estimates (16) and (17), for $n \geq N_0$, we deduce from inequality (15)

$$(18) \quad \int_{-n}^n |v|^2 dx \leq K^2 \int_{\Delta_{-n} \cup \Delta_n} |v|^2 dx + 2C \int_{\Delta_{-n}} m \varphi'_n |v|^2 dx - 2C \int_{\Delta_n} m \varphi'_n |v|^2 dx.$$

Let us consider the function

$$\rho(x) := \begin{cases} -\int_x^0 \frac{ds}{m(s)}, & x \in (-\infty; 0], \\ \int_0^x \frac{ds}{m(s)}, & x \in [0; \infty); \end{cases}$$

which satisfies

$$\rho(x) \rightarrow -\infty \quad \text{as } x \rightarrow -\infty, \quad \text{and} \quad \rho(x) \rightarrow +\infty \quad \text{as } x \rightarrow +\infty,$$

in view of condition (m_2) of Theorem 2. By condition (m_1) of Theorem 2, the derivative has the properties

$$\rho'(x) = \frac{1}{m(x)} > 0 \quad \text{for any } x \in \mathbb{R}, \quad \text{and} \quad \rho' \in W_{2,loc}^1(\mathbb{R}).$$

Then

$$(19) \quad (\varphi_n[\rho(x)])' = (\varphi_n[\rho])'_\rho (\rho(x))'_x = (\varphi_n[\rho])'_\rho \frac{1}{m(x)},$$

and

$$\varphi_n[\rho] \in W_{2,comp}^2(\mathbb{R}) \quad \text{for each } n \in \mathbb{Z}.$$

Let us put

$$\begin{aligned} X_n^- &:= \{x \in \mathbb{R} \mid \rho(x) \in \Delta_{-n}\} \quad \text{for each } n = 0, 1, 2, \dots, \\ X_n^+ &:= \{x \in \mathbb{R} \mid \rho(x) \in \Delta_n\} \quad \text{for each } n = 0, 1, 2, \dots, \end{aligned}$$

and

$$U_n := \bigcup_{k=0}^{n-1} X_n^- \cup X_n^+.$$

Note that under condition (m_2) of Theorem 2, the following relations hold:

$$(20) \quad \lim_{n \rightarrow -\infty} \sup X_n^- = -\infty,$$

$$(21) \quad \lim_{n \rightarrow \infty} \inf X_n^+ = \infty,$$

and

$$(22) \quad \lim_{n \rightarrow \infty} U_n = \lim_{n \rightarrow \infty} \bigcup_{k=0}^{n-1} X_n^- \cup X_n^+ = \mathbb{R}.$$

Let us substitute the functions $\varphi_n(\rho(\cdot))$ with $n \in \mathbb{Z}$ into (13). As when proving inequalities (15) and (18), we infer by (19) with $n \geq N_0$ that

$$\begin{aligned} \int_{U_n} |v(x)|^2 dx &\leq K^2 \int_{X_n^- \cup X_n^+} |v(x)|^2 dx + 2 \int_{X_n^-} m(x)(\varphi_n[\rho(x)])' |v(x)|^2 dx - 2 \int_{X_n^+} m(x)(\varphi_n[\rho(x)])' |v(x)|^2 dx \\ &\leq K^2 \int_{X_n^- \cup X_n^+} |v(x)|^2 dx + 2C \int_{X_n^-} m(x)(\varphi_n[\rho])'_\rho \rho'(x) |v(x)|^2 dx - 2C \int_{X_n^+} m(x)(\varphi_n[\rho])'_\rho \rho'(x) |v(x)|^2 dx \\ &\leq K^2 \int_{X_n^- \cup X_n^+} |v(x)|^2 dx + 2CK \int_{X_n^-} |v(x)|^2 dx + 2CK \int_{X_n^+} |v(x)|^2 dx. \end{aligned}$$

Thus,

$$(23) \quad \int_{U_n} |v(x)|^2 dx \leq K^2 \int_{X_n^- \cup X_n^+} |v(x)|^2 dx + 2CK \int_{X_n^- \cup X_n^+} |v(x)|^2 dx \quad \text{whenever } n \geq N_0.$$

Let us now pass to the limit as $n \rightarrow \infty$ in inequality (23). Since $v \in L^2(\mathbb{R})$ and because relations (20) – (22) hold, we obtain

$$\int_{X_n^- \cup X_n^+} |v(x)|^2 dx \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

and, finally, conclude that

$$\int_{\mathbb{R}} |v(x)|^2 dx \leq 0, \quad \text{i.e.,} \quad v(x) \equiv 0.$$

The proof is complete. □

4. INTERVAL-TYPE THEOREM

Since functions of class $W_{2,loc}^1(\mathbb{R})$ are continuous on the entire axis, then they are locally bounded. Therefore, it follows from condition (m_1) of Theorem 2 that the function r_1 belongs to $L_{loc}^\infty(\mathbb{R})$ and, unlike the function r_0 , cannot have local singularities. To include a wider class of admissible functions r_1 , we now present an alternative m -accretivity criterion for minimal operators. These conditions are imposed only on the behaviour of the function r_1 concerning only sequences of intervals in neighborhoods of $\pm\infty$, and there are no additional conditions outside these intervals.

Self-adjointness conditions of this type for Schrödinger operators of form (1) with $r \equiv 0$ and with a real continuous potential q on \mathbb{R} were first established in Ismagilov's paper [13]. They have the following form. Let Δ_n be a sequence of disjoint intervals of length $|\Delta_n|$ with $n \in \mathbb{Z}$, and let Δ_n tend to $\pm\infty$ as $n \rightarrow \pm\infty$. If

$$q(x) \geq -\gamma_n \quad (\gamma_n > 0) \quad \text{for every } x \in \Delta_n \quad \text{whatever } n \in \mathbb{Z},$$

and if

$$\Sigma_{n=-\infty}^{-1}(\gamma_n + |\Delta_n|^{-2})^{-1} = \Sigma_{n=1}^{\infty}(\gamma_n + |\Delta_n|^{-2})^{-1} = \infty,$$

then the minimal symmetric operator L_0 is self-adjoint.

These results were further developed in [22] and the publications cited therein.

The main result of this section is the following statement.

Theorem 3. *Let the coefficients of the formal differential Schrödinger expression (1) satisfy conditions (2), and let the pre-minimal operators L_{00} and L_{00}^+ be accretive. If there exists a sequence of intervals $\Delta_n := [a_n; b_n]$ with $n \in \mathbb{Z}$ such that*

$$-\infty < a_n < b_n < \infty, \quad b_n \rightarrow -\infty \quad \text{as } n \rightarrow -\infty, \quad \text{and} \quad a_n \rightarrow \infty \quad \text{as } n \rightarrow \infty,$$

and if

(B₁) *there exists a number $\delta > 0$ such that $|\Delta_n| \geq \delta$ for each $n \in \mathbb{N}$,*

(B₂) *there exists a number $C > 0$ such that*

$$\begin{aligned} r_1^+(x) &\leq C|\Delta_{-n}| \quad \text{for every } x \in \Delta_{-n}, \quad \text{whatever } n \in \mathbb{N}, \\ r_1^-(x) &\leq C|\Delta_n| \quad \text{for every } x \in \Delta_n, \quad \text{whatever } n \in \mathbb{N}; \end{aligned}$$

then minimal operators L_0 and L_0^* are m -accretive, and $L_0 = L$ and $L_0^* = L^+$.

Remark 2. It follows from Theorem 3 that if the assumptions given therein are fulfilled for the function r_1 and if

$$(L_{00}u, u) \in \Lambda \quad \text{for every } u \in \text{Dom}(L_{00}), \quad \text{and} \quad (L_{00}^+v, v) \in \bar{\Lambda} \quad \text{for every } v \in \text{Dom}(L_{00}^+),$$

then $\rho(L_0) \supset \mathbb{C} \setminus \Lambda$ and $\rho(L_0^+) \supset \mathbb{C} \setminus \bar{\Lambda}$, where Λ is a closed sector of the half-plane $\text{Re } \lambda \geq 0$ with vertex at the point 0.

Proof of Theorem 3. As in the proof of Theorem 2, let $v \in L^2(\mathbb{R})$ be a solution of the equation $L^+v = 0$. We show that $v \equiv 0$. Without loss of generality, we assume that the intervals $\Delta_n = [a_n; b_n]$ with $n \in \mathbb{Z}$ are mutually disjoint, and that $b_{-1} < 0$ and $a_1 > 0$. Let $\{\varphi_n\}_{n \in \mathbb{N}}$ be a sequence of real-valued compactly supported functions from $W_{2,comp}^2(\mathbb{R})$ that satisfy the following conditions:

- (a₁) $0 \leq \varphi_n(x) \leq 1$ for every $x \in \mathbb{R}$;
- (a₂) $\varphi_n(x) = 1$ for every $x \in [b_{-n}; a_n]$;
- (a₃) $\text{supp } \varphi_n \subset [a_{-n}; b_n]$;
- (a₄) $\varphi_n'(x) \geq 0$ for any $x \in [a_{-n}; b_{-n}]$, and $\varphi_n'(x) \leq 0$ for any $x \in [a_n; b_n]$;
- (a₅) for some number $K > 0$ the inequalities hold

$$|\varphi_n'(x)| \leq \begin{cases} K|\Delta_{-n}|^{-1} & \text{if } x < 0, \\ K|\Delta_n|^{-1} & \text{if } x > 0. \end{cases}$$

Then, by (13) and by assumptions (B₁) and (B₂) of the theorem, and since $\text{supp } \varphi'_n \subset \Delta_{-n} \cup \Delta_n$ for each $n \in \mathbb{N}$, we obtain the following:

$$\begin{aligned}
\int_{b-n}^{a_n} |v|^2 dx &\leq \int_{\mathbb{R}} \varphi_n^2 |v|^2 dx \leq \int_{\mathbb{R}} (\varphi'_n)^2 |v|^2 dx + 2 \int_{\mathbb{R}} r_1 \varphi'_n \varphi_n |v|^2 dx \\
&= \int_{\Delta_{-n} \cup \Delta_n} (\varphi'_n)^2 |v|^2 dx + 2 \int_{\Delta_{-n}} r_1 \varphi'_n \varphi_n |v|^2 dx + 2 \int_{\Delta_n} r_1 \varphi'_n \varphi_n |v|^2 dx \\
&\leq K^2 \delta^2 \int_{\Delta_{-n} \cup \Delta_n} |v|^2 dx + 2 \int_{\Delta_{-n}} (r_1^+ - r_1^-) \varphi'_n \varphi_n |v|^2 dx + 2 \int_{\Delta_n} (r_1^+ - r_1^-) \varphi'_n \varphi_n |v|^2 dx \\
&= K^2 \delta^2 \int_{\Delta_{-n} \cup \Delta_n} |v|^2 dx + 2 \int_{\Delta_{-n}} r_1^+ \varphi'_n \varphi_n |v|^2 dx - 2 \int_{\Delta_n} r_1^- \varphi'_n \varphi_n |v|^2 dx \\
&\quad - 2 \int_{\Delta_{-n}} r_1^- \varphi'_n \varphi_n |v|^2 dx + 2 \int_{\Delta_n} r_1^+ \varphi'_n \varphi_n |v|^2 dx \\
&\leq K^2 \delta^2 \int_{\Delta_{-n} \cup \Delta_n} |v|^2 dx + 2 \int_{\Delta_{-n}} r_1^+ \varphi'_n \varphi_n |v|^2 dx - 2 \int_{\Delta_n} r_1^- \varphi'_n \varphi_n |v|^2 dx \\
&\leq K^2 \delta^2 \int_{\Delta_{-n} \cup \Delta_n} |v|^2 dx + 2CK \int_{\Delta_{-n}} |v|^2 dx + 2CK \int_{\Delta_n} |v|^2 dx \\
&= K^2 \delta^2 \int_{\Delta_{-n} \cup \Delta_n} |v|^2 dx + 2CK \int_{\Delta_{-n} \cup \Delta_n} |v|^2 dx.
\end{aligned}$$

Here, we use the inequalities

$$-2 \int_{\Delta_{-n}} r_1^- \varphi'_n \varphi_n |v|^2 dx \leq 0 \quad \text{and} \quad 2 \int_{\Delta_n} r_1^+ \varphi'_n \varphi_n |v|^2 dx \leq 0.$$

Therefore, we have

$$(24) \quad \int_{a-n}^{b_n} |v|^2 dx \leq K^2 \delta^2 \int_{\Delta_{-n} \cup \Delta_n} |v|^2 dx + 2CK \int_{\Delta_{-n} \cup \Delta_n} |v|^2 dx \quad \text{for each } n \in \mathbb{N}.$$

Passing in (24) to the limit as $n \rightarrow \infty$, we infer that $v \equiv 0$ because $v \in L^2(\mathbb{R})$.

The proof is complete. □

ACKNOWLEDGMENTS

The work of the first named author (Vladimir Mikhailets) was funded by the Isaac Newton Institute of Mathematical Sciences "Solidarity Program" and the London Mathematical Society. The author wishes to thank the Department of Mathematics, King's College London, for its hospitality.

The work of the second named author (Volodymyr Molyboga) was supported by a grant from the Simons Foundation (SFI-PD-Ukraine-00014586).

REFERENCES

- [1] Albeverio, S.; Gesztesy, F.; Høegh Krohn, R.; Holden, H. *Solvable models in quantum mechanics*, 2nd ed., With an appendix by Pavel Exner; AMS Chelsea Publishing: Providence, RI, 2005; xiv+488 pp.; doi: 10.1090/chel/350.
- [2] Albeverio, S.; Kostenko, A.; Malamud, M. Spectral theory of semibounded Sturm-Liouville operators with local interactions on a discrete set. *J. Math. Phys.* **2010**, 51, no. 10, 102102, 24 pp.; doi: 10.1063/1.3490672.
- [3] Clark, S.; Gesztesy, F. On Povzner–Wienholtz-type self-adjointness results for matrix-valued Sturm–Liouville operators. *Proceedings of the Royal Society of Edinburgh* **2003**, 133A, no. 4, 747–758; doi: 10.1017/S0308210500002651.
- [4] Dunford, N.; Schwartz, J. T. *Linear operators. Part II: Spectral theory. Selfadjoint operators in Hilbert space*. With the assistance of William G. Bade and Robert G. Bartle.; Interscience Publishers John Wiley & Sons: New York - London, 1963; ix+pp. 859–1923+7; MR 0188745.
- [5] Glazman, I. *Direct methods of qualitative spectral analysis of singular differential operators*; Daniel Davey & Co., Inc.: New York, 1966; ix+234 pp.; MR 0190800.
- [6] Gesztesy, F.; Nichols, R.; Zinchenko, M. *Sturm-Liouville operators, their spectral theory, and some applications*; American Mathematical Society Colloquium Publications, **67**, American Mathematical Society: Providence, RI, 2024; xv+927 pp.; doi: 10.1090/coll/067.
- [7] Goriunov, A.; Mikhaleits, V. Regularization of singular Sturm-Liouville equations. *Methods Funct. Anal. Topology* **2010**, 16, no. 2, 120–130; MR 2667807.
- [8] Goriunov, A.; Mikhaleits, V.; Pankrashkin, K. Formally self-adjoint quasi-differential operators and boundary-value problems. *Electron. J. Differential Equations* **2013**, No. 101, 16 pp.; MR 3065054.
- [9] Eckhardt, J.; Gesztesy, F.; Nichols, R.; Teschl, G. Weyl–Titchmarsh theory for Sturm–Liouville operators with distributional potentials. *Opuscula Math.* **2013**, 33, n. 3, 467–563; doi: 10.7494/OpMath.2013.33.3.467.
- [10] Everitt, W.; Markus, L. *Boundary value problems and symplectic algebra for ordinary differential and quasi-differential operators*; Mathematical Surveys and Monographs, **61**; American Mathematical Society: Providence, RI, 1999; xii+187 pp.; MR 1647856.
- [11] Hartman, P., Differential equations with non-oscillatory eigenfunctions. *Duke Math. J.* **1948**, 15, 697–709; MR 0027927.
- [12] Hryniv, R.; Mykytyuk, Ya. Self-adjointness of Schrödinger operators with singular potentials. *Methods Funct. Anal. Topology* **2012**, 18, no. 2, 152–159; MR 2978191.
- [13] Ismagilov, R. S. On the self-adjointness of the Sturm-Liouville operator (Russian). *Uspehi Mat. Nauk* **1963**, 18, no. 5(113), 161–166; MR 0155037.
- [14] Kato, T. *Perturbation theory for linear operators*; Classics in Mathematics; Springer-Verlag: Berlin, 1995; xxii+619 pp.; MR 1335452.
- [15] Kostenko, A.; Malamud, M.; Nicolussi, N. A Glazman-Povzner-Wienholtz theorem on graphs. *Adv. Math.* **2022**, 395, Paper No. 108158, 30 pp.; doi: 10.1016/j.aim.2021.108158.
- [16] Mikhaleits, V.; Goriunov, A.; Molyboga, V. Povzner–Wienholtz-type theorems for Sturm–Liouville operators with singular coefficients. *Complex Anal. Oper. Theory* **2022**, 16, no. 8, Paper No. 113, 13 pp.; doi: 10.1007/s11785-022-01291-y.
- [17] Mikhaleits, V.; Molyboga, V. Remarks on Schrödinger operators with singular matrix potentials. *Methods Funct. Anal. Topology* **2013**, 19, no. 2, 161–167; MR 3098494.
- [18] Mikhaleits, V.; Molyboga, V. Schrödinger operators with measure-valued potentials: semiboundedness and spectrum. *Methods Funct. Anal. Topology* **2018**, 24, no. 3, 240–254; MR 3860804.
- [19] Mikhaleits, V.; Sobolev, A. Common eigenvalue problem and periodic Schrödinger operators. *J. Funct. Anal.* **1999**, 165, no. 1, 150–172; doi: 10.1006/jfan.1999.3406.
- [20] Naimark, M. *Linear Differential Operators* (Russian), 2nd ed.; With an appendix by V. È. Ljance; Nauka: Moscow, 1969; 526 pp.; MR 0353061.
- [21] Orochko, Yu. B. The hyperbolic equation method in the theory of operators of Schrödinger type with locally integrable potential. *Uspekhi Mat. Nauk* **1988**, 43, no. 2(260), 43–86, 230; doi: 10.1070/RM1988v043n02ABEH001728.
- [22] Oročko, Ju. B. Sufficient conditions for the selfadjointness of a Sturm-Liouville operator (Russian). *Mat. Zametki* **1974**, 15, 271–280; MR 0375002.
- [23] Phillips, R. S. Dissipative operators and hyperbolic systems of partial differential equations. *Trans. Amer. Math. Soc.* **1959**, 90, 193–254; doi: 10.2307/1993202.
- [24] Povzner, A. Ya. On the expansion of arbitrary functions in eigenfunctions of the operator $-\Delta u + cu$ (Russian). *Mat. Sbornik N.S.* **1953**, 32/74, 109–156; MR 0053330.
- [25] Reed, M.; Simon, B. *Methods of modern mathematical physics. II. Fourier analysis, self-adjointness*; Academic Press [Harcourt Brace Jovanovich, Publishers]: New York - London, 1975; xv+361 pp.; MR 0493420.
- [26] Rellich, F. Halbbeschränkte gewöhnliche Differentialoperatoren zweiter Ordnung. *Math. Ann.* **1951**, 122, 343–368; doi: 10.1007/BF01342848.
- [27] Sears, D. B. Note on the uniqueness of the Green’s functions associated with certain differential equations. *Canad. J. Math.* **1950**, 2, 314–325; doi: 10.4153/cjm-1950-029-9.
- [28] Schmüdgen, K. *Unbounded self-adjoint operators on Hilbert space*; Graduate Texts in Mathematics, 265; Springer: Dordrecht, 2012; xx+432 pp.; doi: 10.1007/978-94-007-4753-1.
- [29] Shin, D. Quasi-differential operators in Hilbert space. (Russian), Rec. Math. [Mat. Sbornik] N.S. **1943**, 13/55, 39–70; MR 0011534.
- [30] Shubin, M. Essential self-adjointness for semi-bounded magnetic Schrödinger operators on non-compact manifolds. *J. Funct. Anal.* **2001**, 186, no. 1, 92–116; doi: 10.1006/jfan.2001.3778.
- [31] Stetkaer-Hansen, H. A generalization of a theorem of Wienholtz concerning essential selfadjointness of singular elliptic operators. *Math. Scand.* **1966**, 19, 108–112; doi: 10.7146/math.scand.a-10798.


- [32] Weidmann, J. *Spectral theory of ordinary differential operators*; Lecture Notes in Mathematics, 1258, Springer-Verlag: Berlin, 1987; vi+303 pp.; doi: 10.1007/BFb0077960.
- [33] Weyl, H. Über gewöhnliche Differentialgleichungen mit Singularitäten und die zugehörigen Entwicklungen willkürlicher Funktionen (German). *Math. Ann.* **1910**, 68, no. 2, 220–269; doi: 10.1007/BF01474161.
- [34] Wienholtz, E. Halbbeschränkte partielle Differentialoperatoren zweiter Ordnung vom elliptischen Typus. *Math. Ann.* **1958**, 135, 50–80; doi: 10.1007/BF01350827.
- [35] Zettl, A. Formally self-adjoint quasi-differential operator. *Rocky Mountain J. Math.* **1975**, 5, no. 3, 453–474; doi: 10.1216/RMJ-1975-5-3-453.
- [36] Zettl, A. *Sturm–Liouville Theory*; Mathematical Surveys and Monographs, 121; American Mathematical Society: Providence, RI, 2005; xii+328 pp.; doi: 10.1090/surv/121.
- [37] Zettl, A. *Recent developments in Sturm–Liouville theory*; De Gruyter Studies in Mathematics, 76, De Gruyter: Berlin, 2021; xii+245 pp.; doi: 10.1515/9783110719000.

VLADIMIR MIKHAILOTS,  [HTTPS://ORCID.ORG/0000-0002-1332-1562](https://orcid.org/0000-0002-1332-1562)

DEPARTMENT OF MATHEMATICS, KING’S COLLEGE LONDON, 2CR2LS, UNITED KINGDOM

INSTITUTE OF MATHEMATICS OF NAS OF UKRAINE, 3, TERESCHENKIVSKA STR., KYIV-4, UKRAINE, 01024

Email address, Vladimir Mikhailots: mikhailots@imath.kiev.ua, vladimir.mikhailots@gmail.com

VOLODYMYR MOLYBOGA,  [HTTPS://ORCID.ORG/0000-0002-5683-5694](https://orcid.org/0000-0002-5683-5694)

INSTITUTE OF MATHEMATICS OF NAS OF UKRAINE, 3, TERESCHENKIVSKA STR., KYIV-4, UKRAINE, 01024

Email address, Volodymyr Molyboga: molyboga@imath.kiev.ua, vm.imath@gmail.com