

ANTI-SELF-DUAL BLOWUPS II

VSEVOLOD SHEVCHISHIN AND GLEB SMIRNOV

ABSTRACT. Let X be a closed, oriented four-manifold with $b_2^+ \leq 3$, and suppose X contains a collection of pairwise disjoint embedded (-2) -spheres. We prove that there is a Riemannian metric on X such that the Poincaré dual of each of these spheres is represented by an anti-self-dual harmonic 2-form. This extends our earlier result for (-1) -spheres. The main new ingredient is an application of Eliashberg's h -principle for overtwisted contact structures, which we use to construct self-dual harmonic forms on four-orbifolds with prescribed local behaviour near the orbifold singular set.

1. Main result. Let (X, g) be a closed, oriented Riemannian 4-manifold, and let $\Lambda^2 = \Lambda^2 T_X^*$ denote the bundle of 2-forms. The Hodge operator $*$: $\Lambda^2 \rightarrow \Lambda^2$ has eigenvalues ± 1 and induces the splitting:

$$\Lambda^2 = \Lambda^+ \oplus \Lambda^-,$$

where Λ^\pm are the ± 1 -eigenbundles. A 2-form φ is self-dual (SD) if $*\varphi = \varphi$ and anti-self-dual (ASD) if $*\varphi = -\varphi$. Let

$$\mathcal{H}_g^2 = \{\varphi \in \Gamma(\Lambda^2) : \Delta\varphi = 0\}$$

be the space of harmonic 2-forms. By Hodge theory, every harmonic 2-form is closed, and the natural map $\mathcal{H}_g^2 \rightarrow H^2(X; \mathbb{R})$ is an isomorphism. Since Δ commutes with $*$, we obtain the splitting:

$$\mathcal{H}_g^2 = \mathcal{H}_{+,g} \oplus \mathcal{H}_{-,g},$$

where $\mathcal{H}_{+,g}$ and $\mathcal{H}_{-,g}$ are the spaces of harmonic SD and ASD forms. $\mathcal{H}_{+,g}, \mathcal{H}_{-,g} \subset H^2(X; \mathbb{R})$ are positive- and negative-definite with respect to the cup product, and they are orthogonal. Their dimensions $b_2^\pm = \dim \mathcal{H}_{\pm,g}$ are independent of g ; they are the maximal dimensions of positive- and negative-definite subspaces of $H^2(X; \mathbb{R})$, hence topological invariants of X .

For each metric g , consider the positive-definite subspace:

$$P(g) = \mathcal{H}_{+,g} \subset H^2(X; \mathbb{R}) \quad \text{of dimension } b_2^+.$$

$g \rightarrow P(g)$ is the period map of X , introduced in [9]. $P(g)$ depends only on the conformal class of g , and the map P is open in the C^∞ -topology on metrics. It remains an open problem whether a given positive-definite subspace $\pi \subset H^2(X; \mathbb{R})$ arises as $P(g)$ for some metric g .

Let us now state our main result:

Theorem 1. *Let X be a closed, oriented 4-manifold with $b_2^+ \leq 3$. Suppose X contains a collection of pairwise disjoint embedded spheres S_i , each with self-intersection (-2) . Then there exists a Riemannian metric g on X such that every g -self-dual harmonic form u satisfies:*

$$\int_{S_i} u = 0 \quad \text{for all } S_i.$$

Equivalently, if $s_i \in H^2(X; \mathbb{Z})$ denotes the Poincaré dual of S_i , then each s_i is represented by an anti-self-dual form.

Scaduto [13] recently proved that the period map P has dense image for every 4-manifold, and is surjective when $b_2^+ = 1$. This does not imply Theorem 1. Indeed, the conditions $s_i \perp P(g)$ define a closed, positive-codimension subset of the Grassmannian of positive b_2^+ -planes in $H^2(X; \mathbb{R})$, and a dense open subset may avoid such a set.

Remark. Consider the following example. Let X be the underlying manifold of a smooth complex K3 surface. The cup product on $H^2(X; \mathbb{R})$ has signature $(3, 19)$. Let Gr_X^+ be the Grassmannian of 3-dimensional positive-definite planes in $H^2(X; \mathbb{R})$. For any class $\delta \in H^2(X; \mathbb{Z})$ with $\delta^2 = -2$, define:

$$\delta^\perp = \{\pi \in Gr_X^+ : \pi \perp \delta\}.$$

Set:

$$D = Gr_X^+ - \bigcup_{\delta^2 = -2} \delta^\perp.$$

The global Torelli theorem for K3 surfaces states that D is precisely the points in Gr_X^+ realized as the period of a Kähler metric on X ; see [15, 21]. Each δ^\perp has codimension 2 in Gr_X^+ , so D is open and dense. Although no point of δ^\perp can be the period of a Kähler metric, it may still occur as the period of a more general Riemannian metric. This indeed occurs: Kummer surfaces, which are K3, contain 16 disjoint embedded rational curves. Thus, Theorem 1 produces metrics for which up to 16 pairwise disjoint integral (-2) -classes are simultaneously represented by ASD forms.

In our previous work [14], we proved the analogue of Theorem 1 for collections of pairwise disjoint embedded (-1) -spheres. Both proofs follow the same overall strategy, but the (-2) -case presents an additional difficulty, which we now describe.

The first step is to construct a Riemannian orbifold (M, g) together with a closed SD form that vanishes at a prescribed point $p \in M$. Blowing up M at p then produces a manifold X with an embedded exceptional sphere. In the (-1) -case, the point p may be chosen arbitrarily, so producing a metric g with a closed SD form vanishing somewhere is relatively straightforward. In contrast, in the (-2) -case, the point p must be the orbifold point of M ; otherwise the exceptional sphere created in the resolution has self-intersection (-1) rather than (-2) . Thus, the metric g on M has to be chosen much more carefully.

The new geometric input is the following existence theorem for closed SD forms with prescribed zeros. Its proof occupies § 5, § 6, and § 7.

Proposition 1. *Let M be a compact, oriented 4-orbifold with isolated singular points p_1, \dots, p_n , each of type $\mathbb{R}^4/\{\pm 1\}$, and assume $b_2^+(M) \geq 1$. Then there exist a Riemannian metric g on M , flat in neighborhoods of all p_i , and a closed, g -self-dual orbifold form ψ such that $\psi(p_i) = 0$ for all i .*

For each p_i , fix an orbifold chart:

$$\pi_i : (B^4, g_{\text{euc}}) \longrightarrow (U_i, g) \cong B^4/\{\pm 1\} \ni p_i,$$

which is an isometry. An orbifold form means that on $U_i - \{p_i\}$, the pullback $\pi_i^* \psi$ is antipodal-invariant and extends smoothly over $0 \in B^4$; $\psi(p_i) = 0$ means that $\pi_i^* \psi$ vanishes at $0 \in B^4$.

The proof of Proposition 1 proceeds as follows. To begin with, we reduce the problem to constructing a near-symplectic form on M with prescribed behavior near the points p_i . This is done via a theorem of Auroux-Donaldson-Katzarkov (Proposition 1 in [2]). Then, following Taubes

[20], we construct such a near-symplectic form via Eliashberg's classification of overtwisted contact structures in dimension 3 [3].

With Proposition 1 in hand, our argument proceeds as follows. In §2, we introduce an iterative neck-stretching scheme for closed SD forms on manifolds with a long cylindrical neck. In §3, we apply this scheme to the blowup of M at p . This shows that, after inserting a sufficiently long neck, one can find a metric g for which $P(g)$ is arbitrarily close to being orthogonal to a prescribed collection of (-2) -classes. In §4, we then use Proposition 1 to deform such almost-orthogonal metrics so that those (-2) -classes become genuinely ASD, thereby proving Theorem 1. Although these arguments closely parallel our earlier work [14], we reprove certain steps here for completeness.

Combining Theorem 1 with the main result of [14], one obtains the following mixed statement.

Theorem. *Let X be a closed, oriented 4-manifold with $b_2^+ \leq 3$. Suppose X contains a collection of pairwise disjoint embedded spheres S_i , each with self-intersection (-1) or (-2) . Then there exists a Riemannian metric g on X such that every g -self-dual harmonic form u satisfies:*

$$\int_{S_i} u = 0 \quad \text{for all } S_i.$$

This statement does not require any new ideas beyond those used in the pure (-1) - and (-2) -cases. Our neck-stretching scheme treats all spheres simultaneously. We therefore omit a separate proof.

Remark. A closed ASD 2-form whose cohomology class is integral is the curvature of a line bundle with an ASD connection. Thus, our neck-stretching construction is formally similar to Taubes' existence proofs for ASD connections [18, 16, 17] on higher-dimensional bundles. However, unlike Taubes' positive-index problems, the condition that a given set of (-2) -classes be represented by ASD forms is not open in the C^∞ -topology on metrics; the index of our problem is negative.

2. A neck-stretching argument. The gluing argument in this section is inspired by the existing literature (see, e.g., [1, 11, 18]), but we present it in a form tailored to our later applications.

Let X_1 and X_2 be compact, oriented 4-manifolds with common boundary

$$\partial X_1 = \partial X_2 = Y.$$

Fix $\delta > 0$ small. On each X_i choose a Riemannian metric g_i which, on a collar neighborhood

$$(-\delta, 1] \times Y \subset X_i,$$

splits as a product:

$$g_i = dt \otimes dt + g_Y. \tag{2.1}$$

Throughout let us take $Y = \mathbb{RP}^3$ equipped with the metric g_Y induced from the unit round metric g_{S^3} on S^3 via the double cover $S^3 \rightarrow \mathbb{RP}^3$. All results remain valid if $H^1(Y; \mathbb{R}) = 0$.

Let

$$P_i = [0, 1] \times Y \subset X_i$$

denote the smaller collar of the boundary.

Let us form the non-compact manifold X_i^∞ by attaching the semi-infinite cylinder $[0, \infty) \times Y$ along P_i , and extending the product metric (2.1) over the cylindrical end. The resulting metric is denoted g_i^∞ .

Fix $T > 0$ and consider the finite cylinder:

$$N = [0, T] \times Y,$$

equipped with the product metric (2.1). Introduce the extended cylinder:

$$N' = [-1, T+1] \times Y,$$

and attach it to X_1 and X_2 as follows: the collar P_1 is identified with the subcylinder $[-1, 0] \times Y \subset N'$, and P_2 with $[T, T+1] \times Y \subset N'$. For each T , this yields a closed 4-manifold:

$$X_T = X_1 \cup N \cup X_2,$$

carrying a metric g_T which equals g_i on X_i and the product metric on $N = [0, T] \times Y$.

Inside N , let us single out the unit cylinders:

$$Q_1 = [0, 1] \times Y, \quad Q_2 = [T-1, T] \times Y.$$

Let

$$\mathcal{D} = d + d^*, \quad \Delta = \mathcal{D}^2$$

denote the Hodge operator and the Hodge Laplacian.

Let $L^2(X_i^\infty)$ denote the space of L^2 -integrable differential forms on X_i^∞ with respect to the metric g_i . Let $\mathcal{H}^+(X_i^\infty)$ denote the space of SD harmonic forms on X_i^∞ that are also L_2 , i.e.,

$$\mathcal{H}^+(X_i^\infty) = \{g_i^\infty\text{-self-dual}\} \cap \ker \Delta \cap L^2(X_i^\infty).$$

For a domain A , use $\|\cdot\|_{k,A}$ to denote the L_k^2 -Sobolev norm on A .

Introduce cut-off functions ρ_1, ρ_2 such that:

- (1) $\rho_2 = 0$ on $X_1 - P_1$, $\rho_2 = 1$ on $X_T - X_1$, and ρ_2 interpolates between 0 and 1 on P_1 .
- (2) $\rho_1 = 0$ on $X_2 - P_2$, $\rho_1 = 1$ on $X_T - X_2$, and ρ_1 interpolates between 0 and 1 on P_2 .

Fix a form $\psi \in \mathcal{H}^+(X_1^\infty)$. Construct two sequences of smooth SD forms,

$$v^{(i)}, u^{(i)}, \quad i \geq 1,$$

such that:

$$v^{(i)} \in L^2(X_1^\infty) \text{ for } i \text{ odd}, \quad v^{(i)} \in L^2(X_2^\infty) \text{ for } i \text{ even}, \quad u^{(i)} \in L^2(X_T) \text{ for all } i,$$

via the recursion:

- (a) $v^{(1)} = \psi$.
- (b) If i is odd, restrict $v^{(i)}$ to $X_1 \cup N \cup P_2$ and set:

$$u^{(i)} = \rho_1 v^{(i)}.$$

If i is even, restrict $v^{(i)}$ to $X_2 \cup N \cup P_1$ and set:

$$u^{(i)} = \rho_2 v^{(i)}.$$

- (c) Suppose i is odd (the even case is symmetric). Construct an SD form $v^{(i+1)} \in L^2(X_2^\infty)$ solving

$$\mathcal{D}v^{(i+1)} = -\mathcal{D}u^{(i)}.$$

The construction produces elliptic estimates of the form:

$$\|v^{(i+1)}\|_{k, X_2 \cup Q_2} \leq A_k \|\Delta u^{(i)}\|_{k-2, P_2}, \tag{2.2}$$

with constants A_k independent of T and i . For even i , an identical estimate holds on $X_1 \cup Q_1$.

Let us now explain step (c) in detail for odd i . The even case is similar.

To begin with, assume $v^{(i-1)}$ and $u^{(i-1)}$ have already been constructed. Fix numbers $0 < T_1 < T_2$ and choose a smooth, positive function $\chi: [0, \infty) \times Y \rightarrow \mathbb{R}$ such that:

$$\chi(t, y) = \begin{cases} 1, & t \in [0, T_1], \\ e^{-2t}, & t \geq T_2. \end{cases}$$

Extend χ to all of X_1^∞ by setting it equal to 1 on X_1 . Consider the conformal rescaling:

$$\hat{g}_1 = \chi g_1^\infty.$$

(X_1^∞, \hat{g}_1) has an asymptotically Euclidean end. Indeed, setting $r = e^{-t}$ yields:

$$e^{-2t}(dt \otimes dt + g_{S^3}) = dr \otimes dr + r^2 g_{S^3}.$$

Thus, the region $[T_2, \infty) \times Y$ endowed with \hat{g}_1 is isometric to the punctured closed ball of radius e^{-T_2} modulo the antipodal map. Adding the point at infinity produces a compact Riemannian orbifold, smooth away from the added point, where the metric is locally isometric to $\mathbb{R}^4/\{\pm 1\}$.

Use \hat{X}_1 to denote the orbifold compactification of X_1^∞ .

Choose a neighborhood $U \subset \hat{X}_1$ of the orbifold point together with an isometry:

$$B \xrightarrow{\pi} B/\{\pm 1\} \xrightarrow{\tau} U,$$

where π identifies antipodal points and τ is the chosen isometry. A differential form ϕ is smooth on U if $(\pi \circ \tau)^* \phi$ extends smoothly across the origin in B . A form on \hat{X}_1 is smooth if it is smooth away from the orbifold point and smooth on U . On the domain of such smooth forms, the operator \mathcal{D} remains elliptic and self-adjoint, so standard Hodge theory applies.

$\mathcal{D}u^{(i-1)}$ is supported in $P_1 \cup P_2$. Let us restrict $\mathcal{D}u^{(i-1)}$ to $X_1 \cup N$ and extend it by zero to the remainder of \hat{X}_1 . Let us continue to denote this extension by $\mathcal{D}u^{(i-1)}$.

Let $\mathcal{H}(\hat{X}_1)$ denote the space of harmonic 2-forms on (\hat{X}_1, \hat{g}_1) . For any $\varphi \in \mathcal{H}(\hat{X}_1)$,

$$\int_{\hat{X}_1} \langle \varphi, \Delta u^{(i-1)} \rangle = \int_{\hat{X}_1} \langle \mathcal{D}\varphi, \mathcal{D}u^{(i-1)} \rangle = 0,$$

because $\Delta\varphi = 0$ implies $\mathcal{D}\varphi = 0$. Thus, $\Delta u^{(i-1)}$ is L^2 -orthogonal to $\mathcal{H}(\hat{X}_1)$. Hodge theory on the compact orbifold (\hat{X}_1, \hat{g}_1) then gives a unique SD form $\hat{v}^{(i)}$ such that:

$$\Delta \hat{v}^{(i)} = -\Delta u^{(i-1)}, \quad \hat{v}^{(i)} \perp \mathcal{H}(\hat{X}_1).$$

This can be strengthened:

Lemma 1. $\mathcal{D}\hat{v}^{(i)} = -\mathcal{D}u^{(i-1)}$ for $\hat{v}^{(i)}$ defined above.

Proof. To begin with, observe that $du^{(i-1)}$ is an exact 3-form on \hat{X}_1 . Indeed:

$$du^{(i-1)} = d(\rho_2 v^{(i-1)}) = d\rho_2 \wedge v^{(i-1)}.$$

Since $H^2(Y; \mathbb{R}) = 0$, there exists a 1-form α on P_1 with $d\alpha = v^{(i-1)}$. Hence:

$$du^{(i-1)} = -d(d\rho \wedge \alpha) \quad \text{on } P_1.$$

Consider $\hat{v}^{(i)} + u^{(i-1)}$. By construction:

$$\mathcal{D}^2(\hat{v}^{(i)} + u^{(i-1)}) = 0.$$

Thus:

$$\Delta \mathcal{D}(\hat{v}^{(i)} + u^{(i-1)}) = \Delta d(\hat{v}^{(i)} + u^{(i-1)}) + \Delta d^*(\hat{v}^{(i)} + u^{(i-1)}) = 0.$$

In particular, $d(\hat{v}^{(i)} + u^{(i-1)})$ is harmonic. Since it is also exact, it must vanish. Both $\hat{v}^{(i)}$ and $u^{(i-1)}$ are self-dual; therefore:

$$\mathcal{D}(\hat{v}^{(i)} + u^{(i-1)}) = d(\hat{v}^{(i)} + u^{(i-1)}) - *d(\hat{v}^{(i)} + u^{(i-1)}) = 0,$$

and the lemma follows. \square

Elliptic regularity now gives, for each k ,

$$\|\hat{v}^{(i)}\|_{k, \hat{X}_1} \leq A_k \|\Delta u^{(i-1)}\|_{k-2, \hat{X}_1} = A_k \|\Delta u^{(i-1)}\|_{k-2, P_1},$$

with constants A_k depending only on \hat{g}_1 .

Restrict $\hat{v}^{(i)}$ to the cylindrical-end manifold:

$$v^{(i)} = \hat{v}^{(i)}|_{X_1^\infty}.$$

Choosing T_2 large enough, assume $\chi = 1$ on $X_1 \cup Q_1$. On that region, the metrics g_1^∞ and \hat{g}_1 agree, and hence:

$$\|v^{(i)}\|_{k, X_1 \cup Q_1} \leq \|\hat{v}^{(i)}\|_{k, \hat{X}_1} \leq A_k \|\Delta u^{(i-1)}\|_{k-2, P_1},$$

which is precisely (2.2).

The L^2 -norm of 2-forms is conformally invariant; therefore, if $\hat{v}^{(i)}$ is L^2 for \hat{g}_1 , then it also L^2 for g_1^∞ .

Finally, the equation

$$\mathcal{D}v^{(i)} = -\mathcal{D}u^{(i-1)}$$

holds on (X_1^∞, \hat{g}_1) by construction. To see that it also holds for g_1^∞ , note that the kernel of \mathcal{D} is locally conformally invariant, so the only issue is the region where $\mathcal{D}u^{(i-1)}$ is supported. But $\mathcal{D}u^{(i-1)}$ is supported in P_1 , where $\chi = 1$ and $g_1^\infty = \hat{g}_1$. This completes step (c).

Iterating this procedure one obtains sequences $\{u^{(i)}\}$ and $\{v^{(i)}\}$ as above. Formally, one expects:

$$\mathcal{D}\left(\sum_i u^{(i)}\right) = 0.$$

More precisely, for each i :

- the support of $\mathcal{D}u^{(i)}$ is contained in $P_1 \cup P_2$;
- on P_2 we have $\mathcal{D}u^{(i)} = -\mathcal{D}u^{(i-1)}$ for even i , while on P_1 we have $\mathcal{D}u^{(i)} = -\mathcal{D}u^{(i-1)}$ for odd i .

Lemma 2. *Let $k > 0$. For all large enough T , the series*

$$u = \sum_{i=1}^{\infty} u^{(i)}$$

converges in the L_k^2 -norm on (X_T, g_T) , and its tail satisfies:

$$\|u - u^{(1)}\|_{k, X_T} = O(e^{-2T}), \quad \|u - u^{(1)} - u^{(2)}\|_{k, X_T} = O(e^{-4T}).$$

Proof. It is enough to find constants $C_k > 0$, independent of T and i , such that:

$$\|u^{(i)}\|_{k, X_T} \leq C_k e^{-2T} \|u^{(i-1)}\|_{k, X_T}. \quad (2.3)$$

The needed convergence then follows by summing a geometric series.

To this end, let us study harmonic forms on the cylindrical neck. On $N = [0, T] \times Y$, the Hodge Laplacian splits as:

$$\Delta = -\partial_t^2 + \Delta_3,$$

where Δ_3 is the Hodge Laplacian on Y , acting in the y -variables only. Since ψ is harmonic on X_1^∞ , its restriction to N admits a Fourier expansion:

$$\psi = \sum_{\lambda} (\alpha_{\lambda} + * \alpha_{\lambda}) e^{-\lambda t}, \quad (2.4)$$

where λ ranges over square roots (positive and negative) of eigenvalues of Δ_3 acting on 2-forms, the α_{λ} are Δ_3 -eigenforms with eigenvalue λ^2 , and $*$ is the Hodge star on Y .

Eigenvalues and eigenforms of Δ_3 are known (see, e.g., [4]). $\mathcal{D}\psi = 0$ forces $d\alpha_{\lambda} = 0$. Hence, only closed eigenforms occur. The smallest eigenvalue of the Hodge Laplacian on closed 2-forms on the unit sphere S^3 is 4, and these eigenforms descend to \mathbb{RP}^3 . Thus, the smallest eigenvalue for closed 2-forms on Y is also 4. Consequently, every λ in (2.4) satisfies $\lambda \geq 2$. (Only the fact that the smallest eigenvalue is positive will be used below.)

The L^2 condition on X_1^∞ rules out negative λ in (2.4), so we may rewrite:

$$\psi = \sum_{\lambda \geq 2} (\alpha_{\lambda} + * \alpha_{\lambda}) e^{-\lambda t}.$$

For $s \in [0, T]$, define the translated unit cylinder:

$$Q_1 + s = [s, s+1] \times Y \subset N.$$

Since the α_{λ} are independent of t ,

$$\|\psi\|_{0, Q_1+s}^2 = \sum_{\lambda \geq 2} \int_s^{s+1} 2e^{-2\lambda t} \left(\int_Y \langle \alpha_{\lambda}, \alpha_{\lambda} \rangle \right) dt \leq e^{-4s} \|\psi\|_{0, Q_1}^2.$$

Thus:

$$\|\psi\|_{0, Q_1+s} \leq e^{-2s} \|\psi\|_{0, Q_1}. \quad (2.5)$$

Choose $\delta > 0$ small so that g_T remains product on the longer collars $[-\delta, 1+\delta] \times Y$ and $[T-1-\delta, T+\delta] \times Y$ for all large T . Interior elliptic estimates for the translation-invariant operator Δ imply, for each k ,

$$\|\psi\|_{k, Q_1+s}^2 \leq B'_k \left(\|\psi\|_{0, Q_1+s+\delta}^2 + \|\psi\|_{0, Q_1+s-\delta}^2 \right), \quad (2.6)$$

with a constant B'_k independent of s and T . Combining (2.5) and (2.6), we find:

$$\|\psi\|_{k, Q_1+s}^2 \leq B_k e^{-4s} \|\psi\|_{k, Q_1}^2$$

for a suitable B_k independent of s and T . Summing over $s = 0, 1, \dots, T-1$, we obtain:

$$\|\psi\|_{k, N}^2 = \sum_{j=0}^{T-1} \|\psi\|_{k, Q_1+j}^2 \leq D_k \|\psi\|_{k, Q_1}^2,$$

and at the far end of the neck,

$$\|\psi\|_{k, P_2}^2 = \|\psi\|_{k, Q_1+T}^2 \leq B_k e^{-4T} \|\psi\|_{k, Q_1}^2.$$

B_k, D_k depend only on (Y, g_Y) and the choice of δ . Hence, the same estimates hold with $v^{(i)}$ in place of ψ :

$$\begin{aligned} \|v^{(i)}\|_{k,N}^2 &\leq D_k \|v^{(i)}\|_{k,Q_1}^2, \quad \|v^{(i)}\|_{k,P_2}^2 \leq B_k e^{-4T} \|v^{(i)}\|_{k,Q_1}^2, \quad \text{if } i \text{ is odd,} \\ \|v^{(i)}\|_{k,N}^2 &\leq D_k \|v^{(i)}\|_{k,Q_2}^2, \quad \|v^{(i)}\|_{k,P_1}^2 \leq B_k e^{-4T} \|v^{(i)}\|_{k,Q_2}^2, \quad \text{if } i \text{ is even.} \end{aligned} \quad (2.7)$$

On Q_1 (for odd i) and on Q_2 (for even i), we have $u^{(i)} = v^{(i)}$. Since $u^{(i)}$ and $v^{(i)}$ also agree on N , (2.7) implies:

$$\|u^{(i)}\|_{k,N}^2 \leq D_k \|u^{(i)}\|_{k,Q_1}^2 \quad \text{for } i \text{ odd,} \quad \|u^{(i)}\|_{k,N}^2 \leq D_k \|u^{(i)}\|_{k,Q_2}^2 \quad \text{for } i \text{ even.} \quad (A)$$

ρ_1, ρ_2 are smooth, so for we obtain:

$$\|u^{(i)}\|_{k,P_2}^2 \leq E_k \|v^{(i)}\|_{k,P_2}^2 \quad \text{for } i \text{ odd,} \quad \|u^{(i)}\|_{k,P_1}^2 \leq E_k \|v^{(i)}\|_{k,P_1}^2 \quad \text{for } i \text{ even,}$$

Combining this with (2.7), we find:

$$\|u^{(i)}\|_{k,P_2}^2 \leq F_k e^{-4T} \|u^{(i)}\|_{k,Q_1}^2 \quad \text{for } i \text{ odd,} \quad \|u^{(i)}\|_{k,P_1}^2 \leq F_k e^{-4T} \|u^{(i)}\|_{k,Q_2}^2 \quad \text{for } i \text{ even,} \quad (B)$$

where $F_k = B_k E_k$.

Finally, using (2.2) together with continuity of Δ , we obtain:

$$\begin{aligned} \|u^{(i)}\|_{k,X_1 \cup Q_1} &\leq A_k \|\Delta u^{(i-1)}\|_{k-2,P_1} \leq G_k \|u^{(i-1)}\|_{k,P_1}, \quad \text{if } i \text{ is odd,} \\ \|u^{(i)}\|_{k,X_2 \cup Q_2} &\leq A_k \|\Delta u^{(i-1)}\|_{k-2,P_2} \leq G_k \|u^{(i-1)}\|_{k,P_2}, \quad \text{if } i \text{ is even,} \end{aligned} \quad (C)$$

for some G_k .

Let us now estimate $\|u^{(i)}\|_{k,X_T}$ for odd i (the even case is symmetric). We compute:

$$\begin{aligned} \|u^{(i)}\|_{k,X_T}^2 &= \|u^{(i)}\|_{k,X_1}^2 + \|u^{(i)}\|_{k,N}^2 + \|u^{(i)}\|_{k,P_2}^2 \stackrel{(A)}{\leq} \|u^{(i)}\|_{k,X_1}^2 + D_k \|u^{(i)}\|_{k,Q_1}^2 + \|u^{(i)}\|_{k,P_2}^2 \stackrel{(B)}{\leq} \\ &\leq \|u^{(i)}\|_{k,X_1}^2 + D_k \|u^{(i)}\|_{k,Q_1}^2 + e^{-4T} F_k \|u^{(i)}\|_{k,Q_1}^2 \leq H_k \|u^{(i)}\|_{k,X_1 \cup Q_1}^2 \stackrel{(C)}{\leq} \\ &\leq H_k G_k^2 \|u^{(i-1)}\|_{k,P_1}^2 \stackrel{(B)}{\leq} e^{-4T} F_k H_k G_k^2 \|u^{(i-1)}\|_{k,Q_2}^2 \leq e^{-4T} C_k^2 \|u^{(i-1)}\|_{k,X_T}^2. \end{aligned}$$

This completes the proof. \square

Let us now isolate the contribution of $u^{(1)}$ to the full series u . Along the neck, write the Fourier expansion of ψ as:

$$\psi = \sum_{\lambda \geq 2} \psi_\lambda, \quad \psi_\lambda = a_\lambda e^{-\lambda t}, \quad (2.8)$$

where each ψ_λ is an SD eigenform of $-\partial_t^2 + \Delta_3$ with parameter λ .

Let $v^{(2)}$ be as above. Decompose it as follows:

$$v^{(2)} = \sum_{\lambda \geq 2} v_\lambda^{(2)},$$

where $v_\lambda^{(2)} \in L^2(X_2^\infty)$ solves

$$\mathcal{D}v_\lambda^{(2)} = -\mathcal{D}(\rho_1 \psi_\lambda).$$

Define $u_\lambda^{(2)} = \rho_2 v_\lambda^{(2)}$ on X_T so that

$$u^{(2)} = \sum_{\lambda \geq 2} u_\lambda^{(2)}.$$

Let $c > 0$ be such that c^2 is the next eigenvalue of Δ_3 on closed 2-forms after the minimal eigenvalue 4. Let us show that $u^{(2)}$ is well-approximated by the part corresponding to the lowest eigenvalue.

Lemma 3. *For each k , we have:*

$$\left\| \sum_{\lambda > 2} u_\lambda^{(2)} \right\|_{k, X_2} = O(e^{-cT}),$$

for $c > 2$ as above.

Proof. By the definition of $u_\lambda^{(2)}$, we have:

$$\mathcal{D} \left(\sum_{\lambda \geq c} u_\lambda^{(2)} \right) = - \mathcal{D} \left(\rho_1 \sum_{\lambda \geq c} \psi_\lambda \right).$$

Using (C), we find:

$$\left\| \sum_{\lambda \geq c} u_\lambda^{(2)} \right\|_{k, X_2} \stackrel{(C)}{\leq} G_k \left\| \rho_1 \sum_{\lambda \geq c} \psi_\lambda \right\|_{k, P_2} \leq G_k \sqrt{E_k} \left\| \sum_{\lambda \geq c} \psi_\lambda \right\|_{k, P_2}.$$

For the tail $\sum_{\lambda \geq c} \psi_\lambda$, repeat the decay argument used to prove (2.5), but using that now $\lambda \geq c > 2$. This gives:

$$\left\| \sum_{\lambda \geq c} \psi_\lambda \right\|_{k, P_2} \leq B_k e^{-cT} \left\| \sum_{\lambda \geq c} \psi_\lambda \right\|_{0, Q_1} = O(e^{-cT}),$$

and the lemma follows. \square

3. Resolution formula. Let (M, g) be a Riemannian 4-orbifold, and let $p \in M$ be an isolated orbifold point modelled on $\mathbb{R}^4/\{\pm 1\}$ with the standard flat metric. Thus there exists a neighbourhood U of p and an isometry:

$$U \simeq (B/\{\pm 1\}, g_0),$$

where $B \subset \mathbb{R}^4$ is a ball centered at the origin, g_0 is the flat metric (the action of $\{\pm 1\}$ is orthogonal).

Choose a linear complex structure on $\mathbb{R}^4 \cong \mathbb{C}^2$ such that g_0 is Kähler. Then view U as a complex orbifold domain. Let ω denote the Kähler form on U associated to g , and let $r: M \rightarrow \mathbb{R}$ be the distance to p with respect to g . On U one has:

$$\omega = -dd^{\mathbb{C}}r^2.$$

After rescaling g , we may assume:

$$U = \{x \in M \mid r(x) < 1\}.$$

The unitary group $U(2)$ preserves (\mathbb{C}^2, g_0) and commutes with the action of $\{\pm 1\}$; hence, it induces an isometric action on U that leaves both r and ω invariant.

Let $Y \subset U$ be defined as:

$$Y = \{x \in M \mid r(x) = 1\},$$

and let g_Y be the induced metric. Geometrically, Y is the unit sphere in \mathbb{C}^2 modulo the antipodal map.

Choose $a > 0$ so small that $a \cdot e < 1$ ($\ln e = 1$), and set:

$$N = \{x \in U \mid a \leq r(x) \leq a \cdot e\}.$$

To apply Lemma 2, introduce the annuli:

$$P_1 = \{x \in U \mid a \cdot e \leq r(x) \leq a \cdot e^2\}, \quad P_2 = \{x \in U \mid a \cdot e^{-1} \leq r(x) \leq a\},$$

which agree with the notation in §2.

The metric on N has the form:

$$dr \otimes dr + r^2 g_Y.$$

Hence, N is conformally equivalent to the cylinder $[0, 1] \times Y$ with the product metric:

$$dt \otimes dt + g_Y.$$

Let us modify M as follows. Consider the neighbourhood of p defined as:

$$U_a = \{x \in U \mid r(x) < a\} \subset M.$$

This neighbourhood contains the cuff P_2 but not the neck N . Let

$$\sigma: X_2 \longrightarrow U_a$$

be the minimal resolution of U_a , obtained by blowing up the orbifold point p . The exceptional set is an embedded sphere $C \subset X_2$ with self-intersection (-2) .

Consider a smaller neighbourhood of p defined as:

$$U_\delta = \{x \in U \mid r(x) < \delta\} \subset M.$$

By standard complex-analytic results (see, e.g., Ch. 1 of [6]), the complex surface X_2 admits a strictly plurisubharmonic function $h: X_2 \rightarrow \mathbb{R}$ such that:

$$h = r^2 \quad \text{on } X_2 - \sigma^{-1}(U_\delta).$$

Define a Kähler form:

$$\omega' = -dd^c h \quad \text{on } X_2,$$

and let g' be the associated Riemannian metric. By construction, g and g' agree away from U_δ . Let us now build a manifold X by cutting out U_a from M and gluing in (X_2, g') along the common boundary:

$$X = (M - U_a) \cup_{\partial X_2} X_2.$$

Introduce a conformal transformation of the metric on X . Let $\chi: X \rightarrow \mathbb{R}$ be a positive function such that the χ -rescaled metric on $P_1 \cup N \cup P_2$ takes the form:

$$dt \otimes dt + g_Y,$$

and such that $\chi = 1$ away from a small neighbourhood of $P_1 \cup N \cup P_2$. For $T > 0$, let (X_T, g_T) denote the space obtained from X by replacing the factor $[0, 1]$ in N by $[0, T]$. When T is large, (X_T, g_T) contains a long cylindrical region.

Lemma 4. *Let M , X , and X_T be as above, and let $C \subset X_2$ be the exceptional (-2) -sphere. Assume that M carries an SD harmonic form ψ . Let $\langle \psi, \omega \rangle_p$ denote the pointwise inner product with respect to g , evaluated at p . Then, for all large enough T , there exists an SD harmonic form u on X_T such that:*

$$\|u - \psi\|_{C^0, X_1} \longrightarrow 0 \quad \text{as } T \rightarrow \infty,$$

and

$$\int_C u = A \langle \omega, \psi \rangle_p e^{-2T} + O(e^{-cT}) \tag{3.1}$$

for some constants $A > 0$ and $c > 2$ independent of T .

To clarify the quantity $\langle \omega, \psi \rangle_p$: choose a uniformizing chart $\pi: (B, g_0) \rightarrow (U, g)$. Both ω and ψ are smooth on U ; thus their lifts $\pi^*\omega$ and $\pi^*\psi$ are smooth, $\{\pm 1\}$ -invariant forms on B . Define $\langle \omega, \psi \rangle_p$ by $\langle \omega, \psi \rangle_p = \langle \pi^*\omega, \pi^*\psi \rangle_{0 \in B}$, which is independent of the chosen uniformizing chart.

Proof. In the neighbourhood U of p , the form ψ admits a Fourier expansion:

$$\psi = \sum_{\lambda \geq 2} \psi_\lambda, \quad \psi_\lambda = a_\lambda r^\lambda,$$

where each ψ_λ is an SD harmonic form on U . Smoothness at the orbifold point forces all exponents to be positive. Introducing the cylindrical coordinate t via $r = e^{-t}$, the series takes the form used in (2.8).

Let P_1, P_2 be as above, and let χ be the function such that the χg satisfies:

$$dt \otimes dt + g_Y \quad \text{on } P_1 \cup N \cup P_2,$$

and such that χ equals 1 away from a small neighbourhood of $P_1 \cup N \cup P_2$.

Replace the original neck N by a longer cylindrical region $[0, T] \times Y$, and analytically continue ψ from P_1 to P_2 along this neck. On P_2 , the continued form has the expansion:

$$\psi = \sum_{\lambda \geq 2} \psi_\lambda e^{-\lambda T},$$

and this series converges on all of M_2 . For $\lambda > 2$, the norm of ψ_λ tends to 0 as $r \rightarrow 0$. Hence, the leading contribution to $\langle \psi, \omega \rangle$ at p comes from the $\lambda = 2$ mode, giving:

$$\langle \psi, \omega \rangle_p = e^{-2T} \langle \psi_2, \omega \rangle_p.$$

Choose holomorphic coordinates (z, w) on \mathbb{C}^2 , and set:

$$\Omega = dz \wedge dw.$$

The form Ω is invariant under the action of $\{\pm 1\}$, so it descends to U . The eigenspace of the Laplacian on Y with eigenvalue 4 is 3-dimensional, and we may write:

$$\psi_2 = a_1 \omega_1 + a_2 \omega_2 + a_3 \omega_3, \quad a_i \in \mathbb{R},$$

where:

$$\omega_1 = \omega, \quad \omega_2 = \operatorname{Re}(\Omega), \quad \omega_3 = \operatorname{Im}(\Omega).$$

These satisfy:

$$\langle \omega_i, \omega_j \rangle = 2\delta_{ij} \quad \text{on } U,$$

so in particular:

$$\langle \psi_2, \omega \rangle = 2a_1.$$

Replacing U_a by X_2 produces X_T . Using Lemma 2, construct an SD harmonic form u on X_T such that:

$$u = \rho_1 \psi + u^{(2)} + O(e^{-4T}),$$

with $d\rho_1$ supported in P_2 . Using Lemma 3, when restricted to X_2 , we obtain:

$$u^{(2)} = u_2^{(2)} + O(e^{-cT}), \quad \mathcal{D}u_2^{(2)} = -e^{-2T} \mathcal{D}(\rho_1 \psi_2),$$

for some $c > 2$. To solve

$$\mathcal{D}u_2^{(2)} = -e^{-2T} \mathcal{D}(\rho_1 \psi_2),$$

observe that ψ_2 extends from P_2 to a closed SD form on all of X_2 :

$$\eta = a_1 \omega' + a_2 \operatorname{Re}(\sigma^* \Omega) + a_3 \operatorname{Im}(\sigma^* \Omega).$$

Indeed, the holomorphic form $\sigma^*\Omega$ is g' -self-dual, and ω' is Kähler for g' ; both remain self-dual after introducing the conformal factor χ .

Thus:

$$u_2^{(2)} = e^{-2T} (1 - \rho_1) \eta.$$

Since $\rho_1 = 0$ near the exceptional sphere C , we obtain:

$$u = e^{-2T} \eta + O(e^{-cT}) \quad \text{near } C.$$

We compute:

$$\int_C u = e^{-2T} \int_C \eta + O(e^{-cT}) = e^{-2T} \int_C a_1 \omega' + O(e^{-cT}) = e^{-2T} \langle \psi_2, \omega \rangle_p A + O(e^{-cT}),$$

for some constant $A > 0$ independent of T . This completes the proof. \square

4. Proof of Theorem 1. It suffices to treat the case $b_2^+ = 3$; the general case is no harder. Let (M, g) be an oriented Riemannian 4-orbifold with isolated singular points p_1, \dots, p_n , each of type $\mathbb{R}^4/\{\pm 1\}$. By Proposition 1, we may choose g so that there exists a g -self-dual harmonic 2-form ψ_1 which vanishes at every p_α . Since $b_2^+(M) = 3$, we can choose further SD harmonic forms ψ_2, ψ_3 so that ψ_1, ψ_2, ψ_3 is a basis for SD harmonic forms.

For each $\alpha = 1, \dots, n$, fix a neighbourhood V_α of p_α with an isometry

$$V_\alpha \simeq (B/\{\pm 1\}, g_0),$$

where $B \subset \mathbb{R}^4$ is a ball centered at the origin and g_0 is the flat metric. Choose linear complex structures J_α on B such that g_0 is Kähler. J_α is invariant under the antipodal map, so it descends to V_α . Let ω_α be the resulting Kähler form on V_α .

By rotating J_α , we may arrange that, for each α ,

$$\langle \omega_\alpha, \psi_1 \rangle_{p_\alpha} = \langle \omega_\alpha, \psi_2 \rangle_{p_\alpha} = 0.$$

For each α , let $D_\alpha \subset \mathbb{R}^3$ be a 3-disc with coordinates $(s_{\alpha,1}, s_{\alpha,2}, s_{\alpha,3})$. Set:

$$D^{3n} = D_1 \times \dots \times D_n \subset \mathbb{R}^{3n}.$$

Let g_s , $s \in D^{3n}$, be a family of orbifold metrics with $g_0 = g$. Let $a_i \in H^2(M; \mathbb{R})$ denote the cohomology class of ψ_i , and let $\psi_{i,s}$ be the g_s -self-dual part of the g_s -harmonic representative of a_i . For each α and i , define:

$$\pi_{\alpha,i}(s) = \langle \omega_\alpha, \psi_{i,s} \rangle_{p_\alpha}, \quad \alpha = 1, \dots, n, \quad i = 1, 2, 3,$$

where the inner product is computed using g_s . By construction, $\pi_{\alpha,i}(0) = 0$ for all α, i .

Lemma 5. *There exist neighbourhoods $U_\alpha \subset V_\alpha$ of p_α and a family of metrics g_s , $s \in D^{3n}$, such that $g_s = g$ on each U_α for all $s \in D^{3n}$, and such that the map*

$$s \rightarrow (\pi_{\alpha,i}(s))_{\alpha,i},$$

is a local diffeomorphism near $s = 0$.

Proof. The proof is identical to that of Lemma 4 in [14], which relies on [8, 7]. \square

For simplicity, assume $n = 1$ and drop index α ; the general case is no harder. Let (M, g_s) be as in Lemma 5. Let us apply the neck-stretching procedure of §3 inside U . This produces a family $(M, g_{T,s})$ which agrees with g_s on U (technically, on a slightly smaller neighbourhood). Let us next resolve the double point p using the complex structure J . Let C denote the exceptional (-2) -sphere, and let the resulting smooth family of 4-manifolds be denoted by $X_{T,s}$.

By Lemma 4, there exist SD harmonic forms $u_{i,s}$ on $X_{T,s}$ such that:

$$\int_C u_{i,s} = A \pi_i(s) e^{-2T} + O(e^{-cT}), \quad A > 0, \quad c > 2. \quad (4.1)$$

For large T , the error term in (4.1) is dominated by the leading contribution; hence, the map

$$s \rightarrow \left(\int_C u_{1,s}, \int_C u_{2,s}, \int_C u_{3,s} \right)$$

is surjective onto a neighbourhood of the origin in \mathbb{R}^3 . This completes the proof.

5. Near-contact forms. In this section we consider contact forms with isolated, nondegenerate singularities on 3-manifolds. The goal is to show how Eliashberg's h -principle gives a complete classification of such forms up to deformation.

Let Y be a 3-manifold. A 1-form λ on Y is called a *near-contact form* if:

- (1) λ is transverse to the zero section of T_Y^* , so its zero set $\lambda^{-1}(0) = \{p_1, \dots, p_k\}$ is a discrete (for simplicity, finite) set of points;
- (2) there exists a volume form vol on Y such that, writing

$$\lambda \wedge d\lambda = f \text{ vol}.$$

the function f is positive on $Y - \{p_1, \dots, p_k\}$, and each p_i is a nondegenerate local minimum of f . (This condition then holds for any volume form inducing the same orientation as $\lambda \wedge d\lambda$.)

Lemma 6. *Suppose that p is a transverse zero of λ and that λ is contact in a small punctured neighbourhood of p ; then $d\lambda$ also vanishes at p .*

Proof. Choose a connection ∇ near p . Since λ is contact in a punctured neighbourhood of p , we have:

$$\nabla(\lambda \wedge d\lambda)|_p = 0.$$

Using

$$\nabla(\lambda \wedge d\lambda) = \nabla\lambda \wedge d\lambda + \lambda \wedge \nabla d\lambda,$$

we obtain:

$$\nabla\lambda|_p \wedge d\lambda|_p = 0.$$

The transverse vanishing of λ at p now implies $d\lambda|_p = 0$. This completes the proof. \square

If p is a transverse zero of a 1-form λ and $d\lambda$ also vanishes at p , then the bilinear form:

$$A(\xi, \eta) = \nabla_\xi \lambda(\eta) \quad \text{for } \xi, \eta \in T_Y|_p, \quad (5.1)$$

is symmetric and nondegenerate. Since $\lambda|_p = 0$, the form A does not depend on ∇ .

Lemma 7. *If a near-contact 1-form λ vanishes at p , then the form A defined by (5.1) is not definite.*

Proof. Choose a flat metric g near p , let ∇ be its flat connection, and let vol_g be its volume form. Let Y be the vector field dual to $d\lambda$ with respect to vol_g ; that is,

$$\iota_Y \text{vol}_g = d\lambda.$$

Then:

$$L_Y \text{vol}_g = (\text{div } Y) \text{vol}_g = 0.$$

Thus, the endomorphism field $\nabla Y: T_Y \rightarrow T_Y$ is traceless everywhere. In particular, the linear map:

$$B(\xi) = \nabla_\xi Y \in T_Y|_p \quad \text{for all } \xi \in T_Y|_p,$$

satisfies $\text{tr } B = 0$. We compute:

$$\lambda \wedge d\lambda = \lambda(Y) \text{vol}_g.$$

Since λ and Y both vanish at p , it follows that:

$$\nabla_\xi \nabla_\eta \lambda(Y)|_p = A(\xi, B\eta) + A(\eta, B\xi) \quad \text{for all } \xi, \eta \in T_Y|_p.$$

Assume for contradiction that A is definite. Choose g so that $g = A$ at p . Set:

$$H(\xi, \xi) = \nabla_\xi^2 \lambda(Y)|_p = 2g(\xi, B\xi).$$

For any g -orthonormal basis $\{e_1, e_2, e_3\}$, we obtain:

$$\text{tr}_g H = \sum_{i=1}^3 H(e_i, e_i) = 2 \text{tr } B = 0.$$

Hence, the function $f = \lambda(Y)$ cannot have a nondegenerate maximum or minimum at p . This contradiction completes the proof. \square

Define the index of a transverse zero p of λ as:

$$\text{ind}_p \lambda = \text{sgn det } A.$$

For a nondegenerate zero p of a near-contact form λ , this index provides a topological obstruction to deforming one such form into another (through near-contact forms with a nondegenerate zero at p). Another obstruction is the orientation induced by $\lambda \wedge d\lambda$ on a punctured neighbourhood of p . We now show that there are no further obstructions, even for compactly supported deformations.

Lemma 8. *Let λ and λ' be near-contact 1-forms defined on a neighbourhood of a point $p \in Y$. Assume that λ and λ' both vanish at p , that*

$$\text{ind}_p(\lambda) = \text{ind}_p(\lambda'), \tag{5.2}$$

and that $\lambda \wedge d\lambda$ and $\lambda' \wedge d\lambda'$ induce the same orientation on a punctured neighbourhood of p . Let W be any neighbourhood of p on which both forms are defined. Then there exist nested neighbourhoods

$$p \in U \subset V \subset W$$

and a family of near-contact forms λ_t , $t \in [0, 1]$, such that:

- (1) $\lambda_0 = \lambda$ on W , $\lambda_t = \lambda$ on $W - V$ for all $t \in [0, 1]$, and $\lambda_1 = \lambda'$ on U ;
- (2) for each $t \in [0, 1]$, the form λ_t is near-contact on V , and p is the unique zero of λ_t in V .

Proof. To begin with, identify the neighbourhood W of p with the unit ball $B_1 \subset \mathbb{R}^3$, and let p correspond to the origin. Choose a flat metric g in \mathbb{R}^3 , let ∇ be its flat connection, let vol_g be its volume form, and let $r: \mathbb{R}^3 \rightarrow \mathbb{R}$ be the distance from the origin.

Using (5.2), we may assume that:

$$\nabla \lambda|_p = \nabla \lambda'|_p.$$

Then:

$$\nabla^2 \left([(1-t)\lambda + t\lambda'] \wedge d[(1-t)\lambda + t\lambda'] \right) \Big|_p = (1-t) \nabla^2(\lambda \wedge d\lambda) \Big|_p + t \nabla^2(\lambda' \wedge d\lambda') \Big|_p.$$

Let f and f' be defined as:

$$\lambda \wedge d\lambda = f \text{vol}_g, \quad \lambda' \wedge d\lambda' = f' \text{vol}_g.$$

Then:

$$(1-t) \nabla^2(\lambda \wedge d\lambda) \Big|_p + t \nabla^2(\lambda' \wedge d\lambda') \Big|_p = [(1-t) \nabla^2 f + t \nabla^2 f'] \text{vol}_g \Big|_p.$$

Since $\lambda \wedge d\lambda$ and $\lambda' \wedge d\lambda'$ induce the same orientation near p , let us choose that local orientation so that both f and f' are nonnegative. Each has a nondegenerate minimum at p , hence their convex combination does also. Thus, there exists $c > 0$ such that for all t :

$$(1-t)f + tf' \geq cr^2 \quad \text{near } p. \quad (5.3)$$

Choose $0 < \varepsilon, R < 1$. Let B_R and $B_{\varepsilon R}$ be the balls of radii R and εR . Let $\rho = \rho(r)$ be a nonincreasing cut-off function such that:

$$\rho|_{B_{\varepsilon R}} = 1, \quad \rho|_{B_1 - B_R} = 0.$$

Define:

$$\lambda_t = \lambda + t\rho(\lambda' - \lambda).$$

For $W = B_1$, $V = B_R$, and $U = B_{\varepsilon R}$, property (1) of the lemma is immediate. Let us show ε and R can be chosen so that (2) holds. We compute:

$$\begin{aligned} \lambda_t \wedge d\lambda_t &= \lambda \wedge d\lambda + t\rho(\lambda' - \lambda) \wedge d\lambda + t\lambda \wedge d(\rho(\lambda' - \lambda)) + t^2\rho(\lambda' - \lambda) \wedge d(\rho(\lambda' - \lambda)) = \\ &= (1-t\rho)\lambda \wedge d\lambda + t\rho\lambda' \wedge d\lambda' - t\rho(\lambda' - \lambda) \wedge d(\lambda' - \lambda) \\ &\quad - t d\rho \wedge \lambda \wedge (\lambda' - \lambda) + t^2\rho^2(\lambda' - \lambda) \wedge d(\lambda' - \lambda) = \\ &= [(1-\rho t)f + \rho t f'] \text{vol}_g - t d\rho \wedge \lambda \wedge (\lambda' - \lambda) - t\rho(1-t\rho)(\lambda' - \lambda) \wedge d(\lambda' - \lambda). \end{aligned}$$

Using $0 \leq \rho \leq 1$ and (5.3), choose $R > 0$ small enough so that:

$$(1-\rho t)f + \rho t f' \geq cr^2 \quad \text{on } B_R. \quad (5.4)$$

For a domain A , write $\|\cdot\|_{k,A}$ for the C^k -norm on A .

Using:

$$\lambda|_p = \lambda'|_p = 0, \quad \nabla(\lambda' - \lambda)|_p = 0, \quad d\lambda|_p = d\lambda'|_p = 0,$$

there exists $C > 0$ such that for all r ,

$$\|\lambda\|_{0,B_r} \leq Cr, \quad \|\lambda'\|_{0,B_r} \leq Cr, \quad \|\lambda' - \lambda\|_{0,B_r} \leq Cr^2, \quad \|d\lambda\|_{0,B_r} \leq Cr, \quad \|d\lambda'\|_{0,B_r} \leq Cr. \quad (5.5)$$

Let f_t be defined as:

$$\lambda_t \wedge d\lambda_t = f_t \text{vol}_g.$$

On B_R :

$$\begin{aligned}
f_t &\geq [(1-\rho t)f + \rho t f'] - t\|d\rho\|_{0,B_r}\|\lambda\|_{0,B_r}\|\lambda' - \lambda\|_{0,B_r} - t\rho(1-t\rho)\|\lambda' - \lambda\|_{0,B_r}\|d(\lambda' - \lambda)\|_{0,B_r} \geq \\
&\stackrel{(5.4)}{\geq} cr^2 - \|d\rho\|_{0,B_r}\|\lambda\|_{0,B_r}\|\lambda' - \lambda\|_{0,B_r} - \|\lambda' - \lambda\|_{0,B_r}\|d(\lambda' - \lambda)\|_{0,B_r} \geq \\
&\stackrel{(5.5)}{\geq} cr^2 - C^2 r^3 \|d\rho\|_{0,B_r} - 2C^2 r^3.
\end{aligned}$$

Fix R . For each $\delta > 0$, there exists $\varepsilon > 0$, depending on δ but not on R , and a nonincreasing $\rho = \rho(r)$ such that:

$$\rho = 1 \text{ on } r \leq \varepsilon R, \quad \rho = 0 \text{ on } r \geq R, \quad \partial_r \rho \geq -\delta/r.$$

In the logarithmic coordinate $t = \ln r$, this is immediate: we want $\rho = 1$ for $t \leq \ln R - \ln(1/\varepsilon)$ and $\rho = 0$ for $t \geq \ln R$, with $\partial_t \rho \geq -\delta$.

With such a choice:

$$f_t \geq cr^2 - (c/2)r^2 - 2C^2 r^3 \quad \text{on } B_R.$$

Thus, for $R > 0$ small enough, $f_t \geq \frac{c}{3}r^2$ on B_R . This completes the proof. \square

Taubes [20] constructs a near-contact form on \mathbb{R}^3 with a zero at $0 \in \mathbb{R}^3$ such that every sufficiently small punctured neighbourhood of 0 contains an overtwisted disc. For completeness, we give a proof of this result by constructing a sequence of such forms.

Lemma 9. *There exists a near-contact form λ defined in a neighbourhood of $0 \in \mathbb{R}^3$ such that λ has a zero at 0, and such that for each small enough neighbourhood $0 \in U \subset \mathbb{R}^3$, the punctured set $U - \{0\}$ contains an overtwisted disc.*

Proof. Choose a flat metric g in \mathbb{R}^3 , let ∇ be its flat connection, and let vol_g be its volume form.

Fix any near-contact form λ defined in a neighbourhood of $0 \in \mathbb{R}^3$ such that λ has a zero at 0 and no other zeros. Assume that the orientations induced by vol_g and $\lambda \wedge d\lambda$ agree.

Consider the sphere $S_\varepsilon \subset \mathbb{R}^3$ of radius ε . We shall show that there exists a homogeneous cubic polynomial C such that, for each small enough ε , the 1-form

$$\lambda + dC,$$

when restricted to S_ε , contains an overtwisted disc. Since C is homogeneous cubic and

$$(\lambda + dC) \wedge d(\lambda + dC) = \lambda \wedge d\lambda + dC \wedge d\lambda,$$

the form $\lambda + dC$ remains near-contact in a neighbourhood of the origin.

To proceed, write λ as:

$$\lambda = da + \mu,$$

where $a: \mathbb{R}^3 \rightarrow \mathbb{R}$ is given by

$$a(x) = \frac{1}{2}\langle Ax, x \rangle, \quad x \in \mathbb{R}^3,$$

with A defined as in (5.1). By changing the sign of λ if needed, we may assume that in some orthonormal frame the quadratic form a is given by

$$a(x) = \frac{1}{2}(x_1^2 + x_2^2 - x_3^2).$$

Define the dilation $\varphi_\varepsilon: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ as:

$$\varphi_\varepsilon(x) = \varepsilon x, \quad x \in \mathbb{R}^3.$$

Consider the unit sphere $S_1 \subset \mathbb{R}^3$, and let V be a small neighbourhood of S_1 . Since both μ and $\nabla\mu$ vanish at the origin $0 \in \mathbb{R}^3$, there exists a constant $D > 0$, depending only on μ , such that for each ε ,

$$\|\varphi_\varepsilon^* \mu\|_{1,V} \leq D\varepsilon^3. \quad (5.6)$$

We compute:

$$\varphi_\varepsilon^*(\lambda + dC) = \varepsilon^2 \left(da + \frac{\varphi_\varepsilon^* \mu}{\varepsilon^2} + \varepsilon dC \right).$$

In particular, the $\varepsilon^{-2} \varphi_\varepsilon^*(\lambda + dC)$ is an ε -small C^1 -perturbation of da . We now choose this perturbation in a controlled way.

Restrict a to the unit sphere S_1 . The critical set of $a|_{S_1}$ is as follows:

- (1) The two points $x_3 = \pm 1$, which are nondegenerate critical points.
- (2) The circle $x_3 = 0$, which consists entirely of critical points. This circle is Morse-Bott for $a|_{S_1}$. Indeed:

$$\nabla_{x_3} da = -dx_3.$$

Choose C so that the 2-form

$$dC \wedge \nabla_{x_3} da = -dC \wedge dx_3,$$

when restricted to the unit sphere S_1 , does not vanish at the points of the big circle $x_3 = 0$. It suffices to take $C = C(x_1, x_2)$, independent of x_3 , and arranged so that $dC \neq 0$ unless $x_1 = x_2 = 0$.

Since

$$\varphi_\varepsilon^*(\mu + dC) \wedge \nabla_{x_3} \varphi_\varepsilon^* da = \varepsilon^2 \varphi_\varepsilon^* \mu \wedge \nabla_{x_3} da + \varepsilon^5 dC \wedge \nabla_{x_3} da,$$

it follows from (5.6) that, after multiplying C by a sufficiently large constant if needed, the 2-form

$$\varphi_\varepsilon^*(\mu + dC) \wedge \nabla_{x_3} \varphi_\varepsilon^* da,$$

when restricted to S_1 , does not vanish at any point of the circle $x_3 = 0$.

With this choice of C , and for ε small enough, the 1-form

$$\varepsilon^{-2} \varphi_\varepsilon^*(\lambda + dC) = da + \frac{\varphi_\varepsilon^* \mu}{\varepsilon^2} + \varepsilon dC,$$

when restricted to S_1 , has exactly two transverse zeros p_+ and p_- . These are small perturbations of the points $x_3 = 1$ and $x_3 = -1$, respectively.

Set:

$$\lambda_\varepsilon = da + \frac{\varphi_\varepsilon^* \mu}{\varepsilon^2} + \varepsilon dC.$$

We will show that for each sufficiently small but nonzero ε , the restriction $\lambda_\varepsilon|_{S_1}$ admits an overtwisted disc. Consider the Euler field:

$$E = x_1 \partial_{x_1} + x_2 \partial_{x_2} + x_3 \partial_{x_3},$$

and define ω by $\iota_E \text{vol}_g = \omega$. Its restriction to S_1 is nondegenerate. Restrict ω and λ_ε to S_1 , and let X_ε be the ω -dual of $\lambda_\varepsilon|_{S_1}$; that is, $\iota_X \omega = \lambda_\varepsilon|_{S_1}$. Since λ_ε is C^1 -close to da , the vector field X_ε is C^1 -close to X_0 .

X_ε has exactly two zeros p_+ and p_- on S_1 , both transverse. Because a has nondegenerate minima at $x_3 = \pm 1$, the linearization of X_0 at these points has a pair of imaginary eigenvalues; therefore,

the linearization of X_ε at p_\pm cannot have purely real eigenvalues, so neither p_+ nor p_- is a saddle. We will show that:

$$\operatorname{div} X_\varepsilon(p_+) < 0, \quad \operatorname{div} X_\varepsilon(p_-) < 0 \quad \text{for all small enough but nonzero } \varepsilon. \quad (5.7)$$

Let f_ε be defined by:

$$\lambda_\varepsilon \wedge d\lambda_\varepsilon = f_\varepsilon \operatorname{vol}_g.$$

We compute:

$$f_\varepsilon \omega = f_\varepsilon \iota_E \operatorname{vol}_g = \iota_E(\lambda_\varepsilon \wedge d\lambda_\varepsilon) = \lambda_\varepsilon(E) d\lambda_\varepsilon - \lambda_\varepsilon \wedge \iota_E d\lambda_\varepsilon.$$

Restricting to S_1 :

$$f_\varepsilon \omega = \lambda_\varepsilon(E) (\operatorname{div} X) \omega - \lambda_\varepsilon \wedge \iota_E d\lambda_\varepsilon.$$

Restricting to p_+ and p_- (where $\lambda_\varepsilon = 0$):

$$f_\varepsilon(p_+) = \lambda_\varepsilon(E)(p_+) \operatorname{div} X_\varepsilon(p_+), \quad f_\varepsilon(p_-) = \lambda_\varepsilon(E)(p_-) \operatorname{div} X_\varepsilon(p_-).$$

Since $f_\varepsilon > 0$ everywhere on S_1 and

$$\lambda_\varepsilon(E)(p_+) \rightarrow -1, \quad \lambda_\varepsilon(E)(p_-) \rightarrow -1 \quad \text{as } \varepsilon \rightarrow 0,$$

we obtain (5.7). Consequently, both p_+ and p_- are sinks of X_ε , and X_ε has no other zeros on S_1 . Hence, X_ε must admit a limiting cycle. This completes the proof. \square

Recall the following theorem of Eliashberg (Theorem 3.1.1 in [3]).

Theorem ([3]). *Let M be a compact 3-manifold and let $A \subset M$ be a closed subset such that $M - A$ is connected. Let K be a compact space and $L \subset K$ a closed subspace. Let $\{\xi_t\}_{t \in K}$ be a family of 2-plane distributions which are contact for all $t \in L$ and are contact near A for all $t \in K$. Assume there exists an embedded disc $\Delta \subset M - A$ such that ξ_t is contact near Δ and (Δ, ξ_t) is an overtwisted disc for all $t \in K$. Then there exists a family of contact structures $\{\xi'_t\}_{t \in K}$ on M such that ξ'_t agrees with ξ_t near A for all $t \in K$, and agrees with ξ_t everywhere for all $t \in L$.*

Proposition 2. *Let Y be a closed 3-manifold, and let λ_0 and λ_1 be near-contact forms on Y with the same zero set:*

$$\lambda_0^{-1}(0) = \lambda_1^{-1}(0) = \{p_1, \dots, p_k\}, \quad k > 0.$$

Suppose λ_0 and λ_1 are homotopic as nonzero 1-forms on $Y - \{p_1, \dots, p_k\}$, and that $\lambda_0 \wedge d\lambda_0$ and $\lambda_1 \wedge d\lambda_1$ induce the same orientation on Y . Then there exists a near-contact homotopy λ_t , $t \in [0, 1]$, between λ_0 and λ_1 on the whole of Y .

Proof. Since λ_0 and λ_1 are homotopic as nonvanishing 1-forms on $Y - \{p_1, \dots, p_k\}$, we have for each zero p_i :

$$\operatorname{ind}_{p_i}(\lambda_0) = \operatorname{ind}_{p_i}(\lambda_1).$$

Using Lemmas 9 and 8, we may assume that there exist neighbourhoods $U_i \ni p_i$ such that $\lambda_0|_{U_i} = \lambda_1|_{U_i}$, and that (U_i, λ_0) contains an overtwisted disc. Choose closed neighbourhoods $A_i \subset U_i$ of p_i such that $U_i - A_i$ still contains an overtwisted disc. Set $A = A_1 \cup \dots \cup A_k$; shrinking the A_i if necessary, we may assume $Y - A$ is connected. We can now construct a family of 1-forms α_t , $t \in [0, 1]$, such that:

- (1) $\alpha_0 = \lambda_0$, $\alpha_1 = \lambda_1$, and $\alpha_t = \lambda_0 = \lambda_1$ on each U_i ;
- (2) for all t , the form α_t is nowhere zero on $Y - \{p_1, \dots, p_k\}$.

The lemma now follows from Eliashberg's theorem. \square

6. Near-symplectic forms. In this section we review the notion of a near-symplectic form; see [2, 20, 12, 10, 5] for background. Our goal is to prove Proposition 3.

Let X be a 4-manifold. A 2-form ω on X is called a *near-symplectic form* if:

- (1) $d\omega = 0$;
- (2) ω is symplectic away from its zero set, i.e., for each $x \in X$ with $\omega_x \neq 0$, $\omega_x \wedge \omega_x \neq 0$;
- (3) the zero set $Z(\omega) = \{x \in X \mid \omega_x = 0\}$ is a smooth 1-dimensional submanifold (a union of embedded circles and arcs), and for each $x \in Z$ and $\xi \in T_X|_x$,

$$(\nabla_\xi \omega)_x \wedge (\nabla_\xi \omega)_x \neq 0.$$

If X has boundary, Z is additionally required to be transverse to ∂X .

This definition is equivalent to the following. Let ω be a closed 2-form with zero set $Z = \omega^{-1}(0)$. Then ω is near-symplectic iff Z is a smooth 1-manifold and there exists a volume form vol on X such that:

$$\omega \wedge \omega = f \text{vol},$$

where $f > 0$ on $X - Z$ and Z is a Morse-Bott critical submanifold of f ; that is, $df = 0$ along Z and the Hessian of f is nondegenerate on the normal bundle of Z .

Endow X with a Riemannian metric and let ω be a closed SD form. Viewing ω as a section of $\Lambda^2_+ T_X^*$, assume it is transverse to the zero section; that is, for each $x \in Z = \omega^{-1}(0)$, $\nabla \omega_x$ has rank 3. Then Z is a smooth 1-manifold, and ω is necessarily near-symplectic: for each $x \in Z$ and $\xi \in T_X|_x$, the SD form $\nabla_\xi \omega_x$ is nonzero; hence, its wedge square is also nonzero.

Auroux, Donaldson, and Katzarkov proved a converse (Proposition 1 in [2]):

Theorem ([2]). *Suppose ω is a near-symplectic form on X . Then there is a Riemannian metric g on X such that ω is self-dual with respect to g .*

A smooth hypersurface $\Sigma \subset X$ is called ω -convex if there exists a vector field V defined near Σ such that $L_V \omega = \omega$, V is transverse to Σ , and V points outward along Σ from the compact region it bounds. Such a vector field V is called an ω -Liouville vector field. An ω -convex surface Σ is necessarily transverse to Z , and the primitive $\lambda = \iota_V \omega$ restricts to a near-contact form on Σ . The same applies to ω -concave surfaces, obtained by replacing $L_V \omega = \omega$ with $L_V \omega = -\omega$.

Lemma 10 (Near-symplectic cobordism). *Let λ_t , $t \in [0, 1]$, be a smooth family of near-contact forms on a closed 3-manifold Y . Then there exists a near-symplectic form ω on $W = [0, 1] \times Y$ and ω -Liouville vector fields V_0 and V_1 , defined near $\{0\} \times Y$ and $\{1\} \times Y$, respectively, such that V_0 is transverse to $\{0\} \times Y$ and points into W , V_1 is transverse to $\{1\} \times Y$ and points out of W , and*

$$\iota_{V_0} \omega|_{\{0\} \times Y} = c_1 \lambda_0, \quad \iota_{V_1} \omega|_{\{1\} \times Y} = c_2 \lambda_1,$$

for some constants $c_1, c_2 > 0$. In particular, $\{0\} \times Y$ is ω -concave and $\{1\} \times Y$ is ω -convex.

Proof. To begin with, assume that all λ_t vanish at the same finite set of points $p_1, \dots, p_k \in Y$, and that for each p_i the linearization $(\nabla \lambda_t)_{p_i}$ is independent of t . We also assume the family is constant near the endpoints: there exists $\varepsilon > 0$ such that $\lambda_t = \lambda_0$ for $t \in [0, \varepsilon]$ and $\lambda_t = \lambda_1$ for $t \in [1 - \varepsilon, 1]$.

We use the same parameter $t \in [0, 1]$ as the coordinate on the first factor of $W = [0, 1] \times Y$. For each t , regard λ_t as a 1-form on the slice $\{t\} \times Y$. These assemble to a smooth 1-form λ on W , with no dt -component:

$$\lambda(\partial_t) = 0, \quad \lambda|_{\{t\} \times Y} = \lambda_t.$$

We seek a function $h = h(t)$ such that the 2-form

$$d(h\lambda)$$

is near-symplectic on W . Using the decomposition

$$d = dt \wedge \partial_t + d_Y,$$

we compute:

$$d(h\lambda) = dh \wedge \lambda + h d\lambda = \dot{h} dt \wedge \lambda_t + h dt \wedge \dot{\lambda}_t + h d_Y \lambda_t,$$

where $\dot{\lambda}_t = \partial_t \lambda_t$. Thus:

$$d(h\lambda) \wedge d(h\lambda) = 2h dt \wedge \left(\dot{h} \lambda_t \wedge d_Y \lambda_t + h \dot{\lambda}_t \wedge d_Y \lambda_t \right).$$

Choose a volume form vol on Y and a compatible connection ∇ . Define smooth families of functions $f_t, g_t: Y \rightarrow \mathbb{R}$ as:

$$\lambda_t \wedge d_Y \lambda_t = f_t \text{ vol}, \quad \dot{\lambda}_t \wedge d_Y \lambda_t = g_t \text{ vol}.$$

Setting $h(t) = e^{Ct}$ with $C > 0$, we find:

$$d(h\lambda) \wedge d(h\lambda) = 2e^{2Ct} dt \wedge (Cf_t + g_t) \text{ vol}.$$

Let $p_i \in Y$ be a zero of λ_t . By Lemma 6, $(d_Y \lambda_t)_{p_i} = 0$. Since $(\nabla \lambda_t)_{p_i}$ is independent of t , it follows that $(\nabla \dot{\lambda}_t)_{p_i} = 0$. Thus, for each zero p_i and each t ,

$$g_t(p_i) = 0, \quad (dg_t)_{p_i} = 0, \quad (\nabla^2 g_t)_{p_i} = 0.$$

On the other hand, the near-contact condition on λ_t gives $f_t(p_i) = 0$, $(df_t)_{p_i} = 0$, and the Hessian $(\nabla^2 f_t)_{p_i}$ is positive-definite. Hence, $Cf_t + g_t$ has a nondegenerate minimum at each zero p_i . It follows that there exist neighbourhoods $U_i \ni p_i$ such that for each $C > 1$ and each $t \in [0, 1]$, the function $Cf_t + g_t$ is positive on $U_i - \{p_i\}$. Set $U = U_1 \cup \dots \cup U_k$. Since g_t is uniformly bounded and $Y - U$ is compact, it follows that for all large enough $C > 0$, the function $Cf_t + g_t$ is positive on $Y - \{p_1, \dots, p_k\}$ for all $t \in [0, 1]$.

Define a function $f: W \rightarrow \mathbb{R}$ as:

$$f(t, y) = e^{2Ct} (Cf_t(y) + g_t(y)).$$

Then $f > 0$ away from the arcs $[0, 1] \times \{p_i\}$, and each such arc is a Morse-Bott critical submanifold of f . Hence, $d(e^{Ct}\lambda)$ is near-symplectic on W .

Near $t = 1$ the family λ_t is constant, so $d(e^{Ct}\lambda_t)$ coincides with $d(e^{Ct}\lambda_1)$ on a collar of $\{1\} \times Y$. Moreover,

$$L_{\partial_t} d(e^{Ct}\lambda_t) = C d(e^{Ct}\lambda_t),$$

so $C^{-1}\partial_t$ is a Liouville vector field for $d(e^{Ct}\lambda_t)$ near ∂W . This completes the proof. \square

Taubes [19] and Perutz [12] show that a near-symplectic form ω canonically orients its zero set $Z = \omega^{-1}(0)$. For completeness, we recall their construction.

Let ω be a near-symplectic form on X , and let $Z = \omega^{-1}(0)$ be its 1-dimensional zero set. For each $x \in Z$, the range of $\nabla \omega_x$ is a 3-plane $P_x \subset \Lambda^2 T_X^*|_x$, on which the wedge pairing is positive-definite. Let $N_x \subset T_X|_x$ be a complementary subspace to $T_Z|_x$. Then $\nabla \omega_x: N_x \rightarrow P_x$ is an isomorphism. Let $\partial_Z \in T_Z|_x$ be a tangent vector. Define the bilinear form:

$$A(\xi, \eta) = (\iota_{\partial_Z}(\nabla_\xi \omega))_x(\eta), \quad \xi, \eta \in N_x. \quad (6.1)$$

Since P_x is positive-definite, A is nondegenerate, and thus the sign of $\det A$ is defined. The canonical orientation on Z is defined by requiring that ∂_Z be chosen so that $\det A > 0$.

$d\omega = 0$ and $(\nabla_{\partial_Z}\omega)_x = 0$ imply that A is symmetric.

Recall an additional property of A , found in [19, 12]. Orient X so that $\omega \wedge \omega \geq 0$. Let $x \in Z$ and let P_x be the range of $\nabla\omega_x$. The maximal positive-definite 3-plane P_x determines a conformal class of metrics g_x at $T_X|_x$ such that $P_x = \Lambda_+^2 T_X^*|_x$. Choose $N_x \subset T_X|_x$ to be the g_x -orthogonal complement of $T_Z|_x$. With ∂_Z and A as above, the form A is g_x -traceless: for each g_x -orthonormal basis e_1, e_2, e_3 of N_x ,

$$\sum_{i=1}^3 A(e_i, e_i) = 0.$$

Let us now specialize to \mathbb{R}^4 and consider near-symplectic forms whose coefficients are homogeneous of degree 1. Let ω be such a form. Since ω is homogeneous of degree 1 and near-symplectic, its zero set $Z = \omega^{-1}(0)$ is a line in \mathbb{R}^4 . Let us classify these forms up to deformation; this is a reformulation of [19, 12].

Lemma 11. *Let \mathbb{R}^4 be oriented and let $\ell \subset \mathbb{R}^4$ be a line. Consider the space of near-symplectic forms ω on \mathbb{R}^4 with homogeneous coefficients of degree 1, such that $\omega \wedge \omega \geq 0$ and whose zero set is exactly ℓ . Then this space has precisely two connected components, distinguished by the orientation they induce on ℓ .*

Proof. To begin with, choose a linear metric g on \mathbb{R}^4 such that the range of $\nabla\omega_0$ consists of g -self-dual forms. Since ω has homogeneous linear coefficients, it follows that $\omega_x = (\nabla_x\omega)_0$ for all $x \in \mathbb{R}^4$; in particular, ω is g -self-dual. The space of linear metrics is convex, so it suffices to classify homogeneous linear self-dual forms ω on \mathbb{R}^4 whose zero set is ℓ . We show that this space has two connected components.

Let $N \subset \mathbb{R}^4$ be the g -orthogonal complement of ℓ at the origin. Fix $\partial_\ell \in \ell$, and let ∇ be the flat connection associated to g . As in (6.1), define a bilinear form:

$$A(\xi, \eta) = (\iota_{\partial_\ell}(\nabla_\xi\omega))_0(\eta), \quad \xi, \eta \in N.$$

Then A is nondegenerate, symmetric, and traceless. For a homogeneous linear self-dual form ω vanishing exactly on ℓ , the form A determines ω uniquely. Conversely, any nondegenerate symmetric traceless form A on N arises from such an ω .

The space of nondegenerate symmetric traceless bilinear forms on N has two connected components, distinguished by the sign of $\det A$. This completes the proof. \square

Let us now specialize to the case when X is a ball in \mathbb{R}^4 .

Proposition 3. *Let $B \subset \mathbb{R}^4$ be the closed unit ball with boundary $\Sigma = \partial B$, and let ω and ω' be near-symplectic forms defined in a neighbourhood of B such that $\omega \wedge \omega$ and $\omega' \wedge \omega'$ induce the same orientation on B . Suppose that Σ is both ω -convex and ω' -convex, and let λ and λ' be the induced near-contact forms on Σ . Let*

$$Z = \omega^{-1}(0), \quad Z' = \omega'^{-1}(0).$$

Assume that Z and Z' are unions of arcs (i.e. have no circular components) and that Z and Z' have the same number of connected components. Then λ and λ' are homotopic through near-contact forms on Σ .

Moreover, suppose that the antipodal \mathbb{Z}_2 -action on B preserves ω and ω' , and that there exist \mathbb{Z}_2 -invariant ω - and ω' -Liouville vector fields near Σ making Σ ω - and ω' -convex. Then the above homotopy may be chosen \mathbb{Z}_2 -equivariant, so the induced near-contact forms on the quotient $\Sigma/\mathbb{Z}_2 \cong \mathbb{RP}^3$ are homotopic.

Proof. For simplicity, we treat only the non-equivariant statement; the antipodal case is identical, with all constructions carried out \mathbb{Z}_2 -equivariantly.

Let L be a closed manifold and define its cone:

$$C(L) = ([0, 1] \times L) / \sim, \quad (t_1, x_1) \sim (t_2, x_2) \iff x_1 = x_2, t_1 = t_2 = 1.$$

Let p_* be the apex and $L_0 = \{0\} \times L \subset C(L)$.

A link in $C(L)$ is a finite union of disjoint embedded arcs $\gamma_i \subset C(L) - \{p_*\}$ with endpoints in L_0 , each transverse to L_0 ; an oriented link is such a link with an orientation on each component. Two oriented links are isotopic with free endpoints in L_0 if they are connected by a smooth 1-parameter family of oriented links as above. It is standard that if $\dim L \geq 3$, any two oriented links in $C(L)$ are isotopic with free endpoints in L_0 ; this depends only on dimension of L , not on $\pi_1(L)$.

Now strengthen this as follows. Let $D_1, \dots, D_k \subset L_0$ be pairwise disjoint closed 3-discs. Each cone $C(D_i)$ embeds in $C(L)$, as does the wedge

$$C(D_1) \vee \dots \vee C(D_k) \subset C(L),$$

which is a strong deformation retract of $C(L)$. Let $Z = \gamma_1 \cup \dots \cup \gamma_k \subset C(L)$ be a link. We may isotope Z so that there are disjoint $D_i \subset L_0$ such that $\gamma_i \subset C(D_i)$. Then $C(L) - Z$ deformation retracts onto the wedge

$$(C(D_1) - \gamma_1) \vee \dots \vee (C(D_k) - \gamma_k).$$

Each $C(D_i)$ is a pointed 4-ball; therefore, $C(D_i) - \gamma_i$ retracts onto a pointed 2-sphere. Hence, $C(L) - Z$ deformation retracts (relative to p_*) onto a wedge of k 2-spheres. Again, only the dimension of L is used.

Let us apply this with $L = \Sigma = \partial B$, so that $C(L)$ is B , and p_* is the origin of B . Let ω and ω' be near-symplectic forms on B with zero sets

$$Z = \omega^{-1}(0), \quad Z' = \omega'^{-1}(0).$$

Both Z and Z' are oriented links in B and have the same number of components. By the above link isotopy statement, there is an isotopy of Z' to Z . Thus, we may assume that $Z' = Z$ and the canonical orientations on Z induced by ω and ω' coincide.

Set:

$$Z \cap \Sigma = \{p_1, \dots, p_k\}, \quad \Sigma = \partial B.$$

Let V and V' be Liouville vector fields for ω and ω' , defined in tubular neighbourhoods of Σ and transverse to Σ . Set:

$$\lambda = \iota_V \omega, \quad \lambda' = \iota_{V'} \omega'.$$

Since $\omega \wedge \omega$ and $\omega' \wedge \omega'$ are cooriented in B , the 3-forms $\lambda \wedge d\lambda$ and $\lambda' \wedge d\lambda'$ are cooriented on Σ . We shall show that λ and λ' are homotopic as non-vanishing 1-forms on $\Sigma - \{p_1, \dots, p_k\}$. By Lemma 2, this yields a homotopy through near-contact forms.

In fact, we construct a homotopy between ω and ω' through nondegenerate 2-forms on $B - Z$; this induces the desired homotopy between λ and λ' .

Consider the bundle $\mathcal{N} \rightarrow B$ of nondegenerate 2-forms on B . It has two connected components, distinguished by the orientation induced on B . Let $\mathcal{N}^+ \subset \mathcal{N}$ be the connected component of those forms which induce the same orientation as ω . Since ω and ω' are nondegenerate and compatible with the orientation, they define sections $\omega, \omega': B - Z \rightarrow \mathcal{N}^+|_{B-Z}$.

$B - Z$ deformation retracts onto a wedge of pieces $C(D_i) - \gamma_i$, and each $C(D_i) - \gamma_i$ retracts onto a small 2-sphere S_i linking γ_i . Thus, it suffices to show that ω and ω' are homotopic as sections of $N^+ \rightarrow S_i$ for each S_i .

Fix i . Let U_i be an arbitrary small neighbourhood of an interior point of γ_i . By a small C^1 -perturbation, arrange that ω and ω' coincide with their linear parts in U_i . Deform S_i so that $S_i \subset U_i$. Since ω and ω' induce the same orientation on γ_i , it follows from Lemma 11 that there is a near-symplectic deformation of ω to ω' on $U_i - \gamma_i$. This completes the proof. \square

Proposition 3 and Lemma 10 allow us to introduce the following near-symplectic surgery. Let $B \subset \mathbb{R}^4$ be the unit ball with boundary $\Sigma = \partial B$. Let ω, ω' be near-symplectic forms defined in a neighbourhood of B such that ω and ω' induce the same orientation on B . Suppose that Σ is both ω - and ω' -convex, and that both $Z = \omega^{-1}(0)$ and $Z' = \omega'^{-1}(0)$ are unions of arcs and have the same number of connected components. Then there exist a smaller ball $B_0 \subset B$ and a near-symplectic form $\hat{\omega}$ defined in a neighbourhood of B such that $\hat{\omega} = \omega$ near ∂B and such that $(B_0, \hat{\omega})$ is diffeomorphic to $(B, c\omega')$ for some constant $c > 0$.

Moreover, suppose that the antipodal \mathbb{Z}_2 -action on B preserves ω and ω' , and there exist \mathbb{Z}_2 -invariant ω - and ω' -Liouville vector fields near Σ making Σ ω - and ω' -convex. Then $\hat{\omega}$ can be chosen \mathbb{Z}_2 -invariant.

7. Proof of Proposition 1. It suffices to treat the case $n = 1$; the general case is no harder. Let (M, g) be an oriented Riemannian 4-orbifold with a double point p . Choose a neighbourhood V of p equipped with an isometry $V \simeq (B/\{\pm 1\}, g_0)$, where $B \subset \mathbb{R}^4$ is a ball centered at the origin and g_0 is the flat metric. Let ψ be a closed SD form on M . We may assume $\psi(p) \neq 0$, otherwise there is nothing to prove. By Darboux' theorem, (V, ψ) is diffeomorphic to the quotient $(B, \omega_0)/\{\pm 1\}$, where ω_0 is the standard symplectic form on B .

Proposition 1 follows from the following stronger statement.

Proposition 4. *Let $B \subset \mathbb{R}^4$ be the unit ball centered at the origin, and let g_0 be the flat metric. Then there exists a near-symplectic form ω in a neighbourhood of B such that ω vanishes at $0 \in B^4$, ω is invariant under the antipodal map, ω coincides with the standard symplectic form in a neighbourhood of ∂B , and ω is g_0 -self-dual in a neighbourhood of the origin.*

If this is the case, replace ψ on V by such an ω . Then the theorem of Auroux-Donaldson-Katzarkov applies to keep ψ self-dual for some metric that agrees with g_0 near p . The rest of this section is the proof of Proposition 4.

Lemma 12. *Let $F = (f_1, f_2, f_3): \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be defined as:*

$$\begin{aligned} f_1(x) &= 2x_1x_3 - 2x_2x_4 - x_3x_4, \\ f_2(x) &= -x_1(4x_2 + x_3), & H(x_1, x_4) &= 3x_1^2 - x_1x_4 - x_4^2. \\ f_3(x) &= x_2^2 + x_3^2 - H(x_1, x_4), \end{aligned}$$

Then for each $\varepsilon \geq 0$,

$$F^{-1}(0, 0, -\varepsilon) = \{x \in \mathbb{R}^4 : x_2 = x_3 = 0, H(x_1, x_4) = \varepsilon\}.$$

For $\varepsilon > 0$, dF has full rank on $F^{-1}(0, 0, -\varepsilon)$, and for $\varepsilon = 0$, it has full rank on $F^{-1}(0, 0, 0) - \{0\}$.

Set:

$$\omega = f_1\omega_1 + f_2\omega_2 + f_3\omega_3, \tag{7.1}$$

where

$$\omega_1 = dx_1 \wedge dx_2 + dx_3 \wedge dx_4, \quad \omega_2 = dx_1 \wedge dx_3 - dx_2 \wedge dx_4, \quad \omega_3 = dx_1 \wedge dx_4 + dx_2 \wedge dx_3.$$

Then ω is closed and self-dual with respect to the standard flat metric and orientation. In particular, ω is near-symplectic on $\mathbb{R}^4 - \{0\}$, and $\omega + \varepsilon\omega_3$ is near-symplectic on \mathbb{R}^4 for all $\varepsilon > 0$.

Proof. All statements follow by a direct computation. \square

Lemma 13. *Let ω be as in (7.1). Then there exists a small, antipodal-invariant perturbation ω_ε of ω such that $\omega_\varepsilon = \omega$ outside a neighbourhood of the origin, ω_ε is near-symplectic on all of \mathbb{R}^4 , and its zero set $\omega_\varepsilon^{-1}(0)$ has exactly two non-compact connected components.*

Proof. Let

$$E = \sum_{i=1}^4 x_i \partial_{x_i}$$

be the Euler field. Let ω_3 be as in (7.1). Set:

$$\lambda = \frac{1}{4} \iota_E \omega, \quad \mu = \frac{1}{2} \iota_E \omega_3,$$

so that $d\lambda = \omega$ and $d\mu = \omega_3$.

Let $\rho: \mathbb{R}^4 \rightarrow \mathbb{R}$ be a radial cut-off function such that $\rho(x) = \rho(-x)$, $\rho = 1$ on a ball U about the origin, and $\rho = 0$ outside a slightly larger ball. Set:

$$\omega_\varepsilon = d(\lambda + \varepsilon\rho\mu).$$

Since λ, μ, ρ are invariant under $x \mapsto -x$, so is ω_ε . On the region $\rho = 1$ (in particular, on U),

$$\omega_\varepsilon = \omega + \varepsilon\omega_3,$$

which is near-symplectic for all $\varepsilon > 0$. On $\rho = 0$, $\omega_\varepsilon = \omega$, which is also near-symplectic. For all small ε , ω_ε remains near-symplectic on the support of $d\rho$. Hence, ω_ε is near-symplectic on all of \mathbb{R}^4 .

$\omega^{-1}(0)$ is a union of two lines through the origin, and the zero set of $\omega + \varepsilon\omega_3$ is a small smoothing of this cross into two branches of the hyperbola $H(x_1, x_4) = \varepsilon$. Hence, for all small $\varepsilon > 0$, $\omega_\varepsilon^{-1}(0) \cap U$ consists exactly of 2 arcs, each with two ends on ∂U . Let K be a closed neighbourhood of $\mathbb{R}^4 - U$, strictly larger than $\mathbb{R}^4 - U$. Then $\omega_\varepsilon - \omega$ has compact support in K and is C^1 -small. Hence, for all small $\varepsilon > 0$, $\omega_\varepsilon^{-1}(0) \cap (\mathbb{R}^4 - U)$ consists exactly of 4 rays. These rays attach to the two arcs to produce two connected components. This completes the proof. \square

Proof of Proposition 4. Consider \mathbb{R}^4 with the standard symplectic form ω_{st} and the unit ball $B \subset \mathbb{R}^4$. Let us construct the desired near-symplectic form by a sequence of modifications of ω_{st} . The first step uses the following result of Taubes. See [20].

Theorem ([20]). *There exists a near-symplectic form ω on \mathbb{R}^4 such that its zero set $\omega^{-1}(0)$ is a circle near the origin and ω coincides with a standard symplectic form outside a neighbourhood of the origin.*

Let us apply Taubes' construction away from the origin. Pick a point $x \in \mathbb{R}^4 - \{0\}$ and a small neighbourhood $U \ni x$. Taubes' argument allows us to modify ω_{st} inside U to create a single zero circle in U . To keep antipodal invariance, perform the same modification near $-x$. Denote the resulting form by ω .

$Z = \omega^{-1}(0)$ is a circle $C \subset U$ and its antipodal image. Pick a point $p \in C$. Perform a small C^1 -perturbation of ω near p so that ω coincides with its linear part there. This immediately yields a ball neighbourhood $D \ni p$ whose boundary $\Sigma = \partial D$ is ω -convex. To keep antipodal invariance, perform the same perturbation near $-p$.

Choose a ball neighbourhood D_0 of the origin $0 \in B$ with ω -convex boundary $\Sigma_0 = \partial D_0$. Pick a point $a \in \Sigma_0$ and a point $b \in \Sigma$ with $\omega_b \neq 0$. Join a to b by a path δ . To keep antipodal invariance, also introduce the antipodal points $-a, -b$ and path $-\delta$.

Form the ω -convex connected sum of $D, D_0, (-D)$ along paths δ and $-\delta$. That is, choose an antipodal-invariant, ω -convex neighbourhood

$$X \supset D \cup D_0 \cup (-D).$$

(X, ω) is a near-symplectic 4-ball with convex boundary embedded in (B, ω) , and the embedding is antipodal-invariant. $\omega^{-1}(0) \cap X$ consists of 2 arcs and no circular components.

Since X is a ball, it admits a near-symplectic form ω_ε as in Lemma 13. By construction, ∂X is ω_ε -convex. Both $\omega|_X$ and ω_ε are antipodal-invariant and may be assumed cooriented. Then, as explained at the end of §6, one can modify ω only near $X \subset B$ so that

$$\omega = c\omega_\varepsilon \quad \text{on } X$$

for some constant $c > 0$, and ω remains antipodal-invariant.

Finally, as in Lemma 13, deform $c\omega_\varepsilon$ to a homogeneous SD form (with respect to a flat metric) that vanishes at the fixed point of the antipodal map. The resulting form ω on B satisfies all properties required in Proposition 4, and the proof follows.

References.

- [1] M. Atiyah, V. Patodi, and I. Singer. Spectral asymmetry and riemannian geometry. I. *Mathematical Proceedings of the Cambridge Philosophical Society*, 77(1):43–69, 1975.
- [2] S. Auroux, D.; Donaldson and L. Katzarkov. Singular lefschetz pencils. *Geom. Topol.*, 9:1043–1114, 2005.
- [3] Y. Eliashberg. Classification of overtwisted contact structures on 3-manifolds. *Invent. Math.*, 98(3):623–637, 1989.
- [4] G. Folland. Harmonic analysis of the de Rham complex on the sphere. *Crelle's Journal*, 1989(398):130–143, 1989.
- [5] D. Gay and R. Kirby. Constructing symplectic forms on 4-manifolds which vanish on circles. *Geom. Topol.*, 8(2):743–777, 2004.
- [6] P. Griffiths, and J. Harris, *Principles of algebraic geometry*. Wiley classics library. Wiley, New York, NY, 1994.
- [7] K. Honda. Transversality theorems for harmonic forms. *Rocky Mountain Journal of Mathematics*, 34(2), 2004.
- [8] C. LeBrun. Yamabe constants and the perturbed Seiberg–Witten equations. *Comm. Anal. Geom.*, 5(3):535–553, 1997.
- [9] C. LeBrun. Einstein metrics, four-manifolds, and differential topology. *Surveys in Differential Geometry*, 8(1):235–255, 2003.
- [10] Y. Lekili. Wrinkled fibrations on near-symplectic manifolds. *Geom. Topol.*, 13(1):277–318, 2009.
- [11] L. Nicolaescu. On the Cappell–Lee–Miller gluing theorem. *Pacific Journal of Mathematics*, 206(1):159–185, 2002.
- [12] T. Perutz. Zero-sets of near-symplectic forms. *J. Symp. Geom.*, 4(3):237–257, 2006.
- [13] C. Scaduto. Metric stretching and the period map for smooth 4-manifolds, 2023. preprint.
- [14] V. Shevchishin and G. Smirnov. Anti-self-dual blowups. To appear in *Math. Res. Lett.*
- [15] Y.-T. Siu. A simple proof of the surjectivity of the period map of K3 surfaces. *Manuscripta Mathematica*, 35(3):311–321, 1981.
- [16] C. Taubes. Self-dual Yang-Mills connections on non-self-dual 4-manifolds. *J. Diff. Geom.*, 17(1), 1982.

- [17] C. Taubes. Self-dual connections on 4-manifolds with indefinite intersection matrix. *J. Diff. Geom.*, 19(2), 1984.
- [18] C. Taubes. Gauge theory on asymptotically periodic 4-manifolds. *J. Diff. Geom.*, 25(3), 1987.
- [19] C. Taubes. Seiberg-Witten invariants and pseudo-holomorphic subvarieties for self-dual, harmonic 2-forms. *Geom. Topol.*, 3(1):167–210, 1999.
- [20] C. Taubes. A proof of a theorem of Luttinger and Simpson about the number of vanishing circles of a near-symplectic form on a 4-dimensional manifold. *Math. Res. Lett.*, 13(4):557–570, 2006.
- [21] A. Todorov. Applications of the Kahler-Einstein-Calabi-Yau metric to moduli of K3 surfaces. *Invent. Math.*, 61:251–266, 1980.

FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF WARMIA AND MAZURY,
UL. ŚLONECZNA 54, 10-710 OLSZTYN, POLAND

Email address: vsevolod@matman.uwm.edu.pl, shevchishin@gmail.com

MATHEMATICAL SCIENCES INSTITUTE, AUSTRALIAN NATIONAL UNIVERSITY, CANBERRA, AUSTRALIA

Email address: gleb.smirnov@anu.edu.au