
Evaluating A/B Testing Methodologies via Sample Splitting: Theory and Practice

Ryan Kessler* James McQueen Miikka Rokkanen
Amazon *Amazon* *Amazon*

Abstract

We develop a theoretical framework for sample splitting in A/B testing environments, where data for each test are partitioned into two splits to measure methodological performance when the true impacts of tests are unobserved. We show that sample-split estimators are generally biased for full-sample performance but consistently estimate sample-split analogues of it. We derive their asymptotic distributions, construct valid confidence intervals, and characterize the bias-variance trade-offs underlying sample-split design choices. We validate our theoretical results through simulations and provide implementation guidance for A/B testing products seeking to evaluate new estimators and decision rules.

1 Introduction

Measuring the impact of proposed improvements to the way in which A/B tests are analyzed and used to inform decision making is critical for large-scale A/B testing products. Over the last few decades, there have been several proposed improvements to A/B testing methodologies, including new estimators of impacts (for example, [1], [2], and [3]) and new rules for deciding when to launch (for example, [4], [5] and [6]). The scope of proposed improvements continues to expand, with more recent work exploring AI-driven approaches (for example, [7] and [8]). A/B testing products must determine which of these methodologies deliver meaningful value relative to their implementation costs.

The key challenge is that the true impacts of A/B tests are unobservable. Each test yields only a noisy estimate of the true impact, making it difficult to tell whether a proposed methodology actually leads to an overall improvement in estimates or decision making across tests. While simulation methods can be used to evaluate performance, they require strong assumptions about the true underlying data generating process.

Sample splitting offers a potential solution to this problem ([9]), but its theoretical foundation remains undeveloped. Under sample splitting, each A/B test is divided into two, non-overlapping splits: one to implement the proposed methodology, the other to construct an independent, unbiased proxy for the true impact. Comparing these two quantities across A/B tests allows us to measure the average performance of the proposed methodology, without making strong distributional assumptions ([9]). However, the statistical properties of these sample-split estimators have yet to be characterized formally: their bias, variance, and asymptotic distributions remain unknown, and there are no established methods for conducting statistical inference. Without such foundations, products lack guidance on exactly how to use sample splitting, limiting its adoption in practice.

In this paper, we introduce a theoretical framework for sample splitting in A/B testing environments, characterizing both what these methods can measure and how to conduct valid inference. We show that while sample splitting enables performance evaluation when true impacts are unobserved, it does so at a cost. Because methodologies are implemented on only a split of the data, sample-split estimators are generally biased for what we ultimately care about — how the methodologies perform on the full sample. We show that sample-split estimators consistently estimate sample-split analogues of full-sample performance, derive their asymptotic distributions, and show how to construct valid asymptotic confidence intervals. We demonstrate that while a careful choice of how the sample is split

*Email: rykessle@amazon.com, jmcq@amazon.com, miikka@amazon.com. We thank Dmitry Taubinsky for inspiring discussions and Thomas Richardson and James Robins for helpful comments and suggestions.

can reduce bias for full-sample performance, this improvement generally comes at the cost of increased variance. We validate our main theoretical results through simulations.

Our theoretical framework has important practical implications for A/B testing products. First, it shows that, in some cases, products can dramatically reduce the computational costs of sample splitting by simulating sample-split estimates rather than repeatedly partitioning unit-level data. Second, it provides a principled approach to statistical inference, enabling products to conduct hypothesis testing (for example, to determine whether one methodology significantly outperforms another) and statistical power calculations (for example, to determine how many A/B tests are needed to reliably detect a given performance difference between methodologies). Third, it demonstrates that common sample splitting practices — such as splitting data equally or using hundreds of random partitions — are often suboptimal. These insights, together with the step-by-step algorithms we provide, enable products to implement principled sample splitting efficiently at scale, making it a practical and powerful tool for evaluating proposed methodological improvements.

More generally, our work contributes to the literature on cross-validation and resampling methods ([10]). Like cross-validation, sample splitting separates the application of a methodology from its evaluation by partitioning data. The key difference is that in cross-validation the prediction target is observed directly, whereas in our setting the prediction target is itself an estimate subject to sampling variation. To our knowledge, [9] were the first to apply the idea of sample splitting in the A/B testing context, building upon earlier work in the estimation of heterogeneous treatment effects by [11] and [12]. [9] demonstrate the potential of sample splitting in evaluating the performance of A/B testing methodologies but leave its statistical properties largely unformalized.² We extend their work by introducing a theoretical framework for sample splitting in A/B testing environments, characterizing both what these methods can measure and how to conduct valid inference.

2 Motivating Theory

2.1 Context

We consider a set of $i \in \{1, \dots, I\}$ feature changes, each evaluated through an A/B test where the change is exposed to a random subset of units. Each feature change has a true impact Δ_i , which is a random variable with realized values denoted by δ_i . These true impacts Δ_i are independently and identically distributed (iid) according to some unknown probability distribution G . Conditional on the true impact $\Delta_i = \delta_i$, the A/B test generates random data \hat{X}_i independently distributed according to a probability distribution F , with realized values \hat{x}_i .

We define an *A/B testing methodology* $\kappa(\hat{x}_i)$ to be a mapping of the data \hat{x}_i to a real-valued output. The methodology could be an estimator of the true impact δ_i or a decision rule that determines whether to launch the given treatment ($\kappa(\hat{x}_i) = 1$) or not ($\kappa(\hat{x}_i) = 0$). Given the randomization of the feature change and the central limit theorem, we have an unbiased estimator $\kappa(\hat{X}_i) = \hat{\Delta}_i$ of δ_i that is normally distributed, with realized values $\hat{\delta}_i$ and sampling variance τ_i^2 :

$$\hat{\Delta}_i | \Delta_i = \delta_i, \tau_i^2 \stackrel{\text{ind}}{\sim} N(\delta_i, \tau_i^2) \quad (1)$$

While the sampling variances τ_i^2 are typically estimated from data, we follow standard practice in the A/B testing literature and treat them as known (see, for example, [6]). In most large-scale A/B testing environments, τ_i^2 is estimated with far greater precision than $\hat{\Delta}_i$, making this a pragmatic simplification that helps focus attention on the first-order uncertainty in $\hat{\Delta}_i$.³

We measure the performance of the methodology $\kappa(\hat{x}_i)$ given a true impact of $\Delta_i = \delta_i$ via a real-valued *performance function* $p(\kappa(\hat{x}_i), \delta_i)$. As illustrated in the examples below, the specification of the performance function depends on the type of methodology being evaluated.

²The authors acknowledge that their work is guided primarily by intuition and heuristics rather than principled theory. For example, they note the use “*t*-test heuristics” (page 7) and suggest that the confidence intervals they report “should be interpreted cautiously” (page 11).

³In an A/B test with normally distributed outcomes and a total of N units divided equally across treatment and control, the sampling variance of $\hat{\Delta}_i$ decreases at a rate of $1/N$, while the sampling variances of standard estimators of τ_i^2 decrease at a rate of $1/N^2$. So, for example, when $N = 1M$ the sampling variance of $\hat{\Delta}_i$ will be at least 1M times larger than the sampling variance of standard estimators of τ_i^2 . Similar results hold for heavy-tailed outcomes when winsorization is applied, as is common in large-scale A/B testing.

Example 1 (Measuring performance of an estimator): We may want to measure the performance of a shrinkage estimator $\kappa(\hat{x}_i) = w_i \cdot \hat{\delta}_i$ for $w_i \in (0, 1)$ under squared error in which case we could specify the performance function $p(\kappa(\hat{x}_i), \delta_i) = (\kappa(\hat{x}_i) - \delta_i)^2$.

Example 2 (Measuring performance of a decision rule): We may want to measure the performance of the decision rule $\kappa(\hat{x}_i) = 1(\hat{\delta}_i > \tau_i \cdot \Phi^{-1}(0.975))$ to launch treatments when the estimated impact is positive and statistically significant at the 5 percent level. In this case, we could specify the performance function $p(\kappa(\hat{x}_i), \delta_i) = \kappa(\hat{x}_i) \cdot \delta_i - (1 - \kappa(\hat{x}_i)) \cdot \delta_i$ to capture the idea that we get δ_i when we launch and $-\delta_i$ when we do not. We hereafter refer to this measure of performance and its counterpart $p(\kappa(\hat{x}_i), \delta_i) = \kappa(\hat{x}_i) \cdot \delta_i$ as *decision value* and *launch-only decision value*, respectively.

In what follows, for any random variable K_i , we write F_K to denote its distribution induced by the sampling distribution F and G_K to denote its distribution induced by the distribution of true impacts G . We write $E_F(\cdot)$ and $\text{Var}_F(\cdot)$ for the expectation and variance with respect to randomness induced by F , and $E_G(\cdot)$ and $\text{Var}_G(\cdot)$ for the expectation and variance with respect to randomness induced by G . Finally, we write $E(\cdot)$ and $\text{Var}(\cdot)$ for the expectation and variance with respect to both sources of randomness, integrating over both F and G . We denote by $\phi(\cdot)$ and $\Phi(\cdot)$ the probability density function and cumulative distribution function of the standard normal distribution, respectively.

2.2 Ideal Estimands

Our goal is to quantify how well different methodologies perform across A/B tests. To this end, we begin by defining a set of *ideal estimands* that capture how we would measure performance if the true impacts δ_i were observable. These estimands serve as our conceptual targets and help clarify the sense in which our estimators (introduced in section 2.3.4) may be biased for what we ultimately want to measure.

For each A/B test i , the expected performance of a methodology κ under performance measure p given a true impact of $\Delta_i = \delta_i$ is given by:

$$Y_i(\kappa) = E_F \left(p \left(\kappa(\hat{X}_i), \delta_i \right) \right) \quad (2)$$

Here $\kappa(\hat{X}_i)$ is the methodology's output (estimate or launch decision) given data \hat{X}_i . The function $p(\kappa(\hat{X}_i), \delta_i)$ measures how well this output performs given a true impact δ_i . Taking the expectation over the sampling variability induced by F averages out the randomness in the data \hat{X}_i , yielding the expected performance $Y_i(\kappa)$ for a given true impact.

While $Y_i(\kappa)$ captures the expected performance of κ for a given test, it varies across tests due to differences in the true impacts Δ_i :

$$Y_i(\kappa) \stackrel{\text{iid}}{\sim} G_Y, \text{ with } E_G(Y_i(\kappa)) = \theta(\kappa), \text{Var}_G(Y_i(\kappa)) = \lambda^2(\kappa) \quad (3)$$

One ideal estimand is therefore the average performance of the methodology across tests, $\theta(\kappa)$. The quantity $\theta(\kappa)$ accounts for both the sampling variation in each A/B test and the variation in true impacts across tests, making it a natural target when evaluating methodological performance at scale.

While average performance can be helpful in benchmarking a given methodology, we are often more interested in differences in average performance across methodologies. Another ideal estimand is therefore:

$$\theta(\kappa^1, \kappa^2) = \frac{\theta(\kappa^2) - \theta(\kappa^1)}{\theta(\kappa^1)} \quad (4)$$

where κ^1 and κ^2 are different methodologies, and we normalize by $\theta(\kappa^1)$ so that $\theta(\kappa^1, \kappa^2)$ can be interpreted as the percent change in performance of methodology κ^2 relative to κ^1 .

Example 3 (Illustrating performance estimands for estimators): Assume that $\tau_i = \tau$ for all $i = 1, \dots, I$ and that $G \sim N(0, \sigma^2)$. Consider evaluating the relative performance of two estimators under squared error $p(\kappa(\hat{x}_i), \delta_i) = (\kappa(\hat{x}_i) - \delta_i)^2$: the unbiased estimator $\kappa^1(\hat{x}_i) = \hat{\delta}_i$ and the Bayes estimator $\kappa^2(\hat{x}_i) = \sigma^2(\sigma^2 + \tau^2)^{-1} \hat{\delta}_i$. In this case, we have that:

$$\theta(\kappa^1, \kappa^2) = \frac{\frac{\sigma^2 \tau^2}{\sigma^2 + \tau^2} - \tau^2}{\tau^2} = \frac{\sigma^2}{\sigma^2 + \tau^2} - 1 < 0 \quad (5)$$

consistent with the fact that the Bayes estimator minimizes mean squared error ([13]).

Example 4 (Illustrating performance estimands for decision rules): Assume that $\tau_i = \tau$ for all $i = 1, \dots, I$ and that $G \sim N(0, \sigma^2)$. Consider evaluating the relative performance of two decision rules under launch-only decision value $p(\kappa(\hat{x}_i), \delta_i) = \kappa(\hat{x}_i) \cdot \delta_i$: a frequentist decision rule $\kappa^1(\hat{x}_i) = 1(\hat{\delta}_i > \tau \cdot \Phi^{-1}(0.975))$ which launches treatments when the estimated impact is positive and statistically significant at the 5 percent level and a Bayes decision rule $\kappa^2(\hat{x}_i) = 1(\sigma^2(\sigma^2 + \tau^2)^{-1}\hat{\delta}_i > 0)$ which launches treatments when the posterior mean is positive. In this case, we have that:

$$\theta(\kappa^1, \kappa^2) = \frac{\frac{\sigma^2}{\sqrt{\sigma^2 + \tau^2}} \left(\phi(0) - \phi\left(\frac{\tau \cdot \Phi^{-1}(0.975)}{\sqrt{\sigma^2 + \tau^2}}\right) \right)}{\frac{\sigma^2}{\sqrt{\sigma^2 + \tau^2}} \phi\left(\frac{\tau \cdot \Phi^{-1}(0.975)}{\sqrt{\sigma^2 + \tau^2}}\right)} = \frac{\phi(0)}{\phi\left(\frac{\tau \cdot \Phi^{-1}(0.975)}{\sqrt{\sigma^2 + \tau^2}}\right)} - 1 > 0 \quad (6)$$

consistent with the fact that the Bayes decision rule maximizes mean launch-only decision value ([6]).

Note that these examples assume that the distribution of true impacts G is normal only to help make the estimands concrete. Our sample splitting framework, introduced below, is designed to evaluate performance of methodologies without requiring any assumptions about G .

2.3 Sample Splitting

We evaluate the performance of methodologies via sample splitting, where we partition each A/B test into two splits: one to implement the proposed methodology, the other to construct an independent, unbiased proxy for the true impact. This approach enables performance measurement when true impacts are unobserved ([9]). However, because the proposed methodology is implemented on only a split of the sample, we cannot directly estimate the ideal estimands in section 2.2. In what follows, we outline a general approach to generating sample-split data, which applies to any methodology. We then show that this approach can be simplified for “plug-in methodologies” that depend on the data only through a set of sufficient statistics. With this data foundation, we then define sample-split analogues of the ideal estimands, introduce corresponding unbiased estimators, and outline an approach to statistical inference.

2.3.1 Data Generation: General Approach

We implement sample splitting as follows. For each A/B test, we randomly partition the units assigned to each experience (control and treatment) into two splits, labeled a and b , with a fraction $\alpha \in (0, 1)$ allocated to split a and the remaining $(1 - \alpha)$ to split b . For a given random partition s , let \hat{X}_{ias} and \hat{X}_{ibs} be the data associated with units in splits a and b , with $(\hat{x}_{ias} \cup \hat{x}_{ibs}) = \hat{x}_i$ and $(\hat{x}_{ias} \cap \hat{x}_{ibs}) = \emptyset$. We use split a for *training*, applying both the methodology of interest and the unbiased estimator $\hat{\Delta}_i$ of δ_i . This yields an estimate or launch decision $\kappa(\hat{x}_{ias})$ and an estimate $\hat{\delta}_{ias}$. We use split b for *evaluation*, applying the unbiased estimator $\hat{\Delta}_i$ of δ_i to obtain a proxy for the true impact $\hat{\delta}_{ibs}$. Because we treat τ_i^2 as known, we do not estimate sampling variances of $\hat{\Delta}_{ias}$ and $\hat{\Delta}_{ibs}$ based on data but instead compute them deterministically based on τ_i^2 and the split fraction α . Specifically, $\tau_{ias}^2 = \tau_i^2 / \alpha$ and $\tau_{ibs}^2 = \tau_i^2 / (1 - \alpha)$, reflecting the standard inverse relationship between sampling variance and sample size.

Because the methodology $\kappa(\hat{X}_{ias})$ and the proxy for the true impact $\hat{\Delta}_{ibs}$ are based on disjoint samples of units, they are statistically independent conditional on the true impact $\Delta_i = \delta_i$:

$$\kappa(\hat{X}_{ias}) \perp \hat{\Delta}_{ibs} \mid \Delta_i = \delta_i \quad (7)$$

Additionally, because the partitions are determined at random, the proxy for the true impact $\hat{\Delta}_{ibs}$ is (under equation (1)) unbiased and normally distributed conditional on the true impact $\Delta_i = \delta_i$:

$$\hat{\Delta}_{ibs} \mid \Delta_i = \delta_i, \tau_i^2 \stackrel{\text{ind}}{\sim} N(\delta_i, \tau_{ibs}^2) \quad (8)$$

Together, equations (7) and (8) provide the statistical foundation for our sample splitting methodology. The conditional independence property in equation (7) ensures that noise affecting the methodology’s output does not also influence its validation, guarding against spurious estimates of performance. The normality and unbiasedness properties in equation (8) ensure that split b provides a valid proxy for the true impact δ_i .

A single partition suffices to estimate performance, but results can vary substantially across different partitions. To reduce this variability, we repeat our sample splitting procedure S times, generating new random partitions that each divide units into splits a and b . This yields estimates $\{\kappa(\hat{x}_{ias}), \hat{\delta}_{ias}, \tau_{ias}^2, \hat{\delta}_{ibs}, \tau_{ibs}^2\}_{s=1}^S$ which we then average for each test in hopes of obtaining more stable results. Importantly, while splits a and b remain independent within each partition s (maintaining the property in equation (7)), they are not independent across partitions, as they are based on overlapping data. Our approach to estimation and inference, outlined in section 2.3.4, takes this dependence into account.

2.3.2 Data Generation: Plug-in Methodologies

While the previous section illustrated how sample-split data can be generated by repeatedly partitioning the unit-level data of a test, many methodologies depend on the unit-level data only through the sufficient statistics $(\hat{\delta}_{ias}, \tau_{ias}^2)$. We refer to these as “plug-in methodologies”, as they simply plug the estimates $(\hat{\delta}_{ias}, \tau_{ias}^2)$ into deterministic functions, without changing the way in which the estimates themselves are constructed. Plug-in methodologies include shrinkage estimators $\kappa(\hat{x}_{ias}) = w_i \cdot \hat{\delta}_{ias}$ for $w_i \in (0, 1)$ and threshold decision rules $\kappa(\hat{x}_{ias}) = 1(\hat{\delta}_{ias} > \tau_{ias} \cdot c)$ for launch threshold $c \geq 0$.

For these methodologies, we can simplify sample-split data generation. Instead of repeatedly partitioning unit-level data and re-computing estimates, we can directly sample $(\hat{\Delta}_{ias}, \hat{\Delta}_{ibs})$ from their joint distribution given the full-sample estimates $(\hat{\delta}_i, \tau_i^2)$ and true impact δ_i . This shortcut, formalized in the proposition below, eliminates the need for unit-level data in partitioning while preserving the statistical guarantees that sample splitting provides in performance evaluation.

Proposition 1 (Evaluating plug-in methodologies without unit-level partitioning): Let splits a and b be formed by randomly partitioning units in an A/B test with proportions α and $(1 - \alpha)$, respectively. Under equation (1), the joint distribution of $(\hat{\Delta}_{ias}, \hat{\Delta}_{ibs})$ conditional on the full-sample estimates $(\hat{\delta}_i, \tau_i^2)$ and true impact δ_i is given by:

$$\begin{pmatrix} \hat{\Delta}_{ias} \\ \hat{\Delta}_{ibs} \end{pmatrix} \Big| \hat{\Delta}_i = \hat{\delta}_i, \Delta_i = \delta_i, \tau_i^2 \stackrel{\text{ind}}{\sim} N \left(\begin{pmatrix} \hat{\delta}_i \\ \hat{\delta}_i \end{pmatrix}, \begin{pmatrix} \frac{\tau_i^2}{\alpha} - \tau_i^2 & -\tau_i^2 \\ -\tau_i^2 & \frac{\tau_i^2}{1-\alpha} - \tau_i^2 \end{pmatrix} \right) \quad (9)$$

For a given test, the process of repeatedly partitioning units and re-estimating $(\hat{\delta}_{ias}, \hat{\delta}_{ibs})$ can therefore be approximated by resampling from the conditional distribution in equation (9) given the full-sample estimates $(\hat{\delta}_i, \tau_i^2)$. Given draws from this distribution, plug-in methodologies (being deterministic functions of $(\hat{\delta}_{ias}, \tau_{ias}^2)$) can then be applied, yielding estimates $\{\kappa(\hat{x}_{ias}), \hat{\delta}_{ias}, \tau_{ias}^2, \hat{\delta}_{ibs}, \tau_{ibs}^2\}_{s=1}^S$.⁴

Proof: See appendix A □

Proposition 1 is similar in spirit to [14], who generate “doppelganger” observations without access to unit-level data by deconvolving the overall estimates for a test.

By eliminating the need for unit-level data in partitioning, proposition 1 lowers the computational cost of evaluating plug-in methodologies. Without the shortcut, evaluating plug-in methodologies requires processing N unit observations for each of the I tests S times. With it, we simply draw from the conditional distribution S times for each of the I tests. In contexts with millions of units and thousands of tests, the savings can be substantial, enabling us to increase the scale of our analyses by orders of magnitude and obtain significantly more precise estimates of performance.

However, not all methodologies can be evaluated via this shortcut. The shortcut relies on the methodology treating the estimates $(\hat{\delta}_{ias}, \tau_{ias}^2)$ as sufficient statistics for the underlying data \hat{x}_{ias} . Some methodologies — such as covariate adjustment, stratification, and outlier handling — affect how these estimates are constructed and therefore still require repeated unit-level partitioning.

⁴Equation (9) implies that, conditional on full-sample estimates $(\hat{\delta}_i, \tau_i^2)$, $\hat{\Delta}_{ias}$ and $\hat{\Delta}_{ibs}$ are perfectly negatively correlated across partitions. This follows from the fact that $\hat{\Delta}_i = \alpha \hat{\Delta}_{ias} + (1 - \alpha) \hat{\Delta}_{ibs}$, so increases in one require decreases in the other.

2.3.3 Estimands

To evaluate performance of methodologies under sample splitting, we introduce generalized, sample-splitting analogues of the ideal estimands introduced in 2.2. These estimands reflect that performance depends not only on the methodology κ and performance function p , but also on the share of the sample α used in implementing the methodology. Given α , the expected performance of a methodology κ under performance measure p given a true impact of $\Delta_i = \delta_i$ is given by:

$$Y_i(\kappa, \alpha) = E_F \left(p \left(\kappa \left(\hat{X}_{ias} \right), \delta_i \right) \right) \quad (10)$$

This generalization nests the ideal case: as $\alpha \rightarrow 1$ the methodology is implemented on the full sample and we recover the ideal performance measure in equation (2).

Just like its ideal counterpart $Y_i(\kappa)$, $Y_i(\kappa, \alpha)$ varies across tests due to differences in true impacts Δ_i :

$$Y_i(\kappa, \alpha) \stackrel{\text{iid}}{\sim} G_Y, \text{ with } E_G(Y_i(\kappa, \alpha)) = \theta(\kappa, \alpha), \text{Var}_G(Y_i(\kappa, \alpha)) = \lambda^2(\kappa, \alpha) \quad (11)$$

We denote realizations of $Y_i(\kappa, \alpha)$ by $y_i(\kappa, \alpha)$.

As in section 2.2, our sample-splitting estimands are then the average performance of the methodology across tests, $\theta(\kappa, \alpha)$, and percent differences in average performance across methodologies:

$$\theta(\kappa^1, \kappa^2, \alpha) = \frac{\theta(\kappa^2, \alpha) - \theta(\kappa^1, \alpha)}{\theta(\kappa^1, \alpha)} \quad (12)$$

Example 5 (Illustrating performance estimands for estimators, continued): Returning to example 3, we now consider the sample-split analogue of the difference in mean squared error across the estimators. For any $\alpha < 1$, sample splitting overestimates the performance benefits of the Bayes estimator. Replacing τ^2 with τ^2/α , we have that:

$$\theta(\kappa^1, \kappa^2, \alpha) = \frac{\sigma^2}{\sigma^2 + \frac{\tau^2}{\alpha}} - 1 < \frac{\sigma^2}{\sigma^2 + \tau^2} - 1 = \theta(\kappa^1, \kappa^2) \quad (13)$$

For example, when $\sigma^2 = \tau^2 = 1$, the Bayes estimator reduces mean squared error by 50 percent. Sample splitting with $\alpha = 0.50$ would, however, suggest a reduction of 66 percent.

Example 6 (Illustrating performance estimands for decision rules, continued): Returning to example 4, we now consider the sample-split analogue of the difference in launch-only decision value across the decision rules. For any $\alpha < 1$, sample splitting overestimates the performance benefits of the Bayes decision rule. Replacing τ^2 with τ^2/α , we have that:

$$\theta(\kappa^1, \kappa^2, \alpha) = \frac{\phi(0)}{\phi\left(\frac{\sqrt{\frac{\tau^2}{\alpha}} \cdot \Phi^{-1}(0.975)}{\sqrt{\sigma^2 + \frac{\tau^2}{\alpha}}}\right)} - 1 > \frac{\phi(0)}{\phi\left(\frac{\tau \cdot \Phi^{-1}(0.975)}{\sqrt{\sigma^2 + \tau^2}}\right)} - 1 = \theta(\kappa^1, \kappa^2) \quad (14)$$

For example, when $\sigma^2 = \tau^2 = 1$, the Bayes decision rule increases the launch-only decision value by 160 percent. Sample splitting with $\alpha = 0.50$ would, however, suggest a benefit of 260 percent.

As these examples illustrate, the sample-split estimands can differ substantially from their ideal counterparts, with the magnitude of this difference governed by the split fraction α . In subsequent sections, we discuss how to estimate these sample-split estimands and the tradeoffs underlying different choices of α .

2.3.4 Estimators and Inference

Sample splitting enables us to estimate the performance of methodologies by replacing the unobserved true impacts δ_i with the independent, unbiased proxies $\hat{\Delta}_{ibs}$ from split b . This leads to natural estimators of the sample-split estimands $Y_i(\kappa, \alpha)$ introduced in the previous section.

For each A/B test i and partition s , we consider an estimator $\hat{Y}_{is}(\kappa, \alpha)$ whose average across partitions $\hat{Y}_i(\kappa, \alpha, S) = \frac{1}{S} \sum_{s=1}^S \hat{Y}_{is}(\kappa, \alpha)$ satisfies:

$$\begin{aligned}\hat{Y}_i(\kappa, \alpha, S) | Y_i(\kappa, \alpha) = y_i(\kappa, \alpha) &\stackrel{\text{ind}}{\sim} F_{\hat{Y}}, \text{ with } E_F(\hat{Y}_i(\kappa, \alpha, S)) = y_i(\kappa, \alpha), \\ \text{Var}_F(\hat{Y}_i(\kappa, \alpha, S)) &= \Omega_i^2(\kappa, \alpha, S)\end{aligned}\quad (15)$$

The variance $\Omega_i^2(\kappa, \alpha, S)$ reflects both sampling variability in the full-sample data \hat{X}_i given the true impact $\Delta_i = \delta_i$ and variability in the sample-split data $(\hat{X}_{ias}, \hat{X}_{ibs})$ given the full-sample data.

Given $\hat{Y}_i(\kappa, \alpha, S)$ satisfying equation (15), we estimate the average performance of methodologies across tests and differences in average performance across methodologies. We estimate the average performance across tests $\theta(\kappa, \alpha)$ via:

$$\hat{\theta}(\kappa, \alpha, S) = \frac{1}{I} \sum_{i=1}^I \hat{Y}_i(\kappa, \alpha, S) \quad (16)$$

We estimate the difference in average performance across methodologies $\theta(\kappa^1, \kappa^2, \alpha)$ via:

$$\hat{\theta}(\kappa^1, \kappa^2, \alpha, S) = \frac{\hat{\theta}(\kappa^2, \alpha, S) - \hat{\theta}(\kappa^1, \alpha, S)}{\hat{\theta}(\kappa^1, \alpha, S)} \quad (17)$$

Because each $\hat{Y}_i(\kappa, \alpha, S)$ is unbiased for $y_i(\kappa, \alpha)$, the estimator $\hat{\theta}(\kappa, \alpha, S)$ is unbiased for $\theta(\kappa, \alpha)$. We formalize this via the following proposition.

Proposition 2 (Unbiasedness): Under equations (11) and (15), the estimator of average performance $\hat{\theta}(\kappa, \alpha, S)$ is unbiased for $\theta(\kappa, \alpha)$.

Proof: See appendix A □

The challenge is therefore to construct estimators $\hat{Y}_{is}(\kappa, \alpha)$ that satisfy equation (15) for different performance functions p . Table 1 presents such estimators for common choices of p , including bias, mean squared error, and decision value. We provide proofs of the unbiasedness condition in (15) for each estimator in appendix A, relying on the sample splitting conditions in equations (7) and (8).

Table 1: Sample-split estimands and unbiased estimators by performance measure

Methodology	Measure of performance p	Sample-split estimand $Y_i(\kappa, \alpha)$	Unbiased estimator $\hat{Y}_{is}(\kappa, \alpha)$
Estimator	Bias	$E_F(\kappa(\hat{X}_{ias}) - \delta_i)$	$\kappa(\hat{X}_{ias}) - \hat{\delta}_{ibs}$
Estimator	Squared error	$E_F((\kappa(\hat{X}_{ias}) - \delta_i)^2)$	$(\kappa(\hat{X}_{ias}) - \hat{\delta}_{ibs})^2 - \tau_{ibs}^2$
Decision rule	Decision value	$E_F(\kappa(\hat{X}_{ias}) \cdot \delta_i - (1 - \kappa(\hat{X}_{ias})) \cdot \delta_i)$	$\kappa(\hat{X}_{ias}) \cdot \hat{\delta}_{ibs} - (1 - \kappa(\hat{X}_{ias})) \cdot \hat{\delta}_{ibs}$
Decision rule	Launch-only decision value	$E_F(\kappa(\hat{X}_{ias}) \cdot \delta_i)$	$\kappa(\hat{X}_{ias}) \cdot \hat{\delta}_{ibs}$

Notes: This table presents sample-split estimands $Y_i(\kappa, \alpha)$ and corresponding unbiased estimators $\hat{Y}_{is}(\kappa, \alpha)$ for different measures of performance.

Although $\hat{\theta}(\kappa, \alpha, S)$ is unbiased for $\theta(\kappa, \alpha)$, the estimator of the difference in average performance across methodologies $\hat{\theta}(\kappa^1, \kappa^2, \alpha, S)$ is generally biased for $\theta(\kappa^1, \kappa^2, \alpha)$. The bias arises from the nonlinearity of the percent change: even when $\hat{\theta}(\kappa^1, \alpha, S)$ and $\hat{\theta}(\kappa^2, \alpha, S)$ are unbiased for $\theta(\kappa^1, \alpha)$ and $\theta(\kappa^2, \alpha)$, respectively, their ratio $\hat{\theta}(\kappa^1, \kappa^2, \alpha, S)$ need not be unbiased for $\theta(\kappa^1, \kappa^2, \alpha)$.

Having established the bias properties of the estimators $\hat{\theta}(\kappa, \alpha, S)$ and $\hat{\theta}(\kappa^1, \kappa^2, \alpha, S)$, we now turn to their large-sample behavior. In practice, we care not only about whether the estimators are close to their true values on average, but also about their sampling variability and the rate at which they converge to their true values as the number of tests I grows large.

We summarize our results via the two following propositions. For brevity, we focus the propositions on $\hat{\theta}(\kappa, \alpha, S)$; analogous results for $\hat{\theta}(\kappa^1, \kappa^2, \alpha, S)$ follow immediately from standard asymptotic theory.

Proposition 3 (Asymptotic normality): Under equations (11) and (15) and the regularity condition that $\sup_i E(\hat{Y}_i(\kappa, \alpha)^2) < \infty$, the estimator $\hat{\theta}(\kappa, \alpha)$ is asymptotically normally distributed:

$$\sqrt{I} \left(\hat{\theta}(\kappa, \alpha, S) - \theta(\kappa, \alpha) \right) \xrightarrow{d} N(0, \zeta^2(\kappa, \alpha, S)) \quad (18)$$

where $\zeta^2(\kappa, \alpha, S) = \lambda^2(\kappa, \alpha) + E_G(\Omega_i^2(\kappa, \alpha, S))$.

Proof: See appendix A □

Proposition 3 shows that the estimator $\hat{\theta}(\kappa, \alpha, S)$ is asymptotically normally distributed. Its asymptotic variance $\zeta^2(\kappa, \alpha, S)$ captures heterogeneity in true performance across tests via $\lambda^2(\kappa, \alpha)$ and the average sampling variance of estimated performance across tests via $E_G(\Omega_i^2(\kappa, \alpha, S))$.

Proposition 4 shows that $\zeta^2(\kappa, \alpha, S)$ can be consistently estimated, enabling valid asymptotic inference.

Proposition 4 (Valid asymptotic inference): Under equations (11) and (15) and the regularity condition that $\sup_i E(\hat{Y}_i(\kappa, \alpha, S)^2) < \infty$, the estimator:

$$\hat{\zeta}^2(\kappa, \alpha, S) = \frac{1}{I} \sum_{i=1}^I (\hat{Y}_i(\kappa, \alpha, S) - \hat{\theta}(\kappa, \alpha, S))^2 \quad (19)$$

is consistent for the asymptotic variance $\zeta^2(\kappa, \alpha, S) = \lambda^2(\kappa, \alpha) + E_G(\Omega_i^2(\kappa, \alpha, S))$. Therefore, the $(1 - v)$ confidence intervals:

$$\hat{\theta}(\kappa, \alpha, S) \pm \frac{\hat{\zeta}(\kappa, \alpha, S)}{\sqrt{I}} \cdot \Phi^{-1}\left(1 - \frac{v}{2}\right) \quad (20)$$

will cover $\theta(\kappa, \alpha)$ with probability $(1 - v)$ as $I \rightarrow \infty$.

Proof: See appendix A □

Together, these results enable principled estimation of the performance of methodologies via sample splitting. The unbiasedness of the estimators allows us to reliably estimate average performance under sample splitting, while the asymptotic normality enables us to quantify uncertainty in our estimates. The asymptotic confidence intervals can be used both for hypothesis testing (for example, to determine whether one methodology significantly outperforms another) and statistical power calculations (for example, to determine how many A/B tests are needed to reliably detect a given performance difference between methodologies). Algorithm 1 provides a step-by-step guide towards implementation.

Algorithm 1 Estimating average performance of a methodology κ

1. Identify I historical A/B tests and choose split fraction α and number of partitions S
2. Generate sample-split data:
 - (a) If evaluating a plug-in methodology, simulate data, as described in section 2.3.2
 - (b) Otherwise, generate data via unit-level partitioning, as described in section 2.3.1
3. Choose measure of performance $\hat{Y}_{is}(\kappa, \alpha)$ from table 1 and compute average across partitions $\hat{Y}_i(\kappa, \alpha, S) = \frac{1}{S} \sum_{s=1}^S \hat{Y}_{is}(\kappa, \alpha)$ for each test
4. Construct estimate of average performance $\hat{\theta}(\kappa, \alpha, S) = \frac{1}{I} \sum_{i=1}^I \hat{Y}_i(\kappa, \alpha, S)$ and its sampling variance $\hat{\zeta}^2(\kappa, \alpha, S) = \frac{1}{I} \sum_{i=1}^I (\hat{Y}_i(\kappa, \alpha, S) - \hat{\theta}(\kappa, \alpha, S))^2$
5. Construct $(1 - v)$ confidence intervals for average performance:

$$\hat{\theta}(\kappa, \alpha, S) \pm \frac{\hat{\zeta}(\kappa, \alpha, S)}{\sqrt{I}} \cdot \Phi^{-1}\left(1 - \frac{v}{2}\right)$$

Note, however, that care must be taken in applying these asymptotic results in smaller samples of A/B tests. The normal approximation in proposition 3 requires a sufficient number of A/B tests to be reliable. Similarly, the variance estimator in proposition 4 may be unstable in small samples, potentially leading to poor coverage of the resulting confidence intervals. Our simulation results in the next section suggest that these approximations work well with thousands of tests, but practitioners should exercise caution when working with significantly smaller samples.

3 Simulations

In this section, we simulate a stylized A/B testing environment in which the difference in average performance between two methodologies is known. These simulations serve three purposes. First, they validate theoretical predictions outlined in previous sections. Second, they assess the adequacy of the asymptotic approximations underlying our approach to estimation and inference in finite samples. Third, they allow us to assess how performance of our estimators varies as a function of key sample-split design parameters.

3.1 Data Generating Process

We simulate from the following data generating process (DGP):

$$\Delta_i \stackrel{\text{iid}}{\sim} N(0, \sigma^2) \quad (21)$$

$$\epsilon_i \stackrel{\text{iid}}{\sim} \chi_1^2 \quad (22)$$

$$\tau_i^2 = \tau^2 + \epsilon_i \quad (23)$$

$$\hat{\Delta}_i | \Delta_i = \delta_i, \tau_i^2 \stackrel{\text{ind}}{\sim} N(\delta_i, \tau_i^2) \quad (24)$$

$$\begin{pmatrix} \hat{\Delta}_{ias} \\ \hat{\Delta}_{ibs} \end{pmatrix} \Big| \hat{\Delta}_i = \hat{\delta}_i, \Delta_i = \delta_i, \tau_i^2 \stackrel{\text{ind}}{\sim} N \left(\begin{pmatrix} \hat{\delta}_i \\ \hat{\delta}_i \end{pmatrix}, \begin{pmatrix} \frac{\tau_i^2}{\alpha} - \tau_i^2 & -\tau_i^2 \\ -\tau_i^2 & \frac{\tau_i^2}{1-\alpha} - \tau_i^2 \end{pmatrix} \right) \quad (25)$$

We vary five key parameters: the variance of true impacts σ^2 , baseline sampling variance τ^2 , split fraction α , number of A/B tests I , and number of random partitions S .

This DGP captures three key features of standard A/B testing environments. First, we capture heteroskedasticity across tests by modeling the sampling variance of the estimated impacts via $\tau_i^2 = \tau^2 + \epsilon_i$, where τ is a constant and ϵ_i is a chi-squared random variable with 1 degree of freedom. Second, we capture different signal-to-noise ratios by considering different pairwise combinations of the variance of true impacts σ^2 and the baseline sampling variance τ^2 . Third, we capture the real-world correlations across partitions s for a given test by drawing $(\hat{\Delta}_{ias}, \hat{\Delta}_{ibs})$ from their joint distribution given the full-sample estimates $(\hat{\delta}_i, \tau_i^2)$.

As a benchmark specification, we consider $I = 5K$ A/B tests, $S = 30$ partitions, a split fraction $\alpha = 0.5$, a variance of true impacts $\sigma^2 = 1$, and a baseline sampling variance $\tau^2 = 2$.

3.2 Evaluation

We use the simulations to evaluate the relative performance of different point estimators and decision rules. For point estimators, we compare the performance of the unbiased estimator $\kappa^1(\hat{x}_i) = \hat{\delta}_i$ to that of the Bayes estimator $\kappa^2(\hat{x}_i) = \sigma^2(\sigma^2 + \tau_i^2)^{-1} \hat{\delta}_i$ under squared error, building upon example 3. For decision rules, we compare the performance of the frequentist decision rule $\kappa^1(\hat{x}_i) = 1(\hat{\delta}_i > \tau_i \cdot \Phi^{-1}(0.975))$ to that of a Bayes decision rule $\kappa^2(\hat{x}_i) = 1(\sigma^2(\sigma^2 + \tau_i^2)^{-1} \hat{\delta}_i > 0)$ under launch-only decision value, building upon example 4.


For each methodology type, we measure the bias of the estimator $\hat{\theta}(\kappa^1, \kappa^2, \alpha, S)$, both relative to the ideal estimand $\theta(\kappa^1, \kappa^2)$ and the sample-split estimand $\theta(\kappa^1, \kappa^2, \alpha)$. We also measure the variance of $\hat{\theta}(\kappa^1, \kappa^2, \alpha, S)$ and the coverage of its 95 percent confidence interval. Finally, we assess how the bias and variance vary as a function of the number of tests I , number of partitions S , and split fraction α .

3.3 Results

Figure 1 confirms the theoretical predictions outlined in previous sections. Sample splitting estimators are biased for the ideal estimands but are consistent for the sample-split estimands, resulting in negligible bias in the sample of $I = 5K$ A/B tests. The confidence intervals achieve nominal coverage, covering the sample-split estimand approximately 95 percent of the time.


Figure 2 highlights a bias-variance tradeoff underlying the choice of the split fraction α . As more data is allocated to the training split (that is, as $\alpha \rightarrow 1$), bias for the ideal estimand falls to zero but the variance of the estimated difference in performance rises. Under the baseline DGP, this produces a

Figure 1: Bias and coverage of 95 percent confidence intervals at different (τ^2, σ^2)



U -shaped curve in terms of mean-squared error, with the minimizing α greater than 0.5. We present results for alternative specifications of the DGP in appendix B.


Figure 2: Bias-variance tradeoff underlying choice of split fraction α



Together, the results in figures 1 and 2 underscore both the value and limitations of sample splitting. Sample splitting makes estimation possible by using a split of the data to construct an independent, unbiased proxy for the true impact, thereby overcoming the fundamental challenge that true impacts are unobserved. However, because we implement the methodology on only a split of the data, we cannot construct unbiased estimators of the quantity we ultimately care about: the performance of the methodology if it had been implemented on the full sample. We can reduce bias for this ideal estimand by increasing α but this comes at the cost of increasing the variance of our estimates of performance.

Figure 3 illustrates how the variance of sample-split estimators depends on both the number of A/B tests I and partitions S . While increasing either dimension reduces variance, the marginal benefits of increasing I and S differ substantially. Increasing the number of tests I consistently reduces variance. By contrast, increasing the number of partitions S yields diminishing returns, with minimal improvements beyond $S = 30$ under the baseline DGP. This pattern reflects fundamental differences between additional tests and additional partitions: while increasing the number of tests results in new, independent information about performance, increasing the number of partitions eventually just provides different realizations of the same underlying information. We present results for alternative specifications of the DGP in appendix B.

Figure 3: Variance as a function of the number of tests I and number of partitions S



4 Takeaways for Practitioners

We conclude by distilling our theoretical and simulation findings into four concrete recommendations for sample-splitting practitioners:

1. *Understand the fundamental tradeoff:* Sample splitting enables performance evaluation when true impacts are unobserved, but this comes at a cost. Because methodologies are implemented on only a split of the data, sample-split estimates are generally biased for what we ultimately care about — how the methodologies perform on the full sample. While this bias can be reduced by increasing the split fraction α , it cannot be eliminated entirely without sacrificing our ability to conduct valid inference.
2. *Split wisely, not necessarily evenly:* The standard split fraction of $\alpha = 0.5$ may be suboptimal. Choosing a split fraction $\alpha > 0.5$ can reduce bias in sample-split performance estimates without an offsetting increase in their variance.
3. *Skip unit-level partitioning for plug-in methods:* If you are evaluating a plug-in methodology that applies functions to the estimates $(\hat{\delta}_{ias}, \tau_{ias}^2)$ without changing the way in which they are constructed, you may not need to repeatedly partition the unit-level data underlying each A/B test. Instead, you may be able to draw sample-split estimates from their joint distribution given the full-sample estimates, as suggested in proposition 1. This can significantly lower the computational costs of sample splitting, enabling you to increase the scale of your analyses by orders of magnitude and obtain significantly more precise estimates of performance.
4. *Be careful when the number of tests I is small:* Our theoretical results are valid in the limit as the number of tests I grows large. When I is small, asymptotic approximations may perform poorly, resulting in unreliable estimates of performance and confidence intervals with poor coverage. Even when I is sufficient for reliable asymptotic approximations, you may still lack power to detect differences in performance between methodologies. In this case, alternative approaches to performance evaluation (such as simulation) may be more appropriate.
5. *Don't waste time and compute — $S = 30$ is often sufficient:* Using multiple partitions reduces variance, but most of the gains accrue across small S , leaving relatively little benefit to $S > 30$. Unlike increasing the number of A/B tests (I), which brings new independent information to estimation, increasing the number of partitions (S) eventually just provides different realizations of the same underlying information.

References

- [1] Alex Deng, Ya Xu, Ron Kohavi, and Toby Walker. Improving the sensitivity of online controlled experiments by utilizing pre-experiment data. In *Proceedings of the Sixth ACM International Conference on Web Search and Data Mining*, WSDM '13, page 123–132, New York, NY, USA, 2013. Association for Computing Machinery. ISBN 9781450318693. doi: 10.1145/2433396.2433413. URL <https://doi.org/10.1145/2433396.2433413>.
- [2] Alex Deng, Lo-Hua Yuan, Naoya Kanai, and Alexandre Salama-Manteau. Zero to hero: Exploiting null effects to achieve variance reduction in experiments with one-sided triggering. In *Proceedings of the Sixteenth ACM International Conference on Web Search and Data Mining*, WSDM '23,

- page 823–831, New York, NY, USA, 2023. Association for Computing Machinery. ISBN 9781450394079. doi: 10.1145/3539597.3570413. URL <https://doi.org/10.1145/3539597.3570413>.
- [3] Alex Deng, Luke Hagar, Nathaniel T. Stevens, Tatiana Xifara, and Amit Gandhi. Metric decomposition in A/B tests. In *Proceedings of the 30th ACM SIGKDD Conference on Knowledge Discovery and Data Mining*, KDD '24, page 4885–4895, New York, NY, USA, 2024. Association for Computing Machinery. ISBN 9798400704901. doi: 10.1145/3637528.3671556. URL <https://doi.org/10.1145/3637528.3671556>.
 - [4] David Goldberg and James E. Johndrow. A decision theoretic approach to A/B testing, 2017. URL <https://arxiv.org/abs/1710.03410>.
 - [5] Eduardo M. Azevedo, Alex Deng, José L. Montiel Olea, and E. Glen Weyl. Empirical bayes estimation of treatment effects with many A/B tests: An overview. *AEA Papers and Proceedings*, 109:43–47, May 2019. doi: 10.1257/pandp.20191003. URL <https://www.aeaweb.org/articles?id=10.1257/pandp.20191003>.
 - [6] Eduardo M. Azevedo, Alex Deng, José Luis Montiel Olea, Justin Rao, and E. Glen Weyl. A/B testing with fat tails. *Journal of Political Economy*, 128(12):4614–000, 2020. doi: 10.1086/710607. URL <https://doi.org/10.1086/710607>.
 - [7] R. Maria Del Rio-Chanona, Akhil Raman, Midhun Srikant, Amanda Askill, Yuntao Bai, Yoshua Bengio, Miles Brundage, Jack Clark, Sean Gerrish, Gillian Hadfield, Atoosa Kasirzadeh, Victoria Krakovna, David Krueger, Jan Leike, James Miller, Saurabh Mishra, Gauri Sastry, Richard Sutton, Helen Toner, and Amanda Askill. Simulating human behavior with AI agents, 2023. URL <https://hai.stanford.edu/assets/files/hai-policy-brief-simulating-human-behavior-with-ai-agents.pdf>.
 - [8] R. Maria del Rio-Chanona, Marco Pangallo, and Cars Hommes. Can generative AI agents behave like humans? Evidence from laboratory market experiments, 2025. URL <https://arxiv.org/abs/2505.07457>.
 - [9] Nilesh Tripuraneni, Dhruv Madeka, Dean Foster, Dominique Perrault-Joncas, and Michael I. Jordan. Meta-analysis of randomized experiments with applications to heavy-tailed response data, 2023. URL <https://arxiv.org/abs/2112.07602>.
 - [10] M Stone. Cross-validators choice and assessment of statistical predictions. *Journal of the Royal Statistical Society: Series B (Methodological)*, 36(2):111 – 147, 1971.
 - [11] Susan Athey and Guido Imbens. Recursive partitioning for heterogeneous causal effects. *Proceedings of the National Academy of Sciences*, 113(27):7353–7360, 2016. doi: 10.1073/pnas.1510489113. URL <https://www.pnas.org/doi/abs/10.1073/pnas.1510489113>.
 - [12] Victor Chernozhukov, Mert Demirer, Esther Duflo, and Iván Fernández-Val. Fisher–schultz lecture: Generic machine learning inference on heterogeneous treatment effects in randomized experiments, with an application to immunization in india. *Econometrica*, 93(4):1121–1164, 2025. doi: <https://doi.org/10.3982/ECTA19303>. URL <https://onlinelibrary.wiley.com/doi/abs/10.3982/ECTA19303>.
 - [13] George Casella and Roger L. Berger. *Statistical Inference*. Wadsworth Group, 2002.
 - [14] Randall Lewis. Doppelgangers v. twins for out-of-sample evaluation. *Internal Amazon Working Paper*, 2025.

A Appendix: Proof of Claims

Proposition 1 (Evaluating plug-in methodologies without unit-level partitioning): Let splits a and b be formed by randomly partitioning units in an A/B test with proportions α and $(1 - \alpha)$, respectively. Under equation (1), the joint distribution of $(\hat{\Delta}_{ias}, \hat{\Delta}_{ibs})$ conditional on the full-sample estimates $(\hat{\delta}_i, \tau_i^2)$ and true impact δ_i is given by:

$$\begin{pmatrix} \hat{\Delta}_{ias} \\ \hat{\Delta}_{ibs} \end{pmatrix} \Big| \hat{\Delta}_i = \hat{\delta}_i, \Delta_i = \delta_i, \tau_i^2 \stackrel{\text{ind}}{\sim} N \left(\begin{pmatrix} \delta_i \\ \hat{\delta}_i \end{pmatrix}, \begin{pmatrix} \frac{\tau_i^2}{\alpha} - \tau_i^2 & -\tau_i^2 \\ -\tau_i^2 & \frac{\tau_i^2}{1-\alpha} - \tau_i^2 \end{pmatrix} \right) \quad (26)$$

For a given test, the process of repeatedly partitioning units and re-estimating $(\hat{\delta}_{ias}, \hat{\delta}_{ibs})$ can therefore be approximated by resampling from the conditional distribution in equation (9) given the full-sample estimates $(\hat{\delta}_i, \tau_i^2)$. Given draws from this distribution, plug-in methodologies (being deterministic functions of $(\hat{\delta}_{ias}, \tau_{ias}^2)$) can then be applied, yielding estimates $\{\kappa(\hat{x}_{ias}), \hat{\delta}_{ias}, \tau_{ias}^2, \hat{\delta}_{ibs}, \tau_{ibs}^2\}_{s=1}^S$.

Proof: By central limit theorem and the random partitioning of units into disjoint splits, we have that:

$$\begin{pmatrix} \hat{\Delta}_{ias} \\ \hat{\Delta}_{ibs} \end{pmatrix} \Big| \Delta_i = \delta_i, \tau_i^2 \stackrel{\text{ind}}{\sim} N \left(\begin{pmatrix} \delta_i \\ \delta_i \end{pmatrix}, \begin{pmatrix} \frac{\tau_i^2}{\alpha} & 0 \\ 0 & \frac{\tau_i^2}{1-\alpha} \end{pmatrix} \right) \quad (27)$$

Note that $\hat{\Delta}_i = \alpha \hat{\Delta}_{ias} + (1 - \alpha) \hat{\Delta}_{ibs}$ and that, for any multivariate normal vector V with mean μ_V and variance Σ_V , any affine transformation MV is also multivariate normal, with mean $M\mu_V$ and variance $M\Sigma_V M'$. Defining:

$$M = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ \alpha & 1 - \alpha \end{pmatrix} \quad (28)$$

we therefore have that:

$$\begin{pmatrix} \hat{\Delta}_{ias} \\ \hat{\Delta}_{ibs} \\ \hat{\Delta}_i \end{pmatrix} \Big| \Delta_i = \delta_i, \tau_i^2 \stackrel{\text{ind}}{\sim} N \left(\begin{pmatrix} \delta_i \\ \delta_i \\ \delta_i \end{pmatrix}, \begin{pmatrix} \frac{\tau_i^2}{\alpha} & 0 & \tau_i^2 \\ 0 & \frac{\tau_i^2}{1-\alpha} & \tau_i^2 \\ \tau_i^2 & \tau_i^2 & \tau_i^2 \end{pmatrix} \right) \quad (29)$$

Finally, note that if:

$$\begin{pmatrix} V \\ W \end{pmatrix} \Big| \Delta_i = \delta_i, \tau_i^2 \stackrel{\text{ind}}{\sim} N \left(\begin{pmatrix} \mu_V \\ \mu_W \end{pmatrix}, \begin{pmatrix} \Sigma_{VV} & \Sigma_{VW} \\ \Sigma_{VW} & \Sigma_{WW} \end{pmatrix} \right) \quad (30)$$

then $V|W = w, \Delta_i = \delta_i, \tau_i^2 \stackrel{\text{ind}}{\sim} N(\mu_V + \Sigma_{VW}\Sigma_{WW}^{-1}(w - \mu_W), \Sigma_{VV} - \Sigma_{VW}\Sigma_{WW}^{-1}\Sigma_{WV})$. Applying this result with $V = (\hat{\Delta}_{ias}, \hat{\Delta}_{ibs})'$ and $W = \hat{\Delta}_i$ yields:

$$\mu_V + \Sigma_{VW}\Sigma_{WW}^{-1}(w - \mu_W) = \begin{pmatrix} \delta_i \\ \delta_i \end{pmatrix} + \begin{pmatrix} \tau_i^2 \\ \tau_i^2 \end{pmatrix} \begin{pmatrix} 1 \\ \frac{1}{\tau_i^2} \end{pmatrix} (\delta_i - \delta_i) = \begin{pmatrix} \delta_i \\ \delta_i \end{pmatrix} \quad (31)$$

$$\Sigma_{VV} - \Sigma_{VW}\Sigma_{WW}^{-1}\Sigma_{WV} = \begin{pmatrix} \frac{\tau_i^2}{\alpha} & 0 \\ 0 & \frac{\tau_i^2}{1-\alpha} \end{pmatrix} - \begin{pmatrix} \tau_i^2 \\ \tau_i^2 \end{pmatrix} \begin{pmatrix} 1 \\ \frac{1}{\tau_i^2} \end{pmatrix} (\tau_i^2 \quad \tau_i^2) = \begin{pmatrix} \frac{\tau_i^2}{\alpha} - \tau_i^2 & -\tau_i^2 \\ -\tau_i^2 & \frac{\tau_i^2}{1-\alpha} - \tau_i^2 \end{pmatrix} \quad (32)$$

which therefore establishes the claim \square

Lemma 1 (Test-level unbiasedness): Under equations (7) and (8) the sample-split estimators $\hat{Y}_{is}(\kappa, \alpha)$ listed in table 1 are unbiased for their corresponding sample-split estimands $Y_i(\kappa, \alpha)$.

Proof: Assume that equations (7) and (8) hold. For bias, the claim then follows immediately from the unbiasedness of $\hat{\Delta}_{ibs}$. For squared error, we follow [9], noting that:

$$E_F \left(\left(\kappa \left(\hat{X}_{ias} \right) - \hat{\Delta}_{ibs} \right)^2 \right) - \tau_{ibs}^2 = E_F \left(\left(\kappa \left(\hat{X}_{ias} \right) - \delta_i + \delta_i - \hat{\Delta}_{ibs} \right)^2 \right) - \tau_{ibs}^2 \quad (33)$$

$$= E_F \left(\left(\kappa \left(\hat{X}_{ias} \right) - \delta_i \right)^2 + \left(\delta_i - \hat{\Delta}_{ibs} \right)^2 \right) - \tau_{ibs}^2 \quad (34)$$

$$= E_F \left(\left(\kappa \left(\hat{X}_{ias} \right) - \delta_i \right)^2 \right) \quad (35)$$

where equation (34) follows from the independence of $\kappa(\hat{X}_{ias})$ and $\hat{\Delta}_{ibs}$ conditional on $\Delta_i = \delta_i$ and the unbiasedness of $\hat{\Delta}_{ibs}$. The claim then follows in equation (35) from the definition of the sampling variance of $\hat{\Delta}_{ibs}$. For launch-only decision value, we follow [9], noting that:

$$E_F \left(\kappa \left(\hat{X}_{ias} \right) \cdot \hat{\Delta}_{ibs} \right) = E_F \left(\kappa \left(\hat{X}_{ias} \right) \right) E_F \left(\hat{\Delta}_{ibs} \right) \quad (36)$$

$$= E_F \left(\kappa \left(\hat{X}_{ias} \cdot \delta_i \right) \right) \quad (37)$$

where equations (36) and (37) follow from the independence of $\kappa(\hat{X}_{ias})$ and $\hat{\Delta}_{ibs}$ conditional on $\Delta_i = \delta_i$ and the unbiasedness of $\hat{\Delta}_{ibs}$. The same argument applies to decision value \square

Proposition 2 (Unbiasedness): Under equations (11) and (15), the estimator of average performance $\hat{\theta}(\kappa, \alpha, S)$ is unbiased for $\theta(\kappa, \alpha)$.

Proof: Under equations (11) and (15), we have that:

$$E \left(\hat{\theta}(\kappa, \alpha, S) \right) = E_G \left(E_F \left(\hat{\theta}(\kappa, \alpha, S) \right) \right) \quad (38)$$

$$= E_G \left(E_F \left(\frac{1}{I} \sum_{i=1}^I \hat{Y}_i(\kappa, \alpha, S) \right) \right) \quad (39)$$

$$= E_G \left(\frac{1}{I} \sum_{i=1}^I E_F \left(\hat{Y}_i(\kappa, \alpha, S) \right) \right) \quad (40)$$

$$= E_G \left(Y_i(\kappa, \alpha) \right) \quad (41)$$

$$= \theta(\kappa, \alpha) \quad (42)$$

where equation (38) follows from the law of iterated expectations, equation (39) follows from the definition of $\hat{\theta}(\kappa, \alpha, S)$, equation (40) follows from the linearity of expectations, equation (41) follows from equation (15), and equation (42) follows from equation (11) \square

Lemma 2 (Consistency): Under equations (11) and (15) and the regularity condition that $\sup_i E(\hat{Y}_i(\kappa, \alpha, S)^2) < \infty$, the estimator $\hat{\theta}(\kappa, \alpha, S)$ is consistent for $\theta(\kappa, \alpha)$.

Proof: Under equations (11) and (15) and the regularity condition that $\sup_i E(\hat{Y}_i(\kappa, \alpha, S)^2) < \infty$:

$$\hat{\theta}(\kappa, \alpha, S) \xrightarrow{p} E \left(\hat{Y}_i(\kappa, \alpha, S) \right) \quad (43)$$

$$= E_G \left(E_F \left(\hat{Y}_i(\kappa, \alpha, S) \right) \right) \quad (44)$$

$$= E_G \left(Y_i(\kappa, \alpha) \right) \quad (45)$$

$$= \theta(\kappa, \alpha) \quad (46)$$

where equation (43) follows from the law of large numbers, equation (44) follows from the law of iterated expectations, equation (45) follows from equation (15), and equation (46) follows from equation (11) \square

Proposition 3 (Asymptotic normality): Under equations (11) and (15) and the regularity condition that $\sup_i E(\hat{Y}_i(\kappa, \alpha, S)^2) < \infty$, the estimator $\hat{\theta}(\kappa, \alpha, S)$ is asymptotically normally distributed:

$$\sqrt{I} \left(\hat{\theta}(\kappa, \alpha, S) - \theta(\kappa, \alpha) \right) \xrightarrow{d} N \left(0, \zeta^2(\kappa, \alpha, S) \right) \quad (47)$$

where $\zeta^2(\kappa, \alpha, S) = \lambda^2(\kappa, \alpha) + E_G(\Omega_i^2(\kappa, \alpha, S))$.

Proof: First, note that:

$$\text{Var}\left(\hat{Y}_i(\kappa, \alpha, S)\right) = \text{Var}_G\left(E_F\left(\hat{Y}_i(\kappa, \alpha, S)\right)\right) + E_G\left(\text{Var}_F\left(\hat{Y}_i(\kappa, \alpha, S)\right)\right) \quad (48)$$

$$= \text{Var}_G\left(Y_i(\kappa, \alpha)\right) + E_G\left(\Omega_i^2(\kappa, \alpha, S)\right) \quad (49)$$

$$= \lambda^2(\kappa, \alpha) + E_G\left(\Omega_i^2(\kappa, \alpha, S)\right) \quad (50)$$

where equation (48) follows from the law of total variance, equation (49) follows from equation (15), and equation (50) follows from equation (11).

Given the independence of $\hat{Y}_i(\kappa, \alpha, S)$ across tests, the regularity condition that $\sup_i E(\hat{Y}_i(\kappa, \alpha, S)^2) < \infty$, and lemma 1, the result then follows from the Lindeberg-Feller central limit theorem \square

Proposition 4 (Valid asymptotic inference): Under equations (11) and (15) and the regularity condition that $\sup_i E(\hat{Y}_i(\kappa, \alpha, S)^2) < \infty$, the estimator:

$$\hat{\zeta}^2(\kappa, \alpha, S) = \frac{1}{I} \sum_{i=1}^I (\hat{Y}_i(\kappa, \alpha, S) - \hat{\theta}(\kappa, \alpha, S))^2 \quad (51)$$

is consistent for the asymptotic variance $\zeta^2(\kappa, \alpha, S) = \lambda^2(\kappa, \alpha) + E_G(\Omega_i^2(\kappa, \alpha, S))$. Therefore, the $(1 - v)$ confidence intervals:

$$\hat{\theta}(\kappa, \alpha, S) \pm \frac{\hat{\zeta}(\kappa, \alpha, S)}{\sqrt{I}} \cdot \Phi^{-1}\left(1 - \frac{v}{2}\right) \quad (52)$$

will cover $\theta(\kappa, \alpha)$ with probability $(1 - v)$ as $I \rightarrow \infty$.

Proof: We can decompose the estimator $\hat{\zeta}^2(\kappa, \alpha, S)$ as follows:

$$\hat{\zeta}^2(\kappa, \alpha, S) = \frac{1}{I} \sum_{i=1}^I \left((\hat{Y}_i - \theta)^2 - 2(\hat{Y}_i - \theta)(\hat{\theta} - \theta) + (\hat{\theta} - \theta)^2 \right) \quad (53)$$

$$= \frac{1}{I} \sum_{i=1}^I (\hat{Y}_i - \theta)^2 - \frac{2(\hat{\theta} - \theta)}{I} \sum_{i=1}^I (\hat{Y}_i - \theta) + (\hat{\theta} - \theta)^2 \quad (54)$$

$$= \frac{1}{I} \sum_{i=1}^I (\hat{Y}_i - \theta)^2 - (\hat{\theta} - \theta)^2 \quad (55)$$

where we suppress the (κ, α, S) and (κ, α) notation in the above for simplicity. Under equations (11) and (15) and the regularity condition that $\sup_i E(\hat{Y}_i(\kappa, \alpha, S)^2) < \infty$, the first term converges in probability to $\zeta^2(\kappa, \alpha, S)$ by the law of large numbers. The claim that $\hat{\zeta}^2(\kappa, \alpha, S) \xrightarrow{p} \zeta^2(\kappa, \alpha, S)$ then follows from lemma 2 and the continuous mapping theorem, which together imply that the second term $(\hat{\theta}(\kappa, \alpha, S) - \theta(\kappa, \alpha))^2$ converges in probability to zero. Given this result, Slutsky's theorem then implies that:

$$\frac{\sqrt{I}(\hat{\theta}(\kappa, \alpha, S) - \theta(\kappa, \alpha))}{\hat{\zeta}(\kappa, \alpha, S)} \xrightarrow{d} N(0, 1) \quad (56)$$

from which the claim regarding confidence intervals follows immediately \square

B Appendix: Supplemental Results

Figure 4: Bias-variance tradeoff underlying choice of split fraction α (robustness)




Figure 5: Variance as a function of the number of tests I and number of partitions S (robustness)

