

ISOTOPES OF BIRACKS AND ZHANG TWISTS OF ALGEBRAS

XIAOLAN YU AND YANFEI ZHANG

ABSTRACT. In this paper, we introduce the notion of an \mathbb{N}^p -graded birack and construct its isotope. Every involutive \mathbb{N}^p -graded birack gives rise to an \mathbb{N}^p -graded Yang-Baxter algebra. We study the relation between isotopes of involutive \mathbb{N}^p -graded biracks and Zhang twists of \mathbb{N}^p -graded Yang-Baxter algebras. As an example, Yang-Baxter algebras determined by distributive solutions are proved to be Zhang twists of polynomial algebras.

INTRODUCTION

The Yang–Baxter equation is a cornerstone of mathematical physics that arises in the study of integrable systems, quantum groups, and quantum information theory. Its origins trace back to foundational papers by Yang [19] and Baxter [1]. Let V be a vector space. A solution of the Yang–Baxter equation is a linear mapping $R : V \otimes V \rightarrow V \otimes V$ such that the equality

$$(\text{id} \otimes R)(R \otimes \text{id})(\text{id} \otimes R) = (R \otimes \text{id})(\text{id} \otimes R)(R \otimes \text{id})$$

of operators hold in the space $\text{End}(V \otimes V \otimes V)$. The problem of describing all possible solutions is highly complex and therefore there were some simplifications introduced.

A particularly nice class of solutions is provided by set-theoretic solutions proposed by Drinfeld [5]. Let X be a set and $r : X \times X \rightarrow X \times X$ a bijective mapping. r is called a set-theoretic solution of the Yang-Baxter equation if the following braid relation holds

$$(\text{id} \times r)(r \times \text{id})(\text{id} \times r) = (r \times \text{id})(\text{id} \times r)(r \times \text{id}).$$

It is clear that a set-theoretic solution extends to a linear one, but more important than this is that set-theoretic solutions lead to their own remarkable algebraic and combinatoric structure. During the last three decades, the study

2020 *Mathematics Subject Classification.* 16T25, 20N05, 16W50, 16S80.

Key words and phrases. Yang-Baxter Equation; Set-theoretic solution; One-sided quasi-group; Birack; Isotope; Zhang twist.

of set-theoretic solutions and related structures has notably intensified, see for instance [3], [4], [6], [7], [12], [10], [11], [13], [14], [16].

On the algebra side, each set-theoretic solution (X, r) associates a quadratic Yang-Baxter algebra $A(X, r)$ as proposed in [11, Section 6]. If X is finite and r is nondegenerate and involutive, then $A(X, r)$ has remarkable algebraic, homological and combinatorial properties [8], [13]. Our motivation of this paper is to try to find more ways to study properties of Yang-Baxter algebras.

It is known that there is a one-to-one correspondence between non-degenerate, involutive set-theoretic solutions of the Yang-Baxter equation and involutive biracks. A birack $(X, \circ, \backslash_\circ, \bullet, /_\bullet)$ is an algebra such that $(X, \circ, \backslash_\circ)$ is a left quasigroup, $(X, \bullet, /_\bullet)$ is a right quasigroup, and some additional identities are satisfied. Such correspondence allows us to study solutions of the Yang-Baxter equations via involutive biracks.

In quasigroup theory (see, e.g., [17, Section II.2]), isotopy is a standard method for transforming one quasigroup into another. In their work of involutive set-theoretic solutions of the Yang-Baxter equation of multipermutation level 2, Jedlička, Pilitowska, and Zamojska-Dzienio used a special form of isotope to construct non-distributive solutions from distributive solutions [15]. Inspired by their work, in Section 2, we introduce the notions of an \mathbb{N}^p -graded birack and its isotope, and then discuss their properties. Our main theorem is presented in Section 3. Each involutive \mathbb{N}^p -graded birack $(X, \circ, \backslash_\circ, \bullet, /_\bullet)$ determines an \mathbb{N}^p -graded Yang-Baxter algebra $A(X, \mathbf{R}^\circ)$. We find that under some conditions, the algebra $A^*(X, \mathbf{R}^*)$ determined by an isotope $(X, *, \backslash_*, \diamond, /_\diamond)$ of the birack $(X, \circ, \backslash_\circ, \bullet, /_\bullet)$ is isomorphic to a Zhang twist of the algebra $A(X, \mathbf{R}^\circ)$ (Theorem 3.2). The Zhang twist is a twist of graded algebras introduced by J.J. Zhang in order to describe graded algebras whose graded module categories are equivalent [20]. Many important properties of graded algebras, including Gelfand–Kirillov dimension, the Noetherian property, being a domain, global dimension, and Artin–Schelter regularity are preserved under Zhang twist. As an example our main theorem, we show that the Yang–Baxter algebras arising from distributive solutions are Zhang twists of polynomial algebras. This class of algebras is believed to share properties similar to those of polynomial algebras. We will discuss them in detail in a subsequent paper.

1. PRELIMINARIES

Throughout \mathbb{k} is a base field of characteristic zero. All vector spaces and algebras are over \mathbb{k} .

1.1. Quasigroups and biracks. A *left quasigroup* is an algebra $(X, \circ, \backslash_\circ)$ with two binary operations: \circ and \backslash_\circ , such that for every $x, y \in X$, the following conditions holds

$$(1.1) \quad x \circ (x \backslash_\circ y) = y = x \backslash_\circ (x \circ y).$$

A *right quasigroup* is defined analogously as an algebra $(X, \bullet, /\bullet)$ with two binary operations: \bullet and $/\bullet$, such that for every $x, y \in X$, the following conditions holds

$$(1.2) \quad (y /\bullet x) \bullet x = y = (y \bullet x) /\bullet x.$$

Let $(X, \circ, \backslash_\circ)$ be a left quasigroup. For each $x \in X$, the *left translation* $\mathcal{L}_x : X \rightarrow X$ by x is defined as

$$\mathcal{L}_x(a) = x \circ a,$$

for any $a \in X$. Condition (1.1) guarantees that each \mathcal{L}_x is bijective, with $\mathcal{L}_x^{-1}(a) = x \backslash_\circ a$. Similarly, Condition (1.2) gives that each *right translation* $\mathcal{R}_x : X \rightarrow X$ by x defined by $\mathcal{R}_x(a) = a \bullet x$ is bijective with $\mathcal{R}_x^{-1}(a) = a /\bullet x$.

For a left quasigroup $(X, \circ, \backslash_\circ)$, a bijection $f : X \rightarrow X$ is called an *automorphism* of $(X, \circ, \backslash_\circ)$ if

$$f(x \circ y) = f(x) \circ f(y),$$

for every $x, y \in X$. It is easy to see that a bijection $f : X \rightarrow X$ is an automorphism of $(X, \circ, \backslash_\circ)$ if and only if

$$f\mathcal{L}_x f^{-1} = \mathcal{L}_{f(x)},$$

for every $x \in X$.

An automorphism of a right quasigroup can be defined analogously. A bijection $f : X \rightarrow X$ is an automorphism of $(X, \bullet, /\bullet)$ if and only if

$$f\mathcal{R}_x f^{-1} = \mathcal{R}_{f(x)},$$

for every $x \in X$.

A left quasigroup $(X, \circ, \backslash_\circ)$ is called

- *non-degenerate* if the mapping

$$\begin{array}{ccc} X & \longrightarrow & X \\ x & \longmapsto & x \backslash_\circ x \end{array}$$

is a bijection;

- *right cyclic* if for every $x, y, z \in X$:

$$(x \setminus_{\circ} y) \setminus_{\circ} (x \setminus_{\circ} z) = (y \setminus_{\circ} x) \setminus_{\circ} (y \setminus_{\circ} z),$$

or equivalently

$$\mathcal{L}_x \mathcal{L}_{x \setminus_{\circ} y} = \mathcal{L}_y \mathcal{L}_{y \setminus_{\circ} x}.$$

An algebra $(X, \circ, \setminus_{\circ}, \bullet, /_{\bullet})$ with four binary operations is called a *birack*, if $(X, \circ, \setminus_{\circ})$ is a left quasigroup, $(X, \bullet, /_{\bullet})$ is a right quasigroup and the following holds for any $x, y, z \in X$,

$$\begin{aligned} x \circ (y \circ z) &= (x \circ y) \circ ((x \bullet y) \circ z), \\ (x \circ y) \bullet ((x \bullet y) \circ z) &= (x \bullet (y \circ z)) \circ (y \bullet z), \\ (x \bullet y) \bullet z &= (x \bullet (y \circ z)) \bullet (y \bullet z). \end{aligned}$$

If the set X is finite, we will call the birack *finite*.

Convention 1.1. All biracks we study in this paper are finite.

A birack is called *involution* if it additionally satisfies

$$\begin{aligned} (x \circ y) \circ (x \bullet y) &= x \\ (x \circ y) \bullet (x \bullet y) &= y \end{aligned}$$

for every $x, y \in X$.

The following one to one correspondence between involutive biracks and non-degenerate right cyclic left quasigroups is well-known (see e.g. [2, Proposition 1.5]).

Lemma 1.2. (1) *Let $(X, \circ, \setminus_{\circ}, \bullet, /_{\bullet})$ be an involutive birack. Then $(X, \circ, \setminus_{\circ})$ is a non-degenerate right cyclic left quasigroup.*
 (2) *Conversely, let $(X, \circ, \setminus_{\circ})$ be a non-degenerate right cyclic left quasigroup. For every $x, y \in X$, define*

$$\begin{aligned} x \bullet y &= \mathcal{R}_y(x) = (x \circ y) \setminus_{\circ} x, \\ x /_{\bullet} y &= \mathcal{R}_y^{-1}(x). \end{aligned}$$

Then the algebra $(X, \circ, \setminus_{\circ}, \bullet, /_{\bullet})$ is an involutive birack.

The operations \setminus_{\circ} and $/_{\bullet}$ are connected by

$$(x \setminus_{\circ} x) /_{\bullet} (x \setminus_{\circ} x) = x \quad \text{and} \quad (x /_{\bullet} x) \setminus_{\circ} (x /_{\bullet} x) = x.$$

A birack $(X, \circ, \setminus_{\circ}, \bullet, /_{\bullet})$ is said to satisfy the condition **lri** provided that

$$(x \circ y) \bullet x = y = x \circ (y \bullet x),$$

all $x, y \in X$. That is, $\mathcal{R}_x = \mathcal{L}_x^{-1}$ for all $x \in X$.

Remark 1.3. An involutive birack $(X, \circ, \backslash, \bullet, /)$ satisfies Condition **lr1** if and only if

$$(1.3) \quad (x \circ y) \backslash x = y \backslash x,$$

for any $x, y \in X$. In fact, in an involutive birack $x \bullet y = (x \circ y) \backslash x$. Hence,

$$\mathcal{R}_y(x) = \mathcal{L}_y^{-1}(x) \iff (x \circ y) \backslash x = y \backslash x.$$

1.2. Set-theoretic solution of the Yang-Baxter equation. By a *quadratic set*, we mean a pair (X, r) , where X is a nonempty set and $r : X \times X \rightarrow X \times X$ a bijective mapping. We write the image of (x, y) under r as

$$r(x, y) = ({}^x y, x^y).$$

The above equation defines a “left action” $\mathcal{L} : X \times X \rightarrow X$, and a “right action” $\mathcal{R} : X \times X \rightarrow X$, on X as:

$$\mathcal{L}_x(y) = {}^x y, \quad \mathcal{R}_y(x) = x^y,$$

for all $x, y \in X$.

A quadratic set (X, r) is called:

- *non-degenerate* if the mappings \mathcal{L}_x and \mathcal{R}_x are bijections, for all $x \in X$;
- *involutive* if $r^2 = \text{id}_{X^2}$;
- *square free* if $r(x, x) = (x, x)$, for every $x \in X$;
- a *quantum binomial set* if it is nondegenerate, involutive and square-free.

A *set-theoretic solution of the Yang-Baxter equation* (YBE) is a quadratic set (X, r) , such that the following braid relation holds

$$(\text{id} \times r)(r \times \text{id})(\text{id} \times r) = (r \times \text{id})(\text{id} \times r)(r \times \text{id}).$$

In this case, (X, r) is also called a *braided set*. A solution is called *non-degenerate* (*involutive*, or *square free*) if it is non-degenerate (involutive, or square free) as a quadratic set. If X is a finite set, then (X, r) is called a finite solution.

Convention 1.4. In the followings, by a *solution* (of the YBE) we mean a set-theoretic, non-degenerate and involutive solution of the YBE. We consider only *finite* solutions.

Each quadratic set (X, r) determines a set of quadratic defining relations $\mathbf{R}(r)$ defined by

$$xy = y'x' \in \mathbf{R}(r) \text{ whenever } r(x, y) = (y', x').$$

The unital semigroup $S(X, r) = \langle X; \mathbf{R}(r) \rangle$, with a set of generators X and a set of defining relations $\mathbf{R}(r)$, is called the *semigroup associated with* (X, r) . The *group* $G(X, r)$ *associated with* (X, r) is defined analogously. The *algebra associated with* (X, r) is defined as $A(X, r) = \mathbb{k}\langle X \rangle / (\mathbf{R}(r))$. When (X, r) is a solution of the YBE, $S(X, r)$, resp. $G(X, r)$, $A(X, r)$ is called the *Yang-Baxter (YB) semigroup*, resp. the *YB group*, the *YB algebra*.

Let (X, r) be a nondegenerate quadratic set. It is well-known that (X, r) is a set-theoretic solution of the YBE if and only if the following condition holds for all $x, y, z \in X$ (e.g. [10, Lemma 2.5]).

$$\mathbf{11} : {}^x({}^y z) = {}^{xy}({}^{xy} z), \quad \mathbf{r1} : (x^y)^z = (x^y z)^{y^z}, \quad \mathbf{lr3} : (xy)^{(x^y(z))} = (x^{yz})(y^z).$$

For every non-degenerate braided set, condition **11** implies that the mapping $x \rightarrow \mathcal{L}_x$ extends canonically to a group homomorphism

$$\mathcal{L} : G(X, r) \longrightarrow \text{Sym}(X),$$

where $\text{Sym}(X)$ is symmetric group on X . The homomorphism \mathcal{L} defines the *canonical left action* of $G(X, r)$ on the set X . Similarly, **r1** gives the *canonical right action* of $G(X, r)$ on the set X .

The *YB permutation group* $\mathcal{G}(X, r)$ of a finite solution (X, r) is the subgroup of the symmetric group $\text{Sym}(X)$ on X generated by $\{\mathcal{L}_x : x \in X\}$.

It is known (see e.g. [2, Lemma 1.2]) that there is a one-to-one correspondence between solutions of the YBE and involutive biracks.

Lemma 1.5. (1) *If $(X, \circ, \backslash \circ, \bullet, / \bullet)$ is an involutive birack, then defining*

$$r(x, y) = (x \circ y, x \bullet y).$$

We obtain that (X, r) is a solution of the YB equation.

(2) *Let (X, r) be a solution of the YB equation. The bijective mapping r naturally defines a left action \mathcal{L} and a right action \mathcal{R} on X as:*

$$r(x, y) = (\mathcal{L}_x(y), \mathcal{R}_y(x)).$$

Then defining

$$\begin{aligned} x \circ y &= \mathcal{L}_x(y), & x \backslash \circ y &= \mathcal{L}_x^{-1}(y), \\ x \bullet y &= \mathcal{R}_y(x), & x / \bullet y &= \mathcal{R}_y^{-1}(x). \end{aligned}$$

We obtain an involutive birack $(X, \circ, \backslash \circ, \bullet, / \bullet)$.

Such a correspondence allows us to treat each solution as an involutive birack.

2. ISOTOPES OF QUASIGROUPS

This section serves as a preparation for Section 3. We introduce the notion of an \mathbb{N}^p -graded birack and generalize the isotope construction of [15] to this graded setting.

We say a left quasigroup $(X, \circ, \backslash \circ)$ is \mathbb{N}^p -graded if there is a partition on X :

$$X = X_1 \cup X_2 \cup \cdots \cup X_p$$

and for any $x \in X, y \in X_i$,

$$(2.1) \quad x \circ y \in X_i.$$

If $x \in X_i$, then the degree vector is assigned as:

$$\deg(x) = (0, \dots, 0, 1, 0, \dots, 0) \in \mathbb{N}^p,$$

where the 1 is in the i -th position. Condition (2.1) means that all left translation is degree-preserving.

An \mathbb{N}^p -graded automorphism f of $(X, \circ, \backslash \circ)$ is an automorphism of $(X, \circ, \backslash \circ)$ such that $f(X_i) \subseteq X_i$ for all $1 \leq i \leq p$.

Let $\phi := \{\phi_s \mid s = 1, \dots, p\}$ be a sequence of commuting degree-preserving bijection of the set X , i.e., for every $1 \leq s \leq p$, $\phi_s(x) \in X_i$ for any $x \in X_i$, $1 \leq i \leq p$. For a degree vector $\alpha = (a_1, \dots, a_p)$, set $\phi^\alpha = \phi_1^{a_1} \cdots \phi_p^{a_p}$. Define on the set X new binary operations:

$$x * y := x \circ \phi^{|x|}(y) = \mathcal{L}_x \phi^{|x|}(y),$$

$$x \backslash_* y := \phi^{-|x|}(x \backslash \circ y) = \phi^{-|x|} \mathcal{L}_x^{-1}(y).$$

The algebra $(X, *, \backslash_*)$ is called the ϕ -isotope of $(X, \circ, \backslash \circ)$.

It is easy to see that

$$x * (x \backslash_* y) = \mathcal{L}_x \phi^{|x|} \phi^{-|x|} \mathcal{L}_x^{-1}(y) = y$$

and

$$x \backslash_* (x * y) = \phi^{-|x|} \mathcal{L}_x^{-1} \mathcal{L}_x \phi^{|x|}(y) = y.$$

So the ϕ -isotope $(X, *, \backslash_*)$ is also a left quasigroup. Moreover, the grading of $(X, \circ, \backslash \circ)$ induces an \mathbb{N}^p -grading on $(X, *, \backslash_*)$.

Lemma 2.1. *Let $(X, \circ, \backslash \circ)$ be a non-degenerate \mathbb{N}^p -graded left quasigroup and let $\phi := \{\phi_s \mid s = 1, \dots, p\}$ be a sequence of commuting degree-preserving bijection of the set X . Then the ϕ -isotope $(X, *, \backslash_*)$ is also non-degenerate.*

Proof. The left quasigroup $(X, \circ, \backslash_\circ)$ is \mathbb{N}^p -graded, say

$$(2.2) \quad X = X_1 \cup X_2 \cup \cdots \cup X_p.$$

All left translations is degree-preserving. By the non-degeneracy of the left quasigroup, for each subset X_s , the mapping

$$\begin{aligned} X_s &\longrightarrow X_s \\ x &\longmapsto \mathcal{L}_x^{-1}(x) = x \backslash_\circ x \end{aligned}$$

is bijective. Each ϕ_s is bijective, so the mapping

$$\begin{aligned} X_s &\longrightarrow X_s \\ x &\longmapsto \phi_s \mathcal{L}_x^{-1}(x) = x \backslash_* x \end{aligned}$$

is also bijective. (2.2) is a partition of X . Therefore, the mapping

$$\begin{aligned} T_\phi : X &\longrightarrow X \\ x &\longmapsto \phi_s \mathcal{L}_x^{-1}(x) = x \backslash_* x \end{aligned}$$

is bijective. The ϕ -isotope $(X, *, \backslash_*)$ is also non-degenerate. \square

Lemma 2.2. *Let $(X, \circ, \backslash_\circ)$ be an \mathbb{N}^p -graded right cyclic left quasigroup, and $\phi := \{\phi_s \mid s = 1, \dots, p\}$ be a sequence of commuting \mathbb{N}^p -graded automorphisms of $(X, \circ, \backslash_\circ)$ such that*

$$(2.3) \quad \mathcal{L}_{\phi_s(x)} = \mathcal{L}_x.$$

*for every $1 \leq s \leq p$ and any $x \in X$. Then the ϕ -isotope $(X, *, \backslash_*)$ of $(X, \circ, \backslash_\circ)$ is also right cyclic.*

Proof. We need to proof that

$$(x \backslash_* y) \backslash_* (x \backslash_* z) = (y \backslash_* x) \backslash_* (y \backslash_* z),$$

for every $x, y, z \in X$. Note that $\{\phi_s \mid s = 1, \dots, p\}$ is a sequence of automorphism of $(X, \circ, \backslash_\circ)$. Then $\mathcal{L}_{\phi_s(x)} = \phi_s \mathcal{L}_x \phi_s^{-1}$ for every $x \in X$ and $1 \leq s \leq p$. In this case, Condition (2.3) is equivalent to

$$(2.4) \quad \phi_s \mathcal{L}_x = \mathcal{L}_x \phi_s.$$

That is, every ϕ_s commutes with all left translations.

We have the following equations hold

$$\begin{aligned}
& (x \setminus_* y) \setminus_* (x \setminus_* z) \\
&= (\phi^{-|x|} \mathcal{L}_x^{-1}(y)) \setminus_* (\phi^{-|x|} \mathcal{L}_x^{-1}(z)) \\
&= \phi^{-|\phi^{-|x|}(\mathcal{L}_x^{-1}(y))|} \mathcal{L}_{\phi^{-|x|} \mathcal{L}_x^{-1}(y)}^{-1} \phi^{-|x|} \mathcal{L}_x^{-1}(z) \\
&= \phi^{-|y|} \mathcal{L}_{\phi^{-|x|} \mathcal{L}_x^{-1}(y)}^{-1} \phi^{-|x|} \mathcal{L}_x^{-1}(z) \\
&\stackrel{(2.3)}{=} \phi^{-|y|} \mathcal{L}_{\mathcal{L}_x^{-1}(y)}^{-1} \phi^{-|x|} \mathcal{L}_x^{-1}(z) \\
&\stackrel{(2.4)}{=} \phi^{-(|x|+|y|)} \mathcal{L}_{\mathcal{L}_x^{-1}(y)}^{-1} \mathcal{L}_x^{-1}(z) \\
&= \phi^{-(|x|+|y|)} \mathcal{L}_{\mathcal{L}_y^{-1}(x)}^{-1} \mathcal{L}_y^{-1}(z) \quad ((X, \circ, \setminus_\circ) \text{ is right cyclic}) \\
&\stackrel{(2.4)}{=} \phi^{-|x|} \mathcal{L}_{\mathcal{L}_y^{-1}(x)}^{-1} \phi^{-|y|} \mathcal{L}_y^{-1}(z) \\
&\stackrel{(2.3)}{=} \phi^{-|x|} \mathcal{L}_{\phi^{-|y|} \mathcal{L}_y^{-1}(x)}^{-1} (x) \phi^{-|y|} \mathcal{L}_y^{-1}(z) \\
&= \phi^{-|\phi^{-|y|} \mathcal{L}_y^{-1}(x)|} \mathcal{L}_{\phi^{-|y|} \mathcal{L}_y^{-1}(x)}^{-1} (x) \phi^{-|y|} \mathcal{L}_y^{-1}(z) \\
&= (\phi^{-|y|} \mathcal{L}_y^{-1}(x)) \setminus_* (\phi^{-|y|} \mathcal{L}_y^{-1}(z)) \\
&= (y \setminus_* x) \setminus_* (y \setminus_* z).
\end{aligned}$$

The third and the last third equation follow from the fact that $(X, \circ, \setminus_\circ)$ is an \mathbb{N}^p -graded left quasigroup and $\phi := \{\phi_s \mid s = 1, \dots, p\}$ is a sequence of \mathbb{N}^p -graded automorphism of $(X, \circ, \setminus_\circ)$. \square

We know that involutive biracks are in one-to-one correspondence with non-degenerate right cyclic left quasigroups. Now we are ready to define the isotope of an \mathbb{N}^p -birack.

A birack $(X, \circ, \setminus_\circ, \bullet, \setminus_\bullet)$ is called \mathbb{N}^p -graded if there is a partition on X :

$$X = X_1 \cup X_2 \cup \dots \cup X_p$$

and for any $x \in X, y \in X_i$,

$$x \circ y \in X_i, \text{ and } y \bullet x \in X_i.$$

Remark 2.3. If $(X, \circ, \setminus_\circ, \bullet, /_\bullet)$ is an involutive birack, then $(X, \circ, \setminus_\circ)$ is a non-degenerate right cyclic left quasigroup. The birack $(X, \circ, \setminus_\circ, \bullet, /_\bullet)$ is an \mathbb{N}^p -graded birack if and only if the $(X, \circ, \setminus_\circ)$ is an \mathbb{N}^p -graded left quasigroup.

Let $(X, \circ, \setminus_\circ, \bullet, /_\bullet)$ be an \mathbb{N}^p -graded involutive birack. Then $(X, \circ, \setminus_\circ)$ is an \mathbb{N}^p -graded non-degenerate right cyclic left quasigroup. Let $\phi := \{\phi_s \mid s = 1, \dots, p\}$ be a sequence of commuting degree-preserving bijection of the set X such that the ϕ -isotope $(X, *, \setminus_*)$ of $(X, \circ, \setminus_\circ)$ is right cyclic. Then the left quasigroup $(X, *, \setminus_*)$ uniquely determines the involutive birack $(X, *, \setminus_*, \diamond, /_\diamond)$ as follows

$$(2.5) \quad x * y = \mathcal{L}_x \phi^{|x|}(y),$$

$$(2.6) \quad x \diamond y = (x * y) \setminus_* x = \phi^{-|y|} \mathcal{L}_{\mathcal{L}_x \phi^{|x|}(y)}^{-1}(x) = \phi^{-|y|}(x \bullet \phi^{|x|}(y)).$$

The birack $(X, *, \setminus_*, \diamond, / \diamond)$ is called the ϕ -isotope of $(X, \circ, \setminus_\circ, \bullet, / \bullet)$. It is easy to see that $(X, *, \setminus_*, \diamond, / \diamond)$ is also an \mathbb{N}^p -graded birack.

Lemma 2.4. *Let $(X, \circ, \setminus_\circ, \bullet, / \bullet)$ be an \mathbb{N}^p -graded involutive birack satisfying Condition **lri**, and $\phi := \{\phi_s \mid s = 1, \dots, p\}$ be a sequence of commuting degree-preserving bijection of the set X . Consider the ϕ -isotope $(X, *, \setminus_*, \diamond, / \diamond)$ of the set X . Then $(X, *, \setminus_*, \diamond, / \diamond)$ satisfies Condition **lri** if and only if $(X, \circ, \setminus_\circ, \bullet, / \bullet)$ satisfies Condition (2.3). In this case,*

$$y \diamond x = x \setminus_* y = \phi^{-|x|} \mathcal{L}_x^{-1}(y),$$

for any $x, y \in X$.

Proof. It is clear that if $(X, *, \setminus_*, \diamond, / \diamond)$ satisfies Condition **lri**, then

$$y \diamond x = x \setminus_* y = \phi^{-|x|} \mathcal{L}_x^{-1}(y),$$

for any $x, y \in X$.

Indeed, for every $1 \leq s \leq p$, $y \in X_s$ and $x \in X$, we have

$$\begin{aligned} y \diamond x = x \setminus_* y &\iff \phi^{-|x|} \mathcal{L}_{\mathcal{L}_y \phi^{|y|}(x)}^{-1}(y) = \phi^{-|x|} \mathcal{L}_x^{-1}(y) \\ &\iff \phi^{-|x|}(y \bullet \phi^{|y|}(x)) = \phi^{-|x|}(y \bullet x) \\ &\iff y \bullet \phi^{|y|}(x) = y \bullet x \\ &\iff \mathcal{R}_{\phi^{|y|}(x)} = \mathcal{R}_x \\ &\iff \mathcal{L}_{\phi_s(x)} = \mathcal{L}_x. \end{aligned}$$

Therefore, $(X, *, \setminus_*, \diamond, / \diamond)$ satisfies Condition **lri** if and only if $(X, \circ, \setminus_\circ, \bullet, / \bullet)$ satisfies Condition (2.3) \square

Proposition 2.5. *Let $(X, \circ, \setminus_\circ, \bullet, / \bullet)$ be an \mathbb{N}^p -graded involutive birack satisfying Condition **lri**, and $\phi := \{\phi_s \mid s = 1, \dots, p\}$ be a sequence of commuting graded automorphisms of $(X, \circ, \setminus_\circ, \bullet, / \bullet)$ satisfying Condition (2.3). Then the ϕ -isotope $(X, *, \setminus_*, \diamond, / \diamond)$ of $(X, \circ, \setminus_\circ, \bullet, / \bullet)$ is also an \mathbb{N}^p -graded involutive birack satisfying Condition **lri**.*

Proof. $(X, \circ, \setminus_\circ, \bullet, / \bullet)$ is an \mathbb{N}^p -graded involutive birack. Then $(X, \circ, \setminus_\circ)$ is a non-degenerate right cyclic left quasigroup by Lemma 1.2. By Lemma 2.1 and 2.2, the ϕ -isotope $(X, *, \setminus_*)$ is also non-degenerate right cyclic. It determines the \mathbb{N}^p -graded involutive birack $(X, *, \setminus_*, \diamond, / \diamond)$. By Lemma 2.4, $(X, *, \setminus_*, \diamond, / \diamond)$ also satisfies Condition **lri**. \square

3. ISOTOPES AND ZHANG TWISTS

In this section, we will show the relation between the isotopes of involutive biracks and the Zhang twists of algebras.

Let $(X, \circ, \backslash_\circ, \bullet, /_\bullet)$ be an involutive birack. It canonically determines an algebra $A = A(X, \mathbf{R}^\circ)$ generated by X with quadratic defining relations \mathbf{R}° defined by

$$xy = (x \circ y)(x \bullet y)$$

for any $x, y \in X$.

If $(X, \circ, \backslash_\circ, \bullet, /_\bullet)$ is an \mathbb{N}^p -graded birack, then the grading on X naturally defines a grading on A . The algebra A is an \mathbb{N}^p -graded algebra.

Remark 3.1. For an $(\mathbb{N}^p$ -graded) involutive birack $(X, \circ, \backslash_\circ, \bullet, /_\bullet)$, the $(\mathbb{N}^p$ -graded) algebra $A = A(X, \mathbf{R}^\circ)$ is just the YB algebra of the solution corresponding to the involutive birack.

Let B be any \mathbb{N}^p -graded algebra. Let $\phi := \{\phi_s \mid s = 1, \dots, p\}$ be a sequence of commuting \mathbb{N}^p -graded algebra automorphisms of B . Then the *Zhang twist* B^ϕ of B by ϕ is defined as follows: $B^\phi = B$ as a \mathbb{N}^p -graded vector space, and the new multiplication \star of B^ϕ is determined by

$$a \star b = a\phi^{|a|}(b)$$

where $\phi^{|a|} = \phi_1^{a_1} \cdots \phi_p^{a_p}$ if the degree of a is $|a| = (a_1, \dots, a_p)$.

Let $(X, \circ, \backslash_\circ, \bullet, /_\bullet)$ be an \mathbb{N}^p -graded involutive birack satisfying condition **lri**. It determines an \mathbb{N}^p -graded algebra $A = A(X, \mathbf{R}^\circ)$. Let $\phi := \{\phi_s \mid s = 1, \dots, p\}$ be a sequence of commuting graded automorphism of $(X, \circ, \backslash_\circ, \bullet, /_\bullet)$ satisfying Condition (2.3). It is easy to see that $\{\phi_s \mid s = 1, \dots, p\}$ naturally induce a sequence of commuting graded automorphism on the algebra A . By Proposition 2.5, the ϕ -isotope $(X, *, \backslash_*, \diamond, /_\diamond)$ of $(X, \circ, \backslash_\circ, \bullet, /_\bullet)$ is also an \mathbb{N}^p -graded involutive birack satisfying condition **lri**. It determines an \mathbb{N}^p -graded algebra $A^* = A^*(X, \mathbf{R}^*)$.

Theorem 3.2. *Keep the notations as above, the algebra $A^* = (A^*, X, \mathbf{R}^*)$ obtained by the isotope $(X, *, \backslash_*, \diamond, /_\diamond)$ is isomorphic to the Zhang twist A^ϕ of A by ϕ .*

Proof. For the algebra $A^* = (A^*, X, \mathbf{R}^*)$, the relation \mathbf{R}^* is defined by

$$\begin{aligned} xy &= (x * y)(x \diamond y) = (x * y)(y \backslash_* x) \\ &= \mathcal{L}_x(\phi^{|x|}(y))\phi^{-|y|}(\mathcal{L}_y^{-1}(x)). \end{aligned}$$

We need to prove that A^ϕ is isomorphic to the algebra generated by X with relation \mathbf{R}^* . We have

$$\begin{aligned}
x \star y &= x\phi^{|x|}(y) \\
&= (x \circ \phi^{|x|}(y))(\phi^{|x|}(y) \setminus \circ x) \\
&= \mathcal{L}_x(\phi^{|x|}(y))\mathcal{L}_{\phi^{|x|}(y)}^{-1}(x) \\
&\stackrel{(\mathbf{lri})}{=} \mathcal{L}_x(\phi^{|x|}(y))\mathcal{L}_y^{-1}(x) \\
&= \mathcal{L}_x(\phi^{|x|}(y)) \star \phi^{-|\mathcal{L}_x(\phi^{|x|}(y))|}(\mathcal{L}_y(x)) \\
&= \mathcal{L}_x(\phi^{|x|}(y)) \star \phi^{-|y|}(\mathcal{R}_y(x)).
\end{aligned}$$

The last equation follows from that each ϕ_s ($1 \leq s \leq p$) and all left translation preserve degree. Therefore, the Zhang twist of A^ϕ is isomorphic to the algebra A^* . \square

Since the Zhang twist preserves many important properties of graded algebras, with the help of Theorem 3.2, we can reduce the study of a structurally complicated YB algebra to that of a simpler one via isotopy. The following is an example, we will show that YB algebras of distributive solutions are Zhang twists of polynomial algebras.

Example 3.3. A birack $(X, \circ, \setminus \circ, \bullet, / \bullet)$ is called *distributive*, if for every $x, y, z \in X$,

$$\begin{aligned}
x \circ (y \circ z) &= (x \circ y) \circ (x \circ z), \\
(y \bullet z) \bullet x &= (y \bullet x) \bullet (z \bullet x).
\end{aligned}$$

If the birack $(X, \circ, \setminus \circ, \bullet, / \bullet)$ involutive, then it is distributive if and only if for every $x, y, z \in X$,

$$(3.1) \quad x \circ (y \circ z) = (x \circ z) \circ (x \circ z),$$

i.e., $\mathcal{L}_x\mathcal{L}_y = \mathcal{L}_{x \circ y}\mathcal{L}_x$ [15, Corollary 5.7]. Following from [15, Corollary 5.4] and [12, Lemma 7.1], we know that an involutive distributive birack satisfies Condition **lri**.

A solution of YBE is called *distributive*, if it corresponds to a distributive involutive birack.

Let $(X, \circ, \setminus \circ, \bullet, / \bullet)$ be an involutive distributive birack. There is an equivalence \sim defined on X :

$$x \sim y \quad \text{if and only if} \quad \mathcal{L}_x = \mathcal{L}_y.$$

This equivalence gives a partition on X . That is,

$$(3.2) \quad X = \bigcup_{x \in X} [x],$$

where $[x] = \{a \in X \mid x \sim a\}$. By Lemma 3.1 and Corollary 5.4 in [15], an involutive distributive birack is *2-reductive*, i.e.,

$$(3.3) \quad (x \circ y) \circ z = y \circ z \quad (\iff \mathcal{L}_{\mathcal{L}_x(y)} = \mathcal{L}_y)$$

for any $x, y, z \in X$. Then for any $y \in [x]$ and $x \in X$, $\mathcal{L}_x(y) \in [x]$, (3.2) gives a grading on the birack. Assume that X is partitioned into p equivalence classes, numbering them as X_1, \dots, X_p . The birack $(X, \circ, \backslash_\circ, \bullet, /_\bullet)$ is an \mathbb{N}^p -graded birack.

In each equivalence class X_s , select a representative element x_s . We have $\mathcal{L}_y = \mathcal{L}_{x_s}$, for any $y \in X_s$. We denote the inverse of these equal mappings as ϕ_s , i.e. $\phi_s = \mathcal{L}_{x_s}^{-1}$. Directly from (3.1) and (3.3), we obtain that all left translations are automorphisms of the birack satisfying Condition (2.3). Moreover, the permutation group generated by the left transformations are abelian. So $\phi := \{\phi_s \mid s = 1, \dots, p\}$ is a sequence of commuting graded automorphisms of the birack satisfying Condition (2.3). The ϕ -isotope of the birack $(X, \circ, \backslash_\circ, \bullet, /_\bullet)$ is just the projection birack, i.e. the birack whose all left and right translations are the identity. Since involutive distributive biracks satisfy Condition **lri**, we only need to check for the left translations. Indeed, for any $x, y \in X$,

$$x * y = \mathcal{L}_x \phi^{|x|}(y) = \mathcal{L}_x \mathcal{L}_x^{-1}(y) = y.$$

The projection birack corresponds the trivial solution of the YBE, whose YB algebra is just the polynomial algebra. Therefore, we obtain the following proposition.

Proposition 3.4. *The YB algebra of a distributive solution (X, r) of the YBE is a Zhang twist of the polynomial algebra over the set X .*

Acknowledgement. The authors are supported by a grant from NSFC (No. 12371017).

REFERENCES

- [1] R. J. Baxter, *Partition function of the eight-vertex lattice model*, Ann. Physics **70** (1972), 193–228.
- [2] P. Dehornoy, *Set-theoretic solutions of the Yang-Baxter equation, RC-calculus, and Garside germs*, Adv. Math. **282** (2015), 93–127.
- [3] A. Doikou, *Set-theoretic Yang-Baxter equation, braces and Drinfeld twists*, J. Phys A **54** (2021), 415201.
- [4] A. Doikou, A. Smoktunowicz, *From braces to Hecke algebras and quantum groups*, J. Algebra Appl. **22** (2023), 2350179.

- [5] V. G. Drinfel'd, *On some unsolved problems in quantum group theory*, Quantum groups (Leningrad, 1990), 1–8, Lecture Notes in Math., 1510, Springer, Berlin, 1992.
- [6] P. Etingof, T. Schedler, A. Soloviev, *Set-theoretical solutions to the quantum Yang–Baxter equation*, Duke Math. J. **100** (1999), 169–209.
- [7] T. Gateva–Ivanova, *A combinatorial approach to the set-theoretic solutions of the Yang–Baxter equation*, J. Math. Phys. **45** (2004), 3828–3858.
- [8] T. Gateva–Ivanova, *Quadratic algebras, Yang–Baxter equation, and Artin–Schelter regularity*, Adv. Math. **230** (2012), 2152–2175.
- [9] T. Gateva–Ivanova, *Set-theoretic solutions of the Yang–Baxter equation, braces and symmetric groups*, Adv. Math. **338** (2018), 649–701.
- [10] T. Gateva–Ivanova, S. Majid, *Matched pairs approach to set theoretic solutions of the Yang–Baxter equation*, J. Algebra **319** (2008), 1462–1529.
- [11] T. Gateva–Ivanova, S. Majid, *Quantum spaces associated to multipermutation solutions of level two*, Alg. Rep.. Theor. **14** (2011), 341–376.
- [12] T. Gateva–Ivanova, *Set-theoretic solutions of the Yang–Baxter equation, braces and symmetric groups*, Adv. Math. **338** (2018), 649–701.
- [13] T. Gateva–Ivanova, M. Van den Bergh, *Semigroups of I-type*, J. Algebra **206** (1998), 97–112.
- [14] L. Guarnieri, L. Vendramin, *Skew braces and the Yang–Baxter equation*, Math. Comput. **86** (2017), 2519–2534.
- [15] P. Jedlička, A. Pilitowska, and A. Zamojska-Dzienio, *The construction of multipermutation solutions of the Yang–Baxter equation of level 2*, J. Combin. Theory Ser. A **176** (2020), 105295.
- [16] J.-H. Lu, M. Yan, Y.-C. Zhu, *On the set-theoretical Yang–Baxter equation*, Duke Math. J. **104** (2000), 1–18.
- [17] H. Pflugfelder, *Quasigroups and Loops: Introduction*, Heldermann Verlag, Berlin, 1990.
- [18] W. Rump, *A decomposition theorem for square-free unitary solutions of the quantum Yang–Baxter equation*, Adv. Math. **193** (2005), 40–55.
- [19] C. N. Yang, *Some exact results for the many-body problem in one dimension with repulsive delta-function interaction*, Phys. Rev. Lett. **19** (1967), 1312–1315.
- [20] J.J. Zhang, *Twisted graded algebras and equivalences of graded categories*, Proc. London Math. Soc. (3), **72** (1996), 281–311.

XIAOLAN YU

DEPARTMENT OF MATHEMATICS, HANGZHOU NORMAL UNIVERSITY, HANGZHOU, ZHEJIANG
310036, CHINA

Email address: xlyu@hznu.edu.cn

YANFEI ZHANG

DEPARTMENT OF MATHEMATICS, HANGZHOU NORMAL UNIVERSITY, HANGZHOU, ZHEJIANG
310036, CHINA

Email address: zyfeig@163.com