

BI-ISOLATED D.C.E. DEGREES AND Σ_1 INDUCTION

YONG LIU AND CHENG PENG

ABSTRACT. A Turing degree is d.c.e. if it contains a set that is the difference of two c.e. sets. A d.c.e. degree \mathbf{d} is *isolated* if there exists a c.e. degree $\mathbf{a} < \mathbf{d}$ such that every c.e. degree below \mathbf{d} is also below \mathbf{a} ; \mathbf{d} is *upper isolated* if there exists a c.e. degree $\mathbf{a} > \mathbf{d}$ such that every c.e. degree above \mathbf{d} is also above \mathbf{a} ; \mathbf{d} is *bi-isolated* if it is both isolated and upper isolated. In this paper, we prove the existence of bi-isolated d.c.e. degrees in models of IS_1 .

1. INTRODUCTION

The Ershov hierarchy [10, 11, 12] for Δ_2^0 degrees has been extensively studied in the literature. A set A is n -c.e. (*n-computably enumerable*) if there exists a computable function f such that, for all x , $\lim_s f(x, s) = A(x)$, $f(x, 0) = 0$, and $|\{s \mid f(x, s+1) \neq f(x, s)\}| \leq n$. In this way, the 1-c.e. sets coincide with the usual definition of the c.e. sets; 2-c.e. sets coincide with the d.c.e. sets, which is the difference of two c.e. sets. A degree is an n -c.e. degree if it contains an n -c.e. set, and we are mostly interested in comparing the structure of n -c.e. degrees with that of m -c.e. degrees to see whether they differ. Along this line, earlier results show that a maximal degree (that is, an incomplete degree with only $\mathbf{0}'$ above it) exists among the d.c.e. degrees [1, 2], but not among the c.e. degrees [18]. In other words, while the c.e. degrees are dense, the d.c.e. degrees are not. However, various kinds of weak density results do hold (for example, [5, 7, 6]). In particular, Cooper and Yi [7] proved that for any c.e. degree \mathbf{a} and d.c.e. degree \mathbf{d} with $\mathbf{a} < \mathbf{d}$, there exists some d.c.e. degree \mathbf{c} strictly between \mathbf{a} and \mathbf{d} . They also proved that it is necessary for \mathbf{c} to be d.c.e. by introducing the notion of isolated d.c.e. degrees and proving that such a degree exists.

Definition 1.1. A d.c.e. degree \mathbf{d} is *isolated* if there exists a c.e. degree $\mathbf{a} < \mathbf{d}$ such that every c.e. degree below \mathbf{d} is also below \mathbf{a} .

Efremov [8, 9] and Wu [19] then studied similar notions and showed, respectively, the existence of an upper isolated d.c.e. degree and the existence of a bi-isolated d.c.e. degree.

Definition 1.2. A d.c.e. degree \mathbf{d} is *upper isolated* if there exists a c.e. degree $\mathbf{a} > \mathbf{d}$ such that every c.e. degree above \mathbf{d} is also above \mathbf{a} . A d.c.e. degree \mathbf{d} is *bi-isolated* if it is both isolated and upper isolated.

2020 *Mathematics Subject Classification.* 03D28, 03F30, 03H15.

Key words and phrases. bi-isolated d.c.e. degree, reverse recursion theory, inductive strength.

Peng's research was partially funded by the Science and Technology Project of the Hebei Education Department (No. QN2023009) and the National Natural Science Foundation of China (No. 12271264). Yong Liu's research was partially funded by Nanjing Xiaozhuang University (No. 2022NXY39).

Their results were further strengthened by Liu [17], who showed that there exists an isolated maximal d.c.e. degree, which is clearly bi-isolated.

From the perspective of reverse mathematics, one seeks to clarify the proof-theoretic strength required to establish a given theorem. In the context of reverse recursion theory, Paris and Kirby [14] introduced a hierarchy of fragments of Peano arithmetic, which particularly considers different levels of induction schemes and bounding schemes. For our purposes, we consider schemes not beyond $\text{I}\Sigma_1$ and provide only a brief overview; for a comprehensive treatment, see Chong, Li, and Yang [3].

We work in the language of first-order arithmetic $\mathcal{L} = \{0, 1, +, \times, \exp, <\}$, where \exp denotes the exponential function $x \mapsto 2^x$. To study recursion theory, one often works in the base theory $\text{P}^- + \text{I}\Sigma_0 + \text{B}\Sigma_1 + \text{Exp}$, where P^- denotes PA without induction schemes, $\text{I}\Sigma_0$ denotes the induction scheme for Σ_0 formulas, $\text{B}\Sigma_1$ denotes the bounding scheme for Σ_1 formulas, and Exp denotes the axiom $\forall x \exists y (y = \exp(x))$. Over this base theory, Li [15] showed that the existence of a proper d.c.e. degree below $\mathbf{0}'$ is equivalent to $\text{I}\Sigma_1$. To construct a d.c.e. degree with additional properties, one may have to use infinite injury priority methods, which apparently demand more inductive strength. However, it was proved [16] that isolated and upper isolated d.c.e. degrees exist in models of $\text{I}\Sigma_1$, leaving open the question of whether bi-isolated d.c.e. degrees exist in models of $\text{I}\Sigma_1$. We answer this question affirmatively in this paper.

Theorem 1.3. *Bi-isolated d.c.e. degrees exist in every model of $\text{I}\Sigma_1$.*

In light of Li's result [15], our theorem can be rephrased as:

Corollary 1.4. $\text{P}^- + \text{I}\Sigma_0 + \text{B}\Sigma_1 + \text{Exp} \vdash \text{I}\Sigma_1 \leftrightarrow \text{There exists a bi-isolated proper d.c.e. degree below } \mathbf{0}'.$ \square

Wu's proof [19] of the existence of a bi-isolated degree relies on the existence of non-cupping degrees: a noncomputable c.e. degree \mathbf{a} is *non-cupping* (or, *non-cupppable* as in [20]) if there does not exist a c.e. degree $\mathbf{b} < \mathbf{0}'$ such that $\mathbf{a} \vee \mathbf{b} = \mathbf{0}'$. However, the existence of such degrees is equivalent to $\text{I}\Sigma_2$ over the base theory $\text{P}^- + \text{B}\Sigma_2$ [20]; non-cupping degrees do not exist in every model of $\text{I}\Sigma_1$. Therefore, the original path to proving the existence of bi-isolated d.c.e. degrees does not work in models of $\text{I}\Sigma_1$. In this paper, we provide a direct proof of the existence of bi-isolated d.c.e. degrees in models of $\text{I}\Sigma_1$, using only finite injury priority methods.

The key observation is that the strategy for satisfying the isolation requirements is *essentially* similar to the standard Friedberg-Muchnik requirements, and we combine techniques from [16] to prove the existence of bi-isolated d.c.e. degrees in models of $\text{I}\Sigma_1$. In the classical Friedberg-Muchnik construction, each requirement searches for a specific computation to preserve and, once found, immediately acts to ensure its preservation. If such a computation does not exist, the requirement is simply satisfied automatically. Analogously—though with somewhat more complexity—a single isolation requirement (denoted as an N -requirement below) must search for multiple potential computations and aim to preserve at least one of them, often with a delay as we wait for a suitable candidate to appear. If a candidate does not appear, we proceed to construct an alternative functional $W = \Delta^A$ as described below.

Since we are working in models of IS_1 , we can use the finite injury argument in the usual way for the most part, except that the following lemma for $n = 1$ is needed to verify that our construction works in models of IS_1 .

Lemma 1.5 (H. Friedman). *Let $n \geq 1$ and $\mathcal{M} \models \text{P}^- + \text{IS}_n$. Then every bounded Σ_n set is \mathcal{M} -finite, and every partial Σ_n function maps a bounded set to a bounded set.*

2. PROOF OF THEOREM 1.3

We build a d.c.e. set D and a c.e. set $A \leq_T D$ such that the following requirements will be met:

- $N_e : W_e = \Psi_e^D \rightarrow W_e = \Delta^A$,
- $R_e : D = \Phi_e^{W_e} \rightarrow K = \Gamma^{W_e}$, and
- $P_e : D \neq \Theta_e^A$.

Here, $\{\Psi_e\}_{e \in M}$, $\{\Phi_e\}_{e \in M}$, and $\{\Theta_e\}_{e \in M}$ are fixed lists of Turing functionals in our model M . We take A to be the Lachlan set $L(D)$ of D (to be defined below). The set K is taken to be $\{2e + 1 \mid \Phi_e(e) \downarrow\}$, a subset of odd numbers.

These requirements are sufficient. As the Lachlan set of D , $A \leq_T D$ (see below). The P_e -requirements imply $A <_T D$. The N_e -requirements show that D is isolated from below by A . To see $D <_T K$, suppose otherwise that $D \equiv_T K$. Applying the Sacks Splitting Theorem to get $K = D_0 \oplus D_1$, we have

$$K \equiv_T D_0 \oplus D_1 \leq_T A <_T K,$$

a contradiction. Therefore, we have $A <_T D <_T K$. The R_e -requirements show that D is isolated from above by K .

2.1. Preliminaries. We will use a priority tree to organize the construction: The priority tree is $\mathcal{T} = \{0\}^{<M}$, where we assign α to N_e if $|\alpha| = 3e$, R_e if $|\alpha| = 3e + 1$, and P_e if $|\alpha| = 3e + 2$. At stage s , we let the root of \mathcal{T} act. When α acts, it decides whether $\alpha \hat{\ } 0$ acts or we stop the current stage. If α acts at stage s , then we say s is an α -stage. An α -stage s is an α -expansionary stage if $\ell_\alpha(s) > \ell_\alpha(t)$ for each α -expansionary stage $t < s$, where

$$\ell_\alpha(s) = \begin{cases} \max\{y : \forall x \leq y (W_{e,s}(x) = \Psi_{e,s}^{D_s}(x) \downarrow)\}, & \text{if } |\alpha| = 3e, \\ \max\{y : \forall x \leq y (D_s(x) = \Phi_{e,s}^{W_{e,s}}(x) \downarrow)\}, & \text{if } |\alpha| = 3e + 1. \end{cases}$$

For a given enumeration $\{D_s\}_{s \in M}$ of a d.c.e. set D , we define

$$D_s^\sharp(x) = |\{t \leq s \mid D_t(x) \neq D_{t-1}(x)\}|.$$

Note that $D_s(x) = D_s^\sharp(x) \bmod 2$. Let σ be an M -finite sequence. The d.c.e. set D is *unrestorable to σ* at stage s if $\exists x (D_s(x) \neq \sigma(x) \wedge D_s^\sharp(x) = 2)$, and *restorable* otherwise. The *Lachlan set* of D is a c.e. set

$$L(D) = \{\langle x, s \rangle \mid D_{s-1}^\sharp(x) = 0 \wedge D_s^\sharp(x) = 1 \wedge D(x) = 0\} \leq_T D.$$

We write $x^* = \langle x, s \rangle$ with s such that $D_{s-1}^\#(x) = 0$ and $D_s^\#(x) = 1$. An enumeration of $L(D)$ is to enumerate x^* into $L(D)$ at stage s for the least s such that $D_s^\#(x) = 2$.

We will also use some miscellaneous notations for future use: Let σ be an M -finite sequence and let $X \subseteq M$ be a set, viewed as a binary sequence. We write $\sigma \subseteq X$ if σ is an initial segment of X .

We sometimes use the standard notation $\beta[s]$ to indicate that the expression β is evaluated with respect to stage s . Sometimes, we omit $[s]$ when the context is clear.

2.2. P_e -Strategy in Isolation. Let α be the P_e -node. *In this section, we assume that α is not going to be initialized.* This is a standard Friedberg-Muchnik strategy.

At stage s ,

- (P1) If α is *satisfied*, we let $\alpha \hat{=} 0$ act.
- (P2) If the diagonalizing witness w has not been picked, we pick a fresh one.
- (P3) If $\Theta_{e,s}^{A_s}(w) \uparrow$, we let $\alpha \hat{=} 0$ act.
- (P4) If $\Theta_{e,s}^{A_s}(w) \downarrow = 1$, we claim that α is *satisfied*, initialize all nodes below α , and stop the current stage. (We shall prevent w from entering D .)
- (P5) If $\Theta_{e,s}^{A_s}(w) \downarrow = 0$, we enumerate w into D and claim that α is *satisfied*, initialize all nodes below α , and stop the current stage. (We do not let others extract w from D .)

In Items (P4) and (P5), we want to preserve the computation $\Theta_e^A(w) \downarrow$ found at stage s , where the use is $l = \theta_e^A(w)$. Recall that A is the Lachlan set of D . This computation is injured only if some $x^* < l$ is enumerated into A (necessarily, $D_s(x) = 1$), which happens when x is extracted from D . However, we will see that *all extractions are due to an N -node*, and whenever an extraction occurs, all nodes below it are initialized (see Remark 2.4). Under this assumption, we have the following lemma.

Lemma 2.1. *Let α be a P_e -node. Suppose α is not initialized after some stage s_0 . Then the P_e -requirement is satisfied.*

2.3. R_e -Strategy in Isolation. Let α be the R_e -node. *In this section, we assume that α is not going to be initialized.* The node α will build a local functional Γ such that $\Gamma^{W_e}(x) = K(x)$ for all $x \in M$, provided that $\Phi_e^W = D$. To ensure the correctness of each $\Gamma^{W_e}(x) = K(x)$, before defining $\Gamma^{W_e}(x)$, we pick a fresh number $d_{e,x}$, called the *agitator* for x , and wait for an α -expansionary stage s when $d_{e,x} < \ell_\alpha(s)$. Then we define $\Gamma^{W_e}(x) = K_s(x)$ with use $\gamma(x) = \varphi_e^{W_e}(d_{e,x})$. Whenever $0 = \Gamma^{W_e}(x) \neq K(x) = 1$, we enumerate $d_{e,x}$ into D to undefine $\gamma_e^{W_e}(x)$, which will occur at the next α -expansionary stage, if there is one.

The agitator $d_{e,x}$ is either *undefined*, *defined*, *active*, *enumerated*, or *obsolete*, as described below.

R_e -strategy: For all $x \in M$, the agitator $d_{e,x}$ is initially *undefined*. At stage s ,

- (R1) If s is not an α -expansionary stage or $\ell_\alpha(s) < d_{e,x}$ for some $d_{e,x}$ that is defined, we let $\alpha \hat{0}$ act (and skip the instructions below).

Below, we assume s is an α -expansionary stage and $\ell_\alpha(s) \geq d_{e,x}$ for all $d_{e,x}$ that are defined.

- (R2) For each x such that $D_s(d_{e,x}) = 1$, if x is even, we extract $d_{e,x}$; if x is odd, we keep $d_{e,x}$ in D . In both cases, we let $d_{e,y}$ be *undefined* for all $y \geq x$.
- (R3) Let x be the least (if any) such that $\Gamma^{W_{e,s}}(x) \downarrow \neq K_s(x)$. We enumerate $d_{e,x}$ into D so that $D(d_{e,x}) = 1$. We stop the current stage.
- (R4) Let y be the least (which always exists) such that $\Gamma^{W_{e,s}}(y) \uparrow$. For each x with $y \leq x < \ell_\alpha(s)$,
- (R4a) If $K_s(x) = 1$, we define $\Gamma^{W_{e,s}}(x) = 1$ with use $\gamma(x) = 0$. $d_{e,x}$ is claimed to be *obsolete*.
- (R4b) If $K_s(x) = 0$ but $d_{e,x}$ is undefined, we let $d_{e,x}$ be *defined* with a fresh number.
- (R4c) If $K_s(x) = 0$ and $d_{e,x}$ is defined, we define $\Gamma^{W_{e,s}}(x) = 0$ with use $\gamma(x) = \varphi_e^W(d_{e,x})$. We say $d_{e,x}$ is *active*.

Then we let $\alpha \hat{0}$ act.

Note that we are defining the Γ -functional ourselves in an orderly manner, so if $\Gamma^{W_e}(x) \uparrow$ at a stage, then $\Gamma^{W_e}(y) \uparrow$ for all $y > x$ at the same stage. Thus, Items (R4a) and (R4c) are legitimate. We note that for $d_{e,x}$ with odd x , an R -node is responsible for enumerating it into D as in Item (R3) (recall, K is a subset of odd numbers), and never extracts it; an N -node is responsible for extracting it as in Item (C2b) below. For $d_{e,x}$ with even x , an N -node is responsible for enumerating it into D as in Item (N4) below and an R -node is responsible for extracting it as in Item (R2).

If we focus on a particular $d_{e,x}$ with x odd, a typical lifespan is as follows: At first, it is undefined. In Item (R4b), $d_{e,x}$ is *defined* with a fresh number. At the next α -expansionary stage, it is active in Item (R4c). At the following α -expansionary stage, x is enumerated into K and therefore $d_{e,x}$ is enumerated into D in Item (R3) ($\Gamma(x)$ is incorrect). Then, at the next α -expansionary stage, it is undefined in Item (R2) (this is when we undefine $\Gamma(x)$), and then in Item (R4a), it becomes *obsolete*. Thus, $\Gamma^W(x) = K(x)$ is always correct.

Let us preview the conflicts between the R -node α and the N -node β , with $\alpha \subseteq \beta$. Essentially, β wants to protect a *diagonalizing computation* $\sigma \subseteq D$ with $\Psi^\sigma(y) = 0 \neq W(y) = 1$. This computation is injured if α enumerates some $d_{e,x} < |\sigma|$ into D with x odd. The best β can expect is that, for some fixed even number k , those $d_{e,x}$ with $x < k$ are allowed to injure β (as will be seen, this happens only M -finitely many times), but for $x > k$, we would like to have $d_{e,x} > |\sigma|$. To achieve this, β enumerates $d_{e,k}$ into D . There are two cases:

- α is waiting for its expansionary stage. Meanwhile, α poses no threat to other nodes. Then β simply searches for another diagonalizing computation, which will not be threatened by α .
- α has an expansionary stage. Then $\Gamma(k)$ is undefined and, in Item (R2) of the α -strategy, $d_{e,k}$ is extracted and undefined. Then in Item (R4b), $d_{e,k}$ is defined with a fresh number $> |\sigma|$.

We return to the discussion of the R_e -strategy.

Lemma 2.2. *Let α be the R_e -node. Suppose α is not initialized after some stage s_0 . Suppose $\Phi_e^{W_e} = D$. Then we have $\Gamma^{W_e} = K$.*

Proof. Since $\Phi_e^W = D$, there are M -infinitely many α -expansionary stages. Let $x \in M$ be given.

Case 1. Item (R4a) occurs. Then $\Gamma^{W_e}(x) = K(x) = 1$.

Case 2. Item (R4a) does not occur, and for some α -expansionary stage $s_1 > s_0$, we have $\Gamma^{W_e, s_1}(x) \downarrow \neq K_{s_1}(x)$. Then Item (R3) occurs, and so $d_{e,x}$ is enumerated into D at s_1 . Let s_2 be the next α -expansionary stage; we have¹

$$0 = D_{s_1}(d_{e,x}) = \Phi_{e,s_1}^{W_e, s_1}(d_{e,x}) \downarrow \neq \Phi_{e,s_2}^{W_e, s_2}(d_{e,x}) \downarrow = D_{s_2}(d_{e,x}) = 1.$$

Hence, $\Gamma^{W_e}(x) \uparrow$ at stage s_2 . Then Item (R4a) occurs, contradicting the hypothesis.

Case 3. Item (R4a) does not occur and, for all α -expansionary stages s , whenever $\Gamma^{W_e, s}(x) \downarrow$, it equals $K_s(x)$. Let $s_1 > s_0$ be the stage after which $d_{e,x}$ is never undefined. (The existence of such a stage involves the conflicts with N -nodes and will be proved in Lemma 2.12.) Therefore, $d_{e,x}$ is fixed after s_1 . Since $d_{e,x}$ can be enumerated or extracted at most once, there is another stage $s_2 \geq s_1$ after which $D(d_{e,x}) = D_{s_2}(d_{e,x})$. We note that $\Phi_e^{W_e}(d_{e,x}) \downarrow$ implies there exists a stage $s_3 > s_2$ after which $\Phi_e^{W_e}(d_{e,x})$ is never undefined again. Let $s_4 > s_3$ be the next α -expansionary stage. $\Gamma^{W_e}(x)$ is defined with use $\gamma(x) = \varphi_e^{W_e}(d_{e,x})$ at the end of s_4 (as in Item (R4c)). After s_4 , $\Gamma^{W_e}(x)$ is never undefined. Thus, $\Gamma^{W_e}(x) = K(x)$.

Hence, in all cases, $\Gamma^{W_e}(x) = K(x)$. As x is arbitrary, $\Gamma^{W_e} = K$. \square

Together with Lemma 2.8 and Lemma 2.2, we obtain the following lemma:

Lemma 2.3. *For each $e \in M$, R_e is satisfied.* \square

2.4. N_e -Requirement in Isolation. Recall that $A = L(D)$.

The idea for the N_e -node α is to define and maintain $W_e = \Delta^A$ during its expansionary stages. Under the assumption that there are infinitely many α -expansionary stages, there are three possible outcomes:

- (O1) $\Delta^A = W_e$ is total and correct.

¹See Remark 2.5 for a future modification

(O2) If, for some x , $\Delta^A(x) \neq W_e(x)$, then D is restored to some diagonalizing computation σ with $\Psi_e^\sigma(x) \neq W_e(x)$.

(O3) If, for some x , $\Delta^A(x)$ diverges, then $\Psi_e^D(x)$ diverges.

In all cases, the N_e -requirement is satisfied. The main focus for α is to keep $\Delta^A(x)$ correct. To elaborate, suppose we are to define $\Delta^A(x)$ at some α -expansionary stage s . We let $\sigma = D_s \upharpoonright \psi_{e,s}(x)$ and $\tau = A_s \upharpoonright l$ for some fresh number l . In particular, for those z with $\sigma(z) = 1$, we have $z^* < k$. We have $\Psi_e^\sigma(x) = W_{e,s}(x)$, and we define $\Delta^\tau(x) = W_{e,s}(x)$ and $(x; \sigma, \tau)$ as the *diagonalizing pair*. At a later stage $t > s$, we will have three cases:

- (1) D_t becomes unrestorable to σ . Let z be the number such that $D_t(z) \neq \sigma(z)$, with $D_t^\sharp(z) = 2$. Then z^* is enumerated into A_t and therefore $A_t \notin [\tau]$. Therefore, $\Delta^{A_t}(x) \uparrow$. In this case, we cancel the diagonalizing pair and start anew. If this happens infinitely many times, then we have Item (O3).
- (2) If D_t is still restorable to σ and $\Delta^{A_t}(x) \uparrow$ (due to some noise), we define $\Delta^{A_t}(x) = W_{e,t}(x)$ with the same use $|\tau|$ and update the diagonalizing pair to $(x; \sigma, A_t \upharpoonright |\tau|)$ (the previous one is canceled). In other words, if Δ^A is not incorrect yet, we have to ensure the totality of Δ^A , corresponding to Item (O1).
- (3) If D_t is still restorable to σ and $\Delta^{A_t}(x) \downarrow \neq W_{e,t}(x)$, then we claim that σ is a diagonalizing computation, and we restore D_t to σ so that $W_{e,t}(x) \neq \Psi_e^{D_t}(x)$. This corresponds to Item (O2).

The above strategy is referred to as one cycle; the purpose of this cycle is to maintain its version of Δ^A or to produce a diagonalizing computation. We may need multiple cycles, and the k -th cycle is referred to as a $\text{Cycle}(k)$ -module.

Cycle(k)-module at stage s :

- (C1) If s is not an α -expansionary stage, we let α^0 act and skip the instructions below.

Below, we assume s is an α -expansionary stage.

- (C2) Let $x \leq s$ be the least (which always exists) such that $\Delta^{A_s}(x) \uparrow$ or $\Delta^{A_s}(x) \downarrow \neq W_{e,s}(x)$.

(C2a) If $\Delta^{A_s}(x) \uparrow$, for each y with $x \leq y < s$,

(C2a.i) if there is an $(y; \sigma, \tau)$ for some σ and τ has been defined such that D_s is restorable to σ , then we define $\Delta^{A_s}(y) = W_{e,s}(y)$ with the same use as $l = |\tau|$ and define $(y; \sigma, A_s \upharpoonright l)$ as the diagonalizing pair (the old one is canceled).

(C2a.ii) otherwise, we define $\Delta^{A_s}(y) = W_{e,s}(y)$ with a fresh use z and define $(y; D_s \upharpoonright \psi_{e,s}(y), A_s \upharpoonright z)$ as the diagonalizing pair.

Then we let α^0 act.

- (C2b) If $\Delta^{A_s}(x) \downarrow \neq W_{e,s}(x)$, let $(x; \sigma, \tau)$ with $\tau \subseteq A_s$ be the diagonalizing pair. We restore D_s to σ and claim that σ is the *diagonalizing computation* for x , denoted as $\sigma = \text{DiagComp}(k)$.

Note that Item (C2a.ii) happens if the diagonalizing pair is not defined or D is un-restorable to σ . Since we define the Δ -functional in an orderly manner, $\Delta^{A_s}(x) \uparrow$ implies $\Delta^{A_s}(y) \uparrow$ for $y > x$. Once a diagonalizing computation $\sigma = \text{DiagComp}(k)$ is found, it may still have conflicts with R -nodes above α . Recall from the discussion above Lemma 2.2.

2.5. Conflicts between N_e -Strategy and R_i -Strategy. Let α be an N_e -node. There are e many R -nodes above α , namely the R_i -nodes for $i < e$. Let $\sigma = \text{DiagComp}(k)$ be the diagonalizing computation for x found in the $\text{Cycle}(k)$ -module. We restore D to σ immediately. Then, for each $i < e$,

- if $d_{i,x}$ with $x < 2(e-i)$ is enumerated or extracted, then α is initialized. It will be shown that this happens to α only M -finitely many times.
- for $x > 2(e-i)$, we would like to have $d_{i,x} > |\sigma|$ so future enumeration of this number will not injure the diagonalizing computation σ .

To achieve the second point, α itself enumerates $d_{i,2(e-i)}$ for each $i < e$ with $d_{i,2(e-i)}$ defined, into D , in order to undefine $\Gamma_i^{W_e}(d_{i,2(e-i)})$. The diagonalizing computation σ is injured by the enumeration of $d_{i,2(e-i)}$. Once the R_i -node has an expansionary stage, $\Gamma_i^{W_e}(d_{i,2(e-i)})$ becomes undefined, $d_{i,2(e-i)}$ is extracted, and for each $x \geq 2(e-i)$, the R_i -node will pick a fresh number $d_{i,x} > |\sigma|$. Once this happens for all $i < e$, D is restored to σ and it is cleared of any future conflicts with R_i -nodes for $i < e$. Before this is done, α starts the $\text{Cycle}(k+1)$ -module.

To summarize, $\text{Cycle}(k)$ can be in *Phase 1*, *Phase 2*, *accomplished*, or *initialized*. In *Phase 1*, $\text{Cycle}(k)$ searches for a diagonalizing computation. Once it is found, $\text{Cycle}(k)$ enters *Phase 2*. Then, we enumerate all the $d_{i,2(e-i)}$ (if defined) for $i < e$ into D .

Let $d_i^* = d_{i,2(e-i)}$ record the current version of $d_{i,2(e-i)}$ for $i < e$. We define

$$\Upsilon_k(s) = \{i < e \mid D_s(d_i^*) = 1\}.$$

If for some stage s , $\Upsilon_k(s) = \emptyset$, then D is restored to σ and $\text{Cycle}(k)$ is *accomplished*. If $\text{Cycle}(k)$ is initialized, then its version of Δ^A is discarded and Υ_k is canceled.

N_e -strategy: Let α be the N_e -node. Initially, $\text{CycCount}(e) = 0$ and $\text{Cycle}(k)$ is initialized for all $k \in M$. At stage s (with $s^* < s$ the last α -stage),

- (N1) If, for some $d_{i,x}$ with $i < e$ and $x < 2(e-i)$, we have $D_{s^*}(d_{i,x}) \neq D_s(d_{i,x})$, then initialize α and all nodes below α . Stop the current stage.
- (N2) If, for some (least) k , $\Upsilon_k(s) \neq \Upsilon_k(s^*)$, initialize all $\text{Cycle}(j)$ -modules for $j > k$ and all nodes below α .
- (N3) If, for some (least) k , $\Upsilon_k(s) = \emptyset$, then declare $\text{Cycle}(k)$ to be *accomplished*, and let $\alpha \hat{=} 0$ act.

(N4) Otherwise, let k be the least such that $\text{Cycle}(k)$ is *initialized* or in *Phase 1*. Let $\text{Cycle}(k)$ (continued) act. If Item (C2b) occurs in the $\text{Cycle}(k)$ -module, proceed as follows:

- For each $i < e$,
 - If $d_{i,2(e-i)}$ is active, enumerate it into D .
 - If $d_{i,2(e-i)}$ is defined, make it undefined.
 - If $d_{i,2(e-i)}$ is already enumerated, do nothing.
- Define $\Upsilon_k(s)$ as above. $\text{Cycle}(k)$ now enters *Phase 2*.
- Add 1 to $\text{CycCount}(e)$.
- Initialize all nodes below α and stop the current stage.

Remark 2.4. The extraction of $d_{i,2(e-i)}$ that happens in Item (R2) for the R_i -strategy will cause Item (N2) to occur, and all nodes below the N_e -node are therefore initialized. This phenomenon confirms the assumption discussed above Lemma 2.1.

Remark 2.5. In the proof of Lemma 2.2, where α is the R_e -node, we deliberately omit a subtle case to make the main idea clear. In fact, after s_1 , when $d_{e,x}$ is enumerated, an N_j -node with $j > e$ can extract it at some stage $s_{1.5} > s_1$ when Item (C2b) occurs. According to Item (N4), we immediately enumerate $d_{e,2(j-e)}$ into D . We have $d_{e,2(j-e)} < x$; otherwise, the N_j -node would be initialized at stage s_1 when the R_e -node enumerates $d_{e,x}$ into D . In such a case, we have $\Phi_e^W(d_{e,2(j-e)})[s_1] \downarrow \neq \Phi_e^W(d_{e,2(j-e)})[s_2] \downarrow$, which implies $\Gamma^W(2(j-e)) \uparrow$ and hence $\Gamma^W(x) \uparrow$ at stage $s_2 > s_{1.5}$, leading to the same contradiction as in the proof.

Lemma 2.6. For each $e \in M$, $\text{CycCount}(e) \leq 2^e$.

Proof. We prove this by induction on e . For $e = 0$, the N_0 -node has no R -nodes above it. Therefore, once $\text{Cycle}(0)$ enters *Phase 2*, it becomes *accomplished* and no more cycles are needed. Therefore, $\text{CycCount}(0) \leq 1 = 2^0$.

Let α be the N_e -node. If $\text{Cycle}(0)$ never enters *Phase 2*, there will be no other cycles. So we assume $\text{Cycle}(0)$ enters *Phase 2* at some stage s_0 . Note that if $i \in \Upsilon_0(s)$, then $d_{i,2(e-i)}$ is still enumerated at stage s and the R_i -node is still waiting for an expansionary stage and therefore poses no more threat to α . In other words, initially, α has e many R -nodes that have potential conflicts with α . If $|\Upsilon_0(s)| = j$ for some stage s , then there are only $e - j$ many R -nodes that have potential conflicts with α . α now has a situation analogous to that of an N_{e-j} -node with $\text{CycCount}(e-j) \leq 2^{e-j}$. Therefore, there will be at most 2^{e-j} more cycles that can ever enter *Phase 2* after s , while $|\Upsilon_0(s)| = j$ remains unchanged.

Since $|\Upsilon_0(s)|$ is non-increasing, ranging from e down to 0, and whenever $|\Upsilon_0(s)| = 0$, $\text{Cycle}(0)$ becomes *accomplished* and no more cycles are needed. Therefore,

$$\begin{aligned} (1) \quad & \text{CycCount}(e) \leq 1 + \text{CycCount}(0) + \text{CycCount}(1) + \cdots + \text{CycCount}(e-1) \\ (2) \quad & \leq 1 + 2^0 + 2^1 + \cdots + 2^{e-1} = 2^e, \end{aligned}$$

where the first 1 is for $\text{Cycle}(0)$.

This completes the proof. \square

Lemma 2.7. *Let α be the N_e -node. Suppose that α is not initialized after s_0 . Then, the N_e -requirement is satisfied.*

Proof. Suppose that $W_e = \Psi_e^D$. Then α has M -infinitely many expansionary stages. Since $\text{CycCount}(e) \leq 2^e$ (Lemma 2.6), there exists a stage $s_1 > s_0$ after which the status of all cycles of α remains unchanged. If, for some k , we have $\Upsilon_k(s_1) = \emptyset$, let $\sigma = \text{DiagComp}(k)$ be the diagonalizing computation for x . Then $\sigma \subseteq D$ and

$$\Phi_e^\sigma(x) \downarrow \neq W_e(x),$$

contradicting the hypothesis that $W_e = \Psi_e^D$. Therefore, no cycles are accomplished. Let $\text{Cycle}(k)$ be the (unique) cycle which is in *Phase 1* at stage s_1 . Item (C2b) does not occur in the $\text{Cycle}(k)$ -module. Fix an arbitrary $x \in M$. Recall that Item (C2a.ii) occurs if the diagonalizing pair is not defined (which happens only once), or if D is unrestorable to σ . Therefore, if Item (C2a.ii) happens M -infinitely often, then D is unrestorable to σ M -infinitely often. This necessarily implies $\Psi_e^D(x) \uparrow$, contradicting the hypothesis that $W_e = \Psi_e^D$. Therefore, there exists a stage $s_2 > s_1$ after which Item (C2a.ii) no longer happens, so $\Delta^A(x) \downarrow$.

Hence, $\Delta^A = W_e$, and this completes the proof. \square

2.6. Construction and Verification. The construction proceeds stage by stage as follows:

At stage s , let the root of the priority tree \mathcal{T} act. Suppose α is to act.

- If $|\alpha| = s$, stop the current stage.
- If α is an N_e -node, let α act according to the N_e -strategy.
- If α is an R_e -node, let α act according to the R_e -strategy.
- If α is a P_e -node, let α act according to the P_e -strategy.

It remains to prove the following:

Lemma 2.8. *For each α , there exists some stage s_0 after which α is not initialized.*

In models of full arithmetic, the lemma is straightforward because all relevant inductions are available and the number of initializations can be shown to be finite. However, when working within $I\Sigma_1$, we must provide a concrete, uniform bound on the number of times each node α can be initialized in order to carry out the argument within the weaker

induction available. Therefore, to establish the lemma in $I\Sigma_1$, it suffices to effectively bound the number of initializations for each α .

Definition 2.9. Let $f(e)$, $g(e)$, and $h(e)$ denote $1 +$ the number of times initialization occurs to N_e , R_e , and P_e , respectively.

Estimated in a coarse manner, we have the following recurrence relations:

Lemma 2.10. *For all $e \in M$,*

- (1) $f(0) = 1$
- (2) $g(e) \leq e(\text{CycCount}(e) + 1)f(e)$
- (3) $h(e) = g(e)$
- (4) $f(e + 1) \leq 2h(e) + e + 1$

Proof. (1) The N_0 -node is never initialized, so $f(0) = 1$.

(2) While the N_e -node is not initialized, there are at most $\text{CycCount}(e)$ cycles, and for each cycle, Item (N2) happens at most e times. Therefore, $g(e) \leq e(\text{CycCount}(e) + 1)f(e)$.

(3) The R_e -node does not initialize nodes below it. In particular, the P_e -node is not initialized by the R_e -node, so $h(e) = g(e)$.

(4) While the P_e -node is not initialized, Item (P4) or Item (P5) happens at most once. We also have to count the number of times Item (N1) occurs for the N_{e+1} -node. Most cases are absorbed in the initializations of N_e , R_e , and P_e . The other cases are caused by Item (R3). There are at most $e + 1$ odd numbers $< 2(e + 1)$. Thus, Item (R3) happens at most $e + 1$ times *globally*, as K is global. Therefore $f(e + 1) \leq 2h(e) + e + 1$.

□

Lemma 2.11. *For all $e \in M$, $f(e) \leq 2^{e^2}$ and $h(e) = g(e) \leq e(2^e + 1)2^{e^2}$.*

Proof. We only need to prove the first part, and this is proved by induction on e . $f(0) \leq 1 = 2^{0^2}$.

From (2) and (4) in Lemma 2.10, we have

$$f(e + 1) \leq 2e(\text{CycCount}(e) + 1)f(e) + e + 1.$$

From the induction hypothesis $f(e) \leq 2^e$ and the basic inequality $e < e + 1 \leq 2^e \leq 2^{e^2}$, we have

$$\begin{aligned}
 f(e+1) &\leq 2e(2^e + 1)2^{e^2} + e + 1 \\
 &\leq (e2^{e+1} + 2e)2^{e^2} + 2^{e^2} \\
 &= (e2^{e+1} + e + e + 1)2^{e^2} \\
 &\leq (e2^{e+1} + 2^e + 2^e)2^{e^2} \\
 &\leq (e+1)2^{e+1}2^{e^2} \\
 &\leq 2^{2e+1}2^{e^2} = 2^{(e+1)^2}.
 \end{aligned}$$

□

Proof of Lemma 2.8. Let $\alpha \in \mathcal{T}$. Let

$$A_\alpha = \{i \mid \alpha \text{ is initialized at least } i \text{ times}\},$$

By Lemma 2.10, A_α is bounded. Since it is also Σ_1 , A_α is M -finite (Lemma 1.5). Let $f : A_\alpha \rightarrow M$ map i to the stage s when α is initialized for the i -th time; then f is Σ_1 -definable, and $\mathbf{B}\Sigma_1$ implies that the range of f is bounded by some stage s_0 . Then, for all stages $s > s_0$, α is not initialized. □

Lemma 2.12. *For each $e, x \in M$, there exists a stage s after which $d_{e,x}$ is not undefined.*

Proof. We remark that $d_{e,x}$ can be undefined only in Item (R2). Therefore, we only need to consider which node is responsible for enumerating some $d_{e,y}$ with $y \leq x$ into D .

Suppose x is even, so $x = 2i$ for some i . Thus, the N_{e+i} -node is responsible for enumerating $d_{e,x}$ into D . Let s_0 be the stage after which both R_e - and N_{e+i} -nodes are not initialized, by Lemma 2.8. By the choice of s_0 and Item (N1), no $d_{e,y}$ with $y < x$ is enumerated into or extracted from D after s_0 . After s_0 , only the N_{e+i} -node can possibly enumerate $d_{e,x}$ into D for the last time. Therefore, there exists a stage $s_1 > s_0$ after which $d_{e,x}$ is not undefined.

Suppose x is odd, so $x = 2i + 1$ for some i . Let s_0 be the stage after which R_e is not initialized and $d_{e,2i}$ is always defined. Since x is odd, $d_{e,x}$ is enumerated into D only when x enters K , which happens at most once. Therefore, there exists a stage $s_1 > s_0$ after which $d_{e,x}$ is not undefined.

This completes the proof. □

We remark that our bi-isolated set D can be made low, since the lowness requirements are not more difficult to satisfy than the N_e -requirements, as can be seen in [16].

3. OPEN PROBLEMS

In the literature [4], there is an alternative definition of isolated degrees: a d.c.e. degree \mathbf{d} is *isolated* if there exists a c.e. degree $\mathbf{a} < \mathbf{d}$ and there is no other c.e. degree strictly between \mathbf{a} and \mathbf{d} . Note that the requirement that \mathbf{d} be a *proper* d.c.e. degree

is not included in this definition or in Definition 1.1. Therefore, the two definitions are equivalent because of the Sacks Density Theorem [18]. While Groszek, Mytilinaios, and Slaman [13] showed that BS_2 suffices to prove the Sacks Density Theorem, the exact strength remains unclear:

Question 3.1. *Does the Sacks Density Theorem fail in some model of IS_1 ?*

Depending on the answer to this question, the two versions of isolated d.c.e. degrees may differ from each other in some model of IS_1 .

REFERENCES

- [1] S. Barry Cooper, Leo Harrington, Alistair H. Lachlan, Steffen Lempp, and Robert I. Soare. The d.r.e. degrees are not dense. *Annals of Pure and Applied Logic*, 55(2):125–151, 1991.
- [2] S. Barry Cooper, Leo Harrington, Alistair H. Lachlan, Steffen Lempp, and Robert I. Soare. Corrigendum to “the d.r.e. degrees are not dense” [ann. pure appl. logic 55 (1991) 125–151]. *Annals of Pure and Applied Logic*, 168(12):2164–2165, 2017.
- [3] Chi Tat Chong, Wei Li, and Yue Yang. Nonstandard models in recursion theory and reverse mathematics. *Bulletin of Symbolic Logic*, 20:170–200, 2014.
- [4] S. Barry Cooper. *Degrees of Unsolvability*. Thesis, 1971.
- [5] S. Barry Cooper. The density of the low n -r.e. degrees. *Archive for Mathematical Logic*, 31(1):19–24, 1991.
- [6] S. Barry Cooper, Steffen Lempp, and Philip Watson. Weak density and cupping in the d-r.e. degrees. *Israel Journal of Mathematics*, 67(2):137–152, 1989.
- [7] S. Barry Cooper and Xiaoding Yi. Isolated d.r.e. degrees. *Preprint series, University of Leeds, Department of Pure Mathematics*, 1995.
- [8] A. A. Efremov. d-r. e. degrees that are isolated from above. ii. *Izvestiya Vysshikh Uchebnykh Zavedenii. Matematika*, (7):18–25, 1998.
- [9] A. A. Efremov. d-r.e. degrees that are isolated from above. i. *Izvestiya Vysshikh Uchebnykh Zavedenii. Matematika*, (2):20–28, 1998.
- [10] Yu. L. Ershov. On a hierarchy of sets. i. *Algebra and Logic*, 7:25–43, 1968.
- [11] Yu. L. Ershov. On a hierarchy of sets, ii. *Algebra and Logic*, 7(4):212–232, 1968.
- [12] Yu. L. Ershov. On a hierarchy of sets. iii. *Algebra and Logic*, 9(1):20–31, 1970.
- [13] Marcia J. Groszek, Michael E. Mytilinaios, and Theodore A. Slaman. The sacks density theorem and Σ_2 -bounding. *Journal of Symbolic Logic*, 61(2):450–467, 1996.
- [14] L. A. S. Kirby and J. B. Paris. Initial segments of models of peano’s axioms. *Set Theory and Hierarchy Theory V*, pages 211–226. Springer Berlin Heidelberg.
- [15] Wei Li. Δ_2 degrees without Σ_1 induction. *Israel Journal of Mathematics*, 201:989–1012, 2014.
- [16] Yiqun Liu, Yong Liu, and Cheng Peng. Isolated d.c.e. degrees and Σ_1 induction. *Annals of Pure and Applied Logic*, 177(3):103678, 2026.
- [17] Yong Liu. Isolated maximal d.r.e. degrees. *Annals of Pure and Applied Logic*, 170(4):515–538, 2019.
- [18] Gerald E. Sacks. The recursively enumerable degrees are dense. *Annals of Mathematics*, 80(2):300–312, 1964.
- [19] Guohua Wu. Bi-isolation in the d.c.e. degrees. *Journal of Symbolic Logic*, 69(2):409–420, 2004.
- [20] Yue Yang. Σ_2 Induction and cuppable degrees, page 215–228. De Gruyter, Berlin, Boston, 1999.

(Yong Liu) SCHOOL OF INFORMATION ENGINEERING, NANJING XIAOZHUANG UNIVERSITY, CHINA

Email address: liuyong@njxzc.edu.cn

(Peng) INSTITUTE OF MATHEMATICS, HEBEI UNIVERSITY OF TECHNOLOGY, CHINA

Email address: pengcheng@hebut.edu.cn