

Generalized Beth–Uhlenbeck entropy formula from the Φ –derivable approach

D. Blaschke,^{1,2,3,*} G. Röpke,^{4,†} and G. Baym^{5,‡}

¹*Institute of Theoretical Physics, University of Wrocław, Max Born place 9, 50-204 Wrocław, Poland*

²*Helmholtz-Zentrum Dresden-Rossendorf (HZDR), Bautzner Landstrasse 400, 01328 Dresden, Germany*

³*Center for Advanced Systems Understanding (CASUS), Untermarkt 20, 02826 Görlitz, Germany*

⁴*Institut für Physik, Universität Rostock, Albert-Einstein-Strasse 23-24, 18059 Rostock, Germany*

⁵*Department of Physics, University of Illinois, 1110 W. Green Street, Urbana, IL 61801, USA*

(Dated: December 4, 2025)

We derive a generalized Beth-Uhlenbeck formula for the entropy of a dense fermion system with strong two-particle correlations, including scattering states and bound states. We work within the Φ –derivable approach to the thermodynamic potential. The formula takes the form of an energy-momentum integral over a statistical distribution function times a unique spectral density. In the near mass-shell limit, the spectral density reduces, contrary to naïve expectations, not to a Lorentzian but rather to a "squared Lorentzian" shape. The relation of the Beth-Uhlenbeck formula to the Φ –derivable approach is exact at the two-loop level for Φ . The formalism we develop, which extends the Beth-Uhlenbeck approach beyond the low-density limit, includes Mott dissociation of bound states, in accordance with Levinson's theorem, and the self-consistent back reaction of correlations in the fermion propagation. We discuss applications to further systems, such as quark matter and nuclear matter.

I. INTRODUCTION

Fermionic systems that can form bound states are very common in physics. Well known examples include plasmas, in particular the hydrogen plasma and the electron-hole plasma in semiconductors, ruled by the Coulomb interaction. Other examples are nuclear matter and the quark-gluon plasma, where the constituents interact via the strong QCD interaction. All these systems share in common the formation of bound states and their dissolution at high densities. Although treating the correlations in such interacting quantum systems is very challenging, they can be treated by similar approaches within the framework of quantum statistical mechanics.

Various methods have been worked out in quantum statistics to treat many-particle systems, such as ionic plasmas and warm dense matter, the electron-hole plasma in semiconductors, the quark-gluon plasma, nuclear matter, etc. In particular, the Matsubara Green's function method and the use of Feynman diagrams provide the tools to investigate these interacting many-particle systems. Partial summations can be performed to introduce quasiparticles and self-energies, screening of interactions, vertex corrections, formation of bound states, and other collective excitations, leading to an improved perturbative approach and successful approximation schemes. See the book of Kadanoff and Baym [1] for the quantum statistical approach to the Coulomb many-particle problem via Green's functions methods in equilibrium and nonequilibrium.

As pointed out by Baym and Kadanoff [2, 3] and others, the selection of diagrams that define special approximations must be taken with care. One has to obey the conservation laws, beyond simply at the vertices in diagrams, obey exact sum rules and Ward identities, and avoid double counting of contributions. As quasiparticles become shifted and broadened, one has to take into account effects that contribute in the same order of magnitude because of possible compensation effects. While of high interest, it is a big challenge in linear response theory and kinetic theory to describe the evolution of many-particle systems while guaranteeing conservation of particle number, energy, and momentum, etc. The Φ -functional approach provides a prescription [3] to construct such approximations. This concept of conserving approximations has been further elaborated by, e.g., Riedel [4], Vanderheyden and Baym [5], and Blaizot et al. [6].

Another earlier approach to the thermodynamic properties of non-relativistic plasmas was worked out using the Beth-Uhlenbeck equation for the second virial coefficient [7, 8]. To treat higher densities, this low-density expansion requires improvements, which can be done by introducing quasiparticles. Zimmermann and Stolz [9] showed then that the contribution of the scattering states must be modified, and a $\sin^2 \delta$ – term appears in the Beth-Uhlenbeck formula to compensate the mean-field contributions, which are already taken into account in the quasiparticle shift. A detailed discussion of this generalized Beth-Uhlenbeck formula is found in Schmidt et al. [10].

Our aim in this paper is to show the equivalence of the Φ -functional and the generalized Beth-Uhlenbeck approaches. We use a simple model of interaction for which we construct the spectral functions, and with the help of generalized optical theorems we

*Electronic address: blaschke@ift.uni.wroc.pl

†Electronic address: roepke@uni-rostock.de

‡Electronic address: gbaym@illinois.edu

show that the appearance of the $\sin^2 \delta$ term corresponds to the squared Breit-Wigner profile in the spectral function.

We follow the approach of Vanderheyden and Baym [5] who start from relativistic field theories described by the QED Lagrangian. Other systems where these methods are of interest include Fermi liquids such as ^3He , as considered by Riedel [4], and the quark-gluon plasma, see Blaizot et al. [6]. Our approach, demonstrated in a simple case, can be generalized to more complex systems.

We mention here related examples that show the necessity of using consistent approximations in plasma physics. References [11, 12], following the Green's function approach, derived an effective in-medium Schrödinger equation for plasmas, showing that when the Fock shift of single-particle energies are considered, one also has to take Pauli blocking of the interaction into account. When introducing screening for the interaction potential, one has also to include the Montroll-Ward self-energy terms (the rainbow diagrams); and when improving the two-particle propagators in the polarisation loop by medium effects, one must also improve the vertex [13]. In addition to plasma physics, associated work has been carried out for the NJL model of chiral quark-meson model [14–19]. A formulation of the back-reaction problem within the path integral approach has been given for the NJL model [20] where it was shown how the form of a Φ -derivable thermodynamical potential could be obtained for quark matter with composite mesons and how the diagrammatic expansion for the Φ -functional is related to the $1/N_c$ expansion of the NJL model.

II. Φ -DERIVABLE APPROACH TO THE ELECTRON-POSITRON PLASMA IN QED IN THE TWO-LOOP APPROXIMATION

A fundamental approach to the theory of plasmas starts from quantum electrodynamics (QED), with fermions coupled to the electromagnetic field, the approach of Vanderheyden and Baym [5] in discussing self-consistent approximations in relativistic plasmas. Methods similar to those proposed in that paper can be applied to other plasmas, e.g. the electron-proton plasma.

A. Thermodynamical potential for the electron-positron-photon system with a “sunset” Φ -functional

Applying the Φ -derivable formulation [3] to QED thermodynamics, we closely follow Ref. [5] where the thermodynamic potential reads

$$\beta\Omega = \Phi[G, D] - \text{Tr}\Sigma G + \text{Tr} \ln(-\gamma_0 G) + \frac{1}{2}\text{Tr}\Pi D - \frac{1}{2}\text{Tr} \ln(-D), \quad (1)$$

where G is the electron Green's function with self-energy Σ , and D is the photon Green's function with self-energy or polarization operator Π ; γ_0 is the Dirac matrix. The functional $\Phi[G, D]$ is the sum of all skeleton diagrams of the thermodynamic potential. This ansatz guarantees that the stationary condition for the thermodynamic potential against variations of the Green's functions,

$$\frac{\delta\Omega}{\delta G} = \frac{\delta\Omega}{\delta D} = 0, \quad (2)$$

is equivalent to the statement that the propagators G for electrons and D for photons fulfill the Dyson equations

$$\begin{aligned} G^{-1} &= G_0^{-1} - \Sigma, \\ D^{-1} &= D_0^{-1} - \Pi, \end{aligned} \quad (3)$$

where G_0 and D_0 are the bare electron and photon propagators, and Σ and Π are calculated as functional derivatives,

$$\Sigma = \frac{\delta\Phi}{\delta G}, \quad \Pi = 2 \frac{\delta\Phi}{\delta D}. \quad (4)$$

We assume the Coulomb gauge.

Using the spectral representations for the Green's functions, Eq. (3),

$$G(\omega_n, \mathbf{p}) = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \frac{A(\omega, \mathbf{p})}{\omega_n - \omega}, \quad (5)$$

$$D_T(\omega_n, \mathbf{q}) = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \frac{B_T(\omega, \mathbf{q})}{\omega_n - \omega}, \quad (6)$$

$$D_L(\omega_n, \mathbf{q}) = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \frac{B_L(\omega, \mathbf{q})}{\omega_n - \omega} + \frac{1}{\mathbf{q}^2}, \quad (7)$$

with the fermionic Matsubara frequencies $\omega_n = (2n + 1)\pi iT + \mu$ for electrons, where μ is the electron chemical potential, and bosonic Matsubara frequencies $\omega_n = 2\pi inT$ for photons.

The frequency sums in (1) can be evaluated by contour integration and yield

$$\Omega = T\Phi[G, D] + \sum_{\mathbf{p}} \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} f(\omega) \text{Tr}_{\text{Dirac}} \text{Im} [\Sigma G + \ln(-\gamma_0 G^{-1})] + \sum_{\mathbf{q}} \oint \frac{d\omega}{2\pi} n(\omega) \sum_{l=L,T} g_l \text{Im} [\Pi_l D_l + \ln(-D_l^{-1})], \quad (8)$$

with the statistical factors $f(\omega) = \{\exp[\beta(\omega - \mu)] + 1\}^{-1}$ for fermions and $n(\omega) = \{\exp(\beta\omega) - 1\}^{-1}$ for bosons. (The slash through the boson integral indicates the principal value.) The degeneracy factors are $g_L = 1$ for longitudinal modes and $g_T = 2$ for transverse modes.

From this expression for the thermodynamic potential all other equations of state can be obtained by differentiation. The entropy S and particle number N are obtained as derivatives with respect to the temperature and the chemical potential:

$$S = -\left. \frac{\partial \Omega}{\partial T} \right|_{\mu}, \quad N = -\left. \frac{\partial \Omega}{\partial \mu} \right|_T. \quad (9)$$

Because of the stationarity (2), we can ignore the spectral densities of the propagators when differentiating $\Phi[G, D]$, and need only differentiate the statistical factors. Let us consider the entropy as the derivative with respect to the temperature,

$$S = -\left. \frac{\partial \Omega}{\partial T} \right|_{\mu, V, A, B} = S_f + S_b + S', \quad (10)$$

with the fermionic and bosonic contributions,

$$S_f \equiv -\sum_{\mathbf{p}} \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \frac{\partial f(\omega)}{\partial T} \text{Tr}_{\text{Dirac}} [\text{Im} \Sigma \text{Re} G + \text{Im} \ln(-\gamma_0 G^{-1})], \quad (11)$$

$$S_b \equiv -\sum_{\mathbf{q}, l} g_l \oint \frac{d\omega}{2\pi} \frac{\partial n(\omega)}{\partial T} [\text{Im} \Pi_l \text{Re} D_l + \text{Im} \ln(-D_l^{-1})], \quad (12)$$

respectively. The remaining term,

$$S' \equiv -\left. \frac{\partial(T\Phi)}{\partial T} \right|_{A, B} - \sum_{\mathbf{p}} \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \frac{\partial f(\omega)}{\partial T} \text{Tr}_{\text{Dirac}} [\text{Im} G \text{Re} \Sigma] - \sum_{\mathbf{q}, l} g_l \oint \frac{d\omega}{2\pi} \frac{\partial n(\omega)}{\partial T} \text{Im} D_l \text{Re} \Pi_l, \quad (13)$$

has the important property that it vanishes in the two-loop approximation for Φ , as illustrated diagrammatically in Fig. 1 for fermions (solid lines) interacting via the exchange of a boson (dashed line). (In Ref. [5] the same diagram is denoted as “one-loop.”) This vanishing has been shown for QED in Ref. [5] and for QCD in Ref. [6]. It is based on the cancellation of the derivative of the Φ -functional and the terms with the trace over the product of the imaginary part of the propagator (spectral density) of a particle and the real part of its selfenergy, see Eq. (13). The cancellation holds also for both the entropy and particle density in the two-loop approximation, $S' = N' = 0$. We demonstrate this cancellation for the simple case of a scalar field ϕ with a ϕ^3 interaction vertex in Appendix A.




FIG. 1: The two-loop functional Φ , where the solid line is the electron propagator and the dashed line the photon propagator.

Figure 2 shows the corresponding self-energy diagrams obtained as variational derivatives, Eqs. (4).



FIG. 2: Self-energy diagrams derived from the Φ in Fig. 1.

In the following we focus on the bosonic entropy, S_b , and rewrite Eq. (12) using the identity

$$\frac{\partial n(\omega)}{\partial T} = -\frac{\partial \sigma_n(\omega)}{\partial \omega}, \quad (14)$$

where $\sigma_n(\omega) \equiv -n(\omega) \ln n(\omega) + (1+n(\omega)) \ln(1+n(\omega))$ is the contribution to the entropy of a photon mode of frequency ω . After partial integration over ω one obtains

$$S_b = \sum_{\mathbf{q}} \int_0^\infty \frac{d\omega}{2\pi} \sigma_n(\omega) B_s(\omega, \mathbf{q}), \quad (15)$$

where

$$B_s(\omega, \mathbf{q}) = \sum_{l=L,T} g_l B_{s,l}(\omega, \mathbf{q}), \quad (16)$$

with

$$B_{s,l}(\omega, \mathbf{q}) = \frac{\partial}{\partial \omega} [\text{Re} D_l L_l - 2 \text{Im} \ln(-D_l^{-1})], \quad (17)$$

and $L_l(\omega, \mathbf{q}) \equiv -2 \text{Im} \Pi_l(\omega + i0^+, \mathbf{q})$ the imaginary part of the photon self-energy.

To interpret the contents of this formula, we expand the inverse propagator of the Dyson equation (3) in the vicinity of the plasmon poles ω_l which are defined by its zeroes, $\text{Re} D_l^{-1}(\omega = \omega_l, \mathbf{q}) = 0$, to obtain $\text{Re} D_l^{-1} \simeq (\omega - \omega_l)/Z_l$ and $\text{Im} D_l^{-1} = L_l/2$, where $Z_l = d \text{Re} D / d\omega|_{\omega_l}$. In this approximation the propagator is

$$D_l = Z_l \frac{(\omega - \omega_l) + i Z_l L_l / 2}{(\omega - \omega_l)^2 + (Z_l L_l / 2)^2}. \quad (18)$$

The frequency derivatives in Eq. (17) can now be evaluated to give

$$\begin{aligned} \frac{\partial}{\partial \omega_q} \text{Re} D_l L_l &= \frac{Z_l L_l}{(\omega - \omega_l)^2 + (Z_l L_l / 2)^2} - \frac{2(\omega - \omega_l)^2 L_l}{[(\omega - \omega_l)^2 + (Z_l L_l / 2)^2]^2}, \\ \frac{\partial}{\partial \omega_q} \text{Im} \ln(-D_l^{-1}) &= \frac{Z_l L_l / 2}{(\omega - \omega_l)^2 + (Z_l L_l / 2)^2}. \end{aligned}$$

As discussed in [5], the entropy spectral function (17) in the vicinity of the longitudinal and transverse plasmon modes of frequencies $\omega_L(q)$ and $\omega_T(q)$, takes the form of a simple squared Lorentzian

$$B_{s,l}(\omega_q, \mathbf{q}) \simeq \frac{4(Z_l L_l / 2)^3}{[(\omega_q - \omega_l(q))^2 + (Z_l L_l / 2)^2]^2}, \quad (19)$$

When compared with the spectral densities of photons B_l , which have a Lorentzian form close to the poles, $B_l \simeq (Z_l L_l) / [(\omega_q - \omega_l(q))^2 + (Z_l L_l / 2)^2]$, we see that the $B_{s,l}$ have a stronger peak and smaller wings.

The entropy formula (15) with the spectral function (17) holds for the rather wide class of two-loop diagrams for the Φ -functional (as, e.g., given in Fig. 1) since for these diagrams the cancellation $S' = 0$ of Eq. (13) applies, as it has been stated in Refs. [5, 6]. In the next section we show that the equation of state for the entropy (15) in this approximation takes the form of a generalized Beth-Uhlenbeck equation.

B. Generalized Beth-Uhlenbeck formula

In this section, we suppress the mode index $l = L, T$ for brevity and denote the frequency derivative by a prime. We rewrite the second term on the right side of Eq. (17) as

$$[\text{Im} \ln(-D^{-1})]' = -\text{Im} (D \Pi') = \underbrace{(\text{Re} D' \text{Im} \Pi - \text{Im} D \text{Re} \Pi')}_{2 \text{Im}(D \text{Im} \Pi D^{*'} \text{Im} \Pi)} - \underbrace{(\text{Im} \Pi \text{Re} D' + \text{Re} D \text{Im} \Pi')}_{{(\text{Im} \Pi \text{Re} D)'}}, \quad (20)$$

where we prove the relation

$$\text{Re} D' \text{Im} \Pi - \text{Im} D \text{Re} \Pi' = 2 \text{Im}(D \text{Im} \Pi D^{*'} \text{Im} \Pi), \quad (21)$$

which can be called "derivative optical theorem", in Appendix B. The right side of Eqs. (21) is similar to the final result for the thermodynamic potential derived by Dashen, Ma and Bernstein [21] in their Eq. (4.32). It can be viewed as its generalisation from the low-density limit to the self-consistent quasiparticle level which includes the effects of phase space occupation via Π_I . For its evaluation in the case of deuterons in nuclear matter, see Ref. [10].

We now introduce the phase function δ of the polar representation of the propagators $D = |D| \exp(i\delta)$ to find the relationships

$$D \text{Im} \Pi = \sin \delta e^{i\delta}, \quad D^{*'} \text{Im} \Pi = -i\delta' \sin \delta e^{-i\delta}, \quad (22)$$

which together with Eq. (21) allow to rewrite Eq. (17) in terms of the phase shift δ as

$$- [\text{Re} D \text{Im} \Pi + \text{Im} \ln(-D^{-1})]' = 2\delta' \sin^2 \delta. \quad (23)$$

Inserting (23) into (17) we arrive at the main result of this paper, the generalized Beth-Uhlenbeck equation of state for the entropy (15)

$$S_b = 2V \int \frac{d^3 q}{(2\pi)^3} \int_0^\infty \frac{d\omega}{\pi} \sigma_b(\omega) \sin^2 \delta(\omega, \mathbf{q}) \frac{\partial \delta(\omega, \mathbf{q})}{\partial \omega}. \quad (24)$$

This result has interesting consequences for the phenomenology of fermion-boson systems. Similar to the standard Beth-Uhlenbeck equation of state, in the two-loop approximation for Φ one obtains a good quasiparticle picture for the statistical system as a superposition of elementary (e.g., quarks) as well as composite (e.g., hadronic) degrees of freedom which in general are all off-shell, including the elementary ones, as a result of the self-consistency of the approach. Ideal quasiparticles with on-mass-shell behaviour are obtained when the corresponding phase function degenerates to a step function at the mass of the particle.

The spectral function (23) of the generalized Beth-Uhlenbeck formula (24) has the shape of the "squared Lorentzian" (19) when for the phase shift the resonance shape with the plasmon pole at $\omega = \omega_R$ and the width $\Gamma = Z_R \text{Im} \Pi$ is assumed

$$\delta(\omega) = -\arctan \left[\frac{\Gamma}{\omega - \omega_R} \right]. \quad (25)$$

This form corresponds to a Breit-Wigner (or Lorentz-) profile for the derivative of the phase shift

$$\frac{\partial \delta(\omega)}{\partial \omega} = -\frac{\Gamma}{(\omega - \omega_R)^2 + \Gamma^2}, \quad (26)$$

then the spectral weight in the generalized Beth-Uhlenbeck formula for the entropy does *not* have the Breit-Wigner shape expected from the standard Beth-Uhlenbeck approach, but rather it is a "squared Lorentzian" profile

$$\sin^2 \delta(\omega) \frac{\partial \delta(\omega)}{\partial \omega} = \frac{\Gamma^3}{[(\omega - \omega_R)^2 + \Gamma^2]^2}. \quad (27)$$

This result, which is obtained here for the simplifying ansatz (26) for the phase shift, has been found before within the Φ -derivable approach in Ref. [5] for QED (see Eq. (19) of the previous subsection) and in Ref. [22] for the entropy of the Lee model. Similar results were obtained also in other approaches and in a different context, for instance within the kinetic theory approach to radiation in non-equilibrium plasmas [23]. For applications in thermodynamic systems, the effect of considering the second virial coefficient of quasiparticles in the medium rather than free on-shell particles is a sharpening of the spectral distribution describing the resonant scattering (correlation).

III. DISCUSSION

The close relation between the conserving approximations in terms of Φ , and the consistent generalization of the Beth-Uhlenbeck formula is important because we need the introduction of quasiparticles to cover a wide range of densities – as known from mean-field approximations in Coulomb systems, relativistic mean-field approximations (RMF) in nuclear systems, and self-consistent (gap) equations in the quark-gluon plasma. Such a treatment of correlations, for instance in the Beth-Uhlenbeck

formula, must be performed carefully to avoid double counting. We have presented here the simplest approximation for the self-energy contribution to the quasiparticle picture, the self-consistent Hartree-Fock approximation. If we consider higher approximations for the self-energy to determine the quasiparticle contribution in the generalized Beth-Uhlenbeck formula, the corresponding corrections have to be performed also for the two-particle contribution of this formula.

The generalized Beth-Uhlenbeck approach has the advantage over the standard one in that it is applicable also beyond the low-density limit, when due to phase space occupation, the spectral properties of correlations described by scattering phase shifts are strongly modified compared to the case of interactions in free space. The quasiparticle picture is valid also in the high-density limit, where strongly correlated bound states are suppressed by Pauli blocking – the Mott effect. In the strongly degenerate limit, the interacting fermion system is described as a Landau Fermi-liquid. Importantly, the self-consistent mean-field (Hartree-Fock) approach discussed here works not only at low densities but is also applicable as a self-consistently generalized Beth-Uhlenbeck formula at high densities.

While the subdivision of the density into an ‘free’ quasiparticle part and the remaining correlated part is a formal issue, and in the final bookkeeping only the total density is relevant, we have to avoid double counting. The discussion of free and bound density, ionization degree, etc. in plasma physics is an old problem, often discussed with ill-defined concepts. After separation of the single-quasiparticle term in the spectral function, the correlated contribution contains bound state contributions as well as contributions from the continuum.

This concept was worked out for the proton-neutron system in Ref. [10], where the self-consistent two-particle contribution contains a possible bound state (deuteron) and the contribution of the continuum of scattering states. Here empirical RMF expressions were used instead of the self-consistent mean-field shift of the quasiparticle energy; the Mott effect was found, and the contribution of scattering states was calculated using a separable potential approach. One can introduce a generalized phase shift, as a function of temperature and density, to cover both the bound state and the scattering contribution.

An important issue in using the formalism described in our work is the question of correlations higher than second order, for instance, if bound states are formed with more than two particles, such as molecules in a plasma. We can formulate a cluster-virial [24] where the atoms are the ‘elementary’ particles. The effective (Lennard-Jones) interaction leads to the formation of molecules. This chemical picture should be obtained within a physical picture of electrons and ions as the elementary particles, as can be done by introducing the corresponding ladder diagrams. A systematic approach, valid also at high densities, has to be formulated consistently, including screening and Pauli blocking.

We mention as well the application of the Beth-Uhlenbeck approach to quark-hadron system, where three quarks can bind to form a nucleon, and mesons are bound states of a quark and an antiquark, described as “plasmons” of the relativistic quark plasma within the Nambu–Jona-Lasinio model [16, 17]. This model field theory is suitable for describing the dynamical chiral symmetry breaking transition which generates a quark mass gap in the vacuum and which leads to the pion being simultaneously the Goldstone boson of this broken symmetry and a bound state. At finite temperatures and chemical potentials, restoration of chiral symmetry triggers the Mott dissociation of the pion which loses its character as a Goldstone boson and becomes a resonance in the quark plasma with a mass pole shifted to the complex energy plane, where the imaginary part of the self energy describes the finite decay width (inverse lifetime) of this state. An achievement of the Beth-Uhlenbeck formulation with phase shifts has been a generalization to finite temperatures of the Levinson theorem for mesons [18] and diquarks [14, 15], as a prerequisite for describing baryons as quark-diquark bound states within a cluster virial expansion. The Beth-Uhlenbeck approach has recently been applied in Ref. [25] to the (2+1)D Gross-Neveu model as a model field theory to describe the Mott dissociation of excitons in graphene-type materials. We note that phenomenological approaches that account for a finite lifetime of hadronic states in a resonance gas by simply introducing their spectral broadening violate the in-medium Levinson theorem! See also Ref. [26].

In Ref. [27] the Beth-Uhlenbeck approach to scalar-pseudoscalar mesons was extended to the three-flavor case with kaons and their chiral partner states. First extensions to the generalized Beth-Uhlenbeck scheme, the subject of the present work have been presented in [28, 29]. This scheme has recently successfully described the transition from a full hadron resonance gas to the quark-gluon plasma, where Mott dissociation of hadrons and the simultaneous occurrence of quarks and gluons was driven by the chiral restoration transition [30]. This work demonstrated that the description of the quark-hadron thermodynamics within the generalized Beth-Uhlenbeck-type model is in excellent agreement with data from *ab-initio* lattice QCD thermodynamics and can be used to interpret them, e.g., by quantifying the composition of the system for given values of the temperature and chemical potentials.

For kinetic processes, it is relevant how correlations are treated and how quasiparticles are formulated and included. The Boltzmann equation, in particular the Landau collision term, contributes to the entropy production.

Of interest also is the Dyson-Bethe-Salpeter equation of motion for the two-particle Green function, see Ref. [31]. The kernel in Equation (4.3) on that report is divided in a static part and a dynamic part, with the static part, Eq. (4.7) there defining the mean-field contributions. This is also introduced by the cluster-mean field approach [32, 33].

IV. CONCLUSIONS

We have shown here, how a generalized Beth-Uhlenbeck equation of state, with the typical $\sin^2 \delta$ correction factor relative to the standard approach, can be obtained within the Φ -derivable approach to model quantum field theories, assuming a two-loop Φ -functional (yielding "sunset" diagrams). Such a choice is quite general and finds applications, e.g., in describing the Mott dissociation of nuclear clusters in nuclear matter or of hadrons in quark matter.

Acknowledgments

With great pleasure we dedicate this paper to Michael Bonitz on his sixty-fifth birthday. He created the International Workshop series on "Kadanoff-Baym Equations: Progress and Perspectives for Many-Body Physics" at the University of Rostock, where in the 1999 Edition the authors came together for the first time in person and G.B. presented a fresh look on his Φ -derivable approach from the perspective of the then just published Ref. [5].

G.B. and D.B. are grateful to the WE-Heraeus Foundation for supporting their participation in the Seminar on "Optics and its Applications in Quantum Technologies" in Yerevan (Armenia), where the main idea for this contribution was conceived. The work of D.B. was supported by the Polish NCN under grant No. 2021/43/P/ST2/03319. G.R. acknowledges an honorary stipend from the Foundation for Polish Science within the Alexander von Humboldt program under grant No. DPN/JJL/402-4773/2022 which supports his visits at the University of Wroclaw where this work was finalized.

-
- [1] L. P. Kadanoff and G. Baym, *Quantum Statistical Mechanics*, 1962.
 - [2] G. Baym and L. P. Kadanoff, "Conservation laws and correlation functions," *Phys. Rev.* **124**, 287 (1961).
 - [3] G. Baym, "Self consistent approximations in many body systems," *Phys. Rev.* **127**, 1391 (1962).
 - [4] E. Riedel, "The entropy expression of a Fermi liquid in shielded potential approximation. Application to ^3He ," *Zeitschr. Phys.* **210**, 403 (1968).
 - [5] B. Vanderheyden and G. Baym, "Self-consistent approximations in relativistic plasmas: Quasiparticle analysis of the thermodynamic properties," *J. Statist. Phys.* **93**, 843 (1998).
 - [6] J. P. Blaizot, E. Iancu and A. Rebhan, "Approximately self-consistent resummations for the thermodynamics of the quark gluon plasma. I. Entropy and density," *Phys. Rev. D* **63**, 065003 (2001).
 - [7] G. Uhlenbeck, E. Beth, "The quantum theory of the non-ideal gas. I. Deviations from the classical theory," *Physica* **3**, 729 (1936).
 - [8] E. Beth, G. Uhlenbeck, "The quantum theory of the non-ideal gas. II. Behaviour at low temperatures," *Physica* **4**, 915 (1937).
 - [9] R. Zimmermann, H. Stolz, "The Mass Action Law in Two-Component Fermi Systems Revisited. Excitons and Electron-Hole Pairs," *physica status solidi (b)* **131**, 151 (1985).
 - [10] M. Schmidt, G. Röpke, H. Schulz, "Generalized Beth-Uhlenbeck approach for hot nuclear matter," *Annals Phys.* **202**, 57 (1990).
 - [11] G. Röpke, K. Kilimann, W. D. Kraeft, D. Kremp and R. Zimmermann, "The Influence of Dynamical Effects on the Two-Particle States (Excitons) in the Electron-Hole Plasma," *Phys. Stat. Sol. (b)* **88**, K59 (1978).
 - [12] R. Zimmermann, K. Kilimann, W. D. Kraeft, D. Kremp and G. Röpke, "Dynamical screening and self-energy of excitons in the electron-hole plasma," *Phys. Stat. Sol. (b)* **90**, 175 (1978).
 - [13] G. Röpke and R. Der, "The influence of two-particle states (excitons) on the dielectric function of the electron-hole plasma," *phys. stat. sol. (b)* **92**, 501 (1979).
 - [14] D. Blaschke, M. Buballa, A. Dubinin, G. Röpke and D. Zablocki, "Generalized Beth-Uhlenbeck approach to mesons and diquarks in hot, dense quark matter," *Annals Phys.* **348**, 228 (2014).
 - [15] D. Blaschke, A. Dubinin and M. Buballa, "Polyakov-loop suppression of colored states in a quark-meson-diquark plasma," *Phys. Rev. D* **91**, no. 12, 125040 (2015).
 - [16] J. Hüfner, S. P. Klevansky, P. Zhuang and H. Voss, "Thermodynamics of a quark plasma beyond the mean field: A generalized Beth-Uhlenbeck approach," *Annals Phys.* **234**, 225 (1994).
 - [17] P. Zhuang, J. Hüfner and S. P. Klevansky, "Thermodynamics of a quark - meson plasma in the Nambu-Jona-Lasinio model," *Nucl. Phys. A* **576**, 525 (1994).
 - [18] A. Wergieluk, D. Blaschke, Y. L. Kalinovsky and A. Friesen, "Pion dissociation and Levinson's theorem in hot PNJL quark matter," *Phys. Part. Nucl. Lett.* **10**, 660 (2013).
 - [19] K. Yamazaki and T. Matsui, "Quark-Hadron Phase Transition in the PNJL model for interacting quarks," *Nucl. Phys. A* **913**, 19 (2013).
 - [20] D. Blaschke and D. Ebert, "Variational path-integral approach to back-reactions of composite mesons in the Nambu-Jona-Lasinio model," *Nucl. Phys. B* **921**, 753 (2017).
 - [21] R. Dashen, S. K. Ma and H. J. Bernstein, "S Matrix formulation of statistical mechanics," *Phys. Rev.* **187**, 345-370 (1969).
 - [22] W. Weinhold, "Thermodynamik mit Resonanzzuständen", Dissertation (unpublished), TU Darmstadt (1998).
 - [23] V. G. Morozov and G. Röpke, "Kinetic theory of radiation in non-equilibrium relativistic plasmas," *Annals Phys.* **324**, 1261 (2009).
 - [24] G. Röpke, N.-U. Bastian, D. Blaschke, T. Klähn, S. Typel and H. H. Wolter, "Cluster virial expansion for nuclear matter within a quasi-particle statistical approach," *Nucl. Phys. A* **897**, 70 (2013).

- [25] B. Mahato, D. Blaschke and D. Ebert, "Beth-Uhlenbeck equation for the thermodynamics of fluctuations in a generalized (2+1)D Gross-Neveu model," Phys. Rev. D **112**, 036013 (2025).
- [26] Z. Yu, G. Baym, and C. J. Pethick, "Calculating energy shifts in terms of phase shifts," J. Phys. B: At. Mol. Opt. Phys. **44**, 195207-1-6, (2011); arXiv:0911.2853.
- [27] D. Blaschke, A. Dubinin, A. Radzhabov and A. Wergieluk, "Mott dissociation of pions and kaons in hot, dense quark matter," Phys. Rev. D **96**, no.9, 094008 (2017).
- [28] D. Blaschke, A. Dubinin and L. Turko, "Mott-hadron resonance gas and lattice QCD thermodynamics," Phys. Part. Nucl. **46**, no. 5, 732 (2015).
- [29] D. Blaschke, A. Dubinin and L. Turko, "Generalized Beth-Uhlenbeck approach to the equation of state for quark-hadron matter," Acta Phys. Polon. Supp. **10**, 473-480 (2017).
- [30] D. Blaschke, M. Cierniak, O. Ivanytskyi and G. Röpke, "Thermodynamics of quark matter with multi-quark clusters in an effective Beth-Uhlenbeck type approach," Eur. Phys. J. A **60**, no.1, 14 (2024).
- [31] P. Schuck, D.S. Delion, J. Dukelsky, M. Jemai, E. Litvinova, G. Röpke, M. Tohyama, "Equation of Motion Method for strongly correlated Fermi systems and Extended RPA approaches," Physics Reports **929**, 1-84 (2021).
- [32] G. Röpke, "Bound states and superfluidity in strongly coupled Fermion systems," Ann. Physik (Leipzig) **3**, 145 (1994).
- [33] Jiasen Jin et al., "Cluster Mean-Field Approach to the Steady-State Phase Diagram of Dissipative Spin Systems," Phys. Rev. X **6**, 031011 (2016).

Appendix A: Proving $S' = 0$ in scalar field theory with a ϕ^3 self-interaction at two-loop order

In this Appendix, we demonstrate the cancellation expressed by the vanishing of S' , Eq. (13) for the simple case of a scalar field ϕ with a ϕ^3 interaction vertex. The two-particle irreducible Feynman diagram for the Φ - functional in two-loop approximation is shown in Fig. 3.




FIG. 3: Feynman diagram for the Φ - functional for scalar ϕ^3 theory at two-loop order (sunset diagram).

We write S' for this case,

$$S' \equiv -\frac{\partial(T\Phi/V)}{\partial T} \Big|_D + \int \frac{d^4k}{(2\pi)^4} \left\{ \frac{\partial n(\omega)}{\partial T} \text{Re} \Pi \text{Im} D \right\}, \quad (\text{A1})$$

where the first term is

$$-\frac{T}{V} \Phi = \frac{g^2}{12} T^2 \sum_{\omega_1, \omega_2} \int \frac{d^3k_1 d^3k_2}{(2\pi)^6} D(\omega_1, |k_1|) D(\omega_2, |k_2|) D(-\omega_1 - \omega_2, |-k_1 - k_2|). \quad (\text{A2})$$

Using the spectral representation

$$D(\omega, k) = \int_{-\infty}^{\infty} \frac{dk_0}{2\pi} \frac{\rho(k_0, k)}{k_0 - \omega} \quad (\text{A3})$$

and evaluating Matsubara sums, we obtain

$$-\frac{T}{V} \Phi = \frac{g^2}{12} T^2 \sum_{\omega_1, \omega_2} \int \frac{d^4k d^4k' d^4k''}{(2\pi)^9} \delta^{(3)}(\mathbf{k} + \mathbf{k}' + \mathbf{k}'') \rho(k) \rho(k') \rho(k'') \frac{-1}{\omega_1 - k_0} \frac{-1}{\omega_2 - k'_0} \frac{1}{\omega_1 + \omega_2 + k''_0}. \quad (\text{A4})$$

Partial fraction decomposition of the three energy denominators and Matsubara summation over ω_1, ω_2 yields

$$\frac{1}{k_0 + k'_0 + k''_0} \{ [n(k''_0) + 1][n(k_0) + n(k'_0) + 1] + n(k_0)n(k'_0) \}. \quad (\text{A5})$$

Performing the temperature derivative and renaming variables under the integrals we obtain

$$\partial_T [n(k_0 + n(k'_0) + n(k''_0)) + n(k'_0)n(k_0) + n(k'_0)n(k''_0) + n(k_0)n(k''_0)] \rightarrow 3\partial_T n(k_0) [1 + n(k'_0) + n(k''_0)]. \quad (\text{A6})$$

To evaluate the second term in Eq. (A1), we perform the Matsubara sums

$$\begin{aligned} \text{Re}\Pi(\omega, q) &= -\frac{g^2}{2} \int \frac{d^3k}{(2\pi)^3} \int \frac{dk_0}{2\pi} \int \frac{dk'_0}{2\pi} \rho(k_0, |k|) \rho(k'_0, |\mathbf{k} + \mathbf{q}|) \sum_{\omega_1} \frac{1}{\omega_1 - k_0} \frac{1}{\omega_1 + \omega - k'_0} \\ &= -\frac{g^2}{2} \int \frac{d^3k}{(2\pi)^3} \int \frac{dk_0}{2\pi} \int \frac{dk'_0}{2\pi} \rho(k_0, |k|) \rho(k'_0, |\mathbf{k} + \mathbf{q}|) \frac{1 + n(k_0) + n(k'_0)}{\omega + k_0 + k'_0} \end{aligned} \quad (\text{A7})$$

and obtain

$$\begin{aligned} &\int \frac{d^4q}{(2\pi)^4} \frac{\partial n(k_0)}{\partial T} \text{Re}\Pi(\omega, q) \text{Im}D(\omega, q) = \\ &= -\frac{g^2}{2 \cdot 2} \int \frac{d^4q}{(2\pi)^4} \int \frac{d^4k}{(2\pi)^4} \int \frac{d^4k'}{2\pi} \delta^{(3)}(\mathbf{q} + \mathbf{k} + \mathbf{k}') \rho(q) \rho(k) \rho(k') \partial_T n(q_0) [1 + n(k_0) + n(k'_0)] \frac{1}{q_0 + k_0 + k'_0}. \end{aligned} \quad (\text{A8})$$

This proves the cancellation of S' for the scalar theory with cubic selfinteraction in the 2-loop approximation (sunset diagram) for the Φ -functional.

This cancellation holds as well for the density $n = N/V = -\partial\Omega/\partial\mu$, see [5].

Appendix B: Generalized optical theorems

The following derivation parallels similar steps performed for the thermodynamic \mathcal{T} -matrix in [9]. In this Appendix, we denote the real and imaginary parts of complex functions $F(z, \mathbf{q})$ by subscripts R and I , respectively, and suppress their arguments so that $F_I = \text{Im}F(\omega + i0^+, \mathbf{q})$ and $F_R = \text{Re}F(\omega, \mathbf{q})$. We decompose the propagator and the photon self-energy (polarization function) into real and imaginary part, $D = D_R + iD_I$ and $\Pi = \Pi_R + i\Pi_I$, resp.

Assuming that the inverse exists, we can write out the two identities $D = D^* D^{*-1} D$ and $D^* = D^* D^{-1} D$ as,

$$D_R + iD_I = D^*(D_R^{-1} - iD_I^{-1})D, \quad (\text{B1})$$

$$D_R - iD_I = D^*(D_R^{-1} + iD_I^{-1})D. \quad (\text{B2})$$

Adding and subtracting these equations yields

$$D_R = D^* D_R^{-1} D, \quad (\text{B3})$$

$$D_I = -D^* D_I^{-1} D, \quad (\text{B4})$$

which together with $D^{-1} = D_0^{-1} - \Pi$, can be combined to

$$D_I = D^* \Pi_I D = D \Pi_I D^*, \quad (\text{B5})$$

which is the off-shell optical theorem.

Differentiating (B4) both sides with respect to the energy ω (denoting this derivative with a prime) and using the fact, that in the Coulomb gauge the free photon propagator $D_0 = 1/\mathbf{q}^2$ is a real and independent of energy, so that $(D_R^{-1})' = -\Pi_R'$ and $D_I^{-1} = -\Pi_I$, we obtain

$$\begin{aligned} D_R' &= D^{*'} D_R^{-1} D + D^* (D_R^{-1})' D + D^* D_R^{-1} D' \\ &= D^{*'} \underbrace{(D_R^{-1} + iD_I^{-1} - iD_I^{-1})}_{D^{-1}} D + D^* (D_R^{-1})' D + D^* \underbrace{(D_R^{-1} - iD_I^{-1} + iD_I^{-1})}_{D^{*-1}} D' \\ &= \underbrace{D^{*'} + D'}_{2D_R'} - iD^{*'} D_I^{-1} D + iD^* D_I^{-1} D' + D^* (D_R^{-1})' D \\ &= D^* \Pi_R' D - iD^{*'} \Pi_I D + iD^* \Pi_I D', \end{aligned} \quad (\text{B6})$$

which can be called a derivative optical theorem.

Multiplying both sides of Eq. (B6) by Π_I and using cyclic invariance of the product of matrices under the trace (integral over energy and momentum) and the optical theorem (B5) one obtains the following.

$$D'_R \Pi_I = \underbrace{D^* \Pi'_R D \Pi_I}_{\Pi'_R D_I} + \underbrace{i D^* \Pi_I S' \Pi_I - i D^{*'} \Pi_I D \Pi_I}_{2\text{Im}[\Pi_I D \Pi_I D^{*'}]} , \quad (\text{B7})$$

which completes the proof of the relationship

$$D'_R \Pi_I - \Pi'_R D_I = 2\text{Im} [\Pi_I D \Pi_I D^{*'}] . \quad (\text{B8})$$