

THE DIRICHLET HEAT TRACE FOR DOMAINS WITH CURVED CORNERS

SHI-ZHUO LOOI AND DAVID SHER

ABSTRACT. We study the short-time asymptotics of the Dirichlet heat trace on planar curvilinear polygons. For such domains we show that the coefficient of $t^{1/2}$ in the expansion splits into a boundary integral of κ^2 and a sum of local corner contributions, one for each vertex. Each curved corner contribution depends only on the interior angle α and on the limiting curvatures κ_{\pm} on the adjacent sides. Using a conformal model and a parametrix construction on the sector heat space, we express this contribution in the form $c_{1/2}(\alpha) r_0(\alpha, \kappa_+, \kappa_-)$, where $c_{1/2}(\alpha)$ is given by a Hadamard finite part of an explicit trace over the exact sector. For right-angled corners we compute $c_{1/2}(\pi/2) = 1/(16\sqrt{\pi})$ and obtain a closed formula for the $t^{1/2}$ coefficient. As an application we extend a previous result in the literature by showing that any admissible curvilinear polygon that is Dirichlet isospectral to a polygon must itself be a polygon with straight sides.

1. INTRODUCTION

A classical problem in spectral geometry is to determine what geometric information is encoded in the spectrum of the Laplacian on a Riemannian manifold [8]. This is frequently approached using heat trace methods. Specifically, one examines the short-time asymptotic expansion of the trace of the heat kernel. This trace is spectrally determined, and therefore the coefficients in the asymptotic expansion are *spectral invariants* – geometric quantities determined by the spectrum. In 1967, McKean and Singer computed the first three heat invariants for a smoothly bounded domain in \mathbb{R}^2 [7]. Using these, they showed that the area, perimeter, and Euler characteristic of such a domain are all spectral invariants. It is natural to ask whether one can deduce similar information for a less smooth domain, and indeed this has been studied in depth [2, 11, 10].

We now explain the setting of this paper. Let $\Omega \subseteq \mathbb{R}^2$ be a curvilinear polygon, with piecewise smooth boundary, smooth except for a finite number n of corners P_j of interior angles $\alpha_j > 0$, $j = 1, \dots, n$. Let κ be the inward-pointing curvature of the boundary, and let $\gamma(s)$ be an arc-length parametrization of the boundary, oriented counterclockwise. For each j , let $\kappa_{j,\pm}$ be the limits of $\kappa(\gamma(s))$ as s approaches $\gamma^{-1}(P_j)$ from above and below respectively.

It is well known that the trace of the Dirichlet heat kernel on Ω , $H_{\Omega}^D(t)$, has an asymptotic expansion as $t \rightarrow 0$, beginning with a t^{-1} term. The t^{-1} , $t^{-1/2}$, and t^0 terms of this expansion have been computed, first in the exact polygonal setting in unpublished work of Ray, as cited by [2] and [12], and then extended to the general curvilinear setting [10]. Our interest here is in the next term, which we expect at order $t^{1/2}$. In [4], based on results in the literature such as [1], the authors give the coefficient of this $t^{1/2}$ term in the “straight corners” case, where a neighborhood of each corner P_j is isometric to an exact sector. Without this assumption, the curvature of the sides may interact with the angle at the corner. As we show, this interaction produces a term at order $t^{1/2}$.

Theorem 1. *With notation as above, the Dirichlet heat trace $H_\Omega^D(t)$ has an expansion as $t \rightarrow 0$ given by*

$$(1) \quad H_\Omega^D(t) = \frac{|\Omega|}{4\pi t} - \frac{|\partial\Omega|}{8\sqrt{\pi t}} + \frac{1}{12\pi} \left(\int_{\partial\Omega} \kappa ds + \sum_{j=1}^n \frac{\pi^2 - \alpha_j^2}{2\alpha_j} \right) \\ + \sqrt{t} \left(\frac{1}{256\sqrt{\pi}} \int_{\partial\Omega} \kappa^2 ds + \sum_{j=1}^n \mathcal{C}_{1/2}(\alpha_j, \kappa_{j,+}, \kappa_{j,-}) \right) + O(t \log t),$$

where $\mathcal{C}_{1/2}$ is a function only of the angle α and the limit curvatures κ_\pm .

We can say substantially more about $\mathcal{C}_{1/2}(\alpha, \kappa_+, \kappa_-)$ whenever $\alpha \neq \pi$. Define

$$(2) \quad r_0 = r_0(\alpha, \kappa_+, \kappa_-) = \frac{1}{2} \csc \alpha \sqrt{\kappa_+^2 + \kappa_-^2 - 2\kappa_+ \kappa_- \cos \alpha}.$$

Theorem 2. *For $\alpha \neq \pi$, the function $\mathcal{C}_{1/2}(\alpha, \kappa_+, \kappa_-)$ has the form*

$$(3) \quad \mathcal{C}_{1/2}(\alpha, \kappa_+, \kappa_-) = c_{1/2}(\alpha) r_0(\alpha, \kappa_+, \kappa_-)$$

where $c_{1/2}(\alpha)$ is a function depending only on α , whose general expression is given by a Hadamard renormalized trace on an exact sector:

$$(4) \quad c_{1/2}(\alpha) = - \text{fp} \iint_{(0,\infty) \times (0,\alpha)} \int_0^1 (1-\sigma)^{-1} \sigma^{-1/2} \iint_{(0,\infty) \times (0,\alpha)} \\ a \left(\frac{R}{\sqrt{1-\sigma}}, \theta, \frac{\sqrt{\sigma} \rho}{\sqrt{1-\sigma}}, \phi \right) b \left(\rho, \phi, \frac{R}{\sqrt{\sigma}}, \theta \right) \rho d\rho d\phi R dR d\theta.$$

Here a and b are functions related to the exact heat kernel on a sector of angle α , explicitly given by (31) and (46), and fp denotes the Hadamard finite part at $R = \infty$ of the improper integral in R .

Moreover, $c_{1/2}(\alpha)$ has the special value

$$(5) \quad c_{1/2}\left(\frac{\pi}{2}\right) = \frac{1}{16\sqrt{\pi}}.$$

Remark. *A few comments:*

- The $O(t \log t)$ error in (1) is not necessarily optimal, and in fact we expect that it should be $O(t)$.
- Each of a and b is a function only of α and the variables of integration, see (31), (46). Thus (4) is indeed independent of κ_\pm .
- For inverse spectral results, it is crucial that (5) is nonzero. We can only verify this explicitly for $\alpha = \pi/2$. It would be of significant interest to find out for which α we have $c_{1/2}(\alpha) \neq 0$.
- Theorem 2 does not work for $\alpha = \pi$ because the conformal model that we use breaks down in that case, as can be seen from the fact that r_0 is undefined when $\alpha = \pi$. So the case $\alpha = \pi$, where the boundary has an abrupt change in second derivative, remains open.

1.1. Inverse spectral applications. In [4], the authors prove that any curvilinear polygon with straight corners is in fact isospectral to a polygon. They require straight corners because of a lack of a formula in the curved corner case. We can strengthen their result.

Definition. A curvilinear polygon Ω is admissible if, for each j , either $c_{1/2}(\alpha_j) > 0$ or Ω is straight near P_j .

Proposition 1. Suppose that an admissible curvilinear polygon Ω is Dirichlet isospectral to a polygon. Then Ω is a polygon.

Proof. For a polygon, the $t^{1/2}$ term in the heat trace vanishes. So it must vanish for Ω . But under our assumptions, by Theorems 1 and 2, the $t^{1/2}$ coefficient in $H_\Omega^D(t)$ is given by

$$(6) \quad \frac{1}{256\sqrt{\pi}} \int_{\partial\Omega} \kappa^2 ds + \sum_{j=1}^n c_{1/2}(\alpha_j) \sqrt{\kappa_{+,j}^2 + \kappa_{-,j}^2}.$$

Since this coefficient is non-negative and only zero for an exact polygon, Ω is an exact polygon. \square

Remark. Observe from (5) that any polygon where all non-straight corners are right angles is admissible.

1.2. Plan of the paper. In section 2 we prove Theorem 1. This is done by examining the geometric microlocal description of the heat kernel given in [10]. We show that there is a well-defined corner contribution $\mathcal{C}_{1/2}\sqrt{t}$, and then prove that any two corners with the same angle α and limit curvatures κ_\pm yield the same contribution, regardless of the rest of Ω . To take advantage of this uniformity, in section 3 we introduce a simple conformal model for a curvilinear corner and compute the contribution $\mathcal{C}_{1/2}\sqrt{t}$ in that model. This computation proceeds via a parametrix construction as in [10] and allows us to obtain both (3) and (4). Finally, in section 4, we prove (5) by computing a sufficient number of terms in the Dirichlet heat trace expansion for a semicircle and comparing with (1).

ACKNOWLEDGEMENTS

D.S. is grateful for support from the AMS-Simons research enhancement grant 501949-9208. S.L. is grateful for support from NSF grant DMS-2346799.

We thank Javier Gómez-Serrano for suggesting the problem to us.

2. THE CORNER CONTRIBUTION

Our approach to understanding the heat kernel on Ω is based on the methods of geometric microlocal analysis [9, 10]. The strategy is to first resolve the singularities of the heat kernel at $t = 0$ by constructing a “double heat space,” Ω_h^2 , via a sequence of geometric blow-ups. The heat kernel lifts under these blow-ups to a well-behaved (“polyhomogeneous conormal”) function on the manifold with corners Ω_h^2 [10]. Roughly speaking, polyhomogeneous conormal functions are those whose asymptotic expansions at the boundary hypersurfaces are compatible and jointly polyhomogeneous at higher-codimension corners; the precise definition may be found in [9].

The boundary of Ω_h^2 consists of boundary hypersurfaces corresponding to distinct asymptotic regimes. Among these hypersurfaces are:

- The interior diagonal (td), modeling heat flow far from the boundary.
- The side faces (sf), modeling heat flow near a smooth edge.
- The front faces (ff), modeling heat flow at a corner.

The method of proof in [10], which is important for our work here, is to use geometric microlocal analysis to perform a parametrix construction. Specifically, one constructs a parametrix $K(t, z, z')$, an approximate solution to the heat equation, by specifying its leading order behavior at each of the three boundary hypersurfaces td, sf, and ff. This should be done in such a way so that when the heat operator is applied to the parametrix $K(t, z, z')$, the result $R(t, z, z')$ is lower order at td, sf, and ff. In order to arrange this, we choose leading order behavior that solves the relevant model problems for the heat equation at each boundary hypersurface. We also must ensure that the specified behaviors at td, sf, and ff are consistent with each other at the intersections of the boundary hypersurfaces, so that there actually exists a polyhomogeneous conormal function on Ω_h^2 with all of the correct behavior. Once we have a good enough parametrix, we can recover the true heat kernel via the Volterra series

$$H_\Omega^D = K - K \star R + K \star R \star R - \cdots.$$

To analyze this Volterra series, we use a composition formula for polyhomogeneous conormal functions on Ω_h^2 , which in our setting is [10, Theorem 3.16].

To be specific about what we mean when we talk about orders: set $T = \sqrt{t}$. We measure orders at each of td, sf, and ff in powers of T ; a function which has leading order behavior T^{-2} has order -2 . “Lower order” means larger powers of T . Throughout, “vanishes to infinite order at td (or sf, or ff)” means the expansion has no terms of any finite T -order there.

The model cone heat kernel has order -2 at td, sf, and ff. We write $\text{ord}_{\text{ff}}(\cdot)$ for this T -order. The key result of [10, Theorem 3.16] for our setting is that if B vanishes to infinite order at td, then $\text{ord}_{\text{ff}}(A \star B) = \text{ord}_{\text{ff}}(A) + \text{ord}_{\text{ff}}(B) + 4$.

As a first use of these techniques, we show that the corner contribution $\mathcal{C}_{1/2}(\alpha, \kappa_+, \kappa_-)$ indeed depends only on α and κ_\pm . Suppose that there exist two domains Ω and Ω' , each with one corner, with the same α and the same κ_\pm . By [10], the heat kernels H_Ω^D and $H_{\Omega'}^D$ are polyhomogeneous conormal on Ω_h^2 and $(\Omega')_h^2$ respectively. Since α is the same, the faces ff of these respective heat spaces are geometrically identical. We now state a lemma which informally says that: under these conditions, $H_\Omega^D - H_{\Omega'}^D$ has leading order at worst t^0 at ff.

Lemma 1. *Let Ω and Ω' be domains with a single corner having the same opening angle α and the same one-sided boundary curvatures κ_\pm at that corner. Then $H_\Omega^D - H_{\Omega'}^D$ has order $\leq T^0$ (at worst $T^0 = t^0$) at ff. Equivalently, their ff coefficients of orders $T^{-2} = t^{-1}$ and $T^{-1} = t^{-1/2}$ agree.*

Proof. We use $H_{\Omega'}^D$ to build a parametrix K for H_{Ω}^D . We will build this parametrix K for H_{Ω}^D by prescribing its leading behavior at the three faces. Specifically:

- At td, we specify that K should have the usual interior expansion, which in this setting is simply the Euclidean heat kernel on \mathbb{R}^2 , which has order -2 with no lower order terms.
- At sf, we specify that K should have the same expansion as the Dirichlet heat kernel on a smooth domain with boundary whose boundary coincides with the appropriate smooth part of $\partial\Omega$. Such a heat kernel is known to have a polyhomogeneous conormal expansion by [6], and that expansion is consistent with the Euclidean heat kernel at td. The first term, at order -2 , is universal. The second term, at order -1 , follows from the boundary calculus construction in [6]; by a standard scaling argument, this coefficient is a universal multiple of the curvature $\kappa(z')$.
- At ff, we specify that K should have the same t^{-1} (order -2) and $t^{-1/2}$ (order -1) terms as $H_{\Omega'}^D$. These are consistent with the interior expansion at td, since they are consistent for $H_{\Omega'}^D$.

We must check that the ff specifications are consistent with the sf ones. But this is the point of having κ_{\pm} equal. Since the heat kernel $H_{\Omega'}^D$ is polyhomogeneous conormal, our ff specifications are consistent with the Euclidean behavior at td and also with the first two terms at sf on $(\Omega')_h^2$. But since κ_{\pm} are identical between Ω and Ω' , these terms on $(\Omega')_h^2$ have the same restrictions to $\text{sf} \cap \text{ff}$ as the first two terms at sf on Ω_h^2 . Thus our ff and sf specifications are consistent.

This consistency means that there exists a parametrix $K(t, z, z')$ with the specified expansions at td, sf, and ff. This parametrix solves the model problems to infinite order at td and sf and to two orders at ff. So $R(t, z, z') = (\partial_t - \Delta_{\Omega})K(t, z, z')$ has leading order -2 , as opposed to -4 , at ff, and leading order ∞ at td and sf. Using the composition formula of [10], we conclude that for each $n \geq 1$, $K \star (R^{\star n})$ has leading order at worst t^0 at ff.

$$\text{ord}_{\text{ff}}(K \star R^{\star n}) \geq (-2) + n(-2) + 4n = 2n - 2 \geq 0 \quad (n \geq 1).$$

Thus

$$H_{\Omega}^D - K = \sum_{n \geq 1} (-1)^n K \star R^{\star n}$$

has order at worst 0 at ff, and since the same is true for $H_{\Omega'}^D - K$, so $H_{\Omega}^D - H_{\Omega'}^D = O(T^0)$ at ff. The Lemma is proven. \square

We can now prove Theorem 1.

Proof of Theorem 1. As in [10], the heat trace is obtained by restricting H_{Ω}^D to the spatial diagonal and then integrating over $z \in \Omega$. The technical tool used is Melrose's pushforward theorem. We will not repeat all the details here, but the upshot is that each face td, sf, and ff gives a separate contribution to the expansion (1). A priori, the expansions can interact to give logarithmic terms, but for the same reasons as in [10, p. 49], keeping in mind that we now have two orders at ff rather than one, there are no logarithmic terms until at least $O(T^2 \log T) = O(t \log t)$ (and potentially

much later, if at all). The contribution from td is

$$\frac{|\Omega|}{4\pi t} + O(t^\infty).$$

The contributions from sf are

$$-\frac{|\partial\Omega|}{8\sqrt{\pi t}} + \frac{1}{12\pi} \int_{\partial\Omega} \kappa ds + \frac{\sqrt{t}}{256\sqrt{\pi}} \int_{\partial\Omega} \kappa^2 ds + O(t).$$

And the contributions from ff, since the first is already known and the second depends only on α and κ_\pm , are

$$\sum_{j=1}^n \frac{\pi^2 - \alpha_j^2}{24\pi\alpha_j} + \sqrt{t} \sum_{j=1}^n \mathcal{C}_{1/2}(\alpha_j, \kappa_{j,+}, \kappa_{j,-}) + O(t),$$

for some unknown function $\mathcal{C}_{1/2}(\alpha, \kappa_+, \kappa_-)$. Summing the three faces gives Theorem 1. \square

Note that we will perform a similar parametrix construction, in detail, in the next section, when we build a conformal model to find an expression for $\mathcal{C}_{1/2}(\alpha, \kappa_+, \kappa_-)$.

3. THE CONFORMAL MODEL AND THE CONSTRUCTION

The goal of this section is to find an expression for the contribution $\mathcal{C}_{1/2}$. Since it depends only on α and on κ_\pm , we can work with a specific domain with those values. We take advantage of a conformal transformation to build such a domain, and then we construct its heat kernel.

3.1. The conformal model domain. We define a conformal mapping from \mathbb{C} to \mathbb{C} . Let w be the complex coordinate on the target space and z be the complex coordinate on the source space. The Euclidean metric on the target \mathbb{C} is $g_{Euc} = dw d\bar{w}$. Fix $z_0 \in \mathbb{C}$ and let $\Phi = \Phi_{z_0} : \mathbb{C} \rightarrow \mathbb{C}$ be the holomorphic map given by

$$w = \Phi(z) = z + z_0 z^2.$$

Near the origin, it is a small perturbation of the identity map.

To compute the pullback metric $\Phi^*(g_{Euc})$, we use the standard fact that for a holomorphic map Φ , $dw = \Phi'(z)dz$ and $d\bar{w} = \overline{\Phi'(z)}d\bar{z}$. Therefore,

$$\begin{aligned} \Phi^*(g_{Euc}) &= \Phi^*(dw d\bar{w}) \\ &= (\Phi'(z)dz)(\overline{\Phi'(z)}d\bar{z}) \\ &= \Phi'(z)\overline{\Phi'(z)}dz d\bar{z} \\ &= |\Phi'(z)|^2 dz d\bar{z} \end{aligned}$$

Thus, the pullback metric is:

$$\Phi^*(g_{Euc}) = |1 + 2z_0 z|^2 dz d\bar{z}.$$

Let $F(z) = |1 + 2z_0 z|^2$ be the conformal factor derived above. An immediate calculation yields

$$(7) \quad F(z) = 1 + 4\Re(z_0 z) + 4|z_0|^2 |z|^2.$$

If we write $z = x + iy$ and $z_0 = u_0 + iv_0$, we obtain

$$(8) \quad F(x + iy) = 1 + 4(u_0 x - v_0 y) + 4(u_0^2 + v_0^2)(x^2 + y^2).$$

On the other hand, if we write $z = re^{i\theta}$ and $z_0 = r_0 e^{i\theta_0}$ (this r_0 matches (2); see (12), below), we obtain

$$(9) \quad F(re^{i\theta}) = 1 + 4r_0 r \cos(\theta + \theta_0) + 4r_0^2 r^2.$$

We see that F , however it is written, is smooth, though of course not holomorphic, from $\mathbb{C} \rightarrow \mathbb{R}$. It is also smooth down to zero in polar coordinates (r, θ) .

For later use, we also record the gradient of F at the origin:

$$(10) \quad \nabla F(0, 0) = \langle 4u_0, -4v_0 \rangle = 4r_0 \langle \cos \theta_0, -\sin \theta_0 \rangle,$$

where the latter expression is in polar coordinates but still measuring x and y -derivatives,

We now revert to \mathbb{R}^2 . Let Ω_0 be the flat sector $\{z \in \mathbb{C} \mid 0 < \arg(z) < \alpha\}$. The map Φ takes Ω_0 to a curvilinear domain Ω with the same corner angle α . The signed curvatures of the two boundary edges of Ω at the corner are given by:

- κ_1 (on the image of the $\theta = 0$ ray): $2\Im(z_0)$
- κ_2 (on the image of the $\theta = \alpha$ ray): $-2\Im(z_0 e^{i\alpha})$

Here the orientation is chosen so that the signed curvatures measure curvature in the inward-pointing direction, so that a convex domain will have $\kappa_1 \geq 0$ and $\kappa_2 \geq 0$. A domain with symmetry about $\arg(z) = \alpha/2$ will have $\kappa_1 = \kappa_2$.

By choosing the complex parameter z_0 , we can match any desired curvatures κ_+, κ_- , as long as $\alpha \neq \pi$. Solve the following linear system for $z_0 = u_0 + iv_0$:

$$\begin{aligned} \kappa_+ &= 2v_0 \\ \kappa_- &= -2\Im((u_0 + iv_0)(\cos \alpha_0 + i \sin \alpha_0)) = -2(u_0 \sin \alpha_0 + v_0 \cos \alpha_0) \end{aligned}$$

This yields $z_0 = u_0 + iv_0$, where

$$(11) \quad \begin{cases} u_0 = -\frac{1}{2}\kappa_+ \cot \alpha + \frac{1}{2}\kappa_- \csc \alpha; \\ v_0 = \frac{1}{2}\kappa_+. \end{cases}$$

In terms of the curvatures, we observe that

$$(12) \quad r_0 = \frac{1}{2} \csc \alpha \sqrt{\kappa_+^2 - 2\kappa_+ \kappa_- \cos \alpha + \kappa_-^2}; \quad \theta_0 = \arg(u_0 + iv_0).$$

The pullback by Φ of the Euclidean metric on Ω is

$$F \cdot g_{Eucl, \Omega_0},$$

where we now interpret F as a function with domain \mathbb{R}^2 rather than \mathbb{C} . By the formula for the Laplacian in two dimensions under a conformal change of metric $g = Fg_0$, the new Laplacian is $\Delta_g = F^{-1}\Delta_{g_0}$. Therefore, the pullback Laplacian is

$$\Phi^*(\Delta_\Omega) = F^{-1}\Delta_{\Omega_0}.$$

We want to understand the heat kernel for the operator Δ_Ω on the curved domain Ω . By pulling back, this is equivalent to studying the heat kernel for the operator $F^{-1}\Delta_{\Omega_0}$ on Ω_0 . The corresponding heat equation is $\partial_t H = F^{-1}\Delta_{\Omega_0} H$. We define our heat operator on the model space Ω_0 as:

$$\mathcal{L} = \partial_t - F^{-1}\Delta_{\Omega_0}.$$

We now construct the heat kernel for \mathcal{L} . We use the techniques of [10], working at td, then at sf and ff. This corresponds to building a parametrix for the heat kernel, first at the interior diagonal, and then at the diagonal both near the boundary and near a corner. Once we have a sufficiently accurate parametrix, we iterate away the error via a Volterra series to produce the true heat kernel.

3.2. The heat kernel at td. It is proved in [10] that the heat kernel for \mathcal{L} , being the pullback of the heat kernel for the Laplacian on the curvilinear polygon Ω , is polyhomogeneous conormal on the double space of [10]. We also know that its full expansion at td is the same as that for the heat kernel of the operator $\Phi^*(\Delta_{\mathbb{R}^2}) = F^{-1}\Delta_{\mathbb{R}^2}$. This expansion may be written in terms of the (interior) boundary defining function $T = \sqrt{t}$ as

$$T^{-2}a_{-2} + T^{-1}a_{-1} + T^0a_0 + \dots,$$

where a_{-2} , a_{-1} , a_0 , etc. are smooth functions on the face td. We refer to a_k as the td-coefficients of the heat kernel; in particular, a_0 denotes the coefficient of order T^0 at td, whose restriction along $\text{td} \cap \text{ff}$ will determine $\mathcal{C}_{1/2}(\alpha, \kappa_+, \kappa_-)$.

From [10], these functions are smooth down to ff as well. And T vanishes to first order at ff as well as at td. Thus the term $T^k a_k$ has order k at ff.¹ Therefore, the only terms which are relevant to the leading and subleading terms at ff are $T^{-2}a_{-2} + T^{-1}a_{-1}$.

We now use the work of Grieser [6] to compute a_{-2} and a_{-1} . We use his notation throughout this subsection: the coordinates are (t, x, y) , where $x \in \mathbb{R}^2$ is an interior coordinate on the first factor and $y \in \mathbb{R}^2$ on the second. The initial parametrix for the heat kernel is

$$(13) \quad K_1(t, x, y) = \frac{1}{4\pi t} \exp\left(-\frac{F(y)|x-y|^2}{4t}\right).$$

Computing the application of the heat operator $(\partial_t - F^{-1}(x)\Delta_{\mathbb{R}^2})$ to K_1 , we obtain the error term

$$(14) \quad R_1(t, x, y) := (\partial_t - F^{-1}(x)\Delta_{\mathbb{R}^2})K_1(t, x, y) = \left(1 - \frac{F(y)}{F(x)}\right) \frac{\partial}{\partial t} K_1(t, x, y).$$

The prefactor is $O(\sqrt{t})$ near the diagonal, so R_1 is $t^{1/2}$ less singular than $\partial_t K_1$.

Notation (td symbols). In the rescaled variables (t, X, y) with $X = (x - y)/\sqrt{t}$, any heat kernel A will admit an asymptotic of the form

$$(15) \quad A(t, x, y) \sim \sum_{q \in d + \frac{1}{2}\mathbb{N}_0} t^q \Phi_q(A)(X, y)$$

as $t \downarrow 0$ with X fixed. The coefficient $\Phi_q(A)$ is the *td-symbol of order q* of the operator A ; if A has order d , then $\Phi_d(A)$ is the principal td-symbol of A . Referring to Grieser [6] for the definition of

¹This is the reason that the heat expansion does not pick up logarithmic terms from the pushforward theorem coming from the intersection $\text{td} \cap \text{ff}$. When we convert to densities, the $T^k a_k$ term has order $k + 2$ at ff and since this is not k , there are no log terms. Similar considerations rule out logarithmic terms from the other intersections.

the various operator spaces Ψ_H^d , we have $K_1(t, x, y) \in \Psi_H^{-1}$. Roughly, this means that $K_1(t, x, y)$ has an expansion of the form (15) with $d = -1$, and with coefficients decaying to infinite order at $|X| = \infty$.

To find the principal td-symbol of K_1 , we compute, with $X = (x - y)/\sqrt{t}$,

$$K_1(t, X\sqrt{t} + y, y) = \frac{1}{4\pi t} \exp\left(-\frac{F(y)}{4} X^2\right),$$

so

$$(16) \quad \Phi_{-1}(K_1)(X, y) = \frac{1}{4\pi} \exp\left(-\frac{F(y)}{4} X^2\right).$$

We also analyze $R_1(t, X, y)$. Writing it out we obtain

$$(17) \quad R_1(t, X, y) = \left(\frac{F(X\sqrt{t} + y) - F(y)}{F(X\sqrt{t} + y)}\right) \left(\frac{F(y)}{4} X^2 - 1\right) \frac{1}{t} K_1(t, X, y).$$

The leading order term of this expression at $t = 0$ is

$$(18) \quad \left(\frac{\sqrt{t}X \cdot \nabla(F)(y)}{F(y)}\right) \left(\frac{F(y)}{4} X^2 - 1\right) \frac{1}{t} K_1(t, X, y).$$

Therefore $R_1 \in \Psi_H^{-1/2}$ with

$$(19) \quad \Phi_{-1/2}(R_1)(X, y) = \left(\frac{X \cdot \nabla(F)(y)}{F(y)}\right) \left(\frac{F(y)}{4} X^2 - 1\right) \frac{1}{4\pi} \exp\left(-\frac{F(y)}{4} X^2\right).$$

By the heat kernel construction in Grieser [6], the leading and subleading terms of the true heat kernel at td will be the same as those of

$$K_1 - K_1 * R_1.$$

K_1 has zero subleading term in the coordinates (t, X, y) , so for the subleading term we just need to understand $K_1 * R_1$, in particular just the *leading* term of $K_1 * R_1$, which is $\Phi_{-3/2}(K_1 * R_1)$. This can be computed using [6, Proposition 2.6]. Observe that the integration in Z is with respect to $F(y) dZ$, the metric at y . We obtain, including the minus sign,

$$(20) \quad \Phi_{-3/2}(-K_1 * R_1)(X, y) = - \int_0^1 \int_{\mathbb{R}^2} (1 - \sigma)^{-1} \sigma^{-3/2} \frac{1}{4\pi} \exp\left(-\frac{F(y)}{4} \frac{(X - Z)^2}{1 - \sigma}\right) \\ \frac{Z \cdot \nabla(F)(y)}{\sqrt{\sigma} F(y)} \left(\frac{F(y)}{4} \frac{Z^2}{\sigma} - 1\right) \frac{1}{4\pi} \exp\left(-\frac{F(y)}{4} \frac{Z^2}{\sigma}\right) F(y) dZ d\sigma.$$

This integral can be computed, which we now do. Let

$$c = \frac{F(y)}{4}; \quad V = \nabla(F)(y); \quad \tilde{Z} = \frac{Z}{\sqrt{\sigma}}; \quad \tilde{X} = \frac{X}{\sqrt{\sigma}}.$$

We obtain

$$(21) \quad \Phi_{-3/2}(-K_1 * R_1) = -\frac{1}{16\pi^2} \int_0^1 (1 - \sigma)^{-1} \sigma^{-1/2} \\ \int_{\mathbb{R}^2} \tilde{Z} \cdot V (c\tilde{Z}^2 - 1) \exp(-c\tilde{Z}^2 - c\sigma \frac{(\tilde{X} - \tilde{Z})^2}{1 - \sigma}) d\tilde{Z} d\sigma.$$

As usual, we can complete the square to do the inner integral. Returning to X , we get

$$\exp(-cX^2) \int_{\mathbb{R}^2} \tilde{Z} \cdot V(c\tilde{Z}^2 - 1) \exp(-\frac{c}{1-\sigma}(\tilde{Z} - \sqrt{\sigma}X)^2) d\tilde{Z}.$$

Let $\hat{Z} = \tilde{Z} - \sqrt{\sigma}X$:

$$\exp(-cX^2) \int_{\mathbb{R}^2} (\hat{Z} + \sqrt{\sigma}X) \cdot V(c(\hat{Z} + \sqrt{\sigma}X)^2 - 1) \exp(-\frac{c}{1-\sigma}\hat{Z}^2) d\hat{Z}.$$

Any of the terms in the prefactor

$$((\hat{Z} + \sqrt{\sigma}X) \cdot V)(c(\hat{Z} + \sqrt{\sigma}X)^2 - 1)$$

which are odd in one of the coordinates of \hat{Z} yield zero when integrated. Up to such terms, that pre-factor is

$$(\sqrt{\sigma}X \cdot V)(c\hat{Z}^2 + c\sigma X^2 - 1) + 2c\sqrt{\sigma}(\hat{z}_1^2 x_1 v_1 + \hat{z}_2^2 x_2 v_2).$$

Using the usual formulas for Gaussian integrals, we get that the inner integral in (21) is

$$\begin{aligned} \exp(-cX^2) [(\sqrt{\sigma}X \cdot V)(c\sigma X^2 - 1) \frac{\pi(1-\sigma)}{c} + (\sqrt{\sigma}X \cdot V) \frac{2\pi(1-\sigma)^2}{c}] \\ = c^{-1}\pi\sigma^{1/2}(1-\sigma) \exp(-cX^2)(X \cdot V)(c\sigma X^2 + 1 - 2\sigma), \end{aligned}$$

and therefore

$$(22) \quad \Phi_{-3/2}(-K_1 * R_1) = -\frac{1}{16c\pi} \exp(-cX^2)(X \cdot V) \int_0^1 (c\sigma X^2 + 1 - 2\sigma) d\sigma,$$

Performing the simple integration, we obtain

$$(23) \quad \Phi_{-3/2}(-K_1 * R_1) = -\frac{1}{32\pi} \exp(-cX^2)(X \cdot V)X^2.$$

Inserting the variables, we have

$$(24) \quad \Phi_{-3/2}(-K_1 * R_1) = -\frac{1}{32\pi} X^2 (X \cdot \nabla F(y)) \exp(-\frac{F(y)}{4} X^2).$$

We can now write down the two leading orders of the heat kernel at td: they are

$$(25) \quad T^{-2} \frac{1}{4\pi} \exp(-\frac{F(y)}{4} X^2) - T^{-1} \frac{1}{32\pi} X^2 (X \cdot \nabla F(y)) \exp(-\frac{F(y)}{4} X^2).$$

As one may check, this expression is in the kernel of $T^2\mathcal{L}$ up to $O(T^0)$.

Remark. *Two observations:*

- *If F is constant, the subleading term is zero. In fact all subleading terms are zero – this is the Euclidean setting, where the initial parametrix is equal to the true heat kernel.*
- *In all cases, $\Phi_{-3/2}(K_1 * R_1)(0, y) = 0$. This means that the restriction of $K_1 * R_1$ to the spatial diagonal has no term of order T^{-1} at td. This is unsurprising, as we do not expect any term of the form $T^{-1} = t^{-1/2}$ in the interior heat trace expansion.*

3.3. A parametrix for the heat kernel. We take our heat operator \mathcal{L} and apply it to a parametrix

$$H^{(1)} := G \cdot H_{\text{cone}},$$

where H_{cone} is the Dirichlet heat kernel for an exact sector of angle α , and where G is a scalar function chosen as a corrector to allow $H^{(1)}$ to satisfy the heat equation to two orders at td. Specifically,

$$G := (1 - \frac{1}{8}TX^2X \cdot \nabla F(y)) \exp[-\frac{F(y) - 1}{4}X^2],$$

which, when multiplied by the Euclidean heat kernel (the model for H_{cone} at td), yields (25). We do not claim that G is polyhomogeneous on the double space; it may blow up at td depending on the sign of $F(y) - 1$ (because $X = |z - z'|/T$). However, G is polyhomogeneous away from td and $G \cdot H_{\text{cone}}$ is polyhomogeneous everywhere.

Now examine G at ff, the corner front face. This face is the most singular stratum of the heat space, where $t \rightarrow 0$ and both points $x, y = 0$. The function G is equal to 1 at ff because $T = 0$ at ff, and $F = 1$ at ff. As a consequence, $H^{(1)}$ is in the kernel of the heat operator \mathcal{L} to leading order at ff; we say that it “solves the model problem” to one order at ff. It also solves the model problem to two orders at td. And since $T = 0$ at sf and $\exp[-\frac{1}{4}(F(y) - 1)X^2]H_{\text{cone}}$ agrees to first order at sf with the initial parametrix $\exp[-\frac{1}{4}(F(y) - 1)X^2]H_{\mathbb{R}_+^2}$ used by Grieser at sf [6], $H^{(1)}$ also solves the model problem to one order at sf.

Now define:

$$R^{(1)} := \mathcal{L}(H^{(1)}).$$

If $H^{(1)}$ were not correctly chosen, we would expect $R^{(1)}$ to be two orders worse than $H^{(1)}$ at each of td, ff, and sf, which means order -4 at each. However, since $H^{(1)}$ solves the model problem to two orders at td and one order at ff, and also one order at sf, $R^{(1)}$ has leading orders given by -2 at td, -3 at ff, and -3 at sf.

In order to use the composition theorem of [10, Theorem 3.16], which requires one of the two operators to vanish to infinite order at td, we work with a slightly improved parametrix $H^{(2)}$ given by solving away the full expansion at td. This equals $H^{(1)}$ plus a term which is order *zero* at td, ff, and sf. In particular, we note that if we have

$$R^{(2)} := \mathcal{L}(H^{(2)}),$$

then $R^{(2)} - R^{(1)}$ has order -2 at ff. $H^{(2)} - H^{(1)}$ also has order -2 at ff. Moreover, $R^{(2)}$ vanishes to infinite order at td and it has leading order -3 at both ff and sf.

By Duhamel’s principle and our usual Neumann series sum, the true heat kernel for our conformal metric is

$$H = H^{(2)} - H^{(2)} * R^{(2)} + \sum_{k=2}^{\infty} (-1)^k H^{(2)} * (R^{(2)})^k.$$

We are interested in finding its term of order -1 at ff (the sub-leading term). By the composition formula in [10], using in particular that

$$\text{ord}_{\text{ff}}(A * B) = \text{ord}_{\text{ff}}(A) + \text{ord}_{\text{ff}}(B) + 4,$$

the terms with $k \geq 2$ have no order -1 term at ff, so the order -1 term of our true heat kernel at ff is the same as that of

$$H^{(2)} - H^{(2)} * R^{(2)}.$$

This in turn² is the same as that of

$$H^{(1)} - H^{(1)} * R^{(1)} = GH_{cone} - GH_{cone} * (\mathcal{L}(GH_{cone})).$$

Here $H^{(1)}$ has order -2 at ff, so we need its leading and subleading terms. $R^{(1)}$ has leading ff order -3 . Thus the leading ff term of the convolution $H^{(1)} * R^{(1)}$ comes from $T^{-2} * T^{-3}$, which lands at -1 at ff, by the ff composition rule.

A critical fact here is that by the composition formula in [10], the leading term of $H^{(1)} * R^{(1)}$ at ff depends only on the leading terms of $H^{(1)}$ and $R^{(1)}$ at ff, not on their leading terms at other boundary faces (in particular at sf). This is helpful for the following reasons.

- First, the integral in the convolution $H^{(1)} * R^{(1)}$ is done with respect to the metric Fg_{Eucl} , but since we are computing at the ff face, which corresponds to the vertex where $r = 0$, we can evaluate $F(z)$ at that point. As established, $F(0) = 1$. So, for the purpose of calculating the leading term of the convolution, the integration measure simplifies from the Riemannian volume form to the standard Euclidean one.
- Second, when computing the second convolution factor $\mathcal{L}(GH_{cone})$, we can ignore any term below its leading order -3 . Similarly, for the first convolution factor, GH_{cone} , we only need its leading order term, which is of order T^{-2} . All smoother terms can be ignored as they will not affect the leading term of the result.

This leads to the following simplification. Consider $GH_{cone} * (\mathcal{L}(GH_{cone}))$. We only need to worry about the leading order of each term, so G may be replaced with 1 and we need only analyze

$$H_{cone} * (\mathcal{L}(GH_{cone})).$$

Let us compute $\mathcal{L}(GH_{cone})$. It is

$$(26) \quad \mathcal{L}(GH_{cone}) = \mathcal{L}(G)H_{cone} + G\mathcal{L}(H_{cone}) - 2F^{-1}\nabla G \cdot \nabla(H_{cone}).$$

Using the fact that $\mathcal{L}_{cone}H_{cone} = 0$, this becomes

$$(27) \quad \mathcal{L}(GH_{cone}) = \mathcal{L}(G)H_{cone} - G(F^{-1} - 1)\Delta(H_{cone}) - 2F^{-1}\nabla G \cdot \nabla(H_{cone}).$$

We may replace the \mathcal{L} with \mathcal{L}_{cone} in the first term, as $(\mathcal{L} - \mathcal{L}_{cone})(G)$ has order zero at ff. We may also drop the G in the second term and the F^{-1} in the third term, as they are both 1 to leading order at ff. Thus up to terms which are irrelevant for our calculation,

$$\mathcal{L}(GH_{cone}) \sim \mathcal{L}_{cone}(G)H_{cone} - (F^{-1} - 1)\Delta(H_{cone}) - 2\nabla G \cdot \nabla(H_{cone}).$$

²The justification for using the simpler parametrix $H^{(1)}$ to compute the subleading term at the ff face is done by an analysis of operator orders within the composition calculus of [10]. Let $D = H^{(2)} - H^{(1)}$ be the correction term that improves the parametrix at the td face. By construction, the leading order of D at the ff face is at least T^0 . The difference between the first two terms of the respective Neumann series is $\Delta H = D - H^{(1)} * \mathcal{L}(D) - D * R^{(1)} - D * \mathcal{L}(D)$. A direct application of the composition formula shows that each term in this expression has a leading order of T^0 or higher at the ff face. For instance, the most singular term, $H^{(1)} * \mathcal{L}(D)$, has an order of at least $-2 + (-2) + 4 = 0$. Since the entire difference ΔH is no more singular than T^0 , the expressions $H^{(1)} - H^{(1)} * R^{(1)}$ and $H^{(2)} - H^{(2)} * R^{(2)}$ must agree for all terms of order T^{-1} and below. We are therefore justified in using the simpler expression to compute the desired subleading coefficient.

In conclusion, to two orders at ff, we have

$$(28) \quad H \sim GH_{cone} - H_{cone} * (\mathcal{L}_{cone}(G)H_{cone} - (F^{-1} - 1)\Delta(H_{cone}) - 2\nabla G \cdot \nabla H_{cone}),$$

where the convolution $-$ and all derivatives here $-$ are taken with respect to the Euclidean metric on the cone. The geometric correction factor G must be retained in the stand-alone factor $G H_{cone}$ (before taking the trace): its T^{+1} contribution times the T^{-2} of H_{cone} produces one of the two T^{-1} pieces in the kernel at ff we are calculating.

3.4. Computations with G . We now turn to the explicit calculation of the terms in Equation (28). We are interested in the first two terms in their expansion at ff.

Throughout, we replace Grieser's x and y by z and z' . Let z have polar coordinates (r, θ) and let z' have polar coordinates (r', θ') , and then let

$$R = r/T, R' = r'/T.$$

With this notation, natural local coordinates near the interior of the front face ff are $(T, R, \theta, R', \theta')$, where T is a boundary defining function. In these coordinates, we compute the first two terms in the expansion (28), of orders T^{-2} and T^{-1} . To do this we compute two terms in the expansion of GH_{cone} . We also compute the leading T^{-2} term in the expansion of H_{cone} and the leading T^{-3} term in the expansion of the second convolution factor, as by the composition formula in [10] their convolution will give a term of order $T^{-2-3+4} = T^{-1}$ at ff. To this end, we denote

$$(29) \quad a(R, \theta, R', \theta') = T^2 H_{cone}|_{ff};$$

$$(30) \quad b(R, \theta, R', \theta') = r_0^{-1}(T^3(\mathcal{L}_{cone}(G)H_{cone} - (F^{-1} - 1)\Delta(H_{cone}) - 2\nabla G \cdot \nabla H_{cone}))|_{ff}.$$

Our goal is to find expressions for a and b .

First, we have a formula for H_{cone} due to Cheeger [2]:

$$H_{cone} = \frac{1}{2t} \exp\left[-\frac{1}{4t}(R^2 + (R')^2)\right] \sum_{j=1}^{\infty} I_{\mu_j}\left(\frac{RR'}{2t}\right) \varphi_j(\theta) \varphi_j(\theta'),$$

where $\varphi_j(\theta)$ is the L^2 -normalized eigenfunction

$$\varphi_j(\theta) = \sqrt{\frac{2}{\alpha}} \sin\left(\frac{\pi j \theta}{\alpha}\right).$$

Since by scaling we have

$$H_{cone} = T^{-2} a(R, \theta, R', \theta'),$$

we conclude that

$$(31) \quad a(R, \theta, R', \theta') = \frac{1}{2} \exp\left[-\frac{R^2 + (R')^2}{4}\right] \sum_{j=1}^{\infty} I_{\mu_j}\left(\frac{RR'}{2}\right) \varphi_j(\theta) \varphi_j(\theta').$$

For all other terms we must analyze G . Recall that G is a function of $T = \sqrt{t}$, the source point z' , and the scaled, relative position vector $X = (z - z')/T$. We have the relations

$$\begin{aligned} z &= (r \cos \theta, r \sin \theta) = (RT \cos \theta, RT \sin \theta) \\ z' &= (r' \cos \theta', r' \sin \theta') = (R'T \cos \theta', R'T \sin \theta'). \end{aligned}$$

We can directly compute the components of the scaled vector X :

$$(32) \quad X = (R \cos \theta - R' \cos \theta', R \sin \theta - R' \sin \theta').$$

Define the distance function $D = D(R, \theta, R', \theta')$ by

$$(33) \quad D^2 := |X|^2 = (R \cos \theta - R' \cos \theta')^2 + (R \sin \theta - R' \sin \theta')^2 = R^2 + (R')^2 - 2RR' \cos(\theta - \theta').$$

For future use we also define

$$(34) \quad S = S(R, \theta, R', \theta') = (R \cos(\theta + \theta_0) + R' \cos(\theta' + \theta_0)).$$

With these relations, we can write the full expression for G in the front-face coordinates:

$$(35) \quad G(T, R, \theta, R', \theta') = \left[1 - \frac{1}{8} T D^2 \langle R \cos \theta - R' \cos \theta', R \sin \theta - R' \sin \theta' \rangle \cdot \nabla F(R' T, \theta') \right] \exp\left[-\frac{1}{4} (F(R' T, \theta')) D^2\right].$$

Now we compute the expansion of G at ff in powers of T . Since F is smooth, we have by Taylor series that

$$G \approx 1 - \frac{1}{8} T D^2 \langle R \cos \theta - R' \cos \theta', R \sin \theta - R' \sin \theta' \rangle \cdot \nabla F(0, 0) - \frac{1}{4} T D^2 \langle R' \cos \theta', R' \sin \theta' \rangle \cdot \nabla F(0, 0) + O(T^2).$$

By combining the vector coefficients and plugging in $\nabla F(0, 0) = 4r_0 \langle \cos \theta_0, -\sin \theta_0 \rangle$, this expression simplifies:

$$(36) \quad G = 1 - T D^2 \left[\frac{1}{8} \langle R \cos \theta + R' \cos \theta', R \sin \theta + R' \sin \theta' \rangle \right] \cdot (4r_0 \langle \cos \theta_0, -\sin \theta_0 \rangle) + O(T^2)$$

$$(37) \quad = 1 - \frac{r_0}{2} T D^2 \langle R \cos \theta + R' \cos \theta', R \sin \theta + R' \sin \theta' \rangle \cdot \langle \cos \theta_0, -\sin \theta_0 \rangle + O(T^2)$$

$$(38) \quad = 1 - \frac{r_0}{2} T D^2 (R(\cos \theta \cos \theta_0 - \sin \theta \sin \theta_0) + R'(\cos \theta' \cos \theta_0 - \sin \theta' \sin \theta_0)) + O(T^2)$$

$$(39) \quad = 1 - \frac{r_0}{2} T D^2 S + O(T^2).$$

From this we immediately conclude that, at ff,

$$(40) \quad GH_{cone} = T^{-2} a(R, \theta, R', \theta') - \frac{r_0}{2} T^{-1} D^2 S a(R, \theta, R', \theta') + O(1).$$

One nice feature of this expression is that, since D^2 is zero on the spatial diagonal, the sub-leading term of GH_{cone} gives no contribution to the heat trace.

The last step is to find an expression for $b(R, \theta, R', \theta')$ by simplifying (30). As a first step, we examine the middle term in (30). We have

$$F^{-1}(r, \theta) = 1 - 4r_0 r \cos(\theta + \theta_0) + O(r^2),$$

so in our front face coordinates,

$$(F^{-1} - 1) = -4r_0 T R \cos(\theta + \theta_0) + O(T^2).$$

We use the notation

$$\Delta_{R,\theta} = \partial_R^2 + \frac{1}{R} \partial_R + \frac{1}{R^2} \partial_\theta^2, \quad \nabla_{R,\theta} = (\partial_R, \frac{1}{R} \partial_\theta).$$

With this notation, we have $\nabla_{r,\theta} = T^{-1}\nabla_{R,\theta}$ and $\Delta_{r,\theta} = T^{-2}\Delta_{R,\theta}$, and so

$$\Delta H_{cone} = T^{-2}\Delta_{R,\theta}H_{cone},$$

which by conic scaling is

$$T^{-4}\Delta_{R,\theta}a(R, \theta, R', \theta').$$

Therefore

$$-(F^{-1} - 1)\Delta H_{cone} = 4r_0T^{-3}R\cos(\theta + \theta_0)\Delta_{R,\theta}a(R, \theta, R', \theta'),$$

and $T^3r_0^{-1}$ times this is the contribution that is made to $b(R, \theta, R', \theta')$:

$$(41) \quad -T^3r_0^{-1}(F^{-1} - 1)\Delta H_{cone} = 4R\cos(\theta + \theta_0)\Delta_{R,\theta}a.$$

The other terms in (30) involve derivatives of G , so we must compute these using (36). At ff,

$$\nabla G = -\frac{a_0}{2}\nabla_{R,\theta}(D^2S) + O(T); \quad \Delta G = -\frac{a_0}{2}T^{-1}\Delta_{R,\theta}(D^2S) + O(1).$$

On the other hand, ∂_t lifts to

$$\frac{1}{2}T^{-2}(T\partial_T - R\partial_R - R'\partial_{R'})$$

and so

$$(42) \quad \partial_t G = -\frac{a_0}{4}T^{-1}D^2S + \frac{a_0}{4}T^{-1}(R\partial_R + R'\partial_{R'})(D^2S) + O(1).$$

All in all,

$$(43) \quad \mathcal{L}G = -\frac{a_0}{4}T^{-1}(1 - R\partial_R - R'\partial_{R'} - 2\Delta_{R,\theta})(D^2S) + O(1).$$

Lemma 2. *With $\Delta_{R,\theta}$ and $\nabla_{R,\theta}$ the polar Laplacian/gradient in (R, θ) ,*

$$\Delta_{R,\theta}(D^2) = 4, \quad \Delta_{R,\theta}S = 0, \quad \Delta_{R,\theta}(D^2S) = 8R\cos(\theta + \theta_0), \quad (R\partial_R + R'\partial_{R'})(D^2S) = 3D^2S.$$

Proof. This is a direct computation, using for instance that $\Delta_{R,\theta}R^2 = 4$ and S is harmonic in (R, θ) . \square

Putting all of this together, we obtain

$$(44) \quad b(R, \theta, R', \theta') = \left(\frac{1}{2}D^2S + 4R\cos(\theta + \theta_0) \right) a(R, \theta, R', \theta') \\ + 4R\cos(\theta + \theta_0)\Delta_{R,\theta}a(R, \theta, R', \theta') \\ + \nabla_{R,\theta}(D^2S) \cdot \nabla_{R,\theta}a(R, \theta, R', \theta').$$

For later use, we rewrite the gradient term by using the identity

$$(45) \quad \nabla u \cdot \nabla v = \frac{1}{2}\Delta(uv) - \frac{1}{2}v\Delta u - \frac{1}{2}u\Delta v.$$

This yields, after some computations,

$$\begin{aligned}
(46) \quad b(R, \theta, R', \theta') &= \left(\frac{1}{2}D^2S\right) a(R, \theta, R', \theta') \\
&+ 4R \cos(\theta + \theta_0) \Delta_{R, \theta} a(R, \theta, R', \theta') \\
&+ \frac{1}{2} \Delta_{R, \theta} (D^2S \cdot a(R, \theta, R', \theta')) - \frac{1}{2} (D^2S) \Delta_{R, \theta} a(R, \theta, R', \theta').
\end{aligned}$$

Either (44) or (46) may be used.

3.4.1. The little trace and the convolution. To find the heat trace, we first restrict our kernel to the diagonal to find the so-called “little trace”. By (40), the restriction to the diagonal of the T^{-1} coefficient of GH_{cone} is zero, and therefore the first term in (28) does not contribute to the T^{-1} term of interest. The only contribution comes from the second term in (28), and that contribution is the convolution of the leading orders of the two factors. This convolution is

$$- \int_0^t \int_0^\infty \int_0^{\alpha_0} (t-s)^{-1} a\left(\frac{r}{\sqrt{t-s}}, \theta, \frac{r'}{\sqrt{t-s}}, \theta'\right) s^{-3/2} b\left(\frac{r'}{\sqrt{s}}, \theta', \frac{r''}{\sqrt{s}}, \theta''\right) r' d\theta' dr' ds.$$

Switching to front face variables yields that the term of interest, H_{sub} , is, where \cong denotes having the same heat trace coefficient at order T^{-1} ,

$$\begin{aligned}
(47) \quad H_{\text{sub}} &\cong -T^{-1} \int_0^1 \int_0^\infty \int_0^{\alpha_0} (1-\sigma)^{-1} \sigma^{-3/2} \\
&\times a\left(\frac{R}{\sqrt{1-\sigma}}, \theta, \frac{R'}{\sqrt{1-\sigma}}, \theta'\right) b\left(\frac{R'}{\sqrt{\sigma}}, \theta'', \frac{R''}{\sqrt{\sigma}}, \theta''\right) R' d\theta' dR' d\sigma.
\end{aligned}$$

Setting $(R'', \theta'') = (R, \theta)$ gives the little trace. If you also make the change of variables $\tilde{R}' = R'/\sqrt{\sigma}$, then $R' dR' = \sigma \tilde{R}' d\tilde{R}'$. Then

$$\begin{aligned}
(48) \quad (\text{tr } H_{\text{sub}})(R, \theta) &= -T^{-1} \int_0^1 \int_0^\infty \int_0^{\alpha_0} (1-\sigma)^{-1} \sigma^{-1/2} \\
&\times a\left(\frac{R}{\sqrt{1-\sigma}}, \theta, \frac{\sqrt{\sigma} \tilde{R}'}{\sqrt{1-\sigma}}, \theta'\right) b\left(\tilde{R}', \theta', \frac{R}{\sqrt{\sigma}}, \theta\right) \tilde{R}' d\theta' d\tilde{R}' d\sigma.
\end{aligned}$$

To find the contribution to the heat trace, we first restrict this kernel to the diagonal by setting $(R', \theta') = (R, \theta)$, then we integrate with respect to $r dr d\theta = T^2 R dR d\theta$. By Melrose’s pushforward theorem (see [9]; see also [5, Thm 3.10] for a precise statement), the trace contribution is given by a *renormalized* integral. Writing down this renormalized integral yields (4).

4. THE CASE $\alpha = \pi/2$

Our goal here is to compute through the $t^{1/2}$ term in the short-time expansion of the heat trace of the Laplacian on a domain Ω which is a half-disk of radius 1, with Dirichlet boundary conditions. This heat trace is denoted

$$(49) \quad \text{Tr } e^{-t\Delta_\Omega}.$$

The spectrum of the Dirichlet Laplacian on Ω consists of $\{j_{m,k}^2\}_{m,k \in \mathbb{N}}$, each with multiplicity one. This may be compared to the spectrum of the Dirichlet Laplacian on a disk D , which consists of all of these eigenvalues but with multiplicity two, as well as the set $\{j_{0,k}^2\}_{k \in \mathbb{N}}$, each with multiplicity

one. As a result, if we let

$$(50) \quad H(t) := \sum_{k=1}^{\infty} e^{-tj_{0,k}^2},$$

then

$$(51) \quad \text{Tr } e^{-t\Delta_{\Omega}} = \frac{1}{2} \text{Tr } e^{-t\Delta_D} - \frac{1}{2} H(t).$$

We will take advantage of this.

The heat trace $\text{Tr } e^{-t\Delta_D}$ is well understood and has a full polyhomogeneous asymptotic expansion in half-integer powers as $t \rightarrow 0$. The beginning of this expansion is, from [4],

$$(52) \quad \text{Tr } e^{-t\Delta_D} = \frac{1}{4} t^{-1} - \frac{\sqrt{\pi}}{4} t^{-1/2} + \frac{1}{6} + \frac{\sqrt{\pi}}{128} t^{1/2} + O(t).$$

For $H(t)$, we use the asymptotics of Bessel function zeroes due to McMahon [3, (10.21.19)]:

$$(53) \quad j_{0,k} = (k - \frac{1}{4})\pi + \frac{1}{8(k - \frac{1}{4})\pi} + O(k^{-3}),$$

which imply

$$(54) \quad j_{0,k}^2 = (k - \frac{1}{4})^2 \pi^2 + \frac{1}{4} + O(k^{-2}).$$

Based on this we define the function

$$(55) \quad \tilde{H}(t) = \sum_{k=1}^{\infty} \exp[-t((k - \frac{1}{4})^2 \pi^2 + \frac{1}{4})]$$

and estimate its difference with $H(t)$.

Lemma 3. *With all notation as above, as $t \rightarrow 0$,*

$$(56) \quad H(t) - \tilde{H}(t) = O(t).$$

This is helpful because we also have the following lemma.

Lemma 4. *As $t \rightarrow 0$,*

$$(57) \quad \tilde{H}(t) = \frac{1}{2\sqrt{\pi}} t^{-1/2} + c - \frac{1}{8\sqrt{\pi}} t^{1/2} + O(t).$$

Here c is a constant which is irrelevant to our purposes.

From these two Lemmas and (51) we immediately deduce this proposition.

Proposition 2. *As $t \rightarrow 0$,*

$$(58) \quad \text{Tr } e^{-t\Delta_{\Omega}} = \frac{1}{8} t^{-1} - (\frac{\sqrt{\pi}}{8} + \frac{1}{4\sqrt{\pi}}) t^{-1/2} - \frac{1}{2} c + (\frac{\sqrt{\pi}}{256} + \frac{1}{16\sqrt{\pi}}) t^{1/2} + O(t).$$

Remark. *We could reverse engineer c from the known expansion from [10] but there is no need.*

We are now in a position to state what the contribution of each corner of the half-disk to the $t^{1/2}$ term is. The $t^{1/2}$ term above is the sum of a curvature integral over the boundary and a pair of identical corner contributions. From [4] we see that the $\frac{\sqrt{\pi}}{256}$ is precisely the curvature integral. Dividing the remainder in half shows

Theorem 3. *The contribution to the $t^{1/2}$ coefficient in the heat trace of a half-disk of radius 1 from each of its two corners is*

$$(59) \quad \frac{1}{32\sqrt{\pi}} t^{1/2}.$$

It remains only to prove the Lemmas.

4.1. Proofs of the Lemmas. First we prove Lemma 3.

Proof. Rearrangement of the various expressions gives

$$(60) \quad H(t) - \tilde{H}(t) = e^{-t/4} \sum_{k=1}^{\infty} e^{-t(k-\frac{1}{4})^2 \pi^2} (1 - e^{-t \cdot O(k^{-2})}).$$

Here $O(k^{-2})$ is shorthand for $j_{0,k}^2 - (k - \frac{1}{4})^2 \pi^2 - \frac{1}{4}$, which is bounded in absolute value by Ck^{-2} for some universal constant C . Since $k \geq 1$ and $t \leq 1$ is all that is relevant, we have that for $t \leq 1$, $|1 - e^{-t \cdot O(k^{-2})}| \leq \tilde{C}tk^{-2}$ for a universal constant \tilde{C} . This implies

$$(61) \quad |H(t) - \tilde{H}(t)| \leq \tilde{C}t \sum_{k=1}^{\infty} e^{-t(k-\frac{1}{4})^2 \pi^2} k^{-2}.$$

Estimating each exponential from above by 1 yields that the right-hand side is $O(t)$ as desired. \square

Now we prove Lemma 4.

Proof. First write

$$(62) \quad \tilde{H}(t) = e^{-t/4} \hat{H}(t),$$

where

$$(63) \quad \hat{H}(t) = \sum_{k=1}^{\infty} \exp[-t(k - \frac{1}{4})^2 \pi^2].$$

We know from [NRS] that $\text{Tr } e^{-t\Delta_\Omega}$ has a full polyhomogeneous asymptotic expansion, and therefore by (51), so does $H(t)$. By Lemma 3, $\tilde{H}(t)$ has at least a polyhomogeneous expansion up to $O(t)$, and therefore so does $\hat{H}(t)$. The upshot of this is that one may define a zeta function

$$(64) \quad \hat{\zeta}(s) = \frac{1}{\Gamma(s)} \int_0^\infty \hat{H}(t) t^{s-1} dt,$$

and that this zeta function, initially defined for $\Re(s) \gg 0$, has a meromorphic continuation at least to the half-plane $\Re(s) > -1$. Moreover, by the usual argument (agreement for $\Re(s) \gg 0$),

$$(65) \quad \hat{\zeta}(s) = \sum_{k=1}^{\infty} ((k - \frac{1}{4})\pi)^{-2s}.$$

But now we recognize the zeta function $\hat{\zeta}(s)$ as a form of a Hurwitz zeta function. Specifically,

$$(66) \quad \hat{\zeta}(s) = \pi^{-2s} \zeta\left(2s, \frac{3}{4}\right),$$

where $\zeta(s, a)$ is the Hurwitz zeta function. The function $\zeta(s, a)$ has a single, simple pole at $s = 1$, with residue 1 [3, 25.11(i)]. So $\hat{\zeta}(s)$ has a single, simple pole at $s = 1/2$, and its residue there is $\frac{1}{2\pi}$. Thus the function

$$(67) \quad \int_0^\infty \hat{H}(t) t^{s-1} dt = \Gamma(s) \hat{\zeta}(s)$$

has poles at $s = 1/2$ and also at each non-positive integer; moreover its residue at $s = 1/2$ is $\Gamma(1/2)/(2\pi) = \frac{1}{2\sqrt{\pi}}$.

We can now reverse-engineer the expansion of $\hat{H}(t)$ as $t \rightarrow 0$, which is morally taking an inverse Mellin transform. We already know such a partial expansion exists up to order $O(t)$, and now we know that its coefficients are at orders $t^{-1/2}$, t^0 , and higher integer powers of t which are absorbed in $O(t)$. The coefficient of $t^{-1/2}$ is precisely the residue of $\Gamma(s)\hat{\zeta}(s)$ at $1/2$, which is $\frac{1}{2\sqrt{\pi}}$. So, for some constant c which is not relevant for our purposes,

$$(68) \quad \hat{H}(t) = \frac{1}{2\sqrt{\pi}} t^{-1/2} + c + O(t).$$

Multiplying by $e^{-t/4} \sim 1 - \frac{t}{4} + O(t^2)$,

$$(69) \quad \tilde{H}(t) = \frac{1}{2\sqrt{\pi}} t^{-1/2} + c - \frac{1}{8\sqrt{\pi}} t^{1/2} + O(t),$$

completing the proof. □

REFERENCES

- [1] T. Branson and P. Gilkey. The asymptotics of the Laplacian on a manifold with boundary. *Communications in partial differential equations*, 15(2):245–272, 1990.
- [2] J. Cheeger. Spectral geometry of singular Riemannian spaces. *J. Differential Geom.*, 18(4):575–657, 1983.
- [3] *NIST Digital Library of Mathematical Functions*. <https://dlmf.nist.gov/>, Release 1.2.4 of 2025-03-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [4] A. Enciso and J. Gómez-Serrano. Spectral determination of semi-regular polygons. *Journal of Differential Geometry*, 122(3):399–419, November 2022.
- [5] D. Grieser. Basics of the b-calculus. In *Approaches to Singular Analysis: A Volume of Advances in Partial Differential Equations*, pages 30–84. Springer, 2001.
- [6] D. Grieser. Notes on heat kernel asymptotics. Lecture notes, 2004. Available at <https://web.math.ku.dk/grubb/notes/heat.pdf>.
- [7] H. McKean Jr and I. Singer. Curvature and the eigenvalues of the laplacian. *Journal of Differential Geometry*, 1(1-2):43–69, 1967.
- [8] M. Kac. Can one hear the shape of a drum? *The american mathematical monthly*, 73(4P2):1–23, 1966.
- [9] R. Melrose. *The Atiyah-Patodi-Singer Index Theorem*. A K Peters/CRC Press, 1st edition, 1993.
- [10] M. Nursultanov, J. Rowlett, and D. A. Sher. The heat kernel on curvilinear polygonal domains in surfaces. *Annales mathématiques du Québec*, 49(1):1–61, 2025.
- [11] M. van den Berg and S. Srisatkunarah. Heat equation for a region in \mathbb{R}^2 with a polygonal boundary. *J. London Math. Soc. (2)*, 37(1):119–127, 1988.
- [12] M. van den Berg and S. Srisatkunarah. Heat flow and brownian motion for a region in \mathbb{R}^2 with a polygonal boundary. *Prob. Theory and Related Fields*, 86:41–52, 1990.

CALIFORNIA INSTITUTE OF TECHNOLOGY, THE DIVISION OF PHYSICS, MATHEMATICS AND ASTRONOMY, 1200 E CALIFORNIA BLVD, PASADENA CA 91125

Email address: `looi@caltech.edu`

DEPAUL UNIVERSITY, DEPARTMENT OF MATHEMATICAL SCIENCES, 2320 N. KENMORE AVE., CHICAGO IL 60614, USA

Email address: `dsher@depaul.edu`