

# TOTAL 2-CUT COMPLEXES OF POWERS OF CYCLE GRAPHS AND CARTESIAN PRODUCTS OF CERTAIN GRAPHS

PRATIKSHA CHAUHAN, SAMIR SHUKLA, AND KUMAR VINAYAK

**ABSTRACT.** For a positive integer  $k$ , the *total  $k$ -cut complex* of a graph  $G$ , denoted as  $\Delta_k^t(G)$ , is the simplicial complex whose facets are  $\sigma \subseteq V(G)$  such that  $|\sigma| = |V(G)| - k$  and the induced subgraph  $G[V(G) \setminus \sigma]$  does not contain any edge. These complexes were introduced by Bayer et al. in [4] in connection with commutative algebra. In the same paper, they studied the homotopy types of these complexes for various families of graphs, including cycle graphs  $C_n$ , squared cycle graphs  $C_n^2$ , and Cartesian products of complete graphs and path graphs  $K_m \square P_2$  and  $K_2 \square P_n$ .

In this article, we extend the work of Bayer et al. for these families of graphs. We focus on the complexes  $\Delta_2^t(G)$  and determine the homotopy types of these complexes for three classes of graphs: (i)  $p$ -th powers of cycle graphs  $C_n^p$  (ii)  $K_m \square P_n$  and (iii)  $K_m \square C_n$ . Using discrete Morse theory, we show that these complexes are homotopy equivalent to wedges of spheres. We also give the number and dimension of spheres appearing in the homotopy type.

Our result on powers of cycle graphs  $C_n^p$  proves a conjecture of Shen et al. about the homotopy type of the complexes  $\Delta_2^t(C_n^p)$ .

## 1. INTRODUCTION

All graphs are assumed to be finite and simple, meaning that they have no loops or multiple edges. The vertex set and edge set of a graph  $G$  are denoted by  $V(G)$  and  $E(G)$ , respectively.

Graph complexes are simplicial complexes constructed from graphs by encoding their combinatorial structure into higher-dimensional simplices. In topological combinatorics, graph complexes are one of the main objects of study. Various topological invariants of these complexes, such as homotopy type, connectivity, homology, and their refinements, provide powerful tools for understanding combinatorial properties of graphs. A classical and one of the most celebrated examples of graph complexes is the neighbourhood complex, introduced by L. Lovász in 1978 [29]. Lovász gave a lower bound for the chromatic number of a graph in terms of the topological connectivity of its neighbourhood complex, and using this bound, he proved the Kneser conjecture [27].

Graph complexes are prevalent across the literature, and have applications in various areas of mathematics, engineering, and data analysis. They are central to graph theory and combinatorics, where the topology of these complexes is related to the combinatorics of the underlying graphs (see [1, 13, 31, 32, 33, 37]), and have found many uses in data analysis and machine learning through simplicial representation of networks and topological data analysis (see [9, 17, 20]). Graph complexes also appear extensively in commutative algebra, where they are intricately related to monomial ideals and resolutions (see [14, 15, 24, 35, 36, 40, 42]). In representation theory, they are used in the study of symmetric group actions on simplicial complexes and associated homological representations (see [2, 5, 8]). Graph complexes have connections to geometric group theory, where they serve as combinatorial models for classifying spaces and group actions (see [21, 22, 43]), and also have applications in random topology (see [6, 26, 34, 39]). For comprehensive treatments of graph complexes, we refer the reader to the books [25], [28] and [30].

Recently, Bayer et al. in [4] and [3] introduced two new families of graph complexes: total cut complexes and cut complexes. For  $k \geq 1$ , the *total  $k$ -cut complex* of a graph  $G$ , denoted as  $\Delta_k^t(G)$ , is the simplicial complex whose facets (maximal simplices) are  $\sigma \subseteq V(G)$  such that  $|\sigma| = |V(G)| - k$  (where  $|\cdot|$  denotes cardinality) and the induced subgraph  $G[V(G) \setminus \sigma]$  does not contain any edge (see

2020 *Mathematics Subject Classification.* Primary: 55P10, 57M15, 55U05, Secondary: 05C69, 05E45, 57Q70.

*Key words and phrases.* Total cut complex, Cut complex, Powers of cycle graphs, Cartesian product of graphs, Circulant graphs, Homotopy, Discrete Morse theory.

Example 1.1). The  $k$ -cut complex of a graph  $G$ , denoted as  $\Delta_k(G)$ , is the simplicial complex whose facets are  $\sigma \subseteq V(G)$  such that  $|\sigma| = |V(G)| - k$  and the induced subgraph  $G[V(G) \setminus \sigma]$  is disconnected. One of the main motivations behind these complexes is a famous theorem of Ralf Fröberg connecting commutative algebra and graph theory through topology (see [19, Theorem 1], [16, p. 274]).

It can be easily observed that  $\Delta_2^t(G) = \Delta_2(G)$ . In fact, it is mentioned in [4] that both the complexes  $\Delta_k^t(G)$  and  $\Delta_k(G)$  can be considered as a generalization of the complex  $\Delta_2^t(G)$ .

**Example 1.1.** Let  $G$  be the graph given in Figure 1(A). Then the set of maximal simplices of  $\Delta_2^t(G)$  is  $\{\{1, 2, 3\}, \{1, 3, 5\}, \{2, 3, 4\}, \{2, 4, 5\}\}$  (see Figure 1(B)).



FIGURE 1

In this article, we consider the total 2-cut complexes (hence 2-cut complexes) of three families of graphs: (i) powers of cycle graphs, (ii) Cartesian product of complete graphs and path graphs, and (iii) Cartesian product of complete graphs and cycle graphs.

For a graph  $G$  and a positive integer  $p$ , the  $p$ -th power graph of  $G$  is a graph  $G^p$  with  $V(G^p) = V(G)$  and  $\{u, v\} \in E(G^p)$  if and only if there exists a path between  $u$  and  $v$  of length at most  $p$  in  $G$  (here, the length of a path is the number of edges in the path). Clearly,  $G^1 = G$ .

Let  $\Gamma$  be a group and let  $S \subset \Gamma$  be a subset not containing the identity element. The *Cayley graph* of  $\Gamma$  with respect to  $S$  is the graph  $G(\Gamma, S)$  having vertex set  $\Gamma$ , and  $\{u, v\} \in E(G(\Gamma, S))$  if and only if  $uv^{-1} \in S \cup S^{-1}$ , where  $S^{-1} = \{x^{-1} \mid x \in S\}$ . For  $n \geq 2$  and  $S \subset \{1, 2, \dots, n-1\}$ , the *circulant graph*  $C_n(S)$  is a Cayley graph of  $\mathbb{Z}_n$ , the cyclic group of order  $n$ .

For  $n \geq 3$ , let  $C_n$  denote the cycle graph on  $n$  vertices  $\{0, 1, \dots, n-1\}$ . Observe that for  $p \geq 1$ ,  $V(C_n^p) = V(C_n) = \{0, 1, \dots, n-1\}$  and  $E(C_n^p) = \{\{i, i+j \pmod{n}\} \mid 0 \leq i \leq n-1 \text{ and } 1 \leq j \leq p\}$ . It is easy to check that for  $n \geq 2p+1$ ,  $C_n^p$  is a circulant graph  $C_n(\{1, 2, \dots, p\})$ .

In [4], the authors studied  $\Delta_k^t(C_n)$  (Theorem 4.15) and  $\Delta_2^t(C_n^2)$  (Proposition 4.19). They proved that these complexes (if nonvoid) are homotopy equivalent to wedges of spheres. In [38], the authors determined the homotopy type of  $\Delta_3^t(C_n^2)$  (Theorem 1.5) and, for  $k \geq 3$ , the homotopy type of  $\Delta_k^t(C_n^2)$  when  $n = 3k+1, 3k+2$  (Theorem 1.6).

The  $k$ -cut complexes of powers of cycle graphs have been studied by many authors. In [3, Proposition 7.11], the authors proved that for  $k \geq 3$ ,  $\Delta_k(C_n)$  is shellable<sup>1</sup>, and in [12, Theorem 1.4], it has been proved that  $\Delta_3(C_n^2)$  is shellable for  $n \geq k+6$ . In [11], the authors proved that  $\Delta_3(C_n^p)$  is shellable for  $n \geq 6p-3$ . Hence all of these complexes are wedges of spheres up to homotopy type.

For general  $k \geq 2$  and  $p \geq 2$ , the topology of the complexes  $\Delta_k^t(C_n^p)$  and  $\Delta_k(C_n^p)$  is not known. For  $k=2$ , Shen et al. made the following conjecture about the homotopy type of the complexes  $\Delta_k^t(C_n^p)$ .

**Conjecture 1.2** ([38, Conjecture 5.1]). *For  $p \geq 3$  and  $n \geq 3p+1$ ,  $\Delta_2^t(C_n^p) \simeq \mathbb{S}^{n-4}$ .*

One of the main results of this article is that Conjecture 1.2 is true. It is easy to check that for  $n < 2p+2$ , the complex  $\Delta_2^t(C_n^p)$  is void. For  $n \geq 2p+2$ , we prove the following.

**Theorem 1.3** (Theorem 3.10). *Let  $n \geq 2p+2$ . Then*

$$\Delta_2^t(C_n^p) \simeq \begin{cases} \mathbb{S}^{\frac{n-4}{2}} & \text{if } n = 2p+2, \\ \mathbb{S}^{n-4} & \text{if } n \geq 3p+1. \end{cases}$$

<sup>1</sup>A simplicial complex  $\Delta$  is called *shellable* if its facets can be arranged in a linear order  $F_1, F_2, \dots, F_t$  in such a way that for all  $2 \leq j \leq t$ , the subcomplex  $(\bigcup_{i=1}^{j-1} F_i) \cap F_j$  is pure and of dimension  $|F_j| - 2$ .

For  $2p + 3 \leq n \leq 3p$ , using SageMath, we computed the homology groups of the complex  $\Delta_2^t(C_n^p)$  for certain values of  $n$  and  $p$ . Based on our calculation, we conjecture about the homotopy type of  $\Delta_2^t(C_n^p)$  for all  $2p + 3 \leq n \leq 3p$  (see Conjecture 4.1 of Section 4).

The *Cartesian product* of graphs  $G$  and  $H$ , denoted by  $G \square H$ , is the graph where  $V(G \square H) = V(G) \times V(H)$  and  $\{(u_1, v_1), (u_2, v_2)\} \in E(G \square H)$  if and only if either  $u_1 = u_2$  and  $\{v_1, v_2\} \in E(H)$ , or  $v_1 = v_2$  and  $\{u_1, u_2\} \in E(G)$ .

For a positive integer  $n$ , let  $K_n$  and  $P_n$  denote the complete graph and the path graph on  $n$  vertices, respectively. In [4], it is proved that  $\Delta_2^t(K_m \square P_2)$  (Theorem 4.8) and  $\Delta_2^t(K_2 \square P_n)$  (Theorem 4.16) are homotopy equivalent to wedges of spheres. Further, in [10, Theorem 3.2], the authors proved that for  $2 \leq k \leq n$ ,  $\Delta_k^t(K_2 \square P_n)$  is homotopy equivalent to a wedge of spheres. In this article, we extend the results of [4] to all  $K_m$  and all  $P_n$ , and prove the following.

**Theorem 1.4** (Theorem 3.12). *For  $m, n \geq 2$ ,  $\Delta_2^t(K_m \square P_n) \simeq V_{(m-1)(n-1)} \mathbb{S}^{mn-4}$ .*

We also investigate the total 2-cut complexes of  $K_m \square C_n$  and determine their homotopy types.

**Theorem 1.5** (Theorem 3.14). *For  $m \geq 2$  and  $n \geq 4$ ,  $\Delta_2^t(K_m \square C_n) \simeq V_{n(m-1)+1} \mathbb{S}^{mn-4}$ .*

Beyond the total 2-cut complexes, we investigate the total 3-cut and 3-cut complexes of  $K_m \square P_n$  and  $K_m \square C_n$ . We computed their homology groups for small values of  $m$  and  $n$  using SageMath. These computations lead us to propose conjectures about their homotopy types (see Conjectures 4.3 to 4.6).

This paper is organized as follows: In Section 2, we recall the necessary preliminaries related to graph theory, simplicial complexes and discrete Morse theory. Section 3 is divided into three subsections (Sections 3.1 to 3.3) where we prove Theorems 1.3 to 1.5, respectively. Finally, Section 4 proposes several conjectures and questions based on our SageMath computations.

## 2. PRELIMINARIES

This section presents some basic definitions and results used in this article.

**2.1. Graph.** A *graph*  $G$  is a pair  $(V(G), E(G))$ , where  $V(G)$  is its vertex set and  $E(G) \subseteq \binom{V(G)}{2}$  is the edge set. For any  $u, v \in V(G)$ , we say that  $u$  and  $v$  are adjacent if  $\{u, v\} \in E(G)$ . We write  $u \sim v$  for adjacency and  $u \not\sim v$  for non-adjacency. A *subgraph*  $H$  of  $G$  is a graph with  $V(H) \subseteq V(G)$  and  $E(H) \subseteq E(G)$ . For a subset  $U \subseteq V(G)$ , the *induced subgraph*  $G[U]$  is the subgraph with  $V(G[U]) = U$  and  $E(G[U]) = \{\{a, b\} \in E(G) \mid a, b \in U\}$ .

For  $u, v \in V(G)$ , a *path* from  $u$  to  $v$  is a sequence of distinct vertices  $u = v_0, v_1, \dots, v_n = v$  such that  $v_i \sim v_{i+1}$  for all  $0 \leq i \leq n-1$ . The *length* of a path is the number of edges in the path. A graph is *connected* if there exists a path between each pair of its vertices; otherwise, it is *disconnected*.

The *complete graph*  $K_n$  on  $n$  vertices is the graph in which any two distinct vertices are adjacent. For  $n \geq 1$ , the *path graph*  $P_n$  is the graph with  $V(P_n) = \{0, 1, \dots, n-1\}$  and  $E(P_n) = \{\{i, i+1\} \mid 0 \leq i \leq n-2\}$ . For  $n \geq 3$ , the *cycle graph*  $C_n$  is the graph with  $V(C_n) = \{0, 1, \dots, n-1\}$  and  $E(C_n) = \{\{i, i+1\} \mid 0 \leq i \leq n-2\} \cup \{\{0, n-1\}\}$ .

We refer the reader to [7] and [41] for more details about the graph terminologies used in this article.

**2.2. Simplicial complex.** A *finite abstract simplicial complex*  $\Delta$  is a collection of finite sets such that if  $\tau \in \Delta$  and  $\sigma \subset \tau$ , then  $\sigma \in \Delta$ . The elements of  $\Delta$  are called *simplices* of  $\Delta$ . If  $\sigma \subset \tau$ , we say that  $\sigma$  is a *face* of  $\tau$ . The *dimension* of a simplex  $\sigma$  is equal to  $|\sigma| - 1$ . The *dimension* of an abstract simplicial complex is the maximum of the dimensions of its simplices. If a simplex has dimension  $d$ , it is said to be *d-dimensional*. The 0-dimensional simplices are called *vertices* of  $\Delta$ . An abstract simplicial complex which is an empty collection of sets is called the *void* abstract simplicial complex, and is denoted by  $\emptyset$ . A simplex that is not a face of any other simplex is called a *maximal simplex* or *facet*. A *subcomplex*  $\Delta'$  of  $\Delta$  is a simplicial complex such that  $\sigma \in \Delta'$  implies  $\sigma \in \Delta$ .

In this article, we consider any simplicial complex as a topological space, namely, its geometric realization (see [28] for details). For terminologies related to algebraic topology, we refer to [23].

**2.3. Discrete Morse theory.** Discrete Morse theory was introduced by Robin Forman [18] in 1998 as a combinatorial adaptation of classical Morse theory. For more details, we refer the reader to [25, Chapter 4] and [28, Chapter 11], which serve as the primary sources for the definitions and results in this section.

Throughout this section, let  $X$  be a set and let  $\Sigma$  be a finite family of finite subsets of  $X$ .

**Definition 2.1.** A *matching* on  $\Sigma$  is a family  $\mathcal{M}$  of pairs  $\{\sigma, \tau\}$  with  $\sigma, \tau \in \Sigma$  such that no set is contained in more than one pair in  $\mathcal{M}$ .

A set  $\sigma \in \Sigma$  is said to be *matched* in  $\mathcal{M}$  if it is contained in some pair of  $\mathcal{M}$ ; for convenience, we write  $\sigma \in \mathcal{M}$ . Otherwise,  $\sigma$  is said to be *critical* or *unmatched* with respect to  $\mathcal{M}$ , and we write  $\sigma \notin \mathcal{M}$ .

**Definition 2.2.** A matching  $\mathcal{M}$  is said to be *partial* if  $\{\sigma, \tau\} \in \mathcal{M}$  implies either  $\sigma \prec \tau$  (i.e.,  $\sigma \subset \tau$  and no  $\rho \in \Sigma$  satisfies  $\sigma \subset \rho \subset \tau$ ), or  $\sigma \succ \tau$ .

Note that  $\mathcal{M}$  is a partial matching on  $\Sigma$  if and only if there exists  $\Sigma' \subset \Sigma$  and an injective map  $\mu : \Sigma' \rightarrow \Sigma \setminus \Sigma'$  such that  $\mu(\sigma) \succ \sigma$  for all  $\sigma \in \Sigma'$ .

**Definition 2.3.** A partial matching  $\mathcal{M}$  on  $\Sigma$  is called *acyclic* if there does not exist a cycle

$$\mu(\sigma_1) \succ \sigma_1 \prec \mu(\sigma_2) \succ \sigma_2 \prec \mu(\sigma_3) \succ \sigma_3 \prec \cdots \prec \mu(\sigma_t) \succ \sigma_t \prec \mu(\sigma_1), \quad t \geq 2.$$

**Definition 2.4.** Let  $x \in X$ . A matching  $\mathcal{M}_x$  is called an *element matching* on  $\Sigma$  using  $x$  if every pair in  $\mathcal{M}_x$  is of the form  $\{\sigma \setminus \{x\}, \sigma \cup \{x\}\}$  for some  $\sigma \in \Sigma$ .

It is easy to observe that any element matching is a partial matching.

**Lemma 2.5** ([25, Lemma 4.1]). *Let  $x \in X$ . Define*

$$\mathcal{M}_x := \{ \{ \sigma \setminus \{x\}, \sigma \cup \{x\} \} \mid \sigma \setminus \{x\}, \sigma \cup \{x\} \in \Sigma \}.$$

*Let  $\mathcal{M}'$  be an acyclic matching on  $\Sigma' := \{ \sigma \in \Sigma \mid \sigma \notin \mathcal{M}_x \}$ . Then  $\mathcal{M} := \mathcal{M}_x \cup \mathcal{M}'$  is an acyclic matching on  $\Sigma$ .*

Let  $\{x_0, x_1, \dots, x_{n-1}\}$  be a subset of  $X$ . We define a sequence of element matchings  $\mathcal{M}_{x_0}, \mathcal{M}_{x_1}, \dots, \mathcal{M}_{x_{n-1}}$  on  $\Sigma$  as follows.

Let  $\mathcal{C}_0 := \Sigma$ . For each  $0 \leq i \leq n-1$ , define

$$\begin{aligned} \mathcal{M}_{x_i} &:= \{ \{ \sigma \setminus \{x_i\}, \sigma \cup \{x_i\} \} \mid \sigma \setminus \{x_i\}, \sigma \cup \{x_i\} \in \mathcal{C}_i \}; \\ \mathcal{C}_{i+1} &:= \{ \sigma \in \mathcal{C}_i \mid \sigma \notin \mathcal{M}_{x_i} \}. \end{aligned} \tag{1}$$

Observe that  $\mathcal{C}_{i+1} = \{ \sigma \in \Sigma \mid \sigma \notin \mathcal{M}_{x_j} \text{ for all } 0 \leq j \leq i \}$ . Moreover, the matchings  $\mathcal{M}_{x_0}, \mathcal{M}_{x_1}, \dots, \mathcal{M}_{x_{n-1}}$  are pairwise disjoint.

**Remark 2.6.** *Let  $0 \leq i \leq n-1$ . Then the following results hold:*

- (i)  $\mathcal{C}_{i+1} \subseteq \mathcal{C}_i \subseteq \mathcal{C}_{i-1} \subseteq \dots \subseteq \mathcal{C}_1 \subseteq \mathcal{C}_0 = \Sigma$ .
- (ii) For any  $\sigma \in \mathcal{C}_i$ ,  $\sigma \in \mathcal{C}_{i+1}$  if and only if  $\sigma \setminus \{i\} \notin \mathcal{C}_i$  or  $\sigma \cup \{i\} \notin \mathcal{C}_i$ .
- (iii) Let  $\sigma \in \mathcal{C}_i$  and let  $i \leq j \leq n-1$ . If  $\sigma \setminus \{r\} \notin \mathcal{C}_i$  or  $\sigma \cup \{r\} \notin \mathcal{C}_i$  for all  $i \leq r \leq j$ , then  $\sigma \in \mathcal{C}_{j+1}$ .

To define a matching on a simplicial complex, we consider it simply as the set of its simplices (or faces).

**Theorem 2.7** ([25, Theorem 4.14]). *Let  $\Delta$  be a simplicial complex and let  $\mathcal{M}$  be an acyclic matching on  $\Delta$  such that the empty set is not critical. Then  $\Delta$  is homotopy equivalent to a cell complex with one cell of dimension  $d \geq 0$  for each critical face of  $\Delta$  of dimension  $d$  plus one additional 0-cell.*

Theorem 2.7 implies the following result.

**Corollary 2.8** ([25, Theorem 4.8]). *If an acyclic matching on a simplicial complex  $\Delta$  has all its critical faces in the same dimension  $d$ , then  $\Delta$  is homotopy equivalent to a wedge of spheres of dimension  $d$ .*

## 3. PROOFS

In this section, we determine the homotopy type of  $\Delta_2^t(C_n^p)$  (Theorem 1.3),  $\Delta_2^t(K_m \square P_n)$  (Theorem 1.4), and  $\Delta_2^t(K_m \square C_n)$  (Theorem 1.5).

Let  $G$  be a graph, and let  $u, v \in V(G)$ . Recall that the notation  $u \sim v$  denotes that the vertices  $u$  and  $v$  are adjacent in  $G$ , and  $u \not\sim v$  denotes that  $u$  and  $v$  are not adjacent. Let  $A \subseteq V(G)$ . For  $B \subseteq A$ , we say that  $B$  is a disconnected 2-set in  $A$  if  $B = \{u, v\}$  for some  $u \not\sim v$ . Throughout this article,  $A^c$  denotes the complement of  $A$  in  $V(G)$ .

We begin by establishing some basic results that will be used later. The following remark is a direct implication of the definition of the total 2-cut complex.

**Remark 3.1.** *Let  $G$  be a graph, and let  $\sigma \subseteq V(G)$ . Then  $\sigma \in \Delta_2^t(G)$  if and only if there exist  $u, v \in \sigma^c$  such that  $u \not\sim v$ , i.e.,  $\{u, v\}$  is a disconnected 2-set in  $\sigma^c$ . Equivalently,  $\sigma \notin \Delta_2^t(G)$  if and only if the induced subgraph  $G[\sigma^c]$  is a complete graph.*

**Proposition 3.2.** *Let  $G$  be a graph and  $u, v \in V(G)$  with  $u \neq v$ . Define*

$$\begin{aligned}\mathcal{M}_v &:= \{\{\sigma \setminus \{v\}, \sigma \cup \{v\}\} \mid \sigma \setminus \{v\}, \sigma \cup \{v\} \in \Delta_2^t(G)\}; \\ \mathcal{C} &:= \{\sigma \in \Delta_2^t(G) \mid \sigma \notin \mathcal{M}_v\}.\end{aligned}$$

*Then, for any  $\sigma \in \Delta_2^t(G)$ , the following statements hold:*

- (i)  $\sigma \in \mathcal{C}$  if and only if  $\sigma \cup \{v\} \notin \Delta_2^t(G)$ . Equivalently,  $\sigma \in \mathcal{C}$  if and only if every disconnected 2-set in  $\sigma^c$  contains  $v$ .
- (ii) If  $\sigma \in \mathcal{C}$  and  $\sigma \cup \{u\} \in \Delta_2^t(G)$ , then  $\sigma \cup \{u\} \in \mathcal{C}$ .
- (iii) If  $\sigma \in \mathcal{C}$  and  $u \sim w$  for all  $w \in (\sigma \cup \{v\})^c$ , then  $\sigma \setminus \{u\} \in \mathcal{C}$ .

*Proof.* (i) Since  $\Delta_2^t(G)$  is a simplicial complex,  $\sigma \cup \{v\} \in \Delta_2^t(G)$  implies  $\sigma \setminus \{v\} \in \Delta_2^t(G)$ . It follows that  $\mathcal{M}_v = \{\{\sigma \setminus \{v\}, \sigma \cup \{v\}\} \mid \sigma \cup \{v\} \in \Delta_2^t(G)\}$ . By the definition of  $\mathcal{C}$ ,  $\sigma \in \mathcal{C}$  if and only if  $\sigma \notin \mathcal{M}_v$ , which means that  $\{\sigma \setminus \{v\}, \sigma \cup \{v\}\} \notin \mathcal{M}_v$ . Since  $\{\sigma \setminus \{v\}, \sigma \cup \{v\}\} \notin \mathcal{M}_v$  if and only if  $\sigma \cup \{v\} \notin \Delta_2^t(G)$ , it follows that  $\sigma \in \mathcal{C}$  if and only if  $\sigma \cup \{v\} \notin \Delta_2^t(G)$ .

By Remark 3.1,  $\sigma \in \Delta_2^t(G)$  and  $\sigma \cup \{v\} \notin \Delta_2^t(G)$  if and only if every disconnected 2-set in  $\sigma^c$  contains  $v$ . Therefore,  $\sigma \in \mathcal{C}$  if and only if every disconnected 2-set in  $\sigma^c$  contains  $v$ .

- (ii) Let  $\sigma \in \mathcal{C}$  and  $\sigma \cup \{u\} \in \Delta_2^t(G)$ . If  $u \in \sigma$ , then clearly  $\sigma \cup \{u\} = \sigma \in \mathcal{C}$ . Hence, assume that  $u \notin \sigma$ . By (i),  $\sigma \in \mathcal{C}$  implies  $\sigma \cup \{v\} \notin \Delta_2^t(G)$ . Hence  $(\sigma \cup \{u\}) \cup \{v\} \notin \Delta_2^t(G)$ . Therefore,  $\sigma \cup \{u\} \in \mathcal{C}$  by (i).
- (iii) Let  $\sigma \in \mathcal{C}$ . Then  $\sigma \cup \{v\} \notin \Delta_2^t(G)$  by (i). By Remark 3.1,  $u_1 \sim u_2$  for all  $u_1, u_2 \in (\sigma \cup \{v\})^c$ . Now, assume further that  $u \sim w$  for all  $w \in (\sigma \cup \{v\})^c$ . Then  $v_1 \sim v_2$  for all  $v_1, v_2 \in ((\sigma \setminus \{u\}) \cup \{v\})^c$ . By Remark 3.1,  $(\sigma \setminus \{u\}) \cup \{v\} \notin \Delta_2^t(G)$ . Therefore,  $\sigma \setminus \{u\} \in \mathcal{C}$  by (i).  $\square$

To prove Theorems 1.3 to 1.5, we construct acyclic matchings on the total 2-cut complex  $\Delta_2^t(G)$  of the graph  $G$  under consideration. This is done by selecting an appropriate subset of vertices and defining a sequence of element matchings as in Equation (1). Our aim is to characterize the critical faces with respect to the matching. Then, using Theorem 2.7 and Corollary 2.8, we determine the homotopy type of  $\Delta_2^t(G)$ .

**3.1. Powers of cycle graphs.** In this section, we prove Theorem 1.3. We assume throughout that  $p \geq 1$  is an integer. Recall that the  $p$ -th power of a cycle graph  $C_n^p$  is the graph on the vertex set  $V(C_n^p) = \{0, 1, \dots, n-1\}$ , where for any distinct vertices  $u, v \in V(C_n^p)$ , we have  $u \sim v$  in  $C_n^p$  if and only if  $v \equiv u \pm t \pmod{n}$  for some  $1 \leq t \leq p$ .

Before proving Theorem 1.3, we establish several key results. We begin by defining a sequence of element matchings  $\mathcal{M}_0, \mathcal{M}_1, \dots, \mathcal{M}_{n-1}$  on  $\Delta_2^t(C_n^p)$  and the sets  $\mathcal{C}_0, \mathcal{C}_1, \dots, \mathcal{C}_n$ , using vertices  $0, 1, \dots, n-1$ , as in Equation (1). By Lemma 2.5,  $\mathcal{M} := \bigsqcup_{i=0}^{n-1} \mathcal{M}_i$  is an acyclic matching on  $\Delta_2^t(C_n^p)$ . To proceed, we characterize the sets  $\mathcal{C}_{i+1}$  for  $0 \leq i \leq n-1$ , thus identifying the critical faces with respect to the matching  $\mathcal{M}$ .

Note that  $\mathcal{C}_0 = \Delta_2^t(C_n^p)$  and  $\mathcal{C}_1 = \{\sigma \in \Delta_2^t(C_n^p) \mid \sigma \notin \mathcal{M}_0\}$ . To characterize  $\mathcal{C}_1$ , we apply Proposition 3.2 with  $v = 0$ . We characterize  $\mathcal{C}_{i+1}$  inductively in Proposition 3.3 for all  $1 \leq i \leq p$ . For each

$1 \leq i \leq p$ , we define a subset  $\mathcal{X}_i \subseteq V(C_n^p)$  as follows:

$$\mathcal{X}_i := \{i + p + 1, i + p + 2, \dots, i + n - p - 1\}.$$

Clearly, for  $n \geq 2p + 2$  and for all  $1 \leq i \leq p$ , we have  $\mathcal{X}_i \neq \emptyset$ .

**Proposition 3.3.** *Let  $n \geq 2p + 2$  and  $1 \leq i \leq p$ . If  $\sigma \in \mathcal{C}_i$ , then  $\sigma \in \mathcal{C}_{i+1}$  if and only if  $\{1, 2, \dots, i\} \subseteq \sigma$  and  $\mathcal{X}_i \not\subseteq \sigma$ .*

*Proof.* The proof proceeds by induction on  $i$  for  $1 \leq i \leq p$ . We first prove the base case  $i = 1$ . Let  $\sigma \in \mathcal{C}_1$ . It suffices to show that  $\sigma \in \mathcal{C}_2$  if and only if  $1 \in \sigma$  and  $\mathcal{X}_1 \not\subseteq \sigma$ .

- Suppose  $\sigma \in \mathcal{C}_2$ . We first show that  $1 \in \sigma$ . Assume, for contradiction, that  $1 \notin \sigma$ . Since  $\sigma \in \mathcal{C}_1 \subseteq \Delta_2^t(C_n^p)$ , there exists a disconnected 2-set in  $\sigma^c$  by Remark 3.1. Moreover, by Proposition 3.2 (i), every disconnected 2-set in  $\sigma^c$  contains 0. Hence, there exists  $v \in V(C_n^p)$  such that  $v \approx 0$  and  $\{0, v\} \subseteq \sigma^c$ . Since  $0 \sim 1$ , we get  $v \neq 1$ . Therefore,  $\{0, v\} \subseteq (\sigma \cup \{1\})^c$ , and by Remark 3.1,  $\sigma \cup \{1\} \in \Delta_2^t(C_n^p)$ . By Proposition 3.2 (ii),  $\sigma \cup \{1\} \in \mathcal{C}_1$ . Further, since  $\sigma \setminus \{1\} = \sigma \in \mathcal{C}_1$ , it follows by Remark 2.6 (ii) that  $\sigma \notin \mathcal{C}_2$ , a contradiction. Thus,  $1 \in \sigma$ .

Now, we show that  $\mathcal{X}_1 \not\subseteq \sigma$ . Suppose  $\mathcal{X}_1 \subseteq \sigma$ . Then  $\mathcal{X}_1 \subseteq \sigma \cup \{0\}$ . By the definition of  $C_n^p$ ,  $1 \approx u$  if and only if  $u \in \{p + 2, p + 3, \dots, n - p\} = \mathcal{X}_1$ . Hence  $1 \sim v$  for all  $v \in (\sigma \cup \{0\})^c$ . By Proposition 3.2 (iii),  $\sigma \setminus \{1\} \in \mathcal{C}_1$ . Since  $\sigma \cup \{1\} = \sigma \in \mathcal{C}_1$ , we get  $\sigma \notin \mathcal{C}_2$  by Remark 2.6 (ii), a contradiction. Therefore,  $\mathcal{X}_1 \not\subseteq \sigma$ .

- Conversely, assume that  $1 \in \sigma$  and  $\mathcal{X}_1 \not\subseteq \sigma$ . Since  $1 \approx u$  if and only if  $u \in \mathcal{X}_1$ , there exists  $v \in \sigma^c$  such that  $1 \approx v$ . Moreover, since  $1 \sim 0$ , we have  $v \neq 0$ . It follows that  $\{1, v\}$  is a disconnected 2-set in  $((\sigma \setminus \{1\}) \cup \{0\})^c$ . By Remark 3.1,  $(\sigma \setminus \{1\}) \cup \{0\} \in \Delta_2^t(C_n^p)$ , and hence  $\sigma \setminus \{1\} \notin \mathcal{C}_1$  by Proposition 3.2 (i). Since  $\sigma \in \mathcal{C}_1$ , we conclude that  $\sigma \in \mathcal{C}_2$  by Remark 2.6 (ii).

Therefore, the result holds for  $i = 1$ .

Let  $1 < i \leq p$  and assume the result holds for all  $1 \leq i' < i$ ; that is, if  $\sigma \in \mathcal{C}_{i'}$ , then  $\sigma \in \mathcal{C}_{i'+1}$  if and only if  $\{1, 2, \dots, i'\} \subseteq \sigma$  and  $\mathcal{X}_{i'} \not\subseteq \sigma$ . We now prove the result for  $i$ . Let  $\sigma \in \mathcal{C}_i$ . We show that  $\sigma \in \mathcal{C}_{i+1}$  if and only if  $\{1, 2, \dots, i\} \subseteq \sigma$  and  $\mathcal{X}_i \not\subseteq \sigma$ .

- Suppose  $\sigma \in \mathcal{C}_{i+1}$ . By Remark 2.6 (i),  $\mathcal{C}_{i+1} \subseteq \mathcal{C}_i \subseteq \dots \subseteq \mathcal{C}_1$ . Hence  $\sigma \in \mathcal{C}_j$  and  $\sigma \in \mathcal{C}_{j+1}$  for all  $1 \leq j < i$ . By the induction hypothesis, for each such  $j$ , we have  $\{1, 2, \dots, j\} \subseteq \sigma$  and  $\mathcal{X}_j \not\subseteq \sigma$ .

We first prove that  $\{1, 2, \dots, i\} \subseteq \sigma$ . Since  $\{1, 2, \dots, j\} \subseteq \sigma$  for all  $1 \leq j < i$ , we have  $\{1, 2, \dots, i - 1\} \subseteq \sigma$ .

Suppose  $i \notin \sigma$ . Since  $\sigma \in \mathcal{C}_1$ , every disconnected 2-set in  $\sigma^c$  contains 0 by Proposition 3.2 (i). Moreover,  $1 < i \leq p$  implies that  $0 \sim i$ . It follows that there exists  $v \in \sigma^c \setminus \{i\}$  such that  $v \approx 0$  and  $\{0, v\} \subseteq (\sigma \cup \{i\})^c$ . Hence  $\sigma \cup \{i\} \in \Delta_2^t(C_n^p)$  by Remark 3.1. By Proposition 3.2 (ii),  $\sigma \in \mathcal{C}_1$  and  $i \neq 0$  implies that  $\sigma \cup \{i\} \in \mathcal{C}_1$ .

Since  $\{1, 2, \dots, j\} \subseteq \sigma$  for all  $1 \leq j < i$ , we get  $\{1, 2, \dots, j\} \subseteq \sigma \cup \{i\}$  for all  $1 \leq j < i$ . Note that  $n \geq 2p + 2$  implies  $p + 1 \leq n - p - 1$ . Further, since  $1 < i \leq p$ , for any  $1 \leq j < i$ , we have  $1 < i \leq p < j + p + 1 \leq j + n - p - 1 < i + n - p - 1 \leq n - 1$ . This gives  $i \notin \mathcal{X}_j$  for all  $1 \leq j < i$ . Therefore,  $\mathcal{X}_j \not\subseteq \sigma$  for all  $1 \leq j < i$  implies  $\mathcal{X}_j \not\subseteq \sigma \cup \{i\}$  for all  $1 \leq j < i$ .

We have  $\sigma \cup \{i\} \in \mathcal{C}_1$ , and for each  $1 \leq j < i$ ,  $\{1, 2, \dots, j\} \subseteq \sigma \cup \{i\}$  and  $\mathcal{X}_j \not\subseteq \sigma \cup \{i\}$ . Since  $\sigma \cup \{i\}$  satisfies the conditions of the induction hypothesis for each  $1 \leq j < i$ , by iterative application of the induction hypothesis, we have  $\sigma \cup \{i\} \in \mathcal{C}_{j+1}$  for all  $1 \leq j < i$ . In particular, for  $j = i - 1$ , we get  $\sigma \cup \{i\} \in \mathcal{C}_i$ . Then  $\sigma \setminus \{i\} = \sigma \in \mathcal{C}_i$  implies that  $\sigma \notin \mathcal{C}_{i+1}$  by Remark 2.6 (ii), a contradiction. Hence  $i \in \sigma$ .

Since  $\{1, 2, \dots, i - 1\} \subseteq \sigma$ , we conclude that  $\{1, 2, \dots, i\} \subseteq \sigma$ .

We now show that  $\mathcal{X}_i \not\subseteq \sigma$ . Suppose  $\mathcal{X}_i \subseteq \sigma$ . Since  $1 < i \leq p$ , we have  $0 \notin \mathcal{X}_i$ . Moreover, by the definition of  $C_n^p$ ,  $i \approx u$  if and only if  $u \in \mathcal{X}_i$ . It follows that  $\mathcal{X}_i \subseteq \sigma \cup \{0\}$ , and  $i \sim v$  for all  $v \in (\sigma \cup \{0\})^c$ . By Proposition 3.2 (iii), we get  $\sigma \setminus \{i\} \in \mathcal{C}_1$ .

For each  $1 \leq j < i$ , we have  $\{1, 2, \dots, j\} \subseteq \sigma$  and  $\mathcal{X}_j \not\subseteq \sigma$ . Clearly,  $\{1, 2, \dots, j\} \subseteq \sigma \setminus \{i\}$  and  $\mathcal{X}_j \not\subseteq \sigma \setminus \{i\}$  for each  $1 \leq j < i$ . Since  $\sigma \setminus \{i\} \in \mathcal{C}_1$ , applying the induction hypothesis iteratively to  $\sigma \setminus \{i\}$ , we have  $\sigma \setminus \{i\} \in \mathcal{C}_{j+1}$  for all  $1 \leq j < i$ . Taking  $j = i - 1$ , we get



$\sigma \setminus \{i\} \in \mathcal{C}_i$ . Since  $\sigma \cup \{i\} = \sigma \in \mathcal{C}_i$ , Remark 2.6 (ii) implies  $\sigma \notin \mathcal{C}_{i+1}$ , a contradiction. Hence  $\mathcal{X}_i \not\subseteq \sigma$ .

- Conversely, assume  $\{1, 2, \dots, i\} \subseteq \sigma$  and  $\mathcal{X}_i \not\subseteq \sigma$ . Since  $1 < i \leq p$ , we have  $i \approx u$  if and only if  $u \in \mathcal{X}_i$ , and  $i \sim 0$ . Hence there exists  $v \in \sigma^c \setminus \{0\}$  such that  $i \approx v$ , which implies that  $\{i, v\}$  is a disconnected 2-set in  $((\sigma \setminus \{i\}) \cup \{0\})^c$ . So,  $(\sigma \setminus \{i\}) \cup \{0\} \in \Delta_2^t(C_n^p)$ . Therefore,  $\sigma \setminus \{i\} \notin \mathcal{C}_1$  by Proposition 3.2 (i). Since  $\mathcal{C}_i \subseteq \mathcal{C}_1$  by Remark 2.6 (i),  $\sigma \setminus \{i\} \notin \mathcal{C}_i$ . By Remark 2.6 (ii),  $\sigma \in \mathcal{C}_i$  implies  $\sigma \in \mathcal{C}_{i+1}$ .

This completes the proof.  $\square$

Although Proposition 3.3 holds for all  $n \geq 2p + 2$ , we apply it here only to the case  $n = 2p + 2$ . For  $n \geq 3p + 1$ , we employ a different approach.

**Lemma 3.4.** *Let  $\tau = \{1, 2, \dots, p\}$ . For  $n = 2p + 2$ ,  $\mathcal{C}_{p+1} = \{\tau\}$ .*

*Proof.* By the definition of  $C_{2p+2}^p$ ,  $\{0, p+1\}$  is the only disconnected 2-set containing 0 in  $V(C_{2p+2}^p)$ .

First, assume that  $\sigma \in \mathcal{C}_{p+1}$ . We prove  $\sigma = \tau$ . By Remark 2.6 (i),  $\mathcal{C}_{p+1} \subseteq \mathcal{C}_p \subseteq \dots \subseteq \mathcal{C}_1 \subseteq \mathcal{C}_0 = \Delta_2^t(C_n^p)$ . Hence  $\sigma \in \mathcal{C}_i$  and  $\sigma \in \mathcal{C}_{i+1}$  for all  $1 \leq i \leq p$ . By Proposition 3.3, for each  $1 \leq i \leq p$ , we obtain  $\{1, 2, \dots, i\} \subseteq \sigma$  and  $i + p + 1 \notin \sigma$ . It follows that  $\{1, 2, \dots, p\} \subseteq \sigma$  and  $\{p+2, p+3, \dots, 2p+1\} \subseteq \sigma^c$ . Since  $\sigma \in \mathcal{C}_1$ , Proposition 3.2 (i) implies that every disconnected 2-set in  $\sigma^c$  contains 0. Hence  $\{0, p+1\} \subseteq \sigma^c$ . This means that  $\sigma = \{1, 2, \dots, p\}$ , and thus  $\sigma = \tau$ .

We now prove that  $\tau \in \mathcal{C}_{p+1}$ . Observe that  $\{0, p+1\}$  is the only disconnected 2-set in  $\tau^c$ . Hence  $\tau \in \Delta_2^t(C_{2p+2}^p)$  and  $\tau \cup \{0\} \notin \Delta_2^t(C_{2p+2}^p)$ . By Proposition 3.2 (i),  $\tau \in \mathcal{C}_1$ . Since  $\{1, 2, \dots, p\} \subseteq \tau$  and  $\{p+2, p+3, \dots, 2p+1\} \subseteq \tau^c$ , it follows that  $\{1, 2, \dots, i\} \subseteq \tau$  and  $i + p + 1 \notin \tau$  for each  $1 \leq i \leq p$ . Applying Proposition 3.3 iteratively, we obtain  $\tau \in \mathcal{C}_{p+1}$ .

Hence  $\mathcal{C}_{p+1} = \{\tau\}$ .  $\square$

We have set up the necessary results for  $n = 2p + 2$ . Now, we address the case  $n \geq 3p + 1$ . For  $\sigma \in \Delta_2^t(C_n^p)$ , define

$$m_\sigma := \max(\sigma^c).$$

By Remark 3.1, if  $\sigma \in \Delta_2^t(C_n^p)$ , then  $C_n^p[\sigma^c]$  is not a complete graph. From the definition of  $C_n^p$ , we then obtain the following.

**Remark 3.5.** *Let  $\sigma \in \Delta_2^t(C_n^p)$ . Then  $m_\sigma \in \{p+1, p+2, \dots, n-1\}$ .*

For  $\sigma \in \Delta_2^t(C_n^p)$ , we have  $m_\sigma - p \geq 1$  by Remark 3.5. From Remark 2.6 (i),  $\mathcal{C}_{m_\sigma-p} \subseteq \mathcal{C}_1$ . The proposition below establishes the reverse inclusion  $\mathcal{C}_1 \subseteq \mathcal{C}_{m_\sigma-p}$ .

**Proposition 3.6.** *Let  $n \geq 3p + 1$ . If  $\sigma \in \mathcal{C}_1$ , then  $\sigma^c \subseteq \{0, m_\sigma - p, m_\sigma - p + 1, \dots, m_\sigma\}$  and  $\sigma \in \mathcal{C}_{m_\sigma-p}$ .*

*Proof.* Suppose  $\sigma \in \mathcal{C}_1$ . We first show that  $\sigma^c \subseteq \{0, m_\sigma - p, m_\sigma - p + 1, \dots, m_\sigma\}$ . Since  $\mathcal{C}_1 \subseteq \Delta_2^t(C_n^p)$ , we have  $\sigma \in \Delta_2^t(C_n^p)$ . Hence  $m_\sigma \in \{p+1, p+2, \dots, n-1\}$  by Remark 3.5. Clearly,  $m_\sigma \neq 0$ . By Proposition 3.2 (i), every disconnected 2-set in  $\sigma^c$  contains 0 (as  $\sigma \in \mathcal{C}_1$ ). Therefore, since  $m_\sigma \in \sigma^c$ , it follows that if there exists  $v \in V(C_n^p)$  such that  $v \neq 0$  and  $v \approx m_\sigma$ , then  $v \in \sigma$ . We consider the following cases depending on the value of  $m_\sigma$ :

- Suppose  $m_\sigma \in \{p+1, p+2, \dots, n-p-1\}$ . Then  $1 \leq m_\sigma - p < m_\sigma < m_\sigma + p \leq n-1$ . Since  $\max(\sigma^c) = m_\sigma$ , it follows that  $\{m_\sigma + 1, m_\sigma + 2, \dots, m_\sigma + p\} \subseteq \sigma$ . Let  $u \in \{1, 2, \dots, m_\sigma - p - 1\} \cup \{m_\sigma + p + 1, m_\sigma + p + 2, \dots, n-1\}$ . Then  $u \neq 0$  and  $u \approx m_\sigma$ , which implies  $u \in \sigma$ . Hence  $\sigma^c \subseteq \{0, m_\sigma - p, m_\sigma - p + 1, \dots, m_\sigma\}$ .
- Suppose  $m_\sigma = n - p$ . Then  $1 < n - 2p = m_\sigma - p < m_\sigma < m_\sigma + p - 1 = n - 1$  and  $m_\sigma + p \pmod{n} = 0$ . It follows that  $\{m_\sigma + 1, m_\sigma + 2, \dots, n-1\} \subseteq \sigma$ . Now, let  $u \in \{1, 2, \dots, m_\sigma - p - 1\}$ . Observe that  $u \neq 0$  and  $u \approx m_\sigma$ . Hence  $u \in \sigma$ , which implies  $\{1, 2, \dots, m_\sigma - p - 1\} \subseteq \sigma$ . Therefore  $\sigma^c \subseteq \{0, m_\sigma - p, m_\sigma - p + 1, \dots, m_\sigma\}$ .
- Suppose  $m_\sigma \in \{n - p + 1, n - p + 2, \dots, n-1\}$ . Since  $\sigma \in \Delta_2^t(C_n^p)$  and every disconnected 2-set in  $\sigma^c$  contains 0, there exists  $w \in \sigma^c$  such that  $w \approx 0$ . By the definition of  $C_n^p$ ,  $w \in \{p+1, p+2, \dots, n-p-1\}$ .

Observe that  $1 \leq m_\sigma + p \pmod{n} < p+1 \leq w \leq n-p-1 < m_\sigma \leq n-1$ . Since  $n \geq 3p+1$ , we have  $m_\sigma + p \pmod{n} < m_\sigma - 2p$ . Moreover,  $p \geq 1$  implies  $m_\sigma - 2p < m_\sigma - p < m_\sigma$ . Therefore  $m_\sigma + p \pmod{n} < w, m_\sigma - p < m_\sigma$ . If  $w < m_\sigma - p$ , then  $w \approx m_\sigma$ , which implies  $w \in \sigma$  (as  $w \neq 0$ ), a contradiction. So  $w \geq m_\sigma - p$ .

It follows that  $1 \leq m_\sigma + p \pmod{n} < m_\sigma - 2p \leq w - p < w < w + p \leq n - 1$ . Let  $u \in \{1, 2, \dots, m_\sigma + p \pmod{n}\}$ . Then  $u \neq 0$  and  $w \approx u$ , which implies  $\{u, w\}$  is a disconnected 2-set not containing 0. Since  $w \in \sigma^c$ , we get  $u \in \sigma$ , and thus  $\{1, 2, \dots, m_\sigma + p \pmod{n}\} \subseteq \sigma$ . Now, let  $u' \in \{m_\sigma + p \pmod{n} + 1, m_\sigma + p \pmod{n} + 2, \dots, m_\sigma - p - 1\}$ . Then  $u' \neq 0$  and  $u' \approx m_\sigma$ , which implies  $u' \in \sigma$ . Hence  $\{m_\sigma + p \pmod{n} + 1, m_\sigma + p \pmod{n} + 2, \dots, m_\sigma - p - 1\} \subseteq \sigma$ . Moreover,  $\max(\sigma^c) = m_\sigma$  implies that  $\{m_\sigma + 1, m_\sigma + 2, \dots, n - 1\} \subseteq \sigma$ . This means that  $\sigma^c \subseteq \{0, m_\sigma - p, m_\sigma - p + 1, \dots, m_\sigma\}$ .

Therefore,  $\sigma^c \subseteq \{0, m_\sigma - p, m_\sigma - p + 1, \dots, m_\sigma\}$ .

We now prove that  $\sigma \in \mathcal{C}_{m_\sigma - p}$ . If  $m_\sigma = p + 1$ , then  $m_\sigma - p = 1$ . Since  $\sigma \in \mathcal{C}_1$ , we are done.

Now, suppose  $m_\sigma \in \{p + 2, p + 3, \dots, n - 1\}$ . Let  $1 \leq i \leq m_\sigma - p - 1$ . Since  $\sigma^c \subseteq \{0, m_\sigma - p, m_\sigma - p + 1, \dots, m_\sigma\}$ ,  $i \in \sigma$ . We first show that  $(\sigma \setminus \{i\}) \cup \{0\} \in \Delta_2^t(C_n^p)$ . Note that  $((\sigma \setminus \{i\}) \cup \{0\})^c = (\sigma \cup \{0\})^c \cup \{i\}$ . By Remark 3.1, it suffices to find  $v \in (\sigma \cup \{0\})^c$  such that  $i \approx v$ .

- Suppose  $m_\sigma \in \{p + 2, p + 3, \dots, n - p\}$ . Then  $2 \leq m_\sigma - p < m_\sigma < m_\sigma + p \leq n$ . Since  $1 \leq i \leq m_\sigma - p - 1$ , by the definition of  $C_n^p$ ,  $i \approx m_\sigma$ . Therefore,  $v = m_\sigma$  such that  $i \approx v$ .
- Suppose  $m_\sigma \in \{n - p + 1, n - p + 2, \dots, n - 1\}$ . Since  $n \geq 3p + 1$  and  $p \geq 1$ , we have  $m_\sigma + p \pmod{n} < m_\sigma - 2p < m_\sigma - p$ . So  $1 \leq m_\sigma + p \pmod{n} \leq m_\sigma - p - 1$ . Since  $1 \leq i \leq m_\sigma - p - 1$ , we have  $1 \leq i \leq m_\sigma + p \pmod{n}$  or  $m_\sigma + p \pmod{n} < i \leq m_\sigma - p - 1$ .

If  $m_\sigma + p \pmod{n} < i < m_\sigma - p$ , then  $i \approx m_\sigma$ . Therefore, we have  $v = m_\sigma$  with  $i \approx v$ .

Now, assume that  $1 \leq i \leq m_\sigma + p \pmod{n}$ . Since  $\sigma \in \mathcal{C}_1$ , it follows by Proposition 3.2 (i) that every disconnected 2-set in  $\sigma^c$  contains 0. Further,  $\sigma \in \mathcal{C}_1 \subseteq \Delta_2^t(C_n^p)$  implies that there exists  $u \in \sigma^c \setminus \{0\} = (\sigma \cup \{0\})^c$  such that  $0 \approx u$ . By definition of  $C_n^p$ , we have  $p + 1 \leq u \leq n - p - 1$ . Moreover,  $\sigma^c \subseteq \{0, m_\sigma - p, m_\sigma - p + 1, \dots, m_\sigma\}$  implies that  $m_\sigma - p \leq u \leq m_\sigma$ . This means that  $m_\sigma - p \leq u \leq n - p - 1$ , and thus  $m_\sigma - 2p \leq u - p < u < u + p \leq n - 1$ . Since  $1 \leq m_\sigma + p \pmod{n} < m_\sigma - 2p$  and  $1 \leq i \leq m_\sigma + p \pmod{n}$ , we get  $i \notin \{u - p, u - p + 1, \dots, u, \dots, u + p\}$ , and hence  $i \approx u$ . Thus, we have  $v = u$  with  $i \approx v$ .

In all cases,  $(\sigma \setminus \{i\}) \cup \{0\} \in \Delta_2^t(C_n^p)$ . Hence  $\sigma \setminus \{i\} \notin \mathcal{C}_1$  for all  $1 \leq i \leq m_\sigma - p - 1$  by Proposition 3.2 (i), and therefore  $\sigma \in \mathcal{C}_{m_\sigma - p}$  by Remark 2.6 (iii) (as  $\sigma \in \mathcal{C}_1$ ).

This completes the proof.  $\square$

Proposition 3.6 states that if  $\sigma \in \mathcal{C}_1$ , then  $\sigma \in \mathcal{C}_{m_\sigma - p}$ . The following result provides a sufficient condition for when  $\sigma \notin \mathcal{C}_{m_\sigma - p + 1}$ .

**Proposition 3.7.** *Let  $n \geq 3p + 1$  and let  $\sigma \in \mathcal{C}_1$ . Suppose that there exists a disconnected 2-set  $\{0, v\}$  in  $\sigma^c$  such that  $v \neq m_\sigma - p$ . Then  $\sigma \notin \mathcal{C}_{m_\sigma - p + 1}$ .*

*Proof.* Since  $\sigma \in \mathcal{C}_1$ , it follows by Proposition 3.6 that  $\sigma \in \mathcal{C}_{m_\sigma - p}$  and  $\sigma^c \subseteq \{0, m_\sigma - p, m_\sigma - p + 1, \dots, m_\sigma\}$ . We have either  $m_\sigma - p \in \sigma$  or  $m_\sigma - p \notin \sigma$ .

First, suppose that  $m_\sigma - p \in \sigma$ . Then  $m_\sigma - p \notin (\sigma \cup \{0\})^c$ . Hence  $(\sigma \cup \{0\})^c \subseteq \{m_\sigma - p + 1, m_\sigma - p + 2, \dots, m_\sigma\}$ , and thus  $m_\sigma - p \sim u$  for all  $u \in (\sigma \cup \{0\})^c$ . Since  $\sigma \in \mathcal{C}_1$ , we get  $\sigma \setminus \{m_\sigma - p\} \in \mathcal{C}_1$  by Proposition 3.2 (iii). Observe that  $m_{(\sigma \setminus \{m_\sigma - p\})} = m_\sigma$ . By Proposition 3.6,  $\sigma \setminus \{m_\sigma - p\} \in \mathcal{C}_1$  implies  $\sigma \setminus \{m_\sigma - p\} \in \mathcal{C}_{m_\sigma - p}$ . Since  $\sigma \cup \{m_\sigma - p\} = \sigma \in \mathcal{C}_{m_\sigma - p}$ , we have  $\sigma \notin \mathcal{C}_{m_\sigma - p + 1}$  by Remark 2.6 (ii).

Now, let  $m_\sigma - p \notin \sigma$ . Since there exists a disconnected 2-set  $\{0, v\}$  in  $\sigma^c$  such that  $v \neq m_\sigma - p$ , we get  $\sigma \cup \{m_\sigma - p\} \in \Delta_2^t(C_n^p)$ . By Proposition 3.2 (ii),  $\sigma \cup \{m_\sigma - p\} \in \mathcal{C}_1$  (as  $\sigma \in \mathcal{C}_1$ ). Since  $m_\sigma - p \neq m_\sigma$ , we get  $m_{(\sigma \cup \{m_\sigma - p\})} = m_\sigma$ . It follows that  $\sigma \cup \{m_\sigma - p\} \in \mathcal{C}_{m_\sigma - p}$  by Proposition 3.6. We have  $\sigma \setminus \{m_\sigma - p\} = \sigma \in \mathcal{C}_{m_\sigma - p}$ . By Remark 2.6 (ii),  $\sigma \notin \mathcal{C}_{m_\sigma - p + 1}$ .  $\square$

**Proposition 3.8.** *Let  $n \geq 3p + 1$ . For  $\sigma \in \Delta_2^t(C_n^p)$ , we have the following.*

- (i) *If  $\{0, m_\sigma - p\}$  is the only disconnected 2-set in  $\sigma^c$ , then  $m_\sigma \in \{n - p, n - p + 1, \dots, n - 1\}$ .*
- (ii)  *$\sigma \in \mathcal{C}_{n-p}$  if and only if  $\{0, m_\sigma - p\}$  is the only disconnected 2-set in  $\sigma^c$ .*



*Proof.* (i) By Remark 3.5,  $m_\sigma \in \{p+1, p+2, \dots, n-1\}$ . If  $m_\sigma \in \{p+1, p+2, \dots, n-p-1\}$ , then  $0 \approx m_\sigma$ . Since  $\{0, m_\sigma\} \subseteq \sigma^c$ , we get a contradiction to the fact that  $\{0, m_\sigma - p\}$  is the only disconnected 2-set in  $\sigma^c$ . Hence  $m_\sigma \in \{n-p, n-p+1, \dots, n-1\}$ .

(ii) Suppose  $\sigma \in \mathcal{C}_{n-p}$ . Since  $p+1 \leq m_\sigma \leq n-1$ , we have  $1 < m_\sigma - p + 1 \leq n-p$ . Hence  $\mathcal{C}_{n-p} \subseteq \mathcal{C}_{m_\sigma-p+1} \subseteq \mathcal{C}_1$  implies that  $\sigma \in \mathcal{C}_{m_\sigma-p+1}$  and  $\sigma \in \mathcal{C}_1$ . Now,  $\sigma \in \mathcal{C}_1$  implies that every disconnected 2-set in  $\sigma^c$  contains 0. If  $\{0, v\}$  is a disconnected 2-set in  $\sigma^c$  with  $v \neq m_\sigma - p$ , then  $\sigma \notin \mathcal{C}_{m_\sigma-p+1}$  by Proposition 3.7, a contradiction. Therefore,  $\{0, m_\sigma - p\}$  is the only disconnected 2-set in  $\sigma^c$ .

Conversely, assume that  $\{0, m_\sigma - p\}$  is the only disconnected 2-set in  $\sigma^c$ . By Remark 3.1,  $\sigma \cup \{0\}$  and  $\sigma \cup \{m_\sigma - p\} \notin \Delta_2^t(C_n^p)$ . Hence  $\sigma \in \mathcal{C}_1$ , and thus  $\sigma \in \mathcal{C}_{m_\sigma-p}$  by Proposition 3.6. Moreover, since  $\mathcal{C}_{m_\sigma-p} \subseteq \Delta_2^t(C_n^p)$ ,  $\sigma \cup \{m_\sigma - p\} \notin \mathcal{C}_{m_\sigma-p}$ . Therefore,  $\sigma \in \mathcal{C}_{m_\sigma-p+1}$  by Remark 2.6 (ii).

Since  $\{0, m_\sigma - p\}$  is the only disconnected 2-set in  $\sigma^c$ ,  $m_\sigma \in \{n-p, n-p+1, \dots, n-1\}$  by (i). If  $m_\sigma = n-1$ , then  $m_\sigma - p + 1 = n-p$ , and thus  $\sigma \in \mathcal{C}_{m_\sigma-p+1}$  implies that  $\sigma \in \mathcal{C}_{n-p}$ , as required.

Now, assume that  $m_\sigma \in \{n-p, n-p+1, \dots, n-2\}$ . Let  $m_\sigma - p + 1 \leq j \leq n-p-1$ . Then  $m_\sigma \geq n-p$  and  $n \geq 3p+1$  implies that  $p+1 < j < n-p$ , and hence  $j \approx 0$ . Since  $\{0, m_\sigma - p\}$  is the only disconnected 2-set in  $\sigma^c$ , we have  $j \in \sigma$ . By Proposition 3.6,  $\sigma \in \mathcal{C}_1$  implies that  $\sigma^c \subseteq \{0, m_\sigma - p, m_\sigma - p + 1, \dots, m_\sigma\}$ . Since  $m_\sigma - p < m_\sigma - p + 1 \leq j \leq n-p-1 < m_\sigma$ , it follows that  $j \sim u$  for all  $u \in (\sigma \cup \{0\})^c$ . Therefore,  $\sigma \in \mathcal{C}_1$  implies that  $\sigma \setminus \{j\} \in \mathcal{C}_1$  by Proposition 3.2 (iii). Note that  $m_{(\sigma \setminus \{j\})} = m_\sigma$  (as  $j < m_\sigma$ ). Moreover,  $\{0, j\} \subseteq (\sigma \setminus \{j\})^c$  such that  $0 \not\sim j$  and  $j \neq m_\sigma - p$ . Hence  $\sigma \setminus \{j\} \notin \mathcal{C}_{m_\sigma-p+1}$  by Proposition 3.7. Since  $\sigma \in \mathcal{C}_{m_\sigma-p+1}$  and  $m_\sigma - p + 1 \leq j \leq n-p-1$ , we get  $\sigma \in \mathcal{C}_{n-p}$  by Remark 2.6 (iii).  $\square$

**Lemma 3.9.** *Let  $n \geq 3p+1$ . Then  $\mathcal{C}_{n-p+1} = \{\tau\}$ , where  $\tau^c = \{0, n-2p, n-p\}$ .*

*Proof.* Since  $n \geq 3p+1$ ,  $0 \approx n-2p$ . Hence  $\tau \in \Delta_2^t(C_n^p)$ . We first show that  $\tau \in \mathcal{C}_{n-p+1}$ . Observe that  $\{0, m_\tau - p\} = \{0, n-2p\}$  is the only disconnected 2-set in both  $\tau^c$  and  $(\tau \cup \{n-p\})^c$ . Therefore,  $\tau \in \mathcal{C}_{n-p}$  by Proposition 3.8 (ii). Clearly,  $\tau \cup \{n-p\} \in \Delta_2^t(C_n^p)$  and  $\tau \cup \{0, n-p\} \notin \Delta_2^t(C_n^p)$ . Hence  $\tau \cup \{n-p\} \in \mathcal{C}_1$ . Since  $\{0, n-2p\}$  is a disconnected 2-set in  $(\tau \cup \{n-p\})^c$  with  $n-2p = m_{(\tau \cup \{n-p\})} \neq m_{(\tau \cup \{n-p\})} - p$ , it follows from Proposition 3.7 that  $\tau \cup \{n-p\} \notin \mathcal{C}_{n-3p+1}$ . Further,  $\mathcal{C}_{n-p} \subseteq \mathcal{C}_{n-3p+1}$  (as  $n-3p+1 < n-p$ ) implies that  $\tau \cup \{n-p\} \notin \mathcal{C}_{n-p}$ . Therefore,  $\tau \in \mathcal{C}_{n-p}$  implies  $\tau \in \mathcal{C}_{n-p+1}$  by Remark 2.6 (ii).

Let  $\sigma \in \mathcal{C}_{n-p+1}$ . We prove that  $\sigma = \tau$ . Since  $\mathcal{C}_{n-p+1} \subseteq \mathcal{C}_{n-p} \subseteq \dots \subseteq \mathcal{C}_1$ ,  $\sigma \in \mathcal{C}_1 \cap \mathcal{C}_{n-p}$ . From Proposition 3.6,  $\sigma^c \subseteq \{0, m_\sigma - p, m_\sigma - p + 1, \dots, m_\sigma\}$ , and by Proposition 3.8 (ii),  $\{0, m_\sigma - p\}$  is the only disconnected 2-set in  $\sigma^c$ . Hence  $m_\sigma \in \{n-p, n-p+1, \dots, n-1\}$  by Proposition 3.8 (i).

Suppose  $m_\sigma \in \{n-p+1, n-p+2, \dots, n-1\}$ . Then  $m_\sigma - p < n-p < m_\sigma$ . Since  $n-p \sim 0$  and  $\sigma^c \subseteq \{0, m_\sigma - p, m_\sigma - p + 1, \dots, m_\sigma\}$ , it follows that  $n-p \sim u$  for all  $u \in \sigma^c$ . Therefore,  $\{0, m_\sigma - p\}$  is the only disconnected 2-set in  $\sigma^c$  implies that  $\{0, m_\sigma - p\}$  is the only disconnected 2-set in both  $(\sigma \setminus \{n-p\})^c$  and  $(\sigma \cup \{n-p\})^c$ . Moreover,  $m_{(\sigma \setminus \{n-p\})} = m_{(\sigma \cup \{n-p\})} = m_\sigma$ . By Proposition 3.8 (ii),  $\sigma \setminus \{n-p\}, \sigma \cup \{n-p\} \in \mathcal{C}_{n-p}$ . Hence  $\sigma \notin \mathcal{C}_{n-p+1}$  by Remark 2.6 (ii), a contradiction. So  $m_\sigma = n-p$ .

Since  $\sigma^c \subseteq \{0, m_\sigma - p, m_\sigma - p + 1, \dots, m_\sigma\}$ , we get  $\sigma^c \subseteq \{0, n-2p, n-2p+1, \dots, n-p\}$ . Moreover,  $p \geq 1$  and  $n \geq 3p+1$  imply  $p+1 \leq n-2p \leq n-p-1$ . This means that  $0 \approx u$  for all  $u \in \sigma^c \setminus \{0, n-p\}$ . We have  $\{0, m_\sigma - p\} = \{0, n-2p\}$  is the only disconnected 2-set in  $\sigma^c$  and  $m_\sigma = n-p \in \sigma^c$ . It follows that  $\sigma^c = \{0, n-2p, n-p\}$ . Therefore  $\sigma = \tau$ .  $\square$

**Theorem 3.10.** *Let  $n \geq 2p+2$ . Then  $\Delta_2^t(C_n^p) \simeq \begin{cases} \mathbb{S}^{\frac{n-4}{2}} & \text{if } n = 2p+2, \\ \mathbb{S}^{n-4} & \text{if } n \geq 3p+1. \end{cases}$*

*Proof.* By our construction,  $\mathcal{M} = \bigsqcup_{i=0}^{n-1} \mathcal{M}_i$  is an acyclic matching on  $\Delta_2^t(C_n^p)$  with  $\mathcal{C}_n$  as the set of all critical faces. Observe that for any  $0 \leq i \leq n-1$ , if  $\mathcal{C}_i$  is a singleton set, then  $\mathcal{C}_i = \mathcal{C}_n$ .

- Let  $n = 2p + 2$ . Then Lemma 3.4 implies  $\mathcal{C}_{p+1} = \{\sigma\}$ , where  $\sigma = \{1, 2, \dots, p\}$ . Therefore  $\mathcal{C}_n = \{\sigma\}$ . Note that the dimension of  $\sigma$  is  $p - 1 = \frac{n-4}{2}$ .
- Let  $n \geq 3p + 1$ . By Lemma 3.9,  $\mathcal{C}_{n-p+1} = \{\tau\}$ , where  $\tau^c = \{0, n-2p, n-p\}$ . Hence  $\mathcal{C}_n = \{\tau\}$ . Moreover, the dimension of  $\tau$  is  $n - 4$ .

Result follows from Theorem 2.7 and Corollary 2.8.  $\square$

**3.2. Cartesian product of complete graphs and path graphs.** In this section, we prove Theorem 1.4. For a positive integer  $q$ , let  $[q] = \{1, 2, \dots, q\}$  and  $[0, q] = [q] \cup \{0\}$ . The Cartesian product  $K_m \square P_n$  has vertex set

$$V(K_m \square P_n) = V(K_m) \times V(P_n) = \{(i, j) \mid i \in [0, m-1] \text{ and } j \in [0, n-1]\}.$$

For any distinct  $(i_1, j_1), (i_2, j_2) \in V(K_m \square P_n)$ , we have  $(i_1, j_1) \sim (i_2, j_2)$  if and only if  $i_1 = i_2$  and  $|j_1 - j_2| = 1$ , or  $j_1 = j_2$ . Equivalently,  $(i_1, j_1) \approx (i_2, j_2)$  if and only if  $i_1 = i_2$  and  $|j_1 - j_2| > 1$ , or  $i_1 \neq i_2$  and  $j_1 \neq j_2$ .

For notational convenience, we denote the graph  $K_m \square P_n$  by  $\mathcal{G}_{mn}$ , and write any vertex of the form  $(0, j) \in V(\mathcal{G}_{mn})$  simply as  $(j)_0$ . We now define a sequence of element matchings  $\mathcal{M}_{(0)_0}, \mathcal{M}_{(1)_0}, \dots, \mathcal{M}_{(n-1)_0}$  on  $\Delta_2^t(\mathcal{G}_{mn})$  and the sets  $\mathcal{C}_0, \mathcal{C}_1, \dots, \mathcal{C}_n$ , using vertices  $(0)_0, (1)_0, \dots, (n-1)_0$ , as in Equation (1). By Lemma 2.5,  $\mathcal{M} := \bigsqcup_{i=0}^{n-1} \mathcal{M}_{(i)_0}$  is an acyclic matching on  $\Delta_2^t(\mathcal{G}_{mn})$ . Clearly,  $\mathcal{C}_0 = \Delta_2^t(\mathcal{G}_{mn})$  and  $\mathcal{C}_1 = \{\sigma \in \Delta_2^t(\mathcal{G}_{mn}) \mid \sigma \notin \mathcal{M}_{(0)_0}\}$ .

**Proposition 3.11.** *Let  $\sigma \in \mathcal{C}_1$ .*

- (i) *If  $\sigma^c = \{(0)_0, (j)_0\}$  for some  $j \in [n-1]$ , then  $\sigma, \sigma \setminus \{(j-1)_0\} \notin \mathcal{C}_j$ , and hence  $\sigma, \sigma \setminus \{(j-1)_0\} \notin \mathcal{C}_n$ .*
- (ii) *If  $\{(1)_0, (2)_0, \dots, (j)_0\} \subseteq \sigma, \{(0)_0, (i, j)\} \subseteq \sigma^c$  and  $\sigma \setminus \{(j)_0\} \in \mathcal{C}_1$  for some  $i \in [m-1], j \in [n-1]$ , then  $\sigma, \sigma \setminus \{(j)_0\} \notin \mathcal{C}_n$ .*

*Proof.* (i) Suppose  $\sigma^c = \{(0)_0, (j)_0\}$ . Since  $\sigma \in \mathcal{C}_1 \subseteq \Delta_2^t(\mathcal{G}_{mn})$ ,  $(0)_0 \approx (j)_0$  by Remark 3.1. Hence  $j \in [n-1] \setminus \{1\}$ . Since  $\mathcal{C}_n \subseteq \mathcal{C}_j$ , it suffices to prove  $\sigma, \sigma \setminus \{(j-1)_0\} \notin \mathcal{C}_j$ .

We have  $(\sigma \cup \{(0)_0\})^c = \{(j)_0\}$  and  $(j-1)_0 \sim (j)_0$ . By Proposition 3.2 (iii),  $\sigma \setminus \{(j-1)_0\} \in \mathcal{C}_1$ .

If  $j = 2$ , then  $\sigma \cup \{(1)_0\} = \sigma \in \mathcal{C}_1$  and  $\sigma \setminus \{(1)_0\} \in \mathcal{C}_1$  imply  $\sigma, \sigma \setminus \{(1)_0\} \notin \mathcal{C}_2 = \mathcal{C}_j$  by Remark 2.6 (ii).

Now, assume  $3 \leq j \leq n-1$ . Let  $1 \leq r \leq j-2$ . Then  $(r)_0 \in \sigma, \sigma \setminus \{(j-1)_0\}$ . Since  $j-r \geq 2$ ,  $(j)_0 \approx (r)_0$ . This implies  $(\sigma \setminus \{(r)_0\}) \cup \{(0)_0\}, (\sigma \setminus \{(r)_0, (j-1)_0\}) \cup \{(0)_0\} \in \Delta_2^t(\mathcal{G}_{mn})$ . Hence  $\sigma \setminus \{(r)_0\}, \sigma \setminus \{(r)_0, (j-1)_0\} \notin \mathcal{C}_1$  by Proposition 3.2 (i). From Remark 2.6 (iii), we obtain  $\sigma, \sigma \setminus \{(j-1)_0\} \in \mathcal{C}_{j-1}$ . Since  $\sigma \cup \{(j-1)_0\} = \sigma$ , Remark 2.6 (ii) implies  $\sigma, \sigma \setminus \{(j-1)_0\} \notin \mathcal{C}_j$ .

- (ii) Let  $i \in [m-1], j \in [n-1]$  and suppose that  $\{(1)_0, (2)_0, \dots, (j)_0\} \subseteq \sigma, \{(0)_0, (i, j)\} \subseteq \sigma^c$  and  $\sigma \setminus \{(j)_0\} \in \mathcal{C}_1$ . Since  $\mathcal{C}_n \subseteq \mathcal{C}_{j+1}$  by Remark 2.6 (i), we prove  $\sigma, \sigma \setminus \{(j)_0\} \notin \mathcal{C}_{j+1}$ .

For  $j = 1$ ,  $\sigma \cup \{(1)_0\} = \sigma \in \mathcal{C}_1$  and  $\sigma \setminus \{(1)_0\} \in \mathcal{C}_1$  imply  $\sigma, \sigma \setminus \{(1)_0\} \notin \mathcal{C}_2 = \mathcal{C}_{j+1}$ .

Now, suppose  $2 \leq j \leq n-1$ . Let  $1 \leq r \leq j-1$ . Then  $(r)_0 \in \sigma, \sigma \setminus \{(j)_0\}$  and  $(r)_0 \approx (i, j)$  (as  $i \neq 0$  and  $j \neq r$ ). Hence  $(\sigma \setminus \{(r)_0\}) \cup \{(0)_0\}, (\sigma \setminus \{(r)_0, (j)_0\}) \cup \{(0)_0\} \in \Delta_2^t(\mathcal{G}_{mn})$ . From Proposition 3.2 (i),  $\sigma \setminus \{(r)_0\}, \sigma \setminus \{(r)_0, (j)_0\} \notin \mathcal{C}_1$ . Therefore  $\sigma, \sigma \setminus \{(j)_0\} \in \mathcal{C}_j$  by Remark 2.6 (iii), and thus  $\sigma, \sigma \setminus \{(j)_0\} \notin \mathcal{C}_{j+1}$  by Remark 2.6 (ii).  $\square$

**Theorem 3.12.** *For  $m, n \geq 2$ ,  $\Delta_2^t(K_m \square P_n) \simeq V_{(m-1)(n-1)} \mathbb{S}^{mn-4}$ .*

*Proof.* We first prove that  $\sigma \in \mathcal{C}_n$  if and only if  $\sigma^c = \{(0)_0, (i, j-1), (i, j)\}$  for some  $i \in [m-1]$  and  $j \in [n-1]$ .

First, assume  $\sigma \in \mathcal{C}_n$ . Since  $\mathcal{C}_n \subseteq \mathcal{C}_1 \subseteq \Delta_2^t(\mathcal{G}_{mn})$ ,  $\sigma \in \mathcal{C}_1 \cap \Delta_2^t(\mathcal{G}_{mn})$ . By Proposition 3.2 (i), every disconnected 2-set in  $\sigma^c$  contains  $(0)_0$ . It follows that there  $\sigma^c = \{(0)_0, (i, j)\} \cup X$  for some  $(i, j) \in V(\mathcal{G}_{mn})$  and  $X \subseteq V(\mathcal{G}_{mn})$  such that  $(i, j) \approx (0)_0$  and  $\mathcal{G}_{mn}[X \cup \{(i, j)\}]$  is a complete graph. Since  $(i, j) \in V(\mathcal{G}_{mn})$ , we have  $i \in [0, m-1]$ .

- Suppose  $i = 0$ . Then  $(i, j) = (j)_0 \approx (0)_0$  implies  $j \in [n-1] \setminus \{1\}$ . Since  $\mathcal{G}_{mn}[X \cup \{(j)_0\}]$  is a complete graph, either (i)  $X = \emptyset$ , or (ii)  $X = \{(j-1)_0\}$ , or (iii)  $X \subseteq \{(i', j) \mid i' \in [m-1]\}$  with  $X \neq \emptyset$ .

We have  $\sigma^c = \{(0)_0, (j)_0\} \cup X$ . If  $X = \emptyset$ , then  $\sigma^c = \{(0)_0, (j)_0\}$ , and if  $X = \{(j-1)_0\}$ , then  $(\sigma \cup \{(j-1)_0\})^c = \{(0)_0, (j)_0\}$ . In either case, Proposition 3.11 (i) contradicts  $\sigma \in \mathcal{C}_n$ .

Now, assume  $X \subseteq \{(i', j) \mid i' \in [m-1]\}$  with  $X \neq \emptyset$ . Then there exists  $t \in [m-1]$  such that  $(t, j) \in X$ . Observe that  $\{(1)_0, (2)_0, \dots, (j)_0\} \subseteq \sigma \cup \{(j)_0\}$  and  $\{(0)_0, (t, j)\} \subseteq (\sigma \cup \{(j)_0\})^c$ . Since  $\sigma \in \mathcal{C}_1$ , applying Proposition 3.11 (ii) to  $\sigma \cup \{(j)_0\}$  gives  $\sigma \notin \mathcal{C}_n$ , again a contradiction. Hence  $i \neq 0$ .

- Suppose  $i \in [m-1]$ . Then  $(i, j) \approx (0)_0$  implies  $j \in [n-1]$ . Since  $\mathcal{G}_{mn}[X \cup \{(i, j)\}]$  is a complete graph, either (i)  $X = \emptyset$ , or (ii)  $X = \{(i, j-1)\}$ , or (iii)  $X \subseteq \{(i', j) \mid i' \in [0, m-1] \setminus \{i\}\}$  with  $X \neq \emptyset$ .

If  $X = \emptyset$ , then  $\sigma^c = \{(0)_0, (i, j)\}$  and  $\{(1)_0, (2)_0, \dots, (j)_0\} \subseteq \sigma$ . Since  $\sigma \in \mathcal{C}_1$ ,  $(\sigma \cup \{(0)_0\})^c = \{(i, j)\}$  and  $(j)_0 \sim (i, j)$ , we have  $\sigma \setminus \{(j)_0\} \in \mathcal{C}_1$  by Proposition 3.2 (iii). Hence  $\sigma \notin \mathcal{C}_n$  by Proposition 3.11 (ii), a contradiction.

Now, suppose  $X \subseteq \{(i', j) \mid i' \in [0, m-1] \setminus \{i\}\}$  with  $X \neq \emptyset$ . We have  $\sigma^c = \{(0)_0, (i, j)\} \cup X$ . If  $(j)_0 \in X$ , then  $\{(1)_0, (2)_0, \dots, (j)_0\} \subseteq \sigma \cup \{(j)_0\}$ . Moreover,  $\{(0)_0, (i, j)\} \subseteq (\sigma \cup \{(j)_0\})^c$  and  $\sigma \in \mathcal{C}_1$ . Applying Proposition 3.11 (ii) to  $\sigma \cup \{(j)_0\}$ , we get  $\sigma \notin \mathcal{C}_n$ , a contradiction. If  $(j)_0 \notin X$ , then  $\{(1)_0, (2)_0, \dots, (j)_0\} \subseteq \sigma$ . Observe that  $\mathcal{G}_{mn}[(\sigma \setminus \{(j)_0\}) \cup \{(0)_0\}]$  is a complete graph. Hence  $\sigma \setminus \{(j)_0\} \in \mathcal{C}_1$  by Proposition 3.2 (iii). By Proposition 3.11 (ii),  $\sigma \notin \mathcal{C}_n$ , a contradiction.

Therefore  $X = \{(i, j-1)\}$ .

This means that  $\sigma^c = \{(0)_0, (i, j-1), (i, j)\}$  for some  $i \in [m-1]$  and  $j \in [n-1]$ .

Conversely, assume that  $\sigma^c = \{(0)_0, (i, j-1), (i, j)\}$  for some  $i \in [m-1]$  and  $j \in [n-1]$ . Since  $(0)_0 \approx (i, j)$  and  $(i, j-1) \sim (i, j)$ , it follows that  $\sigma \in \Delta_2^t(\mathcal{G}_{mn})$  and  $\sigma \cup \{0\} \notin \Delta_2^t(\mathcal{G}_{mn})$ . Therefore  $\sigma \in \mathcal{C}_1$  by Proposition 3.2 (i).

Let  $1 \leq r \leq n-1$ . Then  $(r)_0 \in \sigma$ . Observe that if  $r = j$ , then  $(r)_0 \approx (i, j-1)$ ; and if  $r \neq j$ , then  $(r)_0 \approx (i, j)$ . Hence  $(\sigma \setminus \{(r)_0\}) \cup \{(0)_0\} \in \Delta_2^t(\mathcal{G}_{mn})$ . By Proposition 3.2 (i),  $\sigma \setminus \{(r)_0\} \notin \mathcal{C}_1$ . Hence  $\sigma \in \mathcal{C}_n$  by Remark 2.6 (iii).

We conclude that  $\mathcal{C}_n = \{\sigma \in \Delta_2^t(\mathcal{G}_{mn}) \mid \sigma^c = \{(0)_0, (i, j-1), (i, j)\} \text{ for some } i \in [m-1], j \in [n-1]\}$ . Note that  $|\mathcal{C}_n| = (m-1)(n-1)$  and each  $\sigma \in \mathcal{C}_n$  has dimension  $mn-4$ . Hence the result follows by Theorem 2.7 and Corollary 2.8.  $\square$

**3.3. Cartesian product of complete graphs and cycle graphs.** In this section, we prove Theorem 1.5. For a positive integer  $q$ , we use the notation  $[q]$  and  $[0, q]$  from the previous section. The Cartesian product  $K_m \square C_n$  has vertex set

$$V(K_m \square C_n) = V(K_m) \times V(C_n) = \{(i, j) \mid i \in [0, m-1] \text{ and } j \in [0, n-1]\}.$$

For any distinct  $(i_1, j_1), (i_2, j_2) \in V(K_m \square C_n)$ ,  $(i_1, j_1) \sim (i_2, j_2)$  if and only if  $i_1 = i_2$  and  $|j_1 - j_2| \in \{1, n-1\}$ , or  $j_1 = j_2$ . Equivalently,  $(i_1, j_1) \approx (i_2, j_2)$  if and only if  $i_1 = i_2$  and  $1 < |j_1 - j_2| < n-1$ , or  $i_1 \neq i_2$  and  $j_1 \neq j_2$ .

The proof of Theorem 1.5 follows the same approach as that of Theorem 1.4. The only difference arises from the adjacency of vertices 0 and  $n-1$  in the cycle graph  $C_n$ .

Let  $\mathcal{G}'_{mn}$  denote the graph  $K_m \square C_n$ . We write  $(j)_0$  for a vertex of the form  $(0, j) \in V(\mathcal{G}'_{mn})$ . We define a sequence of element matchings  $\mathcal{M}_{(0)_0}, \mathcal{M}_{(1)_0}, \dots, \mathcal{M}_{(n-1)_0}$  on  $\Delta_2^t(\mathcal{G}'_{mn})$  and the sets  $\mathcal{C}_0, \mathcal{C}_1, \dots, \mathcal{C}_n$ , using vertices  $(0)_0, (1)_0, \dots, (n-1)_0$ , as in Equation (1). Then  $\mathcal{M} := \bigsqcup_{i=0}^{n-1} \mathcal{M}_{(i)_0}$  is an acyclic matching on  $\Delta_2^t(\mathcal{G}'_{mn})$ . We have  $\mathcal{C}_0 = \Delta_2^t(\mathcal{G}'_{mn})$  and  $\mathcal{C}_1 = \{\sigma \in \Delta_2^t(\mathcal{G}'_{mn}) \mid \sigma \notin \mathcal{M}_{(0)_0}\}$ .

The following result is analogous to Proposition 3.11.

**Proposition 3.13.** *Let  $\sigma \in \mathcal{C}_1$ .*

- (i) *If  $\sigma^c = \{(0)_0, (j)_0\}$  for some  $j \in [n-1]$ , then  $\sigma, \sigma \setminus \{(j-1)_0\} \notin \mathcal{C}_j$ , and hence  $\sigma, \sigma \setminus \{(j-1)_0\} \notin \mathcal{C}_n$ .*
- (ii) *If  $\{(1)_0, (2)_0, \dots, (j)_0\} \subseteq \sigma$ ,  $\{(0)_0, (i, j)\} \subseteq \sigma^c$  and  $\sigma \setminus \{(j)_0\} \in \mathcal{C}_1$  for some  $i \in [m-1]$ ,  $j \in [n-1]$ , then  $\sigma, \sigma \setminus \{(j)_0\} \notin \mathcal{C}_n$ .*

*Proof.* (i) We use a similar argument as in the proof of Proposition 3.11 (i). Note that in the proof of Proposition 3.11 (i),  $(0)_0 \approx (j)_0$  if and only if  $j \in [n-1] \setminus \{1\}$ , and if  $3 \leq j \leq n-1$ , then for any  $1 \leq r \leq j-2$ , we have  $j-r \geq 2$ , which implies  $(j)_0 \approx (r)_0$ . Here,  $(0)_0 \approx (j)_0$  if and only if  $j \in [n-2] \setminus \{1\}$ , and if  $3 \leq j \leq n-2$ , then for any  $1 \leq r \leq j-2$ , we have  $2 \leq j-r \leq n-3$ , which implies  $(j)_0 \approx (r)_0$ . Using these results and following the proof of Proposition 3.11 (i), we are done.

(ii) The proof is identical to the proof of Proposition 3.11 (ii). □

**Theorem 3.14.** For  $m \geq 2$  and  $n \geq 4$ ,  $\Delta_2^t(K_m \square C_n) \simeq \bigvee_{n(m-1)+1} \mathbb{S}^{mn-4}$ .

*Proof.* Let

$$\begin{aligned} A &= \{\sigma \subseteq V(\mathcal{G}'_{mn}) \mid \sigma^c = \{(0)_0, (n-2)_0, (n-1)_0\}\}, \\ B &= \{\sigma \subseteq V(\mathcal{G}'_{mn}) \mid \sigma^c = \{(0)_0, (i, 0), (i, n-1)\} \text{ for some } i \in [m-1]\}, \\ C &= \{\sigma \subseteq V(\mathcal{G}'_{mn}) \mid \sigma^c = \{(0)_0, (i, j-1), (i, j)\} \text{ for some } i \in [m-1], j \in [n-1]\}. \end{aligned}$$

Note that  $A$ ,  $B$ , and  $C$  are pairwise disjoint. We show that  $\mathcal{C}_n = A \sqcup B \sqcup C$ .

Assume first that  $\sigma \in \mathcal{C}_n$ . Then  $\sigma \in \mathcal{C}_1 \cap \Delta_2^t(\mathcal{G}'_{mn})$ . Therefore, Proposition 3.2 (i) implies that every disconnected 2-set in  $\sigma^c$  contains  $(0)_0$ . This means that  $\sigma^c = \{(0)_0, (i, j)\} \cup X$  for some  $(i, j) \in V(\mathcal{G}'_{mn})$  and  $X \subseteq V(\mathcal{G}'_{mn})$  such that  $(i, j) \approx (0)_0$  and  $\mathcal{G}'_{mn}[X \cup \{(i, j)\}]$  is a complete graph. Since  $(i, j) \in V(\mathcal{G}'_{mn})$ , we have  $i \in [0, m-1]$ .

- Suppose  $i = 0$ . Then  $(i, j) = (j)_0 \approx (0)_0$  implies  $j \in [n-2] \setminus \{1\}$ . Since  $\mathcal{G}'_{mn}[X \cup \{(j)_0\}]$  is a complete graph,  $X$  satisfies one of the following conditions: (i)  $X = \emptyset$ , (ii)  $X = \{(j-1)_0\}$ , (iii)  $X = \{(n-1)_0\}$  when  $j = n-2$ , or (iv)  $X \subseteq \{(i', j) \mid i' \in [m-1]\}$  with  $X \neq \emptyset$ .

Using Proposition 3.13 and following the proof of Theorem 3.12, cases (i), (ii), and (iv) contradict  $\sigma \in \mathcal{C}_n$ . Hence, the only possibility is  $X = \{(n-1)_0\}$  when  $j = n-2$ . It follows that  $\sigma^c = \{(0)_0, (n-2)_0, (n-1)_0\}$ , and thus  $\sigma \in A$ .

- Suppose  $i \in [m-1]$ . In this case,  $(i, j) \approx (0)_0$  implies  $j \in [n-1]$ . Since  $\mathcal{G}'_{mn}[X \cup \{(i, j)\}]$  is a complete graph,  $X$  satisfies one of the following conditions: (i)  $X = \emptyset$ , (ii)  $X = \{(i, j-1)\}$ , (iii)  $X = \{(i, 0)\}$  when  $j = n-1$ , or (iv)  $X \subseteq \{(i', j) \mid i' \in [0, m-1] \setminus \{i\}\}$  with  $X \neq \emptyset$ .

Again, following the proof of Theorem 3.12 and using Proposition 3.13, cases (i) and (iv) contradict  $\sigma \in \mathcal{C}_n$ . Hence, either  $X = \{(i, j-1)\}$ , or  $X = \{(i, 0)\}$  when  $j = n-1$ . This means that either  $\sigma^c = \{(0)_0, (i, j-1), (i, j)\}$  for some  $i \in [m-1]$  and  $j \in [n-1]$ , or  $\sigma^c = \{(0)_0, (i, 0), (i, n-1)\}$  for some  $i \in [m-1]$ . Therefore  $\sigma \in B \sqcup C$ .

We conclude that  $\sigma \in A \sqcup B \sqcup C$ .

Conversely, let  $\sigma \in A \sqcup B \sqcup C$ .

- Suppose  $\sigma \in A$ . Then  $\sigma^c = \{(0)_0, (n-2)_0, (n-1)_0\}$ . Further,  $(0)_0 \approx (n-2)_0$  and  $(n-2)_0 \sim (n-1)_0$  implies  $\sigma \in \Delta_2^t(\mathcal{G}'_{mn})$  and  $\sigma \cup \{0\} \notin \Delta_2^t(\mathcal{G}'_{mn})$ . Therefore  $\sigma \in \mathcal{C}_1$  by Proposition 3.2 (i).

Let  $1 \leq r \leq n-3$ . Then  $(r)_0 \in \sigma$ . Since  $1 < (n-1) - r < n-1$ ,  $(r)_0 \approx (n-1)_0$ . Hence  $(\sigma \setminus \{(r)_0\}) \cup \{(0)_0\} \in \Delta_2^t(\mathcal{G}'_{mn})$ , and thus  $\sigma \setminus \{(r)_0\} \notin \mathcal{C}_1$  by Proposition 3.2 (i). From Remark 2.6 (iii),  $\sigma \in \mathcal{C}_{n-2}$ .

Now, since  $(0)_0 \sim (n-1)_0$ , we have  $\sigma \cup \{(n-2)_0\} \notin \Delta_2^t(\mathcal{G}'_{mn})$ . Then  $\mathcal{C}_{n-2} \subseteq \Delta_2^t(\mathcal{G}'_{mn})$  implies  $\sigma \cup \{(n-2)_0\} \notin \mathcal{C}_{n-2}$ . By Remark 2.6 (ii),  $\sigma \in \mathcal{C}_{n-1}$ .

We have  $\sigma \cup \{(n-1)_0\} = \{(0)_0, (n-2)_0\}$ . Clearly,  $\sigma \cup \{(0)_0, (n-1)_0\} \notin \Delta_2^t(\mathcal{G}'_{mn})$ . By Proposition 3.2 (i),  $\sigma \cup \{(n-1)_0\} \in \mathcal{C}_1$ . Applying Proposition 3.13 (i) to  $\sigma \cup \{(n-1)_0\}$  (for  $j = n-2$ ), we get  $\sigma \cup \{(n-1)_0\} \notin \mathcal{C}_{n-2}$ . Since  $\mathcal{C}_{n-1} \subseteq \mathcal{C}_{n-2}$ ,  $\sigma \cup \{(n-1)_0\} \notin \mathcal{C}_{n-1}$ . Thus  $\sigma \in \mathcal{C}_n$  by Remark 2.6 (ii).

- Suppose  $\sigma \in B$ . Then  $\sigma^c = \{(0)_0, (i, 0), (i, n-1)\}$  for some  $i \in [m-1]$ . Since  $(0)_0 \approx (i, n-1)$  and  $(i, 0) \sim (i, n-1)$ , it follows that  $\sigma \in \Delta_2^t(\mathcal{G}'_{mn})$  and  $\sigma \cup \{0\} \notin \Delta_2^t(\mathcal{G}'_{mn})$ . By Proposition 3.2 (i),  $\sigma \in \mathcal{C}_1$ .

Let  $1 \leq r \leq n-1$ . Then  $(r)_0 \in \sigma$ . Since  $i \neq 0$  and  $r \neq 0$ ,  $(r)_0 \approx (i, 0)$ . Hence  $(\sigma \setminus \{(r)_0\}) \cup \{(0)_0\} \in \Delta_2^t(\mathcal{G}'_{mn})$ . By Proposition 3.2 (i),  $\sigma \setminus \{(r)_0\} \notin \mathcal{C}_1$ . Therefore  $\sigma \in \mathcal{C}_n$  by Remark 2.6 (iii).

- Suppose  $\sigma \in C$ . Then  $\sigma^c = \{(0)_0, (i, j-1), (i, j)\}$  for some  $i \in [m-1], j \in [n-1]$ . Using the same argument as in the converse part of the proof of Theorem 3.12, we get  $\sigma \in \mathcal{C}_n$ .

We conclude that  $\mathcal{C}_n = A \sqcup B \sqcup C$ . Note that  $|A| = 1$ ,  $|B| = m-1$ , and  $|C| = (m-1)(n-1)$ . Thus,  $|A| + |B| + |C| = 1 + (m-1) + (m-1)(n-1) = n(m-1) + 1$ . Further, each  $\sigma \in A \sqcup B \sqcup C$  has dimension  $mn-4$ . Hence the result follows by Theorem 2.7 and Corollary 2.8.  $\square$

#### 4. CONCLUSION AND FUTURE DIRECTIONS

This article focuses on total 2-cut complexes of the  $p$ -th powers of cycle graphs  $C_n^p$ , the Cartesian product of complete graphs and path graphs  $K_m \square P_n$ , and the Cartesian product of complete graphs and cycle graphs  $K_m \square C_n$ .

We have determined the homotopy type of  $\Delta_2^t(C_n^p)$  for  $n = 2p+2$  and  $n \geq 3p+1$ , thus proving Conjecture 1.2. For  $2p+3 \leq n \leq 3p$ , we investigated  $\Delta_2^t(C_n^p)$  using SageMath by computing their homology groups (with coefficients in  $\mathbb{Z}$ ). Table 1 summarizes the homology data for  $\Delta_2^t(C_n^p)$  for small values of  $n$  and  $p$ . Blank entries in the table indicate that the corresponding complex  $\Delta_2^t(C_n^p)$  is void. An entry of the form  $i : \mathbb{Z}^\beta$  denotes that the  $i^{\text{th}}$  homology group is  $\mathbb{Z}^\beta$ . Entries above and below the colored region correspond to  $n = 2p+2$  and  $n \geq 3p+1$  respectively, which we have already proved in Theorem 1.3. Entries in the colored region correspond to  $2p+3 \leq n \leq 3p$ .

$\begin{smallmatrix} p \\ n \end{smallmatrix}$	3	4	5	6	7	8
8	2 : $\mathbb{Z}$					
9	4 : $\mathbb{Z}^2$					
10	6 : $\mathbb{Z}$	3 : $\mathbb{Z}$				
11	7 : $\mathbb{Z}$	5 : $\mathbb{Z}$				
12	8 : $\mathbb{Z}$	7 : $\mathbb{Z}^3$	4 : $\mathbb{Z}$			
13	9 : $\mathbb{Z}$	9 : $\mathbb{Z}$	7 : $\mathbb{Z}$			
14	10 : $\mathbb{Z}$	10 : $\mathbb{Z}$	8 : $\mathbb{Z}$	5 : $\mathbb{Z}$		
15	11 : $\mathbb{Z}$	11 : $\mathbb{Z}$	10 : $\mathbb{Z}^4$	8 : $\mathbb{Z}^2$		
16	12 : $\mathbb{Z}$	12 : $\mathbb{Z}$	12 : $\mathbb{Z}$	10 : $\mathbb{Z}$	6 : $\mathbb{Z}$	
17	13 : $\mathbb{Z}$	13 : $\mathbb{Z}$	13 : $\mathbb{Z}$	11 : $\mathbb{Z}$	9 : $\mathbb{Z}$	
18	14 : $\mathbb{Z}$	14 : $\mathbb{Z}$	14 : $\mathbb{Z}$	13 : $\mathbb{Z}^5$	12 : $\mathbb{Z}$	7 : $\mathbb{Z}$
19	15 : $\mathbb{Z}$	15 : $\mathbb{Z}$	15 : $\mathbb{Z}$	15 : $\mathbb{Z}$	13 : $\mathbb{Z}$	11 : $\mathbb{Z}$
20	16 : $\mathbb{Z}$	16 : $\mathbb{Z}$	16 : $\mathbb{Z}$	16 : $\mathbb{Z}$	14 : $\mathbb{Z}$	13 : $\mathbb{Z}^3$
21	17 : $\mathbb{Z}$	17 : $\mathbb{Z}$	17 : $\mathbb{Z}$	17 : $\mathbb{Z}$	16 : $\mathbb{Z}^6$	15 : $\mathbb{Z}$
22	18 : $\mathbb{Z}$	18 : $\mathbb{Z}$	18 : $\mathbb{Z}$	18 : $\mathbb{Z}$	18 : $\mathbb{Z}$	16 : $\mathbb{Z}$
23	19 : $\mathbb{Z}$	19 : $\mathbb{Z}$	19 : $\mathbb{Z}$	19 : $\mathbb{Z}$	19 : $\mathbb{Z}$	17 : $\mathbb{Z}$
24	20 : $\mathbb{Z}$	20 : $\mathbb{Z}$	20 : $\mathbb{Z}$	20 : $\mathbb{Z}$	20 : $\mathbb{Z}$	19 : $\mathbb{Z}^7$
25	21 : $\mathbb{Z}$	21 : $\mathbb{Z}$	21 : $\mathbb{Z}$	21 : $\mathbb{Z}$	21 : $\mathbb{Z}$	21 : $\mathbb{Z}$

TABLE 1. Non-zero homology  $\Delta_2^t(C_n^p)$

Based on the data in Table 1, we propose the following conjecture about the homotopy type of  $\Delta_2^t(C_n^p)$  for  $2p+3 \leq n \leq 3p$ .

**Conjecture 4.1.** *Let  $2p+3 \leq n \leq 3p$ , and let  $r$  be a positive integer such that  $2p + \frac{p}{r+1} < n \leq 2p + \frac{p}{r}$ .*

- (i) *If  $n = 2p + \frac{p}{r}$ , then  $\Delta_2^t(C_n^p) \simeq \bigvee_{n-2p-1} \mathbb{S}^{n-2r-3}$ .*
- (ii) *If  $2p + \frac{p}{r+1} < n < 2p + \frac{p}{r}$ , then  $\Delta_2^t(C_n^p) \simeq \mathbb{S}^{n-2r-4}$ .*

Observe that for fixed  $p$  and each  $n$  with  $2p+3 \leq n \leq 3p$ , there exists a unique positive integer  $r$  such that  $2p + \frac{p}{r+1} < n \leq 2p + \frac{p}{r}$ . Hence, Conjecture 4.1 addresses all  $n$  satisfying  $2p+3 \leq n \leq 3p$ .

The powers of cycle graphs  $C_n^p$  are circulant graphs  $C_n(\{1, 2, \dots, p\})$  and hence Cayley graphs of  $\mathbb{Z}_n$ . This raises the following natural question.

**Question 4.2.** What are the homotopy types of total cut complexes and cut complexes of general circulant graphs? More broadly, what about arbitrary Cayley graphs?

We have also determined the homotopy types of  $\Delta_2^t(K_m \square P_n) = \Delta_2(K_m \square P_n)$  and  $\Delta_2^t(K_m \square C_n) = \Delta_2(K_m \square C_n)$ . Using SageMath, we further investigated the total 3-cut and 3-cut complexes of these Cartesian products. We computed their homology groups (with coefficients in  $\mathbb{Z}$ ) for some small

values of  $m$  and  $n$ , which are presented in Tables 2, 3, 4 and 5. The notation  $i : \mathbb{Z}^\beta$  denotes that the  $i^{\text{th}}$  homology group is  $\mathbb{Z}^\beta$ , and  $*$  indicates that the corresponding entry is not known. The patterns observed in these tables lead us to Conjectures 4.3 to 4.6.

$\begin{smallmatrix} n \\ m \end{smallmatrix}$	3	4	5	6	7
2	$0 : \mathbb{Z}$	$2 : \mathbb{Z}^3$	$4 : \mathbb{Z}^6$	$6 : \mathbb{Z}^{10}$	$8 : \mathbb{Z}^{15}$
3	$3 : \mathbb{Z}^4$	$6 : \mathbb{Z}^{12}$	$9 : \mathbb{Z}^{24}$	$12 : \mathbb{Z}^{40}$	$15 : \mathbb{Z}^{60}$
4	$6 : \mathbb{Z}^9$	$10 : \mathbb{Z}^{27}$	$14 : \mathbb{Z}^{54}$	$*$	$*$
5	$9 : \mathbb{Z}^{16}$	$14 : \mathbb{Z}^{48}$	$*$	$*$	$*$
6	$12 : \mathbb{Z}^{25}$	$18 : \mathbb{Z}^{75}$	$*$	$*$	$*$
7	$15 : \mathbb{Z}^{36}$	$*$	$*$	$*$	$*$

TABLE 2. Non-zero homology of  $\Delta_3^t(K_m \square P_n)$ 

**Conjecture 4.3.** For  $m \geq 2$  and  $n \geq 3$ ,  $\tilde{H}_i(\Delta_3^t(K_m \square P_n)) = \begin{cases} \mathbb{Z}^{\frac{1}{2}(n^2-3n+2)(m-1)^2} & \text{if } i = mn - 6, \\ 0 & \text{otherwise.} \end{cases}$

$\begin{smallmatrix} n \\ m \end{smallmatrix}$	3	4	5	6
3	$4 : \mathbb{Z}^2$ $5 : \mathbb{Z}^6$	$7 : \mathbb{Z}^3$ $8 : \mathbb{Z}^{21}$	$10 : \mathbb{Z}^4$ $11 : \mathbb{Z}^{45}$	$13 : \mathbb{Z}^5$ $14 : \mathbb{Z}^{78}$
4	$7 : \mathbb{Z}^6$ $8 : \mathbb{Z}^{12}$	$11 : \mathbb{Z}^9$ $12 : \mathbb{Z}^{40}$	$15 : \mathbb{Z}^{12}$ $16 : \mathbb{Z}^{84}$	$*$
5	$10 : \mathbb{Z}^{12}$ $11 : \mathbb{Z}^{20}$	$15 : \mathbb{Z}^{18}$ $16 : \mathbb{Z}^{65}$	$*$	$*$
6	$13 : \mathbb{Z}^{20}$ $14 : \mathbb{Z}^{30}$	$*$	$*$	$*$

TABLE 3. Non-zero homology of  $\Delta_3(K_m \square P_n)$ 

**Conjecture 4.4.** For  $m, n \geq 3$ ,  $\tilde{H}_i(\Delta_3(K_m \square P_n)) = \begin{cases} \mathbb{Z}^{\frac{1}{2}(m^2-3m+2)(n-1)} & \text{if } i = mn - 5, \\ \mathbb{Z}^{\frac{m}{2}(mn-m-2)(n-2)} & \text{if } i = mn - 4, \\ 0 & \text{otherwise.} \end{cases}$

$\begin{smallmatrix} n \\ m \end{smallmatrix}$	3	4	5	6	7	8	9	10
2	void complex	$2 : \mathbb{Z}^9$	$4 : \mathbb{Z}^{14}$	$6 : \mathbb{Z}^{22}$	$8 : \mathbb{Z}^{29}$	$10 : \mathbb{Z}^{37}$	$12 : \mathbb{Z}^{46}$	$14 : \mathbb{Z}^{56}$
3	$3 : \mathbb{Z}$	$6 : \mathbb{Z}^{30}$	$9 : \mathbb{Z}^{48}$	$12 : \mathbb{Z}^{73}$	$15 : \mathbb{Z}^{99}$	$*$	$*$	$*$
4	$6 : \mathbb{Z}^3$	$10 : \mathbb{Z}^{63}$	$14 : \mathbb{Z}^{102}$	$*$	$*$	$*$	$*$	$*$
5	$9 : \mathbb{Z}^6$	$14 : \mathbb{Z}^{108}$	$*$	$*$	$*$	$*$	$*$	$*$
6	$12 : \mathbb{Z}^{10}$	$18 : \mathbb{Z}^{165}$	$*$	$*$	$*$	$*$	$*$	$*$
7	$15 : \mathbb{Z}^{15}$	$*$	$*$	$*$	$*$	$*$	$*$	$*$

TABLE 4. Non-zero homology of  $\Delta_3^t(K_m \square C_n)$ 

**Conjecture 4.5.** (i) For  $m \geq 3$  and  $n = 3$ ,  $\tilde{H}_i(\Delta_3^t(K_m \square C_n)) = \begin{cases} \mathbb{Z}^{\frac{1}{2}(m^2-3m+2)} & \text{if } i = mn - 6, \\ 0 & \text{otherwise.} \end{cases}$

(ii) For  $m \geq 2$  and  $n \in \{4, 5\}$ ,  $\tilde{H}_i(\Delta_3^t(K_m \square C_n)) = \begin{cases} \mathbb{Z}^{\frac{m-1}{2}((m-1)n^2-(m-3)n-2)} & \text{if } i = mn - 6, \\ 0 & \text{otherwise.} \end{cases}$

(iii) For  $m \geq 2$  and  $n \geq 6$ ,  $\tilde{H}_i(\Delta_3^t(K_m \square C_n)) = \begin{cases} \mathbb{Z}^{\frac{1}{2}((m-1)^2n^2-(m^2-4m+3)n+2)} & \text{if } i = mn - 6, \\ 0 & \text{otherwise.} \end{cases}$



$\begin{smallmatrix} n \\ m \end{smallmatrix}$	4	5	6	7	8
2	$3 : \mathbb{Z}$ $4 : \mathbb{Z}^4$	$6 : \mathbb{Z}^{11}$	$8 : \mathbb{Z}^{25}$	$10 : \mathbb{Z}^{43}$	$12 : \mathbb{Z}^{65}$
3	$7 : \mathbb{Z}^6$ $8 : \mathbb{Z}^{12}$	$10 : \mathbb{Z}^5$ $11 : \mathbb{Z}^{31}$	$13 : \mathbb{Z}^6$ $14 : \mathbb{Z}^{64}$	*	*
4	$11 : \mathbb{Z}^{15}$ $12 : \mathbb{Z}^{24}$	$15 : \mathbb{Z}^{15}$ $16 : \mathbb{Z}^{61}$	*	*	*
5	$15 : \mathbb{Z}^{28}$ $16 : \mathbb{Z}^{40}$	*	*	*	*

TABLE 5. Non-zero homology of  $\Delta_3(K_m \square C_n)$ 

**Conjecture 4.6.** (i) For  $m \geq 2$  and  $n = 4$ ,  $\tilde{H}_i(\Delta_3(K_m \square C_n)) = \begin{cases} \mathbb{Z}^{2m^2-5m+3} & \text{if } i = mn - 5, \\ \mathbb{Z}^{2m(m-1)} & \text{if } i = mn - 4, \\ 0 & \text{otherwise.} \end{cases}$

(ii) For  $m = 2$  and  $n \geq 5$ ,  $\tilde{H}_i(\Delta_3(K_m \square C_n)) = \begin{cases} \mathbb{Z}^{2n^2-8n+1} & \text{if } i = mn - 4, \\ 0 & \text{otherwise.} \end{cases}$

## ACKNOWLEDGEMENT

The first author is supported by HTRA fellowship by IIT Mandi, India. The second author is supported by the seed grant project IITM/SG/SMS/95 by IIT Mandi, India.

## REFERENCES

- [1] E. Babson and D. N. Kozlov. Complexes of graph homomorphisms. *Israel J. Math.*, 152:285–312, 2006.
- [2] M. Bate, B. Everitt, S. Ford, and E. Ramos. Homology of matching complexes and representations of symmetric groups. *arXiv preprint arXiv:2312.13750*, 2023.
- [3] M. Bayer, M. Denker, M. J. Milutinović, R. Rowlands, S. Sundaram, and L. Xue. Topology of cut complexes of graphs. *SIAM J. Discrete Math.*, 38(2):1630–1675, 2024.
- [4] M. Bayer, M. Denker, M. J. Milutinović, R. Rowlands, S. Sundaram, and L. Xue. Total cut complexes of graphs. *Discrete Comput. Geom.*, 73(2):500–527, 2024.
- [5] A. Björner, L. Lovász, S. T. Vrećica, and R. T. Živaljević. Chessboard complexes and matching complexes. *J. London Math. Soc. (2)*, 49(1):25–39, 1994.
- [6] O. Bobrowski and M. Kahle. Topology of random geometric complexes: a survey. *J. Appl. Comput. Topol.*, 1(3-4):331–364, 2018.
- [7] J. A. Bondy and U. S. R. Murty. *Graph theory*, volume 244 of *Graduate Texts in Mathematics*. Springer, New York, 2008.
- [8] S. Bouc. Homologie de certains ensembles de 2-sous-groupes des groupes symétriques. *J. Algebra*, 150(1):158–186, 1992.
- [9] G. Carlsson. Topology and data. *Bull. Amer. Math. Soc.*, 46(2):255–308, 2009.
- [10] H. Chandrakar, N. R. Hazra, D. Rout, and A. Singh. Topology of total cut and cut complexes of grid graphs. *arXiv preprint arXiv:2408.07646*, 2024.
- [11] P. Chauhan and S. Shukla. Shellability of 3-cut complexes of power of cycle graphs. *In preparation*, 2025.
- [12] P. Chauhan, S. Shukla, and K. Vinayak. Shellability of 3-cut complexes of squared cycle graphs. *J. Homotopy Relat. Struct.*, 20(1):163–193, 2025.
- [13] P. Deshpande, S. Shukla, and A. Singh. Distance  $r$ -domination number and  $r$ -independence complexes of graphs. *European J. Combin.*, 102:Paper No. 103508, 14, 2022.
- [14] A. Dochtermann and A. Engström. Algebraic properties of edge ideals via combinatorial topology. *Electron. J. Combin.*, 16(2):Research Paper 2, 24, 2009.
- [15] A. Dochtermann and A. Engström. Cellular resolutions of cointerval ideals. *Math. Z.*, 270(1-2):145–163, 2012.
- [16] J. A. Eagon and V. Reiner. Resolutions of Stanley-Reisner rings and Alexander duality. *J. Pure Appl. Algebra*, 130(3):265–275, 1998.
- [17] H. Edelsbrunner and J. L. Harer. *Computational topology: an introduction*. American Mathematical Society, 2010.
- [18] R. Forman. Morse theory for cell complexes. *Adv. Math.*, 134(1):90–145, 1998.
- [19] R. Fröberg. On Stanley-Reisner rings. In *Topics in algebra, Part 2 (Warsaw, 1988)*, volume 26, Part 2 of *Banach Center Publ.*, pages 57–70. PWN, Warsaw, 1990.
- [20] C. Giusti, R. Ghrist, and D. S. Bassett. Clique topology reveals intrinsic geometric structure in neural correlations. *Proc. Natl. Acad. Sci.*, 112(44):13455–13460, 2015.
- [21] M. Gromov. Hyperbolic groups. In *Essays in group theory*, volume 8 of *Math. Sci. Res. Inst. Publ.*, pages 75–263. Springer, New York, 1987.
- [22] R. K. Gupta, S. Sarkar, and S. Shukla. On the Vietoris-Rips complexes of integer lattices. *arXiv preprint arXiv:2511.04238*, 2025.

- [23] A. Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002.
- [24] J. Herzog and T. Hibi. *Monomial ideals*, volume 260 of *Graduate Texts in Mathematics*. Springer-Verlag London, Ltd., London, 2011.
- [25] J. Jonsson. *Simplicial complexes of graphs*, volume 1928 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2008.
- [26] M. Kahle. Sharp vanishing thresholds for cohomology of random flag complexes. *Ann. of Math. (2)*, 179(3):1085–1107, 2014.
- [27] M. Kneser. Aufgabe 300. *Jber. Deutsch. Math.-Verein*, 58, 1955.
- [28] D. Kozlov. *Combinatorial algebraic topology*, volume 21 of *Algorithms and Computation in Mathematics*. Springer, Berlin, 2008.
- [29] L. Lovász. Kneser’s conjecture, chromatic number, and homotopy. *J. Combin. Theory Ser. A*, 25(3):319–324, 1978.
- [30] J. Matoušek. *Using the Borsuk-Ulam theorem*. Universitext. Springer-Verlag, Berlin, 2003. Lectures on topological methods in combinatorics and geometry, Written in cooperation with Anders Björner and Günter M. Ziegler.
- [31] T. Matsushita. Dominance complexes and vertex cover numbers of graphs. *J. Appl. Comput. Topol.*, 7(2):363–368, 2023.
- [32] R. Meshulam. The clique complex and hypergraph matching. *Combinatorica*, 21(1):89–94, 2001.
- [33] R. Meshulam. Domination numbers and homology. *J. Combin. Theory Ser. A*, 102(2):321–330, 2003.
- [34] R. Meshulam and N. Wallach. Homological connectivity of random  $k$ -dimensional complexes. *Random Structures Algorithms*, 34(3):408–417, 2009.
- [35] E. Miller and B. Sturmfels. *Combinatorial Commutative Algebra*, volume 227 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2005.
- [36] V. Reiner and J. Roberts. Minimal resolutions and the homology of matching and chessboard complexes. *J. Algebraic Combin.*, 11(2):135–154, 2000.
- [37] R. Santhanam and S. Shukla. Vertex cut of a graph and connectivity of its neighbourhood complex. *Discrete Math.*, 346(8):Paper No. 113432, 11, 2023.
- [38] Y. Shen, Z. Song, F. Yu, L. W. Zhou, and J. Zhuang. Homotopy type of total cut complexes of squared cycle graphs. *arXiv preprint arXiv:2510.22574*, 2025.
- [39] S. Shukla and D. Yogeshwaran. Spectral gap bounds for the simplicial Laplacian and an application to random complexes. *J. Combin. Theory Ser. A*, 169:105134, 20, 2020.
- [40] R. P. Stanley. *Combinatorics and commutative algebra*, volume 41 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, second edition, 1996.
- [41] D. B. West. *Introduction to graph theory*. Prentice Hall, Inc., Upper Saddle River, N.J., second edition, 2001.
- [42] R. Woodroffe. Matchings, coverings, and Castelnuovo-Mumford regularity. *J. Commut. Algebra*, 6(2):287–304, 2014.
- [43] M. C. B. Zaremsky. Bestvina-Brady discrete Morse theory and Vietoris-Rips complexes. *Amer. J. Math.*, 144(5):1177–1200, 2022.

SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, IIT MANDI, INDIA  
 Email address: d22037@students.iitmandi.ac.in

SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, IIT MANDI, INDIA  
 Email address: samir@iitmandi.ac.in

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KENTUCKY, USA  
 Email address: kumar.vinayak@uky.edu