

DETECT DUALITY OBSTRUCTION OF CALIBRATIONS IN SMOOTH CATEGORY

YONGSHENG ZHANG

ABSTRACT. This paper consists of three parts: a. exhibit a new gluing result which can dramatically simplify extensions of calibration pairs; b. observe that every Lawlor cone can support coflat calibrations singular only at the origin; c. show that there exist many Lawlor cones which cannot support any smooth calibrations. As an application, we extend our previous work on detecting duality obstruction of calibrations in the smooth category.

1. INTRODUCTION

In [Zha17, Zha18] the author developed a very “soft” framework about extending calibration pairs which play effective roles in detecting duality obstruction and realizing problem about singularities (to be explained in below). By combining these methods the framework has been carried forward to attack several long-term standing conjectures regarding area-minimizing currents, e.g. [Liu24, Liua, Liub].

The reason for the calibration method being so useful is due to the fundamental theorem of calibrated geometry. Detailed explanations about the theorem will be given in §2. A special situation of the theorem says that any calibrated closed smooth submanifold is mass-minimizing in its homology class. So, the **duality question** in the smooth category is the following:

(★) If a closed smooth submanifold M in some Riemannian manifold (X, g) is already mass-minimizing in its homology class, then can we have some smooth calibration form to calibrate M ?

In Example 3 of [Zha17] we observe some intriguing duality obstruction phenomenon for the first time. It leads us to some smooth closed hypersurface M of dimension $m \geq 7$ in some Riemannian manifold (X, g) of dimension $m + 1$ which minimizes mass among its homology class $[M]$ but cannot be calibrated by any smooth calibration form. Actually, beyond [Zha17], examples of high codimension with the duality obstruction can be gained. Among other constructions,

(*) for simplicity, consider the Cartesian product $(X \times \mathbb{S}^\ell, g \oplus g_0)$ where (\mathbb{S}^ℓ, g_0) is the standard unit sphere with $\ell \geq 1$. Take $q \in (\mathbb{S}^\ell, g_0)$. Then the submanifold $M \times \{q\}$ gives us a homologically mass-minimizing smooth submanifold of codimension $\ell + 1$ and it cannot be calibrated by any smooth calibration form.

Date: December 5, 2025.

2020 Mathematics Subject Classification. Primary 53C38.

Key words and phrases. Calibration, duality obstruction, realization problem.

This obstruction in (*) comes from some delicate balance between geometry and topology. And it has been further pointed out on page 431 of [FH17] that our obstruction even exists in local around the submanifold in the category of continuous calibration forms. To distinguish this local obstruction (*) from global ones, the author used the following example to explain to others since 2020 and it should be faithfully mentioned that such kind of illustration can trace back at least to [FH17]. Let $Y = S^1 \times S^{2020}$ and choose a closed embedded curve γ_2 to be a representative of twice the generator e of $H_1(Y; \mathbb{Z})$. Then as in [Zha17] we can conformally change background metric such that γ_2 becomes the unique length minimizer in its homology class. As a result, any length minimizer γ_1 in class e cannot be calibrated by any smooth calibration since otherwise the integral current $2\gamma_1$ is also calibrated and thus length minimizing in $[\gamma_2]$ as well which contradicts with the uniqueness in the above. Here the obstruction comes from global in the sense that one cannot apply the idea in a sufficiently small neighborhood of γ_2 for the duality obstruction phenomenon.

In this paper, we extend our (localizable) duality obstruction for question (★) to genuine high codimension situations beyond that in (*). The idea is to find minimizing cones which support coflat calibrations singular only at the origins but cannot be calibrated by any smooth calibration form, and then apply our extension result for calibration pairs in [Zha17].

To be able to include more general local models which we can start with, we establish the following two results.

Theorem 1.1. *Let L be a closed embedded submanifold in $\mathbb{S}^N \subset \mathbb{R}^{N+1}$. If the cone $C(L) = \{tp : t \in (0, \infty), p \in L\}$ can be shown area-minimizing by Lawlor's criterion, then $C(L)$ supports coflat calibrations singular only at the origin.*

Remark 1.2. *When a closed submanifold $L \subset \mathbb{S}^N$ is not totally geodesic, the cone $C(L)$ has only one singular point at the origin and it is called a regular cone.*

Remark 1.3. *We say a current supports a calibration if it is calibrated by the calibration, see Definition 2.5.*

Theorem 1.4. *Let L_i be a closed embedded minimal submanifold in \mathbb{S}^{N_i} for $i = 1, \dots, n$. Suppose that L_1 is of codimension one. Then the cone $C(L_1 \times \dots \times L_n)$ over their minimal product $L_1 \times \dots \times L_n$ supports no smooth calibration.*

In conjunction with recent configuration result for area-minimizing cones in [Zha], these two theorems in particular imply the following.

Corollary 1.5. *Given any closed embedded minimal hypersurface L in $\mathbb{S}^N \subset \mathbb{R}^{N+1}$, let $L^{\times n}$ denote the minimal product of n copies of L . Then, when n is sufficiently large, the cone $C(L^{\times n})$ over $L^{\times n}$ supports coflat calibrations singular only at the origin but no smooth calibration forms.*

The paper is organized as follows. To be self-contained, we review some basics and useful tools from calibrated geometry in §2 and moreover establish a much better gluing result in §3 which can significantly simplify/improve the gluing procedures in the framework. A proof of Theorem 1.1 will be included in §4, and a

shortcut for the realization result in [Zha18] will be found. In §5 we give a proof of Theorem 1.4 by virtue of a result by J. Simons. In §6 we extend our duality obstruction beyond (*).

2. BASICS OF CALIBRATIONS

Let us first review some necessary definitions and basic properties in the theory of calibrations.

2.1. Definitions and basic properties. Let us begin with smooth forms.

Definition 2.1. Let ϕ be a smooth m -form on a Riemannian manifold (X, g) . At a point $x \in X$ the **comass** of ϕ_x is defined to be

$$\|\phi\|_{x,g}^* = \max \{ \phi_x(\vec{V}_x) : \vec{V}_x \text{ is a unit simple } m\text{-vector at } x \}.$$

Here “simple” means $\vec{V}_x = e_1 \wedge e_2 \cdots \wedge e_m$ for some $e_i \in T_x X$.

A remark in [Zha17] about the comass that we shall use is the following.

Lemma 2.2. At a point x where $\phi_x \neq 0$, we have

$$\begin{aligned} \|\phi\|_{x,g}^* &= \max \{ \phi(\vec{V}_x) : \vec{V}_x \text{ is a simple } m\text{-vector at } x \text{ with } \|\vec{V}_x\|_g = 1 \} \\ &= \max \{ 1/\|\vec{V}_x\|_g : \vec{V}_x \text{ is a simple } m\text{-vector at } x \text{ with } \phi(\vec{V}_x) = 1 \} \\ &= 1/\min \{ \|\vec{V}_x\|_g : \vec{V}_x \text{ is a simple } m\text{-vector at } x \text{ with } \phi(\vec{V}_x) = 1 \}. \end{aligned}$$

The dual complex of smooth forms gives us currents. To be more precise, by $(\mathcal{E}'_*(X), d)$ we mean the dual complex of the *de Rham* complex of X . Elements of $\mathcal{E}'_m(X)$ are m -dimensional *de Rham* currents (with compact support) and d is the adjoint of exterior differentiation.

Definition 2.3. The **mass** $\mathbf{M}(T)$ of $T \in \mathcal{E}'_m(X)$ is defined to be

$$\sup \{ T(\psi) : \psi \text{ smooth } m\text{-form with } \sup_{x \in X} \|\psi\|_{x,g}^* \leq 1 \}.$$

When T has finite mass, there exists a unique Radon measure $\|T\|$ satisfying

$$\int_X f \cdot d\|T\| = \sup \{ T(\psi) : \|\psi\|_{x,g}^* \leq f(x) \}$$

for any nonnegative continuous function f on X . Moreover, the Radon-Nikodym Theorem asserts the existence of a $\|T\|$ measurable tangent m -vector field \vec{T} a.e. with vectors $\vec{T}_x \in \Lambda^m T_x X$ of unit length in the dual norm of the comass norm, with

$$(2.1) \quad T(\psi) = \int_X \psi_x(\vec{T}_x) d\|T\|(x) \text{ for any smooth } m\text{-form } \psi,$$

or briefly $T = \vec{T} \cdot \|T\|$ a.e. $\|T\|$.

The groundbreaking paper [FF60] by Federer and Fleming introduced the concepts of normal currents and integral currents. Let $\mathbb{M}_m(X) = \{T \in \mathcal{E}'_m(X) : \mathbf{M}(T) < \infty\}$. Then $N_m(X) = \{T \in \mathbb{M}_m(X) : dT \in \mathbb{M}_{m-1}(X)\}$ is the space of m -dimensional

normal currents. Note that $(N_*(X), d)$ form a chain complex and moreover by de Rham, Federer and Fleming, we have natural isomorphisms

$$(2.2) \quad H_*(\mathcal{E}'_*(X)) \cong H_*(X; \mathbb{R}) \cong H_*(N_*(X)).$$

In this paper we adopt the following terminologies.

Definition 2.4. A smooth form ϕ on (X, g) is called a **calibration** if $\sup_X \|\phi\|_g^* = 1$ and $d\phi = 0$. Such a triple (X, ϕ, g) is called a **calibrated manifold**.

Definition 2.5. Let ϕ be a calibration on (X, g) . We say that a current T of local finite mass is **calibrated** by ϕ , if $\phi_x(\vec{T}_x) = 1$ a.a. $x \in X$ for $\|T\|$. In this case we say T **supports** calibration ϕ .

The fundamental theorem of calibrated geometry below says that every calibrated current is mass-minimizing in its homology class. By default the homology is taken for normal currents but in fact it holds for competitors of *de Rham* currents.

Theorem 2.6 ([HL82a]). If T is a calibrated current with compact support in (X, ϕ, g) and T' is any compactly supported current homologous to T (i.e., $T - T'$ is a boundary and in particular $dT = dT'$), then

$$\mathbf{M}(T) \leq \mathbf{M}(T')$$

with equality if and only if T' is calibrated as well.

The subclass $\mathcal{R}_m(X) \subset \mathbb{M}_m(X)$ of rectifiable m -currents is the closure of the integral Lipschitz m -chains in the \mathbf{M} -topology. Then $I_m(X) := \{T \in \mathcal{R}_m(X) : dT \in \mathcal{R}_{m-1}(X)\}$ is the set of integral m -currents. Federer and Fleming [FF60] also established the corresponding isomorphism

$$(2.3) \quad H_*(I_*(X)) \cong H_*(X; \mathbb{Z}).$$

So a calibrated integral current minimizes the mass not only in the homology class of integral currents but also that for normal currents. If one merely focuses on its minimality in (2.3) then it is called (homologically) area-minimizing.

In practice, sometimes one cannot avoid calibrations with singularities.

Definition 2.7. Let ϕ be a calibration of degree m on $X - S_\phi$, where S_ϕ is a closed subset of X of Hausdorff m -measure zero. Then ϕ is called a **coflat calibration** on X . We say ϕ **calibrates** a current, if it is calibrated by ϕ on $X - S_\phi$. Similarly, in such a case we say the current **supports** the coflat calibration ϕ .

A coflat version (Theorem 4.9 in [HL82a]) of the fundamental theorem of calibrated geometry is also true. Namely, a current calibrated by a coflat calibration ϕ in $X - S_\phi$ is homologically mass-minimizing in the sense of Theorem 2.6. More generally, the same property holds even for certain non-continuous ‘‘calibrations’’ with larger dimensional singular set, e.g. see Lawlor’s calibrations in §4, which provide a sufficient criterion for a regular cone to be area-minimizing.

Although our methods in [Zha17, Zha18] are quite soft, however there exists certain local rigidity in the sense that after blowing-up the local models always

live in some Euclidean spaces with standard metrics. Hence, to begin with suitable local models is important to us.

According to a fundamental result (Theorem 5.4.3 in [Fed69]) there always exists at least one tangent cone for area-minimizing integral current, and moreover, the tangent cone is area-minimizing. In this paper we mainly focus on the situation of regular area-minimizing cones in standard Euclidean space (\mathbb{R}^n, g_E) .

2.2. Four lemmas. A beautiful canonical decomposition lemma for an m -form with respect to certain m -plane in \mathbb{R}^n is the following (its proof in [HL82b] inspires a remark for $m = 1$ and our Lemma 2.12, so we include it in full details).

Lemma 2.8 (Lemma 2.12 in [HL82b]). *Let $\xi \in \Lambda^m \mathbb{R}^n$ be a simple m -vector with $V = \text{span } \xi$. Suppose $\phi \in \Lambda^m \mathbb{R}^n$ satisfies $\phi(\xi) = 1$. Then there exists a unique oriented complementary subspace W to V with the following property. For any basis v_1, \dots, v_n of \mathbb{R}^n such that $\xi = v_1 \wedge \dots \wedge v_m$ and v_{m+1}, \dots, v_n is basis for W , one has that*

$$(2.4) \quad \phi = v_1^* \wedge \dots \wedge v_m^* + \sum a_I v_I^*,$$

where $a_I = 0$ whenever $i_{m-1} \leq m$. Here $I = \{i_1, \dots, i_m\}$ with $i_1 < \dots < i_m$.

Remark 2.9. *Note that the decomposition does not depends on ambient metric.*

Proof. Define linear map

$$\lambda : \Lambda^{m-1} V \longrightarrow \text{Hom}(\mathbb{R}^n, \mathbb{R})$$

by $\eta \mapsto \eta \lrcorner \phi$, namely, $\lambda(\eta)(v) = \phi(\eta \wedge v)$. Set $\eta_i = (-1)^{m+i} v_1 \wedge \dots \wedge \hat{v}_i \wedge \dots \wedge v_m$ for $i = 1, \dots, m$. Consider forms $\lambda_i = \lambda(\eta_i)$. Then we have

$$(2.5) \quad \lambda_i(v_j) = (-1)^{m+i} \phi(v_1 \wedge \dots \wedge \hat{v}_i \wedge \dots \wedge v_m \wedge v_j) = \delta_{ij}$$

for $1 \leq i \leq m$ and $1 \leq j \leq m$. Since $(\lambda_1 \wedge \dots \wedge \lambda_m)(\xi) = 1$, it follows that

$$(2.6) \quad W \triangleq \bigcap_{i=1}^m \ker(\lambda_i)$$

is a subspace complementary to $V = \text{span } \xi$.

Choose v_{m+1}, \dots, v_n to be any basis of W . Then $\phi(v_1 \wedge \dots \wedge \hat{v}_i \wedge \dots \wedge v_m \wedge v_j) = \pm \lambda_i(v_j) = 0$ for $1 \leq i \leq m$ and $m+1 \leq j \leq n$. Hence the proof gets completed. \square


A powerful lemma to nicely control comass of a form by altering metics is the following.

Lemma 2.10 (Lemma 2.14 in [HL82b]). *Let ϕ , $V = \text{span } \xi$, and W be given as in Lemma 2.8. Consider an inner product $\langle \cdot, \cdot \rangle$ on \mathbb{R}^n such that $V \perp W$ and $\|\xi\| = 1$. Choose any constant $C^2 > \binom{n}{m} \|\phi\|^*$ and define a new inner product on $\mathbb{R}^n = V \oplus W$ by setting $\langle \cdot, \cdot \rangle' = \langle \cdot, \cdot \rangle_V + C^2 \langle \cdot, \cdot \rangle_W$. Then under this new metric we have*

$$\|\phi\|^* = 1 \text{ and } \phi(\xi) = \|\xi\| = 1.$$

Remark 2.11. *If $\phi(\xi) = \vartheta$ (positive) not necessarily one, one can apply Lemma 2.8 to $\vartheta^{-1} \phi$ for $\|\phi\|^* = \vartheta$, $\|\xi\| = 1$ and $\phi(\xi) = \vartheta$ by choosing $C^2 > \vartheta^{-1} \binom{n}{m} \|\phi\|^*$.*

In fact the ideas of Lemmas 2.8 and 2.10 also work for $m = 1$. To illustrate how they apply, we consider the situation in the figure. Although we do not need a



metric a priori, for simplicity, let ξ be of unit length and t a vector with $\langle t, t \rangle_{g_E} > 1$ and $\langle \xi, t \rangle_{g_E} = 1$ in (\mathbb{R}^n, g_E) . Then we can construct a form $\phi(\cdot) = \langle t, \cdot \rangle_{g_E}$ which has the property $\phi(\xi) = 1$.

Now directly define $W \triangleq \ker(\phi)$. Then $\{v \in \mathbb{R}^n : \phi(v) = \langle \xi, v \rangle = 1\} = \{\xi\} + W$. Define $g'(\cdot, \cdot) = \langle \cdot, \cdot \rangle_V + C^2 \langle \cdot, \cdot \rangle_W$ as in Lemma 2.10. Then, by Lemma 2.2 we have

$$\begin{aligned} \|\phi\|_{g'}^* &= 1 / \min\{\|v\|_{g'} : v \text{ is a vector at } 0 \text{ with } \phi(v) = 1\} \\ &= 1 / \sqrt{\|\xi\|^2 + C^2 \langle v - \xi, v - \xi \rangle_W} \\ &\leq 1 \end{aligned}$$

with equality if and only if $v = \xi$. Note that here any $C^2 > 0$ is sufficient for the conclusion $\|\phi\|_{g'}^* = 1$ and $\phi(\xi) = \|\xi\|_{g'} = 1$.

It should be particularly emphasized that the distribution W in the above for generating metric $g'(\cdot, \cdot) = \langle \cdot, \cdot \rangle_V + C^2 \langle \cdot, \cdot \rangle_W$ to make $\|\phi\|_{g'}^* = 1$ and $\phi(\xi) = \|\xi\|_{g'} = 1$ is rigid and not allowed to vary at all. For example, if one prefers to use $W = V^\perp$ and enlarge C^2 for $g' = g_E|_V + C^2 g_E|_{V^\perp}$, then no matter how large C^2 one chooses it is simply impossible to have $\|\phi\|_{g'}^* = 1$. Similar phenomenon also occurs for simple m -form with constant coefficients when $m > 1$.

Although Lemma 2.8 is independent on metric, sometimes when a prior metric gets involved we can have a coupled decomposition. An immediate application to the case where ϕ is a calibration with constant coefficients in (\mathbb{R}^n, g_E) and $\phi(\xi) = 1$ for some unit simple m -vector ξ can be derived.

Lemma 2.12 (Decomposition for calibration form). *Let v_1, \dots, v_n be an orthonormal basis of (\mathbb{R}^n, g_E) with $\xi = v_1 \wedge \dots \wedge v_m$ and $V = \text{span } \xi$. Assume that ϕ is a calibration with constant coefficients which calibrates the oriented V . Then W in Lemma 2.8 has to be $\text{span}\{v_{m+1}, \dots, v_n\}$ and as in Lemma 2.8 we have*

$$(2.7) \quad \phi = v_1^* \wedge \dots \wedge v_m^* + \sum a_I v_I^*,$$

where $a_I = 0$ whenever $i_{m-1} \leq m$. Here $I = \{i_1, \dots, i_m\}$ with $i_1 < \dots < i_m$.

Proof. For $m \geq 2$, as in the proof of Lemma 2.8, linear maps λ_i for $i = 1, \dots, m$ can be defined. Now we must have

$$(2.8) \quad \lambda_i(v_j) = (-1)^{m+i} \phi(v_1 \wedge \dots \wedge \hat{v}_i \wedge \dots \wedge v_m \wedge v_j) = \delta_{ij}$$

for $1 \leq i \leq m$ and $1 \leq j \leq n$ (not merely for $1 \leq j \leq m$). Otherwise, if $\lambda_i(v_j) = a \neq 0$ in (2.8) for some $1 \leq i \leq m < j \leq n$, then consider the unit simple m -vector

$$\tilde{\xi} = v_1 \wedge \dots \wedge \left(\frac{1}{\sqrt{1+a^2}} v_i + \frac{a}{\sqrt{1+a^2}} v_j \right) \wedge \dots \wedge v_m$$

and its pairing $\phi(\tilde{\xi}) = \sqrt{1+a^2} > 1$ contradicting with the assumption that ϕ is a calibration form.

Due to the dimension reason, $W = \bigcap_{i=1}^m \ker(\lambda_i) = \text{span}\{v_{m+1}, \dots, v_n\}$.

When $m = 1$, instead of constructing λ_i we determine W using ϕ directly and the same kind of arguments apply for the conclusion. \square

A key lemma that we establish in [Zha17] is the following.

Lemma 2.13 (Comass control for gluing procedure). *For any m -form ϕ , positive functions a and b , and metrics g_1 and g_2 , it follows*

$$(2.9) \quad \|\phi\|_{ag_1+bg_2}^* \leq \frac{1}{\sqrt{a^m \cdot \frac{1}{\|\phi\|_{g_1}^{*2}} + b^m \cdot \frac{1}{\|\phi\|_{g_2}^{*2}}}}$$

where $\frac{1}{0}$ and $\frac{1}{+\infty}$ are identified with $+\infty$ and 0 respectively.

However, an apparent dissatisfactory is that even with $\|\phi\|_{g_1}^{*2} = 1$ and $\|\phi\|_{g_2}^{*2} = 1$ (except for $m = 1$) one cannot draw the conclusion that $\|\phi\|_{(1-s)g_1+sg_2}^{*2} \leq 1$ for $s \in (0, 1)$. We shall obtain a better lemma regarding this in the next section.

3. NEW GLUING RESULT FOR THE EXTENSION OF CALIBRATION PAIRS

In this section we explain how to extend calibration pairs. Case with some more general situation was studied, but we shall restrict ourselves to the simplest situation. Let (S, o) be an oriented connected compact submanifold with no boundary and with only one singular point o . Then the extension result of calibration pair means the following.

Theorem 3.1 (Theorem 4.6 in [Zha17]). *Given (S, o) in (X, g) with $[S] \neq [0] \in H_m(X; \mathbb{R})$. If $B_\epsilon(o; g) \cap S$ is calibrated by a calibration in some ϵ -ball $(B_\epsilon(o; g), g)$ centered at o , then there exists a metric \hat{g} coinciding with g on $B_{\frac{\epsilon}{3}}(o; g)$ such that S can be calibrated in (X, \hat{g}) by some calibration.*

Actually, we can simplify the gluing procedures in [Zha17] based on a satisfactory new result in §3.1 which can greatly improve the key control Lemma 2.13.

3.1. A new gluing result. In this subsection, we assume that $\phi (\neq 0)$ is an m -form with constant coefficients in \mathbb{R}^n and that $\|\phi\|_{g_1}^{*2} \leq 1$ and $\|\phi\|_{g_2}^{*2} \leq 1$ for two metrics g_1, g_2 (with constant coefficients) on the vector space \mathbb{R}^n .

A naive question is whether or not the comass of ϕ remains no larger than one for the family of metrics $g(s) = (1-s)g_1 + sg_2$. We confirm this affirmatively.

Theorem 3.2 (Nice control for gluing metrics). *Assume that ϕ, g_1, g_2 and $g(s)$ are given as above. Then $\|\phi\|_{g(s)}^* \leq 1$.*

Proof. There are three steps and the first two are the same as in the proof of Lemma 2.13 in [Zha17]. Let us first observe that, according to Lemma 2.2, we only need to focus on the smallest size of elements in

$$(3.1) \quad \{\text{simple } m\text{-vector } Q \text{ at } 0 \text{ with } \phi(Q) = 1\}$$

under metric $g(s)$.

The second is that, for each $P = \text{span } Q$, there exists an orthonormal basis $\{e_1, \dots, e_m\}$ with respect to $g_1|_P$ such that $g_2|_P$ is diagonalized as $\text{diag}\{\lambda_1, \dots, \lambda_m\}$ under the basis and

$$Q = \mathfrak{t} e_1 \wedge \dots \wedge e_m$$

where \mathfrak{t} is a nonzero real number ($|\mathfrak{t}| \geq 1$ since $\|\phi\|_{g_E}^* \leq 1$). Hence,

$$(3.2) \quad T(Q; s) \triangleq \left(\|Q\|_{g(s)}^*\right)^2 = \mathfrak{t}^2(1-s+s\lambda_1)\cdots(1-s+s\lambda_m).$$

Note that by the assumptions $\|\phi\|_{g_1}^* \leq 1$ and $\|\phi\|_{g_2}^* \leq 1$ it implies that both $T(Q; 0)$ and $T(Q; 1)$ are no less than one.

Lastly, let us consider the function $F(X_1, \dots, X_m) = (\Pi_{i=1}^m X_i)^{-1}$. A pleasant property of F is that it is strictly convex over the interior of the first quadrant of \mathbb{R}^m . This can be checked very easily as

$$\begin{aligned} \text{Hess } F &= F(X_1, \dots, X_m) \cdot \begin{bmatrix} \frac{2}{X_1^2} & \frac{1}{X_1 X_2} & \cdots & \frac{1}{X_1 X_m} \\ \frac{1}{X_1 X_2} & \frac{2}{X_2^2} & \cdots & \frac{1}{X_2 X_m} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{X_1 X_m} & \frac{1}{X_2 X_m} & \cdots & \frac{2}{X_m^2} \end{bmatrix} \\ &= F(X_1, \dots, X_m) \cdot \left\{ \text{diag}\left(\frac{1}{X_1^2}, \dots, \frac{1}{X_m^2}\right) + \left(\frac{1}{X_1^2}, \dots, \frac{1}{X_m^2}\right)^T \left(\frac{1}{X_1^2}, \dots, \frac{1}{X_m^2}\right) \right\} \end{aligned}$$

is strictly positive.

By choosing $X_i(s) = (1-s)1 + s\lambda_i$, note that $F(X_1(s), \dots, X_m(s)) = \frac{\mathfrak{t}^2}{T(Q; s)}$ and

$$\begin{aligned} F(X_1(s), \dots, X_m(s)) &\leq (1-s)F(1, \dots, 1) + sF(\lambda_1, \dots, \lambda_m) \\ &= (1-s)\frac{\mathfrak{t}^2}{T(Q; 0)} + s\frac{\mathfrak{t}^2}{T(Q; 1)} \\ &\leq (1-s)\mathfrak{t}^2 + s\mathfrak{t}^2 = \mathfrak{t}^2. \end{aligned}$$


Therefore, we get $T(Q; s) \geq 1$ for all $s \in [0, 1]$ and thus $\|\phi\|_{g(s)}^* \leq 1$. \square

Remark 3.3. *If for some Q we have $\|Q\|_{g_1}^* = \|Q\|_{g_2}^* = 1$, then the strict convexity leads to strict inequality $\|Q\|_{g(s)}^* < 1$ for $s \in (0, 1)$ unless $\lambda_i = 1$ for all $1 \leq i \leq m$ which implies $\|Q\|_{g(s)}^* \equiv 1$ for $s \in [0, 1]$. In particular, this tells that $P = \text{span } Q$ is calibrated by ϕ with respect to $g(s)$ for some $s \in (0, 1)$ if and only if P is calibrated by ϕ with respect to both g_1 and g_2 . The latter in fact forces P to be calibrated by ϕ with respect to $g(s)$ for all $s \in (0, 1)$.*

3.2. Simplified proof of Theorem 3.1. Suppose that $\phi(\neq 0)$ is a calibration m -form with constant coefficients in (\mathbb{R}^n, g_E) . Namely, $\|\phi\|_{g_E}^* \leq 1$.

Let $\xi \in \Lambda^m \mathbb{R}^n$ be a unit simple m -vector with $V = \text{span } \xi$. Suppose that $1 \geq \phi(\xi) = \vartheta > 0$. Then there exists a unique subspace W in Lemma 2.8. By Lemma 2.10 and Remark 2.11 we have $\|\phi\|_{g'}^* = \phi(\xi) = \vartheta$ with respect to some g' . Hence, by Lemma 3.2 a nicely behaved gluing for g_E and g' guarantees that $\|\phi\|_{(1-s)g_E + sg'}^* \leq 1$ for $s \in [0, 1]$.

For an application to show how the result is efficient, we give a simplified proof of Theorem 3.1 according to the following figure. Let \mathbf{r} be the induced distance



from o in S by the a priori metric g . Then regions Γ_1 , Ω and Γ_2 with height h here can be defined by certain ranges of \mathbf{r} .

Note that the nearest projection π to $S \cap (\Gamma_1 \cup \Omega \cup \Gamma_2)$ can be well-defined on $\Gamma \triangleq \Gamma_1 \cup \Omega \cup \Gamma_2$ when h is small. Denote the local calibration form in assumption by φ . Since the region Ω can deform to a set of dimension strictly less than m , it follows that $\varphi = d\psi$ on Ω for some smooth form ψ of degree $m - 1$. The gluing of forms in [Zha17] $\Phi \triangleq d(\tau\psi + (1 - \tau)\pi^*(\psi|_S))$, where τ decreases from one to zero, provides a smooth form on $\Xi \triangleq B_{\frac{\epsilon}{2}}(o; g) \cup \Gamma$. A remark here is that when ϵ is small enough $B_{\frac{\epsilon}{2}}(o; g) \cap S$ is approximately the set of points on S which have $\mathbf{r} = \frac{\epsilon}{2}$, e.g. see [Sim83] about the uniqueness of tangent cone at o .

Denote the oriented unit horizontal m -vector by \vec{Z} in Γ with the property that $\vec{Z}|_S$ agrees with the orientation of S and $\text{span } \vec{Z}$ is perpendicular to the fiber directions of π with respect to g . Clearly, \vec{Z} is smooth and so is $\Phi(\vec{Z})$. Shrink h if necessary to make $\Phi(\vec{Z}) > 0$ in Γ . Let $Z = \left(\Phi(\vec{Z})\right)^{-1} \vec{Z}$. Then $\Phi(Z) \equiv 1$ in Γ . Set

$\tilde{g} = \left(\Phi(\vec{Z})\right)^{\frac{2}{m}} g$ in Γ and then $\|Z\|_{\tilde{g}} = 1$. By Lemmas 2.8 and 2.10 for $V = \text{span } Z$ and the compactness of $\bar{\Gamma}$, one can get a smooth metric $g' = \tilde{g}|_Z + C^2 \tilde{g}|_W$ with large C^2 such that $\|\Phi\|_{g'}^* \equiv 1$ in Γ . The smoothness of g' is due to the smoothness of Φ and Z which induces the smoothness of W and that of $\tilde{g}|_W$.

Now we do the metric gluing for g and g' in Γ_1 using function $\mathbf{s} = \mathbf{s}(\mathbf{r})$ which decreases from one to zero. Since $\Phi = \varphi$ in Γ_1 , we have $\|\Phi\|_g^* \leq 1$ in Γ_1 . By applying Theorem 3.2 pointwise, we obtain $\|\Phi\|_{\mathbf{s}g + (1-\mathbf{s})g'}^* \leq 1$ in Γ_1 . Note that $\mathbf{s}g + (1-\mathbf{s})g'$ actually forms a smooth metric \check{g} in Ξ .

As a result, we get a smooth calibration pair Φ and \check{g} in Ξ . Since $\Phi(\vec{Z}) = 1$ in $S \cap \Gamma$, it follows that g' and g induce the same metric of the submanifold $S \cap \Gamma$. Hence, by Remark (3.3) for $Q = Z = \vec{Z}$ at every point in $S \cap \Gamma_1$, we can conclude that S is calibrated by Φ with respect to \check{g} .

A further extension from the local calibration pair (Φ, \check{g}) on Ξ to a global calibration pair $(\hat{\Phi}, \hat{g})$ on X and meanwhile making S calibrated by $\hat{\Phi}$ can be made based on the non-vanishing homology assumption. One can follow the step in [Zha17] and we omit it here.

Remark 3.4. *In fact by Lemma 2.12 the distribution of W pointwise in $S \cap \Gamma_1$ and in Γ_2 is exactly fiber directions of π .*

Remark 3.5. *The method can also be applied to coflat calibration form with singular set being \emptyset .*

4. EVERY LAWLOR CONE SUPPORTS CALIBRATIONS SINGULAR ONLY AT THE ORIGIN

In this section, we show that every Lawlor cone can support coflat calibrations which are singular only at the origin of the ambient Euclidean space (\mathbb{R}^N, g_E) . Here by Lawlor cone we mean a minimal cone which can be shown area-minimizing according to Lawlor's curvature criterion (in practice by the c-control or F-control in below). Such a cone naturally supports non-continuous dual calibrations. Based on some analysis in [Zha18], we are able to construct some coflat calibrations resolving the discontinuity and making the singular set exactly be $\{o\}$. Besides, a shortcut for the realization result in [Zha18] will be explained.

First of all let us review some relevant materials from [Law91].

4.1. Introduction to Lawlor's curvature criterion. The field of area-minimizing cones has been successively studied for decades due to their special central role in several famous questions in differential geometry and geometric measure theory, e.g. see [Fle62, DG65, Alm66, Sim68, BDGG69, Law72, HL82a, HS85, FK85, Che88, Law89, Law91, Sma99].

The theory of calibrations mentioned in §2 is an effective way to study these objects. Sometimes certain objects naturally associate with calibrations and further relate to dual area-nonincreasing projections. As in [HS85], Hardt and Simons showed that for any given area-minimizing regular hypercone C in \mathbb{R}^N there exists

a dilation-invariant foliation of the entire space \mathbb{R}^N by minimal hypersurfaces with C being the only singular leave. Hence, the oriented unit volume form of the foliation induces a coflat calibration singular at the origin o and also possibly in $C \sim o$. Furthermore, the normal vector field integrates a dilation-invariant foliation by curves along which an area-nonincreasing projection to the cone C is then obtained.

In [Law91] Lawlor established a similar structure in some suitable angular neighborhood of a minimal cone $C(L^k)$ instead of in the whole space \mathbb{R}^N . If the boundary of the neighborhood can be mapped to the origin by the projection, then one can map everything outside the neighborhood to the origin. Thus an area-nonincreasing projection is gained and the area-minimality of the cone is proved.

To be more precise, [Law91] introduced a foliation by dilation-invariant curves and for further simplicity use a uniform symmetric curve γ through $x \in L$, namely, for every $x \in L$ and any unit normal vector v_x of $L \subset \mathbb{S}^N$ at x , the curve $\gamma(x, v_x)$ that Lawlor considered has uniform expression in the polar coordinate $r = r(\theta)$. The sufficient condition for the projection along foliation by $\{\mathbb{R}_+ \cdot \gamma\}$ to be area-nonincreasing is the inequality

$$(4.1) \quad \frac{dr}{d\theta} \leq r \sqrt{r^{2k+2} \cos^{2k} \theta \inf_{x \in L, v_x \in N_x L} \left(\det \left(\mathbf{I} - (\tan \theta) \mathbf{h}_{ij}^{v_x} \right) \right)^2 - 1}$$

with $r(0) = \|x\| = 1$. Here \mathbf{h}_{ij}^v is the second fundamental form at x for v . To map the boundary of the region where the projection is defined, one wishes to have $\lim_{\theta \uparrow \theta_0} r(\theta) = \infty$ for some vanishing angle θ_0 . To get the narrowest, (4.1) with equality was studied. Set $t = \tan \theta$, $h(t) = (r(\theta) \cos \theta)^{-k-1}$ and

$$p(t) = \inf_{x \in L, v_x \in N_x L} \det \left(\mathbf{I} - t \mathbf{h}_{ij}^{v_x} \right) = 1 + p_2 t^2 + \dots \text{ with negative } p_2.$$

Then the equivalent version of (4.1) in creating calibration form was obtained in [Law91]

$$(4.2) \quad \left(h(t) - \frac{t}{k+1} h'(t) \right)^2 + \left(\frac{h'(t)}{k+1} \right)^2 \leq (p(t))^2, \quad h(0) = 1.$$

Note that the candidate calibration has the nice form $d(g\psi)$ for some peculiar smooth form ψ (see [Law91] for more details) and the (4.2) is just the requirement for comass no larger than one. As for the boundary condition, $\lim_{\theta \uparrow \theta_0} r(\theta) = \infty$ exactly corresponds to the situation that h hits the t -axis at $t_0 = \tan \theta_0$ for the first time. Moreover, once the normal radius $R(L)$ of L in \mathbb{S}^N is larger than twice the vanishing angle, one can get a well-defined area-nonincreasing projection to $C(L)$.

Lawlor's curvature criterion is the following.

Theorem 4.1 ([Law91]). *Given a regular minimal cone $C(L) \subset \mathbb{R}^{N+1}$, if the vanishing angle θ_0 exists and $\theta_0 \leq \frac{R(L)}{2}$, then $C(L)$ is area-minimizing.*

In practice, Lawlor used the control

$$(4.3) \quad (1 - \alpha t) e^{\alpha t} < F(\alpha, t, k+1) \leq p(t)$$


where $\alpha = \sup_{x \in L, v_x \in N_x L} \|\mathbf{h}_{ij}^{v_x}\|$ and $F(\alpha, t, k+1) = \left(1 - \alpha t \sqrt{\frac{k}{k+1}}\right) \left(1 + \frac{\alpha t}{\sqrt{k(k+1)}}\right)^k$ to derive two kinds of vanishing angles by employing (4.3) to replace $p(t)$

$$\theta_c(k+1, \alpha) > \theta_F(k+1, \alpha) \geq \theta_0(C(L)).$$

So, if either θ_F or θ_c is less than $\frac{R(L)}{2}$, then $C(L)$ is area-minimizing.

4.2. Proof of Theorem 1.1. By a Lawlor cone we mean the smallest vanishing angle $\theta_0(C(L))$ is strictly less than $\frac{R(L)}{2}$ in the practical sense. All area-minimizing cones gained by applying Lawlor's curvature criterion with F-control or c-control satisfy $\theta_0(C(L)) < \frac{R(L)}{2}$. Even if there existed some cone $C(L)$ with $\theta_0(C(L)) = \frac{R(L)}{2}$ in the theoretical sense, one cannot detect it through Lawlor's curvature criterion as the solution $h_0(t)$ which touches the t -axis at $t = \tan \theta_0$ has an unstable behavior around $t = 0$ (in the stage of second order) and simply becomes unobservable using Lawlor's vanishing angle table or any other numerical methods.

As a result, each Lawlor cone must have some solution h to (4.2) with equality everywhere which touches the t -axis at some $t = \tan \theta_3 > \tan \theta_0$ as in the following graph.



Proof of Theorem 1.1. For point (t, y) for $0 < y < \sqrt{t^2 + 1}p(t)$, the inequality (4.2) leads to

$$(4.4) \quad \frac{k}{t^2 + 1} \left(ty - \sqrt{(t^2 + 1)p^2(t) - y^2} \right) \leq h'(t) \leq \frac{k}{t^2 + 1} \left(ty + \sqrt{(t^2 + 1)p^2(t) - y^2} \right).$$

Since $\frac{\partial}{\partial y} \left(ty - \sqrt{(t^2 + 1)p^2(t) - y^2} \right) = t + \frac{y}{\sqrt{(t^2 + 1)p^2(t) - y^2}} > 0$, in order to reach zero fastest, the solution h_0 must satisfy

$$(4.5) \quad h'(t) = \frac{k}{t^2 + 1} \left(th - \sqrt{(t^2 + 1)p^2(t) - h^2} \right),$$


with $h(0) = 1$. At the stage of second order, $h_0(t) = 1 - a_{\max}t^2 + \dots$ and $h_3(t) = 1 - a_{\min}t^2 + \dots$ where $a_{\max, \min} = \frac{k}{4} \left(k - 2 \pm \sqrt{(k - 2)^2 + 8p_2} \right)$. Hence, as in [Zha18] we can choose $a_{\min} < a < a_{\max}$ and sufficiently small δ such that for, all $t \in (0, \tan \delta]$,

$1 - at^2$ satisfies (4.2) which lies between h_0 and h_3 . Moreover it can further extend to h_1 by (4.5).

Note that h_0 is strictly decreasing, that is to say $h(t) < p(t)$ in the interval $(0, \tan \theta_0)$. So, one can arrange a and δ to make h_1 strictly decreasing as well. Now one gets $h_1(t) < p(t)$ and around the graph of h_1 in the figure the inequality (4.4) tells that the slope $h'(t)$ of a solution to the inequality of (4.2) is allowed to vary from a strictly negative number to a strictly positive number as illustrated in the figure. Then one can deform h_1 to h_2 which smoothly joins $1 - at^2$ and the t -axis with its graph touching the axis at $\tan \theta_2$ as close as one wishes to the point $\tan \theta_1$, and moreover satisfies the calibration inequality (4.2). The reason is that in any small interval $[\delta, \epsilon + \delta)$ one can extend $1 - at^2$ smoothly and make the extension fulfill (4.5) near $t = \epsilon + \delta$. This allows us to further extend it by (4.5) and it will hit the t -axis at a point $\tan \hat{\theta}$ close to $\tan \theta_1$. Similarly as what has been done around $t = \delta$, one can deform it around $\tan \hat{\theta}$ to make it smoothly tangent to the t -axis at $\tan \theta_2$ such that the resulting h_2 satisfies (4.2) and $\tan \theta_2$ can be as close to $\tan \hat{\theta}$ as one wants.

As a result, the corresponding calibration $\phi = d(h\psi)$ is smooth in $C(L) \sim o$ and also in the boundary of its θ_2 -angular neighborhood $U(\theta_2)$ with vanishing $d(h\psi)$ outside $U(\theta_2)$. Thus, we finish the proof. \square

Remark 4.2. *We would like to point out that one can also consider smooth extensions as follows.*



They cause no problem for gaining smooth calibrations away from o for the conclusion in Theorem 1.1 but the latter cannot induce a global area-nonincreasing projection to $C(L)$ faithfully corresponding to the figure while the former can.

4.3. A shortcut for the realization result in [Zha18]. How to get some closed homologically area-minimizing singular hypersurface in some Riemannian manifold remained a long-term standing question until [Sma99] where N. Smale found first family of such examples utilizing many tools from geometric measure theory and geometric analysis.

Based on [Zha16] where the author showed every homogeneous area-minimizing hypercone supports coflat calibrations singular only at the origin, Theorem 3.1 ([Zha17]) can create examples of closed homologically area-minimizing singular hypersurfaces in some Riemannian manifolds.

In [Zha18] we study more general realization question:

Can any area-minimizing cone be realized as a tangent cone at a point of some homologically area-minimizing compact singular submanifold?

And we answered affirmatively for every oriented Lawlor cone $C(L)$.

The ideas of [Zha18] include a reduction argument of the question to local by Allard's monotonicity, some modification on Lawlor's dual non-continuous calibration for a calibration which is smooth in $L(C) \sim 0$, and a series of mollifications on Lipschitz potential of the non-continuous calibration by virtue of Nash's embedding theorem. Suitable combinations among them also play a role for other related questions.

Let us recall that, as in [Sma99], we consider in [Zha18]

$$\Sigma_C \triangleq (C \times \mathbb{R}) \cap \mathbb{S}^N$$

in $\mathbb{R}^N \times \mathbb{R}$. Let M be an embedded oriented connected compact m -dimensional submanifold in some N -dimensional oriented compact manifold T with $[M] \neq [0] \in H_m(T; \mathbb{Z})$. Within smooth balls around a point of M and a regular point of Σ_C respectively one can connect T and \mathbb{S}^N , M and Σ_C simultaneously through one connected sum. Denote by X and S the resulting manifold and submanifold (singular at two points p_1 and p_2). Then $[S] \neq [0] \in H_m(X; \mathbb{Z})$.

Now we would like to point out here that Theorem 1.1 and Theorem 3.1 (for C being a Lawlor cone and S having exactly two singular points p_1, p_2 , the argument in §3.2 is still valid since Φ and \check{g} in Γ_2 are uniquely determined by the initial g cf. Remark 3.4 so things match together well) immediately open a shortcut to conclude that every oriented Lawlor cone $C(L)$ can be realized to the realization question.

5. MINIMAL PRODUCT AND A PROOF OF THEOREM 1.4

We will give a proof of Theorem 1.4 in this section. Actually, the result holds for continuous calibration as well.

To begin let us briefly recall the minimal product construction.

5.1. Minimal product. Given minimal submanifolds $\{L_i\}$ in \mathbb{S}^{N_i} respectively for $i = 1, \dots, n$, a nice algorithm to generate new minimal submanifold in sphere is the following

$$(5.1) \quad L_1 \times \cdots \times L_n \triangleq (\lambda_1 L_1, \dots, \lambda_n L_n) \subset \mathbb{S}^{N_1 + \cdots + N_n + n - 1}.$$

where $\lambda_i = \sqrt{\frac{k_i}{k}}$ with k_i the dimension of L_i and $k = \sum_{i=1}^n k_i > 0$.

This structure may be already known by some experts for quite long period. Special situations with all inputs being full spheres or all being \mathbb{R} -spaces have been studied, e.g. see [Law91, OS21] and references therein. A simple proof of the structure for general minimal submanifolds through Takahashi Theorem can be

found in the book [Xin03]¹. In 2016 (arXiv chronology), the structure was rediscovered independently by [TZ20] and [CH18] using different methods. Readers may choose either familiar way to understand it and the complete information of second fundamental form can be found in [TZ20].

5.2. Proof of Theorem 1.4. A preparation for the proof is the following wonderful result by Jim Simons (also see an equivalent version in [DG65]).

Let M be a minimal hypersurface of \mathbb{S}^n . Choose an orientation and denote the unit normal vector at $x \in M$ by $N(x)$. The parallel replacement of $N(x)$ at the origin is called x^* . Use M^* to stand for the image of M under the mapping $x \mapsto x^*$.

Theorem 5.1 (Version in [Sim68]). *Suppose that M is a closed minimal hypersurface in \mathbb{S}^n . Then either M^* is a single point, in which case $M = \mathbb{S}^{n-1}$, or M^* lies in no open hemisphere of \mathbb{S}^n .*

Now we begin our proof of Theorem 1.4.

Proof of Theorem 1.4. Assume that each $L_i \subset \mathbb{S}^{N_i}$ is a closed minimal submanifold and L_1 of codimension one in \mathbb{S}^{N_1} . Suppose that ϕ is a continuous calibration defined in the entire space (\mathbb{R}^N, g_E) which calibrates the cone $C(L_1 \times \cdots \times L_n) \subset \mathbb{R}^N$ over minimal product $L_1 \times \cdots \times L_n$.

The first observation is that the restriction ϕ_o of ϕ at the origin calibrates $C(L_1 \times \cdots \times L_n)$. This is due to the continuity of calibration ϕ and that the tangent cone of a cone is itself. And one can apply argument by contradiction to get the observation rigorously.

Based on the observation, we know that $C(L_1 \times \cdots \times L_n)$ must be calibrated by ϕ_o which is a calibration with constant coefficients. Fix $x_i \in L_i$ and let ξ_i be the oriented unit tangent k_i -vector of $T_{x_i}L_i$ for $i = 2, \dots, n$. Then we shall focus on the calibration (k_1+1) -form $\Phi_o \triangleq (\xi_2 \wedge \cdots \wedge \xi_n) \lrcorner \phi_o$ with constant coefficients. Denote $\mathcal{P}^*(\Phi_0)$ to be the pull-back of Φ_0 to the linear subspace \mathbb{R}^{N_1+1} via the inclusion $\mathcal{P} : \mathbb{R}^{N_1+1} \hookrightarrow \mathbb{R}^N$ by $\mathcal{P}(x) = (x, 0, \dots, 0)$. Note that for $x \in L_1$ we have the oriented unit tangent k_1 -vector $\xi = \xi(x)$ of $T_x L_1$ and $\xi \wedge x$ is the volume form of $C(L_1) \subset \mathbb{R}^{N_1+1}$ (reverse the orientation if need be, here and in the sequel).

Without loss of generality, assume that L_1 is connected (or restrict ourselves to one connected component of L_1). We aim to show that evaluation of $\mathcal{P}^*(\Phi_0)$ with $\xi \wedge x$ is a common nonzero constant for all $x \in L_1$. First note that evaluation of $\mathcal{P}^*(\Phi_0)$ with $\xi \wedge x$ has the same value of Φ_0 with $\xi \wedge x$. Moreover, Φ_o (or $-\Phi_o$) calibrates all the $(k_1 + 1)$ -vectors $\{\xi \wedge p\}$ where $p = p(x) = (\lambda_1 x, \lambda_2 x_2, \dots, \lambda_n x_n)$ with x running in L_1 . So, for each $x \in L_1$, we can extend an orthonormal basis $\{v_1, \dots, v_{k_1}\}$ of $\text{span } \xi$ to an orthonormal basis

$$(5.2) \quad \{v_1, \dots, v_{k_1}, v_{k_1+1}, v_{k_1+2}, \dots, v_N\}$$

¹This reference came into the author's notice in 2020, thanks to the presentation copy (new edition) from Professor Xin to the author in 2019.

of \mathbb{R}^N with $v_{k_1+1} = p$ and $v_{k_1+2} = p^\perp$ a unit orthogonal vector to p in the plane $\text{span}\{p, x\}$. Now by applying Lemma 2.12 we have the decomposition

$$(5.3) \quad \Phi_o = \pm v_1^* \wedge \cdots \wedge v_{k_1+1}^* + \sum a_I v_I^*,$$

where $a_I = 0$ whenever $i_{k_1} \leq k_1 + 1$. For nonzero a_I , the corresponding multi-index I must have at least two indices strictly larger than $k_1 + 1$. Note that among elements of (5.2) only v_{k_1+1} and v_{k_1+2} have nonzero evaluations with x but the coefficient a_I vanishes for $I = 1\ 2\ \cdots\ k_1\ (\widehat{k_1 + 1})\ (k_1 + 2)$. As a result, it follows by (5.3) that

$$(5.4) \quad \begin{aligned} (\mathcal{P}^*(\Phi_o))(\xi \wedge x) &= \Phi_o(\xi \wedge x) \\ &= (\pm v_1^* \wedge \cdots \wedge v_{k_1+1}^*)(\xi \wedge x) \\ &= \pm \lambda_1 \end{aligned}$$

for all $\xi(x) \wedge x$ where $x \in L_1$.

By the assumption, we know that $k_1 + 1 = N_1$. So $\mathcal{P}(\Phi_o)$ is a simple form in \mathbb{R}^{N_1+1} . In particular, by (5.4) and the connectedness of L_1 (or its connected component), the unit normal dual vector to $\mathcal{P}(\Phi_o)$ has constant angle $\arccos \lambda_1$ or $-\arccos \lambda_1$ with respect to the unit normal of $\xi \wedge x$ for all $x \in L_1$. Namely, L_1^* is contained in an open hemisphere of \mathbb{S}^{N_1} but this cannot happen according to Theorem 5.1.

Therefore, we finish the proof by contradiction argument, with the conclusion that whenever someone among $\{L_i\}$ is a minimal hypersurface in sphere, the cone $C(L_1 \times \cdots \times L_n)$ cannot be calibrated by any globally defined continuous calibration form. \square

Remark 5.2. *There are many examples of area-minimizing cones of high codimension which can support calibration with constant coefficients, e.g. see the coassociative cone and special Lagrangian cones in [HL82a]. The spiral minimal product algorithm in [LZ] can produce a plethora of new examples of special Lagrangian cones based on given ones. However, it is unclear to the author at this moment whether the cone over the minimal product (of inputs of positive dimensions) here could support some or not continuous calibration.*

6. DETECT DUALITY OBSTRUCTION IN HIGHER CODIMENSION

Before discussion on the situation of higher codimension, let us review main steps for creating Example 3 in [Zha17] which displays the duality obstruction of calibration in the smooth category for codimension one.

6.1. Review on the duality obstruction in codimension one. Roughly speaking, there are three steps. The first is to construct closed singular submanifold (S, o) of nonzero real homology class in some manifold X with tangent cone C at o being a homogeneous area-minimizing hypercone (say the Simons cone $C_{3,3}$). Then by [Zha16] and Theorem 3.1 we can get a coflat calibration $\hat{\Phi}$ singular only at o with respect to some smooth metric \hat{g} on X such that $S \sim o$ is calibrated by $\hat{\Phi}$.

The second step is to fix a closed smooth submanifold M in $[S]$ which has empty intersection with S and conformally change metric \hat{g} in sufficiently small neighborhood of M such that after the change M and S are the only two mass-minimizing in the class $[S]$. This can be made rigorously by calibration arguments, for details see [Zha17].

The last step is to observe that, under the new metric h , although M is homologically mass-minimizing, it cannot be calibrated by any smooth/continuous calibration form. Since otherwise the singular hypersurface S must be calibrated according to the fundamental theorem of calibrated geometry, Theorem 2.6. As a result, the non-flat tangent cone of S at o then must be calibrated by a calibration with constant coefficients in $(T_o X, h_o)$ and this is impossible.

From the steps in the above, the duality obstruction of calibration in the smooth category is finally due to codimension one.

6.2. Generalization to higher codimension. Let us mention that the first two steps in the above can be done without difficulty for higher codimension.

The reason is that one can apply the construction in §4.3 with $T^N = S_1^1 \times \cdots \times S_N^1$ and the submanifold being the torus $T^m = S_1^1 \times \cdots \times S_m^1 \times \{q_{m+1}\} \times \cdots \times \{q_N\}$ where $q_j \in S_j^1$ for $m < j \leq N$. Hence, for any m -dimensional cone $C \subset \mathbb{R}^N$ which can be calibrated by some coflat calibration singular only at the origin, the connect sum of T^N and \mathbb{S}^N around regular points of T^m and Σ_C gives us X and S in the first step.

As already explained in §4.3, Theorem 3.1 can be applied to the resulting S with two singular points. Let $(\hat{\Phi}, \hat{g})$ be the calibration pair on X and $\hat{\Phi}$ calibrates S . Further, one can choose M to be $S_1^1 \times \cdots \times S_m^1 \times \{q'_{m+1}\} \times \cdots \times \{q'_N\}$ where some $q'_j \neq q_j$. Then by exactly the same reason one can conformally change \hat{g} such that M and S are the only two mass-minimizing in the class $[S]$.

Again, by the fundamental theorem of calibrated geometry, if M is calibrated by some smooth/continuous calibration form, then the tangent cone C of S at singular point p_1 (or p_2) has to be calibrated by a calibration with constant coefficients.

Now we are ready to state what kind of local model $C \subset \mathbb{R}^N$ can lead to duality obstruction in higher codimension in the last step. Actually there are quite many. The following is a recent configuration result for mass-minimizing cones in Euclidean spaces.

Theorem 6.1 ([Zha]). *Given embedded closed minimal submanifolds $\{L_i\}$ respectively in Euclidean spheres with $i = 1, 2, \dots, n$, every cone over the minimal product L of sufficiently many copies among these $\{L_i\}$ is area-minimizing.*

Another neat configuration result is Corollary 4.2 of [Zha].

Corollary 6.2. *Let L be a closed smooth minimal submanifold in some \mathbb{S}^n . Then the cone over minimal product $L \times \mathbb{S}^d$ is area-minimizing for sufficiently large d .*

Both results are proved by Lawlor's criterion. So our Theorem 1.1 here asserts that there exist coflat calibration forms singular only at the origin for each Lawlor cone in Theorem 6.1 and Corollary 6.2.

If a Lawlor cone $C(L_1^{\times s_1} \times \cdots \times L_n^{\times s_n})$ from Theorem 6.1 has some input, say L_1 , a minimal hypersurface in \mathbb{S}^{N_1} with $s_1 > 0$, then by our Theorem 1.4 we know that it cannot be calibrated by any smooth/continuous calibration form.

Let L be an arbitrary closed minimal submanifold in \mathbb{S}^N then $C(L \times \mathbb{S}^{2025} \times \mathbb{S}^d)$ is a Lawlor cone when d is sufficiently large. Note that the minimal product $\mathbb{S}^{2025} \times \mathbb{S}^d$ is a minimal hypersurface in \mathbb{S}^{2026+d} . So by our Theorem 1.4 viewing $L \times \mathbb{S}^{2025} \times \mathbb{S}^d = L \times (\mathbb{S}^{2025} \times \mathbb{S}^d)$ we know that $C(L \times \mathbb{S}^{2025} \times \mathbb{S}^d)$ cannot be calibrated by any smooth/continuous calibration form.

Therefore, based on the such Lawlor cones as local models we can follow the above three steps to detect the duality obstruction of calibrations in the smooth category for the situation of higher codimension. In summary, the key points are 1. existence of coflat calibration singular only at the origin in local model by Theorem 1.1 in the second step and 2. the contradiction by Theorem 1.4 in the last step.

REFERENCES

- [Alm66] Frederick J. Almgren, Jr., *Some interior regularity theorems for minimal surfaces and an extension of Bernstein's theorem*, Ann. Math. **84** (1966), 277–292.
- [BDGG69] Enrico Bombieri, Ennio De Giorgi, and E. Giusti, *Minimal cones and the Bernstein problem*, Invent. Math. **7** (1969), 243–268.
- [Che88] Benny N. Cheng, *Area-minimizing cone-type surfaces and coflat calibrations*, Indiana Univ. Math. J. **37** (1988), 505–535.
- [CH18] Jaigyoung Choe and Jens Hoppe, *Some minimal submanifolds generalizing the Clifford torus*, Math. Nach. **291** (2018), 2536–2542.
- [DG65] Ennio De Giorgi, *Una estensione del teorema di Bernstein*, Ann. Sc. Norm. Sup. Pisa **19** (1965), 79–85.
- [FF60] Herbert Federer and Wendell H. Fleming, *Normal and integral currents*, Ann. Math. **72** (1960), 458–520.
- [Fed69] Herbert Federer, *Geometric Measure Theory*, Springer-Verlag, New York, 1969.
- [FK85] Dirk Ferus and Hermann Karcher, *Non-rotational minimal spheres and minimizing cones*, Comment. Math. Helv. **60** (1985), 247–269.
- [Fle62] Wendell H. Fleming, *On the oriented Plateau problem*, Rend. Circolo Mat. Palermo **9** (1962), 69–89.
- [FH17] Michael Freedman and Matthew Headrick, *Bit threads and holographic entanglement*, Comm. Math. Phys. **352** (2017), 407–438.
- [HS85] Robert Hardt and Leon Simon, *Area minimizing hypersurfaces with isolated singularities*, J. Reine. Angew. Math. **362** (1985), 102–129.
- [HL82a] F. Reese Harvey and H. Blaine Lawson, Jr., *Calibrated geometries*, Acta Math. **148** (1982), 47–157.
- [HL82b] F. R. Harvey and H. B. Lawson, Jr., *Calibrated foliations*, Amer. J. Math. **104** (1982), 607–633.
- [Law89] Gary R. Lawlor, *The angle criterion*, Invent. Math. **95** (1989), 437–446.
- [Law91] ———, *A Sufficient Criterion for a Cone to be Area-Minimizing*, Vol. 91, Mem. of the Amer. Math. Soc., 1991.
- [Law72] H. Blaine Lawson, Jr., *The equivariant Plateau problem and interior regularity*, Trans. Amer. Math. Soc. **173** (1972), 231–249.
- [LZ] Haizhong Li and Yongsheng Zhang, *Spiral Minimal Products*, arXiv: 2306.03328.
- [Liu24] Zhenhua Liu, *Every finite graph arises as the singular set of a compact 3-D calibrated area minimizing surface*, Comm. Pure Appl. Math. **77** (2024), 3670–3707.

- [Liua] ———, *Area-minimizing submanifolds are not generically smooth*, arXiv:2206.08315.
- [Liub] ———, *On a conjecture of Almgren: area-minimizing submanifolds with fractal singular sets*, arXiv:2110.13137.
- [OS21] Shinji Ohno and Takashi Sakai, *Area-minimizing cones over minimal embeddings of R -spaces*, Josai Math. Monogr. **13** (2021), 69–91.
- [Sim83] Leon Simon, *Asymptotics for a Class of Non-Linear Evolution Equations, with Applications to Geometric Problems*, Ann. of Math. **118** (1983), 525–571.
- [Sim68] James Simons, *Minimal varieties in riemannian manifolds*, Ann. of Math. **88** (1968), 62–105.
- [Sma99] Nathan Smale, *Singular homologically area minimizing surfaces of codimension one in Riemannian manifolds*, Invent. Math. **135** (1999), 145–183.
- [TZ20] Zizhou Tang and Yongsheng Zhang, *Minimizing cones associated with isoparametric foliations*, J. Differential Geom. **115** (2020), 367–393.
- [Xin03] Yuanlong Xin, *Minimal submanifolds and related topics*, Nankai Tracts in Mathematics, World Scientific Publishing, 2003 (and Second Edition in 2018).
- [Zha16] Yongsheng Zhang, *On Lawson’s area-minimizing hypercones*, Acta Math. Sin. (Engl. Ser.) **32** (2016), 1465–1476.
- [Zha17] ———, *On extending calibration pairs*, Adv. Math. **308** (2017), 645–670.
- [Zha18] ———, *On realization of tangent cones of homologically area-minimizing compact singular submanifolds*, J. Diff. Geom. **109** (2018), 177–188.
- [Zha] ———, *Some configuration results for area-minimizing cones*, arXiv:2510.17240.

Email address: yongsheng.chang@gmail.com