

STOCHASTIC DENSITY FUNCTIONAL THEORY THROUGH THE LENS OF MULTILEVEL MONTE CARLO METHOD*

XUE QUAN[†] AND HUAJIE CHEN[‡]

Abstract. The stochastic density functional theory (sDFT) has exhibited advantages over the standard Kohn-Sham DFT method and has become an attractive approach for large-scale electronic structure calculations. The sDFT method avoids the expensive matrix diagonalization by introducing a set of random orbitals and approximating the density matrix via Chebyshev expansion of a matrix-valued function. In this work, we study the sDFT with a plane-wave discretization, and discuss variance reduction algorithms in the framework of multilevel Monte Carlo (MLMC) methods. In particular, we show that the density matrix evaluation in sDFT can be decomposed into many levels by increasing the plane-wave cutoffs or the Chebyshev polynomial orders. This decomposition renders the computational cost independent of the discretization size or temperature. To demonstrate the efficiency of the algorithm, we provide rigorous analysis of the statistical errors and present numerical experiments on some material systems.

Key words. stochastic density functional theory; plane-wave discretization; Chebyshev polynomial expansion; multilevel Monte Carlo method.

1. Introduction. Kohn-Sham density functional theory (DFT) [20, 23] is among the most widely used electronic structure models in molecular simulations and materials science, which achieves an excellent balance between accuracy and computational cost. The cost of standard Kohn-Sham DFT scales cubically with system size because of repeated diagonalization of the Kohn-Sham Hamiltonian. To overcome this computational bottleneck, much effort has been devoted to the development of so-called linear-scaling DFT methods [2, 22, 26, 34, 38]. However, most methods rely on Kohn’s “nearsightedness” principle [22] and are typically applicable to non-metallic systems.

An alternative approach termed stochastic DFT (sDFT) [3, 14] has been introduced to accelerate the DFT calculations. The key idea is to approximate the density matrix by applying the Chebyshev polynomial expansion of the projection operator to a set of stochastic orbitals. In particular, one approximates the density matrix by

$$f(H) \approx p_M^2(H) = \sum_{i=1}^n (p_M(H)\mathbf{e}_i)(p_M(H)\mathbf{e}_i)^{\mathbf{h}} \approx \sum_{\chi \in \mathbb{S}} (p_M(H)\chi)(p_M(H)\chi)^{\mathbf{h}},$$

where $H \in \mathbb{C}^{n \times n}$ denotes the discrete Hamiltonian matrix within certain discretization, p_M is an M -th order Chebyshev approximation of $f^{\frac{1}{2}}$, $\{\mathbf{e}_i\}$ is a complete basis set, and \mathbb{S} is a set of stochastic orbitals, where each $\chi \in \mathbb{S}$ satisfies $\chi \in \mathbb{C}^n$ and $\mathbb{E}[\chi\chi^{\mathbf{h}}] = I$. The sDFT method circumvents the expensive orthogonalization procedure in standard Kohn-Sham DFT, reducing the computation to merely matrix-vector multiplications $p_M(H)\chi$.

Since the stochastic sampling of the random orbitals introduces significant statistical noise, the efficiency of sDFT critically depends on effective variance reduction techniques. Many strategies have been developed in recent years, including the embedded-fragmentation scheme [12, 28], the energy-window sDFT [11], the tempering sDFT [29], and hybrid methods [24, 37]. However, to our knowledge, there is no

* This work was supported by the National Natural Science Foundation of China under grant No. 12371431.

[†]Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China (xquan@mail.bnu.edu.cn).

[‡]School of Mathematical Sciences, Beijing Normal University, Beijing 100875, China (chen.huajie@bnu.edu.cn).

mathematical theory for variance reduction techniques in sDFT. Recently we see a first mathematical analysis for sDFT [6], which establishes nearly-optimal scaling in both the thermodynamic and complete basis set limits, by exploiting a discretization-independent stochastic estimator, a stochastic mirror descent reformulation of the self-consistent field iteration, and a pole expansion calculation.

The purpose of this work is to develop a variance reduction framework for sDFT and provide a rigorous analysis for the associated statistical errors. We employ the multilevel Monte Carlo (MLMC) method, a powerful noise reduction strategy that uses a hierarchical structure: most samples are taken on computationally inexpensive coarse levels, while only a small number of samples are taken on the more expensive fine levels. The MLMC method was first introduced in [16, 17], and has since found widespread applications in numerous fields [1, 4, 36], particularly in matrix trace estimation problems [15, 18] that are closely related to this work. We investigate the use of MLMC within sDFT and provide insights for existing variance reduction strategies. In particular, we study the sDFT with a plane-wave discretization and develop two distinct MLMC approaches for evaluating the Kohn–Sham map. These two approaches leverage hierarchical constructions in plane-wave cutoffs and Chebyshev expansion orders, making the cost independent of the discretization size and the temperature, respectively.

Outline. The rest of this work is organized as follows. In Section 2, we briefly review the Kohn–Sham DFT with a plane-wave discretization, and present a density matrix based formulation for the problem. In Section 3, we describe the sDFT approach and provide some basic variance analysis. In Section 4, we present a general MLMC framework to evaluate the Kohn–Sham map in sDFT, and investigate two multilevel decomposition strategies by exploiting the hierarchy of plane-wave discretization and polynomial expansion, respectively. In Section 5, we present numerical simulations on some material systems to demonstrate the efficiency of MLMC methods. In Section 6, we provide some concluding remarks.

Notations. For $a \in \mathbb{C}$, we will denote the complex conjugate by \bar{a} . For a matrix $A \in \mathbb{C}^{m \times n}$, we will denote its conjugate transpose by A^h . For a random variable X , we will use $\mathbb{E}[X]$ and $\mathbb{V}[X]$ to denote the expectation and variance, respectively. With slight abuse of notation, we will use $|\cdot|$ to denote the Euclidean norm of a vector, the volume of a bounded domain in \mathbb{R}^d , and the cardinality of a finite set. Throughout this paper, we will use the symbol C to denote a generic positive constant that may change from one line to the next. The dependence of C on model parameters will normally be clear from the context or stated explicitly.

2. Kohn–Sham density functional theory. In this section, we review the Kohn–Sham DFT with a plane-wave discretization. In particular, the discrete Kohn–Sham equation is formulated as a fixed-point problem with respect to the density matrix, which provides the foundation of the sDFT method.

2.1. Kohn–Sham equation. Consider a many-particle system with N electrons lying in a d -dimensional domain $\Omega \subset \mathbb{R}^d$. In this work, we shall focus on systems with periodic boundary condition, whose domain can be written as $\Omega = A[0, 1)^d$ with $A \in \mathbb{R}^{d \times d}$ a non-singular matrix. There is a corresponding Bravais lattice $\mathbb{L} := A\mathbb{Z}^d$ and a dual lattice $\mathbb{L}^* := 2\pi A^{-T}\mathbb{Z}^d$.

Let $v_{\text{ext}} : \Omega \rightarrow \mathbb{R}$ be an external potential originating from nuclear attractions. By neglecting the spin variables, the electron Hamiltonian in Kohn–Sham density

functional theory is given by

$$(2.1) \quad \mathcal{H}^{\text{KS}}[\rho] := -\frac{1}{2}\Delta_{\mathbf{r}} + v_{\text{ext}}(\mathbf{r}) + \int_{\Omega} \frac{\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}' + v_{\text{xc}}(\rho(\mathbf{r})),$$

where $\rho : \Omega \rightarrow \mathbb{R}$ is the electron density satisfying $\int_{\Omega} \rho(\mathbf{r}) d\mathbf{r} = N$, and $v_{\text{xc}} : \mathbb{R} \rightarrow \mathbb{R}$ represents the so-called exchange-correlation potential within the local density approximation (LDA) [25]. In (2.1), the first term represents the kinetic energy, the second term is the external potential, the third term gives the mean field electrostatic interaction between the electrons, and the last term describes the correlations between electrons. Throughout this paper, we restrict our analysis to the pseudopotential [25] framework, so that v_{ext} is assumed to be sufficiently smooth and it is natural to employ plane-wave discretizations.

The ground state of the system can be obtained by solving the following Kohn-Sham equation: Find $(\lambda_i, \psi_i) \in \mathbb{R} \times H_{\text{per}}^1(\Omega)$, such that $\int_{\Omega} \psi_i^*(\mathbf{r}) \psi_j(\mathbf{r}) d\mathbf{r} = \delta_{ij}$ and

$$(2.2) \quad \mathcal{H}^{\text{KS}}[\rho] \psi_i = \lambda_i \psi_i \quad i = 1, 2, \dots,$$

where the electron density is given by the eigenpairs $\rho(\mathbf{r}) = \sum_i f(\lambda_i) |\psi_i(\mathbf{r})|^2$ with some smearing function $f : \mathbb{R} \rightarrow (0, 1)$. In this work, we will mainly consider the most widely used smearing function, the Fermi-Dirac function

$$(2.3) \quad f(\lambda) = f_{\mu}(\lambda) := \frac{1}{1 + \exp(\beta(\lambda - \mu))},$$

where $\beta > 0$ determines the smearing width, and $\mu \in \mathbb{R}$ is the Fermi level such that the condition $\sum_i f_{\mu}(\lambda_i) = N$ is satisfied. Note that when $\beta \rightarrow \infty$, f_{μ} becomes a Heaviside function $1_{(-\infty, \mu)}$, in which case only the N lowest eigenvalues of $H^{\text{KS}}[\rho]$ need to be solved.

Note that the Kohn-Sham equation (2.2) is a nonlinear eigenvalue problem, as the operator $\mathcal{H}^{\text{KS}}[\rho]$ depends on the eigenpairs $\{(\lambda_i, \psi_i)\}_{i \in \mathbb{N}_+}$ to be solved. In practice, a self-consistent field (SCF) iteration is usually used to solve the nonlinear problem. At each iteration step, the Hamiltonian is constructed with an input electron density, then an output electron density is obtained by solving the linearized eigenvalue problem and some charge-mixing technique [25, 27, 33]. The iteration procedure is terminated when the difference between two consecutive densities is negligible.

2.2. Plane-wave discretizations. In this work, we consider periodic boundary conditions for the Kohn-Sham equation and employ a plane-wave method for numerical discretization. We denote by $e_{\mathbf{G}}(\mathbf{r}) = |\Omega|^{-1/2} e^{i\mathbf{G} \cdot \mathbf{r}}$ the plane-wave basis with wavevector $\mathbf{G} \in \mathbb{L}^*$. The family $\{e_{\mathbf{G}}\}_{\mathbf{G} \in \mathbb{L}^*}$ forms an orthonormal basis set of $L_{\text{per}}^2(\Omega)$. Given an energy cut-off $E_c > 0$, we define the following finite dimensional subspace

$$(2.4) \quad X_{E_c}(\Omega) := \left\{ u \in L_{\text{per}}^2(\Omega) : u(\mathbf{r}) = \sum_{\mathbf{G} \in \mathbb{L}^*, \frac{1}{2}|\mathbf{G}|^2 \leq E_c} c_{\mathbf{G}} e_{\mathbf{G}}(\mathbf{r}) \right\},$$

the dimension of which is denoted by $n := \dim(X_{E_c}(\Omega)) = |\{\mathbf{G} \in \mathbb{L}^* : |\mathbf{G}|^2 \leq 2E_c\}|$.

With the plane-wave discretization, the eigenfunctions $\psi_i(\mathbf{r})$ are approximated by linear combinations of the basis functions $e_{\mathbf{G}}(\mathbf{r})$ in $X_{E_c}(\Omega)$, whose coefficients are denoted by $\mathbf{u}_i := \{u_{i,\mathbf{G}}\}_{|\mathbf{G}|^2 \leq 2E_c} \in \mathbb{C}^n$. Then (2.2) is approximated by the following nonlinear algebraic eigenvalue problem

$$(2.5) \quad H[\rho_{\mathbf{u}}] \mathbf{u}_i = \varepsilon_i \mathbf{u}_i \quad i = 1, \dots, n,$$

where the electron density $\rho_{\mathbf{u}}$ is associated to the approximate eigenfunction as

$$(2.6) \quad \begin{aligned} \rho_{\mathbf{u}}(\mathbf{r}) &= \sum_{i=1}^n f(\varepsilon_i) \left| \sum_{|\mathbf{G}|^2 \leq 2E_c} \mathbf{u}_{i,\mathbf{G}} e_{\mathbf{G}}(\mathbf{r}) \right|^2 \\ &= \sum_{|\mathbf{G}|^2 \leq 2E_c} \sum_{|\mathbf{G}'|^2 \leq 2E_c} \left(\sum_{i=1}^n f(\varepsilon_i) \mathbf{u}_{i,\mathbf{G}} \mathbf{u}_{i,\mathbf{G}'} \right) \overline{e_{\mathbf{G}}(\mathbf{r})} e_{\mathbf{G}'}(\mathbf{r}), \end{aligned}$$

and the matrix elements of the discrete Hamiltonian $H[\rho_{\mathbf{u}}] \in \mathbb{C}^{n \times n}$ are given by

$$(2.7) \quad H[\rho_{\mathbf{u}}]_{\mathbf{G}\mathbf{G}'} = \int_{\Omega} \overline{e_{\mathbf{G}}(\mathbf{r})} \mathcal{H}^{\text{KS}}[\rho_{\mathbf{u}}] e_{\mathbf{G}'}(\mathbf{r}) \, d\mathbf{r}.$$

The *a priori* error estimates for the approximations of eigenvalue problems can be found in [30] for linear problems and in [7, 8, 9] for nonlinear Kohn-Sham equations.

2.3. Kohn-Sham map of density matrix. In the rest of this section, we shall rewrite the discrete Kohn-Sham equation (2.5) as a fixed-point problem with respect to the density matrix. The density matrix formalism is fundamental for introducing the stochastic DFT methods.

Let $\mathcal{D} := \{P \in \mathbb{C}^{n \times n} : P = P^h\}$ be the space for Hermite matrices, endowed with the Frobenius inner product $(A, B)_{\mathbb{F}} := \text{Tr}(A^h B)$. The admissible set of discrete density matrices for an N -electron system is given by

$$(2.8) \quad \mathcal{D}_N := \{P \in \mathcal{D} : 0 \preceq P \preceq 1, \text{Tr}(P) = N\}.$$

For any $P \in \mathcal{D}_N$, it can be written in the form of

$$(2.9) \quad P = \sum_{i=1}^n f_i \mathbf{v}_i \mathbf{v}_i^h \quad \text{with } 0 \leq f_i \leq 1, \sum_{i=1}^n f_i = N \text{ and } \mathbf{v}_i^h \mathbf{v}_j = \delta_{ij}.$$

Note that the density matrix P contains all information of the orbitals \mathbf{v}_i and occupation numbers f_i . Given a density matrix $P \in \mathcal{D}_N$, we observe from (2.6) and (2.9) that the associated electron density can be expressed by

$$(2.10) \quad \rho_P(\mathbf{r}) := \sum_{|\mathbf{G}|^2 \leq 2E_c} \sum_{|\mathbf{G}'|^2 \leq 2E_c} P_{\mathbf{G}\mathbf{G}'} \overline{e_{\mathbf{G}}(\mathbf{r})} e_{\mathbf{G}'}(\mathbf{r}).$$

With an “input” density matrix $P \in \mathcal{D}_N$, we write the discrete Kohn-Sham Hamiltonian with a little abuse of notation

$$(2.11) \quad H[P] := H[\rho_P].$$

Denoting the eigenpairs of $H[P]$ by $\{(\varepsilon_i, \mathbf{u}_i)\}_{1 \leq i \leq n}$, we have an “output” density matrix $\sum_{i=1}^n f(\varepsilon_i) \mathbf{u}_i \mathbf{u}_i^h$ by using (2.9). We describe the map from an input density matrix to an output density matrix by the Kohn-Sham map $\Phi : \mathcal{D}_N \rightarrow \mathcal{D}_N$, with

$$(2.12) \quad \Phi(P) := f(H[P]).$$

Then the ground state density matrix P_* is a fixed-point of the map Φ , that is, it solves the the following equation

$$(2.13) \quad P_* = \Phi(P_*).$$

The simplest method for seeking the fixed point solution of (2.13) is to use the iteration $P_{k+1} = \Phi(P_k)$, until $\|P_{k+1} - P_k\|_F$ is sufficiently small. This iteration (together with some mixing scheme) is equivalent to performing SCF iterations to solve the Kohn-Sham equation.

The primary computational bottleneck in Kohn-Sham DFT is the evaluation of the Kohn-Sham map $\Phi(P)$. The direct evaluation requires a diagonalization of the Hamiltonian $H[P]$, whose cost scales cubically as $\mathcal{O}(n^3)$. In the zero temperature limit, i.e. when $\beta \rightarrow \infty$ in (2.3), only the lowest N eigenpairs are required and the cost also scales cubically as $\mathcal{O}(nN^2)$ due to the orthogonalization of N eigenvectors.

3. Stochastic DFT. The stochastic DFT method cures the cubic scaling by employing two key strategies to reduce the cost of evaluating (2.12): (i) random orbital approximation, and (ii) Chebyshev polynomial expansion. In this section, we provide details of these approximations and present a basic error analysis.

Without loss of generality, we assume in the following that the spectrum of $H[P]$ lies in $[-1, 1]$. This assumption is not essential as one can always apply a shift and rescaling to the discrete Hamiltonian by $a(H[P] - bI)$ (and also to the smearing function by $f(\frac{\cdot}{a} + b)$) with appropriate constants a and b .

3.1. Random orbital approximation. We first introduce the notion of random orbital. We call $\chi \in \mathbb{C}^n$ a random orbital if the entries of χ are independent random variables and satisfy the condition

$$(3.1) \quad \mathbb{E}[\chi\chi^h] = I$$

with $I \in \mathbb{R}^{n \times n}$ the identity matrix. Standard choices for χ satisfying (3.1) could be (a) $\chi_j \in \{-1, 1, -i, i\}$ with equal probability $\frac{1}{4}$; (b) $\chi_j = \exp(i\theta)$ with θ uniformly distributed in $[0, 2\pi]$; and (c) χ_j follows the normal distribution.

Let $P \in \mathcal{D}_N$ and $\{\mathbf{e}_i\}$ be the standard basis set of \mathbb{R}^n , with \mathbf{e}_i having 1 at the i -th entry and 0 otherwise. By using (2.12) and (3.1), we can write the Kohn-Sham map of the density matrix as an expectation in the following form

$$(3.2) \quad \begin{aligned} \Phi(P) &= f(H[P]) = \sum_{i=1}^n f(H[P])\mathbf{e}_i\mathbf{e}_i^h \\ &= \mathbb{E}\left(f(H[P])\chi\chi^h\right) \end{aligned}$$

$$(3.3) \quad = \mathbb{E}\left[\Phi(P, \chi, f^{\frac{1}{2}})\right] \quad \text{with} \quad \Phi(P, \chi, f^{\frac{1}{2}}) := \left(f^{\frac{1}{2}}(H[P])\chi\right)\left(f^{\frac{1}{2}}(H[P])\chi\right)^h.$$

Here the summation over a complete basis set $\{\mathbf{e}_i\}$ is replaced by the expectation with respect to random orbitals, enabling potential reduction of the computational cost through stochastic methods. Since the Fermi-Dirac distribution satisfies $f(x) \geq 0$, the matrix-valued function $f^{\frac{1}{2}}(H[P])$ is well-defined. Note that while equations (3.2) and (3.3) are mathematically equivalent, the form (3.3) is preferred in sDFT calculations because its sample complexity is less sensitive to the discretization size [6].

We can then approximate (3.3) by employing a stochastic sampling approach. Let \mathbb{S} be a set of random orbitals satisfying (3.1), obtained through some practical sampling techniques. Then the expectation in (3.3) can be approximated by

$$(3.4) \quad \Phi_{\mathbb{S}}(P) := \frac{1}{|\mathbb{S}|} \sum_{\chi \in \mathbb{S}} \Phi(P, \chi, f^{\frac{1}{2}}).$$

The accuracy of the stochastic approximation $\Phi_{\mathbb{S}}(P)$ is implied by the Chebyshev's inequality [5]

$$(3.5) \quad \mathbb{P} \left(\|\Phi_{\mathbb{S}}(P) - \Phi(P)\|_{\text{F}} \geq \epsilon \sqrt{\frac{\mathbb{V}[\Phi(P, \chi, f^{\frac{1}{2}})]}{|\mathbb{S}|}} \right) \leq \epsilon^{-2},$$

where $\mathbb{V}[\Phi(P, \chi, f^{\frac{1}{2}})]$ denotes the variance of the random matrix

$$(3.6) \quad \mathbb{V}[\Phi(P, \chi, f^{\frac{1}{2}})] := \sum_{i,j=1}^n \mathbb{V}[(\Phi(P, \chi, f^{\frac{1}{2}}))_{ij}] = \mathbb{E} \left[\|\Phi(P, \chi, f^{\frac{1}{2}}) - \mathbb{E}[\Phi(P, \chi, f^{\frac{1}{2}})]\|_{\text{F}}^2 \right].$$

We observe from (3.5) that the accuracy of the stochastic approximation depends on (i) the number of random orbitals in \mathbb{S} ; and (ii) the variance of the random matrix. Therefore, the variance (3.6) is the key to the efficiency of the sDFT calculations. We provide an estimate for this variance in the following lemma, whose proof is given in Appendix A.1.

LEMMA 3.1. *Let $\chi \in \mathbb{C}^n$ be the random orbital satisfying (3.1). Then*

$$(3.7) \quad \begin{aligned} \mathbb{V}[\Phi(P, \chi, f^{\frac{1}{2}})] &= \sum_{\substack{i,j=1 \\ i \neq j}}^n (\Phi(P))_{ii} (\Phi(P))_{jj} + \sum_{\substack{i,j=1 \\ i \neq j}}^n (\Phi(P))_{ij}^2 \mathbb{E}[\bar{\chi}_i^2] \mathbb{E}[\chi_j^2] \\ &\quad + \sum_{i=1}^n (\Phi(P))_{ii}^2 (\mathbb{E}[|\chi_i|^4] - 1). \end{aligned}$$

Moreover, if χ additionally satisfies¹

$$(3.8) \quad \mathbb{E}[\chi_i^2] = 0 \quad \text{and} \quad \mathbb{E}[|\chi_i|^4] = 1 \quad \text{for any } 1 \leq i \leq n,$$

then

$$(3.9) \quad \mathbb{V}[\Phi(P, \chi, f^{\frac{1}{2}})] = \sum_{\substack{i,j=1 \\ i \neq j}}^n (\Phi(P))_{ii} (\Phi(P))_{jj} = (\text{Tr}(\Phi(P)))^2 - \sum_{i=1}^n (\Phi(P))_{ii}^2.$$

Remark 3.2 (L^2 -error of the electron density). For $P \in \mathcal{D}_N$, let $\rho_{\mathbb{S}}$ and ρ denote the electron densities associated to $\Phi_{\mathbb{S}}(P)$ and $\Phi(P)$ respectively. We have from (2.10) that the stochastic error in L^2 -norm for the density can be estimated by

$$(3.10) \quad \begin{aligned} \|\rho_{\mathbb{S}} - \rho\|_{L^2(\Omega)} &= \left(\int_{\Omega} \left| \sum_{\mathbf{G}} \sum_{\mathbf{G}'} \left((\Phi_{\mathbb{S}}(P))_{\mathbf{G}\mathbf{G}'} - (\Phi(P))_{\mathbf{G}\mathbf{G}'} \right) \overline{e_{\mathbf{G}}(\mathbf{r})} e_{\mathbf{G}'}(\mathbf{r}) \right|^2 d\mathbf{r} \right)^{\frac{1}{2}} \\ &\leq \frac{1}{\sqrt{|\Omega|}} \|\Phi_{\mathbb{S}}(P) - \Phi(P)\|_{\text{F}}. \end{aligned}$$

¹Among the three random orbital examples provided below equation (3.1), both (a) and (b) satisfy this additional condition, while (c) does not – χ_i follows a normal distribution with the fourth moment $\mathbb{E}[|\chi_i|^4] = 3$.

Note that (3.9) and $\text{Tr}(\Phi(P)) = N$ imply that $\mathbb{V}[\phi(P, \chi, f^{\frac{1}{2}})] \sim \mathcal{O}(N^2)$. This together with (3.5), (3.10) and $|\Omega| \sim \mathcal{O}(N)$ can indicate that the L^2 -error of the electron density scales like

$$\|\rho_{\mathbb{S}} - \rho\|_{L^2(\Omega)} \sim \mathcal{O}\left(\sqrt{N/|\mathbb{S}|}\right).$$

Therefore, the number of random orbitals required in \mathbb{S} should scale as $\mathcal{O}(N)$ to achieve the same accuracy as the system size N increases. We mention that a different scaling for $|\mathbb{S}|$ can be obtained when a different accuracy criterion is considered. In many physics contexts [3, 11], the error averaged to per electron is used as the metric, under which the number of required random orbitals can be significantly smaller.

3.2. Chebyshev polynomial expansion. To evaluate the random matrix in (3.3), one needs to apply the matrix-valued function $f^{\frac{1}{2}}(H[P])$ on each random orbital $\chi \in \mathbb{S}$. To avoid direct diagonalization of the Hamiltonian $H[P]$, $f^{\frac{1}{2}}$ is approximated on $[-1, 1]$ by an M -th order polynomial expansion:

$$(3.11) \quad f^{\frac{1}{2}}(x) \approx p_M(x) := \sum_{m=0}^M c_m T_m(x) \quad x \in [-1, 1],$$

where $T_m(\cdot)$ is the m -th Chebyshev polynomial and c_m is the corresponding coefficient. Then (3.3) and (3.4) can be further approximated by

$$(3.12) \quad \Phi_M(P) := p_M^2(H[P]) = \mathbb{E}\left[\phi(P, \chi, p_M)\right]$$

$$(3.13) \quad \approx \frac{1}{|\mathbb{S}|} \sum_{\chi \in \mathbb{S}} \phi(P, \chi, p_M) =: \Phi_{\mathbb{S}, M}(P).$$

For each random orbital $\chi \in \mathbb{S}$, the product $p_M(H[P])\chi$ can be efficiently computed through recursive matrix-vector multiplications using Chebyshev polynomial recurrence relations. In practice, these matrix-vector multiplications are carried out using the Fast Fourier Transform (FFT), which allows the computational cost to scale linearly with the matrix size n .

Note that $f^{\frac{1}{2}}$ is analytic on $[-1, 1]$ but possesses singularities in the complex plane. The locations of these singularities depend on both the smearing parameter β and the Hamiltonian scaling constant a . Consequently, the Chebyshev polynomial approximation error decays exponentially with the polynomial order M [35]

$$(3.14) \quad \|p_M - f^{\frac{1}{2}}\|_{L^\infty([-1, 1])} \leq C \exp(-\alpha M),$$

where the decay rate α is determined by the constants β and a .

We now obtain an efficient approximation of the Kohn-Sham map $\Phi(P)$ for a given density matrix $P \in \mathcal{D}_N$, as $\Phi_{\mathbb{S}, M}(P)$ defined in (3.13). This is the essence of sDFT, that avoids the expensive eigensolver by replacing the complete basis set with stochastic orbitals and approximating the smearing function with Chebyshev polynomial expansion. The approximation error of $\Phi_{\mathbb{S}, M}$ is governed by the sampling size $|\mathbb{S}|$, the variance estimated in Lemma 3.1, and the polynomial order M .

Remark 3.3 (Cost scaling). The total computational cost for evaluating $\Phi_{\mathbb{S}, M}(P)$ scales as $\mathcal{O}(nM|\mathbb{S}|)$. Since $|\mathbb{S}| = \mathcal{O}(N)$ (see Remark 3.2), the complexity becomes $\mathcal{O}(nNM)$. This cost depends on the system size N , the degrees of freedom n (which is proportional to N in most of the DFT calculations), and the smearing parameter

β . In the following section, we shall introduce two variance reduction strategies: one that makes the computational cost independent of n , and the other that makes it independent of β .

4. Multilevel Monte Carlo methods for sDFT. In this section, we shall first present a general multilevel Monte Carlo (MLMC) framework to modify the standard sDFT scheme for evaluating the Kohn-Sham map in (3.13),

$$\Phi_{\mathbb{S},M}(P) = \frac{1}{|\mathbb{S}|} \sum_{\chi \in \mathbb{S}} \Phi(P, \chi, p_M) = \frac{1}{|\mathbb{S}|} \sum_{\chi \in \mathbb{S}} \left(p_M(H[P])\chi \right) \left(p_M(H[P])\chi \right)^h,$$

and then investigate two strategies of the multilevel decomposition, by exploiting the hierarchy of plane-wave discretization and polynomial expansion respectively.

4.1. General framework. To evaluate $\Phi_{\mathbb{S},M}$ within the MLMC framework, we shall first construct a sequence of approximations of $p_M(H[P])$, denoted by

$$\widehat{\mathcal{Q}}^{(\ell)} \in \mathbb{C}^{n \times n}, \quad \ell = 0, 1, \dots, L.$$

Specifically, we set $\widehat{\mathcal{Q}}^{(L)} := p_M(H[P])$ for the highest level and define $\widehat{\mathcal{Q}}^{(-1)} := \mathbf{0}$ for simplicity of subsequent notations. As ℓ increases, the approximations $\widehat{\mathcal{Q}}^{(\ell)}$ exhibit a hierarchical structure, with increasing accuracy but also increasing computational cost. This yields a sequence of increasingly precise approximations for the random matrix $\Phi(P, \chi, p_M)$ in (3.13) as

$$(4.1) \quad \widehat{\Phi}_{\chi}^{(\ell)} := (\widehat{\mathcal{Q}}^{(\ell)} \chi) (\widehat{\mathcal{Q}}^{(\ell)} \chi)^h, \quad \ell = 0, 1, \dots, L.$$

We then obtain a multilevel decomposition of $\Phi_M(P)$ as

$$(4.2) \quad \Phi_M(P) = \mathbb{E}_{\chi^{(0)}} [\widehat{\Phi}_{\chi^{(0)}}^{(0)}] + \sum_{\ell=1}^L \mathbb{E}_{\chi^{(\ell)}} [\widehat{\Phi}_{\chi^{(\ell)}}^{(\ell)} - \widehat{\Phi}_{\chi^{(\ell)}}^{(\ell-1)}],$$

which gives the fundamental form for the MLMC method. At each level ℓ ($\ell = 0, 1, \dots, L$), we construct a distinct set of random orbitals $\mathbb{S}^{(\ell)}$ to approximate the expectation value. The size of $\mathbb{S}^{(\ell)}$ varies across levels to account for the different variances of the random matrices. After sampling the sets of random orbitals for all levels, we can approximate $\Phi_M(P)$ in (4.2) with

$$(4.3) \quad \Phi_{\{\mathbb{S}^{(\ell)}\},M}(P) := \frac{1}{|\mathbb{S}^{(0)}|} \sum_{\chi^{(0)} \in \mathbb{S}^{(0)}} \widehat{\Phi}_{\chi^{(0)}}^{(0)} + \sum_{\ell=1}^L \left(\frac{1}{|\mathbb{S}^{(\ell)}|} \sum_{\chi^{(\ell)} \in \mathbb{S}^{(\ell)}} (\widehat{\Phi}_{\chi^{(\ell)}}^{(\ell)} - \widehat{\Phi}_{\chi^{(\ell)}}^{(\ell-1)}) \right).$$

The multilevel strategy (4.3) outperforms the single level approach (3.13) by taking most of the samples at the computationally inexpensive low levels, and only a few samples at the costly high levels. The efficiency of the MLMC methods in sDFT lies in two key aspects: (i) Design appropriate hierarchical approximations $\{\widehat{\mathcal{Q}}^{(\ell)}\}_{\ell=0}^L$ for $p_M(H[P])$ so that the variance decreases as ℓ increases; and (ii) Select optimal sampling sizes $|\mathbb{S}^{(\ell)}|$ to balance the computational cost and the approximation error. In the remainder of this subsection, we present the variance estimates and the strategy for optimizing the computational cost within a general MLMC framework.

With a hierarchical approximation $\{\widehat{\mathcal{Q}}^{(\ell)}\}_{\ell=0}^L$, the variance at each level is

$$(4.4) \quad \mathcal{V}_{\ell} := \mathbb{V}[\widehat{\Phi}_{\chi}^{(\ell)} - \widehat{\Phi}_{\chi}^{(\ell-1)}].$$

The following lemma provides a general estimate of the variance \mathcal{V}_ℓ for $\ell = 1, \dots, L$, whose proof is given in Appendix A.2.

LEMMA 4.1. *Let $\chi \in \mathbb{C}^n$ be the random orbital satisfying (3.1) and (3.8). Then there exists a constant C independent of n, N, M , such that*

$$(4.5) \quad \mathcal{V}_\ell \leq CN \|\widehat{\mathcal{Q}}^{(\ell-1)} - p_M(H[P])\|_{\mathbb{F}}^2 \quad \forall 1 \leq \ell \leq L.$$

Intuitively, the estimate (4.5) shows that as ℓ increases, if $\widehat{\mathcal{Q}}^{(\ell)}$ provides a better approximation of $p_M(H[P])$, then the variance decays and hence fewer random orbitals are required at higher levels.

Then we shall investigate the design of an optimal sampling strategy across all levels. Assume that at the ℓ -th level, the cost to evaluate $\widehat{\Phi}_\chi^{(\ell)} - \widehat{\Phi}_\chi^{(\ell-1)}$ for each random orbital $\chi \in \mathbb{S}^{(\ell)}$ is \mathcal{C}_ℓ . We see from (4.3) that the total cost and the total variance of the MLMC method are given respectively by $\mathcal{C} = \sum_{\ell=0}^L |\mathbb{S}^{(\ell)}| \mathcal{C}_\ell$ and $\mathcal{V} = \sum_{\ell=0}^L \mathcal{V}_\ell / |\mathbb{S}^{(\ell)}|$. To achieve a target accuracy $\mathcal{V}/\sqrt{N} = \epsilon$ (see Remark 3.2) while minimizing the total cost \mathcal{C} , the number of stochastic orbitals $|\mathbb{S}^{(\ell)}|$ at each level should be selected as

$$(4.6) \quad |\mathbb{S}^{(\ell)}| \propto \epsilon^{-2} \frac{1}{N} \sqrt{\frac{\mathcal{V}_\ell}{\mathcal{C}_\ell}} \left(\sum_{\ell=0}^L \sqrt{\mathcal{V}_\ell \mathcal{C}_\ell} \right).$$

Then the total cost for an MLMC approximation of the Kohn-Sham map, i.e. evaluation of (4.3), is estimated by

$$(4.7) \quad \mathcal{C} \propto \epsilon^{-2} \frac{1}{N} \left(\sum_{\ell=0}^L \sqrt{\mathcal{V}_\ell \mathcal{C}_\ell} \right)^2.$$

Note that both the variance \mathcal{V}_ℓ and the computational cost \mathcal{C}_ℓ depend on the hierarchical construction of $\widehat{\mathcal{Q}}^{(\ell)}$. Therefore, we can optimize the hierarchical structure by minimizing the total cost in (4.7), giving rise to the problem

$$(4.8) \quad \min_{\{\widehat{\mathcal{Q}}^{(\ell)}\}} \sum_{\ell=0}^L \sqrt{\mathcal{V}_\ell \mathcal{C}_\ell}.$$

In the rest of this section, we shall develop two MLMC approaches for sDFT calculations leveraging hierarchical structures derived from the plane-wave discretization and Chebyshev polynomial expansion, respectively. Note that the number of levels L is fixed in the following discussions and numerical experiments. Nevertheless our theories and methods can be extended to accommodate a varying L .

Remark 4.2 (Cost reduction by MLMC). In our following hierarchical constructions, the variance \mathcal{V}_ℓ decreases more rapidly than the cost \mathcal{C}_ℓ increases as the level ℓ grows. As a result, the product $\mathcal{V}_\ell \mathcal{C}_\ell$ decreases with ℓ , implying that the total cost of the MLMC approximation is dominated by $\mathcal{V}_0 \mathcal{C}_0$. Since the cost of standard sDFT can be approximated as $\mathcal{C}_L \mathcal{V}_0$, the MLMC approach reduces the cost of sDFT by a factor of $\mathcal{C}_0/\mathcal{C}_L$ (see also [17]).

4.2. Multilevel on energy cutoffs. In DFT calculations, the primary numerical parameter that determines the accuracy and computational cost is the energy cutoff E_c for the plane-wave basis set (2.4). We first exploit the energy cutoff parameter to construct hierarchical approximations of $p_M(H[P])$.

Let

$$0 < E_c^{(0)} < \dots < E_c^{(L)} = E_c$$

be a sequence of energy cutoffs. At the ℓ -th level, the energy cutoff $E_c^{(\ell)}$ defines a finite dimensional subspace $X_{E_c^{(\ell)}}(\Omega)$, whose degrees of freedom is given by $n^{(\ell)} := \dim(X_{E_c^{(\ell)}}(\Omega)) = \mathcal{O}((E_c^{(\ell)})^{\frac{d}{2}})$. For each level $\ell = 0, \dots, L$, we construct the corresponding Hamiltonian matrix $H^{(\ell)}[P] \in \mathbb{C}^{n^{(\ell)} \times n^{(\ell)}}$, whose matrix elements follow the same definition as (2.7). The idea is to approximate $p_M(H[P])$ by $p_M(H^{(\ell)}[P])$ at the ℓ -th level. Since $p_M(H^{(\ell)}[P]) \in \mathbb{C}^{n^{(\ell)} \times n^{(\ell)}}$ has a smaller matrix size than the full one $p_M(H[P]) \in \mathbb{C}^{n \times n}$, it must be expanded by zero-padding to ensure compatibility with the MLMC formula (4.3). In particular, matrix elements of $\widehat{Q}^{(\ell)} \in \mathbb{C}^{n \times n}$ at the ℓ -th level are defined by

$$(4.9) \quad (\widehat{Q}^{(\ell)})_{\mathbf{G}\mathbf{G}'} := \begin{cases} p_M(H^{(\ell)}[P])_{\mathbf{G}\mathbf{G}'} & \text{if } |\mathbf{G}|^2 \leq 2E_c^{(\ell)} \text{ and } |\mathbf{G}'|^2 \leq 2E_c^{(\ell)}, \\ 0 & \text{otherwise.} \end{cases}$$

The random matrix $\widehat{\Phi}_\chi^{(\ell)}$ in (4.1) can be obtained accordingly.

At the ℓ -th level, the evaluation of $\widehat{\Phi}_\chi^{(\ell)} - \widehat{\Phi}_\chi^{(\ell-1)}$ in (4.3) for each random orbital $\chi \in \mathbb{S}^{(\ell)}$ is achieved by recursive matrix-vector production, and the cost scales as

$$(4.10) \quad \mathcal{C}_\ell \propto n^{(\ell)} + n^{(\ell-1)} \propto (E_c^{(\ell)})^{\frac{d}{2}} + (E_c^{(\ell-1)})^{\frac{d}{2}},$$

where $n^{(-1)} = E_c^{(-1)} = 0$. We shall then use Lemma 4.1 to estimate the variance \mathcal{V}_ℓ with the multilevel construction (4.9). The following result shows that the variance decays exponentially with respect to the increasing energy cutoffs, the proof of which is given in Appendix A.3.

THEOREM 4.3. *Let χ be the random orbital satisfying (3.1) and (3.8), $\widehat{Q}^{(\ell)}$ be given by (4.9), $\widehat{\Phi}_\chi^{(\ell)}$ be given by (4.1), and \mathcal{V}_ℓ be given by (4.4). Then there exist constants C and γ depending on the smearing parameter β , such that*

$$(4.11) \quad \mathcal{V}_\ell \leq CN \left(\exp(-2\gamma\sqrt{E_c^{(\ell-1)}}) + \exp(-2\alpha M) \right) \quad \forall \ell = 1, \dots, L.$$

The second term in the estimate (4.11) is not relevant to the hierarchical construction (4.9), but comes from the polynomial approximation error. In practice, we can take sufficiently large M so that the variance becomes dominated by the first part of (4.11). Note that according to Lemma 3.1, we have $\mathcal{V}_0 = \mathcal{O}(N^2)$. Combining these with the cost \mathcal{C}_ℓ in (4.10), the optimization problem (4.8) can be reformulated as the following problem that finds the optimal energy cutoffs across all levels

$$(4.12) \quad \min_{\{E_c^{(\ell)}\}} \left\{ \sqrt{N}(E_c^{(0)})^{\frac{d}{4}} + \sum_{\ell=1}^L \exp(-\gamma\sqrt{E_c^{(\ell-1)}}) \sqrt{(E_c^{(\ell)})^{\frac{d}{2}} + (E_c^{(\ell-1)})^{\frac{d}{2}}} \right\}.$$

Note that we have omitted various constant factors in (4.12). For simplicity of implementations, we can choose energy cutoffs that follow an algebraic growth rate across levels

$$(4.13) \quad E_c^{(\ell)} = E_0 + (E_c - E_0) \left(\frac{\ell + s}{L + s} \right)^p \quad \text{with } p > 0.$$

Here, $E_0 > 0$ is a small energy cutoff for level $\ell = 0$ to capture the key characteristics of the system, and the parameter s is selected empirically. Then we can optimize the rate parameter p in (4.13) to determine the optimal cost for this MLMC strategy (see our numerical experiments in Section 5).

Remark 4.4 (Cost scaling for multilevel on energy cutoffs). We observe that the second term of the objective function in (4.12) can be bounded by a constant independent of E_c :

$$\sum_{\ell=1}^L e^{-\gamma\sqrt{E_c^{(\ell-1)}}} \sqrt{(E_c^{(\ell)})^{\frac{d}{2}} + (E_c^{(\ell-1)})^{\frac{d}{2}}} \leq C \int_{E_c^{(0)}}^{E_c} e^{-\gamma\sqrt{x}} x^{\frac{d}{4}} dx \leq CT \left(\frac{d}{2} + 2 \right),$$

where $\Gamma(\cdot)$ denotes the Gamma function. As a result, the total cost of the MLMC scheme with multilevel energy cutoffs is dominated by the cost at level $\ell = 0$, which is $\mathcal{O}(N^{-1}\mathcal{V}_0\mathcal{C}_0)$. Together with the fact $\mathcal{V}_0 = \mathcal{O}(N^2)$ and $\mathcal{C}_0 = n^{(0)}M$, the total cost scales as $\mathcal{O}(n^{(0)}NM)$, which does not depend on the discretization size n (and hence also independent of the finest energy cutoff E_c).

4.3. Multilevel on polynomial orders. The polynomial order M serves as another crucial numerical parameter that governs both accuracy and computational cost in sDFT. This motivates us to construct hierarchical approximations of $p_M(H[P])$ by selecting a sequence of increasing polynomial orders.

Let

$$0 < M^{(0)} < \dots < M^{(L)} = M.$$

We approximate $p_M(H[P])$ by

$$(4.14) \quad \widehat{\mathcal{Q}}^{(\ell)} := p_{M^{(\ell)}}(H[P]),$$

which gives rise to the corresponding random matrix $\widehat{\Phi}_\chi^{(\ell)}$ in (4.1). By using the recursive matrix-vector production, the computational cost of $\widehat{\Phi}_\chi^{(\ell)} - \widehat{\Phi}_\chi^{(\ell-1)}$ in (4.3) for each random orbital $\chi \in \mathbb{S}^{(\ell)}$ scales as

$$(4.15) \quad \mathcal{C}_\ell \propto M^{(\ell)}.$$

We next analyze the variance \mathcal{V}_ℓ at each level. The following result establishes that the variance \mathcal{V}_ℓ decays exponentially as ℓ increases, whose proof is given in Appendix A.4.

THEOREM 4.5. *Let χ be the random orbital satisfying (3.1) and (3.8), $\widehat{\mathcal{Q}}^{(\ell)}$ be given by (4.14), $\widehat{\Phi}_\chi^{(\ell)}$ be given by (4.1), and \mathcal{V}_ℓ be given by (4.4). Then there exist constants C and α depending on β , such that*

$$(4.16) \quad \mathcal{V}_\ell \leq CN \exp(-2\alpha M^{(\ell-1)}) \quad \forall \ell = 1, \dots, L.$$

Combining (4.15), (4.16) and $\mathcal{V}_0 = \mathcal{O}(N^2)$, we approximately obtain the optimization problem (4.8) for polynomial orders

$$(4.17) \quad \min_{\{M^{(\ell)}\}} \left\{ \sqrt{NM^{(0)}} + \sum_{\ell=1}^L \exp(-\alpha M^{(\ell-1)}) \sqrt{M^{(\ell)}} \right\}.$$

Analogous to the energy cutoff selection approach in (4.13), we assume an algebraic growth of $M^{(\ell)}$, by taking

$$(4.18) \quad M^{(\ell)} = M_0 + \left\lceil (M - M_0) \left(\frac{\ell + t}{L + t} \right)^q \right\rceil \quad \text{with } q > 0.$$

Here, $M_0 \in \mathbb{Z}_+$ is a small polynomial order to roughly capture the characteristics of the function f , and the parameter t is selected empirically. We then optimize the rate parameter q in (4.18) to determine the optimal cost for this MLMC strategy (see our numerical experiments in Section 5).

Remark 4.6 (Cost scaling for multilevel on polynomial orders). Similar to Remark 4.4, the cost of the multilevel on polynomial orders is $\mathcal{O}(N^{-1}\mathcal{V}_0\mathcal{C}_0)$. Combined with $\mathcal{V}_0 = \mathcal{O}(N^2)$ and $\mathcal{C}_0 = nM^{(0)}$, this multilevel approach yields a total computational cost of $\mathcal{O}(nNM^{(0)})$, which is independent of the polynomial order M (and hence also independent of the smearing parameter β).

5. Numerical experiments. In this section, we present numerical simulations of several 2D material systems using sDFT. Specifically, we consider three graphene-type systems of increasing complexity: perfect graphene lattice, graphene with a Stone–Wales defect, and boron/nitrogen-doped graphene. The corresponding atomic configurations are shown in Figure 1. For each system, the system size N is varied from 16 to 128 through the construction of supercells with different sizes.

All simulations are carried out on a workstation equipped with an Intel Xeon W-3275M 56-core processor and 1.5 TB of RAM. The computations are performed based on the packages DFTK.jl [19] (a library of Julia routines for plane-wave DFT algorithms) and SDFT.jl [32] (our Julia package for sDFT computations). For the DFT models, we employ the norm-conserving pseudopotentials in separable (Kleinman–Bylander) form [21] and the PW-LDA exchange–correlation functional [31].

We first test the variance of the random matrix $\phi(P, \chi, f^{\frac{1}{2}})$ by theoretical prediction (from Lemma 3.1) and numerical tests (from averaging 10 independent simulations). We choose the energy cutoff at $E_c = 20$ Ha and the smearing parameter at $\beta = 10^3$ Ha $^{-1}$ for all systems, and show the growth of the variance with respect to the system size in Figure 2. The results demonstrate excellent agreement between the numerically sampled variance and the theoretical prediction from (3.9) across all three different types of systems. Moreover, we see that the variance exhibits a quadratic increase with respect to the system size.

We further test the stochastic error of the electron density

$$\Delta\rho := \|\rho_{\mathbb{S}} - \rho_{\text{ref}}\|_{L^2(\Omega)},$$

where the reference density ρ_{ref} is obtained from a standard DFT calculation using the complete basis set $\{\mathbf{e}_i\}$. In Figure 3, we compare the stochastic error for unit cells of sizes 2×2 , 3×3 and 4×4 . We observe that the error scales linearly with $(N/|\mathbb{S}|)^{1/2}$ across all systems, which is consistent with our analysis in Remark 3.2.

We finally test the performance of the MLMC methods in sDFT for the three systems, with two multilevel strategies introduced in Section 4. We quantify the computational cost in terms of the number of floating-point operations determined by (4.7). For the multilevel strategy with energy cutoffs, we fix the smearing parameter at $\beta = 10^2$ Ha $^{-1}$ and employ an algebraic growth in (4.13) with parameters $E_0 = 6.0$ Ha and $s = 0.1$. We show in Figure 4 the variance of both $\hat{\phi}_{\chi}^{(\ell)}$ and $\hat{\phi}_{\chi}^{(\ell)} - \hat{\phi}_{\chi}^{(\ell-1)}$ across levels with the cutoff energy $E_c = 40$ Ha. It is observed that $\mathbb{V}[\hat{\phi}_{\chi}^{(\ell)} - \hat{\phi}_{\chi}^{(\ell-1)}]$ decays exponentially as ℓ increases, while the variance $\mathbb{V}[\hat{\phi}_{\chi}^{(\ell)}]$ remains nearly unchanged across levels. This is consistent with our discussion in Theorem 4.3, indicating that the MLMC scheme requires significantly less random orbitals than the standard sampling scheme. In Figure 5, we compare the computational costs with three different energy cutoffs $E_c = 20, 30, 40$ Ha (on the finest level) with corresponding numbers of levels




FIG. 1. Atomic configurations of the graphene-type systems. (a) Perfect lattice. (b) Stone-Wales defect. (c) Boron (blue) / nitrogen (red) doped graphene.




FIG. 2. The variance of the random matrix $\phi(P, \chi, f^{\frac{1}{2}})$ with respect to the number of electrons.




FIG. 3. The statistical error of electron density with respect to $(N/|S|)^{1/2}$.

$L = 2, 3, 4$, respectively. We observe that the computational costs align along a line of N^2M , indicating that the cost is independent of E_c with the MLMC scheme. Given that the degrees of freedom scales as $n \propto E_c^{3/2}N$, the cost scaling $\mathcal{O}(N^2M)$ is also consistent with our analysis in Remark 4.4.

For the multilevel strategy with polynomial orders, we fix the energy cutoff at $E_c = 20$ Ha, and employ an algebraic growth in (4.18) with parameters $M_0 = 85$ and $t = 0$. In Figure 6, we show the variance of random matrices with the smearing parameter $\beta = 10^4$ Ha $^{-1}$, and observe that $\mathbb{V}[\hat{\phi}_\chi^{(\ell)} - \hat{\phi}_\chi^{(\ell-1)}]$ decays exponentially with

ℓ , while $\mathbb{V}[\widehat{\Phi}_\chi^{(\ell)}]$ remains nearly constant. In Figure 7, we compare the computational costs against nN for different smearing parameters $\beta = 10^2, 10^3, 10^4 \text{ Ha}^{-1}$, with corresponding numbers of levels $L = 2, 3, 4$, respectively. We see that all costs align along a line, indicating that the cost of the multilevel approach is independent of the smearing parameter β .

6. Conclusions. In this work, we study the sDFT method under a plane-wave discretization and explore various variance reduction strategies within the MLMC framework. In particular, we develop two hierarchical approaches based on plane-wave cutoffs and Chebyshev expansion orders, which render the computational cost independent of the discretization size and the temperature, respectively. Our analysis also provides theoretical insights into existing variance-reduction techniques in sDFT, such as the energy-window method [11] and the tempering method [29].

Appendix A. Proofs.

A.1. Proof of Lemma 3.1. Let χ be the random orbital satisfying (3.1). Denote by $A := f^{\frac{1}{2}}(H[P])$, $B := AA^h$, and $B_\chi := (A\chi)(A\chi)^h$. Then to show (3.7), it is only necessary to prove

$$(A.1) \quad \mathbb{V}[B_\chi] = \sum_{i,j=1, i \neq j}^n B_{ii}B_{jj} + \sum_{i,j=1, i \neq j}^n B_{ij}^2 \mathbb{E}[\bar{\chi}_i^2] \mathbb{E}[\chi_j^2] + \sum_{i=1}^n B_{ii}^2 (\mathbb{E}[|\chi_i|^4] - 1).$$

We first express the matrix elements of B and B_χ explicitly

$$B_{ij} = \sum_k A_{ik} \bar{A}_{jk} \quad \text{and} \quad (B_\chi)_{ij} = ((A\chi)(A\chi)^h)_{ij} = \sum_{k,\ell} A_{ik} \bar{A}_{j\ell} \chi_k \bar{\chi}_\ell.$$

By using (3.1), we can derive the expectation and variance of $(B_\chi)_{ij}$ as

$$\begin{aligned} \mathbb{E}[(B_\chi)_{ij}] &= \sum_{k,\ell} A_{ik} \bar{A}_{j\ell} \mathbb{E}[\chi_k \bar{\chi}_\ell] = \sum_k A_{ik} \bar{A}_{jk} \quad \text{and} \\ \mathbb{V}[(B_\chi)_{ij}] &= \mathbb{E}[|(B_\chi)_{ij}|^2] - |\mathbb{E}[(B_\chi)_{ij}]|^2 \\ &= \sum_{k,\ell,s,t} \bar{A}_{ik} A_{j\ell} A_{is} \bar{A}_{jt} \mathbb{E}[\bar{\chi}_k \chi_\ell \chi_s \bar{\chi}_t] - \sum_{k,s} \bar{A}_{ik} A_{jk} A_{is} \bar{A}_{js} \\ (A.2) \quad &= \sum_{k \neq \ell, s \neq t} \bar{A}_{ik} A_{j\ell} A_{is} \bar{A}_{jt} \mathbb{E}[\bar{\chi}_k \chi_\ell \chi_s \bar{\chi}_t] + \sum_k |A_{ik}|^2 |A_{jk}|^2 (\mathbb{E}[|\chi_k|^4] - 1). \end{aligned}$$

To estimate the first term of (A.2), we have from (3.1) that $\mathbb{E}[\bar{\chi}_k \chi_\ell \chi_s \bar{\chi}_t] \neq 0$ only if $(k, \ell) = (s, t)$ or $(k, \ell) = (t, s)$. This indicates

$$\begin{aligned} &\sum_{k \neq \ell, s \neq t} \bar{A}_{ik} A_{j\ell} A_{is} \bar{A}_{jt} \mathbb{E}[\bar{\chi}_k \chi_\ell \chi_s \bar{\chi}_t] \\ &= \sum_{k \neq \ell} |A_{ik}|^2 |A_{j\ell}|^2 \mathbb{E}[\bar{\chi}_k \chi_\ell \chi_k \bar{\chi}_\ell] + \sum_{k \neq \ell} \bar{A}_{ik} A_{j\ell} A_{i\ell} \bar{A}_{jk} \mathbb{E}[\bar{\chi}_k \chi_\ell \chi_\ell \bar{\chi}_k] \\ (A.3) \quad &= \sum_{k \neq \ell} |A_{ik}|^2 |A_{j\ell}|^2 + \sum_{k \neq \ell} \bar{A}_{ik} A_{j\ell} A_{i\ell} \bar{A}_{jk} \mathbb{E}[\bar{\chi}_k^2] \mathbb{E}[\chi_\ell^2]. \end{aligned}$$




FIG. 4. Variance across levels for multilevel energy cutoffs.




FIG. 5. Computational cost with respect to N^2M , for multilevel energy cutoffs.

FIG. 6. Variance across levels for multilevel polynomial orders.

FIG. 7. Computational cost with respect to nN , for multilevel polynomial orders.

Note that $H[P] \in \mathcal{D}$ implies $A \in \mathcal{D}$, therefore we have $A = A^h$ and

$$(A.4) \quad \sum_{i,j} |A_{ik}|^2 |A_{j\ell}|^2 = \left(\sum_i |A_{ki}|^2 \right) \left(\sum_j |A_{\ell j}|^2 \right) = B_{kk} B_{\ell\ell},$$

$$(A.5) \quad \sum_{i,j} \bar{A}_{ik} A_{j\ell} A_{i\ell} \bar{A}_{jk} = \left(\sum_i A_{ki} \bar{A}_{\ell i} \right) \left(\sum_j A_{kj} \bar{A}_{\ell j} \right) = B_{k\ell}^2.$$

By taking into accounts (A.2)-(A.5), we obtain

$$\mathbb{V}[B_\chi] = \sum_{i,j} \mathbb{V}[(B_\chi)_{ij}] = \sum_{k \neq \ell} B_{kk} B_{\ell\ell} + \sum_{k \neq \ell} B_{k\ell}^2 \mathbb{E}[\bar{\chi}_k^2] \mathbb{E}[\chi_\ell^2] + \sum_k B_{kk}^2 \left(\mathbb{E}[|\chi_k|^4] - 1 \right)$$

which completes the proof of (A.1) (and hence (3.7)).

The result (3.9) follows immediately from (3.7) if the random orbital χ_i additionally satisfy the conditions in (3.8). This completes the proof. \square

A.2. Proof of Lemma 4.1. For simplicity of the presentation, we will denote $p_M(H[P])$ by \mathcal{P}_M in this proof. We have from (4.4) that

$$(A.6) \quad \mathcal{V}_\ell = \mathbb{V}[\hat{\Phi}_\chi^{(\ell)} - \hat{\Phi}_\chi^{(\ell-1)}] \leq \left(\sqrt{\mathbb{V}[\hat{\Phi}_\chi^{(\ell)} - \phi(P, \chi, p_M)]} + \sqrt{\mathbb{V}[\hat{\Phi}_\chi^{(\ell-1)} - \phi(P, \chi, p_M)]} \right)^2.$$

By using the formulas (3.3) for $\phi(P, \chi, p_M)$ and (4.1) for $\hat{\Phi}_\chi^{(\ell)}$, we have

$$\begin{aligned} & \mathbb{V}[\hat{\Phi}_\chi^{(\ell)} - \phi(P, \chi, p_M)] \\ &= \mathbb{V}\left[(\hat{\mathcal{Q}}^{(\ell)} \chi)(\hat{\mathcal{Q}}^{(\ell)} \chi)^h - (\mathcal{P}_M \chi)(\mathcal{P}_M \chi)^h\right] \\ &= \mathbb{V}\left[(\hat{\mathcal{Q}}^{(\ell)} \chi) \left((\hat{\mathcal{Q}}^{(\ell)} - \mathcal{P}_M) \chi \right)^h + \left((\hat{\mathcal{Q}}^{(\ell)} - \mathcal{P}_M) \chi \right) (\mathcal{P}_M \chi)^h\right] \\ &\leq \left(\sqrt{\mathbb{V}\left[(\hat{\mathcal{Q}}^{(\ell)} \chi) \left((\hat{\mathcal{Q}}^{(\ell)} - \mathcal{P}_M) \chi \right)^h\right]} + \sqrt{\mathbb{V}\left[\left((\hat{\mathcal{Q}}^{(\ell)} - \mathcal{P}_M) \chi \right) (\mathcal{P}_M \chi)^h\right]} \right)^2. \end{aligned}$$

By using a proof similar to that of Lemma 3.1, we can obtain

$$\begin{aligned} \mathbb{V}\left[(\hat{\mathcal{Q}}^{(\ell)} \chi) \left((\hat{\mathcal{Q}}^{(\ell)} - \mathcal{P}_M) \chi \right)^h\right] &\leq \text{Tr}\left((\hat{\mathcal{Q}}^{(\ell)})^2\right) \text{Tr}\left((\hat{\mathcal{Q}}^{(\ell)} - \mathcal{P}_M)^2\right) \quad \text{and} \\ \mathbb{V}\left[\left((\hat{\mathcal{Q}}^{(\ell)} - \mathcal{P}_M) \chi \right) (\mathcal{P}_M \chi)^h\right] &\leq \text{Tr}(\Phi_M(P)) \text{Tr}\left((\hat{\mathcal{Q}}^{(\ell)} - \mathcal{P}_M)^2\right). \end{aligned}$$

Since $\text{Tr}\left((\hat{\mathcal{Q}}^{(\ell)})^2\right) = \mathcal{O}(N)$ and $\text{Tr}(\Phi_M(P)) = \mathcal{O}(N)$, we have

$$\mathbb{V}[\hat{\Phi}_\chi^{(\ell)} - \phi(P, \chi, p_M)] \leq CN \|\hat{\mathcal{Q}}^{(\ell)} - \mathcal{P}_M\|_{\mathbb{F}}^2.$$

This together with (A.6) implies

$$\mathcal{V}_\ell \leq CN \left(\|\hat{\mathcal{Q}}^{(\ell)} - \mathcal{P}_M\|_{\mathbb{F}} + \|\hat{\mathcal{Q}}^{(\ell-1)} - \mathcal{P}_M\|_{\mathbb{F}} \right)^2 \leq CN \|\hat{\mathcal{Q}}^{(\ell-1)} - \mathcal{P}_M\|_{\mathbb{F}}^2,$$

where the second inequality follows from the fact that the accuracy of $\hat{\mathcal{Q}}^{(\ell)}$ increases with ℓ . This completes the proof. \square

A.3. Proof of Theorem 4.3. Let $\widehat{\mathcal{Q}}^{(\ell-1)}$ be given by (4.9). Using Lemma 4.1, it is only necessary for us to prove

$$(A.7) \quad \left\| \widehat{\mathcal{Q}}^{(\ell-1)} - p_M(H[P]) \right\|_{\mathbb{F}}^2 \leq C \left(e^{-2\gamma\sqrt{E_c^{(\ell-1)}}} + e^{-2\alpha M} \right) \quad \forall 1 \leq \ell \leq L.$$

In the rest of this proof, we will denote $H[P]$ by H for simplicity of the presentation. Let $\widetilde{H^{(\ell-1)}}$ be the zero-padded matrix of $H^{(\ell-1)}$ over $\mathbb{C}^{n \times n}$, that is

$$(A.8) \quad \widetilde{H^{(\ell-1)}}_{\mathbf{G}\mathbf{G}'} := \begin{cases} H_{\mathbf{G}\mathbf{G}'} & \text{if } |\mathbf{G}|^2, |\mathbf{G}'|^2 \leq 2E_c^{(\ell-1)}, \\ 0 & \text{otherwise.} \end{cases}$$

Then we can decompose the left-hand side of (A.7) into three parts

$$\begin{aligned} & \left\| \widehat{\mathcal{Q}}^{(\ell-1)} - p_M(H) \right\|_{\mathbb{F}} \\ & \leq \left\| p_M(H) - f^{\frac{1}{2}}(H) \right\|_{\mathbb{F}} + \left\| p_M(H^{(\ell-1)}) - f^{\frac{1}{2}}(H^{(\ell-1)}) \right\|_{\mathbb{F}} + \left\| f^{\frac{1}{2}}(\widetilde{H^{(\ell-1)}}) - f^{\frac{1}{2}}(H) \right\|_{\mathbb{F}} \\ & =: I_1 + I_2 + I_3, \end{aligned}$$

where the first two terms I_1 and I_2 correspond to the polynomial approximation error, and the last term I_3 captures the energy cutoff error.

We first estimate I_1 . Note that H is a scaled Hamiltonian such that all the eigenvalues lie in $[-1, 1]$. Therefore, we can obtain from (3.14) that

$$(A.9) \quad I_1 = \left\| p_M(H) - f^{\frac{1}{2}}(H) \right\|_{\mathbb{F}} \leq C \left\| p_M - f^{\frac{1}{2}} \right\|_{L^\infty([-1, 1])} \leq C e^{-\alpha M}.$$

By using a similar argument, we can estimate the second term I_2 as

$$(A.10) \quad I_2 \leq C e^{-\alpha M}.$$

For the last term I_3 , we have

$$(A.11) \quad I_3 \leq C e^{-\gamma\sqrt{E_c^{(\ell-1)}}},$$

which will be proved in the rest of this section. The argument is based on two key observations: (i) the exponential convergence of $f^{\frac{1}{2}}(H^{(\ell-1)})$ with the energy cutoff $E_c^{(\ell-1)}$, and (ii) the exponential off-diagonal decay of its matrix elements. We will state and prove them in Lemma A.1 and Lemma A.2 respectively in the following.

Combining the estimates of (A.9), (A.10) and (A.11), we can obtain (A.7), which completes the proof of Theorem 4.3. \square

LEMMA A.1. *There exist positive constants C and γ depending on β such that*

$$(A.12) \quad \left| f^{\frac{1}{2}}(\widetilde{H^{(\ell-1)}})_{\mathbf{G}\mathbf{G}'} - f^{\frac{1}{2}}(H)_{\mathbf{G}\mathbf{G}'} \right| \leq C e^{-\gamma\sqrt{E_c^{(\ell-1)}}}$$

for any $1 \leq \ell \leq L$ and $|\mathbf{G}|^2, |\mathbf{G}'|^2 \leq E_c^{(\ell-1)}$.

Proof. Let $\mathfrak{d}(\cdot)$ denote the spectrum of a matrix. Then we have from (A.8) that $\mathfrak{d}(\widetilde{H^{(\ell-1)}}) = \mathfrak{d}(H^{(\ell-1)}) \cup \{0\}$. Let Γ be a contour that encloses the spectrum of $\widetilde{H^{(\ell-1)}}$ and H , and avoids all the singularities of $f^{\frac{1}{2}}$, while satisfying

$$(A.13) \quad \min \left\{ \text{dist}(\Gamma, \mathfrak{d}(H)), \text{dist}(\Gamma, \mathfrak{d}(\widetilde{H^{(\ell-1)}})) \right\} \geq \sigma_\beta.$$

Here σ_β depends on β .

Using the smoothness assumption for the potentials of the Kohn-Sham Hamiltonian in (2.1), we have that there exists some $\gamma_h > 0$ such that

$$(A.14) \quad \left| \widetilde{H^{(\ell-1)}}_{\mathbf{G}\mathbf{G}'} \right| \leq C e^{-\gamma_h |\mathbf{G}-\mathbf{G}'|} \quad \text{and} \quad \left| H_{\mathbf{G}\mathbf{G}'} \right| \leq C e^{-\gamma_h |\mathbf{G}-\mathbf{G}'|}.$$

Then by using (A.13), (A.14) and a Combes-Thomas type estimate [13] (see also similar arguments in [10, Lemma 6]), we have that there exists constants $C > 0$ and $\gamma_{h,\beta} > 0$ depending on β , such that for any $z \in \Gamma$,

$$(A.15) \quad \left| (z - \widetilde{H^{(\ell-1)}})_{\mathbf{G}\mathbf{G}'}^{-1} \right| \leq C e^{-\gamma_{h,\beta} |\mathbf{G}-\mathbf{G}'|}, \quad \left| (z - H)_{\mathbf{G}\mathbf{G}'}^{-1} \right| \leq C e^{-\gamma_{h,\beta} |\mathbf{G}-\mathbf{G}'|}.$$

Note that (A.8) and (A.14) implies

$$\left| (H - \widetilde{H^{(\ell-1)}})_{\mathbf{G}''\mathbf{G}'''} \right| \leq \begin{cases} 0 & \text{if } |\mathbf{G}''|^2, |\mathbf{G}'''|^2 \leq 2E_c^{(\ell-1)}, \\ C e^{-\gamma_h |\mathbf{G}''-\mathbf{G}'''} & \text{otherwise.} \end{cases}$$

This together with (A.15) and $|\mathbf{G}|^2, |\mathbf{G}'|^2 \leq 2E_c^{(\ell-1)}$ implies that

$$\begin{aligned} & \left| (z - \widetilde{H^{(\ell-1)}})_{\mathbf{G}\mathbf{G}'}^{-1} - (z - H)_{\mathbf{G}\mathbf{G}'}^{-1} \right| \\ & \leq \sum_{\substack{|\mathbf{G}''|^2 \leq 2E_c \\ |\mathbf{G}'''|^2 \leq 2E_c}} \left| (z - \widetilde{H^{(\ell-1)}})_{\mathbf{G}\mathbf{G}''}^{-1} (H - \widetilde{H^{(\ell-1)}})_{\mathbf{G}''\mathbf{G}'''} (z - H)_{\mathbf{G}'''\mathbf{G}'}^{-1} \right| \\ & \leq C \left(\sum_{\substack{2E_c^{(\ell-1)} < |\mathbf{G}''|^2 \leq 2E_c \\ |\mathbf{G}'''|^2 \leq 2E_c}} + \sum_{\substack{|\mathbf{G}''|^2 \leq 2E_c^{(\ell-1)} \\ 2E_c^{(\ell-1)} < |\mathbf{G}'''|^2 \leq 2E_c}} \right) e^{-(\gamma_{h,\beta} |\mathbf{G}-\mathbf{G}''| + \gamma_h |\mathbf{G}''-\mathbf{G}'''| + \gamma_{h,\beta} |\mathbf{G}'''-\mathbf{G}'|)} \\ & \leq C \sum_{2E_c^{(\ell-1)} < |\mathbf{G}''|^2 \leq 2E_c} e^{-\gamma (|\mathbf{G}-\mathbf{G}''| + |\mathbf{G}'-\mathbf{G}''|)} \leq C e^{-\gamma \sqrt{E_c^{(\ell-1)}}}, \end{aligned}$$

where the constants C and γ depend on β . This together with the contour integral representation implies

$$\begin{aligned} & \left| f^{\frac{1}{2}}(\widetilde{H^{(\ell-1)}})_{\mathbf{G}\mathbf{G}'} - f^{\frac{1}{2}}(H)_{\mathbf{G}\mathbf{G}'} \right| \\ & \leq C \oint_{\Gamma} |f^{\frac{1}{2}}(z)| \cdot \left| (z - \widetilde{H^{(\ell-1)}})_{\mathbf{G}\mathbf{G}'}^{-1} - (z - H)_{\mathbf{G}\mathbf{G}'}^{-1} \right| dz \leq C e^{-\gamma \sqrt{E_c^{(\ell-1)}}} \oint_{\Gamma} |f^{\frac{1}{2}}(z)| dz. \end{aligned}$$

Since the Hamiltonian H is bounded from below and $f^{\frac{1}{2}}$ exhibits an exponential decay, we have that $\oint_{\Gamma} |f^{\frac{1}{2}}(z)| dz$ is bounded by some constant independent of $E_c^{(\ell)}$. Therefore, the estimate (A.12) holds and the proof is completed. \square

LEMMA A.2. *There exist positive constants C and γ depending on β , such that*

$$(A.16) \quad \left| f^{\frac{1}{2}}(H^{(\ell-1)})_{\mathbf{G}\mathbf{G}'} \right| \leq C e^{-\gamma (\min\{|\mathbf{G}|, |\mathbf{G}'|\} + |\mathbf{G}-\mathbf{G}'|)}$$

for any $1 \leq \ell \leq L+1$.

Proof. Without loss of generality, we assume that $|\mathbf{G}| \leq |\mathbf{G}'|$. Let $\bar{R} := |\mathbf{G}|/2$, we will prove (A.16) by considering two cases $|\mathbf{G} - \mathbf{G}'| < \bar{R}/2$ and $|\mathbf{G} - \mathbf{G}'| \geq \bar{R}/2$, respectively.

For the first case with $|\mathbf{G} - \mathbf{G}'| < \bar{R}/2$, we have $|\mathbf{G}'| \geq |\mathbf{G}| - \bar{R}/2 > \bar{R}$. Define

$$H_{\bar{R}}^{(\ell-1)} := \begin{cases} H_{\mathbf{G}\mathbf{G}'} & \text{if } \bar{R}^2/4 < |\mathbf{G}|^2, |\mathbf{G}'|^2 \leq 2E_c^{(\ell-1)}, \\ 0 & \text{otherwise.} \end{cases}$$

We then observe that

$$(A.17) \quad \left| f^{\frac{1}{2}}(H^{(\ell-1)})_{\mathbf{G}\mathbf{G}'} \right| \leq \left| f^{\frac{1}{2}}(H^{(\ell-1)})_{\mathbf{G}\mathbf{G}'} - f^{\frac{1}{2}}(H_{\bar{R}}^{(\ell-1)})_{\mathbf{G}\mathbf{G}'} \right| + \left| f^{\frac{1}{2}}(H_{\bar{R}}^{(\ell-1)})_{\mathbf{G}\mathbf{G}'} \right|.$$

To estimate the first term of (A.17), we can employ a similar proof to that of Lemma A.1 and obtain

$$\left| f^{\frac{1}{2}}(H^{(\ell-1)})_{\mathbf{G}\mathbf{G}'} - f^{\frac{1}{2}}(H_{\bar{R}}^{(\ell-1)})_{\mathbf{G}\mathbf{G}'} \right| \leq Ce^{-\gamma\bar{R}}.$$

This together with $|\mathbf{G} - \mathbf{G}'| < \bar{R}/2 = |\mathbf{G}|/4$ leads to

$$(A.18) \quad \left| f^{\frac{1}{2}}(H^{(\ell-1)})_{\mathbf{G}\mathbf{G}'} - f^{\frac{1}{2}}(H_{\bar{R}}^{(\ell-1)})_{\mathbf{G}\mathbf{G}'} \right| \leq Ce^{-\gamma(|\mathbf{G}|+|\mathbf{G}-\mathbf{G}'|)}.$$

To estimate the second term of (A.17), we recall that the Hamiltonian is scaled and shifted by constants a and b respectively (introduced at the beginning of Section 3). Then we have from the decay estimate (A.14) that there exists a constant $C_V > 0$ such that $\sum_{\mathbf{G}' \neq \mathbf{G}''} |(H_{\bar{R}}^{(\ell-1)})_{\mathbf{G}'\mathbf{G}''}| \leq |a|C_V$. By using the Gershgorin circle theorem, we have

$$\mathfrak{d}(H_{\bar{R}}^{(\ell-1)}) \subset \bigcup_{\bar{R}^2/4 < |\mathbf{G}''|^2 \leq 2E_c^{(\ell-1)}} \left\{ \lambda \in \mathbb{R} : \left| \lambda - a \left(\frac{1}{2} |\mathbf{G}''|^2 - b \right) \right| < |a|C_V \right\}.$$

Therefore, there exists a contour Γ enclosing the spectrum of $H_{\bar{R}}^{(\ell-1)}$, such that it avoids all the singularities of $f^{\frac{1}{2}}$, and satisfies $\text{dist}(\Gamma, \mathfrak{d}(H_{\bar{R}}^{(\ell-1)})) > \sigma_\beta$ and

$$\left| \frac{1}{a} \text{Re}(z) + b \right| \geq \min_{\bar{R}^2/4 < |\mathbf{G}''|^2 \leq 2E_c^{(\ell-1)}} \frac{1}{2} |\mathbf{G}''|^2 - C_V - 1 > \frac{1}{8} \bar{R}^2 - C_V - 1 > \frac{1}{8} \bar{R} - \bar{C}_V$$

for any $z \in \Gamma$, where the constant \bar{C}_V only depends on C_V . Note that $f^{\frac{1}{2}}$ decays exponentially, i.e., $|f^{\frac{1}{2}}(z)| \leq Ce^{-\zeta_\beta |\frac{z}{a} + b|}$ for some ζ_β depending on β . This together with the estimate (A.15) implies

$$(A.19) \quad \left| f^{\frac{1}{2}}(H_{\bar{R}}^{(\ell-1)})_{\mathbf{G}\mathbf{G}'} \right| \leq C \oint_{\Gamma} |f^{\frac{1}{2}}(z)| \cdot \left| (z - H_{\bar{R}}^{(\ell-1)})_{\mathbf{G}\mathbf{G}'}^{-1} \right| dz \\ \leq Ce^{-\zeta_\beta (\frac{\bar{R}}{8} - \bar{C}_V)} e^{-\gamma(|\mathbf{G}-\mathbf{G}'|)} \leq Ce^{-\gamma(|\mathbf{G}|+|\mathbf{G}-\mathbf{G}'|)}.$$

Combining (A.17), (A.18), (A.19), we obtain the estimate (A.16).

We then consider the second case with $|\mathbf{G} - \mathbf{G}'| \geq \bar{R}/2$. By using (A.15) and the contour integral representation, we have

$$\left| f^{\frac{1}{2}}(H^{(\ell-1)})_{\mathbf{G}\mathbf{G}'} \right| \leq Ce^{-\gamma|\mathbf{G}-\mathbf{G}'|},$$

which together with $|\mathbf{G} - \mathbf{G}'| \geq \bar{R}/2 = |\mathbf{G}|/4$ leads to (A.16). \square

We are now ready to prove the estimate for I_3 .

Proof of (A.11). We decompose the basis into two ranges $|\mathbf{G}|^2 \leq \frac{1}{8}E_c^{(\ell-1)}$ and $\frac{1}{8}E_c^{(\ell-1)} < |\mathbf{G}|^2 < 2E_c$, and then obtain

$$(A.20) \quad I_3 \leq \left(\sum_{\substack{|\mathbf{G}|^2 \leq \frac{1}{8}E_c^{(\ell-1)} \\ |\mathbf{G}'|^2 \leq \frac{1}{8}E_c^{(\ell-1)}}} + 2 \sum_{\substack{\frac{1}{8}E_c^{(\ell-1)} < |\mathbf{G}|^2 \leq 2E_c \\ |\mathbf{G}'|^2 \leq 2E_c}} \right) \left| f^{\frac{1}{2}}(\widetilde{H^{(\ell-1)}})_{\mathbf{G}\mathbf{G}'} - f^{\frac{1}{2}}(H)_{\mathbf{G}\mathbf{G}'} \right|$$

$$=: T_1 + T_2.$$

To estimate T_1 , we use (A.12) from Lemma A.1, which gives

$$\left| f^{\frac{1}{2}}(\widetilde{H^{(\ell-1)}})_{\mathbf{G}\mathbf{G}'} - f^{\frac{1}{2}}(H)_{\mathbf{G}\mathbf{G}'} \right| \leq C e^{-\gamma \sqrt{E_c^{(\ell-1)}}} \leq C e^{-\gamma(\sqrt{E_c^{(\ell-1)}} - |\mathbf{G}| - |\mathbf{G}'|)},$$

Therefore,

$$(A.21) \quad T_1 \leq C \sum_{\substack{|\mathbf{G}|^2 \leq \frac{1}{8}E_c^{(\ell-1)} \\ |\mathbf{G}'|^2 \leq \frac{1}{8}E_c^{(\ell-1)}}} e^{-\gamma(\sqrt{E_c^{(\ell-1)}} - |\mathbf{G}| - |\mathbf{G}'|)} \leq C e^{-\gamma \sqrt{E_c^{(\ell-1)}}}.$$

We then use (A.16) from Lemma A.2 to obtain the estimate for T_2 ,

$$(A.22) \quad T_2 \leq \sum_{\substack{\frac{1}{8}E_c^{(\ell-1)} < |\mathbf{G}|^2 \leq 2E_c \\ |\mathbf{G}'|^2 \leq 2E_c, |\mathbf{G}'| \leq |\mathbf{G}|}} e^{-\gamma(|\mathbf{G}'| + |\mathbf{G} - \mathbf{G}'|)} + \sum_{\substack{\frac{1}{8}E_c^{(\ell-1)} < |\mathbf{G}|^2 \leq 2E_c \\ |\mathbf{G}'|^2 \leq 2E_c, |\mathbf{G}'| > |\mathbf{G}|}} e^{-\gamma(|\mathbf{G}| + |\mathbf{G} - \mathbf{G}'|)}$$

$$\leq C e^{-\gamma \sqrt{E_c^{(\ell-1)}}}.$$

Combining (A.20), (A.21) and (A.22), we obtain (A.11) and complete the proof. \square

A.4. Proof of Theorem 4.5. Let $\widehat{\mathcal{Q}}^{(\ell-1)}$ be given by (4.14). Using Lemma 4.1, it suffices to prove that there exist constants C and α depending on β , such that

$$(A.23) \quad \|\widehat{\mathcal{Q}}^{(\ell-1)} - p_M(H[P])\|_{\mathbb{F}}^2 \leq C e^{-2\alpha M^{(\ell-1)}} \quad \forall 1 \leq \ell \leq L.$$

Note that

$$(A.24) \quad \|\widehat{\mathcal{Q}}^{(\ell-1)} - p_M(H[P])\|_{\mathbb{F}} \leq C \|p_{M^{(\ell-1)}} - p_M\|_{L^\infty([-1,1])}.$$

By using the error estimate (3.14) for Chebyshev polynomial approximation, we have

$$\|p_{M^{(\ell-1)}} - p_M\|_{L^\infty([-1,1])} \leq \|p_{M^{(\ell-1)}} - f^{\frac{1}{2}}\|_{L^\infty([-1,1])} + \|p_M - f^{\frac{1}{2}}\|_{L^\infty([-1,1])}$$

$$\leq C(e^{-\alpha M^{(\ell-1)}} + e^{-\alpha M}) \leq C e^{-\alpha M^{(\ell-1)}}.$$

This together with (A.24) implies (A.23) and hence completes the proof. \square

- [1] D. F. ANDERSON AND D. J. HIGHAM, Multilevel monte carlo for continuous time markov chains, with applications in biochemical kinetics, Multiscale Model. Simul., 10 (2012), pp. 146–179.
- [2] R. BAER AND M. HEAD-GORDON, Sparsity of the density matrix in Kohn-Sham density functional theory and an assessment of linear system-size scaling methods, Phys. Rev. Lett., 79 (1997), pp. 3962–3965.
- [3] R. BAER, D. NEUHAUSER, AND E. RABANI, Self-averaging stochastic Kohn-Sham density-functional theory, Phys. Rev. Lett., 111 (2013), 106402.
- [4] A. BARTH, C. SCHWAB, AND N. ZOLLINGER, Multi-level Monte Carlo finite element method for elliptic PDEs with stochastic coefficients., Numer. Math., 119 (2011), pp. 123–161.
- [5] Z. W. BIRNBAUM, J. RAYMOND, AND H. S. ZUCKERMAN, A generalization of Tshebyshev’s inequality to two dimensions, Ann. Math. Statist., 18 (1947), pp. 70 – 79.
- [6] Y. CAI AND M. LINDSEY, Toward optimal-scaling DFT: stochastic Hartree theory in the thermodynamic and complete basis set limits at arbitrary temperature, 2025, <https://arxiv.org/abs/2504.15816>.
- [7] E. CANCÈS, R. CHAKIR, AND Y. MADAY., Numerical analysis of nonlinear eigenvalue problems, J. Sci. Comput., 45 (2010), pp. 90–117.
- [8] E. CANCÈS, R. CHAKIR, AND Y. MADAY, Numerical analysis of the planewave discretization of some orbital-free and Kohn-Sham models, ESAIM: M2AN., 46 (2012), pp. 341–388.
- [9] H. CHEN, X. GONG, L. HE, Z. YANG, AND A. ZHOU, Numerical analysis of finite dimensional approximations of Kohn-Sham models, Adv. Compu. Math., 38 (2011), pp. 1–32.
- [10] H. CHEN AND C. ORTNER, QM/MM methods for crystalline defects. part 2: Consistent energy and force-mixing, Multiscale Model. Simul., 15 (2017), pp. 184–214, <https://doi.org/10.1137/15M1041250>.
- [11] M. CHEN, R. BAER, D. NEUHAUSER, AND E. RABANI, Energy window stochastic density functional theory, J. Chem. Phys., 151 (2019), 114116.
- [12] M. CHEN, R. BAER, D. NEUHAUSER, AND E. RABANI, Overlapped embedded fragment stochastic density functional theory for covalently-bonded materials, J. Chem. Phys., 150 (2019), 034106.
- [13] J. COMBES AND L. THOMAS, Asymptotic behaviour of eigenfunctions for multiparticle Schrödinger operators, Commun. Math. Phys., 34 (1973), pp. 251–270.
- [14] M. D. FABIAN, B. SHPIRO, E. RABANI, D. NEUHAUSER, AND R. BAER, Stochastic density functional theory, WIREs Comput. Mol. Sci., 9 (2019), p. e1412.
- [15] A. FROMMER, M. N. KHALIL, AND G. RAMIREZ-HIDALGO, A multilevel approach to variance reduction in the stochastic estimation of the trace of a matrix, SIAM Journal on Scientific Computing, 44 (2022), pp. A2536–A2556.
- [16] M. B. GILES, Multilevel Monte Carlo path simulation, Oper. Res., 56 (2008), pp. 607–617.
- [17] M. B. GILES, Multilevel Monte Carlo methods, Acta Numer., 24 (2015), pp. 259–328.
- [18] E. HALLMAN AND D. TROESTER, A multilevel approach to stochastic trace estimation, Linear Algebra Appl., 638 (2022), pp. 125–149.
- [19] M. F. HERBST, A. LEVITT, AND E. CANCÈS, DFTK: A Julian approach for simulating electrons in solids, Proc. JuliaCon Conf., 3 (2021), p. 69, <https://doi.org/10.21105/jcon.00069>.
- [20] P. HOHENBERG AND W. KOHN, Inhomogeneous electron gas, Phys. Rev., 136 (1964), pp. B864–B871.
- [21] L. KLEINMAN AND D. M. BYLANDER, Efficacious form for model pseudopotentials, Phys. Rev. Lett., 48 (1982), pp. 1425–1428, <https://doi.org/10.1103/PhysRevLett.48.1425>.
- [22] W. KOHN, Density functional and density matrix method scaling linearly with the number of atoms, Phys. Rev. Lett., 76 (1996), pp. 3168–3171.
- [23] W. KOHN AND L. SHAM, Self-consistent equations including exchange and correlation effects, Phys. Rev., 140 (1965), pp. A1133–A1138.
- [24] Q. LIU AND M. CHEN, Plane-wave-based stochastic-deterministic density functional theory for extended systems, Phys. Rev. B, 106 (2022), p. 125132.
- [25] R. MARTIN, Electronic Structure: Basic Theory and Practical Methods, Cambridge University Press, 2004.
- [26] F. MAURI, G. GALLI, AND R. CAR, Orbital formulation for electronic-structure calculations with linear system-size scaling, Phys. Rev. B, 47 (1993), pp. 9973–9976.
- [27] R. MCWEENY, The density matrix in self-consistent field theory. I. Iterative construction of the density matrix., Proc. R. Soc. Lond. A, 235 (1956), pp. 496–509.
- [28] D. NEUHAUSER, R. BAER, AND E. RABANI, Communication: Embedded fragment stochastic density functional theory, J. Chem. Phys., 141 (2014), 041102.
- [29] M. NGUYEN, W. LI, Y. LI, R. BAER, E. RABANI, AND D. NEUHAUSER, Tempering stochastic density functional theory, J. Chem. Phys., 155 (2021), 204105.
- [30] J. OSBORN, Spectral approximation for compact operators, Math. Comp., 29 (1975), pp. 712–

- 725.
- [31] P. PERDEW AND Y. WANG, Accurate and simple analytic representation of the electron-gas correlation energy, Phys. Rev. B., 45 (1992), 13244.
 - [32] X. QUAN AND H. CHEN, SDFT.jl, <https://github.com/xuequan818/SDFT.jl>.
 - [33] C. ROOTHAAN, New developments in molecular orbital theory, Rev. Mod. Phys., 23 (1951), pp. 69–89.
 - [34] F. SHIMOJO, A. NAKANO, R. K. KALIA, AND P. VASHISHTA, Electronic processes in fast thermite chemical reactions: A first-principles molecular dynamics study, Phys. Rev. E, 77 (2008), p. 066103.
 - [35] L. N. TREFETHEN, Approximation Theory and Approximation Practice, Extended Edition, SIAM, Philadelphia, 2013.
 - [36] E. ULLMANN AND I. PAPAIOANNOU, Multilevel estimation of rare events, SIAM/ASA J. Uncertain. Quantif., 3 (2015), pp. 922–953.
 - [37] A. J. WHITE AND L. A. COLLINS, Fast and universal kohn-sham density functional theory algorithm for warm dense matter to hot dense plasma, Phys. Rev. Lett., 125 (2020), 055002.
 - [38] W. YANG, Direct calculation of electron density in density-functional theory, Phys. Rev. Lett., 66 (1991), pp. 1438–1441.