

MONOCHROMATIC PRODUCTS IN RANDOM INTEGER SETS

ROGER LIDÓN, DARÍO MARTÍNEZ, PATRICK MORRIS, AND MIQUEL ORTEGA

ABSTRACT. A well-known consequence of Schur's theorem is that for $r \in \mathbb{N}$, if n is sufficiently large, then any r -colouring of $[n]$ results in monochromatic $a, b, c \in [n]$ such that $ab = c$. In this paper we are interested in the threshold at which the binomial random set $[n]_p$ almost surely inherits this Ramsey-type property. In particular for $r = 2$ colours, we show that this threshold lies between $n^{-1/9-o(1)}$ and $n^{-1/11}$. Whilst analogous questions for solutions to (sets of) linear equations are now well understood, our work suggests that both the behaviour of the thresholds and the proof methods needed to determine them differ substantially in the non-linear setting.

1. INTRODUCTION

A collection of (ordered) subsets $\mathcal{S} \subset \mathbb{N}$ is *partition-regular*¹ if for any $r \in \mathbb{N}$, any r -colouring of $[n] := \{1, 2, \dots, n\}$ with $n \in \mathbb{N}$ is sufficiently large, results in a monochromatic set $S \in \mathcal{S}$. Determining which arithmetic configurations lead to collections of sets \mathcal{S} that are partition-regular is a major theme in modern combinatorics. The earliest result in this area of arithmetic Ramsey theory (and indeed in Ramsey theory as a whole) is due to Schur [30] over a century ago, who showed that the collection of all *sums* $\{(a, b, c) \in \mathbb{N}^3 : a + b = c\}$ is partition-regular. In 1933, Rado [23] proved a far-reaching generalisation of Schur's result, characterising for which matrices A the system of linear homogeneous equations $A\mathbf{x} = 0$ has a solution set that is partition-regular. This celebrated result in particular implies both Schur's theorem and the famous theorem of van der Waerden [32] showing that for any r , the collection of r -term arithmetic progressions $\{(a, a + d, \dots, a + (r - 1)d) : a, d \in \mathbb{N}\}$ is partition-regular. Further work completely settled partition-regularity for collections of sets defined by finite sets of equations, see e.g. the survey of Hindman [20].

1.1. Random sets of integers. In the 1990s, researchers started to strengthen statements showing partition-regularity by exploring the Ramsey properties of *sparse* host sets. In the context of monochromatic sums, we define a set $A \subseteq \mathbb{N}$ to be *r -Schur* if any r -colouring of A results in monochromatic $(a, b, c) \in A^3$ with $a + b = c$. When $r = 2$, we simply refer to the set A as being *Schur*. Whilst Schur's theorem [30] guarantees that $[n]$ is Schur for all n sufficiently large, it turns out that much sparser subsets of $[n]$ are typically also Schur. This was shown in pioneering work of Graham, Rödl and Ruciński [16]. The set $[n]_p$ denotes the binomial random subset of $[n]$ obtained by keeping each integer in $[n]$ with probability $p = p(n)$ independently and we say that a property $\mathcal{A} \subseteq 2^{[n]}$ holds in $[n]_p$ asymptotically almost surely (a.a.s. for short) if the probability that it holds tends to 1 as n tends to infinity.

Theorem 1.1 ([16]). *We have that*

$$\lim_{n \rightarrow \infty} \Pr([n]_p \text{ is Schur}) = \begin{cases} 0 & \text{if } p = o(n^{-1/2}); \\ 1 & \text{if } p = \omega(n^{-1/2}). \end{cases} \quad (0)$$

$$(1)$$

Date: December 5, 2025.

P. Morris was supported by the European Union's Horizon Europe Marie Skłodowska-Curie grant RAND-COMB-DESIGN - project number 101106032 [16]. M. Ortega was supported by the project PID2023-147202NB-I00, funded by MICIU/AEI/10.13039/501100011033/, as well as the FPI grant PRE2021-099120.

¹This definition is often stated in an infinitary fashion, asking for any r -colouring of \mathbb{N} to give monochromatic $S \in \mathcal{S}$. When every member $S \in \mathcal{S}$ is finite, these two formulations are equivalent by the so-called *compactness principle*.

For any property $\mathcal{A} \subseteq 2^{[n]}$, a function $p_* = p_*(n)$ is the² *threshold* for \mathcal{A} if whenever $p = \omega(p_*)$, we have that $[n]_p$ a.a.s. has the property \mathcal{A} whilst whenever $p = o(p_*)$, we have that $[n]_p$ a.a.s. does *not* have the property \mathcal{A} . Thus Theorem 1.1 establishes that $n^{-1/2}$ is the threshold for the Schur property. Further work established a sparse random analogue of Rado's theorem [23], determining the threshold for $[n]_p$ to contain monochromatic solutions to sets of homogeneous equations $A\mathbf{x} = 0$ in any colouring, for any A such that the solution set is partition-regular. The (0)-statements for these thresholds as well as (1)-statements for a subclass of so-called *density-regular* matrices A were proven by Rödl and Ruciński [25] who conjectured that their (0)-statements provided the correct threshold in all cases. This took over a decade to verify with Friedgut, Rödl and Schacht [15] finally providing a proof for all (1)-statements. The (1)-statements for all these Rado-type properties in fact hold for any bounded number of colours r and, perhaps surprisingly, the thresholds do not depend on the number of colours r . In particular, it follows from [15] that for *any* $r \geq 2$, the threshold for $[n]_p$ to be r -Schur is at $n^{1/2}$.

Shortly after the breakthrough work of Friedgut, Rödl and Schacht [15] providing (1)-statements for the random Rado theorem, Conlon and Gowers [9] and independently Schacht [27] developed powerful *transference principles* which establish optimal (1)-statements for a wide range of extremal and Ramsey properties in random discrete structures. In particular, these provided new proofs for some of (1)-statements for Rado-type problems discussed above. By adopting an abstract viewpoint of independent sets in hypergraphs, Balogh, Morris and Samotij [7] and independently Saxton and Thomason [26] then developed the theory of *hypergraph containers* which encapsulated many of the applications of the previous transference principles and led to an extremely flexible and elegant tool for establishing (1)-statements in random discrete structures. In particular, Nenadov and Steger [22] showed that one can use hypergraph containers to establish thresholds for Ramsey properties. This has led to a golden age in random Ramsey theory with many recent advances. Restricting only to the arithmetic setting here, we mention thresholds for solutions with repeated entries to Rado-type equations [31], resilience versions of Rado thresholds [17], perturbed Schur thresholds [2, 10], asymmetric random Rado theorems [3, 18], sharp thresholds for Schur and van der Waerden [13, 14, 29], thresholds for Rado type properties in random subsets of the primes and abelian groups [12] and canonical van der Waerden thresholds [4].

1.2. Monochromatic products. All of the random arithmetic Ramsey results mentioned above concern solutions to linear equations. In this paper, we make a first foray into exploring non-linear equations by considering monochromatic products. The fact that the set³ $\{(a, b, c) \in ([n] \setminus \{1\})^3 : ab = c\}$ is partition-regular can be derived from Schur's theorem for sums by considering powers of 2. In general, we say a set $A \subseteq \mathbb{N}$ is r -product-Schur if any r -colouring of S results in monochromatic $(a, b, c) \in A^3$ with $ab = c$ and when $r = 2$, we will just say that S is product-Schur. As the property of being product-Schur is monotone increasing⁴, the Bollobás-Thomason theorem [8] implies that there exists a threshold for the property. Our main theorem gives bounds on this threshold.

Theorem 1.2. *We have that*

$$\lim_{n \rightarrow \infty} \Pr([n]_p \text{ is product-Schur}) = \begin{cases} 0 & \text{if } p \leq n^{-1/9-o(1)}; \\ 1 & \text{if } p \geq \omega(n^{-1/11}). \end{cases} \quad (0) \quad (1)$$

Remark 1.3. We remark that in our definition of being product-Schur we allow for *degenerate* products $a^2 = b$. Our (1)-statement in Theorem 1.2 works in the slightly stronger setting of forcing non-degenerate monochromatic products. That is, if $p = \omega(n^{-1/11})$ then a.a.s. any 2-colouring of $[n]_p$ will result in monochromatic $a, b, c \in [n]_p$ with $ab = c$ and $a \neq b$.

²This is a slight abuse of notation as thresholds are not unique but rather determined up to constant factors.

³We omit the integer 1 here to avoid trivialities.

⁴That is, if A is product-Schur and $A \subseteq B$ then B is also product-Schur.

Theorem 1.2 makes a first step towards answering a recent question of Mattos, Mergoni Cecchelli and Parczyk [21]. Their motivation came from a question of Prendiville at the 2022 British Combinatorial Conference (BCC), which asked for another strengthening of Schur’s theorem to be generalised to the setting of products. Answering a \$100 question of Graham [16], Robertson and Zeilberger [24] and independently Schoen [28] proved that any 2-colouring of $[n]$ gives $(\frac{1}{11} + o(1))n^2$ monochromatic sums and that this is tight (their results are listed with the constant $1/22$ as they consider (a, b, c) and (b, a, c) as the same solution). For more colours, determining asymptotically the minimum number of monochromatic sums is an intriguing open question. Prendiville asked how many monochromatic products are necessarily guaranteed when colouring $[n] \setminus \{1\}$. This was then addressed by Aragão, Chapman, Ortega and Souza [6] and independently by Mattos, Mergoni Cecchelli and Parczyk [21]. For two colours, the first group [6] determined the minimum number of products precisely, showing that it is $(\frac{1}{2\sqrt{2}} + o(1))n^{1/2} \log n$. The second group [21] showed a lower bound of $n^{1/3 - o(1)}$ for this problem and also initiated the study of other variations of Schur’s theorem in the product setting. In particular, they determined precisely the threshold for the appearance of a product in the random set $[n]_p$ and asked the question of determining the threshold for the random set to be product-Schur.

The collection of products in $[n] \setminus \{1\}$ behaves very differently to the collection of sums. Indeed there are $\Theta(n \log n)$ products which is much less than the $\Theta(n^2)$ sums in $[n]$, and the collection is less regular, with the number of products involving an integer $a \in [n] \setminus \{1\}$ depending heavily on the location of a in the interval. In the Ramsey setting, the so-called *Schur multiplicity* results discussed above also show a difference in behaviour between the sum and product case. Indeed for sums when n is large we have that a positive fraction of all sums are monochromatic for any 2-colouring of $[n]$, an example of a phenomenon known as *supersaturation*. For products on the other hand, there are colourings in which only a vanishing fraction of the $\Theta(n \log n)$ products are monochromatic. Given this, it is perhaps not surprising that the proof of our (1)-statement for Theorem 1.2 differs from existing proofs in random Ramsey theory. Indeed, the transference principles [9, 27] and the hypergraph container method [7, 26] discussed above crucially rely on supersaturation and thus seem to be powerless in the setting of monochromatic products. To prove our (1)-statement, we instead appeal to a certain parameterised family of small configurations which have the product-Schur property and show that a.a.s. one of these configurations will appear in our random set $[n]_p$. This is reminiscent of the proof of the (1)-statement for Schur’s theorem in the randomly perturbed model [2, 10] when starting with a dense set of integers. The configurations we use are also similar (in fact contain) those used in [6] to count monochromatic solutions. Known proof methods establishing (0)-statements for random Ramsey results also seem ineffective in the product setting and we establish the (0)-statement of Theorem 1.2 through a careful greedy colouring procedure.

1.3. More colours. We also give initial bounds on thresholds for the random set to be r -product-Schur for $r \geq 3$. In particular, these bounds show that, unlike all of the thresholds [15] for solutions to sets of linear equations as in Rado’s theorem, for monochromatic products, the threshold depends on the number of colours. To state our results, we introduce some definitions. The *Schur number* $S(r)$ is the minimum n such that any r -colouring of $[n]$ results in a monochromatic sum. The *double-sum Schur number* $S'(r)$ is the minimum n such that any r -colouring of $[n]$ results in either a monochromatic sum $(a, b, c) \in [n]^3$ with $a + b = c$ or a monochromatic *shifted sum* $(a, b, c) \in [n]^3$ with $a + b = c - 1$. Double-sum Schur numbers were introduced by Abbott and Hanson [1] and also feature in the work of Mattos, Mergoni Cecchelli and Parczyk [21] bounding the size of the largest r -product-Schur subset of $[n]$.

Theorem 1.4. *For any $2 \leq r \in \mathbb{N}$ we have that*

$$\lim_{n \rightarrow \infty} \Pr([n]_p \text{ is } r\text{-product-Schur}) = \begin{cases} 0 & \text{if } p = o(n^{-1/S'(r)}); \\ 1 & \text{if } p = \omega(n^{-1/S(r)^2}). \end{cases} \quad (0)$$

$$(1)$$

One can check by hand or by computer that $S'(2) = S(2) = 5$ and $S'(3) = S(3) = 15$. The bounds given by Theorem 1.4 are thus weaker than those from Theorem 1.2 for two colours, whilst for three colours Theorem 1.4 shows that the threshold for being 3-product-Schur lies between $n^{-1/14}$ and $n^{-1/196}$, and in particular is larger than the threshold for being 2-product Schur. In fact, for a small number of colours we can do slightly better than the bound given by Theorem 1.4 (0), showing that the threshold for being 3-product Schur is at least $n^{-1/19}$, see Remark 4.2. In general, the values of $S'(r)$ and $S(r)$ may differ, for example $S'(4) = 41 < 45 = S(4)$ but we have that $3S(r-1) - 1 \leq S'(r) \leq S(r)$ as shown by Abbott and Hanson [1, Theorem 4.1]. Calculating (double-sum) Schur numbers is extremely challenging already for $r = 5$ [19] and the best general bounds give that $\Omega(3.17176^r) \leq S(r) \leq r!e$ with the lower bound due to Exoo [11] and the upper bound already shown by Schur himself [30]. Theorem 1.4 thus gives that if p_r^* is the threshold for $[n]_p$ being r -product Schur, then $n^{-1/c_1^r} \leq p_r^* \leq n^{-1/c_2^r \log r}$ for some constants $c_1, c_2 > 0$.

Organisation. We prove the (0)-statement of Theorem 1.2 in Section 2 and the (1)-statement in Section 3. We then prove Theorem 1.4 in Section 4 and give some concluding remarks in Section 5.

Notation. We write $[n] := \{1, \dots, n\}$ and $(m, n] := \{m+1, \dots, n\}$ for $m, n \in \mathbb{N}$. Given two sets $A, B \subset [n]$, we write $A \cdot B = \{ab : a \in A, b \in B\}$ for the set of pairwise products, and likewise for longer products. We let $d : \mathbb{N} \rightarrow \mathbb{N}$ count the number $d(i)$ of divisors of $i \in \mathbb{N}$, and write $D(n) = \max_{1 \leq i \leq n} d(i)$ for the maximum of this function in $[n]$. We say an event occurs asymptotically almost never if it a.a.s. does not occur. Finally, we omit floors and ceilings when they are not necessary and all logarithms are in the natural base.

2. THE (0)-STATEMENT FOR TWO COLOURS

Our goal in this section is to give a proof of the (0)-statement of Theorem 1.2. We first state a more precise version of the theorem. Define

$$\omega(n) := D(n)^{100},$$

The precise exponent of $D(n)$ here is not relevant, we just take it large enough to give us enough spare room in later bounds. The divisor bound tells us that $D(n) = n^{o(1)}$ (see, for example, Theorem 13.12 in [5]), so the (1)-statement is a consequence of the following.

Theorem 2.1. *If $p \leq n^{-1/9} \omega(n)^{-1}$, then a.a.s. we have that $[n]_p$ is product-Schur.*

In order to prove Proposition 2.1 we must show that, with high probability, we may partition our random set $S \sim [n]_p$ into two disjoint colours B, R so that $S = B \cup R$ and B, R are both product-free. We attempt to build such a colouring using the following greedy algorithm.

Algorithm 1 Colouring algorithm for two-coloured case

Input: $S \subset \mathbb{Z}, |S| < \infty$

```

1:  $B \leftarrow \emptyset$ 
2:  $R \leftarrow \emptyset$ 
3: for  $k \in S$  in increasing order do
4:   if  $k \notin (R \cdot R) \cup (R \cdot R \cdot R)$  then
5:      $R \leftarrow R \cup \{k\}$ 
6:   else if  $k \notin B \cdot B$  then
7:      $B \leftarrow B \cup \{k\}$ 
8:   else
9:     Algorithm fails
10:  end if
11: end for
```




Figure 1. Situation in which Algorithm 1 fails. The upper vertex of a triangle is the product of the lower ones, and z, w, f may be either red or one.

As long as the algorithm does not fail, we obtain such a colouring.

Observation 2.2. If Algorithm 1 terminates successfully on a set $S \subset [n]$, then S is not product-Schur.

Proof. Since we only add a new element to B or R if it is not the product of two smaller elements in the set, B and R will be product-free, and if the algorithm successfully terminates we also have that every integer in S receives a colour. \square

Hence, to prove Theorem 2.1 it suffices to see that the algorithm successfully terminates a.a.s.. We now look for possible causes for the failure of our algorithm, which turn out to be the following configurations of integers (see Figure 1 for a diagram of how such configurations look like).

Definition 2.3. We say a tuple $M = (a, b, c, d, e, f, x, y, z, u, v, w)$ is a *forbidden configuration* in S if it satisfies the following:

- (i) The equalities $c = ab = def$, $a = xyz$, and $b = uvw$ hold.
- (ii) We have that $a, b, c, d, e, x, y, u, v \in S$, and $f, z, w \in S \cup \{1\}$.
- (iii) a and b are different from $c, d, e, f, x, y, z, u, v, w$.
- (iv) If we denote $A = \{x, y, z, d, e, f, u, v, w\} \setminus \{1\}$, then A is disjoint to $A \cdot A$ and $A \cdot A \cdot A$.

The following lemma tells us that these are the only possible obstructions for the successful termination of the algorithm.

Lemma 2.4. If Algorithm 1 fails for a set S with $1 \notin S$, there exists a forbidden configuration in S .

Proof. Suppose the algorithm fails to colour $c \in S$. This means that c can be written both as $c = ab$ with $a, b \in B$, and as $c = def$ with $d, e \in R$ and $f \in R \cup \{1\}$. Notice too that $\{a, b\} \cap \{d, e, f\} = \emptyset$, since the algorithm guarantees that $B \cap R = \emptyset$, and by assumption $1 \notin B \subset S$.

As $a, b \in B$, the definition of the algorithm implies that $a, b \in R \cdot R \cup R \cdot R \cdot R$, so there exist $x, y, u, v \in R$, $z, w \in R \cup \{1\}$ such that $a = xyz$ and $b = uvw$. Again, each of these six numbers must be distinct from a and b because of their different colours.

We claim that the tuple $(a, b, c, d, e, f, x, y, z, u, v, w)$ is a forbidden configuration in S . In fact, we have already seen properties (i) to (iii) in Definition 2.3, and it only remains to check that condition (iv) holds. Since in Algorithm 1 we only add a new element to R if it is not the product of two or three smaller terms in R , we can guarantee that $R \cap (R \cdot R) = \emptyset$ and $R \cap (R \cdot R \cdot R) = \emptyset$. Writing $A = \{x, y, z, d, e, f, u, v, w\} \setminus \{1\}$ as in condition (iv), we have that $A \subset R$, so $A \cap (A \cdot A) = \emptyset$ and $A \cap (A \cdot A \cdot A) = \emptyset$. \square

2.1. Lack of forbidden configurations. The previous discussion reduces the proof of Theorem 2.1 to the following key proposition, to which we dedicate the rest of the section.

Proposition 2.5. *For $n \in \mathbb{N}$, a function $p = p(n) \leq n^{-1/9}\omega(n)^{-1}$, and $S \sim [n]_p$ we have that a.a.s. there is a forbidden configuration in S .*

Proof of Theorem 2.1 assuming Proposition 2.5. Run Algorithm 1 on $S \sim [n]_p$. The algorithm will terminate successfully a.a.s. on account of Lemma 2.4, Proposition 2.6 and the fact that $\Pr(1 \in S) = p = o(1)$, so the theorem follows from Observation 2.2. \square

In fact, we further reduce it to the following, which will allow us to worry only about forbidden configurations containing large enough elements.

Proposition 2.6. *For $n \in \mathbb{N}$, $p \leq n^{-1/9}\omega(n)^{-1}$, and $S \sim [n]_p$ we have that a.a.s. there is a forbidden configuration in $S \cap (n^{1/9}, n]$.*

Proof of Proposition 2.5 assuming Proposition 2.6. If there is a forbidden configuration in S that is not a forbidden configuration in $S \cap (n^{1/9}, n]$, S must contain a term that is smaller than $n^{1/9}$. However, by the union bound we have that

$$\Pr\left(S \cap [n^{1/9}] \neq \emptyset\right) \leq \sum_{s=0}^{n^{1/9}} \Pr(s \in S) \leq pn^{1/9} \leq \omega(n)^{-1} = o(1).$$

Therefore,

$$\begin{aligned} \Pr(\text{there is a forbidden configuration in } S) &\leq \Pr\left(S \cap [n^{1/9}] \neq \emptyset\right) + \\ &\quad \Pr\left(\text{there is a forbidden configuration in } S \cap (n^{1/9}, n]\right), \end{aligned}$$

and Proposition 2.5 follows from Proposition 2.6. \square

In order to prove Proposition 2.6 we will make repeated use of the union bound to control the appearance probability of different configurations. Concretely, given a family $\mathcal{A} \subset 2^{[n]}$ of uniform cardinality $r = |\mathcal{A}|$ for every $A \in \mathcal{A}$, and given a random set $S \sim [n]_p$, we will use that

$$\Pr(\exists A \in \mathcal{A} \text{ with } A \subset S) \leq \sum_{A \in \mathcal{A}} \Pr(A \subset S) \leq |\mathcal{A}|p^r. \quad (2)$$

Before going into the full details of the proof, let us sketch why the proposition is true. Suppose that all possible forbidden configurations in S satisfy $f = z = w = 1$ and that the nine numbers $a, b, c, d, e, x, y, u, v \in S$ are all distinct, which is in a sense the generic case. Since all of these nine numbers are divisors of c , the total number of configurations of this form is at most $nD(n)^9$. Hence, union bounding as in (2) gives that the probability of having such a configuration in S is at most $nD(n)^9p^9 = \omega(n)^{-9}D(n)^9 = o(1)$.

However, this is far from a proper proof. To begin with, the elements f, z, w may not be equal to one, but more importantly, the different terms in a forbidden configuration may not be distinct, giving worse bounds on the appearance of a fixed configuration. Therefore, in order to recover the union bound argument, it is necessary to prove that there are far fewer forbidden configurations of this kind. Hence, our strategy will be to split the possible forbidden configurations according to which variables are equal, and bound the total number of configurations of each kind, to be able to apply the union bound.

It is precisely because of this reason that we require $k \notin R \cdot R \cdot R$ in the first condition of Algorithm 1. If we define the more natural algorithm allowing $k \in R \cdot R \cdot R$, this leads to the worse threshold of $p \leq n^{-1/8}\omega(n)^{-1}$ for the algorithm to terminate successfully a.a.s. (in particular, there are too many

forbidden configurations of the form $x = d$, $z = w = f = 1$, and all other variables distinct, for union bound arguments to work).

From the previous discussion, we see that, when union bounding over a certain type of forbidden configuration, the following parameter will be crucial.

Definition 2.7. The *effective size* of a forbidden configuration M is the number of distinct elements different than 1 in M .

We now begin in earnest the proof of Proposition 2.6, which will need a fair amount of casework. From now on, let $M = (a, b, c, d, e, f, x, y, z, u, v, w)$ be a possible forbidden configuration satisfying the conditions in Definition 2.3. In particular, we know that

$$xyzuvw = def, \quad (3)$$

a key relation from which we will extract information in most cases we consider. In the setting of Proposition 2.6, we only consider forbidden configurations where $x, y, u, v, d, e \geq n^{1/9}$, $a, b, c \geq n^{2/9}$ and $f, z, w \geq n^{1/9}$ if they are not 1. Before going into the actual cases, we also isolate a simple lemma that is useful for counting the number of certain configurations.

Lemma 2.8. Let $e_1, \dots, e_k \in \mathbb{N}$ with $1 \leq e_1 \leq e_2 \leq \dots \leq e_k$ and $1 \leq t \leq n$. Let $e := \sum_{i=1}^k e_i$. Then the number of k -tuples (a_1, \dots, a_k) such that $t \leq a_i \leq n$ for all $2 \leq i \leq k$ such that $t \leq a_i \leq n$ for all $1 \leq i \leq k$ with $e_i \geq 2$, and $\prod_{i=1}^k a_i^{e_i} \leq n$ is at most

$$nt^{k-e}D(n)^k$$

Proof. As $\prod_{i=1}^k a_i^{e_i} \leq n$, and $a_i \geq t$ for each $i \in [k]$, any valid choice of (a_1, \dots, a_k) will have $\ell := \prod_{i=1}^k a_i \leq nt^{k-e}$. After choosing ℓ (at most nt^{k-e} choices) the a_i are chosen as divisors of ℓ (at most $D(n)$ choices for each $i \in [k]$). \square

In particular, we will apply it several times in the following form, which, in informal terms, tells us that what we lose by having repetitions in x, y, z, u, v, w is compensated by better bounds on the number of such configurations.

Corollary 2.9. Given a forbidden configuration with effective size s , define $r = r_1 - r_2$, where r_1 is the number of terms among x, y, z, u, v, w which are not 1, and $r_2 = |\{x, y, z, u, v, w\} \setminus \{1\}|$, so that r counts the number of repetitions among terms that are not one. Forbidden configurations such that $s + r \geq 9$ appear asymptotically almost never in $[n]_p \cap (n^{1/9}, n]$.

Proof. Consider fixed values of s, r, r_1, r_2 such that $s + r \geq 9$. Applying Lemma 2.8 to the condition $xyzuvw \leq n$ with $t = n^{1/9}$ implies there are at most $nn^{(r_2-r_1)/9}D(n)^{r_2} = n^{1-r/9}D(n)^{r_2}$ forbidden configurations with this value of s, r, r_1, r_2 . Union bounding as in (2) tells us that these configurations appear with probability at most $n^{1-r/9}D(n)^{r_2}p^s = o(1)$. Again union bounding over all constantly many possible values of s, r, r_1 and r_2 yields the result. \square

We now split forbidden configurations into different cases, where we use somewhat different approaches to count the number of forbidden configurations. Given a forbidden configuration, let s be its effective size, let m count how many numbers among d, e, f belong to $\{x, y, z, u, v, w\}$, and let m_d and m_f denote the number of terms among x, y, z, u, v, w equal to d and f respectively. In Table 1 we see a description of all possible cases we consider.

The only possibilities missing from the table, which are when $a \neq b$ and $d = f \neq e$ or $e = f \neq d$, are symmetric to the case where $a \neq b$ and $d = e \neq f$, with $f \neq 1$, and can be proved by relabeling the terms. As the naming suggests, we have divided these cases into three different groups. As we have already

| Conditions | | Case |
|--------------------------|-----------------------|------|
| $a = b$ | $s \geq 5$ | 1.1 |
| | $s = 4$ | 1.2 |
| | $s = 3$ | 1.3 |
| d, e, f distinct | $m = 0$ | 2.1 |
| | $m = 1$ | 2.2 |
| | $m = 2$ | 2.3 |
| | $m = 3$ | 2.4 |
| $a \neq b$ | $d = e = f$ | 3.1 |
| | $d = e, f = 1$ | 3.2 |
| $d = e \neq f, f \neq 1$ | $m_d = 0, m_f = 0$ | 3.3 |
| | $m_d = 0, m_f \geq 1$ | 3.4 |
| | $m_d = 1, m_f = 0$ | 3.5 |
| | $m_d = 1, m_f \geq 1$ | 3.6 |
| | $m_d \geq 2$ | 3.7 |

Table 1. Forbidden configuration cases

said, for every case we prove that configurations of that kind appear asymptotically almost never. Union bounding over all 14 cases proves Proposition 2.6.

2.1.1. Forbidden configurations of Group 1. All cases in this group satisfy that $a = b$. If $a = b$, then $c = ab = a^2$ must be a perfect square. Hence, there are only $n^{1/2}$ possible choices for c , and all other terms of the configuration divide c , so the total number of forbidden configurations of this kind is at most $n^{1/2}D(n)^{11}$. We split the configurations according to their effective size s , which must be at least 3 because a, c, d are distinct.

Case 1.1 ($s \geq 5$). If the effective size of the configuration is at least 5, a union bound as in (2) bounds the appearance probability for these configurations by $n^{1/2}D(n)^{11}p^5 = o(1)$.

Case 1.2 ($s = 4$). If the effective size is four, since $d, e, f, x, y, z, u, v, w$ are all distinct from a and c , they can only take values 1, g or h for some variables $g, h \geq n^{1/9}$. Plugging this into (3) gives a relation of the form $g^\ell h^r = g^i h^j$ for some i, j, ℓ, r with $\ell + r \in \{4, 5, 6\}$, $i + j \in \{2, 3\}$. Reordering, we have that $g^{\ell-i} = h^{j-r}$. As $i + j < \ell + r$, one of the two sides of the equation is not equal to one, so in fact both sides cannot be equal to one. This implies that g and h are powers of a common factor, and then a itself must be a perfect power. Once we have that a is a perfect power, then $c = a^2$ must be a perfect power of exponent at least 4. Hence, we have $O(n^{1/4})$ possible choices for c (accounting for powers of exponent 4 and greater) and configurations of this family appear at most $O(n^{1/4}D(n)^{11})$ times in $[n]$. A union bound gives that the appearance probability is at most $O(n^{1/4}D(n)^{11}p^4) = o(1)$.

Case 1.3 ($s = 3$). Finally, if the effective size of one such configuration is three, then a and b must be squares or cubes themselves, implying that configurations of this family appear at most $O(n^{1/4}D(n)^{11})$ in $[n]$. Now a union bound is enough to finish.

2.1.2. Forbidden configurations of Group 2. In this group we study forbidden configurations satisfying

$$a \neq b \quad d, e, f \text{ distinct.}$$

Recall that m counts how many numbers among d, e, f belong to $\{x, y, z, u, v, w\} \setminus \{1\}$, and let r be as in Corollary 2.9. Since d, e, f are distinct and different to a, b (on account of their colours), we have that $s + r + m$ is precisely the number of non-one terms in the configuration. We split the configurations according to the value of m .

Case 2.1 ($m = 0$). Since the number of non-one terms in the configuration is at least 9 and $m = 0$, we have that $s + r = s + r + m \geq 9$, and we may apply Corollary 2.9.

Case 2.2 ($m = 1$). Suppose that $f = z = w = 1$. In that case, (3) reduces to $de = xyuv$, and the fact that $m = 1$ tells us that we can simplify a factor on each side of the equality, contradicting condition (iv) in Definition 2.3.

It follows that if $m = 1$ then f, z, w cannot be all equal to one. Therefore, the number of non-one terms in the forbidden configuration is at least 10, so $s + r + 1 = s + r + m \geq 10$, so $s + r \geq 9$ and again we conclude by applying Corollary 2.9.

Case 2.3 ($m = 2$). Suppose that $f = 1$. Then (3) becomes $de = xyzuvw$, and from $m = 2$ we may deduce that the product of four numbers among u, v, w, x, z, w is 1, a contradiction. Hence assume $f \neq 1$. The fact that $m = 2$ tells us that we can simplify two terms on each side of the equality $def = xyzuvw$. The resulting equality will contradict condition (iv) in Definition 2.3 unless z and w are both not equal to one. In that case, the total number of non-one terms in the configuration is 12, so $s + r \geq 12 - m \geq 9$ and we may apply Corollary 2.9.

Case 2.4 ($m = 3$). This case is actually void, as $def = xyzuvw$ cannot hold if $d, e, f \in \{x, y, z, u, v, w\}$, because it would imply that a product of three numbers among x, y, z, u, v, w is equal to one.

2.1.3. *Forbidden configurations of Group 3.* Finally, it only remains to prove that forbidden configurations where

$$a \neq b \quad d, e, f \text{ not distinct}$$

occur asymptotically almost never. We begin by proving the first two cases.

Case 3.1 ($d = e = f$). In this case, $c = def = d^3$ is a cube. Hence, there are at most $n^{1/3}D(n)^{11}$ possible configurations of this form. We also know they have effective size at least four, because a, b, c, d are all distinct. Therefore, they appear with probability at most $n^{1/3}D(n)^{11}p^4 = o(1)$.

Case 3.2 ($d = e, f = 1$). In this case, $c = def = d^2$ is a perfect square. Hence, there are at most $n^{1/2}D(n)^{11}$ configurations of this form in $[n]$. By a union bound, the appearance probability of such a configuration of effective size at least 5 is at most $n^{1/2}D(n)^{11}p^5 = o(1)$. As a, b, c, d are all different, effective size 4 is the only remaining possibility. However, in such a case, a and b must both be equal to either d^2 or d^3 , contradicting $ab = c = d^2$. Therefore, there are no configurations of this form with effective size 4.

In the remaining cases, we assume without loss of generality that $d = e \neq f$ and $f \neq 1$. Recall that m_d and m_f respectively denote the number of values among x, y, z, u, v, w equal to d and f . We split all these forbidden configurations into 5 groups based on the values of m_d and m_f . As usual, we prove that configurations in each case occur asymptotically almost never.

Case 3.3 ($m_d = 0, m_f = 0$). This case follows from Corollary 2.9. Indeed, adopting the notation there we have that $s = 5 + r_2 = 5 + r_1 - r$ and so $s + r \geq 5 + r_1 \geq 9$.

Case 3.4 ($m_d = 0, m_f \geq 1$). After possibly relabeling terms, we may assume that $x = f$. In that case, (3) becomes

$$fyzuvw = d^2f,$$

which implies that $yzuvw = d^2$. We split our analysis according to the size of d .

- If $d \geq n^{3/9}$, from $c = d^2 f$ and Lemma 2.8 we see that there are at most $n^{6/9} D(n)^2$ possible choices for c , and at most $n^{6/9} D(n)^{11}$ possible configurations of this kind. If the effective size is at least six, they appear with probability at most $n^{6/9} D(n)^{11} p^6 = o(1)$. Since a, b, c, d, f are distinct, the only remaining possibility is effective size 5, and in that case x, y, z, u, v, w are all equal to 1 or f , so $c = xyzuvw$ is a perfect power of exponent at least 4. Hence, there are at most $O(n^{1/4})$ choices for c , and union bounding again concludes.
- If $d < n^{3/9}$, we study the probability that the subconfiguration formed by (y, z, u, v, w, d, b) appears. By assumption, we have at most $n^{3/9}$ possible choices for d . Since y, z, u, v, w all divide d^2 , once we fix d there are at most $D(n)^5$ choices for these terms, and then $b = uvw$ is uniquely determined. Hence, there are at most $n^{3/9} D(n)^5$ possible subconfigurations in $(n^{1/9}, n]$, and every subconfiguration has effective size at least three, because b, d, u are distinct, so the probability that such a subconfiguration appears is bounded by $n^{3/9} D(n)^5 p^3 = o(1)$. This also bounds the probability of the whole configuration appearing.

Case 3.5 ($m_d = 1, m_f = 0$). Assume without loss of generality that $u = d$. Note that (3) gives

$$xyzvw = df = \frac{d^2 f}{d} \leq n^{8/9},$$

since $d \geq n^{1/9}$. Fixing some choice r'_1 to be the number of terms among x, y, z, v, w that are not equal to 1 and a choice of $r'_2 = |\{x, y, z, v, w\} \setminus \{1\}|$, we have from Lemma 2.8 (applied with n replaced by $n^{8/9}$) that there are at most $n^{8/9} n^{(r'_2 - r'_1)/9} D(n)^{r'_2}$ choices for x, y, z, v and w . Furthermore, note that once x, y, z, v, w are given, there are at most $D(n)^2$ ways to complete the configuration, since there will be at most $D(n)^2$ choices for d and f , and d, f determine c . The effective size of such a configuration is precisely $s = 5 + r'_2$ with the 5 accounting for a, b, c, d and f . Therefore the probability that a configuration with values r'_1 and r'_2 appears is at most $D(n)^{r'_2+2} n^{(8+r'_2-r'_1)/9} p^s = o(n^{3-r'_1/9}) = o(1)$ as $r'_1 \geq 3$. Union bounding over all choices of r'_1 and r'_2 thus a.a.s. rules out all configurations in this group.

Case 3.6 ($m_d = 1, m_f \geq 1$). In this case, (3) gives $d^2 f = xyzuvw$, and we know that a d and an f in the left hand side will cancel out. Thus d must be the product of two, three or four numbers different from 1 among x, y, z, u, v, w . The first two cases are impossible by condition (iv) in Definition 2.3, so d is the product of four such numbers. Therefore $d \geq n^{4/9}$ and $f \leq n/d^2 \leq n^{1/9}$, which cannot occur.

Case 3.7 ($m_d \geq 2$). In this case, (3) gives again $d^2 f = xyzuvw$. Since $m_d \geq 2$, we deduce that f must be the product of two, three or four numbers among x, y, z, u, v, w , with the other two being equal to d . Thus by condition (iv) in Definition 2.3, f must be the product of four such numbers, implying that $z, w \neq 1$ and $m_f = 0$. Now using that $z, w \neq 1$, we have that the r in Corollary 2.9 counts all of the repetitions among x, y, z, u, v, w . We then have that the effective size of the configuration is $s = 10 - r$ as a, b, c and f are all distinct and none of the terms x, y, z, u, v, w coincide with a, b, c or f (those that are not equal to d divide f and so cannot equal f). The conclusion thus follows from Corollary 2.9. \square

3. THE (1)-STATEMENT FOR TWO COLOURS

In this section, we prove the (1)-statement of Theorem 1.2. To do so, we find a certain small pattern that is product-Schur, and then show that a copy of this pattern appears in $[n]_p$ with high probability when p is large enough. We begin by proving that the desired pattern is product-Schur.

Lemma 3.1. *Given $a, b, c, d \in [n]$, the set*

$$S_{a,b,c,d} = \{a, b, ab, c, ac, abc, a^2bc, a^2b^2c, d, ad, bd, abd, a^2d, a^2bd\} \quad (4)$$

is product-Schur.

Proof. Suppose we have a colouring $S_{a,b,c,d} = B \cup R$ with no monochromatic products. Without loss of generality, assume that $a \in B$.

We claim that if $b \in B$, the subpattern

$$\{a, b, ab, c, ac, abc, a^2bc, a^2b^2c\},$$

must have a monochromatic product. To see this, we split depending on the colour of c :

- If $c \in B$, from the lack of monochromatic products we deduce that

$$a, b, c \in B \implies ab, ac \in R \implies (ab)(ac) = a^2bc \in B,$$

and then

$$a^2bc \in B, a \in B \implies (a^2bc)/a = abc \in R.$$

From $(abc)(ab) = a^2b^2c$ we deduce that $a^2b^2c \in B$, but then $b(a^2bc) = a^2b^2c$ is a solution with all terms belonging to B , a contradiction.

- If $c \in R$, we see that

$$a, b \in B \implies ab \in R,$$

which together with $c \in R$ gives that $(ab)c = abc \in B$. Therefore,

$$a, b, abc \in B \implies a(abc) = a^2bc \in R, (abc)/b = ac \in R,$$

a contradiction because $(ab)(ac) = a^2bc$ is a monochromatic product.

If $b \in R$, the subpattern

$$\{a, b, ab, d, ad, bd, abd, a^2d, a^2bd\}$$

must have a monochromatic product by an analogous reasoning, where fixing the colour of d forces the rest of the colouring, leading to a contradiction in both cases. \square

The (1)-statement of Theorem 1.2 is then a consequence of the following proposition.

Proposition 3.2. *For $p = \omega(n^{-1/9})$, the random set $[n]_p$ a.a.s. contains a subset of the form (4).*

Proof. For convenience, we will only consider sets $S_{a,b,c,d}$ for integers $a \in A, b \in B, c \in C, d \in D$, where

$$\begin{aligned} A &= [(1 - \delta)n^{1/11}, n^{1/11}], \\ B &= [(1 - \delta)n^{2/11}/2, n^{2/11}/2], \\ C &= [(1 - \delta)n^{5/11}, n^{5/11}], \\ D &= [(1 - \delta)n^{7/11}/3, n^{7/11}/3]. \end{aligned}$$

Here δ is any small enough fixed positive constant, say, $\delta = 10^{-3}$. These intervals are chosen so that the different possible terms of $S_{a,b,c,d}$ lie in disjoint intervals. Concretely, we have that

$$\begin{aligned} a \in A &= [(1 - \delta)n^{1/11}, n^{1/11}], & a^2b^2c \in I_{a^2b^2c} &:= [(1 - 5\delta)n^{11/11}/4, n^{11/11}/4], \\ b \in B &= [(1 - \delta)n^{2/11}/2, n^{2/11}/2], & d \in D &= [(1 - \delta)n^{7/11}/3, n^{7/11}/3], \\ ab \in I_{ab} &:= [(1 - 2\delta)n^{3/11}/2, n^{3/11}/2], & ad \in I_{ad} &:= [(1 - 2\delta)n^{8/11}/3, n^{8/11}/3], \\ c \in C &= [(1 - \delta)n^{5/11}, n^{5/11}], & bd \in I_{bd} &:= [(1 - 2\delta)n^{9/11}/6, n^{9/11}/6], \\ ac \in I_{ac} &:= [(1 - 2\delta)n^{6/11}, n^{6/11}], & abd \in I_{abd} &:= [(1 - 3\delta)n^{10/11}/6, n^{10/11}/6], \\ abc \in I_{abc} &:= [(1 - 3\delta)n^{8/11}/2, n^{8/11}/2], & a^2d \in I_{a^2d} &:= [(1 - 3\delta)n^{9/11}/3, n^{9/11}/3], \\ a^2bc \in I_{a^2bc} &:= [(1 - 4\delta)n^{9/11}/2, n^{9/11}/2], & a^2bd \in I_{a^2bd} &:= [(1 - 4\delta)n^{11/11}/6, n^{11/11}/6]. \end{aligned}$$

which are disjoint intervals for any large enough n .

We now prove that if $p = \omega(n^{-1/11})$, we can find at least one instance of $S_{a,b,c,d}$ in $[n]_p$. We will do this by revealing $[n]_p$ step by step, and choosing one of a, b, c, d at each step.

- In the first step, we reveal A . By independence, we have that

$$\Pr(A \cap [n]_p = \emptyset) = \prod_{i \in A} \Pr(i \notin [n]_p) = (1-p)^{\delta n^{1/11}} \leq e^{-\delta p n^{1/11}} = o(1).$$

If $[n]_p \cap A \neq \emptyset$, we set a to be any element in $[n]_p \cap A$ and proceed to the second step.

- In the second step, we reveal B and I_{ab} , which are independent from A by disjointness. With an analogous reasoning, we see that

$$\Pr(\nexists i \in B \cap [n]_p \text{ with } ai \in [n]_p) = \prod_{i \in B} \Pr(i \notin B \text{ or } ai \notin [n]_p) \leq e^{-\delta p^2 n^{2/11}} = o(1).$$

If there exists such an i , we set b to be equal to it and carry out the third step.

- In the third step, we reveal $C, I_{ac}, I_{abc}, I_{a^2bc}$ and $I_{a^2b^2c}$. Again, we have

$$\Pr(\nexists i \in C \cap [n]_p \text{ with } ai, abi, a^2bi, a^2b^2i \in [n]_p) \leq e^{-\delta p^5 n^{5/11}} = o(1),$$

and we let c be such an i if it exists. If there is one, we move to the final step.

- In the last step, we reveal $D, I_{ad}, I_{bd}, I_{abd}, I_{a^2d}$ and I_{a^2bd} . Again,

$$\Pr(\nexists i \in D \cap [n]_p \text{ with } ai, bi, abi, a^2i, a^2bi \in [n]_p) \leq e^{-\delta p^6 n^{7/11}} = o(1),$$

and let d be such an i if it exists.

Union bounding over the complement, we see that the probability that we may not find such a, b, c, d is $o(1)$. Therefore, a.a.s. there exists a set of the form $S_{a,b,c,d}$ in $[n]_p$ as required. \square

4. MORE COLOURS

We now look at an arbitrary number of colours $r \geq 2$. Recall the (0)-statement in Theorem 1.4.

Theorem 4.1. *For $p = o(n^{-1/S'(r)})$ the set $[n]_p$ is r -product-Schur asymptotically almost never.*

Proof. To begin with, note that

$$\Pr([n]_p \cap [n^{1/S'(r)}] = \emptyset) = (1-p)^{n^{1/S'(r)}} \geq e^{-2pn^{1/S'(r)}} = 1 - o(1).$$

Therefore,

$$\Pr([n]_p \cap [n^{1/S'(r)}] \neq \emptyset) = o(1).$$

Now let $\psi: [S(r) - 1] \rightarrow [r]$ be some r -colouring of $[S'(r) - 1]$ avoiding monochromatic sums and monochromatic shifted sums (solutions to $a + b = c - 1$). We colour the intervals of the form $I_i = (n^{i/S'(r)}, n^{(i+1)/S'(r)})$ for $i \in [S'(r) - 1]$ according to ψ . More concretely, we colour all numbers in $I_i \cap [n]_p$ with the colour $\psi(i)$. Since a.a.s. all numbers in $[n]_p$ belong to one of these intervals, all numbers end up coloured a.a.s..

Finally, we observe that, if $a \in I_\alpha$ and $b \in I_\beta$, then $ab \in I_{\alpha+\beta} \cup I_{\alpha+\beta+1}$. Therefore, if a, b, ab have the same colour, then $\psi(\alpha) = \psi(\beta)$ and $\psi(\alpha), \psi(\beta) \in \{\psi(\alpha+\beta), \psi(\alpha+\beta+1)\}$, which is impossible by the definition of ψ . This proves that our colouring contains no monochromatic solutions to $xy = z$ in $[n]_p$. We conclude by noting that this is a full colouring of $[n]_p$ a.a.s.. \square

Remark 4.2. For small values of r we can in fact slightly improve the bound on the threshold with a variation of the idea. Notice that colouring I_1 and I_2 with the same colour does not generate monochromatic tuples, since a.a.s. no monochromatic tuples exist in $[p^{-3+o(1)}]_p$ by a union bound argument.

This trick allows for colourings ψ that avoid all triples of the forms $a + b = c$ and $a + b + 1 = c$ except $1 + 1 = 2$. Computer calculations show that with 3 colours we can colour up to⁵ 18 (while $S'(3) = 13$) and with four colours up to⁶ 54 (when $S'(4) = 40$). This improves the bounds for $r \in \{3, 4\}$ to $n^{-1/19+o(1)}$ and $n^{-1/55+o(1)}$, respectively. However, we have not found a better general construction using this idea.

Finally, we prove the (1)-statement in Theorem 1.4. The idea of the proof is the same we used for the 1-statement of the two colour case, but with a more general pattern that applies to all r . We recall the statement.

Theorem 4.3. *For $p = \omega(n^{-1/S(r)^2})$, the set $[n]_p$ is r -product-Schur a.a.s..*

Proof. Consider the pattern

$$U_x = \{x, x^2, \dots, x^{S(r)}\}$$

for $2 \leq x \leq n^{1/S(r)}$. Note that U_x is not r -product-colourable by the definition of $S(r)$. We now show that, for our range of p , a.a.s. there is $U_x \subset [n]_p$ for some $x \in [n]_p$.

Consider the family of sets $\{U_x \mid x \in X\}$ for the set $X \subset [n^{1/S(r)}]$ built greedily in the following manner. Iterate through values of $x \in [n^{1/S(r)}]$ and add U_x if it is disjoint from all other U_y for y already belonging to the set. For any given $x \in [n^{1/S(r)}]$, there are at most $S(r)^2$ values $y \in [n^{1/S(r)}]$ such that $U_x \cap U_y \neq \emptyset$, so X has size at least $\frac{1}{S(r)^2} n^{1/S(r)}$.

Note that for any given x we have $\mathbb{P}(U_x \subset [n]_p) = p^{S(r)}$, and that the events $U_x \subset [n]_p$ for $x \in X$ are mutually independent on account of the sets U_x being disjoint. Hence, the probability that $U_x \not\subset [n]_p$ for every $x \in X$ satisfies

$$\begin{aligned} \mathbb{P}\left(\bigwedge_{x \in X} U_x \not\subset [n]_p\right) &= \left(1 - p^{S(r)}\right)^{|X|} \\ &\leq \left(1 - p^{S(r)}\right)^{\frac{1}{S(r)^2} n^{1/S(r)}} \\ &\leq \exp\left(-p^{S(r)} \frac{1}{S(r)^2} n^{1/S(r)}\right) = o(1). \end{aligned}$$

Hence, $[n]_p$ contains U_x for some $x \in [n]_p$ a.a.s., and such a pattern is r -product-Schur, so $[n]_p$ also is. \square

5. CONCLUDING REMARKS

In this paper, we showed that the threshold for the random set $[n]_p$ lies between $n^{-1/9-o(1)}$ and $n^{-1/11}$. We believe that our upper bound provides the true location of the threshold.

Conjecture 5.1. *The threshold for $[n]_p$ to be product-Schur lies at $n^{-1/11}$.*

This conjecture is motivated by two sub-conjectures, each of which could provide natural steps towards its resolution. Firstly, we believe that the threshold is determined by the appearance in $[n]_p$ of some bounded size configuration which is itself product-Schur. Secondly, we believe that the configuration used in our proof of Theorem 1.2 (1) is optimal, that is, there is no other bounded size configuration that is product-Schur and appears at a density less than $p = n^{-1/11}$. As a first step towards Conjecture 5.1, it would be interesting to improve on the lower bound of $n^{-1/9-o(1)}$. We believe that it might be possible to slightly improve the lower bound using a more involved greedy colouring scheme, but getting a significant improvement on the lower bound would probably require new ideas.

For the r -product-Schur problem with more colours, we showed that the threshold p_r^* lies between $n^{-S'(r)}$ and $n^{-S(r)^2}$ where $S(r)$ is the Schur number and $S'(r)$ is the double-sum Schur number. We

⁵The construction here is AABBBACCCACCCABBBA.

⁶The construction here is AABBBACCCACCCABBADDDADDDABBBADDDADDDABBBACCCACCCABBBA.

remark that one could also try to use the idea for our 2-colour (0)-statement to get a lower bound in the r -colour case. Indeed, the idea there was to take a greedy colouring and show that it a.a.s. does not fail to colour all the integers. One can show by induction that there are 3^r integers in a (non-degenerate) configuration which witnesses the failure of the algorithm and so one cannot expect a better bound than $p_r^* \geq n^{-1/3^r}$ using this approach. As $S'(r) \geq 3S(r-1) - 1 = \Omega(3.17176^r)$ [1, 11], this does worse than the lower bound we give here for r sufficiently large.

It would be very interesting to close the gap on our knowledge of $\log_n 1/p_r^*$ which we know is between exponential and factorial in r . Given that this is also the limit of our knowledge on the growth of Schur numbers [11, 30], this seems like it could be a considerable challenge. Nonetheless, it may be possible to improve the upper and lower bounds of $\log_n 1/p_r^*$ in terms of the Schur number $S(r)$. Finally we remark that it would be natural to study other nonlinear equations in this random Ramsey setting and we believe our techniques developed here will be useful for this. In particular, some pertinent candidates are $a^2 = bc$ and $ab = a + b$.

Acknowledgements. Part of this work was done as part of the *Barcelona Introduction to Mathematical Research – (BIMR 2024)* and we would like to thank BGSMath and Centre de Recerca Matemàtica (CRM) for organising this summer program. We also thank Shagnik Das, Leticia Mattos and Juanjo Rué for interesting discussions around this topic.

REFERENCES

1. H. Abbott and D. Hanson, *A problem of Schur and its generalizations*, Acta Arithmetica **20** (1972), no. 2, 175–187.
2. E. Aigner-Horev and Y. Person, *Monochromatic Schur triples in randomly perturbed dense sets of integers*, SIAM Journal on Discrete Mathematics **33** (2019), no. 4, 2175–2180.
3. ———, *An asymmetric random Rado theorem: 1-statement*, Journal of Combinatorial Theory, Series A **193** (2023), 105687.
4. J. D. Alvarado, Y. Kohayakawa, P. Morris, G. O. Mota, and M. Ortega, *A canonical van der Waerden theorem in random sets*, Discrete Mathematics Days 2024 (2024), 148.
5. T. M. Apostol, *Introduction to analytic number theory*, 1976.
6. L. Aragão, J. Chapman, M. Ortega, and V. Souza, *On the number of monochromatic solutions to multiplicative equations*, Combinatorica **45** (2025), 64.
7. J. Balogh, R. Morris, and W. Samotij, *Independent sets in hypergraphs*, J. Amer. Math. Soc. **28** (2015), no. 3, 669–709.
8. B. Bollobás and A. Thomason, *Threshold functions*, Combinatorica **7** (1987), no. 1, 35–38.
9. D. Conlon and W. T. Gowers, *Combinatorial theorems in sparse random sets*, Ann. of Math. (2) **184** (2016), no. 2, 367–454.
10. S. Das, C. Knierim, and P. Morris, *Schur properties of randomly perturbed sets*, European Journal of Combinatorics **121** (2024), 103843.
11. G. Exoo, *A lower bound for Schur numbers and multicolor Ramsey numbers*, Electron. J. Comb. **1** (1994), R8.
12. A. Freschi, R. Hancock, and A. Treglown, *Typical Ramsey properties of the primes, abelian groups and other discrete structures*, arXiv preprint arXiv:2405.19113 (2024).
13. E. Friedgut, H. Hàn, Y. Person, and M. Schacht, *A sharp threshold for van der Waerden’s theorem in random subsets*, Discrete Analysis (2016), no. 7, 20 pp.
14. E. Friedgut, E. Kuperwasser, W. Samotij, and M. Schacht, *Sharp thresholds for Ramsey properties*, arXiv preprint arXiv:2207.13982 (2022).
15. E. Friedgut, V. Rödl, and M. Schacht, *Ramsey properties of random discrete structures*, Random Structures & Algorithms **37** (2010), no. 4, 407–436.
16. R. Graham, V. Rödl, and A. Ruciński, *On Schur properties of random subsets of integers*, Journal of Number Theory **61** (1996), no. 2, 388–408.
17. R. Hancock, K. Staden, and A. Treglown, *Independent sets in hypergraphs and Ramsey properties of graphs and the integers*, SIAM Journal on Discrete Mathematics **33** (2019), no. 1, 153–188.
18. R. Hancock and A. Treglown, *An asymmetric random Rado theorem for single equations: The 0-statement*, Random Structures & Algorithms **60** (2022), no. 4, 529–550.
19. M. Heule, *Schur number five*, Proceedings of the AAAI Conference on Artificial Intelligence, vol. 32, 2018.

20. N. Hindman, *Partition regularity of matrices*, Combinatorial Number Theory, De Gruyter Proceedings in Mathematics, Editors: B. Landman, M. Nathanson, J. Nešetřil, R. Nowakowski and C. Pomerance, De Gruyter, 2005, pp. 265–298.
21. L. Mattos, D. Mergoni Cecchelli, and O. Parczyk, *On product Schur triples in the integers*, SIAM Journal on Discrete Mathematics **39** (2025), no. 2, 1082–1095.
22. R. Nenadov and A. Steger, *A short proof of the random Ramsey theorem*, Combinatorics, Probability and Computing **25** (2016), no. 1, 130–144.
23. R. Rado, *Studien zur kombinatorik*, Mathematische Zeitschrift **36** (1933), no. 1, 424–470.
24. A. Robertson and D. Zeilberger, *A 2-coloring of $[1, n]$ can have $(1/22)n^2 + o(n)$ monochromatic Schur triples, but not less!*, The Electronic Journal of Combinatorics (1998), R19.
25. F. Rödl and A. Ruciński, *Rado partition theorem for random subsets of integers*, Proceedings of the London Mathematical Society **74** (1997), no. 3, 481–502.
26. D. Saxton and A. Thomason, *Hypergraph containers*, Invent. Math. **201** (2015), no. 3, 925–992.
27. M. Schacht, *Extremal results for random discrete structures*, Ann. of Math. (2) **184** (2016), no. 2, 333–365.
28. T. Schoen, *The number of monochromatic Schur triples*, European Journal of Combinatorics **20** (1999), no. 8, 855–866.
29. F. Schulenburg, *Threshold results for cycles*, Ph.D. thesis, University of Hamburg, 2016.
30. I. Schur, *Über kongruenz $x^m + y^m \equiv z^m \pmod{p}$* , Jahresbericht der Deutschen Mathematiker-Vereinigung **25** (1917), 114–116.
31. C. Spiegel, *A note on sparse supersaturation and extremal results for linear homogeneous systems*, The Electronic Journal of Combinatorics (2017), P3.38.
32. B. L. van der Waerden, *Beweis einer baudetschen vermutung*, Nieuw Arch. Wiskunde **15** (1927), 212–216.

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT POLITÈCNICA DE CATALUNYA (UPC), CARRER DE PAU GARGALLO 14, 08028 BARCELONA, SPAIN.

Email address: {rlidon2006 | dariogol.com | pmorrismaths | miquel.ortega9}@gmail.com